

H α fluxes and extinction distances for planetary nebulae in the IPHAS survey of the Northern Galactic Plane

Thavisha E. Dharmawardena,^{1,2,3,4} M. J. Barlow,² J. E. Drew,² A. Seales,^{2,5} S. E. Sale,⁶ D. Jones,^{7,8} A. Mampaso,^{7,8} Q. A. Parker,⁹ L. Sabin,¹⁰ R. Wesson²

¹Max Planck Institute for Astronomy, Königstuhl 17, 69117 Heidelberg, Germany.

²Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT, U.K.

³Academia Sinica Institute of Astronomy and Astrophysics, 11F ASMAX, No.1, Sec. 4, Roosevelt Rd, Taipei 10617, Taiwan.

⁴National Central University, No. 300, Zhongda Rd., Zhongli District, Taoyuan City 32001, Taiwan.

⁵The Royal Institution of Great Britain, 21 Albemarle Street, London W1S 4BS, U.K.

⁶Rudolf Peierls Centre for Theoretical Physics, Keble Road, Oxford OX1 3NP, U.K.

⁷Instituto de Astrofísica de Canarias, E-38205 La Laguna, Tenerife, Spain

⁸Departamento de Astrofísica, Universidad de La Laguna, E-38206 La Laguna, Tenerife, Spain

⁹Department of Physics, University of Hong Kong, CYM Physics Building, Hong Kong;

The Laboratory for Space Research, University of Hong Kong, Hong Kong

¹⁰Instituto de Astronomía, Universidad Nacional Autónoma de México, Apdo. Postal 877, C.P. 22860 Ensenada, B.C., México

Received:

ABSTRACT

We report H α filter photometry for 197 northern hemisphere planetary nebulae (PNe) obtained using imaging data from the IPHAS survey. H α +[N II] fluxes were measured for 46 confirmed or possible PNe discovered by the IPHAS survey and for 151 previously catalogued PNe that fell within the area of the northern Galactic Plane surveyed by IPHAS. After correcting for [N II] emission admitted by the IPHAS H α filter, the resulting H α fluxes were combined with published radio free-free fluxes and H β fluxes, in order to estimate mean optical extinctions to 143 PNe using ratios involving their integrated Balmer line fluxes and their extinction-free radio fluxes. Distances to the PNe were then estimated using three different 3D interstellar dust extinction mapping methods, including the IPHAS-based H-MEAD algorithm of Sale (2014). These methods were used to plot dust extinction versus distance relationships for the lines of sight to the PNe; the intercepts with the derived dust optical extinctions allowed distances to the PNe to be inferred. For 17 of the PNe in our sample reliable Gaia DR2 distances were available and these have been compared with the distances derived using three different extinction mapping algorithms as well as with distances from the nebular radius vs. H α surface brightness relation of Frew et al. (2016). That relation and the H-MEAD extinction mapping algorithm yielded the closest agreement with the Gaia DR2 distances.

Key words: (ISM:) planetary nebulae: general; planetary nebulae: individual; (ISM:) dust, extinction

1 INTRODUCTION

Accurate distance estimation to Galactic planetary nebulae has represented a persistent difficulty over the years, with many different methods, both direct and statistical, having been applied - see Frew et al. (2016) for a recent comprehensive summary of the various methods. Although Gaia has now measured parallaxes for a significant number of PN central stars (e.g. Kimeswenger & Barriá 2018; González-Santamaría et al. 2019), for many PNe high nebular surface brightnesses or faint central star magnitudes make reliable

Gaia parallaxes difficult to determine. For such nebulae alternative distance estimation methods must be sought.

The Galactic PN distance estimation method that will be discussed in this paper is based on the determination of interstellar dust extinctions and distances to a sufficient number of field stars nearby on the sky to a PN that a plot of the stellar dust extinctions versus distance allows the distance to the PN to be inferred from its measured dust extinction. Early examples of the use of this technique for PN distance estimation were published by Lutz (1973), Kaler & Lutz (1985) and Gathier et al. (1986). In recent

years a number of deep optical and infrared imaging surveys have enabled the production of extensive 3D reddening maps of the Galactic plane, allowing distances to be estimated to stars and nebulae by plotting reddening vs. distance for their sight lines (e.g. Marshall et al. 2006; Sale et al. 2009, 2014; Sale 2012; Lallement et al. 2014, 2019; Green et al. 2015, 2019).

The Isaac Newton Telescope Photometric H-alpha Survey of the northern Galactic plane (IPHAS; Drew et al. 2005; Barentsen et al. 2014), in addition to obtaining filter photometry in the Sloan r' and i' bands, has produced a large narrow-band $H\alpha$ photometric sample, enabling many emission line stars and nebulae to be identified and measured. Data from IPHAS have been used to identify new examples of young stellar objects (Vink et al. 2008), classical Be stars Raddi et al. (2015), supernova remnants (Sabin et al. 2013) and planetary nebulae (Mampaso et al. 2006; Corradi et al. 2011; Viironen et al. 2009a; Sabin et al. 2014).

From the initial IPHAS data release (González-Solares et al. 2008) to the second data release (Barentsen et al. 2014) the applications of IPHAS data have expanded. Sale et al. (2009), Sale (2012) and Sale et al. (2014) have demonstrated how the information from colours using the r' , i' and (in particular) $H\alpha$ filters enables spectral types and interstellar dust extinctions to be inferred for objects in the IPHAS survey, enabling 3D extinction mapping to be carried out for sight lines covered by the survey. For example, Giannanco et al. (2011) applied the IPHAS-based extinction mapping technique MEAD (Mapping Extinction Against Distance; Sale et al. 2009) to planetary nebulae, using spectroscopic Balmer-line ratios to estimate reddening and distances for 70 planetary nebulae.

In the past, dust extinctions to PNe have often been obtained from the ratio of radio free-free fluxes (unaffected by dust extinction) to $H\beta$ filter fluxes. In this paper we measure $H\alpha$ fluxes for a sample of 197 PNe in the northern Galactic Plane observed by IPHAS and use them, along with published integrated $H\beta$ fluxes that are available for 37 of the nebulae, to determine interstellar dust extinctions to the nebulae. In Section 2 of the paper we present $H\alpha$ flux measurements for 197 PNe observed by IPHAS and correct for the [N II] 6548, 6583 Å contributions to these fluxes. In Section 3 we combine the [N II]-corrected $H\alpha$ fluxes with published $H\beta$ and radio free-free fluxes in order to derive interstellar dust extinctions to 143 PNe. In Section 4 we derive extinction distances to these PNe using a number of 3D extinction mapping techniques and for 17 PNe in the sample that are judged to have reliable Gaia DR2 distances we compare these to the extinction mapping distances.

2 NEBULAR $H\alpha$ FLUXES FOR NORTHERN PLANETARY NEBULAE

2.1 The IPHAS survey

The IPHAS survey (Drew et al. 2005) was carried out using the Wide Field Camera (WFC) on the 2.5m Isaac Newton telescope located in the Observatorio del Roque de los Muchachos on the island of La Palma. It was a digital optical photometric survey of the northern Galactic plane, carried out in three bands, r' , i' and $H\alpha$. The IPHAS footprint extends over $-5^\circ < b < +5^\circ$ and $30^\circ < l < 215^\circ$, forming

an area of approximately 1800 square degrees (Drew et al. 2005; Barentsen et al. 2014).

The survey obtained images using the WFC's $H\alpha$ narrow-band filter with a full width at half maximum (FWHM) bandwidth of 95 Å centred at 6568 Å and broadband Sloan r' (bandwidth 1380 Å centred at 6260 Å) and i' (bandwidth 1535 Å centred at 7670 Å) filters. Images were taken sequentially with all three filters at each pointing, with exposure times of 120s for the $H\alpha$ filter and mostly 30s and 10s for the r' and i' filters, respectively. The WFC detector pixels projected to 0.333 arcsec per pixel on the sky.

Each WFC image is a mosaic from its four CCDs, capturing a sky area of approximately 0.25 square degrees. To fill in the gaps between the CCDs, a set of offset pointings with the same filters and exposure times was obtained after each prime pointing, resulting in 7635 pairs of telescope pointings being used to capture the total IPHAS survey area (Drew et al. 2005; Barentsen et al. 2014).

The second IPHAS Data Release (DR2) was made available in 2014 (Barentsen et al. 2014). At this time the survey had achieved 5σ magnitude limits of 21.2 ± 0.5 in the r' band, 20.0 ± 0.3 in the i' band and 20.3 ± 0.03 in the $H\alpha$ band and had covered $>90\%$ of the 1800 square degree footprint of the IPHAS survey.

2.2 Planetary nebula source selection

The IPHAS survey observed a large number of PNe and PN candidates. We have utilised the Hong Kong/AAO/Strasbourg $H\alpha$ database¹ (HASH; Parker et al. 2016) to select nebulae classified as true, likely or possible PNe from the following sources:

- (i) Viironen et al. (2009a): A list of candidate compact PNe from the IPHAS survey, of which we measured 4 that are listed in the HASH database as true or possible PNe.
- (ii) Sabin et al. (2014): A catalogue of IPHAS-discovered potential PNe based on the spectroscopic and morphological characteristics of the nebulae (for PN classification criteria see Frew & Parker 2010). We measured 42 of these objects that are listed as true, likely or possible PNe by the HASH database, including 5 objects from the bow-shock nebula discovery paper of Sabin et al. (2010).
- (iii) We measured 151 PNe from the Strasbourg-ESO Catalogue (Acker et al. 1992) that had been observed by IPHAS and which are listed as true PNe by the HASH database.

2.3 $H\alpha$ aperture photometry measurements of PNe observed by IPHAS

The reduced IPHAS $H\alpha$ images are available for download via the INT WFC Archive's Data Quality Control (DQC) query page, hosted by the Cambridge Astronomy Survey Unit (CASU)². All sources had multiple observations and all the observations for each source were downloaded, apart from those flagged as having problems. Column 3 of Tables A1 and A2 of Appendix A lists the estimated angular

¹ <http://202.189.117.101:8999/gpne/db MainPage.php>

² <http://apm3.ast.cam.ac.uk/cgi-bin/wfs/dqc.cgi>

dimensions of the nebulae. For nebulae less than ~ 5 arcsec in diameter this was measured as their FWHM in the IPHAS H α filter images, then corrected in quadrature by the FWHM of nearby stars of similar brightness. For larger nebulae, the quoted angular dimensions were measured at 10% of the peak nebular brightnesses in the IPHAS H α filter images.

In order to obtain integrated source counts, aperture photometry was carried out using the Caltech Aperture Photometry Tool (APT)³ and the Starlink Graphical Astronomy and Image Analysis Tool (GAIA)⁴. The shape (circular/elliptical) and size of the APT aperture used for each PN was adjusted to best match the shape of the PN, as was the annular sky aperture, and the median sky-subtracted net source counts measured. For a small number of PNe in crowded fields for which there were bright stars within the PN aperture, we aimed to include similarly bright stars in the annular sky aperture in order to remove to first order the contribution from the contaminants. We utilised GAIA for those nebulae whose angular radii were too large to fit within the APT's radius limit of 200 pixels. For a few very extended low surface brightness nebulae, the IPHAS 120s H α exposures might not detect the full extent of their emission.

The source counts per second, obtained by dividing the net source counts by the observation exposure times, were converted to instrumental magnitudes and then to filter magnitudes by adding the zero-point filter magnitudes provided for each WFCAM exposure.

The H α filter magnitudes were converted to in-filter fluxes using a zero-magnitude flux calibration for the H α filter of 1.57×10^{-7} ergs cm $^{-2}$ s $^{-1}$. This was obtained by multiplying the mean monochromatic flux for Vega in the H α filter of 1.81×10^{-9} ergs cm $^{-2}$ s $^{-1}$ Å $^{-1}$ (Barentsen et al. 2014) by an equivalent width of 84.04 Å for the H α filter that was obtained by integrating across the transmission profile of the filter as a function of wavelength. At the time of the IPHAS DR2 data release, Vega was adopted to have a magnitude of +0.035 at all optical wavelengths. Between DR2 and the most recent release (Greimel et al., in preparation), the calibration changes show a 1σ scatter of ± 0.03 magnitudes.

In Appendix A, our H α filter aperture photometry results are presented in column 5 of Table A1, for the IPHAS-discovered PNe, while column 5 of Table A2 presents our photometry for previously catalogued PNe observed by the IPHAS survey. A number of very bright PNe had saturated pixels in their H α images and were therefore omitted from our sample (e.g. NGC 7027, BD+30°3639, VY 2-2).

2.4 Correction of the H α filter fluxes for [N II] contributions

The [N II] λ6548.03 Å and λ6583.41 Å doublet is situated on either side of the H α λ6562.82 Å line. The INT-WFC H α filter has an effective wavelength of 6568 Å and a transmission

FWHM of 95 Å and its transmission at the wavelengths of the [N II] lines is the same as at H α to within a few percent.

To correct the H α filter fluxes for [N II] line contributions, spectroscopically measured [N II]/H α flux ratios for each PN were available from the literature for 39 of the 46 IPHAS-discovered PNe and for all 151 of the previously catalogued PNe. If the published spectroscopic data provided the flux for only one line of the [N II] doublet, we corrected for the contribution from both lines assuming an intrinsic 6583/6548 flux ratio of 3.0 (Storey & Zeippen 2000). Column 9 of Table A1 and column 6 of Table A2 list the ([N II] 6583+6548)/H α ratios adopted from the literature, and their sources, while the next column lists log H α line fluxes *after* removing the [N II] line contributions. For nebulae where the differences between individual flux measurements were less than 0.01 dex, we set the final uncertainty on the mean flux to be 0.01 dex.

Most spectroscopic [N II]/H α ratios are based on long-slit spectra. Since observed ratios can sometimes vary significantly with location in extended nebulae, we consider our adopted corrections to be more reliable for those nebulae having angular diameters of less than 5–6 arcsec.

Although the [N II] lines are the major source of contamination of the in-filter H α fluxes, the continuum from stars within the PN measurement aperture is another potential contaminant, although this is corrected to first order via the subtraction of the flux in the sky aperture. The underlying nebular continuum is another potential contributor, although for the case of the narrow-band WFC H α filter its contribution is estimated to be minor. Using the NEBCONT routine in the DIPSO package (Howarth et al. 2004) then, for a nebular electron density of 10^3 cm $^{-3}$ and 10^4 K with $n(\text{He}^+)/n(\text{H}^+) = 0.10$, one finds that for a H α flux towards the faint end of the H α flux distribution shown in Figure 1, i.e. 10^{-12} ergs cm $^{-2}$ s $^{-1}$, the in-filter nebular continuum flux in the 95-Å FWHM IPHAS H α filter would contribute 1.9×10^{-14} ergs cm $^{-2}$ s $^{-1}$, or 1.9 percent of the H α flux. The in-filter flux that would be contributed by a 17th-magnitude central star would be less than 1 percent of the same nebular H α flux.

2.5 Comparison with the H α fluxes of Frew et al. (2013)

Frew et al. (2013) presented a catalogue of H α fluxes for 1258 Galactic PNe obtained using data from the Southern H α Sky Survey Atlas (SHASSA) (Gaustad et al. 2001) and the Virginia Tech Spectral-line Survey (VTSS)(VTech 2014)⁵. The VTSS survey used an f/1.2 camera lens with a 58-mm focal length, together with a 512×512 CCD array that projected to 96 arcsec per pixel on the sky. Its integration times were not specified. Its H α filter had a FWHM of 17.5 Å, transmitting mainly the H α line. The SHASSA survey used an f/1.6 camera lens with a focal length of 52 mm with maximum exposure times of 1200 s onto a 1024×1024 CCD array that projected to 48 arcsec per pixel on the sky. Its H α filter had a FWHM of 32 Å and transmissions of 39%, 26% and 78% at the rest wavelengths of the [N II] 6548, 6583 Å and

³ <http://www.aperturephotometry.org/apttool/>

⁴ <http://www.starlink.rl.ac.uk/star/docs/sun214.htm/sun214.html/>

⁵ <http://www.phys.vt.edu/~halpha/>




Figure 1. IPHAS H α log fluxes (in cgs units) compared to the (a) SHASSA (left) and (b) VTSS (right) H α log fluxes of Frew et al. (2013), all after correction for [N II] contributions. The lines of best fit are shown as dashed lines, with slopes of 0.90 ± 0.03 and 0.84 ± 0.06 for the SHASSA and VTSS comparisons respectively.

H α lines, respectively (Gaustad et al. 2001). The transmissions of these lines were equal for the 95-Å FWHM H α filter used by IPHAS. Therefore for the same PN with the same [N II]/H α ratio, the [N II] contribution to the SHASSA H α filter flux measured by Frew et al. (2013) should be less than that made to the IPHAS WFC H α -filter flux measured here.

Figure 1(a) plots our measured IPHAS H α fluxes against the SHASSA H α fluxes measured by Frew et al. (2013) for 29 nebulae in common, both after [N II] corrections have been applied. The SHASSA fluxes are on average fainter than the IPHAS fluxes by 0.08 ± 0.12 dex, consistent with no net difference within the uncertainty limits. The slope of the line of best fit is 0.90 ± 0.03 .

A similar comparison of the [N II]-corrected IPHAS H α fluxes with the Frew et al. (2013) VTSS H α fluxes for 32 nebulae in common showed the VTSS H α fluxes to be 0.17 ± 0.19 dex brighter than the IPHAS H α fluxes, again consistent with no net difference within the uncertainty limits, with the larger dispersion attributable to the larger uncertainties associated with the VTSS fluxes, see Fig. 1(b).

3 NEBULAR EXTINCTIONS

3.1 Determining dust extinctions to the nebulae

Since the radio free-free fluxes of PNe undergo virtually no extinction by interstellar dust, we can estimate absolute extinctions to PNe at optical wavelengths by comparing their measured hydrogen Balmer line fluxes with Balmer line fluxes predicted from their measured radio free-free fluxes using recombination theory. We have therefore used our measured H α fluxes, as well as H β fluxes from the literature, together with literature radio free-free fluxes, to determine dust extinctions at the wavelengths of the two Balmer lines. The logarithmic extinctions at H β ($C_{H\beta}$) were calculated using the formulation given by Milne & Aller (1975):

$$C_{H\beta} = \log_{10} \frac{3.28 \times 10^{-9} \times S(5\text{GHz}) \times t^{-0.4}}{\ln(9900 \times t^{1.5}) \times [1 + (1 - x'')y + 3.7x''y]} - \log_{10} F(H\beta) \quad (1)$$

where $S(5\text{GHz})$ is the radio flux density at 5 GHz, in Jy, t is the nebular electron temperature in units of 10^4 K, y is the abundance ratio of He to H nuclei, by number, and x'' is the abundance ratio of doubly ionized He to all He atoms. For nebulae where values of t , x'' and y were not available in the literature, they were assumed to have values of 1, 0 and 0.11, respectively. $F(H\beta)$ is the observed integrated H β flux, listed in Table B1 from the sources given in Appendix B.

Logarithmic extinctions at H α ($C_{H\alpha}$) were also calculated using this formulation, after multiplying the first term in equation (1) by 2.85 to allow for a theoretical H α /H β flux ratio of 2.85 for nebulae at an electron temperature of 10^4 K (Storey & Hummer 1995) and replacing the second term by $\log(F(H\alpha))$, the [N II]-corrected IPHAS H α flux.

As well as using 5 GHz radio flux densities taken from the literature, we also made use of flux densities measured at frequencies of 1.4 GHz and 30 GHz (listed in Table B1 in Appendix B). For the $C_{H\alpha}$ and $C_{H\beta}$ extinction calculations, we scaled the 1.4 GHz and 30 GHz fluxes to 5 GHz by assuming a $\nu^{-0.1}$ optically thin free-free spectrum.

We also calculated $E(H\alpha - H\beta)$ reddening based on the ratio of the [N II]-corrected IPHAS H α flux to the published H β filter flux, using equation (2) below to obtain the colour excess between the H α and H β wave bands:

$$E(H\alpha - H\beta) = 2.5 \log_{10} \left(\frac{F(H\alpha)/F(H\beta)}{2.85} \right) \quad (2)$$

Using this color excess we were then able to obtain the absolute visual extinction $Av[E(H\alpha - H\beta)]$ using equation (3) below, which adopts the Galactic Reddening Law of Howarth (1983), with $R_V = Av/E(B - V) = 3.10$.

$$Av [E(H\alpha - H\beta)] = 2.669 \times E(H\alpha - H\beta), \quad (3)$$

where $A_V[E(H\alpha-H\beta)]$ is given in magnitudes. We also adopted the Howarth (1983) reddening law for $R_V = 3.1$ to calculate visual extinctions from $C_{H\alpha}$ ($A_V(C_{H\alpha})$) and $C_{H\beta}$ ($A_V(C_{H\beta})$), using equations (4) and (5):

$$A_V(C_{H\alpha}) = 3.140 \times C_{H\alpha} \quad (4)$$

$$A_V(C_{H\beta}) = 2.135 \times C_{H\beta} \quad (5)$$

Table C1 in Appendix C presents our derived $E(H\alpha-H\beta)$ values for 37 PNe, together with the corresponding values of A_V and $C_{H\beta}$. Table D1 in Appendix D presents the A_V values derived for 143 PNe from the various combinations of the three radio frequency fluxes and the $H\alpha$ and $H\beta$ fluxes. In order to calculate the uncertainties on all A_V values, standard error propagation was used. In some cases the literature fluxes did not have any listed uncertainties and in such cases A_V values obtained using those fluxes were discarded. For each PN, using the $A_V(C_{H\alpha})$ and $A_V(C_{H\beta})$ values corresponding to each of the radio frequencies, 1.4 GHz, 5 GHz and 30 GHz, we obtained a weighted mean radio extinction value, $A_V(\text{Radio})$, as listed in the penultimate column of Table D1.

The derived extinctions showing the best inter-agreement (the maximum allowed variation between the A_V values selected was ± 0.9 magnitudes) are shown in bold script in Table D1 and these were used for obtaining the final weighted averaged $A_V(\text{Radio})$ values. In cases when there were discrepancies between the different radio-based A_V values that were significantly larger than the formal flux uncertainties, the A_V value corresponding to the highest radio frequency was adopted, on the grounds that the higher frequency free-free radio emission was more likely to be optically thin.

For 143 PNe, a weighted mean of $A_V(\text{Radio})$ and $A_V[E(H\alpha-H\beta)]$ was used to generate a final adopted A_V and its uncertainty, which are listed in columns 3 of Table 1.

3.2 Maximum expected extinctions along PN lines of sight

To provide an upper limit to the extinction along the line of sight to each PN, we utilised the NASA/IPAC Infrared Archive (IRSA) *Galactic Dust Reddening and Extinction*⁶ website. The site uses the data of Schlegel et al. (1998) who created Galactic dust temperature and dust column maps by combining results from the IRAS 100 μm and COBE/DIRBE surveys. Using these maps and a relation between 100 μm surface brightness and visual extinction they were able to infer total extinctions along any given line of sight out of the Galaxy. Schlafly & Finkbeiner (2011) recalibrated the Schlegel et al. (1998) relations ($A_V(\text{Schlegel}) = 0.86 A_V(\text{Schlegel})$) and so we use the Schlafly & Finkbeiner (2011) values to predict the maximum value of A_V expected along a line of sight. These values are presented in Fig. 2 and in Appendix E.

⁶ <http://irsa.ipac.caltech.edu/applications/DUST/>

4 EXTINCTION DISTANCES TO GALACTIC PLANETARY NEBULAE

The technique that we adopt for determining distances to the northern Galactic PNe in our IPHAS sample is the 3D extinction mapping method, which uses an extinction versus distance relationship derived from field stars nearby on the sky to each PN in order to estimate its distance (see Lutz 1973; Gathier et al. 1986; Giannanco et al. 2011). In this paper we make use of three recent Galactic extinction versus distance mapping tools that are based on independent large stellar photometric databases. In Section 4.1 we derive extinction distances using the IPHAS-based H-MEAD 3D extinction mapping algorithm (Sale et al. 2014). In Section 4.2 we present distances obtained using the Pan-STARRS 1/2MASS-based BAYESTAR2019 3D extinction mapping tool (Green et al. 2019), while in Section 4.3 we make a limited comparison with distances obtained using the Gaia/2MASS-based STILISM 3D reddening mapping tool (Lallemand et al. 2019). In Section 4.4 we compare distances obtained using each of the above tools with reliable Gaia DR2 distances that are available for 17 of the PNe in our sample.

4.1 Distances using the H-MEAD 3D extinction mapping algorithm

This method rests on photometric data from the IPHAS survey in building distance-extinction relationships for A–K stars across the northern Plane. The source for the extinction-distance relations along the relevant lines of sight is the H-MEAD algorithm (Hierarchical Mapper of Extinction Against Distance) described by Sale (2012) and implemented by Sale et al. (2014). H-MEAD uses hierarchical Bayesian principles to determine the **required** properties of all the available A–K stars within any specified square pencil beam (voxel) in order to derive a mean relation between increasing extinction and distance.

To give a little more detail, Sale et al. (2014) used the IPHAS $(r' - i')$ and $(r' - H\alpha)$ colour-colour plane to assign (probabilistically) spectral type and luminosity class to every detected star in the voxel of interest. The algorithm rejects any objects that are not plausible A–K stars. This information constrains each accepted star's likely absolute magnitude and monochromatic extinction at 5495 Å (A_0 , a quantity that is for most purposes nearly identical to the band mean, A_V). These data, along with the IPHAS apparent magnitudes, are then used to calculate distances to all the retained stars, which are then dynamically-binned according to their distances: each bin contains a minimum of eight stars and is at least 100 pc in depth. H-MEAD takes into account sources of inaccuracy, scatter and bias such as photometric error, unresolved ISM substructure within the voxel, binarity and photometric magnitude limits (see also Sale et al. 2009; Giannanco et al. 2011).

The photometric depth of the IPHAS survey enabled the northern Galactic Plane to be extinction-mapped by Sale et al. (2014) with voxel sizes ranging from 5×5 arcmin 2 at the highest stellar densities up to, very occasionally, 60×60 arcmin 2 in the darkest regions. This variation is in response to the requirement of at least 200 stars to work with, within the voxel. The most frequent angular resolu-




Figure 2. H-MEAD distance-extinction relationships for the sight lines to (a) Abell 71 ($l = 85.00, b = +4.49$), (b) K 3-57 ($l = 72.20, b = +0.10$), (c) M 2-50 ($l = 97.68, b = -2.45$) and (d) NGC 7354 ($l = 107.84, b = +2.32$). The solid blue curve in each plot represents the H-MEAD extinction versus distance relation for the sightline to the nebula. The solid and dashed red horizontal lines correspond to the value of A_V derived for the nebula, and its corresponding uncertainties, while the solid and dashed green horizontal lines correspond to the Schlafly & Finkbeiner (2011) A_V limit and its uncertainties.

tion is 10×10 arcmin 2 . The useful distance range is usually from 1 up to 10 kpc, set by the bright and faint limits of the survey photometry, respectively.

The H-MEAD distance-extinction relationship curves from Sale et al. (2014) that we selected as best matching each of the PN sightlines were obtained from the website <http://www.iphas.org/data/extinction>.

Once the appropriate distance-extinction relation for the voxel enclosing the coordinates of a PN had been identified and downloaded, it was simply a matter of plotting it and reading off the distance corresponding to the final averaged A_V of the PN. The uncertainty on this distance was found by obtaining the distances corresponding to the upper and lower limits for A_V . We were able to do this for 143 PNe - examples are shown in Figure 2. The distance-extinction relationships for all the sightlines analyzed are presented in Appendix E. The cut-off distance in all of them is 14 kpc.

The distance-extinction curves typically begin with a sharp increase in A_V and end as a plateau with very little subsequent increase in A_V with distance. A sharp jump in extinction over a small distance range will be due to the presence of a dense interstellar cloud extinguishing the starlight.

There are two reasons for the terminal plateaus commonly seen. The first is that the dust layer is physically thin (100 pc or so above/below the Galactic equatorial plane) and is excited more quickly for sightlines at higher Galactic latitude. Lines of sight at high Galactic latitudes tend to run out of after a few kpc. This can be seen for the PN Abell 71 ($b = +4.49^\circ$) whose distance-extinction relationship is shown in Figure 2. Its curve reaches a plateau at a heliocentric distance of ~ 4 kpc. The second factor that can be at work, particularly at low Galactic latitudes within the Solar Circle, is high extinction limiting the detected stars to within a short distance - forcing H-MEAD to rely on the prior in extending the curve further out. In such cases the prior can impose the plateau (see Sale et al. 2014). The distance-extinction relationship for K 3-57 ($l = 72.20^\circ, b = 00.10^\circ$), shown in Figure 2, is a good example of this.

It is unavoidable that this method of distance estimation does not always deliver a useful result. For some PNe the derived value of A_V is above the maximum A_V of the plateau region of the relevant extinction-distance curve but below the A_V (Schlafly) value for the sightline. Where this happens, it is likely that more heavily reddened field stars

Table 1. Adopted extinctions and distances derived from Sale et al. (2014) H-MEAD and Green et al. (2019)

PN G	Name	A _V (mag)	H-MEAD (kpc)	Green+2019 (kpc)
030.8+03.4	Abell 47	8.27 ± 0.33	> 10.5	> 5.2
031.0+04.1	K 3-6	7.18 ± 0.17	5.30 ^{+0.50} _{-0.26}	> 4.5
031.3-00.5	HaTr 10	3.26 ± 0.19	0.94 ^{+0.19} _{-0.11}	2.33 ^{+0.74} _{-0.50}
031.7+01.7	PC 20	5.45 ± 0.10	4.57 ^{+0.33} _{-0.33}	12.6 ^{+2.31} _{-4.47}
032.0-03.0	K 3-18	3.93 ± 0.18	6.00 ^{+0.40} _{-0.49}	> 3.42
032.5-03.2	K 3-20	2.13 ± 0.19	2.45 ^{+0.77} _{-0.56}	3.63 ^{+0.50} _{-0.59}
032.9-02.8	K 3-19	3.40 ± 0.10	6.35 ^{+0.29} _{-0.36}	> 6.63
033.8-02.6	NGC 6741	1.89 ± 0.03	2.44 ^{+0.07} _{-0.06}	3.58 ^{+0.03} _{-0.03}
035.7-05.0	K 3-26	2.12 ± 0.16	7.11 ^{+6.27} _{-0.74}	> 5.2
035.9-01.1	Sh 2-71	1.44 ± 0.13	0.12 ^{+0.01} _{-0.01}	0.49 ^{+0.02} _{-0.02}
036.9-02.6	HaTr 13	2.82 ± 0.25	2.58 ^{+0.99} _{-0.65}	3.90 ^{+5.60} _{-0.77}
036.9-01.1	HaTr 11	5.00 ± 0.11	8.35 ^{+0.44} _{-0.33}	> 8.3
039.5-02.7	M 2-47	3.08 ± 0.10	2.05 ^{+0.20} _{-0.22}	> 3.6
039.8-02.1	K 3-17	7.93 ± 0.09	5.42 ^{+0.27} _{-0.27}	> 6.3
040.3-00.4	Abell 53	4.12 ± 0.11	0.87 ^{+0.08} _{-0.05}	0.86 ^{+0.01} _{-0.01}
040.4-03.1	K 3-30	2.62 ± 0.11	1.99 ^{+0.23} _{-0.49}	3.99 ^{+0.74} _{-0.08}
041.8+04.4	K 3-15	2.41 ± 0.08	> 9.00	> 4.83
043.0-03.0	M 4-14	2.72 ± 0.06	4.44 ^{+0.44} _{-0.43}	> 6.2
043.1+03.8	M 1-65	1.74 ± 0.05	2.25 ^{+0.11} _{-0.19}	3.22 ^{+0.19} _{-0.25}
043.3+02.2	PM 1-276	3.26 ± 0.11	1.99 ^{+0.20} _{-0.33}	3.03 ^{+0.29} _{-0.07}
045.9-01.9	K 3-33	5.30 ± 0.05	3.21 ^{+0.09} _{-0.18}	> 16.9
046.3-03.1	PB 9	3.39 ± 0.06	6.83 ^{+0.18} _{-0.12}	> 4.3
046.4-04.1	NGC 6803	1.24 ± 0.03	4.25 ^{+0.14} _{-0.25}	3.65 ^{+0.45} _{-0.23}
046.8+02.9	CTSS 4	3.03 ± 0.19	> 9.0	> 4.3
047.1+04.1	K 3-21	2.77 ± 0.19	7.32 ^{+6.68} _{-0.65}	> 4.3
048.0-02.3	PB 10	3.59 ± 0.11	6.73 ^{+0.29} _{-0.23}	> 8.8
048.1+01.1	K 3-29	6.89 ± 0.05	> 10.0	> 9.3
048.5+04.2	K 4-16	2.92 ± 0.06	> 7.0	> 4.5
048.7+02.3	K 3-24	6.07 ± 0.19	> 8.0	> 2.5
048.7+01.9	Hen 2-429	3.50 ± 0.10	6.04 ^{+0.47} _{-0.63}	4.67 ^{+1.76} _{-0.59}
049.4+02.4	Hen 2-428	2.86 ± 0.09	2.90 ^{+0.36} _{-0.30}	3.84 ^{+0.05} _{-0.09}
050.4-01.6	K 4-28	5.15 ± 0.06	5.74 ^{+0.21} _{-0.11}	6.75 ^{+0.04} _{-0.06}
051.0-04.5	PC 22	2.10 ± 0.15	> 7.0	> 5.6
051.0+03.0	Hen 2-430	3.56 ± 0.17	2.71 ^{+0.59} _{-0.44}	3.14 ^{+2.13} _{-0.06}
051.3+01.8	PM 1-295	3.56 ± 0.11	0.97 ^{+0.15} _{-0.02}	1.40 ^{+0.08} _{-0.16}
051.9-03.8	M 1-73	1.57 ± 0.10	3.46 ^{+0.66} _{-0.64}	4.45 ^{+0.66} _{-0.72}
052.2-04.0	M 1-74	1.58 ± 0.12	6.87 ^{+0.63} _{-0.87}	7.21 ^{+1.81}
052.5-02.9	Me 1-1	1.15 ± 0.09	2.42 ^{+0.18} _{-0.40}	2.73 ^{+0.52} _{-0.46}
052.9-02.7	K 3-41	1.44 ± 0.30	2.74 ^{+0.46} _{-0.76}	2.74 ^{+0.49} _{-1.07}
052.9+02.7	K 3-31	4.61 ± 0.08	4.46 ^{+0.21} _{-0.23}	> 2.5
053.2-01.5	K 3-38	4.56 ± 0.08	5.88 ^{+0.17} _{-0.07}	9.74 ^{+0.04} _{-0.04}
053.8-03.0	Abell 63	0.67 ± 0.20	0.92 ^{+0.25} _{-0.41}	0.76 ^{+0.47} _{-0.19}
054.2-03.4	Necklace	2.41 ± 0.19	> 8.0	> 4.3
054.4-02.5	M 1-72	2.03 ± 0.32	6.82 ^{+3.12} _{-1.74}	> 5.5
055.1-01.8	K 3-43	3.48 ± 0.27	6.65 ^{+0.65} _{-0.80}	> 4.1
055.2+02.8	Hen 2-432	3.40 ± 0.08	2.50 ^{+0.31} _{-0.35}	> 3.1
055.3+02.7	Hen 1-1	3.85 ± 0.07	4.15 ^{+0.20} _{-0.38}	> 7.8
055.5-00.5	M 1-71	4.39 ± 0.13	7.01 ^{+0.19} _{-0.23}	9.55 ^{+0.02} _{-0.02}
055.6+02.1	Hen 1-2	2.78 ± 0.07	0.28 ^{+1.11} _{-0.02}	2.50 ^{+0.23} _{-0.20}
056.0+02.0	K 3-35	6.63 ± 0.17	9.96 ^{+4.38} _{-1.21}	> 10.9
056.4-00.9	K 3-42	5.24 ± 0.07	8.38 ^{+0.20} _{-0.19}	10.5 ^{+0.03} _{-0.03}
057.9-01.5	Hen 2-447	4.42 ± 0.05	6.77 ^{+0.05} _{-0.21}	> 8.0
058.9+01.3	K 3-40	3.00 ± 0.08	5.43 ^{+0.15} _{-0.17}	6.64 ^{+0.16} _{-0.64}
059.4+02.3	K 3-37	4.33 ± 0.07	7.20 ^{+0.25} _{-0.21}	> 7.5
059.0+04.6	K 3-34	2.57 ± 0.29	> 7.0	> 4.7
059.9+02.0	K 3-39	4.41 ± 0.07	7.76 ^{+0.26} _{-0.23}	> 7.0
060.4+01.5	PM 1-310	4.96 ± 0.27	> 8.0	> 5.8
060.5-00.3	K 3-45	4.91 ± 0.19	5.33 ^{+0.56} _{-0.65}	10.1 ^{+1.22} _{-0.45}
060.5+01.8	Hen 2-440	3.93 ± 0.05	8.47 ^{+0.28} _{-0.25}	11.1 ^{+0.04} _{-0.04}
060.8-03.6	NGC 6853	0.10 ± 0.14	0.11 ^{+0.07} _{-0.04}	0.96 ^{+0.02} _{-0.08}

Table 1 – continued Adopted extinctions and distances derived from Sale et al. (2014) H-MEAD and Green et al. (2019)

PN G	Name	A _V (mag)	H-MEAD (kpc)	Green+2019 (kpc)
062.4-00.2	M 2-48	3.45 ± 0.15	2.75 ^{+0.29} _{-0.55}	3.30 ^{+0.06} _{-0.03}
063.8-03.3	K 3-54	4.82 ± 0.22	5.30 ^{+0.49} _{-0.21}	> 4.3
064.9-02.1	K 3-53	2.91 ± 1.02	3.88 ^{+2.84} _{-2.57}	3.27 ^{+11.7} _{-2.21}
065.1-03.5	We 1-9	3.16 ± 1.02	1.34 ^{+3.48} _{-0.62}	1.60 ^{+13.3} _{-0.58}
065.9+00.5	NGC 6842	2.51 ± 0.11	4.34 ^{+0.16} _{-0.10}	4.85 ^{+0.04} _{-0.04}
066.9+02.2	K 4-37	4.25 ± 0.19	> 7.0	> 8.8
067.9-00.2	K 3-52	8.46 ± 0.18	> 8.0	> 7.8
068.3-02.7	Hen 2-459	4.61 ± 0.05	5.44 ^{+0.09} _{-0.04}	> 5.0
068.6+01.1	Hen 1-4	2.36 ± 0.11	3.40 ^{+0.66} _{-0.31}	3.10 ^{+0.25} _{-0.07}
068.7+01.9	K 4-41	3.72 ± 0.32	> 8.5	11.5 ^{+0.19} _{-0.20}
068.7+03.0	PC 23	3.66 ± 0.21	> 7.0	> 6.7
068.8-00.0	M 1-75	4.20 ± 0.19	6.20 ^{+0.25} _{-0.67}	7.75 ^{+1.16} _{-4.53}
069.2+03.8	K 3-46	2.15 ± 0.26	4.73 ^{+1.49} _{-1.80}	5.92 ^{+9.03} _{-2.50}
069.2+02.8	K 3-49	3.37 ± 0.07	> 8.0	> 3.5
069.4-02.6	NGC 6894	1.40 ± 0.14	1.01 ^{+0.01} _{-0.01}	1.43 ^{+0.14} _{-0.07}
069.6-03.9	K 3-58	4.11 ± 0.19	> 7.0	> 5.5
069.7+00.0	K 3-55	7.05 ± 0.16	> 8.0	> 8.5
071.6-02.3	M 3-35	4.61 ± 0.08	4.57 ^{+0.30} _{-0.27}	> 5.5
072.1+00.1	K 3-57	5.35 ± 0.33	9.94 ^{+3.88} _{-1.75}	> 5.1
073.0-02.4	K 3-76	4.27 ± 0.29	4.30 ^{+0.45} _{-0.57}	> 5.4
074.5+02.1	NGC 6881	3.60 ± 0.03	5.15 ^{+0.16} _{-0.05}	5.87 ^{+0.05} _{-0.05}
075.6+04.3	ARO 342	2.26 ± 0.19	> 7.0	> 4.7
076.3+01.1	Abell 69	4.20 ± 0.17	5.50 ^{+0.16} _{-0.19}	6.25 ^{+0.12} _{-0.12}
076.4+01.8	KjPn 3	3.42 ± 0.19	5.02 ^{+0.49} _{-0.28}	4.86 ^{+0.46} _{-0.63}
077.5+03.7	KjPn 1	1.50 ± 0.21	2.36 ^{+0.74} _{-0.33}	2.28 ^{+0.35} _{-0.68}
077.7+03.1	KjPn 2	5.51 ± 0.19	6.89 ^{+2.24} _{-0.87}	> 4.6
078.3-02.7	K 4-53	4.88 ± 0.10	9.31 ^{+4.69} _{-2.02}	> 3.9
078.9+00.7	Sd 1	5.06 ± 0.19	2.84 ^{+0.23} _{-0.17}	1.07 ^{+0.01} _{-0.01}
084.2+01.0	K 4-55	2.94 ± 0.18	0.63 ^{+0.10} _{-0.01}	1.17 ^{+0.33} _{-0.38}
084.9+04.4	Abell 71	2.44 ± 0.11	2.26 ^{+0.08} _{-0.04}	2.44 ^{+0.01} _{-0.01}
088.7+04.6	K 3-78	4.47 ± 0.05	7.70 ^{+6.30} _{-1.41}	> 3.2
088.7-01.6	NGC 7048	0.82 ± 0.04	2.03 ^{+0.05} _{-0.06}	1.99 ^{+0.34} _{-0.75}
089.0+00.3	NGC 7026	1.77 ± 0.05	2.15 ^{+0.11} _{-0.11}	2.68 ^{+0.01} _{-0.01}
089.8-00.6	Sh 1-89	3.52 ± 0.20	3.39 ^{+0.40} _{-0.18}	5.99 ^{+0.06} _{-0.40}
091.6-04.8	K 3-84	1.25 ± 0.24	6.83 ^{+7.17} _{-2.01}	> 4.3
091.6+01.8	We 1-11	4.43 ± 0.06	2.16 ^{+0.11} _{-0.04}	2.07 ^{+0.20} _{-0.13}
093.3-00.9	K 3-82	3.72 ± 0.14	4.75 ^{+0.15} _{-0.13}	> 7.8
093.3-02.4	M 1-79	0.84 ± 0.04	2.31 ^{+0.17} _{-0.20}	1.33 ^{+0.14} _{-0.25}
094.5-00.8	K 3-83	4.75 ± 0.04	14.0 ^{+0.47} _{-2.5}	> 9.1
095.1-02.0	M 2-49	3.67 ± 0.06	8.05 ^{+0.78} _{-0.51}	> 5.1
095.2+00.7	K 3-62	5.35 ± 0.04	5.37 ^{+0.04} _{-0.04}	> 8.2
096.3+02.3	K 3-61	3.78 ± 0.05	3.25 ^{+0.20} _{-0.14}	8.59 ^{+2.37} _{-0.12}
097.6-02.4	M 2-50	2.01 ± 0.05	> 6.0	> 8.0
098.1+02.4	K 3-63	2.54 ± 0.08	2.84 ^{+0.39} _{-0.41}	3.70 ^{+0.29} _{-0.19}
098.2+04.9	K 3-60	4.55 ± 0.05	> 5.0	> 2.8
102.8-05.0	Abell 80	0.12 ± 0.06	1.54 ^{+0.43} _{-0.45}	0.33 ^{+0.05} _{-0.03}
103.2+00.6	M 2-51	1.79 ± 0.08	1.60 ^{+0.10} _{-0.07}	2.33 ^{+0.05} _{-0.04}
103.7+00.4	M 2-52	2.91 ± 0.05	4.19 ^{+0.04} _{-0.06}	3.85 ^{+0.02} _{-0.02}
104.1+01.0	Bl 2-1	5.31 ± 0.04	5.26 ^{+0.09} _{-0.09}	> 4.8
104.4-01.6	M 2-53	2.41 ± 0.05	2.09 ^{+0.07} _{-0.05}	> 4.8
107.4-02.6	K 3-87	4.45 ± 0.12	> 6.0	> 3.7
107.4-00.6	K 4-57	3.85 ± 0.19	> 6.0	> 10.7
107.7-02.2	M 1-80	1.64 ± 0.05	4.33 ^{+0.25} _{-0.26}	5.09 ^{+0.18} _{-0.51}
107.8+02.3	NGC 7354	3.70 ± 0.03	2.12 ^{+0.06} _{-0.03}	2.56 ^{+0.49} _{-0.04}
112.5-00.1	KjPn 8	1.80 ± 0.11	2.36 ^{+0.03} _{-0.26}	2.88 ^{+0.24} _{-0.11}
112.5+03.7	K 3-88	4.65 ± 0.06	> 5.0	> 3.3
119.3+00.3	BV 5-1	2.52 ± 0.05	4.56 ^{+0.14} _{-0.07}	> 5.5
121.6+00.0	BV 5-2	2.43 ± 0.06	1.89 ^{+0.15} _{-0.13}	3.40 ^{+0.03} _{-0.04}
121.6+03.5	We 1-1	4.05 ± 0.06	3.28 ^{+0.06} _{-0.18}	> 2.1

Table 1 – continued Adopted extinctions and distances derived from Sale et al. (2014) H-MEAD and Green et al. (2019)

PN G	Name	A _V (mag)	H-MEAD (kpc)	Green+2019 (kpc)
122.1-04.9	Abell 2	1.75 ± 0.05	> 5.0	> 2.9
126.3+02.9	K 3-90	2.76 ± 0.05	3.92 ^{+0.08} _{-0.07}	> 3.9
126.6+01.3	IPHAS PN-1	1.47 ± 1.05	0.10 ^{+2.14} _{-0.03}	0.76 ^{+2.89} _{-0.45}
129.5+04.5	K 3-91	3.51 ± 0.05	> 4.5	> 2.3
130.2+01.3	IC 1747	2.02 ± 0.04	4.20 ^{+0.12} _{-0.21}	4.64 ^{+0.08} _{-0.07}
130.4+03.1	K 3-92	1.96 ± 0.19	4.42 ^{+0.39} _{-0.90}	4.81 ^{+10.1} _{-1.53}
131.5+02.6	Abell 3	0.44 ± 0.06	0.43 ^{+0.02} _{-0.18}	0.26 ^{+0.05} _{-0.02}
132.4+04.7	K 3-93	3.74 ± 0.06	> 4.5	> 2.3
136.1+04.9	Abell 6	2.14 ± 0.19	0.46 ^{+0.22} _{-0.03}	1.00 ^{+0.05} _{-0.05}
138.8+02.8	IC 289	2.33 ± 0.02	0.96 ^{+0.01} _{-0.01}	3.82 ^{+0.03} _{-0.03}
142.1+03.4	K 3-94	2.84 ± 0.15	> 5.0	> 3.5
147.4-02.3	M 1-4	3.23 ± 0.06	2.75 ^{+0.07} _{-0.09}	> 4.0
147.8+04.1	M 2-2	2.95 ± 0.15	4.55 ^{+9.45} _{-0.45}	> 3.0
151.4+00.5	K 3-64	3.28 ± 0.25	9.98 ^{+4.02} _{-3.54}	> 10.2
173.5+03.2	Pu 2	3.95 ± 0.17	3.22 ^{+0.25} _{-0.26}	> 1.5
178.3-02.5	K 3-68	2.73 ± 0.19	8.88 ^{+5.12} _{-4.04}	> 5.3
181.5+00.9	Pu 1	2.55 ± 0.06	10.1 ^{+3.94} _{-2.1}	> 4.7
184.0-02.1	M 1-5	2.84 ± 0.02	> 4.5	> 5.1
184.6+00.6	K 3-70	4.13 ± 0.12	> 5.0	> 5.0
184.8+04.4	K 3-71	3.52 ± 0.27	> 4.0	> 2.6
194.2+02.5	J 900	1.64 ± 0.03	> 4.5	> 10.2
201.7+02.5	K 4-48	3.05 ± 0.10	> 4.0	> 4.4
210.3+01.9	M 1-8	2.48 ± 0.21	> 5.0	> 3.5
212.0+04.3	M 1-9	1.17 ± 0.07	2.90 ^{+0.58} _{-0.76}	> 4.1

along the line of sight were fainter than the IPHAS survey limits. In cases of PNe with A_V falling in the plateau region of the extinction-distance relationship, the uncertainties on their estimated distances will be much larger, as shown by the relationship for M 2-50 in Figure 2. Seventeen PNe were found to have A_V values larger than the maximum A_V for their H-MEAD extinction-distance curve as well as lying above the uncertainty limits for the corresponding A_V(Schlafly) values. Five of those 17 PNe are located between $l=184$ and $l=202$ degrees. In some of these cases this could be due to overestimated extinction values due to errors in the adopted radio fluxes or in the adopted [N II]/H α correction factors. Another possibility is that there might be significant variations of A_V within some of the $10' \times 10'$ arcmin voxels used to sample the sight lines. We consider it unlikely that significant internal dust extinction within the PNe themselves is the cause of such discrepancies. There are only a few confirmed cases of PNe with large internal dust extinctions (e.g. NGC 7027, NGC 6302) - typical bright PNe such as NGC 7009 have been found to have relatively small internal dust columns (Walsh et al. 2016).

The extinction-distance relationship for NGC 7354 shown in Figure 2 is an example of a well-behaved curve, where the calculated A_V for the PN lies on the sharply increasing portion of the curve, and where the A_V(Schlafly) value lies above the A_V limit corresponding to the plateau region of the extinction-distance curve.

The measured distances to all 143 PNe are listed in column 4 of Table 1. Table 1. For PNe which had calculated A_V values greater than the H-MEAD curve's maximum A_V limit, they were assigned a minimum distance corresponding

to just after the onset of the final plateau in the H-MEAD extinction curve.

4.2 Distances using the BAYESTAR2019 3D extinction mapping algorithm

Green et al. (2019) have constructed a 3D dust reddening map for the sky north of declination -30° . By making use of Pan-STARRS 1 optical photometry, 2MASS near-infrared photometry and Gaia DR2 parallaxes and applying a hierarchical Bayesian model (BAYESTAR2019), they inferred distances, reddenings and types for 800 million stars in order to create a 3D dust map extending beyond several kpc, which can be accessed at doi:10.7910/DVN/2EJ9TX. We made use of this map to estimate distances for the PNe in our samples, using reddening values corresponding to the extinctions listed in Table 1. The resulting distance estimates, or lower limits, are listed in the final column of the same table. As was the case with H-MEAD distance estimates, where PNe had calculated A_V values that were greater than the maximum A_V limits of the BAYESTAR2019 curves, they were assigned a minimum distance corresponding to just after the onset of the final plateau in the extinction curves.

4.3 Distances using the STILISM 3D extinction mapping algorithm

We have also used the STILISM 3D reddening mapping algorithm of Lallement et al. (2019) (see https://astro-acri-st.fr/gaia_dev/), to estimate reddening distances to PNe in our sample. Because the algorithm makes use of stars having 5σ or better Gaia DR2 parallaxes, for most directions

this tool does not reach beyond distances of ~ 3 kpc and so provides only lower limits to the distances to the majority of the PNe in our sample. However, STILISM distances, or lower limits, are listed in column 8 of Table 2 for 17 of the PNe in our sample that are judged to have reliable Gaia DR2 distances.

4.4 Comparison with Gaia DR2 distances

For a subset of the PNe in our sample that were judged to have reliable central star distances from Gaia DR2 parallaxes, we can compare these reliable astrometric distances with the distances estimated using the different reddening mapping methods.

The following criteria were used to select the PNe in our sample as having reliable central star parallaxes in the Gaia DR2 archive (Gaia Collaboration et al. 2018): (a) the number of effective Gaia visits, $N_{\text{per}} \geq 8$; (b) a parallax signal-to-noise ratio > 3.4 ; (c) a renormalised unit-weighted error $u/u_0(G, B_P - R_P) < 1.4$, where $u = \sqrt{\chi^2/(N-5)}$ and $u_0(G, B_P - R_P)$ is the magnitude and colour-dependent reference value.⁷ 17 of the PNe in our sample satisfied these criteria. Table 2 lists Gaia DR2 parallaxes⁸ in column 3, followed in column 4 by the distances obtained by inverting the parallaxes and in column 5 by Gaia DR2 distances from Bailer-Jones et al. (2018) that are based on a weak distance prior and which are often adopted for stars with lower signal-to-noise Gaia DR2 parallaxes. For the sample of 17 PNe listed in Table 2, the Gaia DR2 parallaxes are of sufficiently high reliability that the Bailer-Jones et al. (2018) distances agree closely with the inverted parallax distances. One of the PNe listed in Table 2 has a pre-Gaia parallax measurement. Benedict et al. (2009) reported a parallax of 2.47 ± 0.16 mas for the central star of NGC 6853, versus the Gaia DR2 parallax of 2.658 ± 0.044 mas.

For comparison with the above Gaia DR2 PN central star distances, column 6 of Table 2 lists the PN distances obtained using the H-MEAD extinction mapping method of Sale et al. (2014), while column 7 lists the distances obtained using the BAYESTAR2019 extinction mapping method of Green et al. (2019). Column 8 of Table 2 provides Lallemand et al. (2019) STILISM distances for 10 of the 17 PNe that have reliable Gaia distances. Finally, column 9 of Table 2 lists PN distances from Frew et al. (2016), a widely used statistical distance estimator that is based on a H α surface brightness-radius relation for PNe.

Figure 3 plots the Gaia DR2 distances of Bailer-Jones et al. (2018) against the distance estimates obtained from (a) H-MEAD (Sale et al. 2014); (b) Green et al. (2019); (c) Lallemand et al. (2019); and (d) Frew et al. (2016). The Pearson correlation coefficient (r) and the corresponding probability value (p) for a null correlation have been calculated for each plot (lower limits to distances were not used). In descending order of their correlation with the Gaia DR2 distances are the Frew et al. (2016) distances ($r=0.61$, $p=0.012$; 16

⁷ See Gaia mission document GAIA-C3-TN-LU-LL-124-01 by L. Lindegren at www.rssd.esa.int/doc_fetch.php?id=3757412

⁸ As recommended by Lindegren et al. (2018), the published Gaia-DR2 parallaxes have been increased by 0.03 mas to allow for the global zero-point of -0.03 mas.

PNe), the H-MEAD distances ($r=0.58$, $p=0.023$; 15 PNe), the Lallemand et al. (2019) distances ($r=0.57$, $p=0.085$; for 10 PNe at distances < 3 kpc), with the Green et al. (2019) distances yielding $r=0.35$ and a p value of 0.22 for 14 PNe.

Giammanco et al. (2011) used the MEAD reddening-distance algorithm of Sale et al. (2009) to estimate reddening distances for 70 PNe located in IPHAS fields, using reddening estimates that were based mainly on spectroscopic H α /H β ratios taken from the literature. Only seven of the eighteen PNe in Table 2 have distance estimates from Giammanco et al. (2011) that are not upper or lower limits, so we have not estimated their correlation coefficient with the Gaia DR2 distances.

5 DISCUSSION AND CONCLUSIONS


Using aperture photometry we have measured in-band IPHAS H α filter fluxes for 46 PNe discovered by IPHAS and for 151 previously catalogued PNe. These in-band fluxes were corrected for [N II] 6548,6583 Å contributions using published spectroscopic [N II]/H α ratios for 39 of the IPHAS-discovered PNe and for all 151 of the previously catalogued PNe. Radio free-free flux measurements at 1.4, 5 or 30 GHz were available from the literature for 143 of the previously catalogued PNe and these were combined with the [N II]-corrected H α fluxes and with previously published H β filter fluxes to give a weighted mean extinction estimate for each PN from its radio free-free to optical Balmer line flux ratios.

We have used these derived interstellar extinctions with the IPHAS-based H-MEAD interstellar extinction mapping tool of Sale et al. (2014) in order to estimate distances to each PN from the H-MEAD extinction vs. distance plot for each sight line. For comparison purposes, we have also estimated extinction distances to the PNe using the Green et al. (2019) extinction mapping tool, which is based on Pan-STARRS 1 and 2MASS photometry and Gaia parallaxes. Gaia DR2 distances that were judged to be reliable were available for 17 of the PNe in our sample and these distances were compared to the H-MEAD and Green et al. extinction distances for this sub-sample, as well as to a smaller number of extinction distances obtained using the Lallemand et al. (2019) extinction mapping tool. We also compared the Gaia DR2 distances for this sub-sample with the widely used PN distances from Frew et al. (2016) that are based on a relation between nebular radius and nebular H α surface brightness. For the sub-sample of 17 PNe, the Frew et al. and the H-MEAD distances were found to show similar degrees of correlation with Gaia DR2 distances.

Given that 3D extinction mapping tools are expected to become increasingly sophisticated and to achieve higher angular resolution over time, the weakest link may often be the accuracy of reddening estimates, which need to be significantly improved for many objects. In the case of PNe, reddening values based on observed Balmer line ratios cover too small a wavelength range to be considered ideal. Radio free-free to optical Balmer line ratios cover a much larger wavelength baseline and yield total dust extinctions at the wavelengths of the Balmer lines, but currently many PNe either have no measured radio fluxes or have radio fluxes with relatively low signal to noise ratios. More accurate radio

Table 2. Comparison to Gaia DR2 distances

PN G	Name	Gaia DR2 Parallax (mas)	Gaia DR2 Parallax Distance (kpc)	Gaia DR2 Bailer-Jones+ (kpc)	H-MEAD (kpc)	Green +2019 (kpc)	Lallement +2019 (kpc)	Frew +2016 (kpc)
035.9-01.1	Sh 2-71	0.588±0.035	1.62 ^{+0.10} _{-0.08}	1.62 ^{+0.10} _{-0.09}	0.12 ^{+0.01} _{-0.01}	0.49 ^{+0.02} _{-0.02}	0.76 ^{+0.11} _{-0.05}	1.32 ± 0.47
043.1+03.8	M 1-65	0.224±0.040	3.94 ^{+0.74} _{-0.53}	3.90 ^{+0.74} _{-0.55}	2.25 ^{+0.11} _{-0.19}	3.22 ^{+0.19} _{-0.25}	1.68 ^{+0.06} _{-0.03}	6.85 ± 2.01
048.5+04.2	K 4-16	0.191±0.053	4.53 ^{+1.37} _{-0.86}	4.39 ^{+1.32} _{-0.86}	> 7.0	> 4.5	> 3.0	11.1 ± 3.5
051.3+01.8	PM 1-295	0.339±0.030	2.71 ^{+0.24} _{-0.20}	2.72 ^{+0.24} _{-0.21}	1.40 ^{+0.08} _{-0.16}	1.21 ^{+0.03} _{-0.03}	> 3.5	–
052.5-02.9	Me 1-1	0.226±0.037	3.91 ^{+0.67} _{-0.49}	3.87 ^{+0.65} _{-0.49}	2.42 ^{+0.18} _{-0.40}	2.73 ^{+0.52} _{-0.46}	2.91 ^{+0.87} _{-0.57}	6.17 ± 1.87
053.8-03.0	Abell 63	0.361±0.034	2.55 ^{+0.25} _{-0.20}	2.57 ^{+0.26} _{-0.21}	0.92 ^{+0.25} _{-0.41}	0.76 ^{+0.47} _{-0.19}	0.63 ^{+0.47} _{-0.11}	3.79 ± 1.12
060.8-03.6	NGC 6853	2.658± 0.044	0.37 ^{+0.01} _{-0.01}	0.37 ^{+0.01} _{-0.01}	0.11 ^{+0.07} _{-0.04}	0.96 ^{+0.02} _{-0.08}	0.42 ^{+0.10} _{-0.23}	0.31 ± 0.09
065.9+00.5	NGC 6842	0.476±0.046	1.98 ^{+0.20} _{-0.16}	1.99 ^{+0.21} _{-0.17}	4.34 ^{+0.16} _{-0.10}	4.85 ^{+0.04} _{-0.04}	2.15 ^{+0.10} _{-0.09}	2.20 ± 0.64
069.4-02.6	NGC 6894	0.839±0.131	1.15 ^{+0.21} _{-0.15}	1.18 ^{+0.24} _{-0.17}	1.01 ^{+0.01} _{-0.01}	1.43 ^{+0.14} _{-0.07}	1.17 ^{+0.04} _{-0.02}	1.50 ± 0.43
069.7+00.0	K 3-55	0.600±0.071	1.58 ^{+0.20} _{-0.15}	1.61 ^{+0.22} _{-0.17}	> 8.0	> 8.5	> 3.0	3.54 ± 1.32
076.4+01.8	KjPn 3	0.307±0.090	2.97 ^{+1.09} _{-0.63}	2.89 ^{+1.02} _{-0.63}	5.02 ^{+0.49} _{-0.28}	4.86 ^{+0.46} _{-0.63}	2.61 ^{+1.20} _{-1.15}	17.2 ± 4.9
107.8+02.3	NGC 7354	0.454±0.085	2.07 ^{+0.45} _{-0.31}	2.10 ^{+0.50} _{-0.35}	2.12 ^{+0.06} _{-0.03}	2.56 ^{+0.49} _{-0.04}	> 3.0	1.26 ± 0.37
130.2+01.3	IC 1747	0.308±0.051	2.95 ^{+0.51} _{-0.38}	2.95 ^{+0.54} _{-0.40}	4.20 ^{+0.12} _{-0.21}	4.64 ^{+0.08} _{-0.07}	> 3.5	3.08 ± 1.00
131.5+02.6	Abell 3	0.343±0.084	2.68 ^{+0.78} _{-0.49}	2.70 ^{+0.86} _{-0.54}	0.43 ^{+0.02} _{-0.18}	0.26 ^{+0.05} _{-0.02}	0.37 ^{+0.01} _{-0.01}	2.47 ± 0.73
136.1+04.9	Abell 6	1.004± 0.177	0.97 ^{+0.20} _{-0.14}	1.01 ^{+0.25} _{-0.17}	0.46 ^{+0.22} _{-0.03}	1.00 ^{+0.05} _{-0.05}	1.10 ^{+0.10} _{-0.07}	1.30 ± 0.39
138.8+02.8	IC 289	0.628±0.060	1.52 ^{+0.15} _{-0.13}	1.53 ^{+0.16} _{-0.13}	0.96 ^{+0.01} _{-0.01}	3.82 ^{+0.03} _{-0.03}	> 3.5	1.88 ± 0.58
147.8+04.1	M 2-2	0.192±0.054	4.54 ^{+1.46} _{-0.91}	4.26 ^{+1.23} _{-0.82}	4.55 ^{+9.45} _{-0.45}	> 3.0	> 3.0	5.22 ± 1.51

**Figure 3.** Correlations between Gaia-DR2 distances from Bailer-Jones et al. (2018) and those from the following methods. Top left: (a) H-MEAD (Sale et al. 2014), with a Pearson correlation coefficient of $r = 0.58$ (15 PNe); top right: (b) Green et al. (2019), with $r = 0.35$ (14 PNe); bottom left: (c) Lallement et al. (2019), with $r = 0.57$ (10 PNe); and bottom right: (d) Frew et al. (2016), with $r = 0.61$ (16 PNe). Upward-pointing arrows indicate lower limits, which were not used in calculating correlation coefficients. Note that the y-axis scale for plot (d) differs from those of the other three plots.

flux measurements, over a range of radio wavelengths, are needed. For the northern PNe with IPHAS H α fluxes that have been discussed here, the ongoing Very Large Array Sky Survey (VLASS; Lacy et al. 2020) will help reach this goal, by providing 2-4 GHz radio fluxes at an angular resolution of 2.5 arcsec and with a 1σ sensitivity of 0.07 mJy by the completion of the survey.

DATA AVAILABILITY

Calibrated IPHAS images can be obtained from <http://apm3.ast.cam.ac.uk/cgi-bin/wfs/dqc.cgi>. H-MEAD distance-extinction data for fields within the IPHAS footprint can be obtained from <http://www.iphas.org/data/extinction>.

ACKNOWLEDGEMENTS

TD's research has been supported under grants MOST104-2628-M-001-004-MY3 and MOST107-2119-M-001-031-MY3 from the Ministry of Science and Technology of Taiwan, and grant AS-IA-106-M03 from Academia Sinica. Her work was also partially funded by the Sonderforschungsbereich SFB881 "The Milky Way System" of the German Research Foundation (DFG). MB and RW acknowledge support from European Research Council grant SNDUST ERC-2015-AdG-694520. DJ and AM acknowledge support from the State Research Agency (AEI) of the Spanish Ministry of Science, Innovation and Universities (MCIU) and the European Regional Development Fund (FEDER) under grant AYA2017-83383-P.

REFERENCES

- Aaquist O. B., Kwok S., 1990, *A&AS*, 84, 229
 Acker A., Stenholm B., Tylenda R., Raytchev B., 1991, *A&AS*, 90, 89
 Acker A., Marcout J., Ochsenbein F., Stenholm B., Tylenda R., Schohn C., 1992, The Strasbourg-ESO Catalogue of Galactic Planetary Nebulae. Parts I, II.
 Aller L. H., Czyzak S. J., 1979, *Astrophysics and Space Science*, 62, 397
 Bailer-Jones C. A. L., Rybizki J., Fouesneau M., Mantelet G., Andrae R., 2018, *AJ*, 156, 58
 Barentsen G., et al., 2014, *MNRAS*, 444, 3230
 Barker T., 1978, *ApJ*, 221, 145
 Benedict G. F., et al., 2009, *AJ*, 138, 1969
 Bohigas J., 2001, *Revista Mexicana de Astronomia y Astrofisica*, 37, 237
 Cahn J. H., Rubin R. H., 1974, *AJ*, 79, 128
 Cahn J. H., Kaler J. B., Stanghellini L., 1992, *A&AS*, 94, 399
 Cantrell J. R., 2014, Planetary Nebulae, <http://www.chara.gsu.edu/~cantrell/pne/index.htm>
 Capriotti E. R., Daub C. T., 1960, *ApJ*, 132, 677
 Carrasco L., Serrano A., Costero R., 1983, *Rev. Mex. Astron. Astrofis.*, 8, 187
 Collins G. W. I., Daub C. T., O'Dell C. R., 1961, *ApJ*, 133, 471
 Condon J. J., Kaplan D. L., 1998, *ApJS*, 117, 361
 Corradi R. L. M., et al., 2011, *MNRAS*, 410, 1349
 DENIS Consortium 2003, VizieR Online Data Catalog, p. B/denis
 Drew J. E., et al., 2005, *MNRAS*, 362, 753
 Frew D. J., Parker Q. A., 2010, *Publ. Astron. Soc. Australia*, 27, 129
 Frew D. J., Bojićić I. S., Parker Q. A., 2013, *MNRAS*, 431, 2
 Frew D. J., Parker Q. A., Bojićić I. S., 2016, *MNRAS*, 455, 1459
 Gaia Collaboration et al., 2018, *A&A*, 616, A1
 Gathier R., Pottasch S. R., Pel J. W., 1986, *A&A*, 157, 171
 Gaustad J. E., McCullough P. R., Rosing W., Van Buren D., 2001, *The Publications of the Astronomical Society of the Pacific*, 113, 1326
 Giannanco C., et al., 2011, *A&A*, 525, A58
 González-Santamaría I., Manteiga M., Manchado A., Ulla A., Dafonte C., 2019, *A&A*, 630, A150
 González-Solares E. A., et al., 2008, *MNRAS*, 388, 89
 Green G. M., et al., 2015, *ApJ*, 810, 25
 Green G. M., Schlafly E., Zucker C., Speagle J. S., Finkbeiner D., 2019, *ApJ*, 887, 93
 Guerrero M. A., Manchado A., Serra-Ricart M., 1996, *ApJ*, 456, 651
 Henry R. B. C., Kwinter K. B., Jaskot A. E., Balick B., Morrison M. A., Milingo J. B., 2010, *ApJ*, 724, 748
 Higgs L. A., 1971, *Publications of The Astrophysics Branch, NRC*, Ottawa, 1, 1
 Howarth I. D., 1983, *MNRAS*, 203, 301
 Howarth I. D., Murray J., Mills D., Berry D. S., 2004, *Starlink User Note*, 50
 Huggins P. J., Bachiller R., Planesas P., Forveille T., Cox P., 2005, *ApJS*, 160, 272
 Irabor T., et al., 2018, *MNRAS*, 480, 2423
 Isaacman R., 1984, *MNRAS*, 208, 399
 Kaler J. B., 1976, *ApJS*, 31, 517
 Kaler J. B., 1983a, *ApJ*, 264, 594
 Kaler J. B., 1983b, *ApJ*, 271, 188
 Kaler J. B., Lutz J. H., 1985, *PASP*, 97, 700
 Kaler J. B., Shaw R. A., Kwinter K. B., 1990, *ApJ*, 359, 392
 Kaler J. B., Shaw R. A., Browning L., 1997, *PASP*, 109, 289
 Karachentsev I. D., Karachentseva V. E., Huchtmeier W. K., 2001, *A&A*, 366, 428
 Kerber F., Mignani R. P., Guglielmetti F., Wicenec A., 2003, *A&A*, 408, 1029
 Kerton C. R., Brunt C. M., 2003, *A&A*, 399, 1083
 Kimeswenger S., Barría D., 2018, *A&A*, 616, L2
 Kwinter K. B., Henry R. B. C., 2001, *ApJ*, 562, 804
 Kwok S., Aaquist O. B., 1993, *PASP*, 105, 1456
 Lacy M., et al., 2020, *PASP*, 132, 035001
 Lallement R., Vergely J. L., Valette B., Puspitarini L., Eyer L., Casagrande L., 2014, *A&A*, 561, A91
 Lallement R., Babusiaux C., Vergely J. L., Katz D., Arenou F., Valette B., Hottier C., Capitanio L., 2019, *A&A*, 625, A135
 Lindegren L., et al., 2018, *A&A*, 616, A2
 Lopez J. A., Vazquez R., Rodriguez L. F., 1995, *ApJ*, 455, L63
 Loup C., Forveille T., Omont A., Paul J. F., 1993, *A&AS*, 99, 291
 Lumsden S. L., Hoare M. G., Urquhart J. S., Oudmaijer R. D., Davies B., Mottram J. C., Cooper H. D. B., Moore T. J. T., 2013, *ApJS*, 208, 11
 Lutz J. H., 1973, *ApJ*, 181, 135
 Mampaso A., et al., 2006, *A&A*, 458, 203
 Marshall D. J., Robin A. C., Reylé C., Schultheis M., Picaud S., 2006, *A&A*, 453, 635
 Milingo J. B., Kwinter K. B., Henry R. B. C., Souza S. P., 2010, *ApJ*, 711, 619
 Milne D. K., 1979, *A&AS*, 36, 227
 Milne D. K., Aller L. H., 1975, *A&A*, 38, 183
 O'Dell C. R., 1963, *ApJ*, 138, 1018
 Ofek E. O., Frail D. A., 2011, *ApJ*, 737, 45
 Osterbrock D. E., Stockhausen R. E., 1960, *ApJ*, 131, 310
 Parker Q. A., Acker A., Frew D. J., et al., 2006, *MNRAS*, 373, 79
 Parker Q. A., Bojićić I. S., Frew D. J., 2016, in *Journal of*

- Physics Conference Series. p. 032008 ([arXiv:1603.07042](https://arxiv.org/abs/1603.07042)), doi:10.1088/1742-6596/728/3/032008
- Pazderska B. M., Gawroński M. P., Feiler R., et al., 2009, A&A, 498, 463
- Peña M., 2005, Revista Mexicana de Astronomia y Astrofisica, 41, 423
- Peimbert M., Torres-Peimbert S., 1971, Boletin de los Observatorios Tonantzintla y Tacubaya, 6, 21
- Perek L., 1971, Bulletin of the Astronomical Institutes of Czechoslovakia, 22, 103
- Pollacco D. L., Bell S. A., 1997, MNRAS, 284, 32
- Quireza C., Rocha-Pinto H. J., Maciel W. J., 2007, A&A, 475, 217
- Raddi R., Drew J. E., Steeghs D., Wright N. J., Drake J. J., Barentsen G., Fabregat J., Sale S. E., 2015, MNRAS, 446, 274
- Rosado M., Kwitter K. B., 1982, Rev. Mex. Astron. Astrofis., 5, 217
- Sabin L., 2008, PhD thesis, School of Physics and Astronomy, University of Manchester
- Sabin L., Zijlstra A. A., Wareing C., Corradi R. L. M., Mam-paso A., Viironen K., Wright N. J., Parker Q. A., 2010, Publ. Astron. Soc. Australia, 27, 166
- Sabin L., et al., 2013, MNRAS, 431, 279
- Sabin L., et al., 2014, MNRAS, 443, 3388
- Sale S. E., 2012, MNRAS, 427, 2119
- Sale S. E., et al., 2009, MNRAS, 392, 497
- Sale S. E., et al., 2014, MNRAS, 443, 2907
- Schlafly E. F., Finkbeiner D. P., 2011, ApJ, 737, 103
- Schlegel D. J., Finkbeiner D. P., Davis M., 1998, ApJ, 500, 525
- Schwarz H. E., Monteiro H., 2006, ApJ, 648, 430
- Setia Gunawan D. Y. A., de Bruyn A. G., van der Hucht K. A., Williams P. M., 2003, ApJS, 149, 123
- Siódmiak N., Tylenda R., 2001, A&A, 373, 1032
- Skiff B. A., 2014, VizieR Online Data Catalog, p. B/mk
- Stanghellini L., Shaw R. A., Villaver E., 2008, ApJ, 689, 194
- Storey P. J., Hummer D. G., 1995, MNRAS, 272, 41
- Storey P. J., Zeippen C. J., 2000, MNRAS, 312, 813
- Turatto M., Cappellaro E., Sabbadin F., et al., 1990, AJ, 99, 1170
- Turner K. C., Terzian Y., 1984, AJ, 89, 501
- VTech 2014, VTSS: The Virginia Tech Spectral-Line Survey, <http://www.phys.vt.edu/~halph/>
- Viironen K., et al., 2009a, A&A, 502, 113
- Viironen K., et al., 2009b, A&A, 504, 291
- Vink J. S., Drew J. E., Steeghs D., Wright N. J., Martin E. L., Gänsicke B. T., Greimel R., Drake J., 2008, MNRAS, 387, 308
- Walsh J. R., Monreal-Ibero A., Barlow M. J., Ueta T., Wesson R., Zijlstra A. A., 2016, A&A, 588, A106
- Weidmann W. A., Díaz R. J., 2008, PASP, 120, 380
- Zhang Y., Liu X.-W., Luo S.-G., et al., 2005, A&A, 442, 249
- Zijlstra A. A., Pottasch S. R., Bignell C., 1989, A&AS, 79, 329

APPENDIX A: IPHAS H α FLUX MEASUREMENTS

A1 IPHAS H α filter fluxes, [N II]/H α ratios and corrected H α fluxes for IPHAS-discovered PNe

Table A1 presents the IPHAS H α fluxes measured for the IPHAS-discovered PNe in the sample. Columns (1) and (2) list the PN G Galactic coordinates and the IPHAS names for the nebulae. Column (3) lists the angular dimensions of the nebulae, measured at 10% of their peak brightnesses in the IPHAS H α filter images. Column (4) lists the IPHAS run numbers of the images that were used for the nebular aperture photometry, while column (5) lists the measured in-band H α fluxes, in cgs units. Columns (6)-(8) respectively list the logs of the individual fluxes, the log of the mean flux and its uncertainty. Column (9) lists the [N II]/H α flux ratios adopted from the literature, when available, with the letter in brackets next to each ratio representing the literature source, listed below. [N II]-corrected log H α fluxes are given in column (10).

The sources for the [N II]/H α flux ratios are: (a) Parker et al. (2006); (b) Viironen et al. (2009b); (c) Acker et al. (1991); (d) Sabin et al. (2014); (e) Sabin (2008); (f) Mam-paso et al. (2006).

A2 IPHAS H α filter fluxes, [N II]/H α ratios and corrected H α fluxes for previously catalogued PNe

Table A2 presents the IPHAS H α fluxes measured for the previously catalogued PNe. Columns (1) and (2) list PN G Galactic coordinates and catalogue names for the nebulae. Column (3) lists the angular dimensions of the nebulae, measured at 10% of their peak brightnesses in the IPHAS H α filter images. Column (4) lists the IPHAS run numbers of the images that were used for the nebular aperture photometry, while column (5) lists the measured in-band H α fluxes, in cgs units. Column (6) lists the [N II]/H α flux ratios adopted from the literature, with the letter in brackets next to each ratio representing the literature source, listed below. [N II]-corrected log H α fluxes are given in column (7), while columns (8) and (9) present the mean log H α fluxes (obtained by averaging the linear values of the [N II]-corrected fluxes), and their corresponding uncertainties, in dex.

The sources for the [N II]/H α flux ratios were: (a) Acker et al. (1991); (b) Frew et al. (2013); (c) Rosado & Kwitter (1982); (d) Kaler et al. (1997); (e) Milingo et al. (2010); (f) Guerrero et al. (1996); (g) Lopez et al. (1995); (h) Peña (2005); (i) Aller & Czyzak (1979); (j) Barker (1978); (k) Bohigas (2001); (l) Cantrell (2014); (m) Henry et al. (2010); (n) Kaler (1976); (o) Kaler (1983a); (p) Kwitter & Henry (2001); (q) Pollacco & Bell (1997); (r) Schwarz & Monteiro (2006); (s) Zhang et al. (2005); (t) Turatto et al. (1990), (u) Kaler (1983b).

APPENDIX B: H β AND RADIO FLUXES FOR THE SAMPLE PNE

The H β fluxes and the radio fluxes at 1.4 GHz, 5 GHz and 30 GHz, together with their uncertainties, are listed in Table B1. For PNe where fluxes at a given frequency were available from more than one source, we adopted a weighted average.

The H β fluxes are obtained from the following publications: (a) Kaler & Lutz (1985); (b) Kaler (1983b); (c) Collins et al. (1961); (d) Peimbert & Torres-Peimbert (1971); (e) Barker (1978); (f) O'Dell (1963); (g) Osterbrock & Stockhausen (1960); (h) Kaler (1983a); (i) Weidmann & Díaz (2008); (j) Capriotti & Daub (1960); (k) Loup et al. (1993); (l) Henry et al. (2010); (m) Kaler et al. (1990); (n) Carrasco et al. (1983); (o) Perek (1971); (p) Kaler (1976).

The references for the radio fluxes are as follows: (a) Pazderska et al. (2009); (b) Cahn et al. (1992); (c) Zijlstra et al. (1989); (d) Aaquist & Kwok (1990); (e) Siódmiak & Tylenda (2001); (f) Quireza et al. (2007); (g) Isaacman (1984); (h) Cahn & Rubin (1974); (i) Turner & Terzian (1984); (j) Higgs (1971); (k) Milne (1979); (l) Milne & Aller (1975); (m) Condon & Kaplan (1998); (n) Kerber et al. (2003); (o) DENIS Consortium (2003); (p) Ofek & Frail (2011); (q) Skiff (2014); (r) Huggins et al. (2005); (s) Lumsden et al. (2013); (t) Stanghellini et al. (2008); (u) Viironen et al. (2009a); (v) Setia Gunawan et al. (2003); (w) Kerton & Brunt (2003); (x) Karachentsev et al. (2001); (y) Kwok & Aaquist (1993); (z) Irabor et al. (2018).

APPENDIX C: E(H α -H β)-BASED C_{H β} AND A_V VALUES

Table C1 lists for 38 PNe the values of E(H α -H β), C_{H β} and A_V[E(H α -H β)], and their uncertainties, derived by comparing the measured H α fluxes with the literature H β filter fluxes listed in Table B1.

APPENDIX D: A_V VALUES FROM COMPARISONS OF BALMER-LINE AND RADIO FLUXES

Table D1 lists the individual values of A_V, and their uncertainties, that were derived from a comparison of the radio fluxes at 1.4 GHz, 5 GHz and 30 GHz (listed in Table B1) with the H α fluxes listed in Tables A1 and A2 and the H β fluxes listed in Table B1, using the relations given in Section 3.1. The extinction values used to obtain the weighted averaged A_{V(Radio)} values listed in the penultimate column of Table D1 are indicated in bold script in the table.

Table A1. H α +[N II] flux measurements for IPHAS-discovered PNe

PN G (1)	Name IPHAS (2)	Angular Diam ('') (3)	Run (4)	Filter F (5)	Flux (ergs cm $^{-2}$ s $^{-1}$) LogF (6)	Mean (7)	unc. (8)	[N II]/H α Ratio (9)	LogF(H α) (10)
Viironen et al. (2009a)									
044.4-04.1	J192717.94+081429.4	4.6	459511 459673	4.73E-14 4.52E-14	-13.33 -13.34	-13.33	0.01		
050.0+01.7	J191701.33+155947.8	5.0	476362 476365 527663	4.38E-14 4.23E-14 4.36E-14	-13.36 -13.37 -13.36	-13.36	0.01		
127.0-01.0	J012544.68+613611.6	5	367787 367900 367903	2.77E-14 2.79E-14 2.84E-14	-13.56 -13.55 -13.55	-13.55	0.01	0.16(c)	-13.62
181.4-03.7	J053440.77+254238.2	5	372878	8.82E-14	-13.05	-13.05	0.01	0.05(b)	-13.08
Sabin et al. (2014)									
029.5-00.2	XJ184616.3-030625	7	401522 401525	3.87E-14 4.03E-14	-13.41 -13.39	-13.40	0.01	0.71(d)	-13.63
029.9+03.7	XJ183249.6-005638	7	400650 400671	2.26E-13 2.23E-13	-12.65 -12.65	-12.65	0.01	0.51(d)	-12.83
032.3-04.5	XJ190631.5-023236	5	407121 407124	2.39E-15 2.54E-15	-14.62 -14.59	-14.61	0.01	0.49(d)	-14.78
038.9-01.3	XJ190718.1+044056	14.5 x 8	563823	3.34E-14	-13.48	-13.48		0.86(d)	-13.75
040.5+01.9	XJ185815.8+073753	11	407764 407797 407800	5.46E-14 5.36E-14 5.71E-14	-13.26 -13.27 -13.24	-13.26	0.01	1.01(d)	-13.56
040.7+03.4	XJ185322.1+083018 (Ear Nebula)	92 x 67	406710 406713	4.32E-13 4.24E-13	-12.36 -12.37	-12.37	0.01	0.00(d)	-12.37
043.3+03.5	XJ185744.4+105053	121 x 86	407458	3.02E-12	-11.52	-11.52		4.25(d)	-12.24
043.6+01.7	XJ190454.0+101801	23 x 24	410053 572482 572485	6.88E-13 6.31E-13 6.16E-13	-12.16 -12.20 -12.21	-12.19	0.01	1.24(d)	-12.54
043.8+02.1	XJ190338.5+104227	641	476759	8.18E-11	-10.09	-10.09			
044.2+01.9	XJ190518.3+105750	7.3	572470 572473	3.96E-14 3.84E-14	-13.40 -13.42	-13.41	0.01	2.13(d)	-13.91
045.4+02.6	J190447.9+121845	23 x 12	399198 399201	3.00E-13 3.00E-13	-12.52 -12.52	-12.52	0.01	1.87(d)	-12.98
048.1+01.4	XJ191421.1+140936	58 x 49	413381	4.51E-13	-12.35	-12.35	0.01		
048.2+01.9	J191255.5+143248	12.5	412044 413381 413384	4.90E-14 4.22E-14 4.55E-14	-13.31 -13.37 -13.34	-13.34	0.02	1.0(d)	-13.64
048.7-00.2	XJ192152.0+135223	18	571003	1.06E-13	-12.98	-12.98		1.83(d)	-13.43
048.9+04.3	XJ190512.4+161347	67	399234 399237	1.40E-12 1.39E-12	-11.85 -11.86	-11.86	0.01	0.28(d)	-11.97
049.2+00.0	J192153.9+143057	25 x 21	687803	7.97E-14	-13.10	-13.10		6.09(d)	-13.95
049.3-04.0	XJ193637.3+123808	317	459751	3.74E-12	-11.43	-11.43			
049.5-01.4	XJ192751.3+140127	14.5	418221 562419	1.07E-13 1.02E-13	-12.97 -12.99	-12.98	0.01	1.26(d)	-13.33
049.7-00.7	XJ192543.2+143546	281	562362	1.17E-10	-9.93	-9.93			
054.2-03.4	XJ194359.5+170901 (Necklace Nebula)	33 x 38	460367	1.55E-12	-11.81	-11.81	0.01	1.26(e)	-12.16
057.9-00.7	XJ194226.0+214521	36 x 23	460466 460469	2.04E-13 2.08E-13	-12.69 -12.68	-12.69	0.01	2.03(d)	-13.17
059.1+03.5	J192837.8+245025	20 x 12.5	414081	3.71E-13	-12.43	-12.43		0.81(d)	-12.69
059.7-01.0	XJ194727.6+230816	43	460682 460685	6.55E-13 6.67E-13	-12.18 -12.18	-12.18	0.01	1.04(d)	-12.49
062.9+01.3	J194510.7+270930	12.5	460538 460541	5.85E-14 5.68E-14	-13.23 -13.25	-13.24	0.01	2.82(d)	-13.82
063.3+02.2	XJ194240.6+275109	87	460436	2.71E-12	-11.57	-11.57		0.39(d)	-11.71
063.5+00.0	J195126.5+265839	9.9	461153 571498	8.62E-14 7.87E-14	-13.06 -13.10	-13.08	0.02	1.73(d)	-13.52
065.8+05.1	XJ193630.3+312810	392	398786	1.29E-11	-10.89	-10.89			
067.8+01.8	XJ195436.4+313326	14	756015 756018	4.20E-13 4.54E-13	-12.38 -12.34	-12.36	0.02	2.60(d)	-12.92
073.6+02.8	XJ200522.0+365942	12.5	570090 570093	1.30E-13 1.32E-13	-12.89 -12.88	-12.88	0.01	2.06(d)	-13.37

Table A1 – continued H α +[N II] flux measurements for IPHAS-discovered PNe

PN G (1)	Name IPHAS (2)	Angular Diam ('') (3)	Run (4)	Filter Flux (ergs cm $^{-2}$ s $^{-1}$)				[N II]/H α Ratio (9)	LogF(H α) (10)
				F (5)	LogF (6)	Mean (7)	unc. (8)		
077.0+03.6	J200018.6+365935	39	474781	2.86E-12	-11.54	-11.54	0.01	0.70(d)	-11.77
			474784	2.86E-12	-11.54				
114.2+03.7	J232713.0+650923	23	538647	4.41E-13	-12.36	-12.36	0.01	1.04(d)	-12.67
			538650	4.31E-13	-12.37				
114.4+00.0	XJ233841.2+614146	65 x 52	541345	1.18E-12	-11.93	-11.92	0.01	0.26(d)	-12.02
			541712	1.26E-12	-11.90				
			541757	1.17E-12	-11.93				
114.7-01.2	J234403.7+603242	29	527721	2.45E-13	-12.61	-12.60	0.01	1.00(d)	-12.90
			527724	2.54E-13	-12.59				
126.6+01.3	XJ012507.9+635652 (IPHAS PN-1)	44 x 30	367778	1.20E-12	-11.92	-11.93	0.01	3.20(f)	-12.55
			367781	1.15E-12	-11.94				
			367984	1.18E-12	-11.93				
			367897	1.21E-12	-11.92				
127.6-01.1	J013109.0+612259	30	368165	1.15E-13	-12.94	-12.97	0.03	0.00(d)	-12.97
			368168	1.01E-13	-13.00				
132.8+02.0	J022045.1+631135	44 x 32	373190	9.51E-12	-11.02	-11.01	0.01	0.21(d)	-11.09
			373193	1.00E-11	-11.00				
139.0+03.2	J031345.6+613710	129	431413	1.87E-12	-11.73	-11.73		0.58(d)	-11.93
144.1-00.5	J033105.4+553852	34	473618	1.05E-13	-12.98	-12.89	0.09	2.21(d)	-13.41
			473621	1.58E-13	-12.80				
150.0-00.3	J040329.6+520824	43 x 108	599667	1.77E-13	-12.75	-12.75		0.00(d)	-12.75
157.1+04.4	J045627.2+501722	108	485522	3.01E-12	-11.52	-11.52		0.90(d)	-11.80
193.5-03.1	J060328.2+154106	66 x 57	703212	2.75E-13	-12.56	-12.56		0.71(d)	-12.79
195.4-04.0	J060416.2+133251	40 x 17	476270	4.40E-13	-12.36	-12.35	0.01	3.53(d)	-13.01
			476273	4.47E-13	-12.35				

Table A2. [N II] corrections and final H α fluxes for catalogued PNe

PN G (1)	Name (2)	Angular Diam. ('') (3)	Run (4)	Filter Flux (ergs cm $^{-2}$ s $^{-1}$) (5)	[N II]/H α Ratio (6)	LogF(H α) (7)	Mean (8)	unc. (9)
030.8+03.4	Abell 47	15 × 10	400722	1.95E-13	0.60(a)	-12.91	-12.93	0.01
			400725	1.84E-13		-12.94		
031.0+04.1	K 3-6	0.7	400677	6.51E-13	0.20(b)	-12.27	-12.26	0.01
			400680	6.71E-13		-12.25		
031.3-00.5	HaTr 10	21 × 9.2	401663	1.47E-12	6.50(a)	-12.71	-12.71	0.01
			401666	1.44E-12		-12.72		
031.7+01.7	PC 20	2.0 × 1.3	463947	8.59E-13	1.10(a)	-12.39	-12.39	
032.0-03.0	K 3-18	1.0	568881	8.45E-13	0.50(a)	-12.25	-12.25	0.01
			568884	8.27E-13		-12.26		
032.5-03.2	K 3-20	4.0	568869	2.03E-12	0.90(a)	-11.97	-11.97	0.01
			568872	2.03E-12		-11.97		
032.9-02.8	K 3-19	0.7	406755	1.89E-12	0.50(a)	-11.90	-11.86	0.04
			464288	2.24E-12		-11.83		
033.8-02.6	NGC 6741	3.0	406776	5.36E-11	1.45(a)	-10.66	-10.67	0.01
			568875	5.16E-11		-10.68		
035.7-05.0	K 3-26	8 × 5	564360	1.29E-12	0.10(a)	-11.93	-11.93	0.01
			564363	1.29E-12		-11.93		
035.9-01.1	Sh 2-71	52 × 125	408156	7.23E-11	2.50(o,u)	-10.69	-10.73	0.05
			408159	5.86E-11		-10.78		
036.9-01.1	HaTr 11	9.7 × 10.4	478115	9.47E-13	1.70(a)	-12.46	-12.46	
036.9-02.6	HaTr 13	16.5 × 13.0	410137	8.00E-13	1.25(a)	-12.45	-12.43	0.01
			410419	8.56E-13		-12.42		
038.7+01.9	YM 16	271	407061	1.00E-11	5.20(a)	-11.79	-11.75	0.04
			407064	1.21E-11		-11.71		
039.5-02.7	M 2-47	1.3	750300	3.78E-12	0.10(a)	-11.46	-11.46	0.01
			750303	3.92E-12		-11.45		
039.8+02.1	K 3-17	4.9 × 4.3	407434	1.55E-12	0.92(a)	-12.09	-12.09	0.01
			407437	1.55E-12		-12.09		
040.4-03.1	K 3-30	0.7	412717	1.75E-12	0.10(a)	-11.80	-11.80	
041.8+04.4	K 3-15	0.7	402091	1.30E-12	0.27(a)	-11.99	-12.00	0.01
			402094	1.25E-12		-12.01		
043.0-03.0	M 4-14	6.7 × 24	569664	4.38E-12	2.36(a)	-11.89	-11.89	0.01
			569667	4.25E-12		-11.90		
			572512	4.38E-12		-11.89		
043.1+03.8	M 1-65	1.3	407458	7.34E-12	0.50(a)	-11.31	-11.31	0.01
			407503	7.38E-12		-11.31		
043.3+02.2	PM 1-276	14	408168	1.13E-12	0.00(a)	-11.95	-11.95	0.01
			408171	1.12E-12		-11.95		
045.9-01.9	K 3-33	0.6	687128	2.89E-13	0.90(a)	-12.82	-12.82	0.01
			687131	2.90E-13		-12.82		
046.3-03.1	PB 9	4.6	418230	2.27E-12	0.00(a)	-11.64	-11.65	0.01
			418233	2.21E-12		-11.66		
046.4-04.1	NGC 6803	3.0	419085	3.11E-11	0.4(b)	-10.65	-10.65	0.01
			419088	3.18E-11		-10.64		
046.8+02.9	CTSS 4	4.6	562802	3.68E-13	0.71(d,e)	-12.67	-12.66	0.01
			562805	3.82E-13		-12.65		
047.1+04.1	K 3-21	6.9	399123	7.96E-13	2.05(a)	-12.58	-12.58	0.01
			399168	8.13E-13		-12.57		
047.1-04.2	Abell 62	164	571857	2.12E-11	0.18(o,u)	-10.75	-10.76	0.02
			571860	1.95E-11		-10.78		
048.0-02.3	PB 10	1.3	418503	3.53E-12	0.20(b)	-11.53	-11.53	
048.1+01.1	K 3-29	0.7	562674	6.95E-13	0.70(b)	-12.39	-12.39	0.01
			562677	6.89E-13		-12.39		
048.5+04.2	K 4-16	1.3	399210	4.88E-13	0.00(b)	-12.31	-12.31	0.01
			399213	4.83E-13		-12.32		
048.7+01.9	Hen 2-429	2.7	413423	8.09E-12	0.90(a)	-11.37	-11.37	0.01
			413426	8.06E-12		-11.37		
048.7+02.3	K 3-24	4.6 × 1.4	412047	8.35E-13	2.10(a)	-12.57	-12.56	0.01
			412083	8.64E-13		-12.55		

Table A2 – continued [N II] corrections and final H α fluxes for catalogued PNe

PN G (1)	Name (2)	Angular Diam. ('') (3)	Run (4)	Filter Flux (ergs cm $^{-2}$ s $^{-1}$) (5)	[N II]/H α Ratio (6)	LogF(H α) (7)	Mean (8)	unc. (9)
048.76-01.53	DeHt 4	37 × 29	418158 418161 687110	9.08E-13 1.04E-12 7.06E-13	2.00(a)	-12.52 -12.46 -12.63	-12.54	0.05
049.4+02.4	Hen 2-428	7 × 22	562662 562665	4.49E-12 4.62E-12	0.62(a)	-11.56 -11.54	-11.55	0.01
050.4-01.6	K 4-28	0.1	418805	1.13E-13	0.15(a)	-13.01	-13.01	
051.0+03.0	Hen 2-430	0.7	413369 570931	4.20E-12 4.58E-12	0.63(b)	-11.59 -11.55	-11.57	0.02
051.0-04.5	PC 22	15 × 11	460170 460173	2.07E-12 2.63E-12	0.30(a)	-11.80 -11.69	-11.75	0.05
051.3+01.8	PM 1-295	1.0	528339 528342	1.21E-12 1.27E-12	0.51(t)	-12.09 -12.07	-12.08	0.01
051.9-03.8	M 1-73	2.3	460149 460152	1.90E-11 1.87E-11	0.54(b)	-10.91 -10.92	-10.91	0.01
052.2-04.0	M 1-74	1.3	460152 570946 570949	8.16E-12 9.69E-12 8.89E-12	0.20(a)	-11.17 -11.09 -11.13	-11.13	0.02
052.5-02.9	Me 1-1	3.7	460074	2.51E-11	0.94(e)	-10.90	-10.90	
052.9+02.7	K 3-31	0.7	571854	7.02E-13	0.43(a)	-12.31	-12.31	
052.9-02.7	K 3-41	0.7	460074	5.35E-13	0.00(a)	-12.27	-12.27	
053.2-01.5	K 3-38	3.0	457828 457831	8.22E-13 8.16E-13	0.00(a)	-12.09 -12.09	-12.09	0.01
053.8-03.0	Abell 63	19	562793 687734	2.52E-12 2.36E-12	0.01(q)	-11.61 -11.63	-11.62	0.01
054.4-02.5	M 1-72	1.3	625328	8.68E-12	0.64(a)	-11.28	-11.28	
055.1-01.8	K 3-43	7.6	686772 921493 921496	2.95E-13 3.35E-13 5.36E-13	0.07(d,e)	-12.56 -12.50 -12.30	-12.46	0.08
055.2+02.8	Hen 2-432	0.7	413703 413706 571261	1.94E-12 2.09E-12 2.19E-12	0.29(a)	-11.82 -11.79 -11.77	-11.79	0.02
055.3+02.7	Hen 1-1	3.2	413703 413706	1.94E-12 2.05E-12	1.42(a)	-12.10 -12.07	-12.08	0.01
055.5-00.5	M 1-71	1.0	570277 570280	1.14E-11 1.12E-11	0.47(b)	-11.11 -11.12	-11.11	0.01
055.6+02.1	Hen 1-2	1.6	413754 413775	3.19E-12 3.12E-12	0.83(d,e)	-11.76 -11.77	-11.76	0.01
056.0+02.0	K 3-35	0.7	413778 414090	2.08E-12 2.31E-12	4.04(a)	-12.38 -12.34	-12.36	0.02
056.4-00.9	K 3-42	0.7	921405 921408	2.69E-13 2.76E-13	0.27(a)	-12.67 -12.66	-12.67	0.01
057.9-01.5	Hen 2-447	0.7	460562 460565	2.60E-12 2.58E-12	0.56(b)	-11.78 -11.78	-11.78	0.01
058.9+01.3	K 3-40	2.0	420440 420443	1.80E-12 1.85E-12	0.18(a)	-11.82 -11.81	-11.81	0.01
059.0+04.6	K 3-34	8 × 5	413733 571246 571249	1.18E-12 1.26E-12 1.28E-12	2.50(a)	-12.47 -12.44 -12.44	-12.45	0.01
059.4+02.3	K 3-37	1.3	420097 420100	5.06E-13 5.14E-13	0.03(a)	-12.31 -12.30	-12.30	0.01
059.9+02.0	K 3-39	0.7	570229 570232	6.29E-13 6.36E-13	0.53(a)	-12.39 -12.38	-12.38	0.01
060.4+01.5	PM 1-310	0.7	421140 922041	9.73E-14 1.18E-13	1.04(h)	-13.32 -13.24	-13.28	0.04
060.5+01.8	Hen 2-440	0.7	421136 421140	2.82E-12 2.78E-12	0.57(a)	-11.75 -11.75	-11.75	0.01
060.5-00.3	K 3-45	3.9 × 0.7	572264 572267	1.68E-13 1.63E-13	2.02(a)	-13.25 -13.27	-13.26	0.01
060.8-03.6	NGC 6853	331 × 472	571657	2.57E-09	1.30(a)	-8.95	-8.95	

Table A2 – continued [N II] corrections and final H α fluxes for catalogued PNe

PN G (1)	Name (2)	Angular Diam. ('') (3)	Run (4)	Filter Flux (ergs cm $^{-2}$ s $^{-1}$) (5)	[N II]/H α Ratio (6)	LogF(H α) (7)	Mean (8)	unc. (9)
062.4-00.2	M 2-48	1.8×1.5	461361	6.54E-12	3.60(a)	-11.85	-11.85	0.01
			461364	6.40E-12		-11.86		
063.8-03.3	K 3-54	0.7	563008	3.38E-13	0.84(a)	-12.74	-12.74	0.01
			563011	3.38E-13		-12.74		
064.9-02.1	K 3-53	0.7	408279	1.16E-12	0.19(a)	-12.01	-12.01	0.01
			570256	1.16E-12		-12.01		
065.1-03.5	We 1-9	23×19	563071	1.29E-12	0.89(a)	-12.17	-12.11	0.06
			750375	1.70E-12		-12.05		
065.9+00.5	NGC 6842	52.4	474763	8.58E-12	0.07(a)	-11.09	-11.09	0.01
			572390	8.73E-12		-11.09		
066.9+02.2	K 4-37	1.0	399436	6.59E-13	6.59(a)	-13.06	-13.05	0.01
			399523	6.96E-13		-13.04		
067.9-00.2	K 3-52	0.7	408634	5.27E-13	2.09(a)	-12.77	-12.77	0.01
			475209	5.31E-13		-12.76		
068.3-02.7	Hen 2-459	0.7	412200	5.85E-12	1.00(b,d,e)	-11.53	-11.55	0.01
			412203	5.59E-12		-11.55		
			413073	5.32E-12		-11.57		
068.6+01.1	Hen 1-4	22×20	407192	4.99E-12	1.15(l)	-11.63	-11.64	0.01
			407195	4.93E-12		-11.64		
068.7+01.9	K 4-41	1.7	686283	9.25E-13	0.07(a)	-12.06	-12.06	
068.7+03.0	PC 23	0.7	399511	1.46E-12	0.31(a)	-11.95	-11.95	0.01
			399514	1.45E-12		-11.96		
068.8-00.0	M 1-75	17×9	410146	6.65E-12	5.11(d,e,k,l)	-11.96	-11.97	0.01
			410149	6.57E-12		-11.97		
069.2+02.8	K 3-49	0.7	399544	8.05E-13	0.80(a)	-12.35	-12.35	0.01
			570078	8.15E-13		-12.34		
069.2+03.8	K 3-46	26×12	399415	5.23E-12	8.53(a)	-12.26	-12.26	0.01
			399478	4.91E-12		-12.29		
			534190	5.53E-12		-12.24		
069.4-02.6	NGC 6894	37	530437	3.99E-11	0.78(o,u)	-10.65	-10.64	0.01
			530440	4.19E-11		-10.63		
069.6-03.9	K 3-58	9×6	750399	3.24E-12	1.38(a)	-11.87	-11.86	0.01
			750426	3.38E-12		-11.85		
069.7+00.0	K 3-55	5.3	410760	9.79E-13	1.09(d,e)	-12.33	-12.33	0.01
			410763	9.90E-13		-12.32		
071.6-02.3	M 3-35	0.7	455655	5.66E-12	0.03(b)	-11.26	-11.21	0.03
			750405	6.61E-12		-11.19		
			750408	6.84E-12		-11.18		
072.1+00.1	K 3-57	3.0×4.6	412119	1.67E-12	0.59(a)	-11.98	-11.97	0.01
			412511	1.73E-12		-11.96		
073.0-02.4	K 3-76	1.7	459527	4.93E-13	0.00(a)	-12.31	-12.32	0.02
			572330	4.59E-13		-12.34		
074.5+02.1	NGC 6881	1.2×1.1	475983	1.04E-11	0.70(o,u)	-11.21	-11.21	0.01
075.6+04.3	ARO 342	23	471338	9.62E-13	1.41(l)	-12.40	-12.41	0.01
			569457	9.29E-13		-12.41		
			569460	8.92E-13		-12.43		
076.3+01.1	Abell 69	19.1	531486	1.38E-12	5.20(a)	-12.65	-12.65	0.01
			531489	1.39E-12		-12.65		
076.4+01.8	KjPn 3	0.7	568992	1.65E-13	0.00(a)	-12.78	-12.79	0.01
			568995	1.63E-13		-12.79		
077.5+03.7	KjPn 1	1.0	472088	9.45E-13	0.17(a)	-12.09	-12.07	0.03
			472091	9.05E-13		-12.11		
			473554	1.14E-12		-12.01		
077.7+03.1	KjPn 2	1.6×1.1	569004	3.36E-13	1.36(a)	-12.85	-12.85	0.01
			569007	3.32E-13		-12.85		
078.3-02.7	K 4-53	9.5×14.5	366816	1.45E-12	0.62(l)	-12.05	-12.06	0.02
			366867	1.35E-12		-12.08		
078.9+00.7	Sd 1	10	414455	1.12E-12	0.86(a)	-12.22	-12.22	
084.2+01.0	K 4-55	7×9	367619	3.94E-12	5.40(f,k,l)	-12.21	-12.21	
084.9+04.4	Abell 71	108	369201	3.28E-11	1.50(o,u)	-10.88	-10.91	0.02
			369834	2.96E-11		-10.93		

Table A2 – continued [N II] corrections and final H α fluxes for catalogued PNe

PN G (1)	Name (2)	Angular Diam. ('') (3)	Run (4)	Filter Flux (ergs cm $^{-2}$ s $^{-1}$) (5)	[N II]/H α Ratio (6)	LogF(H α) (7)	Mean (8)	unc. (9)
088.7+04.6	K 3-78	1.7	369543 370351 370354	4.81E-13 4.44E-13 4.64E-13	0.00(a)	-12.32 -12.35 -12.33	-12.33	0.01
088.7-01.6	NGC 7048	63	419174 419177	5.78E-11 5.90E-11	2.24(l)	-10.75 -10.74	-10.74	0.01
089.0+00.3	NGC 7026	25 × 8	411588 411591	9.37E-11 9.28E-11	0.62(b,l,o)	-10.24 -10.24	-10.24	0.01
089.8-00.6	Sh 1-89	52 × 9	460260	3.53E-12	3.64(a)	-12.12	-12.12	
091.6+01.8	We 1-11	23 × 22	412607 412610	1.26E-12 1.23E-12	2.00(a)	-12.38 -12.39	-12.38	0.01
091.6-04.8	K 3-84	3.6	463449 463452	2.96E-12 2.92E-12	2.05(a)	-12.01 -12.02	-12.02	0.01
093.3-00.9	K 3-82	23	530076 756036	2.19E-12 2.77E-12	0.29(a)	-11.77 -11.67	-11.72	0.05
093.3-02.4	M 1-79	24 × 42	756069 756072	2.66E-11 2.53E-11	1.07(o,u)	-10.89 -10.91	-10.92	0.01
094.5-00.8	K 3-83	4.2	750456 750459 756042	4.85E-13 5.01E-13 4.90E-13	2.82(a)	-12.90 -12.88 -12.89	-12.89	0.01
095.1-02.0	M 2-49	0.7	419822 419825	2.66E-12 2.82E-12	0.79(d,e)	-11.83 -11.80	-11.82	0.01
095.2+00.7	K 3-62	1.3	647763 647766	1.81E-12 1.86E-12	0.26(d,e)	-11.84 -11.83	-11.84	0.01
096.3+02.3	K 3-61	5.6	413465 413468	9.42E-13 8.95E-13	0.10(a)	-12.07 -12.09	-12.08	0.01
097.6-02.4	M 2-50	2.0	421326 421329	1.40E-12 1.41E-12	0.00(a)	-11.85 -11.85	-11.85	0.01
098.1+02.4	K 3-63	6.0 × 1.3	419524 478931 478934	7.04E-13 8.71E-13 8.54E-13	0.12(o,u)	-12.20 -12.11 -12.12	-12.14	0.03
098.2+04.9	K 3-60	1.0	413124 413127	9.43E-13 9.79E-13	0.19(o,u)	-12.10 -12.09	-12.09	0.01
102.8-05.0	Abell 80	144 × 106	464077 464082	1.63E-11 1.75E-11	1.45(o,u)	-11.18 -11.15	-11.16	0.02
103.2+00.6	M 2-51	47 × 40	537395	2.40E-11	1.49(o,u)	-11.02	-11.02	0.01
103.7+00.4	M 2-52	7.3 × 9.5	462315 462318	6.42E-12 6.27E-12	3.36(k,l)	-11.83 -11.84	-11.84	0.01
104.1+01.0	Bl 2-1	0.7	462351 462354 599592 599595	7.29E-13 7.31E-13 7.40E-13 7.34E-13	0.17(a)	-12.21 -12.20 -12.20 -12.20	-12.20	0.01
104.4-01.6	M 2-53	12 × 8	597700 597703	6.12E-12 5.85E-12	2.12(l)	-11.71 -11.73	-11.72	0.01
107.4-00.6	K 4-57	0.7	570770	1.26E-13	0.06(a)	-12.92	-12.92	
107.4-02.6	K 3-87	2.6	418634 570442	2.97E-13 3.58E-13	0.09(a)	-12.56 -12.48	-12.52	0.04
107.7-02.2	M 1-80	5.3	418634 418862	6.99E-12 7.27E-12	0.61(a)	-11.36 -11.35	-11.35	0.01
107.8+02.3	NGC 7354	18	374924 374972 375594	2.81E-11 2.79E-11 2.65E-11	0.27(b)	-10.65 -10.66 -10.68	-10.66	0.01
112.5+03.7	K 3-88	7 × 8	464459 464752 464755	3.31E-13 3.43E-13 3.49E-13	1.22(a)	-12.83 -12.81 -12.80	-12.81	0.01
112.5-00.1	KjPn 8	3.3	475765 476488	1.49E-12 1.26E-12	3.80(g)	-12.51 -12.58	-12.54	0.03
119.3+00.3	BV 1	31 × 11	414171 414515 414578	7.85E-12 7.57E-12 7.79E-12	4.16(k,l)	-11.82 -11.83 -11.82	-11.82	0.01
121.6+03.5	We 1-1	20	417705 418383 418386	4.74E-13 4.68E-13 4.76E-13	1.07(a)	-12.64 -12.65 -12.64	-12.64	0.01
121.6+00.0	BV 2	3.0 × 8.3	418653 418656 922235	2.53E-12 2.50E-12 2.69E-12	1.05(l,m)	-11.91 -11.91 -11.88	-11.91	0.01

Table A2 – continued [N II] corrections and final H α fluxes for catalogued PNe

PN G (1)	Name (2)	Angular Diam. ('') (3)	Run (4)	Filter Flux (ergs cm $^{-2}$ s $^{-1}$) (5)	[N II]/H α Ratio (6)	LogF(H α) (7)	Mean (8)	unc. (9)
122.1-04.9	Abell 2	27	419262	2.08E-12	0.27(o,u)	-11.79	-11.79	0.01
			419265	2.08E-12		-11.79		
126.3+02.9	K 3-90	6	367533	1.04E-12	0.00(l)	-11.98	-12.01	0.01
			703469	9.58E-13		-12.02		
			703472	9.38E-13		-12.03		
128.0-04.1	Simeiz 22	351 × 48	368180	7.39E-11	5.00(c)	-10.91	-10.87	0.04
			368197	8.86E-11		-10.83		
129.5+04.5	K 3-91	3.6 × 3.0	370422	5.63E-13	2.09(a)	-12.74	-12.75	0.01
			370425	5.48E-13		-12.75		
130.2+01.3	IC 1747	8.0	370158	1.87E-11	0.089(m)	-10.77	-10.77	0.01
			370416	1.80E-11		-10.78		
130.4+03.1	K 3-92	7.8 × 3.3	364766	9.09E-13	1.12(k,l)	-12.37	-12.37	0.01
			370490	8.80E-13		-12.38		
			370493	9.03E-13		-12.37		
131.5+02.6	Abell 3	54	371560	4.29E-12	0.26(o,u)	-11.47	-11.48	0.01
			372310	4.03E-12		-11.50		
			372313	4.32E-12		-11.47		
132.4+04.7	K 3-93	3.0	373840	4.72E-13	1.60(d,e)	-12.74	-12.74	0.01
			373843	4.70E-13		-12.74		
136.1+04.9	Abell 6	166	380703	2.32E-12	0.13(o,u)	-11.69	-11.62	0.07
			380706	3.21E-12		-11.55		
138.8+02.8	IC 289	25	431332	1.47E-11	0.01(b,l)	-10.84	-10.83	0.01
			431335	1.49E-11		-10.83		
142.1+03.4	K 3-94	6.6	474115	1.71E-12	1.72(d,e,k)	-12.20	-12.21	0.01
			474118	1.67E-12		-12.21		
147.4-02.3	M 1-4	2.7	376435	5.04E-12	0.05(b)	-11.32	-11.36	0.02
			428921	4.32E-12		-11.39		
			428924	4.43E-12		-11.38		
147.8+04.1	M 2-2	4.3	482053	6.19E-12	0.17(o,u)	-11.28	-11.27	0.01
			482056	6.52E-12		-11.25		
149.0+04.4	K 4-47	8.3 × 0.9	483031	2.92E-13	2.35(d,e)	-13.06	-13.07	0.01
			483034	2.84E-13		-13.07		
151.4+00.5	K 3-64	9.2	479903	7.96E-13	0.60(a)	-12.30	-12.32	0.01
			479906	7.91E-13		-12.31		
			482095	7.29E-13		-12.34		
153.7-01.4	K 3-65	2.3	538727	3.24E-13	1.25(a)	-12.84	-12.85	0.01
			538730	3.13E-13		-12.86		
160.5-00.5	We 1-2	96	471195	2.36E-12	2.57(d,e)	-12.18	-12.23	0.05
			471198	1.84E-12		-12.29		
163.1-00.8	We 1-3	72	480495	1.10E-11	2.94(d,e)	-11.56	-11.56	
173.5+03.2	Pu 2	17	597407	2.75E-13	0.00(a)	-12.56	-12.56	
178.3-02.5	K 3-68	9.3	372462	6.07E-13	0.00(a)	-12.22	-12.21	0.01
			372465	6.19E-13		-12.21		
181.5+00.9	Pu 1	63	381282	1.79E-12	1.01(a)	-12.05	-12.06	0.01
			381285	1.69E-12		-12.08		
184.0-02.1	M 1-5	1.0	379317	9.58E-12	0.45(b,j)	-11.18	-11.19	0.01
			379320	9.32E-12		-11.19		
184.6+00.6	K 3-70	0.7 × 0.7	381765	1.06E-12	3.60(d,e)	-12.64	-12.64	0.01
			381768	1.04E-12		-12.65		
184.8+04.4	K 3-71	1.7	485492	3.79E-13	0.03(d,e)	-12.44	-12.44	
194.2+02.5	J 900	2.7	545436	3.08E-11	0.18(o,u)	-10.58	-10.58	0.01
			545439	3.09E-11		-10.58		
197.8-03.3	Abell 14	35 × 18.0	477563	1.77E-12	6.20(a)	-12.61	-12.61	0.01
			478381	1.79E-12		-12.60		
201.7+02.5	K 4-48	1.0	603928	2.05E-12	0.96(a)	-11.98	-11.97	0.01
			603931	2.14E-12		-11.96		
201.9-04.6	We 1-4	12.6 × 33	478856	1.81E-12	5.35(k)	-12.54	-12.52	0.03
			478859	2.07E-12		-12.49		
210.0+03.9	We 2-34	222	597464	8.52E-12	4.00(a)	-11.77	-11.77	
210.3+01.9	M 1-8	6.7 × 2.7	599898	7.05E-12	1.46(a)	-11.54	-11.54	
212.0+04.3	M 1-9	0.7	599461	9.43E-12	0.30(a)	-11.14	-11.14	

Table B1. H β fluxes and radio fluxes at 1.4, 5 and 30 GHz

PN G	Name	logF(H β) ergs cm $^{-2}$ s $^{-1}$			S $_{\nu}$ (30 GHz)			S $_{\nu}$ (5 GHz)			S $_{\nu}$ (1.4 GHz)		
		unc.	ref.	mJy	unc.	ref.	mJy	unc.	ref.	mJy	unc.	ref.	
030.8+03.4	Abell 47	—	—	—	—	—	55	11	(d)	7.9	0.6	(m)	
031.0+04.1	K 3-6	—	—	98.6	8.3	(a)	—	—	—	10.9	0.6	(m)	
031.3-00.5	HaTr 10	—	—	—	—	—	—	—	—	5	0.5	(n)	
031.7+01.7	PC 20	—	—	—	—	—	—	—	—	27.4	0.9	(m)	
032.0-03.0	K 3-18	—	—	—	—	—	11	1	(c)	—	—	—	
032.5-03.2	K 3-20	—	—	—	—	—	—	—	—	6.3	0.6	(o)	
032.9-02.8	K 3-19	—	—	—	—	—	22.5	0.06	(y)	12.4	0.7	(m)	
033.8-02.6	NGC 6741	-11.34	0.01	(a)	—	—	197	10	(g, i)	132.8	4.8	(m)	
035.7-05.0	K 3-26	—	—	—	—	—	—	—	—	6.8	0.5	(m)	
035.9-01.1	Sh 2-71	-11.6	0.05	(b)	—	—	66	7	(m)	57.9	3.1	(m)	
036.9-02.6	HaTr 13	—	—	—	—	—	—	—	—	3.6	0.5	(p)	
036.9-01.1	HaTr 11	—	—	—	—	—	—	—	—	17	0.7	(p)	
039.5-02.7	M 2-47	—	—	—	—	—	49	6	(c, h)	39.1	1.7	(m)	
039.8+02.1	K 3-17	—	—	303	9	(a)	398	43	(c, h)	320	9.6	(m)	
040.3-00.4	Abell 53	-12.61	0.05	(b)	54.9	6.5	(a)	76	10	(b)	34.1	1.1	(m)
040.4-03.1	K 3-30	—	—	—	—	—	23	5	(d)	12.7	0.6	(m)	
041.8+04.4	K 3-15	—	—	—	—	—	6.5	0.1	(y)	—	—	—	
043.0-03.0	M 4-14	—	—	—	—	—	—	—	—	11.9	0.6	(m)	
043.1+03.8	M 1-65	—	—	—	—	—	23	1.2	(d, i)	21.3	0.8	(m)	
043.3+02.2	PM 1-276	—	—	—	—	—	—	—	—	15.3	0.6	(q)	
045.9-01.9	K 3-33	—	—	—	—	—	17	3	(d)	9.1	0.5	(m)	
046.3-03.1	PB 9	—	—	—	—	—	40	8	(d)	33.6	1.1	(m)	
046.4-04.1	NGC 6803	-11.18	0.01	(c, p, d)	83.8	6.3	(a)	96.6	7.1	(g, i, l)	69.4	2.1	(m)
046.8+02.9	CTSS 4	—	—	—	—	—	—	—	—	2.5	0.3	(n, m)	
047.1+04.1	K 3-21	—	—	—	—	—	—	—	—	2.5	0.3	(r, m)	
047.1-04.2	Abell 62	—	—	—	—	—	—	—	—	—	—	—	
048.0-02.3	PB 10	—	—	37.1	6.5	(a)	50	10	(d)	50.7	1.6	(m)	
048.1+01.1	K 3-29	—	—	71.2	5.8	(a)	69.1	0.06	(y)	14.2	0.6	(m)	
048.5+04.2	K 4-16	—	—	—	—	—	3	1	(d)	5.2	0.5	(m)	
048.7+02.3	K 3-24	—	—	—	—	—	26	2.6	(h)	9	0.5	(m)	
048.7+01.9	Hen 2-429	—	—	62.1	6.1	(a)	69	6.9	(b)	59	1.8	(m)	
049.4+02.4	Hen 2-428	—	—	—	—	—	—	—	—	28.2	1.3	(m)	
050.4-01.6	K 4-28	—	—	—	—	—	19	4	(d)	5.4	0.5	(m)	
051.0-04.5	PC 22	—	—	—	—	—	12	1.2	(b)	9.7	0.5	(m)	
051.0+03.0	Hen 2-430	—	—	37.7	6.7	(a)	38.5	4	(g)	18.6	0.7	(m)	
051.3+01.8	PM 1-295	—	—	—	—	—	—	—	—	13.9	0.6	(q)	
051.9-03.8	M 1-73	—	—	36.5	4.8	(a)	43	5	(c, g)	48.1	1.5	(m)	
052.2-04.0	M 1-74	-11.75	0.02	(e)	—	—	31	4	(c)	7.2	0.7	(m)	
052.5-02.9	Me 1-1	-11.94	0.03	(f)	—	—	43	5	(c)	36.3	1.2	(m)	
052.9-02.7	K 3-41	—	—	—	—	—	1.7	0.3	(d)	4.7	0.9	(m)	
052.9+02.7	K 3-31	—	—	—	—	—	39	8	(d)	17.3	0.7	(m)	
053.2-01.5	K 3-38	—	—	—	—	—	25.2	2.5	(g)	29.2	1.1	(m)	
053.8-03.0	Abell 63	—	—	—	—	—	—	—	—	4.9	0.5	(n)	
054.2-03.4	Necklace	—	—	—	—	—	—	—	—	5	0.5	(m)	
054.4-02.5	M 1-72	—	—	—	—	—	26	5	(d)	5.7	0.5	(m)	
055.1-01.8	K 3-43	—	—	—	—	—	—	—	—	5.7	0.9	(m)	
055.2+02.8	Hen 2-432	—	—	—	—	—	33.5	5	(d, h, g)	23.2	0.9	(m)	
055.3+02.7	Hen 1-1	—	—	—	—	—	14	1.4	(e, c)	17.3	0.7	(m)	
055.5-00.5	M 1-71	—	—	178.9	6.3	(a)	203.5	13.8	(c, z)	—	—	—	
055.6+02.1	Hen 1-2	—	—	—	—	—	15	1.5	(e)	16.1	1.1	(m)	
056.0+02.0	K 3-35	—	—	60.1	6.4	(a)	40	8	(d)	14.6	0.6	(m)	
056.4-00.9	K 3-42	—	—	—	—	—	18.6	1.7	(z)	11.6	0.8	(m)	
057.9-01.5	Hen 2-447	—	—	65.5	7.1	(a)	62.3	0.06	(y)	23.6	1.2	(m)	
058.9+01.3	K 3-40	—	—	—	—	—	20	4	(d)	17.2	0.7	(m)	
059.4+02.3	K 3-37	—	—	—	—	—	17	3	(d)	14.7	0.6	(m)	
059.0+04.6	K 3-34	—	—	—	—	—	—	—	—	2.9	0.5	(r)	
059.9+02.0	K 3-39	—	—	—	—	—	11.6	0.10	(y)	3.7	0.5	(m)	
060.4+01.5	PM 1-310	—	—	—	—	—	—	—	—	2.5	0.3	(s, m)	
060.5-00.3	K 3-45	—	—	—	—	—	—	—	—	2.5	0.3	(r, m)	
060.5+01.8	Hen 2-440	—	—	—	—	—	40.6	0.06	(y)	26.9	0.9	(m)	
060.8-03.6	NGC 6853	-9.46	0.06	(g)	—	—	1325	19	(m)	—	—	—	

Table B1 – continued H β fluxes and radio fluxes at 1.4, 5 and 30 GHz

PN G	Name	logF(H β) ergs cm $^{-2}$			S $_{\nu}$ (30 GHz)			S $_{\nu}$ (5 GHz)			S $_{\nu}$ (1.4 GHz)		
		s $^{-1}$	unc.	ref.	mJy	unc.	ref.	mJy	unc.	ref.	mJy	unc.	ref.
062.4-00.2	M 2-48	–	–	–	–	–	–	19.6	1.4	(c, z)	17	0.7	(m)
063.8-03.3	K 3-54	–	–	–	–	–	–	7.2	0.9	(d, g)	5.1	0.5	(m)
064.9-02.1	K 3-53	–	–	–	73.8	5.3	(a)	50	10	(d)	10.3	0.5	(m)
065.1-03.5	We 1-9	–	–	–	–	–	–	–	–	–	9.9	0.5	(m)
065.9+00.5	NGC 6842	-11.73	0.03	(f)	–	–	–	127	12.7	(j)	43.9	2.1	(m)
066.9+02.2	K 4-37	–	–	–	–	–	–	–	–	–	2.5	0.3	(r, m)
067.9-00.2	K 3-52	–	–	–	90.2	9.9	(a)	65	13	(d)	17.3	0.7	(m)
068.3-02.7	Hen 2-459	–	–	–	102.5	6.9	(a)	86.6	0.08	(y)	14	0.6	(m)
068.6+01.1	Hen 1-4	–	–	–	–	–	–	–	–	–	16.2	0.7	(n)
068.7+01.9	K 4-41	–	–	–	–	–	–	15	3	(d)	11.8	0.6	(m)
068.7+03.0	PC 23	–	–	–	–	–	–	18	2	(c, d)	11	1.2	(m)
068.8-00.0	M 1-75	–	–	–	–	–	–	26	2.6	(h)	37	1.2	(m)
069.2+03.8	K 3-46	–	–	–	–	–	–	–	–	–	3.3	0.5	(r)
069.2+02.8	K 3-49	–	–	–	–	–	–	5.9	0.05	(y)	–	–	–
069.4-02.6	NGC 6894	-11.41	0.05	(b, c)	–	–	–	61	7	(c)	–	–	–
069.6-03.9	K 3-58	–	–	–	–	–	–	31	3	(c)	5.7	0.5	(m)
069.7+00.0	K 3-55	–	–	–	76.9	7	(a)	90	18	(d)	87.1	2.6	(m)
071.6-02.3	M 3-35	-12.45	0.06	(e)	161.8	6.4	(a)	150	28	(d, i)	29.9	1	(m)
072.1+00.1	K 3-57	–	–	–	–	–	–	60	12	(d)	47.3	1.5	(m)
073.0-02.4	K 3-76	–	–	–	–	–	–	12	2	(d)	–	–	–
074.5+02.1	NGC 6881	-12.26	0.03	(h)	109.1	7.4	(a)	110	0.08	(y)	70.9	2.2	(m)
075.6+04.3	ARO 342	–	–	–	–	–	–	–	–	–	2.5	0.3	(n, m)
076.3+01.1	Abell 69	-14.1	0.2	(i)	–	–	–	5	0.5	(t, m)	2.5	0.3	(n, m)
076.4+01.8	KjPn 3	–	–	–	–	–	–	–	–	–	2.5	0.3	(n, m)
077.5+03.7	KjPn 1	–	–	–	–	–	–	2.8	0.2	(u)	–	–	–
077.7+03.1	KjPn 2	–	–	–	–	–	–	–	–	–	10	1	(r)
078.3-02.7	K 4-53	–	–	–	44.4	6.3	(a)	–	–	–	35.1	1.5	(m)
078.9+00.7	Sd 1	–	–	–	–	–	–	27	2.7	(m)	2.5	0.3	(v, m)
084.2+01.0	K 4-55	–	–	–	–	–	–	–	–	–	6.6	0.6	(r)
084.9+04.4	Abell 71	-11.75	0.02	(i)	–	–	–	83	8.3	(t, m)	5	0.5	(f, m)
088.7+04.6	K 3-78	–	–	–	–	–	–	17	2	(c, d)	15.2	0.6	(m)
088.7-01.6	NGC 7048	-11.41	0.01	(f)	–	–	–	37	4	(c)	46.8	2.1	(m)
089.0+00.3	NGC 7026	-10.9	0.02	(a, j)	208.2	9.5	(a)	277	77	(j)	267	9.5	(m)
089.8-00.6	Sh 1-89	–	–	–	–	–	–	–	–	–	14.7	1.6	(m)
091.6-04.8	K 3-84	–	–	–	–	–	–	–	–	–	3	0.4	(r)
091.6+01.8	We 1-11	–	–	–	–	–	–	–	–	–	13.3	0.6	(m)
093.3-00.9	K 3-82	–	–	–	–	–	–	30	3	(c)	37.3	1.6	(m)
093.3-02.4	M 1-79	-11.73	0.01	(b)	–	–	–	19	1.9	(c)	23.6	1.4	(m)
094.5-00.8	K 3-83	–	–	–	–	–	–	6.5	0.7	(c)	5.1	0.5	(m)
095.1-02.0	M 2-49	–	–	–	–	–	–	74	8	(d, g)	28.3	1	(m)
095.2+00.7	K 3-62	–	–	–	101.3	7.8	(a)	118	0.10	(y)	59.9	1.9	(m)
096.3+02.3	K 3-61	–	–	–	–	–	–	14	1.4	(c)	16.9	0.7	(m)
097.6-02.4	M 2-50	-12.48	0.03	(e)	–	–	–	6.5	0.7	(c)	9.1	1.1	(m)
098.1+02.4	K 3-63	-13.97	0.06	(b)	–	–	–	29	2.9	(c)	8.6	0.5	(m)
098.2+04.9	K 3-60	-13.17	0.18	(b)	–	–	–	43	9	(d)	28.1	0.9	(m)
102.8-05.0	Abell 80	-11.8	0.12	(b)	–	–	–	3	0.3	(b)	9.4	2	(m)
103.2+00.6	M 2-51	-11.97	0.04	(k)	–	–	–	41	4.1	(t, m)	28.4	1.7	(w)
103.7+00.4	M 2-52	–	–	–	–	–	–	14	1.4	(c)	15.4	0.6	(m)
104.1+01.0	B1 2-1	–	–	–	–	–	–	61.2	0.11	(y)	22	0.8	(m)
104.4-01.6	M 2-53	–	–	–	–	–	–	11	1.1	(c)	15.1	0.6	(m)
107.4-02.6	K 3-87	–	–	–	–	–	–	4.5	0.5	(c)	11.5	1.1	(m)
107.4-00.6	K 4-57	–	–	–	–	–	–	–	–	–	2.5	0.3	(n, m)
107.7-02.2	M 1-80	–	–	–	–	–	–	25	3	(c)	18	0.7	(m)
107.8+02.3	NGC 7354	-11.58	0.01	(a, f)	439	13	(a)	597	76	(j)	582	23	(m)
112.5-00.1	KjPn 8	–	–	–	–	–	–	1	0	(m)	2.5	0.3	(n, m)
112.5+03.7	K 3-88	–	–	–	–	–	–	–	–	–	5.8	0.9	(m)
119.3+00.3	BV 5-1	-12.7	0.03	(h)	–	–	–	–	–	–	11.7	1.2	(m)
121.6+00.0	BV 5-2	–	–	–	–	–	–	–	–	–	9.3	1.4	(m)
121.6+03.5	We 1-1	–	–	–	–	–	–	–	–	–	5.6	0.5	(m)
122.1-04.9	Abell 2	-12.37	0.03	(b)	–	–	–	2.3	0.2	(c)	9.4	1.1	(m)

Table B1 – continued H β fluxes and radio fluxes at 1.4, 5 and 30 GHz

PN G	Name	logF(H β)		S $_{\nu}$ (30 GHz)			S $_{\nu}$ (5 GHz)			S $_{\nu}$ (1.4 GHz)				
		s $^{-1}$	ergs cm $^{-2}$	unc.	ref.	mJy	unc.	ref.	mJy	unc.	ref.	mJy	unc.	ref.
126.3+02.9	K 3-90	–	–	–	–	–	–	–	13.9	1.4	(c)	12.8	0.6	(m)
126.6+01.3	IPHAS PN-1	–	–	–	–	–	–	–	–	–	–	4.2	0.5	(u)
129.5+04.5	K 3-91	–	–	–	–	–	–	1.5	0.2	(c)	3.4	0.4	(m)	
130.2+01.3	IC 1747	-11.49	0.03	(a, f)	67.4	3.6	(a)	128	24	(j)	85.9	2.6	(m)	
130.4+03.1	K 3-92	–	–	–	–	–	–	2	0.2	(b)	–	–	–	
131.5+02.6	Abell 3	-12.61	0.1	(b)	–	–	–	2.7	0.3	(c)	7.8	1.3	(m)	
132.4+04.7	K 3-93	–	–	–	–	–	–	–	–	–	3.5	0.6	(m)	
136.1+04.9	Abell 6	-12.43	0.06	(b)	–	–	–	12	1.2	(b)	–	–	–	
138.8+02.8	IC 289	-11.69	0.01	(f)	98.8	4	(a)	212	49	(j)	153.1	5.8	(m)	
142.1+03.4	K 3-94	–	–	–	–	–	–	5.5	0.6	(c)	7.1	0.5	(m)	
147.4-02.3	M 1-4	-12.14	0.05	(h)	59.4	5.5	(a)	84.5	3.9	(d, g)	78.2	2.4	(m)	
147.8+04.1	M 2-2	-12.22	0.02	(b)	–	–	–	54	5	(c)	52.5	1.6	(m)	
149.0+04.4	K 4-47	–	–	–	–	–	–	–	–	–	–	–	–	
151.4+00.5	K 3-64	-13.82	0.2	(l)	–	–	–	–	–	–	–	6.5	0.9	(n)
153.7-01.4	K 3-65	–	–	–	–	–	–	–	–	–	–	–	–	
160.5-00.5	We 1-2	–	–	–	–	–	–	–	–	–	–	–	–	
163.1-00.8	We 1-3	–	–	–	–	–	–	–	–	–	–	–	–	
173.5+03.2	Pu 2	–	–	–	–	–	–	–	–	–	–	6.2	0.5	(n)
178.3-02.5	K 3-68	–	–	–	–	–	–	5	0.5	(c)	8.7	0.5	(m)	
181.5+00.9	Pu 1	–	–	–	–	–	–	7	0.1	(t, m)	2.7	0.9	(x)	
184.0-02.1	M 1-5	-12.05	0.01	(e)	49.3	2.8	(a)	71	9	(d, g)	42.8	1.7	(m)	
184.6+00.6	K 3-70	–	–	–	–	–	–	6	1	(d)	4.7	0.5	(m)	
184.8+04.4	K 3-71	–	–	–	–	–	–	–	–	–	4.9	0.5	(m)	
194.2+02.5	J 900	-11.32	0.02	(m)	86.1	4.8	(a)	110	8	(d, g, i)	108.4	3.3	(m)	
201.7+02.5	K 4-48	-12.94	0.07	(m)	–	–	–	–	–	–	12.1	0.6	(m)	
201.9-04.6	We 1-4	–	–	–	–	–	–	–	–	–	–	–	–	
210.0+03.9	We 2-34	–	–	–	–	–	–	–	–	–	–	–	–	
210.3+01.9	M 1-8	-12.37	0.02	(n)	–	–	–	23	7	(k)	16.6	0.6	(m)	
212.0+04.3	M 1-9	-11.66	0.03	(o)	–	–	–	30	5	(d)	23.9	1.1	(m)	

Table C1. E(H α -H β), C $_{\text{H}\beta}$ and A V [E(H α -H β)] values for catalogued PNe

PN G	Name	E(Hα-Hβ)	unc.	C$_{\text{H}\beta}$	unc.	AV[E(Hα-Hβ)]	unc.
033.8-02.6	NGC 6741	0.54	0.07	0.68	0.09	1.45	0.18
035.9-01.1	Sh 2-71	1.04	0.24	1.3	0.3	2.78	0.66
040.3-00.4	Abell 53	1.27	0.17	1.59	0.21	3.4	0.46
046.4-04.1	NGC 6803	0.19	0.07	0.24	0.09	0.51	0.18
052.2-04.0	M 1-74	0.42	0.09	0.52	0.12	1.11	0.25
052.5-02.9	Me 1-1	1.46	0.12	1.83	0.15	3.9	0.32
060.8-03.6	NGC 6853	0.13	0.2	0.17	0.25	0.36	0.53
065.9+00.5	NGC 6842	0.46	0.12	0.57	0.15	1.23	0.32
069.4-02.6	NGC 6894	0.79	0.17	0.99	0.21	2.11	0.46
071.6-02.3	M 3-35	1.97	0.2	2.46	0.25	5.25	0.55
074.5+02.1	NGC 6881	1.48	0.12	1.85	0.15	3.94	0.32
076.3+01.1	Abell 69	2.48	0.62	3.1	0.78	6.63	1.66
084.9+04.4	Abell 71	0.98	0.11	1.22	0.13	2.61	0.28
088.7-01.6	NGC 7048	0.53	0.07	0.66	0.09	1.41	0.18
089.0+00.3	NGC 7026	0.51	0.09	0.64	0.12	1.37	0.25
093.3-02.4	M 1-79	0.93	0.07	1.17	0.09	2.49	0.18
097.6-02.4	M 2-50	0.43	0.12	0.54	0.15	1.15	0.32
098.1+02.4	K 3-63	0.94	0.22	1.17	0.28	2.5	0.61
098.2+04.9	K 3-60	1.56	0.55	1.95	0.69	4.15	1.48
102.8-05.0	Abell 80	0.46	0.36	0.57	0.45	1.23	0.97
103.2+00.6	M 2-51	1.25	0.15	1.56	0.19	3.33	0.41
107.8+02.3	NGC 7354	1.15	0.07	1.44	0.09	3.08	0.18
119.3+00.3	BV 5-1	1.05	0.12	1.32	0.15	2.81	0.32
122.1-04.9	Abell 2	0.32	0.12	0.4	0.15	0.86	0.32
130.2+01.3	IC 1747	0.65	0.12	0.82	0.15	1.74	0.32
131.5+02.6	Abell 3	1.7	0.3	2.12	0.38	4.53	0.82
136.1+04.9	Abell 6	0.91	0.34	1.14	0.42	2.42	0.91
138.8+02.8	IC 289	1.01	0.07	1.26	0.09	2.69	0.18
147.4-02.3	M 1-4	0.81	0.18	1.02	0.22	2.17	0.47
147.8+04.1	M 2-2	1.25	0.09	1.56	0.12	3.34	0.25
151.4+00.5	K 3-64	2.62	1.15	3.28	1.44	7.00	3.08
184.0-02.1	M 1-5	1.02	0.07	1.28	0.09	2.73	0.18
184.8+04.4	K 3-71	1.82	0.3	2.28	0.38	4.87	0.82
194.2+02.5	J 900	0.71	0.09	0.88	0.12	1.89	0.25
201.7+02.5	K 4-48	1.29	0.22	1.61	0.28	3.43	0.6
210.3+01.9	M 1-8	0.93	0.12	1.16	0.15	2.49	0.32
212.0+04.3	M 1-9	0.16	0.12	0.21	0.15	0.44	0.32

Table D1. Calculated radio $A_V(C_{H\alpha})$, $A_V(C_{H\beta})$ values and weighted averaged $A_V(\text{Radio})$ values for catalogued PNe

PN G	Name	30 GHz				5 GHz				1.4 GHz				Weighted Averaged $A_V(\text{Radio})$	unc.
		$A_V(C_{H\alpha})$	unc.	$A_V(C_{H\beta})$	unc.	$A_V(C_{H\alpha})$	unc.	$A_V(C_{H\beta})$	unc.	$A_V(C_{H\alpha})$	unc.	$A_V(C_{H\beta})$	unc.		
030.8+03.4	Abell 47	—	—	—	—	8.27	0.33	—	—	5.45	0.16	—	—	8.27	0.33
031.0+04.1	K 3-6	7.18	0.17	—	—	—	—	—	—	3.76	0.13	—	—	7.18	0.17
031.3-00.5	HaTr 10	—	—	—	—	—	—	—	—	3.26	0.19	—	—	3.26	0.19
031.7+01.7	PC 20	—	—	—	—	—	—	—	—	5.45	0.10	—	—	5.45	0.10
032.0-03.0	K 3-18	—	—	—	—	3.93	0.18	—	—	—	—	—	—	3.93	0.18
032.5-03.2	K 3-20	—	—	—	—	—	—	—	—	2.13	0.19	—	—	2.13	0.19
032.9-02.8	K 3-19	—	—	—	—	3.68	0.12	—	—	2.69	0.19	—	—	3.40	0.10
033.8-02.6	NGC 6741	—	—	—	—	2.56	0.12	2.21	0.07	1.85	0.03	1.72	0.05	1.90	0.03
035.7-05.0	K 3-26	—	—	—	—	—	—	—	—	2.12	0.16	—	—	2.12	0.16
035.9-01.1	Sh 2-71	—	—	—	—	1.33	0.29	1.80	0.21	0.98	0.22	—	—	1.39	0.13
036.9-02.6	HaTr 13	—	—	—	—	—	—	—	—	2.82	0.25	—	—	2.82	0.25
036.9-01.1	HaTr 11	—	—	—	—	—	—	—	—	5.00	0.11	—	—	5.00	0.11
039.5-02.7	M 2-47	—	—	—	—	3.46	0.22	—	—	2.98	0.11	—	—	3.08	0.10
039.8+02.1	K 3-17	8.19	0.1	—	—	8.32	0.20	—	—	7.85	0.10	—	—	7.93	0.09
040.3-00.4	Abell 53	4.21	0.22	3.95	0.22	4.41	0.24	4.09	0.23	3.15	0.1	—	—	4.16	0.11
040.4-03.1	K 3-30	—	—	—	—	3.51	0.36	—	—	2.52	0.12	—	—	2.62	0.11
041.8+04.4	K 3-15	—	—	—	—	2.41	0.08	—	—	—	—	—	—	2.41	0.08
043.0-03.0	M 4-14	—	—	—	—	—	—	—	—	2.72	0.06	—	—	2.72	0.06
043.1+03.8	M 1-65	—	—	—	—	1.97	0.13	—	—	1.69	0.06	—	—	1.74	0.05
043.3+02.2	PM 1-276	—	—	—	—	—	—	—	—	3.26	0.11	—	—	3.26	0.11
045.9-01.9	K 3-33	—	—	—	—	6.29	0.30	—	—	5.27	0.06	—	—	5.30	0.05
046.3-03.1	PB 9	—	—	—	—	3.79	0.33	—	—	3.37	0.06	—	—	3.39	0.06
046.4-04.1	NGC 6803	1.93	0.16	1.47	0.09	1.88	0.16	1.44	0.09	1.26	0.06	1.02	0.05	1.26	0.03
046.8+02.9	CTSS 4	—	—	—	—	—	—	—	—	3.03	0.19	—	—	3.03	0.19
047.1+04.1	K 3-21	—	—	—	—	—	—	—	—	2.77	0.19	—	—	2.77	0.19
047.1-04.2	Abell 62	—	—	—	—	—	—	—	—	—	—	—	—	—	—
048.0-02.3	PB 10	3.56	0.30	—	—	3.73	0.33	—	—	3.57	0.12	—	—	3.59	0.11
048.1+01.1	K 3-29	7.15	0.17	—	—	6.87	0.06	—	—	4.53	0.07	—	—	6.89	0.05
048.5+04.2	K 4-16	—	—	—	—	2.35	0.53	—	—	2.92	0.06	—	—	2.92	0.06
048.7+02.3	K 3-24	—	—	—	—	6.07	0.19	—	—	4.45	0.07	—	—	6.07	0.19
048.7+01.9	Hen 2-429	3.77	0.19	—	—	3.67	0.19	—	—	3.28	0.14	—	—	3.50	0.10
049.4+02.4	Hen 2-428	—	—	—	—	—	—	—	—	2.86	0.09	—	—	2.86	0.09
050.4-01.6	K 4-28	—	—	—	—	7.04	0.35	—	—	5.15	0.06	—	—	5.15	0.06
051.0-04.5	PC 22	—	—	—	—	2.45	0.31	—	—	1.98	0.18	—	—	2.10	0.15
051.0+03.0	Hen 2-430	3.71	0.30	—	—	3.49	0.20	—	—	2.33	0.08	—	—	3.56	0.17
051.3+01.8	PM 1-295	—	—	—	—	—	—	—	—	3.56	0.11	—	—	3.56	0.11
051.9-03.8	M 1-73	1.60	0.24	—	—	1.58	0.21	—	—	1.56	0.12	—	—	1.57	0.10
052.2-04.0	M 1-74	—	—	—	—	1.89	0.23	1.64	0.16	—	—	—	—	1.72	0.13
052.5-02.9	Me 1-1	—	—	—	—	1.23	0.21	2.09	0.17	0.83	0.10	—	—	0.91	0.09
052.9-02.7	K 3-41	—	—	—	—	1.44	0.30	—	—	2.65	0.06	—	—	1.44	0.30
052.9+02.7	K 3-31	—	—	—	—	5.83	0.34	—	—	4.55	0.08	—	—	4.61	0.08
053.2-01.5	K 3-38	—	—	—	—	4.54	0.19	—	—	4.56	0.09	—	—	4.56	0.08
053.8-03.0	Abell 63	—	—	—	—	—	—	—	—	0.67	0.20	—	—	0.67	0.20
054.2-03.4	Necklace	—	—	—	—	—	—	—	—	2.41	0.19	—	—	2.41	0.19
054.4-02.5	M 1-72	—	—	—	—	2.03	0.32	—	—	—	—	—	—	2.03	0.32
055.1-01.8	K 3-43	—	—	—	—	—	—	—	—	3.48	0.27	—	—	3.48	0.27

Table D1 – continued Calculated radio $A_V(C_{H\alpha})$, $A_V(C_{H\beta})$ values and weighted averaged $A_V(\text{Radio})$ values for catalogued PNe

PN G	Name	30 GHz			5 GHz			1.4 GHz			Weighted Averaged $A_V(\text{Radio})$	unc.			
		$A_V(C_{H\alpha})$	unc.	$A_V(C_{H\beta})$	unc.	$A_V(C_{H\alpha})$	unc.	$A_V(C_{H\beta})$	unc.	$A_V(C_{H\alpha})$	unc.	$A_V(C_{H\beta})$	unc.		
055.2+02.8	Hen 2-432	–	–	–	–	4.01	0.26	–	–	3.33	0.09	–	3.40	0.08	
055.3+02.7	Hen 1- 1	–	–	–	–	3.76	0.19	–	–	3.87	0.08	–	3.85	0.07	
055.5-00.5	M 1-71	–	–	–	–	4.36	0.15	4.50	0.29	–	–	–	4.39	0.13	
055.6+02.1	Hen 1-2	–	–	–	–	2.84	0.19	–	–	2.77	0.08	–	2.78	0.07	
056.0+02.0	K 3-35	6.85	0.20	–	–	6.03	0.33	–	–	4.48	0.08	–	6.63	0.17	
056.4-00.9	K 3-42	–	–	–	–	5.95	0.18	–	–	5.13	0.07	–	5.24	0.07	
057.9-01.5	Hen 2-447	5.13	0.20	–	–	4.82	0.06	–	–	3.32	0.09	–	4.42	0.05	
058.9+01.3	K 3-40	–	–	–	–	3.36	0.33	–	–	2.98	0.08	–	3.00	0.08	
059.4+02.3	K 3-37	–	–	–	–	4.68	0.30	–	–	4.31	0.07	–	4.33	0.07	
059.0+04.6	K 3-34	–	–	–	–	–	–	–	–	2.57	0.29	–	2.57	0.29	
059.9+02.0	K 3-39	–	–	–	–	4.41	0.07	–	–	2.68	0.06	–	4.41	0.07	
060.4+01.5	PM 1-310	–	–	–	–	–	–	–	–	4.96	0.27	–	4.96	0.27	
060.5-00.3	K 3-45	–	–	–	–	–	–	–	–	4.91	0.19	–	4.91	0.19	
060.5+01.8	Hen 2-440	–	–	–	–	4.12	0.06	–	–	3.40	0.09	–	3.93	0.05	
060.8-03.6	NGC 6853	–	–	–	–	–	–	0.08	0.14	–	–	–	0.08	0.14	
062.4-00.2	M 2-48	–	–	–	–	3.46	0.15	–	–	3.09	1.30	–	3.45	0.15	
063.8-03.3	K 3-54	–	–	–	–	4.87	0.23	–	–	4.22	0.82	–	4.82	0.22	
064.9-02.1	K 3-53	–	–	–	–	5.23	0.33	–	–	2.91	0.82	–	2.91	1.02	
065.1-03.5	We 1-9	–	–	–	–	–	–	–	–	3.16	1.02	–	3.16	1.02	
065.9+00.5	NGC 6842	–	–	–	–	3.04	0.19	2.46	0.16	–	–	–	2.70	0.12	
066.9+02.2	K 4-37	–	–	–	–	–	–	–	–	4.25	0.19	–	4.25	0.19	
067.9-00.2	K 3-52	8.65	0.21	–	–	7.96	0.33	–	–	5.98	1.3	–	8.46	0.18	
068.3-02.7	Hen 2-459	5.02	0.15	–	–	4.55	0.06	–	–	1.89	1.03	–	4.61	0.05	
068.6+01.1	Hen 1-4	–	–	–	–	–	–	–	–	2.36	0.11	–	2.36	0.11	
068.7+01.9	K 4-41	–	–	–	–	3.76	0.33	–	–	3.26	1.03	–	3.72	0.32	
068.7+03.0	PC 23	–	–	–	–	3.66	0.21	–	–	–	–	–	3.66	0.21	
068.8-00.0	M 1-75	–	–	–	–	4.20	0.19	–	–	–	–	–	4.20	0.19	
069.2+03.8	K 3-46	–	–	–	–	–	–	–	–	2.15	0.26	–	2.15	0.26	
069.2+02.8	K 3-49	–	–	–	–	3.37	0.07	–	–	–	–	–	3.37	0.07	
069.4-02.6	NGC 6894	–	–	–	–	1.17	0.21	1.47	0.22	–	–	–	1.32	0.15	
069.6-03.9	K 3-58	–	–	–	–	4.11	0.19	–	–	–	–	–	4.11	0.19	
069.7+00.0	K 3-55	7.06	0.18	–	–	7.03	0.33	–	–	–	–	–	7.05	0.16	
071.6-02.3	M 3-35	4.54	0.12	4.76	0.14	4.19	0.32	4.53	0.31	–	–	2.92	0.16	4.60	0.08
072.1+00.1	K 3-57	–	–	–	–	5.35	0.33	–	–	–	–	–	5.35	0.33	
073.0-02.4	K 3-76	–	–	–	–	4.27	0.29	–	–	–	–	–	4.27	0.29	
074.5+02.1	NGC 6881	3.87	0.15	3.89	0.07	3.64	0.06	3.74	0.07	2.87	0.10	3.21	0.09	3.59	0.03
075.6+04.3	ARO 342	–	–	–	–	–	–	–	–	2.26	0.19	–	2.26	0.19	
076.3+01.1	Abell 69	–	–	–	–	4.11	0.19	4.91	0.61	2.99	0.19	4.15	0.61	4.18	0.18
076.4+01.8	KjPn 3	–	–	–	–	–	–	–	–	3.42	0.19	–	3.42	0.19	
077.5+03.7	KjPn 1	–	–	–	–	1.50	0.21	–	–	–	–	–	1.50	0.21	
077.7+03.1	KjPn 2	–	–	–	–	–	–	–	–	5.51	0.19	–	5.51	0.19	
078.3-02.7	K 4-53	5.49	0.25	–	–	–	–	–	–	4.75	0.11	–	4.88	0.10	
078.9+00.7	Sd 1	–	–	–	–	5.06	0.19	–	–	1.64	0.19	–	5.06	0.19	
084.2+01.0	K 4-55	–	–	–	–	–	–	–	–	2.94	0.18	–	2.94	0.18	
084.9+04.4	Abell 71	–	–	–	–	2.35	0.21	2.43	0.14	–	–	–	2.41	0.11	

Table D1 – continued Calculated radio $A_V(C_{H\alpha})$, $A_V(C_{H\beta})$ values and weighted averaged $A_V(\text{Radio})$ values for catalogued PNe


PN G	Name	30 GHz			5 GHz			1.4 GHz			Weighted Averaged $A_V(\text{Radio})$	unc.	
		$A_V(C_{H\alpha})$	unc.	$A_V(C_{H\beta})$	unc.	$A_V(C_{H\alpha})$	unc.	$A_V(C_{H\beta})$	unc.	$A_V(C_{H\alpha})$	unc.	$A_V(C_{H\beta})$	unc.
088.7+04.6	K 3-78	–	–	–	–	4.78	0.22	–	–	4.45	0.06	–	–
088.7-01.6	NGC 7048	–	–	–	–	0.55	0.20	0.82	0.12	0.70	0.06	0.92	0.06
089.0+00.3	NGC 7026	–	–	–	–	2.08	0.45	1.85	0.31	1.85	0.07	1.70	0.08
089.8-00.6	Sh 1-89	–	–	–	–	–	–	–	–	3.52	0.20	–	–
091.6-04.8	K 3-84	–	–	–	–	–	–	–	–	1.25	0.24	–	–
091.6+01.8	We 1- 11	–	–	–	–	–	–	–	–	4.43	0.06	–	–
093.3-00.9	K 3-82	–	–	–	–	3.62	0.30	–	–	3.75	0.16	–	–
093.3-02.4	M 1-79	–	–	–	–	0.38	0.19	1.06	0.11	0.51	0.06	1.14	0.08
094.5-00.8	K 3-83	–	–	–	–	5.21	0.19	–	–	4.71	0.06	–	–
095.1-02.0	M 2-49	–	–	–	–	5.15	0.2	–	–	3.67	0.06	–	–
095.2+00.7	K 3-62	5.89	0.16	–	–	5.86	0.06	–	–	4.76	0.06	4.9	0.25
096.3+02.3	K 3-61	–	–	–	–	3.70	0.19	–	–	3.79	0.06	–	–
097.6-02.4	M 2-50	–	–	–	–	1.83	0.19	1.62	0.16	2.12	0.06	1.81	0.18
098.1+02.4	K 3-63	–	–	–	–	4.38	0.23	3.78	0.23	2.55	0.09	2.53	0.18
098.2+04.9	K 3-60	–	–	–	–	5.28	0.35	4.92	0.70	4.53	0.06	4.41	0.46
102.8-05.0	Abell 80	–	–	–	–	–	–	–	–	0.11	0.06	0.47	0.50
103.2+00.6	M 2-51	–	–	–	–	1.83	0.20	2.31	0.18	1.15	0.15	1.85	0.14
103.7+00.4	M 2-52	–	–	–	–	2.95	0.19	–	–	2.91	0.06	–	–
104.1+01.0	Bl 2-1	–	–	–	–	6.11	0.06	–	–	4.54	0.06	–	–
104.4-01.6	M 2-53	–	–	–	–	2.17	0.19	–	–	2.43	0.06	–	–
107.4-02.6	K 3-87	–	–	–	–	3.55	0.27	–	–	4.67	0.13	–	–
107.4-00.6	K 4-57	–	–	–	–	–	–	–	–	3.85	0.19	–	–
107.7-02.2	M 1-80	–	–	–	–	2.22	0.22	–	–	1.60	0.06	–	–
107.8+02.3	NGC 7354	3.95	0.10	3.67	0.03	4.12	0.23	3.79	0.14	3.92	0.09	3.65	0.06
112.5-00.1	KjPn 8	–	–	–	–	1.58	0.12	–	–	2.66	0.25	–	–
112.5+03.7	K 3-88	–	–	–	–	–	–	–	–	4.65	0.06	–	–
119.3+00.3	BV 5-1	–	–	–	–	–	–	–	–	2.50	0.06	2.60	0.16
121.6+00.0	BV 5-2	–	–	–	–	–	–	–	–	2.43	0.06	–	–
121.6+03.5	We 1-1	–	–	–	–	–	–	–	–	4.05	0.06	–	–
122.1-04.9	Abell 2	–	–	–	–	0.05	0.19	0.31	0.16	1.80	0.06	1.50	0.17
126.3+02.9	K 3-90	–	–	–	–	3.03	0.19	–	–	2.74	0.06	–	–
126.6+01.3	IPHAS PN-1	–	–	–	–	–	–	–	–	1.47	1.05	–	–
129.5+04.5	K 3-91	–	–	–	–	2.64	0.19	–	–	3.58	0.06	–	–
130.2+01.3	IC 1747	1.88	0.06	1.83	0.09	2.60	0.32	2.32	0.24	2.42	0.09	2.20	0.24
130.4+03.1	K 3-92	–	–	–	–	1.96	0.19	–	–	–	–	–	1.96
131.5+02.6	Abell 3	–	–	–	–	–	–	0.85	0.32	0.40	0.07	–	0.41
132.4+04.7	K 3-93	–	–	–	–	–	–	–	–	3.74	0.06	–	3.74
136.1+04.9	Abell 6	–	–	–	–	2.03	0.36	2.16	0.23	–	–	–	2.12
138.8+02.8	IC 289	2.12	0.11	2.30	0.03	2.92	0.38	2.84	0.24	–	–	2.42	0.06
142.1+03.4	K 3-94	–	–	–	–	2.84	0.15	–	–	–	–	–	2.84
147.4-02.3	M 1- 4	3.44	0.19	3.03	0.11	3.68	0.12	3.19	0.15	–	–	3.00	0.14
147.8+04.1	M 2- 2	–	–	–	–	2.74	0.18	–	–	–	–	–	2.74
149.0+04.4	K 4-47	–	–	–	–	–	–	–	–	–	–	–	–
151.4+00.5	K 3-64	–	–	–	–	–	–	–	–	3.25	0.25	4.45	1.3
153.7-01.4	K 3-65	–	–	–	–	–	–	–	–	–	–	–	–
160.5-00.5	We 1-2	–	–	–	–	–	–	–	–	–	–	–	–
163.1-00.8	We 1-3	–	–	–	–	–	–	–	–	–	–	–	–


Table D1 – continued Calculated radio $A_V(C_{H\alpha})$, $A_V(C_{H\beta})$ values and weighted averaged $A_V(\text{Radio})$ values for catalogued PNe


PN G	Name	30 GHz			5 GHz			1.4 GHz			Weighted Averaged $A_V(\text{Radio})$	unc.	
		$A_V(C_{H\alpha})$	unc.	$A_V(C_{H\beta})$	unc.	$A_V(C_{H\alpha})$	unc.	$A_V(C_{H\beta})$	unc.	$A_V(C_{H\alpha})$	unc.	$A_V(C_{H\beta})$	unc.
173.5+03.2	Pu 2	–	–	–	–	–	–	–	–	3.95	0.17	–	3.95
178.3-02.5	K 3-68	–	–	–	–	2.73	0.19	–	–	–	–	–	2.73
181.5+00.9	Pu 1	–	–	–	–	2.71	0.07	–	–	1.24	0.53	–	2.55
184.0-02.1	M 1-5	2.98	0.13	2.90	0.02	3.23	0.23	3.07	0.14	–	–	2.48	0.06
184.6+00.6	K 3-70	–	–	–	–	4.32	0.29	–	–	–	–	–	4.32
184.8+04.4	K 3-71	–	–	–	–	–	–	–	–	–	–	3.76	0.32
194.2+02.5	J 900	2.09	0.13	1.62	0.05	1.50	0.13	1.68	0.11	–	–	1.55	0.07
201.7+02.5	K 4-48	–	–	–	–	–	–	–	–	2.30	0.12	3.14	0.20
201.9-04.6	We 1- 4	–	–	–	–	–	–	–	–	–	–	–	–
210.0+03.9	We 2- 34	–	–	–	–	–	–	–	–	–	–	–	–
210.3+01.9	M 1-8	–	–	–	–	2.47	0.49	2.48	0.36	–	–	2.06	0.1
212.0+04.3	M 1-9	–	–	–	–	1.79	0.29	1.35	0.22	1.30	0.12	1.03	0.11
												1.21	0.07


**APPENDIX E: H-MEAD EXTINCTION
VERSUS DISTANCE PLOTS**

H-MEAD extinction versus distance plots for the lines of sight to all of the previously catalogued PNe observed by IPHAS, ordered by increasing PN G, are shown in Figures E1-E19. The solid blue curve in each plot represents the H-MEAD extinction versus distance relation for the sightline to the nebula. The solid and dashed red horizontal lines correspond to the value of A_V derived for the nebula, and its corresponding uncertainties, while the solid and dashed green horizontal lines correspond to the Schlafly & Finkbeiner (2011) A_V limit and its uncertainties.

**Figure E1.** H-MEAD Extinction vs. Distance plots for Abell 47, K 3-6, HaTr 10, PC 20, K 3-18, K 3-20, K 3-19 and NGC 6741.

**Figure E2.** H-MEAD Extinction vs. Distance plots for K 3-26, Sh 2-71, HaTr 13, HaTr 11, M 2-47, K 3-17, Abell 53 and K 3-30.

**Figure E3.** H-MEAD Extinction vs. Distance plots for K 3-15, M 4-14, M 1-65, PM 1-276, K 3-33, PB 9 NGC 6803 and CTSS 4.

**Figure E4.** H-MEAD Extinction vs. Distance plots for K 3-21, PB 10, K 3-29, K 4-16, K 3-24, Hen 2-429, Hen 2-428 and K 4-28.




Figure E5. H-MEAD Extinction vs. Distance plots for PNe PC 22, Hen 2-430, PM 1-295, M 1-73, M 1-74, Me 1-1, K 3-41 and K 3-31. MNRAS 000, 000–000 (0000)




Figure E6. H-MEAD Extinction vs. Distance plots for K 3-38, Abell 63, Necklace nebula, M1-72, K 3-43, Hen2-432, Hen 1-1 and M 1-71.




Figure E7. H-MEAD Extinction vs. Distance plots for Hen 1-2, K 3-35, K 3-42, Hen 2-447, K 3-40, K 3-37, K 3-34 and K 3-39.




Figure E8. H-MEAD Extinction vs. Distance plots for PM 1-310, K 3-45, Hen 2-440, NGC 6853, M 2-48, K 3-54, K 3-53 and We 1-9.




Figure E9. H-MEAD Extinction vs. Distance plots for NGC 6842, K 4-37, K 3-52, Hen 2-459, Hen 1-4, K 4-41, PC 23 and M 1-75.
MNRAS **000**, 000–000 (0000)




Figure E10. H-MEAD Extinction vs. Distance plots for K 3-46, K 3-49, NGC 6894, K 3-58, K 3-55, K 3-35, K 3-57 and K 3-76.





Figure E11. H-MEAD Extinction vs. Distance plots for NGC 6881, ARO 342, Abell 69, KjPn 3, KjPn 1, KjPn 2, K 4-53 and Sd 1.
MNRAS 000, 000–000 (0000)

**Figure E12.** H-MEAD Extinctions vs. Distance plots for K 4-55, Abell 71, K 3-78, NGC 7048, NGC 7026, Sh 1-89, K 3-84 and We 1-11.





Figure E13. H-MEAD Extinction vs. Distance plots for K 3-82, M 1-79, K 3-83, M 2-49, K 3-62, K 3-61, M 2-50 and K 3-63.

**Figure E14.** H-MEAD Extinction vs. Distance plots for K 3-60, Abell 80, M 2-51, M 2-52, Bl 2-1, M 2-53, K 3-87 and K 4-57.







Figure E15. H-MEAD Extinction vs. Distance plots for M 1-80, NGC 7354, KjPn 8, K 3-88, BV 5-1, BV 5-2, We 1-1 and Abell 2. MNRAS 000, 000–000 (0000)

**Figure E16.** H-MEAD Extinction vs. Distance plots for K 3-90, IPHAS PN-1, K 3-91, IC 1747, K 3-92, Abell 3, K 3-93 and Abell 6.

**Figure E17.** H-MEAD Extinction vs. Distance plots for IC 289, K 3-94, M 1-4, M 2-2, K 3-64, Pu 2, K 3-68 and Pu 1.

**Figure E18.** H-MEAD Extinction vs. Distance plots for M 1-5, K 3-70, K 3-71, J 900, K 4-48, M 1-8 and M 1-9.