

On Well-Structured Convex-Concave Saddle Point Problems and Variational Inequalities with Monotone Operators

Anatoli Juditsky ^{*} Arkadi Nemirovski [†]

Abstract

For those acquainted with **CVX** (aka disciplined convex programming) of M. Grant and S. Boyd [9], the motivation of this work is the desire to extend the scope of **CVX** beyond convex minimization—to convex-concave saddle point problems and variational inequalities with monotone operators. To attain this goal, given a family \mathcal{K} of cones (e.g., Lorentz, semidefinite, geometric, etc), we introduce the notions of \mathcal{K} -conic representation of a convex-concave saddle point problem and of variational inequality with monotone operator. We demonstrate that given such a representation of the problem of interest, the latter can be reduced straightforwardly to a conic problem on a cone from \mathcal{K} and thus can be solved by (any) solver capable to handle conic problems on cones from \mathcal{K} (e.g., **Mosek** or **SDPT3** in the case of semidefinite cones). We also show that \mathcal{K} -representations of convex-concave functions and monotone vector fields admit a fully algorithmic calculus which helps to recognize the cases when a saddle point problem or variational inequality can be converted into a conic problem on a cone from \mathcal{K} and to carry out such conversion.

1 Introduction

Along with emergence of powerful computational tools, one of important components of what is sometimes referred to as Interior-Point Revolution in Convex Optimization was the rise of (informal) notion of a “well-structured convex problem”—convex optimization problem which can be reformulated as a conic problem on a cone from a “nice” family \mathcal{K} , most notably, the families of finite direct products of nonnegative rays (Linear Programming), Lorentz cones (Second Order Conic Programming), and semidefinite cones (Semidefinite Programming). Problems of the latter “magic families” cover basically all needs arising in applications of Convex Optimization to decision making, engineering, statistics, etc. At the same time, reformulating problem of interest as LP/SOCP/SDP makes the problem amenable for numerical processing by a “universal” SDP solver, so that end user should not care much about number-crunching. Today, there exist powerful universal SDP solvers such as **SDPT3** [19] or industrial-grade **Mosek** [2], with permanently improving performance and reliability.

The “maiden form” of an optimization problem arising in applications usually is not the “nice conic” one, even when at the end of the day the problem can be cast into, say, an SDP program. Such transformation is carried out with the help of a special “calculus” (see, e.g., [3, 4, 5]) which allows to recognize that the problem at hand can be reformulated as, say, an SDP. This calculus, as any other, has two components: (a) “raw materials”—a collection of sets and functions possessing explicit representations via SDP (what exactly “representation” means, will be explained in Section 2), and (b) “calculus rules” expressing SDP representation of the result of a specific operation with functions/sets via SDP representations of the operands. It turns out that calculus rules for SDP representations cover all basic convexity-preserving operations with functions (like taking linear combinations with nonnegative coefficients and convexity-preserving compositions) and sets (like taking finite intersections and images/inverse images under affine mappings). Moreover, the calculus rules happen to be fully algorithmic and thus can be implemented on a compiler. This possibility is

^{*}LJK, Université Grenoble Alpes, 700 Avenue Centrale, 38401 Domaine Universitaire de Saint-Martin-d’Hères, France, anatoli.juditsky@univ-grenoble-alpes.fr

[†]ISyE, Georgia Institute of Technology, Atlanta, Georgia 30332, USA, nemirovs@isye.gatech.edu.

utilized, for instance, in the CVX software of M. Grant and S. Boyd [9] which is the second-to-none in terms of its scope and user-friendliness tool for numerical processing of convex optimization problems.

So far, we were speaking about (constrained) convex minimization, and this is the usual scope of the calculus of well-structured convex problems. For example, given a number of SDP-representable functions, the calculus allows to obtain SDP representation of their maximum and thus convert to SDP program, in an automated fashion, the problem of minimizing this maximum over a feasible set given by an SDP representation. However, such calculus does *not* allow to get an SDP representation of the maximum of *infinite* family of SDP-representable functions, even when this family is perfectly well organized. As a result, convex-concave saddle point problems, well-structured from the “viewpoint of a human,” cannot be “fed to CVX” as they are, and reducing such a problem to the conic form amenable for the existing powerful convex optimization software requires ad hoc work on case-to-case basis. Similar difficulties arise when instead of convex-concave saddle point problems someone is looking to solve numerically other problems with convex structure, most notably, variational inequalities (VI) with monotone operators. Monotone VI’s can be thought of as the most general problems with convex structure: on one hand, other problems of this type (constrained convex minimization, convex-concave saddle points, convex Nash equilibrium problems) under extremely mild boundedness and regularity assumptions can be reduced to monotone VI’s (for details, see e.g., [5, Section 5.6]). On the other hand, efficient *black box oriented* algorithms of convex minimization, like First Order or Ellipsoid methods, can be extended to monotone VI’s (cf., e.g., [14]) without harming their efficiency estimates. There exists also rich literature on extending polynomial time Interior-Point algorithms onto linear and monotone complementarity problems, see, e.g., [11, 12, 7, 16, 17, 18] and references therein. In particular, it is well known that VI with *affine* monotone operator and domain admitting conic representation can be straightforwardly reduced to a conic optimization problem. In [13, Section 7.4] this fact was somehow extended to nonlinear monotone VI’s from certain restricted class—those admitting an explicit “convex representation” allowing to reduce the VI of interest to an affine monotone VI on a properly modified domain; that reference develops also a rudimentary calculus of convex representations of monotone VI’s. Some results¹ on reformulations of VI’s as convex optimization problems can be found in [1]. That being said, to the best of our knowledge, beyond the classes of VI’s with affine monotone operators and related bilinear saddle point problems there is no much understanding of what is a “well structured” convex-concave saddle point problem or “genuine”—not potential—monotone VI, and of how such a problem should be represented in order for the representation to admit a meaningful algorithmic calculus, on one hand, and to allow for straightforward conversion of the problem into a nice conic program, say, SDP one, on the other.

The goal of what follows is to introduce the notion of “conic representations” of problems with convex structure, specifically, convex-concave saddle point problems and monotone VI’s, via a given family \mathcal{K} of cones (e.g., direct products of nonnegative rays/Lorentz/Semidefinite cones). As we will see, given such a representation, the problem of interest can be straightforwardly converted to a conic problem on a cone from \mathcal{K} , and representations in question admit a fully algorithmic calculus, similar to the calculus of conic representations of convex functions/sets. In Section 3 we develop conic representations of convex-concave functions, and then in Section 4 extend this framework to representations of monotone vector fields. Derivations in Section 4 can be viewed as a “well-structured” (i.e., conic representation-oriented) version of convex representation of a monotone operator introduced in [13, Section 7.4].

2 Preliminaries

Consider a family \mathcal{K} of regular (i.e., closed, convex, pointed, and with nonempty interior) cones in Euclidean spaces such that the family

1. contains nonnegative ray,
2. is closed w.r.t. taking finite direct products of its members, and
3. is closed w.r.t. passing from a cone to its dual.

¹dealing with the search for strong solutions to VI’s with not necessarily monotone operator F under rather restrictive assumptions that $x^T F(x)$ and $-F(x)$ are convex on polyhedral set $\mathcal{X} \subset \mathbf{R}_+^n$

From now on we adopt the following convention: *we operate with once for ever fixed family \mathcal{K} of cones satisfying Assumptions 1–3, and whenever in the sequel we consider a cone, say \mathbf{K} , this cone is assumed to belong to \mathcal{K}* . Besides this, \mathbf{K}^* stands for the cone dual to \mathbf{K} .

- Given a cone $\mathbf{K} \in \mathcal{K}$ we call a constraint in variables $x \in \mathbf{R}^n$ of the form

$$Ax \leq_{\mathbf{K}} a \tag{1}$$

\mathcal{K} -conic; as usual, $a \leq_{\mathbf{K}} b$ means that $b - a \in \mathbf{K}$. Note that a system

$$A_k x \leq_{\mathbf{K}_k} a_k, 1 \leq k \leq K,$$

of finitely many \mathcal{K} -conic constraints is equivalent to a single \mathcal{K} -conic constraint²

$$[A_1; \dots; A_K]x \leq_{\mathbf{K}} [a_1; \dots; a_K]$$

where $\mathbf{K} = \mathbf{K}_1 \times \dots \times \mathbf{K}_K$ belongs to \mathcal{K} along with $\mathbf{K}_k, k \leq K$, since \mathcal{K} is closed w.r.t. taking direct products. For similar reasons, augmenting a \mathcal{K} -conic constraint $Ax \leq_{\mathbf{K}} a$ in variables x by a system S of finitely many linear equalities and nonstrict linear equalities, we get a conic constraint on x . Indeed, S can be written equivalently as $Bx \leq b$, so that the augmented constraint reads

$$[A; B]x \leq_{\mathbf{K} \times \mathbf{R}_+^m} [a; b], m = \dim b,$$

and the cone $\mathbf{K} \times \mathbf{R}_+^m$ belongs to \mathcal{K} since the latter family contains \mathbf{K}, \mathbf{R}_+ and is closed w.r.t. taking finite direct products.

- We call conic constraint (1) *essentially strictly feasible*, if \mathbf{K} can be decomposed as $\mathbf{K} = \mathbf{R}_+^m \times \mathbf{L}$ with regular cone \mathbf{L} , so that (1) is of the form

$$[P; Q]x \leq_{\mathbf{R}_+^m \times \mathbf{L}} [p; q],$$

and there exists \bar{x} such that $P\bar{x} \leq p$ and $Q\bar{x} <_{\mathbf{L}} q$, the latter meaning that $q - Q\bar{x} \in \text{int } \mathbf{L}$.

- We say that a set $X \subset \mathbf{R}^n$ is \mathcal{K} -representable if for properly selected A, B, c and $\mathbf{K} \in \mathcal{F}$ one has

$$X = \{x : \exists u : Ax + Bu \leq_{\mathbf{K}} c\};$$

whenever this is the case, the \mathcal{K} -conic constraint $Ax + Bu \leq_{\mathbf{K}} c$ in the right hand side is called \mathcal{K} -representation of X . We call a function $f : \mathbf{R}^n \rightarrow \mathbf{R} \cup \{+\infty\}$ \mathcal{K} -representable if the epigraph $\text{Epi}\{f\} = \{(x, t) : t \geq f(x)\}$ is \mathcal{K} -representable and we refer to a \mathcal{K} -representation of $\text{Epi}\{f\}$ as the \mathcal{K} -representation of f . In other words, asserting that \mathcal{K} -conic constraint $Ax + tb + Bu \leq_{\mathbf{K}} c$ represents f is the same as saying that the following equivalence takes place:

$$t \geq f(x) \Leftrightarrow \exists u : Ax + tb + Bu \leq_{\mathbf{K}} c.$$

Observe that the level sets $\{x : f(x) \leq a\}, a \in \mathbf{R}$, of \mathcal{K} -representable function f are \mathcal{K} -representable:

$$\{t \geq f(x) \Leftrightarrow \exists u : Ax + tb + Bu \leq_{\mathbf{K}} c\} \Rightarrow \{x : f(x) \leq a\} = \{x : \exists u : Ax + Bu \leq_{\mathbf{K}} c - ab\}.$$

Thus, given \mathcal{K} -representations $\{x : \exists u : Ax + Bu \leq_{\mathbf{K}} c\}$ of $X \subset \mathbf{R}^n$ and $\{t \geq f(x) \Leftrightarrow \exists v : Cx + td + Ev \leq_{\mathbf{L}} g\}$ of $f : \mathbf{R}^n \rightarrow \mathbf{R} \cup \{+\infty\}$, the problem

$$\min_{x \in X} f(x) \tag{P}$$

is equivalent to the conic problem

$$\min_{x, t, u, v} \{t : [Ax + Bu; Cx + td + Ev] \leq_{\mathbf{K} \times \mathbf{L}} [c; g]\} \tag{Q}$$

²We follow the ‘‘Matlab convention’’ for matrices: $[A, B]$ and $[A; B]$ denote, respectively, the horizontal and vertical concatenations of two matrices of compatible dimensions.

on a cone from \mathcal{K} , where equivalence of (\mathcal{P}) and (\mathcal{Q}) precisely means that x is a feasible solution to (\mathcal{P}) with a finite value of the objective if and only if there exist t, u, v such that $t \geq f(x)$ and x, t, u, v is feasible for (\mathcal{Q}) .

\mathcal{K} -representable functions/sets admit fully algorithmic calculus: all basic convexity-preserving operations with sets and functions, (e.g., taking finite intersections, images/inverse images under affine mappings, and direct products of sets, or taking linear combinations with nonnegative coefficients and convexity-preserving compositions of functions) as applied to \mathcal{K} -representable operands, produce \mathcal{K} -representable results, with representations of the results readily given by those of the operands; for details, see, e.g., [4, 5]. When a solver for conic problems on cones from \mathcal{K} is available, this calculus allows one to recognize that in the problem (\mathcal{P}) of interest objective f and domain X are \mathcal{K} -representable and find the corresponding \mathcal{K} -representations,³ so the problem of interest can be converted into a conic problem (\mathcal{Q}) on a cone from \mathcal{K} and solved by the solver at hand.

From now on, we operate with a once for ever fixed family \mathcal{K} of cones satisfying the above Assumptions 1–3, which allows for the convention as follows: whenever in the sequel a cone arises, this cone, if the otherwise is explicitly stated, belongs to \mathcal{K} . Besides this, \mathbf{K}^* stands for the cone dual to \mathbf{K} .

3 Well-structured convex-concave saddle point problems

As far as its paradigm and set of rules are concerned, “calculus of \mathcal{K} -representability” for, say, \mathcal{K} specified as \mathcal{SDP} (finite direct products of semidefinite cones) covers all “basic” needs of “well-structured” convex minimization. There is, however, an exception—the case in which the objective in the convex problem of interest

$$\min_{x \in \mathcal{X}} \bar{\Psi}(x) \tag{P}$$

is given implicitly:

$$\bar{\Psi}(x) = \sup_{y \in \mathcal{Y}} \psi(x, y) \tag{2}$$

where \mathcal{Y} is convex set and $\psi : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbf{R}$ is convex-concave (i.e., convex in $x \in \mathcal{X}$ and concave in $y \in \mathcal{Y}$) and continuous. Problem (P) with objective given by (2) is called “primal problem associated with the convex-concave saddle point problem $\min_{x \in \mathcal{X}} \max_{y \in \mathcal{Y}} \psi(x, y)$,” and problems of this type do arise in some applications of well-structured convex optimization. We are about to define a *saddle point* version of \mathcal{K} -representability along with the corresponding calculus which allows to convert “ \mathcal{K} -representable convex-concave saddle point problems” into usual conic problems on cones from \mathcal{K} .

3.1 Conic representability of convex-concave function—definition

Let \mathcal{X}, \mathcal{Y} be nonempty convex sets given by \mathcal{K} -representations:

$$\mathcal{X} = \{x : \exists \xi : Ax + B\xi \leq_{\mathbf{K}_x} c\}, \quad \mathcal{Y} = \{y : \exists \eta : Cy + D\eta \leq_{\mathbf{K}_y} e\} \quad [\mathbf{K}_x \in \mathcal{K}, \mathbf{K}_y \in \mathcal{K}].$$

Let us say that a convex-concave continuous function $\psi(x, y) : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbf{R}$ is \mathcal{K} -representable on $\mathcal{X} \times \mathcal{Y}$, if it admits representation of the form

$$\forall (x \in \mathcal{X}, y \in \mathcal{Y}) : \psi(x, y) = \inf_{f, t, u} \{f^T y + t : Pf + tp + Qu + Rx \leq_{\mathbf{K}} s\} \tag{3}$$

where $\mathbf{K} \in \mathcal{K}$. We call representation (3) *essentially strictly feasible*, if the conic constraint

$$Pf + tp + Qu \leq_{\mathbf{K}} s - Rx$$

in variables f, t, u is essentially strictly feasible for every $x \in \mathcal{X}$.

³Note that calculus rules are fully algorithmic and thus can be implemented by a compiler, the most famous example being the CVX software [9] which can handle “semidefinite representability” (family \mathcal{K} comprised of direct products of semidefinite cones), and as a byproduct—conic quadratic representability.

3.2 Main observation

Assume that \mathcal{Y} is compact and is given by essentially strictly feasible \mathcal{K} -representation

$$\mathcal{Y} = \{y : \exists \eta : Cy + D\eta \leq_{\mathbf{K}_y} e\}. \quad (4)$$

Then problem (P) can be processed as follows: for $x \in \mathcal{X}$ we have

$$\begin{aligned} \bar{\Psi}(x) &= \max_{y \in \mathcal{Y}} \inf_{f,t,u} [f^T y + t : Pf + tp + Qu + Rx \leq_{\mathbf{K}} s] \\ &= \inf_{f,t,u} \left\{ \max_{y \in \mathcal{Y}} [f^T y + t : Pf + tp + Qu + Rx \leq_{\mathbf{K}} s] \right\} \quad \left[\begin{array}{l} \text{Sion-Kakutani Theorem; recall} \\ \text{that } \mathcal{Y} \text{ is convex and compact} \end{array} \right] \\ &= \inf_{f,t,u} \left\{ \max_{y,\eta} [f^T y : Cy + D\eta \leq_{\mathbf{K}_y} e] + t : Pf + tp + Qu + Rx \leq_{\mathbf{K}} s \right\} \\ &= \inf_{f,t,u} \left\{ \min_{\lambda} [\lambda^T e : C^T \lambda = f, D^T \lambda = 0, \lambda \geq_{\mathbf{K}_y^*} 0] + t : Pf + tp + Qu + Rx \leq_{\mathbf{K}} s \right\} \\ &\quad \left[\begin{array}{l} \text{by strong conic duality, see, e.g., [5, Theorem 1.4.4];} \\ \text{recall that (4) is essentially strictly feasible,} \end{array} \right] \end{aligned}$$

so that the problem of interest

$$\min_{x \in \mathcal{X}} \bar{\Psi}(x) \quad (a)$$

reduces to the explicit \mathcal{K} -conic problem

$$\min_{x,\xi,f,t,u,\lambda} \left\{ \begin{array}{l} Pf + tp + Qu + Rx \leq_{\mathbf{K}} s, \\ e^T \lambda + t : C^T \lambda = f, D^T \lambda = 0, \lambda \geq_{\mathbf{K}_y^*} 0, \\ Ax + B\xi \leq_{\mathbf{K}_x} c \end{array} \right\}. \quad (b)$$

Here, “reduction” means that the x -component of a feasible solution $\zeta = (x, \xi, f, t, u, \lambda)$ to (b) is a feasible solution to (a) with the value of the objective of the latter problem at x being \leq the value of the objective of (b) at ζ , and the optimal values in (a) and (b) are the same. Thus, as far as building feasible approximate solutions of a prescribed accuracy $\epsilon > 0$ in terms of the objective are concerned, problem (a) reduces to the explicit conic problem (b). Note, however, that (a) and (b) are not “exactly the same”—it may happen that (a) is solvable while (b) is not so. “For all practical purposes,” this subtle difference is of no importance since in actual computations exactly optimal solutions usually are not reachable anyway.

Discussion. Note that for continuous convex-concave function $\psi : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbf{R}$ the set

$$\mathcal{Z} = \{[f; t; x] : x \in \mathcal{X}, f^T y + t \geq \psi(x, y) \forall y \in \mathcal{Y}\}$$

clearly is convex, and by the standard Fenchel duality we have

$$\forall (x \in \mathcal{X}, y \in \mathcal{Y}) : \psi(x, y) = \inf_{f,t} [f^T y + t : [f; t; x] \in \mathcal{Z}]. \quad (5)$$

\mathcal{K} -representability of ψ on $\mathcal{X} \times \mathcal{Y}$ means that (5) is preserved when replacing the set \mathcal{Z} with its properly selected \mathcal{K} -representable subset. Given that \mathcal{Z} is convex, this assumption seems to be not too restrictive; taken together with \mathcal{K} -representability of \mathcal{X} and \mathcal{Y} , it can be treated as the definition of \mathcal{K} -representability of the convex-concave function ψ . The above derivation shows that convex-concave saddle point problem with \mathcal{K} -representable domain and cost function (more precisely, the primal minimization problem (P) induced by this saddle point problem) can be represented in explicit \mathcal{K} -conic form, at least when the \mathcal{K} -representations of the cost and of (compact) \mathcal{Y} are essentially strictly feasible.

Note also that if \mathcal{X} and \mathcal{Y} are convex sets and a function $\psi(x, y) : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbf{R}$ admits representation (3), then ψ automatically is convex in $x \in \mathcal{X}$ and concave in $y \in \mathcal{Y}$.

3.3 Symmetry

Assume that representation (3) is essentially strictly feasible. Then for all $x \in \mathcal{X}, y \in \mathcal{Y}$ we have by conic duality

$$\begin{aligned}\psi(x, y) &= \inf_{f, t, u} \{ f^T y + t : Pf + tp + Qu + Rx \leq_{\mathbf{K}} s \} \\ &= \sup_{\bar{u} \in \mathbf{K}^*} \{ \bar{u}^T [Rx - s] : P^T \bar{u} + y = 0, p^T \bar{u} + 1 = 0, Q^T \bar{u} = 0 \},\end{aligned}$$

whence, setting

$$\bar{\mathcal{X}} = \mathcal{Y}, \bar{\mathcal{Y}} = \mathcal{X}, \bar{x} = y, \bar{y} = x, \bar{\psi}(\bar{x}, \bar{y}) = -\psi(\bar{y}, \bar{x}) = -\psi(x, y),$$

we have

$$\begin{aligned}(\forall \bar{x} \in \bar{\mathcal{X}}, \bar{y} \in \bar{\mathcal{Y}}) : \\ \bar{\psi}(\bar{x}, \bar{y}) &= -\psi(x, y) = \inf_{\bar{u} \in \mathbf{K}^*} \{ -\bar{u}^T [Rx - s] : P^T \bar{u} + y = 0, p^T \bar{u} + 1 = 0, Q^T \bar{u} = 0 \} \\ &= \inf_{\bar{f}, \bar{t}, \bar{u}} \left\{ \bar{f}^T \bar{y} + \bar{t} : \underbrace{\left[\begin{array}{l} \bar{f} = -R^T \bar{u}, \bar{t} = s^T \bar{u}, Q^T \bar{u} = 0, \\ p^T \bar{u} + 1 = 0, P^T \bar{u} + \bar{x} = 0, \bar{u} \in \mathbf{K}^* \end{array} \right]}_{\Leftrightarrow \bar{P} \bar{f} + \bar{t} \bar{p} + \bar{Q} \bar{u} + \bar{R} \bar{x} \leq_{\bar{\mathbf{K}}} \bar{s}} \right\}\end{aligned}$$

with $\bar{\mathbf{K}} \in \mathcal{K}$. We see that a (essentially strictly feasible) \mathcal{K} -representation of convex-concave function ψ on $\mathcal{X} \times \mathcal{Y}$ induces straightforwardly a \mathcal{K} -representation of the “symmetric entity”—the convex-concave function $\bar{\psi}(y, x) = -\psi(x, y)$ on $\mathcal{Y} \times \mathcal{X}$, with immediate consequences for converting the optimization problem

$$\sup_{y \in \mathcal{Y}} \left[\bar{\Psi}(y) := \inf_{x \in \mathcal{X}} \psi(x, y) \right] \tag{D}$$

into the standard conic form.

3.4 Calculus of conic representations of convex-concave functions

Representations of the form (3) admit a calculus.

3.4.1 Raw materials

Raw materials for the calculus are given by

1. Functions $\psi(x, y) = a(x)$, where $a(x)$, $\text{Dom } a \supset \mathcal{X}$, is \mathcal{K} -representable:

$$t \geq a(x) \Leftrightarrow \exists u : \bar{R}x + t\bar{p} + \bar{Q}u \leq_{\bar{\mathbf{K}}} \bar{s}$$

In this case

$$\psi(x, y) = \inf_{f, t, u} \left\{ f^T y + t : \underbrace{f = 0, \bar{R}x + t\bar{p} + \bar{Q}u \leq_{\bar{\mathbf{K}}} \bar{s}}_{\Leftrightarrow Pf + tp + Qu + Rx \leq_{\mathbf{K}} s \text{ with } \mathbf{K} \in \mathcal{K}} \right\}.$$

2. Functions $\psi(x, y) = -b(y)$, where $b(y)$, $\text{Dom } b \supset \mathcal{Y}$, is \mathcal{K} -representable:

$$t \geq b(y) \Leftrightarrow \exists u : \bar{R}y + t\bar{p} + \bar{Q}u \leq_{\bar{\mathbf{K}}} \bar{s}$$

with essentially strictly feasible \mathcal{K} -representation. In this case

$$\begin{aligned}\psi(x, y) &= -b(y) = -\inf_{t, u} \{ t : \bar{R}y + t\bar{p} + \bar{Q}u \leq_{\bar{\mathbf{K}}} \bar{s} \} \\ &= -\sup_{u \in \bar{\mathbf{K}}^*} \left\{ -[\bar{R}^T u]^T y - s^T u : -u^T \bar{p} = 1, \bar{Q}^T u = 0 \right\} \text{ [by conic duality]} \\ &= \inf_{f, t, u} \left\{ f^T y + t : \underbrace{f = R^T u, t = s^T u, \bar{p}^T u + 1 = 0, \bar{Q}^T u = 0, u \geq_{\bar{\mathbf{K}}^*} 0}_{\Leftrightarrow Pf + tp + Qu \leq_{\mathbf{K}} s \text{ with } \mathbf{K} \in \mathcal{K}} \right\}.\end{aligned}$$

3. Bilinear functions:

$$\psi(x, y) \equiv a^T x + b^T y + x^T A y + c \Rightarrow \psi(x, y) = \min_{f, t} \left\{ f^T y + t : \underbrace{f = A^T x + b, t = a^T x + c}_{\Leftrightarrow Pf + tp + Rx \leq s} \right\}.$$

4. “Generalized bilinear functions.” Let $\mathbf{U} \in \mathcal{K}$ and E be the embedding Euclidean space of \mathbf{U} .

(a) Let $\bar{\mathcal{X}}$ be a nonempty \mathcal{K} -representable set, and let continuous mapping $F(x) : \bar{\mathcal{X}} \rightarrow E$ possess \mathcal{K} -representable \mathbf{U} -epigraph⁴

$$\text{Epi}_{\mathbf{U}} F := \{(x, z) \in \bar{\mathcal{X}} \times E, z \geq_{\mathbf{U}} F(x)\} = \{(x, z) : \exists u : \bar{R}x + \bar{S}z + \bar{T}u \leq_{\bar{\mathbf{K}}} \bar{s}\}.$$

Then the function

$$\bar{\psi}(x, y) = y^T F(x) : \bar{\mathcal{X}} \times \mathbf{U}^* \rightarrow \mathbf{R}$$

is \mathcal{K} -representable on $\bar{\mathcal{X}} \times \mathbf{U}^*$:

$$\begin{aligned} \forall (x \in \bar{\mathcal{X}}, y \in \mathbf{U}^*) : \\ \bar{\psi}(x, y) &= y^T F(x) = \inf_f \{f^T y : f \geq_{\mathbf{U}} F(x)\} \\ &= \inf_{f, u} \{f^T y : \bar{R}x + \bar{S}z + \bar{T}u \leq_{\bar{\mathbf{K}}} \bar{s}\}. \end{aligned}$$

(b) Let \mathcal{Y} be a nonempty \mathcal{K} -representable set, and let continuous mapping $G(y) : \mathcal{Y} \rightarrow E$ possess \mathcal{K} -representable \mathbf{U}^* -hypograph,

$$\begin{aligned} \text{Hypo}_{\mathbf{U}^*} G &:= \{(y, w) \in \mathcal{Y} \times E : w \leq_{\mathbf{U}^*} G(y)\} \\ &= \{(y, w) : \exists u : \underline{R}y + \underline{S}w + \underline{Q}u \geq_{\underline{\mathbf{K}}} \underline{s}\} \quad [\underline{\mathbf{K}} \in \mathcal{K}], \end{aligned}$$

the representation being essentially strictly feasible. Then the function

$$\underline{\psi}(\underline{x}, y) = \underline{x}^T G(y) : \mathbf{U} \times \mathcal{Y} \rightarrow \mathbf{R}$$

is \mathcal{K} -representable on $\mathbf{U} \times \mathcal{Y}$:

$$\begin{aligned} \forall (\underline{x} \in \mathbf{U}, y \in \mathcal{Y}) : \\ \underline{\psi}(\underline{x}, y) &= \underline{x}^T G(y) = \sup \{\underline{x}^T w : w \leq_{\mathbf{U}^*} G(y)\} \quad [\text{due to } \underline{x} \in \mathbf{U}] \\ &= \sup_{w, u} \{\underline{x}^T w : \underline{R}y + \underline{S}w + \underline{Q}u \geq_{\underline{\mathbf{K}}} \underline{s}\} \\ &= \inf_{\lambda} \{[\underline{s} - \underline{R}y]^T \lambda : \underline{S}^T \lambda = \underline{x}, \underline{Q}^T \lambda = 0, \lambda \in \underline{\mathbf{K}}^*\} \quad [\text{by conic duality}] \quad (6) \\ &= \inf_{f, t, u = [\lambda; w]} \left\{ f^T y + t : \underbrace{\begin{bmatrix} f + \underline{R}^T \lambda = 0, \underline{s}^T \lambda = t, \\ \underline{S}^T \lambda = \underline{x}, \underline{Q}^T \lambda = 0, \lambda \in \underline{\mathbf{K}}^* \end{bmatrix}}_{\Leftrightarrow Pf + tp + Qu + R\underline{x} \leq_{\underline{\mathbf{K}}} s} \right\} \quad [\underline{\mathbf{K}} \in \mathcal{K}]. \end{aligned}$$

(c) Let \mathcal{Y} and $G(\cdot)$ be as in item 4b) with $G(\mathcal{Y}) \subset \mathbf{U}^*$, let \mathcal{X} be a nonempty \mathcal{K} -representable set, and let

$$F(x) : \mathcal{X} \rightarrow \mathbf{U}$$

be continuous \mathbf{U} -convex mapping with \mathcal{K} -representable \mathbf{U} -epigraph:

$$\begin{aligned} \text{Epi}_{\mathbf{U}} F &:= \{(x, z) : x \in \mathcal{X}, z \geq_{\mathbf{U}} F(x)\} \\ &= \{x, z : \exists v : \hat{R}x + \hat{S}z + \hat{Q}v \leq_{\hat{\mathbf{K}}} \hat{s}\} \quad [\hat{\mathbf{K}} \in \mathcal{K}] \end{aligned} \quad (7)$$

⁴This implies, in particular that the \mathbf{U} -epigraph of F is convex, or, which is the same, that F is \mathbf{U} -convex:

$$\forall (x', x'' \in \bar{\mathcal{X}}, \lambda \in [0, 1]) : F(\lambda x' + (1 - \lambda)x'') \leq_{\mathbf{U}} \lambda F(x') + (1 - \lambda)F(x'').$$

Then the function

$$\psi(x, y) = F^T(x)G(y) : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbf{R}$$

is continuous convex-concave and admits \mathcal{K} -representation as follows:

$$\begin{aligned} \forall(x \in \mathcal{X}, y \in \mathcal{Y}) : \\ \psi(x, y) &= F^T(x)G(y) \\ &= \inf_z \{z^T G(y) : z \geq_{\mathbf{U}} F(x)\} \quad [\text{since } G(y) \in \mathbf{U}^*] \\ &= \inf_{z,v} \left\{ z^T G(y) : \widehat{R}x + \widehat{S}z + \widehat{Q}v \leq_{\widehat{\mathbf{K}}} \widehat{s} \right\} \quad [\text{by (7)}] \\ &= \inf_{z,v} \left\{ \inf_{f,t} \{f^T y + t : Pf + tp + Qu + Rz \leq_{\mathbf{K}} s\} : \widehat{R}x + \widehat{S}z + \widehat{Q}v \leq_{\widehat{\mathbf{K}}} \widehat{s} \right\} \end{aligned}$$

due to (6)—note that on the domain on which $\inf_{z,v}$ is taken we have $z \geq_{\mathbf{U}} F(x) \in \mathbf{U}$, making (6) applicable. We conclude that

$$\psi(x, y) = \inf_{f,t,u=[z;v]} \left\{ f^T y + t : \underbrace{\begin{bmatrix} Pf + tp + Qu + Rz \leq_{\mathbf{K}} s, \\ \widehat{R}x + \widehat{S}z + \widehat{Q}v \leq_{\widehat{\mathbf{K}}} \widehat{s} \end{bmatrix}}_{\Leftrightarrow \widetilde{P}f + \widetilde{t}\widetilde{p} + \widetilde{Q}u + \widetilde{R}x \leq_{\widetilde{\mathbf{K}}} \widetilde{s}} \right\} \quad [\widetilde{\mathbf{K}} \in \mathcal{K}].$$

3.4.2 Basic calculus rules

Basic calculus rules are as follows.

1. [Direct summation] Let $\theta_i > 0$, $i \leq I$, and let

$$\begin{aligned} \forall(x^i \in \mathcal{X}^i, y^i \in \mathcal{Y}^i, i \leq I) : \\ \psi_i(x^i, y^i) = \inf_{f_i, t_i, u_i} \{f_i^T y^i + t_i : P_i f_i + t_i p_i + Q_i u_i + R_i x^i \leq_{\mathbf{K}_i} s_i\} . \end{aligned}$$

Then

$$\begin{aligned} \forall(x = [x^1; \dots; x^I] \in \mathcal{X} = \mathcal{X}_1 \times \dots \times \mathcal{X}_I, y = [y^1; \dots; y^I] \in \mathcal{Y} = \mathcal{Y}_1 \times \dots \times \mathcal{Y}_I) : \\ \psi(x, y) := \sum_i \theta_i \psi_i(x^i, y^i) \\ = \inf_{f,t,u=\{f_i, t_i, u_i, i \leq I\}} \left\{ f^T y + t : \underbrace{\begin{bmatrix} f = [\theta_1 f_1; \dots; \theta_I f_I], t = \sum_i \theta_i t_i, \\ P_i f_i + t_i p_i + Q_i u_i + R_i x^i \leq_{\mathbf{K}_i} s_i, 1 \leq i \leq I \end{bmatrix}}_{\Leftrightarrow Pf + tp + Qu + Rx \leq_{\mathbf{K}} s \text{ with } \mathbf{K} \in \mathcal{K}} \right\} \end{aligned}$$

2. [Affine substitution of variables] Let

$$\begin{aligned} \forall(\xi \in \mathcal{X}^+, \eta \in \mathcal{Y}^+) : \\ \psi_+(\xi, \eta) = \inf_{f_+, t_+, u_+} \{f_+^T \eta + t_+ : P_+ f_+ + t_+ p_+ + Q_+ u_+ + R_+ \xi \leq_{\mathbf{K}_+} s_+\} , \end{aligned}$$

and

$$x \mapsto Ax + b : \mathcal{X} \rightarrow \mathcal{X}^+, y \mapsto By + c : \mathcal{Y} \rightarrow \mathcal{Y}^+.$$

Then

$$\begin{aligned} \forall(x \in \mathcal{X}, y \in \mathcal{Y}) : \\ \psi(x, y) &:= \psi_+(Ax + b, By + c) \\ &= \inf_{f_+, t_+, u_+} \{f_+^T (By + c) + t_+ : P_+ f_+ + t_+ p_+ + Q_+ u_+ + R_+ [Ax + b] \leq_{\mathbf{K}_+} s_+\} \\ &= \inf_{f,t,u=[f_+; t_+; u_+]} \left\{ f^T y + t : \underbrace{\begin{bmatrix} f = B^T f_+, t = t_+ + f_+^T c \\ P_+ f_+ + t_+ p_+ + Q_+ u_+ + R_+ Ax \leq_{\mathbf{K}_+} s_+ - R_+ b \end{bmatrix}}_{\Leftrightarrow Pf + tp + Qu + Rx \leq_{\mathbf{K}} s \text{ with } \mathbf{K} \in \mathcal{K}} \right\} . \end{aligned}$$

3. [Taking conic combinations] This rule, evident by itself, is a combination of the two preceding rules: Let $\theta_i > 0$ and $\psi_i(x, y) : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbf{R}$, $i \leq I$, be such that

$$\forall(x \in \mathcal{X}, y \in \mathcal{Y}) : \psi_i(x, y) = \inf_{f_i, t_i, u_i} \{f_i^T y^i + t_i : P_i f_i + t_i p_i + Q_i u_i + R_i x^i \leq_{\mathbf{K}_i} s_i\}.$$

Then

$$\begin{aligned} \forall(x \in \mathcal{X}, y \in \mathcal{Y}) : \psi(x, y) &:= \sum_i \theta_i \psi_i(x, y) \\ &= \inf_{f, t, u = \{f_i, t_i, u_i, i \leq I\}} \left\{ f^T y + t : \underbrace{\left[\begin{array}{l} f = \sum_i \theta_i f_i, t = \sum_i \theta_i t_i, \\ P_i f_i + t_i p_i + Q_i u_i + R_i x^i \leq_{\mathbf{K}_i} s_i, 1 \leq i \leq I \end{array} \right]}_{\Leftrightarrow Pf + tp + Qu + Rx \leq_{\mathbf{K}} s \text{ with } \mathbf{K} \in \mathcal{K}} \right\}. \end{aligned}$$

4. [Projective transformation in x -variable] Let

$$\forall(x \in \mathcal{X}, y \in \mathcal{Y}) : \psi(x, y) = \inf_{f, t, u} \{f^T y + t : Pf + tp + Qu + Rx \leq_{\mathbf{K}} s\}.$$

Then

$$\begin{aligned} (\forall(\alpha, x) : \alpha > 0, \alpha^{-1}x \in \mathcal{X}, \forall y \in \mathcal{Y}) \\ \bar{\psi}((\alpha, x), y) := \alpha \psi(\alpha^{-1}x, y) = \inf_{f, t, u} \left\{ f^T y + t : Pf + tp + Qu + Rx - \alpha s \leq_{\mathbf{K}} 0 \right\}. \end{aligned}$$

5. [Superposition in x -variable] Let \mathcal{X}, \mathcal{Y} be \mathcal{K} -representable, $\bar{\mathcal{X}}$ be a \mathcal{K} -representable subset of some \mathbf{R}^n , and let $\mathcal{K} \ni \mathbf{U}$ be a cone in \mathbf{R}^n . Furthermore, assume that

$$\bar{\psi}(\bar{x}, y) : \bar{\mathcal{X}} \times \mathcal{Y} \rightarrow \mathbf{R}$$

is a continuous convex-concave function which is \mathbf{U} -nondecreasing in $\bar{x} \in \bar{\mathcal{X}}$, i.e.

$$\forall(y \in \mathcal{Y}, \bar{x}', \bar{x}'' \in \bar{\mathcal{X}} : \bar{x}' \leq_{\mathbf{U}} \bar{x}'') : \bar{\psi}(\bar{x}'', y) \geq \bar{\psi}(\bar{x}', y),$$

and admits \mathcal{K} -representation on $\bar{\mathcal{X}} \times \mathcal{Y}$:

$$\forall(\bar{x} \in \bar{\mathcal{X}}, y \in \mathcal{Y}) : \bar{\psi}(\bar{x}, y) = \inf_{f, t, \bar{u}} \{f^T y + t : \bar{P}f + t\bar{p} + \bar{Q}\bar{u} + \bar{R}\bar{x} \leq_{\bar{\mathbf{K}}} \bar{s}\}.$$

Let also

$$x \mapsto X(x) : \mathcal{X} \mapsto \bar{\mathcal{X}}$$

be a \mathbf{U} -convex mapping such that the intersection of the \mathbf{U} -epigraph of the mapping with $\mathcal{X} \times \bar{\mathcal{X}}$ admits \mathcal{K} -representation:

$$\{(x, \bar{x}) : x \in \mathcal{X}, \bar{x} \in \bar{\mathcal{X}}, \bar{x} \geq_{\mathbf{U}} X(x)\} = \{(x, \bar{x}) : \exists v : Ax + B\bar{x} + Cv \leq_{\bar{\mathbf{K}}} d\}.$$

Then the function $\psi(x, y) = \bar{\psi}(X(x), y)$ admits \mathcal{K} -representation on $\mathcal{X} \times \mathcal{Y}$:

$$\begin{aligned} \forall(x \in \mathcal{X}, y \in \mathcal{Y}) : \psi(x, y) &= \bar{\psi}(X(x), y) = \inf_{\bar{x}} \{\bar{\psi}(\bar{x}, y) : \bar{x} \in \bar{\mathcal{X}} \& \bar{x} \geq_{\mathbf{U}} X(x)\} \\ &= \inf_{f, t, \bar{u}, \bar{x}} \left\{ f^T y + t : \begin{array}{l} \bar{x} \in \bar{\mathcal{X}}, \bar{x} \geq_{\mathbf{U}} X(x) \\ \bar{P}f + t\bar{p} + \bar{Q}\bar{u} + \bar{R}\bar{x} \leq_{\bar{\mathbf{K}}} \bar{s} \end{array} \right\} \\ &= \inf_{f, t, \bar{u}, \bar{x}, v} \left\{ f^T y + t : \begin{array}{l} Ax + B\bar{x} + Cv \leq_{\bar{\mathbf{K}}} d \\ \bar{P}f + t\bar{p} + \bar{Q}\bar{u} + \bar{R}\bar{x} \leq_{\bar{\mathbf{K}}} \bar{s} \end{array} \right\} \\ &= \inf_{f, t, u = \{\bar{u}, \bar{x}, v\}} \left\{ f^T y + t : \underbrace{\left[\begin{array}{l} Ax + B\bar{x} + Cv \leq_{\bar{\mathbf{K}}} d \\ \bar{P}f + t\bar{p} + \bar{Q}\bar{u} + \bar{R}\bar{x} \leq_{\bar{\mathbf{K}}} \bar{s} \end{array} \right]}_{\Leftrightarrow Pf + tp + Qu + Rx \leq_{\mathbf{K}} s \text{ with } \mathbf{K} \in \mathcal{K}} \right\}. \end{aligned}$$

6. [Partial maximization] Let

$$\begin{aligned} & \forall (x \in \mathcal{X}, y = [w; z] \in \mathcal{Y}) : \\ & \psi(x, [w; z]) = \inf_{[g;h],\tau,u} \{g^T w + h^T z + \tau : Gg + Hh + \tau p + Qu + Rx \leq_{\mathbf{K}} s\} \quad [\mathbf{K} \in \mathcal{K}], \end{aligned}$$

and let \mathcal{Y} be compact and given by \mathcal{K} -representation:

$$\mathcal{Y} = \{[w; z] : \exists v : Aw + Bz + Cv \leq_{\mathbf{L}} r\}$$

such that the conic constraint $Bz + Cv \leq_{\mathbf{L}} r - Aw$ in variables z, v is essentially strictly feasible for every $w \in \mathcal{W} = \{w : \exists z : [w; z] \in \mathcal{Y}\}$. Then the function

$$\bar{\psi}(x; w) := \max_z \{\psi(x, [w; z]) : [w; z] \in \mathcal{Y}\} : \mathcal{X} \times \mathcal{W} \rightarrow \mathbf{R}$$

is \mathcal{K} -representable provided it is continuous:⁵

$$\begin{aligned} & \forall (x \in \mathcal{X}, w \in \mathcal{W}) : \\ & \bar{\psi}(x; w) = \max_z \left\{ \inf_{[g;h],\tau,u} [g^T w + h^T z + \tau : Gg + Hh + \tau p + Qu + Rx \leq_{\mathbf{K}} s] : [w; z] \in \mathcal{Y} \right\} \\ & = \inf_{[g;h],\tau,u} \left\{ \max_z [g^T w + h^T z + \tau : [w; z] \in \mathcal{Y}] : Gg + Hh + \tau p + Qu + Rx \leq_{\mathbf{K}} s \right\} \\ & \quad \text{[by the Sion-Kakutani Theorem; note that for } w \in \mathcal{W} \\ & \quad \text{the set } \{z : [w; z] \in \mathcal{Y}\} \text{ is nonempty, convex and compact]} \\ & = \inf_{[g;h],\tau,u} \left\{ \max_{z,v} [g^T w + h^T z + \tau : Bz + Cv \leq_{\mathbf{L}} r - Aw] : Gg + Hh + \tau p + Qu + Rx \leq_{\mathbf{K}} s \right\} \\ & = \inf_{[g;h],\tau,u} \left\{ \min_{\xi} [g^T w + (r - Aw)^T \xi + \tau : B^T \xi = h, C^T \xi = 0, \xi \geq_{\mathbf{L}^*} 0] : \right. \\ & \quad \left. Gg + Hh + \tau p + Qu + Rx \leq_{\mathbf{K}} s \right\} \quad \text{[by conic duality]} \\ & = \inf_{f,t,\bar{u}=[g,h,\tau,u,\xi]} \left\{ f^T w + t : \underbrace{\begin{bmatrix} f = g - A^T \xi, t = r^T \xi + \tau, B^T \xi = h, C^T \xi = 0, \\ \xi \geq_{\mathbf{L}^*} 0, Gg + Hh + \tau p + Qu + Rx \leq_{\mathbf{K}} s \end{bmatrix}}_{\Leftrightarrow \bar{P}f + t\bar{p} + \bar{Q}\bar{u} + \bar{R}x \leq_{\bar{\mathbf{K}}} \bar{s} \text{ with } \bar{\mathbf{K}} \in \mathcal{K}} \right\}. \end{aligned}$$

Note that the last three rules combine with symmetry to induce “symmetric” rules on perspective transformation and superposition in y -variable and partial minimization in x -variable.

3.5 Illustrations

A. Our first illustration is motivated by a statistical application of saddle point optimization—near-optimal recovery of linear forms in Discrete observation scheme, see [10, Section 3.1]. Let

$$\psi(x, y) = \ln \left(\sum_i e^{x_i y_i} \right) : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbf{R},$$

\mathcal{X} and \mathcal{Y} be \mathcal{K} -representable, and let $\mathcal{Y}, 0 \notin \mathcal{Y}$, be a compact subset of the nonnegative orthant. Because for $z > 0$

$$\ln z = \inf_u z e^u - u - 1,$$

for $y \geq 0$ we clearly have

$$\begin{aligned} \ln(\sum_i e^{x_i y_i}) &= \inf_u [(\sum_i e^{x_i y_i}) e^u - u - 1] = \inf_{f,u} [\sum_i y_i f_i - u - 1 : f_i \geq e^{x_i + u}] \\ &= \inf_{f,t,u} \{f^T u + t : f_i \geq e^{x_i + u} \forall i \ \& \ t \geq -u - 1\} \end{aligned}$$

⁵Representation to follow holds true without the latter assumption; we make it to stay consistent with our general definition of representability, where the convex-concave function in question is assumed to be continuous.

The resulting representation is a \mathcal{K} -representation, provided that the closed w.r.t. taking finite direct products and passing to the dual cone family \mathcal{K} of regular cones contains \mathbf{R}_+ , the exponential cone

$$\mathbf{E} = \text{cl} \{[t; s; r] : t \geq se^{r/s}, s > 0\},$$

and, therefore, its dual cone

$$\mathbf{E}^* = \text{cl} \{[\tau; \sigma; -\rho] : \tau > 0, \rho > 0, \sigma \geq \rho \ln(\rho/\tau) - \rho\}.$$

B. Let now

$$\psi(x, y) = \left(\sum_{i=1}^n \theta_i^p(x) y_i \right)^{1/p}$$

where $p > 1$, $\theta_i(x)$ are nonnegative \mathcal{K} -representable real-valued functions on \mathcal{K} -representable set \mathcal{X} , and \mathcal{Y} is a \mathcal{K} -representable subset of the nonnegative orthant. In this case, as is easily seen, for all $(x \in \mathcal{X}, y \in \mathcal{Y})$ it holds

$$\psi(x, y) = \inf_{[f; t]} \left\{ f^T y + t : t \geq 0, f \geq 0, t^{\frac{p-1}{p}} f_i^{\frac{1}{p}} \geq \kappa \theta_i(x), i \leq n \right\} \quad [\kappa = p^{-1}(p-1)^{\frac{p-1}{p}}]$$

which can immediately be converted into \mathcal{K} -representation, provided \mathcal{K} contains 3D Lorentz cone $\mathbf{L}^3 = \{x \in \mathbf{R}^3 : x_3 \geq \sqrt{x_1^2 + x_2^2}\}$ and p is rational, see [4, Section 3.3] or [5, Section 2.3.5].

C. In our next example, $\mathcal{X} \subset \mathbf{R}^{m \times n}$ and $\mathcal{Y} \subset \mathbf{S}_+^m$ are nonempty convex sets, and

$$\psi(x, y) = 2\sqrt{\text{Tr}(x^T y x)} : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbf{R}.$$

Taking into account that for $a \geq 0$ one has $2\sqrt{a} = \inf_{s>0} [a/s + s]$, we have

$$\begin{aligned} \forall (x \in \mathcal{X}, y \in \mathcal{Y}) : \\ \psi(x, y) &= 2\sqrt{\text{Tr}(y[xx^T])} = \inf_g \left\{ 2\sqrt{\text{Tr}(yg)} : g \succeq xx^T \right\} \\ &= \inf_{f, s} \left\{ \text{Tr}(yf) + s : s > 0, fs \succeq xx^T \right\} \\ &= \inf_{f, s} \left\{ \text{Tr}(yf) + s : \left[\begin{array}{c|c} f & x \\ \hline x^T & sI_n \end{array} \right] \succeq 0 \right\}. \end{aligned}$$

The resulting representation is \mathcal{K} -representation, provided that \mathcal{K} contains semidefinite cones.

This is how **C** works in Robust Markowitz Portfolio Selection (cf, e.g., [6, 8])

$$\min_{x \in \mathcal{X}} \max_{y \in \mathcal{Y}} \left[-r^T x + 2\rho \sqrt{x^T y x} \right] \quad [\rho > 0]$$

(here $x \in \mathbf{R}^n$ is the composition of portfolio, r is the vector of expected returns, and y is the uncertain covariance matrix of the returns). Assuming for the sake of definiteness that \mathcal{Y} is cut off \mathbf{S}_+^n by the constraints

$$a_i^T y a_i + b_i y + y b_i^T \preceq p_i, i \leq I, y_- \leq y \leq y_+$$

(where \leq for matrices acts entrywise) and applying our machinery on the top of the above semidefinite representation of $2\sqrt{x^T y x}$, the saddle point problem reduces to

$$\min_{x, s, \alpha_i, \mu_{\pm}} \left\{ \begin{array}{l} -r^T x + \rho [s + \sum_i \text{Tr}(\alpha_i p_i) + \text{Tr}(\mu_+ y_+ - \mu_- y_-)] : \\ \left[\begin{array}{c|c} \sum_i [a_i \alpha_i a_i^T + \alpha_i b_i + b_i^T \alpha_i] + \mu_+ - \mu_- & x \\ \hline x^T & s \end{array} \right] \succeq 0 \\ \alpha_i \succeq 0, i \leq I, \mu_{\pm} \succeq 0, x \in \mathcal{X} \end{array} \right\}.$$

D. In our concluding example, \mathcal{K} contains the products of semidefinite cones, $\mathcal{X} = \mathbf{R}^{m \times n}$, $\mathcal{Y} = \mathbf{S}_+^n$, and

$$\psi(x, y) = \text{Tr}(x^T x y^{1/2}) : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbf{R}.$$

This is a “generalized bilinear function”; in terms of item 4.c of Section 3.4.1, we have $F(x) = x^T x$, $G(y) = y^{1/2}$, $\mathbf{U} = \mathbf{U}^* = \mathbf{S}_+^n$, and

$$\begin{aligned} \text{Epi}_{\mathbf{U}} F &:= \{(x, z) : z \succeq x^T x\} = \left\{ (x, z) : \begin{bmatrix} z & x^T \\ x & I_m \end{bmatrix} \succeq 0 \right\}, \\ \text{Hypo}_{\mathbf{U}^*} G &:= \{(y, w) : y \in \mathbf{S}_+^n, w \preceq y^{1/2}\} = \left\{ (y, w) : \exists v : \begin{bmatrix} y & v \\ v & I_n \end{bmatrix} \succeq 0, v \succeq 0, w \preceq v \right\}. \end{aligned}$$

With these data, the construction from item 4.c of Section 3.4.1 leads straightforwardly to the following semidefinite representation of ψ :

$$\forall (x \in \mathbf{R}^{m \times n}, y \in \mathbf{S}_+^n) : \psi(x, y) := \text{Tr}(x^T x y^{1/2}) = \inf_{f, t, u=(z, \beta, \gamma)} \left\{ \text{Tr}(fy) + t : \begin{array}{l} f \in \mathbf{S}^n, \beta \in \mathbf{R}^{n \times n}, \gamma \in \mathbf{S}^n, z \in \mathbf{S}^n \\ t = \text{Tr}(\gamma), z \preceq \beta + \beta^T \\ \begin{bmatrix} f & \beta \\ \beta^T & \gamma \end{bmatrix} \succeq 0, \begin{bmatrix} z & x^T \\ x & I_m \end{bmatrix} \succeq 0 \end{array} \right\}.$$

4 Well-structured variational inequalities with monotone operators

4.1 Preliminaries

Let \mathcal{X} be a nonempty convex subset of Euclidean space E . A vector field $F(x) : x \rightarrow E$ is called *monotone*, if

$$\langle F(x) - F(y), x - y \rangle \geq 0 \quad \forall x, y \in \mathcal{X}.$$

Variational Inequality $\text{VI}(F, \mathcal{X})$ reads

$$\text{Find } x_* \in \mathcal{X} : \langle F(y), x_* - y \rangle \leq 0 \quad \forall y \in \mathcal{X}. \quad \text{VI}(F, \mathcal{X})$$

Solutions which are sought in $\text{VI}(F, \mathcal{X})$ are called *weak solutions*; they do exist whenever \mathcal{X} is a compact set and F is monotone on \mathcal{X} . *Strong solutions* are points $x_* \in \mathcal{X}$ such that

$$\langle F(x_*), x - x_* \rangle \geq 0 \quad \forall x \in \mathcal{X};$$

from monotonicity of F it follows that every strong solution is a weak one. The opposite is true provided that F is continuous on \mathcal{X} .

A natural (in)accuracy measure for a candidate approximate solution to $\text{VI}(F, \mathcal{X})$ is the *dual gap function*

$$\epsilon_{\text{VI}}[x|\mathcal{X}] = \sup_{y \in \mathcal{X}} \langle F(y), x - y \rangle;$$

this function is nonnegative and is zero exactly at weak solutions to $\text{VI}(F, \mathcal{X})$.

Our current goal is to develop a framework for converting “well-structured” VI’s with monotone vector fields into the usual conic problems, thus bringing them within the grasp of Interior-Point polynomial time methods and software like CVX. What follows can be seen as a streamlined “well-structured” (i.e., conic representation-oriented) extension of what in [13, Section 7.4] was called a convex representation of a monotone operator.

4.2 Conic representability of monotone vector fields and monotone VI's in conic form

4.2.1 Conic representation of a monotone vector field

Let $F(x) : \mathcal{X} \rightarrow E$ be a monotone and continuous vector field on nonempty convex subset \mathcal{X} of Euclidean space E . Consider the set

$$\mathcal{F}[F, \mathcal{X}] = \{[t; g; x] \in \mathbf{R} \times E \times E : x \in \mathcal{X}, t - \langle g, y \rangle \geq \langle F(y), x - y \rangle \forall y \in \mathcal{X}\} \quad (8)$$

and let us make several straightforward observations.

4.2.1.A. $\mathcal{F}[F, \mathcal{X}]$ is a convex set which contains all triples $[\langle F(x), x \rangle; F(x); x]$, $x \in \mathcal{X}$; this set is closed provided \mathcal{X} is so. Besides this, $\mathcal{F}[F, \mathcal{X}]$ is t -monotone:

$$[t; g; x] \in \mathcal{F}[F, \mathcal{X}] \text{ and } t' \geq t \Rightarrow [t'; g; x] \in \mathcal{F}[F, \mathcal{X}].$$

Indeed, $\mathcal{F}[F, \mathcal{X}]$ is the intersection of solution set of the system of nonstrict linear constraints

$$t - \langle g, y \rangle - \langle F(y), x \rangle \geq -\langle F(y), y \rangle, y \in \mathcal{X}$$

in variables g, t, x (this set is closed and convex) with the convex set $\{[t; g; x] : x \in \mathcal{X}\}$ (which is closed if so is \mathcal{X}). By monotonicity, for every $x \in \mathcal{X}$ we have

$$\langle F(x), x - y \rangle \geq \langle F(y), x - y \rangle \forall y \in \mathcal{X},$$

so that $[\langle F(x), x \rangle; F(x); x] \in \mathcal{F}[F, \mathcal{X}]$; and t -monotonicity is evident.

4.2.1.B. For $\epsilon \geq 0$, let

$$\mathcal{X}_*(\epsilon) = \{[g; t] \in E \times \mathbf{R} : \sup_{y \in \mathcal{X}} [t - \langle g, y \rangle] \leq \epsilon\}, \quad (9)$$

so that $\mathcal{X}_*(\epsilon)$ is a nonempty closed convex set. Then for every $\epsilon \geq 0$, ϵ -solutions to $\text{VI}(F, \mathcal{X})$ —points $x \in \mathcal{X}$ such that

$$\epsilon_{\text{VI}}[x|F] := \sup_{y \in \mathcal{X}} \langle F(y), x - y \rangle \leq \epsilon$$

are exactly the points

$$x : \exists(t, g) : [t; g; x] \in \mathcal{F}[F, \mathcal{X}] \text{ and } [t; g] \in \mathcal{X}_*(\epsilon); \quad (10)$$

Indeed, let x, t, g be such that $[t; g; x] \in \mathcal{F}[F, \mathcal{X}]$ and $[t; g] \in \mathcal{X}_*(\epsilon)$. From the first inclusion it follows that $x \in \mathcal{X}$ and

$$t - \langle g, y \rangle \geq \langle F(y), x - y \rangle \forall y \in \mathcal{X},$$

while from the second inclusion it follows that $t - \langle g, y \rangle \leq \epsilon$ for all $y \in \mathcal{X}$. Taken together, these two relations imply that $x \in \mathcal{X}$ and $\langle F(y), x - y \rangle \leq \epsilon$ for all $y \in \mathcal{X}$, i.e., $\epsilon_{\text{VI}}[x|F] \leq \epsilon$. Vice versa, if $x \in \mathcal{X}$, $\epsilon \geq 0$, and $\epsilon_{\text{VI}}[x|F] \leq \epsilon$, then the triple $t = \epsilon, g = 0, x$ clearly belongs to $\mathcal{F}[F, \mathcal{X}]$ and $[t; g] \in \mathcal{X}_*(\epsilon)$.

\mathcal{K} -representation of (F, \mathcal{X}) . Given a continuous monotone vector field $F : \mathcal{X} \rightarrow E$ on a nonempty convex subset \mathcal{X} of Euclidean space E , let us call a conic constraint⁶

$$Xx + Gg + tT + Uu \leq_{\mathbf{K}} a \quad (11)$$

in variables $t \in \mathbf{R}$, $g \in E$, $x \in E$, and $u \in \mathbf{R}^k$ a \mathcal{K} -representation of (F, \mathcal{X}) , if the set

$$\mathcal{T} := \{[t; g; x] : \exists u : Xx + Gg + tT + Uu \leq_{\mathbf{K}} a\} \quad (12)$$

possesses the following two properties:

⁶Recall that we have fixed a family \mathcal{K} of regular cones in Euclidean spaces, and by our standing convention, all cones to be considered below belong to \mathcal{K} .

(i) \mathcal{T} is contained in $\mathcal{F}[F, \mathcal{X}]$ and, along with the latter set, is “ t -monotone:”

$$[t; g; x] \in \mathcal{T} \text{ and } t' \geq t \Rightarrow [t'; g; x] \in \mathcal{T}$$

(note that when $\mathcal{T} \neq \emptyset$, t -monotonicity is equivalent to $T \leq_{\mathbf{K}} 0$);

and

(ii) \mathcal{T} contains the set

$$\{[\langle F(x), x \rangle; F(x); x], x \in \mathcal{X}\} \quad [\subset \mathcal{F}[F, \mathcal{X}]]$$

If the set (12) satisfies (i) and the relaxed version of (ii), specifically,

(ii') \mathcal{T} contains the set

$$\{[t; F(x); x] : t > \langle F(x), x \rangle, x \in \mathcal{X}\} \quad [\subset \mathcal{F}[F, \mathcal{X}]]$$

we say that (11) is an *almost \mathcal{K} -representation* of (F, \mathcal{X}) .

Let us make an immediate observation:

Remark 4.1 *In the situation described in the beginning of this section, let \mathcal{X} be closed, and let \mathcal{Y} be a convex set such that*

$$\text{Conv}\{[x^T F(x); F(x); x], x \in \mathcal{X}\} \subset \mathcal{Y} \subset \text{cl Conv}\{[x^T F(x); F(x); x], x \in \mathcal{X}\}$$

Then every \mathcal{K} -representation of \mathcal{Y} represents (F, \mathcal{X}) .

4.2.2 Conic form of conic-representable monotone VI

Our main observation is as follows:

Proposition 4.1 *Let $\mathcal{X} \subset E$ be nonempty convex compact given by essentially strictly feasible \mathcal{K} -representation:*

$$\mathcal{X} = \{x : \exists v : Ax + Bv \leq_{\mathbf{L}} b\}, \quad [\mathbf{L} \in \mathcal{K}], \quad (13)$$

so that $\mathcal{X}_(\epsilon)$, see (9), by conic quality, admits \mathcal{K} -representation as follows:*

$$\mathcal{X}_*(\epsilon) = \{[t; g] : \exists \lambda : A^T \lambda + g = 0, B^T \lambda = 0, t + b^T \lambda \leq \epsilon, \lambda \geq_{\mathbf{L}^*} 0\}. \quad (14)$$

Let, moreover, $F : \mathcal{X} \rightarrow E$ be a continuous monotone vector field, and let (11) be an almost \mathcal{K} -representation of (F, \mathcal{X}) . Then for every $\epsilon > 0$ the system of conic constraints

$$\begin{aligned} Xx + Gg + tT + Uu &\leq_{\mathbf{K}} a & (a) \\ A^T \lambda + g &= 0 & (b.1) \\ B^T \lambda &= 0 & (b.2) \\ t + b^T \lambda &\leq \epsilon & (b.3) \\ \lambda &\geq_{\mathbf{L}^*} 0 & (b.4) \end{aligned} \quad (15)$$

in variables x, g, t, u , and λ is feasible, and x -component of any feasible solution belongs to \mathcal{X} and is an ϵ -solution to $\text{VI}(F, \mathcal{X})$:

$$\epsilon_{\text{VI}}[x|F] \leq \epsilon.$$

Therefore, finding an ϵ -solution to $\text{VI}(F, \mathcal{X})$ reduces to finding a feasible solution to an explicit \mathcal{K} -conic constraint.

When (11) is a \mathcal{K} -representation of (F, \mathcal{X}) , the above conclusion holds true for all $\epsilon \geq 0$.

Proof. Let us fix $\epsilon > 0$, and let \bar{x} solve $\text{VI}(F, X)$ (a solution exists since \mathcal{X} is compact). Then $\langle F(\bar{x}), \bar{x} - y \rangle \leq 0$ for all $y \in \mathcal{X}$ due to continuity of F on \mathcal{X} . Consequently, given $\epsilon > 0$, setting $\bar{t} = \langle F(\bar{x}), \bar{x} \rangle$, $\bar{g} = F(\bar{x})$, we have $[\bar{t} + \epsilon; \bar{g}; \bar{x}] \in \mathcal{X}_*(\epsilon)$, implying by (14) that there exists $\bar{\lambda}$ such that $t = \bar{t} + \epsilon$, $g = \bar{g}$, $\lambda = \bar{\lambda}$ solve (15.b.1-4). Besides this, by (ii') there exists \bar{u} such that $x = \bar{x}$, $g = \bar{g}$, $t = \bar{t} + \epsilon$, $u = \bar{u}$ solve (15.a). Thus, (15) is feasible.

Next, if x, g, t, u, λ is a feasible solution to (15), then by (15.a) one has $[t; g; x] \in \mathcal{T}$, with \mathcal{T} given by (12), whence $[t; g; x] \in \mathcal{F}[F, \mathcal{X}]$, implying that $x \in \mathcal{X}$, see (8). Besides this, (15.b.1-4) say that $[t; g] \in \mathcal{X}_*(\epsilon)$. Thus, $[t; g; x] \in \mathcal{F}[F, \mathcal{X}]$ and $[t; g] \in \mathcal{X}_*(\epsilon)$, implying by 4.2.1.B that $\epsilon_{\text{VI}}[x|F] \leq \epsilon$.

Finally, when (11) is a \mathcal{K} -representation of (F, \mathcal{X}) , the above reasoning works for $\epsilon = 0$. \square

4.3 Calculus of conic representations of monotone vector fields

\mathcal{K} -representations of pairs (F, \mathcal{X}) (\mathcal{X} is a nonempty convex subset of Euclidean space E , $F : \mathcal{X} \rightarrow E$ is a continuous monotone vector field) admit a calculus; for verification of claims to follow, see Appendix A.

4.3.1 Raw materials

Raw materials for the calculus of \mathcal{K} -representable monotone vector fields include:

1. [Affine monotone vector field] Let \mathcal{K} contain all Lorentz cones. Then affine monotone vector field $F(x) = Ax + a$ on $\mathcal{X} = E = \mathbf{R}^n$ is \mathcal{K} -represented by conic constraint (cf. [13, Proposition 7.4.3])

$$t \geq x^T \bar{A}x + a^T x, g = Ax + a \quad (16)$$

in variables t, g, x , where $\bar{A} = \frac{1}{2}[A + A^T]$.

2. [Gradient field of continuously differentiable \mathcal{K} -representable convex function] Let \mathcal{X} be nonempty convex compact set, and $f(x) : \mathcal{X} \rightarrow \mathbf{R}$ be continuously differentiable convex function with \mathcal{K} -representable epigraph,

$$\{(x, s) : s \geq f(x), x \in \mathcal{X}\} = \{(x, s) : \exists u : Ax + sp + Qu \leq_{\mathbf{K}} a\}, \quad (17)$$

the representation being essentially strictly feasible. Then the conic constraint

$$t \geq s + r, Ax + sp + Qu \leq_{\mathbf{K}} a, r \geq a^T \lambda, A^T \lambda = g, p^T \lambda = -1, Q^T \lambda = 0, \lambda \geq_{\mathbf{K}^*} 0 \quad (18)$$

in variables t, g, x, s, r, λ , and u represents $(F(\cdot) := f'(\cdot), \mathcal{X})$ (cf. [13, Proposition 7.4.4]).

3. [Monotone vector field associated with continuously differentiable \mathcal{K} -representable convex-concave function ψ] Let \mathcal{K} contain the 3D Lorentz cone, and let $\mathcal{U} \subset \mathbf{R}^{n_u}$ and $\mathcal{V} \subset \mathbf{R}^{n_v}$ be \mathcal{K} -representable nonempty compact sets:

$$\begin{aligned} \mathcal{U} &= \{u : \exists \alpha : Au + B\alpha \leq_{\mathbf{K}} a\}, & (a) \\ \mathcal{V} &= \{v : \exists \beta : Cv + D\beta \leq_{\mathbf{L}} b\}, & (b) \end{aligned} \quad (19)$$

both representations being essentially strictly feasible. Assume that $\psi(u, v) : \mathcal{U} \times \mathcal{V} \rightarrow \mathbf{R}$ is a continuously differentiable convex-concave function which is \mathcal{K} -representable on $\mathcal{U} \times \mathcal{V}$ with essentially strictly feasible representation (see Section 3.1); that is, ψ admits representation

$$\forall (u \in \mathcal{U}, v \in \mathcal{V}) : \psi(u, v) = \inf_{f, \tau, \xi} \{f^T v + \tau : Pf + \tau p + Q\xi + Ru \leq_{\mathbf{M}} c\} \quad (20)$$

such that the conic constraint $Pf + \tau p + Q\xi \leq_{\mathbf{M}} c - Ru$ in variables f, τ, ξ is essentially strictly feasible for every $u \in \mathcal{U}$. Let

$$F(u, v) = [\nabla_u \psi(u, v); -\nabla_v \psi(u, v)] : \mathcal{X} := \mathcal{U} \times \mathcal{V} \rightarrow E := \mathbf{R}^{n_u} \times \mathbf{R}^{n_v}$$

be the vector field associated with $\mathcal{U}, \mathcal{V}, \psi$; it is well known that this field is monotone. Then

(i) The set

$$\mathcal{Z} = \left\{ \begin{array}{l} (t \in \mathbf{R}, g = [h; e] \in E, x = [u; v] \in \mathcal{X}) : \exists r, s : \\ r > \max_{\zeta \in \mathcal{V}} [\zeta^T e + \psi(u, \zeta)] \quad (a) \\ s > \max_{\omega \in \mathcal{U}} [\omega^T h - \psi(\omega, v)] \quad (b) \\ t \geq r + s \quad (c) \end{array} \right\} \quad (21)$$

satisfies the relations

$$\begin{array}{l} \mathcal{Z} \subset \mathcal{F}[F, \mathcal{X}] \quad (a) \\ x \in \mathcal{X}, g = F(x), t > x^T F(x) \Rightarrow [t; g; x] \in \mathcal{Z}. \quad (b) \end{array} \quad (22)$$

(ii) Besides this, \mathcal{Z} is nothing but the projection on the plane of $(t, g = [h; e], x = [u; v])$ -variables of the solution set of the conic constraint

$$\left[\begin{array}{l} t \geq r + s, r \geq \theta + \gamma^T b + \tau, s \geq \theta' + c^T \delta + a^T \epsilon, \\ Au + B\alpha \leq_{\mathbf{K}} a, Cv + D\beta \leq_{\mathbf{L}} b, Pf + \tau p + Q\xi + Ru \leq_{\mathbf{M}} c \\ C^T \gamma = f + e, D^T \gamma = 0, P^T \delta + v = 0, p^T \delta = -1, Q^T \delta = 0, R^T \delta + A^T \epsilon = h, B^T \epsilon = 0, \\ \gamma \geq_{\mathbf{L}^*} 0, \delta \geq_{\mathbf{M}^*} 0, \epsilon \geq_{\mathbf{K}^*} 0, \eta \geq 0, \theta \geq 0, \eta\theta \geq 1, \eta' \geq 0, \theta' \geq 0, \eta'\theta' \geq 1 \end{array} \right] \quad (23)$$

in variables $t, g = [h; e], x = [u; v], r, s, f, \tau, \xi, \alpha, \beta, \gamma, \eta, \theta, \delta, \epsilon, \eta'$, and θ' .⁷

Taken together, (i) and (ii) say that the conic constraint (23) almost represents (F, \mathcal{X}) .

4. [Univariate monotone rational vector field] Let $\mathcal{X} = [a, b] \subset \mathbf{R}$ ($-\infty < a < b < \infty$), and let $\alpha(x)$ and $\beta(x)$ be real algebraic polynomials such that $\beta(x) > 0$ on \mathcal{X} . Suppose that the univariate vector field given on \mathcal{X} by the function $F(x) = \frac{\alpha(x)}{\beta(x)}$ is nondecreasing on \mathcal{X} , and that \mathcal{K} contain all semidefinite cones. Then the set

$$\mathcal{Y} := \text{Conv}\{[xF(x); F(x); x], x \in \mathcal{X}\}$$

admits explicit \mathcal{K} -representation which, by Remark 4.1, represents (F, \mathcal{X}) .

4.3.2 Calculus rules

Calculus rules are as follows.

1. [Restriction on a \mathcal{K} -representable set] Let conic constraint

$$Xx + Gg + tT + Uu \leq_{\mathbf{K}} a \quad (24)$$

in variables t, g, x, u represent (almost represent) (F, \mathcal{X}) , and let set $\mathcal{Y} \subset E$ be \mathcal{K} -representable:

$$\mathcal{Y} = \{y \in E : \exists v : Ay + Bv \leq_{\mathbf{L}} b\} \quad (25)$$

and have a nonempty intersection $\mathcal{Z} = \mathcal{Y} \cap \mathcal{X}$ with \mathcal{X} . Denoting by \bar{F} the restriction of F on \mathcal{Z} , the conic constraint

$$\{Xx + Gg + tT + Uu \leq_{\mathbf{K}} a, Ax + Bv \leq_{\mathbf{L}} b\} \quad (26)$$

in variables t, g, x, u, v represents (resp., almost represent) (\bar{F}, \mathcal{Z}) (cf. [13, Proposition 7.4.5]).

2. [Direct summation] For $k \leq K$, let \mathcal{X}_k be nonempty convex subsets of Euclidean spaces E_k and $F_k : \mathcal{X}_k \rightarrow E_k$ be continuous monotone vector fields, and let

$$X_k x + G_k g_k + t_k T_k + U_k u_k \leq_{\mathbf{K}_k} a_k, 1 \leq k \leq K \quad (27)$$

⁷When one of the authors was a freshman at Math. Department of Moscow State University, his mate left lecture on Linear Algebra writing in his notebook: "The lecture was terminated due to shortage of indices." The list of variables in (23) comes close to this natural limit...

be \mathcal{K} -representations of (F_k, \mathcal{X}_k) . Denote

$$F([x_1; \dots; x_K]) = [F_1(x_1); \dots; F_K(x_K)] \text{ and } \mathcal{X} = \mathcal{X}_1 \times \dots \times \mathcal{X}_K.$$

Then the conic constraint

$$\begin{aligned} X_k x_k + G_k g_k + t_k T_k + U_k u_k &\leq_{\mathbf{K}_k} a_k, k \leq K & (a) \\ t &= \sum_k t_k & (b) \end{aligned} \quad (28)$$

in variables $t, g := [g_1; \dots; g_K], x := [x_1; \dots; x_K], t_1, \dots, t_K, u_1, \dots, u_K$ \mathcal{K} -represents (F, \mathcal{X}) .

When conic constraints (27) almost represent (F_k, \mathcal{X}_k) , (28) almost represents (F, \mathcal{X}) (cf. [13, Proposition 7.4.6]).

3. [Taking conic combinations] Let \mathcal{X} be a nonempty convex subset of Euclidean space E , and let F_1, \dots, F_K be continuous monotone vector fields on X with (F_k, \mathcal{X}) admitting \mathcal{K} -representations

$$X_k x + G_k g + t T_k + U_k u_k \leq_{\mathbf{K}_k} a_k, 1 \leq k \leq K. \quad (29)$$

Let, further, $\alpha_k > 0$ be given, and let

$$F(x) = \sum_k \alpha_k F_k(x) : \mathcal{X} \rightarrow E.$$

The conic constraint

$$\begin{aligned} X_k x + G_k g_k + t_k T_k + U_k u_k &\leq_{\mathbf{K}_k} a_k, 1 \leq k \leq K & (a) \\ g &= \sum_k \alpha_k g_k & (b) \\ t &= \sum_k \alpha_k t_k & (c) \end{aligned} \quad (30)$$

in variables $t, g, x, \{t_k, g_k, u_k, k \leq K\}$ is a \mathcal{K} -representation of (F, \mathcal{X}) .

When (29) are almost representations of (F_k, \mathcal{X}) , (30) is almost representation of (F, \mathcal{X}) .

4. [Affine substitution of variables] Let \mathcal{X} be a nonempty convex subset of E , F be a continuous monotone vector field on X , with (F, \mathcal{X}) given by \mathcal{K} -representation

$$Xx + Gg + tT + Uu \leq_{\mathbf{K}} a. \quad (31)$$

Assume that $\xi \mapsto A\xi + a$ is an affine mapping from Euclidean space Λ to E , and

$$\Xi = \{\xi : A\xi + a \in \mathcal{X}\}.$$

Then vector field $\Phi(\xi) = A^T F(A\xi + a) : \Xi \rightarrow \Lambda$ is continuous and monotone, and the conic constraint

$$\tau = t - \langle g, a \rangle, \gamma = A^T g, X(A\xi + a) + Gg + tT + Uu \leq_{\mathbf{K}} a \quad (32)$$

in variables τ, γ, ξ, g, t , and u represents (Φ, Ξ) .

When (31) nearly represents (F, \mathcal{X}) , conic constraint (32) nearly represents (Φ, Ξ) .

4.4 Illustrations

4.4.1 “Academic” illustration

Let \mathcal{K} contain Lorentz cones, $M \in \mathbf{R}^{n \times n}$ be such that $\overline{M} = \frac{1}{2}[M + M^T] \succeq 0$, and let $\mathcal{X} \subset \mathbf{R}_+^n$ be nonempty and \mathcal{K} -representable:

$$\mathcal{X} = \{x : \exists u_0 : A_0 x + B_0 u_0 \leq_{\mathbf{K}_0} a_0\}. \quad (33)$$

Suppose that the operator

$$F(x) = Mx - [f_1(x); \dots; f_n(x)] \quad (34)$$

is monotone on \mathcal{X} , and that functions $f_i(x)$ and $-\sum_i x_i f_i(x)$ are \mathcal{K} -representable on \mathcal{X} :

$$\begin{aligned} \{(x, s) : x \in \mathcal{X}, s \geq f_i(x)\} &= \{(x, s) : \exists u_{1i} : A_{1i}x + sB_{1i} + C_{1i}u_{1i} \leq_{\mathbf{K}_{1i}} a_{1i}\} & (a_i) \\ \{(x, s) : x \in \mathcal{X}, s \geq -\sum_i x_i f_i(x)\} &= \{(x, s) : \exists u_2 : A_2x + sB_2 + C_2u_2 \leq_{\mathbf{K}_2} a_2\} & (b) \end{aligned} \quad (35)$$

Observe, that when $M \succeq \alpha I_m$ with $\alpha > 0$, F definitely is monotone provided that $f(x) := [f_1(x); \dots; f_n(x)]$ is Lipschitz continuous on \mathcal{X} with Lipschitz constant w.r.t. $\|\cdot\|_2$ bounded with α .

Given $x \in \mathcal{X}$ and $g \in \mathbf{R}^n$, let us consider the function

$$f_{x,g}(y) = g^T y + F^T(y)(x - y) = -y^T \overline{M} y + \sum_i [g_i y_i + x_i [M y]_i - f_i(y)] : \mathcal{X} \rightarrow \mathbf{R}.$$

We are in the situation where $x_i \geq 0$ for $x \in \mathcal{X}$, so that

$$\begin{aligned} & \{(y, r) : y \in \mathcal{X}, f_{x,g}(y) \geq r\} \\ = & \left\{ (y, r) : \exists u_0, s_0, s_{1i}, s_2 : \begin{aligned} & A_0 y + B_0 u_0 \leq_{\mathbf{K}_0} a_0, \quad y^T \overline{M} y - s_0 \leq 0 \\ & s_{1i} + f_i(y) \leq 0, \quad 1 \leq i \leq n, \quad s_2 - \sum_i y_i f_i(y) \leq 0 \\ & \sum_i [g_i + [M^T x]_i] y_i - s_0 + \sum_i x_i s_{1i} + s_2 \geq r \end{aligned} \right\} \\ = & \left\{ (y, r) : \exists u_0, s_0, s_{1i}, s_2, u_{1i}, u_2 : \begin{aligned} & A_0 y + B_0 u_0 \leq_{\mathbf{K}_0} a_0, \quad y^T \overline{M} y - s_0 \leq 0, \quad A_2 y - s_2 B_2 + C_2 u_2 \leq_{\mathbf{K}_2} a_2 \\ & A_{1i} y - s_{1i} B_{1i} + C_{1i} u_{1i} \leq_{\mathbf{K}_{1i}} a_{1i}, \quad 1 \leq i \leq n \\ & \sum_i [g_i + [M^T x]_i] y_i - s_0 + \sum_i x_i s_{1i} + s_2 \geq r \end{aligned} \right\} \end{aligned}$$

Taking into account that $\overline{M} \succeq 0$, the constraint $y^T \overline{M} y - s_0 \leq 0$ can be represented by strictly feasible conic constraint

$$A_3 y + s_0 B_3 \leq_{\mathbf{K}_3} a_3$$

on Lorentz cone \mathbf{L}^3 , we get

$$\begin{aligned} \max_{y \in \mathcal{X}} f_{x,g}(y) &= \sup_{y, u_0, s_0, s_{1i}, s_2, u_{1i}, u_2} \left\{ \sum_i [g_i + [M^T x]_i] y_i - s_0 + \sum_i x_i s_{1i} + s_2 : \right. \\ & \left. \begin{aligned} & A_0 y + B_0 u_0 \leq_{\mathbf{K}_0} a_0, \quad A_3 y + s_0 B_3 \leq_{\mathbf{K}_3} a_3 \\ & A_{1i} y - s_{1i} B_{1i} + C_{1i} u_{1i} \leq_{\mathbf{K}_{1i}} a_{1i}, \quad 1 \leq i \leq n, \quad A_2 y - s_2 B_2 + C_2 u_2 \leq_{\mathbf{K}_2} a_2 \end{aligned} \right\}. \end{aligned} \quad (36)$$

Assume that the system of conic constraints (36) in variables $y, s_0, u_0, s_{1i}, s_2, u_{1i}$ and u_2 is essentially strictly feasible. Then, by conic quality,

$$\begin{aligned} \max_{y \in \mathcal{X}} f_{x,g}(y) &= \min_{\mu, \nu, \xi_i, \eta} \left\{ a_0^T \mu + a_3^T \nu + \sum_i a_{1i}^T \xi_i + a_2^T \eta : \right. \\ & \left. \begin{aligned} & M^T x + g = A_0^T \mu + A_3^T \nu + \sum_i A_{1i}^T \xi_i + A_2^T \eta \\ & x_i + B_{1i}^T \xi_i = 0, \quad 1 \leq i \leq n, \quad B_2^T \eta = -1, \quad B_3^T \nu = -1 \\ & B_0^T \mu = 0, \quad C_{1i}^T \xi_i = 0, \quad 1 \leq i \leq n, \quad C_2^T \eta = 0, \\ & \mu \geq_{\mathbf{K}_0^*} 0, \quad \nu \geq_{\mathbf{K}_3^*} 0, \quad \xi_i \geq_{\mathbf{K}_{1i}^*} 0, \quad 1 \leq i \leq n, \quad \eta \geq_{\mathbf{K}_2^*} 0 \end{aligned} \right\}. \end{aligned}$$

Now, recalling what $f_{x,g}(y)$ is, we end up with \mathcal{K} -representation of $\mathcal{F}[F, \mathcal{X}]$:

$$\begin{aligned} \mathcal{F}[F, \mathcal{X}] &:= \{[t; g; x] : x \in \mathcal{X}, t - g^T y \geq F^T(y)[x - y] \quad \forall y \in \mathcal{X}\} \\ &= \left\{ [t; g; x] : \exists u_0, \mu, \nu, \xi_i, \eta : \right. \\ & \left. \begin{aligned} & t \geq a_0^T \mu + a_3^T \nu + \sum_i a_{1i}^T \xi_i + a_2^T \eta, \quad A_0 x + B_0 u_0 \leq_{\mathbf{K}_0} a_0 \\ & M^T x + g = A_0^T \mu + A_3^T \nu + \sum_i A_{1i}^T \xi_i + A_2^T \eta \\ & x_i + B_{1i}^T \xi_i = 0, \quad 1 \leq i \leq n, \quad B_2^T \eta = -1, \quad B_3^T \nu = -1 \\ & B_0^T \mu = 0, \quad C_{1i}^T \xi_i = 0, \quad 1 \leq i \leq n, \quad C_2^T \eta = 0 \\ & \mu \geq_{\mathbf{K}_0^*} 0, \quad \nu \geq_{\mathbf{K}_3^*} 0, \quad \xi_i \geq_{\mathbf{K}_{1i}^*} 0, \quad 1 \leq i \leq n, \quad \eta \geq_{\mathbf{K}_2^*} 0 \end{aligned} \right\}, \end{aligned}$$

so that the conic constraint

$$\left[\begin{array}{l} t \geq a_0^T \mu + a_3^T \nu + \sum_i a_{1i}^T \xi_i + a_2^T \eta, A_0 x + B_0 u_0 \leq_{\mathbf{K}_0} a_0 \\ x_i + B_{1i}^T \xi_i = 0, 1 \leq i \leq n, B_2^T \eta = -1, B_3^T \nu = -1 \\ B_0^T \mu = 0, C_{1i}^T \xi_i = 0, 1 \leq i \leq n, C_2^T \eta = 0 \\ \mu \geq_{\mathbf{K}_0^*} 0, \nu \geq_{\mathbf{K}_3^*} 0, \xi_i \geq_{\mathbf{K}_{1i}^*} 0, 1 \leq i \leq n, \eta \geq_{\mathbf{K}_2^*} 0 \end{array} \right]$$

in variables $t, g, x, u_0, \mu, \nu, \xi_i$ and η is a \mathcal{K} -representation of (F, \mathcal{X}) .

4.4.2 Nash Equilibrium

The “covering story” for this example is as follows:

$n \geq 2$ retailers intend to enter certain market, say, one of red herrings. To this end they should select their “selling capacities” (say, rent areas at malls) $x_i, 1 \leq i \leq n$, in given ranges $\mathcal{X}_i = [0, X_i]$ ($0 < X_i < \infty$). With the selections $x = [x_1; \dots; x_n] \in \mathcal{X} = \mathcal{X}_1 \times \dots \times \mathcal{X}_n \subset \mathbf{R}_+^n$, the monthly loss of the i -th retailer is

$$\phi_i(x) = c_i x_i - \frac{x_i}{\sum_{j=1}^n x_j + a} b,$$

where $c_i x_i, c_i > 0$, is the price of the capacity $x_i, b > 0$ is the dollar value of the demand on red herrings, and $a > 0$ is the total selling capacity of the already existing retailers; the term $-\frac{x_i}{\sum_{j=1}^n x_j + a} b$ is the minus the revenue of the i -th retailer under the assumption that the total demand is split between retailers proportionally to their selling capacities.

We want to find is a Nash equilibrium – a point $x^* = [x_1^*; \dots; x_n^*] \in \mathcal{X}$ such that every one of the functions $\phi_i(x_i^*, \dots, x_{i-1}^*, x_i, x_{i+1}^*, \dots, x_n^*)$ attains its minimum over $x_i \in \mathcal{X}_i$ at $x_i = x_i^*$, so that for the i -th retailer there is no incentive to deviate from capacity x_i^* provided that all other retailers j stick to capacities x_j^* , and this is so for all i .

As is immediately seen, the Nash Equilibrium problem in question is convex, meaning that $\phi_i(x)$ are convex in x_i and concave in $x^i = (x_i, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$ and, on the top of it, $\sum_i \phi_i(x)$ is convex on \mathcal{X} . It is well known (for justification, see, e.g., [14]) that for such a problem, the vector field

$$F(x) = \left[\frac{\partial \phi_1(x)}{\partial x_1}; \frac{\partial \phi_2(x)}{\partial x_2}; \dots; \frac{\partial \phi_n(x)}{\partial x_n} \right] : \mathcal{X} \rightarrow \mathbf{R}^n$$

is monotone, and that Nash Equilibria are exactly the weak=strong solutions to $\text{VI}(F, \mathcal{X})$. Specifying \mathcal{K} as the family of direct products of Lorentz cones, we are about to demonstrate that F admits an explicit \mathcal{K} -representation on \mathcal{X} , allowing to reduce the problem of finding Nash Equilibrium to an explicit Second Order conic problem. This is how the construction goes (to save notation, in what follows we set $b = a = 1$, which always can be achieved by appropriate scaling of the capacities and loss functions):

1. Observe that all we need is a \mathcal{K} -representation $\mathcal{R}(t, g, x, u)$ of (F, \mathcal{X}) —a \mathcal{K} -conic constraint in variables $t \in \mathbf{R}, g \in \mathbf{R}^n, x \in \mathbf{R}^n$ and additional variables u —satisfying the requirements specified in Section 4.2.1. By Proposition 4.1, given \mathcal{R} , for $\epsilon \geq 0$ we can write down an explicit \mathcal{K} -conic constraint $\mathcal{C}_\epsilon(t, g, x, u, v)$ in variables t, g, x and additional variables u, v , with the size of the constraint (dimension of the associated cone and the total number of variables) independent of ϵ such that the constraint is feasible, and the x -component of any feasible solution to the constraint is an ϵ -solution to $\text{VI}(F, \mathcal{X})$:

$$x \in \mathcal{X} \ \& \ \epsilon_{\text{VI}}[x|F] = \max_{y \in \mathcal{X}} F^T(y)[x - y] \leq \epsilon.$$

As a result, finding ϵ -solution to $\text{VI}(F, \mathcal{X})$ reduces to solving an explicit solvable Second Order feasibility conic problem of the size independent of ϵ .

2. Thus, all we need is to find an explicit \mathcal{K} -representation of (F, \mathcal{X}) . This can be done as follows:

(a) Consider the convex-concave function

$$\psi(u, v) = -\frac{u}{u+v+1} : \mathbf{R}_+^2 \rightarrow \mathbf{R}$$

along with the associated monotone vector field

$$\Phi(u, v) = \left[\frac{\partial \psi(u, v)}{\partial u}; -\frac{\partial \psi(u, v)}{\partial v} \right] = \left[-\frac{v+1}{(u+v+1)^2}; -\frac{u}{(u+v+1)^2} \right].$$

Let also A_i be $2 \times n$ matrix with i -th column $[1; 0]$ and all remaining columns equal to $[0; 1]$, and let $G(x) = \nabla \left(\frac{1}{\sum_j x_j + 1} \right)$. As is immediately seen, we have

$$\forall x \in \mathbf{R}_+^n : \Psi(x) := 2c + \sum_{i=1}^n A_i^T \Phi(A_i x) + G(x) = 2F(x),$$

so that a \mathcal{K} -representation of F is readily given by \mathcal{K} -representations of the constant monotone vector field $\equiv c$, of Φ , and of G via our calculus (rules on affine substitution of argument and on summation).

A \mathcal{K} -representation of the constant vector field $\equiv c$ is trivial; it is given by the system of linear equalities $t = c^T x, g = c$ on variables $(t, g, x) \in \mathbf{R} \times \mathbf{R}^n \times \mathbf{R}^n$. A \mathcal{K} -representation of the gradient vector field $G(x)$ is given by calculus rule 2 in Section 4.3.1; we are in the case where (17) reads

$$\begin{aligned} \{(x, s) : s \geq f(x), x \in X\} &= \left\{ (x, s) : -x \leq 0, x \leq X := [X_1; \dots; X_n], \right. \\ &\quad \left. \underbrace{-[0; \sum_i x_i - s; \sum_i x_i + s] \leq_{\mathbf{L}^3} [2; 1; 1]} \right\}, \\ &\Leftrightarrow x + 1 + s \geq 0, \\ &\quad (\sum_i x_i + 1 + s)^2 \geq (\sum_i x_i + 1 - s)^2 + 4 \\ &\Leftrightarrow s(\sum_i x_i + 1) \geq 1 \end{aligned}$$

so that (18), as seen from immediate computation, reduces to the system of constraints

$$\begin{aligned} t \geq r + s, 0 \leq x \leq X, r \geq \sum_i \max[X_i(g_i + \theta), 0] + \theta - 2\nu \\ \underbrace{[2; \sum_i x_i + 1 - s; \sum_i x_i + 1 + s] \geq_{\mathbf{L}^3} 0}_{\Leftrightarrow s(\sum_j x_j + 1) \geq 1}, \underbrace{[2\nu; \theta - 1; \theta + 1] \geq_{\mathbf{L}^3} 0}_{\Leftrightarrow \nu^2 \leq \theta} \end{aligned}$$

in variables t, g, x, θ, ν .

(b) It remains to find a \mathcal{K} -representation of the vector field Φ on a rectangle $\Xi = \mathcal{U} \times \mathcal{V}$, $\mathcal{U} = [0, U]$, $\mathcal{V} = [0, V]$ with given $U > 0$, $V > 0$ (they should be large enough in order for $A_i x, x \in \mathcal{X}$, to take values in the rectangle).

Let us use the construction described in item 3 of Section 4.3.1. By (i) of item 3, the set

$$\overline{\mathcal{Z}} = \left\{ (t \in \mathbf{R}, g = [h; e] \in \mathbf{R}^2, x = [u; v] \in \Xi) : \exists r, s : \begin{array}{l} r \geq \max_{\zeta \in \mathcal{V}} [\zeta e + \psi(u, \zeta)] \\ s \geq \max_{\omega \in \mathcal{U}} [\omega^T h - \psi(\omega, v)] \\ t \geq r + s \end{array} \right\}$$

satisfies the relations

$$\begin{aligned} \overline{\mathcal{Z}} &\subset \mathcal{F}[\Phi, \Xi] \\ x \in \mathcal{X}, g = \Phi(x), t \geq x^T \Phi(x) &\Rightarrow [t; g; x] \in \overline{\mathcal{Z}}, \end{aligned}$$

so that all we need to get a \mathcal{K} -representation of Φ is to represent $\overline{\mathcal{Z}}$ as a projection onto the plane of t, g, x -variables of the solution set of \mathcal{K} -conic constraint in variables t, g, x and additional

variables. To this end note that for all $(u, v) \in \Xi = \mathcal{U} \times \mathcal{V}$ one has

$$\psi(u, v) = \min_{f, t, s, \tau} \left\{ f v + t : \underbrace{\begin{bmatrix} \tau \geq 0, 1 \geq f \geq 0, s \geq 0, t \leq 1, s + \tau \leq 1 \\ [2s; u - f; u + f] \in \mathbf{L}^3 \\ [2(1 - s); t - f; t - f + 2] \in \mathbf{L}^3 \\ [2; u - \tau + 1; u + \tau + 1] \in \mathbf{L}^3 \end{bmatrix}}_{\Leftrightarrow (f, t, s, \tau) \in \Pi_+} \right\} \quad (37)$$

and

$$-\psi(u, v) = \min_{f, t, s} \left\{ f u + t : \underbrace{\begin{bmatrix} 1 \geq f \geq 0, 0 \leq s \leq 1, t \leq 1 \\ [2s; v + 1 - f; v + 1 + f] \in \mathbf{L}^3 \\ [2(1 - s); t - 1; t + 1] \in \mathbf{L}^3 \end{bmatrix}}_{\Leftrightarrow (f, t, s) \in \Pi_-} \right\}, \quad (38)$$

see Section B for justification. Since Π_{\pm} are nonempty convex compact sets, by the Sion-Kakutani Theorem one has from (37):

$$\begin{aligned} \max_{\zeta \in \mathcal{V}} [\zeta e + \psi(u, \zeta)] &= \max_{0 \leq \zeta \leq V} \min_{f, t, s, \tau} \{ [f + e]\zeta + t : (f, t, s, \tau) \in \Pi_+ \} \\ &= \min_{f, t, s, \tau} \left\{ \max_{0 \leq \zeta \leq V} [f + e]\zeta + t : (f, t, s, \tau) \in \Pi_+ \right\} \\ &= \min_{f, t, s, \tau} \{ \max[0, V(f + e)] + t : (f, t, s, \tau) \in \Pi_+ \}, \end{aligned}$$

and from (38):

$$\begin{aligned} \max_{\omega \in \mathcal{U}} [\omega h - \psi(\omega, v)] &= \max_{0 \leq \omega \leq U} \min_{f, t, s} \{ [f + h]\omega + t : (f, t, s) \in \Pi_- \} \\ &= \min_{f, t, s} \left\{ \max_{0 \leq \omega \leq U} [f + h]\omega + t : (f, t, s) \in \Pi_- \right\} \\ &= \min_{f, t, s} \{ \max[0, U(f + h)] + t : (f, t, s) \in \Pi_- \}. \end{aligned}$$

As a result,

$$\overline{\mathcal{Z}} = \left\{ (t \in \mathbf{R}, g = [h; e] \in \mathbf{R}^2, 0 \leq x := [u; v] \leq [U; V]) : \exists r, s, f_{\pm}, t_{\pm}, s_{\pm}, \tau : \right. \\ \left. \begin{aligned} t &\geq r + s, r \geq \max[0, V(f_+ + e)] + t_+, s \geq \max[0, U(f_- + h)] + t_- \\ (f_+, t_+, s_+, \tau) &\in \Pi_+, (f_-, t_-, s_-, \tau) \in \Pi_- \end{aligned} \right\}.$$

Recalling what are Π_{\pm} , we can straightforwardly represent $\overline{\mathcal{Z}}$ as projection onto the space of t, g, x -variables of a set given by a \mathcal{K} -conic inequality, ending up with an explicit \mathcal{K} -representation of Φ .

Remark 4.2 The just outlined construction can be used in the case of a general convex Nash Equilibrium problem where, given n convex compact sets $\mathcal{X}_i \subset \mathbf{R}^{k_i}$ and n continuously differentiable functions $\phi_i(x_1, \dots, x_n) : \mathcal{X} := \mathcal{X}_1 \times \dots \times \mathcal{X}_n \rightarrow \mathbf{R}$ with ϕ_i convex in x_i , concave in $(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$ and such that $\sum_i \phi_i$ is convex, one is looking for Nash Equilibria—points $x^* = [x_1^*; \dots; x_n^*] \in \mathcal{X}$ such that for every i , the function $\phi_i(x_1^*, \dots, x_{i-1}^*, x_i, x_{i+1}^*, \dots, x_n^*)$ attains its minimum over $x_i \in \mathcal{X}_i$ at $x_i = x_i^*$. As it was already mentioned, these equilibria are exactly the weak=strong solutions to $\text{VI}(F, \mathcal{X})$ where the monotone vector field F is given by $F(x_1, \dots, x_n) = \left[\frac{\partial \phi_1(x)}{\partial x_1}; \dots; \frac{\partial \phi_n(x)}{\partial x_n} \right]$. Observing that the monotone vector fields

$$F^i(x) = \left[-\frac{\partial \phi_i(x)}{\partial x_1}; \dots; -\frac{\partial \phi_i(x)}{\partial x_{i-1}}; \frac{\partial \phi_i(x)}{\partial x_i}; -\frac{\partial \phi_i(x)}{\partial x_{i+1}}; \dots; -\frac{\partial \phi_i(x)}{\partial x_n} \right]$$

associated with convex-concave functions ϕ_i and the monotone vector field $F^0(x) = \nabla(\sum_i \phi_i(x))$ are linked to F by the relation $2F = \sum_{i=0}^n F^i$, we see that in order to get a \mathcal{K} -representation of F , it suffices to have at our disposal \mathcal{K} -representations of F^i , $0 \leq i \leq n$. These latter representations, in good cases, can be built according to recipes presented in items 2 and 3 of Section 4.3.1.

The latter remark puts in proper perspective the “red herring” illustration which by itself is of no actual interest: setting $s = \sum_i x_i + 1$, $x \in \mathcal{X}$ is a Nash Equilibrium if and only if

$$c_i - \frac{1}{s} + \frac{x_i}{s^2} = \frac{\partial \phi_i(x)}{\partial x_i} \begin{cases} \geq 0, & x_i = 0 \\ = 0, & 0 < x_i < X_i \\ \leq 0, & x_i = X_i. \end{cases}$$

As a result, finding the equilibrium reduces to solving on the segment $s \in [1, \sum_i X_i + 1]$ the univariate equation

$$\sum_i x_i(s) + 1 = s, \quad x_i(s) = \begin{cases} 0, & s(1 - sc_i) < 0 \\ s(1 - sc_i), & 0 \leq s(1 - sc_i) \leq X_i \\ X_i, & s(1 - sc_i) > X_i. \end{cases}$$

Acknowledgment

This work was supported by Multidisciplinary Institute in Artificial intelligence MIAI @ Grenoble Alpes (ANR-19-P3IA-0003).

References

- [1] Aghassia, M., Bertsimas, D., Perakis, G. “Solving asymmetric variational inequalities via convex optimization” – *Operations Research Letters* 34 (2006) 481–490
- [2] Andersen, E. D. *MOSEK optimization suite*, MOSEK ApS, 2019. <https://docs.mosek.com/9.2/intro.pdf>
- [3] Boyd, S., Vandenberghe, L. *Convex Optimization*, Cambridge University Press, 2004. https://web.stanford.edu/~boyd/cvxbook/bv_cvxbook.pdf
- [4] Ben-Tal, A., Nemirovski, A. *Lectures on Modern Convex Optimization: Analysis, Algorithms and Engineering Applications*, MPS-SIAM Series on Optimization, SIAM, Philadelphia, 2001.
- [5] Ben-Tal, A., Nemirovski, A. *Lectures on Modern Convex Optimization* [ISyE GaTech], <http://www2.isye.gatech.edu/~nemirovs/LMCOLN2021WithSol.pdf>
- [6] Ghaoui, L., Oks, M., Oustry, F. Worst-case value-at-risk and robust portfolio optimization: A conic programming approach – *Operations research*, **51**(4) 2003, 543–556.
- [7] Facchinei, F., Pang, J.-S. *Finite Dimensional Variational Inequalities and Complementarity Problems, Vol. 1 and 2*, Springer, 2003.
- [8] Goldfarb, D., Iyengar, G. Robust portfolio selection problems – *Mathematics of operations research*, **28**(1) 2003, 1–38.
- [9] Grant, M., and Boyd, S. *CVX: Matlab software for disciplined convex programming, version 2.2*, CVX Research, Inc, 2020. <http://cvxr.com/cvx/doc/>
- [10] Juditsky, A., Nemirovski, A. *Statistical Inference via Convex Optimization*, Princeton University Press, 2020. <http://www2.isye.gatech.edu/~nemirovs/StatOptPUPWeb.pdf>
- [11] Kojima, M., Megiddo, N., Noma, T., Yoshise, A. *A Unified Approach to Interior Point Algorithms for Linear Complementarity Problems*, Lecture Notes in Computer Science 538, Springer, 1991.
- [12] Monteiro, R.D.C., Zanjacomo, P. “General Interior-Point Maps and Existence of Weighted Paths for Nonlinear Semidefinite Complementarity Problems” – *Mathematics of Operations Research* **25** 2000, 381–399.

- [13] Nesterov, Yu., Nemirovskii, A. *Interior-Point Polynomial Algorithms in Convex Programming*, SIAM, 1994.
- [14] Nemirovski, A., Onn, S., Rothblum, U. “Accuracy certificates for computational problems with convex structure” – *Mathematics of Operations Research* **35:1** 2010, 52–78. https://www2.isye.gatech.edu/~nemirovs/MOR_AccuracyCertificates.pdf
- [15] Nesterov, Yu. “Squared Functional Systems and Optimization Problems” – In: Frenk, H., Roos, C., Terlaky, T., Zhang, S. (Eds.) *High Performance Optimization*, Springer 2000, 405–440.
- [16] Petra C. G., Potra F. A. “A homogeneous model for monotone mixed horizontal linear complementarity problems” – *Computational Optimization and Applications* **72** 2019, 241–267.
- [17] Ralph, D., Wright, S. J. “Superlinear convergence of an interior-point method for monotone variational inequalities” – In Ferris, M. C., Pang, J. S. (Eds.) *Complementarity and Variational Problems: State of the Art*, SIAM Publications, Philadelphia, 1997, 345–385.
- [18] Sim, C. K. “Interior point method on semi-definite linear complementarity problems using the Nesterov–Todd (NT) search direction: polynomial complexity and local convergence” – *Computational Optimization and Applications* **74** 2019, 583–621.
- [19] Toh, K., Todd, M., Tütüncü, R. H. On the implementation and usage of SDPT3—a Matlab software package for semidefinite-quadratic-linear programming, version 4.0 – In Anjos, M. F., Lasserre, J. B. (Eds.) *Handbook on semidefinite, conic and polynomial optimization*, Springer, Boston, MA, 2012, 715–754.

A Derivations for Section 4.3

A.1 Verification of “raw materials”

1. *[Affine monotone vector field]* Note first that the symmetric part \bar{A} of A is $\succeq 0$ due to the monotonicity of F on E , so that (16) can be rewritten as a conic constraint on a direct product of properly selected Lorentz cone and nonnegative orthant, and this direct product belongs to \mathcal{K} . Second, when t, g, x solve (16), for every $y \in E$ one has $x^T Ax = x^T \bar{A}x$, whence

$$t - y^T g = t - y^T [Ax + a] = \underbrace{t - x^T [Ax + a]}_{=t - x^T \bar{A}x - x^T a \geq 0} - [y - x]^T [Ax + a] \geq [x - y]^T [Ax + a] \geq [x - y]^T [Ay + a],$$

that is, $[t; g; x] \in \mathcal{F}[F, E]$, as required in item (i) of a representation of (F, E) . Furthermore, when $x \in E$, setting $g = F(x) = Ax + a$ and $t = x^T F(x) = x^T [Ax + a] = x^T [\bar{A}x + a]$, we see that $[t; g; x]$ satisfy (16), as required in item (ii) of a representation of (F, E) .

2. *[Gradient field of continuously differentiable \mathcal{K} -representable convex function]* Let

$$f_*(y) = \sup_{x \in \mathcal{X}} [x^T y - f(x)]$$

be the Fenchel transform of f :

$$\begin{aligned} f_*(y) &= \sup_{x \in \mathcal{X}} [x^T y - f(x)] = \sup_{x \in \mathcal{X}, s \geq f(x)} [x^T y - s] \\ &= \sup_{x, u} [x^T y - s : Ax + sp + Qu \leq_{\mathbf{K}} a] \\ &= \min_{\lambda} \{ a^T \lambda : A^T \lambda = y, p^T \lambda = -1, Q^T \lambda = 0, \lambda \geq_{\mathbf{K}^*} 0 \} \quad [\text{by conic duality}], \end{aligned}$$

that is,

$$r \geq f_*(g) \Leftrightarrow \exists \lambda : r \geq a^T \lambda, A^T \lambda = g, p^T \lambda = -1, Q^T \lambda = 0, \lambda \geq_{\mathbf{K}^*} 0. \quad (39)$$

Next, let

$$\begin{aligned} \mathcal{Z} &= \{(t, g, x) : \exists s, r : s \geq f(x), r \geq f_*(g), s + r \leq t\} & (a) \\ &= \left\{ (t, g, x) : \exists s, r, u, \lambda : \begin{array}{l} t \geq s + r, Ax + sp + Qu \leq_{\mathbf{K}} a \\ r \geq a^T \lambda, A^T \lambda = g, p^T \lambda = -1, Q^T \lambda = 0, \lambda \geq_{\mathbf{K}^*} 0 \end{array} \right\} & (b) \end{aligned} \quad (40)$$

where (40.b) is due to (17) and (39). Thus, \mathcal{Z} is the projection of the solution set of (18) on the space of t, g, x -variables, and we should check that

(i) $\mathcal{Z} \subset \mathcal{F}[f'(\cdot), \mathcal{X}]$, and

(ii) when $x \in \mathcal{X}$, $g = f'(x)$, and $t = x^T f'(x)$, we have $[t; g; x] \in \mathcal{Z}$.

(i): Let $(t, g, x) \in \mathcal{Z}$. We have $t \geq s + r$ for properly selected $s \geq f(x)$ and $r \geq f_*(g)$, whence for $y \in \mathcal{X}$ it holds

$$t - g^T y \geq f(x) + f_*(g) - g^T y \geq f(x) - f(y) \geq (x - y)^T f'(y),$$

that is, $[t; g; x] \in \mathcal{F}[f'(\cdot), \mathcal{X}]$. (i) is proved.

(ii): Given $x \in \mathcal{X}$, let us set $g = f'(x)$, $t = x^T g$, $s = f(x)$, $r = f_*(g)$. We have

$$r = f_*(g) = \sup_{x'} \{g^T x' - f(x') : x' \in \mathcal{X}\} = g^T x - f(x)$$

(the concluding equality is due to $g = f'(x)$); thus, $t = r + s$. Invoking (40.a), we conclude that $(t, g, x) \in \mathcal{Z}$. (ii) is proved.

3. [Monotone vector field associated with \mathcal{K} -representable convex-concave function ψ]

Verifying (i):

1^o. Let us show (22.a). Let $(t, g = [h; e], x = [u; v]) \in \mathcal{Z}$, and let r, s be reals which, taken together with t, g, x , form a feasible solution to the system of constraints in (21). For $y = [w; z] \in \mathcal{X}$ we have

$$\begin{aligned} t - y^T g &\geq r + s - w^T h - z^T e \quad [\text{by (21.c)}] \\ &\geq [z^T e + \psi(u, z)] + [w^T h - \psi(w, v)] - w^T h - z^T e \quad [\text{by (21.a,b)}] \\ &= \psi(u, z) - \psi(w, v) = [\psi(u, z) - \psi(w, z)] + [\psi(w, z) - \psi(w, v)] \\ &\geq [u - w]^T \nabla_u \psi(w, z) - [v - z]^T \nabla_v \psi(w, z) \quad [\text{because } \psi \text{ is convex-concave}] \\ &= [x - y]^T F(y), \end{aligned}$$

implying that $\mathcal{Z} \subset \mathcal{F}[F, \mathcal{X}]$.

2^o. Let us now verify (22.b). Given $x = [u; v] \in \mathcal{X}$, let us set

$$g = F(x) = [h; e], \quad h = \nabla_u \psi(u, v), \quad e = -\nabla_v \psi(u, v); \quad \bar{t} = x^T F(x) = h^T u + e^T v,$$

and let $t > \bar{t}$. The function $\zeta^T e + \psi(u, \zeta)$ of $\zeta \in \mathcal{V}$ is concave, and its gradient w.r.t. ζ taken at the point $\zeta = v$ vanishes, implying that

$$\bar{r} := v^T e + \psi(u, v) \geq \zeta^T e + \psi(u, \zeta) \quad \forall \zeta \in \mathcal{V}.$$

Similarly, the function $\omega^T h - \psi(\omega, v)$ of $\omega \in \mathcal{U}$ is concave, and its gradient w.r.t. ω taken at the point $\omega = u$ vanishes, implying that

$$\bar{s} := u^T h - \psi(u, v) \geq \omega^T h - \psi(\omega, v) \quad \forall \omega \in \mathcal{U}.$$

Observing that $\bar{r} + \bar{s} = \bar{t} < t$, we can find $r > \bar{r}$ and $s > \bar{s}$ in such a way that $r + s \leq t$. Looking at the constraints (21.a-c), we conclude that whenever $x = [u; v] \in \mathcal{X}$ and $t > x^T F(x)$, the triple $(t, g = F(x), x)$ can be augmented by r, s to satisfy all constraints (21.a-c), that is, $(t, g, x) \in \mathcal{Z}$, as claimed in (22.b).

Verifying (ii): Clearly, all we need in order to justify claim in (ii) is to show that the set

$$\mathcal{Z}^+ = \left\{ (g = [h; e], x = [u; v], r, s) : u \in \mathcal{U}, v \in \mathcal{V}, r > \max_{z \in \mathcal{V}} [z^T e + \psi(u, z)], s > \max_{w \in \mathcal{U}} [w^T h - \psi(w, v)] \right\}$$

is the projection of the solution set of the conic constraint

$$\begin{bmatrix} r \geq \theta + \gamma^T b + \tau, \eta \geq 0, \theta \geq 0, \eta\theta \geq 1 \\ s \geq \theta' + c^T \delta + a^T \epsilon, \eta' \geq 0, \theta' \geq 0, \eta'\theta' \geq 1 \\ Au + B\alpha \leq_{\mathbf{K}} a, Cv + D\beta \leq_{\mathbf{L}} b, Pf + \tau p + Q\xi + Ru \leq_{\mathbf{M}} c \\ C^T \gamma = f + e, D^T \gamma = 0, \gamma \geq_{\mathbf{L}^*} 0, \delta \geq_{\mathbf{M}^*} 0, \epsilon \geq_{\mathbf{K}^*} 0 \\ P^T \delta + v = 0, p^T \delta = -1, Q^T \delta = 0, R^T \delta + A^T \epsilon = h, B^T \epsilon = 0 \end{bmatrix} \quad (41)$$

in variables $x = [u; v], g = [h; e], r, s, \alpha, \beta, \gamma, f, \tau, \xi, \delta, \epsilon, \eta, \theta, \eta'$, and θ' on the plane of $g = [h; e], x = [u; v], r, s$ -variables. Here is the proof.

1^o. Recall that \mathcal{V} is convex and compact. Thus, we have

$$\begin{aligned} & \forall (x = [u; v] \in \mathcal{X}, g = [h; e]) : \\ & \max_{z \in \mathcal{V}} [z^T e + \psi(u, z)] = \max_{z \in \mathcal{V}} \left[z^T e + \inf_{f, \tau, \xi} [f^T z + \tau : Pf + \tau p + Q\xi + Ru \leq_{\mathbf{M}} c] \right] \quad [\text{by (20)}] \\ & = \max_{z \in \mathcal{V}} \inf_{f, \tau, \xi} [(f + e)^T z + \tau : Pf + \tau p + Q\xi + Ru \leq_{\mathbf{M}} c] \\ & = \inf_{f, \tau, \xi} \left\{ \max_{z \in \mathcal{V}} [(f + e)^T z + \tau : Pf + \tau p + Q\xi + Ru \leq_{\mathbf{M}} c] \right\} \quad [\text{by the Sion-Kakutani Theorem}] \\ & = \inf_{f, \tau, \xi} \left[\max_{z, \beta} [(f + e)^T z + \tau : Cz + D\beta \leq_{\mathbf{L}} b] : Pf + \tau p + Q\xi + Ru \leq_{\mathbf{M}} c \right] \quad [\text{by (19.b)}] \\ & = \inf_{f, \tau, \xi, \gamma} \left\{ \gamma^T b + \tau : C^T \gamma = f + e, D^T \gamma = 0, \gamma \geq_{\mathbf{L}^*} 0, Pf + \tau p + Q\xi + Ru \leq_{\mathbf{M}} c \right\} \\ & \quad [\text{by conic duality; recall that (19.b) is essentially strictly feasible}.] \end{aligned}$$

Together with (19) the latter relation results in

$$\begin{aligned} & \{ (x = [u; v], g = [h; e], r) : x \in \mathcal{X}, r > \max_{z \in \mathcal{V}} [z^T e + \psi(u, z)] \} \\ & = \left\{ (x = [u; v], g = [h; e], r) : \exists f, \tau, \xi, \alpha, \beta, \gamma : \right. \\ & \quad \left. \begin{aligned} & r > \gamma^T b + \tau, Au + B\alpha \leq_{\mathbf{K}} a, Cv + D\beta \leq_{\mathbf{L}} b, \gamma \geq_{\mathbf{L}^*} 0 \\ & C^T \gamma = f + e, D^T \gamma = 0, Pf + \tau p + Q\xi + Ru \leq_{\mathbf{M}} c \end{aligned} \right\} \\ & = \left\{ (x = [u; v], g = [h; e], r) : \exists f, \tau, \xi, \alpha, \beta, \gamma, \eta, \theta : \right. \\ & \quad \left. \begin{aligned} & r \geq \gamma^T b + \tau + \theta, \eta \geq 0, \theta \geq 0, \eta\theta \geq 1 \\ & Au + B\alpha \leq_{\mathbf{K}} a, Cv + D\beta \leq_{\mathbf{L}} b, \gamma \geq_{\mathbf{L}^*}, \\ & C^T \gamma = f + e, D^T \gamma = 0, Pf + \tau p + Q\xi + Ru \leq_{\mathbf{M}} c \end{aligned} \right\}. \quad (42) \end{aligned}$$

2^o. When $v \in \mathcal{V}$ and $h \in \mathbf{R}^{n_u}$ we have

$$\begin{aligned} & \max_{w \in \mathcal{U}} [w^T h - \psi(w, v)] = \max_{w \in \mathcal{U}} \left[w^T h + \sup_{f, \tau, \xi} [-f^T v - \tau : Pf + \tau p + Q\xi + Rw \leq_{\mathbf{M}} c] \right] \quad [\text{by (20)}] \\ & = \sup_{w, f, \tau, \xi, \alpha} [w^T h - f^T v - \tau : Pf + \tau p + Q\xi + Rw \leq_{\mathbf{M}} c, Aw + B\alpha \leq_{\mathbf{K}} a] \quad [\text{by (19.a)}] \\ & = \min_{\delta, \epsilon} \left\{ c^T \delta + a^T \epsilon : \begin{aligned} & P^T \delta + v = 0, p^T \delta = -1, Q^T \delta = 0, R^T \delta + A^T \epsilon = h \\ & B^T \epsilon = 0, \delta \geq_{\mathbf{M}^*} 0, \epsilon \geq_{\mathbf{K}^*} 0 \end{aligned} \right\} \\ & \quad \left[\begin{aligned} & \text{by conic duality; recall that the } \mathcal{K}\text{-representations (19.a)} \\ & \text{of } \mathcal{U} \text{ and (20) of } \psi \text{ are essentially strictly feasible.} \end{aligned} \right] \end{aligned}$$

Taken together with (19), the latter relation results in

$$\begin{aligned}
& \{ (x = [u; v], g = [h; e], s) : x \in \mathcal{X}, s > \max_{w \in \mathcal{U}} [h^T w - \psi(w, u)] \} \\
&= \left\{ (x = [u; v], g = [h; e], s) : \exists \alpha, \beta, \delta, \epsilon : \right. \\
&\quad \left. \begin{aligned} & s > c^T \delta + a^T \epsilon, Au + B\alpha \leq_{\mathbf{K}} a, Cv + D\beta \leq_{\mathbf{L}} b, \delta \geq_{\mathbf{M}^*} 0, \epsilon \geq_{\mathbf{K}^*} 0 \\ & P^T \delta + v = 0, p^T \delta = -1, Q^T \delta = 0, R^T \delta + A^T \epsilon = h, B^T \epsilon = 0 \end{aligned} \right\} \\
&= \left\{ (x = [u; v], g = [h; e], s) : \exists \alpha, \beta, \delta, \epsilon, \eta', \theta' : \right. \\
&\quad \left. \begin{aligned} & s \geq \theta' + c^T \delta + a^T \epsilon, \eta' \geq 0, \theta' \geq 0, \eta' \theta' \geq 1 \\ & Au + B\alpha \leq_{\mathbf{K}} a, Cv + D\beta \leq_{\mathbf{L}} b, \delta \geq_{\mathbf{M}^*} 0, \epsilon \geq_{\mathbf{K}^*} 0 \\ & P^T \delta + v = 0, p^T \delta = -1, Q^T \delta = 0, R^T \delta + A^T \epsilon = h, B^T \epsilon = 0 \end{aligned} \right\}. \quad (43)
\end{aligned}$$

Finally, (42) and (43) together imply that

$$\begin{aligned}
\mathcal{Z}^+ &:= \left\{ (g = [h; e], x = [u; v], r, s) : u \in \mathcal{U}, v \in \mathcal{V}, r > \max_{z \in \mathcal{V}} [z^T e + \psi(u, z)], s > \max_{w \in \mathcal{U}} [w^T h - \psi(v, u)] \right\} \\
&= \left\{ (g = [h; e], x = [u; v], r, s) : \exists f, \tau, \xi, \alpha, \beta, \gamma, \eta, \theta, \delta, \epsilon, \eta', \theta' : \right. \\
&\quad \left. \begin{aligned} & r \geq \theta + \gamma^T b + \tau, \eta \geq 0, \theta \geq 0, \eta \theta \geq 1 \\ & s \geq \theta' + c^T \delta + a^T \epsilon, \eta' \geq 0, \theta' \geq 0, \eta' \theta' \geq 1 \\ & Au + B\alpha \leq_{\mathbf{K}} a, Cv + D\beta \leq_{\mathbf{L}} b, Pf + \tau p + Q\xi + Ru \leq_{\mathbf{M}} c, \gamma \geq_{\mathbf{L}^*} 0, \delta \geq_{\mathbf{M}^*} 0, \epsilon \geq_{\mathbf{K}^*} 0 \\ & C^T \gamma = f + e, D^T \gamma = 0, P^T \delta + v = 0, p^T \delta = -1, Q^T \delta = 0, R^T \delta + A^T \epsilon = h, B^T \epsilon = 0 \end{aligned} \right\},
\end{aligned}$$

as claimed in (41).

4. [Univariate monotone rational vector field] For evident reasons, it suffices to consider the case of $\mathcal{X} = [0, 1]$; recall that $\beta(t) > 0$ on \mathcal{X} . Let degrees of α and β be μ and ν , respectively, and let

$$\kappa = \max[\mu, \nu] + 1.$$

1⁰. Consider the curves

$$\delta(t) = [tF(t); F(t); t] = \frac{1}{\beta(t)} [t\alpha(t); \alpha(t); t\beta(t)] : [0, 1] \rightarrow \mathbf{R}^3, \gamma(t) = \frac{1}{\beta(t)} [1; t; t^2; \dots; t^\kappa] : [0, 1] \rightarrow \mathbf{R}^{\kappa+1}$$

For properly selected matrix A we have

$$\delta(t) = A\gamma(t), \quad 0 \leq t \leq 1,$$

whence

$$\mathcal{Y} := \text{Conv}\{\delta(t), 0 \leq t \leq 1\} = A\mathcal{Z}, \quad \mathcal{Z} = \text{Conv}\{\gamma(t), 0 \leq t \leq 1\}.$$

We intend to build a semidefinite representation (SDR) of \mathcal{Y} (i.e., \mathcal{K} -representation with \mathcal{K} comprised of finite direct products of semidefinite cones). Semidefinite representability (and \mathcal{K} -representability in general) of a set is preserved when taking linear images: an SDR

$$\mathcal{Z} = \{z : \exists u : \mathcal{A}(z, u) \succeq 0\} \quad [\mathcal{A}(z, u) \text{ is affine in } [z; u] \text{ symmetric matrix}]$$

of \mathcal{Z} implies the representation

$$\mathcal{Y} := A\mathcal{Z} = \{y : \exists [z; u] : y = Az, \mathcal{A}(z, u) \succeq 0\},$$

and the system of the right hand side constraints can be written down as a Linear Matrix Inequality in variable y and additional variables z, u . Thus, all we need is to build an SDR of \mathcal{Z} .

2⁰. We shall get SDR of \mathcal{Z} from SDR of the “support cone”

$$\mathcal{P} = \{[p; q] \in \mathbf{R}^{\kappa+1} \times \mathbf{R} : \min_{z \in \mathcal{Z}} p^T z - q \geq 0\} = \{[p; q] : \min_{0 \leq t \leq 1} p^T \gamma(t) \geq q\}$$

of \mathcal{Z} .

Given $p = [p_0; p_1; \dots; p_{\kappa+1}] \in \mathbf{R}^{\kappa+1}$, let, with a slight abuse of notation, $p(t) = \sum_{i=0}^{\kappa} p_i t^i$ be the polynomial with coefficients p_i , $0 \leq i \leq \kappa$. We have

$$\begin{aligned} [p; q] \in \mathcal{P} &\Leftrightarrow \frac{p(t)}{\beta(t)} \geq q \forall t \in [0, 1] \\ &\Leftrightarrow \frac{(1+\tau^2)^\kappa p(\tau^2/(1+\tau^2))}{(1+\tau^2)^\kappa \beta(\tau^2/(1+\tau^2))} \geq q \forall \tau \in \mathbf{R} \\ &\Leftrightarrow \underbrace{(1+\tau^2)^\kappa p(\tau^2/(1+\tau^2)) - q(1+\tau^2)^\kappa \beta(\tau^2/(1+\tau^2))}_{=:\pi_{p,q}(\tau)} \geq 0 \forall \tau \in \mathbf{R}. \end{aligned}$$

Note that $\pi_{p,q}(\tau)$ is a polynomial of τ of degree $\leq 2\kappa$, and the vector $\pi_{p,q}$ of coefficients of polynomial is linear in $[p; q]$: $\pi_{p,q} = P[p; q]$. We see that \mathcal{P} is the inverse image of the cone $\mathcal{P}_{2\kappa}$ of coefficients of polynomials of degree $\leq 2\kappa$ which are nonnegative on the entire axis:

$$\mathcal{P} = \{[p; q] : P[p; q] \in \mathcal{P}_{2\kappa}\}.$$

As was observed by Yu. Nesterov [15], the cone $\mathcal{P}_{2\kappa}$ is the linear image of the semidefinite cone $\mathbf{S}_+^{\kappa+1}$:

$$\mathcal{P}_{2\kappa} = \{\pi \in \mathbf{R}^{2\kappa+1} : \exists x = [x_{ij}]_{0 \leq i, j \leq \kappa} \in \mathbf{S}_+^{\kappa+1} : [\mathcal{Q}x]_\ell := \sum_{\substack{0 \leq i, j \leq \kappa, \\ i+j=\ell}} x_{ij} = \pi_\ell, 0 \leq \ell \leq 2\kappa\},$$

and we arrive at a semidefinite representation of \mathcal{P} :

$$\mathcal{P} = \{[p; q] : \exists x \in \mathbf{S}_+^{\kappa+1} : x \succeq 0 \ \& \ \mathcal{Q}x = P[p; q]\}. \quad (44)$$

We claim that this representation is essentially strictly feasible. Indeed, let $\bar{p} = [1; 1; \dots; 1] \in \mathbf{R}^{\kappa+1}$ and $\bar{q} = 0$. Then

$$\pi_{\bar{p}, \bar{q}}(\tau) = (1+\tau^2)^\kappa [1 + \tau^2/(1+\tau^2) + [\tau^2/(1+\tau^2)]^2 + \dots + [\tau^2/(1+\tau^2)]^\kappa] = \sum_{i=0}^{\kappa} c_i \tau^{2i}, \quad c_i > 0 \forall i,$$

implying that with $\bar{x} = \text{Diag}\{c_0, c_1, \dots, c_\kappa\} \in \mathbf{S}_+^{\kappa+1}$ one has

$$P[\bar{p}; \bar{q}] = \mathcal{Q}\bar{x} \ \& \ \bar{x} \succ 0.$$

That is, \bar{p}, \bar{q} , and \bar{x} satisfy all constraints in (44) and strictly satisfy the non-polyhedral constraint $x \succeq 0$, as required by essentially strict feasibility.

3⁰. Now we are done: by its origin, \mathcal{Z} is a convex compact set and as such is convex and closed, implying by duality that

$$\begin{aligned} \mathcal{Z} &= \{z \in \mathbf{R}^{\kappa+1} : p^T z - q \geq 0 \ \forall [p; q] \in \mathcal{P}\} \\ &= \left\{ z \in \mathbf{R}^{\kappa+1} : \inf_{p, q, x} \{p^T z - q\} : P[p; q] - \mathcal{Q}x = 0, x \succeq 0 \right\} \quad [\text{by (44)}] \\ &= \left\{ z \in \mathbf{R}^{\kappa+1} : \exists \lambda \in \mathbf{R}^{2\kappa+1} : P^T \lambda = [z; -1], \mathcal{Q}^* \lambda \succeq 0 \right\} \\ &\quad [\text{by semidefinite duality; } \lambda \mapsto \mathcal{Q}^* \lambda : \mathbf{R}^{2\kappa+1} \rightarrow \mathbf{S}_+^{\kappa+1} \\ &\quad \text{is the conjugate of } x \mapsto \mathcal{Q}x : \mathbf{S}_+^{\kappa+1} \rightarrow \mathbf{R}^{2\kappa+1}], \end{aligned}$$

and we arrive at the desired SDR of \mathcal{Z} .

A.2 Verification of calculus rules

1. *[Restriction on a \mathcal{K} -representable set]* Let (24) represent (F, \mathcal{X}) . Suppose that $[t; g; x]$ can be augmented by u, v to solve (26). Then (t, g, x, u) solve (24), implying that $[t; g; x] \in \mathcal{F}[F, \mathcal{X}]$ by definition of \mathcal{K} -representability of (F, \mathcal{X}) , and (x, v) solve (26), implying that $x \in \mathcal{Y}$. Taking together, these inclusions clearly imply that $[t; g; x] \in \mathcal{F}[\bar{F}, \bar{\mathcal{Z}}]$, as required in item (i) of the definition of \mathcal{K} -representation of $(\bar{F}, \bar{\mathcal{Z}})$. Next, when $x \in \mathcal{Z}$, the triple $[t := \langle F(x), x \rangle; g := F(x); x]$ can be augmented by u to solve (24) (by item (ii) of the definition of \mathcal{K} -representation of (F, \mathcal{X})), and because $x \in \mathcal{Y}$, x can be augmented by v to solve (25). Thus, $t := \langle F(x), x \rangle$, $g := F(x)$, x can be augmented by (u, v) to solve (26), as required in item (ii) of \mathcal{K} -representability of $(\bar{F}, \bar{\mathcal{Z}})$. Thus, (26) indeed represents $(\bar{F}, \bar{\mathcal{Z}})$. A completely similar reasoning shows that if (24) almost represents (F, \mathcal{X}) , then (26) almost represents $(\bar{F}, \bar{\mathcal{Z}})$.
2. *[Direct summation]* When $x = [x_1; \dots; x_K] \in \mathcal{X}$ and $g = F(x) = [g_1; \dots; g_K]$, $g_k = F_k(x_k)$, $t_k = \langle x_k, g_k \rangle$ and $t = \sum_k t_k = \langle g, x \rangle$, there exist u_k such that (28.a) take place (by item (ii) of the definition of representation as applied to the representations in (27)), and (28.b) takes place as well, as required in item (ii) of the definition of a representation of (F, \mathcal{X}) . On the other hand, when $x = [x_1; \dots; x_K]$, $g = [g_1; \dots; g_K]$, and t can be augmented by u_k and t_k to solve (28.a) and (28.b), we have $[t_k; g_k; x_k] \in \mathcal{F}[F_k, \mathcal{X}_k]$, $k \leq K$, whence for every $y = [y_1; \dots; y_K] \in \mathcal{X}$ it holds

$$t_k - \langle g_k, y_k \rangle \geq \langle F_k(y_k), x_k - y_k \rangle, \quad k \leq K.$$

When summing up the above inequalities over k , we get

$$t - \langle g, y \rangle \geq \langle F(y), x - y \rangle \quad \forall y \in \mathcal{X},$$

that is, $[t; g; x] \in \mathcal{F}[F, \mathcal{X}]$, as required in item (i) of the definition of a representation of (F, \mathcal{X}) .

The above reasoning, with evident modifications, shows the claim in the case of almost representations.

3. *[Taking conic combinations]* Let t, g, x can be augmented by $u_k, t_k, g_k, k \leq K$ to solve (30). Then $[t_k; g_k; x] \in \mathcal{F}[F_k, \mathcal{X}]$, implying that

$$t_k - \langle g_k, y \rangle \geq \langle F_k(y), x - y \rangle \quad \forall y \in X;$$

multiplying both sides by α_k and summing up over k , we get

$$t - \langle g, y \rangle \geq \langle F(y), x - y \rangle \quad \forall y \in X,$$

that is, $[t; g; x] \in \mathcal{F}[F, \mathcal{X}]$. On the other hand, given $x \in \mathcal{X}$, let us set $g_k = F_k(x)$ and $t_k = \langle F_k(x), x \rangle$. Since the k -th conic constraint in (30) represents (F_k, \mathcal{X}) , there exist $u_k, k \leq K$, such that all relations (30.a) take place. Setting $t = \sum_k \alpha_k t_k$, $g = \sum_k \alpha_k g_k$, we, on one hand, satisfy (30.b-c), and, on the other hand, obtain

$$t = \sum_k \alpha_k \langle F_k(x), x \rangle = \langle F(x), x \rangle, \quad g = \sum_k \alpha_k g_k = F(x),$$

the bottom line being that $[\langle F(x), x \rangle; F(x); x]$ can be augmented by u_k, t_k, g_k to solve (30). Thus, (30) indeed is a \mathcal{K} -representation of (F, \mathcal{X}) .

The above reasoning, with evident modifications, works in the case of almost representations.

4. *[Affine substitution of variables]* Assume that $\tau, \gamma, \xi, g, t, u$ solve (32). Then $t, g, x := A\xi + a$, u satisfy

$$Xx + Gg + tT + Uu \leq_{\mathbf{K}} a,$$

implying that $x \in \mathcal{X}$ and

$$t - \langle g, y \rangle \geq \langle F(y), x - y \rangle \quad \forall y \in \mathcal{X}.$$

Recall that $x = A\xi + a$, $x \in \mathcal{X}$, implies that $\xi \in \Xi$. Now, when $\eta \in \Xi$, setting $y = A\eta + a$, we have $y \in \mathcal{X}$ and

$$\begin{aligned} \tau - \langle \gamma, \eta \rangle &= t - \langle g, a \rangle - \langle A^T g, \eta \rangle = t - \langle g, y \rangle \geq \langle F(y), x - y \rangle \\ &= \langle F(A\eta + a), A(\xi - \eta) \rangle = \langle \Phi(\eta), \xi - \eta \rangle, \end{aligned}$$

and since $\eta \in \Xi$ is arbitrary, we get $[\tau; \gamma; \xi] \in \mathcal{F}[\Phi, \Xi]$, as required in item (i) of the definition of a representation of (Φ, Ξ) . On the other hand, when $\xi \in \Xi$, $\gamma = \Phi(\xi)$, and $\tau = \langle \Phi(\xi), \xi \rangle$, setting $x = A\xi + a$, we get $x \in \mathcal{X}$. Next, when setting $g = F(x)$, $t = \langle F(x), x \rangle$, we obtain $\gamma = A^T g$ and

$$\tau = \langle F(x), x - a \rangle = t - \langle g, a \rangle.$$

Besides this, by the origin of t, g, x and item (ii) of the definition of \mathcal{K} -representation, as applied to (31), there exists u such that t, g, x, u satisfy (31). The bottom line is that ξ, γ, τ can be augmented by t, g, u to solve (32), that is, (32) meets item (ii) of definition of \mathcal{K} -representation of (Φ, Ξ) . The above reasoning, with evident modifications, works in the case of almost representations.

B Verifying (37) and (38)

A. Let us prove that for all $u \in \mathcal{U} = [0, U]$, $v \in \mathcal{V} = [0, V]$ one has

$$\begin{aligned} \psi(u, v) &= -\frac{u}{u+v+1} \\ &= \min_{f, t, s} \left\{ fv + t : 1 \geq f \geq 0, 0 \leq s \leq \sqrt{uf}, s + \frac{1}{u+1} \leq 1, t - f \geq (1-s)^2 - 1, t \leq 1 \right\} \\ &= \min_{f, t, s; \tau} \left\{ fv + t : \begin{array}{l} 1 \geq f \geq 0, s \geq 0, t \leq 1, \tau \geq 0, s + \tau \leq 1 \\ \underbrace{[2s; u - f; u + f] \in \mathbf{L}^3}_{\Leftrightarrow s^2 \leq uf \text{ when } u + f \geq 0} \\ \underbrace{[2(1-s); t - f; t - f + 2] \in \mathbf{L}^3}_{\Leftrightarrow (1-s)^2 \leq t - f + 1 \text{ when } t - f + 2 \geq 0} \\ \underbrace{[2; u - \tau + 1; u + \tau + 1] \in \mathbf{L}^3}_{\Leftrightarrow (u+1)\tau \geq 1 \text{ when } u + \tau + 1 \geq 0} \end{array} \right\}. \end{aligned}$$

Indeed, for $u \in \mathcal{U}$ and $v \in \mathcal{V}$ we have

$$\begin{aligned} &\min_{f, t, s} \left\{ fv + t : 1 \geq f \geq 0, 0 \leq s \leq \sqrt{uf}, s + \frac{1}{u+1} \leq 1, t - f \geq (1-s)^2 - 1, t \leq 1 \right\} \\ &= \min_f \left\{ fv + f - 2\bar{s}(f) + \bar{s}(f)^2 : 1 \geq f \geq 0, \bar{s}(f) = \min \left[\frac{u}{u+1}, \sqrt{uf} \right] \right\} \\ &= \min \left[\min_{0 \leq f \leq u/(u+1)^2} \left\{ fv + f - 2\sqrt{uf} + uf \right\}, \min_{1 \geq f \geq u/(u+1)^2} \left\{ fv + f - \frac{2u}{u+1} + \frac{u^2}{(u+1)^2} \right\} \right] \\ &= \min \left[-\frac{u}{u+v+1}, \frac{u(v+1) - 2u - u^2}{(u+1)^2} \right] = u \min \left[-\frac{1}{v+(u+1)}, \frac{v-(u+1)}{(u+1)^2} \right] = \frac{u}{v+u+1} \min \left[-1, \frac{v^2 - (u+1)^2}{(u+1)^2} \right] \\ &= \psi(u, v), \end{aligned}$$

as claimed.

B. Now let us show that for all $u \in \mathcal{U} = [0, U]$, $v \in \mathcal{V} = [0, V]$

$$\begin{aligned} -\psi(u, v) &:= \frac{u}{u+v+1} \\ &= \min_{f, t, s} \left\{ fu + t : 1 \geq f \geq 0, 0 \leq s \leq \sqrt{(v+1)f}, s \leq 1, 1 \geq t \geq (1-s)^2 \right\} \\ &= \min_{f, t, s} \left\{ fu + t : \begin{array}{l} 1 \geq f \geq 0, 0 \leq s \leq 1, t \leq 1 \\ \underbrace{[2s; v+1 - f; v+1 + f] \in \mathbf{L}^3}_{\Leftrightarrow s^2 \leq (v+1)f \text{ when } v+1+f \geq 0} \\ \underbrace{[2(1-s); t-1; t+1] \in \mathbf{L}^3}_{\Leftrightarrow (1-s)^2 \leq t} \end{array} \right\} \end{aligned}$$

Indeed, for $u \in \mathcal{U}$ and $v \in \mathcal{Q}$ we have

$$\begin{aligned}
& \min_{f,t,s} \left\{ fu + t : 1 \geq f \geq 0, 0 \leq s \leq \sqrt{(v+1)f}, s \leq 1, 1 \geq t \geq (1-s)^2 \right\} \\
&= \min_{f,s} \left\{ fu + (1-s)^2 : 1 \geq f \geq 0, 0 \leq s \leq \sqrt{(v+1)f}, s \leq 1 \right\} \\
&= \min_f \left\{ fu + (1-\bar{s}(f))^2 : 1 \geq f \geq 0, \bar{s}(f) = \min[\sqrt{(v+1)f}, 1] \right\} \\
&= \min \left[\min_{0 \leq f \leq 1/(v+1)} \{ fu + 1 - 2\sqrt{(v+1)f} + (v+1)f \}, \min_{1 \geq f \geq 1/(v+1)} \{ fu \} \right] \\
&= \min \left[\frac{u}{u+v+1}, \frac{u}{v+1} \right] = \frac{u}{u+v+1} = -\psi(u, v),
\end{aligned}$$

as claimed. □