

# SOME PROPERTIES OF ALMOST SUMMING OPERATORS

RENATO MACEDO AND JOEDSON SANTOS

ABSTRACT. In this paper we extend the scope of three important results of the linear theory of absolutely summing operators. The first one was proved by Bu and Kranz in [4] and it asserts that a continuous linear operator between Banach spaces takes almost unconditionally summable sequences into Cohen strongly  $q$ -summable sequences for any  $q \geq 2$ , whenever its adjoint is  $p$ -summing for some  $p \geq 1$ . The second of them states that  $p$ -summing operators with hilbertian domain are Cohen strongly  $q$ -summing operators ( $1 < p, q < \infty$ ), this result is due to Bu [3]. The third one is due to Kwapien [8] and it characterizes spaces isomorphic to a Hilbert space using 2-summing operators. We will show that these results are maintained replacing the hypothesis of the operator to be  $p$ -summing by almost summing. We will also give an example of an almost summing operator that fails to be  $p$ -summing for every  $1 \leq p < \infty$ .

## 1. INTRODUCTION

The space of all continuous linear operators from a Banach space  $X$  to a Banach space  $Y$  will be denoted by  $\mathcal{L}(X, Y)$ . The closed unit ball of  $X$  is denoted by  $B_X$  and its topological dual by  $X^*$ . The symbol  $u^*$  represents the adjoint of a continuous linear operator  $u$ . Below,  $r_i$  are the Rademacher functions and if  $1 \leq p \leq \infty$  we denote by  $p^*$  the conjugate of  $p$ , i.e.,  $\frac{1}{p} + \frac{1}{p^*} = 1$ . For a Banach space  $X$ , the spaces

$$\begin{aligned} \ell_p(X) &:= \{(x_i)_{i=1}^\infty \subset X : \|(x_i)_{i=1}^\infty\|_p := (\sum_{i=1}^\infty \|x_i\|^p)^{1/p} < \infty\}, \\ \ell_p^w(X) &:= \{(x_i)_{i=1}^\infty \subset X : \|(x_i)_{i=1}^\infty\|_{w,p} := \sup_{x^* \in B_{X^*}} \|(x^*(x_i))_{i=1}^\infty\|_p < \infty\}, \\ \ell_p\langle X \rangle &:= \{(x_i)_{i=1}^\infty \subset X : \|(x_i)_{i=1}^\infty\|_{C,p} := \sup_{(x_i^*)_{i=1}^\infty \in B_{\ell_p^w(X^*)}} \|(x_i^*(x_i))_{i=1}^\infty\|_1 < \infty\} \text{ and} \end{aligned}$$

$$Rad(X) := \{(x_i)_{i=1}^\infty \subset X : \|(x_i)_{i=1}^\infty\|_{Rad(X)} := \left( \int_0^1 \|\sum_{i=1}^\infty r_i(t)x_i\|^2 dt \right)^{1/2} < \infty\}$$

are, respectively, the spaces of *absolutely*, *weakly*, *Cohen strongly (or strongly)  $p$ -summable* and *almost unconditionally summable  $X$ -valued sequences* (see [6, 7]).

If  $1 \leq p < \infty$ , we say that a linear operator  $u : X \rightarrow Y$  is *absolutely  $p$ -summing* (or  *$p$ -summing*) if  $(u(x_i))_{i=1}^\infty \in \ell_p(Y)$  whenever  $(x_i)_{i=1}^\infty \in \ell_p^w(X)$ . The class of absolutely  $p$ -summing linear operators from  $X$  to  $Y$  will be represented by  $\Pi_p(X, Y)$  (see [7]).

In [6] Cohen introduced a class of operators which characterizes the  $p^*$ -summing adjoint operators. If  $1 < p \leq \infty$ , we say that a linear operator  $u$  from  $X$  to  $Y$  is *Cohen strongly  $p$ -summing* (or *strongly  $p$ -summing*) if  $(u(x_i))_{i=1}^\infty \in \ell_p\langle Y \rangle$  whenever  $(x_i)_{i=1}^\infty \in \ell_p(X)$ . The class of Cohen strongly  $p$ -summing linear operators from  $X$  to  $Y$  will be denoted by  $\mathcal{D}_p(X, Y)$ . An equivalent formulation

---

*Key words and phrases.* absolutely  $p$ -summing operators, Cohen strongly  $p$ -summing operators, almost  $p$ -summing operators.

2020 Mathematics Subject Classification: Primary 46B45, 47B01; Secondary 47B10, 47L20.

Renato Macedo is partially supported by Capes and Joedson Santos is supported by CNPq, Grant 2019/0014 Paraíba State Research Foundation (FAPESQ) and Public Call n. 03 Produtividade em Pesquisa PROPESQ/PRPG/UFPB PIA13194-2020.

asserts that  $u \in \mathcal{L}(X, Y)$  is Cohen strongly  $p$ -summing if there is a constant  $C \geq 0$  such that

$$\sum_{i=1}^m |y_i^*(u(x_i))| \leq C \cdot \|(x_i)_{i=1}^m\|_p \cdot \|(y_i^*)_{i=1}^m\|_{p^*, w}$$

for all  $x_1, \dots, x_m \in X$ ,  $y_1^*, \dots, y_m^* \in Y^*$  and all positive integers  $m$ .

According to [2], a linear operator  $u \in \mathcal{L}(X; Y)$  is called to be *almost  $p$ -summing*,  $1 \leq p < \infty$ , if there is a constant  $C \geq 0$  such that

$$(1.1) \quad \left( \int_0^1 \left\| \sum_{i=1}^m r_i(t)u(x_i) \right\|^2 dt \right)^{\frac{1}{2}} \leq C \cdot \|(x_i)_{i=1}^m\|_{w, p}$$

for every  $m \in \mathbb{N}$  and  $x_1, \dots, x_m \in X$ . The smallest  $C$  which satisfies the inequality is named the *almost summing norm* of  $u$  denoted by  $\pi_{al.s.p}(u)$ . The class of all almost summing operators from  $X$  to  $Y$  is denoted by  $\Pi_{al.s.p}(X, Y)$ . When  $p = 2$ , these operators are simply called *almost summing* and we write  $\Pi_{al.s}$  instead of  $\Pi_{al.s,2}$  (see [7, Chapter 12]). By [7, Proposition 12.5],

$$(1.2) \quad \bigcup_{1 \leq p < \infty} \Pi_p(X, Y) \subseteq \Pi_{al.s}(X, Y).$$

Now we will give an example of an almost summing operator that fails to be  $p$ -summing for every  $1 \leq p < \infty$ . We provide this example because we have found no information in this direction in the literature to quote.

**Example 1.1.** Consider the following Banach space

$$Y = \left( \sum_{k=3}^{\infty} \ell_k \right)_1 := \left\{ (x_k)_{k=3}^{\infty}; x_k \in \ell_k \text{ and } \|(x_k)_{k=3}^{\infty}\|_1 = \sum_{k=3}^{\infty} \|x_k\|_k \right\}$$

and let

$$T_k : \ell_1 \rightarrow \ell_k$$

be operators such that  $T_k$  fails to be absolutely  $k$ -summing for all  $k = 3, 4, 5, \dots$  (see [1, Proposition 5.2(v)]). On the other hand, for all  $k = 3, 4, 5, \dots$  (see [1, Proposition 5.2(iii)]) we have

$$\Pi_{k+1}(\ell_1, \ell_k) = \mathcal{L}(\ell_1, \ell_k)$$

and thus

$$T_k \in \Pi_{k+1}(\ell_1, \ell_k) \subset \Pi_{al.s}(\ell_1, \ell_k).$$

For all  $k = 3, 4, 5, \dots$ , let

$$i_k : \ell_k \rightarrow Y$$

be the natural inclusions

$$i_k \left( (a_j)_{j=1}^{\infty} \right) = \left( 0, \dots, 0, (a_j)_{j=1}^{\infty}, 0, 0, \dots \right),$$

with  $(a_j)_{j=1}^{\infty}$  at the  $(k-2)$ -th position. Note that  $\|i_k\| = 1$ .

Let

$$T : \ell_1 \rightarrow Y$$

be

$$T(z) = \sum_{k=3}^{\infty} \frac{(i_k \circ T_k)(z)}{C_k},$$

where

$$C_k = 2^k \cdot \pi_{al.s}(T_k).$$

Note that  $T$  is well-defined, linear and continuous. In fact, as  $\|T_k\| \leq \pi_{al.s}(T_k)$  we obtain

$$\begin{aligned} \sum_{k=3}^{\infty} \left\| \frac{i_k \circ T_k}{C_k} \right\| &= \sum_{k=3}^{\infty} \frac{1}{C_k} \|i_k \circ T_k\| \\ &\leq \sum_{k=3}^{\infty} \frac{1}{C_k} \|i_k\| \|T_k\| \\ &= \sum_{k=3}^{\infty} \frac{\|T_k\|}{2^k \cdot \pi_{al.s}(T_k)} \\ &< \infty. \end{aligned}$$

Thus, we conclude that

$$T := \sum_{k=3}^{\infty} \frac{i_k \circ T_k}{C_k}$$

is well-defined in  $\mathcal{L}(\ell_1, Y)$ .

Since  $T_k$  fails to be absolutely  $k$ -summing, it is simple to prove that

$$T \notin \bigcup_{k \in \{3,4,\dots\}} \Pi_k(\ell_1, Y) = \bigcup_{p \geq 1} \Pi_p(\ell_1, Y).$$

In fact, it suffices to note that

$$\|T_k(x)\|_k \leq C_k \cdot \|T(x)\|_1.$$

It remains to prove that  $T$  is almost summing. Since  $T_k \in \Pi_{al.s}(\ell_1, \ell_k)$ , by the ideal property, we have

$$i_k \circ T_k \in \Pi_{al.s}(\ell_1, Y).$$

Note that (since  $\|i_k\| = 1$ ),

$$\begin{aligned} \sum_{k=3}^{\infty} \pi_{al.s} \left( \frac{i_k \circ T_k}{C_k} \right) &= \sum_{k=3}^{\infty} \frac{1}{C_k} \cdot \pi_{al.s}(i_k \circ T_k) \\ &\leq \sum_{k=3}^{\infty} \frac{1}{C_k} \cdot \|i_k\| \cdot \pi_{al.s}(T_k) \\ &= \sum_{k=3}^{\infty} \frac{1}{2^k \cdot \pi_{al.s}(T_k)} \cdot \pi_{al.s}(T_k) \\ &< \infty. \end{aligned}$$

Since  $\Pi_{al.s}(\ell_1, Y)$  is a Banach space, we conclude that

$$\sum_{k=3}^{\infty} \frac{i_k \circ T_k}{C_k}$$

converges in  $\Pi_{al.s}(\ell_1, Y)$ .

A Banach space  $X$  has *type*  $p$ , with  $1 \leq p \leq 2$ , if there is a constant  $C \geq 0$  such that, for all  $m \in \mathbb{N}$  and  $x_1, \dots, x_m \in X$ ,

$$\left( \int_0^1 \left\| \sum_{i=1}^m r_i(t)x_i \right\|^2 dt \right)^{\frac{1}{2}} \leq C \cdot \|(x_i)_{i=1}^m\|_p.$$

The infimum of the  $C$  is denoted by  $T_p(X)$ .

We should mention here Khinchin's inequality and Kahane's inequality:

**Khinchin inequality.** Let  $0 < p < \infty$ . Then, there are positive constants  $A_p$  and  $B_p$  for which

$$A_p \left( \sum_{i=1}^m |x_i|^2 \right)^{\frac{1}{2}} \leq \left( \int_0^1 \left| \sum_{i=1}^m r_i(t)x_i \right|^p dt \right)^{\frac{1}{p}} \leq B_p \left( \sum_{i=1}^m |x_i|^2 \right)^{\frac{1}{2}}$$

holds, regardless of the choice of finitely many scalars  $x_1, \dots, x_m$ .

**Kahane inequality.** Let  $0 < p, q < \infty$ . Then, there is a constant  $K_{p,q} > 0$  for which

$$\left( \int_0^1 \left\| \sum_{i=1}^m r_i(t)x_i \right\|^q dt \right)^{\frac{1}{q}} \leq K_{p,q} \left( \int_0^1 \left\| \sum_{i=1}^m r_i(t)x_i \right\|^p dt \right)^{\frac{1}{p}}$$

holds, regardless of the choice of a Banach space  $X$  and of finitely many vectors  $x_1, \dots, x_m \in X$ .

Using strong tools as Pietsch domination theorem and Khinchin and Kahane inequalities, the main result proved by Bu and Kranz in [4] was:

**Theorem 1.2.** [4, Theorem 1] *Let  $X$  and  $Y$  be Banach spaces and  $u$  be a continuous linear operator from  $X$  to  $Y$ . If  $u^*$  is  $p$ -summing for some  $p \geq 1$ , then for any  $q \geq 2$ ,  $u$  takes almost unconditionally summable sequences in  $X$  into members of  $\ell_q(Y)$ .*

Let  $X$  be a Hilbert space. Cohen in [5] has shown that

$$(1.3) \quad \Pi_2(X, Y) \subseteq \mathcal{D}_2(X, Y) \text{ for all Banach space } Y.$$

In [3], Bu showed that (1.3) is valid with no restrictions of the parameters  $p, q \in (1, \infty)$  instead of  $p = q = 2$ . Cohen [5] also asked if (1.3) characterizes spaces isomorphic to a Hilbert spaces. Kwapien [8] proved that this question has a positive answer. These important results are as follows:

**Theorem 1.3.** [3, Main Theorem] *Let  $1 < p, q < \infty$ , and let  $X$  be a Hilbert space and  $Y$  be Banach space. Then*

$$\Pi_p(X, Y) \subseteq \mathcal{D}_q(X, Y).$$

**Theorem 1.4.** [8] *The following properties of Banach space  $X$  are equivalent.*

- (i) *The space  $X$  is isomorphic to a Hilbert space.*
- (ii) *For every Banach space  $Y$ ,  $\Pi_2(X, Y) \subseteq \mathcal{D}_2(X, Y)$ .*

The paper is organized as follows: In Section 2 our main result is an improvement of the result of Bu and Kranz [4] through a simpler argument than the original one. In Section 3, we extend the statement of the main result of Bu in [3, Main Theorem]. Finally, in Section 4, we show a Kwapien type theorem using almost summing operators to characterize spaces isomorphic to a Hilbert space.

## 2. EXTENDING BU-KRANZ THEOREM

In the next theorem we will show that the Bu-Kranz Theorem is valid for almost  $p^*$ -summing operators instead of  $p$ -summing operators. The proof of this result is very simple and we will not use the Pietsch Domination Theorem, Khinchin's inequality and Kahane's inequality. Our main result is the following:

**Theorem 2.1.** *Let  $X$  and  $Y$  be Banach spaces and  $u$  be a continuous linear operator from  $X$  to  $Y$ . If  $u^*$  is almost  $p^*$ -summing for some  $p \geq 1$ , then  $u$  takes almost unconditionally summable sequences in  $X$  into members of  $\ell_p\langle Y \rangle$ .*

*Proof.* Let  $u \in \mathcal{L}(X, Y)$ . For any  $x_1, \dots, x_m \in X$ , any  $y_1^*, \dots, y_m^* \in Y^*$  and taking  $\theta_i = \text{sign}[y_i^*(u(x_i))]$ , with  $i = 1, \dots, m$ , it follows that

$$\begin{aligned}
\sum_{i=1}^m |y_i^*(u(x_i))| &= \left| \sum_{i=1}^m \theta_i y_i^*(u(x_i)) \right| \\
&= \left| \sum_{i=1}^m u^*(\theta_i y_i^*)(x_i) \right| \\
&= \left| \int_0^1 \sum_{i=1}^m r_i(t) u^*(\theta_i y_i^*) \left( \sum_{i=1}^m r_i(t) x_i \right) dt \right| \\
&\leq \left( \int_0^1 \left\| \sum_{i=1}^m r_i(t) u^*(\theta_i y_i^*) \right\|^2 dt \right)^{1/2} \cdot \left( \int_0^1 \left\| \sum_{i=1}^m r_i(t) x_i \right\|^2 dt \right)^{1/2} \\
&\stackrel{(1.1)}{\leq} \pi_{al.s.p^*}(u^*) \cdot \|(\theta_i y_i^*)_{i=1}^m\|_{p^*,w} \cdot \|(x_i)_{i=1}^m\|_{Rad(X)} \\
&= \pi_{al.s.p^*}(u^*) \cdot \|(y_i^*)_{i=1}^m\|_{p^*,w} \cdot \|(x_i)_{i=1}^m\|_{Rad(X)}.
\end{aligned}$$

Therefore, for each  $(x_i)_{i=1}^\infty \in Rad(X)$  and each  $(y_i^*)_{i=1}^\infty \in \ell_{p^*}^w(Y^*)$ ,

$$\sum_{i=1}^\infty |y_i^*(u(x_i))| < \infty.$$

That is, for each  $(x_i)_{i=1}^\infty \in Rad(X)$ ,  $(u(x_i))_{i=1}^\infty \in \ell_p\langle Y \rangle$ . □

**Corollary 2.2.** *Let  $X$  and  $Y$  be Banach spaces and  $u$  be a continuous linear operator from  $X$  to  $Y$ . If  $u^*$  is almost summing, then for any  $q \geq 2$ ,  $u$  takes almost unconditionally summable sequences in  $X$  into members of  $\ell_q\langle Y \rangle$ .*

*Proof.* Let  $u \in \mathcal{L}(X, Y)$  such that  $u^* \in \Pi_{al.s}(Y^*, X^*)$ . Taking  $p = 2$  in previous theorem, we have that

$$\sum_{i=1}^\infty |y_i^*(u(x_i))| < \infty$$

for each  $(x_i)_{i=1}^\infty \in Rad(X)$  and each  $(y_i^*)_{i=1}^\infty \in \ell_2^w(Y^*)$ .

On the other hand,  $\ell_{q^*}^w(Y^*) \subseteq \ell_2^w(Y^*)$  for  $q^* \leq 2$ . Thus,

$$\sum_{i=1}^\infty |y_i^*(u(x_i))| < \infty$$

for each  $(x_i)_{i=1}^\infty \in \text{Rad}(X)$  and each  $(y_i^*)_{i=1}^\infty \in \ell_{q^*}^w(Y^*)$ .

Consequently, for each  $(x_i)_{i=1}^\infty \in \text{Rad}(X)$ ,  $(u(x_i))_{i=1}^\infty \in \ell_q\langle Y \rangle$ . □

Note that from (1.2) it is obvious that Corollary 2.2 extends the statement of Bu-Kranz Theorem.

**Corollary 2.3.** *Let  $X$  and  $Y$  be Banach spaces such that  $X$  has type  $p$ , with  $1 \leq p \leq 2$ , and  $u$  be a continuous linear operator from  $X$  to  $Y$ . If  $u^*$  is almost  $p^*$ -summing, then  $u$  is Cohen strongly  $p$ -summing.*

*Proof.* Let  $u$  be a continuous linear operator from  $X$  to  $Y$  such that  $u^*$  is almost  $p^*$ -summing. Given  $(x_i)_{i=1}^\infty \in \ell_p(X)$  and since  $X$  has type  $p$ , by [7, Proposition 12.4], we have that  $(x_i)_{i=1}^\infty \in \text{Rad}(X)$ . By Theorem 2.1,  $(u(x_i))_{i=1}^\infty \in \ell_p\langle Y \rangle$ . Hence  $u$  is Cohen strongly  $p$ -summing. □

One of the interesting problems of the theory of absolutely summing operators is to investigate when the adjoint of an operator is  $p$ -summing. A nice result of Cohen in [6] ensures that a linear operator  $u : X \rightarrow Y$  is Cohen strongly  $p$ -summing if, and only if,  $u^* : Y^* \rightarrow X^*$  is  $p^*$ -summing. The next result approaches this kind of problem and provides a type of reciprocal of Bu-Kranz Theorem when  $X$  has type  $p$ , with  $1 \leq p \leq 2$ .

**Theorem 2.4.** *Let  $X$  and  $Y$  be Banach spaces such that  $X$  has type  $p$ , with  $1 \leq p \leq 2$ , and  $u$  be a continuous linear operator from  $X$  to  $Y$ . If  $u$  takes almost unconditionally summable sequences in  $X$  into members of  $\ell_p\langle Y \rangle$ , then  $u^*$  is  $p^*$ -summing (or  $u$  is Cohen strongly  $p$ -summing).*

*Proof.* We just need to show that if  $u$  takes elements of  $\text{Rad}(X)$  into elements of  $\ell_p\langle Y \rangle$ , then  $u^*$  takes elements of  $\ell_{p^*}^w(Y^*)$  into elements of  $\ell_{p^*}(X^*)$ .

Let  $(x_i)_{i=1}^\infty \in \ell_p(X)$  and  $(y_i^*)_{i=1}^\infty \in \ell_{p^*}^w(Y^*)$ . As  $X$  has type  $p$ , by [7, Proposition 12.4] we have that  $(x_i)_{i=1}^\infty \in \text{Rad}(X)$  and by hypothesis  $(u(x_i))_{i=1}^\infty \in \ell_p\langle Y \rangle$ . Thus

$$\sum_{i=1}^{\infty} |y_i^*(u(x_i))| < \infty.$$

So

$$\sum_{i=1}^{\infty} |u^*(y_i^*)(x_i)| < \infty.$$

By arbitrariness of  $(x_i)_{i=1}^\infty \in \ell_p(X)$ ,  $(u^*(y_i^*))_{i=1}^\infty \in \ell_{p^*}(X^*)$ . □

**Corollary 2.5.** [4, Theorem 4] *Let  $X$  and  $Y$  be Banach spaces such that  $X$  has type 2 and  $u \in \mathcal{L}(X, Y)$ . Then  $u$  takes almost unconditionally summable sequences in  $X$  into members of  $\ell_2\langle Y \rangle$  if, and only if,  $u^*$  is 2-summing (or  $u$  is Cohen strongly 2-summing).*

*Proof.* Since  $X$  has type 2 it follows from [7, Proposition 11.10] that  $X^*$  has cotype 2 and by [7, Corollary 12.7] we have that  $\Pi_{al.s}(Y^*, X^*) = \Pi_2(Y^*, X^*)$ . Then the result follows of the Theorem 2.1 and Theorem 2.4. □

Theorem 2.4 tell us that the hypothesis

(\*)  $u \in \mathcal{L}(X, Y)$  takes almost unconditionally summable sequences in  $X$  into members of  $\ell_p\langle Y \rangle$ ,

is a sufficient condition for  $u^*$  to be  $p^*$ -summing, whenever  $X$  has type  $p$ . On the other hand, it is also known that there are operators with a  $p$ -summing adjoint which are not themselves  $p$ -summing. So the hypothesis (\*) is not a sufficient condition for  $u$  to be  $p$ -summing for some  $p \geq 1$ . However, for some Banach spaces  $X$  and  $Y$  we will show that the hypothesis (\*) is also a sufficient condition for  $u$  to be 1-summing. For this we shall recall the notion of  $\mathcal{L}_p$ -spaces. If  $\lambda > 1$  and  $1 \leq p \leq \infty$ , a Banach space  $Y$  is an  $\mathcal{L}_p$ -space if every finite dimensional subspace  $E$  of  $Y$  is contained in a finite dimensional subspace  $F$  of  $Y$  for which there is an isomorphism  $v : F \rightarrow \ell_p^{\dim F}$  such that  $\|v\|\|v^{-1}\| \leq \lambda$ . Also for this purpose we use the following theorem recently proved in [9].

**Theorem 2.6.** [9, Theorem 1.6] *Let  $X$  be a Banach space,  $Y$  be an  $\mathcal{L}_p$ -space with  $2 \leq p < \infty$  and  $u \in \mathcal{L}(X, Y)$ . If  $u^*$  is almost  $p$ -summing, then  $u$  is 1-summing.*

**Theorem 2.7.** *Let  $X$  and  $Y$  be Banach spaces such that  $X$  has type 2 and  $Y$  is an  $\mathcal{L}_2$ -space and  $u$  be a continuous linear operator from  $X$  to  $Y$ . If  $u$  takes almost unconditionally summable sequences in  $X$  into members of  $\ell_2\langle Y \rangle$ , then  $u$  is 1-summing.*

*Proof.* Suppose that  $u \in \mathcal{L}(X, Y)$  takes almost unconditionally summable sequences in  $X$  into members of  $\ell_2\langle Y \rangle$ . As  $X$  has type 2, Corollary 2.5 ensures that  $u^*$  is 2-summing. By (1.2),  $u^*$  is almost summing. Since  $Y$  is an  $\mathcal{L}_2$ -space, it follows from Theorem 2.6 that  $u$  is 1-summing.  $\square$

**Remark 2.8.** *From [6, Theorem 3.2.3(ii)] we know that if  $Y$  is an  $\mathcal{L}_2$ -space, then  $\mathcal{D}_2(X, Y) \subseteq \Pi_2(X, Y)$  for all Banach space  $X$ . Corollary 2.5 and Theorem 2.7 improve this inclusion, they ensure that  $\mathcal{D}_2(X, Y) \subseteq \Pi_1(X, Y)$  whenever  $X$  has type 2 and  $Y$  is an  $\mathcal{L}_2$ -space.*

### 3. EXTENDING BU'S THEOREM

The next result is a variant of Bu's Theorem to  $\mathcal{L}_{p^*}$ -spaces, with  $2 \leq p < \infty$ , and almost  $p$ -summing operators instead of Hilbert spaces and  $p$ -summing operators.

**Theorem 3.1.** *Let  $2 \leq p < \infty$ ,  $1 < q \leq \infty$ ,  $X$  and  $Y$  be Banach spaces such that  $X$  is an  $\mathcal{L}_{p^*}$ -space. Then*

$$\Pi_{al.s.p}(X, Y) \subseteq \mathcal{D}_q(X, Y).$$

*Proof.* Let  $u \in \Pi_{al.s.p}(X, Y)$ . Fix  $x_1, \dots, x_m \in X$  and  $y_1^*, \dots, y_m^* \in Y^*$ . Since  $X$  is an  $\mathcal{L}_{p^*}$ -space, there is a finite dimensional subspace  $F$  of  $X$  containing  $span\{x_1, \dots, x_m\}$  and an isomorphism  $v : F \rightarrow \ell_{p^*}^n$  such that  $\|v\|\|v^{-1}\| \leq \lambda$ .

Note that

$$vx_i = \sum_{k=1}^n \nu_{i,k} e_k, \quad i = 1, \dots, m,$$

where  $(e_k)_{k=1}^n$  is the canonical basis of  $\ell_{p^*}^n = (\ell_p^n)^*$ . Then, using the monotonicity of the  $\ell_p$  norms, the Hölder inequality and the Khinchin inequality it follows that

$$\begin{aligned}
\sum_{i=1}^m |y_i^*(u(x_i))| &= \sum_{i=1}^m |y_i^*(uv^{-1}v(x_i))| \\
&= \sum_{i=1}^m \left| y_i^* \left( uv^{-1} \left( \sum_{k=1}^n \nu_{i,k} e_k \right) \right) \right| \\
&\leq \sum_{i=1}^m \left( \sum_{k=1}^n |\nu_{i,k}|^{p^*} \right)^{1/p^*} \cdot \left( \sum_{k=1}^n |y_i^*(uv^{-1}(e_k))|^p \right)^{1/p} \\
&\leq \sum_{i=1}^m \|vx_i\| \cdot \left( \sum_{k=1}^n |y_i^*(uv^{-1}(e_k))|^2 \right)^{1/2} \\
&\leq \sum_{i=1}^m \|v\| \|x_i\| \cdot \frac{1}{A_{q^*}} \left( \int_0^1 \left| \sum_{k=1}^n r_k(t) y_i^*(uv^{-1}(e_k)) \right|^{q^*} dt \right)^{1/q^*} \\
&\leq \|v\| \cdot \frac{1}{A_{q^*}} \cdot \left( \sum_{i=1}^m \|x_i\|^q \right)^{1/q} \cdot \left( \sum_{i=1}^m \int_0^1 \left| \sum_{k=1}^n r_k(t) y_i^*(uv^{-1}(e_k)) \right|^{q^*} dt \right)^{1/q^*} \\
&= \|v\| \cdot \frac{1}{A_{q^*}} \cdot \|(x_i)_{i=1}^m\|_q \cdot \left( \int_0^1 \sum_{i=1}^m \left| y_i^* \left( \sum_{k=1}^n r_k(t) uv^{-1}(e_k) \right) \right|^{q^*} dt \right)^{1/q^*}.
\end{aligned}$$

But

$$\|(y_i^*)_{i=1}^m\|_{q^*,w} := \sup_{y^{**} \in B_{Y^{**}}} \left( \sum_{i=1}^m |y^{**}(y_i^*)|^{q^*} \right)^{1/q^*} = \sup_{y \in B_Y} \left( \sum_{i=1}^m |y_i^*(y)|^{q^*} \right)^{1/q^*}.$$

Then

$$\begin{aligned}
\sum_{i=1}^m |y_i^*(u(x_i))| &\leq \|v\| \cdot \frac{1}{A_{q^*}} \cdot \|(x_i)_{i=1}^m\|_q \cdot \left( \int_0^1 \left( \left\| \sum_{k=1}^n r_k(t) uv^{-1}(e_k) \right\| \cdot \|(y_i^*)_{i=1}^m\|_{q^*,w} \right)^{q^*} dt \right)^{1/q^*} \\
&\leq \|v\| \cdot \frac{1}{A_{q^*}} \cdot \|(x_i)_{i=1}^m\|_q \cdot \|(y_i^*)_{i=1}^m\|_{q^*,w} \cdot \left( \int_0^1 \left\| \sum_{k=1}^n r_k(t) uv^{-1}(e_k) \right\|^{q^*} dt \right)^{1/q^*}.
\end{aligned}$$

By Kahane's inequality we have

$$\sum_{i=1}^m |y_i^*(u(x_i))| \leq \|v\| \cdot \frac{1}{A_{q^*}} \cdot \|(x_i)_{i=1}^m\|_q \cdot \|(y_i^*)_{i=1}^m\|_{q^*,w} \cdot K_{2,q^*} \cdot \left( \int_0^1 \left\| \sum_{k=1}^n r_k(t) uv^{-1}(e_k) \right\|^2 dt \right)^{1/2}.$$

Since  $u$  is almost  $p$ -summing, it follows that  $uv^{-1}$  also is almost  $p$ -summing and as  $\|(e_k)_{k=1}^n\|_{p,w} = 1$  in  $\ell_p^n$ , we have

$$\begin{aligned} \sum_{i=1}^m |y_i^*(u(x_i))| &\leq \|v\| \cdot \frac{1}{A_{q^*}} \cdot \|(x_i)_{i=1}^m\|_q \cdot \|(y_i^*)_{i=1}^m\|_{q^*,w} \cdot K_{2,q^*} \cdot \pi_{al.s.p}(u) \cdot \|v^{-1}\| \cdot \|(e_k)_{k=1}^n\|_{p,w} \\ &\leq \lambda \cdot \frac{1}{A_{q^*}} \cdot K_{2,q^*} \cdot \pi_{al.s.p}(u) \cdot \|(x_i)_{i=1}^m\|_q \cdot \|(y_i^*)_{i=1}^m\|_{q^*,w}. \end{aligned}$$

Therefore,  $u \in \mathcal{D}_q(X, Y)$ . □

As a corollary, since (1.2) is valid and every Hilbert space is an  $\mathcal{L}_2$ -space, we have an improvement of the Bu's Theorem [3, Main Theorem]:

**Corollary 3.2** (Bu's Theorem improved). *Consider  $1 < q \leq \infty$ . Let  $X$  be an  $\mathcal{L}_2$ -space,  $Y$  be a Banach space. Then*

$$\Pi_{al.s}(X, Y) \subseteq \mathcal{D}_q(X, Y).$$

In [7, Corollary 9.12(b)] there is a characterization of closed subspace of  $L_r$ , with  $1 \leq r < \infty$ , which is: a Banach space  $Y$  is an isomorphic to a closed subspace of  $L_r$  if, and only if, for all Banach space  $X$

$$\mathcal{D}_{r^*}(X, Y) \subseteq \Pi_r(X, Y).$$

In the next corollary, we show a coincidence result among the classes of the Cohen strongly  $r^*$ -summing, absolutely  $r$ -summing and almost summing operators.

**Corollary 3.3.** *Let  $X$  be an  $\mathcal{L}_2$ -space and  $Y$  be an isomorphic to a closed subspace of  $L_r$ , with  $1 \leq r < \infty$ . Then*

$$\Pi_{al.s}(X, Y) = \mathcal{D}_{r^*}(X, Y) = \Pi_r(X, Y).$$

*Proof.* Let  $X$  be an  $\mathcal{L}_2$ -space. So considering  $p = 2$  and  $q = r^*$  in Theorem 3.1 we have that

$$\Pi_{al.s}(X, Y) \subseteq \mathcal{D}_{r^*}(X, Y).$$

Since  $Y$  is an isomorphic to a closed subspace of  $L_r$ , it follows from [7, Corollary 9.12(b)] and (1.2) that  $\mathcal{D}_{r^*}(X, Y) \subseteq \Pi_r(X, Y) \subseteq \Pi_{al.s}(X, Y)$ . Therefore,

$$\Pi_{al.s}(X, Y) = \mathcal{D}_{r^*}(X, Y) = \Pi_r(X, Y). □$$

#### 4. EXTENDING KWAPIEŃ'S THEOREM

The following theorem is the main result of this section. It is an improvement of Kwapien's theorem in [8].

**Theorem 4.1.** *The following properties of Banach space  $X$  are equivalent.*

- (i) *The space  $X$  is isomorphic to a Hilbert space.*
- (ii) *For every  $1 < q \leq \infty$  and every Banach space  $Y$ ,  $\Pi_{al.s}(X, Y) \subseteq \mathcal{D}_q(X, Y)$ .*
- (iii) *For every Banach space  $Y$ ,  $\Pi_{al.s}(X, Y) \subseteq \mathcal{D}_2(X, Y)$ .*

*Proof.* (i)  $\Rightarrow$  (ii) If  $X$  is isomorphic to a Hilbert space, then it is an  $\mathcal{L}_2$ -space. By Theorem 3.1 it follows that  $\Pi_{al.s}(X, Y) \subseteq \mathcal{D}_q(X, Y)$  for all Banach space  $Y$  and all  $1 < q \leq \infty$ .

(ii)  $\Rightarrow$  (iii) It is obvious.

(iii)  $\Rightarrow$  (i) For every Banach space  $Y$ , suppose that  $u \in \Pi_2(X, Y)$ . By (1.2),  $u \in \Pi_{al.s}(X, Y)$ . Now using the hypothesis, it follows that  $u \in \mathcal{D}_2(X, Y)$  and hence Kwapien's theorem implies that  $X$  is isomorphic to a Hilbert space. □

**Corollary 4.2.** *If  $X$  is isomorphic to a Hilbert space and  $Y$  has type 2, then*

$$\Pi_{al.s}(X, Y) = \mathcal{D}_2(X, Y).$$

*Proof.* Since  $X$  is isomorphic to a Hilbert, Theorem 4.1 ensures that  $\Pi_{al.s}(X, Y) \subseteq \mathcal{D}_2(X, Y)$ . Now as  $Y$  has type 2, it follows from [7, Corollary 12.21] that  $\mathcal{D}_2(X, Y) \subseteq \Pi_{al.s}(X, Y)$ . Thus, the proof is complete. □

#### REFERENCES

- [1] G. Bennett, *Schur multipliers*, Duke Math. J., **44** (1977), 603–639.
- [2] G. Botelho, H.A. Braunss and H. Junek, *Almost  $p$ -summing polynomials and multilinear mappings*, Arch. Math., **76** (2001), 109–118.
- [3] Q. Bu, *Some mapping properties of  $p$ -summing operators with hilbertian domain*, Contemp. Math., **328** (2003), 145–149.
- [4] Q. Bu and P. Kranz, *Some mapping properties of  $p$ -summing adjoint operators*, J. Math. Anal. Appl., **303** (2005), 585–590.
- [5] J. S. Cohen, *A characterization of inner-product spaces using absolutely 2-summing operators*, Studia Math., **38** (1970), 271–276.
- [6] J. S. Cohen, *Absolutely  $p$ -summing,  $p$ -nuclear operators and their conjugates*, Math. Ann., **201** (1973), 177–200.
- [7] J. Diestel, H. Jarchow and A. Tonge, *Absolutely Summing Operators*, Cambridge Studies in Advanced Mathematics, Cambridge University Press, 1995.
- [8] S. Kwapien, *A linear topological characterization of inner product space*, Studia Math., **38** (1970). 277–278.
- [9] R. Macedo, D. Pellegrino and J. Santos, *Nonlinear variants of a theorem of Kwapien*, to appear in Isr. J. Math.

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DA PARAÍBA, 58.051-900 - JOÃO PESSOA, BRAZIL.  
*Email address:* `renato.burity@academico.ufpb.br`

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DA PARAÍBA, 58.051-900 - JOÃO PESSOA, BRAZIL.  
*Email address:* `joedson.santos@academico.ufpb.br`