

SOME FIXED-CIRCLE RESULTS IN C^* -ALGEBRA VALUED METRIC SPACES

NILAY DEĞİRMEN

ABSTRACT. In this paper, we consider fixed-circle problem in C^* -algebra valued metric spaces and prove some fixed-circle theorems for self-mappings by defining the notion of fixed-circle on such spaces with geometric interpretation. Furthermore, we give some illustrative examples to substantiate the importance of our newly obtained results.

1. INTRODUCTION AND PRELIMINARIES

The Banach contraction principle [1] is a popular and effective tool used for the existence and uniqueness of solutions of many nonlinear problems arising in physics and engineering sciences. Up to now, researchers generalized the Banach contraction principle in many directions and obtained new results in different metric spaces. With this idea in mind, in 2014, Ma *et al.* [2] introduced the concept of C^* -algebra valued metric space and established some fixed point theorems for self-mappings satisfying the contractive conditions on such spaces. Also, in 2015, Batul and Kamran [3] generalized it by weakening the contractive condition introduced by Ma *et al.* [2]. Going in the same direction, many articles on fixed point results in C^* -algebra valued metric spaces, we refer to [4, 5, 6, 7, 8, 9].

On the other hand, as a geometric approach to the fixed point theory, in [10], Özgür and Taş initiated the investigations concerning a fixed-circle problem in metric spaces and examined some fixed-circle theorems for self-mappings on metric spaces with geometric interpretation by giving some necessary examples to validate their own findings. Since this subject has been developed very fast in recent times due to theoretical mathematical studies and some applications in different fields of mathematical sciences such as neural networks, it has attracted considerable interest from many authors. New solutions of fixed-circle problem was investigated with various aspects and new contractive conditions on both metric spaces and some generalized metric spaces. For fixed-circle results using different techniques, we recommend [11, 12, 13, 14, 15, 16].

In the most existing literature, it has been observed that fixed circles have not been defined in C^* -algebra valued metric spaces. This motivates us to analyze solutions of the fixed-circle problem in such spaces with geometric interpretation. So, our purpose in this study is to define the notion of a fixed circle on a C^* -algebra valued metric space and obtain some fixed-circle theorems for self-mappings on C^* -algebra valued metric spaces. Also, we construct some nontrivial illustrative examples of mappings which have or not fixed circles. Since the theory of

2010 *Mathematics Subject Classification.* 47H10, 54H25, 46L07, 37E10.

Key words and phrases. C^* -algebra, C^* -algebra valued metric space, fixed circle, the existence theorem, the uniqueness theorem.

C^* -algebra is one of the most extensive research areas in operator theory and functional analysis, which is extremely active and having huge applications to theoretical physics and noncommutative geometry, we believe that the results of this article will contribute to studying on fixed-circles with different aspects for C^* -algebra valued metric spaces in the future.

Now, we summarize a number of known definitions and results about C^* -algebras and C^* -algebra valued metric spaces, which will be needed in our subsequent discussions.

We start with the definition of a C^* -algebra and its some properties used in this research.

Definition 1. [17] *A mapping $x \rightarrow x^*$ of a complex algebra \mathbb{A} into \mathbb{A} is called an involution on \mathbb{A} if the following properties hold for all $x, y \in \mathbb{A}$ and $\lambda \in \mathbb{C}$:*

- (i) $(x^*)^* = x$,
- (ii) $(xy)^* = y^*x^*$,
- (iii) $(\lambda x + y)^* = \bar{\lambda}x^* + y^*$.

A complex Banach algebra A with an involution such that for every x in A

$$\|x^*x\| = \|x\|^2$$

is called a C^ -algebra.*

In the rest of paper, \mathbb{A} will denote a unital C^* -algebra with a unit I .

Let $\mathbb{A}_h = \{x \in \mathbb{A} : x = x^*\}$. An element $x \in \mathbb{A}$ is called positive if $x \in \mathbb{A}_h$ and $\sigma(x) \subset \mathbb{R}^+$ where $\sigma(x) = \{\lambda \in \mathbb{R} : x - \lambda I \text{ is non-invertible}\}$, the spectrum of x in \mathbb{A} . If $x \in \mathbb{A}$ is positive, we write it as $\theta \preceq x$, where θ is the zero element in \mathbb{A} . We denote the set of all positive elements of \mathbb{A} by \mathbb{A}_+ . Also, \mathbb{A}_h becomes a partially ordered set by defining $x \preceq y$ to mean $y - x \in \mathbb{A}_+$ [17].

The following statements about involution on \mathbb{A} , positive elements of \mathbb{A} and partial order \preceq on \mathbb{A}_h are true:

- (i) $\|x^*\| = \|x\|$ for all $x \in \mathbb{A}$.
- (ii) If $x \in \mathbb{A}$ is invertible, then x^* is invertible and $(x^*)^{-1} = (x^{-1})^*$.
- (iii) If $x, y, z \in \mathbb{A}_h$, then $x \preceq y$ implies $x + z \preceq y + z$.
- (iv) If $x, y \in \mathbb{A}_+$ and $\alpha, \beta \in \mathbb{R}^+ \cup \{0\}$, then $\alpha x + \beta y \in \mathbb{A}_+$.
- (v) $\mathbb{A}_+ = \{x^*x : x \in \mathbb{A}\}$.
- (vi) If $x, y \in \mathbb{A}_h$ and $z \in \mathbb{A}$, then $x \preceq y$ implies $z^*xz \preceq z^*yz$.
- (vii) If $\theta \preceq x \preceq y$, then $\|x\| \leq \|y\|$ [17].

Using the concept of a positive element in a C^* -algebra, in 2014, Ma and Jiang [2] introduced the notion of a C^* -algebra valued metric space in the following way:

Definition 2. [2] *Let X be a nonempty set. Suppose the mapping $d : X \times X \rightarrow \mathbb{A}$ satisfies:*

- (i) $\theta \preceq d(x, y)$ for all $x, y \in X$ and $d(x, y) = \theta \Leftrightarrow x = y$;
- (ii) $d(x, y) = d(y, x)$ for all $x, y \in X$;
- (iii) $d(x, y) \preceq d(x, z) + d(z, y)$ for all $x, y, z \in X$.

Then, d is called C^ -algebra valued metric on X and (X, \mathbb{A}, d) is called a C^* -algebra valued metric space.*

It is clear that such spaces generalize the concept of metric spaces. The main idea consists in using the set of all positive elements of a unital C^* -algebra instead of the set of real numbers.

Example 1. [2] Let E be a Lebesgue measurable set and $L(L^2(E))$ denote the set of bounded linear operators on Hilbert space $L^2(E)$. Define $d : L^\infty(E) \times L^\infty(E) \rightarrow L(L^2(E))$ by

$$d(f, g) = \pi_{|f-g|}$$

for all $f, g \in L^\infty(E)$, where $\pi_h : L^2(E) \rightarrow L^2(E)$ is the multiplication operator defined by

$$\pi_h(\varphi) = h \cdot \varphi$$

for all $\varphi \in L^2(E)$. Then, d is a C^* -algebra valued metric and $(L^\infty(E), L(L^2(E)), d)$ is a complete C^* -algebra valued metric space.

Example 2. [2] Define $d : \mathbb{R} \times \mathbb{R} \rightarrow M_2(\mathbb{R})$ by

$$d(x, y) = \text{diag}(|x - y|, \alpha|x - y|)$$

for all $x, y \in \mathbb{R}$, where $\alpha \geq 0$ is a constant, the norm on $M_2(\mathbb{R})$ is defined by

$$\left\| \begin{bmatrix} a & b \\ c & d \end{bmatrix} \right\| = \max\{|a|, |b|, |c|, |d|\},$$

and partial ordering on $M_2(\mathbb{R})$ is given by

$$\begin{bmatrix} a_1 & b_1 \\ c_1 & d_1 \end{bmatrix} \preceq \begin{bmatrix} a_2 & b_2 \\ c_2 & d_2 \end{bmatrix} \iff a_1 \leq a_2, b_1 \leq b_2, c_1 \leq c_2 \text{ and } d_1 \leq d_2.$$

Then, d is a C^* -algebra valued metric and $(\mathbb{R}, M_2(\mathbb{R}), d)$ is a C^* -algebra valued metric space.

Example 3. [3] Let $X = [-1, 1] \times [-1, 1]$. Define $d : X \times X \rightarrow \mathbb{R}^2$ by

$$d(x, y) = (|x_1 - y_1|, |x_2 - y_2|)$$

for all $x = (x_1, x_2), y = (y_1, y_2) \in X$, where the norm on \mathbb{R}^2 is defined by

$$\|(a, b)\| = \max\{|a|, |b|\},$$

and partial ordering on \mathbb{R}^2 is given by

$$(a, b) \preceq (c, d) \iff a \leq c \text{ and } b \leq d.$$

Then, d is a C^* -algebra valued metric and (X, \mathbb{R}^2, d) is a C^* -algebra valued metric space.

Example 4. [6] Define $d : \mathbb{R} \times \mathbb{R} \rightarrow M_2(\mathbb{R})$ by

$$d(x, y) = \begin{cases} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, & x \neq y \\ 0, & x = y \end{cases}$$

for all $x, y \in \mathbb{R}$. Then, d is a C^* -algebra valued metric and $(\mathbb{R}, M_2(\mathbb{R}), d)$ is a C^* -algebra valued metric space.

Based on the idea of the Banach contraction principle [1] in classical metric spaces, Ma and Jiang [2] established the following main theorems which implies the existence and uniqueness of fixed point on complete C^* -algebra valued metric spaces.

Theorem 1. [2] *If (X, \mathbb{A}, d) is a complete C^* -algebra valued metric space and $T : X \rightarrow X$ is a C^* -algebra valued contractive mapping on X , that is, there exists an $A \in \mathbb{A}$ with $\|A\| < 1$ such that*

$$(1.1) \quad d(Tx, Ty) \preceq A^* d(x, y) A$$

for all $x, y \in X$, then, there exists a unique fixed point in X .

In the following theorem and Theorem 12, we denote the set $\{a \in \mathbb{A} : ab = ba \text{ for all } b \in \mathbb{A}\}$ by \mathbb{A}' .

Theorem 2. [2] *Let (X, \mathbb{A}, d) be a complete C^* -algebra valued metric space. Suppose the mapping $T : X \rightarrow X$ satisfies for all $x, y \in X$ such that*

$$(1.2) \quad d(Tx, Ty) \preceq A (d(Tx, y) + d(Ty, x))$$

where $A \in \mathbb{A}'_+$ and $\|A\| < \frac{1}{2}$. Then, there exists a unique fixed point in X .

Inspired by preceding observation, many authors stated new types of contractive mappings and studied fixed point theorems. Shehwar et al. [4] gave the extension of Caristi's fixed point theorem [18, 19] for self-mappings defined on C^* -algebra valued metric spaces, which guarantees the existence of fixed point as follows:

Theorem 3. [4] *Let (X, \mathbb{A}, d) be a complete C^* -algebra valued metric space, $\phi : X \rightarrow \mathbb{A}_+$ be a lower semi continuous map and $T : X \rightarrow X$ be such that*

$$(1.3) \quad d(x, Tx) \preceq \phi(x) - \phi(Tx)$$

for all $x \in X$. Then, T has at least one fixed point in X .

Kadelburg and Radenovic [5] introduced the extension of Ćirić's fixed point theorem [20] for self-mappings defined on C^* -algebra valued metric spaces, which guarantees the existence and uniqueness of fixed point as follows:

Theorem 4. [5] *Let (X, \mathbb{A}, d) be a C^* -algebra valued metric space and $T : X \rightarrow X$ be a mapping on X . Suppose that there exists an $A \in \mathbb{A}$ with $\|A\| < 1$ such that for all $x, y \in X$ there exists $u(x, y) \in \{d(x, y), d(x, Tx), d(y, Ty), d(x, Ty), d(y, Tx)\}$ such that*

$$(1.4) \quad d(Tx, Ty) \preceq A^* u(x, y) A.$$

Then, T has a unique fixed point in X .

2. MAIN RESULTS

We begin this section by introducing the concept of a circle on a C^* -algebra valued metric space.

Definition 3. *Let (X, \mathbb{A}, d) be a C^* -algebra valued metric space, $x_0 \in X$ and $r \in \mathbb{A}_+$. Then, the circle with the centered x_0 and the radius r is defined by*

$$C_{x_0, r}^{C^*} = \{x \in X : d(x, x_0) = r\}.$$

Example 5. *Consider the C^* -algebra valued metric space $(L^\infty(E), L(L^2(E)), d)$ given in Example 1 for $E = [0, 1]$. Choose the center x_0 as the function f defined by*

$$f : [0, 1] \rightarrow \mathbb{R}, \quad f(x) = \chi_{[\frac{1}{2}, 1]}(x) = \begin{cases} 1, & x \in [\frac{1}{2}, 1] \\ 0, & x \notin [\frac{1}{2}, 1] \end{cases}$$

and the radius r as the multiplication operator π_h defined by

$$\pi_h : L^2 [0, 1] \rightarrow L^2 [0, 1], \quad \pi_h (\varphi) = h \cdot \varphi$$

for the function $h : [0, 1] \rightarrow \mathbb{R}$, $h(x) = \begin{cases} 1, & x \in (\mathbb{R} \setminus \mathbb{Q}) \cap [0, 1] \\ \infty, & x \in \mathbb{Q} \cap [0, 1] \end{cases}$. Then, it can be easily seen that $f, h \in L^\infty [0, 1]$ and $\pi_h \in L(L^2 [0, 1])$. Thus, we get

$$\begin{aligned} C_{f, \pi_h}^{C^*} &= \{g \in L^\infty [0, 1] : d(g, f) = \pi_h\} \\ &= \{g \in L^\infty [0, 1] : \pi_{|g-f|} = \pi_h\} \\ &= \{g \in L^\infty [0, 1] : |g - f| = h\} \\ &= \{g \in L^\infty [0, 1] : |g(x) - f(x)| = h(x) \text{ for each } x \in [0, 1]\} \\ &= \left\{ g \in L^\infty [0, 1] : |g(x) - 0| = 1 \text{ for each } x \in (\mathbb{R} \setminus \mathbb{Q}) \cap \left[0, \frac{1}{2}\right) \right\} \\ &\quad \cup \left\{ g \in L^\infty [0, 1] : |g(x) - 0| = \infty \text{ for each } x \in \mathbb{Q} \cap \left[0, \frac{1}{2}\right) \right\} \\ &\quad \cup \left\{ g \in L^\infty [0, 1] : |g(x) - 1| = 1 \text{ for each } x \in (\mathbb{R} \setminus \mathbb{Q}) \cap \left[\frac{1}{2}, 1\right] \right\} \\ &\quad \cup \left\{ g \in L^\infty [0, 1] : |g(x) - 1| = \infty \text{ for each } x \in \mathbb{Q} \cap \left[\frac{1}{2}, 1\right] \right\} \\ &= \left\{ g \in L^\infty [0, 1] : g(x) = -1 \text{ or } g(x) = 1 \text{ for each } x \in (\mathbb{R} \setminus \mathbb{Q}) \cap \left[0, \frac{1}{2}\right) \right\} \\ &\quad \cup \left\{ g \in L^\infty [0, 1] : g(x) = -\infty \text{ or } g(x) = \infty \text{ for each } x \in \mathbb{Q} \cap \left[0, \frac{1}{2}\right) \right\} \\ &\quad \cup \left\{ g \in L^\infty [0, 1] : g(x) = 0 \text{ or } g(x) = 2 \text{ for each } x \in (\mathbb{R} \setminus \mathbb{Q}) \cap \left[\frac{1}{2}, 1\right] \right\} \\ &\quad \cup \left\{ g \in L^\infty [0, 1] : g(x) = -\infty \text{ or } g(x) = \infty \text{ for each } x \in \mathbb{Q} \cap \left[\frac{1}{2}, 1\right] \right\}. \end{aligned}$$

For example, the function g defined by

$$g : [0, 1] \rightarrow \mathbb{R}, \quad g(x) = \begin{cases} 1 & x \in (\mathbb{R} \setminus \mathbb{Q}) \cap \left[0, \frac{1}{7}\right) \\ -1 & x \in (\mathbb{R} \setminus \mathbb{Q}) \cap \left(\frac{1}{7}, \frac{1}{2}\right) \\ \infty & x \in \mathbb{Q} \cap \left[0, \frac{1}{2}\right) \\ 2 & x \in (\mathbb{R} \setminus \mathbb{Q}) \cap \left[\frac{1}{2}, \frac{99}{100}\right) \\ 0 & x \in (\mathbb{R} \setminus \mathbb{Q}) \cap \left[\frac{99}{100}, 1\right] \\ -\infty & x \in \mathbb{Q} \cap \left[\frac{1}{2}, 1\right] \end{cases}$$

is in the circle $C_{f, \pi_h}^{C^*}$.

Example 6. Consider the C^* -algebra valued metric space $(\mathbb{R}, M_2(\mathbb{R}), d)$ given in Example 2 for $\alpha = 3$. Choose the center $x_0 = 0$ and the radii $r_1 = \begin{bmatrix} 2 & 0 \\ 0 & 6 \end{bmatrix}$,

$r_2 = \begin{bmatrix} 2 & 0 \\ -1 & 5 \end{bmatrix}$. Then, we get

$$\begin{aligned} C_{0,r_1}^{C^*} &= \left\{ x \in \mathbb{R} : d(x, 0) = \begin{bmatrix} 2 & 0 \\ 0 & 6 \end{bmatrix} \right\} \\ &= \left\{ x \in \mathbb{R} : \begin{bmatrix} |x| & 0 \\ 0 & 3|x| \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 6 \end{bmatrix} \right\} \\ &= \{x \in \mathbb{R} : |x| = 2\} \\ &= \{-2, 2\} \end{aligned}$$

and

$$\begin{aligned} C_{0,r_2}^{C^*} &= \left\{ x \in \mathbb{R} : d(x, 0) = \begin{bmatrix} 2 & 0 \\ -1 & 5 \end{bmatrix} \right\} \\ &= \left\{ x \in \mathbb{R} : \begin{bmatrix} |x| & 0 \\ 0 & 3|x| \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ -1 & 5 \end{bmatrix} \right\} \\ &= \emptyset. \end{aligned}$$

Example 7. Consider the C^* -algebra valued metric space (X, \mathbb{R}^2, d) given in Example 3. Choose the center $x_0 = (\frac{-1}{2}, 0)$ and the radius $r = (1, 1)$. Then, we get

$$\begin{aligned} C_{(\frac{-1}{2}, 0), (1, 1)}^{C^*} &= \left\{ x = (x_1, x_2) \in X : d\left(x, \left(\frac{-1}{2}, 0\right)\right) = (1, 1) \right\} \\ &= \left\{ x = (x_1, x_2) \in X : \left|x_1 + \frac{1}{2}\right| = |x_2| = 1 \right\} \\ &= \left\{ \left(\frac{-3}{2}, -1\right), \left(\frac{-3}{2}, 1\right), \left(\frac{1}{2}, -1\right), \left(\frac{1}{2}, 1\right) \right\}. \end{aligned}$$

Example 8. Consider the C^* -algebra valued metric space $(\mathbb{R}, M_2(\mathbb{R}), d)$ given in Example 4. Choose the center $x_0 = 3$ and the radius $r = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$. Then, we get

$$\begin{aligned} C_{3,r}^{C^*} &= \left\{ x \in \mathbb{R} : d(x, 3) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right\} \\ &= \{x \in \mathbb{R} : x \neq 3\} \\ &= \mathbb{R} - \{3\}. \end{aligned}$$

Example 9. Define $d : \mathbb{R} \times \mathbb{R} \rightarrow M_2(\mathbb{R})$ by

$$d(x, y) = \text{diag}(|e^x - e^y|, \alpha |e^x - e^y|)$$

for all $x, y \in \mathbb{R}$, where $\alpha \geq 0$ is a constant. It is easy to verify d is a C^* -algebra valued metric and $(\mathbb{R}, M_2(\mathbb{R}), d)$ is a complete C^* -algebra valued metric space. If we choose the center $x_0 = 1$ and the radii $r_1 = \begin{bmatrix} 2e & 0 \\ 0 & 4e \end{bmatrix}$, $r_2 = \begin{bmatrix} \frac{e}{2} & 0 \\ 0 & e \end{bmatrix}$ we get

for $\alpha = 2$

$$\begin{aligned}
 C_{1,r_1}^{C^*} &= \left\{ x \in \mathbb{R} : d(x, 1) = \begin{bmatrix} 2e & 0 \\ 0 & 4e \end{bmatrix} \right\} \\
 &= \left\{ x \in \mathbb{R} : \begin{bmatrix} |e^x - e| & 0 \\ 0 & 2|e^x - e| \end{bmatrix} = \begin{bmatrix} 2e & 0 \\ 0 & 4e \end{bmatrix} \right\} \\
 &= \{x \in \mathbb{R} : |e^x - e| = 2e\} \\
 &= \{1 + \ln 3\}
 \end{aligned}$$

and

$$\begin{aligned}
 C_{1,r_2}^{C^*} &= \left\{ x \in \mathbb{R} : d(x, 1) = \begin{bmatrix} \frac{e}{2} & 0 \\ 0 & e \end{bmatrix} \right\} \\
 &= \left\{ x \in \mathbb{R} : \begin{bmatrix} |e^x - e| & 0 \\ 0 & 2|e^x - e| \end{bmatrix} = \begin{bmatrix} \frac{e}{2} & 0 \\ 0 & e \end{bmatrix} \right\} \\
 &= \left\{ x \in \mathbb{R} : |e^x - e| = \frac{e}{2} \right\} \\
 &= \{1 + \ln 3 - \ln 2, 1 - \ln 2\}.
 \end{aligned}$$

2.1. The existence of fixed circles. In this part, we introduce the notion of a fixed circle on a C^* -algebra valued metric space. Then, we prove some fixed-circle theorems that guarantee the existence of fixed-circles for a self-mapping satisfying some conditions on C^* -algebra valued metric spaces.

Now, we define the new concept as follows:

Definition 4. Let (X, \mathbb{A}, d) be a C^* -algebra valued metric space, $C_{x_0,r}^{C^*}$ be a circle on X and $T : X \rightarrow X$ be a self-mapping. If $Tx = x$ for all $x \in C_{x_0,r}^{C^*}$, then the circle $C_{x_0,r}^{C^*}$ is called as the fixed circle of T .

The first solution of fixed-circle problem in C^* -algebra valued metric spaces is given using the inequality (1.3) in Theorem 3 which is an extension of Caristi's fixed point theorem for mappings defined on C^* -algebra valued metric spaces as follows:

Theorem 5. Let (X, \mathbb{A}, d) be a C^* -algebra valued metric space, $C_{x_0,r}^{C^*}$ be any circle on X . Define the mapping $\varphi : X \rightarrow \mathbb{A}_+$ as

$$(2.1) \quad \varphi(x) = d(x, x_0)$$

for all $x \in X$. If T is a self-mapping defined on X satisfying the conditions

$$(2.2) \quad d(x, Tx) \preceq \varphi(x) - \varphi(Tx)$$

and

$$(2.3) \quad r \preceq d(Tx, x_0)$$

for all $x \in C_{x_0,r}^{C^*}$, then the circle $C_{x_0,r}^{C^*}$ is a fixed circle of T .

Proof. Let x be any point in the circle $C_{x_0,r}^{C^*}$. Then, using the (2.2), (2.1), (2.3) and the definition of the relation \preceq , we get

$$\begin{aligned}
 d(x, Tx) &\preceq \varphi(x) - \varphi(Tx) \\
 &= d(x, x_0) - d(Tx, x_0) \\
 &= r - d(Tx, x_0) \\
 &\preceq r - r = 0
 \end{aligned}$$

and so $d(x, Tx) = 0$ which means that $Tx = x$. Then, we obtain that $C_{x_0, r}^{C^*}$ is a fixed circle of T . \square

Remark 1. *Theorem 5 guarantees the existence of at least one fixed circle of a self-mapping on a C^* -algebra valued metric space, while Theorem 3 guarantees the existence of at least one fixed point of a self-mapping on a C^* -algebra valued metric space. We note that if the circle $C_{x_0, r}^{C^*}$ has only one element, Theorem 5 is a special case of Theorem 3.*

Remark 2. *The inequality (2.2) says that Tx is not in the exterior of the circle $C_{x_0, r}^{C^*}$ for each $x \in C_{x_0, r}^{C^*}$. In the same way, the inequality (2.3) says that Tx is not in the interior of the circle $C_{x_0, r}^{C^*}$ for each $x \in C_{x_0, r}^{C^*}$. It follows that $T(C_{x_0, r}^{C^*}) \subset C_{x_0, r}^{C^*}$ under the conditions (2.2) and (2.3).*

The following example illustrates Theorem 5.

Example 10. *Consider the C^* -algebra valued metric space $(L^\infty(E), L(L^2(E)), d)$ given in Example 1 for $E = [0, 1]$ and the circle $C_{f, \pi_h}^{C^*}$ given in Example 5. Define the function*

$$g_0 : [0, 1] \rightarrow \mathbb{R}, \quad g_0(x) = \begin{cases} 1, & x \in (\mathbb{R} \setminus \mathbb{Q}) \cap [0, 1] \\ \infty, & x \in \mathbb{Q} \cap [0, 1] \end{cases}.$$

It is easy to show that $g_0 \in L^\infty[0, 1]$. Let us define the self-mapping $T : L^\infty[0, 1] \rightarrow L^\infty[0, 1]$ as

$$Tg = \begin{cases} g, & g \in C_{f, \pi_h}^{C^*} \\ g_0, & g \notin C_{f, \pi_h}^{C^*} \end{cases}.$$

Then, with a direct computation it can be seen that the self-mapping T satisfies the conditions (2.2) and (2.3). That is to say that the circle $C_{f, \pi_h}^{C^}$ is a fixed circle of T .*

Now, we give an example of a self-mapping which satisfies the condition (2.2) and does not satisfy the condition (2.3).

Example 11. *Consider the C^* -algebra valued metric space $(\mathbb{R}, M_2(\mathbb{R}), d)$ given in Example 2 for $\alpha = 3$ and the circle $C_{0, r_1}^{C^*}$ given in Example 6. Let us define the self-mapping $T : \mathbb{R} \rightarrow \mathbb{R}$ as*

$$Tx = \begin{cases} \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}.$$

With a simple verification it can be shown that T satisfies the condition (2.2) and does not satisfy the condition (2.3). Notice that the circle $C_{0, r_1}^{C^}$ is not a fixed circle of T .*

Now, we present an example in order to show that Teorem 5 fail out, if the condition (2.3) is satisfied but the condition (2.2) is not.

Example 12. Consider the C^* -algebra valued metric space (X, \mathbb{R}^2, d) given in Example 3 and the circle $C_{(\frac{-1}{2}, 0), (1, 1)}^{C^*}$ given in Example 7. Let us define the self-mapping $T : X \rightarrow X$ as

$$Tx = \begin{cases} (\frac{-3}{2}, -2), & x = (\frac{-3}{2}, -1) \\ (\frac{-3}{2}, 2), & x = (\frac{-3}{2}, 1) \\ (\frac{1}{2}, -2), & x = (\frac{1}{2}, -1) \\ (\frac{1}{2}, 2), & x = (\frac{1}{2}, 1) \\ (1, 5), & x \notin C_{(\frac{-1}{2}, 0), (1, 1)}^{C^*} \end{cases}.$$

Then, T satisfies the condition (2.3) and does not satisfy the condition (2.2). Notice that the circle $C_{(\frac{-1}{2}, 0), (1, 1)}^{C^*}$ is not a fixed circle of T .

We now state another existence theorem for fixed circles of a self-mapping on a C^* -algebra valued metric space as follows:

Theorem 6. Let (X, \mathbb{A}, d) be a C^* -algebra valued metric space, $C_{x_0, r}^{C^*}$ be any circle on X and the mapping φ be as in (2.1). If T is a self-mapping defined on X satisfying the conditions

$$(2.4) \quad d(x, Tx) \preceq \varphi(x) + \varphi(Tx) - 2r$$

and

$$(2.5) \quad d(Tx, x_0) \preceq r$$

for all $x \in C_{x_0, r}^{C^*}$, then the circle $C_{x_0, r}^{C^*}$ is a fixed circle of T .

Proof. Let x be any point in the circle $C_{x_0, r}^{C^*}$. Then, using the (2.4), (2.1), (2.5) and the definition of the relation \preceq , we get

$$\begin{aligned} d(x, Tx) &\preceq \varphi(x) + \varphi(Tx) - 2r \\ &= d(x, x_0) + d(Tx, x_0) - 2r \\ &= d(Tx, x_0) - r \\ &\preceq r - r = 0 \end{aligned}$$

and so $d(x, Tx) = 0$ which implies that $Tx = x$. Then, we deduce that $C_{x_0, r}^{C^*}$ is a fixed circle of T . \square

Remark 3. The inequality (2.4) means that Tx is not in the interior of the circle $C_{x_0, r}^{C^*}$ for each $x \in C_{x_0, r}^{C^*}$. In the same fashion, the inequality (2.5) means that Tx is not in the exterior of the circle $C_{x_0, r}^{C^*}$ for each $x \in C_{x_0, r}^{C^*}$. These two results show that $T(C_{x_0, r}^{C^*}) \subset C_{x_0, r}^{C^*}$ under the conditions (2.4) and (2.5).

We continue with an example which satisfy the requirements of Theorem 6 as follows:

Example 13. Consider the C^* -algebra valued metric space $(\mathbb{R}, M_2(\mathbb{R}), d)$ for $\alpha = 2$ and the circle $C_{1, r_2}^{C^*}$ given in Example 9. Let us define the self-mapping $T : \mathbb{R} \rightarrow \mathbb{R}$ as

$$Tx = \begin{cases} x, & x \in C_{1, r_2}^{C^*} \\ 1 - \ln 3, & x \notin C_{1, r_2}^{C^*} \end{cases}.$$

Clearly, T satisfies the conditions (2.4) and (2.5), and we derive that the circle $C_{1, r_2}^{C^*}$ is a fixed circle of T .

Now, we introduce an example of a self-mapping which satisfies the condition (2.4) and does not satisfy the condition (2.5).

Example 14. Consider the C^* -algebra valued metric space (X, \mathbb{R}^2, d) given in Example 3, the circle $C_{(\frac{-1}{2}, 0), (1, 1)}^{C^*}$ given in Example 7 and the self-mapping $T : X \rightarrow X$ given in Example 12. Then, T satisfies the condition (2.4) and does not satisfy the condition (2.5). Notice that the circle $C_{(\frac{-1}{2}, 0), (1, 1)}^{C^*}$ is not a fixed circle of T .

Now, we furnish an example in order to evidence that Theorem 6 fail out, if the condition (2.5) is satisfied but the condition (2.4) is not.

Example 15. Consider the C^* -algebra valued metric space $(\mathbb{R}, M_2(\mathbb{R}), d)$ given in Example 4 and the circle $C_{3,r}^{C^*}$ given in Example 8. Let us define the self-mapping $T : \mathbb{R} \rightarrow \mathbb{R}$ as

$$Tx = \begin{cases} 3, & x \in C_{3,r}^{C^*} \\ 0, & x \notin C_{3,r}^{C^*} \end{cases} .$$

It is not hard to prove that T satisfies the condition (2.5) and does not satisfy the condition (2.4). Notice that the circle $C_{3,r}^{C^*}$ is not a fixed circle of T .

The following theorem presents a new solution for fixed-circle problem obtained by the help of the inequality (1.3) in Theorem 3.

Theorem 7. Let (X, \mathbb{A}, d) be a C^* -algebra valued metric space, $C_{x_0,r}^{C^*}$ be any circle on X and the mapping φ be as in (2.1). If T is a self-mapping defined on X satisfying the conditions

$$(2.6) \quad d(x, Tx) \preceq \varphi(x) - \varphi(Tx)$$

and

$$(2.7) \quad r \preceq A^*d(x, Tx)A + d(Tx, x_0)$$

for all $x \in C_{x_0,r}^{C^*}$ and some $A \in \mathbb{A}$ with $\|A\| < 1$, then the circle $C_{x_0,r}^{C^*}$ is a fixed circle of T .

Proof. Let x be any point in the circle $C_{x_0,r}^{C^*}$. Let $x \neq Tx$. Then, using the (2.6), (2.1) and (2.7) we get

$$\begin{aligned} 0 &\preceq d(x, Tx) \preceq \varphi(x) - \varphi(Tx) \\ &= d(x, x_0) - d(Tx, x_0) \\ &= r - d(Tx, x_0) \\ &\preceq A^*d(x, Tx)A + d(Tx, x_0) - d(Tx, x_0) \\ &= A^*d(x, Tx)A \end{aligned}$$

and so

$$\begin{aligned} 0 &\leq \|d(x, Tx)\| \leq \|A^*d(x, Tx)A\| \\ &\leq \|A^*\| \|d(x, Tx)\| \|A\| \\ &= \|A\|^2 \|d(x, Tx)\| \\ &< \|d(x, Tx)\| . \end{aligned}$$

This yields a contradiction with our assumption. Therefore, we conclude that $x = Tx$ for all $x \in C_{x_0,r}^{C^*}$ and hence $C_{x_0,r}^{C^*}$ is a fixed circle of T . \square

The following example support Theorem 7.

Example 16. Consider the C^* -algebra valued metric space $(\mathbb{R}, M_2(\mathbb{R}), d)$ for $\alpha = 2$ and the circle $C_{1,r_2}^{C^*}$ given in Example 9 and the self-mapping $T : \mathbb{R} \rightarrow \mathbb{R}$ given in Example 13. Then, with a direct computation it can be seen that the self-mapping T satisfies the conditions (2.6) and (2.7) for $A = \begin{bmatrix} \frac{1}{5} & 0 \\ 0 & -\frac{3}{5} \end{bmatrix} \in M_2(\mathbb{R})$ with $\|A\| = \frac{3}{5} < 1$. Note that the circle $C_{1,r_2}^{C^*}$ is a fixed circle of T .

Now, we give an example of a self-mapping which satisfies the condition (2.6) and does not satisfy the condition (2.7).

Example 17. Consider the C^* -algebra valued metric space $(\mathbb{R}, M_2(\mathbb{R}), d)$ given in Example 4, the circle $C_{3,r}^{C^*}$ given in Example 8 and the self-mapping $T : \mathbb{R} \rightarrow \mathbb{R}$ given in Example 15. It can be easily seen T satisfies the condition (2.6). But since there does not exist $A \in M_2(\mathbb{R})$ with $\|A\| < 1$ such that

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \preceq A^* \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} A,$$

the condition (2.7) does not hold. Obviously, the circle $C_{3,r}^{C^*}$ is not a fixed circle of T .

Now, we present an example in order to show that whether the Theorem 7 is valid, if the condition (2.7) is satisfied but the condition (2.6) is not.

Example 18. Consider the C^* -algebra valued metric space $(\mathbb{R}, M_2(\mathbb{R}), d)$ given in Example 2 for $\alpha = 3$ and the circle $C_{0,r_1}^{C^*}$ given in Example 6. Let us define the self-mapping $T : \mathbb{R} \rightarrow \mathbb{R}$ as

$$Tx = \begin{cases} 2, & x \in C_{0,r_1}^{C^*} \\ -5, & x \notin C_{0,r_1}^{C^*} \end{cases}.$$

It is not hard to show that T satisfies the condition (2.7) for $A = \begin{bmatrix} \frac{-1}{2} & 0 \\ 0 & \frac{1}{2} \end{bmatrix} \in M_2(\mathbb{R})$ with $\|A\| = \frac{1}{2} < 1$ and does not satisfy the condition (2.6). Notice that the circle $C_{0,r_1}^{C^*}$ is not a fixed circle of T .

We finish this section by give our last existence theorem for fixed circles of a self-mapping on a C^* -algebra valued metric space as follows:

Theorem 8. Let (X, \mathbb{A}, d) be a C^* -algebra valued metric space, $C_{x_0,r}^{C^*}$ be any circle on X and the mapping φ be as in (2.1). T is a self-mapping defined on X satisfying the condition

$$(2.8) \quad A^* d(x, Tx) A \preceq \varphi(x) - \varphi(Tx)$$

for all $x \in X$ where $A \in \mathbb{A}$ is an invertible element and $\|A^{-1}\| < 1$ if and only if T fixes the circle $C_{x_0,r}^{C^*}$ and $T = I_X$.

Proof. Suppose that T be a self-mapping defined on X satisfying the condition (2.8). Let x be any point in X . Let $x \neq Tx$. Then, using the (2.8), (2.1) and (iii)

given in Definition 2 we get

$$\begin{aligned}
A^*d(x, Tx)A &\preceq \varphi(x) - \varphi(Tx) \\
&= d(x, x_0) - d(Tx, x_0) \\
&\preceq d(x, Tx) + d(Tx, x_0) - d(Tx, x_0) \\
&= d(x, Tx)
\end{aligned}$$

and so

$$d(x, Tx) \preceq (A^*)^{-1}d(x, Tx)A^{-1} = (A^{-1})^*d(x, Tx)A^{-1}.$$

It follows that

$$\begin{aligned}
\|d(x, Tx)\| &\leq \left\| (A^{-1})^*d(x, Tx)A^{-1} \right\| \\
&= \left\| (A^{-1})^* \right\| \|d(x, Tx)\| \|A^{-1}\| \\
&= \|A^{-1}\|^2 \|d(x, Tx)\| \\
&< \|d(x, Tx)\|.
\end{aligned}$$

But it is impossible. Hence, we can write $x = Tx$ for all $x \in X$ and $T = I_X$.

Conversely, suppose that T fixes the circle $C_{x_0, r}^{C^*}$ and $T = I_X$. Then, since $Tx = x$ for all $x \in X$, the condition (2.8) holds for any invertible element $A \in \mathbb{A}$ with $\|A^{-1}\| < 1$. This completes the proof. \square

Remark 4. *Theorem 8 says that if a self-mapping fixes a circle by satisfying the conditions (2.2) and (2.3) (or the conditions (2.4) and (2.5)), but does not satisfy the condition (2.8), then the self-mapping is different from the identity map.*

2.2. The uniqueness of fixed circles. In this part, we discuss the uniqueness of fixed circles in the existence theorems proved in subsection 2.1.

Before giving our uniqueness theorems, we emphasize that the fixed circles $C_{x_0, r}^{C^*}$ in Theorem 5, Theorem 6 and Theorem 7 is not unique with the following result.

Proposition 1. *Let (X, \mathbb{A}, d) be a C^* -algebra valued metric space and $C_{x_1, r_1}^{C^*}, C_{x_2, r_2}^{C^*}, \dots, C_{x_n, r_n}^{C^*}$ be any given circles. Then, there exists at least one self-mapping T of X such that T fixes all the circles $C_{x_1, r_1}^{C^*}, C_{x_2, r_2}^{C^*}, \dots, C_{x_n, r_n}^{C^*}$.*

Proof. Let the self-mapping $T : X \rightarrow X$ be defined as

$$Tx = \begin{cases} x, & x \in \bigcup_{i=1}^n C_{x_i, r_i}^{C^*} \\ \alpha, & x \notin \bigcup_{i=1}^n C_{x_i, r_i}^{C^*} \end{cases}$$

where $\alpha \in X$ is a constant satisfying $d(\alpha, x_i) \neq r_i$ and the mapping $\varphi_i : X \rightarrow [0, \infty)$ be defined as

$$\varphi_i(x) = d(x, x_i)$$

for $i = 1, 2, \dots, n$. Then it is not hard to verify that the conditions (2.2) and (2.3) in Theorem 5 are satisfied for the circles $C_{x_1, r_1}^{C^*}, C_{x_2, r_2}^{C^*}, \dots, C_{x_n, r_n}^{C^*}$. Therefore, all the circles $C_{x_1, r_1}^{C^*}, C_{x_2, r_2}^{C^*}, \dots, C_{x_n, r_n}^{C^*}$ are fixed circles of the self-mapping T by Theorem 5. \square

Firstly, we focus on the uniqueness of fixed circles in Theorem 5 using the inequality (1.1) in Theorem 1 in the following theorem.

Theorem 9. *Let (X, \mathbb{A}, d) be a C^* -algebra valued metric space, $C_{x_0, r}^{C^*}$ be any circle on X and the T be a self-mapping satisfying the conditions (2.2) and (2.3) given in Theorem 5. If T satisfies the contraction condition*

$$(2.9) \quad d(Tx, Ty) \preceq A^* d(x, y) A$$

for all $x \in C_{x_0, r}^{C^*}$, $y \in X - C_{x_0, r}^{C^*}$ and some $A \in \mathbb{A}$ with $\|A\| < 1$, then the circle $C_{x_0, r}^{C^*}$ is unique fixed circle of T .

Proof. Assume that $C_{x_1, \delta}^{C^*}$ is another fixed circle of T . Let a and b be any points in $C_{x_0, r}^{C^*}$ and $C_{x_1, \delta}^{C^*}$, respectively. Then, we get by (2.9)

$$d(a, b) = d(Ta, Tb) \preceq A^* d(a, b) A$$

and so,

$$\|d(a, b)\| \leq \|A^* d(a, b) A\| \leq \|A\|^2 \|d(a, b)\| < \|d(a, b)\|.$$

But this is impossible. Hence, the self-mapping T fixes only circle $C_{x_0, r}^{C^*}$. \square

Now, we determine the uniqueness condition for the fixed circles in Theorem 6 using the condition (1.2) in Theorem 2.

Theorem 10. *Let (X, \mathbb{A}, d) be a C^* -algebra valued metric space, $C_{x_0, r}^{C^*}$ be any circle on X and the T be a self-mapping satisfying the conditions (2.4) and (2.5) given in Theorem 6. If T satisfies the contraction condition*

$$(2.10) \quad d(Tx, Ty) \preceq A(d(Tx, y) + d(Ty, x))$$

for all $x \in C_{x_0, r}^{C^*}$, $y \in X - C_{x_0, r}^{C^*}$ and some $A \in \mathbb{A}'_+$ with $\|A\| < \frac{1}{2}$, then the circle $C_{x_0, r}^{C^*}$ is unique fixed circle of T .

Proof. Assume that $C_{x_1, \delta}^{C^*}$ is another fixed circle of T . Let a and b be any points in $C_{x_0, r}^{C^*}$ and $C_{x_1, \delta}^{C^*}$, respectively. Then, we get by (2.10)

$$d(a, b) = d(Ta, Tb) \preceq A(d(Ta, b) + d(Tb, a)),$$

so that

$$\begin{aligned} \|d(a, b)\| &\leq \|A(d(Ta, b) + d(Tb, a))\| \\ &\leq \|A\| \|d(a, b) + d(b, a)\| \\ &= 2 \|A\| \|d(a, b)\| \\ &< \|d(a, b)\| \end{aligned}$$

which is a contradiction which means that $a = b$. This shows that the self-mapping T fixes only circle $C_{x_0, r}^{C^*}$. \square

Finally, we state our last uniqueness theorem for the fixed circles in Theorem 7 using the condition (1.4) in Theorem 4.

Theorem 11. *Let (X, \mathbb{A}, d) be a C^* -algebra valued metric space, $C_{x_0, r}^{C^*}$ be any circle on X and the T be a self-mapping satisfying the conditions (2.6) and (2.7) given in Theorem 7. If T satisfies the contraction condition that there exists $u(x, y) \in \{d(x, y), d(x, Tx), d(y, Ty), d(x, Ty), d(y, Tx)\}$ such that*

$$(2.11) \quad d(Tx, Ty) \preceq A^* u(x, y) A$$

for all $x \in C_{x_0, r}^{C^*}$, $y \in X - C_{x_0, r}^{C^*}$ and some $A \in \mathbb{A}$ with $\|A\| < 1$, then the circle $C_{x_0, r}^{C^*}$ is unique fixed circle of T .

Proof. Assume that $C_{x_1, \delta}^{C^*}$ is another fixed circle of T . Let a and b be any points in $C_{x_0, r}^{C^*}$ and $C_{x_1, \delta}^{C^*}$, respectively. Then, we get by (2.11)

$$d(a, b) = d(Ta, Tb) \preceq A^*u(a, b)A,$$

so that

$$\begin{aligned} \|d(a, b)\| &\leq \|A^*u(a, b)A\| \\ &\leq \|A\|^2 \|u(a, b)\| \\ &< \|u(a, b)\| \\ &\leq \max\{\|d(a, b)\|, \|d(a, Tb)\|, \|d(b, Tb)\|, \|d(a, Ta)\|, \|d(b, Ta)\|\} \\ &= \|A\|^2 \max\{\|d(a, b)\|, 0\} \\ &= \|d(a, b)\| \end{aligned}$$

which is a contradiction. Hence $a = b$. This implies that the self-mapping T fixes only circle $C_{x_0, r}^{C^*}$. \square

Remark 5. 1) *The uniqueness result in Theorem 5 can be also stated using the contraction conditions (2.10) given in Theorem 10 or (2.11) given in Theorem 11 instead of the contraction condition (2.9).*

2) *The uniqueness result in Theorem 6 can be also stated using the contraction conditions (2.9) given in Theorem 9 or (2.11) given in Theorem 11 instead of the contraction condition (2.10).*

3) *The uniqueness result in Theorem 7 can be also stated using the contraction conditions (2.9) given in Theorem 9 or (2.10) given in Theorem 10 instead of the contraction condition (2.11).*

3. CONCLUSION AND FUTURE WORKS

In this article, we discuss the existence and uniqueness of the fixed-circle for self-mappings satisfying some special conditions on C^* -algebra valued metric spaces. We also furnish some examples to show effectiveness of our theoretical results. Using similar approaches and new contractive conditions, it can be studied new fixed-circle results on C^* -algebra valued metric spaces. Since with this article we develop a new and different perspective instead of the classical fixed point theory in C^* -algebra valued metric spaces, we hope that our results will support researchers for future works and applications to other related areas.

REFERENCES

- [1] Banach, S. (1922). Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales. *Fund. Math.*, 3(1), 133-181.
- [2] Ma, Z., Jiang, L. and Sun, H. (2014). C^* -algebra valued metric spaces and related fixed point theorems. *Fixed Point Theory and Applications*, 2014:206.
- [3] Batul, S. and Kamran, T. (2015). C^* -valued contractive type mappings. *Fixed Point Theory and Applications*, 2015:142.
- [4] Dur-e-Shehwar, Batul, S., Kamran, T. and Ghiura, A. (2016). Caristi's fixed point theorem on C^* -algebra valued metric spaces. *Journal of Nonlinear Sciences and Applications*, 9(2), 584-588.
- [5] Kadelburg, Z. and Radenović, S. (2016). Fixed point results in C^* -algebra valued metric spaces are direct consequences of their standard metric counterparts. *Fixed Point Theory and Applications*, 2016:53.

- [6] Banerjee, A. K. and Paul, A. (2018). On I and I^* -Cauchy conditions in C^* -algebra valued metric spaces. arXiv:1811.05662.
- [7] Moeini, B., Kumar, P. and Aydi, H. (2018). Zamfirescu type contractions on C^* -algebra valued metric spaces. J. Math. Anal, 9(1), 150-161.
- [8] Kumar, M., Imdad, M. and Asim, M. (2020). Some fixed point theorems under E.A. and (CLR) properties on C^* -algebra valued metric spaces. Information Sciences Letters, 9(2), 75-82.
- [9] Asim, M., George, R. and Imdad, M. (2021). Suzuki type multivalued contractions in C^* -algebra valued metric spaces with an application [J]. AIMS Mathematics, 6(2), 1126-1139.
- [10] Özgür, N. Y. and Taş, N. (2019). Some fixed-circle theorems on metric spaces. Bulletin of the Malaysian Mathematical Sciences Society, 42(4), 1433-1449.
- [11] Özgür, N. Y. and Taş, N. (2017). Fixed-circle problem on S -metric spaces with a geometric viewpoint. arXiv:1704.08838.
- [12] Mlaiki, N., Taş, N. and Özgür, N. Y. (2018). On the fixed-circle problem and Khan type contractions. Axioms, 7(4), 80.
- [13] Taş, N., Özgür, N. Y. and Mlaiki, N. (2018). New types of F_C -contractions and the fixed-circle problem. Mathematics, 6(10), 188.
- [14] Tas, N. and Özgür, N. Y. (2018). Some fixed-point results on parametric N_b -metric spaces. Communications of the Korean Mathematical Society, 33(3), 943-960.
- [15] Aydi, H., Taş, N., Özgür, N. Y. and Mlaiki, N. (2019). Fixed-discs in rectangular metric spaces. Symmetry, 11(2), 294.
- [16] Taş, N. (2020). Bilateral-type solutions to the fixed-circle problem with rectified linear units application. Turk. J. Math, 44(4), 1330-1344.
- [17] Murphy, G. J. (2014). C^* -algebras and operator theory. Academic press, Boston.
- [18] Caristi, J. (1976). Fixed point theorems for mappings satisfying inwardness conditions. Transactions of the American Mathematical Society, 215, 241-251.
- [19] Khamsi, M. A. and Kirk, W. A. (2001). An introduction to metric spaces and fixed point theory. John Wiley & Sons, Inc., New York.
- [20] Ćirić, L. B. (1974). A generalization of Banach's contraction principle. Proceedings of the American Mathematical society, 45(2), 267-273.

(N. Değirmen) ONDOKUZ MAYIS UNIVERSITY FACULTY OF ART AND SCIENCES DEPARTMENT OF MATHEMATICS SAMSUN, TURKEY

Email address, N. Değirmen: nilay.sager@omu.edu.tr