

ON A CONJECTURE OF IIZUKA

AZIZUL HOQUE

ABSTRACT. For a given odd positive integer n and an odd prime p , we construct an infinite family of quadruples of imaginary quadratic fields $\mathbb{Q}(\sqrt{d})$, $\mathbb{Q}(\sqrt{d+1})$, $\mathbb{Q}(\sqrt{d+4})$ and $\mathbb{Q}(\sqrt{d+4p^2})$ with $d \in \mathbb{Z}$ such that the class number of each of them is divisible by n . Subsequently, we show that there is an infinite family of quintuples of imaginary quadratic fields $\mathbb{Q}(\sqrt{d})$, $\mathbb{Q}(\sqrt{d+1})$, $\mathbb{Q}(\sqrt{d+4})$, $\mathbb{Q}(\sqrt{d+36})$ and $\mathbb{Q}(\sqrt{d+100})$ with $d \in \mathbb{Z}$ whose class numbers are all divisible by n . Our results provide a complete proof of Iizuka's conjecture (in fact a generalization of it) for the case $m = 1$. Our results also affirmatively answer a weaker version of (a generalization of) Iizuka's conjecture for $m \geq 4$.

1. INTRODUCTION

It has been proved that there are infinitely many real (resp. imaginary) quadratic fields with class numbers divisible by a given positive integer (see [1, 3, 21, 24]). An analogous problem for tuples of quadratic fields arises from Scholz's Spiegelungssatz [19]. In [13], Komatsu studied this problem for a pair of quadratic fields and proved that there are infinitely many pairs of quadratic fields $\mathbb{Q}(\sqrt{d})$ and $\mathbb{Q}(\sqrt{md})$ with $m, d \in \mathbb{Z}$ whose class numbers are divisible by 3. Later, he generalized this result in [14] to n -divisibility of the class numbers of pairs of imaginary quadratic fields. On the other hand, Iizuka [11] studied a slight variant of this problem and construct an infinite family of pairs of imaginary quadratic fields $\mathbb{Q}(\sqrt{d})$ and $\mathbb{Q}(\sqrt{d+1})$ with $d \in \mathbb{Z}$ whose class numbers are divisible by 3. Further, he posed the following conjecture in the same paper.

Conjecture 1.1. *For any prime number p and any positive integer m , there is an infinite family of $m + 1$ successive real (or imaginary) quadratic fields*

$$\mathbb{Q}(\sqrt{d}), \mathbb{Q}(\sqrt{d+1}), \dots, \mathbb{Q}(\sqrt{d+m})$$

with $d \in \mathbb{Z}$ whose class numbers are divisible by p .

Date: August 18, 2021.

2020 Mathematics Subject Classification. 11R11; 11R29.

Key words and phrases. Imaginary quadratic field, Class number, Iizuka's conjecture, Exponent.

In [4], Chattopadhyay and Muthukrishnan extended Iizuka's result from pairs to certain triples of imaginary quadratic fields following the methods used in [11]. In other words, they gave an affirmative answer of a weaker version of Conjecture 1.1 for $p = 3$. It follows from a recent result of Iizuka, Konomi and Nakano [12] that Conjecture 1.1 is true for $m = 1$ when $p = 3, 5, 7$.

Very recently, Krishnamoorthy and Pasupulati [15] cleverly used [16, Theorem 1] and an extended version of [9, Theorem 3.2] to settled Conjecture 1.1 for $m = 1$.

In this paper, for a given odd prime p we construct an infinite family of quadruples of imaginary quadratic fields $\mathbb{Q}(\sqrt{d})$, $\mathbb{Q}(\sqrt{d+1})$, $\mathbb{Q}(\sqrt{d+4})$ and $\mathbb{Q}(\sqrt{d+4p^2})$ with $d \in \mathbb{Z}$ whose class numbers are all divisible by a given odd integer $n \geq 3$. This extends the results of [4, 12, 15] in both directions; from pairs to quadruples/quintuples of fields and from primes to odd integers. It also gives a proof of a weaker version of Conjecture 1.1 for any prime $p \geq 3$ (in fact for any odd integer $n \geq 3$). The precise statement of our first result is the following:

Theorem 1.1. *For any odd positive integer n and any odd prime p , there are infinitely many quadruples of imaginary quadratic fields $\mathbb{Q}(\sqrt{d})$, $\mathbb{Q}(\sqrt{d+1})$, $\mathbb{Q}(\sqrt{d+4})$ and $\mathbb{Q}(\sqrt{d+4p^2})$ whose class numbers are all divisible by n .*

Note that we can construct an infinite family of quintuples or higher tuples of imaginary quadratic fields whose class numbers are all divisible by n by choosing different values for the prime p . For instance, utilizing Corollary 3.1 and following the proof of Theorem 1.1, we get the following:

Theorem 1.2. *For a given odd integer $n \geq 1$, there are infinitely many quintuples of imaginary quadratic fields*

$$\left(\mathbb{Q}(\sqrt{d}), \mathbb{Q}(\sqrt{d+1}), \mathbb{Q}(\sqrt{d+4}), \mathbb{Q}(\sqrt{d+36}), \mathbb{Q}(\sqrt{d+100}) \right)$$

with $d \in \mathbb{Z}$ whose class numbers are all divisible by n .

Our method relies on the prominent result of Bilu, Hanrot and Voutier [2] concerning the primitive divisors of Lehmer numbers as well as on the solvability of certain Lebesgue-Ramanujan-Nagell type equations considered in [8]. This method does not allow us to include an imaginary quadratic field of the form $\mathbb{Q}(\sqrt{d+m})$ in the tuple, when m is non-square integer.

2. n -DIVISIBILITY OF THE CLASS-NUMBERS OF $\mathbb{Q}(\sqrt{1-4U^n})$ AND $\mathbb{Q}(\sqrt{1-V^n})$

Here, we recall some results concerning the n -divisibility of class numbers of the imaginary quadratic fields $\mathbb{Q}(\sqrt{1-4U^n})$ and $\mathbb{Q}(\sqrt{1-V^n})$. These results will be used in the proof of Theorem 1.1.

Theorem A. *If $n \geq 3$ is an odd integer, then for any integer $U \geq 2$ the ideal class groups of the imaginary quadratic fields $\mathbb{Q}(\sqrt{1-4U^n})$ contain an element of order n .*

In 1978, Gross and Rohrlich gave the outline of a proof of this theorem (see [6, Theorem 5.3 and Remark 2]). Their method of proof was based upon the affine points on the Fermat curve $x^p + y^p = 1$ over the imaginary quadratic field $\mathbb{Q}(\sqrt{1-4U^n})$. Later, Louboutin [16] gave a complete proof of this theorem using number theoretic technique. It follows from Siegel's theorem (see [20]) that for each integer $d > 1$ there are at most finitely many positive integers U such that $1 - 4U^n = -dX^2$. This ensures the infinitude of the above family of imaginary quadratic fields.

The n -divisibility of the class numbers of the family of imaginary quadratic fields $\mathbb{Q}(\sqrt{1-V^n})$ was studied by Nagell [18, Theorem 25] for any odd integers $V \geq 3$ and $n \geq 3$. Later, Murty [17, Theorems 1 and 2] proved that the class group of the imaginary quadratic field $\mathbb{Q}(\sqrt{1-V^n})$ has an element of order n when either $V^n - 1$ is square-free or its square part is $< V^{n/2}/8$ for any odd integers $V \geq 5$ and $n \geq 3$. However, it follows from the fact $DX^2 + 1 = V^n$ has no integer solution when both V and n are odd, except for $(V, n) = (5, 3), (7, 3), (13, 3)$ (see [7]), that the above conditions are no longer required. This fact also confirms that there are infinitely many such imaginary quadratic fields. Finally, this was elucidated by Cohn in [5, Corollary 1] as follows:

Theorem B. *Assume that $n \geq 3$ and $V \geq 3$ are odd integers. Then the class number of the imaginary quadratic field $\mathbb{Q}(\sqrt{1-V^n})$ is divisible by n , except for $(n, V) = (5, 3)$.*

3. THE DIVISIBILITY OF THE CLASS NUMBER OF $\mathbb{Q}(\sqrt{p^2 - \ell^n})$

Many special cases of the divisibility of the class number of the imaginary quadratic field $\mathbb{Q}(\sqrt{x^2 - y^n})$ have been studied with some restrictions on x, y and n . One of such restrictions is that y is an odd prime (see [3] and the references therein), and hence none of the known results can be used to complete the proof of Theorem (1.1). Thus, we consider a family of imaginary quadratic fields of the above form where y is not a prime and it will be a useful ingredient in the proof of Theorem 1.1. Here, we mainly prove:

Theorem 3.1. *Let $\ell > 1$ and $n > 1$ be odd integers, and p an odd prime such that $\ell \equiv 3 \pmod{4}$, $\gcd(\ell, p) = 1$ and $p^2 < \ell^n$. Assume that $-d$ is the square-free part of $p^2 - \ell^n$. If $p \not\equiv \pm 1 \pmod{d}$, then the class number of $\mathbb{Q}(\sqrt{p^2 - \ell^n})$ is divisible by n .*

Theorem 3.1 extends [3, Theorem 1.1], where the authors assumed that ℓ is an odd prime. This primality condition on ℓ restricts us to apply [3, Theorem 1.1] in the proof of Theorem 1.1. Further, following the proof of [3, Theorem 1.2], we make the following remark.

Remark 3.1. *The family of imaginary quadratic fields discussed in Theorem 3.1 has infinitely many members.*

Now for $p = 3, 5$, the condition ' $p \not\equiv \pm 1 \pmod{d}$ ' can be removed by applying 3.7 except for $(\ell, n) = (3, 3)$, and thus we have the following straightforward corollary.

Corollary 3.1. *Let $\ell > 1$ and $n > 1$ be as in Theorem 3.1. For $x = 3, 5$ with $\gcd(\ell, x) = 1$, the class number of $\mathbb{Q}(\sqrt{x^2 - \ell^n})$ is divisible by n except the case $(\ell, n) = (3, 3)$.*

The proof of Theorem 3.1 relies on the prominent result of Bilu, Hanrot and Voutier [2] on existence of primitive divisors of Lehmer numbers.

3.1. Lehmer numbers and their primitive divisors. A pair (α, β) of algebraic integers is said to be a Lehmer pair if $(\alpha + \beta)^2$ and $\alpha\beta$ are two non-zero coprime rational integers, and α/β is not a root of unity. For a given positive integer n , the Lehmer numbers correspond to the pair (α, β) are defined as

$$\mathcal{L}_n(\alpha, \beta) = \begin{cases} \frac{\alpha^n - \beta^n}{\alpha - \beta}, & \text{if } n \text{ is odd,} \\ \frac{\alpha^n - \beta^n}{\alpha^2 - \beta^2}, & \text{if } n \text{ is even.} \end{cases}$$

It is known that all Lehmer numbers are non-zero rational integers. Two Lehmer pairs (α_1, β_1) and (α_2, β_2) are said to be equivalent if $\alpha_1/\alpha_2 = \beta_1/\beta_2 \in \{\pm 1, \pm\sqrt{-1}\}$. A prime divisor p of $\mathcal{L}_n(\alpha, \beta)$ is said to be primitive if $p \nmid (\alpha^2 - \beta^2)^2 \mathcal{L}_1(\alpha, \beta) \mathcal{L}_2(\alpha, \beta) \cdots \mathcal{L}_{n-1}(\alpha, \beta)$. The following classical result was proved in [2, Theorem 1.4].

Theorem C. *The Lehmer number $\mathcal{L}_n(\alpha, \beta)$ has primitive divisors for any integer $n > 30$.*

Given a Lehmer pair (α, β) , let $a = (\alpha + \beta)^2$ and $b = (\alpha - \beta)^2$. Then $\alpha = (\sqrt{a} \pm \sqrt{b})/2$ and $\beta = (\sqrt{a} \mp \sqrt{b})/2$. The pair (a, b) is called the parameters corresponding to the Lehmer pair (α, β) . The following lemma is extracted from [22, Theorem 1].

Lemma 3.1. *Let t be an odd integer such that $7 \leq t \leq 29$. If the Lehmer numbers $\mathcal{L}_t(\alpha, \beta)$ have no primitive divisor, then up to equivalence, the parameters (a, b) of the corresponding pair (α, β) are as follows:*

- (i) $(a, b) = (1, -7), (1, -19), (3, -5), (5, -7), (13, -3), (14, -22)$, when $t = 7$;
- (ii) $(a, b) = (5, -3), (7, -1), (7, -5)$, when $t = 9$;
- (iii) $(a, b) = (1, -7)$, when $t = 13$;
- (iv) $(a, b) = (7, -1), (10, -2)$, when $t = 15$.

Let F_k (resp. L_k) denote the k -th term in the Fibonacci (resp. Lucas) sequence defined by $F_0 = 0, F_1 = 1$, and $F_{k+2} = F_k + F_{k+1}$ (resp. $L_0 = 2, L_1 = 1$, and $L_{k+2} = L_k + L_{k+1}$), where $k \geq 0$ is an integer. The following lemma is a part of [2, Theorem 1.3].

Lemma 3.2. *For $p = 3, 5$, let the Lehmer numbers $\mathcal{L}_p(\alpha, \beta)$ have no primitive divisor. Then up to equivalence, the parameters (a, b) of the corresponding pair (α, β) are:*

- (i) For $p = 3$, $(a, b) = \begin{cases} (1+t, 1-3t) & \text{with } t \neq 1, \\ (3^k+t, 3^k-3t) & \text{with } t \not\equiv 0 \pmod{3}, (k, t) \neq (1, 1); \end{cases}$
- (ii) For $p = 5$, $(a, b) = \begin{cases} (F_{k-2\varepsilon}, F_{k-2\varepsilon} - 4F_k) & \text{with } k \geq 3, \\ (L_{k-2\varepsilon}, L_{k-2\varepsilon} - 4L_k) & \text{with } k \neq 1; \end{cases}$

where $t \neq 0$ and $k \geq 0$ are any integers and $\varepsilon = \pm 1$.

3.2. Two important lemmas. Given an integer $D \equiv 0, 1 \pmod{4}$, assume that $h^*(D)$ is the class number of binary quadratic primitive forms with discriminant D . Also for a square-free integer d , let $h(d)$ denote the class number of $\mathbb{Q}(\sqrt{d})$. Then we have (cf. [10, §16.13; p. 444] the following:

Lemma 3.3. *Let $d \equiv 2 \pmod{4}$ be a square-free positive integer. Then $h(-d) = h^*(-4d)$.*

The following lemma is a special case of [8, Theorem 6.2] when $(D_1, D_2) = (1, -d)$.

Lemma 3.4. *Let $d > 3$ and $\ell > 1$ be integers such that $\gcd(\ell, 2d) = 1$. If the equation*

$$x^2 + dy^2 = \ell^z, \quad x, y, z \in \mathbb{N}, \quad \gcd(x, y) = 1$$

has a solution, then all the solutions (x, y, z) of this equation can be expressed as

$$x + y\sqrt{-d} = \varepsilon(a + \mu b\sqrt{-d})^t, \quad z = st,$$

where a, b, s, t are positive integers satisfying

$$a^2 + db^2 = \ell^s, \quad \gcd(a, b) = 1 \quad \text{and} \quad s \mid h^*(-4d)$$

and $\varepsilon, \mu \in \{-1, 1\}$.

3.3. Proof of Theorem 3.1. Let d be the square-free part of $\ell^n - p^2$. Then $p^2 - \ell^n = -dr^2$ for some $r \in \mathbb{N}$, and thus $(x, y, z) = (p, r, n)$ is a positive integer solution of the equation

$$x^2 + dy^2 = \ell^z, \quad \gcd(x, y) = 1.$$

Thus by Lemma 3.4, we get

$$p + r\sqrt{-d} = \varepsilon(a + \mu b\sqrt{-d})^t, \quad \varepsilon, \mu \in \{-1, 1\} \quad (3.1)$$

with

$$n = st, \quad s, t \in \mathbb{N}, \quad (3.2)$$

where a and b are positive integers satisfying

$$a^2 + db^2 = \ell^s, \quad \gcd(a, b) = 1 \quad (3.3)$$

and

$$s \mid h^*(-4d). \quad (3.4)$$

Since $\ell \equiv 3 \pmod{4}$ and n is odd, so that $p^2 - \ell^n = -dr^2$ gives $d \equiv 2 \pmod{4}$ and r is odd. Also both s and t are odd as n is odd. Further reading (3.3) modulo 4, we get $a^2 + 2b^2 \equiv 3 \pmod{4}$ as $\ell \equiv 3 \pmod{4}$ and s is odd, which ensures that both a and b are odd.

We now equate the real parts from both sides in (3.1) to get

$$p = \varepsilon a \sum_{j=0}^{\frac{t-1}{2}} \binom{t}{2j} a^{t-2j-1} (-db^2)^j. \quad (3.5)$$

This implies $a \mid p$ and thus $a = 1, p$. If $a = 1$, then it becomes

$$\sum_{j=0}^{\frac{t-1}{2}} \binom{t}{2j} (-db^2)^j = p\varepsilon = \pm p. \quad (3.6)$$

Reading (3.6) modulo d , we get $p \equiv \pm 1 \pmod{d}$, which contradicts to the assumption. Therefore $a = p$ and thus (3.5) becomes

$$\sum_{j=0}^{\frac{t-1}{2}} \binom{t}{2j} p^{t-2j-1} (-db^2)^j = \varepsilon = \pm 1. \quad (3.7)$$

As $a = p$, so that (3.1) reduces to

$$p + r\sqrt{-d} = \varepsilon(p + \mu b\sqrt{-d})^t, \quad \varepsilon, \mu \in \{-1, 1\}. \quad (3.8)$$

We now assume that $\alpha = \mu b\sqrt{-d} + p$ and $\beta = \mu b\sqrt{-d} - p$. Then both α and β are algebraic integers. Clearly, $(\alpha + \beta)^2 = -4db^2$ and $\alpha\beta = -p^2 - db^2 = -\ell^s$ (by (3.3)) are coprime rational integers. Furthermore, it follows from the following identity

$$\frac{4db^2}{\ell^s} = \frac{(\alpha + \beta)^2}{\alpha\beta} = \frac{\alpha}{\beta} + \frac{\beta}{\alpha} + 2$$

that

$$\ell^s \left(\frac{\alpha}{\beta} \right)^2 + 2(\ell^s - 2db^2) \frac{\alpha}{\beta} + \ell^s = 0.$$

Since $\ell > 1$ and $\gcd(\ell^s, 2(\ell^s - 2db^2)) = \gcd(\ell^s, 4db^2) = \gcd(p^2 + db^2, 4db^2) = 1$, so that the last equation shows that $\frac{\alpha}{\beta}$ is not a root of unity. Therefore (α, β) is a Lehmer pair with parameters $(-4db^2, 4p^2)$ and thus the corresponding Lehmer number for t is

$$\mathcal{L}_t(\alpha, \beta) = \frac{\alpha^t - \beta^t}{\alpha - \beta}$$

as t is odd. Utilizing (3.8), we get

$$|\mathcal{L}_t(\alpha, \beta)| = 1.$$

This confirms that the Lehmer number $\mathcal{L}_t(\alpha, \beta)$ has no primitive divisor, and hence Theorem C and Lemma 3.1 (utilizing the fact that $(-4db^2, 4p^2)$ is the parameters) ensure that $t \in \{1, 3, 5\}$.

In case of $t = 5$, we get by Lemma 3.2 that $-4db^2 = F_{k-2\varepsilon}$ or $-4db^2 = L_{k-2\varepsilon}$. Clearly, none of these is possible.

Finally for $t = 3$, (3.7) implies that $p^2 - 3db^2 = \pm 1$. Reading it modulo 4, we see that '+' sign is not possible, and thus $p^2 - 3db^2 = -1$. This is not possible by reading it modulo 3.

Therefore $t = 1$, and thus (3.2) and (3.4) together imply that $n \mid h^*(-4d)$. Thus, we complete the proof by Lemma 3.3.

4. PROOF OF THEOREM 1.1

We first fix an odd integer $n \geq 3$. We now define the set

$$\mathcal{N}_n = \{k \in \mathbb{N} : n \mid h(1 - 4k^n)\}.$$

Then by Theorem A the set \mathcal{N}_n is infinite.

Now for any $k \in \mathcal{N}_n$, we set $d = 4(1 - 4k^n)^n$. Then $\mathbb{Q}(\sqrt{d}) = \mathbb{Q}(\sqrt{1 - 4k^n})$ as n is odd. Thus, by Theorem A there are infinitely many such d satisfying $n \mid h(d)$. In other words, $\mathcal{F}(\mathcal{N}_n) = \{\mathbb{Q}(\sqrt{1 - 4k^n}) : k \in \mathcal{N}_n\}$ is an infinite set.

Now we assume that $U = 4k^n - 1$ with $k \in \mathcal{N}_n$. Then $1 - 4U^n = 1 - 4(4k^n - 1)^n = 4(1 - 4k^n)^n + 1 = d + 1$, and thus by Theorem A, we have $n \mid h(d + 1)$.

Again for $k \in \mathcal{N}_n$, let us assume that $V = 4k^n - 1$. Then $V \geq 3$ and is odd, and thus by Theorem B, we get $n \mid h(1 - V^n)$. Since $4(1 - V^n) = 4 - 4(4k^n - 1)^n = 4 + 4(1 - 4k^n)^n = d + 4$ and $\mathbb{Q}(\sqrt{4(1 - V^n)}) = \mathbb{Q}(\sqrt{(1 - V^n)})$, so that $n \mid h(d + 4)$.

Finally for any $k \in \mathcal{N}_n$, let $\ell = 4k^n - 1$. Then $\ell \equiv 3 \pmod{4}$ and hence by utilizing Theorem 3.1, we have $n \mid h(p^2 - \ell^n)$ for any odd prime p satisfying $p \not\equiv \pm 1 \pmod{d}$. Here, d is the square-free part of $\ell^n - p^2$. Now $4(p^2 - \ell^n) = 4p^2 - 4\ell^n = 4p^2 - 4(4k^n - 1)^n = d + 4p^2$, which implies that $\mathbb{Q}(\sqrt{d + 4p^2}) = \mathbb{Q}(\sqrt{4(p^2 - \ell^n)})$, and thus $n \mid h(d + 4p^2)$. This completes the proof of Theorem 1.1.

5. CONCLUDING REMARKS

In [23], Xie and Chao studied Conjecture 1.1 and proved the following result using Yamamoto's [24] construction.

Theorem D. *For any odd positive integer n and any positive integer m , there are infinitely many pairs of imaginary fields $\mathbb{Q}(\sqrt{d})$ and $\mathbb{Q}(\sqrt{d + m})$ whose class groups have an element of order n respectively.*

Theorem D can be viewed as a weaker variant of a generalization of Conjecture 1.1. For $m = 1$, it provides a generalization of the main result of [15] though [23] appeared before [15]. In other words, Theorem D gives a complete proof of the following generalization of Conjecture 1.1 for $m = 1$ and a proof of a weaker version of the same for $m \geq 2$.

Conjecture 5.1. *For any odd integer $n \geq 3$ and any integer $m \geq 1$, there is an infinite family of $m + 1$ successive imaginary (or real) quadratic fields*

$$\mathbb{Q}(\sqrt{d}), \mathbb{Q}(\sqrt{d + 1}), \dots, \mathbb{Q}(\sqrt{d + m})$$

with $d \in \mathbb{Z}$ whose class numbers are all divisible by n .

Theorem 1.1 offers a constructive proof of Conjecture 5.1 for $m = 1$. This theorem also offers a proof of a weaker version of Conjecture 5.1 for $m = 4$, which has missed the families of imaginary quadratic fields $\mathbb{Q}(\sqrt{d + 2})$ and $\mathbb{Q}(\sqrt{d + 3})$ from the complete proof. When $m = 4p^2$ with p an odd prime, Theorem 1.1 presents a proof of a weaker version of Conjecture 5.1. We complete this paper by the following remark.

Remark 5.1. *For a given positive integer m , let p_m denote the largest prime less than or equal to m and $\pi(m)$ the prime-counting function. Then for a given positive odd integer*

n , our construction gives an infinite family of at least $(\pi(m) + 2)$ -tuples of imaginary quadratic fields,

$$\left(\mathbb{Q}(\sqrt{d}), \mathbb{Q}(\sqrt{d+1}), \mathbb{Q}(\sqrt{d+4}), \mathbb{Q}(\sqrt{d+36}), \dots, \mathbb{Q}(\sqrt{d+4p_m^2}) \right)$$

with $d \in \mathbb{Z}$ whose class numbers are all divisible by n .

ACKNOWLEDGEMENTS

The author is grateful to Professor K. Chakraborty and Professor Yasuhiro Kishi for their valuable comments on the paper. The author is thankful to Professor Y. Iizuka for providing a copy of [12]. The author gratefully acknowledges the anonymous referee for his/her valuable remarks that immensely improved the results as well as the presentation of the paper. This work was supported by SERB-NPDF (PDF/2017/001958), Govt. of India.

REFERENCES

- [1] N. C. Ankeny and S. Chowla, *On the divisibility of the class number of quadratic fields*, Pacific J. Math. **5** (1955), 321–324.
- [2] Y. Bilu, G. Hanrot and P. M. Voutier, *Existence of primitive divisors of Lucas and Lehmer numbers (with an appendix by M. Mignotte)*, J. Reine Angew. Math. **539** (2001), 75–122.
- [3] K. Chakraborty, A. Hoque, Y. Kishi and P. P. Pandey, *Divisibility of the class numbers of imaginary quadratic fields*, J. Number Theory **185** (2018), 339–348.
- [4] J. Chattopadhyay and S. Muthukrishnan, *On the simultaneous 3-divisibility of class numbers of triples of imaginary quadratic fields*, Acta Arith. **197** (2021), no. 1, 105–110.
- [5] J. H. E. Cohn, *On the Diophantine equation $x^n = Dy^2 + 1$* , Acta Arith. **106** (2003), no. 1, 73–83.
- [6] B. H. Gross and D. E. Rohrlich, *Some results on the Mordell-Weil group of the Jacobian of the Fermat curve*, Invent. Math. **44** (1978), 201–224.
- [7] E. Herrmann, I. Járási and A. Pethő, *Note on J. H. E. Cohn’s paper “The Diophantine equation $x^n = Dy^2 + 1$ ”*, Acta Arith. **113** (2004), no. 1, 69–76.
- [8] C. Heuberger and M. Le, *On the generalized Ramanujan-Nagell equation $x^2 + D = p^Z$* , J. Number Theory **78** (1999), no. 2, 312–331.
- [9] A. Hoque and K. Chakraborty, *Divisibility of class numbers of certain families of quadratic fields*, J. Ramanujan Math. Soc. **34** (2019), no. 3, 281–289.
- [10] L. K. Hua, *Introduction to number theory*, Springer-Verlag, New York, 1982.
- [11] Y. Iizuka, *On the class number divisibility of pairs of imaginary quadratic fields*, J. Number Theory **184** (2018), 122–127.
- [12] Y. Iizuka, Y. Konomi and S. Nakano, *An application of the arithmetic of elliptic curves to the class number problem for quadratic fields*, Tokyo J. Math. (2021). doi: 10.3836/tjm/1502179314

- [13] T. Komatsu, *An infinite family of pairs of quadratic fields $\mathbb{Q}(\sqrt{D})$ and $\mathbb{Q}(\sqrt{mD})$ whose class numbers are both divisible by 3*, Acta Arith. **104** (2002), 129–136.
- [14] T. Komatsu, *An infinite family of pairs of imaginary quadratic fields with ideal classes of a given order*, Int. J. Number Theory **13** (2017), no. 2, 253–260.
- [15] S. Krishnamoorthy and S. Pasupulati, *Note on the p -divisibility of class numbers of an infinite family of imaginary quadratic fields*, Glasgow Math. J. (2021). doi: 10.1017/S001708952100015X
- [16] S. R. Louboutin, *On the divisibility of the class number of imaginary quadratic number fields*, Proc. Amer. Math. Soc. **137** (2009), no. 12, 4025–4028.
- [17] M. R. Murty, *The ABC conjecture and exponents of class groups of quadratic fields*, Contemp. Math. **210** (1998), 85–95.
- [18] T. Nagell, *Über die Klassenzahl imaginär quadratischer, Zahlkörper*, Abh. Math. Sem. Univ. Hambg. **1** (1922), 140–150.
- [19] A. Scholz, *Über die Beziehung der Klassenzahlen quadratischer Körper zueinander*, J. Reine Angew. Math. **166** (1932), 201–203.
- [20] C. L. Siegel, *Über einige Anwendungen Diophantischer Approximationen*, Abh. Preuss. Akad. Wiss. Phys. Math. Kl. **1** (1929), 1-70; Ges. Abh., Band **1**, 209–266.
- [21] K. Soundararajan, *Divisibility of class numbers of imaginary quadratic fields*, J. London Math. Soc. **61** (2000), 681–690.
- [22] P. M. Voutier, *Primitive divisors of Lucas and Lehmer sequences*, Math. Comp. **64** (1995), 869–888.
- [23] C. -F. Xie and C. F. Chao, *On the divisibility of class numbers of imaginary quadratic fields $(\mathbb{Q}(\sqrt{D}), \mathbb{Q}(\sqrt{D+m}))$* , Ramanujan J. **53** (2020), 517–528.
- [24] Y. Yamamoto, *On unramified Galois extensions of quadratic number fields*, Osaka J. Math. **7** (1970), 57–76.

Email address: ahoque.ms@gmail.com