

ZETA AND FREDHOLM DETERMINANTS OF SELF-ADJOINT OPERATORS

LUIZ HARTMANN AND MATTHIAS LESCH

ABSTRACT. Let L be a self-adjoint invertible operator in a Hilbert space such that L^{-1} is p -summable. Under a certain discrete dimension spectrum assumption on L , we study the relation between the (regularized) Fredholm determinant, $\det_p(I+z \cdot L^{-1})$, on the one hand and the zeta regularized determinant, $\det_\zeta(L+z)$, on the other. One of the main results is the formula

$$\frac{\det_\zeta(L+z)}{\det_\zeta(L)} = \exp\left(\sum_{j=1}^{p-1} \frac{z^j}{j!} \cdot \frac{d^j}{dz^j} \log \det_\zeta(L+z)|_{z=0}\right) \cdot \det_p(I+z \cdot L^{-1}).$$

We show that the derivatives $\frac{d^j}{dz^j} \log \det_\zeta(L+z)|_{z=0}$ can be expressed in terms of (regularized) zeta values and heat trace coefficients of L . Furthermore, we give a general criterion in terms of the heat trace coefficients (and which is, *e.g.*, fulfilled for large classes of elliptic operators) which guarantees that the constant term in the asymptotic expansion of the Fredholm determinant, $\log \det_p(I+z \cdot L^{-1})$, equals the zeta determinant of L .

CONTENTS

1. Introduction	1
2. Generalities	4
3. Operators with meromorphic ζ -function	8
4. The Fredholm determinant and the ζ -determinant	18
References	22

1. INTRODUCTION

The classical theory of trace ideals (see [Sim05]) allows to generalize the notion of determinant to operators which differ from the identity by a p -summable operator. More concretely, let first A be a trace class operator in a separable Hilbert

Date: February 28, 2022.

2020 Mathematics Subject Classification. Primary 58J52 ; Secondary 47B10.

Key words and phrases. Discrete dimension spectrum, Zeta-determinant, Fredholm determinant.

Partially support by FAPESP (2018/23202-1, 2021/09534-4) and we gratefully acknowledge the financial support of the Hausdorff Center for Mathematics, Bonn.

space. Then

$$\det(I + z \cdot A) = \sum_{k=0}^{\infty} \operatorname{Tr}(\Lambda^k A) \cdot z^k = \prod_{n=1}^{\infty} (1 + z \cdot \lambda_n(A))$$

is an entire function of z with zeros exactly in $-1/\lambda_n(A)$, where $\lambda_n(A)$ are the non-zero eigenvalues of A counted with multiplicity. This supports the intuition that $z \mapsto \det(I + z \cdot A)$ plays the role of the “characteristic polynomial” of A . The value of $\det(I + z \cdot A)$ at z is called Fredholm determinant of $I + z \cdot A$.

The notion of Fredholm determinant can be lifted to operators A which are only in some Schatten class $\mathbf{B}^p(\mathcal{H})$ for some $p \geq 1$ (see [Sim77]). Then $z \mapsto \det_{N+1}(I + z \cdot A)$, $N + 1 \geq p$, a regularized version of the Fredholm determinant, is an entire function of order at most $N + 1$, again with zeros exactly in $-1/\lambda_n(A)$. The regularization process to obtain \det_{N+1} is the classical Weierstraß method of convergence generating factors which lead to canonical Weierstraß products of finite genus.

A completely different notion of regularized determinants appears in the theory of elliptic operators on manifolds. Here one encounters *unbounded* self-adjoint operators L with a meromorphic zeta function (ζ -function)

$$\zeta(s; L) = \sum_{\lambda \in \operatorname{spec} L \setminus \{0\}} \lambda^{-s}$$

and, by some heuristics, one puts

$$\det_{\zeta} L := \exp(-\zeta'(0; L))$$

and calls this the *zeta regularized determinant* (ζ -regularized determinant) of L . This definition goes back to the celebrated work by Ray and Singer [RS71]. It has deep applications, *e.g.*, the Cheeger-Müller theorem on the analytic torsion.

The definition of $\zeta(s; L)$ extends easily to m -sectorial¹ operators. We will only need this for the *normal* operator $L + z$ where $-z$ is in the resolvent set of L . The ζ -function

$$\zeta(s; L + z) = \sum_{\lambda \in \operatorname{spec} L \setminus \{0\}} (\lambda + z)^{-s}, \quad (1.1)$$

a priori defined for $\Re s$ large and $-z$ in the resolvent set of L is then a holomorphic function of two variables (see also Proposition 3.8 below).

It is then not hard to see that, say for an elliptic self-adjoint differential operator L on a closed manifold, the function $z \mapsto \det_{\zeta}(L + z)$ extends to an entire function of finite order with zeros exactly in $-\lambda_n(L)$, where $\lambda_n(L)$ are the eigenvalues of L counted with multiplicity.

¹An operator L in a Hilbert space is m -sectorial if its spectrum lies in a sector $z_0 + \{z \in \mathbb{C} \mid |\arg z| < \delta\}$ for some $z_0 > 0$ and some $\delta < \pi$ and such that outside the sector the resolvent satisfies an estimate $\|(L - z)^{-1}\| \leq C|z|^{-1}$ for $|z| \geq r > 0$, for some $r > 0$. The argument function is normalized such that $\arg 1 = 0$. Cf. [Kat95, Sec. V.3.10].

In [Les98, Proposition 4.6] it was shown that for certain Sturm-Liouville operators one has

$$\frac{\det_{\zeta}(L+z)}{\det_{\zeta}(L)} = \det(I+z \cdot L^{-1}), \quad (1.2)$$

that is there is a simple relation between the ζ -regularized determinant on the one hand and the Fredholm determinant on the other hand.

One of the goals of this paper is to find the most general purely functional analytic setting which guarantees Eq. (1.2) to hold. The result is the Theorem 4.5 in Section 4.2 below which generalizes Eq. (1.2) to

$$\frac{\det_{\zeta}(L+z)}{\det_{\zeta}(L)} = \exp\left(\sum_{j=1}^{p-1} \frac{z^j}{j!} \cdot \frac{d^j}{dz^j} \log \det_{\zeta}(L+z)|_{z=0}\right) \cdot \det_p(I+z \cdot L^{-1}), \quad (1.3)$$

Here L is a self-adjoint bounded below operator in the Hilbert space \mathcal{H} with p -summable resolvent, $p \in \mathbb{Z}$, $p \geq 1$. Furthermore, roughly speaking it is needed that $\zeta(s; L)$ is meromorphic in \mathbb{C} with 0 being a regular point and $s = 1, \dots, p-1$ being at most simple poles. The numbers $\frac{d^j}{dz^j} \log \det_{\zeta}(L+z)|_{z=0}$ are “global” invariants of L in the sense that they cannot be extracted from the heat expansion of L , respectively as poles of $\zeta(s; L)$. However, these numbers can be expressed in terms of (partie finie) zeta values and heat trace coefficients of L (see Proposition 3.8). The non-trivial information of Eq. (1.3), however, is that

$$\log \det_{\zeta}(L+z) - \log \det_{\zeta}(L) - \log \det_p(I+z \cdot L^{-1})$$

is a polynomial of degree at most $p-1$, vanishing at $z=0$. For $p=1$ this specializes to Eq. (1.2). The equality (1.3) contains another important information. Namely, if $\log \det_{\zeta}(L+z)$ has an asymptotic expansion as $\Re z \rightarrow \infty$ then so does $\log \det_p(I+z \cdot L^{-1})$.

It should have become clear that in general the ζ -determinant is notoriously difficult to compute because it is a relatively rigid spectral invariant which cannot be determined “locally” from, e.g., the resolvent or heat trace expansion. However, for *some* operators (see [HLV17] and the references therein) explicit computations are possible. Then the natural question arises whether one can say anything meaningful about the ζ -determinant of some perturbation of a sufficiently nice operator. In this paper we intensively study the behavior of the ζ -determinant $\det_{\zeta}(L+z)$ as a function of the resolvent parameter.

Equation (1.3) leads us naturally to the second topic of this paper. Namely, we would like to explore further the relation between the ζ -regularized determinant and the asymptotic expansion of $\det_{N+1}(I+z \cdot L^{-1})$ as $\Re z \rightarrow \infty$. To motivate this, we formally take the logarithmic derivative of $z \mapsto \det_{\zeta}(L+z)$ to obtain

$$\mathrm{Tr}\left((L+z)^{-1}\right);$$

this exists only if L^{-1} is trace class. If L^{-1} is in some Schatten class then differentiating often enough gives up to a constant factor

$$\mathrm{Tr}\left((L+z)^{-N-1}\right).$$

If the latter has an asymptotic expansion as $z \rightarrow \infty$ (as is the case for the elliptic differential operators as above) the integration should lead to an asymptotic expansion of $\log \det_{\zeta}(L + z)$, respectively $\log \det_{N+1}(I + z \cdot L^{-1})$, as $\Re z \rightarrow \infty$. For elliptic (pseudo)differential operators on compact manifolds (resp. elliptic boundary value problems if a boundary is present) the existence of the asymptotic expansion of $\log \det_{\zeta}(L + z)$ as $\Re z \rightarrow \infty$ can be proven by genuinely microlocal methods, see [BFK92, Appendix].

We show in this paper that no further microlocal methods are needed. Rather the expansion of $\log \det_{\zeta}(L + z)$ as $\Re z \rightarrow \infty$ is equivalent to the resolvent trace expansion of $\text{Tr}(L + z)^{-p}$ as $\Re z \rightarrow \infty$ which is equivalent to the short time asymptotic expansion of the heat trace (Theorem 3.7, Section 3.3). This contains the expansion result of [BFK92, Appendix] as special case.

Together with (1.3) we consequently show in a purely functional analytic setting that the existence of the heat, resp. resolvent trace expansion (plus a mild assumption on the absence of log-terms which is made mostly for convenience) implies the existence of an asymptotic expansion as $\Re z \rightarrow \infty$ of $\log \det_p(I + z \cdot L^{-1})$. In Corollary 4.7, at the end, we give a precise expression of the coefficients of the expansion in terms of the heat expansion coefficients, resp. ζ -function values. An interesting observation is that the constant term in the expansion of $\log \det_p(I + z \cdot L^{-1})$ equals $\log \det_{\zeta} L$. This generalizes the main result of [Fri89]. In the case of certain Sturm-Liouville operators, similar results were obtained before in [HLV17, p. 3439] and in [LT98]. Recently, there has been quite some interest in concrete computations of ζ -determinants and Fredholm determinant. We mention, e.g., [GK19a, GK19b, GK19c] and the references therein.

2. GENERALITIES

2.1. Notation. We will denote by $\mathbb{Z}, \mathbb{R}, \mathbb{C}$ the integer, real, and complex numbers, respectively, $\mathbb{Z}_+ := \{0, 1, 2, \dots\}$ denotes the non-negative integers and by $\mathbb{N} = \{1, 2, 3, \dots\}$ we denote the natural numbers.

For an analytic function $f : \{z \in \mathbb{C} \mid 0 < |z - a| < \epsilon\} \rightarrow \mathbb{C}$ in a punctured neighbourhood of $a \in \mathbb{C}$, we denote by $\text{Res}_a^m f$ the coefficient of $(z - a)^{-m}$ in the Laurent expansion of f about a , i.e., the Laurent expansion reads

$$f(z) = \sum_{m=-\infty}^{\infty} (\text{Res}_a^{-m} f) \cdot (z - a)^m. \quad (2.1)$$

For the ordinary residue, we write $\text{Res}_a f \equiv \text{Res}_a^1 f = a_{-1}$.

The letters o, O will denote the Landau (small and big) O -symbols.

2.2. Regularized integral and partie finie. In this paper we will use freely the useful partie finie regularized integral, cf. [Les97, Sec. 2.1], [Les98, LT98], [LV11, Sec. 1], [LV15, Sec. 1], [HLV17, Sec. 1]. However, to fix some notation and for the convenience of the reader we present here the cheat sheet version.

Let $f : (0, \infty) \rightarrow \mathbb{C}$ be a locally integrable function with asymptotic expansions

$$\begin{aligned} f(x) &\sim \sum_{\alpha \in \mathcal{A}} \sum_{k=0}^{k_{\alpha}^0} a_{\alpha k}^0 x^{\alpha} \log^k x, \quad \text{as } x \rightarrow 0, \\ f(x) &\sim \sum_{\beta \in \mathcal{B}} \sum_{k=0}^{k_{\beta}^{\infty}} a_{\beta k}^{\infty} x^{\beta} \log^k x, \quad \text{as } x \rightarrow \infty. \end{aligned} \quad (2.2)$$

Here, \mathcal{A}, \mathcal{B} are discrete sets of complex numbers with the property that the intersection with each finite vertical strip is finite. Moreover $+\infty$ is the only accumulation point of $\Re \alpha$, $\alpha \in \mathcal{A}$, and $-\infty$ is the only accumulation point of $\Re \beta$, $\beta \in \mathcal{B}$. One could also work with partial asymptotic expansions but for our purposes complete asymptotic expansions suffice. The class of such functions f satisfying Eq. (2.2) will be called “regular functions with complete asymptotics on \mathbb{R}_+ ”. The sets \mathcal{A}, \mathcal{B} may vary from function to function. For convenience we adopt the convention that for f the coefficients $a_{\alpha k}^0, a_{\alpha k}^{\infty}$ are defined for all $\alpha \in \mathbb{C}$ and all $k \in \mathbb{Z}_+$ with the understanding that $a_{\alpha k}^* = 0$ for all but the countably many $\alpha \in \mathcal{A}$, resp. \mathcal{B} , and that for each fixed $\alpha \in \mathbb{C}$ the coefficients $a_{\alpha k}^*$ vanish for all but finitely many k . The expansion may then be written briefly as

$$\sum_{\alpha, k} a_{\alpha k}^{0/\infty} x^{\alpha} \log^k x.$$

If $f : (0, \infty) \rightarrow \mathbb{C}$ is a regular function with complete asymptotics on \mathbb{R}_+ we write

$$\text{LIM}_{x \rightarrow 0} f(x) := a_{00}^0, \quad \text{LIM}_{x \rightarrow \infty} f(x) := a_{00}^{\infty}, \quad (2.3)$$

for the *regularized limits* of f as $x \rightarrow 0$, resp. $x \rightarrow \infty$. In other words the regularized limit of f as $x \rightarrow 0$ ($x \rightarrow \infty$) is the coefficient of $x^0 \log^0 x$ in the asymptotic expansion of f as $x \rightarrow 0$ ($x \rightarrow \infty$).

If f is a regular function with complete asymptotics on \mathbb{R}_+ then so is $F(x) := \int_1^x f(x) dx$ and hence one puts

$$\int_0^{\infty} f(x) dx := \text{LIM}_{x \rightarrow \infty} F(x) - \text{LIM}_{x \rightarrow 0} F(x) = \text{LIM}_{R \rightarrow \infty} \text{LIM}_{\varepsilon \rightarrow 0} \int_{\varepsilon}^R f(x) dx, \quad (2.4)$$

where the order of LIM after the second $=$ is inconsequential. The regularized integral is a convenient extension of the integral to the class of regular functions with complete asymptotics.

It has its peculiarities, though. We emphasize that changes of variables require some care. For dilations one has the following formula [Les97, Lemma 2.1.4]:

$$\lambda \int_0^{\infty} f(\lambda u) du = \int_0^{\infty} f(x) dx + \sum_{k=0}^{\infty} \frac{a_{-1, k}^{\infty} - a_{-1, k}^0}{k+1} \log^{k+1} \lambda. \quad (2.5)$$

Another peculiarity is

$$\int_0^{\infty} x^{\alpha} \cdot \log^k x dx = 0, \quad (2.6)$$

for all $\alpha \in \mathbb{C}$, $k \in \mathbb{Z}_+$ [Les97, Eq. 2.1.12].

2.2.1. *Mellin transform and relation to Hadamard's partie finie.* For a regular function with complete asymptotics f one defines for any $c > 0$ the partial Mellin transforms by

$$\begin{aligned} (M_c^+ f)(z) &:= \int_c^\infty x^{z-1} f(x) dx, & \Re z \ll 0, \\ (M_c^- f)(z) &:= \int_0^c x^{z-1} f(x) dx, & \Re z \gg 0. \end{aligned} \quad (2.7)$$

It turns out [Les97, Sec. 2.1] that M_c^\pm extends meromorphically to \mathbb{C} and the Mellin transform $(Mf)(z) := (M_c^+ f)(z) + (M_c^- f)(z)$ is well-defined independently of the choice of c . Furthermore, it is a meromorphic function with poles of order at most $\max(k_\alpha^\infty, k_\alpha^0) + 1$ in $-\alpha$ [Les97, Prop-Def. 2.1.2]. There are explicit formulas for the principal part of Mf about $-\alpha$ in terms of $a_{\alpha,k}^{0/\infty}$, cf. [Les97, Eq. 2.1.8 and 2.1.9]. This may be viewed as a version of Watson's Lemma (see [Wat18, pg. 133]) for the Mellin transform.

Outside the poles of Mf one now finds the convenient global representation

$$(Mf)(z) = \int_0^\infty x^{z-1} f(x) dx \quad (2.8)$$

for all $z \in \mathbb{C}$ except the set $-\mathcal{A} \cup (-\mathcal{B})$.

The Mellin transform can be used for an alternative (equivalent) approach to the regularized integral: namely, if F is a meromorphic function in a neighbourhood of $a \in \mathbb{C}$ one defines the *finite part* of F at a by

$$\text{Pf}_{z=a} F(z) := (\text{Pf} F)(a) := \text{Res}_{z=a} \left((z-a)^{-1} F(z) \right), \quad (2.9)$$

where the residue notation is explained in Section 2.1. In other words $(\text{Pf} F)(a)$ is the constant term in the Laurent expansion of $F(z)$ about a . Needless to say, if F is regular at a then $(\text{Pf} F)(a) = F(a)$. Fortunately, it turns out that the finite part of the Mellin transform at 1 of a regular function with complete asymptotics gives its regularized integral:

Proposition 2.1 ([Les97, Prop. 2.1.7]). *Let $f : (0, \infty) \rightarrow \mathbb{C}$ be a regular function with complete asymptotics. Then*

$$\int_0^\infty f(x) dx = \text{Pf}_{z=1} Mf(z) = \text{Pf}_{z=0} \int_0^1 x^z f(x) dx + \text{Pf}_{z=0} \int_1^\infty x^z f(x) dx,$$

where the integrals are a priori defined for $\Re z$ large, resp. small, and then analytically continued to \mathbb{C} (a neighbourhood of $z = 0$ would suffice).

With this we have the formula, valid for all $z \in \mathbb{C}$,

$$(\text{Pf} Mf)(z) = \int_0^\infty x^{z-1} f(x) dx. \quad (2.10)$$

2.2.2. *Example.* We illustrate the concepts introduced before for the derivatives of the Γ -function. This will also be needed later. For future reference we record the following facts about Γ :

$$\Gamma(s) = \frac{1}{s} - \gamma + O(s), \quad \frac{1}{\Gamma(s)} = s + \gamma \cdot s^2 + O(s^3), \quad \text{as } s \rightarrow 0, \quad (2.11)$$

$$\text{Res}_{-n} \Gamma = \frac{(-1)^n}{n!}, \quad (\text{Pf } \Gamma)(-n) = (\text{Res}_{-n} \Gamma) \cdot (L_n - \gamma). \quad (2.12)$$

Here, $L_n := \sum_{j=1}^n \frac{1}{j}$ and $\gamma := \lim_{n \rightarrow \infty} \left(\sum_{j=1}^n \frac{1}{j} - \log n \right)$ denotes the Euler-Mascheroni constant. Recall from Section 2.1 that $\text{Res}_\xi f$ denotes the residue of the analytic function f about the isolated singularity ξ . For $n \in \mathbb{Z}_+$, $n > 0$,

$$\frac{1}{\Gamma(n+s)} = \frac{1}{(n-1)!} + \frac{1}{(n-1)!} \cdot (\gamma - L_{n-1})s + O(s^2), \quad \text{as } s \rightarrow 0, \quad (2.13)$$

$$\frac{1}{\Gamma(-n+s)} = (-1)^n n! \cdot s + (-1)^n n! \cdot (\gamma - L_n)s^2 + O(s^3), \quad \text{as } s \rightarrow 0. \quad (2.14)$$

We have

$$(\text{Pf } \partial^k \Gamma)(\alpha) = \int_0^\infty x^{\alpha-1} \log^k x e^{-x} dx, \quad (2.15)$$

for all $\alpha \in \mathbb{C}$. Clearly, the equality is valid for $\Re \alpha > 0$ as an ordinary integral. Hence, by analytic continuation and Eq. (2.10) it is valid for all α . The change of variables formula will be used to show the next lemma.

Lemma 2.2. *For $\alpha, z \in \mathbb{C}$ with $\Re z > 0$ we have*

$$\begin{aligned} & z^\alpha \cdot \int_0^\infty x^{\alpha-1} \log^k x e^{-xz} dx \\ &= \int_0^\infty x^{\alpha-1} \log^k(x/z) e^{-x} dx + \delta_{-\alpha, \mathbb{Z}_+} \frac{(-1)^{k+1} (\text{Res}_\alpha \Gamma) \log^{k+1} z}{k+1} \\ &= \sum_{j=0}^k \binom{k}{j} (-1)^j (\text{Pf } \partial^{k-j} \Gamma)(\alpha) \cdot \log^j z + \delta_{-\alpha, \mathbb{Z}_+} \frac{(-1)^{k+1} (\text{Res}_\alpha \Gamma) \log^{k+1} z}{k+1}, \end{aligned}$$

where

$$\delta_{-\alpha, \mathbb{Z}_+} = \begin{cases} 1, & \text{if } -\alpha \in \mathbb{Z}_+, \\ 0, & \text{otherwise.} \end{cases}$$

In the proof we will need the combinatorial formula

$$\sum_{j=0}^k \binom{k}{j} \frac{1}{j+1} (-a)^{k-j} a^{j+1} = \frac{(-1)^k a^{k+1}}{k+1} \quad (2.16)$$

which follows by noting that $\binom{k}{j} \frac{1}{j+1} = \frac{1}{k+1} \binom{k+1}{j+1}$ and applying the binomial theorem.

Proof. We apply the change of variables formula Eq. (2.5) to the function $f(x) = (x/z)^{\alpha-1} \log^k(x/z) e^{-x}$. Certainly this is a regular function with complete asymptotics. Since $\Re z > 0$ it decays exponentially as $x \rightarrow \infty$ and therefore we have $a_{\alpha k}^\infty = 0$. It remains to find the terms $x^{-1} \log^k x$ in the asymptotic expansion as $x \rightarrow 0$, or in the notation of Eq. (2.2), $a_{-1,k}^0$. The expansion as $x \rightarrow 0$ is obtained from the Taylor expansion of e^{-x} . Thus if $-\alpha \notin \mathbb{Z}_+$ then $a_{-1,k}^0 = 0$ as well. If $-\alpha \in \mathbb{Z}_+$ then we find for the terms of the form $x^{-1} \log^j x$ as $x \rightarrow 0$

$$\begin{aligned} & (x/z)^{\alpha-1} \sum_{j=0}^k \binom{k}{j} (-\log z)^{k-j} \log^j x \frac{(-1)^{\alpha}}{(-\alpha)!} x^{-\alpha} = \\ & x^{-1} z^{1-\alpha} (\text{Res}_\alpha \Gamma) \sum_{j=0}^k \binom{k}{j} (-\log z)^{k-j} \log^j x. \end{aligned}$$

Thus the correction term in the formula (2.5) equals

$$(\text{Res}_\alpha \Gamma) z^{1-\alpha} \frac{(-1)^{k+1} \log^{k+1} z}{k+1},$$

where we have used the Eq. (2.15) and (2.16). The rest is now straightforward. \square

3. OPERATORS WITH MEROMORPHIC ζ -FUNCTION

3.1. Set up. For the trace ideals in the algebra of bounded operators on a separated Hilbert space we refer to [Sim05] as a standard reference.

We now describe the class of operators we are going to consider in the remainder of this paper. Let \mathcal{H} be a separable complex Hilbert space and let $L : \mathcal{D}(L) \subset \mathcal{H} \rightarrow \mathcal{H}$ be a self-adjoint operator, which is bounded below. It is not a big loss of generality to assume for convenience that

$$\langle Lu, u \rangle \geq 0, \quad u \in \mathcal{D}(L) \subset \mathcal{H}. \quad (3.1)$$

We consider the heat semigroup e^{-tL} associated to L for any $t \geq 0$. Since $L = L^* \geq 0$ we have that $\text{spec } L$ is contained in the nonnegative real axis. We assume additionally that

$$(L + I)^{-1} \in \mathbf{B}^p(\mathcal{H}), \quad (3.2)$$

for some $p \geq 1$. Here, $\mathbf{B}^p(\mathcal{H})$ denotes the Schatten class of order p . The condition (3.2) implies that the sum

$$\zeta(s; L) := \sum_{\lambda \in \text{spec } L \setminus \{0\}} \lambda^{-s}, \quad (3.3)$$

converges absolutely and locally uniformly for $\Re(s) > p$. The function $\zeta(s; L)$ is called the *zeta function* (*ζ -function*) associated to L . The Mellin transform of $e^{-\lambda t}$,

$$\int_0^\infty t^{s-1} e^{-t\lambda} dt = \Gamma(s) \cdot \lambda^{-s}, \quad \lambda > 0,$$

implies the representation

$$\begin{aligned}\zeta(s; L) &= \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} \cdot (\operatorname{Tr}(e^{-tL}) - \dim \ker L) dt, \\ &= \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} \cdot \operatorname{Tr}(e^{-tL}) dt,\end{aligned}\tag{3.4}$$

for $\Re(s) > p$. The second line follows from Eq. (2.6). Following Connes-Moscovici (cf. [CM95, Definition II.1] and [Les13, Definition 2.1]) we define:

Definition 3.1. *The operator L is said to have discrete dimension spectrum, if $\zeta(s; L)$ extends to a meromorphic function in the complex plane \mathbb{C} such that on finite vertical strips one has a uniform estimate $|\Gamma(s) \cdot \zeta(s; L)| = O(|s|^{-N})$, when $\Im(s) \rightarrow \infty$, for each $N \in \mathbb{N}$. We will denote by $\Sigma(L)$ the set of poles of the function $\Gamma(s) \cdot \zeta(s; L)$.*

The discrete dimension spectrum assumption is, via the formula (3.4), in fact equivalent to a strengthened form of a complete asymptotic expansion of the trace of the heat semigroup. To explain this, we first note that by Eq. (3.1) and (3.2) one has

$$\operatorname{Tr}(e^{-tL}) \sim \dim \ker L + O(e^{-t\lambda_{\min}}), \quad \text{as } t \rightarrow \infty,\tag{3.5}$$

with $\lambda_{\min} > 0$ denoting the smallest positive eigenvalue of L . Hence the asymptotic expansion as $t \rightarrow \infty$ in the sense of Eq. (2.2) is automatic. Now let us *assume* that

$$\operatorname{Tr}(e^{-tL}) \sim \sum_{\alpha \in \mathcal{A}} \sum_{k=0}^{k_\alpha} A_{\alpha k}^H t^\alpha \log^k t, \quad \text{as } t \rightarrow 0+,\tag{3.6}$$

with a discrete set $\mathcal{A} \subset \mathbb{C}$ such that the intersection with each finite vertical strip is finite and such that $\Re \alpha, \alpha \in \mathcal{A}$ has only $+\infty$ as accumulation point. The assumption (3.6) means in other words that the heat semigroup is a regular function of t with complete asymptotics in the sense of Section 2.2. Furthermore, by Eq. (3.4), $\Gamma(s) \cdot \zeta(s; L)$ is nothing but the Mellin-transform of the trace of the heat semigroup. It then follows from the discussion in Section 2.2.1 that $\zeta(s; L)$ extends meromorphically to \mathbb{C} with poles in $-\mathcal{A}$. In fact, from Proposition 2.1 we infer the universally valid formula

$$\left(\operatorname{Pf}(\Gamma(\cdot) \cdot \zeta(\cdot; L)) \right)(s) = \int_0^\infty t^{s-1} \operatorname{Tr}(e^{-tL}) dt, \quad s \in \mathbb{C},$$

resp., for all $s \in \mathbb{C} \setminus (-\mathcal{A})$ we have

$$\zeta(s; L) = \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} \operatorname{Tr}(e^{-tL}) dt.$$

As a word of warning, we emphasize that one cannot expect this to hold for the finite parts in the poles, cf. the discussion in the next section below.

For $\alpha \in \mathcal{A}$ the principal part of $\zeta(s; L)$ about $-\alpha$ is determined by

$$\frac{1}{\Gamma(s)} \sum_{k=0}^{k_\alpha} (-1)^k k! \frac{A_{\alpha k}^{H'}}{(s + \alpha)^{k+1}},\tag{3.7}$$

where (see *e.g.*, [BL99, Sec. 2]).

$$A_{\alpha k}^{H'} := \begin{cases} A_{\alpha k}^H, & (\alpha, k) \neq (0, 0), \\ A_{\alpha k}^{H'} = A_{00}^H - \dim \ker L, & (\alpha, k) = (0, 0). \end{cases} \quad (3.8)$$

As a consequence, if $\alpha \notin \mathbb{Z}_+$ the order of the pole $-\alpha$ is $k_\alpha + 1$ with residue

$$\operatorname{Res}_{-\alpha}^{k_\alpha+1} \zeta(\cdot; L) = (-1)^{k(\alpha)} \frac{A_{\alpha k_\alpha}^H (k_\alpha)!}{\Gamma(-\alpha)}, \quad (3.9)$$

and if $\alpha \in \mathbb{Z}_+$ then, since $\Gamma(s)^{-1}$ has a simple zero at $-\alpha$, the order of the pole $-\alpha$ is k_α with residue

$$\operatorname{Res}_{-\alpha}^{k_\alpha} \zeta(\cdot; L) = (-1)^{k_\alpha+\alpha} \alpha! (k_\alpha)! A_{\alpha k_\alpha}^{H'}. \quad (3.10)$$

Recall that $\operatorname{Res}_a^k f$ denotes the coefficient of $(s-a)^{-k}$ in the Laurent expansion of f about a , see Section 2.1.

While the complete asymptotic expansion implies that $\zeta(s; L)$ has a meromorphic extension to the whole plane, in order to obtain the expansion from the meromorphic ζ -function, one needs the finite vertical strip decay assumption in order to be able to shift vertical integration contours from the inverse Mellin transform (*cf.* [BL99, Sec. 2]). This condition, being slightly stronger than just meromorphicity, is reflected in a stronger asymptotic expansion condition. The result, being a standard application of complex analysis, reads as follows (see *e.g.*, [BL99, Lemma 2.2] and the references therein).

Proposition 3.2. *Let L be a bounded below discrete operator in a Hilbert space with resolvent of \mathfrak{p} -Schatten class. The operator L has discrete dimension spectrum if, and only if, $\operatorname{Tr}(e^{-tL})$ has an asymptotic expansion (3.6) that can be differentiated, i.e., for $N \in \mathbb{Z}_+$, $K > 0$, we have*

$$\left| \partial_t^N \left(\operatorname{Tr}(e^{-tL}) - \sum_{\Re \alpha \leq N+K} \sum_{k=0}^{k_\alpha} A_{\alpha k}^H t^\alpha \log^k t \right) \right| \leq C_{N,K} t^K, \quad \text{as } t \rightarrow 0+. \quad (3.11)$$

From the asymptotic expansion of the trace of the heat semigroup one can derive various other trace expansions. Well-known is the relation to the resolvent expansion which we will also briefly review in Section 3.3 below. Less well-known is probably the expansion of the *zeta determinant* (ζ -determinant) which we are going to explain first.

3.2. The ζ -determinant and its expansion in terms of the resolvent parameter.

3.2.1. Standing assumptions. During this subsection, unless otherwise said, L denotes a discrete bounded below (Eq. (3.1)) self-adjoint operator in the Hilbert space \mathcal{H} whose resolvent lies in the \mathfrak{p}^{th} -Schatten class. Furthermore, we assume Eq. (3.6) to hold and from now on we write $A_{\alpha k}^H(L)$ to indicate the dependence of the operator L on the heat coefficients; recall that Eq. (3.5) is automatic. By Eq. (3.10), 0 is a pole of $\zeta(s; L)$ of order $k_0 := k(0)$. In this section, we would like to ensure that $\zeta(s; L+z)$ is regular at $s=0$ for all $\Re z > 0$ and therefore in this section we assume in addition that $k_{-n} = 0$ for all $n \in \mathbb{Z}_+$. In view of Eq. (3.7) and

(3.9) this is equivalent to $A_{-n,k}^H(L) = 0$ for all $n \in \mathbb{Z}_+$ and all $k > 0$. In terms of $\zeta(s; L)$ this means that $\zeta(s; L)$ is regular at 0 and that in the positive integers there are at most poles of at most order 1.

3.2.2. *The ζ -determinant of $L+z$.* Under our standing assumptions $\zeta(s; L)$ is regular at $s = 0$.

Definition 3.3. *The ζ -regularized determinant of L is defined by*

$$\det_\zeta(L) := \exp(-\zeta'(0; L)).$$

Our first result relates the ζ -determinant of L to the heat trace.

Proposition 3.4. *Let L satisfy the Standing assumptions 3.2.1. Then $\zeta(s; L)$ is regular at $s = 0$ and one has*

$$\zeta(0; L) = A_{00}^H(L) - \dim \ker L, \quad (3.12)$$

$$-\log \det_\zeta(L) = \gamma \cdot (A_{00}^H(L) - \dim \ker L) + \int_0^\infty t^{-1} \operatorname{Tr}(e^{-tL}) dt. \quad (3.13)$$

Here, γ denotes the Euler-Mascheroni constant (see Eq. (2.11), (2.12)) and $A_{00}^H(L)$ denotes the coefficient of $t^0 \log^0 t$ in the asymptotic expansion of $\operatorname{Tr}(e^{-tL})$ as $t \rightarrow 0+$, see Eq. (3.6) and (3.8).

While Eq. (3.12) has been known for a long time, Eq. (3.13) is more subtle. At least for Dirac type operators, Eq. (3.13) was already known to K. P. Wojciechowski, cf. [Woj99, Eq. (3.5) and (3.6)]. These formulas are the key to a simple derivation of the asymptotic expansion of $\zeta(0; L+z)$ and $\log \det_\zeta(L+z)$ as $\Re z \rightarrow \infty$. The latter will lead to yet another equivalent characterization of the discrete dimension spectrum assumption. It is on the same footing as the resolvent expansion.

Proof. As $s \rightarrow 0$ we have for the Mellin transform

$$\int_0^\infty t^{s-1} \operatorname{Tr}(e^{-tL}) dt = \frac{A_{00}^H(L) - \dim \ker L}{s} + \int_0^\infty t^{-1} \operatorname{Tr}(e^{-tL}) dt + O(s).$$

We use, as before, $A_{00}^{H'}(L) := A_{00}^H(L) - \dim \ker L$. By Section 2.2.1 and Eq. (2.11) we obtain as $s \rightarrow 0$

$$\begin{aligned} \zeta(s; L) &= (s + \gamma \cdot s^2 + O(s^3)) \cdot \left(\frac{A_{00}^{H'}(L)}{s} + \int_0^\infty t^{-1} \operatorname{Tr}(e^{-tL}) dt + O(s) \right), \\ &= A_{00}^{H'}(L) + \left(\gamma A_{00}^{H'}(L) + \int_0^\infty t^{-1} \operatorname{Tr}(e^{-tL}) dt \right) \cdot s + O(s^2) \end{aligned}$$

and the claim follows. \square

We will later need the behavior of $\zeta(s; L)$ near $n > 0$, $n \in \mathbb{Z}_+$. Under the Standing assumptions 3.2.1, we apply the same strategy as in the proof of the previous proposition and obtain for the poles of $\zeta(s; L)$

$$\zeta(s; L) \sim \sum_{n=1}^{\infty} \frac{A_{-n,0}^H}{\Gamma(s)} \cdot \frac{1}{s-n} + \text{Entire}(s).$$

Hence for an integer $n > 0$,

$$\zeta(s+n; L) \sim \frac{A_{-n,0}^H}{(n-1)!} \cdot \frac{1}{s} + \text{Pf}_{s=n} \zeta(s; L) + O(s), \quad \text{as } s \rightarrow 0. \quad (3.14)$$

Lemma 3.5. *Let L satisfy the Standing assumptions 3.2.1. Then we have for $\Re z > 0$*

$$\zeta(0; L+z) = A_{00}^H(L+z) = \sum_{n=0}^{\infty} (\text{Res}_{-n} \Gamma) \cdot A_{-n,0}^H(L) \cdot z^n,$$

Furthermore, the ζ -function of $L+z$ is also regular at 0 and it has at most simple poles in the positive integers. In other words, for all $\Re z > 0$ the operator $L+z$ satisfies the Standing assumptions 3.2.1.

Note that the previous sum is finite.

Proof. Let $\Re z > 0$. Then $\text{Tr}(e^{-t(L+z)}) = O(e^{-tz})$, as $t \rightarrow \infty$. As $t \rightarrow 0+$ we have

$$\begin{aligned} \text{Tr}(e^{-t(L+z)}) &= e^{-tz} \cdot \text{Tr}(e^{-tL}) \\ &\sim \sum_{n=0}^{\infty} \frac{(-z)^n}{n!} t^n \cdot \sum_{\alpha,k} A_{\alpha k}^H(L) \cdot t^\alpha \log^k t \\ &\sim \sum_{\alpha} \sum_{n=0}^{\infty} \sum_k (\text{Res}_{-n} \Gamma) \cdot A_{\alpha-n,k}^H(L) \cdot z^n \cdot t^\alpha \log^k t. \end{aligned}$$

This gives in fact the general formula

$$A_{\alpha k}^H(L+z) = \sum_{n=0}^{\infty} (\text{Res}_{-n} \Gamma) \cdot A_{\alpha-n,k}^H(L) \cdot z^n.$$

From this we read off the claim about the constant term as well as the fact that since $A_{\alpha k}^H(L) = 0$ for $-\alpha \in \mathbb{Z}_+, k > 0$, this is also true for the $A_{\alpha k}^H(L+z)$. \square

Lemma 3.6. *Let L satisfy the Standing assumptions 3.2.1. Then as $\Re z \rightarrow \infty$*

$$\int_0^{\infty} t^{-1} \text{Tr}(e^{-t(L+z)}) dt \sim \sum_{\alpha,k} A_{\alpha k}^H(L) \int_0^{\infty} e^{-tz} \cdot t^{\alpha-1} \log^k t dt + O(e^{-\Re z}).$$

Proof. We split the integral into \int_0^1 and \int_1^{∞} . Since $\text{Tr}(e^{-t(L+z)}) = O(e^{-\Re z})$, for $t \geq 1$ the contribution of \int_1^{∞} is $O(e^{-\Re z})$ as well, and hence it contributes only to the error term of the claimed expansion.

Next we fix a large N and write

$$\text{Tr}(e^{-tL}) = \sum_{\Re \alpha \leq N} \sum_{k=0}^{k_\alpha} A_{\alpha k}^H(L) \cdot t^\alpha \log^k t + O(t^N),$$

as $t \rightarrow 0+$. The contribution of the remainder to \int_0^1 is $O(z^{-N})$. So it remains to look at each individual summand which contributes

$$\int_0^1 e^{-tz} \cdot t^{\alpha-1} \log^k t dt = \int_0^{\infty} e^{-tz} \cdot t^{\alpha-1} \log^k t dt - \int_1^{\infty} e^{-tz} \cdot t^{\alpha-1} \log^k t dt.$$

The contribution of the last summand is again of exponential decay in z . Thus we reach the conclusion. \square

Combining Lemmas 2.2, 3.5, and 3.6 we arrive at

Theorem 3.7. *Let L be a discrete self-adjoint bounded below operator in the Hilbert space \mathcal{H} with resolvent of p^{th} -Schatten class. Moreover, assume that the trace of the heat semigroup has an asymptotic expansion (3.6) with $A_{\alpha k}^H(L) = 0$ for $\alpha \leq 0$ and $k > 0^2$. Then $\log \det_{\zeta}(L + z)$ has an asymptotic expansion as $\Re z \rightarrow \infty$*

$$\log \det_{\zeta}(L + z) \sim_{\Re z \rightarrow \infty} - \sum_{\substack{\alpha, k \\ -\alpha \notin \mathbb{Z}_+}} (-1)^k \sum_{j \geq k} A_{\alpha j}^H(L) \cdot \binom{j}{k} \cdot (\text{Pf } \partial^{j-k} \Gamma)(\alpha) \cdot z^{-\alpha} \log^k z \quad (3.15)$$

$$+ A_{00}^H(L) \cdot \log z \quad (3.16)$$

$$+ \sum_{n=1}^{\infty} A_{-n, 0}^H(L) \cdot (\text{Res}_{-n} \Gamma) \cdot (z^n \log z - L_n \cdot z^n), \quad (3.17)$$

where $L_n = \sum_{j=1}^n \frac{1}{j}$.

Note that the coefficient of $z^0 \log^0 z$ in this expansion vanishes. Thus we obtain a different proof of [HLV17, Lemma 2.2].

For (general) pseudodifferential operators on a compact manifold (with or without boundary) the result was proved by genuinely pseudodifferential methods in [BFK92, Appendix]. Our standing assumption is in general only fulfilled for *differential operators*. Our result shows, however, that even if there are no log-terms in the heat expansion, nevertheless the terms t^{-n} , $n \in \mathbb{Z}_+$, in the heat expansion cause a term $z^n \log z$ in the expansion of $\log \det_{\zeta}$. Insofar we cannot confirm the claim made on the bottom of page 63 in [BFK92]. Of course the Standing assumption 3.2.1 can be relaxed to cover [BFK92]'s result completely. If there is a pole at 0 of the ζ -function we define the regularized determinant as the finite part of $-\zeta'$. This is not a problem at all. However, formulas become very clumsy.

Proof. The result follows by combining Lemmas 2.2, 3.5, and 3.6, Proposition 3.4 and Eq. (2.11), (2.12): We apply (3.12) to $L + z$ to obtain

$$\log \det_{\zeta}(L + z) = -\gamma \cdot A_{00}^H(L + z) - \int_0^{\infty} t^{-1} \text{Tr}(e^{-t(L+z)}) dt.$$

By Lemma 3.5 the first summand expand as

$$-\gamma \cdot A_{00}^H(L + z) = - \sum_{n=0}^{\infty} (\text{Res}_{-n} \Gamma) \cdot A_{-n, 0}^H(L) \cdot \gamma \cdot z^n. \quad (3.18)$$

To the second summand we apply Lemma 3.6 and obtain

$$- \int_0^{\infty} t^{-1} \text{Tr}(e^{-t(L+z)}) dt \sim_{\Re z \rightarrow \infty} - \sum_{\alpha, k} A_{\alpha k}^H(L) \int_0^{\infty} e^{-tz} t^{\alpha-1} \log^k t dt. \quad (3.19)$$

²i.e. the Standing assumptions 3.2.1 are fulfilled.

To each regularized integral on the right hand side of Eq. (3.19) we apply Lemma 2.2:

$$-\int_0^\infty e^{-tz} t^{\alpha-1} \log^k t dt = -\sum_{j=0}^k \binom{k}{j} (-1)^j (\text{Pf } \partial^{k-j} \Gamma)(\alpha) \cdot z^{-\alpha} \log^j z \quad (3.20)$$

$$+ \delta_{-\alpha, \mathbb{Z}_+} \frac{(-1)^k (\text{Res}_\alpha \Gamma)}{k+1} z^{-\alpha} \log^{k+1} z. \quad (3.21)$$

Due to the assumption, $A_{\alpha k}^H = 0$ if $-\alpha \in \mathbb{Z}_+$ and $k > 0$ the last summand appears only with $k = 0$.

Combining (3.18) - (3.21) we arrive at

$$\begin{aligned} \log \det_\zeta(L+z) &\sim_{\Re z \rightarrow \infty} \\ &-\sum_{\substack{\alpha, k \\ -\alpha \notin \mathbb{Z}_+}} A_{\alpha k}^H(L) \sum_{j=0}^k \binom{k}{j} (-1)^j (\text{Pf } \partial^{k-j} \Gamma)(\alpha) \cdot z^{-\alpha} \log^j z \end{aligned} \quad (3.22)$$

$$-\sum_{n=0}^{\infty} A_{-n,0}^H(L) \cdot \left(\gamma \cdot (\text{Res}_{-n} \Gamma) + (\text{Pf } \Gamma)(-n) \right) \cdot z^{-n} \quad (3.23)$$

$$+\sum_{n=0}^{\infty} A_{-n,0}^H(L) \cdot (\text{Res}_{-n} \Gamma) \cdot z^n \log z. \quad (3.24)$$

Substituting $(j, k) \rightarrow (k, j)$, Eq. (3.22) gives (3.15). Using Eq. (2.11) and (2.12) one checks that

$$\text{Res}_0 \Gamma = 1, \quad \gamma \cdot \text{Res}_0 \Gamma + (\text{Pf } \Gamma)(0) = 0,$$

and for $n > 0$

$$(\text{Pf } \Gamma)(-n) = (\text{Res}_{-n} \Gamma) \cdot (L_n - \gamma).$$

Thus Eq. (3.23), (3.24) together give Eq. (3.16) and (3.17) completing the proof. \square

We are now able to describe the Taylor expansion of $\log \det_\zeta(L+z)$ about $z = 0$.

Proposition 3.8. *Let L satisfy the Standing assumptions 3.2.1. Then $\zeta(s; L+z)$ is holomorphic for (s, z) in a neighbourhood of $(0, 0)$ and for $n \in \mathbb{Z}_+$, $n > 0$,*

$$\frac{d^n}{dz^n} \log \det_\zeta(L+z) \Big|_{z=0} = (-1)^{n-1} (n-1)! \left(\text{Pf}_{s=n} \zeta(s; L) + \frac{A_{-n,0}^H}{(n-1)!} \cdot L_{n-1} \right), \quad (3.25)$$

where $L_{n-1} = \sum_{j=1}^{n-1} \frac{1}{j}$. Therefore, the Taylor expansion of $\log \det_\zeta(L+z)$ about $z = 0$ reads

$$\begin{aligned} \log \det_\zeta(L+z) &= \log \det_\zeta(L) + \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} \cdot (\text{Pf}_{s=n} \zeta(s; L)) \cdot z^n \\ &\quad + \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n!} \cdot L_{n-1} \cdot A_{-n,0}^H \cdot z^n. \end{aligned}$$

Note that the last sum is in fact finite.

Proof. By Hartogs's theorem [Hör90, Theorem 2.2.8], to prove that the zeta function of $L + z$ is holomorphic in a neighbourhood of $(0, 0)$ it is enough to verify that $\zeta(s; L + z)$ is holomorphic in s and z , separately. For fixed z with $|z| < \lambda_{\min}$, Lemma 3.5 shows that $s \mapsto \zeta(s; L + z)$ is regular near $s = 0$. For fixed s , s small enough, one invokes the heat expansion to see that $z \mapsto \zeta(s; L + z)$ is regular near $z = 0$. Therefore, $\zeta(s; L + z)$ is holomorphic in a neighbourhood of $(0, 0)$ and we can commute the derivative in z with the derivative in s of $\zeta(s; L + z)$. Thus, let $n \in \mathbb{Z}_+$, $n > 0$.

$$\begin{aligned} \frac{d^n}{dz^n} \log \det_{\zeta}(L + z)|_{z=0} &= -\frac{d^n}{dz^n} \left(\frac{d}{ds} \zeta(s; L + z)|_{s=0} \right) |_{z=0} \\ &= -\frac{d}{ds} \left(\frac{d^n}{dz^n} \text{Tr}((L + z)^{-s})|_{z=0} \right) |_{s=0} \\ &= -\frac{d}{ds} \left((-1)^{n-1} \left(\prod_{j=1}^{n-1} (s + j) \right) \cdot s \cdot \zeta(n + s; L) \right) |_{s=0}. \end{aligned}$$

Now we use Eq. (3.14) and obtain Eq. (3.25). \square

3.3. The heat expansion and the resolvent expansion. The method of proof of Theorem 3.7 in fact can be summarized in yet another Watson Lemma type expansion lemma. As an application we present a concise exposition of the equivalence between the heat and the resolvent expansion in full generality.

In this section we still assume the Standing assumptions 3.2.1³. For the self-adjoint operator L the resolvent can be expressed in terms of the heat operator as

$$(L + z)^{-1} := \int_0^{\infty} e^{-zt} e^{-tL} dt, \quad \text{for } \Re z > 0, \quad (3.26)$$

and more generally,

$$(L + z)^{-N} := \frac{1}{(N-1)!} \int_0^{\infty} t^{N-1} e^{-zt} e^{-tL} dt, \quad \text{for } \Re z > 0. \quad (3.27)$$

Under the Assumption (3.2), we have for $N \geq p$

$$\text{Tr}((L + z)^{-N}) = \frac{1}{(N-1)!} \int_0^{\infty} t^{N-1} e^{-zt} \text{Tr}(e^{-tL}) dt. \quad (3.28)$$

Assuming Eq. (3.6) the expansion of $\text{Tr}((L + z)^{-N})$ as $\Re z \rightarrow \infty$ follows along the lines of the proof of the Theorem 3.7 and the lemmas used therein. However, we think it is worth to single out the following regularized integral version of Watson's Lemma (cf. [Wat18, Pg. 133]).

Lemma 3.9. *Let $q : (0, \infty) \rightarrow \mathbb{C}$ be a locally integrable function. Assume that as $t \rightarrow 0+$ q has an asymptotic expansion*

$$q(t) \sim \sum_{\alpha, k} A_{\alpha k}^q t^{\alpha-1} \log^k t, \quad \text{as } t \rightarrow 0+, \quad (3.29)$$

³Actually the assumption on the vanishing of $A_{-n, k}^H(L)$ for $n \in \mathbb{Z}_+$ and $k > 0$ is not needed and may be dropped in this section.

and that there exists $r_0 > 0$ such that

$$\int_1^\infty e^{-r_0 t} |q(t)| dt < \infty. \quad (3.30)$$

Then we have an asymptotic expansion

$$\int_0^\infty e^{-zt} q(t) dt \sim \sum_{\alpha, k} A_{\alpha k}^q \int_0^\infty e^{-tz} t^{\alpha-1} \log^k t dt + O((\Re z)^{-N}), \text{ as } \Re z \rightarrow \infty, \quad (3.31)$$

$$\sim \sum_{\alpha, k} (-1)^k \sum_{j \geq k} A_{\alpha j}^q \binom{j}{k} (\text{Pf } \partial^{j-k} \Gamma)(\alpha) \cdot z^{-\alpha} \cdot \log^k z \quad (3.32)$$

$$+ \sum_{n=0}^\infty \sum_{k=0}^{k_n} A_{-n, k}^q \frac{(-1)^{k-1} \text{Res}_{-n} \Gamma}{k+1} \cdot z^n \cdot \log^{k+1} z + O((\Re z)^{-N}), \quad (3.33)$$

for any $N > 0$.

Remark 3.10. (1) In Eq. (3.29) we chose to write the exponent of t as $\alpha - 1$ instead of α to have a better comparability to Lemmas 2.2, 3.5, 3.6 and Theorem 3.7.

(2) The notation $\sum_{\alpha, k}$ has the usual meaning as elaborated in Section 2.2, cf. also

Eq. (3.6).

(3) The expansion (3.31) is valid as $\Re z \rightarrow \infty$ in particular it holds in any sector $\Lambda_\epsilon := \{z \in \mathbb{C} \mid |\arg z| < \epsilon\}$ for some $0 < \epsilon < \pi/2$. Furthermore, in any such sector we have

$$O((\Re z)^{-N}) = O(|z|^{-N}), \quad \text{as } |z| \rightarrow \infty,$$

in Λ_ϵ for any $N > 0$.

Proof. The proof follows the same scheme as the proof of Lemma 3.6 and of Theorem 3.7.

First we split the integral into \int_0^1 and \int_1^∞ . Due to the assumption (3.30) we find

$$\left| \int_1^\infty e^{-zt} q(t) dt \right| \leq e^{-(\Re z - r_0)t} \int_1^\infty e^{-r_0 t} |q(t)| dt = O\left(e^{-(\Re z - r_0)t}\right) = O((\Re z)^{-N}).$$

Fixing a large K and writing

$$q(t) = \sum_{\Re \alpha \leq K} \sum_{k=0}^{k_\alpha} A_{\alpha k}^q t^{\alpha-1} \log^k t + O(t^K) =: q_0(t) + O(t^K),$$

we note that

$$\begin{aligned} \left| \int_0^1 e^{-zt} (q(t) - q_0(t)) dt \right| &\leq C \cdot \int_0^1 e^{-(\Re z)t} t^K dt \\ &= C \cdot \int_0^{(\Re z)t} e^{-u} u^K du \cdot (\Re z)^{-K-1} = O\left((\Re z)^{-K-1}\right). \end{aligned}$$

Therefore it remains to look at each individual summand which, as in the proof of Lemma 3.6, contributes

$$\int_0^1 e^{-tz} \cdot t^{\alpha-1} \log^k t dt = \int_0^\infty e^{-zt} \cdot t^{\alpha-1} \log^k t dt - \int_1^\infty e^{-zt} \cdot t^{\alpha-1} \log^k t dt.$$

The second integral is $O\left(e^{-\Re z - \delta}\right)$, $\Re z \rightarrow \infty$ for any $\delta > 0$. Thus the expansion (3.31) is proved.

As in the proof of Theorem 3.7 we apply Lemma 2.2 to each regularized integral on the right hand side of Eq. (3.31):

$$\begin{aligned} & \int_0^\infty e^{-zt} t^{\alpha-1} \log^k t \, dt \\ &= \sum_{j=0}^k \binom{k}{j} (-1)^j (\text{Pf } \partial^{k-j} \Gamma)(\alpha) \cdot z^{-\alpha} \cdot \log^j z + \delta_{-\alpha, \mathbb{Z}_+} \frac{(-1)^{k-1} \text{Res}_\alpha \Gamma}{k+1} \cdot z^{-\alpha} \cdot \log^{k+1} z. \end{aligned}$$

This gives Eq. (3.32) and (3.33). \square

Proposition 3.11. *Let L be a self-adjoint bounded below operator in the Hilbert space \mathcal{H} with resolvent in the \mathfrak{p}^{th} -Schatten class. Moreover, assume that the trace of the heat semigroup has an asymptotic expansion Eq. (3.6). Let $N \in \mathbb{N}$, $N \geq \mathfrak{p}$. Then $\text{Tr}\left((L+z)^{-N}\right)$ has the following asymptotic expansion as $\Re z \rightarrow \infty$*

$$\begin{aligned} \text{Tr}\left((L+z)^{-N}\right) &\sim \sum_{\alpha, k} (-1)^k \sum_{j \geq k} A_{\alpha k}^H(L) \frac{(\text{Pf } \partial^{j-k} \Gamma)(\alpha + N)}{(N-1)!} \cdot z^{-\alpha-N} \cdot \log^k z \\ &\quad + \sum_{m=0}^\infty \sum_{k=0}^{k_m} A_{-m-N, k}^H(L) \frac{(-1)^{k-1} \text{Res}_{-m-N} \Gamma}{k+1} \cdot z^{m+N} \cdot \log^{k+1} z. \end{aligned}$$

Proof. The result follows directly from the previous lemma and Eq. (3.28). \square

On the other hand, we can write the heat semigroup as a contour integral in terms of the resolvent,

$$e^{-tL} = t^{1-N} \frac{(N-1)!}{2\pi i} \int_C e^{-t\mu} \cdot (L - \mu)^{-N} d\mu, \quad (3.34)$$

where the contour C is depicted in the Figure 1.

Then one has, cf. e.g., [BL96, Lemma 2.2].

Proposition 3.12. *Let L be a self-adjoint bounded below operator in the Hilbert space \mathcal{H} with resolvent in the \mathfrak{p}^{th} -Schatten class. If the resolvent trace has the following asymptotic expansion*

$$\text{Tr}\left((L+z)^{-N}\right) \sim \sum_{\alpha, k} A_{\alpha k}^R(L) \cdot z^{-\alpha-N} \cdot \log^k z, \quad \text{as } \Re z \rightarrow \infty, z \in \Lambda_\epsilon, \quad (3.35)$$

for some $N \geq \mathfrak{p}$, then heat semigroup has the following asymptotic expansion as $t \rightarrow 0+$

$$\text{Tr}(e^{-tL}) \sim \sum_{\alpha, k} (-1)^{k+1} (N-1)! \sum_{j=k}^{k_\alpha} A_{\alpha j}^R(L) \binom{j}{j-k} \left(\text{Pf } \partial^{j-k} \frac{1}{\Gamma}\right) (-\alpha - N) \cdot t^\alpha \log^k t.$$




FIGURE 1. A graphic representation of the curve C.

We conclude this section observing that it is possible to obtain a result similar to Proposition 3.4 using the asymptotic expansion of the resolvent trace. This is done using the definition of the zeta function involving the resolvent trace,

$$\zeta(s; L) = \frac{\sin \pi s}{\pi} \cdot \frac{(N-1)! \cdot \Gamma(1+s-N)}{\Gamma(s)} \int_0^\infty z^{N-1-s} \cdot \text{Tr}((L+z)^{-N}) dz, \quad (3.36)$$

for some $N \geq p$ and $\Re s > N$. Here we used Eq. (3.34) and the definition of Euler Gamma function (cf. [GR07, 8.310-2]) to write the last equation.

4. THE FREDHOLM DETERMINANT AND THE ζ -DETERMINANT

4.1. Fredholm determinant of operators in some Schatten class. We first recall some facts on the Fredholm determinant. Under a discrete dimension spectrum assumption we will derive our main result on the relation between the Fredholm determinant and the zeta-regularized determinant. Our standard reference for Fredholm determinants and trace class operators are [Sim77] and [Sim05].

Recall from [Sim77, Chap. 3] that for a trace class operator A in the Hilbert space \mathcal{H}

$$\begin{aligned} \det(I + z \cdot A) &= \sum_{k=0}^{\infty} \text{Tr}(\Lambda^k A) \cdot z^k \\ &= \prod_{n=1}^{\infty} (1 + z \cdot \lambda_n(A)), \end{aligned} \quad (4.1)$$

where $\Lambda^k A$ is the induced operator on the exterior power $\Lambda^k \mathcal{H}$ and $\{\lambda_n(A)\}$ is the summable sequence of eigenvalues of A . The latter identity is equivalent to Lidskii's theorem [Sim77, Theorem 3.7].

4.1.1. Standing assumptions. During this subsection, we assume that we are given a self-adjoint operator L satisfying Eq. (3.1) and (3.2). Additionally, we assume from

now on that L is invertible. Denote by $\lambda_1 \leq \lambda_2 \leq \dots$ the sequence of eigenvalues of L with multiplicities. If $p = 1$ then L^{-1} is trace class and (4.1) implies

$$\det(I + L^{-1}) = \prod_{q=1}^{\infty} (1 + \lambda_q^{-1}). \quad (4.2)$$

Due to the fact that L^{-1} is trace class the right hand side is an absolutely convergent product.

Proposition 4.1. *Under the Standing assumptions 4.1.1, $z \mapsto \det(I + z \cdot L^{-1})$ is an entire function of order 1 with zeros exactly in $-\text{spec } L$. The order of the zeros $-\lambda$, $\lambda \in \text{spec } L$, equals the multiplicity of the eigenvalue λ . Moreover,*

$$\frac{d}{dz} \log \det(I + z \cdot L^{-1}) = \text{Tr}((L + z)^{-1}). \quad (4.3)$$

Proof. The first claim is standard, cf. [Sim77, Chap. 3], as the product expansion

$$\det(I + z \cdot L^{-1}) = \prod_{q=1}^{\infty} (1 + z \cdot \lambda_q^{-1})$$

is a canonical Weierstraß-product of an entire function of genus 0. Therefore, the logarithmic derivative may be taken term by term and the claim follows. \square

It was shown in [Sim77, Section 6] that the previous construction may be lifted to operators whose resolvent is in the p^{th} -Schatten class. In this case one obtains a canonical Weierstraß-product of higher genus.

Definition 4.2. *Suppose that $L^{-1} \in \mathbf{B}^p(\mathcal{H})$. Then for $N + 1 \geq p$ the $N + 1$ -Fredholm determinant of $I + L^{-1}$ is defined by*

$$\begin{aligned} \det_{N+1}(I + L^{-1}) &:= \prod_{q=1}^{\infty} \left((1 + \lambda_q^{-1}) \cdot \exp\left(\sum_{k=1}^N (-1)^k \frac{\lambda_q^{-k}}{k}\right) \right) \\ &= \det\left((I + L^{-1}) \cdot \exp\left(\sum_{k=1}^N (-1)^k \frac{L^{-k}}{k}\right) \right). \end{aligned}$$

Note that for $p = 1$, $\det_1(I + L^{-1}) = \det(I + L^{-1})$. The next theorem summarizes the main properties of the $N + 1$ -Fredholm determinant.

Theorem 4.3. *Let L be a bounded below invertible self-adjoint operator with $L^{-1} \in \mathbf{B}^p(\mathcal{H})$. Then for $N + 1 \geq p$ the function $z \mapsto \det_{N+1}(I + z \cdot L^{-1})$ is an entire function of order at most $N + 1$. Moreover, $f_{N+1}(z) := \log \det_{N+1}(I + z \cdot L^{-1})$ satisfies*

$$\frac{d}{dz} f_{N+1}(z) = (-z)^N \text{Tr}(L^{-N}(L + z)^{-1}), \quad (4.4)$$

$$f_{N+1}(0) = f'_{N+1}(0) = \dots = f_{N+1}^{(N)}(0) = 0, \quad (4.5)$$

and

$$\frac{d^{N+1}}{dz^{N+1}} f_{N+1}(z) = (-1)^N N! \text{Tr}((L + z)^{-(N+1)}). \quad (4.6)$$

Proof. It follows from the very definition that $f_{N+1}(z)$ is the logarithm of a canonical Weierstraß-product of genus N , see also [Sim77, Theorem 3.3 and Lemma 6.1]. Consequently, the logarithmic derivate is the compactly convergent sum of the logarithmic derivatives of the individual factors:

$$\frac{d}{dz} \log \det_{N+1}(I + z \cdot L^{-1}) = \sum_{q=1}^{\infty} \left((\lambda_q + z)^{-1} + \sum_{k=0}^{N-1} (-1)^{k+1} \left(\frac{z}{\lambda_q}\right)^k \cdot \lambda_q^{-1} \right).$$

The sum inside equals, for $z \neq \lambda_q$,

$$-\frac{1}{\lambda_q} \frac{\left(-\frac{z}{\lambda_q}\right)^N - 1}{-\frac{z}{\lambda_q} - 1} = \frac{\left(-\frac{z}{\lambda_q}\right)^N - 1}{z + \lambda_q},$$

hence

$$\frac{d}{dz} \log \det_{N+1}(I + z \cdot L^{-1}) = (-z)^N \operatorname{Tr}(L^{-N}(L + z)^{-1}),$$

thus Eq. (4.5) and (4.6) are proved. \square

It is a direct consequence of the last theorem that if L has discrete dimension spectrum then the Fredholm determinant of L has an asymptotic expansion, as $\Re z \rightarrow \infty$, that can be differentiated. On the other side, if the Fredholm determinant of L has an asymptotic expansion, as $\Re z \rightarrow \infty$, that can be differentiated then by Theorem 4.3 and Propositions 3.2 and 3.12, L has discrete dimension spectrum. We summarize this discussion in the next corollary (compare with [Spr06, Lemma 2.2]).

Corollary 4.4. *Let L be a self-adjoint bounded below operator in the Hilbert space \mathcal{H} satisfying Eq. (3.1) and (3.2). For L to have discrete dimension spectrum it is necessary and sufficient that the logarithm of the Fredholm determinant has an asymptotic expansion, as $\Re z \rightarrow \infty$, that can be differentiated.*

The explicit knowledge of the coefficients of the asymptotic expansion of the Fredholm determinant as $\Re z \rightarrow \infty$ implies the explicit knowledge of the coefficients of the heat expansion as $t \rightarrow 0+$. However, the converse of this implication is not true, *e.g.*, we will see below that the asymptotic expansion of the Fredholm determinant contains $\log \det_{\zeta} L$ which is a “global” invariant of the heat trace.

4.2. Asymptotic expansion of the Fredholm determinant. In this section we assume that \mathcal{H} is a separable Hilbert space and L is an invertible self-adjoint operator satisfying Eq. (3.1) and (3.2).

Now we come to one of our main results.

Theorem 4.5. *Let L be an invertible self-adjoint bounded below operator in the Hilbert space \mathcal{H} satisfying Eq. (3.1) and (3.2). Furthermore assume that L has discrete dimension spectrum as defined in Def. 3.1 and that the ζ -function $\zeta(s; L)$ of L is regular at 0 and has at most simple poles in $s = 1, \dots, p - 1$. Then*

$$\det_{\zeta}(L + z) = \exp\left(\sum_{j=0}^{p-1} \frac{z^j}{j!} \cdot \frac{d^j}{dz^j} \log \det_{\zeta}(L + z)|_{z=0}\right) \cdot \det_p(I + z \cdot L^{-1}). \quad (4.7)$$

Remark 4.6. Recall that the assumptions on the regularity at 0 and the simple poles in the positive integers are equivalent to the assumption that in the heat expansion $\Lambda_{\alpha,k}^H(L) = 0$ for $-\alpha \in \mathbb{Z}_+$ and $k > 0$, *i.e.*, we are under the Standing assumptions 3.2.1.

Proof. By Proposition 3.8, $\zeta(s; L + z)$ is holomorphic for (s, z) in a neighbourhood of $(0, 0)$. Again we commute the derivative in z with the derivative in s to obtain,

$$\frac{d^p}{dz^p} \log \det_{\zeta}(L + z) = -\frac{d^p}{dz^p} \left(\frac{d}{ds} \zeta(s; L + z)|_{s=0} \right) = -\frac{d}{ds} \left(\frac{d^p}{dz^p} \zeta(s; L + z) \right)|_{s=0}.$$

This implies that

$$\begin{aligned} -\frac{d}{ds} \frac{d^p}{dz^p} \zeta(s; L + z) &= \sum_{\lambda \in \text{spec } L} (-s-1) \dots (-s-p+1) (\lambda + z)^{-s-p} \\ &\quad - (-s) \sum_{\lambda \in \text{spec } L} \frac{d}{ds} \left((-s-1) \dots (-s-p+1) (\lambda + z)^{-s-p} \right). \end{aligned}$$

Therefore, using Theorem 4.3 we obtain

$$\begin{aligned} \frac{d^p}{dz^p} \log \det_{\zeta}(L + z) &= (-1)^{p-1} (p-1)! \text{Tr}(L + z)^{-p} \\ &= \frac{d^p}{dz^p} \log \det_p(I + z \cdot L^{-1}). \end{aligned}$$

Now the result follows from the series representation of the function $\log \det_{\zeta}(L + z)$ at $z = 0$. \square

The value at zero is the best we can obtain from the heat expansion (*cf.* Proposition 3.4). The ζ -regularized determinant will not only depend on the heat expansion. However, the asymptotic expansion of the Fredholm determinant delivers sufficient information to obtain the derivative of the ζ -regularized determinant. In [Fri89], the constant term of the asymptotic expansion of the Fredholm determinant as $\Re z \rightarrow \infty$ is determined for an elliptic differential operator, and it is equal to minus the logarithm of the ζ -regularized determinant of the operator. A similar result for a self-adjoint operator in Hilbert space with trace class resolvent is proved in [HLV17, pag. 3439]. Theorem 4.5 provides an extension of these results as we see in the next corollary.

Corollary 4.7. *Let L an invertible self-adjoint bounded below operator in the Hilbert space \mathcal{H} satisfying Eq. (3.1) and (3.2). Furthermore assume that L has discrete dimension spectrum as defined in Def. 3.1 and that the ζ -function of L is regular at 0 and has at most simple poles in $s = 1, \dots, p-1$. Then the logarithm of the Fredholm determinant*

of $I + z \cdot L^{-1}$ has an asymptotic expansion as follows

$$\begin{aligned}
& \log \det_p(I + z \cdot L^{-1}) \\
& \sim \sum_{\alpha, k} A_{\alpha k}^F(L) \cdot z^{-\alpha} \cdot \log^k z, \quad \text{as } \Re z \rightarrow \infty \\
& \sim - \sum_{\substack{\alpha, k \\ -\alpha \notin \mathbb{Z}_+}} (-1)^k \sum_{j \geq k} A_{\alpha j}^H(L) \binom{j}{k} \cdot (\text{Pf } \partial^{j-k} \Gamma)(\alpha) \cdot z^{-\alpha} \log z \\
& \quad - \log \det_\zeta(L) + A_{00}^H(L) \cdot \log z \\
& \quad + \sum_{n=1}^{\infty} A_{-n,0}^H(L) \cdot \frac{(-1)^n}{n!} \cdot z^n \log z + \sum_{n=p}^{\infty} A_{-n,0}^H(L) \cdot \frac{(-1)^{n+1}}{n!} \cdot L_n \cdot z^n \\
& \quad + \sum_{n=1}^{p-1} \frac{(-1)^{n+1}}{n} \cdot \left(\frac{A_{-n,0}^H}{n!} - (\text{Pf}_{s=n} \zeta(s; L)) \right) \cdot z^n, \quad \text{as } \Re z \rightarrow \infty.
\end{aligned}$$

In particular⁴,

$$A_{00}^F(L) = -\log \det_\zeta(L) = \gamma \cdot A_{00}^H(L) + \int_0^\infty t^{-1} \text{Tr}(e^{-tL}) dt, \quad (4.8)$$

$$A_{01}^F(L) = \zeta(0; L) = A_{00}^H(L). \quad (4.9)$$

Proof. From Theorem 4.5 we have that

$$\log \det_p(I + z \cdot L^{-1}) = \log \det_\zeta(L + z) - \sum_{j=0}^{p-1} \frac{z^j}{j!} \cdot \frac{d^j}{dz^j} \log \det_\zeta(L + z)|_{z=0}.$$

Now we apply Theorem 3.7, Proposition 3.8 and obtain the asymptotic expansion of the Fredholm determinant as $\Re z \rightarrow \infty$. \square

REFERENCES

- [BFK92] D. Burghelea, L. Friedlander, and T. Kappeler, *Meyer-Vietoris type formula for determinants of elliptic differential operators*, J. Funct. Anal. **107** (1992), no. 1, 34–65. MR 1165865
- [BL96] Jochen Brüning and Matthias Lesch, *On the spectral geometry of algebraic curves*, J. Reine Angew. Math. **474** (1996), 25–66. MR 1390691
- [BL99] ———, *On the η -invariant of certain nonlocal boundary value problems*, Duke Math. J. **96** (1999), no. 2, 425–468. MR 1666570
- [CM95] A. Connes and H. Moscovici, *The local index formula in noncommutative geometry*, Geom. Funct. Anal. **5** (1995), no. 2, 174–243. MR 1334867
- [Fri89] Leonid Friedlander, *The asymptotics of the determinant function for a class of operators*, Proc. Amer. Math. Soc. **107** (1989), no. 1, 169–178. MR 975642
- [GK19a] Fritz Gesztesy and Klaus Kirsten, *Computing traces, determinants, and ζ -functions for Sturm-Liouville operators: a survey*, Analysis and operator theory, Springer Optim. Appl., vol. 146, Springer, Cham, 2019, pp. 93–148. MR 3965220
- [GK19b] ———, *Effective computation of traces, determinants, and ζ -functions for Sturm-Liouville operators*, J. Funct. Anal. **276** (2019), no. 2, 520–562. MR 3906283

⁴The first two equalities of Eq. (4.8) and (4.9) were proved in [Spr06, Proposition 2.6] assuming the asymptotic expansion of $\log \det_p(I + z \cdot L^{-1})$ as $z \rightarrow \infty$.

- [GK19c] ———, *On traces and modified Fredholm determinants for half-line Schrödinger operators with purely discrete spectra*, *Quart. Appl. Math.* **77** (2019), no. 3, 615–630. MR 3962585
- [GR07] I. S. Gradshteyn and I. M. Ryzhik, *Table of integrals, series, and products*, seventh ed., Elsevier/Academic Press, Amsterdam, 2007, Translated from the Russian, Translation edited and with a preface by Alan Jeffrey and Daniel Zwillinger, With one CD-ROM (Windows, Macintosh and UNIX). MR 2360010
- [HLV17] Luiz Hartmann, Matthias Lesch, and Boris Vertman, *Zeta-determinants of Sturm-Liouville operators with quadratic potentials at infinity*, *J. Differential Equations* **262** (2017), no. 5, 3431–3465. MR 3584897
- [Hör90] Lars Hörmander, *An introduction to complex analysis in several variables*, third ed., North-Holland Mathematical Library, vol. 7, North-Holland Publishing Co., Amsterdam, 1990. MR 1045639
- [Kat95] Tosio Kato, *Perturbation theory for linear operators*, *Classics in Mathematics*, Springer-Verlag, Berlin, 1995, Reprint of the 1980 edition. MR 1335452
- [Les97] Matthias Lesch, *Operators of Fuchs type, conical singularities, and asymptotic methods*, *Teubner-Texte zur Mathematik [Teubner Texts in Mathematics]*, vol. 136, B. G. Teubner Verlagsgesellschaft mbH, Stuttgart, 1997. MR 1449639 (98d:58174)
- [Les98] ———, *Determinants of regular singular Sturm-Liouville operators*, *Math. Nachr.* **194** (1998), 139–170. MR 1653090 (99j:58220)
- [Les13] ———, *A gluing formula for the analytic torsion on singular spaces*, *Anal. PDE* **6** (2013), no. 1, 221–256. MR 3068545
- [LT98] Matthias Lesch and Jürgen Tolksdorf, *On the determinant of one-dimensional elliptic boundary value problems*, *Comm. Math. Phys.* **193** (1998), no. 3, 643–660. MR 1624851 (2000b:58060)
- [LV11] Matthias Lesch and Boris Vertman, *Regular singular Sturm–Liouville operators and their zeta-determinants*, *J. Funct. Anal.* **261** (2011), no. 2, 408–450. MR 2793118
- [LV15] ———, *Regularizing infinite sums of zeta-determinants*, *Math. Ann.* **361** (2015), no. 3-4, 835–862. MR 3319550
- [RS71] D. B. Ray and I. M. Singer, *R-torsion and the Laplacian on Riemannian manifolds*, *Advances in Math.* **7** (1971), 145–210. MR 295381
- [Sim77] Barry Simon, *Notes on infinite determinants of Hilbert space operators*, *Advances in Math.* **24** (1977), no. 3, 244–273. MR 0482328
- [Sim05] ———, *Trace ideals and their applications*, second ed., *Mathematical Surveys and Monographs*, vol. 120, American Mathematical Society, Providence, RI, 2005. MR 2154153
- [Spro6] Mauro Spreafico, *Zeta invariants for sequences of spectral type, special functions and the Lerch formula*, *Proc. Roy. Soc. Edinburgh Sect. A* **136** (2006), no. 4, 863–887. MR 2250451
- [Wat18] G. N. Watson, *The Harmonic Functions Associated with the Parabolic Cylinder*, *Proc. London Math. Soc. (2)* **17** (1918), 116–148. MR 1575566
- [Woj99] Krzysztof P. Wojciechowski, *The ζ -determinant and the additivity of the η -invariant on the smooth, self-adjoint Grassmannian*, *Comm. Math. Phys.* **201** (1999), no. 2, 423–444. MR 1682214

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE SÃO CARLOS (UFSCAR), BRAZIL

Email address: hartmann@dm.ufscar.br, luizhartmann@ufscar.br

URL: <http://www.dm.ufscar.br/profs/hartmann>

UNIVERSITÄT BONN, GERMANY

Email address: ml@matthiaslesch.de, lesch@math.uni-bonn.de

URL: www.matthiaslesch.de, www.math.uni-bonn.de/people/lesch