PULL-BACK OF SINGULAR LEVI-FLAT HYPERSURFACES

ANDRÉS BELTRÁN, ARTURO FERNÁNDEZ-PÉREZ, AND HERNÁN NECIOSUP

ABSTRACT. We study singular real analytic Levi-flat subsets invariant by singular holomorphic foliations in complex projective spaces. We give sufficient conditions for a real analytic Levi-flat subset to be the pullback of a semianalytic Levi-flat hypersurface in a complex projective surface under a rational map or to be the pull-back of a real algebraic curve under a meromorphic function. In particular, we give an application to the case of a singular real analytic Levi-flat hypersurface. Our results improve previous ones due to Lebl and Bretas – Fernández-Pérez – Mol.

1. INTRODUCTION AND STATEMENT OF THE RESULTS

Let M be a complex manifold of $\dim_{\mathbb{C}} M = N \geq 2$, a closed subset $H \subset M$ is a real analytic subvariety if for every $p \in H$, there are real analytic functions with real values $\varphi_1, \ldots, \varphi_k$ defined in a neighborhood $U \subset M$ of p, such that $H \cap U$ is equal to the set where all $\varphi_1, \ldots, \varphi_k$ vanish. A complex subvariety is precisely the same notion, considering holomorphic functions instead of real analytic functions. We say that a real analytic subvariety His *irreducible* if whenever we write $H = H_1 \cup H_2$ for two subvarieties H_1 and H_2 of M, then either $H_1 = H$ or $H_2 = H$. If H is irreducible, it has a welldefined dimension $\dim_{\mathbb{R}} H$. Let H_{reg} denote its regular part, i.e., the subset of points near which H is a real analytic submanifold of dimension equal to $\dim_{\mathbb{R}} H$. A set is *semianalytic* if it is locally constructed from real analytic sets by finite union, finite intersection, and complement. For a real analytic subvariety H, the set $\overline{H_{reg}}$ is a semianalytic subset where the closure is with the standard topology. In general, the inclusion $\overline{H_{reg}} \subset H$ is proper, which happens, for instance in the Whitney umbrella. We really only study the set H_{reg} , in this sense, we consider $Sing(H) := H_{reg} \setminus H_{reg}$ as the singular set of H, this is not the usual definition of the singular set in the literature, see for instance [15].

If $H \subset M$ is a real analytic hypersurface i.e., a real analytic subvariety of real codimension one, then for each $p \in H_{reg}$, there is a unique complex hyperplane $\mathscr{L}_p \subset T_p H_{reg}$. This defines a real analytic distribution $p \mapsto \mathscr{L}_p$

²⁰¹⁰ Mathematics Subject Classification. Primary 32V40 - 32S65.

Key words and phrases. Levi-flat subsets - Holomorphic foliations.

This work was supported by the Pontificia Universidad Católica del Perú project VRI-DGI-2018-0024. The second author is partially supported by CNPq-Brazil Grant Number 302790/2019-5.

2 ANDRÉS BELTRÁN, ARTURO FERNÁNDEZ-PÉREZ, AND HERNÁN NECIOSUP

of complex hyperplanes in TH_{req} . When this distribution is *integrable* in the sense of Frobenius, we say that H is Levi-flat. Here, H_{reg} is foliated by codimension one immersed complex submanifolds. This foliation, denoted by \mathscr{L} , is known as Levi foliation. According to Cartan [4], \mathscr{L} can be extended to a non-singular holomorphic foliation in a neighborhood of H_{reg} in M, but in general, it is not possible to extend \mathscr{L} to a singular holomorphic foliation in a neighborhood of H. There are examples of singular Levi-flat hypersurfaces whose Levi foliations extend to singular holomorphic webs in the ambient space, see for instance [8] and [21]. When there is a singular holomorphic foliation \mathscr{F} in the ambient space M that coincide with the Levi foliation on H_{reg} , we say either that H is *invariant* by \mathscr{F} or that \mathscr{F} is tangent to H. Cerveau and Lins Neto [6] proved that germs of singular foliations of codimension one at $(\mathbb{C}^N, 0)$ tangent to real analytic Levi-flat hypersurfaces have meromorphic (possibly holomorphic) first integrals. We recall that a non-constant function f is the *first integral* for a foliation \mathscr{F} if each leaf of \mathcal{F} is contained in a level set of f. In the global context, the same problem has been studied in [1] and [9].

The aim of this paper is to study holomorphic foliations tangent to real analytic Levi-flat subsets in complex manifolds. An irreducible real analytic subvariety $H \subset M$, where M is an N-dimensional complex manifold, $N \geq 2$, is a Levi-flat subset if it has real dimension 2n+1 and its regular part H_{reg} is foliated by immersed complex manifolds of complex dimension n. Similarly to the case of hypersurfaces, this foliation is called *Levi foliation* of H and will be denoted by \mathscr{L} . The number n is the Levi dimension of H. We use the qualifier "Levi" for the foliation, its leaves, and its dimension. Since we deal with real analytic Levi-flat subsets in complex manifolds we shall consider that H is *coherent*. Coherence implies that H admits a global complexification [11, p. 40]. Here coherent means that its ideal sheaf $\mathcal{I}(H)$ in $\mathcal{A}_{\mathbb{R},M}$, the sheaf of germs of real analytic functions with real values in M, is a coherent sheaf of $\mathcal{A}_{\mathbb{R},M}$ -modules. It follows from Oka's theorem [17, p. 94 Proposition 5] that H is coherent if the sheaf $\mathcal{I}(H)$ is locally finitely generated, the latter means that for every point $p \in H$ there exists an open neighborhood $U \subset M$ and a finite number of functions φ_i , real analytic in U and vanishing on H, such that for any $q \in U$, the germs of φ_j at q generate the ideal $\mathcal{I}(H_q)$, where H_q is the germ of H at q. We remark that not every real analytic subset is coherent as we shall see in Section 3 of this paper.

In [3], singular Levi-flat subsets appear in the result of the lifting of a real analytic Levi-flat hypersurface to the projectivized cotangent bundle of the ambient space through the Levi foliation and in [20], the authors gave a complete characterization of discritical singularities of local Levi-flat subsets in terms of their Segre varieties.

Let Y be a complex projective surface, $T \subset Y$ be a real analytic Leviflat hypersurface, $X \subset \mathbb{P}^N$, $N \geq 3$, be a complex projective subvariety of complex dimension k < N and $\rho : X \dashrightarrow Y$ be a dominant rational map. Then it is easy to show that $H = \overline{\rho^{-1}(T)}$ is a real analytic Levi-flat subset in \mathbb{P}^N and so H is a Levi-flat subset defined via pull-back. Therefore, one natural question is:

Given a real analytic Levi-flat subset $H \subset \mathbb{P}^N$. Under what condition, H is given by the pull-back of a Levi-flat hypersurface in a projective complex surface via a rational map?

In [14], Lebl gave sufficient conditions for a real analytic Levi-flat hypersurface in \mathbb{P}^N to be a pull-back of a real algebraic curve in \mathbb{C} via a meromorphic function. In [2], Bretas et al. proved an analogous result for real analytic Levi-flat subsets in \mathbb{P}^N . The main hypothesis in these articles is that the Levi foliation has infinitely many algebraic leaves. In this paper, we give an answer to the question, assuming that H is invariant by a singular holomorphic foliation on \mathbb{P}^N with quasi-invariant subvarieties (see Section 2). An irreducible complex subvariety $S \subset X$ of complex dimension n is quasi-invariant by a global n-dimensional foliation \mathscr{F} on a complex projective manifold X if it is not \mathscr{F} -invariant, but the restriction to the foliation \mathscr{F} to S is an algebraically integrable foliation of dimension n-1, i.e. every leaf of $\mathscr{F}|_S$ is algebraic. The concept of quasi-invariant subvarieties was introduced by Pereira-Spicer [19] for codimension one holomorphic foliations on complex projective manifolds to prove a variant of the classical Darboux-Jouanolou Theorem. Here we shall use this concept for Levi foliations to prove our main result:

Theorem 1. Let $H \subset \mathbb{P}^N$, $N \geq 3$, be an irreducible real analytic Levi-flat subset of Levi dimension n invariant by an n-dimensional singular holomorphic foliation \mathscr{F} on \mathbb{P}^N . Suppose that H is coherent and n > N/2. If the Levi foliation has infinitely many quasi-invariant subvarieties of complex dimension n, then there exists a unique projective subvariety X of complex dimension n + 1 containing H such that either there exists a rational map $R: X \dashrightarrow \mathbb{P}^1$, and real algebraic curve $C \subset \mathbb{P}^1$ such that $\overline{H_{reg}} \subset \overline{R^{-1}(C)}$ or there exists a dominant rational map $\rho: X \dashrightarrow Y$ on a projective surface Y and a semianalytic Levi-flat subset $T \subset Y$ such that $\overline{H_{reg}} \subset \overline{\rho^{-1}(T)}$.

We emphasize that the hypothesis n > N/2 implies that H is necessarily a real analytic subvariety with singularities. In fact, Ni-Wolfson [18, Theorem 2.4] proved that no nonsingular real analytic Levi-flat subset of the Levi dimension n exist in \mathbb{P}^N , n > N/2.

Applying Theorem 1 to n = N - 1, we get the following corollary:

Corollary 1. Let $H \subset \mathbb{P}^N$, $N \geq 3$, be an irreducible coherent real analytic Levi-flat hypersurface invariant by a codimension one holomorphic foliation \mathscr{F} on \mathbb{P}^N . If the Levi foliation has infinitely many quasi-invariant complex hypersurfaces, then either there exists a rational map $R : \mathbb{P}^N \dashrightarrow \mathbb{P}^1$, and real algebraic curve $C \subset \mathbb{P}^1$ such that $\overline{H_{reg}} \subset \overline{R^{-1}(C)}$ or there exists a dominant rational map $\rho : \mathbb{P}^N \dashrightarrow Y$ on a projective surface Y and a semianalytic Levi-flat subset $T \subset Y$ such that $\overline{H_{reg}} \subset \overline{\rho^{-1}(T)}$.

4 ANDRÉS BELTRÁN, ARTURO FERNÁNDEZ-PÉREZ, AND HERNÁN NECIOSUP

When H is a real analytic hypersurface, the above corollary gives a nice characterization of coherent real analytic Levi-flat hypersurfaces in \mathbb{P}^N , $N \geq 3$, invariant by codimension one holomorphic foliations which admit infinitely many quasi-invariant complex hypersurfaces. Observe that, in order to improve our results, we need to extend the Levi foliation of a Levi-flat subset to a holomorphic foliation in the ambient space. Therefore, another interesting question is:

Given a real analytic Levi-flat subset $H \subset \mathbb{P}^N$ with Levi foliation \mathcal{L} . Under what condition, \mathcal{L} extend to a singular holomorphic foliation on \mathbb{P}^N ?

When H is a local real analytic Levi-flat hypersurface, Lebl solved the above question in the non-dicritical case in [15].

The paper is organized as follows: in Section 2, we define the concept of quasi-invariant subvarieties of a foliation with complex leaves and state the main result of [19], such a result is key to prove Theorem 1. Section 3 is devoted to the study of real analytic Levi-flat subset in complex manifolds, using some results of [3] and [2], we prove the algebraic extension of the intrinsic complexification of H. In Section 4, we prove Theorem 1 and in Section 5 we prove Corollary 1. Finally, in Section 6, we give two examples. The first is an example of a Levi-flat hypersurface where Theorem 1 applies. In the second example, we construct a Levi-flat hypersurface in \mathbb{P}^3 that is not a pull-back of a Levi-flat hypersurface of \mathbb{P}^2 under a rational map. Moreover, this example also is not a pull-back of a real algebraic curve under a meromorphic function.

2. Foliations with complex leaves and quasi-invariant subvarieties

2.1. Foliations with complex leaves. A foliation with complex leaves of complex dimension n is a smooth foliation \mathcal{G} of dimension 2n whose local models are domains $U = W \times B$ of $\mathbb{C}^n \times \mathbb{R}^k$, $W \subset \mathbb{C}^n$, $B \subset \mathbb{R}^k$ and whose local transformations are of the form

(1)
$$\varphi(z,t) = (f(z,t), h(t)),$$

where f is holomorphic with respect to z. A domain U as above is said to be a distinguished coordinate domain of \mathcal{G} and $z = (z_1, \ldots, z_n), t = (t_1, \ldots, t_k)$ are said to be distinguished local coordinates. As examples of such foliations we have the Levi foliations of Levi-flat hypersurfaces of \mathbb{C}^n , see for instance [5] and [10].

If we replace \mathbb{R}^k by \mathbb{C}^k and in (1) we assume $t \in \mathbb{C}^k$ and that f, h are holomorphic with respect to z, t then we get the notion of holomorphic foliation of complex codimension k.

Now we define foliations with singularities. Let M be a complex manifold. A singular foliation with complex leaves \mathcal{G} of dimension n on M is a foliation with complex leaves of dimension n on $M \setminus E$, where E is a real analytic subvariety of M of real dimension < 2n. A point $p \in E$ is called a *removable* singularity of \mathcal{G} of there is a chart (U, φ) around p, compatible with the atlas \mathcal{A} of \mathcal{G} restricted to $M \setminus E$, in the sense that $\varphi \circ \varphi_i^{-1}$ and $\varphi_i \circ \varphi^{-1}$ have the form (1) for all $(U_i, \varphi_i) \in \mathcal{A}$ with $U \cap U_i \neq \emptyset$. The set of non-removable singularities of \mathcal{G} in E is called the *singular set* of \mathcal{G} , and is denoted by $\operatorname{Sing}(\mathcal{G})$.

2.2. Quasi-invariant subvarieties. Let Z be a projective manifold of complex dimension $N \ge 2$ and let \mathscr{G} be a foliation with complex leaves of dimension n on Z.

Definition 2.1. We say that \mathscr{G} is an *algebraically integrable foliation* on Z if every leaf of \mathscr{G} is algebraic, i.e. every leaf of \mathscr{G} is a projective complex subvariety in Z.

Motivated by [19], we define the concept of a *subvariety quasi-invariant* by a real analytic foliation with complex leaves.

Definition 2.2. An irreducible subvariety $S \subset Z$ of complex dimension n is *quasi-invariant* by a foliation \mathscr{G} if it is not \mathscr{G} -invariant, but the restriction of the foliation \mathscr{G} to S is an algebraically integrable foliation.

We note that the restriction foliation $\mathscr{G}|_S$ is a codimension one foliation on S and when $\mathscr{G}|_S$ is an algebraically integrable foliation, we have that every leaf of $\mathscr{G}|_S$ are projective complex hypersurfaces in S. Codimension one holomorphic foliations on Z which admit infinitely many quasi-invariant hypersurfaces have been studied in [19] and its main result is the following.

Theorem 2.1 (Pereira-Spicer [19]). Let \mathscr{F} be a codimension one holomorphic foliation on a projective manifold Z. If \mathscr{F} admits infinitely many quasiinvariant hypersurfaces then either \mathscr{F} is an algebraically integrable foliation, or \mathscr{F} is a pull-back of a foliation of dimension one on a projective surface under a dominant rational map.

3. Real analytic subsets

3.1. Coherent real analytic subsets. We present some of the fundamental results concerning coherent real analytic subsets.

Let H be a real analytic subset in an open set $U \subset \mathbb{C}^n$ and let $\mathcal{I}(H)$ be its ideal sheaf, it is the sheaf of germs of real analytic functions with real values vanishing on H.

Definition 3.1. *H* is said to be coherent if $\mathcal{I}(H)$ is a coherent sheaf of $\mathcal{A}_{\mathbb{R},U}$ -modules, where $\mathcal{A}_{\mathbb{R},U}$ is the sheaf of germs of real analytic functions with real values in *U*.

Proposition 3.1. [17, p. 95] If H is a coherent real analytic subset and the germ H_p of H at p is irreducible, then for q near p, we have

 $\dim_{\mathbb{R}} H_p = \dim_{\mathbb{R}} H_q.$

6 ANDRÉS BELTRÁN, ARTURO FERNÁNDEZ-PÉREZ, AND HERNÁN NECIOSUP

It is well known that locally, a real analytic subset always admits a complexification (see for instance [11, p. 40]) and it is not true for global real analytic subsets. It is shown in [11, p. 54] that the global complexification of a coherent real analytic subset in a complex manifold always exists.

Theorem 3.2. [11, p. 54] A real analytic subset in a complex manifold is coherent if and only if it admits a global complexification.

Now we build an irreducible real analytic hypersurface in \mathbb{P}^3 which is not coherent. Let $[z_0 : z_1 : z_2 : z_3]$ be the homogeneous coordinates in \mathbb{P}^3 and set $H \subset \mathbb{P}^3$ be the complex cone whose equation is

$$H = \{ (z_3\bar{z}_0 + \bar{z}_3 z_0) \left((z_1\bar{z}_0 + \bar{z}_1 z_0)^2 + (z_2\bar{z}_0 + \bar{z}_2 z_0)^2 \right) - (z_1\bar{z}_0 + \bar{z}_1 z_0)^3 = 0 \}.$$

The germ H_p of H at p = [1:0:0:0] is irreducible and of real dimension 5 at p. However, in a neighborhood of [1:0:0:z], $z \neq 0$, H reduces to the complex line $z_1 = z_2 = 0$, which is of real dimension 2. By Proposition 3.1, it follows that H is not coherent.

3.2. Levi-flat subset in complex manifolds. We give a brief resume of definitions and some known results about real analytic Levi-flat subsets in complex manifolds. Let H be an irreducible real analytic Levi-flat subset of Levi dimension n in an N-dimensional complex manifold M. The notion of Levi-flat subset germifies and, in general, we do not distinguish a germ at $(\mathbb{C}^N, 0)$ from its realization in some neighborhood U of $0 \in \mathbb{C}^N$. If $p \in H_{reg}$ then, according to [2, Proposition 3.1], there exists a holomorphic coordinate system $z = (z', z'') \in \mathbb{C}^{n+1} \times \mathbb{C}^{N-n-1}$ such that $z(p) = 0 \in \mathbb{C}^N$ and the germ of H at p is defined by

(2) $H = \{ z = (z', z'') \in \mathbb{C}^{n+1} \times \mathbb{C}^{N-n-1} : \operatorname{Im}(z_{n+1}) = 0, z'' = 0 \},\$

where $z' = (z_1, ..., z_{n+1})$ and $z'' = (z_{n+2}, ..., z_N)$ and the Levi foliation is given by

 $\{z = (z', z'') \in \mathbb{C}^{n+1} \times \mathbb{C}^{N-n-1} : z_{n+1} = c, z'' = 0, \text{ with } c \in \mathbb{R}\}.$

This trivial model is, in fact, a local form for a non-singular real analytic Levi-flat subset. Note that in the local form (2), $\{z''=0\}$ corresponds to the unique local (n + 1)-dimensional complex subvariety of the ambient space containing the germ of H_{reg} at p. These local subvarieties glue together forming a complex variety defined in a whole neighborhood of H_{reg} . It is analytically extendable to a neighborhood of $\overline{H_{reg}}$ by the following theorem:

Theorem 3.3 (Brunella [3]). Let M be an N-dimensional complex manifold and $H \subset M$ be a real analytic Levi-flat subset of Levi dimension n. Then, there exists a neighborhood $V \subset M$ of $\overline{H_{reg}}$ and a unique complex variety $X \subset V$ of dimension n + 1 containing H.

The variety X is the realization in the neighborhood V of a germ of complex analytic variety around H. We denote it — or its germ — by H^i and call it *intrinsic complexification* or *i-complexification* of H. It plays a central role in the theory of real analytic Levi-flat subsets. The notion of intrinsic complexification also appears in [22] with the name of the *Segre* envelope. If H is invariant by a holomorphic foliation on M, the same holds for its *i*-complexification, see for instance [2, Proposition 3.3].

Proposition 3.4. Let $H \subset M$ be a real analytic Levi-flat subset of Levi dimension n, where M is a complex manifold of dimension N. If H is invariant by an n-dimensional holomorphic foliation \mathcal{F} on M, then its icomplexification H^i is also invariant by \mathcal{F} .

As a consequence, if we denote by $\mathscr{F}^i := \mathscr{F}|_{H^i}$ (the restriction of \mathscr{F} to H^i), we have \mathscr{F}^i has codimension one in H^i . The following proposition shows the importance of the assumption of the *coherence* of a Levi-flat subset.

Proposition 3.5. [2, Proposition 3.6] Let M be an N-dimensional complex manifold and $H \subset M$ be an irreducible real analytic Levi-flat subset of Levi dimension n. Suppose that H is coherent. Then, there exist an open neighborhood $V \subset M$ of H and a unique irreducible complex subvariety X of V of complex dimension n + 1 containing H.

The variety X is the small variety of complex dimension n+1 that contains H. Again, let us denote this variety by H^i , the intrinsic complexification of H.

3.3. Levi-flat subsets in complex projective spaces. In this subsection, we state some results of real analytic Levi-flat subset in \mathbb{P}^N . Let $\sigma : \mathbb{C}^{N+1} \to \mathbb{P}^N$ be the natural projection. Suppose that H is a real-analytic subvariety of \mathbb{P}^N . Define the set $\tau(H)$ to be the set of points $z \in \mathbb{C}^{N+1}$ such that $\sigma(z) \in H$ or z = 0. A real analytic subvariety $H \subset \mathbb{P}^N$ is said to be algebraic if $H = \sigma(V)$ for some real algebraic complex cone V in \mathbb{C}^{N+1} . A set V is a complex cone when $p \in V$ implies $\lambda p \in V$ for all $\lambda \in \mathbb{C}$.

The following construction offers several examples of Levi-flat subsets in \mathbb{P}^N .

Proposition 3.6. [2, Proposition 6.1] Let $X \subset \mathbb{P}^N$ be an irreducible (n+1)dimensional algebraic variety, R be a rational function in X and $C \subset \mathbb{P}^1$ be a real algebraic one-dimensional subvariety. Then the set $\overline{R^{-1}(C)}$ is a real algebraic Levi-flat subset of Levi dimension n whose i-complexification is X

When we add the hypothesis that the Levi-flat subset is invariant by a singular holomorphic foliation in the ambient space, we can state a reciprocal result.

Proposition 3.7. [2, Proposition 6.3] Let \mathscr{F} be a singular holomorphic foliation in \mathbb{P}^N tangent to a real analytic Levi-flat subset H of Levi dimension n. Suppose that H is coherent and its i-complexification extends to an algebraic subvariety H^i in \mathbb{P}^N . If \mathscr{F}^i has a rational first integral R, then there exists a real algebraic one-dimensional subvariety $C \subset \mathbb{P}^1$ such that $\overline{H_{reg}} \subset \overline{R^{-1}(C)}$.

Now, since H is coherent, the intrinsic complexification H^i is well-defined as a complex subvariety in a neighborhood of H. Our aim is to extend H^i to an algebraic subvariety in \mathbb{P}^N . To get this, we use the following extension theorem.

Theorem 3.8 (Chow [7]). Let $Z \subset \mathbb{P}^N$ be a complex algebraic subvariety of dimension k and V be a connected neighborhood of Z in \mathbb{P}^N . Then any complex analytic subvariety of dimension higher than N - k in V that intersects Z extends algebraically to \mathbb{P}^N .

Under certain hypotheses, we can prove that the *i*-complexification H^i can be extended to \mathbb{P}^N .

Proposition 3.9. Let $H \subset \mathbb{P}^N$, $N \geq 3$, be an irreducible coherent real analytic Levi-flat subset of Levi dimension n such that n > N/2. If the Levi foliation \mathscr{L} has a quasi-invariant complex algebraic subvariety of complex dimension n, then H^i extends algebraically to \mathbb{P}^N .

Proof. Denote by L such quasi-invariant algebraic complex subvariety with $\dim_{\mathbb{C}} L = n - 1$. Since L algebraic with $L \subset H^i$ and $\dim_{\mathbb{C}} H^i = n + 1 > N - (n-1)$, we can apply Theorem 3.8 to prove that H^i extends algebraically to \mathbb{P}^N .

To end this section, we shall prove the following proposition.

Proposition 3.10. Let $H \subset \mathbb{P}^N$ be an irreducible coherent real analytic Levi-flat subset of Levi dimension n invariant by an n-dimensional singular holomorphic foliation \mathscr{F} in \mathbb{P}^N . Suppose that the i-complexification H^i extends to an algebraic variety in \mathbb{P}^N . If the Levi-foliation \mathcal{L} has infinitely many quasi-invariant algebraic subvarieties of complex dimension n-1. Then, either the foliation $\mathscr{F}^i = \mathscr{F}|_{H^i}$ has a rational first integral in H^i , or \mathscr{F}^i is a pull-back of a foliation on a projective surface under a dominant rational map.

Proof. First of all, we need to desingularize the *i*-complexification H^i . According to Hironaka desingularization theorem, there exist a complex manifold $\widetilde{H^i}$ and a proper bimeromorphic morphism $\pi : \widetilde{H^i} \to H^i$ such that

- (1) $\pi: \widetilde{H^i} \setminus (\pi^{-1}(\operatorname{Sing}(H^i)) \to H^i \setminus \operatorname{Sing}(H^i)$ is a biholomorphism,
- (2) $\pi^{-1}(\operatorname{Sing}(H^i))$ is a simple normal crossing divisor.

Since H^i is compact then $\widetilde{H^i}$ is too. We lift \mathscr{F}^i to an *n*-dimensional singular holomorphic foliation $\widetilde{\mathscr{F}^i}$ on $\widetilde{H^i}$. Since $\dim_{\mathbb{C}} \widetilde{H^i} = n + 1$, we have $\widetilde{\mathscr{F}^i}$ has codimension one on $\widetilde{H^i}$ and the tangency condition between \mathscr{F}^i and H implies that \mathscr{F}^i has infinitely many quasi-invariant closed subvarieties (these are algebraic and of codimension one in $\widetilde{H^i}$). Thus the same holds for $\widetilde{\mathscr{F}^i}$. By Theorem 2.1, either $\widetilde{\mathscr{F}^i}$ has a rational first integral or there exist a dominant rational map $\tilde{\rho} : \widetilde{H^{i}} \dashrightarrow Y$, where Y is a projective complex surface, \mathcal{G} is a foliation by curves on Y and $\widetilde{\mathscr{F}^{i}} = \tilde{\rho}^{*}(\mathcal{G})$. If $\widetilde{\mathscr{F}^{i}}$ admits a rational first integral in $\widetilde{H^{i}}$, then all leaves of $\widetilde{\mathscr{F}^{i}}$ are compact and so their π -images are compact leaves of \mathscr{F}^{i} in H^{i} . Applying Gómez-Mont's theorem [12], we have that there exists a one-dimensional projective manifold S and a rational map $f : H^{i} \dashrightarrow S$ whose fibers contain the leaves of \mathscr{F}^{i} . A rational first integral is obtained by composing f with any non-constant rational map $r : S \dashrightarrow \mathbb{P}^{1}$. If $\widetilde{\mathscr{F}^{i}}$ is a pull-back of a foliation \mathcal{G} on a projective complex surface Y under a dominant rational map $\tilde{\rho} : \widetilde{H^{i}} \dashrightarrow Y$ then \mathscr{F}^{i} is the pull-back of \mathcal{G} under $\rho := \tilde{\rho} \circ \pi^{-1} : H^{i} \dashrightarrow Y$, since π is a birational map. \Box

4. Proof of Theorem 1

With all the above results, we can prove Theorem 1.

Theorem 1. Let $H \subset \mathbb{P}^N$, $N \geq 3$, be an irreducible real analytic Levi-flat subset of Levi dimension n invariant by an n-dimensional singular holomorphic foliation \mathscr{F} on \mathbb{P}^N . Suppose that H is coherent and n > N/2. If the Levi foliation has infinitely many quasi-invariant subvarieties of complex dimension n, then there exists a unique projective subvariety X of complex dimension n + 1 containing H such that either there exists a rational map $R: X \dashrightarrow \mathbb{P}^1$, and real algebraic curve $C \subset \mathbb{P}^1$ such that $\overline{H_{reg}} \subset \overline{R^{-1}(C)}$ or there exists a dominant rational map $\rho: X \dashrightarrow Y$ on a projective surface Y and a semianalytic Levi-flat subset $T \subset Y$ such that $\overline{H_{reg}} \subset \overline{\rho^{-1}(T)}$.

Proof. By Proposition 3.5, there exist an open neighborhood $V \subset \mathbb{P}^N$ of Hand a unique irreducible complex subvariety H^i of V of complex dimension n + 1 containing H. The Proposition 3.4 implies that H^i is invariant by \mathscr{F} and moreover it extends algebraically to \mathbb{P}^N by Proposition 3.9. We denote $\mathscr{F}^i := \mathscr{F}|_{H^i}$ the restrict foliation to H^i . Observe now that \mathscr{F}^i is a foliation of codimension one on H^i which admit infinitely many quasiinvariant subvarieties of complex dimension n - 1. Therefore, either \mathscr{F}^i has a rational first integral in H^i , or \mathscr{F}^i is a pull-back of a foliation on a projective surface under a dominant rational map by Proposition 3.10.

If \mathscr{F}^i has a first integral R then there exists a real algebraic curve $C \subset \mathbb{P}^1$ such that $\overline{H_{reg}} \subset \overline{R^{-1}(C)}$ by Proposition 3.7. Now if we assume that \mathscr{F}^i is a pull-back of a foliation \mathcal{G} on a projective complex surface Y under a dominant rational map $\rho : H^i \dashrightarrow Y$. Then we can take $X = H^i$. Let us prove that there exists a semianalytic Levi-flat subset $T \subset Y$. Indeed, let $z \in H_{reg} \setminus Ind(\rho)$ (here $Ind(\rho)$ denotes the indeterminacy set of ρ). Then there exists a neighborhood $U \subset H^i \setminus Ind(\rho)$ of z and a non-singular real analytic curve $\gamma : (-\epsilon, \epsilon) \to U$ such that $\gamma(0) = z$, $\{\gamma\} \subset H_{reg}$, and such that γ is transverse to the Levi foliation \mathscr{L} on H_{reg} . Let $L_{\gamma(t)}$ be the leaf of \mathscr{L} through $\gamma(t)$. Since $L_{\gamma(t)}$ is also a leaf of \mathscr{F}^i and $\mathscr{F}^i = \rho^*(\mathcal{G})$, then $\rho(L_{\gamma(t)})$ is a leaf of \mathcal{G} . Let us denote $A_t = \overline{\rho(L_{\gamma(t)})} \subset Y$ and define

$$T_z := \bigcup_{t \in (-\epsilon, \epsilon)} A_t \subset V_z,$$

where V_z is a neighborhood of T_z on Y. Note that T_z is a union of complex subvarieties parametrized by t such that each A_t contains leaves of \mathcal{G} , thus T_z is a semianalytic Levi-flat subset on V_z . These local constructions are sufficiently canonical to be patched together when z varies on H_{reg} : if $T_{z_1} \subset V_{z_1}$ and $T_{z_2} \subset V_{z_2}$ are as above, with $V_{z_1} \cap V_{z_2} \neq \emptyset$, then $T_{z_1} \cap V_{z_1} \cap V_{z_2}$ and $T_{z_2} \cap V_{z_1} \cap V_{z_2}$ have some common leaves of \mathcal{G} because \mathcal{G} is a global foliation defined on Y, so T_{z_1} and T_{z_2} can be glued by identifying these leaves. In this way, we get a semianalytic Levi-flat subset T in Y.

Finally, we assert that $\overline{H_{reg}} \subset \overline{\rho^{-1}(T)}$. In fact, let $w \in \overline{H_{reg}}$, then there exists a sequence $z_k \to w$, $z_k \in H_{reg}$, so $\rho(z_k) \in T$ which imply that $z_k \in \rho^{-1}(T)$ and $w \in \overline{\rho^{-1}(T)}$. This finishes the proof.

5. Proof of Corollary 1

Corollary 1. Let $H \subset \mathbb{P}^N$, $N \geq 3$, be an irreducible coherent real analytic Levi-flat hypersurface invariant by a codimension one holomorphic foliation \mathscr{F} on \mathbb{P}^N . If the Levi foliation has infinitely many quasi-invariant complex hypersurfaces, then either there exists a rational map $R : \mathbb{P}^N \dashrightarrow \mathbb{P}^1$, and real algebraic curve $C \subset \mathbb{P}^1$ such that $\overline{H_{reg}} \subset \overline{R^{-1}(C)}$ or there exists a dominant rational map $\rho : \mathbb{P}^N \dashrightarrow Y$ on a projective complex surface Y and a semianalytic Levi-flat subset $T \subset Y$ such that $\overline{H_{reg}} \subset \overline{\rho^{-1}(T)}$.

Proof. If H is an irreducible real analytic Levi-flat hypersurface in \mathbb{P}^N , $N \geq 3$, then the Levi dimension of H is N-1. Moreover

$$N-1 > N/2 \iff N > 2.$$

Thus, we can apply Theorem 1 to H, so there exist a unique projective subvariety X of complex dimension N containing H such that either there exists a rational map $R: X \dashrightarrow \mathbb{C}$, and real algebraic curve $C \subset \mathbb{C}$ such that $\overline{H_{reg}} \subset \overline{R^{-1}(C)}$ or there exists a dominant rational map $\rho: X \dashrightarrow Y$ on a projective complex surface Y and a semianalytic Levi-flat subset $T \subset Y$ such that $\overline{H_{reg}} \subset \overline{\rho^{-1}(T)}$. Since $X \subset \mathbb{P}^N$ has complex dimension N, we must have $X = \mathbb{P}^N$ and hence we conclude the proof. \Box

6. Examples

Example 6.1. We give an example of a real analytic Levi-flat hypersurface in \mathbb{P}^3 where Theorem 1 applies. Let

$$H = \{ [z_0 : z_1 : z_2 : z_3] \in \mathbb{P}^3 : z_0 z_1 \bar{z}_2 \bar{z}_3 - z_2 z_3 \bar{z}_0 \bar{z}_1 = 0 \},\$$

then H is Levi-flat because it is foliated by the complex hypersurfaces

(3)
$$z_0 z_1 = c z_2 z_3$$
, where $c \in \mathbb{R}$.

Let $\mathscr F$ be the codimension one holomorphic foliation on $\mathbb P^3$ of degree two defined by

$$\omega = z_1 z_2 z_3 dz_0 + z_0 z_2 z_3 dz_1 - z_0 z_1 z_3 dz_2 - z_0 z_1 z_2 dz_3,$$

then \mathscr{F} has a rational first integral $R: \mathbb{P}^3 \dashrightarrow \mathbb{P}^1$ given by

$$R[z_0:z_1:z_2:z_3] = [z_0z_1:z_2z_3].$$

Since the leaves of $\mathscr{F}|_H$ coincide with the leaves of the Levi foliation (3), H must be invariant by \mathscr{F} . On the other hand, note that $H = \overline{R^{-1}(C)}$, where

$$C = \{ [t:u] \in \mathbb{P}^1 : t\bar{u} - u\bar{t} = 0 \}.$$

Example 6.2. In the following example, we construct a real analytic Leviflat hypersurface H in \mathbb{P}^3 that is not a pull-back of a Levi-flat hypersurface of \mathbb{P}^2 under a rational map, furthermore, H also is not a pull-back of a real algebraic curve under a meromorphic function.

Consider $z = (z_0, z_1, z_2, z_3), \ \bar{z} = (\bar{z}_0, \bar{z}_1, \bar{z}_2, \bar{z}_3)$ and

$$F(z,\bar{z}) = \det \begin{pmatrix} z_0 & z_1 & z_2 & z_3 & 0 & 0 \\ 0 & z_0 & z_1 & z_2 & z_3 & 0 \\ 0 & 0 & z_0 & z_1 & z_2 & z_3 \\ \bar{z}_0 & \bar{z}_1 & \bar{z}_2 & \bar{z}_3 & 0 & 0 \\ 0 & \bar{z}_0 & \bar{z}_1 & \bar{z}_2 & \bar{z}_3 & 0 \\ 0 & 0 & \bar{z}_0 & \bar{z}_1 & \bar{z}_2 & \bar{z}_3 \end{pmatrix}$$

Define $H = \{[z_0 : z_1 : z_2 : z_3] \in \mathbb{P}^3 : F(z, \overline{z}) = 0\}$, H is a real analytic hypersurface well defined since F is a bihomogeneous polynomial of bi-degree (3,3). Moreover, H is Levi-flat, because it is foliated by the complex hyperplanes

(4)
$$z_0 + cz_1 + c^2 z_2 + c^3 z_3 = 0$$
, where $c \in \mathbb{R}$.

Let \mathscr{W} be the codimension one holomorphic 3-web on \mathbb{P}^3 given by the implicit differential equation $\Omega = 0$,

$$\Omega = \det \begin{pmatrix} z_0 & z_1 & z_2 & z_3 & 0 & 0 \\ 0 & z_0 & z_1 & z_2 & z_3 & 0 \\ 0 & 0 & z_0 & z_1 & z_2 & z_3 \\ dz_0 & dz_1 & dz_2 & dz_3 & 0 & 0 \\ 0 & dz_0 & dz_1 & dz_2 & dz_3 & 0 \\ 0 & 0 & dz_0 & dz_1 & dz_2 & dz_3 \end{pmatrix}$$

Since the leaves of $\mathscr{W}|_H$ and \mathscr{L} are the same, we get H is invariant by \mathscr{W} .

Now, we prove that H is not a pull-back of a Levi-flat hypersurface of \mathbb{P}^2 . To prove this fact, we use the following result of [13, Proposition 4.4]:

Proposition 6.1. Let ω_1 , ω_2 and ω_3 be independent germs of integrable 1forms at $(\mathbb{C}^3, 0)$ with singular sets of codimension at least two. Suppose that there exists a non-zero holomorphic 2-form η , locally decomposable outside its singular set, that is tangent to each ω_i , for i = 1, 2, 3. Then ω_1, ω_2 and ω_3 define foliations that are in a pencil. Furthermore, η is integrable, defining the axis foliation of this pencil.

Suppose by contradiction that H is a pull-back of a Levi-flat hypersurface under a dominant rational map $\rho : \mathbb{P}^3 \dashrightarrow \mathbb{P}^2$. Then pick a point $p \in U_0$, where U_0 is an open subset in \mathbb{P}^3 such that $\rho|_{U_0} : U_0 \subset \mathbb{C}^3 \to \mathbb{C}^2$ is a holomorphic submersion. We may have needed to perhaps move to yet another point $p' \in U_0$ such that U_0 does not intersect the discriminant set of the web \mathscr{W} . We set p = p' and works in a neighborhood of U_0 . Therefore, the germ of \mathscr{W} at p is a decomposable 3-web, defined by the superposition of three independent foliations $\mathscr{F}_1, \mathscr{F}_2$, and \mathscr{F}_3 . We can assume that these foliations are defined by independent germs of integrable 1-forms ω_1, ω_2 , and ω_3 respectively. Since H is given by a pull-back, all the leaves of \mathscr{L} and, hence the leaves of \mathscr{W} in $H \cap U_0$ are tangent to the fibers of $\rho|_{U_0}$, these fibers define a non-zero holomorphic 2-form η_ρ that is tangent to each ω_i , for i = 1, 2, 3. Then, according to Proposition 6.1, ω_1, ω_2 , and ω_3 define foliations that are in a pencil, an absurd. Hence, the assertion is proved.

Now we assert that H is not a pull-back of a real algebraic curve under a meromorphic function. In fact, H is a Levi-flat hypersurface in \mathbb{P}^3 such that there does not exist a point contained in infinitely many leaves of \mathscr{L} , because, the leaves of \mathscr{L} are given by the equation (4) and through at a point only pass three leaves. If H is defined by a pull-back of a meromorphic function, there has to exist a point p of indeterminacy since the dimension is at least 2. Then through at p pass infinitely many leaves of \mathscr{L} . Since Hdoes not satisfy this property, we finish the proof of the assertion.

Acknowledgments. The authors wish to express his gratitude to Maria Aparecida Soares Ruas (ICMC - USP, São Carlos) and Judith Brinkschulte (Universität Leipzig) for many valuable conversations and suggestions.

References

- Beltrán, A., Fernández-Pérez, A., and Neciosup, H.: Existence of dicritical singularities of Levi-flat hypersurfaces and holomorphic foliations. Geometriae Dedicata (2017), doi.org/10.1007/s10711-017-0303-4
- [2] Bretas, J., Fernández-Pérez, A., and Mol, R.: Holomorphic foliations tangent to Levi-flat subsets. J. Geom. Anal. (2019) 29: 1407. https://doi.org/10.1007/s12220-018-0043-1
- [3] Brunella, M.: Singular Levi-flat hypersurfaces and codimension one foliations. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) vol. VI, no. 4 (2007), 661-672.
- [4] Cartan, E.: Sur la géométrie pseudo-conforme des hypersurfaces de l'espace de deux variables complexes. Ann. Mat. Pura Appl., 11 (1) (1933), 17-90.
- [5] Cerveau, D., and Sad, P.: Fonctions et feuilletages Levi-flat. ètude locale. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5), 3(2):427-445, (2004).
- [6] Cerveau, D., and Lins Neto, A.: Local Levi-flat hypersurfaces invariants by a codimension one holomorphic foliation. American Journal of Mathematics, vol. 133 no. 3, (2011), 677-716. doi.org/10.1353/ajm.2011.0018

- [7] Chow, W.L.: On meromorphic maps of algebraic varieties. Ann. Math. 2 (89), 391-403 (1969).
- [8] Fernández-Pérez, A.: On Levi-flat hypersurfaces with generic real singular set. J. Geom. Anal. (2013) 23: 2020. doi.org/10.1007/s12220-012-9317-1
- [9] Fernández-Pérez, A.: Levi-flat hypersurfaces tangent to projective foliations. J Geom. Anal. (2014) 24: 1959. doi.org/10.1007/s12220-013-9404-y
- [10] Fernández-Pérez, A., Mol, R., and Rosas, R.: On singular real analytic Levi-flat foliations. To be published: The Asian Journal of Mathematics, 2021.
- [11] Guaraldo, F., Macri P., and Tancredi, A.: Topics on real analytic spaces. Springer-Verlag, (2013).
- [12] Gómez-Mont, X.: Integrals for holomorphic foliations with singularities having all leaves compact. Ann. Inst. Fourier 39(2), 451-458 (1989).
- [13] Junca, D., and Mol, R.: Holomorphic vector fields tangent to foliations in dimension three. To be published: An. Acad. Brasil. Ciênc., 2021.
- [14] Lebl, J.: Algebraic Levi-flat hypervarieties in complex projective space. J. Geom. Anal. (2012) 22: 410. https://doi.org/10.1007/s12220-010-9201-9
- [15] Lebl, J.: Singular set of a Levi-flat hypersurface is Levi-flat. Math. Ann. (2013) 355: 1177. doi.org/10.1007/s00208-012-0821-1
- [16] Lebl, J.: Singular Levi-flat hypersurfaces in complex projective space induced by curves in the Grassmannian. Internat. J. Math. 26 (2015), no. 5, 1550036, 17 pp.
- [17] Narasimhan, R.: Introduction to the theory of analytic spaces. Lectures Notes in Mathematics. Springer-Verlag Berlin Heidelberg (1966). doi.org/10.1007/BFb0077071
- [18] Ni, L., and Wolfson, J.: The Lefschetz theorem for CR submanifolds and the nonexistence of real analytic Levi flat submanifolds. Communications in Analysis and Geometry. Volume 11, Number 3, 553-564, (2003).
- [19] Pereira, J.V., and Spicer C.: Hypersurfaces quasi-invariant by codimension one foliations. Mathematishe Annalen (2019). doi 10.1007/s00208-019-01833-4
- [20] Pinchuk, S., Shafikov, R., and Sukhov, A.: On dicritical singularities of Levi-flat sets. Ark. Mat., 56 (2018), 395-408. DOI: 10.4310/ARKIV.2018.v56.n2.a12
- [21] Shafikov, R., and Sukhov, A.: Germs of singular Levi-flat hypersurfaces and holomorphic foliations. Comment. Math. Helv. 90 (2015), 479-502.
- [22] Sukhov, A.: Levi-flat world: a survey of local theory. Ufimsk. Mat. Zh., (2017), Volume 9, Issue 3, 172-185.

(A. Beltrán) DPTO. CIENCIAS - SECCIÓN MATEMÁTICAS, PONTIFÍCIA UNIVERSIDAD CATÓLICA DEL PERÚ.

Current address: Av. Universitaria 1801, San Miguel, Lima 32, Peru Email address: abeltra@pucp.pe

(A. Fernández-Pérez) DEPARTAMENTO DE MATEMÁTICA - ICEX, UNIVERSIDADE FEDERAL DE MINAS GERAIS, UFMG

Current address: Av. Antônio Carlos 6627, 31270-901, Belo Horizonte-MG, Brasil. Email address: fernandez@ufmg.br

(H. Neciosup) DPTO. CIENCIAS - SECCIÓN MATEMÁTICAS, PONTIFÍCIA UNIVERSIDAD CATÓLICA DEL PERÚ.

Current address: Av. Universitaria 1801, San Miguel, Lima 32, Peru Email address: hneciosup@pucp.pe