

# A NOTE ON DEGENERATE GENERALIZED LAGUERRE POLYNOMIALS AND LAH NUMBERS

TAEKYUN KIM, DMITRY V. DOLGY, DAE SAN KIM, HYE KYUNG KIM, AND SEONG HO PARK

ABSTRACT. The aim of this paper is to introduce the degenerate generalized Laguerre polynomials as the degenerate version of the generalized Laguerre polynomials and to derive some properties related to those polynomials and Lah numbers, including an explicit expression, a Rodrigues' type formula and expressions for the derivatives. The novelty of the present paper is that it is the first paper on degenerate versions of orthogonal polynomials.

## 1. INTRODUCTION

The generalized Laguerre polynomials are classical orthogonal polynomials which are orthogonal with respect to the gamma distribution  $e^{-x}x^\alpha dx$  on the interval  $(0, \infty)$ . The generalized Laguerre polynomials are widely used in many problems of quantum mechanics, mathematical physics and engineering. In quantum mechanics, the Schrödinger equation for the hydrogen-like atom is exactly solvable by separation of variables in spherical coordinates. The radial part of the wave function is a generalized Laguerre polynomial [14]. In mathematical physics, vibronic transitions in the Franck-Condon approximation can also be described by using Laguerre polynomials [6]. In engineering, the wave equation is solved for the time domain electric field integral equation for arbitrary shaped conducting structures by expressing the transient behaviors in terms of Laguerre polynomials [4].

The aim of this paper is to introduce the degenerate generalized Laguerre polynomials as the degenerate version of the generalized Laguerre polynomials and to derive some properties related to those polynomials and Lah numbers. In more detail, we obtain an explicit formula and a Rodrigues' type formula for the degenerate Laguerre polynomials. We also get explicit expressions for the degenerate generalized Laguerre polynomial for  $\alpha = -1$ , an identity involving Lah numbers, the falling factorial moment of the degenerate Poisson random variable with parameter  $\alpha$ , and expressions for the derivatives of the degenerate generalized Laguerre polynomials.

We should mention here that degenerate versions of many special numbers and polynomials have been explored and many interesting results have obtained in recent years [8,10,12]. Furthermore, these have been done not only for special numbers and polynomials but also for transcendental functions like gamma functions [11]. The novelty of the present paper is that this is the first paper which treats degenerate versions of orthogonal polynomials. For the rest of this section, we will recall some necessary facts that will be used throughout this paper.

The Laguerre polynomial  $L_n(x)$  satisfies the second-order linear differential equation

$$xy'' + (1-x)y' + ny = 0, \quad (\text{see [16]}),$$

while the generalized Laguerre polynomial (or the associated Laguerre polynomial)  $L_n^{(\alpha)}(x)$  satisfies the second-order linear differential equation

$$xy'' + (\alpha + 1 - x)y' + ny = 0, \quad (\alpha \in \mathbb{R}).$$

---

2010 *Mathematics Subject Classification.* 11B83; 42C05; 60E99.

*Key words and phrases.* degenerate generalized Laguerre polynomials; Lah numbers; degenerate exponential function.

The Rodrigues' formula of the Laguerre polynomial  $L_n(x)$  is given by

$$(1) \quad L_n(x) = \frac{e^x}{n!} \frac{d^n}{dx^n} (e^{-x} x^n) = \frac{1}{n!} \left( \frac{d}{dx} - 1 \right)^n x^n,$$

while that of the generalized Laguerre polynomial  $L_n^{(\alpha)}(x)$  is given by

$$(2) \quad \begin{aligned} L_n^{(\alpha)}(x) &= \frac{1}{n!} x^{-\alpha} e^x \frac{d^n}{dx^n} (e^{-x} x^{n+\alpha}) \\ &= x^{-\alpha} \frac{1}{n!} \left( \frac{d}{dx} - 1 \right)^n x^{n+\alpha}, \quad (\text{see [2, 9, 16, 17]}). \end{aligned}$$

The generating function of generalized Laguerre polynomials is given by

$$(3) \quad \sum_{n=0}^{\infty} L_n^{(\alpha)}(x) t^n = \frac{1}{(1-t)^{\alpha+1}} e^{-x \frac{t}{1-t}}, \quad (\text{see [9, 16, 17]}).$$

From (3), we get

$$(4) \quad L_n^{(\alpha)}(x) = \sum_{i=0}^n (-1)^i \binom{n+\alpha}{n-i} \frac{x^i}{i!}, \quad (\text{see [9, 17]}).$$

Note that

$$\begin{aligned} L_0^{(\alpha)}(x) &= 1 \\ L_1^{(\alpha)}(x) &= -x + (\alpha + 1) \\ L_2^{(\alpha)}(x) &= \frac{x^2}{2} - (\alpha + 2)x + \frac{(\alpha + 1)(\alpha + 2)}{2}, \\ L_3^{(\alpha)}(x) &= -\frac{x^3}{6} + \frac{\alpha + 3}{2}x^2 - \frac{(\alpha + 2)(\alpha + 3)}{2}x + \frac{(\alpha + 1)(\alpha + 2)(\alpha + 3)}{6}, \dots \end{aligned}$$

The rising factorial sequence is defined as

$$\langle x \rangle_0 = 1, \quad \langle x \rangle_n = x(x+1) \cdots (x+n-1), \quad (n \geq 1),$$

while the falling factorial sequence is defined as

$$(x)_0 = 1, \quad (x)_n = x(x-1) \cdots (x-n+1), \quad (n \geq 1), \quad (\text{see [1-3, 5, 7-13, 15-18]}).$$

We note that the Lah numbers are defined by

$$(5) \quad \langle x \rangle_n = \sum_{k=0}^n L(n, k) (x)_k, \quad (n \geq 0), \quad (\text{see [3, 5, 8, 13, 15]}).$$

From (5), we can easily derive the following equation.

$$(6) \quad \frac{1}{k!} \left( \frac{t}{1-t} \right)^k = \sum_{n=k}^{\infty} L(n, k) \frac{t^n}{n!}, \quad (\text{see [3, 5, 13, 17]}).$$

For any  $\lambda \in \mathbb{R}$ , the degenerate exponential function is defined by

$$(7) \quad e_{\lambda}^x(t) = \sum_{n=0}^{\infty} (x)_{n, \lambda} \frac{t^n}{n!}, \quad (\text{see [10, 11]})$$

where  $(x)_{0, \lambda} = 1$ ,  $(x)_{n, \lambda} = x(x-\lambda) \cdots (x-(n-1)\lambda)$ ,  $(n \geq 1)$ . For  $x = 1$ , we use the notation by  $e_{\lambda}(t) = e_{\lambda}^1(t)$ .

## 2. DEGENERATE GENERALIZED LAGUERRE POLYNOMIALS

For any  $\alpha \in \mathbb{R}$ , we consider the degenerate generalized Laguerre polynomials given by

$$(8) \quad \frac{1}{(1-t)^{\alpha+1}} e_{\lambda} \left( -x \frac{t}{1-t} \right) = \sum_{n=0}^{\infty} L_{n,\lambda}^{(\alpha)}(x) t^n, \quad |t| < 1.$$

From (7), we note that

$$(9) \quad \begin{aligned} \frac{1}{(1-t)^{\alpha+1}} e_{\lambda} \left( -x \frac{t}{1-t} \right) &= \frac{1}{(1-t)^{\alpha+1}} \sum_{m=0}^{\infty} (1)_{m,\lambda} (-1)^m x^m \frac{1}{m!} \left( \frac{t}{1-t} \right)^m \\ &= \sum_{m=0}^{\infty} (1)_{m,\lambda} (-1)^m x^m \frac{1}{m!} t^m \left( \frac{1}{1-t} \right)^{m+\alpha+1} \\ &= \sum_{m=0}^{\infty} (1)_{m,\lambda} (-1)^m x^m \frac{t^m}{m!} \sum_{l=0}^{\infty} \binom{m+\alpha+l}{l} t^l \\ &= \sum_{n=0}^{\infty} \left( \sum_{m=0}^n (1)_{m,\lambda} (-1)^m x^m \frac{1}{m!} \binom{m+\alpha+n-m}{n-m} \right) t^n \\ &= \sum_{n=0}^{\infty} \left( \sum_{m=0}^n (1)_{m,\lambda} (-1)^m x^m \frac{1}{m!} \binom{n+\alpha}{n-m} \right) t^n. \end{aligned}$$

Therefore, by (8) and (9), we obtain the following theorem.

**Theorem 1.** For  $n \geq 0$ , we have

$$L_{n,\lambda}^{(\alpha)}(x) = \sum_{m=0}^n \binom{n+\alpha}{n-m} (-1)^m (1)_{m,\lambda} \frac{1}{m!} x^m.$$

Now, by using Theorem 1, we observe that

$$(10) \quad \begin{aligned} \frac{d^n}{dx^n} \left[ x^{\alpha} e_{\lambda} \left( -\frac{a}{x} \right) \right] &= \frac{d^n}{dx^n} \left[ \sum_{k=0}^{\infty} (1)_{k,\lambda} (-a)^k \frac{1}{k!} x^{\alpha-k} \right] \\ &= \sum_{k=0}^{\infty} (1)_{k,\lambda} \frac{(-a)^k}{k!} \overbrace{(\alpha-k)(\alpha-k-1)\cdots(\alpha-k-n+1)}^{n\text{-times}} x^{\alpha-k-n} \\ &= (-1)^n x^{\alpha-n} \sum_{k=0}^{\infty} (1)_{k,\lambda} (-a)^k \frac{1}{k!} (k-\alpha)(k-\alpha+1)\cdots(k-\alpha+n-1) x^{-k} \\ &= (-1)^n x^{\alpha-n} n! \sum_{k=0}^{\infty} (1)_{k,\lambda} \frac{(-1)^k}{k!} \left( \frac{a}{x} \right)^k \binom{k+n-\alpha-1}{n} \\ &= (-1)^n x^{\alpha-n} n! \sum_{k=0}^{\infty} (1)_{k,\lambda} \frac{(-1)^k}{k!} \left( \frac{a}{x} \right)^k \sum_{l=0}^n \binom{n-\alpha-1}{n-l} \binom{k}{l} \\ &= (-1)^n x^{\alpha-n} n! \sum_{l=0}^n \binom{n-\alpha-1}{n-l} \sum_{k=l}^{\infty} (1)_{k,\lambda} \frac{(-1)^k}{k!} \left( \frac{a}{x} \right)^k \frac{k!}{l!(k-l)!} \\ &= (-1)^n x^{\alpha-n} n! \sum_{l=0}^n \binom{n-\alpha-1}{n-l} \sum_{k=0}^{\infty} (1)_{k+l,\lambda} (-1)^{k+l} \left( \frac{a}{x} \right)^{k+l} \frac{1}{l!k!} \\ &= (-1)^n x^{\alpha-n} n! \sum_{l=0}^n \binom{n-\alpha-1}{n-l} (-1)^l \left( \frac{a}{x} \right)^l \frac{1}{l!} (1)_{l,\lambda} \sum_{k=0}^{\infty} (1-l\lambda)_{k,\lambda} \frac{(-1)^k}{k!} \left( \frac{a}{x} \right)^k \end{aligned}$$

$$\begin{aligned}
&= (-1)^n x^{\alpha-n} n! \sum_{l=0}^n \binom{n-\alpha-1}{n-l} (-1)^l (1)_{l,\lambda} \left(\frac{a}{x}\right)^l \frac{1}{l!} e_{\lambda}^{1-l\lambda} \left(-\frac{a}{x}\right) \\
&= (-1)^n x^{\alpha-n} n! e_{\lambda} \left(-\frac{a}{x}\right) \sum_{l=0}^n \binom{n-\alpha-1}{n-l} (-1)^l (1)_{l,\lambda} \left(\frac{a}{x-a\lambda}\right)^l \frac{1}{l!} \\
&= (-1)^n x^{\alpha-n} n! e_{\lambda} \left(-\frac{a}{x}\right) L_{n,\lambda}^{(-\alpha-1)} \left(\frac{a}{x-a\lambda}\right).
\end{aligned}$$

Therefore, by (10), we obtain the following theorem.

**Theorem 2.** For  $n \geq 0$ , we have

$$\frac{d^n}{dx^n} \left[ x^{\alpha} e_{\lambda} \left(-\frac{a}{x}\right) \right] = (-1)^n x^{\alpha-n} n! e_{\lambda} \left(-\frac{a}{x}\right) L_{n,\lambda}^{(-\alpha-1)} \left(\frac{a}{x-a\lambda}\right).$$

By using Leibniz rule and Theorem 1, we have

$$\begin{aligned}
(11) \quad &\frac{d^n}{dx^n} [e_{\lambda}(-x)x^{n+\alpha}] \\
&= \sum_{m=0}^n \binom{n}{m} \left[ \frac{d^m}{dx^m} e_{\lambda}(-x) \right] \left[ \frac{d^{n-m}}{dx^{n-m}} x^{n+\alpha} \right] \\
&= \sum_{m=0}^n \binom{n}{m} (-1)^m (1)_{m,\lambda} e_{\lambda}^{1-m\lambda}(-x) \cdot (n+\alpha)_{n-m} x^{n+\alpha-n+m} \\
&= n! e_{\lambda}(-x) x^{\alpha} \sum_{m=0}^n \binom{n+\alpha}{n-m} (-1)^m (1)_{m,\lambda} e_{\lambda}^{-m\lambda}(-x) x^m \frac{1}{m!} \\
&= n! e_{\lambda}(-x) x^{\alpha} \sum_{m=0}^n \binom{n+\alpha}{n-m} (-1)^m (1)_{m,\lambda} \left(\frac{x}{1-\lambda x}\right)^m \frac{1}{m!} \\
&= n! e_{\lambda}(-x) x^{\alpha} L_{n,\lambda}^{(\alpha)} \left(\frac{x}{1-\lambda x}\right).
\end{aligned}$$

Thus, we obtain Rodrigues' type formula for the degenerate generalized Laguerre polynomials.

**Theorem 3** (Rodrigues' type formula). For  $n \geq 0$ , we have

$$\frac{x^{-\alpha}}{n! e_{\lambda}(-x)} \frac{d^n}{dx^n} [e_{\lambda}(-x)x^{n+\alpha}] = L_{n,\lambda}^{(\alpha)} \left(\frac{x}{1-\lambda x}\right).$$

For  $\alpha = -1$ , from Theorem 3, we have

$$(12) \quad \frac{x}{n! e_{\lambda}(-x)} \frac{d^n}{dx^n} [e_{\lambda}(-x)x^{n-1}] = L_{n,\lambda}^{(-1)} \left(\frac{x}{1-x}\right).$$

On the other hand, by (8), we get

$$(13) \quad e_{\lambda} \left(-x \frac{t}{1-t}\right) = \sum_{n=0}^{\infty} L_{n,\lambda}^{(-1)}(x) t^n.$$

From (7), we can derive the following equation.

$$\begin{aligned}
 (14) \quad e_\lambda \left( -x \frac{t}{1-t} \right) &= \sum_{k=0}^{\infty} (-1)^k (1)_{k,\lambda} x^k \frac{1}{k!} \left( \frac{t}{1-t} \right)^k \\
 &= \sum_{k=0}^{\infty} (-1)^k (1)_{k,\lambda} x^k \sum_{n=k}^{\infty} L(n,k) \frac{t^n}{n!} \\
 &= \sum_{n=0}^{\infty} \left( \sum_{k=0}^n (-1)^k (1)_{k,\lambda} x^k L(n,k) \right) \frac{t^n}{n!}.
 \end{aligned}$$

Thus, by (13) and (14), we get

$$(15) \quad L_{n,\lambda}^{(-1)}(x) = \frac{1}{n!} \sum_{k=0}^n (-x)^k (1)_{k,\lambda} L(n,k),$$

where  $L(n,k) = \binom{n-1}{k-1} \frac{n!}{k!}$  is the Lah number.

Therefore, we obtain the following theorem.

**Theorem 4.** For  $n \geq 0$ , we have

$$L_{n,\lambda}^{(-1)}(x) = \frac{1}{n!} \sum_{k=0}^n (-x)^k (1)_{k,\lambda} L(n,k) = \sum_{k=0}^n (1)_{k,\lambda} (-x)^k \frac{1}{k!} \binom{n-1}{k-1}.$$

From Theorem 1, we note that

$$\begin{aligned}
 (16) \quad L_{n,\lambda}^{(\alpha)}(x) &= \sum_{m=0}^n (1)_{m,\lambda} x^m (-1)^m \frac{1}{m!} \binom{n+\alpha}{n-m} \\
 &= \sum_{m=0}^n (1)_{m,\lambda} (-x)^m \frac{(n+\alpha)(n+\alpha-1)\cdots(m+\alpha+1)}{m!(n-m)!} \\
 &= \sum_{m=0}^n (1)_{m,\lambda} (-x)^m \frac{(n+\alpha)(n+\alpha-1)\cdots(m+\alpha+1)(m+\alpha)\cdots(\alpha+1)}{m!(n-m)!(m+\alpha)\cdots(\alpha+1)} \\
 &= \sum_{m=0}^n (1)_{m,\lambda} (-x)^m \frac{1}{m!(n-m)!} \frac{(n+\alpha)_n}{(m+\alpha)_m} \\
 &= \sum_{m=0}^n (1)_{m,\lambda} (-x)^m \frac{1}{m!(n-m)!} \frac{(n+\alpha)_n}{(m+\alpha)_m} \frac{\Gamma(\alpha+1)}{\Gamma(\alpha+1)} \\
 &= \Gamma(n+\alpha+1) \sum_{m=0}^n (1)_{m,\lambda} \frac{(-x)^m}{m!(n-m)!} \frac{1}{\Gamma(m+\alpha+1)}.
 \end{aligned}$$

Thus, by (16), we get

$$(17) \quad L_{n,\lambda}^{(\alpha)}(x) = \frac{\Gamma(n+\alpha+1)}{\Gamma(n+1)} \sum_{m=0}^n (1)_{m,\lambda} \binom{n}{m} (-x)^m \frac{1}{\Gamma(m+\alpha+1)}.$$

In particular,  $\alpha = -1$ , we have

$$\begin{aligned}
 (18) \quad L_{n,\lambda}^{(-1)}(x) &= \frac{1}{n} \sum_{m=0}^n (1)_{m,\lambda} \binom{n}{m} (-x)^m \frac{1}{\Gamma(m)} \\
 &= \sum_{m=1}^n (1)_{m,\lambda} (-x)^m \frac{1}{m!} \binom{n-1}{m-1}.
 \end{aligned}$$

Now, we observe that

$$\begin{aligned}
(19) \quad \frac{d^n}{dx^n} e_\lambda \left( \frac{1}{x} \right) &= \frac{d^n}{dx^n} \sum_{k=0}^{\infty} (1)_{k,\lambda} \frac{1}{k!} \left( \frac{1}{x} \right)^k \\
&= \sum_{k=0}^{\infty} (1)_{k,\lambda} \frac{1}{k!} (-1)^n \langle k \rangle_n x^{-n-k} \\
&= \sum_{k=0}^{\infty} (1)_{k,\lambda} \frac{(-1)^n}{k!} \sum_{l=0}^n L(n,l) \langle k \rangle_l x^{-n-k} \\
&= \sum_{l=0}^n (-1)^n L(n,l) \sum_{k=0}^{\infty} (1)_{k,\lambda} \frac{\langle k \rangle_l}{k!} x^{-n-k} \\
&= (-1)^n x^{-n} \sum_{l=0}^n L(n,l) \sum_{k=l}^{\infty} \frac{(1)_{k,\lambda}}{k!} \langle k \rangle_l x^{-k} \\
&= (-1)^n x^{-n} \sum_{l=0}^n L(n,l) \sum_{k=l}^{\infty} \frac{k!}{k!(k-l)!} (1)_{k,\lambda} x^{-k} \\
&= (-1)^n x^{-n} \sum_{l=0}^n L(n,l) x^{-l} \sum_{k=0}^{\infty} \frac{(1)_{k+l,\lambda}}{k!} x^{-k} \\
&= (-1)^n x^{-n} \sum_{l=0}^n L(n,l) x^{-l} (1)_{l,\lambda} \sum_{k=0}^{\infty} \frac{(1-l\lambda)_{k,\lambda}}{k!} x^{-k} \\
&= (-1)^n x^{-n} \sum_{l=0}^n L(n,l) x^{-l} (1)_{l,\lambda} e_\lambda^{1-l\lambda} \left( \frac{1}{x} \right) \\
&= (-1)^n x^{-n} e_\lambda \left( \frac{1}{x} \right) \sum_{l=0}^n L(n,l) (1)_{l,\lambda} x^{-l} \left( 1 + \frac{\lambda}{x} \right)^{-l} \\
&= (-1)^n x^{-n} e_\lambda \left( \frac{1}{x} \right) \sum_{l=0}^n L(n,l) (1)_{l,\lambda} \left( \frac{1}{x+\lambda} \right)^l.
\end{aligned}$$

Therefore, by (19), we obtain the following theorem.

**Theorem 5.** For  $n \geq 1$ , we have

$$\frac{d^n}{dx^n} e_\lambda \left( \frac{1}{x} \right) = (-1)^n x^{-n} e_\lambda \left( \frac{1}{x} \right) \sum_{l=0}^n L(n,l) (1)_{l,\lambda} \left( \frac{1}{x+\lambda} \right)^l.$$

Since

$$L(n,k) = \binom{n-1}{k-1} \frac{n!}{k!} = \binom{n-1}{k-1} \frac{n!}{(n-k)!k!} (n-k)! = \binom{n-1}{k-1} \binom{n}{k} (n-k)!.$$

Thus, we have the following corollary.

**Corollary 6.** For  $n \geq 1$ , we have

$$\frac{d^n}{dx^n} e_\lambda \left( \frac{1}{x} \right) = x^{-n} (-1)^n e_\lambda \left( \frac{1}{x} \right) \sum_{l=1}^n \binom{n-1}{l-1} \binom{n}{l} (1)_{l,\lambda} (n-l)! \left( \frac{1}{x+\lambda} \right)^l.$$

## 3. DEGENERATE POISSON RANDOM VARIABLES

Let  $X$  be the Poisson random variable with parameter  $\alpha(> 0)$ . Then the probability mass function of  $X$  is given by

$$p(i) = P\{X = i\} = \frac{\alpha^i}{i!} e^{-\alpha}, \quad (i = 0, 1, 2, \dots).$$

It is easy to show that

$$E[(X)_n] = \sum_{k=0}^{\infty} (k)_n p(k) = e^{-\alpha} \sum_{k=n}^{\infty} \frac{\alpha^k}{(k-n)!} = e^{-\alpha} \alpha^n \sum_{k=0}^{\infty} \frac{\alpha^k}{k!} = \alpha^n.$$

Thus, we note that

$$E\left[\binom{X}{n}\right] = \frac{\alpha^n}{n!}, \quad (n = 0, 1, 2, \dots).$$

Let  $X_\lambda$  be the degenerate Poisson random variable with parameter  $\alpha(> 0)$ . Then the probability mass function of  $X_\lambda$  is given by

$$p(i) = P\{X_\lambda = i\} = e_\lambda^{-1}(\alpha) \frac{\alpha^i}{i!} (1)_{i,\lambda}, \quad (i = 0, 1, 2, \dots), \quad (\text{see [12]}).$$

Then the following falling factorial moment is given by

$$\begin{aligned} (20) \quad E[(X_\lambda)_n] &= \sum_{k=0}^{\infty} (k)_n p(k) = \sum_{k=0}^{\infty} (k)_n \frac{e_\lambda^{-1}(\alpha)}{k!} \alpha^k (1)_{k,\lambda} \\ &= e_\lambda^{-1}(\alpha) \sum_{k=n}^{\infty} \frac{k(k-1)\cdots(k-n+1)(k-n)!}{k!(k-n)!} \alpha^k (1)_{k,\lambda} \\ &= e_\lambda^{-1}(\alpha) \sum_{k=0}^{\infty} \alpha^{k+n} \frac{1}{k!} (1)_{k+n,\lambda} \\ &= \alpha^n e_\lambda^{-1}(\alpha) \sum_{k=0}^{\infty} (1)_{n,\lambda} (1-n\lambda)_{k,\lambda} \frac{\alpha^k}{k!} \\ &= \alpha^n e_\lambda^{-1}(\alpha) (1)_{n,\lambda} e_\lambda^{1-n\lambda}(\alpha) = \alpha^n (1)_{n,\lambda} \left(\frac{1}{1+\alpha\lambda}\right)^n. \end{aligned}$$

Assume that  $X_\lambda$  is the Poisson random variable with parameter  $\frac{1}{\alpha}(> 0)$ . Then, by using (20), we obtain

$$\begin{aligned} \frac{d^n}{d\alpha^n} \left( e_\lambda \left( \frac{1}{\alpha} \right) \right) &= (-1)^n \alpha^{-n} \sum_{l=0}^n L(n,l) \sum_{k=0}^{\infty} \frac{(k)_l}{k!} (1)_{k,\lambda} \alpha^{-k} \\ &= (-1)^n \alpha^{-n} \sum_{l=0}^n L(n,l) e_\lambda \left( \frac{1}{\alpha} \right) e_\lambda^{-1} \left( \frac{1}{\alpha} \right) \sum_{k=0}^{\infty} \frac{(k)_l}{k!} (1)_{k,\lambda} \alpha^{-k} \\ &= (-1)^n \alpha^{-n} \sum_{l=0}^n L(n,l) e_\lambda \left( \frac{1}{\alpha} \right) E[(X_\lambda)_l] \\ &= (-1)^n \alpha^{-n} e_\lambda \left( \frac{1}{\alpha} \right) \sum_{l=0}^n L(n,l) (1)_{l,\lambda} \left( \frac{1}{\alpha + \lambda} \right)^l. \end{aligned}$$

## 4. DERIVATIVES OF DEGENERATE LAGUERRE POLYNOMIALS

Let us consider the sequence  $y_{n,\lambda}(x)$  which are given by

$$(21) \quad A(t)e_\lambda\left(-x\frac{t}{1-t}\right) = \sum_{n=0}^{\infty} y_{n,\lambda}(x)t^n,$$

where  $A(t)$  is an invertible series.

Note that  $y_{0,\lambda}(x) = A(0)$  is a constant. We now set  $F_\lambda = F_\lambda(x, t) = A(t)e_\lambda\left(-\frac{x}{1-t}t\right)$ .

From (21), we note that

$$(22) \quad \begin{aligned} \frac{\partial}{\partial x}F_\lambda &= A(t)\left(-\frac{t}{1-t}\right)e_\lambda^{1-\lambda}\left(-\frac{xt}{1-t}\right) \\ &= -\left(\frac{t}{(1-t)-x\lambda t}\right)A(t)e_\lambda\left(-\frac{xt}{1-t}\right). \end{aligned}$$

By (22), we get

$$(23) \quad \frac{\partial}{\partial x}F_\lambda - (1+x\lambda)t\frac{\partial}{\partial x}F_\lambda = -tF_\lambda.$$

From (21) and (23), we can derive the following equation

$$(24) \quad \sum_{n=1}^{\infty} y'_{n,\lambda}(x)t^n - (1+x\lambda)\sum_{n=1}^{\infty} y'_{n-1,\lambda}(x)t^n = -\sum_{n=1}^{\infty} y_{n-1,\lambda}(x)t^n.$$

By comparing the coefficients on both sides of (24), we get

$$(25) \quad y'_{n,\lambda}(x) - (1+x\lambda)y'_{n-1,\lambda}(x) = -y_{n-1,\lambda}(x), \quad (n \geq 1),$$

where  $y'_{n,\lambda}(x) = \frac{d}{dx}y_{n,\lambda}(x)$ .

Now, we observe that

$$(26) \quad \begin{aligned} &= \frac{1}{(1-t)-x\lambda t} = -\frac{t}{1-t}\left(\frac{1}{1-\frac{x\lambda}{1-t}t}\right) = -\sum_{l=0}^{\infty} x^l \lambda^l \left(\frac{t}{1-t}\right)^{l+1} \\ &= -\sum_{l=1}^{\infty} x^{l-1} \lambda^{l-1} \left(\frac{t}{1-t}\right)^l \\ &= -\sum_{l=1}^{\infty} x^{l-1} \lambda^{l-1} t^l \sum_{m=0}^{\infty} \binom{m+l-1}{m} t^m \\ &= -\sum_{k=1}^{\infty} \left(\sum_{l=1}^k x^{l-1} \lambda^{l-1} \binom{n-1}{n-l}\right) t^k. \end{aligned}$$

From (22) and (26), we can derive the following equation (27)

$$(27) \quad \begin{aligned} \sum_{n=1}^{\infty} y'_{n,\lambda}(x)t^n &= \frac{\partial}{\partial x}F_\lambda = -\left(\frac{t}{1-t-x\lambda t}\right)A(t)e_\lambda\left(-\frac{xt}{1-t}\right) \\ &= \sum_{k=1}^{\infty} \left(-\sum_{l=1}^k x^{l-1} \lambda^{l-1} \binom{k-1}{k-l}\right) t^k \sum_{m=0}^{\infty} y_{m,\lambda} t^m \\ &= \sum_{n=1}^{\infty} \left(-\sum_{k=1}^n \sum_{l=1}^k x^{l-1} \lambda^{l-1} \binom{k-1}{k-l}\right) y_{n-k,\lambda}(x) t^n. \end{aligned}$$



Thus, by comparing the coefficients on both sides of (27), we get

$$(28) \quad y'_{n,\lambda}(x) = - \sum_{k=1}^n \sum_{l=1}^k x^{l-1} \lambda^{l-1} \binom{k-1}{k-l} y_{n-k,\lambda}(x).$$

Therefore, we obtain the following theorem.

**Theorem 7.** *Let*

$$A(t)e_{\lambda} \left( -\frac{xt}{1-t} \right) = \sum_{n=0}^{\infty} y_{n,\lambda}(x)t^n,$$

where  $A(t)$  is an invertible series. For  $n \geq 1$ , we have

$$y'_{n,\lambda}(x) = (1+x\lambda)y'_{n-1,\lambda}(x) - y_{n-1,\lambda}(x)$$

and

$$y'_{n,\lambda}(x) = - \sum_{k=1}^n \sum_{l=1}^k x^{l-1} \lambda^{l-1} \binom{k-1}{k-l} y_{n-k,\lambda}(x),$$

where  $y'_n(x) = \frac{d}{dx}y_{n,\lambda}(x)$ .

Let us take  $A(t) = (1-t)^{-\alpha-1}$ . Then we have

$$(29) \quad \sum_{n=0}^{\infty} y_{n,\lambda}(x)t^n = (1-t)^{-\alpha-1} e_{\lambda} \left( -\frac{x}{1-x}t \right) = \sum_{n=0}^{\infty} L_{n,\lambda}^{(\alpha)}(x)t^n.$$

Thus, we note that  $y_{n,\lambda}(x) = L_{n,\lambda}^{(\alpha)}(x)$ , ( $n \geq 0$ ).

Therefore, by Theorem 6 and (29), we obtain the following corollary.

**Corollary 8.** *For  $n \geq 1$ , we have*

$$\frac{d}{dx}L_{n,\lambda}^{(\alpha)}(x) = (1+x\lambda) \frac{d}{dx}L_{n-1,\lambda}^{(\alpha)}(x) - L_{n-1,\lambda}^{(\alpha)}(x)$$

and

$$\frac{d}{dx}L_{n,\lambda}^{(\alpha)}(x) = - \sum_{k=1}^n \sum_{l=1}^k x^{l-1} \lambda^{l-1} \binom{k-1}{k-l} L_{n-k,\lambda}^{(\alpha)}(x).$$

## 5. CONCLUSION

In this paper, we introduced the degenerate generalized Laguerre polynomials, which are the first degenerate versions of the orthogonal polynomials, and derived some results related to those polynomials and Lah numbers. Some of the results are an explicit expression, Rodrigues' type formula and some expressions for the derivatives of the degenerate generalized Laguerre polynomials.

As one of our future projects, we would like to continue to study degenerate versions of many special numbers and polynomials.

**Acknowledgments:** Not applicable

**Funding:** This work was supported by the Basic Science Research Program, the National Research Foundation of Korea, (NRF-2021R1F1A1050151).

**Availability of data and materials:** No data were used to support this study.

**Competing interests:** The authors declare no conflict of interest.

**Authors' contributions:** The authors contributed equally to the work. All authors read and approved the final manuscript.

## REFERENCES

- [1] Bayad, A.; Chikhi, J. *Apostol-Euler polynomials and asymptotics for negative binomial reciprocals*, Adv. Stud. Contemp. Math. (Kyungshang) **24** (2014), no. 1, 33–37.
- [2] Boyadzhiev, K. N. Lah numbers, *Laguerre polynomials of order negative one, and the  $n$ th derivative of  $\exp(1/x)$* , Acta Univ. Sapientiae Math. **8** (2016), no. 1, 22–31.
- [3] Carlitz, L. *Some generating functions for Laguerre polynomials*, Duke Math. J. **35** (1968), 825–827.
- [4] Chung, Y.-S.; Sarkar, T. K.; Jung, B. H.; Salazar-Palma, M.; Ji, Z.; Jang, S.; Kim, K. *Solution of time domain electric field integral equation using the  $L$  polynomials*, in IEEE Transactions on Antennas and Propagation **52** (2004), no. 9, 2319–2328.
- [5] Comtet, L. *Advanced combinatorics*, D. Reidel Publishing Co., Dordrecht, 1974.
- [6] de Jong, M.; Seijo, L.; Meijerink, A.; Rabouw, F. T. *Resolving the ambiguity in the relation between Stokes shift and Huang-Rhys parameter*, Phys. Chem. Chem. Phys. **17** (2015), no. 26, 16959–16969.
- [7] Khan, W. A.; Araci, S.; Acikgoz, M.; Esi, A. *Laguerre-based Hermite-Bernoulli polynomials associated with bilateral series*, Tbilisi Math. J. **11** (2018), no. 2, 111–121.
- [8] Kim, H. K.; Lee, D. S. *Some identities of degenerate  $r$ -extended Lah-Bell polynomials*, Proc. Jangjeon Math. Soc. **24** (2021), no. 1, 47–61.
- [9] Kim, T. *Identities involving Laguerre polynomials derived from umbral calculus*, Russ. J. Math. Phys. **21** (2014), no. 1, 36–45.
- [10] Kim, T.; Kim, D. S. *Degenerate Zero-Truncated Poisson Random Variables*, Russ. J. Math. Phys. **28** (2021), no. 1, 66–72.
- [11] Kim, T.; Kim, D. S. *Note on the Degenerate Gamma Function*, Russ. J. Math. Phys. **27** (2020), no. 3, 352–358.
- [12] Kim, T.; Kim, D. S.; Jang, L.-C.; Kim, H. Y. *A note on discrete degenerate random variables*, Proc. Jangjeon Math. Soc. **23** (2020), no. 1, 125–135.
- [13] Kim, T.; Kim, D. S.; Jang, L.-C.; Lee, H.; Kim, H.-Y. *Complete and incomplete Bell polynomials associated with Lah-Bell numbers and polynomials*, Adv. Difference Equ. 2021, Paper No. 101, 12 pp.
- [14] Mawhin, J.; Ronveaux, A. *Schrödinger and Dirac equations for the hydrogen atom, and Laguerre polynomials*, Arch. Hist. Exact Sci. **64** (2010), 429–460.
- [15] Quaintance, J.; Gould, H. W. *Combinatorial identities for Stirling numbers*, The unpublished notes of H. W. Gould. With a foreword by George E. Andrews. World Scientific Publishing Co. Pte. Ltd., Singapore, 2016.
- [16] Rainville, E. D. *Special functions*, Reprint of 1960 first edition. Chelsea Publishing Co., Bronx, N.Y., 1971.
- [17] Roman, S. *The umbral calculus*, Pure and Applied Mathematics, 111. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, 1984.
- [18] Simsek, Y. *Identities and relations related to combinatorial numbers and polynomials*, Proc. Jangjeon Math. Soc. **20** (2017), no. 1, 127–135.

DEPARTMENT OF MATHEMATICS, KWANGWOON UNIVERSITY, SEOUL 139-701, REPUBLIC OF KOREA  
*Email address:* kwangwoonmath@hanmail.net

DEPARTMENT OF MATHEMATICAL METHODS IN ECONOMY, FAR EASTERN FEDERAL UNIVERSITY, 690950  
 VLADIVOSTOK, RUSSIA  
*Email address:* dvdolgy@gmail.com

DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SEOUL 121-742, REPUBLIC OF KOREA  
*Email address:* dskim@sogang.ac.kr

DEPARTMENT OF MATHEMATICS EDUCATION, DAEGU CATHOLIC UNIVERSITY, GYEONGSAN 38430, REPUBLIC OF KOREA  
*Email address:* hkkim@cu.ac.kr

DEPARTMENT OF MATHEMATICS, KWANGWOON UNIVERSITY, SEOUL 139-701, REPUBLIC OF KOREA  
*Email address:* abcd2938471@kw.ac.kr