

PROJECTIONS IN MODULI SPACES OF KLEINIAN GROUPS

HALA ALAQAD, JIANHUA GONG, AND GAVEN MARTIN

ABSTRACT. A two-generator Kleinian group $\langle f, g \rangle$ can be naturally associated with a discrete group $\langle f, \phi \rangle$ with the generator ϕ of order 2 and where

$$\langle f, \phi f \phi^{-1} \rangle = \langle f, g f g^{-1} \rangle \subset \langle f, g \rangle, \quad [\langle f, g f g^{-1} \rangle : \langle f, \phi \rangle] = 2$$

This is useful in studying the geometry of Kleinian groups since $\langle f, g \rangle$ will be discrete only if $\langle f, \phi \rangle$ is, and the moduli space of groups $\langle f, \phi \rangle$ is one complex dimension less. This gives a necessary condition in a simpler space to determine the discreteness of $\langle f, g \rangle$.

The dimension reduction here is realised by a projection of principal characters of two-generator Kleinian groups. In applications it is important to know that the image of the moduli space of Kleinian groups under this projection is closed and, among other results, we show how this follows from Jørgensen's results on algebraic convergence.

1. INTRODUCTION

It is shown in [7, 8] that every two-generator Kleinian group $\langle f, g \rangle$ is determined uniquely up to conjugacy by its *principal character*, i.e., the triple of complex parameters

$$(1.1) \quad (\gamma, \beta, \tilde{\beta}) = (\gamma(f, g), \beta(f), \beta(g)).$$

Here

$$(1.2) \quad \gamma(f, g) = \text{tr}[f, g] - 2, \quad \beta(f) = \text{tr}^2(f) - 4, \quad \beta(g) = \text{tr}^2(g) - 4.$$

Thus, the space of two-generator Kleinian groups $\langle f, g \rangle$ can be identified as a subspace of the three complex dimensional space \mathbb{C}^3 . We define

$$\mathcal{D} = \left\{ (\gamma, \beta, \tilde{\beta}) : (\gamma, \beta, \tilde{\beta}) \text{ is the principal character of a Kleinian group} \right\}$$

via the mapping

$$\langle f, g \rangle \mapsto (\gamma(f, g), \beta(f), \beta(g)).$$

Things are set up so the group $\langle \text{identity} \rangle \mapsto (0, 0, 0) \in \mathbb{C}^3$.

Applying [7, Lemma 2.29], given a two-generator Kleinian group $\langle f, g \rangle$ with principal character $(\gamma, \beta, \tilde{\beta})$ with $\gamma \neq \beta$, there is a two-generator discrete group $\Gamma_\phi = \langle f, \phi \rangle$ with principal character $(\gamma, \beta, -4)$, i.e., ϕ is elliptic of order 2. However, we can waive the condition $\gamma \neq \beta$ (see Lemma 8 below).

If the group Γ_ϕ is Kleinian (i.e., not virtually abelian), then there are suites of inequalities on the entries of the principal character which need to hold. One such is Jørgensen's inequality, $|\gamma| + |\beta| \geq 1$. Conversely, it is often easier to establish new discreteness conditions of two generator groups with one generator of order two since we know of many polynomial trace identities which yield a polynomial semigroup action on the associated moduli space, and the dynamics of this action

2000 *Mathematics Subject Classification.* Primary 30C60, 30F40; Secondary 20H10, 53A35.

Key words and phrases. Principal character, moduli space, Kleinian group.

Research supported by UAEU UPAR Grant (31S315). This paper is partly contained in the first author's PhD thesis.

can be used to investigate discreteness of groups. Let us give one such example leading to Jørgensen's inequality to motivate our subsequent results. We note

$$(1.3) \quad \operatorname{tr}[f, gfg^{-1}] - 2 = (\operatorname{tr}[f, g] - 2)(\operatorname{tr}[f, gfg^{-1}] + 2 - \operatorname{tr}^2(f)),$$

which we write as $\gamma(f, gfg^{-1}) = \gamma(\gamma - \beta)$ Following [9] we write $p_\beta(z) = z(z - \beta)$ and $p_\beta^{(n)}(z) = (p_\beta \circ p_\beta \circ \dots \circ p_\beta)(z)$ then repeated application of this identity followed by projection yields the sequence

$$(\gamma, \beta, -4) \mapsto (p_\beta(\gamma), \beta, -4) \mapsto \dots \mapsto (p_\beta^{(n)}(\gamma), \beta, -4) \mapsto \dots$$

giving a sequence of traces of commutators $\{p_\beta^{(n)}(\gamma)\}_{n=0}^\infty$ in the original discrete group Γ_ϕ . It is usually not too difficult to prove such sequences cannot be infinite and contain convergent subsequences. Jørgensen's inequality is implied by the assertion that $p_\beta^{(n)}(\gamma) \not\rightarrow 0$, hence $|\gamma| > 1 + |\beta|$. Every such trace identity yields a new inequality and the semigroup structure is clearly seen in general by

$$(\gamma, \beta, -4) \mapsto (p_\beta(\gamma), \beta, -4) \mapsto (q_\beta(p_\beta(\gamma)), \beta, -4)$$

where, for example, $q_\beta(z) = z(1 + \beta - z)^2$ from the trace identity

$$\operatorname{tr}[f, gfg^{-1}fg] - 2 = (\operatorname{tr}[f, g] - 2)(\operatorname{tr}^2(f) - \operatorname{tr}[f, g] - 1)^2, \quad \gamma(f, gfg^{-1}fg) = \gamma(1 + \beta - \gamma)^2$$

In another direction, the ergodicity of the polynomial semigroup acting in the complement of the Riley slice,

$$\mathcal{R} = \operatorname{int}\{(\gamma, 0, -4) : \langle f, g \rangle \text{ is discrete and freely generated by } f \text{ and } g\}$$

is used to prove the supergroup density and the existence of unbounded Nielsen classes of generating pairs, [15]. Note that this is not the usual definition of the Riley slice but is known to be equivalent.

In order to further these studies we establish a number of useful results in this article. Dynamically, the elementary subgroups are sinks for the polynomial semigroup action on a slice and are isolated within the subset corresponding to discrete groups (apart from trivial abelian and dihedral examples). Thus, after we observe elementary properties of a two-generator discrete group with principal character $(\gamma, \beta, -4)$ in §3, we list the possible principal characters in Table 1 representing a two-generator elementary discrete group with the principal character $(\gamma, \beta, -4)$ in §4. Then we prove the two complex dimensional slice space

$$\mathcal{D}_2 = \{(\gamma, \beta) : \text{there is } \tilde{\beta} \text{ so } (\gamma, \beta, \tilde{\beta}) \text{ a principal character of a Kleinian group}\}$$

is closed in §5. Then \mathcal{D}_2 is the projection of the three complex dimensional moduli space \mathcal{D} to \mathbb{C}^2 . Applications of this set being closed are in the generalisations of Jørgensen's inequality quantifying the isolated nature of the elementary groups in the moduli space of discrete groups, [2], [3].

2. PRELIMINARIES

Let $\operatorname{Möb}^+(\overline{\mathbb{C}})$ be the Möbius group of the normalized orientation preserving Möbius transformations on the Riemann sphere $\overline{\mathbb{C}}$:

$$\operatorname{Möb}^+(\overline{\mathbb{C}}) = \left\{ \frac{az + b}{cz + d} : a, b, c, d \in \mathbb{C} \text{ and } ad - bc = 1 \right\},$$

and let $\operatorname{PSL}(2, \mathbb{C})$ be the projective special linear topological quotient group :

$$\operatorname{PSL}(2, \mathbb{C}) = \operatorname{SL}(2, \mathbb{C}) / \{\pm Id\},$$

where $\operatorname{SL}(2, \mathbb{C}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a, b, c, d \in \mathbb{C} \text{ and } ad - bc = 1 \right\}$. Then these two groups

are topologically isomorphic, $\operatorname{Möb}^+(\overline{\mathbb{C}}) \cong \operatorname{PSL}(2, \mathbb{C})$ and so there are at least two different ways of thinking about groups throughout this paper, as either subgroups

of $\text{Möb}^+(\overline{\mathbb{C}})$ or subgroups of $\text{PSL}(2, \mathbb{C})$. Each of these groups has its own topology, however the topological isomorphism shows us that the concept of discreteness is the same.

A subgroup Γ of $\text{Möb}^+(\overline{\mathbb{C}})$ is called a *Kleinian group* if it is a non-elementary discrete group (see [4, 16]). Here a group is *elementary* if it is virtually abelian, i.e. a finite extension of an abelian group. The elementary groups are classified [4, §5.1]. A subgroup of $\text{Möb}^+(\overline{\mathbb{C}})$ is *non-elementary* if the limit set,

$$L(\Gamma) = \text{the set of accumulation points of a generic orbit } \Gamma(z) = \{f(z) : f \in \Gamma\}$$

contains infinitely many points. Notice from [12] that a group Γ is Kleinian if and only if every two-generator subgroup of Γ is discrete. It is this characterization of Kleinian groups motivates a focus on two-generator groups.

The principal character of a representation of the two-generator group $\langle f, g \rangle$ conveniently encodes various other geometric quantities. One of the most important is the *complex hyperbolic distance* $\delta + i\theta$ between two axes $\text{ax}(f)$ and $\text{ax}(g)$ of two elliptic or loxodromic Möbius transformations f and g , where $\text{ax}(f)$ is the hyperbolic line in \mathbb{H}^3 joining the fixed points of f and g in $\overline{\mathbb{C}}$, δ is the hyperbolic distance between those two axes, and θ is the holonomy of the transformation whose axis contains the common perpendicular between those two axes and moves $\text{ax}(f)$ to $\text{ax}(g)$. Thus,

$$(2.1) \quad \sinh^2(\delta + i\theta) = \frac{4\gamma}{\beta\bar{\beta}}$$

The parameter $\beta(f)$ describes the conjugacy class of Möbius transformation in which f lies.

Lemma 1. *If an element f of $\text{Möb}^+(\overline{\mathbb{C}})$ is not the identity, then*

- (a) *f is parabolic if and only if $\beta(f) = 0$.*
- (b) *f is elliptic if and only if $\beta(f) \in [-4, 0)$.*
- (c) *f is loxodromic if and only if $\beta(f) \in \mathbb{C} \setminus [0, 4]$.*

In addition, the parameter $\beta(f)$ gives the order of each elliptic element.

Lemma 2. *If an element f of $\text{Möb}^+(\overline{\mathbb{C}})$ is not the identity, then f is elliptic of order p if and only if*

$$\beta(f) = -4 \sin^2\left(\frac{k\pi}{p}\right), \quad \text{for } 1 \leq k < p \text{ and } (k, p) = 1.$$

We use the following easy lemma: a group is Kleinian if and only if it is *virtually Kleinian*, i.e. has a Kleinian subgroup of finite index.

Lemma 3. *Let H be a finite index subgroup of Γ . Then Γ is Kleinian if and only if H is Kleinian.*

The proof follows directly from the fact that if H has finite index in Γ , then it has a coset decomposition

$$\Gamma = \phi_1 H \cup \phi_2 H \cup \cdots \cup \phi_n H,$$

where $\phi_1 = Id, \phi_2, \dots, \phi_n \in \Gamma$ for some n . Thus any sequence $\{g_n\}$ contains a subsequence $\{g_{n_k}\}$ lying in one coset.

3. SLICING SPACES OF TWO-GENERATOR DISCRETE GROUPS

Suppose that $\langle f, g \rangle$ is two-generator discrete subgroup of $\text{Möb}^+(\overline{\mathbb{C}})$ with $\gamma = \gamma(f, g) \neq 0$ and $\gamma(f, g) = \gamma \neq \beta = \beta(f)$, Then Gehring and Martin [7, Lemma 2.29] states that there exist elliptics ϕ and ψ of order 2 such that $\Gamma_\phi = \langle f, \phi \rangle$ and $\Gamma_\psi = \langle f, \psi \rangle$ are discrete with the related two-generator groups $\Gamma_\phi = \langle f, \phi \rangle$ and

$\Gamma_\psi = \langle f, \psi \rangle$ are discrete with principal characters $(\gamma, \beta, -4)$ and $(\beta - \gamma, \beta, -4)$, respectively.

Notice that $\gamma(f, g) = 0$ implies f and g have a common fixed point on $\overline{\mathbb{C}}$ and hence is elementary (see [4, Theorem 4.3.5] and [7, Identity (1.5)]). Thus, the condition $\gamma \neq 0$ for Kleinian groups. We now consider the other restriction $\gamma \neq \beta$ of [7, Lemma 2.29].

We start with our observation of the properties of a two-generator discrete group with the principal character $(\gamma, \beta, -4)$, which has one generator of order 2.

Lemma 4. *Let $\Gamma = \langle f, g \rangle$ be a discrete group with principal character $(\gamma, \beta, -4)$. If $\gamma = \beta$, then Γ is an elementary discrete group, which is isomorphic to a dihedral group D_p , for some $p \in \{1, 2, \dots, \infty\}$.*

Proof. There are two cases to consider. If f is parabolic, then $\gamma = \beta = 0$ by Lemma 1 and Γ as f and g have a common fixed point which we may assume by conjugacy is $\infty \in \overline{\mathbb{C}}$. Then Γ is a subgroup of the upper-triangular matrices

$$f = \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \text{ and } g = \begin{pmatrix} \pm i & y \\ 0 & \mp i \end{pmatrix}$$

and the result follows. We may now assume f and g have the form

$$f = \begin{pmatrix} \lambda & 0 \\ 0 & \frac{1}{\lambda} \end{pmatrix} \text{ and } g = \begin{pmatrix} \alpha & \mu \\ \delta & \nu \end{pmatrix}, \text{ where } \lambda \neq 0, \pm 1, \alpha\nu - \mu\delta = 1.$$

Computing the commutator,

$$[f, g] = \begin{pmatrix} \alpha\nu - \lambda^2\mu\delta & \alpha\lambda^2\mu - \alpha\mu \\ \frac{1}{\lambda^2}\delta\nu - \delta\nu & \alpha\nu - \frac{1}{\lambda^2}\mu\delta \end{pmatrix},$$

and hence $\gamma = \text{tr}[f, g] - 2 = -\mu\delta(\lambda - \frac{1}{\lambda})^2$, and $\beta = \text{tr}^2(f) - 4 = (\lambda - \frac{1}{\lambda})^2$. The assumption $\gamma = \beta$ now implies that $\mu\delta = -1$. Since $\alpha\nu - \mu\delta = 1, \alpha\nu = 0$. Now $\beta(g) = -4$, so $\text{tr}(g) = 0$, which gives $\alpha + \nu = 0$. Then $\alpha = \nu = 0$ and

$$g = \begin{pmatrix} 0 & \mu \\ -\frac{1}{\mu} & 0 \end{pmatrix}.$$

Thus, $g(z) = -\frac{\mu^2}{z}$ interchanges the fixed points 0 and ∞ of f . So Γ has a finite orbit $\{0, \infty\}$ and hence it is an elementary discrete group, [4]. Computation shows the conjugate $gfg^{-1} = f^{-1}$:

$$gfg^{-1} = \begin{pmatrix} \frac{1}{\lambda} & 0 \\ 0 & \lambda \end{pmatrix} = f^{-1}.$$

Thus Γ is a dihedral group D_p , for some $p \in \mathbb{N} \cup \{\infty\}$ by from Theorem 1. \square

Lemma 5. *Let $\Gamma = \langle f, g \rangle$ be a discrete group with the principal character $(\gamma, -4, -4)$, then Γ is isomorphic to one of the following elementary discrete groups:*

- (a) *The dihedral group D_∞ , if $\text{ax}(f) \cap \text{ax}(g) = \emptyset$ or $\text{ax}(f) \cap \text{ax}(g) \neq \emptyset$ in \mathbb{C} .*
- (b) *A dihedral group D_p for some $p \in \mathbb{N}$, if $\text{ax}(f) \cap \text{ax}(g) \neq \emptyset$ in \mathbb{H}^3 .*
- (c) *A parabolic group, if $\text{ax}(f) \cap \text{ax}(g) = \{\infty\}$.*

Proof. (1) $\text{ax}(f) \cap \text{ax}(g) = \emptyset$: Let α be the common perpendicular between two axes $\text{ax}(f)$ and $\text{ax}(g)$. Since f and g are elliptic of order 2, they are the rotations of order 2 about their axes $\text{ax}(f)$ and $\text{ax}(g)$ in \mathbb{H}^3 , respectively. And each of f and g interchanges the ending points of α and fixes α setwise, hence Γ is elementary. Furthermore, the product fg fixes the ending points of α and fixes α setwise, so the axis $\text{ax}(fg) = \alpha$. It follows that fg is loxodromic.

Next, from the Fricke identity (see [6]), we have

$$(3.1) \quad \gamma(f, g) = \beta(f) + \beta(g) + \beta(fg) - \text{tr}(f)\text{tr}(g)\text{tr}(fg) + 8.$$

Since $\beta(f) = \beta(g) = -4$, $\text{tr}(f) = 0$, and $f^2 = Id$, the previous identity (3.1) gives $\gamma(f, fg) = \beta(fg)$. and by Lemma 4, $\langle f, fg \rangle$ is a dihedral group. Notice that fg is loxodromic with infinite order, hence $\langle f, g \rangle = \langle f, fg \rangle \cong D_\infty$.

(2) $\text{ax}(f) \cap \text{ax}(g) \neq \emptyset$ in \mathbb{H}^3 : Since f and g have a common fixed point (say x_0), Γ is elementary. Since Γ is discrete, the axes of elliptics both of order 2 must intersect at x_0 an angle $\theta = \frac{k\pi}{n}$ for some k and $n \geq 2$, the product fg is a rotation about $\text{ax}(fg)$ with the oriented angle $2\theta = \frac{2k\pi}{p}$ for any $n \geq 2$ and some $p \in \mathbb{N}$, where $\text{ax}(fg)$ is perpendicular to $\text{ax}(f)$ and $\text{ax}(g)$ and passing through x_0 . Therefore, fg is elliptic of order p . Notice that f and g are rotations of order 2 and their rotation axes $\text{ax}(f)$ and $\text{ax}(g)$ meet in $\overline{\mathbb{H}^3}$, we have $gfg^{-1} = f^{-1}$. It follows that $\langle f, g \rangle \cong D_p$.

(3) $\text{ax}(f) \cap \text{ax}(g) \neq \emptyset$ in $\overline{\mathbb{C}}$: We may assume by conjugacy that ∞ is the common fixed point of f and g in $\overline{\mathbb{C}}$. Then f and g fix ∞ , and hence $\text{ax}(f)$ and $\text{ax}(g)$ are vertical hyperbolic lines. Notice that both f and g are rotations with the angles π at distinct centers. Since the sum of two rotation angles is 2π , the product fg is a translation and hence fg is parabolic fixing ∞ . Thus the classification of elementary groups [4, Section 5.1], $\langle f, g \rangle$ is a group of Euclidean similarities of \mathbb{C} . \square

We next note the following elementary result.

Lemma 6. *Let $\Gamma = \langle f, g \rangle$ be a two-generator group with principal character $(\gamma, \beta, -4)$, then $\Gamma^g = \langle f, gfg^{-1} \rangle$ is an index two subgroup of $\langle f, g \rangle$: $\Gamma = \Gamma^g \cup g\Gamma^g$.*

Then we have the following.

Lemma 7. *Suppose that $\langle f, g \rangle$ is a two-generator Kleinian group with principal character $(\gamma, \beta, \tilde{\beta})$. If $\tilde{\beta} \neq -4$, then*

$$\text{Fix}(f) \cap \text{Fix}(gfg^{-1}) = \emptyset.$$

Proof. Since $\langle f, g \rangle$ is a two-generator Kleinian group, $\gamma \neq 0$ and hence f and g cannot share any fixed points, so $\text{Fix}(f) \cap \text{Fix}(g) = \emptyset$. Since $g(\text{Fix}(f)) = \text{Fix}(gfg^{-1})$, there are two cases for us to consider.

If $\text{Fix}(f)$ has one element, say $\text{Fix}(f) = \{z_0\}$, then $\text{Fix}(gfg^{-1}) = \{g(z_0)\}$. But $g(z_0) \neq z_0$. Thus, $\text{Fix}(f) \cap \text{Fix}(gfg^{-1}) = \emptyset$.

If $\text{Fix}(f)$ has two elements, say $\text{Fix}(f) = \{z_1, z_2\}$, then $\text{Fix}(gfg^{-1}) = \{g(z_1), g(z_2)\}$ and

$$(3.2) \quad g(z_1) \neq z_1, g(z_2) \neq z_2.$$

If $\text{Fix}(f) \cap \text{Fix}(gfg^{-1}) \neq \emptyset$, then by [4, Theorem 5.1.2] f and gfg^{-1} cannot share exactly one fixed point, and hence $\text{Fix}(f) = \text{Fix}(gfg^{-1}) = g(\text{Fix}(f))$. Thus, the only possibility is that g interchange the fixed points of f , so $g(z_1) = z_2$ and $g(z_2) = z_1$, which implies that g^2 fixes z_1 and z_2 and are not fixed by g . We conclude that g^2 has at least three fixed points and this implies that g^2 is the identity, g has order 2, and $\tilde{\beta} = -4$. \square

In [10, Remark 2,6] it is shown that if $\langle f, g \rangle$ is a two-generator discrete group, then there are two elliptic elements ϕ and ψ of order 2 such that

$$(3.3) \quad \varphi f \phi^{-1} = gfg^{-1} \quad \text{and} \quad \psi f \psi^{-1} = gf^{-1}g^{-1}.$$

In particular, if f is elliptic or loxodromic, then the axes $\text{ax}(\phi)$ and $\text{ax}(\psi)$ meet at right-angles to one and other, and they bisect the common perpendicular between the axes $\text{ax}(f)$ and $\text{ax}(gfg^{-1})$.

Lemma 8. *Let $\langle f, g \rangle$ be a Kleinian group with principal character $(\gamma, \beta, \tilde{\beta})$. Then there are two elliptic elements ϕ and ψ of order 2 such that $\Gamma_\phi = \langle f, \phi \rangle$ and $\Gamma_\psi =$*

$\langle f, \psi \rangle$ are discrete \mathbb{Z}_2 -extension of Γ^g with principal characters $(\gamma, \beta, -4)$ and $(\beta - \gamma, \beta, -4)$, respectively.

Proof. Note $\Gamma^g = \langle f, gfg^{-1} \rangle$ is a subgroup of both $\Gamma_\phi = \langle f, \phi \rangle$ and $\Gamma_\psi = \langle f, \psi \rangle$. Suppose that the principal characters for Γ_ϕ and Γ_ψ are $(\gamma_1, \beta, -4)$ and $(\gamma_2, \beta, -4)$, respectively. Using [7, Lemma 2.1]

$$\begin{aligned}\gamma(\gamma - \beta) &= \gamma(f, gfg^{-1}) = \gamma(f, \phi f \phi^{-1}) = \gamma_1(\gamma_1 - \beta), \\ \gamma(\gamma - \beta) &= \gamma(f, gfg^{-1}) = \gamma(f, \psi f \psi^{-1}) = \gamma_2(\gamma_2 - \beta).\end{aligned}$$

we have two quadratic equations

$$\begin{aligned}\gamma_1^2 - \beta\gamma_1 - \gamma(\gamma - \beta) &= 0, \\ \gamma_2^2 - \beta\gamma_2 - \gamma(\gamma - \beta) &= 0.\end{aligned}$$

Solving these two quadratic equations, $\gamma_1 = \gamma$, or $\beta - \gamma$ and $\gamma_2 = \gamma$, or $\beta - \gamma$. As [14, Identity (6.9)] shows both possibilities occur, so

$$\{\gamma_1, \gamma_2\} = \{\gamma, \beta - \gamma\}.$$

Thus, after relabeling, the principal characters for Γ_ϕ and Γ_ψ are $(\gamma, \beta, -4)$ and $(\beta - \gamma, \beta, -4)$, respectively. This completes the proof. \square

4. EXCEPTIONAL SET OF PRINCIPAL CHARACTERS.

Now the natural question is to determine whether two discrete groups $\Gamma_\phi = \langle f, \phi \rangle$ and $\Gamma_\psi = \langle f, \psi \rangle$ in Lemma 8 are actually Kleinian. Since both Γ_ϕ and Γ_ψ are \mathbb{Z}_2 -extension of Γ^g , by Lemma 3, we only need to decide if Γ^g is a Kleinian subgroup. First, we list the possible principal characters in Tables 1, 2, and 3 representing two-generator elementary discrete groups. The set of principal characters occurring in these three tables is called the *exceptional set of principal characters*. We recall from [4, Section 5.1].

Theorem 1 (Classification of elementary discrete groups). *Let G be an elementary discrete group of $\text{Möb}^+(\overline{\mathbb{C}})$, then G is isomorphic to one of the following groups, where $p \in \{1, 2, \dots, \infty\}$.*

(a) A cyclic group \mathbb{Z}_p :

$$\langle a : a^p = Id \rangle.$$

(b) A dihedral group $D_p \cong \mathbb{Z}_p \rtimes \mathbb{Z}_2$:

$$\langle a, b : a^p = b^2 = Id, bab^{-1} = a^{-1} \rangle.$$

(c) The group $(\mathbb{Z}_p \times \mathbb{Z}) \rtimes \mathbb{Z}_2$ or $\mathbb{Z}_p \times \mathbb{Z}$:

$$\langle a, b, c : aba^{-1}b^{-1} = b^p = c^2 = Id, cac^{-1} = a^{-1}, cbc^{-1} = b^{-1} \rangle.$$

(d) A Euclidean translation group $\mathbb{Z} \times \mathbb{Z}$.

(e) A Euclidean triangle group $\Delta(2, 3, 6)$, or $\Delta(2, 4, 4)$, or $\Delta(3, 3, 3)$:

$$\langle a, b : a^p = b^q = (ab)^r = Id, \frac{1}{p} + \frac{1}{q} + \frac{1}{r} = 1 \rangle.$$

(f) A finite spherical triangle group $A_4 = \Delta(2, 3, 3)$, or $S_4 = \Delta(2, 3, 4)$, or $A_5 = \Delta(2, 3, 5)$:

$$\langle a, b : a^p = b^q = (ab)^r = Id, \frac{1}{p} + \frac{1}{q} + \frac{1}{r} > 1 \rangle.$$

Let f and g be elliptic of order $p \in \{2, 3, 4, 5\}$ in $\text{Möb}^+(\overline{\mathbb{C}})$, and let θ be the angle subtended at the origin between $\text{ax}(f)$ and $\text{ax}(g)$ and hence the hyperbolic distance between those two axes is $\delta = 0$. Then, $\sin^2(\theta)$ can be computed by using [14, Lemmas 6.19, 6.20, and 6.21], the parameters $\beta(g)$ and $\beta(f) \in \left\{ -3, -4, \frac{\sqrt{5}-5}{2}, -\frac{\sqrt{5}+5}{2} \right\}$,

and the commutator parameter is $\gamma(f, g) = \frac{-\beta(f)\beta(g)}{4} \sin^2(\theta)$ by [11, Identity (2.7)]. With these elementary observations, some spherical trigonometry, and Theorem 1 about the classification of the elementary discrete groups, we obtain the following Tables 1 and 2, which list the possible principal characters (including the cases in the thesis [17]) of two-generator elementary discrete groups with non-zero commutator parameters $\gamma(f, g) \neq 0$. They are either the dihedral groups (if $\gamma(f, g) = \beta(f)$ and $\beta(g) = -4$) or the finite spherical triangle groups $A_4 = \Delta(2, 3, 3)$, $S_4 = \Delta(2, 3, 4)$, and $A_5 = \Delta(2, 3, 5)$.

 Table 1. Principal Characters: 2, p .

p	$\sin^2(\theta)$	γ	groups	principal characters
3	$\frac{2}{3}$	-2	A_4	$(-2, -3, -4)$
3	$\frac{1}{3}$	-1	S_4	$(-1, -3, -4)$
3	$\frac{3-\sqrt{5}}{6}$	$\frac{\sqrt{5}-3}{2}$	A_5	$(\frac{\sqrt{5}-3}{2}, -3, -4)$
3	$\frac{3+\sqrt{5}}{6}$	$-\frac{3+\sqrt{5}}{2}$	A_5	$(-\frac{3+\sqrt{5}}{2}, -3, -4)$
3	1	-3	D_3	$(-3, -3, -4)$
4	$\frac{1}{2}$	-1	S_4	$(-1, -2, -4)$
4	1	-2	D_4	$(-2, -2, -4)$
5	$\frac{5-\sqrt{5}}{10}$	$\frac{\sqrt{5}-3}{2}$	A_5	$(\frac{\sqrt{5}-3}{2}, \frac{\sqrt{5}-5}{2}, -4)$
5	$\frac{5-\sqrt{5}}{10}$	-1	A_5	$(-1, -\frac{5+\sqrt{5}}{2}, -4)$
5	$\frac{5+\sqrt{5}}{10}$	-1	A_5	$(-1, \frac{\sqrt{5}-5}{2}, -4)$
5	$\frac{5+\sqrt{5}}{10}$	$-\frac{3+\sqrt{5}}{2}$	A_5	$(-\frac{3+\sqrt{5}}{2}, -\frac{5+\sqrt{5}}{2}, -4)$
5	1	$\frac{\sqrt{5}-5}{2}$	D_5	$(\frac{\sqrt{5}-5}{2}, \frac{\sqrt{5}-5}{2}, -4)$
5	1	$-\frac{5+\sqrt{5}}{2}$	D_5	$(-\frac{5+\sqrt{5}}{2}, -\frac{5+\sqrt{5}}{2}, -4)$

 Table 2. Principal Characters: 3, p .

p	$\sin^2(\theta)$	γ	groups	principal characters
3	$\frac{4}{9}$	-1	A_5	$(-1, -3, -3)$
3	$\frac{8}{9}$	-2	A_4	$(-2, -3, -3)$
4	$\frac{2}{3}$	-1	S_4	$(-1, -2, -3)$
5	$\frac{10-2\sqrt{5}}{15}$	$\frac{\sqrt{5}-3}{2}$	A_5	$(\frac{\sqrt{5}-3}{2}, \frac{\sqrt{5}-5}{2}, -3)$
5	$\frac{10-2\sqrt{5}}{15}$	-1	A_5	$(-1, -\frac{5+\sqrt{5}}{2}, -3)$
5	$\frac{10+2\sqrt{5}}{15}$	-1	A_5	$(-1, \frac{\sqrt{5}-5}{2}, -3)$
5	$\frac{10+2\sqrt{5}}{15}$	$-\frac{3+\sqrt{5}}{2}$	A_5	$(-\frac{3+\sqrt{5}}{2}, -\frac{5+\sqrt{5}}{2}, -3)$

Notice that the angle between intersecting axes of elliptics of order 4 in a discrete group is always either 0 when they meet on the Riemann sphere $\overline{\mathbb{C}}$ or $\frac{\pi}{2}$. This yields the additional parameter $(-1, -2, -2)$ for the elementary group S_4 when generated by two elements of order 4. Furthermore, the angle between intersecting axes of elliptics of order 5 in a discrete group is either $\arcsin \frac{2}{\sqrt{5}}$ or its complement $\arcsin \frac{-2}{\sqrt{5}}$. After possibly taking powers of the generator of order 5, the three additional principal characters can be obtained in the following Table 3.

Table 3. Principal Characters: 4, 4 and 5, 5.

p, p	$\sin^2(\theta)$	γ	Group	Parameters
4, 4	1	-1	S_4	$(-1, -2, -2)$
5, 5	$\frac{4}{5}$	$\frac{\sqrt{5}-3}{2}$	A_5	$(\frac{\sqrt{5}-3}{2}, \frac{\sqrt{5}-5}{2}, \frac{\sqrt{5}-5}{2})$
5, 5	$\frac{4}{5}$	-1	A_5	$(-1, -\frac{5+\sqrt{5}}{2}, \frac{\sqrt{5}-5}{2})$
5, 5	$\frac{4}{5}$	$-\frac{3+\sqrt{5}}{2}$	A_5	$(-\frac{3+\sqrt{5}}{2}, -\frac{5+\sqrt{5}}{2}, -\frac{5+\sqrt{5}}{2})$

The classification of the elementary discrete groups (Theorem 1) plays an important role in listing the tables above. We make the following additional remarks.

- The axes of elliptics both of order 2 can intersect at an angle $\frac{k\pi}{p}$ for any k and $p \geq 2$ giving the dihedral group D_p with principal character $(-4 \sin^2 \frac{k\pi}{p}, -4, -4)$ (see Lemma 5).
- The axes of elliptics elements of orders 2 and p , $p \geq 3$, can meet at right-angles. In this case, the principal character of dihedral group D_p is $(-4 \sin^2 \frac{\pi}{p}, -4 \sin^2 \frac{\pi}{p}, -4)$.
- The axes of elliptics of order p and q ($p \leq q$) in a discrete group meet on $\overline{\mathbb{C}}$, i.e., meeting with angle 0, if and only if

$$(p, q) \in \{(2, 2), (2, 3), (2, 4), (2, 6), (3, 3), (3, 6), (4, 4), (6, 6)\}$$

For all of these groups $\gamma(f, g) = 0$. In particular, Euclidean triangle groups $\Delta(2, 3, 6)$, $\Delta(3, 3, 3)$, $\Delta(2, 4, 4)$, and cyclic groups have $\gamma(f, g) = 0$. Also, Euclidean translation groups have $\gamma = 0$.

Lemma 9. *Let $\langle f, g \rangle$ be a Kleinian group with the principal character $(\gamma, \beta, -4)$ such that is not one of those exceptional groups listed in Table 1. Then the subgroup $\Gamma^g = \langle f, gfg^{-1} \rangle$ is Kleinian.*

Proof. We need to show that the discrete subgroup Γ^g of Kleinian group $\langle f, g \rangle$ can not be elementary. Let $\gamma = \gamma(f, g)$ and $\beta = \beta(f)$. By Lemma 8 there is an elliptic h of order 2 such that $\Gamma_h = \langle f, h \rangle$ is discrete group containing Γ^g with index two and the principal character for Γ_h is $(\gamma, \beta, -4)$. By the hypothesis, $(\gamma, \beta, -4)$ is not one of those exceptional groups listed in Table 1, then Γ_h is not a finite spherical triangle group A_4, S_4 , and A_5 . Thus, Γ^g can only be elementary if $\Gamma^g = \gamma(\gamma - \beta) = 0$, i.e., $\gamma = 0$ or $\gamma = \beta$. Since $\langle f, g \rangle$ is Kleinian, $\gamma \neq 0$. So it can only be $\gamma = \beta$ and hence it is the dihedral group. In this case, since f is not of order 2, $gfg^{-1} = f^{\pm 1}$. In the case $gfg^{-1} = f$, it gives $g(\text{Fix}(f)) = \text{Fix}(gfg^{-1}) = \text{Fix}(f)$ and hence g is elliptic of order 2 (Lemma 7) fixing or interchanging the fixed points of f in $\overline{\mathbb{C}}$. In the case $gfg^{-1} = f^{-1}$, g might be a power of f . In either case $\langle f, g \rangle$ is not Kleinian, a contradiction. \square

The following theorem guarantees that Γ^g is a Kleinian group if f is not elliptic of order $p \leq 6$.

Theorem 2. *Suppose that $\langle f, g \rangle$ is a Kleinian group. If f is not elliptic of order $p \in \{2, 3, 4, 6\}$, then the subgroup $\Gamma^g = \langle f, gfg^{-1} \rangle$ is Kleinian.*

Proof. (a) Suppose that g is elliptic of order 2. Then Γ^g is a subgroup of index two in Γ and Lemma 3 ensures that Γ^g is also Kleinian. (b) Suppose that the order of g is not 2. We need to show that Γ^g is non-elementary. If Γ^g is elementary, then (see [4, Section 5.1]). (i) Γ^g is elementary of type I; each non-trivial element is elliptic. Since the order of f is not $p \leq 6$ or $p = 5$, neither is the order of gfg^{-1} . If $\gamma(f, gfg^{-1}) \neq 0$, then from Tables 1, 2, and 3, Γ^g is neither the finite spherical triangle groups A_4, S_4 , and A_5 , nor the dihedral groups D_3, D_4 , and D_5 .

If $\gamma(f, gfg^{-1}) = 0$, then $\gamma = \beta$, and $\text{Fix}(f) \cap \text{Fix}(gfg^{-1}) \neq \emptyset$. On the other hand, by Lemma 7, $\text{Fix}(f) \cap \text{Fix}(gfg^{-1}) = \emptyset$, a contradiction.

(ii) Suppose Γ^g is an elementary group of type II; then Γ^g is conjugate to a subgroup of $\text{Möb}(\overline{\mathbb{C}})$ fixing ∞ whose every element is parabolic of the form $z \mapsto z+b$, $b \in \mathbb{C}$. Thus, the group Γ^g is abelian and hence $g(\text{Fix}(f)) = \text{Fix}(gfg^{-1}) = \text{Fix}(f)$, a contradiction to Γ being Kleinian.

(iii) Suppose Γ^g is an elementary group of type III; then both f and gfg^{-1} are elliptic or both are loxodromic. In either case Γ^g is abelian and as above this is a contradiction.

We have shown that Γ^g cannot be elementary if g does not have order 2. Hence in all two cases Γ^g is a Kleinian group. \square

Notice that a Kleinian group $\langle f, g \rangle$ with f elliptic of order $p = 5$ or $p \geq 7$ and $\beta(g) \neq -4$, cannot have $\gamma = \beta$. If so, then f and gfg^{-1} have a common fixed point. But this discrete group cannot be a Euclidean group as they do not have elliptics of order 5 or greater than 6. Thus f and gfg^{-1} have two common fixed points and the same trace. Hence $gfg^{-1} = f^{\pm 1}$ and g must fix or interchange the fixed points of f . The first is not possible, and the second shows $g^2 = \text{identity}$.

5. PROJECTIONS AND PRINCIPAL CHARACTERS

Our methods to establish new inequalities rely on a fundamental result concerning spaces of finitely generated Kleinian groups. Namely that they are closed in the topology of algebraic convergence – a result originally due to Jørgensen [12]. We state this now.

Definition 1. *A sequence of n -generator subgroups $\Gamma_j = \langle f_{j,1}, f_{j,2}, \dots, f_{j,n} \rangle$ is said to be convergent algebraically to a n -generator subgroup $\Gamma = \langle f_1, f_2, \dots, f_n \rangle$ in $\text{Möb}(\overline{\mathbb{C}})$ if for each $k = 1, 2, \dots, n$, the sequence of the corresponding generators $\{f_{j,k}\}$ converges uniformly to f_k in the spherical metric of $\overline{\mathbb{C}}$.*

Theorem 3 (Jørgensen). *If a sequence of n -generator Kleinian subgroups $\Gamma_j = \langle f_{j,1}, f_{j,2}, \dots, f_{j,n} \rangle$ converges algebraically to $\Gamma = \langle f_1, f_2, \dots, f_n \rangle$ in $\text{Möb}(\overline{\mathbb{C}})$, then Γ is a Kleinian group.*

Moreover, the map back is an eventual homomorphism. That is for all sufficiently large j the map $\Gamma \rightarrow \Gamma_j$ defined by $f_k \mapsto f_{j,k}$ extends to a homomorphism of the groups.

There is a higher dimensional analogue to this Jørgensen's theorem, but additional hypotheses are necessary – for instance giving a uniform bound on the torsion in a sequence, or asking that the limit set be in geometric position (see [13, Proposition 5.8]).

Now we show the slice \mathcal{D}_2 is closed. Notice that we are free to normalize a sequence of two-generator groups by conjugation and to pass to a subsequences. First a preliminary result.

Theorem 4. *The slice*

$$\mathcal{D}^* = \{(\gamma, \beta, -4) : (\gamma, \beta, -4) \text{ is the principal character of a Kleinian group}\}$$

is closed in the two complex dimensional space \mathbb{C}^2 in the usual topology.

Proof. Suppose that $\{(\gamma_j, \beta_j, -4)\}$ is a sequence of principal characters of Kleinian groups, say $\{\langle f_j, g_j \rangle\}$. We proceed by considering two cases: up to subsequence, f_i is parabolic for all i or not.

Case (a). Suppose f_j is parabolic for all j . Conjugate each Γ_j so that the first generator is now represented by the matrix $f_j = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. Then f_j is constant and it remains to show that the sequence $\{g_j\}$ also converges. Since $\text{tr}(g_j) = 0$, we suppose the matrix for second generator is

$$g_j = \begin{pmatrix} a_j & b_j \\ c_j & -a_j \end{pmatrix}.$$

We calculate that $\gamma_j = c_j^2$ and hence $c_j \rightarrow c$. Jørgensen's inequality gives $|\gamma_j| \geq 1$ for all j and hence $c_j \rightarrow c \neq 0$.

Set

$$h_j = \begin{pmatrix} 1 & \frac{-a_j + i\sqrt{1+c_j}}{c_j} \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad h_j g_j h_j^{-1} = \begin{pmatrix} i\sqrt{1+c_j} & 1 \\ c_j & -i\sqrt{1+c_j} \end{pmatrix}.$$

Since f_j commutes with h_j , f_j is left unchanged under conjugation. The result follows by Theorem 3.

Case (b). We now suppose that f_j is not parabolic for all j . By conjugation, we may assume that f_j is represented by the matrix

$$f_j = \begin{pmatrix} \lambda_j & 0 \\ 0 & \frac{1}{\lambda_j} \end{pmatrix},$$

where $\lambda_j \neq \pm 1$. Thus the parameter $\beta_j = (\lambda_j - \frac{1}{\lambda_j})^2$, which gives the quadratic equation

$$\lambda_j^2 - \sqrt{\beta_j} \lambda_j - 1 = 0$$

and hence $\lambda_j = \frac{\sqrt{\beta_j} \pm \sqrt{\beta_j + 4}}{2}$. Since $\beta_j \rightarrow \beta$, λ_j converges to $\lambda = \frac{\sqrt{\beta} \pm \sqrt{\beta + 4}}{2}$. Noting that $\lambda \neq 0, \pm 1$ for either choice of λ , we conclude that f_j converges to the element f represented by the matrix

$$f_j \rightarrow f = \begin{pmatrix} \lambda & 0 \\ 0 & \frac{1}{\lambda} \end{pmatrix} \quad \text{and} \quad \beta_j \rightarrow \beta = \left(\lambda - \frac{1}{\lambda}\right)^2.$$

It remains to show that the order 2 elements g_j also converge. Note first that $\gamma_j = -\beta_j b_j c_j$ and so we may assume that $b_j c_j \neq 0$. As in the previous case the matrix for g_j can be written as

$$g_j = \begin{pmatrix} a_j & -\frac{1+a_j^2}{c_j} \\ c_j & -a_j \end{pmatrix}.$$

so that $a_j \neq \pm i$ and $1 + a_j^2 \neq 0$. Consider the conjugacy of the group $\langle f_j, g_j \rangle$ by the following diagonal matrix,

$$\phi_j = \begin{pmatrix} i\sqrt{\frac{c_j}{1+a_j^2}} & 0 \\ 0 & -i\sqrt{\frac{1+a_j^2}{c_j}} \end{pmatrix},$$

and then g_j (replaced by as the conjugacy $\phi_j g_j \phi_j^{-1}$:) has the form

$$g_j = \begin{pmatrix} a_j & 1 \\ -(1+a_j^2) & -a_j \end{pmatrix}.$$

Since f_j commutes with ϕ_j , f_j is left unchanged under the conjugation.

The commutator is now

$$[f_j, g_j] = f_j g_j f_j^{-1} g_j^{-1} = \begin{pmatrix} -a_j^2 + \lambda_j^2(1 + a_j^2) & -a_j + \lambda_j^2 a_j \\ \frac{a_j(1 + a_j^2)}{\lambda_j^2} - a_j(1 + a_j^2) & \frac{1 + a_j^2}{\lambda_j^2} - a_j^2 \end{pmatrix}.$$

Hence the parameter γ_j for the group $\langle f_j, g_j \rangle$ is

$$(5.1) \quad \gamma_j = \frac{a_j^2 (\lambda_j^2 - 1)^2 + (\lambda_j^2 - 1)^2}{\lambda_j^2}.$$

At this point we would like to show $\lambda_j \not\rightarrow 1$, but this can actually happen in limits of Dehn surgeries - see the following example. Thus we have two subcases:

Case (b) (i) Suppose $\lambda_j \not\rightarrow 1$.

Solving for a_j yields

$$a_j^2 = \frac{\gamma_j \lambda_j^2 - (\lambda_j^2 - 1)^2}{(\lambda_j^2 - 1)^2} \rightarrow \frac{\gamma \lambda^2 - (\lambda^2 - 1)^2}{(\lambda^2 - 1)^2}.$$

Let $a^2 = \frac{\gamma \lambda^2 - (\lambda^2 - 1)^2}{(\lambda^2 - 1)^2}$, then g_j converges to the element g of $\text{PSL}(2, \mathbb{C})$ represented by the matrix

$$g = \begin{pmatrix} a & 1 \\ -(1 + a^2) & -a \end{pmatrix}.$$

and $\gamma_j \rightarrow \gamma(f, g) = \gamma$. Applying Theorem 3, the limit group $\Gamma = \langle f, g \rangle$ is Kleinian.

Case (b) (ii) Suppose $\lambda_j \rightarrow 1$. This is equivalent to $\beta_j \rightarrow 0$. For each j we may conjugate f_j and g_j so that f_j has a fixed point at ∞ and g_j has a fixed point at 0. Thus f_j is upper triangular and g_j is lower triangular. We may further conjugate by a diagonal matrix to achieve the upper entry in the matrix representing f_j is 1. With this normalisation we have

$$(5.2) \quad f_j = \begin{pmatrix} \lambda_j & 1 \\ 0 & 1/\lambda_j \end{pmatrix}, \quad g_j = \begin{pmatrix} i & 0 \\ c_j & -i \end{pmatrix}$$

and $\lambda_j \rightarrow 1$. We calculate that

$$(5.3) \quad \gamma_j = \gamma(f_j, g_j) = c_j \left(c_j + \frac{2i(\lambda_j^2 - 1)}{\lambda_j} \right).$$

It follows that $c_j \rightarrow \pm\sqrt{\gamma}$, for one choice of sign. Thus $f_j \rightarrow f$ and $g_j \rightarrow g$

$$(5.4) \quad f = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad g = \begin{pmatrix} i & 0 \\ \sqrt{\gamma} & -i \end{pmatrix}$$

This group is nonelementary by Theorem 3 so $\gamma \neq 0$. One case see this since as soon as $|\beta_j| < \frac{1}{2}$, Jørgensen's inequality gives $|\gamma_j| > \frac{1}{2}$. This completes the proof. \square

Example. We next give an example to show the last case can occur. That is $\beta_j \rightarrow 0$ and $\gamma_j \rightarrow \gamma$ (and $|\gamma| = 1$). Let

$$(5.5) \quad f = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad h = \begin{pmatrix} 1 & 0 \\ a_\infty & 1 \end{pmatrix}$$

Then $\langle f, h \rangle$ is (a representation of) the two bridge figure of eight knot group if $a_\infty = (1 + i\sqrt{3})/2$. The relator in this group is

$$(5.6) \quad h f h^{-1} f h = f h f^{-1} h f$$

If we perform $(p, 0)$ Dehn surgery on the complement of this knot we obtain a Kleinian group generated by two elliptics of order p , f_p and h_p . Then up to conjugacy f_p and h_p have the form

$$f_p = \begin{pmatrix} e^{i\pi/p} & 1 \\ 0 & e^{-i\pi/p} \end{pmatrix}, \quad h_p = \begin{pmatrix} e^{i\pi/p} & 0 \\ a_p & e^{-i\pi/p} \end{pmatrix}$$

Then the relator (5.6) holds if and only if

$$a_p = \frac{1}{2} \left(3 - 2 \cos \frac{2\pi}{p} - \sqrt{-1 - 4 \cos \frac{2\pi}{p} + 2 \cos \frac{4\pi}{p}} \right)$$

or its conjugate. Thus $\langle f_p, h_p \rangle$ is the group obtained from this orbifold Dehn surgery. As such it is discrete and nonelementary. Then, as $p \rightarrow \infty$,

$$\begin{aligned} \beta(f_p) = \beta(h_p) &= -4 \sin^2 \frac{\pi}{p} \rightarrow 0 = \beta(f), \quad \text{and} \\ \gamma(f_p, h_p) &= a_p(-2 + a_p + 2 \cos \frac{2\pi}{p}) \rightarrow a_\infty^2 = \gamma(f, g) \end{aligned}$$

As $\beta(f_p) = \beta(h_p)$ we find an involution g_p such that $h_p = g_p f_p g_p^{-1}$ with $\gamma(f_p, g_p) = \gamma(f_p, h_p)$.

We next have the following lemma.

Lemma 10. *Let $(\gamma_j, \beta_j, \tilde{\beta}_j)$ be a sequence of principal characters of two-generator Kleinian groups $\langle f_j, g_j \rangle$, and let $(\gamma, \beta, \tilde{\beta})$ be the principal character of two-generator group $\langle f, g \rangle$. Suppose that (γ_j, β_j) converges to (γ, β) and f is not elliptic of order $p \in \{2, 3, 4, 6\}$. Then $\gamma \neq 0$ and $\gamma \neq \beta$.*

Proof. Since f is not elliptic of order $p \in \{2, 3, 4, 6\}$, f_j is likewise not elliptic of these orders for all but finitely many j . Applying Theorem 2, $\langle f_j, g_j f_j g_j^{-1} \rangle$ is a sequence of Kleinian groups with corresponding principal character $(\gamma_j(\gamma_j - \beta_j), \beta_j, \beta_j)$ which converge to $(\gamma(\gamma - \beta), \beta, \beta)$. Now $\langle f_j, g_j f_j g_j^{-1} \rangle$ is Kleinian group generated by two elements of the same trace. Thus $\gamma_j(\gamma_j - \beta_j) \neq 0$ and hence $\gamma_j \neq 0$ and $\gamma_j \neq \beta_j$.

Now we may apply a result by C.Cao [5, Theorem 5.1] which gives a universal lower bound on the commutator parameter for such groups.

$$\gamma_j(\gamma_j - \beta_j) \geq 0.198.$$

Thus, $\lim_{j \rightarrow \infty} \gamma_j(\gamma_j - \beta_j) = \gamma(\gamma - \beta) \geq 0.198$. It follows that $\gamma \neq 0$ and $\gamma \neq \beta$. \square

Theorem 5. *The two complex dimensional space*

$\mathcal{D}_2 = \{(\gamma, \beta) : (\gamma, \beta, \tilde{\beta}) \text{ is the principal character of a Kleinian group for some } \tilde{\beta}\}$
is closed in the two complex dimensional space \mathbb{C}^2 in the usual topology.

Proof. Suppose that $\{(\gamma_j, \beta_j)\}$ is a sequence in \mathcal{D}_2 with limit (γ, β) in \mathbb{C}^2 . By definition there is a sequence $\tilde{\beta}_j$ so that $\{(\gamma_j, \beta_j, \tilde{\beta}_j)\}$ is the principal character of a Kleinian group. The we project to the principal character $\{(\gamma_j, \beta_j, -4)\}$ of $\langle f_j, \phi_j \rangle$. If infinitely many of these groups are Kleinian, the result we seek follows from Theorem 4. After passing to a subsequence we are left with the case $\langle f_j, \phi_j \rangle$ is elementary for every j .

Case (i). Suppose f_j is not elliptic of order $p \in \{2, 3, 4, 6\}$ for infinitely many j . Then Lemma 10 tells us that $\gamma_j \notin \{0, \beta_j\}$ and hence $\langle f_j, \phi_j \rangle$ is Kleinian, so this case cannot occur.

Case (ii). We may now suppose, after passing to a subsequence, that each f_j is elliptic of order $p \in \{2, 3, 4, 5, 6\}$, $\beta(f_j) = -4 \sin^2 \frac{k\pi}{p} = \beta$, $(k, p) = 1$. $\langle f_j, \phi_j \rangle$ is a discrete elementary group generated by elliptics of orders p and 2. The order of any such group is less than that of A_5 . If infinitely many of these groups are finite, then in fact $[f_j, \phi]$ is elliptic of order 1, 2, 3, 4, 5, or 6. Thus γ_j lies in a finite set of values. Again, after a subsequence we have $\gamma_j = \gamma$. Then $\{(\gamma_j, \beta_j, \tilde{\beta}_j)\} = \{(\gamma, \beta, \tilde{\beta}_j)\}$ and the latter group is Kleinian by hypothesis. Thus $(\gamma, \beta) \in D_2$.

Case (iii). We are left with the situation that $\langle f_j, \phi_j \rangle$ is generated by elliptics of orders p and 2, infinite and elementary for each j . The classification tells us this only happens for the Euclidean groups and any such group has a common fixed point. Then $\gamma(f_j, \phi_j) = 0 = \gamma(f_j, g_j)$ which is not possible as $\langle f_j, g_j \rangle$ is a Kleinian group. \square

REFERENCES

- [1] Alaqad, H., Gong, J. and Martin, G. J., Chebyshev Polynomials and Inequalities for Kleinian Groups, arXiv 2103.05696v1, 2021, 1-18.
- [2] Alaqad, H., Gong, J. and Martin, G. J., Isolation in the moduli space of discrete groups, to appear.
- [3] Alaqad, H., Gong, J. and Martin, G. J., On Jørgensen's inequality for Kleinian groups, to appear.
- [4] A.F. Beardon, *The geometry of discrete groups*, Axler, S., Gehring, F.W., Ribet, K.A., Eds.; Springer-Verlag: Berlin and New York, Germany and USA ,1983.
- [5] C. Cao, Some trace inequalities for discrete groups of Möbius transformations. *Proc. Amer. Math. Soc.* 1995, *123*, 3807-3815.
- [6] R. Fricke and F. Klein. *Vorlesungen über die theorie der automorphen functionen*, Chapter 2, Teubner, Leipzig, 1897.
- [7] F.W. Gehring and G.J. Martin, Commutators, collars and the geometry of Möbius groups, *J. Anal. Math.* 1994, *63*, 175–219.
- [8] F.W. Gehring and G.J. Martin, Inequalities for Möbius transformation and discrete groups FW Gehring, GJ Martin *J. Reine Angew. Math.*, **418**, 31–76
- [9] F.W. Gehring, and G.J. Martin, Iteration theory and inequalities for Kleinian groups. *Bull. Amer. Math. Soc.* 1989, *21*, 57-63.
- [10] F.W. Gehring, and G.J. Martin, Stability and extremality in Jørgensen's inequality, *Proc. Amer. Math. Soc.* 1993 , *117*, 277-282.
- [11] F. W. Gehring and G. J. Martin. 6-torsion and hyperbolic volume, *Proc. Amer. Math. Soc.*, 1993, *117*, 727-735.
- [12] T. Jørgensen, On discrete groups of Möbius transformations, *Amer. J. Math.* 1976, *98*, 739-749.
- [13] G. J. Martin, On discrete Möbius groups in all dimensions: a generalization of Jørgensen's inequality, *Acta Math.*, 1989, *163*, 253–289.
- [14] G.J. Martin, *The Geometry and Arithmetic of Kleinian Groups*, Handbook of Group Actions, Volume I (Advanced lectures in Mathematics Volume 31), *International Press of Boston, Inc.* 2015, 411-494.
- [15] G.J. Martin, *Nondiscrete parabolic characters of the free group F_2 : supergroup density and Nielsen classes in the complement of the Riley slice*, **103**, (2021), 1402–1414.
- [16] Maskit, B. *Kleinian groups*, Springer-Verlag: Berlin and New York, Germany and USA, 1987; pp.2–28. .
- [17] Zhang, Q. *Parameters of the two generator elementary discrete groups*, Master thesis, Massey University 2006, New Zealand.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNITED ARAB EMIRATE UNIVERSTY
Email address: `hala_a@uaeu.ac.ae`

DEPARTMENT OF MATHEMATICAL SCIENCES, UNITED ARAB EMIRATE UNIVERSTY
Email address: `j.gong@uaeu.ac.ae`

SCHOOL OF NATURAL AND COMPUTATIONAL SCIENCES, MASSEY UNIVERSITY,
Email address: `g.j.martin@massey.ac.nz`