1

Cryogenic Penning-Trap Apparatus for Precision Experiments with Sympathetically Cooled (anti)protons

M. Niemann,¹ T. Meiners,¹ J. Mielke,¹ N. Pulido,¹ J. Schaper,^{5,1} M.J. Borchert,^{5,1}

J.M. Cornejo,¹ A.-G. Paschke,^{1,2} G. Zarantonello,^{1,2} H. Hahn,^{2,1} T. Lang,¹ C. Manzoni,³ M. Marangoni,³ G. Cerullo,³ U. Morgner,¹ J.-A. Fenske,²

A. Bautista-Salvador,^{2,1} R. Lehnert,^{4,1} S. Ulmer,⁵ and C. Ospelkaus^{1,2}

¹Institut für Quantenoptik, Leibniz Universität Hannover, Welfengarten 1, 30167 Hannover, Germany, and Laboratorium für Nano- und Quantenengineering, Leibniz Universität Hannover, Schneiderberg 39, 30167 Hannover, Germany

> ² Physikalisch-Technische Bundesanstalt Bundesallee 100, 38116 Braunschweig, Germany

³IFN-CNR, Dipartimento di Fisica, Politecnico di Milano, Piazza L. da Vinci 32, Milano, 20133, Italy

> ⁴Indiana University Center for Spacetime Symmetries Bloomington, IN 47405, USA

> ⁵ Ulmer Fundamental Symmetries Laboratory, RIKEN Hirosawa, Wako, Saitama 351-0198, Japan

Current precision experiments with single (anti)protons to test CPT symmetry progress at a rapid pace, but are complicated by the need to cool particles to sub-thermal energies. We describe a cryogenic Penning-trap setup for ${}^{9}\text{Be}^{+}$ ions designed to allow coupling of single (anti)protons to laser-cooled atomic ions for sympathetic cooling and quantum logic spectroscopy. We report on trapping and laser cooling of clouds and single ${}^{9}\text{Be}^{+}$ ions. We discuss prospects for a microfabricated trap to allow coupling of single (anti)protons to laser-cooled ${}^{9}\text{Be}^{+}$ ions for sympathetic laser cooling to sub-mK temperatures on ms time scales.

1. Introduction

As a result of CPT symmetry, particles and their antiparticles must have the same mass, lifetime, charge, and magnetic moment. The (anti)proton is an attractive candidate to test CPT symmetry by comparing matter– antimatter conjugates in the baryonic sector, complementary to tests in the lepton sector, e.g., with electrons and positrons.¹ Magnetic-moment comparisons^{2,3} in particular provide a sensitive test for potential new physics.⁴ In Penning-trap precision measurements, magnetic moments can be determined by measuring the ratio of the Larmor frequency to the free cyclotron frequency. Compared to the electron and positron, the bigger mass of the (anti)proton makes resistive cooling to the ground state of the cyclotron motion impossible. The resulting finite temperature complicates the measurement of the Larmor frequency via the continuous Stern–Gerlach effect.⁵ Furthermore, systematic effects can be proportional to the oscillation amplitude of the particle making efficient cooling to sub-thermal energies highly desirable. Such temperatures could be reached through sympathetic cooling with a laser-cooled ion. Implementing this in the same potential well could introduce additional systematic effects and is not possible for the commonly used positively charged laser-cooled ions together with the antiproton. We pursue an approach where the (anti)proton and the laser-cooled ion are confined in spatially separate potential wells and interact remotely via the Coulomb interaction.⁶ This approach has already been demonstrated with pairs of atomic ions in radio-frequency Paul traps.^{7,8}

2. Trapping single ⁹Be⁺ ions

We have built and commissioned a cryogenic Penning-trap system to implement this approach. The setup is based on the BASE CERN setup,⁹ enhanced to allow for laser access for ablation loading, photo-ionization, Doppler and ground-state cooling of ⁹Be⁺ ions, as well as for detection of laser-induced fluorescence. The setup is operated at a magnetic field strength of 5T and cooled using a vibration-isolated cryocooler. It is described in more detail in Ref. 10. We load clouds of ⁹Be⁺ ions by focusing single 532 nm laser pulses with tens of μ J pulse energy onto a solid beryllium target embedded into a trap electrode via an in-vacuum off-axis parabolic mirror. We laser cool the ions and detect their presence through laserinduced fluorescence using a laser beam at 313 nm. We apply a series of waveforms to the trap electrodes in order to subsequently reduce the number of particles, until single ions can be observed as identified by discrete steps in the level of laser-induced fluorescence.

3. Experimental prospects for ⁹Be⁺ ions in Penning traps

We have recently demonstrated spin-motional control of ${}^{9}\text{Be}^{+}$ ions using a spectrally tailored frequency comb.¹¹ While the experimental demonstration was carried out at a comparatively low magnetic field in a microfab-

 $\mathbf{2}$

ricated Paul trap, this type of manipulation lends itself to ground-state cooling of ${}^{9}\text{Be}^{+}$ ions in the 5 T field of the Penning trap, where electronspin energy levels with a level spacing of ~ 140 GHz need to be coupled using a stimulated Raman process. Our immediate plans for the setup comprise the demonstration of ground-state cooling and the coupling of pairs of ${}^{9}\text{Be}^{+}$ ions as a precursor for sympathetic cooling of (anti)protons.

4. (Anti-)protons and quantum logic spectroscopy

We have designed an off-axis proton source for the apparatus, which is awaiting commissioning. Coupling a single (anti)proton to a laser-cooled ⁹Be⁺ ion will likely require a microfabricated trap in order to minimize the distance between the particles and obtain a strong coupling. Compared to the commonly used microfabricated surface-electrode Paul traps, a ringshaped electrode of a micro Penning trap would have to be metal coated both on the front and back face of the disc and on the inside of the ring. We have produced test structures using deep reactive ion etching of a silicon wafer. For metalization, wafers have been coated either using resistive evaporation of Ti and Au under constant rotation of the sample or using direct sputter deposition of Au. On top of these thin metal films, a thick electroplated film of Au has been grown and structured using optical lithography. A sample structure is shown in Fig. 1. The process will be extended by including the spacers to electrically isolate multiple such rings already at the stage of the wafer etching. Electrical isolation will be possible through the use of the lithographic step, which allows us to leave parts of the sample uncoated. Once all of the above ingredients have been implemented and a suitable spin-motional coupling mechanism for the (anti)proton has been implemented, quantum logic spectroscopy 6,12 could be envisioned as a means of probing all relevant transitions of single (anti)protons out of the motional ground state.

Acknowledgments

We acknowledge the support of Wissenschaftlicher Gerätebau at PTB for the construction of the setup and the rotation holder, the support of the PTB cleanroom team concerning the fabrication of the sample structures, and P. Hinze for the SEM pictures. We acknowledge funding from ERC StG "QLEDS," and from DFG through SFB 1227 DQ-mat (project B06), DFG grant "Apparatur für kryogene Ionenfallen," and the clusters of excellence QUEST and Quantum Frontiers. RL acknowledges support from



Fig. 1. Gold electroplated silicon test structure for a cylindrical micro Penning trap.

the Alexander von Humboldt Foundation.

References

4

- 1. H. Dehmelt, Rev. Mod. Phys. 62, 525 (1990).
- 2. G. Schneider et al., Science 358, 1081 (2017).
- 3. C. Smorra *et al.*, Nature **550**, 371 (2017).
- 4. R. Bluhm, V.A. Kostelecký, and N. Russell, Phys. Rev. Lett. ${\bf 79},\,1432\,(1997).$
- 5. A. Mooser et~al., Phys. Rev. Lett. ${\bf 110},$ 140405 (2013).
- 6. D.J. Wineland et al., J. Res. Natl. Inst. Stand. Tech. 103, 259 (1998).
- 7. K.R. Brown et al., Nature 471, 196 (2011).
- 8. M. Harlander et al., Nature 471, 200 (2011).
- 9. C. Smorra et al., Eur. Phys. J. ST 224, 3055 (2015).
- 10. M. Niemann et al., arXiv:1906.09249.
- 11. A.-G. Paschke et al., Phys. Rev. Lett. 122, 123606 (2019).
- 12. D.J. Heinzen and D.J. Wineland, Phys. Rev. A 42, 2977 (1990).