# Planets Across Space and Time (PAST). II: Catalog and Analyses of the LAMOST-Gaia-Kepler Stellar Kinematic Properties

DI-CHANG CHEN,<sup>1,2</sup> JIA-YI YANG,<sup>1,2</sup> JI-WEI XIE,<sup>1,2</sup> JI-LIN ZHOU,<sup>1,2</sup> SUBO DONG,<sup>3</sup> ZHENG ZHENG,<sup>4</sup> JING-HUA ZHANG,<sup>5</sup> CHAO LIU,<sup>6,7</sup> HAI-FENG WANG,<sup>8</sup> MAO-SHENG XIANG,<sup>5,9</sup> WEIKAI ZONG,<sup>10</sup> YANG HUANG,<sup>11</sup> AND ALI LUO<sup>5</sup>

<sup>1</sup>School of Astronomy and Space Science, Nanjing University, Nanjing 210023, China

<sup>2</sup>Key Laboratory of Modern Astronomy and Astrophysics, Ministry of Education, Nanjing 210023, China

<sup>3</sup>Kavli Institute for Astronomy and Astrophysics, Peking University, Beijing 100871, China

<sup>4</sup>Department of Physics and Astronomy, University of Utah, Salt Lake City, UT 84112

<sup>5</sup>National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100012, China

<sup>6</sup>Key Lab of Space Astronomy and Technology, National Astronomical Observatories, CAS, 100101, China

<sup>7</sup> University of Chinese Academy of Sciences, Beijing, 100049, China

<sup>8</sup>South-Western Institute for Astronomy Research, Yunnan University, Kunming, 650500, China; LAMOST Fellow

<sup>9</sup>Max-Planck Institute for Astronomy, Königstuhl 17, D-69117 Heidelberg, Germany

<sup>10</sup>Department of Astronomy, Beijing Normal University, Beijing 100875, China

<sup>11</sup>South-Western Institute for Astronomy Research, Yunnan University, Kunming, 650500, China

### ABSTRACT

The Kepler telescope has discovered over 4,000 planets (candidates) by searching  $\sim 200,000$  stars over a wide range of distance (order of kpc) in our Galaxy. Characterizing the kinematic properties (e.g., Galactic component membership and kinematic age) of these Kepler targets (including the planet candidate hosts) is the first step towards studying Kepler planets in the Galactic context, which will reveal fresh insights into planet formation and evolution. In this paper, the second part of the Planets Across the Space and Time (PAST) series, by combining the data from LAMOST and Gaia and then applying the revised kinematic methods from PAST I, we present a catalog of kinematic properties (i.e., Galactic positions, velocities, and the relative membership probabilities among the thin disk, thick disk, Hercules stream, and the halo) as well as other basic stellar parameters for 35,835 Kepler stars. Further analyses of the LAMOST-Gaia-Kepler catalog demonstrate that our derived kinematic age reveals the expected stellar activity-age trend. Furthermore, we find that the fraction of thin (thick) disk stars increases (decreases) with the transiting planet multiplicity ( $N_{\rm p} = 0, 1, 2$  and 3+) and the kinematic age decreases with  $N_{\rm p}$ , which could be a consequence of the dynamical evolution of planetary architecture with age. The LAMOST-Gaia-Kepler catalog will be useful for future studies on the correlations between the exoplanet distributions and the stellar Galactic environments as well as ages.

Keywords: , ----

### 1. INTRODUCTION

With the discovery of thousands of exoplanets, the scope of planetary research has been expanding from the Solar system/neighborhood to a wider range of our Milky Way Galaxy. Understanding how planetary properties depend on the Galactic environment and age will provide crucial insights on planet formation and evolution. Towards a Galactic census of planets, we have started a research project, dubbed Planets Across Space and Time (PAST). In the Paper I (here after PAST I, Chen et al. 2021) of the PAST series, we revisited the kinematic methods for classification of Galactic components and extended the applicable range of the methods from  $\sim 100 - 200$  pc to  $\sim 1.5$  kpc to cover most of planet candidate host stars. Additional, we revised the Age-Velocity dispersion Relation (AVR), which allows

Corresponding author: Ji-Wei Xie jwxie@nju.edu.cn

us to derive kinematic age with a typical uncertainty of  $\sim 10 - 20\%$  for an ensemble of stars. Here, in the second paper of PAST (PAST II), we apply the revised kinematic methods and AVR to a large homogeneous sample of stars with the observational synergy among LAMOST (the Large Sky Area Multi-Object Fiber Spectroscopic Telescope, also known as Goushoujing Telescope; spectroscopy, Wang et al. 1996; Su & Cui 2004; Cui et al. 2012; Zhao et al. 2012; Luo et al. 2012; Zong et al. 2018), Gaia (astrometry, Gaia Collaboration et al. 2018a,b) and Kepler (photometry, Mathur et al. 2017).

The Kepler mission has provided an unprecedented legacy sample for stellar astrophysics and exoplanet science, thanks to the long-term baseline, high-precision observations of a large amount ( $\sim 200,000$ ) of stars (Brown et al. 2011; Huber et al. 2014; Mathur et al. 2017), which discovered 2,348 confirmed planets and 2,368 candidates (data from NASA Exoplanet Archive, EA hereafter; Akeson et al. 2013), contributing the majority of confirmed exoplanets and candidates. One of advantages of the Kepler sample is that it is suitable for studies on planetary systems in the Galactic context because they spread over different Galactic environments in a wide region up to several kpc in our Milky way (Fig 13 of PAST I, Chen et al. 2021). In order to have a reliable Galactic census of planets, one needs accurate characterizations of the Kepler stars (not just the planet candidate hosts). For most Kepler stars, some stellar parameters can be measured in relatively high precision, e.g., 7% for mass, 4% for radius and 112 K for effective temperature, while the uncertainty of stellar age is relatively large, e.g., 56% on average from isochrone fitting (Berger et al. 2020b). The kinematic properties (e.g., Galactic velocities and thin/thick disk memberships) of Kepler stars, however, still remain to be characterized. Furthermore, as we will show below, kinematic ages derived from Galactic velocity dispersions with AVR can be an important complementary to ages derived from other methods. Therefore, in this paper, we focus on the kinematic characterization of the Kepler stars.

To obtain the Galactic component memberships and kinematic age, we only need the 3D space motions without involving stellar evolutionary mode (e.g. Bensby et al. 2003; Soderblom 2010; Bergemann et al. 2014), which can be derived from radial velocity plus five astrometric parameters, i.e., parallax, celestial coordinates (Right Ascension and Declination) and the corresponding proper motions. Fortunately, the second Data release (DR2) of Gaia (e.g., Gaia Collaboration et al. 2016, 2018a,b) has provided the five astrometric parameters for more than 1.3 billion stars, including ~ 180,000 Kepler stars (Berger et al. 2020b). However, Gaia DR2 provides radial velocities only for bright stars (about 7.2 million with a magnitude of  $G \sim 4 - 13$ ) and thus only include ~ 10% of Kepler stars. Thus to make up for that, we rely on the spectroscopic data from LAMOST, which provides radial velocities for over 30% of all Kepler targets with no bias toward Kepler planet candidate hosts (De Cat et al. 2015; Ren et al. 2016; Fu et al. 2020). Besides, LAMOST also provides other physics parameters (e.g. [Fe/H], [ $\alpha$ /Fe], and the magnetic S index), which will enrich our knowledge of the stars and benefit the studies concerning stellar physical properties.

In this paper, we kinematically characterize 35,835 Kepler stars based on their astrometry and radial velocities provided by Gaia and LAMOST. Specifically, in section 2, we describe how to collect the stellar sample from Kepler, LAMOST and Gaia data. In section 3, we briefly introduce the kinematic methods to identify Galactic components (e.g., thin/thick disk) and to derive kinematic ages using the AVR. In section 4, we conduct a catalog of kinematic properties for 35,835 LAMOST-Gaia-Kepler stars. In section 5, we carry out some relative explorations based on our catalog. Finally, we summarize in section 6.

# 2. DATA COLLECTIONS

This section describes how we constructed the stellar sample from Gaia, Kepler and LAMOST for kinematic characterizing.

### 2.1. Kepler: Exoplanet Transit Surveys

We initialized our sample from the Kepler Data Release 25 (DR25) catalog, which contains 197,096 Kepler target stars as well as 8,054 Kepler Objects of Interest (KOIs) (Mathur et al. 2017). Here, we excluded KOIs flagged by 'False Positive (FAP)', leaving 4,034 planets (candidates). We also removed potential binaries by eliminating stars with Gaia DR2 re-normalized unit-weight error (RUWE) > 1.2 (Rizzuto et al. 2018; Berger et al. 2020a) as additional motions caused by binary orbits could affect the results of kinematic characterization. After these cuts, we are left with 175,280 Kepler stars and 3,620 planets (candidates).

# 2.2. Obtaining Five Astrometric Parameters from Gaia

To obtain the astrometric parameters for Kepler stars, we cross-matched them with Gaia DR2, which includes positions on the sky  $(\alpha, \delta)$ , parallaxes, and proper motions  $(\mu_{\alpha}, \mu_{\delta})$  for more than 1.3 billion stars with a limiting

Selection criteria	$N_{s}$	$N_{\rm p}$
Kepler DR25	197,096	8,054
No FAP		4,034
No binary	$175,\!280$	3,620
Cross-matched with Gaia	$163,\!454$	3,409
Cross-match with LAMOST DR4 $$	55,020	1,202
SNR > 10	$54,\!337$	$1,\!183$
RV reliability cut	49,261	1,098
Distance $< 1.54$ kpc	$35,\!835$	1,060

Table 1. Construction of the LAMOST-Gaia-Kepler star sample.

 $N_{\rm s}$  and  $N_{\rm p}$  are the numbers of stars and planets during the process of sample selection in section 2.

magnitude of G = 21 and a bright limit of G  $\approx$  3 (Gaia Collaboration et al. 2018a). The cross-matching was done by adopting the X match service of the Centre de Donnees astronomiques de Strasbourg (CDS, http://cdsxmatch.ustrasbg.fr, Boch et al. (2012)). After inspecting the distribution of separations, we select the separation limit of the cross-matching as where the distribution of separations displayed a minimum,  $\sim$  1.5 arcseconds. To ensure that the matched stars are of similar brightness, we also make a magnitude cut. The magnitude limit was set by inspecting the distribution of magnitude difference, which is 2 mag in Gaia G mag. For multiple matches satisfied these two criteria for the same star, we kept the one with the smallest angular separation. After these selections, we obtained 163,454 stars and 3,409 candidates.

### 2.3. Obtaining RV from LAMOST

Next, to obtained RV data, we rely on the spectroscopic data from LAMOST (Cui et al. 2012; Zhao et al. 2012). The LAMOST DR4 value-added catalog contains parameters derived from a total of 6.5 million stellar spectra for 4.4 million unique stars (Xiang et al. 2017). RVs,  $T_{\text{eff}}$ ,  $\log g$ , and [Fe/H] have been deduced using both the official LAMOST Stellar parameter Pipeline (LASP; Wu et al. 2011) and the LAMOST Stellar Parameter Pipeline at Peking University (LSP3; Xiang et al. 2015). The typical uncertainties for RVs,  $T_{\text{eff}}$ ,  $\log g$ , and [Fe/H] are 5.0 km s<sup>-1</sup>, 150 K, 0.25 dex, and 0.15 dex respectively.

We cross-matched the foregoing sample with LAMOST DR4 value-added catalog by using CDS with the same procedure detailed in section 2.2. By inspecting the distribution of separations and magnitude difference, the separation limit and magnitude cut were set as 5 arcseconds and 2.3 mag. Besides, we applied a quality cut of SNR > 10. To ensure the reliability of RV, we also cross-matched with the LAMOST DR7 catalog (http://dr7.lamost.org/) and remove the stars when the differences in RVs are larger than three times of the uncertainties. We only kept stars with a distance less than 1.54 kpc to the sun, which is the applicable limit of the revised kinematic characteristics and AVR in PAST I (Chen et al. 2021). Finally we obtained a LAMOST-Gaia-Kepler sample of 35,835 stars and 1,060 planets. In Table 1, we summarize the composition of the sample after each step mentioned above.

### 3. METHODS: CLASSIFICATION OF GALACTIC COMPONENTS AND ESTIMATION OF KINEMATIC AGES

In this section, we describe how we distinguish star into different Galactic components (section 3.2) and estimate stellar ages (section 3.3) with revisiting velocity ellipsoid and AVR from PAST I based on data from LAMOST and Gaia DR2.

## 3.1. Space Velocities and Galactic Orbits

We calculated the 3D Galactocentric cylindrical coordinates  $(R, \theta, Z)$  by adopting a location of the Sun of  $R_{\odot} = 8.34$  kpc (Reid et al. 2014) and  $Z_{\odot} = 27$  pc (Chen et al. 2001). By adopting the formulae and matrix equations presented in Johnson & Soderblom (1987), we calculate the Galactic rectangular velocities relative to the Sun (U, V, W) and their errors for the LASMOT-Gaia-Kepler sample. Here U is positive when pointing to the direction of the Galactic center, V is positive along the direction of the Sun orbiting around the Galactic center, and W is positive when pointing towards the North Galactic Pole. To obtain the Galactic rectangular velocities relative to the local standard of rest (LSR) ( $U_{\text{LSR}}, V_{\text{LSR}}, W_{\text{LSR}}$ ), we adopted the solar peculiar motion [ $U_{\odot}, V_{\odot}, W_{\odot}$ ] = [9.58, 10.52, 7.01] km s<sup>-1</sup> (Tian et al. 2015). Cylindrical velocities  $V_R$ ,  $V_{\theta}$ , and  $V_Z$  are defined as positive with increasing R,  $\theta$ , and Z, with the latter towards the North Galactic Pole.

		$k \;({\rm km \; s^{-1}})$ ——	β		
	value	1 $\sigma$ interval	value	1 $\sigma$ interval	
U	23.66	(23.07, 24.32)	0.34	(0.33, 0.36)	
V	12.49	(12.05, 12.98)	0.43	(0.41, 0.45)	
W	8.50	(8.09, 8.97)	0.54	(0.52, 0.56)	
$V_{\rm tot}$	27.55	(26.84, 28.37)	0.40	(0.38, 0.42)	

Table 2. Fitting parameters of the Age-Velocity dispersion relation from PAST I (Chen et al. 2021).

# 3.2. Classification of Galactic Components

In this section, by adopting the wide-used kinematic approaches (Bensby et al. 2003, 2014), we classify stars into different Galactic components (i.e., thin disk, thick disk, halo and Hercules stream) based on their Galactic positions and velocities. The kinematic methods assumes the Galactic velocities to the LSR ( $U_{\text{LSR}}, V_{\text{LSR}}, W_{\text{LSR}}$ ) have multi-dimensional Gaussian distributions:

$$f(U, V, W) = k \times \exp\left(-\frac{(U_{\text{LSR}} - U_{asym})^2}{2\sigma_U^2} - \frac{(V_{\text{LSR}} - V_{asym})^2}{2\sigma_V^2} - \frac{W_{\text{LSR}}^2}{2\sigma_W^2}\right),\tag{1}$$

where  $\sigma_U$ ,  $\sigma_V$ , and  $\sigma_W$  are the characteristic velocity dispersions, and  $V_{asym}$  and  $U_{asym}$  are the asymmetric drifts for different components (the thin disk, the thick disk, the halo, and the Hercules stream). The normalization coefficient are defined as

$$k = \frac{1}{(2\pi)^{3/2} \sigma_U \sigma_V \sigma_W}.$$
(2)

The relative probabilities between two different components, i.e., the thick-disk-to-thin-disk (TD/D), thick-disk to halo (TD/H), the Hercules-to-thin-disk (Herc/D), and the Hercules-to-thick-disk (Herc/TD) can be calculated as

$$\frac{TD}{D} = \frac{X_{\rm TD}}{X_{\rm D}} \cdot \frac{f_{\rm TD}}{f_{\rm D}}, \qquad \frac{TD}{H} = \frac{X_{\rm TD}}{X_{\rm H}} \cdot \frac{f_{\rm TD}}{f_{\rm H}}, \tag{3}$$

$$\frac{Herc}{D} = \frac{X_{\text{Herc}}}{X_{\text{D}}} \cdot \frac{f_{\text{Herc}}}{f_{\text{D}}}, \quad \frac{Herc}{TD} = \frac{X_{\text{Herc}}}{X_{\text{TD}}} \cdot \frac{f_{\text{Herc}}}{f_{\text{TD}}}, \tag{4}$$

where X is the fraction of stars for a given component.

Here we adopt the revised kinematic characteristics and X factor of different Galactic components in PAST I (Chen et al. 2021) and calculate the above Galactic component membership probabilities for the LAMOST-Gaia-Kepler stars. Then we classified them into different Galactic components by adopting the same criteria as in Bensby et al. (2014), which are:

(1) thin disk: TD/D < 0.5 & Herc/D < 0.5;

- (2) thick disk: TD/D > 2 & TD/H > 1 & Herc/TD < 0.5;
- (3) halo: TD/D > 2 & TD/H < 1 & Herc/TD < 0.5;
- (4) Hercules: Herc/D > 1 & Herc/TD > 1.

#### 3.3. Calculating Kinematic Age and Uncertainty

As described in section 3.6 of PAST I, for a group of stars, the typical kinematic age can be derived by using the Age-velocity dispersion relation (AVR), which gives

$$Age_{kin} \equiv t = \left(\frac{\sigma}{k \, km \, s^{-1}}\right)^{\frac{1}{\beta}} \, Gyr, \tag{5}$$

where  $\sigma$  is the velocity dispersion, which is defined as the root mean square of stellar Galactic velocity.  $k, \beta$  are the fitting coefficients of AVR. By means of error propagation, the relative uncertainty of kinematic age can be estimated

as:

$$\frac{\Delta t}{t} = \sqrt{\left(\frac{\partial \ln t}{\partial \beta} \Delta \beta\right)^2 + \left(\frac{\partial \ln t}{\partial k} \Delta k\right)^2 + \left(\frac{\partial \ln t}{\partial \sigma} \Delta \sigma\right)^2} = \sqrt{\left(\ln \frac{t}{Gyr}\right)^2 \left(\frac{\Delta \beta}{\beta}\right)^2 + \frac{1}{\beta^2} \left(\frac{\Delta k}{k}\right)^2 + \frac{1}{\beta^2} \left(\frac{\Delta \sigma}{\sigma}\right)^2},$$
(6)

where  $\Delta$  represents the absolute uncertainty. Here we adopt the two coefficients  $(k, \beta)$  derived in PAST I (Chen et al. 2021), which are listed in Table 2.

### 4. A CATALOG OF LAMOST-GAIA-KEPLER STARS WITH KINEMATIC CHARACTERIZATIONS

Applying the methods described in section 3 to the LAMOST-Gaia-Kepler sample (section 2), we characterize the kinematic properties, e.g., Galactic orbit, velocities and the relative membership probabilities between different Galactic components (TD/D, TD/H, Her/D and Her/TD) for the 35,835 LAMOST-Gaia-Kepler stars (Table 3). Among these stars, there are 764 stars hosting 1,060 planets.

Based on the derived relative membership probabilities between different Galactic components (TD/D, TD/H, Her/D and Her/TD in Table 3), following the criteria as mentioned in section 3.2, we then classify the 35,835 stars into four Galactic components, i.e., thin disk, thick disk, Hercules stream and halo. For these stars not belonging to the above four components, we assign them into a category dubbed 'in between' referring to Bensby et al. (2014).

The numbers of stars and planets in different Galactic components are summarized in Table 4. As can be seen, about 87.1% (31,218/35,835) of stars in our sample are in thin disk and about 5.1% (1,832/35,835) stars are in thick disk. The fractions of halo and Hercules stream stars are about 0.16% (59/35,835) and 1.5% (545/35,835). There are another ~ 6.1% (2,181/35,835) stars belonging to the 'in between' category.

To display the distribution of velocities, in Figure 1, we plot the Toomre diagram of the LAMOST-Gaia-Kepler stars. As can be seen, most stars with low velocities ( $V_{\text{tot}} \leq 70 \text{km s}^{-1}$ ) are in the thin disk, while those with moderate velocities ( $V_{\text{tot}} \sim 70 - 180 \text{km s}^{-1}$ ) are mainly in thick disk. The velocities of halo stars are all larger than 200 km s<sup>-1</sup>. For the Hercules stream, most of these stars have  $V_{\text{LSR}}$  around  $-50 \pm 9 \text{ km s}^{-1}$  and  $(U_{\text{LSR}}^2 + W_{\text{LSR}}^2)^{1/2} \sim 30-90 \text{ km s}^{-1}$ . This is well consistent with the results derived by previous works (e.g. Feltzing et al. 2003; Adibekyan et al. 2013; Bensby et al. 2014; Bonaca et al. 2017).

Then we compare the distributions of chemical abundances for different components in Figure 2. As expected, the thick disk stars are metal-poorer (~ 0.3 dex) and  $\alpha$ -richer (~ 0.1 dex) than the thin disk stars. The Hercules stream stars have velocities and chemical abundances which are between those of thin and thick disk stars. For the halo, these stars have the highest Galactic velocities, poorest [Fe/H] and richest [ $\alpha$ /Fe]. In Figure 3, we plot the total velocity  $V_{\text{tot}}$ , [Fe/H] and [ $\alpha$ /Fe] as a function of TD/D. As can be seen, with the increasing of TD/D,  $V_{\text{tot}}$  and [ $\alpha$ /Fe] increase, while [Fe/H] decreases. Because there seems no clear trend between velocity dispersions and ages for stars in the halo and Hercules stream (Chen et al. 2021), here we only compare the age distributions for stars in the Galactic disk. The typical ages are  $2.43^{+0.32}_{-0.19}$  Gyr,  $9.83^{+1.30}_{-1.02}$  Gyr for thin and thick disk respectively. The Numbers of stars and planets, kinematic properties and chemical abundances of different Galactic components are summarized in Table 4.

For the sake of completeness, we also add the stellar parameters that used during the process of our kinematic characterization (e.g., parallax, proper motion and RV) and other basic stellar parameters (e.g.,  $T_{\text{eff}}$ , log g, [Fe/H] and  $[\alpha/\text{Fe}]$ ) into the catalog. In what follows, we conduct some further analyses and discussions on this catalog.

# 5. FURTHER CATALOG ANALYSES AND DISCUSSIONS

In this section, based on the LAMOST-Gaia-Kepler catalog, we perform some tests and comparisons to verify the reliability of the kinematic age (section 5.1), explore the evolution of stellar magnetic activity with kinematic ages (section 5.2), and analyze the kinematic properties of Kepler planet candidate host stars (section 5.3).

#### 5.1. Verifying the Kinematic Age

In order to verify the reliability of the kinematic age, we compare it with ages derived from other methods, such as asteroseismology, gyrochronology, and isochrone fitting in this subsection.

With seismic parameters  $\Delta \nu$  and  $\nu_{\text{max}}$  derived from Kepler asteroseismic data, and spectroscopic parameters from Apache Point Observatory Galactic Evolution Experiment (APOGEE) project (Majewski et al. 2017), Pinsonneault et al. (2018) provided asteroseismic ages for 6,676 evolved stars.



6



Figure 1. The Toomre diagram of LAMOST-Gaia-Kepler stars for different Galactic components. Top panel shows the full range of velocities while Bottom panel zooms in on the region where a majority of the sample is located. The diagram are colour-coded to represent different memberships. Dotted lines show constant values of the total Galactic velocity  $V_{\text{tot}} = (U_{\text{LSR}}^2 + V_{\text{LSR}}^2 + W_{\text{LSR}}^2)^{1/2}$ , in steps of 100 and 50 km s<sup>-1</sup> respectively, in the two plots.

Gyrochronology can estimate the age of a main sequence star based on its rotation period (Barnes 2010; Barnes & Kim 2010). We use rotation data for Kepler stars from McQuillan et al. (2013a,b, 2014), and derive gyrochronology ages for 5,851 stars by adopting the method provided by Spada & Lanzafame (2020).

Fitting isochrone grid provided by the latest MIST models (Dotter 2016; Choi et al. 2016; Paxton et al. 2011, 2013, 2015, 2018), Berger et al. (2020b) obtained isotropic ages for 186,301 Kepler stars with a typical uncertainty of ~ 56%. Here, we only consider stars with relatively reliable isochrone ages by applying the selection criteria as suggested



Figure 2. The cumulative distributions of velocity  $V_{\text{tot}}$  (a), [Fe/H] (b) and  $[\alpha/\text{Fe}]$  (c) for LAMOST-Gaia-Kepler stars of different Galactic components, i.e., thin disk, thick disk, halo, and Hercules stream.

by Berger et al. (2020b), i.e., stars with GOF>0.99, having TAMS less than 20 Gyr, and metallicity measured by spectroscopic method. We also remove stars with isochrone age longer than 14 Gyr.

In Figure 4, we compare kinematic age inferred from this work and ages derived with other methods to each other. Subplots in each column and row share the same x-axis and y-axis, respectively. We label the methods to derive age and the relevant literatures in the corresponding coordinates. In each subplot, we cross-match the two samples shown in the x and y axes, and print the number of stars in the matched sample in the left upper corner. We sort each sample in ascending order according to the x-axis, and divide them into 10 bins with approximately equal size. For asteroseismic, gyrochronology, and isochrone ages, the dots and errorbars shown in Figure 4 are median ages and median values of relative error. For the kinematic age, we calculate the age and uncertainty with methods mentioned in section 3.3.

Since ages derived from asteroseismology are mainly for giant stars and gyrochronology applies only to main sequence stars, there is no common star between these two samples and thus comparison has never been made between gyrochronology age and asteroseismic age yet. In the upper panels, isochrone age is compared with asteroseismic age and gyrochronology age. Isochrone age exhibits systematic deviation from asteroseismic age and apparent discrepancy with gyrochronology age for stars younger than 3 Gyrs, though in most of cases they are generally consistent with each other due to the large uncertainty of isochrone age. In the lower panels, we show comparisons between kinematic age and other ages. In general, kinematic age matches well with ages derived from asteroseismic, gyrochronology, and isochrone, with relative large differences only in the young end (<2 Gyrs for asteroseismic age, <3 Gyrs for gyrochronology and isochrone ages). Such a discrepancy is not unexpected because the stellar velocity dispersion at early days might be dominated by the initial condition rather than dynamical evolution and thus age derived from velocity dispersion would be overestimated.

#### 5.2. Stellar Magnetic Activity Evolves with Kinematic Age

Stellar properties, such as rotation period and stellar activity, are indicators of stellar age. We explore the connections between them and kinematic age in this subsection.

We match our LAMOST-Gaia-Kepler sample with rotation periods from McQuillan et al. (2013a,b, 2014) by KIC, obtaining 10,548 common stars. We sort the sample according to the TD/D value, divide them into 10 groups with approximately equal size and calculate kinematic ages and uncertainties for each group using the method mentioned in section 3.3. For each bin, we calculated the median rotation period and median value of relative error, and show them with dots and errobars in the upper panel of Figure 5. As kinematic age increases, rotation period becomes longer, which is in agreement with the gyrochronology theory (Barnes & Kim 2010; Barnes 2010). For comparison, we show the age (4.57 Gyr) and rotation period (16.09 days) of the Sun with a yellow dot. As can be seen, the Sun is a typical star which fits well to the kinematic age–rotation period trend.

In our LAMOST–Gaia–Kepler sample, 20,417 stars have the S-index measurements based on the LAMOST spectra (Zhang et al. 2020). We group the sample into 10 subgroups. For each bin, we calculate the median value of S-index and the uncertainty is set as median value of relative error, which are plotted as dots and errobars in the upper panel



Figure 3. Kinematic and chemical properties (from top to bottom: total velocity  $V_{tot}$ , abundances [Fe/H] and [ $\alpha$ /Fe]) as functions of the relative probability between thick disk to thin disk (TD/D) for the LAMOST-Gaia-Kepler stars. Medians and 1  $\sigma$  dispersions are marked in the plots. Histograms of V<sub>tot</sub>, [Fe/H] and [ $\alpha$ /Fe] are shown in the right panels. Histogram of TD/D is displayed in the topmost. The vertical dashed lines represents where TD/D = 1.

of Figure 5. As stars aging, stellar activity reduces and the median value of S-index decreases as expected. For S-index of the Sun, we adopt the mean value  $\langle S \rangle = 0.1694$  suggested by Egeland et al. (2017), and plot it in the figure with a yellow dot. As can be seen, the Sun is exceptionally quiet as compared to stars in the LAMOST–Gaia–Kepler sample, confirming the results of previous studies (e.g., Reinhold et al. 2020).

Stellar magnetic activity can also be revealed by photometry. In Figure 6, we investigate how the photometric noise levels of stars change with kinematic age. For every target, Kepler DR 25 provides Combined Differential Photometric Precision (CDPP, Christiansen et al. 2012) for 14 different time scales, which characterises the noise level in Kepler lightcurves. In order to minimize other parameters that may affect CDPP other than age, such as evolve state, magnitude, and spectral type, we restrict out sample to stars with  $\log g > 4$ , Kepler magnitude (kepmag) lower than 14, and effective temperature between 4,700K to 6,500K, which contains 14,372 stars. Since CDPP is partly contributed by photon noise, following the method of Bastien et al. (2013), we subtract photon noise in quadrature from CDPP, i.e., CDPP-photon noise. Again, we group the star sample into 10 bins then calculate their corresponding kinematic ages and the median value of CDPP-photon noise. We show the CDPP-photon – Age noise relationship in Figure 6, each line with a different color presents CDPP with a different duration. For longer duration, CDPP-photon noise decrease as stars aging, 15 hours CDPP-photon noise drops from 65 ppm to 25 ppm as kinematic age grows from 0.3 Gyr to 12 Gyr. Gilliland et al. (2011) modelled the stellar noise level as a function of age and found that the total

Table 3.	The catalogue	of the LAMOST-Gaia-	Kepler stellar sample

Parameters obtained from Gaia, LAMOST and NASA esoplanet archive (EA)           1         Gaia ID         Long         Unique Gaia source identifier           2         LAMOST JD         string         LAMOST unique spectral ID           3         Kepler JD         integer         Kepler Input Catalog (KIC) ID           4         Gaia RA         Double         deg         Barycentric right ascension           5         Gaia Dec         Double         deg         Barycentric Declination           6         Gaia parallax         Double         mass         Statuated error of parallax           7         Gaia e parallax         Double         mass         Thandard error of poreper motion in right ascension direction           9         Gaia e-punce         Double         mas yr <sup>-1</sup> Proper motion in declination direction           10         Gaia G mag         Double         mas yr <sup>-1</sup> Proper motion in declination direction           11         Gaia a c-punce         Double         mas yr <sup>-1</sup> Standard error of poreper motion in right ascension direction           12         Gaia G mag         Double         mas yr <sup>-1</sup> Standard error of poreper motion in declination direction           13         Kepler mag         Double         mag         Kepl	Column	Name	Format	Units	description
1       Gaia.ID       Long       Unique Gaia source identifier         2       LAMOST.ID       string       LAMOST unique spectral ID         3       Kepler.ID       integer       Kepler Input Catalog (KIC) ID         4       Gaia RA       Double       deg       Barycentric Declination         5       Gaia Dec       Double       mas       Absolute stellar parallax         7       Gaia eparallax       Double       mas       Absolute stellar parallax         8       Gaia parallax       Double       mas yr <sup>-1</sup> Proper motion in dipt ascension direction         9       Gaia eparata       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         10       Gaia eparata       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         12       Gaia G mag       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         13       Kepler mag       Double       mas yr <sup>-1</sup> Standard error of proper motion in direction         14 $T_{eff}$ Float       K       Effective temperature from LAMOST         14 $T_{eff}$ Float       K       Error of metallicity from LAMOST         15		Parame	ters obtained	l from Gaia,	LAMOST and NASA exoplanet archive (EA)
2       LAMOST JD       string       LAMOST unique spectral ID         3       Kepler.ID       integer       Kepler Input Catalog (KIC) ID         4       Gaia DA       Double       deg       Barycentric Declination         5       Gaia Dare       Double       deg       Barycentric Declination         6       Gaia parallax       Double       mass       Statulated error of parallax         8       Gaia pumde       Double       mass       Statulated error of proper motion in right ascension direction         10       Gaia e-punde       Double       mass yr <sup>-1</sup> Standard error of proper motion in declination direction         11       Gaia e-punde       Double       mass yr <sup>-1</sup> Standard error of proper motion in dight ascension direction         12       Gaia Gaia gandee       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         13       Kepler mag       Double       mas yr <sup>-1</sup> Standard error of proper motion in fight ascension direction         14 $T_{aff}$ Float       K       Effective temperature from LAMOST         15       e.T <sub>eff</sub> Float       K       Effective temperature from LAMOST         16       log g       Float       K       Effective temperature	1	Gaia_ID	Long	`	Unique Gaia source identifier
2       LAMOST.ID       string       LAMOST unique spectral ID         3       Kepler.ID       integer       Kepler Input Catalog (KIC) ID         4       Gaia RA       Double       deg       Barycentric right accession         5       Gaia Dee       Double       mas       Absolute stelar parallax         6       Gaia parallax       Double       mas       Absolute stelar parallax         7       Gaia eparallax       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         9       Gaia eparate       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         10       Gaia or parate       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         12       Gaia G mag       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         13       Kepler mag       Double       mag       Kraft apper apparent magnitude         14       T <sub>eff</sub> Float       K       Error of functive temperature from LAMOST         15 $e_{e}T_{eff}/H$ Float       dex       actif centeris abundance from LAMOST         19 $e_{e}[r/Fe]       Float       dex       actif centeris abundance from LAMOST$					
3       Kepler JD       integer       Kepler Input Catalog (KIC) ID         4       Gaia RA       Double       deg       Barycentric right ascension         5       Gaia parallax       Double       deg       Barycentric Declination         6       Gaia parallax       Double       mas       Atsolute stellar parallax         7       Gaia epmra       Double       mas yr <sup>-1</sup> Proper motion in right ascension direction         9       Gaia epmea       Double       mas yr <sup>-1</sup> Proper motion in declination direction         10       Gaia pomdec       Double       mas yr <sup>-1</sup> Proper motion in declination direction         11       Gaia condec       Double       mas yr <sup>-1</sup> Proper motion in declination direction         12       Gaia G mag       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         12       Gaia C mag       Double       mas yr <sup>-1</sup> Natadra' error of proper motion in declination direction         13       Kepler mag       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         14 $T_{eff}$ Float       K       Effective temperature from LAMOST         15       e.log g       Float       K       Effective	2	LAMOST_ID	string		LAMOST unique spectral ID
3       Kepter LD       integer       Reptor input Catalog (RCO) ID         4       Gaia RA       Double       deg       Barycentric Declination         5       Gaia Dec       Double       mas       Absolute stellar parallax         7       Gaia e-parallax       Double       mas       Standard error of proper motion in right ascension direction         9       Gaia e-para       Double       mas yr <sup>-1</sup> Proper motion in declination direction         10       Gaia aprude       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         11       Gaia e-paral       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         12       Gaia formag       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         13       Kepler mag       Double       mag yr <sup>-1</sup> Standard error of proper motion in LAMOST         14       Terr       Float       K       Effective temperature from LAMOST         16       log g       Float       K       Error of surface gravity from LAMOST         17       e.log g       Float       dex       Aretro of float dual echocity from LAMOST         19       e.log /Flei       Float       dex	9				
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	3	Kepler_ID	integer		Kepler Input Catalog (KIC) ID
5       Gaia Dec       Double       deg       Barycentric Declination         6       Gaia parallax       Double       mas       Absolute stellar parallax         7       Gaia e-parala       Double       mas       Yeahadraf error of proper motion in right ascension direction         9       Gaia e-para       Double       mas yr <sup>-1</sup> Proper motion in declination direction         10       Gaia e-parac       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         11       Gaia e-parac       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         12       Gaia forma       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         13       Kepler mag       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         14       Tert       Float       K       Effective temperature temperature         16       log g       Float       K       Effective temperature         17       e.log g       Float       dex       Metallicity from LAMOST         18       [Fe/H]       Float       dex       Error of aclements abundance from LAMOST         21       e.log /Fel       Float       dex <td>4</td> <td>Gaia RA</td> <td>Double</td> <td>deg</td> <td>Barycentric right ascension</td>	4	Gaia RA	Double	deg	Barycentric right ascension
6       Gaia parallax       Double       mas       Absolute stellar parallax         7       Gaia e parallax       Double       mas       Standard error of parallax         8       Gaia purna       Double       mas yr <sup>-1</sup> Proper motion in right ascension direction         9       Gaia e-purde       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         10       Gaia e-purde       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         12       Gaia G mag       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         13       Kepler mag       Double       mas yr <sup>-1</sup> Standard error of proper motion in declination direction         14 $T_{eff}$ Float       K       Effective temperature from LAMOST         15 $e_{$	5	Gaia Dec	Double	deg	Barvcentric Declination
7Gaia e.parallaxDoublemasStandard error of parallax8Gaia puruaDoublemas yr^{-1}Proper motion in right ascension direction9Gaia e.puraDoublemas yr^{-1}Standard error of proper motion in right ascension direction10Gaia e.puraDoublemas yr^{-1}Standard error of proper motion in declination direction11Gaia e.puraDoublemas yr^{-1}Standard error of proper motion in declination direction12Gaia G magDoublemagKepler apparent magnitude13Kepler magDoublemagKepler apparent magnitude14 $T_{eff}$ FloatKEffective temperature from LAMOST15e.TerfFloatKError of factilicity16log gFloatError of antallicity17e.log gFloatdexMetallicity from LAMOST18[Fe/H]FloatdexError of a clements abundance21e.[ $\alpha$ /Fe]FloatdexError of a clements abundance22vrDoublekm s^{-1}Radia telocity23e.vrDoublekm s^{-1}Radia telocity24NpintegerPlanet (candidate) multiplicity25RDoublekpcGalactocentric Cylindrical arial distance26 $\theta$ Doublekm s^{-1}Galactocentric Cylindrical velocities30VzDoublekm s^{-1}Galactocentric Cylindrical velocities31U_LSR	6	Gaia parallax	Double	mas	Absolute stellar parallax
8Gaia pura Gaia e_puraDouble mas yr^1Proper motion in right ascension direction9Gaia e_pura Gaia pundeDouble mas yr^1Standard error of proper motion in right ascension direction11Gaia e_pundecDouble mas yr^1Standard error of proper motion in declination direction12Gaia G mag Gaia G magDouble magGaia G band apparent magnitude14 $T_{eff}$ Float KK15 $e.T_{eff}$ Float KK16log gFloat KK17 $e.log g$ Float FloatK18[Fe/H]Float FloatError of surface gravity from LAMOST19 $e.JFe/H$ Float floatdex20 $[a/Fe]$ Float floatdex21 $e.log g$ Float floatdex22vrDouble ms^{-1}Radia velocity from LAMOST23 $e.a/Fe]$ Float floatdex24NpFloat floatdex25RDouble hypGalactocentric Cylindrical radial distance260Double kpcGalactocentric Cylindrical azimuth angle27ZDouble km s^{-1}Galactocentric Cylindrical azimuth angle28VgDouble km s^{-1}Galactocentric Cylindrical azimuth angle29VgDouble km s^{-1}Galactocentric Cylindrical azimuth angle28VgDouble km s^{-1}Galactocentric Cylindrical azimuth angle29Vg <td< td=""><td>7</td><td>Gaia e_parallax</td><td>Double</td><td>mas</td><td>Standard error of parallax</td></td<>	7	Gaia e_parallax	Double	mas	Standard error of parallax
9Gaia e.pmra Gaia e.pmdeDouble mas yr^-1Standard error of proper motion in right ascension direction10Gaia e.pmde Gaia e.pmdeDouble mas yr^-1Proper motion in declination direction12Gaia G mag Gaia e.pmdeDouble magStandard error of proper motion in declination direction12Gaia G mag Gaia C magDouble mag Mepler magGaia G band apparent magnitude13Kepler mag Log gDouble mag Mepler mag FloatFleat K K Error of effective temperature from LAMOST15e.T <sub>eff</sub> FloatFloatError of surface gravity from LAMOST16log gFloatdex dexMetallicity from LAMOST17e.log gFloatdex dexMetallicity Error of metallicity10[a/Fe]Float dexdex error of a celements abundance from IAMOST21e.[a/Fe] FloatFloat dex dex error of a celements abundance22vr vr Double km s^{-1}Error of radial velocity Planet (candidate) multiplicity24NpintegerPlanet (candidate) multiplicity25R Double km s^{-1}Double Galactocentric Cylindrical azimuth angle e Galactocentric Cylindrical azimuth angle26 $\theta$ Double km s^{-1}Galactocentric Cylindrical velocities29V_0 Double km s^{-1}Galactocentric Cylindrical velocities30VZ Double km s^{-1}Galactocentric Cylindrical velocity to the LSR31U_{LSR} Double km s^{-1}<	8	Gaia pmra	Double	$\mathrm{mas} \mathrm{vr}^{-1}$	Proper motion in right ascension direction
10Gaia pmdec Gaia e_pmdecDouble Double mas yr^-1Proper motion in declination direction Gaia G mag Gaia G magDouble mag TStandard error of proper motion in declination direction (Gaia G mag) Gaia G mag11Gaia G mag Gaia G magDouble magGaia G had apparent magnitude (Kepler apparent magnitude)12Gaia G mag Double magDouble magKepler apparent magnitude13Kepler mag B Gaia G band B gFloatK14 $T_{eff}$ B GgFloatK15 $c.T_{eff}$ E floatFloatSurface gravity from LAMOST16log gFloatError of surface gravity from LAMOST17e-log gFloatdexMetallicity from LAMOST19e_[Fe/H] E floatFloat dexError of a dementa abundance from LAMOST20[a/Fe] E floatFloat dexError of radial velocity Fror of radial velocity Planet (candidate) multiplicity21 $e_i[a/Fe]$ P integerFloat Planet (candidate) multiplicity22vr P Double degGalactocentric Cylindrical radial distance Galactocentric Cylindrical avertical height25R Double VgDouble km s^{-1}Galactocentric Cylindrical avertical R velocities29Vg Double km s^{-1}Galactocentric Cylindrical avertical R velocities29Vg Double km s^{-1}Galactocentric Cylindrical R velocities29Vg Double km s^{-1}Galactocentric Cylindrical R velocities29Vg Dou	9	Gaia e_pmra	Double	$mas vr^{-1}$	Standard error of proper motion in right ascension direction
Interpret       Double       may $T^{-1}$ Standard error of proper motion in declination direction         12       Gaia G mag       Double       mag       Gaia G band apparent magnitude         13       Kepler mag       Double       mag       Kepler apparent magnitude         14 $T_{eff}$ Float       K       Effective temperature from LAMOST         15 $e.T_{eff}$ Float       K       Error of surface gravity from LAMOST         16       log g       Float       dex       Metallicity from LAMOST         18       [Fe/H]       Float       dex       Metallicity from LAMOST         19 $e.[Fe/H]$ Float       dex       Metallicity from LAMOST         20       [a/Fe]       Float       dex       a elements abundance from LAMOST         21 $e.[a/Fe]$ Float       dex       a elements abundance         22       vr       Double       km s^{-1}       Error of a elements abundance         23 $e.vr$ Double       km s^{-1}       Error of radial velocity         24       Np       integer       Planeters derived in this work         25       R       Double       kpc       Galactocentric Cylindrical azimuth agle	10	Gaia pmdec	Double	$mas vr^{-1}$	Proper motion in declination direction
12       Gaia G mag       Double       mag       Gaia G band apparent magnitude         13       Kepler mag       Double       mag $Kepler$ apparent magnitude         14 $T_{eff}$ Float       K       Effective temperature from LAMOST         15 $e.T_{eff}$ Float       K       Effective temperature         16       log g       Float       K       Error of effective temperature         17 $e.log g$ Float       dex       Metallicity from LAMOST         18       [Fe/H]       Float       dex       Error of metallicity         20       [ $\alpha/Fe$ ]       Float       dex       Error of a clements abundance from LAMOST         21 $e.[Fe/H]$ Float       dex       Error of a clements abundance         22       vr       Double       km s^{-1}       Radial velocity from LAMOST         23 $e.vr$ Double       km s^{-1}       Error of a clements abundance         24       Np       integer       Planet (candidate) multiplicity         24       Np       integer       Galactocentric Cylindrical ariad distance         26 $\theta$ Double       km s^{-1}       Galactocentric Cylindrical $2$ velocitites <tr< td=""><td>11</td><td>Gaia e_pmdec</td><td>Double</td><td><math>mas vr^{-1}</math></td><td>Standard error of proper motion in declination direction</td></tr<>	11	Gaia e_pmdec	Double	$mas vr^{-1}$	Standard error of proper motion in declination direction
13       Kepler mag       Double       mag $Kepler$ apparent magnitude         14 $T_{eff}$ Float       K       Effective temperature from LAMOST         15 $e.T_{eff}$ Float       K       Effective temperature         16       log g       Float       K       Error of surface gravity from LAMOST         17 $e.log g$ Float       dex       Metallicity from LAMOST         18       [Fe/H]       Float       dex       Error of metallicity         20       [a/Fe]       Float       dex       Calements abundance from LAMOST         21 $e.[a/Fe]$ Float       dex       a clements abundance from LAMOST         22       vr       Double       km s <sup>-1</sup> Radia Voicity from LAMOST         23 $e.vr$ Double       km s <sup>-1</sup> Radia Voicity from LAMOST         24 $N_p$ integer       Planet (candidate) multiplicity         Planet dex       Error of radial velocity from LAMOST         25       R       Double       kpc       Galactocentric Cylindrical radial distance         26 $\theta$ Double       kpc       Galactocentric Cylindrical avental magle         27       Z	12	Gaia G mag	Double	mag	Gaia G band apparent magnitude
14 $T_{eff}$ Float       K       Effective temperature from LAMOST         15 $e.T_{eff}$ Float       K       Error of effective temperature         16 $\log g$ Float       K       Error of surface gravity from LAMOST         17 $e.log g$ Float       Error of surface gravity from LAMOST         18       [Fe/H]       Float       dex       Metallicity from LAMOST         19 $e.[Fe/H]$ Float       dex       Callentiation         20       [a/Fe]       Float       dex       Error of a clements abundance         21 $e.[Fe/H]$ Float       dex       Error of radial velocity         21 $e.[G/Fe]$ Float       dex       Error of radial velocity         23 $e.vr$ Double       km s <sup>-1</sup> Radial velocity         24 $N_p$ integer       Planet (candidate) multiplicity         25       R       Double       kpc       Galactocentric Cylindrical artical height         28 $V_R$ Double       km s <sup>-1</sup> Galactocentric Cylindrical $2$ velocities         30 $V_Z$ Double       km s <sup>-1</sup> Galactocentric $2$ velocities         31	13	Kepler mag	Double	mag	Kepler apparent magnitude
15e. $T_{eff}$ FloatKError of effective temperature16 $\log g$ FloatKError of surface gravity from LAMOST17e.log gFloatError of surface gravity from LAMOST18[Fe/H]FloatdexMetallicity from LAMOST19e.[Fe/H]FloatdexError of metallicity20[a/Fe]FloatdexError of a clements abundance from LAMOST21e.[a/Fe]FloatdexError of a clements abundance22vrDoublekm s <sup>-1</sup> Radial velocity from LAMOST23e.vrDoublekm s <sup>-1</sup> Error of radial velocity24 $N_p$ integerPlanet (candidate) multiplicityParameters derived in this work25RDoublekpc26 $\theta$ Doublekm s <sup>-1</sup> 27ZDoublekm s <sup>-1</sup> 28 $V_R$ Doublekm s <sup>-1</sup> 29 $V_{\theta}$ Doublekm s <sup>-1</sup> 31 $U_{LSR}$ Doublekm s <sup>-1</sup> 32e. $U_{LSR}$ Doublekm s <sup>-1</sup> 33 $U_{LSR}$ Doublekm s <sup>-1</sup> 34e. $V_{LSR}$ Doublekm s <sup>-1</sup> 34e. $V_{LSR}$ Doublekm s <sup>-1</sup> 34e. $V_{LSR}$ Doublekm s <sup>-1</sup> 35 $W_{LSR}$ Doublekm s <sup>-1</sup> 36e. $W_{LSR}$ Doublekm s <sup>-1</sup> 37 $TD/D$ Doublekm s <sup>-1</sup> 38 $T_{LSR}$ <td< td=""><td>14</td><td>Т.а</td><td>Float</td><td>K</td><td>Effective temperature from LAMOST</td></td<>	14	Т.а	Float	K	Effective temperature from LAMOST
16log gFloatSurface gravity from LAMOST17e.log gFloatError of surface gravity from LAMOST18[Fe/H]FloatdexMetallicity from LAMOST19e.[Fe/H]FloatdexError of metallicity20[a/Fe]Floatdexa elements abundance from LAMOST21e.[a/Fe]FloatdexError of a clements abundance22vrDoublekm s <sup>-1</sup> Radia velocity from LAMOST23e.vrDoublekm s <sup>-1</sup> Error of radial velocity24NpintegerPlanet (candidate) multiplicity25RDoublekgcGalactocentric Cylindrical azimuth angle26 $\theta$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical velocities29V <sub>R</sub> Doublekm s <sup>-1</sup> Galactocentric Cylindrical R velocities29V <sub>R</sub> Doublekm s <sup>-1</sup> Galactocentric Cylindrical R velocities31ULSRDoublekm s <sup>-1</sup> Galactocentric Cylindrical R velocities33V <sub>L</sub> Doublekm s <sup>-1</sup> Galactocentric Cylindrical R velocities31ULSRDoublekm s <sup>-1</sup> Galactocentric Z velocities32e.ULSRDoublekm s <sup>-1</sup> Galactocentric Z velocity to the LSR33VLSRDoublekm s <sup>-1</sup> Cartesian Galactocentric x velocity to the LSR33VLSRDoublekm s <sup>-1</sup> Cartesian Galactocentric x velocity to the LSR34e.VLSRDoublekm s <sup>-1</sup>	15	e T <sub>eff</sub>	Float	K	Error of effective temperature
17e.log gFloatError of surface gravity from LAMOST18[Fe/H]FloatdexMetallicity from LAMOST19e.[Fe/H]FloatdexError of metallicity20[ $\alpha$ /Fe]Floatdexa elements abundance from LAMOST21e.[ $\alpha$ /Fe]Floatdexerror of $\alpha$ elements abundance22vrDoublekm s <sup>-1</sup> Radial velocity from LAMOST23e.vrDoublekm s <sup>-1</sup> Error of radial velocity24 $N_p$ integerPlanet (candidate) multiplicityParameters derived in this work25RDoublekm s <sup>-1</sup> 26 $\theta$ Doublekm s <sup>-1</sup> 27ZDoublekm s <sup>-1</sup> 28 $V_R$ Doublekm s <sup>-1</sup> 29 $V_{\theta}$ Doublekm s <sup>-1</sup> 20 $V_Z$ Doublekm s <sup>-1</sup> 21 $U_{LSR}$ Doublekm s <sup>-1</sup> 23Galactocentric Cylindrical ardial distance24 $N_R$ Doublekm s <sup>-1</sup> 25RDoublekm s <sup>-1</sup> 26 $\theta$ Doublekm s <sup>-1</sup> 27ZDoublekm s <sup>-1</sup> 28 $V_R$ Doublekm s <sup>-1</sup> 29 $V_{\theta}$ Doublekm s <sup>-1</sup> 30 $V_Z$ Doublekm s <sup>-1</sup> 31 $U_{LSR}$ Doublekm s <sup>-1</sup> 32e.U_LSRDoublekm s <sup>-1</sup> 33 $V_{LSR}$ Doublekm s <sup>-1</sup> 3	16	$\log a$	Float		Surface gravity from LAMOST
17e.log gFloatError of surface gravity from LAMOST18[Fe/H]FloatdexMetallicity from LAMOST19e.[Fe/H]FloatdexError of metallicity20[ $\alpha$ /Fe]Floatdex $\alpha$ elements abundance from LAMOST21e.[ $\alpha$ /Fe]FloatdexError of $\alpha$ elements abundance22vrDoublekm s <sup>-1</sup> Radial velocity from LAMOST23e.vrDoublekm s <sup>-1</sup> Error of radial velocity from LAMOST24NpintegerPlanet (candidate) multiplicityParameters derived in this work25RDoublekpc26 $\theta$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical aximuth angle27ZDoublekm s <sup>-1</sup> 28VRDoublekm s <sup>-1</sup> 29VgDoublekm s <sup>-1</sup> 20U_LSRDoublekm s <sup>-1</sup> 21e.ULSRDoublekm s <sup>-1</sup> 23e.VLSRDoublekm s <sup>-1</sup> 34e.VLSRDoublekm s <sup>-1</sup> 35WLSRDoublekm s <sup>-1</sup> 36e.WLSRDoublekm s <sup>-1</sup> 37TD/DDoublekm s <sup>-1</sup> 38TD/HDoublekm s <sup>-1</sup> 39Herc/DDoublekm s <sup>-1</sup> 40Herc/TDDoublekm s <sup>-1</sup> 41ComponentstringClassification of Galactic components <td>10</td> <td></td> <td>1 1000</td> <td></td> <td></td>	10		1 1000		
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	17	$e_{-}\log g$	Float		Error of surface gravity from LAMOST
18[Fe/H]FloatdexMetallicity from LAMOST19e.[Fe/H]Floatdexerror of metallicity20[a/Fe]Floatdexa elements abundance from LAMOST21e.[a/Fe]Floatdexerror of $\alpha$ elements abundance22vrDoublekm s <sup>-1</sup> Radial velocity from LAMOST23e.vrDoublekm s <sup>-1</sup> Error of radial velocity24 $N_p$ integerPlanet (candidate) multiplicity25RDoubledegGalactocentric Cylindrical radial distance26 $\theta$ DoublekpcGalactocentric Cylindrical a relocities29 $V_d$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical $\pi$ velocities29 $V_d$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical $\pi$ velocities31 $U_{LSR}$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical $\pi$ velocities32 $e_J U_{LSR}$ Doublekm s <sup>-1</sup> Galactocentric $x$ velocity to the LSR33 $V_{LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric $x$ velocity to the LSR34 $e_J V_{LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric $x$ velocity to the LSR35 $W_{LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric $z$ velocity to the LSR36 $e_M V_{LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric $z$ velocity to the LSR36 $e_L V_{LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric $z$ velocity to the LSR36 $e_L V$					
19 $e_{-}[Fe/H]$ FloatdexError of metallicity20 $[\alpha/Fe]$ Floatdex $\alpha$ elements abundance from LAMOST21 $e_{-}[\alpha/Fe]$ FloatdexError of $\alpha$ elements abundance22vrDoublekm s <sup>-1</sup> Radial velocity from LAMOST23 $e_{-}vr$ Doublekm s <sup>-1</sup> Error of radial velocity24 $N_p$ integerPlanet (candidate) multiplicityParameters derived in this work25RDoublekpcGalactocentric Cylindrical azimuth angle27ZDoublekpcGalactocentric Cylindrical azimuth angle28 $V_R$ Doublekm s <sup>-1</sup> 29 $V_{\theta}$ Doublekm s <sup>-1</sup> 30 $V_Z$ Doublekm s <sup>-1</sup> 31 $U_{LSR}$ Doublekm s <sup>-1</sup> 32 $e_{-}U_{LSR}$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical Z velocities33 $V_{LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric x velocity to the LSR34 $e_{-}V_{LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric z velocity to the LSR35 $W_{LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric z velocity to the LSR35 $W_{LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric z velocity to the LSR36 $e_{-}V_{LSR}$ Doublekm s <sup></sup>	18	[Fe/H]	Float	dex	Metallicity from LAMOST
20 $[\alpha/Fe]$ Floatdex $\alpha$ elements abundance from $\alpha$ elements abundance21 $e_{-}[\alpha/Fe]$ FloatdexError of $\alpha$ elements abundance22vrDoublekm s <sup>-1</sup> Radial velocity from LAMOST23 $e_{-}vr$ Doublekm s <sup>-1</sup> Error of radial velocity24 $N_{p}$ integerPlanet (candidate) multiplicity25 $R$ DoublekpcGalactocentric Cylindrical radial distance26 $\theta$ DoublekpcGalactocentric Cylindrical azimuth angle27 $Z$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical $R$ velocities29 $V_{\theta}$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical $R$ velocities30 $V_Z$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical $R$ velocities31 $U_{\text{LSR}}$ Doublekm s <sup>-1</sup> Galactocentric $x$ velocity to the LSR32 $e_{-}U_{\text{LSR}}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric $x$ velocity to the LSR33 $V_{\text{LSR}}$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric $x$ velocity to the LSR35 $W_{\text{LSR}}$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric $x$ velocity to the LSR36 $e_{-}W_{\text{LSR}}$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric $x$ velocity to the LSR36 $e_{-}W_{\text{LSR}}$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric $x$ velocity to the LSR37 $TD/D$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric $x$ velocity to the LS	19	$e_{-}[Fe/H]$	Float	dex	Error of metallicity
21 $e_{-}[\alpha/Fe]$ FloatdexError of $\alpha$ elements abundance22vrDoublekm s <sup>-1</sup> Radial velocity from LAMOST23e_vrDoublekm s <sup>-1</sup> Error of radial velocity24 $N_p$ integerPlanet (candidate) multiplicityParameters derived in this work25RDoublekpc26 $\theta$ Doublekpc27ZDoublekm s <sup>-1</sup> 28 $V_R$ Doublekm s <sup>-1</sup> 29 $V_{\theta}$ Doublekm s <sup>-1</sup> 29 $V_{\theta}$ Doublekm s <sup>-1</sup> 31 $U_{LSR}$ Doublekm s <sup>-1</sup> 32 $e_{ULSR}$ Doublekm s <sup>-1</sup> 33 $V_{LSR}$ Doublekm s <sup>-1</sup> 34 $e_{VLSR}$ Doublekm s <sup>-1</sup> 35 $W_{LSR}$ Doublekm s <sup>-1</sup> 36 $e_{WLSR}$ Doublekm s <sup>-1</sup> 37 $TD/D$ Doublekm s <sup>-1</sup> 38 $TD/H$ Doublekm s <sup>-1</sup> 39 $Herc/D$ Doublekm s <sup>-1</sup> 41ComponentstringClassification of Galactic components	20	$[\alpha/{\rm Fe}]$	Float	dex	$\alpha$ elements abundance from LAMOST
22vrDoublekm s^{-1}Radial velocity from LAMOST23e.vrDoublekm s^{-1}Error of radial velocity24 $N_p$ integerPlanet (candidate) multiplicity24 $N_p$ integerPlanet (candidate) multiplicity25 $R$ DoublekpcGalactocentric Cylindrical radial distance26 $\theta$ DoubledegGalactocentric Cylindrical azimuth angle27 $Z$ DoublekpcGalactocentric Cylindrical vertical height28 $V_R$ Doublekm s^{-1}Galactocentric Cylindrical $\theta$ velocities29 $V_{\theta}$ Doublekm s^{-1}Galactocentric Cylindrical $\theta$ velocities30 $V_Z$ Doublekm s^{-1}Galactocentric Cylindrical $Z$ velocities31 $U_{LSR}$ Doublekm s^{-1}Cartesian Galactocentric $x$ velocity to the LSR32 $e.U_{LSR}$ Doublekm s^{-1}cartesian Galactocentric $x$ velocity to the LSR33 $V_{LSR}$ Doublekm s^{-1}cartesian Galactocentric $y$ velocity to the LSR34 $e.V_{LSR}$ Doublekm s^{-1}cartesian Galactocentric $z$ velocity to the LSR35 $W_{LSR}$ Doublekm s^{-1}cartesian Galactocentric $z$ velocity to the LSR36 $e.W_{LSR}$ Doublekm s^{-1}error of Cartesian Galactocentric $z$ velocity to the LSR37 $TD/D$ Doublekm s^{-1}error of Cartesian Galactocentric $z$ velocity to the LSR38 $TD/H$ Doublekm s^{-1} </td <td>21</td> <td><math>e_{-}[\alpha/Fe]</math></td> <td>Float</td> <td>dex</td> <td>Error of <math>\alpha</math> elements abundance</td>	21	$e_{-}[\alpha/Fe]$	Float	dex	Error of $\alpha$ elements abundance
23e.vrDoublekm s^{-1}Error of radial velocity24 $N_{\rm p}$ integerPlanet (candidate) multiplicity24 $N_{\rm p}$ integerPlanet (candidate) multiplicity25 $R$ DoublekpcGalactocentric Cylindrical radial distance26 $\theta$ DoublekpcGalactocentric Cylindrical vertical height28 $V_R$ Doublekm s^{-1}Galactocentric Cylindrical $R$ velocities29 $V_{\theta}$ Doublekm s^{-1}Galactocentric Cylindrical $R$ velocities30 $V_Z$ Doublekm s^{-1}Galactocentric Cylindrical $Z$ velocities31 $U_{\rm LSR}$ Doublekm s^{-1}Galactocentric $X$ velocity to the LSR32 $e.U_{\rm LSR}$ Doublekm s^{-1}Cartesian Galactocentric $y$ velocity to the LSR33 $V_{\rm LSR}$ Doublekm s^{-1}Cartesian Galactocentric $y$ velocity to the LSR34 $e.V_{\rm LSR}$ Doublekm s^{-1}error of Cartesian Galactocentric $y$ velocity to the LSR35 $W_{\rm LSR}$ Doublekm s^{-1}error of Cartesian Galactocentric $z$ velocity to the LSR36 $e.W_{\rm LSR}$ Doublekm s^{-1}error of Cartesian Galactocentric $z$ velocity to the LSR37 $TD/D$ Doublekm s^{-1}error of Cartesian Galactocentric $z$ velocity to the LSR38 $TD/H$ Doublekm s^{-1}error of Cartesian Galactocentric $z$ velocity to the LSR39 $Herc/D$ Doublethick disc to halo membership probability	22	vr	Double	$\rm km~s^{-1}$	Radial velocity from LAMOST
24 $N_p$ integer       Planet (candidate) multiplicity         Parameters derived in this work         25       R       Double       kpc       Galactocentric Cylindrical radial distance         26 $\theta$ Double       deg       Galactocentric Cylindrical azimuth angle         27       Z       Double       kpc       Galactocentric Cylindrical retrical height         28 $V_R$ Double       km s <sup>-1</sup> Galactocentric Cylindrical $R$ velocities         29 $V_{\theta}$ Double       km s <sup>-1</sup> Galactocentric Cylindrical $R$ velocities         30 $V_Z$ Double       km s <sup>-1</sup> Galactocentric $Q$ velocities         31 $U_{\rm LSR}$ Double       km s <sup>-1</sup> Cartesian Galactocentric $x$ velocity to the LSR         32 $e_J U_{\rm LSR}$ Double       km s <sup>-1</sup> Cartesian Galactocentric $y$ velocity to the LSR         33 $V_{\rm LSR}$ Double       km s <sup>-1</sup> cartesian Galactocentric $y$ velocity to the LSR         34 $e_V_{\rm LSR}$ Double       km s <sup>-1</sup> cartesian Galactocentric $z$ velocity to the LSR         35 $W_{\rm LSR}$ Double       km s <sup>-1</sup> cartesian Galactocentric $z$ velocity to the LSR	23	e_vr	Double	$\rm km \ s^{-1}$	Error of radial velocity
Parameters derived in this work25RDoublekpcGalactocentric Cylindrical radial distance26 $\theta$ DoubledegGalactocentric Cylindrical azimuth angle27ZDoublekpcGalactocentric Cylindrical azimuth angle28 $V_R$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical $R$ velocities29 $V_{\theta}$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical $\ell$ velocities30 $V_Z$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical $Z$ velocity31 $U_{\text{LSR}}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric $x$ velocity to the LSR32 $e_*U_{\text{LSR}}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric $x$ velocity to the LSR33 $V_{\text{LSR}}$ Doublekm s <sup>-1</sup> cartesian Galactocentric $y$ velocity to the LSR34 $e_*V_{\text{LSR}}$ Doublekm s <sup>-1</sup> cartesian Galactocentric $z$ velocity to the LSR35 $W_{\text{LSR}}$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric $z$ velocity to the LSR36 $e_*W_{\text{LSR}}$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric $z$ velocity to the LSR37 $TD/D$ Doublethick disc to thin disc membership probability38 $TD/H$ DoubleHercules stream to thin disc membership probability40 $Herc/TD$ DoubleHercules stream to thick disk membership probability41ComponentstringClassification of Galactic components	24	$N_{ m p}$	integer		Planet (candidate) multiplicity
25RDoublekpcGalactocentric Cylindrical radial distance26 $\theta$ DoubledegGalactocentric Cylindrical azimuth angle27ZDoublekpcGalactocentric Cylindrical vertical height28 $V_R$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical vertical height29 $V_{\theta}$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical $\theta$ velocities30 $V_Z$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical Z velocities31 $U_{LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric x velocity to the LSR32 $e_J U_{LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric x velocity to the LSR33 $V_{LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric y velocity to the LSR34 $e_V_{LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric y velocity to the LSR35 $W_{LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric z velocity to the LSR36 $e_{WLSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric z velocity to the LSR37 $TD/D$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric z velocity to the LSR38 $TD/H$ Doublekm s <sup>-1</sup> thick disc to thin disc membership probability40 $Herc/TD$ DoubleHercules stream to thind disk membership probability41ComponentstringClassification of Galactic components				Paramoto	re derived in this work
25RDoublekpcGalactocentric Cylindrical ration distance26 $\theta$ DoubledegGalactocentric Cylindrical azimuth angle27ZDoublekm s <sup>-1</sup> Galactocentric Cylindrical R velocities28 $V_R$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical R velocities29 $V_{\theta}$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical R velocities30 $V_Z$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical Z velocities31 $U_{\rm LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric x velocity to the LSR32 $e_J U_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric y velocity to the LSR33 $V_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric y velocity to the LSR34 $e_J V_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric y velocity to the LSR35 $W_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric z velocity to the LSR36 $e_W_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric z velocity to the LSR37 $TD/D$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric z velocity to the LSR38 $TD/H$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric z velocity to the LSR39 $Herc/D$ DoubleHercules stream to thin disc membership probability40 $Herc/TD$ DoubleHercules stream to thick disk membership probability41ComponentstringClassification of Galactic components		D	Doublo	r ar annete	Cologica childrigal radial distance
200DoubledegGalactocentric Cylindrical variation angle27ZDoublekpcGalactocentric Cylindrical vertical height28 $V_R$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical R velocities29 $V_{\theta}$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical R velocities30 $V_Z$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical Z velocities31 $U_{\rm LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric x velocity to the LSR32 $e_U_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric x velocity to the LSR33 $V_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric y velocity to the LSR34 $e_V_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric z velocity to the LSR35 $W_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric z velocity to the LSR36 $e_W_{\rm LSR}$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric z velocity to the LSR37 $TD/D$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric z velocity to the LSR38 $TD/H$ Doublekm s <sup>-1</sup> thick disc to halo membership probability40 $Herc/TD$ DoubleHercules stream to thin disc membership probability41ComponentstringClassification of Galactic components	20 26	R A	Double	крс dog	Calactocentric Cylindrical azimuth angle
212DoublekpcGalactocentric Cylindrical R velocities28 $V_R$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical R velocities29 $V_{\theta}$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical $\theta$ velocities30 $V_Z$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical Z velocities31 $U_{\rm LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric x velocity to the LSR32 $e_U_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric y velocity to the LSR33 $V_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric y velocity to the LSR34 $e_V_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric z velocity to the LSR35 $W_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric z velocity to the LSR36 $e_W_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric z velocity to the LSR37 $TD/D$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric z velocity to the LSR38 $TD/H$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric z velocity to the LSR39 $Herc/D$ DoubleHercules stream to thin disc membership probability40 $Herc/TD$ DoubleHercules stream to thick disk membership probability41ComponentstringClassification of Galactic components	20	7	Double	lene	Calactocentric Cylindrical azimuth angle
25 $V_R$ Doublekm sGalactocentric Cylindrical $R$ velocities29 $V_{\theta}$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical $\theta$ velocities30 $V_Z$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical $Z$ velocities31 $U_{\rm LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric $x$ velocity to the LSR32 $e_{\cdot}U_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric $x$ velocity to the LSR33 $V_{\rm LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric $y$ velocity to the LSR34 $e_{\cdot}V_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric $y$ velocity to the LSR35 $W_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric $z$ velocity to the LSR36 $e_{\cdot}W_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric $z$ velocity to the LSR37 $TD/D$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric $z$ velocity to the LSR38 $TD/H$ Doublekm s <sup>-1</sup> thick disc to thin disc membership probability39 $Herc/D$ DoubleHercules stream to thin disc membership probability40 $Herc/TD$ DoubleHercules stream to thick disk membership probability41ComponentstringClassification of Galactic components	21		Double	kpc	Galactocentric Cylindrical Vertical height
29 $V_{\theta}$ DoubleKm sGalactocentric Cylindrical $J$ velocities30 $V_Z$ Doublekm s <sup>-1</sup> Galactocentric Cylindrical $Z$ velocities31 $U_{\rm LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric $x$ velocity to the LSR32 $e_{-}U_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric $x$ velocity to the LSR33 $V_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric $y$ velocity to the LSR34 $e_{-}V_{\rm LSR}$ Doublekm s <sup>-1</sup> cartesian Galactocentric $z$ velocity to the LSR35 $W_{\rm LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric $z$ velocity to the LSR36 $e_{-}W_{\rm LSR}$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric $z$ velocity to the LSR37 $TD/D$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric $z$ velocity to the LSR38 $TD/H$ Doublekm s <sup>-1</sup> thick disc to thin disc membership probability39 $Herc/D$ DoubleHercules stream to thin disc membership probability41ComponentstringClassification of Galactic components	20	$V_R$	Double	$km s^{-1}$	Galactocentric Cylindrical A velocities
30 $V_Z$ Doublekm sGalactocentric Cymunical Z velocities31 $U_{\rm LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric x velocity to the LSR32 $e_U_{\rm LSR}$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric x velocity to the LSR33 $V_{\rm LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric y velocity to the LSR34 $e_V_{\rm LSR}$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric y velocity to the LSR35 $W_{\rm LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric z velocity to the LSR36 $e_W_{\rm LSR}$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric z velocity to the LSR37 $TD/D$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric z velocity to the LSR38 $TD/H$ Doublekm s <sup>-1</sup> thick disc to thin disc membership probability39 $Herc/D$ DoubleHercules stream to thin disc membership probability41ComponentstringClassification of Galactic components	29	$V_{\theta}$	Double	$km s^{-1}$	Calactocentric Cylindrical 7 velocities
31 $D_{\rm LSR}$ Doublekm sCartesian Galactocentric x velocity to the LSR32 $e_{U_{\rm LSR}}$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric x velocity to the LSR33 $V_{\rm LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric y velocity to the LSR34 $e_{V_{\rm LSR}}$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric y velocity to the LSR35 $W_{\rm LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric z velocity to the LSR36 $e_{W_{\rm LSR}}$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric z velocity to the LSR37 $TD/D$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric z velocity to the LSR38 $TD/H$ Doublekm s <sup>-1</sup> thick disc to thin disc membership probability39 $Herc/D$ DoubleHercules stream to thin disc membership probability41ComponentstringClassification of Galactic components	91		Double	$km c^{-1}$	Cartacian Calactecentric a velocities
32E-OLSRDoubleKm send of Cartesian Galactocentric 1 velocity to the LSR33 $V_{\rm LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric 1 velocity to the LSR34 $e_{-V_{\rm LSR}}$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric 1 velocity to the LSR35 $W_{\rm LSR}$ Doublekm s <sup>-1</sup> Cartesian Galactocentric 2 velocity to the LSR36 $e_{-W_{\rm LSR}}$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric 2 velocity to the LSR37 $TD/D$ Doublekm s <sup>-1</sup> error of Cartesian Galactocentric 2 velocity to the LSR38 $TD/H$ Doublethick disc to thin disc membership probability39 $Herc/D$ DoubleHercules stream to thin disc membership probability40 $Herc/TD$ DoubleClassification of Galactic components41ComponentstringClassification of Galactic components	30	o Usa	Double	$km s^{-1}$	Cartesian Galactocentric $x$ velocity to the LSR
33 $V_{\rm LSR}$ Doublekm sCartesian Galactocentric y velocity to the LSR34 $e_{-V_{\rm LSR}}$ Doublekm s^{-1}error of Cartesian Galactocentric y velocity to the LSR35 $W_{\rm LSR}$ Doublekm s^{-1}Cartesian Galactocentric z velocity to the LSR36 $e_{-W_{\rm LSR}}$ Doublekm s^{-1}error of Cartesian Galactocentric z velocity to the LSR37 $TD/D$ Doublekm s^{-1}error of Cartesian Galactocentric z velocity to the LSR38 $TD/H$ Doublethick disc to thin disc membership probability39 $Herc/D$ DoubleHercules stream to thin disc membership probability40 $Herc/TD$ DoubleClassification of Galactic components	0∠ 99	$e_{-}O_{\rm LSR}$	Double	$km s^{-1}$	Contoxion Coloctocontria a valority to the LSR
34 $e_{-V_{\rm LSR}}$ Doublekm s - 1error of Cartesian Galactocentric $y$ velocity to the LSR35 $W_{\rm LSR}$ Doublekm s - 1Cartesian Galactocentric $z$ velocity to the LSR36 $e_{-W_{\rm LSR}}$ Doublekm s - 1Cartesian Galactocentric $z$ velocity to the LSR37 $TD/D$ Doublekm s - 1error of Cartesian Galactocentric $z$ velocity to the LSR38 $TD/H$ Doublethick disc to thin disc membership probability39 $Herc/D$ DoubleHercules stream to thin disc membership probability40 $Herc/TD$ DoubleClassification of Galactic components41ComponentstringClassification of Galactic components	აა ეჟ	V <sub>LSR</sub>	Double	km s -	Cartesian Galactocentric $y$ velocity to the LSR
35 $W_{\rm LSR}$ Doublekm s - 1Cartesian Galactocentric z velocity to the LSR36 $e_W_{\rm LSR}$ Doublekm s^{-1}error of Cartesian Galactocentric z velocity to the LSR37 $TD/D$ Doublekm s^{-1}error of Cartesian Galactocentric z velocity to the LSR38 $TD/H$ Doublethick disc to thin disc membership probability39 $Herc/D$ DoubleHercules stream to thin disc membership probability40 $Herc/TD$ DoubleHercules stream to thick disk membership probability41ComponentstringClassification of Galactic components	34 97	$e_{-}V_{LSR}$	Double	km s -	error of Cartesian Galactocentric $y$ velocity to the LSR
$36$ $e_{-W_{\rm LSR}}$ Doublekm s +error of Cartesian Galactocentric z velocity to the LSR $37$ $TD/D$ Doublethick disc to thin disc membership probability $38$ $TD/H$ Doublethick disc to halo membership probability $39$ $Herc/D$ DoubleHercules stream to thin disc membership probability $40$ $Herc/TD$ DoubleHercules stream to thick disk membership probability $41$ ComponentstringClassification of Galactic components	30	WLSR	Double	$km s^{-1}$	Cartesian Galactocentric $z$ velocity to the LSR
37I D/DDoublethick disc to thin disc membership probability38TD/HDoublethick disc to halo membership probability39Herc/DDoubleHercules stream to thin disc membership probability40Herc/TDDoubleHercules stream to thick disk membership probability41ComponentstringClassification of Galactic components	30	$e_W_{LSR}$	Double	km s	error of Cartesian Galactocentric $z$ velocity to the LSR
38TD/HDoublethick disc to halo membership probability39Herc/DDoubleHercules stream to thin disc membership probability40Herc/TDDoubleHercules stream to thick disk membership probability41ComponentstringClassification of Galactic components	37	TD/D	Double		thick disc to thin disc membership probability
39Herc/DDoubleHercules stream to thin disc membership probability40Herc/TDDoubleHercules stream to thick disk membership probability41ComponentstringClassification of Galactic components	38	TD/H	Double		thick disc to halo membership probability
40Herc/TDDoubleHercules stream to third ise membership probability41ComponentstringClassification of Galactic components	30	Herc/D	Double		Hercules stream to thin disc membership probability
40Herc/TDDoubleHercules stream to thick disk membership probability41ComponentstringClassification of Galactic components	00	11010/12	Double		nervales stream to thin the membership probability
41 Component string Classification of Galactic components	40	Herc/TD	Double		Hercules stream to thick disk membership probability
	41	Component	string		Classification of Galactic components

noise on 6.5 hours timescale which is dominated by stellar activity decreases as stars aging. Our result support their

Table 4. Numbers of stars  $(N_s)$  and planets  $(N_p)$  and kinematic and chemical properties of different Galactic components for the LAMOST-Gaia-Kepler sample.



Figure 4. Comparison among ages derived by various methods. The x-axes for each columns from left to the right are asteroseismic age from Pinsonneault et al. (2018), gyrochronology age derived from method provided by Spada & Lanzafame (2020), and isochrone age from Berger et al. (2020b), y-axes for each row are isochrone age and kinematic age derived in this work, respectively. In every subplot, we compare the ages from the corresponding x and y samples, the dots and errorbars are the median age and median value of relative error, except kinematic ages and uncertainties are calculated by method detailed in section 3.3. We label the size of sample in the upper left corner, and the gray dash line shows where x equals to y.

model from the aspect of observation. For shorter duration, the declining trend between CDPP—photon noise and kinematic age is weaker. This is not unexpected because shorter timescale noises could be contributed by granulation which generally increases with age, and thus compensating the decline trend. For main sequence stars, the typical timescale of granulation is on order of a few hundreds seconds (Gilliland et al. 2011; Kallinger et al. 2014), the effect of granulation will be more evident on CDPP with shorter range, such as 1.5 hours.

## 5.3. Kinematic properties of Kepler planet candidate host Stars

In this section, we explore the differences between the kinematic properties of Kepler planet candidate host stars and those of other stars without Kepler planets, which are meaningful to study the formation and evolution environment of planetary systems. To avoid the influence of the stellar evolutionary stage and spectral type, here we only compare



Figure 5. Stellar rotation period and magnetic activity (S-index) as functions of kinematic age. In the upper panel, the dots present the kinematic age and stellar rotation period. The x-axis errorbars are derived from section 3.3, and errorbars for the y-axis are median of rotation period and median value of relative error. The comparison between kinematic age and S-index is shown in the lower panel, with dots and errorbars derived with the same method. In both of the panels, the number of the sample size are shown in the upper/lower left corner, the yellow dots present the age, rotation period, and S-index of the Sun.

the distribution of Solar-type stars, which are the bulk of LAMOST-Gaia-Kepler sample. The selection criteria are taken as an effective temperature  $T_{\text{eff}}$  in the range of 4700–6500 K and a stellar surface gravity log g > 4.0. Finally, we are left with 563 Kepler planet candidate hosts and 20,795 stars without Kepler planets.

5.3.1. Spatial distribution



Figure 6. Comparison between kinematic age and CDPP-photon noise. Each line presents CDPP for a certain time scale, and shown in the legend.

Table 5. Stellar properties of Kepler planet candidate host stars and stars without Kepler planets.

	All Stars without	neighbor stars without	Kepler planet candidate host stars			
	Kepler planets $(N_{\rm p}=0)$	Kepler planets $(N_{\rm p}=0)$	All	$N_{\rm p} = 1$	$N_{\rm p}=2$	$N_{\rm p} \ge 3$
$M_* (M_{\odot})$	$1.06^{+0.21}_{-0.18}$	$1.04_{-0.19}^{+0.20}$	$1.03^{+0.21}_{-0.16}$	$1.03\substack{+0.20\\-0.16}$	$1.03_{-0.17}^{+0.18}$	$1.05_{-0.17}^{+0.22}$
$R_*~(R_{\odot})$	$1.20\substack{+0.41\\-0.31}$	$1.15_{-0.30}^{+0.38}$	$1.10\substack{+0.40\\-0.23}$	$1.09\substack{+0.39\\-0.23}$	$1.12_{-0.27}^{+0.38}$	$1.13_{-0.26}^{+0.38}$
[Fe/H] (dex)	$-0.02\substack{+0.19\\-0.26}$	$-0.01\substack{+0.19\\-0.27}$	$0.00\substack{+0.18\\-0.21}$	$-0.01\substack{+0.19\\-0.21}$	$0.01\substack{+0.13\\-0.23}$	$-0.02\substack{+0.17\\-0.18}$
$[\alpha/{\rm Fe}]~({\rm dex})$	$0.05\substack{+0.10\\-0.08}$	$0.05\substack{+0.09 \\ -0.07}$	$0.04\substack{+0.09\\-0.08}$	$0.04\substack{+0.09\\-0.08}$	$0.03\substack{+0.09 \\ -0.07}$	$0.05\substack{+0.07 \\ -0.07}$
Fr	$86.9^{+0.23}_{-0.23}\%$	$86.8^{+1.4}_{-1.4}\%$	$88.8^{+1.3}_{-1.3}\%$	$88.6^{+1.6}_{-1.6}\%$	$90.0^{+3.2}_{-3.1}\%$	$91.7^{+3.6}_{-3.5}\%$
гD	(18079/20795)	(480/553)	(502/563)	(357/403)	(90/100)	(55/60)
$F_{\mathrm{TD}}$	$5.2^{+0.2}_{-0.2}\%$	$5.8^{+1.0}_{-1.0}\%$	$4.4^{+0.9}_{-0.9}\%$	$5.5^{+1.2}_{-1.2}\%$	$3.0^{+1.7}_{-1.9}\%$	$0^{+3.1}_{-0}\%$
	(1088/20795)	(32/553)	(25/563)	(22/403)	(3/100)	(0/60)
$F_{ m H}$	$0.17^{+0.02}_{-0.02}\%$	$0.36^{+0.48}_{-0.23}\%$	$0^{+0.3}_{-0}\%$	$0^{+0.5}_{-0}\%$	$0^{+1.8}_{-0}\%$	$0^{+3.1}_{-0}\%$
	(35/20795)	(2/553)	(0/563)	(0/403)	(0/100)	(0/60)
$F_{ m Herc}$	$1.6^{+0.08}_{-0.08}\%$	$0.90^{+0.61}_{-0.39}\%$	$1.8^{+0.6}_{-0.5}\%$	$1.5^{+0.6}_{-0.5}\%$	$3.0^{+1.7}_{-1.9}\%$	$1.7^{+1.72}_{-1.4}\%$
	(329/20795)	(5/553)	(10/563)	(6/403)	(3/100)	(1/60)
$V_{\rm tot}~({\rm km~s^{-1}})$	$44.06^{+31.24}_{-22.32}$	$43.05_{-21.13}^{+32.04}$	$36.37^{+33.70}_{-16.36}$	$37.89^{+33.69}_{-17.88}$	$32.30^{+33.64}_{-13.21}$	$29.36^{+30.87}_{-10.88}$
$\sigma_{\rm tot}~({\rm km~s^{-1}})$	$49.07\substack{+0.84 \\ -0.73}$	$49.99^{+1.25}_{-0.91}$	$45.65^{+1.14}_{-0.77}$	$46.92^{+1.20}_{-1.21}$	$42.48^{+1.64}_{-1.32}$	$41.70^{+2.11}_{-1.87}$
Age_kin (Gyr)	$4.28^{+0.55}_{-0.35}$	$4.49_{-0.36}^{+0.60}$	$3.57^{+0.43}_{-0.27}$	$3.82^{+0.50}_{-0.28}$	$2.98^{+0.36}_{-0.20}$	$2.84_{-0.29}^{+0.34}$

F represents the number fraction of different components: thin disk (D), thick disk (TD), halo (H) and Hercules stream (Herc).



Figure 7. The cumulative distributions of distance to the Sun, Galactic coordinates  $(R, \theta, Z)$ , CDPP 4.5h and  $S_{\text{index}}$  for Kepler planet candidate host stars (dashed red lines, labelled as  $N_p > 0$ ) and all stars without Kepler planets (solid black lines, labelled as  $N_p = 0$  (All)). The distributions of the subsamples divided by the numbers of transit planets are plotted in different colours: blue for  $N_p = 1$ , yellow for  $N_p = 2$  and green for  $N_p \ge 3$ . The wide grey lines represent the distributions of the selected nearest neighbors for every Kepler planet candidate host star from stars without Kepler planets (labelled as  $N_{p=0}$  (Nei)). In each panel, the *p* denotes the *p*-value of the two sample KS test for the distributions of the other samples (the subscripts same as the labels) comparing to the neighbor stars without Kepler planets ( $N_p = 0$  (Nei)).

Here we compare the distribution of spatial position in Figure 7. As can be seen, the stars without Kepler planets (solid black lines) have a wider distribution in the distance to the Sun, Galactic radius (R), azimuth angle  $(\theta)$  and height (Z) than that of Kepler planet candidate host stars (dashed red lines). We did Kolmogorov–Smirnov (KS) tests between the distance to the Sun, Galactic coordinates  $(R, \theta, Z)$  of the Kepler planet candidate host stars and those of stars without Kepler planets. The resulted p values are all smaller than 0.003, demonstrating that there are significant differences between their spatial positions. As shown in Figure 5 and 6, the stellar activity (and therefore noise properties) are correlated to their kinematics. This may affect the detectability of planets in multi-planet systems,



Figure 8. The cumulative distribution of Galactic velocity,  $V_{\text{tot}}$  (left panel) and the relative probabilities of the thick-disk-tothin-disk, TD/D (right panel) for all planet candidate host (dashed red lines) 1 transiting planet candidate hosts (blue lines), 2 transiting planet candidate hosts (yellow lines), 3+ transiting planet candidate hosts (green lines) and neighbor stars without Kepler planets (wide grey lines, labelled as  $N_p = 0$  (Nei)). In each panel, the *p* denotes the *p*-value of the two sample KS test for the distributions of the other samples (the subscripts same as the labels) comparing to the neighbor stars without Kepler planets ( $N_p = 0$  (Nei)).

e.g., by creating a bias against smaller planets around active/noisy stars (typically corresponding to younger stars) Thus in the following comparison in stellar kinematic properties, to avoid the bias caused by spatial distribution and stellar activity/noise, we construct a control sample by adopting the NearestNeighbors function in scikit-learn (Pedregosa et al. 2011) to select the nearest neighbors for every host star from stars without Kepler planets in spatial distribution (i.e. R,  $\theta$ , and Z), CDPP and  $S_{index}$ . Here we select the CDPP of 4.5 hour as the transit duration are 4.3 hours by taking the median values of stellar mass (1.03  $M_{\odot}$ ), radius (1.10  $R_{\odot}$ ) and planetary period (10.37 days) in our sample. In the case that a star without planets is selected as the nearest neighbors of multiple times, we exclude duplicate stars. The neighbor stellar sample without Kepler planets contains 551 stars. As can be seen in Figure 7, the distributions of the neighbor stars without Kepler planets (wide grey lines) are very similar to that of Kepler planet candidate host stars (dashed red lines). We also did KS tests between their spatial distributions and stellar activity/noise. The p-values are 0.9999, 0.9456, 0.9877, 0.9971, 0.9156 and 0.8875 for distance, R,  $\theta$ , Z, CDPP 4.5h and  $S_{index}$  respectively. Such high p-values demonstrate the spatial distributions and stellar activity/noise of the planet candidate host stars and neighbor stars without Kepler planets are statistically indistinguishable. Therefore, the selected neighbor stars form a reliable control sample to minimize the spatial and detection bias.

In the following discussions, we divided the Kepler planet candidate host sample into three subsmaples according to the number of transiting planets (tranets) in each system:  $N_{\rm p} = 1$  (403 stars),  $N_{\rm p} = 2$  (100 stars),  $N_{\rm p} \ge 3$  (60 stars). The p-values of KS test between the distribution of the distance and Galactic coordinates for the neighbor stellar sample without Kepler planets and the three subsamples (blue, yellow, red lines in Figure 7) are all larger than 0.25, which prove that their spatial distribution are statistically indistinguishable. In Table 5, we summarized the typical value of physical properties. The uncertainties are taken as the  $50\pm34.1$  percentiles of the corresponding distributions. As can be seen, their typical values in mass, radius, [Fe/H] and [ $\alpha$ /Fe] are indistinguishable within the  $1-\sigma$  uncertainties. We also make KS test and resulted p-values are all larger than 0.1, demonstrating that there is no significant difference in the distributions of mass, radius, [Fe/H] and [ $\alpha$ /Fe].

So far, we have found that Kepler planet candidate hosts have a narrower range of Galactic positions  $(R, \theta, Z)$  and shorter distance as compared to the general Kepler field stars. This is likely to be caused by the observational bias, namely, planets are more easily found around stars closer to the Earth. To remove this spatial and detection bias, we constructed a control sample by selecting the neighbor stars of planet candidate hosts. As we will show below, although planet candidate hosts have similarities with the control sample in Galactic position, mass, radius, chemical abundance, they (especially those with large planet multiplicity, e.g.,  $N_p \ge 3$ ) do differ significantly in kinematic properties.

# 5.3.2. Galactic components and kinematic age

Firstly, we compare the distributions of total velocity and the relative probabilities of the thick-disk-to-thin-disk, TD/D for Kepler planet candidate host subsamples and neighbor non-Kepler planet candidate host sample. As can be seen in Figure 8, the neighbor stars without Kepler planets (wide grey lines) have velocities and TD/D that are on average larger but are statistically indistinguishable (KS test p-values > 0.05) compared to Kepler planet candidate host stars (dashed red lines). This is consistent with the result of McTier & Kipping (2019), which found that Kepler planet candidate hosts (dominated by singles i.e.,  $N_p = 1$  since they are the majority) have similar Galactic velocities and TD/D generally decrease with the increase of the transiting planet multiplicity  $N_p$ . To quantify this trend, we did KS tests between the neighbor sample without Kepler planets and stars hosting 1, 2, 3+ tranets systems. The resulting p-values (as printed in Figure 8) decreases with increasing  $N_p$ , demonstrating that the difference in the distributions of  $V_{\text{tot}}$  and TD/D becomes greater and greater with the increase of  $N_p$ . Especially for stars hosting 2 and 3+ tranets systems, the p-values are smaller than 0.05, demonstrating statistically smaller Galactic velocities and TD/D (smaller fraction of thick disk).

Next, we compare the number fractions of different Galactic components. The number fractions of different Galactic components are calculated with the following formula:

$$F_{i} = \frac{N_{i}}{N_{D} + N_{TD} + N_{H} + N_{Herc} + N_{IB}},$$
(7)

where *i* represents the different Galactic components, i.e., thin disk (D), thick disk (TD), Halo (H), Hercules stream (Herc) and in between (IB). To obtain the uncertainties, we assume that the observation numbers of different Galactic components obey the Poisson distribution. Then we resample the observation numbers of different Galactic components from the given distribution and calculate the number fractions for 10,000 times. The uncertainty  $(1-\sigma \text{ interval})$  of each parameter is set as the range of  $50 \pm 34.1$  percentiles of the 10,000 calculations. The fractions of different components for the stellar sample without Kepler planets, Kepler planet candidate host sample and the three subsamples are summarized in Table 5. We find that with the increase of planet multiplicity ( $N_p$  from 0 to 3+), the number fraction of the thin disk  $F_D$  increases, while the fraction of thick disk  $F_{TD}$  generally decrease, which are illustrated in the upper and middle panels of Figure 9.

Then, we compare the kinematic ages of stars with and without Kepler planets. The kinematic ages and uncertainties are calculated from the velocity dispersions with Equation 5 and 6. As suggested in PAST I, there is no clear trend between velocity dispersions and ages for stars belonging to Hercules stream and halo. Therefore we only consider stars with Herc/D < 0.5&Herc/TD < 0.5&TD/H > 1 in calculating the kinematic ages. As shown in the bottom panel of Figure 9, the typical ages decrease from  $4.49^{+0.60}_{-0.36}$  Gyr,  $3.82^{+0.50}_{-0.28}$  Gyr,  $2.98^{+0.34}_{-0.29}$  Gyr to  $2.84^{+0.34}_{-0.29}$  Gyr when the number of planets  $N_{\rm p}$  increases from 0 to  $\geq 3$ . In 10,000 sets of resampled data, the kinematic age of neighbor non-Kepler planet candidate host sample is bigger than that of  $N_{\rm p} = 1$ ,  $N_{\rm p} = 2$ , and  $N_{\rm p} \geq 3$  subsamples for 8,639, 9,981, and 9,995 times, corresponding to confidence level of 86.39%, 99.81% and 99.95% respectively. We therefore conclude that the ages of 2 and 3+ tranets systems are statistically smaller than stars without Kepler planets, while there is no significant difference for 1 tranet systems within the precision of our age estimations. We summarized the median value and 1  $\sigma$  interval of physical and kinematic properties in Table 5.

Last but not least, we evaluate the statistical significance of the above trends as seen in Figure 9. As can be seen in Figure 9, with the increase of  $N_{\rm p}$ , the number fraction of the thin disk  $F_{\rm D}$  increases, while the fraction of thick disk  $F_{\rm TD}$  and kinematic age Age<sub>kin</sub> generally decrease. To describe the property of the above trends mathematically, we fit



Figure 9. Fractions of thin disk stars ( $F_D$ , top panel) and thick disk stars ( $F_{TD}$ , middle panel) and kinematic age (Age<sub>kin</sub>, bottom panel) as functions of the transiting planet multiplicity ( $N_p$ ) detected around a star in the LAMOST-Gaia-Kepler sample. The dashed lines represent the best fits of linear model.

the relation between  $F_D$ ,  $F_{TD}$ , Age<sub>kin</sub> as a function of  $N_p$  with two models: constant model ( $y = A_0$ ) and linear model ( $y = A \times N_p + b$ ). For each model, we fit the relationships with the Levenberg-Marquardt algorithm (LMA). In order to compare the fitting of different models, we calculate the Akaike information criterion (AIC, Cavanaugh 1997) for each model. The AIC differences between the constant and linear model ( $\Delta AIC \equiv AIC_{con} - AIC_{lin}$ ) are 14.36, 7.69, and 9.15 for the trend of  $F_D$ ,  $F_{TD}$ , and Age<sub>kin</sub> respectively. Therefore a constant model can be confidently ruled out compared with the linear model with an AIC score difference  $\Delta AIC > 6$ . To obtain the confidence level, we fit the Furthermore, as shown in Figure 4, kinematic age is overestimated as compared to asteroseismic age at the young end, i.e., < 2 - 3 Gyr. If adopting such an age correction, the actual age of multi-planet systems (shown in bottom panel of Figure 9) should be smaller, which will make the trend even stronger.

Based on above tests, we conclude that the trends between  $F_D$ ,  $F_{\text{TD}}$ ,  $Age_{kin}$  and  $N_p$  as shown in Figure 9 are statistically significant. The trends cannot be explained by the spatial bias and other effects related to stellar mass, radius, temperature, and chemical abundances because, as can be seen in Table 5 and Figure 7, the Kepler planet candidate host subsamples and the control sample all have similar distributions in spatial positions and physical properties. Therefore, we believe that these trends may be some footprints left in the evolution of planetary dynamics.

#### 6. SUMMARY

The Kepler mission has detected over 4000 planets (candidates) by monitoring  $\sim 200,000$  stars. Accurately characterizing the properties of these stars is essential to study planet properties and their relations to stellar hosts and environments. Previous studies have investigated some of basic stellar properties, e.g., mass, radius (Berger et al. 2018), metallicity (Dong & Zhu 2013). However, the kinematic properties (e.g. Galactic component memberships and kinematic ages) of these stars are yet to be well characterized.

In this paper, as the Paper II of the PAST series, we construct a LAMOST-Gaia-Kepler catalogue of 35,835 Kepler stars and 764 Kepler planet candidate host stars with kinematic properties as well as other basic stellar properties (section 4, Table 3) by combining data from Gaia DR2 and LAMOST DR4 (section 2, Table 1). By adopting the revised kinematic methods in PAST I, we calculate the kinematics (i.e., Galactic position and velocity) and the Galactic component membership probabilities for the stars and then classify them into different Galactic components. The fractions of stars belonging to the thin disk, thick disk, Halo and Hercules stream are 87.1% (31,218/35,835), 5.1% (1,832/35,835), 0.16% (59/35,835), and 1.5% (545/35,835), respectively. We also explore the kinematics and chemical abundances of different Galactic components. As expected, from the thin disk, Hercules stream, thick disk to the halo, the Galactic velocity is getting larger, the metallicity [Fe/H] decreases and [ $\alpha$ /Fe] increases (Table 4 and Figure 2).

Based on our LAMOST-Gaia-Kepler catalog, we derive the stellar kinematic ages with typical uncertainties of 10-20% by adopting the refined Age-velocity dispersion relation (AVR) of PAST I (Chen et al. 2021). Comparing to ages derived from other methods (section 5.1), we find kinematic age generally matches with asteroseismic, gyrochronology, and isochrone ages (Figure 4). We have carried out some analyses to explore the connection between stellar activity and kinematic age (section 5.2). As expected, as stars age, they spin down and become less active in terms of both the magnetic S index (Figure 5) and the photometric variability (Figure 6), which, in turn, verify the kinematic ages derived in this work.

We have also explored the differences in kinematic properties (e.g., Galactic velocities, thin/thick disk memberships and kinematic ages) between the Kepler planet candidate hosts and stars without Kepler planets (section 5.3). For the aspect of spatial position, the stars without Kepler planets have a wider distribution in the distance to the Sun, Galactic radius (R), azimuth angle ( $\theta$ ) and height (Z) than that of Kepler planet candidate host stars (Figure 7), which could be an observational selection bias as planet systems closer to us are easier to be detected (e.g. McTier & Kipping 2019). Thus for a fair comparison between the kinematic properties of stars with and without Kepler planets, we construct a control sample by selecting the nearest neighbors of planet candidate hosts in the spacial distribution (Figure 7). Comparing to stars in the control sample, we find that planet candidate hosts, especially those with large planet multiplicity  $(N_{\rm p} \ge 3)$ , differ significantly in distributions of the Galactic velocity  $V_{\rm tot}$  and the relative probability between thick and thin disks TD/D (Figure 8). In particular, we find a trend that the fraction of thin (thick) disk stars increases (decreases) with transiting planet multiplicity  $(N_{\rm p})$  and the kinematic age decreases with  $N_{\rm p}$  (Figure 9 and Table 5). This provides insights into the formation and evolution of planetary systems with the Galactic components and stellar age. One possible explanation for the trend is that the long term dynamical evolution can pump up orbital eccentricity/inclination of planets (e.g. Zhou et al. 2007) or even cause planet merge/ejection (e.g. Pu & Wu 2015), which reduces the observed transiting planet multiplicity i.e., N<sub>p</sub>. Specifically, in a subsequent paper of the PAST project (Yang et al. in prep), we will study whether/how planetary occurrence and architecture (e.g. inclination) change with the ages and Galactic environments based on the LAMOST-Gaia-Kepler catalog of this work.

The LAMOST-Gaia-Kepler catalog provides the kinematics, Galactic component-memberships, [Fe/H],  $[\alpha/Fe]$  and ages information for thousands of planets (candidates) down to about Earth radius and tens of thousands of wellcharacterized field stars with no bias toward Kepler planet candidate hosts, which will be useful for more future studies of exoplanets at different positions/components of the Galaxy with different ages. The answers of these questions will deepen our understanding of planet formation and evolution.

# ACKNOWLEDGEMENTS

This work has included data from Guoshoujing Telescope (the Large Sky AreaMulti-Object Fiber Spectroscopic Telescope LAMOST), which is a National Major Scientific Project built by the Chinese Academy of Sciences. Funding for the project has been provided by the National Development and Reform Commission. This work presents results from the European Space Agency (ESA) space mission Gaia. Gaia data are being processed by the Gaia Data Processing and Analysis Consortium (DPAC). Funding for the DPAC is provided by national institutions, in particular the institutions participating in the Gaia MultiLateral Agreement (MLA). The Gaia mission website is https://www.cosmos.esa.int/Gaia. The Gaia archive website is https://archives.esac.esa.int/Gaia. We acknowledge the NASA Exoplanet archive, which is operated by the California Institute of Technology, under contract with the National Aeronautics and Space Administration under the Exoplanet Exploration Program.

This work is supported by the National Key R&D Program of China (No. 2019YFA0405100) and the National Natural Science Foundation of China (NSFC; grant No. 11933001, 11973028, 11803012, 11673011, 12003027). J.-W.X. also acknowledges the support from the National Youth Talent Support Program and the Distinguish Youth Foundation of Jiangsu Scientific Committee (BK20190005). C.L. thanks National Key R&D Program of China No. 2019YFA0405500 and the National Natural Science Foundation of China (NSFC) with grant No. 11835057. H.F.W. is supported by the LAMOST Fellow project, National Key Basic R&D Program of China via 2019YFA0405500 and funded by China Postdoctoral Science Foundation via grant 2019M653504 and 2020T130563, Yunnan province postdoctoral Directed culture Foundation, and the Cultivation Project for LAMOST Scientific Payoff and Research Achievement of CAMS-CAS. M. Xiang & Y. Huang acknowledge the National Natural Science Foundation of China (grant No. 11703035).

# REFERENCES

- Adibekyan, V. Z., Figueira, P., Santos, N. C., et al. 2013, A&A, 554, A44, doi: 10.1051/0004-6361/201321520
- Akeson, R. L., Chen, X., Ciardi, D., et al. 2013, PASP, 125, 989, doi: 10.1086/672273
- Barnes, S. A. 2010, ApJ, 722, 222, doi: 10.1088/0004-637X/722/1/222
- Barnes, S. A., & Kim, Y.-C. 2010, ApJ, 721, 675, doi: 10.1088/0004-637X/721/1/675
- Bastien, F. A., Stassun, K. G., Basri, G., & Pepper, J. 2013, Nature, 500, 427, doi: 10.1038/nature12419
- Bensby, T., Feltzing, S., & Lundström, I. 2003, A&A, 410, 527, doi: 10.1051/0004-6361:20031213
- Bensby, T., Feltzing, S., & Oey, M. S. 2014, A&A, 562, A71, doi: 10.1051/0004-6361/201322631
- Bergemann, M., Ruchti, G. R., Serenelli, A., et al. 2014, A&A, 565, A89, doi: 10.1051/0004-6361/201423456
- Berger, T. A., Huber, D., Gaidos, E., & van Saders, J. L. 2018, ApJ, 866, 99, doi: 10.3847/1538-4357/aada83
- Berger, T. A., Huber, D., Gaidos, E., van Saders, J. L., & Weiss, L. M. 2020a, AJ, 160, 108, doi: 10.3847/1538-3881/aba18a

- Berger, T. A., Huber, D., van Saders, J. L., et al. 2020b, AJ, 159, 280, doi: 10.3847/1538-3881/159/6/280
- Boch, T., Pineau, F., & Derriere, S. 2012, in Astronomical Society of the Pacific Conference Series, Vol. 461,
  Astronomical Data Analysis Software and Systems XXI,
  ed. P. Ballester, D. Egret, & N. P. F. Lorente, 291
- Bonaca, A., Conroy, C., Wetzel, A., Hopkins, P. F., & Kereš, D. 2017, ApJ, 845, 101, doi: 10.3847/1538-4357/aa7d0c
- Brown, T. M., Latham, D. W., Everett, M. E., & Esquerdo, G. A. 2011, AJ, 142, 112,

doi: 10.1088/0004-6256/142/4/112

Cavanaugh, J. E. 1997, Statistics & Probability Letters, 33, 201 ,

doi: https://doi.org/10.1016/S0167-7152(96)00128-9

- Chen, B., Stoughton, C., Smith, J. A., et al. 2001, ApJ, 553, 184, doi: 10.1086/320647
- Chen, D.-C., Xie, J.-W., Zhou, J.-L., et al. 2021, ApJ, 909, 115, doi: 10.3847/1538-4357/abd5be
- Choi, J., Dotter, A., Conroy, C., et al. 2016, ApJ, 823, 102, doi: 10.3847/0004-637X/823/2/102

- Christiansen, J. L., Jenkins, J. M., Caldwell, D. A., et al. 2012, PASP, 124, 1279, doi: 10.1086/668847
- Cui, X.-Q., Zhao, Y.-H., Chu, Y.-Q., et al. 2012, Research in Astronomy and Astrophysics, 12, 1197, doi: 10.1088/1674-4527/12/9/003
- De Cat, P., Fu, J. N., Ren, A. B., et al. 2015, ApJS, 220, 19, doi: 10.1088/0067-0049/220/1/19
- Dong, S., & Zhu, Z. 2013, ApJ, 778, 53, doi: 10.1088/0004-637X/778/1/53
- Dotter, A. 2016, ApJS, 222, 8, doi: 10.3847/0067-0049/222/1/8
- Egeland, R., Soon, W., Baliunas, S., et al. 2017, ApJ, 835, 25, doi: 10.3847/1538-4357/835/1/25
- Feltzing, S., Bensby, T., & Lundström, I. 2003, A&A, 397, L1, doi: 10.1051/0004-6361:20021661
- Fu, J.-N., Cat, P. D., Zong, W., et al. 2020, Research in Astronomy and Astrophysics, 20, 167, doi: 10.1088/1674-4527/20/10/167
- Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016, A&A, 595, A1, doi: 10.1051/0004-6361/201629272
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018a, A&A, 616, A1, doi: 10.1051/0004-6361/201833051
- Gaia Collaboration, Katz, D., Antoja, T., et al. 2018b,
   A&A, 616, A11, doi: 10.1051/0004-6361/201832865
- Gilliland, R. L., Chaplin, W. J., Dunham, E. W., et al. 2011, ApJS, 197, 6, doi: 10.1088/0067-0049/197/1/6
- Huber, D., Silva Aguirre, V., Matthews, J. M., et al. 2014, ApJS, 211, 2, doi: 10.1088/0067-0049/211/1/2
- Johnson, D. R. H., & Soderblom, D. R. 1987, AJ, 93, 864, doi: 10.1086/114370
- Kallinger, T., De Ridder, J., Hekker, S., et al. 2014, A&A, 570, A41, doi: 10.1051/0004-6361/201424313
- Luo, A. L., Zhang, H.-T., Zhao, Y.-H., et al. 2012, Research in Astronomy and Astrophysics, 12, 1243, doi: 10.1088/1674-4527/12/9/004
- Majewski, S. R., Schiavon, R. P., Frinchaboy, P. M., et al. 2017, AJ, 154, 94, doi: 10.3847/1538-3881/aa784d
- Mathur, S., Huber, D., Batalha, N. M., et al. 2017, ApJS, 229, 30, doi: 10.3847/1538-4365/229/2/30
- McQuillan, A., Aigrain, S., & Mazeh, T. 2013a, MNRAS, 432, 1203, doi: 10.1093/mnras/stt536
- McQuillan, A., Mazeh, T., & Aigrain, S. 2013b, ApJL, 775, L11, doi: 10.1088/2041-8205/775/1/L11
- McTier, M. A. S., & Kipping, D. M. 2019, MNRAS, 489, 2505, doi: 10.1093/mnras/stz2088

- Paxton, B., Bildsten, L., Dotter, A., et al. 2011, ApJS, 192, 3, doi: 10.1088/0067-0049/192/1/3
- Paxton, B., Cantiello, M., Arras, P., et al. 2013, ApJS, 208, 4, doi: 10.1088/0067-0049/208/1/4
- Paxton, B., Marchant, P., Schwab, J., et al. 2015, ApJS, 220, 15, doi: 10.1088/0067-0049/220/1/15
- Paxton, B., Schwab, J., Bauer, E. B., et al. 2018, ApJS, 234, 34, doi: 10.3847/1538-4365/aaa5a8
- Pedregosa, F., Varoquaux, G., Gramfort, A., et al. 2011, J. Mach. Learn. Res., 12, 2825–2830
- Pinsonneault, M. H., Elsworth, Y. P., Tayar, J., et al. 2018, ApJS, 239, 32, doi: 10.3847/1538-4365/aaebfd
- Pu, B., & Wu, Y. 2015, ApJ, 807, 44, doi: 10.1088/0004-637X/807/1/44
- Reid, M. J., Menten, K. M., Brunthaler, A., et al. 2014, ApJ, 783, 130, doi: 10.1088/0004-637X/783/2/130
- Reinhold, T., Shapiro, A. I., Solanki, S. K., et al. 2020, Science, 368, 518, doi: 10.1126/science.aay3821
- Ren, A., Fu, J., De Cat, P., et al. 2016, ApJS, 225, 28, doi: 10.3847/0067-0049/225/2/28
- Rizzuto, A. C., Vanderburg, A., Mann, A. W., et al. 2018, AJ, 156, 195, doi: 10.3847/1538-3881/aadf37
- Soderblom, D. R. 2010, ARA&A, 48, 581, doi: 10.1146/annurev-astro-081309-130806
- Spada, F., & Lanzafame, A. C. 2020, A&A, 636, A76, doi: 10.1051/0004-6361/201936384
- Su, D.-Q., & Cui, X.-Q. 2004, ChJA&A, 4, 1, doi: 10.1088/1009-9271/4/1/1
- Tian, H.-J., Liu, C., Carlin, J. L., et al. 2015, ApJ, 809, 145, doi: 10.1088/0004-637X/809/2/145
- Wang, S.-G., Su, D.-Q., Chu, Y.-Q., Cui, X., & Wang, Y.-N. 1996, ApOpt, 35, 5155, doi: 10.1364/AO.35.005155
- Wu, Y., Luo, A. L., Li, H.-N., et al. 2011, Research in Astronomy and Astrophysics, 11, 924, doi: 10.1088/1674-4527/11/8/006
- Xiang, M. S., Liu, X. W., Yuan, H. B., et al. 2015, MNRAS, 448, 822, doi: 10.1093/mnras/stu2692
- —. 2017, MNRAS, 467, 1890, doi: 10.1093/mnras/stx129
- Zhang, J., Bi, S., Li, Y., et al. 2020, ApJS, 247, 9, doi: 10.3847/1538-4365/ab6165
- Zhao, G., Zhao, Y.-H., Chu, Y.-Q., Jing, Y.-P., & Deng, L.-C. 2012, Research in Astronomy and Astrophysics, 12, 723, doi: 10.1088/1674-4527/12/7/002
- Zhou, J.-L., Lin, D. N. C., & Sun, Y.-S. 2007, ApJ, 666, 423, doi: 10.1086/519918
- Zong, W., Fu, J.-N., De Cat, P., et al. 2018, ApJS, 238, 30, doi: 10.3847/1538-4365/aadf81