

Two-Class (r, k) -Coloring: Coloring with Service Guarantees*

Pál András Papp

ETH Zurich, Switzerland

apapp@ethz.ch

Roland Schmid

ETH Zurich, Switzerland

roschmi@ethz.ch

Valentin Stoppiello

ETH Zurich, Switzerland

stoppiev@ethz.ch

Roger Wattenhofer

ETH Zurich, Switzerland

wattenhofer@ethz.ch

Abstract

This paper introduces the *Two-Class (r, k) -Coloring* problem: Given a fixed number of k colors, such that only r of these k colors allow conflicts, what is the minimal number of conflicts incurred by an optimal coloring of the graph?

We establish that the family of Two-Class (r, k) -Coloring problems is NP-complete for any $k \geq 2$ when $(r, k) \neq (0, 2)$. Furthermore, we show that Two-Class (r, k) -Coloring for $k \geq 2$ colors with one ($r = 1$) relaxed color cannot be approximated to any constant factor (\notin APX). Finally, we show that Two-Class (r, k) -Coloring with $k \geq r \geq 2$ colors is APX-complete.

2012 ACM Subject Classification Mathematics of computing \rightarrow Graph coloring; Theory of computation \rightarrow Problems, reductions and completeness

Keywords and phrases graph coloring, conflict coloring, NP-complete, approximation complexity

1 Introduction

Graph coloring is a fundamental mathematical problem with various applications in computer science. Usually, we want to color the nodes of a graph with a minimum number of colors such that no two adjacent nodes have the same color.

In many real world applications, however, the number of available colors is strictly limited. For example, we are given a fixed set of frequencies, and we need to assign a frequency (color) to each wireless transmitter (node in a graph), such that wireless transmitters that interfere with each other (that are connected by an edge in the graph) do not use the same frequency. In cases where the set of colors is fixed, we may need to accept a certain number of conflicts, i.e., neighboring nodes which have the same color.

In practice, we often have coloring problems that additionally require strict service guarantees, e.g. priority service for premium customers, or unobstructed communication channels for maintenance and emergency personnel. Such applications can be modeled by partitioning the set of colors into (i) *proper* colors for guaranteed conflict-free service and (ii) *relaxed* colors which allow conflicts. In general, we can have k colors, with r of these k colors being relaxed and $k - r$ of them being proper; we refer to this setting as the Two-Class (r, k) -Coloring problem.

* The authors of this paper are alphabetically ordered.

43 **Contributions**

44 In this paper, we introduce and motivate the *Two-Class (r, k) -Coloring* problem, that is, the
 45 problem of coloring a given graph with a fixed number of k colors of which r colors are relaxed
 46 to allow conflicts. We establish relations between the Two-Class (r, k) -Coloring problem and
 47 several well-studied problems, such as Maximum Independent Set and Max- k -Cut. For our
 48 main contributions, we show that:

- 49 ■ the family of Two-Class (r, k) -Coloring problems is NP-complete, except for the special
 50 case when $(r, k) = (0, 2)$,
- 51 ■ Two-Class $(1, k)$ -Coloring, i.e., with one relaxed color, cannot be approximated to any
 52 constant factor within polynomial time (\notin APX),
- 53 ■ Two-Class (r, k) -Coloring with multiple relaxed colors is APX-complete, i.e., Two-Class
 54 (r, k) -Coloring can be approximated to some constant (but not to an arbitrary constant)
 55 in polynomial time.

56 **2 Related Work**

57 Graph coloring is one of the most fundamental and well-studied problems of computer science.
 58 A general survey of fundamental graph coloring results is available in [12] or [17]. Even
 59 though it has been studied intensively since the early 1970s, graph coloring still receives
 60 significant attention today, with results ranging from distributed algorithms [2] to modified
 61 variants [12] or heuristic solutions for such problems [10]. However, the vast majority of
 62 these studies only consider proper colorings of graphs, i.e. when monochromatic edges are
 63 not allowed at all.

64 The most well-studied relaxed coloring problem is *Defective Coloring*, introduced in [4].
 65 A defective coloring is a (possibly relaxed) coloring of a graph G such that each node of G
 66 has at most d conflicts. Given a graph G and a fixed number of colors k , the goal of the
 67 problem is to find a k -coloring with the minimal possible d value. Defective coloring is also
 68 known to be NP-complete [5], and has been studied extensively on specific classes of graphs
 69 [1, 3, 8].

70 In the case when all color classes allow conflicts, a solution minimizing the number of
 71 conflicts is equivalently a solution maximizing the number of edges that go between different
 72 color classes. Thus the Two-Class (k, k) -Coloring problem is essentially a reformulation of the
 73 Max- k -Cut problem. The Max- k -Cut problem is also a thoroughly studied problem, which is
 74 known to be both NP-complete and APX-complete [11, 13, 16]. Further theoretical work
 75 mostly investigates the best possible constant-factor approximation for the problem [9, 14, 7].
 76 Many of these studies focus primarily on the weighted version of the problem, and obtain
 77 their results for the unweighted graphs as a special case of this.

78 Another related question is the Maximum k -Colorable Subgraph problem: given a graph,
 79 the task here is to select the largest subset V_0 of the vertices such that the subgraph induced
 80 by V_0 still admits a proper k -coloring [18, 19]. Note that the term “Maximum k -Colorable
 81 Subgraph problem” is quite ambiguous in the literature, as it is often also used to refer to the
 82 problem of finding a k -colorable (not necessarily induced) subgraph with maximally many
 83 edges, which is essentially only another reformulation of Max- k -Cut.

84 Our work is also closely connected to the Maximum Independent Set (MaxIS) problem,
 85 which is known to be NP-complete [11], and also not approximable to any constant [13].

86 **3 Two-Class (r, k) -Coloring**

87 We have seen that several relaxations of the traditional graph coloring problem have been
 88 considered in the literature. In this work, we introduce the problem of *Two-Class (r, k) -*
 89 *Coloring*, which follows a utilitarian approach in the sense that we minimize the global
 90 number of conflicts. To begin with, let us introduce some notation.

91 As usual, a graph $G = (V, E)$ consists of a set of n vertices V (with $|V| = n$) and a set of
 92 undirected edges $E \subseteq \binom{V}{2}$. The degree of a node $v \in V$ is denoted by $d(v)$.

93 For such a graph, a k -coloring is a mapping of its n vertices to a set of k colors.
 94 Traditionally, any set of nodes that is assigned the same color is required to form an
 95 independent set in the graph.

96 ► **Definition 3.1.** *The assignment of the same color to two adjacent vertices is called a*
 97 *conflict. Similarly, the edge connecting these two vertices is called a conflict edge (for being*
 98 *a witness of the conflict). All other edges are called covered edges.*

99 Graph Coloring does not allow conflicts at all. In our problems, however, the k colors
 100 are sorted into two groups, and we only require the colors in one of the groups to form
 101 independent sets.

102 ► **Definition 3.2.** *A color is said to be a proper color if the vertices mapped to this color are*
 103 *required to form an independent set (i.e., have no conflicts). Otherwise, the color is called a*
 104 *relaxed color, which might have (or might not have) conflicts.*

105 Given these definitions, we can now define the colorings that we are interested in:

106 ► **Definition 3.3.** *In the Two-Class (r, k) -Coloring problem (with $r \leq k$), a feasible solution*
 107 *is a k -coloring of the input graph such that we can divide the k colors into a group of r relaxed*
 108 *colors and a group of $k - r$ proper colors.*

109 The natural goal in such a problem is to color the graph with as few conflicts as possible.
 110 However, when discussing the connection to related problems, it is beneficial to look at this
 111 problem in the dual perspective of having as many covered edges as possible. Note that
 112 maximizing the number of covered edges in a coloring indeed produces a solution that also
 113 minimizes the number of conflicts. Thus, we study the following problem in the paper.

114 ► **Definition 3.4.** *The Two-Class (r, k) -Coloring problem is defined as follows:*

- 115 ■ **Decision problem:** *Given a graph G and numbers r, k , and c , is there a feasible solution*
 116 *of (r, k) -Color which covers at least c edges?*
- 117 ■ **Optimization problem:** *Given a graph G and numbers r and k , what is the maximal*
 118 *number of edges in G that can be covered by a feasible solution of (r, k) -Color?*

119 Let us finish with some further remarks on notations. Generally, given an optimization
 120 problem P , we denote the value of the optimal solution on graph G by $\text{opt}_P(G)$. For an
 121 arbitrary feasible solution \mathcal{S} of the problem, the value of \mathcal{S} is denoted by $\text{val}_P(\mathcal{S})$.

122 While we study all the above problems on simple graphs only, we also use multigraphs
 123 as a tool in one of our proofs. A multigraph is a graph that allows parallel edges between
 124 nodes, i.e. where E is a multiset of elements from $\binom{V}{2}$.

125 **Example: Two-Class (r, k) -Coloring is Not Equivalent to MaxIS**

126 Let us briefly compare $(1, 2)$ -Color and the Maximum Independent Set (MaxIS) problem on
 127 the example graph of Figure 1. Both problems can be formulated as a coloring problem with

128 one relaxed (white) and one proper (black) color, so the set of feasible solutions for the two
 129 problems are identical. However, while MaxIS aims to maximize the number of black vertices,
 130 $(1, 2)$ -Color maximizes the number of edges covered by the black vertices instead. Figure 1
 131 shows that these problems may have different optimal solutions: the optimum of $(1, 2)$ -Color
 132 has only 5 black nodes and covers 13 edges (left), while the Maximum Independent Set
 contains 10 nodes and covers 11 edges (right).



(a) Optimal $(1, 2)$ -Color solution covering 13 edges. (b) Maximum independent set of size 10.

■ **Figure 1** The $(1, 2)$ -Color problem is not equivalent to the Maximum Independent Set problem.

133

134 4 Complexity Results (NP-Completeness)

135 In this section, we show that (r, k) -Color is NP-complete for $k > 2$ with $0 \leq r \leq k$, and
 136 also for $k = 2$ with $r \in \{1, 2\}$. To begin with, observe that deciding (r, k) -Color is in NP:
 137 A solution to (r, k) -Color covering at least c edges can be verified in polynomial time by
 138 traversing the set of covered edges.

139 Now, for the case $k > 2$, recall that deciding the traditional graph coloring problem
 140 k -Color is NP-complete for $k > 2$ colors:

141 ► **Theorem 4.1** ([11]). *It is NP-complete to decide whether an input graph G admits a*
 142 *k -coloring for a given number of proper colors $k > 2$.*


143 To show that (r, k) -Color is NP-complete for $k > 2$, we may thus show that a polynomial-
 144 time algorithm for solving (r, k) -Color could be used to compute a solution for k -Color in
 145 polynomial time. To that end, note that setting $c = |E|$ directly yields a decision procedure
 146 for k -Color.

147 ► **Lemma 4.2.** *For $k > 2$, it is NP-complete to decide whether an input graph G admits a*
 148 *solution to (r, k) -Color covering at least c edges.*

149 The case for $k = 2$ must be treated separately, as k -Color for $k = 2$ colors (deciding
 150 bipartiteness of a graph) can be solved in linear time, e.g. using a breadth-first search.
 151 Hence, $(0, 2)$ -Color can be decided in linear time for $c = |E|$. For $c < |E|$, note that if there
 152 exists a proper 2-coloring of the input graph G , then it covers all $|E|$ edges and is hence
 153 a solution to $(0, 2)$ -Color for any $c < |E|$. If there exists no proper 2-coloring of G but a
 154 solution \mathcal{S} to $(0, 2)$ -Color covering at least $c < |E|$ edges, \mathcal{S} would have to induce at least
 155 one conflict. However, in $(0, 2)$ -Color there are no relaxed colors that would allow for any
 156 conflicts. Consequently, such a solution \mathcal{S} may not exist and $(0, 2)$ -Color can be decided for
 157 any c in linear time.

158 Perhaps surprisingly, we will show that (r, k) -Color is also NP-complete for $k = 2$ and
 159 $r > 0$. In other words, relaxing at least one color creates a gap in the computational
 160 complexity (assuming $P \neq NP$).

161 To show that $(r, 2)$ -Color is also NP-hard to decide for $r > 0$ and some given minimum
 162 number of covered edges c , note that $(2, 2)$ -Color is equivalent to the well-known MaxCut



■ **Figure 2** Illustration of the reduction to Maximum Independent Set: Given an original graph G on n nodes, we add a clique C of size n^2 and connect every node of G to every node of C .

163 problem which is known to be NP-complete [11]. For (1,2)-Color, we propose a reduction to
 164 the Maximum Independent Set problem. To begin with, let us recall that:

165 ► **Theorem 4.3** ([11]). *It is NP-complete to decide whether an input graph G contains an*
 166 *independent set of some given size k .*

167 We propose the following reduction: Given an input graph $G = (V_G, E_G)$, we determine
 168 whether there exists an independent set of size k as follows:

- 169 1. add a clique C of size $|V_C| = n^2$ to the graph G and connect each clique-vertex $u \in V_C$
 170 with each graph vertex $v \in V_G$ to obtain the modified graph $G' = (V_{G'}, E_{G'})$, as shown
 171 in Figure 2
- 172 2. decide whether there is a solution for (1,2)-Color on G' covering at least $c = k \cdot n^2$ edges,
- 173 3. output the same answer for whether an independent set of size k exists in G .

174 ► **Lemma 4.4.** *It is NP-hard to decide (1,2)-Color.*

175 **Proof.** We will show that (1,2)-Color is NP-hard by arguing that the reduction given above
 176 correctly decides the Maximum Independent Set problem on any given input graph G in
 177 polynomial time, if we assume that (1,2)-Color can be decided in polynomial time. To begin
 178 with, note that the proposed reduction requires only polynomial time.

179 As for the correctness, we assume the two colors are *black* (proper color, i.e. an independent
 180 set) and *white* (relaxed color) and argue that an optimal solution to the (1,2)-Color problem
 181 covering at least $c = k \cdot n^2$ edges exists if and only if there exists an independent set of size k
 182 in G (that may be colored black to establish the required conflict bound c).

183 If there exists an independent set of size k in G , then we can simply color this independent
 184 set in black and cover at least $k \cdot n^2$ edges from each of the k vertices in V_G to each of the n^2
 185 vertices in C . Hence, there exists a solution to (1,2)-Color covering at least $k \cdot n^2$ edges.

186 For the inverse direction, we first argue that the added clique C can be assumed to be
 187 colored entirely white in an optimal solution. Assume for contradiction that it is strictly
 188 optimal to color at least one node in C black. If the clique C contains one black node, no
 189 other vertex of the graph G' can be colored black as black is a proper color and each clique
 190 vertex is fully connected in the graph G' . Hence, such a coloring may cover at most $n^2 + n - 1$
 191 edges. However, this can only be optimal for a maximum independent set of size 1 in G ;
 192 otherwise, we could simply color an independent set of size two in black and cover at least

193 $2n^2 > n^2 + n - 1$ edges. For a maximum independent set of size 1, G must be a clique itself
 194 and thus, any node $v \in V_G$ covers $n^2 + n - 1$ edges as well. Hence, without loss of generality,
 195 we may assume that there exists an optimal solution of $(1, 2)$ -Color on G' where the added
 196 clique C is colored entirely white.

197 It remains to show the following: If there exists a solution to the $(1, 2)$ -Color problem
 198 covering at least $k \cdot n^2$ edges in G' , there must exist an independent set of size k in G . As
 199 argued above, there must exist at least one optimal solution \mathcal{S} in which C is colored entirely
 200 white. We show that \mathcal{S} assigns at least k vertices in G the color black; in other words,
 201 showing that there exists an independent set of size k .

202 Assume (for contradiction) that any maximum independent set on G had size at most
 203 $k - 1$. Furthermore, note that $|E_G| < n^2$ as G is a simple graph on n vertices. Hence,
 204 the solution \mathcal{S} for $(1, 2)$ -Color could cover at most $(k - 1) \cdot n^2 + |E_G| < k \cdot n^2$ edges – a
 205 contradiction. ◀

206 Ultimately, we combine Lemma 4.2, Lemma 4.4 and the known NP-completeness results of
 207 the MaxCut problem [11] to obtain:

208 ▶ **Theorem 4.5.** *For $k \geq 2$ and $(r, k) \neq (0, 2)$, it is NP-complete to decide whether an input*
 209 *graph G admits a solution to (r, k) -Color covering at least c edges.*

210 ▶ **Remark 4.6.** The reductions presented in Section 4 demonstrate that the decision problems
 211 (r, k) -Color are NP-complete for $k \geq 2$ and $(r, k) \neq (0, 2)$. Similarly, it can be shown that the
 212 naturally corresponding optimization problems are NP-hard. Thus, subsequently, we study
 213 the approximability of these optimization problems.

214 5 Approximability Results

215 In this section, we discuss the approximability of the optimal solution in (r, k) -Color problems.
 216 Our results show that the problems (r, k) -Color essentially behave like Max- k -Cut for $r \geq 2$,
 217 while they behave like Maximum Independent Set for $r = 1$. An illustration of the resulting
 218 approximation complexity classes can be found in Figure 3.


219 We assume some familiarity with the polynomial-time approximation complexity classes
 220 APX and PTAS. Intuitively, PTAS contains all problems that can be approximated to any
 221 constant factor, while APX contains all problems that can be approximated to some constant
 222 factor. Problems outside of APX can only be approximated to factors that depend on n .

223 We apply two kinds of reductions in this section. Both reductions reduce a (generally
 224 more well-known) problem P_1 to another problem P_2 . Given a problem P_1 on an input graph
 225 G , our reduction first transforms G into an instance of problem P_2 (in our case, another
 226 input graph G'). Then, assuming that an oracle provides us with a solution \mathcal{S} of problem
 227 P_2 for the input graph G' , we show how to derive a solution \mathcal{S}' for problem P_1 on graph G .
 228 Throughout this section, we leave it to the reader to verify that both transformations are
 229 computable in polynomial time.

230 ▶ **Definition 5.1.** *A reduction is called a continuous reduction if there exists a constant α*
 231 *such that for any input graph G and any possible solution \mathcal{S} obtained on G' , we have*

$$232 \frac{\text{opt}_{P_1}(G)}{\text{val}_{P_1}(\mathcal{S}')} \leq \alpha \cdot \frac{\text{opt}_{P_2}(G')}{\text{val}_{P_2}(\mathcal{S})}.$$

233 Continuous reductions are known to preserve APX-membership; hence, if $P_1 \notin \text{APX}$ is
 234 known, then a continuous reduction from P_1 to P_2 proves that P_2 is not in APX either.



■ **Figure 3** The Two-Class (r, k) -Coloring problems can be separated in two different approximation complexity classes. Two-Class (r, k) -Coloring is APX-complete for $r \geq 2$, while Two-Class $(1, k)$ -Coloring cannot be approximated to any constant factor within polynomial time (\notin APX).

235 An L -reduction (linear reduction), on the other hand, preserves membership in PTAS,
 236 and also preserves APX-completeness.

237 ► **Definition 5.2.** We say that a reduction is an L -reduction if there are constants α, β such
 238 that the following two conditions hold:

239 ■ For any graph G , we have

240
$$\text{opt}_{P_2}(G') \leq \alpha \cdot \text{opt}_{P_1}(G).$$

241 ■ For any graph G and possible solution \mathcal{S} on G' , we have

242
$$\text{opt}_{P_1}(G) - \text{val}_{P_1}(\mathcal{S}') \leq \beta \cdot (\text{opt}_{P_2}(G') - \text{val}_{P_2}(\mathcal{S})).$$

243 Note that these are simplified forms of the definitions, using the fact that we only study
 244 maximization problems in the paper. For a more detailed discussion of classes and tools in
 245 approximation complexity, we refer the reader to [6].

246 **5.1 Approximations for $(1, k)$ -Color**

247 We show that the problems $(1, k)$ -Color are intuitively more similar to the Maximum In-
 248 dependent Set (MaxIS) problem than to MaxCut. To that end, we first discuss $(1, 2)$ -Color
 249 separately and show that there exists a continuous reduction from MaxIS to the $(1, 2)$ -Color
 250 problem. Since it is known that MaxIS \notin APX, this implies $(1, 2)$ -Color \notin APX.

251 In fact, the reduction from MaxIS to $(1, 2)$ -Color follows in a rather straightforward way
 252 from a further analysis of the NP-completeness reduction presented in Section 4. Recall that
 253 we transformed the input graph $G = (V_G, E_G)$ on n nodes into a graph $G' = (V_{G'}, E_{G'})$ on

254 $n^2 + n$ nodes by adding a clique C of size n^2 . Given an algorithm to find an approximate
 255 solution \mathcal{S} to problem $(1, 2)$ -Color in G' in polynomial time, we derive an approximation for
 256 the MaxIS problem on the graph G by selecting the nodes in V_G that have been colored with
 257 the proper color. Once again, we assume the two colors are black (proper color) and white
 258 (relaxed color). The independent set of black nodes is denoted by \mathcal{I} .

259 We distinguish two cases. First, assume that the algorithm returns a coloring on G'
 260 where $|\mathcal{I}| = 1$. Recall that such a coloring can cover at most $n^2 + n - 1$ edges, as this is the
 261 highest possible degree in G' . For the MaxIS approximation, we then choose $\mathcal{I} = \{v\}$ some
 262 arbitrary $v \in V_G$.

263 Otherwise, at least two nodes are colored black. Recall that this implies that all black
 264 nodes are chosen from the original input graph G , as any node of the clique C is connected to
 265 all other nodes in G' . The independent set \mathcal{I} covers at most $|\mathcal{I}| \cdot (n - |\mathcal{I}|)$ edges within G (that
 266 is, of E_G), and exactly $|\mathcal{I}| \cdot n^2$ edges between \mathcal{I} and C . Note that $|\mathcal{I}| \cdot (n - |\mathcal{I}|) \leq |E_G| < n^2$.
 267 In total, this amounts to at most $|\mathcal{I}| \cdot n^2 + n^2$ edges.

268 Since this bound equally holds for the case with $|\mathcal{I}| = 1$ above, we can establish

$$269 \quad \text{val}_{(1,2)\text{-Color}}(\mathcal{S}) < |\mathcal{I}| \cdot n^2 + n^2 = (|\mathcal{I}| + 1) \cdot n^2.$$

270 On the other hand, by choosing a maximum independent set in G (and covering n^2 edges
 271 to the clique with each of its nodes), it follows that

$$272 \quad \text{opt}_{(1,2)\text{-Color}}(G') \geq \text{opt}_{\text{MaxIS}}(G) \cdot n^2.$$

273 Combining these two observations and interpreting the set of black nodes \mathcal{I} as the approximate
 274 solution for the MaxIS problem on G , we get

$$275 \quad \frac{\text{opt}_{(1,2)\text{-Color}}(G')}{\text{val}_{(1,2)\text{-Color}}(\mathcal{S})} \geq \frac{\text{opt}_{\text{MaxIS}}(G) \cdot n^2}{(|\mathcal{I}| + 1) \cdot n^2} = \frac{\text{opt}_{\text{MaxIS}}(G)}{|\mathcal{I}| + 1} \stackrel{|\mathcal{I}| \geq 1}{\geq} \frac{1}{2} \cdot \frac{\text{opt}_{\text{MaxIS}}(G)}{|\mathcal{I}|} = \frac{1}{2} \cdot \frac{\text{opt}_{\text{MaxIS}}(G)}{\text{val}_{\text{MaxIS}}(\mathcal{I})},$$

276 so the condition of continuous reduction is indeed satisfied for $\alpha = 2$. Hence, the existence of
 277 a δ -approximation for the $(1, 2)$ -Color problem would allow us to derive a $\delta/2$ -approximation
 278 for the MaxIS problem. Knowing that MaxIS \notin APX, we conclude that $(1, 2)$ -Color \notin APX.

279 Generalization to $(1, k)$ -Color

280 We now present a continuous reduction from $(1, 2)$ -Color to $(1, k)$ -Color. Together with our
 281 previous result, this shows that $(1, k)$ -Color can also not be approximated to any constant
 282 factor.

283 ► **Lemma 5.3.** *There exists a continuous reduction from $(1, 2)$ -Color to $(1, k)$ -Color for any
 284 constant $k \geq 2$.*

285 **Proof.** Given a solution \mathcal{S} of the $(1, k)$ -Color problem on graph G , we show how to transform
 286 \mathcal{S} into a solution \mathcal{S}' for the $(1, 2)$ -Color problem on the same graph G , while only losing a
 287 constant approximation factor. Note that k is assumed to be a constant.

288 Generally, \mathcal{S} can be regarded as a partitioning of the nodes into independent sets
 289 I_1, \dots, I_{k-1} , and a relaxed color set V_k . Without loss of generality, we can assume that the
 290 sets I_1, \dots, I_{k-1} are sorted in descending order by the number of their outgoing edges $\text{out}(I_j)$,
 291 i.e., the number of edges that have exactly one endpoint in I_j .

292 In order to turn this into a solution \mathcal{S}' for the $(1, 2)$ -Color problem on the same graph G ,
 293 let us select the independent set I_1 as the proper color class of \mathcal{S}' , and let the relaxed color
 294 class consist of all the remaining nodes, i.e. $V_0 := \left(\bigcup_{j=2}^{k-1} I_j\right) \cup V_k$. As I_1 is an independent

295 set, the partitioning (I_1, V_0) is indeed a feasible solution of $(1, 2)$ -Color. Note that the value
 296 of this solution is by definition $\text{val}_{(1,2)\text{-Color}}(\mathcal{S}') = \text{out}(I_1)$.

297 In the original solution \mathcal{S} , since we only have one relaxed color, all covered edges have at
 298 least one of their endpoints in the sets I_1, \dots, I_{k-1} , and thus

$$299 \quad \text{val}_{(1,k)\text{-Color}}(\mathcal{S}) \leq \sum_{j=1}^{k-1} \text{out}(I_j) \leq (k-1) \cdot \text{out}(I_1),$$

300 because we assumed that I_1 is the independent set with the maximum number of outgoing
 301 edges $\text{out}(I_1)$. This shows that

$$302 \quad \frac{1}{k-1} \cdot \text{val}_{(1,k)\text{-Color}}(\mathcal{S}) \leq \text{val}_{(1,2)\text{-Color}}(\mathcal{S}').$$

303 Since the set of feasible colorings in $(1, 2)$ -Color is a subset of those in $(1, k)$ -Color

$$304 \quad \text{opt}_{(1,2)\text{-Color}}(G) \leq \text{opt}_{(1,k)\text{-Color}}(G)$$

305 follows straightforwardly. Thus we have

$$306 \quad \frac{\text{opt}_{(1,2)\text{-Color}}(G)}{\text{val}_{(1,2)\text{-Color}}(\mathcal{S}')} \leq \frac{\text{opt}_{(1,k)\text{-Color}}(G)}{\frac{1}{k-1} \cdot \text{val}_{(1,k)\text{-Color}}(\mathcal{S})} = (k-1) \cdot \frac{\text{opt}_{(1,k)\text{-Color}}(G)}{\text{val}_{(1,k)\text{-Color}}(\mathcal{S})},$$

307 showing that this is indeed a continuous reduction for $\alpha = (k-1)$. \blacktriangleleft

308 Since we have already seen that $(1, 2)$ -Color \notin APX, this implies the following theorem.

309 \blacktriangleright **Theorem 5.4.** *For any constant $k \geq 2$, the problem $(1, k)$ -Color \notin APX.*

310 5.2 Approximations for (r, k) -Color with $r \geq 2$

311 For this case, recall that (k, k) -Color is essentially a reformulation of the Max- k -Cut problem.
 312 Since Max- k -Cut is known to be APX-complete [16], the same naturally holds for (k, k) -Color.
 313 Let us consider some other $r \in \{2, \dots, k-1\}$.

314 We begin by showing that (r, k) -Color is also contained in APX. It is known that using the
 315 probabilistic method and derandomization, one can obtain a simple deterministic algorithm
 316 that finds a partitioning (V_1, V_2) which cuts at least $\frac{|E|}{2}$ edges [15]. This already ensures
 317 that the algorithm in question returns a $\frac{1}{2}$ -approximation for the MaxCut problem.

318 Furthermore, this technique also allows us to approximate the optimum solution of each
 319 (r, k) -Color (with $r \geq 2$) to a factor of $\frac{1}{2}$. Given a problem (r, k) -Color on G , we can simply
 320 run this method to obtain a partitioning (V_1, V_2) with at least $\frac{|E|}{2}$ edges cut, and assign the
 321 first two relaxed colors to these partitions (not using the remaining $k-2$ colors at all). As
 322 we have $\text{opt}_{(r,k)\text{-Color}}(G) \leq |E|$ in any case, this is already a $\frac{1}{2}$ -approximation algorithm for
 323 the problem, showing that (r, k) -Color can always be approximated to a constant factor.

324 \blacktriangleright **Lemma 5.5.** *For $k \geq r \geq 2$, we have (r, k) -Color \in APX.*

325 In order to show that (r, k) -Color is also complete in APX, we present an L-reduction to
 326 transform a solution of (r, k) -Color into a solution of $(2, 2)$ -Color, i.e. MaxCut. The APX-
 327 completeness of Max- k -Cut has also been shown through a reduction to MaxCut in [16]. We
 328 essentially show that the same graph transformation can be applied in our case as in the
 329 reduction of [16], regardless of the difference between Max- k -Cut and (r, k) -Color, i.e. the
 330 extra restriction that some of the classes must be proper colors. For this, we first describe
 331 the reduction of [16].

332 ► **Lemma 5.6.** *There exists an L-reduction from $(2, 2)$ -Color to (k, k) -Color (based on the*
 333 *transformation idea described in [16]).*

334 **Proof.** We describe the reduction in two steps. We first present a reduction to the (r, k) -Color
 335 problem on multigraphs (i.e. graphs allowing parallel edges). We then discuss how to transform
 336 such a multigraph into a simple graph without parallel edges.

337 Given a MaxCut problem on an input graph G with n nodes and m edges, let us create a
 338 multigraph G'_M as follows. We add a clique C of $k - 2$ further nodes to G , with each edge
 339 between the nodes of C having a multiplicity of $2m$. Furthermore, we connect each node
 340 $v \in G$ to each node $u \in C$, with an edge of multiplicity $2d(v)$. The resulting multigraph
 341 consist of $n + k - 2$ nodes and $|E_{G'_M}| = m + (k - 2) \cdot 4m + \binom{k-2}{2} \cdot 2m$ edges (with multiplicity).
 342

343 Note that this already satisfies the first condition of the L-reduction. As we discussed
 344 earlier, an optimum solution of MaxCut on G will certainly cover at least $\frac{m}{2}$ edges. On the
 344 other hand, G'_M has only $|E_{G'_M}| \in \mathcal{O}(k^2 m)$ edges. Thus, we have

$$345 \quad \frac{1}{2} \cdot \text{opt}_{(r,k)\text{-Color}}(G'_M) \leq \mathcal{O}(k^2) \cdot \text{opt}_{\text{MaxCut}}(G),$$

346 which implies the first condition of the L-reduction for some $\alpha = 2 \cdot \mathcal{O}(k^2)$.

347 We now show that the second condition also holds for $\beta = 1$. Note that if we color the
 348 nodes of G with two relaxed colors in an arbitrary way, and use each of the remaining $k - 2$
 349 (proper or relaxed) colors to color one node of C , then we immediately obtain a coloring
 350 with at most m uncovered edges.

351 Consider a solution \mathcal{S} to the (r, k) -Color problem, that is, a k -coloring of the nodes of
 352 G'_M . If such a coloring was to assign the same color to any two nodes of C , then it would
 353 lose at least $2m$ edges, and thus the right side of the second condition would be at least m ,
 354 implying that the condition holds regardless of our choice of \mathcal{S}' . Thus, it remains to show
 355 the condition for the case when \mathcal{S} colors the nodes of C with $k - 2$ different colors.

356 Note that if any $v \in G$ is assigned one of the $k - 2$ colors used in C , it may be recolored to
 357 one of the remaining 2 colors (not used in C). Thereby, the value of a solution \mathcal{S} will increase
 358 by at least $2d(v)$, while introducing at most $d(v)$ new conflicts. Hence, after recoloring all
 359 such vertices in G , we can assume to obtain a solution \mathcal{S} where C is colored with $k - 2$
 360 different colors, and only the remaining 2 colors are used in G .

361 Given a k -coloring \mathcal{S} of G'_M , we can then simply define the solution \mathcal{S}' of MaxCut as the
 362 restriction of \mathcal{S} to the nodes of G . Since both \mathcal{S} and the optimal coloring covers all edges of
 363 G'_M outside of G , and they both use only 2 colors in G , their difference is in fact determined
 364 by how many edges they cover in G . Hence, we have

$$365 \quad \text{opt}_{(r,k)\text{-Color}}(G'_M) - \text{val}_{(r,k)\text{-Color}}(\mathcal{S}) = \text{opt}_{\text{MaxCut}}(G) - \text{val}_{\text{MaxCut}}(\mathcal{S}'),$$

366 and thus the second condition indeed holds with $\beta = 1$.

367 As a second step, we transform the multigraph G'_M into a simple graph G' with the same
 368 properties. For this, we define an auxiliary graph K on $k + 2$ nodes. Let us denote the nodes
 369 of K by w_1, w_2, \dots, w_{k+2} , and let K be the graph obtained by taking a clique on these $k + 2$
 370 nodes, and then deleting the edges (w_1, w_2) , (w_2, w_3) and (w_3, w_4) from this clique. There
 371 are two crucial properties of the resulting graph K : (i) it is k -colorable without any conflicts,
 372 (ii) every valid k -coloring of K assigns different colors to nodes w_2 and w_3 .

373 We then use these copies of the graph K to replace the parallel edges of G'_M . For each
 374 edge (u, v) in G'_M which is not contained in G , we can replace (u, v) by a separate instance
 375 of K , with the nodes u and v taking the roles of nodes w_2 and w_3 . Since we only insert k
 376 new nodes and $\mathcal{O}(k^2)$ new edges for each original edge (u, v) of G'_M , the number of nodes

377 and edges in the resulting graph G' will still remain polynomial in n . In the graph obtained,
 378 each K will behave like an edge in the following sense: If nodes u and v have different color,
 379 then it is possible to color K without a conflict; but if u and v have the same color, then
 380 there will be at least one conflict in K . Thus, if there were originally t parallel edges between
 381 u and v , then assigning the same color to u and v implies that there will certainly be at least
 382 t conflicts within the t copies of K between the two nodes.

383 For the purpose of the reduction, the obtained graph G' behaves exactly like the multi-
 384 graph G'_M . Since the number of edges in G' is in $\mathcal{O}(k^4m)$, the first condition still holds. In
 385 any solution \mathcal{S} , if a copy of K has more than one conflict, we can always recolor the nodes w_1 ,
 386 w_4, w_5, \dots, w_{k+1} such that there are only 0 or 1 conflicts in the end, depending on whether
 387 w_2 and w_3 have the same color or not. Hence, we can assume that each K represents exactly
 388 as many conflicts as the parallel edge it has replaced. Thus, the reduction described for the
 389 multigraph case also works for G' without modification. ◀

390 The transformation described so far is a valid reduction to (k, k) -Color, i.e. when all the
 391 available colors are relaxed. We now show that it also remains a valid reduction for (r, k) -Color.

392 ▶ **Lemma 5.7.** *There exists an L -reduction from $(2, 2)$ -Color to (r, k) -Color for any $r \geq 2$.*

393 **Proof.** For the first part of the reduction (to the multigraph G'_M), we can argue as before
 394 that all nodes of C obtain a different color in \mathcal{S} . We then use the same technique to ensure
 395 that the nodes of G will only get the remaining two colors. This technique still increases
 396 the number of covered edges in each step, but might lead to an invalid coloring in the end if
 397 one (or both) of the two colors used in G are among the proper colors. However, if one of
 398 the proper colors used in G has a conflict, then we can simply swap this proper color with a
 399 relaxed color that is currently used only on a single node in C . After at most 2 such swaps,
 400 we can make sure that both colors used to color G are relaxed colors. We can consider the
 401 resulting coloring as the original solution \mathcal{S} , and apply the same reduction as before.

402 As for the transformation to a simple graph, whenever two neighboring nodes u and v
 403 have different colors, it is always possible to color the whole copy of graph K between them
 404 without any conflicts at all, so it does not matter whether the colors used on the specific
 405 nodes of K are proper or relaxed colors. If u and v have the same color c_1 , then there is at
 406 least one relaxed color $c_2 \neq c_1$. We can then color the copy of K by using c_2 on nodes w_1
 407 and w_4 , and each of the remaining $k - 2$ colors on one of the remaining $k - 2$ nodes; in this
 408 coloring, the only conflict will be on the edge (w_1, w_4) . Thus it is still always possible to
 409 color each copy of K with exactly 0 or 1 conflicts (depending on whether u and v have the
 410 same color), regardless of the colors of u and v being proper or relaxed. ◀

411 Altogether, this shows the following.

412 ▶ **Theorem 5.8.** *For any $k \geq r \geq 2$, the problem (r, k) -Color is APX-complete.*

413 **6 Conclusion – and a Generalization**

414 In the paper, we analyzed the problem of Two-Class (r, k) -Coloring, which aims to maximize
 415 the number of covered edges when coloring a graph with r relaxed and $k - r$ proper colors.
 416 We have seen that $(1, 2)$ -Color is similar but not identical to the Maximum Independent Set
 417 problem. We have shown that the problem (r, k) -Color is NP-complete for any $k \geq 2$ (except
 418 when $(r, k) = (0, 2)$). Furthermore, we also proved that the problems (r, k) -Color with $r \geq 2$
 419 are APX-complete, and that $(1, k)$ -Color cannot be approximated to any constant factor in
 420 polynomial time.

421 Throughout this paper, we have established the close relation between the (r, k) -Color
 422 and the Max- k -Cut problem. Note that the cost function used for the optimization is the
 423 same for both problems, while they only differ in their set of feasible solutions.

424 In practice, however, one may think of other reasonable cost functions. For instance,
 425 Defective Coloring exclusively focuses on the node with the highest number of conflicts, and
 426 does not evaluate the coloring on the rest of the graph.

427 Considering application areas, e.g. modeling interference in frequency allocation problems,
 428 both approaches might be reasonable. A provider would generally want to keep the total
 429 number of conflicts in the graph small, as this corresponds to the overall service quality. On
 430 the other hand, a provider would probably also try to minimize the maximal number of
 431 conflicts at a single node in order to avoid upsetting any particular customer.

432 In fact, an interesting generalization allows us to model both settings (and the trade-off
 433 between them) at the same time. Given a graph $G = (V, E)$ and a k -coloring of G , let us
 434 denote the number of conflicts at node $v \in V$ (i.e., the number of conflict edges incident to v)
 435 by $\kappa(v)$. Then, given a real parameter $p > 0$, we define the *Generalized Conflict Coloring*
 436 problem, where the cost of a coloring \mathcal{S} is defined as

$$437 \quad \text{cost}(\mathcal{S}) \quad := \quad \sum_{v \in V} \kappa(v)^p.$$

438 The goal of the Generalized Conflict Coloring problem is to find a coloring \mathcal{S} on k colors that
 439 minimizes this cost. This generalized cost function can be combined with any restrictions on
 440 the set of feasible colorings, for example, the family of (r, k) -Color problems.

441 In the following, we motivate that this problem is a natural generalization of various
 442 known coloring problems. In the case when $p = 1$, the aim of the problem is to minimize
 443 $\sum_{v \in V} \kappa(v)$, which is simply two times the number of monochromatic edges (conflicts) in the
 444 graph. Hence the cost of a coloring \mathcal{S} is exactly $2 \cdot (|E| - \text{val}_{(k,k)\text{-Color}}(\mathcal{S}))$, and thus the
 445 problem is identical to Two-Class (k, k) -Coloring (i.e., to Max- k -Cut).

446 On the other hand, as p goes to infinity, the sum in the cost will be dominated only by
 447 the highest $\kappa(v)$ value in the graph, and thus a coloring \mathcal{S}_1 will be better than a coloring \mathcal{S}_2
 448 exactly if the maximal $\kappa(v)$ is smaller in \mathcal{S}_1 . This is precisely what is minimized in Defective
 449 Coloring; hence, Defective Coloring is also obtained as a subcase of Generalized Conflict
 450 Coloring in the limit for $p \rightarrow \infty$.

451 Finally, if $p \rightarrow 0$, the difference between the distinct $\kappa(v)$ values will diminish for all
 452 $\kappa(v) > 0$, since all are approaching 1 for p small enough. Thus, in the limit we obtain

$$453 \quad \lim_{p \rightarrow 0} \text{cost}(\mathcal{S}) \quad = \quad \sum_{\substack{v \in V \\ \kappa(v) > 0}} 1.$$

454 In other words, the cost of a coloring \mathcal{S} becomes the number of nodes for which $\kappa(v) > 0$.
 455 Minimizing this corresponds to finding a coloring which leaves the highest possible number
 456 of nodes without a conflict.

457 Furthermore, intermediate values of p allow us to study the trade-off between these
 458 problems. For instance, if $p = 2$ is selected, then we obtain a setting between Max- k -Cut and
 459 Defective Coloring, where we have to optimize both for a small number of total conflicts and
 460 a reasonably balanced distribution of these conflicts among nodes. Altogether, we conclude
 461 that the further study of this Generalized Conflict Coloring problem is not only interesting
 462 from a theoretical point of view, but also motivated by practical applications.

References

- 463 —
- 464 1 Dan Archdeacon. A note on defective colorings of graphs in surfaces. *Journal of Graph Theory*,
465 11(4):517–519, 1987.
- 466 2 Leonid Barenboim and Michael Elkin. Distributed graph coloring: Fundamentals and recent
467 developments. *Synthesis Lectures on Distributed Computing Theory*, 4(1):1–171, 2013.
- 468 3 Lenore Cowen, Wayne Goddard, and C Esther Jesurum. Defective coloring revisited. *Journal*
469 *of Graph Theory*, 24(3):205–219, 1997.
- 470 4 Lenore J Cowen, Robert H Cowen, and Douglas R Woodall. Defective colorings of graphs in
471 surfaces: partitions into subgraphs of bounded valency. *Journal of Graph Theory*, 10(2):187–195,
472 1986.
- 473 5 LJ Cowen, W Goddard, and CE Jesurum. Coloring with defect. In *Proceedings of the eighth*
474 *annual ACM-SIAM symposium on Discrete algorithms*, pages 548–557. Society for Industrial
475 and Applied Mathematics, 1997.
- 476 6 Pierluigi Crescenzi. A short guide to approximation preserving reductions. In *Proceedings of*
477 *Computational Complexity. Twelfth Annual IEEE Conference*, pages 262–273. IEEE, 1997.
- 478 7 Etienne de Klerk, Dmitrii V Pasechnik, and Joost P Warners. On approximate graph colouring
479 and max-k-cut algorithms based on the θ -function. *Journal of Combinatorial Optimization*,
480 8(3):267–294, 2004.
- 481 8 Nancy Eaton and Thomas Hull. Defective list colorings of planar graphs. *Bull. Inst. Combin.*
482 *Appl*, 25(79-87):40, 1999.
- 483 9 Alan Frieze and Mark Jerrum. Improved approximation algorithms for maxk-cut and max
484 bisection. *Algorithmica*, 18(1):67–81, 1997.
- 485 10 Philippe Galinier and Alain Hertz. A survey of local search methods for graph coloring.
486 *Computers & Operations Research*, 33(9):2547–2562, 2006.
- 487 11 Michael R Garey and David S Johnson. *Computers and intractability : a guide to the theory*
488 *of NP-completeness*. A series of books in the mathematical sciences. Freeman, New York, 24th
489 pr. edition, 2003.
- 490 12 Tommy R Jensen and Bjarne Toft. *Graph coloring problems*, volume 39. John Wiley & Sons,
491 2011.
- 492 13 Viggo Kann. *On the approximability of NP-complete optimization problems*. PhD thesis, Royal
493 Institute of Technology Stockholm, 1992.
- 494 14 Viggo Kann, Sanjeev Khanna, Jens Lagergren, and Alessandro Panconesi. On the hardness of
495 approximating max k-cut and its dual. In *ISTCS*, pages 61–67, 1996.
- 496 15 Michael Mitzenmacher and Eli Upfal. *Probability and computing: Randomization and proba-*
497 *bilistic techniques in algorithms and data analysis*. Cambridge university press, 2017.
- 498 16 Christos H Papadimitriou and Mihalis Yannakakis. Optimization, approximation, and com-
499 plexity classes. *Journal of computer and system sciences*, 43(3):425–440, 1991.
- 500 17 Panos M Pardalos, Thelma Mavridou, and Jue Xue. The graph coloring problem: A bib-
501 liographic survey. In *Handbook of combinatorial optimization*, pages 1077–1141. Springer,
502 1998.
- 503 18 Mihalis Yannakakis. Node-and edge-deletion np-complete problems. In *Proceedings of the*
504 *tenth annual ACM symposium on Theory of computing*, pages 253–264. ACM, 1978.
- 505 19 Mihalis Yannakakis and Fanica Gavril. The maximum k-colorable subgraph problem for
506 chordal graphs. *Information Processing Letters*, 24(2):133–137, 1987.