Reverse mathematics of rings

by

Jordan Mitchell Barrett

A thesis submitted to Victoria University of Wellington in fulfilment of the requirements for the degree of Master of Science in Mathematics.

Victoria University of Wellington 2021

Abstract

Using the tools of reverse mathematics in second-order arithmetic, as developed by Friedman, Simpson, and others, we determine the axioms necessary to develop various topics in commutative ring theory. Our main contributions to the field are as follows. We look at fundamental results concerning primary ideals and the radical of an ideal, concepts previously unstudied in reverse mathematics. Then we turn to a fine-grained analysis of four different definitions of Noetherian in the weak base system $RCA_0 + I\Sigma_2$. Finally, we begin a systematic study of various types of integral domains: PIDs, UFDs and Bézout and GCD domains.

Acknowledgments

The document you are now reading would not have been possible without the support of numerous people. First and foremost, I'd like to thank my supervisor Dan Turetsky, who provided an invaluable source of support, ideas and knowledge throughout my Masters. I appreciate the generous scholarship from Victoria University of Wellington which made my studies possible. Thanks also to Long Qian and Valentino Vito for friendship, laughs and thoughtful discussions.

I am grateful to Rod, Noam, and especially Martino, for their support over the years. Thanks to the other members of the logic group (Thomas, Diamant, Linus, Andre) and the proof assistants group (Marco, Julian) for keeping me sane.

Thanks to everyone else in the School of Mathematics and Statistics that made it a lovely place to learn and grow. There are too many to name, but I'd particularly like to mention Joseph, Liam, Amber, Sahas, Steve, Astrid, Lisa, Evelyn and Matthew. A special mention to Caitlin, Alec and the other admin staff who keep things running so smoothly.

Thanks to all my friends, especially Kaspar, my bandmates in Solid Walls of Sound, and the wonderful peeps in Vic Uke. Finally, thanks are owed to my family for their endless love and support throughout. Mum, Jacob, Lottie & Maddie, Dad & Mandy, Ani—I love you all.

Contents

1 Introduction				1	
	1.1	Notational convention	ns	6	
2	Rin	g theory		8	
3	Log	ical prerequisites		13	
	3.1	Second-order arithmet	tic	13	
	3.2	Reverse mathematics		18	
	3.3	Computability		21	
4	Rad	icals of ideals and prin	nary ideals	29	
	4.1	The RAD principle		33	
5	Noe	therian rings	n rings 39		
	5.1	The a.c.c.p		50	
6	Inte	gral domains		51	
	6.1	Bézout and GCD dom	nains	55	
	6.2 UFDs and PIDs			60	
		6.2.1 UFDs		60	
		6.2.2 PIDs		62	
			initions of PIDs	63	
		-	ut PIDs	70	
		6.2.5 Gauss' lemma		74	

CONTENTS		iii
7	Conclusion	78
A	Linear algebra	81
В	Zorn's lemma	85
Bibliography		90

Chapter 1

Introduction

This thesis concerns the intersection of two distinct areas of mathematics: commutative ring theory, and mathematical logic. The first, commutative ring theory, also known as *commutative algebra*, has its roots in classical algebraic number theory and algebraic geometry [Kle98].

19th century number theory was concerned with problems such as solvability of Diophantine equations, or of polynomial congruences, over the integers \mathbb{Z} . A common technique that emerged was to extend \mathbb{Z} by an *algebraic integer*, obtaining new domains such as $\mathbb{Z}[\sqrt{2}i]$, $\mathbb{Z}[i]$, or $\mathbb{Z}[\omega]$ for ω a primitive root of unity. If these new domains had *unique factorisation*, then one could draw conclusions about the original equations or congruences.

Unfortunately, these domains failed to have unique factorisation in many important cases. Kummer's idea was to further add "ideal primes" to the domain to restore unique factorisation [Kum47a; Kum47b]. While brilliant, Kummer's ideas were vague, and Dedekind later put Kummer's work on a rigorous footing by giving the modern definition of an *ideal* in a ring [Ded71]. Two particular features of Dedekind's work—a focus on axiomatic methods, as well as an acceptance of nonconstructive procedures—marked a new style of mathematics which would come to dominate the 20th century, ultimately paving the way for mathematical logic.

On the other hand, algebraic geometry is concerned with algebraic varieties. Given a fixed set of polynomials $\mathcal{P} \subseteq \mathbb{R}[x_1,\ldots,x_n]$, the variety $V_{\mathcal{P}}$ is the set of points in \mathbb{R}^n satisfying the equations $p(x_1,\ldots,x_n)=0$ for all $p\in\mathcal{P}$. These are the higher-dimensional generalisation of algebraic curves (which are the case n=2). Now, to any variety V, we can assign the set I(V) of polynomials which vanish on V; this is an ideal in $\mathbb{R}[x_1,\ldots,x_n]$. Hence, we can study algebraic varieties by studying ideals in polynomial rings. This correspondence was exploited to great effect by Hilbert in his basis theorem [Hil90] and Nullstellensatz [Hil93], and later by Lasker [Las05] and Macauley [Mac13].

Attempting to create a general theory encompassing all these ideas, Fraenkel gave the first abstract definition of a ring in [Fra15], and Sono gave the modern definition soon after [Son17]. This opened the door to the pioneering work of Noether, which established abstract ring theory as a subject. Specifically, Noether [Noe21] generalised the results of Hilbert, Lasker and Macauley to what are now called *Noetherian rings*, and later recast the work of Dedekind et al in an abstract setting [Noe27].

Meanwhile, a mathematical revolution had been brewing. Traditionally, mathematics had been concerned with finite objects and constructive procedures, and grounded in reality [Eve69; MN82]. The work of Dedekind, Hilbert, Peano, Cantor and others in the late 1800s marked a departure from this, thereby ushering in modern, abstract pure mathematics. This new style of mathematics was distinguished by its focus on abstraction and the axiomatic method, and acceptance of nonconstructive proofs—those which prove the existence of an object without actually constructing an example.

Another feature of this new mathematics was the acceptance of *completed infinity* rather than just *potential infinity*—the idea that infinite sets could be manipulated as mathematical objects in their own right. This idea perhaps appeared first in Dedekind's work on ideals [Ded71; Kle98]. Cantor was the first to systematically study infinity, founding the field of set

theory with his seminal work on cardinals [Can74] and ordinals [Can83]. As set theory developed, paradoxes arose (most notably Russell's), and the need for a careful and rigorous foundation for mathematics became clear. One such foundation was provided by ZFC in the 1920s [Zer30].

Cantor's work provided new impetus to mathematical logic, a small subfield of mathematics developed by Boole, De Morgan, and Peano in the mid-to-late 1800s [Boo54; DeM47; Pea89]. Around this time, the ideas of computation, mathematical truth and mathematical proof were formalised for the first time. By the 1930s, logic was a thriving area of mathematics—highlights included Gödel's (in)completeness theorems [Göd29; Göd31], Turing's negative solution to the *Entscheidungsproblem* [Tur37], Tarski's development of model theory [Vau86], and Hilbert's work on proof theory [HB34] and geometry [Hil99].

A later development in logic was reverse mathematics, initiated by Harvey Friedman in the late 1960s [Fri67; Fri69]. Reverse mathematics asks, for a given theorem of mathematics φ , "what axioms are really necessary to prove φ ?" More broadly, it studies the logical implications between foundational principles of mathematics. An early example was the discovery of non-Euclidean geometries, thereby proving the independence of the parallel postulate from Euclid's other axioms [Lob29; Bol32]. Another early result, more in the style of reverse mathematics, was the demonstration that over ZF, the axiom of choice, Zorn's lemma, and the well-ordering principle are all pairwise equivalent [Bir40; FB58; Tra62].

Traditionally, reverse mathematics is done in second-order arithmetic, in which there are two types of objects: natural numbers n, m, k, \ldots , and sets of natural numbers A, B, C, \ldots , and quantification is allowed over both types of objects. Restricting oneself to natural numbers may seem unnecessary limiting, but this is not so. In fact, most mathematics deals with countable or "essentially countable" objects (such as separable metric spaces), and so can be formalised in second-order arithmetic. This includes virtually all "classical" mathematics, or that taught in undergradu-

ate courses [Sim09, p. xiv].

In practice, reverse mathematics involves attempting to prove a theorem φ of "ordinary" mathematics in a weak subsystem \mathcal{S} of second-order arithmetic. But, supposing we can do this, how do we know we've found the optimal (weakest) system? The empirical phenomenon is thus:

"When the theorem is proved from the right axioms, the axioms can be proved from the theorem."

—Harvey Friedman [Fri74]

This is the "reverse" part of reverse mathematics. Having proved φ from \mathcal{S} , to show this is optimal, we want to demonstrate a *reversal* of φ : a proof of \mathcal{S} from φ . This means that φ cannot be proved in a weaker system \mathcal{S}' , because if it could, then \mathcal{S}' would also prove \mathcal{S} via φ , meaning \mathcal{S}' is not actually a weaker system after all.

The utility of reverse mathematics is abundant. Apart from its obvious use in finding the "best" proof of a given statement φ , it also gives us a way to quantify how nonconstructive or noncomputable φ is. The idea is that stronger subsystems correspond to more nonconstructive power, so the "constructiveness" of φ is inversely proportional to the strength of the systems $\mathcal S$ in which φ can be proved [FSS83]. Similarly, many theorems guarantee a solution to a given problem—reverse mathematics then tells us how complex the solution could be relative to the problem, which can be made precise in terms of computability.

Here is an example of reverse mathematics in ring theory. The usual way to prove that every commutative ring has a prime ideal is to prove that it has a maximal ideal (Krull's theorem), and then prove every maximal ideal is prime. However, Friedman, Simpson and Smith showed that the existence of maximal ideals is equivalent to the system ACA₀, whereas the existence of prime ideals is equivalent to the strictly weaker system WKL₀ [FSS83]. This shows the usual proof strategy is not optimal—there

is a "better" way to prove the existence of prime ideals, which doesn't require the stronger assumption that maximal ideals exist. In terms of computability, this shows that maximal ideals can be more "noncomputable" than prime ideals—more precisely, given a computable ring, its maximal ideals could all be as complex as the *halting problem*, while we can always compute a prime ideal from a *PA degree*.

In this thesis, we study the reverse-mathematical content of various theorems of ring theory. We will begin by reviewing basic ideas from ring theory (§2), and from logic, computability and reverse mathematics (§3). We then proceed to study the following key ideas from commutative algebra:

- *Primary ideals* (§4), which are the ideals I such that whenever $ab \in I$, then $a \in I$ or $b^n \in I$ for some n.
- The *radical* \sqrt{I} of an ideal I (§4), which is the set of r such that some r^n is in I.
- *Noetherian rings* (§5), which have many equivalent definitions; one of the more popular is that every ideal is finitely generated.
- Several classes of *integral domains* (§6), including PIDs, UFDs, Bézout and GCD domains, their properties, and the relations between them.

In relation to the philosophy of reverse mathematics, we find that all the results we examine are provable in ACA₀.¹ We will show that many important results, such as the equivalence of different notions of Noetherian, actually *require* ACA₀. Thus, we conclude that ACA₀ is the right axiom system in which to develop (most) of classical commutative algebra. This thesis includes many new, original results—some of the more important ones are:

¹This seems to be true for algebra in general, with the notable exception of some structure theorems and (ordinal) invariant results [Sim09].

- Theorem 4.10: WKL₀ is equivalent to " \sqrt{I} maximal $\implies I$ primary".
- Theorem 5.4: $RCA_0 + I\Sigma_2$ proves the equivalence of weak and strict chain conditions on Σ_1 -ideals.
- Theorems 5.10 and 5.11, showing that WKL₀ or ACA₀ are equivalent to the agreement of several definitions of Noetherian.
- Theorem 6.26: ACA₀ proves that if R is an integral domain in which every prime Σ_1 -ideal is principal, then R is a Σ_1 -PID.
- Theorem 6.30 and Corollary 6.35: ACA₀ is equivalent to "every PID admits a Dedekind–Hasse norm".
- Theorem 6.39, where we construct a PID whose set of primes is Π_2 complete.

1.1 Notational conventions

The following notational conventions will apply to this thesis:

- We will use \mathbb{N} to denote (the underlying set of) the model of arithmetic we are working inside, and ω to denote the standard model $\{0, 1, 2, \ldots\}$. Generally, the distinction will not be important.
- For a ring R, we will use $R[\bar{x}]$ as an abbreviation for $R[x_0, x_1, \ldots]$, the polynomial ring over R in infinitely many indeterminates.
- In mathematics, pairs, tuples and sequences are commonly denoted using parentheses, e.g. (x,y), (a_0,a_1,\ldots) . However, in ring theory, it is also common to use parentheses to denote the *ideal* (A) generated by a collection of elements $A\subseteq R$. To avoid confusion, we will try to consistently use angle brackets $\langle x,y\rangle$, $\langle a_0,a_1,\ldots\rangle$ to denote a pair, tuple or sequence, and reserve parentheses for ideals.

7

- ε or $\langle \rangle$ will denote the empty sequence or tuple.
- The SANS SERIF font will generally be reserved for subsystems and axioms of second-order arithmetic.
- We will use $\varphi_0, \varphi_1, \ldots, \varphi_e, \ldots$ to denote a standard listing of the partial computable functions, and $W_0, W_1, \ldots, W_e, \ldots$ to denote a listing of the c.e. sets.
- We may use $A^{\complement} = \{x : x \notin A\}$ to denote the (absolute) complement of a set A, particularly for sets of natural numbers.
- For mathematical statements φ and ψ , we use $\varphi \vdash \psi$ (" φ proves ψ ") to mean there is a proof of ψ from φ . This notation extends to formal systems, e.g. $\mathcal{S} \vdash \varphi$ means there is a proof of φ in the formal system \mathcal{S} .
- For a statement φ and a structure \mathcal{M} , we use $\mathcal{M} \models \varphi$ (" \mathcal{M} models φ ") to mean the statement φ is true in \mathcal{M} . Similarly, $\mathcal{M} \models \mathcal{S}$ means that all axioms of the formal system \mathcal{S} are true in \mathcal{M} .

Chapter 2

Ring theory

Here, we quickly review the basic notions of ring theory that we will need, as covered in any basic algebra textbook [AM94]. For us, "ring" will mean "commutative ring with unity", unless explicitly stated otherwise.

Definition 2.1. A *ring* is a set R, equipped with constants $0_R, 1_R \in R$, and binary operations $+, \cdot$ on R (called *addition* and *multiplication*, respectively), such that:

- (i) R is an abelian group under addition, with additive identity 0_R .
- (ii) Multiplication is associative: $(a \cdot b) \cdot c = a \cdot (b \cdot c)$ for all $a, b, c \in R$.
- (iii) Multiplication is commutative: $a \cdot b = b \cdot a$ for all $a, b \in R$.
- (iv) 1_R is a two-sided multiplicative identity.
- (v) Multiplication distributes both ways over addition: $a \cdot (b+c) = (a \cdot b) + (a \cdot c)$ and $(a+b) \cdot c = (a \cdot c) + (b \cdot c)$ for all $a,b,c \in R$.

As usual, we will often omit the dot for multiplication, and instead denote it by juxtaposition, i.e. ab instead of $a \cdot b$.

Given a ring R, we can construct a larger ring R[x] of "polynomials over R" in the variable x. We do this by "freely" adding the variable x, i.e.

9

asserting no relationship between x and elements of R. A formal construction follows.

Definition 2.2. For a ring R, the polynomial ring R[x] is defined as follows:

• The underlying set of R[x] is the collection

$$\{\langle a_0, \dots, a_n \rangle \in R^{<\omega} : a_n \neq 0_R\} \cup \{\langle \rangle \}$$

We denote $\langle a_0, \ldots, a_n \rangle$ by $a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0$.

- $0_{R[x]} = 0_R = \langle \rangle$ and $1_{R[x]} = 1_R = \langle 1_R \rangle$.
- Addition and multiplication in R[x] are defined as follows:

$$\left(\sum_{i=0}^{n} a_i x^i\right) + \left(\sum_{i=0}^{m} b_i x^i\right) = \sum_{i=0}^{\max\{n,m\}} (a_i + b_i) x^i$$

$$\left(\sum_{i=0}^{n} a_{i} x^{i}\right) \cdot \left(\sum_{i=0}^{m} b_{i} x^{i}\right) = \sum_{i=0}^{m+n} \left[\sum_{j=0}^{i} a_{j} b_{i-j}\right] x^{i}$$

Given a polynomial ring R[x], we could repeat the construction to get (R[x])[y], which we write simply as R[x,y] for brevity. Iterating this construction, we get an increasing sequence $R[x_0], R[x_0, x_1], R[x_0, x_1, x_2], \ldots$ We will use $R[\bar{x}] = R[x_0, x_1, \ldots]$ to refer to the limit of this sequence.

Definition 2.3. In a ring R, an *ideal* $I \subseteq R$ is a subset of R such that for all $a, b \in I$ and $r, s \in R$, we have $ar + bs \in I$.

Ideals are important since they give us a way to create new rings:

Definition 2.4. Given a ring R and ideal $I \subseteq R$, the *quotient ring* R/I is defined as follows:

- The underlying set is the quotient of R by the equivalence relation $r \sim s \iff r s \in I$. We denote the equivalence class of r by r + I.
- $0_{R/I} = 0_R + I$ and $1_{R/I} = 1_R + I$.

• (a+I)+(b+I)=(a+b)+I and (a+I)(b+I)=(ab)+I. These operations are well-defined.

We recall some important ways of creating ideals.

Definition 2.5. Let R be a ring and $A \subseteq R$ an arbitrary subset. The *ideal* generated by A is the set

$$(A) := \{a_1r_1 + \dots + a_nr_n : a_1, \dots, a_n \in A, r_1, \dots, r_n \in R\}$$

If $A = \{a_1, \dots, a_n\}$ is a finite set, we write (a_1, \dots, a_n) and say this ideal is finitely generated. If $A = \{a\}$, we say (a) is principal.

Definition 2.6. Given two ideals $I, J \subseteq R$, the *ideal quotient of* I *by* J is the set

$$I: J := \{r \in R : (\forall j \in J)(rj \in I)\}\$$

The most common case of Definition 2.6 is when J=(a) is a principal ideal. In this case, we will abuse notation and write I:a instead of I:(a). I:a also admits a simpler definition here, as $I:a=\{r\in R:ra\in I\}$.

We now recall an important subclass of the commutative rings.

Definition 2.7. An element $a \in R$ is a *zero-divisor* if there is $b \neq 0_R$ such that ab = 0.

Definition 2.8. A ring R is an *integral domain* if it has no nonzero zero-divisors, i.e. whenever ab = 0, then a = 0 or b = 0.

Integral domains satisfy cancellation of multiplication: if $a \neq 0$ and ab = ac, then b = c. Indeed, this is an alternative characterisation of integral domains.

Definition 2.9. Given an integral domain R, its *field of fractions* Frac(R) is the ring defined as follows:

• The underlying set of Frac(R) is the quotient of $\{(r,s) \in R^2 : s \neq 0_R\}$ by the equivalence relation $(r,s) \sim (r',s') \iff rs' = r's$. We denote the equivalence class of (r,s) by r/s.

- $0_{\operatorname{Frac}(R)} := 0_R/1_R$ and $1_{\operatorname{Frac}(R)} := 1_R/1_R$.
- (r/s) + (r'/s') = (rs' + r's)/(ss') and (r/s)(r'/s') = (rr')/(ss'). These operations are well-defined.

We can verify that $\operatorname{Frac}(R)$ is indeed a ring, and in fact, it is a field—every nonzero element has an inverse. We can naturally view R as a subring of $\operatorname{Frac}(R)$ via the embedding $r \mapsto r/1$.

There is an important generalisation of Definition 2.9, which covers both the case when R is not an integral domain, and when we don't want every element of R to be a denominator.

Definition 2.10. Suppose $M \subseteq R$ is multiplicatively closed, contains 1_R , and contains no zero-divisors. The *localisation of* R *at* M, Loc (R, M), is the ring whose underlying set is the quotient of $\{(r, m) \in R^2 : m \neq 0_R\}$ by the equivalence relation $(r, m) \sim (r', m') \iff rm' = r'm$, and where the operations are defined as in Definition 2.9.

Intuitively, Loc (R, M) is obtained from R by allowing division by the elements of M. As before, R is naturally a subring of Loc (R, M) via the embedding $r \mapsto r/1$.

Definition 2.11.

- (i) A multiplicatively closed set M is saturated if whenever $ab \in M$, then both $a, b \in M$.
- (ii) $\overline{M} := \{r \in R : (\exists s \in R) (rs \in M)\}$ is the smallest saturated set containing M, and is called the *saturation* of M.

The concept of saturation is important because:

Theorem 2.12. Given multiplicatively closed sets $M, N \subseteq R$, we have that $\operatorname{Loc}(R, M) \cong \operatorname{Loc}(R, N)$ canonically iff $\overline{M} = \overline{N}$. In particular, it is always true that $\operatorname{Loc}(R, M) \cong \operatorname{Loc}(R, \overline{M})$.

Essentially, when we localise at M, we are really localising at its saturation \overline{M} . To see why, take $r \in \overline{M}$, i.e. there is $s \in R$ with $rs \in M$. Then, s/rs is an element of Loc (R,M). But $r \cdot (s/rs) = 1$, so we have introduced an inverse for r, even if $r \notin M$. If we check that M is saturated, then we know that we've *only* added inverses for elements of M, and nothing else.

An important case of localisation is the so-called "localisation at a prime ideal", which is really the localisation at the complement of a prime ideal.

Definition 2.13. An ideal $I \subseteq R$ is *prime* if whenever $ab \in I$, then $a \in I$ or $b \in I$.

Example 2.14. Suppose $P \subseteq R$ is a prime ideal. Then, $M := R \setminus P$ is multiplicatively closed, so we can take the localisation Loc(R, M). This is sometimes called the *localisation of* R at P and denoted R_P .

Another important example of localisation is the total quotient ring, which generalises the field of fractions construction to non-integral domains.

Definition 2.15. For any ring R, the set M of all non-zero-divisors of R is multiplicatively closed and contains 1_R . The *total quotient ring* Frac(R) is the localisation Loc(R, M).

When R is an integral domain, 0_R is the only zero-divisor, so Definition 2.15 reduces to Definition 2.9.

Chapter 3

Logical prerequisites

Here, we review the necessary background material from model theory [Mar02], first- and second-order arithmetic [HP17; Sim09], reverse mathematics [Sim09], and computability [Soa87; Soa16].

3.1 Second-order arithmetic

For us, *arithmetic* will refer to the model-theoretic study of the theory of the natural numbers, $\operatorname{Th}(\mathbb{N})$. Our reverse-mathematical studies will be done in the traditional setting of *second-order arithmetic*. In first-order arithmetic, we are only allowed to quantify over *elements* of \mathbb{N} , while in second-order arithmetic, we may also quantify over *subsets* of \mathbb{N} . This greatly increases the expressive power of our logic (for example, we can define well-foundedness, or completeness of \mathbb{R}).

We work in a two-sorted model theory, whose sorts are *numbers*, denoted with lowercase letters n, m, k, \ldots , and *sets*, denoted in uppercase A, B, C, \ldots Our language is $\mathcal{L}_2 := \{0, 1, +, \cdot, <, \in\}$, where the symbols have the expected types, e.g. 0 is a number, + takes two numbers and returns another, \in is a binary relation between a number and a set, etc.

Numerical \mathcal{L}_2 -terms are defined inductively: 0, 1, and variable symbols x_i are numerical \mathcal{L}_2 -terms, and if s, t are numerical \mathcal{L}_2 -terms, then (s + t),

14

 $(s \cdot t)$ are too. Numerical \mathcal{L}_2 -terms represent (possibly nonstandard) natural numbers, and we will use k to abbreviate the numerical term

$$k := \underbrace{1 + 1 + \dots + 1}_{k \text{ times}}$$

The only set \mathcal{L}_2 -terms are variables X_i . We also define \mathcal{L}_2 -formulae inductively:

Definition 3.1. The collection of \mathcal{L}_2 -formulae is defined as follows:

- (i) If s, t are numerical terms, and X is a set variable symbol, then (s = t), (s < t) and $(s \in X)$ are formulae.
- (ii) If φ , ψ are formulae, then $(\neg \varphi)$, $(\varphi \land \psi)$, $(\varphi \lor \psi)$, $(\varphi \to \psi)$ and $(\varphi \leftrightarrow \psi)$ are formulae.
- (iii) If φ is a formula, then $(\forall x) \varphi$ and $(\exists x) \varphi$ are formulae.
- (iv) If φ is a formula, then $(\forall X) \varphi$ and $(\exists X) \varphi$ are formulae.

A theory S is simply a set of formulae. We may also use the terms *sub-system* or *formal system*, especially when we are considering the members of S as axioms.

There are a wide variety of \mathcal{L}_2 -formulae, which we will now classify based on their "complexity". Our chosen measure of complexity will be based on how many times the quantifiers alternate. This defines a structure known as the *arithmetical hierarchy*.

The lowest level of complexity consists of formulae containing only bounded quantifiers: those of the form $(\forall x)(x < k \to \psi)$ or $(\exists x)(x < k \to \psi)$ for some numerical term. We will often abbreviate these to $(\forall x < k) \psi$ and $(\exists x < k) \psi$ respectively. From there, universal formulae are given Π classifications, and existential formulae given Σ classifications.

Definition 3.2 (arithmetical hierarchy). Let φ be an \mathcal{L}_2 -formula. We say φ is Δ_0^1 or *arithmetical* if it contains no set quantifiers. If φ is arithmetical, we assign further classifications to it as follows:

Figure 3.1: The arithmetical hierarchy.

- (i) φ is called Σ^0_0 and Π^0_0 if it only contains bounded quantifiers.
- (ii) φ is called Σ_{n+1}^0 if it is of the form $\varphi = \exists x_1 \cdots \exists x_n \ \psi$, where ψ is Π_n^0 .
- (iii) φ is called Π_{n+1}^0 if it is of the form $\varphi = \forall x_1 \cdots \forall x_n \psi$, where ψ is Σ_n^0 .
- (iv) φ is called Δ_n^0 if it is Σ_n^0 , and logically equivalent to a Π_n^0 formula.

Often, we will drop the superscript 0, and just speak of Σ_n or Π_n formulae. Now we look at some important principles of arithmetic, which will be pertinent in our study of reverse mathematics.

Definition 3.3. Let $\varphi(x)$, $\psi(x,y)$ be formulae of arithmetic.

(i) The *induction principle for* φ , $|\varphi$, is the statement

$$\varphi(0) \wedge (\forall n) [\varphi(n) \to \varphi(n+1)] \to (\forall n) \varphi(n)$$

(ii) The *strong induction principle for* φ , $\mathsf{I}'\varphi$, is the statement

$$(\forall n) ([(\forall m < n) \varphi(m)] \rightarrow \varphi(n)) \rightarrow (\forall n) \varphi(n)$$

(iii) The *least number principle for* φ , $L\varphi$, is the statement

$$(\exists n) \varphi(n) \to (\exists n') (\varphi(n') \wedge (\forall m < n') \neg \varphi(m))$$

(iv) The bounding principle for ψ , B ψ , is the statement

$$(\forall k) [(\forall n < k)(\exists m) \psi(n, m) \to (\exists \ell)(\forall n < k)(\exists m < \ell) \psi(n, m)]$$

All of the above principles trivially hold in the standard model of arithmetic ω . However, recall that we have *nonstandard* models of arithmetic, which satisfy the same basic axioms as ω . These nonstandard models are linear orders of type $\omega + \mathbb{Z} \cdot K$ for some linear order K [Hen50]. In one of these models, the above principles could fail. For example, L φ fails if φ is true exactly on the $\mathbb{Z} \cdot K$ part, while B ψ could fail if the bound k is nonstandard.

For a *class* of formulae Γ , we define $|\Gamma|$ to be the theory consisting of all the statements $|\varphi|$ for all (appropriate) $\varphi \in \Gamma$. The theories $|\Gamma|$, $|\Gamma|$, $|\Gamma|$ are defined analogously. We will usually take $|\Gamma|$ to be a classification in the arithmetical hierarchy: for example, $|\Sigma|_2$ is induction for all $|\Sigma|_2$ formulae.

These arithmetical principles are closely related to each other:

Theorem 3.4. For every formula of arithmetic $\varphi(x)$, $I'\varphi \equiv L(\neg \varphi)$.

Theorem 3.5 [PK78; HP17]. Over $PA^- + I\Sigma_0$, for all $n \in \mathbb{N}$:

- (i) $|\Sigma_n, \Pi_n, |'\Sigma_n, |'\Pi_n, L\Sigma_n, L\Pi_n$ are all equivalent.
- (ii) $\mathsf{L}\Delta_{n+1}$, $\mathsf{B}\Sigma_{n+1}$, $\mathsf{B}\Pi_n$, $\mathsf{B}\Delta_{n+1}$ are all equivalent, and imply $\mathsf{I}\Delta_{n+1}$.
- (iii) $\mathsf{I}\Sigma_{n+1} \implies \mathsf{B}\Sigma_{n+1} \implies \mathsf{I}\Sigma_n$, and these implications are strict.

Theorem 3.6 [Sla04]. Over $PA^- + I\Sigma_0 + \exp$, $I\Delta_{n+1}$ is equivalent to $L\Delta_{n+1}$ (and hence to $B\Sigma_{n+1}$, $B\Pi_n$, $B\Delta_{n+1}$).

We will not define PA⁻ and exp, but they are extremely weak base theories, which will be subsumed by our chosen base theory RCA₀. Hence, we can simply assume the equivalences in Theorems 3.5 and 3.6, and our arithmetical principles are split into two families of equivalence classes, as shown in Figure 3.2.

We now prove a lemma about induction, which will be useful later.

Definition 3.7. For a formal system S, the *inductive formulae* I(S) *for* S is the collection of formulae φ such that $S \models I'\varphi$.




Figure 3.2: The induction, bounding, and least number principles. Vertical positioning denotes equivalence over $PA^- + I\Sigma_0$, and the horizontal dotted lines denote equivalence over $PA^- + I\Sigma_0 + \exp$.

Lemma 3.8. If $\mathcal{I}(S)$ is closed under negation, then it is closed under arbitrary Boolean combinations.

Proof. Let $\Gamma := \mathcal{I}(\mathcal{S})$: then $\mathcal{S} \models I'\Gamma$ (and indeed, Γ is maximal with this property). Since Γ is closed under negation, $\mathcal{S} \models L\Gamma$ too by Theorem 3.4. Since the connectives \neg and \land are complete for propositional logic [Smi03, p. 98], and we already know Γ is closed under \neg , it suffices to prove closure under \land .

So, pick $\varphi(x), \psi(x) \in \Gamma$: we want to show $\mathcal{S} \models \mathsf{I}'(\varphi \land \psi)$. For each n,

assume

$$(\forall m < n)[\varphi(m) \land \psi(m)] \to \varphi(n) \land \psi(n)$$

By contradiction, suppose there is a with $\neg(\varphi(a) \land \psi(a))$, and without loss of generality, suppose $\neg\varphi(a)$. By L Γ , we can assume a is minimal. By the inductive assumption, there is b < a with $\neg(\varphi(b) \land \psi(b))$, hence $\neg\psi(b)$ by minimality of a. Again by L Γ , assume b is minimal. Then, by the inductive assumption, there is c < b with $\neg(\varphi(c) \land \psi(c))$. But this contradicts minimality of a and b.

Lemma 3.8 allows us to induct on arbitrary Boolean combinations of Σ_1 and Π_1 formulae in RCA₀, which will prove useful.

3.2 Reverse mathematics

For a system S of second-order arithmetic, a *model* M of S consists of a set \mathbb{N} and a collection of subsets of \mathbb{N} , with appropriate interpretations for the \mathcal{L}_2 -symbols, so that all the axioms of S hold. All the systems S we consider will include the basic axioms of first-order arithmetic, hence \mathbb{N} will be restricted to range over nonstandard models $\omega + \mathbb{Z} \cdot K$, where the \mathcal{L}_2 -symbols are given the usual meanings. If $K = \emptyset$, i.e. $\mathbb{N} = \omega$, we call M an ω -model, and then M is determined by a collection of subsets of ω .

We are almost ready to define the subsystems RCA₀, WKL₀, ACA₀ of second-order arithmetic, which will form the basis of our work. Some of these subsystems will include the following second-order comprehension principle:

Definition 3.9. Let $\varphi(x)$ be a formula of arithmetic. The *comprehension principle for* φ is the sentence $C\varphi := (\exists X)(\forall n)(n \in X \leftrightarrow \varphi(n))$.

Essentially, $C\varphi$ asserts that the set $X=\{n:\varphi(n)\}$ exists. As before, given a class Γ of formulae, we write $C\Gamma$ for the theory consisting of $C\varphi$ for all (appropriate) $\varphi\in\Gamma$.

Definition 3.10. RCA₀ is the subsystem consisting of the basic axioms of first-order arithmetic, $I\Sigma_1^0$, and $C\Delta_1^0$.

RCA₀ is the system in which reverse mathematics is usually done. Although RCA₀ doesn't give us Σ_1 comprehension, we do get *bounded* Σ_1 comprehension:

Definition 3.11. Let $\varphi(x)$ be a formula of arithmetic. The *bounded comprehension principle for* φ is the sentence $\mathsf{BC}\varphi := (\forall k)(\exists X)(\forall n)[n \in X \leftrightarrow (\varphi(n) \land n < k)].$

BC φ asserts that for every k, the set $X_k = \{n < k : \varphi(n)\}$ exists.

Lemma 3.12 [Sim09, Thm II.3.9]. RCA₀ proves BC Σ_1 , i.e. BC φ for every Σ_1 formula φ .

Definition 3.13. ACA₀ is the subsystem consisting of the basic axioms of first-order arithmetic, $I\Delta_0^1$, and $C\Delta_0^1$.

WKL₀ has a slightly different definition. Recall that (finitary) Cantor space $2^{<\omega}$ is the set of all finite binary strings. A *tree* is a set $T \subseteq 2^{<\omega}$ that is closed under taking initial segments. A *path through* T is an infinite binary sequence α such that all initial segments are in T.

Definition 3.14. *Weak König's lemma* is the statement that every infinite tree $T \subseteq 2^{<\omega}$ contains a path. WKL₀ is the subsystem consisting of RCA₀ plus weak König's lemma.

Every infinite tree $T\subseteq 2^{<\omega}$ has a path which is arithmetical relative to T [Sim09, Example I.8.8]. Hence, ACA $_0$ implies WKL $_0$, which in turn implies RCA $_0$. In fact, all these implications are strict. Along with the stronger systems ATR $_0$ and Π^1_1 -CA $_0$, these make up the "big five" subsystems of second-order arithmetic. RCA $_0$ has a standard ω -model REC, consisting of the subsets of ω definable by a Δ_1 formula. Similarly, ACA $_0$ has a standard ω -model ARITH, consisting of the subsets of ω definable by an arithmetical

formula. We also have the full ω -model $\mathcal{P}(\omega)$, which is a model of all these axiom systems.

The idea of reverse mathematics is: for a known theorem φ of mathematics, find the weakest formal system \mathcal{S} such that $\mathcal{S} \vdash \varphi$. Given such a proof, we show \mathcal{S} is optimal by demonstrating a *reversal* of φ : a proof $\varphi \vdash \mathcal{S}$. In practice, no single theorem can axiomatise all of mathematics, and so we have to supplement φ with a base theory \mathcal{B} (i.e. a reversal is actually a proof $\mathcal{B} + \varphi \vdash \mathcal{S}$). Throughout this report, we will take $\mathcal{B} = \mathsf{RCA}_0$, unless stated otherwise.

If we prove $S \vdash \varphi$, and then reverse this (over \mathcal{B}), we say that φ is equivalent to S (over \mathcal{B}). A remarkable empirical result of reverse mathematics is that almost all theorems of mathematics are equivalent (over RCA₀) to one of the big five systems, though there are exceptions. In the case of abstract algebra, most results can be proven in ACA₀, except (ordinal) invariant results, which are usually equivalent to ATR₀, and structure theorems, which often fall at the level of Π_1^1 -CA₀. To demonstrate, we now review some reverse-mathematical studies of ring theory in the literature.

Theorem 3.15. The following are provable in RCA₀:

- (i) A field has no nontrivial proper ideals [DLM07].
- (ii) If $I \subseteq R$ is an ideal and R/I is a field, then I is maximal [DLM07].
- (iii) An ideal $I \subseteq R$ is prime if and only if R/I is an integral domain [Con10].
- (iv) Maximal ideals are prime [Con10].
- (v) Euclidean domains are PIDs [Sat16].

Theorem 3.16 (RCA₀). WKL₀ is equivalent to each of the following statements:

(i) Every commutative ring has a prime/radical ideal [FSS83; FSS85].

- (ii) If $I \subseteq R$ is a maximal ideal, then R/I is a field [DLM07].
- (iii) In Artinian rings, prime ideals are maximal [Con10].
- (iv) Every Artinian integral domain is a field [Con10].
- (v) $r \in R$ is nilpotent if and only if r belongs to every prime ideal [Sat16].
- (vi) (Local) Artinian rings are Noetherian [Con10; Con19].
- (vii) Every Artinian ring is a finite direct product of local Artinian rings [Con19].

Theorem 3.17 (RCA $_0$). ACA $_0$ is equivalent to each of the following statements:

- (i) Every commutative ring/domain has a maximal ideal [FSS83].
- (ii) Every commutative ring has a minimal prime ideal [Hat89].
- (iii) If *R* has no nontrivial, proper, principal ideals, it is a field [DLM07].
- (iv) An integral domain is a UFD iff it has the a.c.c.p. and all irreducibles are prime [GM17].
- (v) $r \in R$ belongs to every maximal ideal of R if and only if for all $s \in R$, 1 rs is a unit [Sat16].

3.3 Computability

Here, we review the basic notions of computability theory that we will need. The presentation will be very brief, so the reader unfamiliar with computability theory is urged to consult a textbook on the subject [Soa87; Soa16].

We will assume the Church–Turing thesis, and work with an informal notion of computability. Therefore, we say a set $A\subseteq \omega$ is *computable* if

there is an algorithm which, given n, always terminates and tells us if $n \in A$. Similarly, a function $f : \omega \to \omega$ is (total) computable if there is an algorithm which, given n, always halts and outputs f(n).

By this definition, most sets and functions arising in mathematics are computable. We can construct non-computable sets: the archetypal example is the halting problem \varnothing' , which is the set of pairs $\langle e,n\rangle$ for which the eth computable function halts on input n. Nonetheless, \varnothing' is still computably enumerable, or c.e.: there is an algorithm which lists its elements. There are several equivalent definitions of c.e.:

Proposition 3.18. The following are equivalent for a set $A \subseteq \omega$:

- (i) *A* is c.e., i.e. there is an algorithm which lists the elements of *A*.
- (ii) *A* is the domain of a partial computable function.
- (iii) *A* is the range of a partial computable function.
- (iv) *A* is empty, or the range of a total computable function.
- (v) *A* is finite, or the range of a total computable injection.
- (vi) A has a computable enumeration: a uniformly computable sequence $A_0 \subseteq A_1 \subseteq A_2 \subseteq \cdots$ such that $A = \bigcup_{s \in \omega} A_s$. We can further require that $|A_s| = s$.

There is a close correspondence between computability and the arithmetical hierarchy: a set $A \subseteq \omega$ is computable if and only if it can be defined by a Δ_1 formula, and c.e. if and only if it can be defined by a Σ_1 formula.

We recall some of the basic results of computability theory. Every algorithm can be coded by a natural number: for example, by writing it in a fixed (Turing-complete) programming language. This gives a listing φ_e of the partial computable functions. We can furthermore get a *uniformly computable* listing, i.e. the function $\langle e,n\rangle\mapsto \varphi_e(n)$ is computable. Similarly, we can get a uniformly c.e. listing W_e of the c.e. sets.

We can also consider *relativised computations*: those with access to an *oracle* $A \subseteq \omega$, so that the algorithm can ask at any point if any natural number n is in A. We say A is *Turing reducible* to B, and write $A \subseteq_T B$, if A can be computed with B as an oracle. The relation \subseteq_T is a preorder, so we obtain an equivalence relation \equiv_T in the standard way:

$$A \equiv_{\mathrm{T}} B \iff A \leq_{\mathrm{T}} B \text{ and } B \leq_{\mathrm{T}} A$$

The \equiv_{T} -equivalence-classes are called *Turing degrees*, and are partially ordered by \leq_{T} . The degree of all computable sets is called **0**.

For any oracle $A \subseteq \omega$, we can similarly list all the partial A-computable functions φ_e^A . Thus, for any A, we define the *Turing jump* A' as the halting problem relativised to A, i.e.

$$A := \{\langle e, n \rangle : \varphi_e^A(n) \text{ halts} \}$$

Then, $A <_{\mathrm{T}} A'$, i.e. A' is always strictly above A in the Turing degrees. The Turing jump is also a well-defined function on Turing degrees.

We also have a stronger notion of computable reduction: we say A is m-reducible to B ($A \leq_m B$) if there is a computable function $f: \omega \to \omega$ such that $n \in A \iff f(n) \in B$. Informally, the idea is that we can consult the oracle B only *once*. For some level Γ in the arithmetical hierarchy, we say that a Γ set A is Γ complete if for every Γ set B, $B \leq_m A$.

Proposition 3.19.

- (i) \emptyset' is Σ_1 complete.
- (ii) Inf = $\{e : W_e \text{ is infinite}\}\$ is Π_2 complete.

Another important class of Turing degrees are the PA degrees, which are those that can compute a complete consistent extension of Peano arithmetic. By Gödel's famous incompleteness theorem, **0** is not PA. **0**' is PA, but there are also PA degrees strictly below **0**'. There are also many equivalent characterisations of PA degrees:

Proposition 3.20. For a Turing degree d, the following are equivalent:

- (i) d is PA, i.e. it computes a complete consistent extension of PA.
- (ii) d computes a path through any computable (or Π_1) tree $T \subseteq 2^{<\omega}$.
- (iii) For any disjoint c.e. sets A, B, d computes a *separating set* for A and B, i.e. a set C such that $A \subseteq C$, $B \cap C = \emptyset$.

We can also speak about PA degrees relative to some oracle A, or PA degrees *over* A. Proposition 3.20 relativises: d is PA over A iff it computes an extension of "PA plus a predicate for A", iff it computes a path through any A-computable tree, iff it can separate any two disjoint A-c.e. sets.

Having reviewed classical computability, we now point out the close correspondence between computability and the subsystems RCA_0 , WKL_0 , ACA_0 of second-order arithmetic. This connection arises because our chosen base system RCA_0 somehow corresponds to the "computable world". Indeed, Δ_1 comprehension allows us to define *exactly* the subsets of ω which are computable (with parameters/oracles in the model).

Many of the theorems T studied in reverse mathematics have the form

$$T = (\forall X) \big[\varphi(X) \to (\exists Y) \, \psi(X, Y) \big]$$

where $\varphi(X)$ and $\psi(X,Y)$ are properties of the sets X and Y. We could view T as a problem or challenge: given a set X such that $\varphi(X)$, find a set Y such that $\psi(X,Y)$. Now the connection is thus: $\mathsf{RCA}_0 \vdash T$ when we can always choose Y to be X-computable. $\mathsf{WKL}_0 \vdash T$ when we can choose Y computable from a PA degree over X, and $\mathsf{ACA}_0 \vdash T$ if we can choose Y arithmetical in X.

This correspondence is also useful in reversals of T. Over RCA_0 , $T \vdash \mathsf{WKL}_0$ if for every A, we can construct A-computable X so that every suitable Y is PA over A. Similarly, $T \vdash \mathsf{ACA}_0$ if we can construct X so that every Y computes A'.

While this correspondence is completely precise only in ω -models, a nearly identical proof can generally be made to work in nonstandard models. Furthermore, we will generally just take X to be computable (i.e. $A=\varnothing$), as the fully relativised version will again follow nearly identically. So in practice, we will prove something like "there is a computable X with $\varphi(X)$, such that every Y with $\varphi(X,Y)$ has PA degree [computes \varnothing']", and we will take this as evidence that $T \vdash \mathsf{WKL}_0$ [resp. $T \vdash \mathsf{ACA}_0$].

Now, we will construct various computability-theoretic objects, which will be useful later when proving reversals. We saw that a degree d is PA iff it can separate any two disjoint c.e. sets A, B. Our first lemma is that there is a "universal pair" A, B so that *any* separator has PA degree:

Lemma 3.21. There are disjoint c.e. sets A, B such that whenever C has $A \subseteq C$, $B \cap C = \emptyset$, then C has PA degree.

Proof. Fix a coding F_n of all sentences in the language of PA. Let $A = \{n : \mathsf{PA} \vdash F_n\}$ and $B = \{n : \mathsf{PA} \vdash \neg F_n\}$. Then, A, B are c.e. since PA is computably axiomatisable, so we can search for a proof of F_n from PA. $A \cap B = \emptyset$ since PA is consistent.

We claim A, B are as required. Fix a separator C of A and B. There is no reason C should be consistent, but we can use it to compute a complete consistent extension D, as follows. We build D in stages D_s , with $D_0 = \emptyset$. At stage s, let $G_s := (\bigwedge_{F \in D_s} F) \to F_s$. If $G_s \in C$, let $D_{s+1} = D_s \cup \{F_s\}$; else, let $D_{s+1} = D_s \cup \{\neg F_s\}$. By construction, D is a complete consistent extension of PA, hence C is PA.

Lemma 3.21 will be useful in showing a theorem implies WKL₀. To show a theorem implies ACA₀, we will employ a few different techniques. The first is simply to code \varnothing' itself into a computable ring. Often it is not possible to do this directly, so we have a few other methods.

Definition 3.22. Fix a computable enumeration \varnothing'_s of \varnothing' . The *modulus of* \varnothing' is the function $\mu_{\varnothing'}$ mapping n to the least s such that $\varnothing'_s \upharpoonright_n = \varnothing' \upharpoonright_n$.

Theorem 3.23. Suppose $f: \omega \to \omega$ dominates $\mu_{\varnothing'}$: for every n, $f(n) \ge \mu_{\varnothing'}(n)$. Then, $f \ge_T \varnothing'$.

Proof. We compute if $n \in \emptyset'$ by asking if $n \in \emptyset'_{f(n+1)}$. By definition of $\mu_{\emptyset'}$, this computation is always correct.

Another useful trick to code \varnothing' into a construction is using a c.e. set that is so "dense" that its complement dominates μ'_{\varnothing} .

Lemma 3.24. There is a c.e. set *A* such that

- (i) A^{\complement} is infinite.
- (ii) Any infinite subset $B \subseteq A^{\complement}$ computes \varnothing' .

Proof. Fix an enumeration of \varnothing' , and enumerate A starting with $A_0 = \varnothing$. Now, at stage s, suppose $A_s^{\complement} = \{c_0 < c_1 < \cdots\}$. If n enters \varnothing' at stage s, we put $c_n, c_{n+1}, \ldots, c_s$ into A. We now verify that A has the required properties.

- (i) A^{\complement} is infinite: by induction, all the A_s are finite, so each A_s^{\complement} is infinite. Now given $k \in \mathbb{N}$, $A_{\mu_{\varnothing'}(k+1)}^{\complement}$ contains some j > k which will never leave A^{\complement} .
- (ii) Any infinite subset $B \subseteq A^{\complement}$ computes \varnothing' in a very simple way: if $B = \{b_0 < b_1 < \cdots\}$, then the function $n \mapsto b_n$ dominates $\mu_{\varnothing'}$. The proof is as follows. If $A^{\complement} = \{c_0 < c_1 < \cdots\}$, then each $b_i = c_j$ for some j with $i \le j \le b_i$.

Now if any $k \leq i$ entered \emptyset' at a stage $s \geq b_i$, then $c_k, \ldots, c_i, \ldots, c_j = b_i, \ldots, c_{b_i}$ would have been put into A (by construction). As $b_i \notin A$, this can't have happened, so $\emptyset_{b_i} \upharpoonright_i = \emptyset \upharpoonright_i$, i.e. $b_i \geq \mu_{\emptyset'}(i)$.

To finish this section, we will review some useful ideas from a subbranch of computability called *computable structure theory* [Mon21]. Essentially, this is the study of model theory from the point of view of computability, and the primary objects of study are computable structures. For what follows, let \mathcal{T} be a computable theory in a finite language \mathcal{L} .

Definition 3.25. A computable (presentation of a) T-structure \mathcal{M} consists of:

- (i) A computable subset $M \subseteq \mathbb{N}$.
- (ii) For every constant $c \in \mathcal{L}$, an element $c^{\mathcal{M}} \in M$.
- (iii) For every k-ary function $f \in \mathcal{L}$, a computable function $f^{\mathcal{M}} \colon M^k \to M$.
- (iv) For every k-ary relation $R \in \mathcal{L}$, a computable function $R^{\mathcal{M}} \colon M^k \to \{0,1\}$.

such that, with these interpretations of the \mathcal{L} -symbols, \mathcal{M} satisfies all formulae in \mathcal{T} .

A c.e. \mathcal{T} -structure consists of the same data, but we allow M to be c.e. instead.

We will be primarily concerned with the language $\mathcal{L}_{ring} = \{0, 1, +, \cdot\}$, and the \mathcal{L}_{ring} -theory \mathcal{T}_{ring} consisting of the usual commutative ring axioms. A *computable ring* is just a computable \mathcal{T}_{ring} -structure, i.e. a computable set $R \subseteq \mathbb{N}$ with elements $0_R, 1_R \in R$ and computable binary operations $+, \cdot$ on R which form a ring.

When doing reversals, we will often need to construct a computable ring having certain properties. However, the following result shows that it is sufficient to construct a c.e. ring:

Theorem 3.26. Any c.e. \mathcal{T} -structure \mathcal{M} is (computably) isomorphic to a computable \mathcal{T} -structure \mathcal{M}' .

Proof. If \mathcal{M} is finite, this is trivial, so assume \mathcal{M} is infinite, and fix an injective computable enumeration $\varphi \colon \mathbb{N} \to \mathcal{M}$. The inverse $\varphi^{-1} \colon \mathcal{M} \to \mathbb{N}$ is partial computable, since given $m \in \mathcal{M}$, we can search for the n such that $\varphi(n) = m$.

Now, define \mathcal{M}' as the structure with support \mathbb{N} , and \mathcal{L} -symbols interpreted

$$c^{\mathcal{M}'} := \varphi^{-1}(c^{\mathcal{M}})$$

$$f^{\mathcal{M}'}(n_1, \dots, n_k) := \varphi^{-1} \big[f^{\mathcal{M}} \big(\varphi(n_1), \dots, \varphi(n_k) \big) \big]$$

$$R^{\mathcal{M}'}(n_1, \dots, n_k) := R^{\mathcal{M}} \big(\varphi(n_1), \dots, \varphi(n_k) \big)$$

Then, \mathcal{M}' is a computable \mathcal{T} -structure, and by construction, φ is a computable \mathcal{T} -isomorphism $\mathcal{M}' \to \mathcal{M}$.

When using Theorem 3.26 in practice, we will often abuse notation, and identify \mathcal{M} and \mathcal{M}' via the computable isomorphism φ . This means we will treat c.e. rings as if they are computable. We should note that the isomorphism will also preserve many computable subobjects of \mathcal{M} . For example, if $I \subseteq \mathcal{M}$ is a computable ideal, then so is $\varphi^{-1}(I) \subseteq \mathcal{M}'$.

A particularly useful case of Theorem 3.26 is localising a computable ring by a c.e. subset:

Corollary 3.27. Suppose R is a computable ring, and $M \subseteq R$ is a multiplicatively closed c.e. subset containing 1_R but no zero-divisors. Then Loc(R, M) is (computably) isomorphic to a computable ring S.

Proof. Define Loc (R, M) exactly as in Definition 2.10. This is a computable quotient of a c.e. structure, hence we get a c.e. structure. Now apply Theorem 3.26.

Chapter 4

Radicals of ideals and primary ideals

We now begin our study of ring theory in second-order arithmetic. We can use the standard definitions of rings, polynomial rings, ideals, etc. verbatim in second-order arithmetic. The definition of quotient ring might cause some concern, since we naively define R/I as a set of sets. However, we can amend this by defining the elements of R/I to be minimal representatives of their equivalence class [Sim09, Defn III.5.2].

Definition 4.1 (RCA₀). Let $I \subseteq R$ be an ideal. The *quotient ring* R/I is the set

$$R/I := \{r \in R : (\forall s <_{\mathbb{N}} r)(s \in R \to r - s \notin I)\}$$

which exists by Δ_1 comprehension. RCA $_0$ can define a function $q \colon R \to R/I$, called the *quotient map*, so that for every $r \in R$, q(r) is the unique element of R/I such that $r - q(r) \in I$. The ring operations on R/I are then the operations induced by q from R.

A similar trick can be used to construct the field of fractions or localisations in RCA₀. As usual, we use r+I (or simply r) to denote q(r). In RCA₀, it follows from Definition 4.1 that r+I=s+I iff $r-s\in I$. We will also make wide use of the following theorem:

Theorem 4.2 (ideal correspondence theorem; RCA₀). For a ring R and ideal $I \subseteq R$, the quotient map $q: R \to R/I$ is an isomorphism between the ideals of R containing I, and the ideals of R/I.

Furthermore, the quotient map preserves many properties of ideals, such as maximality, primality, primary-ness, being the radical of another ideal, etc. As a result, this frequently gives an equivalence between statements of the form $\forall R \ \forall I \subseteq R \ \varphi(R,I)$ and $\forall R \ \varphi(R,\{0\})$, with the equivalence provable in RCA₀.

By fixing the parameters a and b, the following lemmas can be proved by Δ_0 induction.

Lemma 4.3 (RCA₀). For all a, b in a commutative ring R, $(ab)^n = a^n b^n$.

Lemma 4.4 (binomial theorem; RCA₀). For all a, b in a commutative ring R, $(a+b)^n = \sum_{j=0}^n \binom{n}{j} a^j b^{n-j}$.

We now examine some properties of ideals and the relationships between them, and show that most of these relationships are provable in RCA_0 .

Definition 4.5. A ring R is *reduced* if it has no nontrivial nilpotent elements, i.e. whenever $x^n = 0$ for some n > 0, then x = 0.

Definition 4.6. Let $I \subseteq R$ be an ideal.

- (i) *I* is *prime* if whenever $ab \in I$, then $a \in I$ or $b \in I$.
- (ii) *I* is *primary* if whenever $ab \in I$, then $a \in I$ or $b^n \in I$ for some n.
- (iii) The radical of I is defined $\sqrt{I} = \{a \in R : \exists n > 0 \text{ such that } a^n \in I\}.$
- (iv) I is *semiprime* or *radical* if whenever $a^n \in I$ for some n > 0, then $a \in I$ (equivalently, $I = \sqrt{I}$).
- (v) The adjoint of I is defined $I^{\not\perp} = \{a \in R : \exists b \notin I \text{ such that } ab \in I\}.$

(vi) I is *primal* if I^{\perp} forms an ideal.

Theorem 4.7. The following are provable in RCA₀:

- (i) \sqrt{I} is an ideal.
- (ii) $I \subseteq R$ is semiprime if and only if R/I is reduced.
- (iii) If *I* is primary, then \sqrt{I} is prime¹.
- (iv) I is prime iff it is primary and semiprime.
- (v) I is primary iff every zero divisor in R/I is nilpotent.
- (vi) If $I \neq R$ is primary, then I is primal (and furthermore, $I^{\not\perp} = \sqrt{I}$).
- (vii) If I is primal, then I^{\perp} is prime.

Proof.

(i) We prove only closure under addition. Suppose $a,b\in \sqrt{I}$, i.e. $a^n,b^m\in I$. Let k:=n+m-1. Using the binomial formula, we can write $(a+b)^k$ as

$$a^{k} + \cdots + {k \choose n} a^{n} b^{m-1} + {k \choose m} a^{n-1} b^{m} + \cdots + b^{k}$$

$$a^{n} () + b^{m} () \in I$$

hence $a + b \in \sqrt{I}$.

- (ii) By the ideal correspondence theorem, this is equivalent to saying $\{0\}$ is semiprime iff R is reduced, which is trivial.
- (iii) Suppose $ab \in \sqrt{I}$: then $(ab)^n = a^nb^n \in I$, so either $a^n \in I$, whence $a \in \sqrt{I}$, or $(b^n)^k = b^{nk} \in I$, whence $b \in \sqrt{I}$.

 $^{^{1}}$ We also say that I is *quasi-primary*.

- (iv) (\Rightarrow) Any prime ideal is trivially primary (take n=1), and semiprime by Δ_0 induction on $\varphi(n): a^n \in I \to a \in I$ for a fixed parameter $a \in I$.
 - (\Leftarrow) If $ab \in I$, then by primary-ness, either $a \in I$ or $b^n \in I$, whence $b \in I$ by semiprimality.
- (v) By the ideal correspondence theorem, this is equivalent to saying $\{0\}$ is primary iff every zero divisor in R is nilpotent, which is immediate from the definition of primary.
- (vi) By (i), it is enough to prove the assertion in brackets. Pick $a \in I^{\not\perp}$, i.e. there is $b \notin I$ with $ab \in I$. Since I primary, either $b \in I$ (which is a contradiction), or $a \in \sqrt{I}$ as required. Now if $a \in \sqrt{I}$, by $\mathsf{L}\Sigma_1$, there is a minimal n > 0 such that $a^n \in I$. Then a^{n-1} witnesses that $a \in I^{\not\perp}$.
- (vii) Suppose $ab \in I^{\not\perp}$, i.e. there is $c \notin I$ such that $abc \in I$. If $bc \in I$, then c witnesses that $b \in I^{\not\perp}$. If $bc \notin I$, then bc witnesses that $a \in I^{\not\perp}$. \square

Now, we present some basic results about operations and relations between two ideals and their radicals, which are provable in RCA₀.

Definition 4.8. Ideals $I, J \subseteq R$ are *comaximal* if every $r \in R$ can be written as r = i + j for $i \in I$, $j \in J$.

It suffices (under RCA₀) to show that 1 = i + j.

Theorem 4.9. The following are provable in RCA₀:

- (i) If \sqrt{I} and \sqrt{J} are comaximal, then I and J are comaximal.
- (ii) $\sqrt{I \cap J} = \sqrt{I} \cap \sqrt{J}$.

Proof.

(i) Suppose 1 = a + b where $a^n \in I$, $b^m \in J$. Let k := n + m - 1. Then, using the binomial formula, we can write $1 = 1^k = (a + b)^k$ as

$$a^{k} + \cdots + {k \choose n} a^{n} b^{m-1} + {k \choose m} a^{n-1} b^{m} + \cdots + b^{k}$$

 $a^{n} () + b^{m} ()$

Thus, 1 = i + j for suitable $a^n \mid i, b^m \mid j$.

(ii) (⊆) Trivial.

$$(\supseteq)$$
 If $a \in \sqrt{I} \cap \sqrt{J}$, then $a^n \in I$, $a^m \in J$, so $a^{\max\{n,m\}} \in I \cap J$.

To conclude this section, we analyse the following characterisation of the radical:

$$\sqrt{I} = \bigcap_{\substack{P \supseteq I \\ P \text{ prime}}} P$$

However, we require ACA₀ to show the LHS exists, and Π_1^1 -CA₀ to show the RHS exists. So, we will analyse the part of this theorem that doesn't require comprehension for either side. The two containments can be written respectively as:

- I and \sqrt{I} are contained in exactly the same prime ideals.
- If $x \notin \sqrt{I}$, then there is a prime ideal $P \supseteq I$ such that $x \notin P$.

By the ideal correspondence theorem, these are respectively equivalent to:

- If r is nilpotent, it belongs to every prime ideal of R.
- If *r* belongs to every prime ideal of *R*, then it is nilpotent.

[Sat16, p. 62] showed that the first statement is provable in RCA_0 , and the second is equivalent to WKL_0 .

4.1 The RAD principle

So far, we have seen that almost all basic facts about radicals and primary ideals are provable in RCA₀. The notable exception is the following:

Theorem 4.10 (RCA₀). WKL₀ is equivalent to "for all ideals $I \subseteq R$, if \sqrt{I} is maximal, then I is primary".

Proof.

(\Rightarrow) We prove the contrapositive of the consequent. Suppose I is not primary: then there are $a,b\in R$ such that $ab\in I$, $a\notin I$, $b\notin \sqrt{I}$. Using WKL₀, we will construct an ideal $J\supsetneq \sqrt{I}$.

Define $T\subseteq 2^{<\omega}$ as the set of all $F\subseteq \{0,1,\ldots,n-1\}$ such that the following hold:²

- (i) If $0_R < n$, then $0_R \in F$.
- (ii) For all $c, d \in F$, if c + d < n, then $c + d \in F$.
- (iii) For all $c \in F$, d < n, if cd < n, then $cd \in F$.
- (iv) For all c < n, if $c^n \in I$, then $c \in F$.
- (v) If b < n, then $b \in F$.
- (vi) If $1_R < n$, then $1_R \notin F$.

T is computable since for a finite string, we can simply check all the above conditions exhaustively. A path J through T is a proper ideal containing b and \sqrt{I} : since $b \notin \sqrt{I}$, it follows that $J \supsetneq \sqrt{I}$. Also, T is downwards closed, hence a tree.

T is infinite since every level is nonempty. Given n, let

$$A_n := \begin{cases} \{c < n : c^n \in I\} \cup \{b\} & b < n \\ \{c < n : c^n \in I\} & b \ge n \end{cases}$$

Enumerating $A_n = \{c_1, \ldots, c_k\}$, let

$$F_n := \{c < n : (\exists d_1, \dots, d_k \in R) (c = c_1 d_1 + \dots + c_k d_k)\} = (A_n) \upharpoonright_n$$

which exists by bounded Σ_1 comprehension (Lemma 3.12).

²Here, we identify binary strings of length n with subsets $F \subseteq \{0, 1, \dots, n-1\}$.

We claim $F_n \in T$ at level n. Conditions (i)–(v) are evidently satisfied. For condition (vi), note that it can only fail if $1_R < n$ and $1_R \in F$, i.e. $1_R = c_1d_1 + \cdots + c_kd_k$ for some $d_1, \ldots, d_k \in R$. Expanding

$$1_R = (1_R)^{nk} = (c_1d_1 + \dots + c_kd_k)^{nk}$$

note that each term is divisible by either b or some $c^n \in I$, i.e. $1_R \in (I, b)$. Then $a = a1_R \in (I, ab) = I$, giving a contradiction.

(\Leftarrow) We construct a computable ring R and a computable, proper ideal $I \subsetneq R$ which is not primary, such that any proper ideal $J \supsetneq \sqrt{I}$ has PA degree. Thus, the assumption that \sqrt{I} is not maximal gives a larger ideal $J \supsetneq \sqrt{I}$ of PA degree, which computes WKL₀.

Fix disjoint c.e. sets A, B as in Lemma 3.21. We will build R and I such that any proper ideal $J \supseteq \sqrt{I}$ computes a separator of A and B.

Let $R = \mathbb{Z}[x, y, z_0, z_1, \ldots]$. To begin, let I = (xy) (the ideal generated by xy). This is to ensure I is not primary (with witness xy). Now if j enters A at stage s, add z_j^s to the list of generators for I. If j enters B at stage s, add $(z_j - 1)^s$ to the list of generators for I.

Then, I is computable - to work out if $p \in I$, we only need to run the construction up to stage $s = \deg(p)$. At this stage, I is finitely generated, so we can compute if $p \in I$. We can assume nothing is enumerated when s = 0, meaning I (and \sqrt{I}) are proper ideals.

Now, let $J \supseteq \sqrt{I}$ be a proper ideal of R. Let $X_J = \{n \in \mathbb{N} : z_n \in J\}$. Then, X_J separates A and B: $A \subseteq X_J$, and $B \cap X_J = \emptyset$, because if there were $n \in B \cap X_J$, then $z_n, (z_n - 1) \in J$ so $1 \in J$, contradicting that J is proper. Since X_J is J-computable, it follows that J has PA degree.

In the (\Leftarrow) construction of Theorem 4.10, note that \sqrt{I} itself is of PA degree. Thus, the computational power lies in comprehension for an ideal containing \sqrt{I} , possibly not strictly. By asserting that \sqrt{I} itself must be

computable, the focus shifts more to the (non-)primary-ness of I. In other words, we want to establish the reverse-mathematical strength of the statement

"if \sqrt{I} exists and is maximal, then I is primary" (RAD) or equivalently,

"if I is not primary and \sqrt{I} exists, then \sqrt{I} is not maximal"

By the ideal correspondence theorem, RAD is also equivalent to:

"if $\sqrt{0}$ exists and is maximal, then $\{0\}$ is primary" "if $\{0\}$ is not primary and $\sqrt{0}$ exists, then $\sqrt{0}$ is not maximal"

From Theorem 4.10, it follows that:

Proposition 4.11. WKL₀ proves RAD.

Conversely, we might try to argue that RAD implies WKL₀. One strategy is to use a construction similar to [DLM07, Thm 3.2]. As before, we fix disjoint, c.e. sets $A, B \subseteq \mathbb{N}$ such that any separator is of PA degree. Starting with some "ring of coefficients" U, we set $R_0 := U[x_0, x_1, \ldots]$. Then, we enumerate R as a c.e. subring of the total quotient ring of R_0 , adding x_n/f for $n \in A$, $f \in U[x_0, \ldots, x_{n-1}]$ a non-zero-divisor, and $(x_n - 1)/f$ for $n \in B$, $f \in U[x_0, \ldots, x_{n-1}]$ a non-zero-divisor.

We want to choose U so that R satisfies the following:

- (a) $I = \{0\}$ is not primary (equivalently, there are non-nilpotent zero divisors).
- (b) The nilradical $\sqrt{0}$ is computable.
- (c) Any proper ideal $J \supsetneq \sqrt{0}$ has PA degree.

To ensure R satisfies condition (c), we will force it to satisfy:

(d) Any proper ideal $J \supseteq \sqrt{0}$ contains a non-zero-divisor.

Then, given an ideal $J \supseteq \sqrt{0}$, we have a non-zero-divisor $f \in J$, so $x_n = f \cdot (x_n/f) \in J$ for sufficiently large $n \in A$, and similarly $x_n \notin J$ for sufficiently large $n \in B$. Thus, J computes a separator for A and B up to finitely many differences. Sensible though it may seem, this strategy is doomed to fail because of the following:

Proposition 4.12. $\{0\} \subseteq R$ is primary iff any proper ideal $J \supseteq \sqrt{0}$ contains a non-zero-divisor.

Proof [MSE21].

- (\Rightarrow) By Theorem 4.7.(v), every zero-divisor is nilpotent, so since J contains a non-nilpotent, it contains a non-zero-divisor.
- (\Leftarrow) Assuming any proper ideal $J \supsetneq \sqrt{0}$ contains a non-zero-divisor, we will show any non-nilpotent is a non-zero-divisor. Let x be non-nilpotent: then the ideal $(x,\sqrt{0})\supsetneq\sqrt{0}$ contains a non-zero-divisor by assumption. In other words, there are $a,b\in R$ so that ax+b is a non-zero-divisor, and $b^n=0$.

By induction, we show that for every d>0, $(ax+b)^d$ is not a zero-divisor. The base case d=1 is true by assumption. Now, suppose $(ax+b)^d$ is not a zero-divisor, but $(ax+b)^{d+1}$ is. Let $r\neq 0$ be such that $(ax+b)^{d+1}r=0$. Then, $(ax+b)^d(ax+b)r=0$. By assumption, ax+b is not a zero-divisor, so $(ax+b)r\neq 0$: this shows that $(ax+b)^d$ is a zero-divisor, a contradiction.

In particular $(ax + b)^n$ is a non-zero-divisor. Expanding $(ax + b)^n$ using the binomial formula, and using the fact that $b^n = 0$, we have $(ax + b)^n = cx$ for some $c \neq 0$. Since cx is not a zero-divisor, it follows that x is not a zero-divisor, as required.

Currently, we know that RAD lies between RCA₀ and WKL₀, but its exact reverse-mathematical strength remains open. It's possible that RAD falls into the "zoo" of reverse-mathematical principles lying strictly between

 RCA_0 and WKL_0 , which have been keenly studied in recent years [San18; RMZoo].

Question 4.13. What is the exact reverse-mathematical strength of RAD?

Chapter 5

Noetherian rings

As we proceed, we will consider ideals $\mathcal{I} \subseteq R$ which may not be *computable* relative to R, but only c.e. relative to R.

Definition 5.1. A Σ_1 -ideal $\mathcal{I} \subseteq R$ is a sequence $\langle a_0, a_1, a_2, \ldots \rangle$ of elements of R, such that for all $i, j \in \mathbb{N}$ and $r, s \in R$, there exists $k = k(i, j, r, s) \in \mathbb{N}$ such that $a_k = a_i r + a_j s$.

To be precise, we will define basic notions such as subset and equality for Σ_1 -ideals. Note that these are defined *non-uniformly*—we don't require there to exist a function witnessing the inclusion/equality.

Definition 5.2. Let $\mathcal{I} = \langle a_0, a_1, \ldots \rangle$, $\mathcal{J} = \langle b_0, b_1, \ldots \rangle$ be Σ_1 -ideals in R, and $r \in R$.

- (i) We say $r \in \mathcal{I}$ if there exists m such that $r = a_m$.
- (ii) We say $\mathcal{I} \subseteq \mathcal{J}$ if for all n, $a_n \in \mathcal{J}$.
- (iii) We say $\mathcal{I} \subsetneq \mathcal{J}$ if $\mathcal{I} \subseteq \mathcal{J}$ and $\mathcal{J} \nsubseteq \mathcal{I}$.
- (iv) We say $\mathcal{I} = \mathcal{J}$ if $\mathcal{I} \subseteq \mathcal{J}$ and $\mathcal{J} \subseteq \mathcal{I}$.

Definition 5.3.

(i) For any $n \in \mathbb{N}$ and $a_0, \ldots, a_{n-1} \in R$, the set

$$(a_0, \dots, a_{n-1}) := \{a_0r_0 + \dots + a_{n-1}r_{n-1} : r_0, \dots, r_{n-1} \in R\}$$

is a Σ_1 -ideal, called the Σ_1 -ideal generated by a_0, \ldots, a_{n-1} .

(ii) A Σ_1 -ideal \mathcal{I} is *finitely generated* if there exist $n \in \mathbb{N}$ and $a_0, \ldots, a_{n-1} \in \mathbb{R}$ such that $\mathcal{I} = (a_0, \ldots, a_{n-1})$, in the sense of Definition 5.2.(iv).

Typical examples of Σ_1 -ideals are principal ideals $(a) \subseteq R$, which in general are not computable. Σ_1 -ideals are thus the right notion of ideal for PIDs.

As [Sat16] remarks, defining Noetherian rings is difficult, because there are many different, classically equivalent notions which are not equivalent over RCA₀. For a countable commutative ring R, [Sat16] considers the following eight definitions of Noetherian:¹

- (i) Every Δ_1 -ideal $I \subseteq R$ is finitely generated.
- (ii) R has no strictly increasing chain of Δ_1 -ideals $I_0 \subsetneq I_1 \subsetneq \cdots$.
- (iii) Every chain of Δ_1 -ideals $I_0 \subseteq I_1 \subseteq \cdots$ in R eventually stabilises.
- (iv) R contains no sequence a_0, a_1, \ldots such that for all $i, a_i \notin (a_0, \ldots, a_{i-1})$.
- (v) For all a_0, a_1, \ldots in R, there is i such that $\{a_i : j \in \mathbb{N}\} \subseteq (a_0, \ldots, a_{i-1})^2$
- (vi) R has no strictly increasing chain of Σ_1 -ideals $\mathcal{I}_0 \subsetneq \mathcal{I}_1 \subsetneq \cdots$.
- (vii) Every chain of Σ_1 -ideals $\mathcal{I}_0 \subseteq \mathcal{I}_1 \subseteq \cdots$ in R eventually stabilises.
- (viii) Every Σ_1 -ideal $\mathcal{I} \subseteq R$ is finitely generated.

¹[Sat16] actually considered their negations, as well as two further conditions which don't characterise Noetherian, but instead a (weaker) condition called a.c.c.p..

²Simpson [Sim88] called property (v) "Hilbertian", and noted the equivalence $(v)\Leftrightarrow(viii)$.

We conduct a full reverse-mathematical analysis of these conditions. Firstly, we reduce them to just five conditions, by showing that, over RCA₀, (ii) and (iii) are equivalent, and (v), (vii) and (viii) are equivalent.

- (ii) \Rightarrow (iii): By contradiction, suppose there is a chain of Δ_1 -ideals $I_0 \subseteq I_1 \subseteq \cdots$ in R which never stabilises. Fix an enumeration a_0, a_1, \ldots of R. Define $n_0 = 0$, j_0 least such that $a_{j_0} \in I_{n_0}$ and $\langle n_{k+1}, j_{k+1} \rangle$ the least pair³ such that $a_{j_{k+1}} \in I_{n_{k+1}} \setminus I_{n_k}$. Then, $I_{n_0} \subsetneq I_{n_1} \subsetneq \cdots$ is a strictly increasing, uniformly Δ_1 chain of ideals.
- (iii)⇒(ii): Trivial.
- (v) \Rightarrow (viii): Let a_0, a_1, \ldots be an enumeration of \mathcal{I} . By assumption, there is i such that $\mathcal{I} \subseteq (a_0, \ldots, a_{i-1})$, whence $\mathcal{I} = (a_0, \ldots, a_{i-1})$ is finitely generated.
- (viii) \Rightarrow (vii): Given a chain of Σ_1 -ideals $\mathcal{I}_0 \subseteq \mathcal{I}_1 \subseteq \cdots$ in R, $\mathcal{I} := \bigcup_{n=0}^{\infty} \mathcal{I}_n$ is a Σ_1 -ideal. Suppose $\mathcal{I} = (b_0, \ldots, b_{m-1})$. Then, in particular, each $b_i \in \mathcal{I}$, so $(\forall i < m)(\exists k_i)(b_i \in I_{k_i})$. The predicate " $b_i \in I_{k_i}$ " is Σ_1 , hence by $\mathsf{B}\Sigma_1$ (see Definition 3.3), we conclude that $(\exists n)(\forall i < m)(\exists k_i < n)(b_i \in I_{k_i})$. Hence all b_i are in I_n , whence the chain stabilises at n.
- (vii) \Rightarrow (v): $(a_0) \subseteq (a_0, a_1) \subseteq (a_0, a_1, a_2) \subseteq \cdots$ is a nested chain of Σ_1 -ideals, so there is some i such that for all j, $a_j \in (a_0, \ldots, a_j) = (a_0, \ldots, a_{i-1})$.
- (vii) \Rightarrow (vi) follows trivially in RCA₀. The converse also seems computably true; however, the induction necessary to prove it appears to go beyond $I\Sigma_1$.

Theorem 5.4 (RCA₀ + $I\Sigma_2$). (vi) \Rightarrow (vii).

Proof. By contradiction, suppose there is a uniformly c.e. chain of Σ_1 -ideals $\mathcal{I}_0 \subseteq \mathcal{I}_1 \subseteq \cdots$ in R which never stabilises. We will build a strictly increasing, uniformly c.e. subsequence $\mathcal{J}_0 \subsetneq \mathcal{J}_1 \subsetneq \cdots$ via a "moving marker" priority argument [Soa16, §4.3.2].

³Such a pair always exists, by assumption. We can find it simply by brute force search.

Each \mathcal{J}_n will have a "marker" m_n pointing to the ideal \mathcal{I}_{m_n} , which it will copy: any element enumerated into \mathcal{I}_{m_n} is also enumerated into \mathcal{J}_n . Additionally, each \mathcal{J}_n , n > 0 keeps track of a "witness" $x_n \in \mathcal{J}_n$ which it believes is not in \mathcal{J}_{n-1} . If we see x_n enter \mathcal{J}_{n-1} , then we increment m_n , and enumerate \mathcal{J}_n till we find a new witness. We will assume that every element enumerated into \mathcal{J}_n also enters \mathcal{J}_m for m > n.

To begin, let $m_0 = 0$, and enumerate the first element of \mathcal{I}_0 into \mathcal{J}_0 . At stage s > 0, suppose we have finitely enumerated $\mathcal{J}_0, \ldots, \mathcal{J}_{s-1}$, and defined markers m_0, \ldots, m_{s-1} and witnesses x_1, \ldots, x_{s-1} . Define \mathcal{J}_s as the current value of \mathcal{J}_{s-1} , and $m_s = m_{s-1} + 1$. For each $i \leq s$, enumerate the next element of \mathcal{I}_{m_i} into $\mathcal{J}_i, \ldots, \mathcal{J}_s$, and let x_s be some element enumerated⁴ into \mathcal{J}_{s-1} .

Now, for $i=1,\ldots,s$ in increasing order, we check if x_i has been enumerated into \mathcal{J}_{i-1} during this stage. If not, nothing need be done. If so, increment m_i by 1, and start enumerating the new \mathcal{I}_{m_i} into $\mathcal{J}_i,\ldots,\mathcal{J}_s$. Since \mathcal{J}_{i-1} is finite so far, we will eventually see an element not in \mathcal{J}_{i-1} : this is our new witness x_i . We stop enumerating \mathcal{I}_{m_i} when x_i is found. This concludes stage s.

Now, we prove by induction on n that every marker m_n eventually stabilises. Formally, we induct on the Σ_2 formula

$$\varphi(n) \coloneqq (\exists s)(\forall t > s)(m_{n,t} = m_{n,s})$$

where $m_{i,s}$ is the value of the *i*th marker at stage s.

Firstly, note that $m_0=0$ never changes. Now, suppose that m_0,\ldots,m_{n-1} have stabilised already. There are infinitely many elements in $\bigcup \mathcal{I}_k \setminus \mathcal{I}_{m_{n-1}}$: let k^* be least such that \mathcal{I}_{k^*} contains one. Then, m_n will stabilise at some point $\geq k^*$, since after this point, we will eventually enumerate an element of $\bigcup \mathcal{I}_k \setminus \mathcal{I}_{m_{n-1}}$, and thus take a witness $x_n \notin \mathcal{I}_{m_{n-1}}$, which will never be discarded. Therefore, the x_n witness that the chain $\mathcal{J}_0 \subseteq \mathcal{J}_1 \subseteq \cdots$ is strictly increasing.

⁴Choosing $x_s \in \mathcal{J}_{s-1}$ ensures that the process in the next paragraph will be carried out for i = s.

We leave open the question of whether $I\Sigma_2$ is actually necessary to prove Theorem 5.4, and instead assume a base theory of $RCA_0 + I\Sigma_2$ for the rest of this chapter. Over $RCA_0 + I\Sigma_2$, we then get four distinct notions of Noetherian, which we name Δ_1 -Noetherian, Δ_1 -a.c.c., sequentially Noetherian and Σ_1 -Noetherian respectively.

Definition 5.5. A ring R is Δ_1 -Noetherian if every Δ_1 -ideal $I \subseteq R$ is finitely generated.

Definition 5.6 (RCA₀). A ring R has the Δ_1 -a.c.c. if either of the following equivalent conditions holds:

- (ii) R has no strictly increasing chain of Δ_1 -ideals $I_0 \subsetneq I_1 \subsetneq \cdots$.
- (iii) Every chain of Δ_1 -ideals $I_0 \subseteq I_1 \subseteq \cdots$ in R eventually stabilises.

Definition 5.7. A ring R is sequentially Noetherian if it contains no sequence a_0, a_1, \ldots such that for all $i, a_i \notin (a_0, \ldots, a_{i-1})$.

Definition 5.8 (RCA₀ + $I\Sigma_2$). A ring R is Σ_1 -Noetherian if any of the following equivalent conditions holds:

- (v) For all a_0, a_1, \ldots in R, there is i such that $\{a_j : j \in \mathbb{N}\} \subseteq (a_0, \ldots, a_{i-1})$.
- (vi) R has no strictly increasing chain of Σ_1 -ideals $\mathcal{I}_0 \subsetneq \mathcal{I}_1 \subsetneq \cdots$.
- (vii) Every chain of Σ_1 -ideals $\mathcal{I}_0 \subseteq \mathcal{I}_1 \subseteq \cdots$ in R eventually stabilises.
- (viii) Every Σ_1 -ideal $\mathcal{I} \subseteq R$ is finitely generated.

We first establish the trivial (true in RCA₀) relations between the different notions of Noetherian.

Proposition 5.9 (RCA $_0$).

- (i) Every Σ_1 -Noetherian ring is Δ_1 -Noetherian.
- (ii) Every Σ_1 -Noetherian ring is sequentially Noetherian.




Figure 5.1: The different definitions of Noetherian, and logical implications between them over $RCA_0 + I\Sigma_2$. Green arrows are conjectured.

- (iii) Every sequentially Noetherian ring has the $\Delta_1\mbox{-a.c.c.}.$
- (iv) Every $\Sigma_1\text{-Noetherian ring has the }\Delta_1\text{-a.c.c.}.$

Proof.

- (i) Trivial from the last definition of $\Sigma_{1}\text{-Noetherian}.$
- (ii) Trivial from the first definition of Σ_1 -Noetherian.
- (iii) Repeat the construction in the proof of (ii)⇒(iii) on page 41. Note

that $(a_{j_k})_{k\in\mathbb{N}}$ is a "bad" sequence: $a_{j_\ell}\in I_{n_k}$ for $\ell\leq k$. Hence, we have $(a_{j_0},\ldots,a_{j_k})\subseteq I_{n_k}$, but $a_{j_{k+1}}\notin I_{n_k}$.

(iv) Follows from the previous two items.

 ACA_0 is enough to prove the equivalence of *all* the given definitions of Noetherian, so it is an upper bound for all arrows in Figure 5.1. We now determine the strength of the other implications.

Theorem 5.10 (RCA $_0$). The following are equivalent:

- (i) ACA₀.
- (ii) Every ring with the Δ_1 -a.c.c. is Σ_1 -Noetherian.
- (iii) Every ring with the Δ_1 -a.c.c. is Δ_1 -Noetherian.
- (iv) Every sequentially Noetherian ring is Σ_1 -Noetherian.
- (v) Every sequentially Noetherian ring is Δ_1 -Noetherian.

Proof. (i) \Rightarrow (ii) was observed above, and (ii) \Rightarrow (iii), (iii) \Rightarrow (v), (ii) \Rightarrow (iv), and (iv) \Rightarrow (v) are trivial in light of Proposition 5.9. Therefore, it just remains to prove (v) \Rightarrow (i). We will use the contrapositive of (v), and construct a computable ring R with a computable, non-finitely-generated ideal $I \subseteq R$, such that every "independent sequence" a_0, a_1, \ldots computes \varnothing '.

We use the ring R from [Con10, Thm 6.1]. In short, start with the ring $R_0 := \mathbb{Q}[\bar{x}]/(x_ix_j:i,j\in\mathbb{N})$, which consists solely of linear polynomials with the multiplication

$$\left(q + \sum a_i x_i\right) \left(r + \sum b_i x_i\right) = qr + \sum (ra_i + qb_i)x_i$$

Then, we enumerate A as in Lemma 3.24, and when we see n enter A, quotient R_0 by $x_n = kx_{n+1}$ for an appropriate choice of $k \in \mathbb{Q}$. We can choose the k in such a way to ensure the final ring R is computable—see [Con10] for details.

The ideal $I=(x_0,x_1,\ldots)\subseteq R$ is computable, as it consists of all polynomials with zero constant term, and not finitely generated since A is co-infinite. Now, given an independent sequence $a_0,a_1,\ldots\in R$, we can obtain an independent sequence $a'_0,a'_1,\ldots\in I$ as follows. Set $a'_i=a_i$ till we find a_n with nonzero constant term q. Then, for all $i\geq n$, we set $a'_i:=qa_{i+1}-r_{i+1}a_n$, where r_j is the constant term of a_j .

Now, I is an ω -dimensional vector space over \mathbb{Q} with basis $\{x_n : n \notin A\}$, and the a_i' are a linearly independent sequence in I. For each n, let f(n) be the largest variable appearing in a_0', \ldots, a_{n+1}' . By independence, f(n) must be greater than the nth element of A^{\complement} . Thus, f dominates $\mu_{\varnothing'}$, and so a_0', a_1', \ldots computes \varnothing' .

Theorem 5.11. WKL₀ is equivalent to "every ring with the Δ_1 -a.c.c. is sequentially Noetherian".

Proof. We will actually work with the contrapositive of the given statement, i.e. "if R has a sequence a_0, a_1, \ldots such that for all $i, a_i \notin (a_0, \ldots, a_{i-1})$, then R has a strictly increasing chain of Δ_1 -ideals".

- (⇒) Let $C \subseteq 2^{\omega}$ consist of the sequences of sets $I = (I_0, I_1, I_2, ...)$ such that
 - (a) For all $k \in \mathbb{N}$, I_k is an ideal.
 - (b) For all $k \in \mathbb{N}$, $I_k \subseteq I_{k+1}$.
 - (c) For all $k \in \mathbb{N}$, $a_k \in I_k$ but $a_{k+1} \notin I_k$.

By writing the above conditions in first-order logic, we can verify that C is a Π_1^0 class. Classically, C is nonempty, since it contains the sequence $(a_0) \subsetneq (a_0, a_1) \subsetneq (a_0, a_1, a_2) \subsetneq \cdots$. Therefore, WKL₀ gives a member of C, which is a strictly increasing chain by condition (c).⁵

⁵This can be done more rigorously à la Theorem 4.10, by building a computable tree $T \subseteq 2^{<\omega}$ such that $\mathcal{C} = [T]$, and verifying every level of T is nonempty.

(\Leftarrow) We build a computable ring R, with a computable bad sequence a_0, a_1, \ldots , such that every strictly increasing chain of computable ideals $I_0 \subsetneq I_1 \subsetneq \cdots$ is of PA degree. As in Theorem 4.10, fix disjoint c.e. sets A, B such that any separator has PA degree. Without loss of generality, we may assume that the complement of $A \cup B$ contains a computable increasing sequence a_0, a_1, \ldots, a_n

The construction is identical to [DLM07, Thm 3.2]. In short, we first set $R_0 := \mathbb{Z}[x_0, x_1, \ldots]$, and add in elements of its field of fractions $\operatorname{Frac}(R_0)$. Begin enumerating A and B, and:

- If n enters A, add to R all elements of the form $\frac{x_n}{p(x_0,\ldots,x_{n-1})}$.
- If n enters B, add to R all elements of the form $\frac{x_n-1}{p(x_0,\ldots,x_{n-1})}$.

Let $R \subseteq \operatorname{Frac}(\mathbb{Z}[\bar{x}])$ be the subring generated by all the above additions. Then, R is c.e., and hence computably isomorphic to a computable ring [DLM07, p. 8].

We claim the sequence $x_{n_0}, x_{n_1}, x_{n_2}, \ldots$ is bad. By Π_1 induction on i, we will show that for all $k \geq i$, $x_{n_k} \notin (x_{n_0}, \ldots, x_{n_{i-1}})$. The base case i = 0 follows since $x_{n_k} \notin \{0\}$. The case i = 1 follows since $n_k \notin A$, so $x_{n_k}/x_{n_0} \notin R$ and thus $x_{n_k} \notin (x_{n_0})$.

Now, assume the inductive hypothesis for i-1. Fixing $k \geq i$, we have, in particular, that $x_{n_k} \notin (x_{n_0}, \ldots, x_{n_{i-2}})$. By contradiction, we will show that $x_{n_k} \notin (x_{n_0}, \ldots, x_{n_{i-1}})$. That is, we suppose that $x_{n_k} \in (x_{n_0}, \ldots, x_{n_{i-1}})$, so we can write $x_{n_k} = \sum_{j < i} r_j x_{n_j}$ for some $r_j \in R$. We will rewrite $x_{n_k} = \sum_{j < i-1} r'_j x_{n_j}$ for some $r'_j \in R$, showing $x_{n_k} \in (x_{n_0}, \ldots, x_{n_{i-2}})$ and contradicting our inductive assumption.

As an arbitrary element of R, r_{i-1} must have the form

$$r_{i-1} = f + \sum_{\ell \in A'} g_{\ell} x_{\ell} / p_{\ell} + \sum_{\ell \in B'} h_{\ell} (x_{\ell} - 1) / q_{\ell}$$

⁶If not, redefine $A' = \{2a : a \in A\}$ and $B' = \{2b : b \in B\}$: then $1, 3, 5, \ldots$ is such a sequence.

where $f \in \mathbb{Z}[\bar{x}]$, $A' \subseteq A$ and $B' \subseteq B$ are finite, and p_{ℓ}, q_{ℓ} are elements of $\mathbb{Z}[x_0, \dots, x_{\ell-1}]$. We will show that each summand in r_{i-1} can be "moved" into a different r_j , hence we can write x_{n_k} as a linear combination of $x_{n_0}, \dots, x_{n_{i-2}}$. Let c_{ℓ}/d_{ℓ} be an arbitrary summand in r_{i-1} , for $c_{\ell}, d_{\ell} \in \mathbb{Z}[\bar{x}]$.

- If $d_{\ell} = d'_{\ell} x_{n_{i-1}}$ for some $d_{\ell} \in \mathbb{Z}[\bar{x}]$, then we can take c_{ℓ}/d_{ℓ} out of r_{i-1} , and put $c_{\ell}/d'_{\ell} x_{n_i}$ into r_j for some j < i-1.
- Otherwise, the term $(c_\ell/d_\ell)x_{n_{i-1}}$ contains a factor of $x_{n_{i-1}}$, which must be cancelled out by another r_j , j < i-1, since x_{n_k} contains no factor of $x_{n_{i-1}}$. Hence, $c_\ell = c'_\ell x_{n_j}$, and we can take c_ℓ/d_ℓ out of r_{i-1} , and put $c'_\ell x_{n_{i-1}}/d_\ell$ into r_j .

This rewrite shows that $x_{n_k} \in (x_{n_0}, \dots, x_{n_{i-2}})$, contradicting our inductive assumption. Hence, $x_{n_0}, x_{n_1}, x_{n_2}, \dots$ is a computable bad sequence, as required.

As shown in [DLM07, Thm 3.2], every nontrivial proper ideal of R has PA degree. If $I_0 \subsetneq I_1 \subsetneq \cdots$ is a strictly increasing chain, then I_1 is a nontrivial proper ideal of R: hence the chain has PA degree. \square

Conjecture 5.12 (RCA $_0$). The following are equivalent:

- (i) ACA₀.
- (ii) Every Δ_1 -Noetherian ring is Σ_1 -Noetherian.
- (iii) Every Δ_1 -Noetherian ring is sequentially Noetherian.
- (iv) Every Δ_1 -Noetherian ring has the Δ_1 -a.c.c..

We previously observed (i) \Rightarrow (ii), and by Proposition 5.9, (ii) \Rightarrow (iii) and (iii) \Rightarrow (iv) are trivial. Therefore, it just remains to prove (iv) \Rightarrow (i). Essentially, this would require us to construct a computable ring R with a uniformly computable, nonstabilising chain of ideals $I_0 \subseteq I_1 \subseteq I_2 \subseteq \cdots$, such

that every non-finitely-generated ideal $J \subseteq R$ computes \varnothing' . Following Conidis [Con10; Con14], one idea would be to take A as in Lemma 3.24, and construct a ring so that every non-finitely-generated ideal computes an infinite subset of A^{\complement} . However, we have not yet found a construction that works.

Noetherian rings have been studied before in reverse math, but the definitions have not been standard until now. For example, Conidis [Con10; Con19; Con21] took Δ_1 -a.c.c. as his definition of "Noetherian", while Simpson [Sim88] used Σ_1 -Noetherian. We hope the results of this section allow a finer analysis of Noetherian rings in reverse mathematics. In particular, one could analyse the reverse-mathematical strength of previously studied theorems, but with a different notion of Noetherian. For example, Conidis [Con10; Con19] proved that "every Artinian ring has the Δ_1 -a.c.c" is equivalent to WKL₀. From this result, we can deduce:

Corollary 5.13. WKL₀ is equivalent to

Proof. In one direction, given an Artinian ring R, WKL₀ proves that R has the Δ_1 -a.c.c. [Con10; Con19], and then that R is sequentially Noetherian by Theorem 5.11.

In the other direction, RCA₀ proves that sequentially Noetherian implies Δ_1 -a.c.c. (Proposition 5.9), hence over RCA₀, (*) implies "every Artinian ring has the Δ_1 -a.c.c", which implies WKL₀.

Simpson [Sim88] showed that, over RCA₀, the theorem "for every field K and $n \in \mathbb{N}$, $K[x_1, \cdots, x_n]$ is Σ_1 -Noetherian" is equivalent to $WO(\omega^\omega) :=$ " ω^ω is well-ordered". Proposition 5.9 shows that RCA₀ + $WO(\omega^\omega)$ also proves this statement for the other notions of Noetherian, but we don't know if these reverse. We would also like to see a study of the more general version of Hilbert's basis theorem: "if R is X-Noetherian, then R[x] is Y-Noetherian", where X, Y are chosen from our four notions of Noetherian.

5.1 The a.c.c.p.

Classically, there is a weakening of the Noetherian chain condition, which only requires every ascending chain of *principal* ideals to stabilise. This is called the *ascending chain condition on principal ideals*, or a.c.c.p. for short. The a.c.c.p. is of interest because it is often sufficient to prove many of the consequences of Noetherian-ness.

In RCA₀, there are two sensible notions of a.c.c.p., as noted by [Sat16]. Again, assuming RCA₀ + $I\Sigma_2$ as our base theory, we can show that they are equivalent.

Theorem 5.14 (RCA₀ + $I\Sigma_2$). The following are equivalent for a ring R:

- (i) There is no sequence a_0, a_1, \ldots in R such that $(a_0) \subsetneq (a_1) \subsetneq \cdots$.
- (ii) Every sequence $(a_0) \subseteq (a_1) \subseteq \cdots$ eventually stabilises.

Proof. (ii) \Rightarrow (i) is trivial. For (i) \Rightarrow (ii), given a nonstabilising sequence $(a_0) \subseteq (a_1) \subseteq \cdots$, we can use the same priority argument as in the proof of Theorem 5.4 to construct a strictly increasing sequence.

Definition 5.15 (RCA₀ + $I\Sigma_2$). R has the *a.c.c.p.* if the conditions of Theorem 5.14 are satisfied.

 $RCA_0 + I\Sigma_2$ clearly proves that a Σ_1 -Noetherian ring has the a.c.c.p.. ACA_0 appears to be necessary to show the other notions of Noetherian imply a.c.c.p.. However, we do not prove this—we leave it as an open question.

Chapter 6

Integral domains

Definition 6.1 (RCA₀). An *integral domain* is a ring R with no nonzero zero-divisors.

Integral domains are those rings satisfying one of the most basic laws of arithmetic—the cancellation of multiplication. There are a wide variety of different subclasses of integral domains, each generalising properties of our favourite rings: \mathbb{Z} , fields, polynomial rings, etc. These form a complex web of implications—an extensive diagram is shown on page 52.

The aim of this chapter is to initiate a systematic study of the different integral domain properties in reverse mathematics. Some special classes of integral domain have already seen study—Euclidean domains [DK11; Sat16], PIDs [Sat16], and UFDs [Bur13; GM17]. We extend this study by:

- considering additional classes of integral domains (Bézout and GCD domains).
- analysing reverse-mathematically the containments between these different classes of integral domains.

The new types of integral domains we consider often have several (classically) equivalent definitions. Hence, part of the analysis is to determine how hard it is to prove these equivalences, and if they are not equivalent in RCA_0 , to determine which is the right notion in RCA_0 .




Figure 6.1: Some of the different types of integral domain, and the logical implications between them [IDMap]. Modified and reproduced with permission.

First, we will review the existing reverse-mathematical work on integral domains. The first work in this area appears to be the study of Euclidean domains, initiated in [DK11] and continued in [Sat16, §6.4].

Definition 6.2 [DK11; Sat16]. An integral domain R is a *Euclidean domain* if there is a function $f: R \setminus \{0_R\} \to \mathbb{N}$ such that for all $a, d \in R$ with $d \neq 0_R$, there are $q, r \in R$ such that a = dq + r and f(r) < f(d) if $r \neq 0_R$.

[DK11] were concerned with the reverse mathematical strength of

every Euclidean domain has a *minimal* Euclidean function (MEF)

They determined that MEF proves ACA₀, and conjectured that MEF is equivalent to ACA₀⁺, the system consisting of ACA₀ plus the assertion that the ω th Turing jump of any set exists. Meanwhile, [Sat16] proved in RCA₀ that Euclidean domains satisfy a version of Bézout's lemma.

Proposition 6.3 (RCA₀). R/I is an integral domain iff $I \subseteq R$ is a prime ideal.

Proof. Trivial by the ideal correspondence theorem (Theorem 4.2). \Box

Definition 6.4. Let R be a ring, and fix $r, s \in R$.

- (i) r is a *unit* if there exists $s \in R$ such that $rs = 1_R$.
- (ii) r is *irreducible* if $r \neq 0_R$, $r \nmid 1_R$, and for all $a, b \in R$, whenever r = ab, then at least one of a, b is a unit.
- (iii) r is *prime* if $r \neq 0_R$, $r \nmid 1_R$, and if for all $a, b \in R$, whenever $r \mid ab$, then $r \mid a$ or $r \mid b$.
- (iv) r, s are associates (written $r \sim s$) if r = su for a unit $u \in R$.

These notions can also be characterised by conditions on the corresponding principal ideals, and these equivalences are trivially provable in RCA₀.

54

Proposition 6.5 (RCA₀).

- (i) r is a unit iff (r) = R.
- (ii) r is irreducible iff (r) is maximal among proper principal ideals.
- (iii) r is prime iff (r) is a nonzero proper prime ideal.
- (iv) In an integral domain, $r \sim s$ iff (r) = (s).

RCA₀ can prove basic properties of the associate relation:

Proposition 6.6 (RCA_0).

- (i) \sim is an equivalence relation.
- (ii) If $a \sim b$ and $c \sim d$, then $ac \sim bd$.
- (iii) In an integral domain, if $a \sim b$ and $ac \sim bd$, then $c \sim d$.

Proof.

- (i) Reflexivity follows since 1_R is a unit. For symmetry, if $a \sim b$, then a = bu where $uv = 1_r$, so b = buv = av, whence $b \sim a$. Transitivity follows since the product of units is a unit.
- (ii) We have a = bu, $c = dv \implies ac = bd(uv) \implies ac \sim bd$.
- (iii) We have a=bu, ac=bdv. Then $buc=bdv \implies uc=dv \implies c=d(u^{-1}v)$, so $c\sim d$.

RCA₀ can also prove many basic facts about primes and irreducibles. Henceforth, given elements $a, b \in R$, we will use " $(a/b) \in R$ " as a shorthand for "there exists $c \in R$ such that a = bc", and use "(a/b)" as a name for c.

Proposition 6.7 (RCA $_0$).

(i) If p is prime and $p \mid a_1 \cdots a_n$, then there is $j \leq n$ such that $p \mid a_j$.

- (ii) If r is irreducible, a not a unit, and $a \mid r$, then $a \sim r$.
- (iii) In an integral domain R, every prime element is irreducible. *Proof.*
 - (i) Fixing a_1, a_2, \ldots , we proceed by induction on

$$\varphi(n) := (p \mid a_1 \cdots a_n \to (\exists j \le n)(p \mid a_j))$$

This formula is $\Sigma_1 \to \Sigma_1$, so by Theorem 3.8, RCA $_0$ can carry out this induction.

- (ii) Since $a \mid r$, we have $(r/a) \in R$ and r = a(r/a). Then, (r/a) must be a unit since r is irreducible.
- (iii) Suppose p is prime and p=ab. Then, $p\mid ab$ in particular, so $p\mid a$ or $p\mid b$ by assumption. WLOG, suppose $p\mid a$, so $(a/p)\in R$. Then, a=p(a/p)=ab(a/p). Hence, b(a/p)=1 and b is a unit.

The converse of Proposition (iii) is not true in general. Integral domains for which the converse holds are called *AP domains*.¹

Definition 6.8. An integral domain R is an AP domain if every irreducible element in R is prime.

6.1 Bézout and GCD domains

[Sat16, Thm 6.34] proved in RCA₀ that a version of Bézout's lemma holds in Euclidean domains. This inspired us to take up a reverse-mathematical analysis of *Bézout domains*—those in which Bézout's identity holds. However, we will first analyse the weaker notion of *GCD domains*—those in which every pair of elements has a gcd. GCD domains can also be characterised by existence of lcms, or in terms of ideals.

¹AP is an abbreviation for "Atoms are Prime", "atom" being an older term for "irreducible element".

Definition 6.9. Let R be a ring, and fix $b \in R$ and a finite subset $A \subseteq R$.

- (i) *b* is the *greatest common divisor* of *A*, written b = gcd(A), if:
 - (a) For all $a \in A$, $b \mid a$, and
 - (b) For every c satisfying property (i)(a), $c \mid b$.
- (ii) *b* is the *least common multiple* of *A*, written b = lcm(A), if:
 - (a) For all $a \in A$, $a \mid b$, and
 - (b) For every c satisfying property (ii)(a), $b \mid c$.

We first need a simple lemma about gcds.

Lemma 6.10 (RCA₀). For all $a, b, c \in R$, if gcd(a, b) and gcd(ac, bc) both exist, then $gcd(ac, bc) = gcd(a, b) \cdot c$.

Proof. Let $d := \gcd(a,b)$ and $e := \gcd(ac,bc)$: we will show that $e \sim dc$. In one direction, since $d \mid a,b$, we have $dc \mid ac,bc \implies dc \mid e$. This implies $(e/dc) \in R$. Conversely, since $e \mid ac,bc$, we have $ed \mid adc,bdc \implies (e/dc)d \mid a,b$. Therefore, $(e/dc)d \mid d$, so $e \mid dc$ as required.

Theorem 6.11 (RCA $_0$). For a ring R, the following are equivalent:

- (i) Any two elements of *R* have a gcd.
- (ii) Any two elements of R have an lcm.
- (iii) For all a, b, there is a unique minimal principal Σ_1 -ideal containing (a, b).
- (iv) The intersection of two principal Σ_1 -ideals is principal.

Proof.

- (i) \Rightarrow (ii): Let $d = \gcd(a, b)$. In particular, $d \mid a, b$, so $(a/d), (b/d) \in R$. We claim $a(b/d) = \operatorname{lcm}(a, b)$. Clearly $a \mid a(b/d)$, but also a(b/d)d = ab = (a/d)bd, so $b \mid a(b/d) = (a/d)b$.
 - Now, suppose $c \in R$ is such that $a, b \mid c$. Then $ab \mid ac, bc$, so $ab \mid \gcd(ac, bc) = dc$ by Lemma 6.10. Hence $a(b/d) \mid c$ as required.
- (ii) \Rightarrow (i): Let $\ell = \text{lcm}(a, b)$. In particular, $a, b \mid \ell$, so $(\ell/a), (\ell/b) \in R$. Also, since $a \mid ab$ and $b \mid ab$, we have $\ell \mid ab$ by definition, so $(ab/\ell) \in R$.

We claim $(ab/\ell) = \gcd(a,b)$. We have $(ab/\ell)(\ell/b)b = ab$, which implies $(ab/\ell)(\ell/b) = a$, so $(ab/\ell) \mid a$, and similarly $(ab/\ell) \mid b$. Now, suppose $e \mid a,b \implies (a/e), (b/e) \in R$. Clearly $a \mid a(b/e)$, but also $a(b/e)e = ab = (a/e)be \implies a(b/e) = (a/e)b$, hence $b \mid a(b/e) = (a/e)b$. By definition of ℓ , $\ell \mid a(b/e) \implies e\ell \mid ab = (ab/\ell)\ell \implies e \mid (ab/\ell)$ as required.

(i) \Leftrightarrow (iii) and (ii) \Leftrightarrow (iv) are straightforward since $a \mid b \iff (b) \subseteq (a)$. We have

$$(a,b) \subseteq (\gcd(a,b))$$
 and $(a) \cap (b) = (\operatorname{lcm}(a,b))$

Definition 6.12. An integral domain R is a GCD domain if R satisfies any of the conditions in Theorem 6.11.

We can extend each of the conditions in Theorem 6.11 from two elements to an arbitrary finite number, but in each case, we seem to require Σ_3 induction.

Proposition 6.13 (RCA₀ + $I\Sigma_3$). An integral domain R is a GCD domain iff any of the following holds:

- (v) Any finite subset of *R* has a gcd.
- (vi) Any finite subset of R has an lcm.
- (vii) Every finitely generated Σ_1 -ideal in R is contained in a unique minimal principal Σ_1 -ideal.

58

(viii) Finite intersections of principal Σ_1 -ideals are principal.

Proof. The equivalences (i) \Leftrightarrow (v), (ii) \Leftrightarrow (vi), (iii) \Leftrightarrow (vii), (iv) \Leftrightarrow (viii) all follow by induction. In each case, the formula we induct over is Σ_3 .

GCD domains also satisfy one of the most fundamental properties of PIDs and UFDs: every irreducible element is prime.

Proposition 6.14 (RCA₀). GCD domains are AP domains.

Proof. Suppose R is a GCD domain, $r \in R$ is irreducible, and $r \mid ab$. Let $d = \gcd(ar, ab)$. In particular, we have $d \mid ar \implies (ar/d) \in R$. The case ab = 0 is trivial, so suppose $ab \neq 0$. Then $a, b \neq 0$ since we are in an integral domain, and hence $d \neq 0$.

Now, since $a \mid ar$, $a \mid ab$ and d is the gcd, we must have $a \mid d$, so $(d/a) \in R$. Then, $dr = (d/a)ar = (d/a)(ar/d)d \implies r = (d/a)(ar/d)$. Since r is irreducible, either (ar/d) or (d/a) is a unit. If (ar/d) is a unit, then $ar \sim d$, so $ar \mid ab \implies r \mid b$. If (d/a) is a unit, then $a \sim d$, and $r \mid ar, ab \implies r \mid d \implies r \mid a$.

Bézout domains are a special type of GCD domains, in which the gcd of A is required to be a *linear combination* of elements of A. There are several definitions of Bézout, which turn out to be equivalent in RCA₀. In particular, this additional condition lowers the amount of induction required to prove the equivalence between the case |A|=2 and the case for arbitrary finite A. Thus, the equivalence between these two cases goes through in RCA₀.

Theorem 6.15. For a commutative ring R with unity, the following are equivalent over RCA₀:

- (i) Every pair $a, b \in R$ has a gcd d, and there are $x, y \in R$ s.t. ax + by = d.
- (ii) Every finite set $A \subseteq R$ has a gcd d, and $d \in (A)$.
- (iii) For all $a, b \in R$, there exists d such that (a) + (b) = (d).

- (iv) Every finitely generated Σ_1 -ideal is principal. *Proof.*
- (i) \Rightarrow (iii): Suppose $d \mid a, d \mid b$ (i.e. $(a/d), (b/d) \in R$), and d = ax + by, for some $x, y \in R$. We claim that (a) + (b) = (d). Picking some $ak + b\ell \in (a) + (b)$, we have

$$ak + b\ell = d(a/d)k + d(b/d)\ell = d((a/d)k + (b/d)\ell) \in (d)$$

Conversely, picking $dk \in (d)$, we have

$$dk = (ax + by)k = a(xk) + b(yk) \in (a) + (b)$$

(iii) \Rightarrow (iv): Fix a finitely generated Σ_1 -ideal $\mathcal{I} = (a_1, \dots, a_n)$. We induct on the formula

$$\varphi(k) = (\exists s) [(a_1, \dots, a_k) = (s_k)]$$

which is Σ_1 , since it can be written as

$$\varphi(k) = (\exists s)(\exists b_1, \dots, b_k, c_1, \dots, c_k)$$
$$(s = b_1 a_1 + \dots + b_k a_k \land (\forall j \le k)(a_j = c_j s))$$

- $\varphi(0)$ is witnessed by $s=0_R$, and $\varphi(1)$ by $s=a_1$. By induction, assume $(a_1,\ldots,a_k)=(s')$. Then, by assumption, there is s such that $(s')+(a_{k+1})=(s)$, whence $(a_1,\ldots,a_k,a_{k+1})=(s)$.
- (iv) \Rightarrow (ii): The ideal (A) is finitely generated, so by assumption, fix d such that (A) = (d). We claim $d = \gcd(A)$. For every $a \in A$, we have $a \in (A) = (d)$, hence $d \mid a$, so d is indeed a common divisor of A.

Since $d \in (d) = (A)$, this immediately implies there exist $b_1, \ldots, b_n \in R$ such that $d = a_1b_1 + \cdots + a_nb_n$, where $A = \{a_1, \ldots, a_n\}$. It then follows that $d = \gcd(A)$. Fix any other common divisor $e \mid A$: then $(a_i/e) \in R$ for each $i \leq n$. Hence

$$d = \sum_{i=1}^{n} a_i b_i = \sum_{i=1}^{n} e(a_i/e)b_i = e \cdot \sum_{i=1}^{n} (a_i/e)b_i$$

so $e \mid d$ as required.

$$(ii)\Rightarrow (i)$$
: Trivial.

Definition 6.16. An integral domain R is $B\'{e}zout$ if any of the equivalent conditions in Theorem 6.15 holds.

From definition (ii) of Bézout and (v) of GCD domain, it follows trivially (in RCA_0) that any Bézout domain is a GCD domain.

6.2 UFDs and PIDs

In this section, we discuss two important types of integral domains: *unique* factorisation domains (UFDs) and principal ideal domains (PIDs). These were some of the earliest types of integral domains considered, and they both have rich and well-developed theories with a lot of overlap. Consequently, we have had to order the results in this section carefully to make it clear that there is no circularity.

6.2.1 UFDs

In what follows, let $[n] := \{1, \dots, n\}$.

Definition 6.17. An integral domain R is a *unique factorisation domain* (UFD) if every nonzero element $r \in R$ can be written $r = uq_1 \cdots q_n$ for a unit u and irreducibles q_1, \ldots, q_n , and this factorisation is unique, i.e. for any other irreducible factorisation $r = u'q'_1 \cdots q'_m$, we have n = m, and there is a bijective map $h: [n] \to [n]$ such that for all $i \le n$, q_i and $q'_{h(i)}$ are associates.

UFDs can also be characterised in terms of the existence of *prime* factorisations, and then uniqueness automatically follows. RCA₀ can prove this characterisation, but first we need some lemmas about UFDs.

Lemma 6.18 (RCA $_0$). UFDs are AP domains.

Proof. Let $q \in R$ be irreducible: we will show q is prime. If $q \mid ab$, then $(ab/q) \in R$. Substituting in the unique factorisation of (ab/q) into q(ab/q) = ab, it follows that q is in the unique factorisation of ab. Multiplying the factorisations of a and b respectively gives another factorisation of ab. Hence, up to units, q is in the unique factorisation of a or b, so $q \mid a$ or $q \mid b$ as required.

Lemma 6.19 (RCA₀). In an integral domain R, suppose that p_1, \ldots, p_n are prime, q_1, \ldots, q_m are irreducible, and $p_1 \cdots p_n \sim q_1 \cdots q_m$. Then, n = m, and there is a bijective map $h \colon [n] \to [n]$ such that for all $i \le n$, p_i and $q_{h(i)}$ are associates.

Proof. Fixing prime p_1, \ldots, p_n and irreducible q_1, \ldots, q_m with $p_1 \cdots p_n \sim q_1 \cdots q_m$, we induct up to n on the Σ_1 formula

$$\varphi(k) = (\exists \text{ injective } h \colon [k] \to [m]) (\forall i \leq n) (p_i \sim q_{h(i)})$$

See [Bur13, p. 50] for details.

Theorem 6.20 (RCA₀). For an integral domain R, the following are equivalent.

- (i) *R* is a UFD, in the sense of Definition 6.17.
- (ii) Every nonzero element $r \in R$ factors into primes and a unit.

Proof.

- (i) \Rightarrow (ii): Follows from Lemma 6.18.
- (ii)⇒(i): By Proposition 6.7.(iii), a prime factorisation is an irreducible factorisation. Uniqueness follows from Lemma 6.19.

By Lemma 6.18, we will use "prime" and "irreducible" interchangeably when discussing UFDs.

Proposition 6.21. UFDs are GCD domains.

Proof. Suppose R is a UFD, and fix finite $A \subseteq R$. By assumption, each $a \in A$ has a unique factorisation $a = up_1 \cdots p_n$. By bounded Σ_1 comprehension, for each $a \in A$, RCA₀ can recursively define finite sets $A_0 \supseteq \cdots \supseteq A_n$ by $A_0 = \{1, \ldots, n\}$ and

$$A_{k+1} = A_k \setminus \{j \in A_k : (\exists b \in R)(p_j = b \cdot p_{\min A_k})\}$$

Letting $q_i = p_{\min A_{i-1}}$ and $b_i = |A_i \setminus A_{i-1}|$ as long as $A_{i-1} \neq \emptyset$, we can instead factorise $a = uq_1^{b_1} \cdots q_m^{b_m}$ for non-associate primes q_1, \ldots, q_m and $b_i > 0$. Then $\gcd(A) = q_1^{c_1} \cdots q_m^{c_m}$ where c_i is the minimum of the b_i for all $a \in A$.

In fact, the above proof shows that UFDs satisfy the stronger definition of GCD domain (Proposition 6.13), without the assumption of $I\Sigma_3$. Thus, we can freely assume the existence of arbitrary gcds in a UFD.

6.2.2 PIDs

[Sat16] furthermore considered *principal ideal domains* (PIDs)—integral domains in which every ideal is principal. As mentioned in the previous section, the most natural notion of ideal for PIDs is Σ_1 -ideal, since the principal ideal (a) is Σ_1 in general, and may not be computable for every a.

Definition 6.22 [Sat16]. An integral domain R is a $(\Sigma_1$ -)PID if every Σ_1 -ideal $\mathcal{I} \subseteq R$ is principal.

Right from the definition, we see that RCA_0 proves every PID is Bézout (using definition (iv) of Bézout), and that every PID is Σ_1 -Noetherian. Recall that Σ_1 -Noetherian was the strongest notion of Noetherian, and so RCA_0 also proves that PIDs satisfy all the other definitions of Noetherian, and that all PIDs have the a.c.c.p..

Classically, every PID is a UFD. The usual proof can be broken down into four steps:

(i) Show every PID is a Noetherian AP domain.

- (ii) Show every element r has an irreducible factor.
- (iii) Then, show r can be written as a product of irreducibles.
- (iv) Show that any prime factorisation of r is unique up to order and units.

RCA₀ can carry out step (i): this is Theorem 6.36.(ii), along with the observation that any PID is Noetherian. RCA₀ can also carry out step (iv): this is Lemma 6.19. The usual proofs of steps (ii) and (iii) require us to recognise when an element is irreducible or a unit, hence they require ACA₀ (as these conditions are Π_2 and Σ_1 respectively). [Bur13; GM17] showed that ACA₀ is necessary for step (iii):

Theorem 6.23 [Bur13; GM17]. ACA₀ is equivalent to "in a ring with a.c.c.p., every element has an irreducible factorisation".

We have not yet determined the reverse mathematical strength of step (ii), i.e. "in a ring with a.c.c.p., every element has an irreducible *factor*". We believe that it could be shown equivalent to ACA₀, using a similar argument to [GM17, Thm 1.2], but using perfect binary trees instead of "fishbones". However, the details have not been worked through at the time of writing.

For completeness, here is a proof in ACA₀ that all PIDs are UFDs.

Corollary 6.24 (ACA $_0$). Every PID is a UFD.

Proof. We saw that RCA_0 proves every PID R is AP and Noetherian (hence has the a.c.c.p.). By Theorem 6.23, every element of R factors into irreducibles. By AP-ness, this is also a prime factorisation, hence R is a UFD by Theorem 6.20.

6.2.3 Equivalent definitions of PIDs

PIDs can be classically characterised in two alternative ways. One is the existence of a *Dedekind–Hasse norm*, a slight generalisation of a Euclidean

norm. The other is the (ostensibly weaker) requirement that only every *prime* ideal is principal. We now show that this latter characterisation is provable in ACA_0 .

Lemma 6.25 (RCA₀). Suppose $I \subseteq R$ is an ideal, and $a \notin I$. If (I, a) and I : a are both principal, then I is principal.

Proof. Suppose (I,a)=(b) and I:a=(c). Since (I,a)=(b), we have $(a/b) \in R$, and there are $d \in R$, $i \in I$ such that b=i+ad. We will in fact prove that I=(bc). To see that $bc \in I$, note that $c \in I:a$, so $ac \in I$. Then, $bc=(i+ad)c=ic+(ac)d \in I$.

To see $I \subseteq (bc)$, pick $r \in I$. Then, $r \in (I, a) = (b)$, so $(r/b) \in R$. Since $r \in I$, we get $r(a/b) = (r/b)b(a/b) = (r/b)a \in I$, whence $(r/b) \in I : a = (c)$. Writing $(r/b) = c\ell$, we have $r = bk = bc\ell \in (bc)$.

Theorem 6.26 (ACA₀). Suppose R is an integral domain in which every prime Σ_1 -ideal is principal. Then, R is a Σ_1 -PID.

Proof. By contrapositive. Suppose R is not a PID: then there is a nonprincipal Σ_1 -ideal $\mathcal{I} \subseteq R$. We will construct a nonprincipal prime ideal $\mathcal{P} \supseteq \mathcal{I}$.

Fix a standard listing of all pairs $(a,b) \in R^2$. By recursion, we simultaneously build a tree $T \subseteq 2^{<\omega}$ and associate every finite binary string $\sigma \in T$ with a Σ_1 -ideal \mathcal{I}_σ . As we construct T, we will ensure that for every $\sigma \in T$, \mathcal{I}_σ is nonprincipal. To begin, we let $T = \{\varepsilon\}$ and $\mathcal{I}_\varepsilon := \mathcal{I}$. Now, given $\sigma \in T$, there are two cases:

- (i) If \mathcal{I}_{σ} is prime, set $\mathcal{P} := \mathcal{I}_{\sigma}$, and stop the construction here—we are done.
- (ii) Otherwise, look for the first pair $(a,b) \in R^2$ such that $ab \in \mathcal{I}_{\sigma}$ but $a,b \notin \mathcal{I}_{\sigma}$. Then set $\mathcal{I}_{\sigma \cap 0} := (\mathcal{I}_{\sigma},a)$ and $\mathcal{I}_{\sigma \cap 1} := \mathcal{I}_{\sigma} : a$. Note that $a \in (\mathcal{I}_{\sigma},a)$ and $b \in \mathcal{I}_{\sigma} : a$. Put $\sigma \cap 0$ into T iff (\mathcal{I}_{σ},a) is nonprincipal, and put $\sigma \cap 1$ into T iff $\mathcal{I}_{\sigma} : a$ is nonprincipal.

Note that all the \mathcal{I}_{σ} are Σ_1 relative to \mathcal{I} . To tell whether a $\Sigma_1^{\mathcal{I}}$ ideal is prime is $\Pi_2^{\mathcal{I}}$, and telling if one is principal is $\Sigma_3^{\mathcal{I}}$. ACA₀ proves the existence of \mathcal{I}''' , which is powerful enough to carry out the construction of T and the \mathcal{I}_{σ} . Thus, ACA₀ proves that T and the \mathcal{I}_{σ} exist.

Now, assume that case (i) never happened. The resulting set T is indeed a tree, and by Lemma 6.25, every $\sigma \in T$ has a successor in T. By induction, it follows that T is infinite. By WKL₀, take a path $\alpha \in [T]$, and define

$$\mathcal{P} = \bigcup_{n \in \mathbb{N}} \mathcal{I}_{lpha \restriction_n}$$

 \mathcal{P} must be a prime ideal, since if there were $a,b \in R$ such that $ab \in \mathcal{P}$ but $a,b \notin \mathcal{P}$, we would have forced $a \in \mathcal{P}$ or $b \in \mathcal{P}$ at some stage of the construction. Furthermore, \mathcal{P} is nonprincipal, since if $\mathcal{P} = (p)$, then $p \in \mathcal{I}_{\alpha \restriction_n}$ for some n, whence $\mathcal{I}_{\alpha \restriction_n} = (p)$. The theorem follows.

Now, we consider the characterisation of PIDs in terms of Dedekind–Hasse norms, and show that this is equivalent to ACA_0 over RCA_0 .

Definition 6.27. A *Dedekind–Hasse norm* on an integral domain R is a function $f: R \to \mathbb{N}$ such that:

- (i) $f(r) = 0 \iff r = 0_R$.
- (ii) For all nonzero $a, b \in R$, either $b \mid a$ or there exist $x, y \in R$ such that 0 < f(ax + by) < f(b).
- (iii) For all nonzero $a, b \in R$, $f(a) \le f(ab)$.

A *Dedekind–Hasse domain* (DHD) is an integral domain which admits a Dedekind–Hasse norm.

Item (iii) is not always included in the definition, since given a function $f: R \to \mathbb{N}$ satisfying just (i) and (ii), we can define $f': R \to \mathbb{N}$ satisfying all three by $f'(r) = \min\{f(ra) : a \neq 0_R\}$. However, this process is not computable, and hence we must assert condition (iii).

Classically, a ring is a DHD if and only if it is a PID. One direction of this equivalence is provable in RCA₀:

Theorem 6.28. RCA₀ proves "every DHD is a PID".

Proof [Hen13]. Suppose (R, f) is a DHD, and $\mathcal{I} \subseteq R$ a nonzero Σ_1 -ideal. The image $A := f(\mathcal{I} \setminus \{0\})$ is a nonempty c.e. subset of \mathbb{N} . By $\mathsf{L}\Sigma_1$, A has a least element n. Then, we enumerate \mathcal{I} till we find $b \neq 0_R$ with f(b) = n.

We claim $\mathcal{I}=(b)$. Pick nonzero $a\in I$. Note that for all $x,y\in R$, $ax+by\in I$, so we can't have 0< f(ax+by)< f(b) by choice of b. It follows that $b\mid a$.

Every Euclidean norm is a Dedekind-Hasse norm (choosing $x=1_R$ every time), and hence:

Corollary 6.29 [Sat16]. RCA₀ proves "every Euclidean domain is a PID".

However, ACA_0 is needed for the converse of this theorem. First, we show that it can be proved in ACA_0 .

Theorem 6.30 (ACA₀). Every Σ_1 -PID is a DHD.

Proof [Hen13]. As we saw in Theorem 6.24, ACA₀ proves that every PID is a UFD. In ACA₀, we can tell which elements are irreducible/prime (as this is Π_2^0), so given an element $r \in R$, simply search for its factorisation $p_1 \cdots p_k$. We define $f: R \to \mathbb{N}$ by mapping 0_R to 0, and $r \neq 0_R$ to k+1, where k is the number of irreducibles in the factorisation of r.

We claim f is a Dedekind-Hasse norm. (i) is true by definition, and (iii) follows since f(ab) = f(a) + f(b). For (ii), since every PID is Bézout, $d = \gcd(a,b)$ can be written as a linear combination of a and b. Then, if $b \sim q_1 \cdots q_k \nmid a \sim p_1 \cdots p_n$, then there is some q_j not associate to any p_i . Hence, f(d) < f(b) as required.

For the reversal, we need to construct a computable PID R so that every DHN on R computes \varnothing' . We want to use our usual method of coding a c.e.

set $A \subseteq \mathbb{N}$ into the polynomial ring $\mathbb{Q}[\bar{x}]$. Unfortunately, $\mathbb{Q}[\bar{x}]$ is not a PID. However, we can take a localisation to make it a PID, and still retain the ability to code using the x_i 's. First, we need a lemma about a certain partial order, which we will use in the construction.

Lemma 6.31 (RCA₀). Let \mathbb{N}^{fin} be the collection of sequences in $\mathbb{N}^{\mathbb{N}}$ which are eventually zero. Define a partial order \leq on $\mathbb{N}^{\mathbb{N}}$ by $\alpha \leq \beta \iff \forall i \ \alpha_i \leq \beta_i$. Then, $(\mathbb{N}^{\text{fin}}, \leq)$ has a meet for every nonempty Σ_1 subset.

Proof. Let $A \subseteq \mathbb{N}^{\text{fin}}$ be nonempty. Essentially, the meet α of A is defined by $\alpha_i = \min\{\beta_i : \beta \in A\}$, but it takes some work to show that this exists in RCA₀.

Formally, we will construct α by viewing it as a function $\mathbb{N} \to \mathbb{N}$, i.e. a set of pairs. Since A is nonempty, fix some $\gamma \in A$. Since $\gamma \in \mathbb{N}^{\mathrm{fin}}$, let $n \in \mathbb{N}$ be such that $(\forall m \geq n)(\gamma_m = 0)$. By Δ_0 comprehension, let $K_0 \coloneqq \{(m,a) : m < n, a < \gamma_m\}$. Via bounded Σ_1 comprehension (Lemma 3.12), we can define the following finite subsets of K_0 :

$$K_{=} := \left\{ (m, a) \in K_0 : (\exists \beta \in A)(a = \beta_m) \right\}$$

$$K_{>} := \left\{ (m, a) \in K_0 : (\exists \beta \in A)(a > \beta_m) \right\}$$

Again by Δ_0 comprehension, let $K := K_= \setminus K_>$. We claim K is a function $n \to \mathbb{N}$: if $(m, a), (m, a') \in K$, then there are $\beta, \beta' \in A$ such that $a = \beta_m$, $a' = \beta'_m$, but for all $\delta \in A$, $a, a' \le \delta_m$. In particular, $a \le \beta'_m = a'$ and $a' \le \beta_m = a$, hence a = a'.

Now, fixing m < n, the set $A_m = \{b : (\exists \beta \in A)(b = \beta_m)\}$ is Σ_1 , so it has a least element a_m by $\mathsf{L}\Sigma_1$. Then, $(m,a_m) \in K$. Hence, K is a function $n \to \mathbb{N}$. We finally define α by

$$\alpha := K \cup \{(m,0) : m \ge n\}$$

and this is a function $\mathbb{N} \to \mathbb{N}$, and an element of \mathbb{N}^{fin} . By definition, $\alpha = \bigwedge A$.

In fact, \mathbb{N}^{fin} has a meet for *every* nonempty subset, but RCA₀ only has enough induction to prove this for Σ_1 subsets.

Before proving the next proposition, we observe that every polynomial $p \in R[\bar{x}]$ can be written as $p = \sum_{\alpha \in F} c_{\alpha} x^{\alpha}$ for a unique choice of finite $F \subseteq \mathbb{N}^{\text{fin}}$ and coefficients $c_{\alpha} \neq 0$ (where $x^{\alpha} \coloneqq \prod x_i^{\alpha_i}$ for $\alpha \in \mathbb{N}^{\text{fin}}$). We will call F the *support* of p, and denote it supp(p).

Proposition 6.32 (RCA₀). There is a computable ring $T \subseteq \operatorname{Frac}(\mathbb{Q}[\bar{x}])$ so that:

- (i) T is a Σ_1 -PID.
- (ii) Every x_i is not a unit in T.
- (iii) For all $i \neq j \in \mathbb{N}$, we have $x_i \nmid x_j$.

Proof. Let $M \subseteq \mathbb{Q}[\bar{x}]$ be the set $M := \mathbb{Q}[\bar{x}] \setminus \bigcup_{i=0}^{\infty}(x_i)$. M is multiplicatively closed: take $p, q \in M$. By assumption, $\operatorname{supp}(p)$ contains at least one element α with $\alpha_i = 0$. Among these, take the ones with maximal α_0 , then of those, the ones with maximal α_1 , etc., until we obtain a unique element α . Do the same to obtain a "lexicographically maximal" element $\beta \in \operatorname{supp}(q)$.

We claim that the pointwise sum $\alpha + \beta \in \operatorname{supp}(pq)$. Otherwise, the cross term $c_{\alpha}d_{\beta}x^{\alpha+\beta}$ would have to be cancelled out by another term $c_{\gamma}d_{\delta}x^{\gamma+\delta}$, with $\gamma \neq \alpha$, $\delta \neq \beta$, $\gamma + \delta = \alpha + \beta$. But then either γ would have to be lexicographically above α , or δ above β —contradiction. Hence, pq has a term $c_{\alpha}d_{\beta}x^{\alpha+\beta}$ not containing x_i . As this holds for all $i \in \mathbb{N}$, $pq \in M$.

Let T be the localisation of $\mathbb{Q}[\bar{x}]$ at M. We claim T satisfies all the conclusions of the proposition. For (iii), note that the elements of T have the form p/m for $p \in \mathbb{Q}[\bar{x}]$, $m \in M$. Now, consider the product $x_i p/m$. If $x_i p/m = x_j$ for $j \neq i$, then $x_i p = x_j m$. However, this is not possible, since m is not divisible by x_i . This also implies that (ii) holds.

Now, we show T is a Σ_1 -PID. Given $p \in \mathbb{Q}[\bar{x}]$, let $\beta := \bigwedge \operatorname{supp}(p)$. Then, $p = x^{\beta}m$ for some $m \in M$. Thus, up to units, every element of T is a product of x_i 's, i.e. a monic monomial. Note also that there is a bijective

map between these monic monomials and \mathbb{N}^{fin} (as defined in Lemma 6.31), where $\alpha \in \mathbb{N}^{\text{fin}}$ corresponds to x^{α} . Furthermore, the relation \leq on \mathbb{N}^{fin} corresponds exactly to the divisibility order on T.

Given a nonzero Σ_1 -ideal $I \subseteq T$, let $X_I \coloneqq \{\alpha \in \mathbb{N}^{\text{fin}} : x^\alpha \in I\}$. Since $X_I \subseteq \mathbb{N}^{\text{fin}}$ is nonempty and Σ_1 , we can define $\alpha \coloneqq \bigwedge X_I$ by Lemma 6.31. Now, we claim $I = (x^\alpha)$. The \subseteq direction follows since x^α divides everything in I.

Conversely, we show $x^{\alpha} \in I$. Let n be such that $\alpha_m = 0$ for all $m \geq n$. By definition of α , for every i < n, there is $\alpha^{(i)} \in X_I$ such that $\alpha_i^{(i)} = \alpha_i$. Now, considering the sum $\sum_{i < n} x^{\alpha^{(i)}}$, we can factor out x^{α} , and we are left with an element m of M. Multiplying by 1/m gives $x^{\alpha} \in I$.

T is very useful in reversals of theorems about PIDs, since we can code into the x_i as we usually would. Here's a simple example.

Theorem 6.33. There is a PID so that the set of units computes \emptyset' .

Proof. Localise
$$T$$
 at $\{x_n : n \in \emptyset'\}$. Then, $n \in \emptyset' \iff x_n$ is a unit. \square

Now, we can complete the reversal of "every PID is a DHD".

Lemma 6.34 (RCA₀). If (R, f) is a DHD, then $r \in R$ is a unit iff $f(r) = f(1_R)$.

Proof. First, since $f(a) \le f(ab)$, we have $f(1_R) \le f(1_R r) = f(r)$ for any $r \ne 0_R$, hence 1_R has minimal norm among nonzero elements of R.

- (⇒) If $rs = 1_R$, then $f(r) \le f(rs) = f(1_R)$. Hence, $f(r) = f(1_R)$ by minimality of $f(1_R)$.
- (⇐) Suppose $f(r) = f(1_R)$. If $r \nmid 1_R$, then there are $x, y \in R$ such that $0 < f(x + ry) < f(r) = f(1_R)$, contradicting minimality of $f(1_R)$. \square

Corollary 6.35. "Every PID is a DHD" implies ACA₀.

Proof. Let R be the PID in Theorem 6.33. By assumption, let f be a DHN on R. By Lemma 6.34, f computes the set of units of R, which computes \varnothing' .

In Theorem 6.33, we constructed a computable PID R so that every DHN on R computes \varnothing' (by computing the units of R). However, note that the collection of DHNs on a PID is a Π_2 set in Baire space. Thus, we expect that this result is not optimal in terms of computability, i.e. one could likely find a computable PID so that every DHN computes some $X >_{\mathbb{T}} \varnothing'$. However, we leave this question open.

Another possible direction is to analyse the strength of "every PID is a DHD" for the weakened notion of DHN without condition (iii). Unfortunately, Lemma 6.34 fails badly in this case, and it is not clear that such a DHN can determine the units. Hence, one would need a different way of getting computational power from a DHN.

6.2.4 Theorems about PIDs

[Sat16] proved several basic results about PIDs in RCA₀:

Theorem 6.36 [Sat16, §6.4]. The following are provable in RCA₀:

- (i) Every Euclidean domain is a Σ_1 -PID.
- (ii) Every PID is an AP domain.
- (iii) For *a* irreducible/prime in a PID, (*a*) is a maximal Δ_1 -ideal.

Corollary 6.37. In a PID, every nonzero prime ideal is maximal.

Proof. If $\mathcal{P} = (p)$ is a nonzero prime ideal, then p is prime, so the result follows by Theorem 6.36.(iii).

Here, we prove some more results concerning PIDs. Notice that in the proof of Theorem 6.33, x_n is irreducible/prime iff $n \notin \emptyset'$, hence we get a computable PID in which the primes are Π_1 complete. Thus, as a corollary, we get:

Corollary 6.38 (RCA₀). The following are equivalent:

- (i) ACA₀.
- (ii) For any PID R, the set of units of R exists.
- (iii) For any PID R, the set of primes of R exists.

Since the set of irreducibles in any ring is Π_2 in general, we would expect that this complexity for the primes is not optimal. Indeed, we can improve it to Π_2 complete:

Theorem 6.39. There is a computable PID whose set of irreducibles is Π_2 complete.

Proof. We will build a polynomial ring R with variables x_e , $e \in \mathbb{N}$, such that x_e is prime iff W_e is infinite. Hence, we will have a many-one reduction from the primes of R to $Inf = \{e : W_e \text{ is infinite}\}$, which is Π_2 complete.

Start with $R_0 := \mathbb{Q}[x_i, y_{i,0}, z_{i,0} : i \in \mathbb{N}]/(x_i - y_{i,0}z_{i,0} : i \in \mathbb{N})$ localised at

$$\left[\bigcup(y_{i,0})\cup\bigcup(z_{i,0})\right]^\complement$$

Now, to build R, we begin enumerating all W_e in parallel. When a new element enters W_e at stage s, we do the following:

- (i) Localise at $y_{e,t}$, where $t = \max\{u : y_{e,u} \in R \text{ at stage } s\}$.
- (ii) Freely add elements $y_{e,s}$, $z_{e,s}$ to R, i.e. let $R_{\text{new}} = R_{\text{old}}[y_{e,s}, z_{e,s}]$.
- (iii) Set $x_e = y_{e,s} z_{e,s}$, i.e. quotient R by $(x_e y_{e,s} z_{e,s})$.
- (iv) Localise at $M = \{ p \in R : p \text{ contains } y_{e,s} \text{ or } z_{e,s} \text{ but } p \notin (y_{e,s}) \cup (z_{e,s}) \}.$

Note that for any ring S and element $s \in S$, the natural map $S \to S[y,z]/(s-yz)$ is injective. Hence, combining steps (ii) and (iii) above, we can consider this as a proper expansion of R. Since we only add elements, and never remove/quotient any, it follows that R is c.e.. Thus, we may assume R is computable by Theorem 3.26.

We claim R is a PID. R is a localisation of $\mathbb{Q}[x_e,y_{e,s},z_{e,s}:\langle e,s\rangle\in K]/(x_e-y_{e,s}z_{e,s}:\langle e,s\rangle\in K)$ for some c.e. set K. By step (iv), the non-units of R are all contained in

$$\bigcup_{\langle e,s\rangle\in K} (y_{e,s}) \cup (z_{e,s})$$

So, as in Proposition 6.32, every element of R is (up to a unit) a product of the $y_{e,s}$ and $z_{e,s}$, so R is a PID by the same argument.

For each x_e , its only possible nontrivial splittings are $y_{e,s}z_{e,s}$ where $\langle e,s\rangle\in K$. If W_e is finite, then the last $y_{e,s}$ we add will never be made a unit; hence x_e is properly reducible into $y_{e,s}z_{e,s}$. Conversely, if W_e is infinite, every $y_{e,s}$ we add will eventually be made a unit, so x_e is irreducible.

There is a well-known characterisation of when the polynomial ring R[x] is a PID: this is exactly when R is a field. In fact, R[x] is famously a Euclidean domain in this case. This characterisation is also provable in RCA₀, but first we need the following technical lemma about polynomial division.

Lemma 6.40 (RCA₀). Suppose R is a field, and fix $p, d \neq 0 \in R[x]$ with $deg(p) \geq deg(d)$. Then, there are polynomials $q, r \in R[x]$ such that

- (i) p = dq + r;
- (ii) $\deg(q) \leq \deg(p) \deg(d)$;
- (iii) $\deg(r) < \deg(d)$.

Proof. Write

$$p = a_n x^n + \cdots + a_m x^m + \cdots + a_1 x + a_0$$

$$d = b_m x^m + \cdots + b_1 x + b_0$$

The claim is that there are

$$q = c_{n-m}x^{n-m} + \cdots + c_1x + c_0$$

 $r = e_{m-1}x^{m-1} + \cdots + e_1x + e_0$

with p = dq + r.

Substituting the above expressions into equation (i), we get a system of n+1 linear equations in n+1 variables e_i , c_i , with coefficients in a_i , b_i (see Figure A.1). The corresponding $(n+1)\times(n+1)$ matrix \mathbf{A} is upper-triangular (see Figure A.1), and all entries on the main diagonal are nonzero (since $b_m \neq 0_R$). Thus, we can obtain a solution for e_i , c_i by computing $\mathbf{A}^{-1}\mathbf{a}$, where \mathbf{a} is the (n+1)-vector of coefficients of p. See Appendix \mathbf{A} .

Theorem 6.41 (RCA₀). For an integral domain R, the following are equivalent:

- (i) R is a field.
- (ii) R[x] is a Euclidean domain.
- (iii) R[x] is a Σ_1 -PID.

Proof.

- (i) \Rightarrow (ii): We claim the degree function deg is a Euclidean function on R[x]. Pick $p, p' \in R[x]$ with $p' \neq 0$. There are two cases:
 - $\deg(p) < \deg(p')$: then q := 0, r := p satisfy the definition of Euclidean.
 - $deg(p) \ge deg(p')$: follows from Lemma 6.40.
- (ii)⇒(iii): Already proven (Corollary 6.29).
- (iii) \Rightarrow (i): Pick $a \in R$ and consider the Σ_1 -ideal (a,x) in R[x]. By assumption, there is $b \in R[x]$ such that (b) = (a,x). We must have $b \in R$, else (b) = (a,x) could not contain constants.

Since $x \in (b)$, there is a polynomial $p \in R[x]$ such that bp = x. Now, p must be linear, so $p = cx + d \implies (bc)x + bd = x$. Matching coefficients, we must have $bc = 1_R$, d = 0.

Hence, b is a unit and (b) = (a, x) = R[x]. In particular, $1_R \in (a, x)$, so there are polynomials $q, r \in R[x]$ such that $1_R = aq + xr$. Write q = q'x + k for $k \in R$; then $1_R = ak + x(r + aq')$. Again matching coefficients, we get r + aq' = 0, $ak = 1_R$. Thus, a is a unit. \Box

6.2.5 Gauss' lemma

Now, we return to our study of UFDs. An important theorem about UFDs is that R is a UFD if and only if R[x] is one. One direction is easy: if R[x] is a UFD, then every $r \in R$ has a factorisation in R[x], but by degree considerations, this must actually be a factorisation in R. The other direction is nontrivial, and we analyse its proof here.

Definition 6.42. Let R be a GCD domain, and $p \in R[x]$, $p \neq 0$. The *content* of p, cont(p) is the gcd of its coefficients. We say p is primitive if cont(p) = 1.

For any nonzero $p \in R[x]$, we can take c = cont(p) and factor p = cp' to get a primitive polynomial p'.

Before we prove the theorem, we need a series of lemmas. The name *Gauss' lemma* is commonly given to any of these lemmas.

Lemma 6.43 (RCA₀; [Sin11, Lem 12.1.8]). Let R be a UFD, and K the field of fractions of R. Fix $p, q \in R[x]$.

- (i) If p, q are primitive, then so is pq.
- (ii) $cont(pq) \sim cont(p) cont(q)$.
- (iii) If p is primitive and $q \mid p$, then q is primitive.
- (iv) If p is primitive, then the following are equivalent:
 - (a) p is irreducible in R[x].
 - (b) p is irreducible in K[x].
 - (c) p is prime in R[x].

(d) p is prime in K[x].

Proof.

- (i) Fix a prime $r \in R$. Since p,q are primitive, both must have some term not divisible by r. Picking the terms $a_n x^n$ in p and $b_m x^m$ in q of maximal degree with this property (by $L\Pi_1$), the x^{n+m} term in pq can't be divisible by r either. As this is true for all primes r, we must have cont(pq) = 1.
- (ii) Let $c_p = \text{cont}(p)$ and $c_q = \text{cont}(q)$, and write $p = c_p p'$, $q = c_q q'$. Then $pq = c_p c_q p' q'$, and since p' q' is primitive by (i), the result follows.
- (iii) From (ii), we get $cont(q) \mid cont(p) \mid 1$ as required.
- (iv) (a) \Rightarrow (b): Suppose p is primitive and irreducible in R[x]. We have $\operatorname{cont}(p)=1$ but p is not a unit, so $\deg(p)>0$. By contradiction, suppose p is reducible as p=rs for non-units $r,s\in K[x]$ (i.e. $\deg(r),\deg(s)>0$). By clearing denominators and dividing off contents, we can find $a,b,c,d\in R$ and primitive $r',s'\in R[x]$ so that r=(a/b)r',s=(c/d)s'.

Then bdp = acr's', so taking contents and using that p, r', s' are primitive, (ii) gives $bd \sim ac$, $p \sim r's'$. Now we've properly factored p in R[x], contradicting irreducibility of p.

- (b) \Rightarrow (d): K[x] is a PID by Theorem 6.41, so AP by Theorem 6.36.(ii).
- (d) \Rightarrow (c): Suppose p is primitive and prime in K[x]. Fix $r,s \in R[x]$ such that $p \mid rs$ in R[x]. Then, $p \mid rs$ in K[x] too, so by primeness, $p \mid r$ or $p \mid s$ in K[x]. Without loss of generality, suppose $p \mid r$ in K[x], i.e. r = ph for $h \in K[x]$. Write r = cr' for $c = \operatorname{cont}(r)$, r' primitive. Clearing denominators and taking contents, write h = (a/b)h' for $a, b \in R$, $h' \in R[x]$ primitive.

Then bcr' = aph', so taking contents and using that r', p, h' are primitive, (ii) gives $bc \sim a, r' \sim ph'$. Now we see $r = cr' \sim p(ch')$ and $ch' \in R[x]$, so p is prime in R[x].

(c)
$$\Rightarrow$$
(a): By Proposition 6.7.(iii).

Theorem 6.44 (RCA₀ + $I\Sigma_3$). If R is a UFD, then R[x] is a UFD.

Proof. We proceed by strong induction on

$$\varphi(n) = (\forall p) \big(\deg p = n \land p \text{ primitive}$$

 $\to (\exists u \text{ unit}) (\exists q_1, \dots, q_m \text{ prime}) (p = uq_1 \cdots q_m) \big)$

By Lemma 6.43, the q_i must be primitive, so we can say they are prime/irreducible in a Π_1 way: they can't be factored into polynomials of strictly lower degree. Hence, φ is a Π_3 formula.

If deg(p) = 0, then since p is primitive, it is a unit, so we are done. So suppose deg(p) > 0. If p is irreducible, then it is prime by Lemma 6.43.(iv). Otherwise, p is properly reducible into non-units q, r. By Lemma 6.43.(iii), q, r are primitive, so by induction, they have prime factorisations. The product of these factorisations gives the required factorisation of p.

We have proven that every *primitive* $p \in R[x]$ has a prime factorisation. This implies *every* $p \in R[x]$ has a prime factorisation, since we can just factor out the content of p as p = cp', and combine the factorisations of c and p'. Now the result follows from Theorem 6.20.

One possible strategy to reduce the amount of induction required for Theorem 6.44 would be to find a "nice" coding of R[x], and do strong induction on the code for p. This would take the induction down to $I\Sigma_2$. For this to work, we would need a coding $c\colon R[x]\cong \mathbb{N}$ such that whenever p=qr for non-units q,r, we have c(q),c(r)< c(p).

The problem is that, for general UFDs, this coding can't be computable. This is because it would allow us to compute the irreducibles of R[x] from \varnothing' , because we could simply check all q < p to find factorisations of p, and ask \varnothing' if q is a unit. [DM18] constructed a UFD R such that $\operatorname{Irr}(R)$ is Π_2 -complete, and since $\operatorname{Irr}(R) = R \cap \operatorname{Irr}\left(R[x]\right)$, it follows that $\operatorname{Irr}\left(R[x]\right)$ is also is Π_2 -complete. For this ring, such a coding must join \varnothing' above \varnothing'' ; in particular, it can't be computable (even from \varnothing').

More generally, it seems the only way to reduce the induction in Theorem 6.44 is to bound the quantifiers on p and q_i . However, if we could bound these, then we could compute the factorisation of any element by a finite search of all elements less than the bound. Essentially, reducing the complexity would require us to *a priori* "know" the factorisation of p, which we cannot expect in general.

Chapter 7

Conclusion

In this thesis, we used the tools of reverse mathematics to analyse several topics in ring theory, particularly radicals, Noetherian rings and integral domains. Every theorem we analysed turned out to be provable in ACA_0 , including Theorem 6.26, the usual proof of which uses Zorn's lemma. We also showed that some key results in commutative algebra actually require ACA_0 (i.e. they are equivalent to ACA_0 over RCA_0):

- The equivalence of several notions of Noetherian (Theorem 5.10).
- Every PID admits a Dedekind–Hasse norm (Corollary 6.35).

Furthermore, we expect that most of the other results that were proven in ACA_0 will turn out to be equivalent to ACA_0 . This suggests that ACA_0 is the right axiom system in which to develop countable commutative algebra.

The majority of the results we studied were provable even in RCA_0 . Some were provable in RCA_0 with extra induction:

- The equivalence of weak and strict chain conditions for Σ_1 -ideals (Theorem 5.4, $I\Sigma_2$).
- If R is a GCD domain, then every finite subset has a gcd (Proposition 6.13, $|\Sigma_3\rangle$).

• If *R* is a UFD, then so is R[x] (Theorem 6.44, $I\Sigma_3$).

Hence, if one had philosophical objections to working in a nonconstructive system such as ACA₀, we can still develop the majority of countable commutative algebra "computably" in RCA₀ (+ $I\Sigma_3$), including most basic facts about integral domains (§6). However, RCA₀ still can't prove some important results such as the existence of irreducible factorisations under a.c.c.p. (Theorem 6.23), and the equivalence of different definitions of Noetherian (Theorems 5.10 and 5.11).

Finally, our work has given rise to many open questions. Two particularly interesting, related problems are reversing Corollary 6.24 (PIDs are UFDs) and Theorem 6.26 (all prime Σ_1 -ideals principal \Longrightarrow PID) in ACA₀. The most obvious way is to construct a computable non-UFD in which every enumeration of a nonprincipal ideal computes \varnothing' (resp. a computable non- Σ_1 -PID in which every enumeration of a nonprincipal prime ideal computes \varnothing'). To do this, we need to be able to force complexity on nonprincipal ideals, so that they require \varnothing' to be enumerated. We could do this, for example, by forcing the nonprincipal ideals to be Π_1 -complete - it seems like novel techniques would be needed to do this.

Here are some of the other problems we'd particularly like to see solved:

- Determine the exact reverse-mathematical strength of RAD (page 36).
- Determine whether $I\Sigma_2$ is necessary for Theorem 5.4.
- Prove Conjecture 5.12.
- Use reverse mathematics to analyse important results about Bézout and GCD domains, such as:
 - *R* is Bézout iff it is a *Prüfer* GCD domain.
 - The following are equivalent for a GCD domain: UFD, a.c.c.p,
 Noetherian, atomic.

- The following are equivalent for a Bézout domain: PID, Noetherian, UFD, a.c.c.p., atomic.
- Determine whether $\mathrm{I}\Sigma_3$ is necessary for Theorem 6.44.

Appendix A

Linear algebra

This appendix is devoted to the proof of Lemma 6.40, and proving in RCA₀ the necessary theorems of linear algebra.

We will only need to consider square matrices.

Definition A.1. Let K be a field. An $n \times n$ matrix **A** over K is an array of elements of K:

$$\begin{bmatrix} A_{1,1} & A_{1,2} & \cdots & A_{1,n} \\ A_{2,1} & A_{2,2} & \cdots & A_{2,n} \\ \vdots & \vdots & \ddots & \vdots \\ A_{n,1} & A_{n,2} & \cdots & A_{n,n} \end{bmatrix}$$

Notationally, we will use the same letter to refer to a matrix and its elements, but the matrix A will be in boldface, while its entries $A_{i,j}$ will be italicised. The definitions of matrix multiplication, identity matrix, invertible matrix are as usual. Furthermore, the usual proofs of associativity of matrix multiplication, uniqueness of inverses, etc. go through in RCA₀.

Definition A.2. Let **A** be an $n \times n$ matrix, and $i, j \leq n$. The *minor submatrix* $\mathbf{A}_{i,j}$ is the $(n-1) \times (n-1)$ matrix obtained from **A** by removing the ith row and jth column.

Definition A.3. The *determinant* of an $n \times n$ matrix **A** is given inductively

on n. If n = 1, then $det(\mathbf{A}) = A_{1,1}$. If n > 1, then

$$\det(\mathbf{A}) := \sum_{i=1}^{n} (-1)^{n+i} A_{n,i} \det(\mathbf{A}_{n,i})$$

Although tedious, the usual proof of the Laplace expansion theorem goes through in RCA₀: that is, we could equally well have done cofactor expansion along a different row/column to define the determinant.

Lemma A.4 (RCA₀). If $det(\mathbf{A}) \neq 0_K$, then **A** is invertible.

Proof. As usual, we define the adjugate of **A** as the matrix **B** such that $B_{i,j} = (-1)^{i+j} \det(\mathbf{A}_{j,i})$. Then, we show that $\mathbf{AB} = \mathbf{BA} = \det(\mathbf{A})\mathbf{I}$: the proof uses the aforementioned Laplace expansion theorem.

Definition A.5. A matrix **A** is upper-triangular if $A_{i,j} = 0$ for all i > j.

Lemma A.6 (RCA₀). The determinant of an upper-triangular $n \times n$ matrix **A** is $A_{1,1} \cdot A_{2,2} \cdots A_{n,n}$.

Proof. Fix **A**, and for all $k \leq n$, let

$$\mathbf{A}^{(k)} := \begin{bmatrix} A_{1,1} & A_{1,2} & \cdots & A_{1,k} \\ A_{2,1} & A_{2,2} & \cdots & A_{2,k} \\ \vdots & \vdots & \ddots & \vdots \\ A_{k,1} & A_{k,2} & \cdots & A_{k,k} \end{bmatrix}$$

By Σ_0 induction on k, we prove that for all $k \leq n$, $\det(\mathbf{A}^{(k)}) = A_{1,1} \cdot A_{2,2} \cdots A_{k,k}$. The result follows taking k = n.

The k=1 case follows directly from the definition of determinant. Now, suppose $\det (\mathbf{A}^{(k)}) = A_{1,1} \cdot A_{2,2} \cdots A_{k,k}$. By definition,

$$\det \left(\mathbf{A}^{(k+1)} \right) = A_{k+1,1} \det \left(\mathbf{A}_{k+1,1}^{(k+1)} \right) + \dots + A_{k+1,k+1} \det \left(\mathbf{A}_{k+1,k+1}^{(k+1)} \right)$$

Since **A** is upper-triangular, all terms except the last are zero. Hence, $\det (\mathbf{A}^{(k+1)}) = A_{k+1,k+1} \det (\mathbf{A}^{(k+1)}_{k+1,k+1})$. However, $\mathbf{A}^{(k+1)}_{k+1,k+1} = \mathbf{A}^{(k)}$, whence the claim follows.

Corollary A.7 (RCA₀). Let **A** be an upper-triangular matrix. If every $A_{i,i} \neq 0_K$, then **A** is invertible.

Proof. By Lemma A.6, $det(\mathbf{A}) \neq 0_K$, so the result follows from A.4.

Corollary A.7 is enough to prove Lemma 6.40. The relevant system of equations and matrix are shown on the following page.

Figure A.1: The system of linear equations and corresponding matrix obtained from the proof of Lemma 6.40, in the case when 2m > n. The case $2m \le n$ looks similar.

Appendix B

Zorn's lemma

The motivation for this section originally came from Theorem 6.26. All of the proofs we could find used Zorn's lemma in an essential way; hence, we wondered if it was possible to formalise those arguments in second-order arithmetic. One might be tempted to say no, since Zorn's lemma for suborders of $\langle \mathcal{P}(\mathbb{N}), \subseteq \rangle$ seems to have an essential third-order quality. Our idea was to pull back the inclusion relation along an indexing of, say, the c.e. sets, reducing a third-order problem to a second-order one. As it turns out, this "pull-back" doesn't work, and we'll discuss why below.

However, supposing it did work, we would reduce the problem to a second-order version of Zorn's lemma. That is, given a set A at some level Γ of the arithmetical hierarchy (Γ depends on the complexity of the original index set) and a partial order \preccurlyeq at some level Θ of the arithmetical hierarchy (depending on the complexity of the sets being indexed), we want to show Zorn's lemma holds for (A, \preccurlyeq) .

We now proceed to the formal development in second-order arithmetic. The definition of partial orders (on subsets of \mathbb{N}) is as usual. For a partial ordering \leq , \leq will denote the corresponding strict relation. \leq denotes the usual order relation on \mathbb{N} .

Definition B.1. Let (A, \preceq) be a partial order.

- (i) A *chain* in (A, \preccurlyeq) is a function $f : \mathbb{N} \to A$ which is \prec -increasing: for all $n, f(n) \prec f(n+1)$.
- (ii) Given a chain f in (A, \preccurlyeq) , an *upper bound* for f is an element $a \in A$ such that $f(n) \preccurlyeq a$ (equivalently, $f(n) \prec a$) for all n.
- (iii) A maximal element $a \in (A, \preceq)$ is one such that there is no $b \in A$ with $a \prec b$.

Definition B.2. Let Γ and Θ be classesof subsets of \mathbb{N} and \mathbb{N}^2 , respectively. $\mathsf{ZL}(\Gamma,\Theta)$ is the following statement: for every Γ subset $A\subseteq\mathbb{N}$ and Θ relation \preccurlyeq on \mathbb{N} , if \preccurlyeq is a partial order, and every chain $f\colon\mathbb{N}\to(A,\preccurlyeq)$ has an upper bound, then (A,\preccurlyeq) has a maximal element.

Evidently, if $\Gamma \subseteq \Gamma'$ and $\Theta \subseteq \Theta'$, then $\mathsf{ZL}(\Gamma', \Theta') \vDash \mathsf{ZL}(\Gamma, \Theta)$. Here is the standard proof of arithmetical Zorn's lemma in ACA_0 .

Theorem B.3. ACA₀ proves $ZL(\Delta_0^1, \Delta_0^1)$.

Proof. Suppose A and \leq are Δ_0^1 (arithmetical), and \leq is a partial order. By contradiction, suppose (A, \leq) has no maximal element. We will construct a chain $f: \mathbb{N} \to A$ with no upper bound.

Fix an enumeration $\langle a_0, a_1, \ldots \rangle$ of A. We define f by recursion, starting with $f(0) = a_0$. If $f(n) = a_k$ has been defined, then search for the next $\ell > k$ such that $a_\ell \succ a_k$, and set $f(n+1) = a_\ell$. Since a_k is not maximal, we know we will always find $a_\ell \succ a_k$.

Now, we claim f has no upper bound in A. Suppose it did have an upper bound a_k (i.e. $f(n) \prec a_k$ for all n). Then, we would have found a_k at some stage of constructing f, and thus set $f(j) = a_k$ for some j. But $a_k \not\prec a_k$, so this is a contradiction.

Essentially the same proof shows that $RCA_0 \models ZL(\Sigma_1, \Delta_1)$. With some care, we can improve this to show:

Theorem B.4. RCA₀ proves $\mathsf{ZL}(\Sigma_1, \Sigma_1)$.

Proof. Suppose $A = \{a_0, a_1, \ldots\}$ is c.e., and $\leq \bigcup_n \leq_n$ is a c.e. partial order. We define f by recursion, starting with $f(0) = a_0$. Then, for $i = 0, 1, \ldots$, we continue searching through pairs $\langle n, k \rangle$ to find one such that $a_n \succ_k f(i)$, and set $f(i+1) = a_n$. Since f(i) is not maximal, we know there is $a_n \succ f(i)$, and this will be revealed at some finite stage k. Hence, f is total. As in the proof of Theorem B.3, f has no upper bound in f.

Theorem B.4 is optimal, in a sense:

Theorem B.5 (RCA_0).

- (i) If the usual order relation \leq is Θ , then $\mathsf{ZL}(\Pi_1,\Theta)$ implies ACA_0 .
- (ii) If \mathbb{N} is Γ , then $\mathsf{ZL}(\Gamma, \Pi_1)$ implies ACA_0 .

Proof.

- (i) Let A be as in Lemma 3.24. We consider the carrier set A^{\complement} (which is Π_1) under the usual order relation \leq . Then, since A^{\complement} is infinite, it has no maximal element, but any chain in A^{\complement} computes \varnothing' .
- (ii) We take $A=\mathbb{N}$ and build a Π_1 partial order \preccurlyeq on \mathbb{N} , using a "block merging strategy". The blocks B_n will be intervals in (\mathbb{N}, \leq) such that all $b \in B_n$ are \preccurlyeq -incomparable, and $b \prec c$ for all $b \in B_n$, $c > \max(B_n)$. We begin with $B_n = n$, i.e. $\preccurlyeq = \leq$.

To merge blocks B_n , B_{n+1} means to remove from \leq all pairs (b,c), where $b \in B_n$ and $c \in B_{n+1}$. We enumerate \varnothing' , and if we see n enter \varnothing' at stage s, and $s \in B_k$ for some k > n, then we merge the blocks B_n, \ldots, B_k . So, in the final partial order, we will have $a \leq b$ iff a is in a strictly earlier block than b.

More formally, in RCA₀ we can define markers $m_{n,s}$ by recursion on s, where $m_{n,s}$ marks the start of B_n at stage s. To begin, $m_{n,0} = n$, and when we see n enter \varnothing' at stage s, we find the least j with $m_{j,s} > s$, and set $m_{n+1,s+1} = m_{j,s}$, $m_{n+2,s+1} = m_{j+1,s}$, etc. Then, let $a \leq b \iff (\forall s)(\exists n < b)(a < m_{n,s} \leq b)$, which is Π_1 .

By bounded Σ_1 comprehension, RCA₀ can prove the existence of $\varnothing' \upharpoonright_n$ and $\varnothing'_s \upharpoonright_n$ for every s and n. Hence, the formula

$$\varphi(n) \; \coloneqq \; (\exists s)(\forall m < n)(m \in \varnothing'_s {\restriction}_n \iff m \in \varnothing' {\restriction}_n)$$

is Σ_1 , so RCA₀ can prove $(\forall n) \varphi(n)$ by induction. Now if $\varnothing' \upharpoonright_n$ has stabilised at stage s, it follows that m_n will henceforth be fixed, so RCA₀ proves all the m_n stabilise and all the B_n are finite.

 (\mathbb{N}, \preceq) has no maximal element, since for every $n \in B_k$, we have $n \prec m$ for any $m \in B_{k+1}$. By $\mathsf{ZL}(\Gamma, \Pi_1)$, let f be a chain in (\mathbb{N}, \preceq) . By Π_1 induction, we can prove that f(n) is in block B_n or higher, by inducting on

$$\varphi(n) := (\forall s)[f(n) > m_{n,s}]$$

Hence, for all n, $f(n+1) \ge \mu_{\varnothing'}(n)$, and so f computes \varnothing' .

Corollary B.6 (RCA₀). $ZL(\Gamma, \Theta)$ is equivalent to ACA₀ if

- (i) $\Gamma \supseteq \Pi_1$ and \leq is Θ , or
- (ii) \mathbb{N} is Γ and $\Theta \supseteq \Pi_1$.

Now, say we are working in a model \mathcal{M} of second-order arithmetic, and have a collection of Σ_1 -ideals which we are trying to apply Zorn's lemma to, e.g.:

$$\mathcal{K} = {\mathcal{I} : \mathcal{I} \text{ is a nonprincipal } \Sigma_1\text{-ideal}}$$

Being Σ_1 in \mathcal{M} , each element of \mathcal{K} has an enumeration which exists in \mathcal{M} . So, the idea would be to index all the enumerations in \mathcal{M} , and pull back along the indexing to obtain a first-order partial ordering, to which $\mathsf{ZL}(\Gamma,\Theta)$ can be applied. The problem is that we cannot index all the possible enumerations in the model, as there may be uncountably many, e.g. when $\mathcal{M} = \mathcal{P}(\omega)$ is the full ω -model.

We could attempt to fix this using an *internal* notion of computability. The idea is we have a universal Σ_1 formula $\varphi(e, n, X)$ such that for all Σ_1 formulae $\psi(n, X)$, we can (in RCA₀) find e such that

$$(\forall X)(\forall n) (\varphi(e, n, X) \leftrightarrow \psi(n, X))$$

Defining $W_e^X := \{n: \varphi(e,n,X)\}$ (but not necessarily assuming this set exists), and given some nonprincipal Σ_1 -ideal \mathcal{I} , we can then look at the set

$$K = \{e : W_e^{\mathcal{I}} \text{ is a nonprincipal ideal}\}$$

which is $\Pi_3^{\mathcal{I}}$. We define a relation on K by $e \preccurlyeq e' \iff W_e^{\mathcal{I}} \subseteq W_{e'}^{\mathcal{I}}$, which is $\Pi_2^{\mathcal{I}}$.

So, it seems that we have successfully reduced the problem to a second-order one. However, now another problem arises: since \leq is $\Pi_2^{\mathcal{I}}$, the chains in K that we are trying to defeat are no longer \mathcal{I} -computable, but only \mathcal{I}'' -computable. We can define an internal notion of \mathcal{I}'' -computability, but the union of (internally) \mathcal{I} -c.e. sets indexed by an (internally) \mathcal{I}'' -computable function may not be (internally) \mathcal{I} -c.e. itself—in general, it will only be (internally) $\Sigma_3^{\mathcal{I}}$.

So, to ensure closure under \mathcal{I}'' -computable chains, we could instead look at indices for $\Sigma_3^{\mathcal{I}}$ nonprincipal ideals. However, now the inclusion relation is $\Pi_4^{\mathcal{I}}$, so the chains we need to defeat are $\mathcal{I}^{(4)}$ -computable, so we would need to pass to $\Sigma_5^{\mathcal{I}}$ nonprincipal ideals to ensure closure. One can see that we will never be able to "catch our tail".

It is disappointing that the principles $\mathsf{ZL}(\Gamma,\Theta)$ don't seem to be applicable in reverse mathematics as we might have hoped. Nonetheless, we have left the results in this appendix, as we think they are interesting in their own right. It would be interesting to look at the statements $\mathsf{ZL}(\Gamma,\Theta)$ from the perspective of *Weihrauch reducibility* [BGP21], where one could obtain a more fine-grained analysis than the crude classification we gave in Theorem B.4 and Corollary B.6.

Bibliography

- [AM94] M. F. Atiyah, I. G. MacDonald. *Introduction to Commutative Algebra*. Avalon Publishing, 1994.
- [BGP21] Vasco Brattka, Guido Gherardi, Arno Pauly. Weihrauch Complexity in Computable Analysis. pp. 367–417 in: Vasco Brattka, Peter Hertling (eds.). Handbook of Computability and Complexity in Analysis. Theory and Applications of Computability. Springer, Cham, 2021. arXiv: 1707.03202.
- [Bir40] Garrett Birkhoff. *Lattice Theory*. Colloquium Publications **25**. American Mathematical Society, Providence, RI, 1940.
- [Bol32] János Bolyai. Appendix scientiam spatii absolute veram exhibens. In: Farkas Bolyai. Tentamen juventutem studiosam in elementa matheseos purae. Kali, 1832.
- [Boo54] George Boole. *An Investigation of the Laws of Thought*. Walton and Maberly, London, 1854.
- [Bur13] Valentin B. Bura. Reverse mathematics of divisibility in integral domains. Masters thesis. Victoria University of Wellington, NZ, 2013. HDL: 10063/2719.
- [Can74] Georg Cantor. *Ueber eine Eigenschaft des Inbegriffs aller reellen algebraischen Zahlen*. Journal für die reine und angewandte Mathematik 77, pp. 258–262, 1874.
- [Can83] Georg Cantor. *Ueber unendliche, lineare Punktmannichfaltigkeiten*. Mathematische Annalen **21(4)**, pp. 545–591, 1883.

[Con10] Chris J. Conidis. *Chain conditions in computable rings*. Transactions of the American Mathematical Society **362(12)**, pp. 6523–6550, 2010.

- [Con14] Chris J. Conidis. *Infinite dimensional proper subspaces of computable vector spaces*. Journal of Algebra **406**, pp. 346–375, 2014.
- [Con19] Chris J. Conidis. *The computability, definability, and proof theory of Artinian rings*. Advances in Mathematics **341**, pp. 1–39, 2019.
- [Con21] Chris J. Conidis. Computability theoretic aspects of an antichain theorem for extendible trees of nontrivial rank. 2021. URL: https://www.math.csi.cuny.edu/~conidis/papers/TAC.pdf.
- [Ded71] Richard Dedekind. Supplement X. In: P. G. L. Dirichlet. Vorlesungen über Zahlentheorie. 2nd ed. Vieweg, Braunschweig, 1871.
- [DeM47] Augustus De Morgan. *Formal Logic*. Taylor and Walton, London, 1847.
- [DK11] Rodney G. Downey, Asher M. Kach. *Euclidean functions of computable Euclidean domains*. Notre Dame Journal of Formal Logic **52(2)**, pp. 163–172, 2011.
- [DLM07] Rodney G. Downey, Steffen Lempp, Joseph R. Mileti. *Ideals in computable rings*. Journal of Algebra **314(2)**, pp. 872–887, 2007.
- [DM18] Damir D. Dzhafarov, Joseph R. Mileti. *The complexity of primes in computable unique factorization domains*. Notre Dame Journal of Formal Logic **59(2)**, pp. 139–156, 2018. arXiv: 1407.2626.
- [Eve69] Howard Eves. *An Introduction to the History of Mathematics*. 3rd ed. Holt, Rinehart and Winston, 1969.
- [FB58] Abraham A. Fraenkel, Yehoshua Bar-Hillel. *Foundations of Set Theory*. North Holland, Amsterdam, 1958.

[Fra15] Adolf Fraenkel. Über die Teiler der Null und die Zerlegung von Ringen. Journal für die reine und angewandte Mathematik 1915(145), pp. 139–176, 1915.

- [Fri67] Harvey Friedman. Subsystems of set theory and analysis. PhD thesis. MIT, 1967. HDL: 1721.1/33486.
- [Fri69] Harvey Friedman. *Bar induction and* Π_1^1 -CA. The Journal of Symbolic Logic **34(3)**, pp. 353–362, 1969.
- [Fri74] Harvey Friedman. Some systems of second order arithmetic and their use. pp. 235–242 in: Proceedings of the International Congress of Mathematicians. Vancouver, 1974.
- [FSS83] Harvey Friedman, Stephen G. Simpson, Rick L. Smith. *Countable algebra and set existence axioms*. Annals of Pure and Applied Logic **25(2)**, pp. 141–181, 1983.
- [FSS85] Harvey M. Friedman, Stephen G. Simpson, Rick L. Smith. *Addendum to "Countable algebra and set existence axioms"*. Annals of Pure and Applied Logic **28(3)**, pp. 319–320, 1985.
- [GM17] Noam Greenberg, Alexander Melnikov. *Proper divisibility in computable rings*. Journal of Algebra **474**, pp. 180–212, 2017.
- [Göd29] Kurt Gödel. Über die Vollständigkeit des Logikkalküls. PhD thesis. University of Vienna, 1929.
- [Göd31] Kurt Gödel. Über formal unentscheidbare Sätze der Principia Mathematica und verwandter Systeme I. Monatshefte für Mathematik und Physik **38(1)**, pp. 173–198, 1931.
- [Hat89] Kostas Hatzikiriakou. *Commutative Algebra in Subsystems of Second Order Arithmetic*. PhD thesis. Penn State, 1989.
- [HB34] David Hilbert, Paul Bernays. *Grundlagen der Mathematik*. Springer, Berlin, 1934.
- [Hen13] Henry (455). Dedekind-Hasse valuation. PlanetMath. 2013. URL: https://planetmath.org/dedekindhassevaluation.

[Hen50] Leon Henkin. *Completeness in the theory of types*. The Journal of Symbolic Logic **15(2)**, pp. 81–91, 1950.

- [Hil90] David Hilbert. *Ueber die Theorie der algebraischen Formen*. Mathematische Annalen **36(4)**, pp. 473–534, 1890.
- [Hil93] David Hilbert. *Ueber die vollen Invariantensysteme*. Mathematische Annalen **42(3)**, pp. 313–373, 1893.
- [Hil99] David Hilbert. *Grundlagen der Geometrie*. B.G. Teubner, Leipzig, 1899.
- [HP17] Petr Hájek, Pavel Pudlák. *Metamathematics of First-Order Arithmetic*. 2nd ed. Perspectives in Logic **3**. Cambridge University Press, Cambridge, 2017.
- [IDMap] R. C. Schwiebert. Field and domain properties. Database of Ring Theory. URL: https://ringtheory.herokuapp.com/maps/domain/.
- [Kle98] Israel Kleiner. *From Numbers to Rings: The Early History of Ring Theory*. Elemente der Mathematik **53(1)**, pp. 18–35, 1998.
- [Kum47a] E. E. Kummer. Über die Zerlegung der aus Wurzeln der Einheit gebildeten complexen Zahlen in ihre Primfactoren. Journal für die reine und angewandte Mathematik **1847(35)**, pp. 327–367, 1847.
- [Kum47b] E. E. Kummer. *Zur Theorie der complexen Zahlen*. Journal für die reine und angewandte Mathematik **1847(35)**, pp. 319–326, 1847.
- [Las05] E. Lasker. *Zur Theorie der moduln und Ideale*. Mathematische Annalen **60(1)**, pp. 20–116, 1905.
- [Lob29] Nikolai Lobachevsky. *A concise outline of the foundations of geometry*. University of Kazan Messenger, 1829.

[Mac13] F. S. Macaulay. *On the resolution of a given modular system into primary systems including some properties of Hilbert numbers.*Mathematische Annalen **74(1)**, pp. 66–121, 1913.

- [Mar02] David Marker. *Model Theory: An Introduction*. Graduate Texts in Mathematics. Springer-Verlag, New York, 2002.
- [MN82] George Metakides, Anil Nerode. *The introduction of non-recursive methods into mathematics*. pp. 319–335 in: A. S. Troelstra, D. van Dalen (eds.). *The L. E. J. Brouwer Cente-nary Symposium, Noordwijkerhout*. Studies in Logic and the Foundations of Mathematics **110**. Elsevier, 1982.
- [Mon21] Antonio Montalbán. *Computable Structure Theory: Within the Arithmetic*. Perspectives in Logic. Cambridge University Press, Cambridge, 2021.
- [MSE21] guest123456789. Answer to "Every ideal $J \supseteq \sqrt{0}$ contains a non-zero-divisor". 2021. URL: https://math.stackexchange.com/g/4009045.
- [Noe21] Emmy Noether. *Idealtheorie in Ringbereichen*. Mathematische Annalen **83(1)**, pp. 24–66, 1921.
- [Noe27] Emmy Noether. *Abstrakter Aufbau der Idealtheorie in algebraischen Zahl- und Funktionenkörpern*. Mathematische Annalen **96(1)**, pp. 26–61, 1927.
- [Pea89] Giuseppe Peano. *Arithmetices principia: nova methodo exposita*. Fratres Bocca, Rome, 1889.
- [PK78] J. B. Paris, L. A. S. Kirby. Σ_n -collection schemas in arithmetic. pp. 199–209 in: Angus Macintyre, Leszek Pacholski, Jeff Paris (eds.). *Logic Colloquium '77*. Studies in Logic and the Foundations of Mathematics **96**. Elsevier, Wrocław, 1978.
- [RMZoo] The Reverse Mathematics Zoo. URL: www.computability.org/zoo/.

[San18] Sam Sanders. *Refining the Taming of the Reverse Mathematics Zoo*. Notre Dame Journal of Formal Logic **59(4)**, pp. 579–597, 2018. arXiv: 1602.02270.

- [Sat16] Takashi Sato. *Reverse mathematics and countable algebraic systems*. PhD thesis. Tohoku University, Japan, 2016. HDL: 10097/96942.
- [Sim09] Stephen G. Simpson. Subsystems of Second Order Arithmetic.2nd ed. Perspectives in Logic. Cambridge University Press,Cambridge, 2009.
- [Sim88] Stephen G. Simpson. *Ordinal numbers and the Hilbert basis theorem*. The Journal of Symbolic Logic **53(3)**, pp. 961–974, 1988.
- [Sin11] Balwant Singh. *Basic Commutative Algebra*. World Scientific, Singapore, 2011.
- [Sla04] Theodore A Slaman. Σ_n -bounding and Δ_n -induction. Proceedings of the American Mathematical Society **132(8)**, pp. 2449–2456, 2004.
- [Smi03] Peter Smith. *An Introduction to Formal Logic*. Cambridge University Press, Cambridge, 2003.
- [Soa16] Robert I. Soare. *Turing Computability*. Theory and Applications of Computability. Springer-Verlag, Berlin, Heidelberg, 2016.
- [Soa87] Robert I. Soare. *Recursively Enumerable Sets and Degrees*. Perspectives in Mathematical Logic. Springer-Verlag, Berlin, Heidelberg, 1987.
- [Son17] Masazo Sono. *On congruences*. Memoirs of the College of Science, Kyoto Imperial University **2(3)**, pp. 203–226, 1917.
- [Tra62] Grace Joy Traylor. *On the equivalence of the axiom of choice, Zorn's lemma, and the well-ordering theorem.* Masters thesis. Atlanta University, 1962.

[Tur37] Alan Turing. *On computable numbers, with an application to the Entscheidungsproblem*. Proceedings of the London Mathematical Society **s2-42(1)**, pp. 230–265, 1937.

- [Vau86] Robert L. Vaught. *Alfred Tarski's work in model theory*. The Journal of Symbolic Logic **51(4)**, pp. 869–882, 1986.
- [Zer30] Ernst Zermelo. Über Grenzzahlen und Mengenbereiche: neue Untersuchungen über die Grundlagen der Mengenlehre. Fundamenta Mathematicae **16**, pp. 29–47, 1930.