

# Propensity score regression for causal inference with treatment heterogeneity

Peng Wu<sup>a,b,†</sup>, Shasha Han<sup>c,d,†</sup>, Xingwei Tong<sup>b</sup> and Runze Li<sup>e,\*</sup>

<sup>a</sup>Beijing Technology and Business University, <sup>b</sup>Beijing Normal University,

<sup>c</sup>Chinese Academy of Medical Sciences and Peking Union Medical College,

<sup>d</sup>Peking University, and <sup>e</sup>The Pennsylvania State University

*Abstract: Understanding how treatment effects vary on several key characteristics is critical in the practice of personalized medicine. In such cases, nonparametric estimation of these conditional average treatment effects is often desirable. However, few methods are available owing to the computational difficulty of such estimations. Furthermore, existing nonparametric methods, such as the inverse probability weighting methods, have limitations that hinder their use when the values of propensity scores are close to zero or one. We propose a propensity score regression (PSR) method that allows nonparametric estimation of such conditional average treatment effects in a wide context. The PSR comprises two nonparametric regressions. First, it regresses on the propensity scores together with the characteristics of interest, to obtain an intermediate estimate. Then, it regresses the intermediate estimate on the characteristics of interest only. By including*

---

†contributed equally.

\*correspond to: rzli@psu.edu

---

*propensity scores as regressors in a nonparametric manner, the PSR eases the computational difficulty substantially while remaining less sensitive to the values of propensity scores. We present its several appealing properties, including consistency and asymptotical normality. In particular, we show the existence of an explicit variance estimator, which we use to assess the analytical behavior of the PSR and its precision. The results of our simulation studies indicate that the PSR outperforms existing methods in various settings with extreme values of propensity scores. We apply our method to the national 2009 flu survey (NHFS) data to investigate the effects of seasonal influenza vaccinations and having paid sick leave across different age groups.*

*Key words and phrases:* Heterogeneous treatment effect, nonparametric estimation, propensity score, high-dimensional covariates.

## 1. Introduction

The heterogeneous treatment effect describes the effect variability due to varying characteristics and is widely used in contexts such as personalized medicine, policy design, and customized marketing (Kent et al., 2018; Yin, 2018; Imai and Strauss, 2011; Sato et al., 2019). In many settings, the characteristics of treatment relevance are only a subset of the baseline covariates ( $X = (X^l, X^{-l})$ ). Understanding how a treatment works for individuals that differ on these few core characteristics ( $X^l$ ) is particularly critical for developing tailored treatment decisions. For example, in clinical

---

practice, older patients tend to suffer more from side effects or drug-drug interactions. Thus, the age-dependent drug efficacy of a treatment is used to balance its benefits and risks (Velentgas et al., 2013). In health policy, age-dependent vaccine effectiveness is used to guide targeted vaccination programs (Soiza et al., 2021). Compared with the conditional treatment effects given the full covariates (Nie and Wager, 2021; Wager and Athey, 2018), these conditional treatment effects given the key characteristics are more easily interpretable and are widely used in clinical settings.

However, estimating such conditional treatment effects is challenging, because such methods should be able to flexibly distinguish the heterogeneous effect (which is conditional on  $X^l$ ) from the effect due to the remaining confounding covariates ( $X^{-l}$ ). Because  $X^{-l}$ , which can be high-dimensional, may still confound the effects of treatments on outcomes, conditioning on  $X^l$  is not sufficient. Moreover, the degree of confounding may vary with  $X^l$ , making modeling the conditional outcome particularly challenging. A nonparametric estimation method allows a fully flexible model and is therefore desirable. However, few nonparametric methods for estimating these heterogeneous treatment effects are available, and the weighting-based methods have limitations that hinder their use in a wide context. For example, the inverse probability weighting proposed by Abre-

---

vaya et al. (2015) uses the inverse of propensity scores as weights to adjust outcomes. However, this method can result in unstable estimates when the values of propensity scores are close to zero or one— that is, the weights are very large, as typically observed in weighting methods to population average treatment effects (see e.g., Hahn, 1998; Rubin, 2001; Kang and Schafer, 2007). The augmented inverse probability weighting (AIPW) methods (Lee et al., 2017) also use the inverse of propensity scores as weights, and they require correctly parametric modeling outcomes to achieve efficiency (Seaman and Vansteelandt, 2018). Although the requirement can be relaxed by leveraging machine learning methods (Fan et al., 2022; Zimmert and Lechner, 2019; Semenova and Chernozhukov, 2021), these methods can rely heavily on extrapolation, which is a critical concern in the context with extreme propensity score values (Kang and Schafer, 2007; Tan, 2007; Wu et al., 2022).

In general, alternative methods rely on a two-step estimation: first, they estimate the conditional treatment effects defined on the full covariates, and then they integrate out the obtained estimates to the desired level of granularity. However, it is often difficult to estimate the conditional treatment effect nonparametrically for a high-dimensional covariate (Abrevaya et al., 2015; Lechner, 2019; Zimmert and Lechner, 2019; Wu et al., 2022). For

---

example, in our example based on data from a national 2009 flu survey (NHFS) (see §5), the dimension of the full covariates is as high as 65 (see Supplementary Material). With such a high dimension, typical nonparametric estimation methods, for example, the local linear regression, would suffer from the so-called curse of dimensionality (Fan and Gijbels, 1996).

Following a two-step estimation, we propose a nonparametric propensity score regression (PSR) method, consisting of two nonparametric regressions. First, it regresses the propensity scores together with the covariates of interest. Then, it integrates out the scores by regressing the estimates from the first regression on the covariates of interest only. Propensity scores exhibit a crucial balancing property, namely, the distributions of full covariates between the treatment groups are identical at each level of the propensity scores (including the one-to-one functions of propensity scores). In our context, the balancing property is useful for controlling the confounding due to the remaining covariates  $X^{-l}$ , and for easing the computational difficulty. The PSR uses a continuous and bounded function of the score, and therefore is less sensitive to extreme propensity scores. Furthermore, by using the propensity scores in a nonparametric manner in the first step, the PSR reduces the influences of errors in the propensity scores on the estimates of the second step (Mammen et al., 2012), and thus enjoys increased robustness to

---

such errors. On the other hand, weighting-based methods achieve a covariate balance for a hypothetical super-population constructed by reweighing units in the study population, where small changes in propensity scores could lead to large discrepancies in weights, and even nonparametric estimation can result in highly unstable estimates.

The idea of including propensity scores in regression is not new in the context of parametric estimators of average treatment effects (see e.g., Little and An, 2004; Zhang and Little, 2009; Zhou et al., 2019; Wu et al., 2021). However, these approaches still rely on technical modeling assumptions for the outcome. When using propensity scores in a parametric regression, the key is to correctly specify the elusive relationship between the propensity score and the outcome, which is intrinsically connected to difficulties in specifying outcome models. Unlike these methods, we propose using propensity scores as regressors in a nonparametric manner in order to estimate heterogeneous treatment effects.

We validate the approach theoretically and show its appealing advantages. To obtain the theoretical results, we assume a parametric estimation of propensity scores, but do allow a nonparametric estimation of propensity scores. Note that, even under the parametric assumption, unlike the weighting-based methods, the PSR allows a one-to-one transformation of

---

propensity scores, and the functional form of the propensity score is less important. We present the theoretical properties of the proposed method, including the consistency and asymptotical normality, and an explicit variance estimator, which we use to assess the analytical behavior of the PSR and its precision.

The PSR is not only valuable for exploring treatment heterogeneity for practical guidance, but also useful in understanding treatment heterogeneity with high-dimensional full covariates. For example, we can decompose the full covariates into many subsets, each with only a few covariates, and estimate the heterogeneous treatment effects on these subsets. With this ensemble of such heterogeneous treatment effects, we may approach the full picture of the treatment heterogeneity with high-dimensional full covariates, which is computationally difficult to estimate directly (Abrevaya et al., 2015; Lechner, 2019; Zimmert and Lechner, 2019; Wu et al., 2022; Semenova and Chernozhukov, 2021).

The remainder of the paper proceeds as follows. In §2, we introduce the basic framework and the motivation of the analysis. In §3, we present the PSR method. Here, §3.1 outlines the method and provides the theoretical validation, and §3.2 provides the nonparametric estimator. In §3.3, we show the theoretical properties. We conduct several simulation studies in §4. In

---

§5, we apply our method to the national 2009 flu survey (NHFS) data to investigate the effects of seasonal influenza vaccination and having paid sick leave across different age groups. We conclude with a discussion in §6.

## 2. Motivations

### 2.1 Notation and assumptions

We adopt the framework of the Rubin Causal Model (Rubin, 1974), also called the potential outcome approach to causal inference (Imbens and Rubin, 2015). Consider a study with  $N$  units. Each unit  $i = 1, \dots, N$  is associated with a vector-valued covariate  $X_i = (X_{i1}, \dots, X_{il}, \dots, X_{ip}) \in \mathcal{X} \subseteq \mathfrak{R}^p$ , measured before being exposed to treatment  $D_i$ . The low-dimensional covariates of interest is denoted by  $X_i^l = (X_{i1}, \dots, X_{il}) \in \mathcal{X}^l \subset \mathfrak{R}^p$ . We write  $X_i = (X_i^l, X_i^{-l})$ . The outcome variable  $Y$  is measured on each unit after its treatment exposure. Associated with treatment  $d$ ,  $d = 0, 1$ , is the potential outcome  $Y_i(d)$ , the value of  $Y$  when unit  $i$  is exposed to treatment  $t$ , which implicitly assumes the stable unit treatment value assumption (SUTVA, Rubin, 1980). When referring to a generic unit, we drop the subscript and write  $X, D, Y(0), Y(1), Y, X^l, X^{-l}$ , and so on.

As in the literature (e.g., Abrevaya et al., 2015), we assume the unconfounded assumption, namely,  $D \perp (Y(1), Y(0)) \mid X$ . We denote the

---

propensity score  $e(X) := \mathbb{P}(D = 1|X)$ , assuming that  $0 < e(x) < 1$ , for any  $x \in \mathcal{X}$ . The heterogeneous treatment effect of interest,  $\tau(x^l)$ , is defined on the subspace of the covariates,  $\mathcal{X}^l$ , as

$$\tau(x^l) := \mathbb{E}(Y(1) - Y(0)|X^l = x^l), \quad (2.1)$$

where  $x^l \in \mathcal{X}^l$ . Often, the dimension of  $\mathcal{X}^l$  is much smaller than that of the full covariates space  $\mathcal{X}$ . Therefore,  $\tau(x^l)$  is at a higher level of granularity than the treatment effects conditional on the full covariates.

## 2.2 Two-step estimation

To estimate an estimand at a higher level of granularity, an intuitive way is to estimate the treatment effects at a lower level of granularity first, and then integrate out the obtained estimates into the subspace of interest. Our idea is broadly a two-step estimation. The key intuition is to explore an estimand at a lower level of granularity that can be estimated unbiasedly and nonparametrically. Below, we describe this idea further.

We write  $\tau(x^l)$  using the tower property of conditional expectation as

$$\tau(x^l) = \mathbb{E}[\mathbb{E}(Y(1) - Y(0)|X^l = x^l, X^{-l}) | X^l = x^l].$$

Let  $\eta(x) := \mathbb{E}(Y(1) - Y(0)|X^l = x^l, X^{-l} = x^{-l})$ , where  $x = (x^l, x^{-l})$ . In principle, we can estimate the insider expectation  $\eta(X)$  first, and then integrate  $X^{-l}$  out with respect to the conditional distribution of  $X^{-l}$  given  $X^l = x^l$ . In this case, our task is to estimate the finest estimand  $\eta(x)$  for each  $x \in \mathcal{X}$ , which can be identified as

$$\begin{aligned} \eta(x) &= \mathbb{E}\left(Y(1)|D = 1, X^l = x^l, X^{-l} = x^{-l}\right) - \mathbb{E}\left(Y(0)|D = 0, X^l = x^l, X^{-l} = x^{-l}\right) \\ &= \mathbb{E}\left(Y|D = 1, X^l = x^l, X^{-l} = x^{-l}\right) - \mathbb{E}\left(Y|D = 0, X^l = x^l, X^{-l} = x^{-l}\right). \end{aligned} \tag{2.2}$$

However, when the dimension  $p$  of the full covariates space is large, non-parametric estimation of  $\eta(X)$  can be difficult (Abrevaya et al., 2015; Lechner, 2019; Zimmert and Lechner, 2019; Wu et al., 2022).

This motivates us to explore an alternative estimand for the first-step estimation. Specifically, we aim to find an estimand that lies in a much higher level of granularity than  $\eta(X)$  while still lying in a lower level than  $\tau(X^l)$ , so that in practice it is possible to estimate the new estimand non-parametrically. Notably, the lower auxiliary variable needs to replace the important role that  $X^{-l}$  plays in the identification. As illustrated in Equation (2.2), conditioning on  $X^{-l}$  and  $X^l$ , i.e., the full covariates  $X$ , facilitates identifying  $\eta(X)$  using the observed data, owing to the natural balancing property of the full covariates  $X$ .

### 2.3 Comparison with existing methods using propensity scores

---

The new auxiliary variable needs to rest on a subspace with a dimension much smaller than  $p$ , and should exhibit the aforementioned balancing property. The propensity score, defined as the probability of assignment to the treatment given the full covariates (Rosenbaum and Rubin, 1983), is one candidate. As a summary of covariates, the propensity score reduces the  $p$ -dimensional covariates into a scalar while exhibiting the desired balancing property. Clearly, any one-to-one functions of the propensity score are candidates as well.

### 2.3 Comparison with existing methods using propensity scores

Existing methods use propensity scores to achieve the covariate balance by reweighing the units. For example, Abrevaya et al. (2015) proposed the inverse probability weighting (IPW) estimator

$$\tau^{IPW}(x^l) = \mathbb{E}\left(DY/e(X) - (1 - D)Y/(1 - e(X)) \middle| X^l = x^l\right). \quad (2.3)$$

As in the weighting methods (see e.g., Hahn, 1998; Rubin, 2001; Kang and Schafer, 2007), the IPW estimator in (2.3) is sensitive to the estimated propensity score values and their estimates are highly unstable if the propensity score values are close to zero or one. When the parametric function of the outcome model is knowable, Lee et al. (2017) propose the

### 2.3 Comparison with existing methods using propensity scores

---

AIPW estimator

$$\tau^{AIPW}(x^l) = \mathbb{E} \left( \frac{D(Y - \mu_1(X))}{e(X)} - \frac{(1 - D)(Y - \mu_0(X))}{1 - e(X)} + (\mu_1(X) - \mu_0(X)) \middle| X^l = x^l \right), \quad (2.4)$$

where  $\mu_d(\cdot)$  for  $d = 0, 1$  are specified outcome functions. The AIPW allows for misspecification of the propensity score model if the parametric outcome functions  $\mu_d(\cdot)$  are specified correctly. However, with high-dimensional covariates, correct specifications of outcome functions  $\mu_d(\cdot)$  are not easy. Several prior works have tried estimating  $\mu_d(\cdot)$  using machine learning methods (e.g., Fan et al., 2022; Semenova and Chernozhukov, 2021), but these methods rely heavily on extrapolation (e.g., Kang and Schafer, 2007; Tan, 2007).

Like the IPW, the AIPW is also sensitive to the extreme propensity score values, even when the propensity score models are specified correctly (Rotnitzky and Vansteelandt, 2014). Instead of using propensity scores as weights, we include them as one regressor in nonparametric regression, yielding the PSR method.

The PSR is conceptually different from the propensity score weighting to achieve the balance. The weighting methods use the inverse of propensity scores as weights to construct a hypothetical super-population, in which the distributions of the covariates between the treated units and the control units can be balanced. Clearly, these weights are unbounded around zero

### 2.3 Comparison with existing methods using propensity scores

---

or one, with small changes in propensity scores leading to potentially large discrepancies in weights, particularly when some propensity score values are extreme. The current setting is different. Here,  $\beta(X^l, e)$  defined in equation (3.1) is a bounded function of the propensity score, and is therefore less sensitive to extreme values of the propensity score.

As such, the PSR is analogous to propensity score matching and subclassification, all of which are based on the balancing property of propensity scores in the study population. But unlike matching on propensity scores, which implicitly involves model specifications (e.g., we need to specify the matching criteria and, in general, different criteria lead to different matched sets), the PSR uses propensity scores in a nonparametric manner, that is, using existing nonparametric methodologies. With parametric modeling on propensity scores, the difficulty of matching extreme propensity score values is likely to lead to substantial bias. However, here the estimation is nonparametric and  $\beta(x^l, e)$  is estimated smoothly. Thus, the results should be less sensitive to minor differences between inexact matches. We show this using a variant of the PSR in which the regression procedure is replaced by matching on propensity scores (§4).

Note that with large differences between the propensity scores of the treated units and the control units — for treated units near a propensity

---

of 1.0 *only*, and for control units near a propensity of 0.0 *only* — the PSR may not be suitable either. We note that in a mild situation, where both some treated units and control units near a propensity of 1.0 and 0.0, IPW and AIPW estimators could generate highly sensitive estimates.

### 3. PSR

The PSR includes two nonparametric regressions as follows. For notational simplicity, we refer to “ $e$ ” as the value of the propensity score  $e(X)$  or any one-to-one function of  $e(X)$ . As discussed in §2, the key idea is to explore an intermediate estimand at a lower level of granularity that can be estimated unbiasedly and nonparametrically. We denote  $\beta(X^l, e)$  as the intermediate estimand, which is conditional on the propensity score and the covariates of interest, as

$$\beta(x^l, e) = \mathbb{E} (Y(1) - Y(0) | X^l = x^l, e(X) = e). \quad (3.1)$$

Note that the estimand  $\beta(x^l, e)$  is conditional on the  $(l+1)$ -dimensional variables, where  $l+1$  is substantially smaller than the dimension of the full covariates  $p$ , and therefore can mitigate the problem of high-dimensional covariates. Estimating  $\beta(x^l, e)$  is our central task. Note that the definition

---

of  $\beta(x^l, e)$  includes both potential outcomes  $Y(1)$  and  $Y(0)$ , but only one of them is observed in the real world. By conditioning on the propensity score, we can replace the potential outcomes with the observed outcome  $Y$ . Below, we prove this nonparametrically.

**Proposition 1.** Suppose that  $\mathbb{E}(Y|D, X^l, e)$  is a nonparametric regression function. Then,  $\beta(X^l, e)$  and  $\mathbb{E}(Y(0)|X^l, e)$  are the functional coefficients corresponding to the treatment indicator  $D$  and the intercept, respectively:

$$\mathbb{E}(Y|D, X^l, e) = \beta(X^l, e) \cdot D + \mathbb{E}(Y(0)|X^l, e), \quad (3.2)$$

where  $\beta(x^l, e) = \mathbb{E}(Y | D = 1, X^l = x^l, e(X) = e) - \mathbb{E}(Y | D = 0, X^l = x^l, e(X) = e)$ .

Model (3.2) is a varying coefficient model (Hastie and Tibshirani, 1993) and can be estimated using standard local linear regression techniques (Fan and Zhang, 1999). Note that we do not make any parametric assumptions. The model is general enough to capture any model specification. The counterpart working model is given by

$$Y = \beta(X^l, e) \cdot D + \mathbb{E}(Y(0)|X^l, e) + \xi, \quad (3.3)$$

where  $\mathbb{E}(\xi|D, X^l, e) = 0$ .

Remarkably, here  $\beta(x^l, e)$  is estimated using the full sample of data un-

---

### 3.1 Description of the approach

der both two treatment conditions,  $d = 0, 1$ . Unlike existing methods such as the AIPW, our approach does not extrapolate the unobserved outcomes using predicted values from the other treatment group. Consequently, it is more likely to exhibit good finite-sample performance. Once we have estimates of  $\beta(x^l, e)$ , we simply integrate out the propensity scores  $e$  to obtain the estimates for  $\tau(x^l)$ . To do so, we conduct a second nonparametric estimation based on the projection relationship of  $\tau(x^l)$  and  $\beta(x^l, e)$ .

**Proposition 2.**  $\tau(x^l)$  is, geometrically, the projection of  $\beta(X^l, e)$  into the subspace spanned by  $X^l$ , specifically,  $\tau^{PSR}(x^l) = \mathbb{E}(\beta(X^l, e)|X^l = x^l)$ .

Proposition 2 suggests that  $\tau(x^l)$  can be estimated nonparametrically, for example, using a local linear regression of  $\beta(X^l, e)$  on  $X^l$ .

### 3.1 Description of the approach

When  $e(X)$  is known, the PSR is implemented using two nonparametric regressions, built on Propositions 1 and 2, respectively. When  $e(X)$  is also estimated from the data, the PSR is implemented in a total of three steps:

**Step 0:** Estimate  $e(X)$  in either a parametric or a nonparametric manner.

**Step 1:** Estimate  $\beta(X^l, e)$  by nonparametrically regressing the outcome  $Y$  on the covariate  $X^l$  and the propensity scores  $e(X)$ .

**Step 2:** Estimate  $\tau(X^l)$  by nonparametrically regressing the estimated values of  $\beta(X^l, e)$  on the covariate  $X^l$  only.

For practical use, we are interested in the large-sample properties of  $\hat{\tau}(x^l)$ . Here, we need to determine how the errors of the estimated propensity scores in Step 0 affect the estimates for  $\tau(x^l)$ . Briefly, the PSR is robust to estimation errors of the propensity scores, provided that the influence of these errors has a negligible effect on the second step estimation (Mammen et al., 2012). In our context, this is because the propensity scores are used in a nonparametric manner (Step 1), where the true propensity scores and their estimated counterparts are asymptotically indistinguishable.

Below, we focus on continuous  $X^l$ , but our results hold in the general setting including discrete  $X^l$ . For additional details on the discrete  $X^l$ , please refer to §7 of the Supplementary Material.

### 3.2 Nonparametric estimator

We use the standard local linear regression as the nonparametric method to estimate  $\beta(x^l, e)$  and  $\tau(x^l)$  in Steps 1 and 2, respectively. For notational simplicity, we focus on the case of  $l = 1$  and consider a more general case in the Supplementary Material.

We consider the case in which the propensity scores are known, and

---

### 3.2 Nonparametric estimator

denote the response vector  $\mathbf{Y} = (Y_1, \dots, Y_N)^\top$ , the regressor vector  $\mathbf{\Gamma} = (\Gamma_1, \dots, \Gamma_N)^\top$ , with the regressors  $\Gamma_i = (D_i, 1, D_i(X_i^l - x^l)/h_1, (X_i^l - x^l)/h_1, D_i(e_i - e)/h_2, (e_i - e)/h_2)^\top$  for  $i = 1, \dots, N$ , and the kernel vector  $\mathbf{W} = \text{diag}\{K_{h_1}(X_1^l - x^l)K_{h_2}(e_1 - e), \dots, K_{h_1}(X_N^l - x^l)K_{h_2}(e_N - e)\}$ , with  $K_{h_j}(u) = K(u/h_j)/h_j$  for  $j = 1, 2$ . The standard local linear estimator  $\tilde{\beta}(x^l, e)$  of  $\beta(x^l, e)$  is

$$\tilde{\beta}(x^l, e) = (1, 0, 0, 0, 0, 0)(\mathbf{\Gamma}^\top \mathbf{W} \mathbf{\Gamma})^{-1} \mathbf{\Gamma}^\top \mathbf{W} \mathbf{Y}, \quad (3.4)$$

which is the first component of  $(\mathbf{\Gamma}^\top \mathbf{W} \mathbf{\Gamma})^{-1} \mathbf{\Gamma}^\top \mathbf{W} \mathbf{Y}$ . Furthermore, we denote that the response vector  $\tilde{\boldsymbol{\beta}} = (\tilde{\beta}(X_1^l, e_1), \dots, \tilde{\beta}(X_N^l, e_N))^\top$ , the regressor vector  $\mathbf{G} = (G_1, \dots, G_n)^\top$  with  $G_i = (1, (X_i^l - x^l)/h_3)^\top$ , and  $\mathbf{\Lambda} = \text{diag}\{K_{h_3}(X_1^l - x^l), \dots, K_{h_3}(X_N^l - x^l)\}$ . The local linear estimator of  $\tau(x^l)$  is

$$\tilde{\tau}(x^l) = (1, 0)(\mathbf{G}^\top \mathbf{\Lambda} \mathbf{G})^{-1} \mathbf{G}^\top \mathbf{\Lambda} \tilde{\boldsymbol{\beta}}. \quad (3.5)$$

Next, we consider the case where the propensity scores are estimated from the data. We replace  $\mathbf{\Gamma}, \mathbf{W}$ , and  $\tilde{\boldsymbol{\beta}}$  in (3.4) and (3.5) with  $\hat{\mathbf{W}}, \hat{\mathbf{\Gamma}}$ , and  $\hat{\boldsymbol{\beta}}$ , respectively. Note that we replace the true propensity scores  $e$  by the estimated  $\hat{e}$ . Using the estimated propensity scores, the local linear

estimator of  $\beta(X^l, e)$  and  $\tau(x^l)$  is then given by

$$\hat{\beta}(x^l, e) = (1, 0, 0, 0, 0, 0)(\hat{\Gamma}^\top \hat{\mathbf{W}} \hat{\Gamma})^{-1} \hat{\Gamma}^\top \hat{\mathbf{W}} \mathbf{Y}, \quad (3.6)$$

$$\hat{\tau}(x^l) = (1, 0)(\mathbf{G}^\top \mathbf{\Lambda} \mathbf{G})^{-1} \mathbf{G}^\top \mathbf{\Lambda} \hat{\beta}. \quad (3.7)$$

### 3.3 Theoretical properties

We first present the following key assumption.

*Assumption 1.* The propensity score model can be written as  $e(X) = g(X^\top \alpha)$ , where  $\alpha$  is the true unknown parameter, and  $g(\cdot)$  is a known function (e.g., generalized linear model).

- (i) The estimates of  $\alpha$ , denoted by  $\hat{\alpha}$ , satisfies  $\hat{\alpha} - \alpha = O_p(N^{-1/2})$ ;
- (ii) The second-order derivative of  $g$  is uniformly bounded, that is,  $\sup_t |g''(t)|$  is bounded.

Assumption 1(i) is critical to our main results because it simplifies the asymptomatic analysis on  $\hat{\tau}(X^l)$ . Note that the parametric modeling assumption on propensity scores can be relaxed, see the Discussion for more details.

**Theorem 1.** Under Assumption 1 and regularity Assumption 2 in the Supplementary Material,  $\hat{\beta}(x^l, e)$  in (3.6) and  $\tilde{\beta}(x^l, e)$  in (3.4) are asymptotically

indistinguishable, that is,  $\sup_{(x^l, e) \in \mathcal{X}^l \times (0, 1)} |\hat{\beta}(x^l, e) - \tilde{\beta}(x^l, e)| = O_p(N^{-1/2})$ .

Theorem 1 demonstrates that the effect of the estimation error of the propensity score on the estimator of  $\beta(x^l, e)$  is negligible. Intuitively, this is because the propensity score is used in a nonparametric manner in Step 1, where we need only locally indistinguishable estimates of the propensity scores from the true values. This finding is consistent with the results of (Mammen et al., 2012) in a different setting on studying the coverage rates of the final estimates. The latter study shows that the effect of the first-step estimation error on the second-step estimation is restricted in a smoothed way through the estimation bias in the first step. From Theorem 1, the effect on the estimator of  $\tau(x^l)$  is small (e.g., Gu and Yang, 2015). We next establish the asymptotical normality of  $\hat{\tau}(x^l)$ . Let  $\bar{K}(x) = \int K(t)K(x + h_1 t/h_3)dt$ ,  $\nu = \int K^2(t)dt$ , where  $f(x^l)$  is the density function for  $X^l$ .

**Theorem 2.** Under Assumption 1 and regularity Assumptions 2–3 in the Supplementary Material,  $\hat{\tau}(x^l)$  in (3.7) is a consistent estimator, and

$$\mathbb{V}(x^l)^{-1/2} (\hat{\tau}(x^l) - \tau(x^l)) \xrightarrow{d} N(0, 1),$$

### 3.3 Theoretical properties

---

where  $\mathbb{V}(x^l)$  represents the asymptotic variance and

$$\mathbb{V}(x^l) = \frac{1}{Nh_3 f(x^l)} \left( \nu \cdot \mathbb{V}(\beta(X^l, e)|X^l = x^l) + \int \bar{K}^2(t) dt \cdot \mathbb{E} \left( \frac{(D - e)^2}{e^2(1 - e)^2} \xi^2 | X^l = x^l \right) \right).$$

Note that Theorem 2 relies on Assumption 3(iv)  $\sqrt{N}h_3(h_1^2 + h_2^2 + h_3^2) \rightarrow 0$  as  $N \rightarrow \infty$ . The common bias term, that is,  $O(h_1^2 + h_2^2 + h_3^2)$ , vanishes under the condition. However, our main conclusion still holds if the assumption is relaxed. Importantly, the asymptotic variance  $\mathbb{V}(x^l)$  can be estimated using the plug-in method given by.

$$\hat{\mathbb{V}}(x^l) := \frac{1}{Nh_3 \hat{f}(x^l)} \left( \nu \cdot \hat{\mathbb{V}}(\beta(X^l, e)|X^l = x^l) + \int \bar{K}^2(t) dt \cdot \hat{\mathbb{E}} \left( \frac{(D - e)^2}{e^2(1 - e)^2} \xi^2 | X^l = x^l \right) \right), \quad (3.8)$$

where  $\hat{f}(x^l)$  is the kernel density estimation,  $\hat{\mathbb{V}}(\beta_1(X^l, e)|X^l = x^l)$  can be estimated by conducting a nonparametric regression of  $(\hat{\beta}(X^l, e) - \hat{\tau}(X^l))^2$  on  $X^l$  (Fan and Yao, 1998). Because  $\xi$  is estimated using the residual in model (3.3), we obtain  $\hat{E}[\xi^2(D - e)^2/e^2(1 - e)^2|X^l = x^l]$  by regressing  $(D - \hat{e})^2 \hat{\xi}^2 / \hat{e}^2(1 - \hat{e})^2$  on  $X^l$ . In addition,  $\nu$  and  $\int \bar{K}^2(x) dx$  can be calculated directly. Finally, we have the following conclusion.

**Corollary 1.** *The estimated asymptotic variance is a consistent estimator, namely,  $\hat{\mathbb{V}}(x^l) - \mathbb{V}(x^l) = o_p(1)$ .*

---

## 4. Simulation studies

We conduct extensive simulation studies to assess the finite-sample performance of the PSR, compared with the existing IPW method of Abrevaya et al. (2015), and the AIPW method of Lee et al. (2017). In addition, we consider a matching variant of the PSR, where we use a matching method to estimate  $\beta(X^l, e)$ . Specifically, we replace Step 1 in the PSR by first creating matched pairs using matching on  $(X^l, \hat{e}(X))$ , and then using the matched pairs to impute the missing potential outcomes. Finally, we calculate  $\beta(X^l, e)$  using the imputed potential outcomes. In the matching, we use one-to-one matching and the Mahalanobis metric. In the following, we focus on settings with extreme propensity score values. We also consider an alternative scenario in which propensity scores are distributed far from zero and one.

### 4.1 Simulation Setup

We set  $l = 1$  and the covariate dimension  $p = 5, 20, \text{ and } 50$ .  $X^l \sim \text{Uniform}(-0.5, 0.5)$  and  $X^{-l} = (X_1^{-l}, \dots, X_{p-1}^{-l}) \sim \text{Norm}(0, \Sigma)$ , with  $\Sigma_{j,k} = 2^{-|j-k|}$  for  $1 \leq j, k \leq p-1$ . The assignment of treatment  $D$  follows the logistic model, with  $\mathbb{P}(D = 1|X) = \exp(X^\top \alpha)/(1 + \exp(X^\top \alpha))$ .

In the setting with extreme values of propensity scores, we consider two

## 4.1 Simulation Setup

---

assignment mechanisms: Mechanism A,  $\alpha = (1, -1, -1, 1, -1, 0, \dots, 0)^\top$ , with five nonzero entries; Mechanism B,  $\alpha = (1, 1, 1, 1, 1, 0, \dots, 0)^\top$ , with five nonzero entries. We conduct four simulation settings, in which the heterogeneous treatment effects  $\tau(x^l)$  are linear, quadratic, polynomial, and complex functions of  $x^l$ , respectively. The potential outcomes under the four contexts are modeled as follows:

$$\text{I: } Y(1) = X^l(1 + 2X^l)^2(X^l - 1)^2 + f(X) + \epsilon(1), Y(0) = f(X) + \epsilon(0), \\ f(X) = (X^l)^2 X_1^{-l} X_2^{-l} X_3^{-l} X_4^{-l},$$

$$\text{II: } Y(1) = X^l(1 - X^l) \cos(X^l) \log(X^l + 2) \exp(X^l) + f(X) + \epsilon(1), Y(0) = \\ f(X) + \epsilon(0), f(X) = (X^l)^2 X_1^{-l} X_2^{-l} X_3^{-l} X_4^{-l}.$$

$$\text{III: } Y(1) = X^l + f(X) + \epsilon(1), Y(0) = f(X) + \epsilon(0), f(X) = \{X^l X_1^{-l} + \\ \exp(X_2^{-l} - 3)(\sin(X_3^{-l}) + \cos(X_4^{-l}))\}/2,$$

$$\text{IV: } Y(1) = 5(X^l)^2 + X^l + f(X) + \epsilon(1), Y(0) = f(X) + \epsilon(0), f(X) = \\ (X^l)^2 \{\sum_{j=1}^4 X_j^{-l} / 2^{j+1}\}.$$

The error terms  $\epsilon(1)$  and  $\epsilon(0)$  are independently and identically distributed (i.i.d.) with  $\text{Norm}(0, 1)$ . The true  $\tau(x^l)$  in the four simulation settings are  $x^l(2x^l + 1)^2(x^l - 1)^2$ ,  $x^l(1 - x^l) \cos(x^l) \log(x^l + 2) \exp(x^l)$ ,  $x^l$ , and  $5(x^l)^2 + x^l$ , respectively. Note that the potential outcome models have a complex form, making it difficult to correctly specify the outcome models.

## 4.1 Simulation Setup

Simulations I and II use assignment Mechanism A, and Simulations III and IV use assignment Mechanism B. The distributions of the propensity scores under the two mechanisms are plotted in Figure 1. We observe that many propensity score values are close to zero or one, making the weighting-based propensity score methods highly unstable in the context.




Figure 1: Distribution of the true propensity score under the two treatment assignment mechanisms ( $N = 2000, p = 5$ ).

We estimate the propensity scores using logistic regression. The bandwidths are chosen using existing methods (Li and Racine, 2007; Ruppert et al., 1995), and implemented using the R functions `npscoefbw` from the package `np` (Racine and Hayfield, 2021) and `dpill` from the package `KernSmooth` (Wand et al., 2021). For the competing AIPW method, the outcome regression functions are estimated using linear models. Each simulation is based on 1000 replicates, with sample sizes 500, 1000, and 2000.

We evaluate the performance of the PSR using the sample average

bias (Bias), sample average standard deviation (SD), mean absolute error (MAE), mean squared error (MSE), and average 95% confidence interval coverage proportion (CP95). For the PSR, CP95 is estimated using the asymptotic variance formula (3.8). For the IPW and AIPW methods, CP95 is estimated as in Abrevaya et al. (2015) and Lee et al. (2017), respectively. Finally, for the matching variant of the PSR and random forest methods, CP95 is estimated using 100 bootstraps.

Table 1 summarizes the results of the PSR for cases I–IV. We observe that as the sample size increases, MAE and MSE decrease, and CP95 becomes closer to the nominal value of 0.95. Moreover, the results are similar for different values of  $p$ , suggesting that the PSR is insensitive to the dimension of the covariates.

We also consider an alternative scenario in which the propensity scores are distributed far from zero and one. We replace the data-generation mechanisms in Simulations I–IV with new mechanisms. Specifically, we replace Mechanism A in Simulations I and II with Mechanism C, where  $\alpha$  is set as  $0.25(1, -1, -1, 1, -1, 0, \dots, 0)^\top$  with five nonzero entries, and denote them as Simulations V and VI, respectively. Then, we replace Mechanism B in Simulations III and IV with Mechanism D,  $\alpha = 0.125(1, -1, -1, 1, -1, 0, \dots, 0)^\top$  with five nonzero entries, and denote them as Simulations VII and VIII,

## 4.1 Simulation Setup

Table 1: The performance of the PSR for cases I–IV.

$(N, p)$	Bias (SD) $\times 10^{-2}$	MAE $\times 10^{-2}$	MSE $\times 10^{-2}$	CP95 %	Bias (SD) $\times 10^{-2}$	MAE $\times 10^{-2}$	MSE $\times 10^{-2}$	CP95 %
	Simulation I				Simulation II			
(500, 5)	0.0 (16.9)	13.2	2.8	87.4	0.1 (15.5)	12.3	2.4	86.5
(1000, 5)	-0.4 (12.7)	10.0	1.6	91.8	0.1 (11.8)	9.3	1.4	90.7
(2000, 5)	-0.5 (9.4)	7.5	0.9	94.4	0.2 (8.4)	6.6	0.7	93.6
(500, 20)	0.0 (17.3)	13.7	3.0	87.5	0.4 (15.4)	12.4	2.6	86.5
(1000, 20)	-0.0 (12.7)	10.3	1.7	89.8	-0.0 (11.6)	9.2	1.4	90.6
(2000, 20)	0.0 (9.6)	7.5	0.9	94.3	-0.0 (9.0)	6.8	0.7	93.8
(500, 50)	-0.4 (17.3)	13.7	3.1	87.3	-0.0 (16.5)	12.7	2.6	86.9
(1000, 50)	-0.2 (13.0)	10.2	1.7	91.5	0.0 (11.8)	9.6	1.5	90.4
(2000, 50)	-0.3 (9.4)	7.5	0.9	94.2	0.0 (9.1)	6.8	0.8	93.8
	Simulation III				Simulation IV			
(500, 5)	3.3 (17.4)	13.7	3.1	88.1	2.0 (22.1)	17.2	4.9	88.9
(1000, 5)	2.7 (13.2)	10.6	1.8	91.4	1.4 (16.8)	13.2	2.8	92.2
(2000, 5)	3.2 (10.5)	8.6	1.2	93.9	1.6 (13.0)	10.2	1.7	95.2
(500, 20)	2.8 (17.8)	14.2	3.4	86.9	2.1 (22.3)	17.5	5.1	88.6
(1000, 20)	2.8 (13.9)	10.8	1.9	91.0	1.6 (16.9)	13.2	2.9	92.7
(2000, 20)	2.5 (10.7)	8.3	1.1	93.4	1.5 (12.8)	10.1	1.7	95.1
(500, 50)	2.9 (19.4)	14.6	3.5	85.8	1.8 (22.1)	17.4	4.9	88.3
(1000, 50)	3.1 (14.4)	11.4	2.1	90.2	1.2 (16.7)	13.0	2.8	92.2
(2000, 50)	2.4 (10.6)	8.6	1.2	93.7	1.3 (13.2)	10.4	1.8	94.6

respectively. Figure 2 shows the propensity score distributions of the treatment assignment mechanisms C and D.




Figure 2: Distribution of the true propensity score under the treatment assignment mechanisms C and D ( $N = 2000, p = 5$ ).

We present the results for Simulations V–VIII, and contrast them with

## 4.1 Simulation Setup

Table 2: The performance of the PSR, IPW, AIPW, and matching variant of the PSR under different simulation settings ( $N = 2000, p = 5$ )

Pair	Case	Bias (SD) $\times 10^{-2}$	MAE $\times 10^{-2}$	MSE $\times 10^{-2}$	CP95 %	Case	Bias (SD) $\times 10^{-2}$	MAE $\times 10^{-2}$	MSE $\times 10^{-2}$	CP95 %
PSR method										
(1)	I	-0.5 (9.4)	7.5	0.9	94.4	V	-0.2 (8.9)	6.9	0.8	93.7
(2)	II	0.2 (8.4)	6.6	0.7	93.6	VI	0.0 (7.7)	6.1	0.6	94.0
(3)	III	3.2 (10.5)	8.6	1.2	93.9	VII	1.1 (8.3)	6.7	0.7	93.2
(4)	IV	1.6 (13.0)	10.2	1.7	95.2	VIII	1.1 (12.1)	9.6	1.5	94.1
IPW method										
(1)	I	0.6 (33.9)	14.7	11.5	92.7	V	0.6 (11.5)	8.6	1.3	91.1
(2)	II	0.3 (31.1)	14.2	9.7	93.7	VI	0.0 (10.9)	8.1	1.2	91.8
(3)	III	-0.1 (53.8)	20.7	28.9	94.2	VII	-0.2 (10.7)	8.1	1.1	92.1
(4)	IV	2.2 (49.0)	22.3	24.1	93.9	VIII	2.4 (11.7)	9.1	1.4	91.2
AIPW method										
(1)	I	0.4 (29.8)	17.0	8.9	94.9	V	-0.2 (13.7)	10.4	1.9	94.0
(2)	II	-0.2 (26.9)	17.0	7.2	94.6	VI	-0.0 (13.9)	10.4	1.9	94.3
(3)	III	0.8 (66.7)	25.6	44.6	94.0	VII	-0.1 (13.3)	10.1	1.8	93.8
(4)	IV	0.5 (61.8)	25.9	38.2	95.0	VIII	0.6 (14.0)	10.8	2.0	94.1
Matching variant of PSR										
(1)	I	0.4 (22.7)	17.7	5.2	93.4	V	0.0 (18.0)	13.9	3.2	96.1
(2)	II	0.5 (22.8)	17.8	5.2	93.2	VI	0.3 (17.9)	13.8	3.2	96.2
(3)	III	0.7 (29.3)	23.1	8.6	90.4	VII	0.2 (18.0)	14.0	3.2	95.8
(4)	IV	0.7 (29.2)	23.0	8.5	90.5	VIII	0.7 (17.8)	13.9	3.2	95.8

those of Simulations I–IV in Table 2 ( $N = 2000, p = 5$ ). Each row represents the results for a pair of simulations that differ only in terms of their propensity score mechanisms. We find no significant differences between Bias (SD), MAE, and MSE when PSR is used. However, when we use the IPW, AIPW, and the matching variant of the PSR, the SD, MAE, and MSE in settings with extreme propensity score values are significantly different from those in settings with general propensity scores. This again shows that the PSR is robust to extreme propensity score values. In addition, we find that the matching variant of the PSR performs similarly to our PSR in terms of bias, but has a larger SD, MAE, and MSE, because both methods leverage the idea of propensity score matching. However, unlike

## 4.2 Alternative estimation methods on propensity scores

Table 3: The performance of the PSR under different estimation errors of propensity scores ( $N = 2000, p = 5$ ).

Method	Bias (SD) $\times 10^{-2}$	MAE $\times 10^{-2}$	MSE $\times 10^{-2}$	CP95 %	Bias (SD) $\times 10^{-2}$	MAE $\times 10^{-2}$	MSE $\times 10^{-2}$	CP95 %
	Simulation I				Simulation II			
Logistic	-0.5 (9.4)	7.5	0.9	94.4	0.2 (8.4)	6.6	0.7	93.6
Probit	-0.1 (9.6)	7.5	0.9	94.1	0.1 (8.9)	6.9	0.8	93.9
Random Forest	0.2 (9.6)	7.6	0.9	91.4	0.0 (8.7)	6.9	0.8	92.7
	Simulation III				Simulation IV			
Logistic	3.2 (10.5)	8.6	1.2	93.9	1.6 (13.0)	10.2	1.7	95.2
Probit	3.2 (10.6)	8.7	1.2	93.8	1.7 (13.1)	10.3	1.7	94.6
Random Forest	3.0 (10.6)	8.8	1.2	91.2	2.5 (12.2)	10.0	1.6	91.6

the matching method, the PSR uses propensity scores in a nonparametric manner and smoothly estimates  $\beta(x^l, e)$ , making it less sensitive to minor differences between inexact matches. This explains why the PSR has a smaller SD, MAE, and MSE than those of its matching variant.

## 4.2 Alternative estimation methods on propensity scores

We consider two alternative scenarios when estimating the propensity score. In the first scenario, we estimate the propensity scores using a probit model. In the second scenario, we estimate the propensity scores nonparametrically, using the random forest method (R package **grf**). We compare the two with the baseline scenario in which the propensity scores are estimated using the true logit model. Our results show that the obtained Bias, SD, MAE, and MSE are all very close under the three scenarios.

---

## 5. Application

We demonstrate our method in two studies using data from the National 2009 H1N1 Flu Survey (NHFS). The NHFS was a large one-time telephone survey conducted in the United States from October 2009 through June 2010, by the Centers for Disease Control and Prevention (CDC). The survey asked questions on participants' seasonal influenza vaccination status, whether had been sick with an influenza-like illness in the past month, the number of days they had taken off work owing to influenza, whether they have paid sick leave benefits, the number of times they see a doctor, as well as other relevant information (e.g., influenza-related behaviors, opinions about influenza vaccine safety and effectiveness, the size of the household, and demographic characteristics) (Centers for Disease Control and Prevention, 2010b). The NHFS public dataset has been released by the CDC, National Center for Immunization and Respiratory Diseases (NCRID), and National Center for Health Statistics (NCHS). The datasets are used to analyze the vaccination coverage, vaccination beliefs, and behaviors (Centers for Disease Control and Prevention, 2010a; Ding et al., 2011; Burger et al., 2021). Using a subset of the data comprising adults (i.e., age  $\geq 18$ ) and English-speaking participants, we conducted the following two studies.

## 5.1 Effect of seasonal influenza vaccination on the number of sick days

---

### 5.1 Effect of seasonal influenza vaccination on the number of sick days

We estimate the effect of seasonal influenza vaccinations on the number of sick days taken because of an influenza-like illness. Our primary outcome is the number of days taken off work (after taking “log”) when sick with an influenza-like illness, as reported by the participants during the interview. We consider a subset of the data, including only adult participants who had reported being infected with an influenza-like illness, and excluding participants with missing outcomes or treatments. We consider covariates that have non-missing values for at least 70% of the participants. For each selected covariate, the missing value is treated as a new category. Our final sample comprised 2442 participants and 65 (i.e.,  $p = 65$ ) dimensional covariates, where 1145 individuals have had a seasonal flu vaccination. We include all the informative covariates in the analysis, because conditioning on any given covariates is, in general, better than not conditioning (Rosenbaum, 2002; Rubin, 2009; Ding and Miratrix, 2015b). Nevertheless, the potential bias introduced by the adjustment needs further attention (Pearl, 2015; Ding and Miratrix, 2015a). Descriptive statistics of the sample and the covariates are listed in the Supplementary Material Table S2.

For our approach, we follow the three steps described in Section 3.1.

## 5.1 Effect of seasonal influenza vaccination on the number of sick days




Figure 3: (a) Distribution of the propensity scores. (b) Effects of seasonal vaccination on taking sick days across age groups. Dashed lines refer to 95% confidence interval.

We estimate the propensity scores using a logistic regression and the heterogeneous effects using a standard local linear regression, as described in Section 3.2. We display the results in Figure 3.

As shown in Figure 3(a), the propensity score values vary from 0.001 to 0.994 for the group of people with a seasonal vaccination, and from 0.001 to 0.978 for the group of people without a seasonal vaccination. Therefore, we cannot use methods that are sensitive to propensity scores. Using our PSR method, we show that the effects of seasonal vaccination on taking sick days vary across age groups (Figure 3b). Furthermore, seasonal vaccines decrease the number of sick days off work for the population aged over 60 and under 35, but have a negligible impact on the other age groups. Because seasonal vaccines prevent severe symptoms of influenza (Deiss et al., 2015),

## 5.2 Effect of having paid sick leave on visiting doctors

---

in general, people with seasonal vaccinations are more likely to develop light symptoms and, therefore, are less likely to take leave off work when sick with influenza. However, our results suggest that people aged 36–59 are equally likely to leave work to see doctors, regardless of the severity of the symptoms.

### 5.2 Effect of having paid sick leave on visiting doctors

In the second study, we estimate the effect of having paid sick leave on the number of times people see a doctor regardless of the disease type. Our treatment is whether the adult earns paid sick time off from employment, with  $D = 1$  representing having paid sick leave, and  $D = 0$  representing not having paid sick leave. The primary outcome  $Y$  is the self-reported number of times a person sees a doctor. During the interview, the participants were asked to provide the number of times they had seen a doctor or other health professional about health since August 2009. For the study, we also consider a subset of the data, including only adults whose paid sick leave indicator  $D$  is known. Our final sample comprised 8425 participants and 62 (i.e.,  $p = 62$ ) dimensional covariates, where 5502 have paid sick leave and 2923 do not. Descriptive statistics of the sample and covariates are listed in the Supplementary Table S3. Similarly, we estimate the propensity scores

## 5.2 Effect of having paid sick leave on visiting doctors




Figure 4: (a) Distribution of the propensity scores. (b) Effects of having paid sick leave on seeing a doctor, across age groups. Dashed lines refer to the 95% confidence interval.

using the logistic models and the heterogeneous effects using the standard local linear regression, as described in Section 3.2. We display the results in Fig. 4.

As in the first study, the propensity scores are distributed with values varying from 0.033 to 0.984 for the group of people with paid sick leave, and 0.001 to 0.956 for people without paid sick leave. The effects of having paid sick leave vary substantially across age groups (Fig. 4b). Specifically, having paid sick leave increases the number of times of seeing doctors for people aged over 65, but has no significant effects for people aged 33–40. Interestingly, having paid sick leave motivates people aged under 33 or aged 41–64 to reduce the number of times they see doctors. One possible explanation is that these people may have a bundle of paid time-off benefits

---

that combines sick days, vacation days, and other types of leave, where reducing the number of times they see a doctor may increase their overall benefits (Zhai et al., 2018; Smith and Kim, 2010).

## 6. Discussion

We have proposed a nonparametric PSR method for estimating the heterogeneous treatment effects in a wide context, including settings in which the propensity scores are close to zero or one and the number of full covariates is large. We have established the large-sample properties, and show that it outperforms existing methods in our simulation studies. Although in the main text we consider continuous  $X^l$ , our methods hold regardless of the type of variable  $X^l$ . In §7 of the Supplementary Material, we present theoretical results also for discrete  $X^l$ , where we use the typical kernel smoothing method for the estimation (Aitchison and Aitken, 1976; Li and Racine, 2010).

Note that our theoretical results hold when we replace the parametric model specification of the propensity score with a semiparametric model, such as the single index model. In this case, we may simply use  $X^\top \hat{\alpha}$  instead of  $e(X)$ , because the PSR is built on the balancing property of the propen-

---

sity score, and any one-to-one function of propensity scores has the same balancing property. For a single index model, many available estimators satisfy the assumptions in Assumption 1 with *unknown* link functions; see, for example, Horowitz and Härdle (1996), Ichimur (1993), Klein and Spady (1993), Härdle et al. (1997), and Wang and Yang (2009) for further detail. Furthermore, methods that can improve the balance property (Huang and Chan, 2017; Wei et al., 2017; Tan, 2020; Imai and Ratkovic, 2014; Ning et al., 2020) could be useful for improving the performance of the PSR.

For nonparametric estimation, we use kernel-based methods, for which an appropriate choice of the bandwidths,  $h_1$ ,  $h_2$ , and  $h_3$ , is important to achieve good accuracy. In the simulation, we simply use the existing bandwidth-selection methods (Li and Racine, 2007), and the bandwidths are chosen independently in the corresponding varying coefficient models or local linear models. To improve accuracy, approaches that simultaneously account for the bandwidth choice in estimating  $\beta(X^l, e)$  and  $\tau(X^l)$  are probably helpful.

Finally, we have focused on continuous outcomes. However, the PSR is not restricted to such outcomes. Note that  $\tau(x^l) = \mathbb{E}[\beta(X^l, e)|X^l = x^l]$  can always be estimated using a local linear regression of  $\beta(X^l, e)$  on  $X^l$ , regardless of the type of outcome. However, when the PSR is used

---

for a discrete outcome, particular care is needed for the potential model extrapolation when estimating  $\beta(X^l, e)$ . This is because in the context of discrete outcomes, the estimation of the model (3.3),  $Y = \beta(X^l, e) \cdot D + \mathbb{E}(Y(0)|X^l, e) + \xi$ , is often done separately for each treatment group, instead of two treatment groups together. Note that  $\beta(X^l, e)$  can be rewritten with two components,

$$\beta(X^l, e) = \mathbb{E}(Y(1) - Y(0)|X^l, e) = \mathbb{E}(Y(1)|X^l, e) - \mathbb{E}(Y(0)|X^l, e).$$

As such, alternative methods are needed to estimate  $\beta(X^l, e)$  without separating the two treatment groups. This is left to future work.

### **Supplementary Material**

The online Supplementary Material includes technical proofs, additional numerical results from the simulation study and empirical application, and extensions of the proposed method.

### **Acknowledgments**

The authors thank the assistant editor and anonymous reviewers for their helpful comments and valuable suggestions. This research was supported

---

## REFERENCES

by the National Natural Science Foundation of China (No. 11971064, 12071015, and 12171374) and the Major Project of National Statistical Science Foundation of China (No. 2021LD01).

### References

- Abrevaya, J., Y. C. Hus, and R. P. Lieli (2015). Estimating conditional average treatment effect. *Journal of Business and Economic Statistics* 33, 485–505.
- Aitchison, J. and C. Aitken (1976). Multivariate binary discrimination by the kernel method. *Biometrika* 63(3), 413–420.
- Burger, A. E., E. N. Reither, S.-E. Mamelund, and S. Lim (2021). Black-white disparities in 2009 h1n1 vaccination among adults in the united states: A cautionary tale for the covid-19 pandemic. *Vaccine* 39, 943–951.
- Centers for Disease Control and Prevention (2010a). *Interim results: state-specific influenza A (H1N1) 2009 monovalent vaccination coverage-United States, October 2009-January 2010*. Centers for Disease Control and Prevention.
- Centers for Disease Control and Prevention (2010b). *National 2009 H1N1 Flu Survey Questionnaire [2010 Q1]*. Centers for Disease Control and Prevention.
- Deiss, R. G., J. C. Arnold, W. J. Chen, S. Echols, M. P. Fairchok, C. Schofield, P. J. Danaher, E. McDonough, M. Ridoré, D. Mor, T. H. Burgess, and E. V. Millar (2015). Vaccine-associated reduction in symptom severity among patients with influenza a/h3n2 disease. *Vaccine* 33, 7160–7167.
- Ding, H., T. A. Santibanez, D. J. Jamieson, C. M. Weinbaum, G. L. Euler, L. A. Grohskopf, P.-J. Lu, and J. A. Singleton (2011). Influenza vaccination coverage among pregnant women—national 2009 h1n1 flu survey (nhfs). *American Journal of Obstetrics and Gynecology* 204, 96–106.
- Ding, P. and L. W. Miratrix (2015a). Reply to professor pearl’s comment. *Journal of Causal Inference* 3, 251–252.
- Ding, P. and L. W. Miratrix (2015b). To adjust or not to adjust? sensitivity analysis of m-bias and butterfly-bias. *Journal of Causal Inference* 3, 41–57.

## REFERENCES

---

- Fan, J. and I. Gijbels (1996). *Local Polynomial Modelling and Its Applications*. Chapman and Hall/CRC.
- Fan, J. and Q. W. Yao (1998). Efficient estimation of conditional variance functions in stochastic regression. *Biometrika* 85, 645–660.
- Fan, J. and W. Zhang (1999). Statistical estimation in varying coefficient models. *Annals of Statistics* 27, 1491–1518.
- Fan, Q., Y. C. Hsu, R. P. Lieli, and Y. Zhang (2022). Estimation of conditional average treatment effects with high-dimensional data. *Journal of Business and Economic Statistics* 40, 313–327.
- Gu, L. and L. Yang (2015). Oracally efficient estimation for single-index link function with simultaneous confidence band. *Electronic Journal of Statistics* 9, 1540–1561.
- Hahn, J. (1998). On the role of the propensity score in efficient semiparametric estimation of average treatment effects. *Econometrica* 66, 315–331.
- Härdle, W., V. Spokoiny, and S. Sperlich (1997). Semiparametric single index versus fixed link function modelling. *Annals of Statistics* 25, 212–243.
- Hastie, T. and R. Tibshirani (1993). Varying-coefficient models. *Journal of the Royal Statistical Society, Series B* 55, 757–796.
- Horowitz, J. L. and W. Härdle (1996). Direct semiparametric estimation of single-index models with discrete covariates. *Journal of the American Statistical Association* 91, 1632–1640.
- Huang, M. Y. and K. C. G. Chan (2017). Joint sufficient dimension reduction and estimation of conditional and average treatment effects. *Biometrika* 104, 583–596.
- Ichimur, H. (1993). Semiparametric least squares (sls) and weighted sls estimation of single-index models. *Journal of Econometrics* 58, 71–120.
- Imai, K. and M. Ratkovic (2014). Covariate balancing propensity score. *Journal of the Royal Statistical Society, Series B* 76, 243–263.
- Imai, K. and A. Strauss (2011). Estimation of heterogeneous treatment effects from randomized experiments, with application to the optimal planning of the get-out-the-vote campaign. *Political Analysis* 19, 1–19.
- Imbens, G. W. and D. B. Rubin (2015). *Causal inference for statistics social and biomedical science*. Cambridge University Press.

## REFERENCES

---

- Kang, J. D. and J. L. Schafer (2007). Demystifying double robustness: a comparison of alternative strategies for estimating a population mean from incomplete data. *Statistical Science* 22, 523–539.
- Kent, D. M., E. Steyerberg, and D. van Klaveren (2018). Personalized evidence based medicine: predictive approaches to heterogeneous treatment effects. *The British Medical Journal* 363, k4245.
- Klein, R. and R. H. Spady (1993). An efficient semiparametric estimator for binary response models. *Econometrica* 61, 387–421.
- Lechner, M. (2019). Modified causal forests for estimating heterogeneous causal effects. [arxiv.org/abs/1812.09487v2](https://arxiv.org/abs/1812.09487v2).
- Lee, S., R. Okui, and Y. J. Whang (2017). Doubly robust uniform confidence band for the conditional average treatment effect function. *Journal of Applied Econometrics* 32, 1207–1225.
- Li, Q. and J. S. Racine (2007). *Nonparametric econometrics*. Princeton University Press.
- Li, Q. and J. S. Racine (2010). Smooth varying-coefficient estimation and inference for qualitative and quantitative data. *Econometric Theory* 26(6), 1607–1637.
- Little, R. and H. An (2004). Robust likelihood-based analysis of multivariate data with missing values. *Statistica Sinica* 14, 949–968.
- Mammen, E., C. Rothe, and M. Schienle (2012). Nonparametric regression with nonparametrically generated covariates. *The Annals of Statistics* 40, 1132–1170.
- Nie, X. and S. Wager (2021). Quasi-oracle estimation of heterogeneous treatment effects. *Biometrika* 108, 299–319.
- Ning, Y., P. Sida, and K. Imai (2020). Robust estimation of causal effects via a high-dimensional covariate balancing propensity score. *Biometrika* 107, 533–554.
- Pearl, J. (2015). Comment on ding and miratrix: “to adjust or not to adjust?”. *Journal of Causal Inference* 3, 59–60.
- Racine, J. S. and T. Hayfield (2021). *np: Nonparametric Kernel Smoothing Methods for Mixed Data Types*. <https://CRAN.R-project.org/package=np>.
- Rosenbaum (2002). *Observational Studies*. Springer.
- Rosenbaum, P. R. and D. B. Rubin (1983). The central role of the propensity score in observa-

## REFERENCES

---

- tional studies for causal effects. *Biometrika* 70, 41–55.
- Rotnitzky, A. and S. Vansteelandt (2014). *Double-robust methods*, pp. 185–212. in "Handbook of Missing Data Methodology". CRC Press, Boca Raton, FL.
- Rubin, D. B. (1974). Estimating causal effects of treatments in randomized and nonrandomized studies. *Journal of educational psychology* 66, 688–701.
- Rubin, D. B. (1980). Discussion of randomization analysis of experimental data in the fisher randomization test by basu. *Journal of the American Statistical Association* 75, 591–593.
- Rubin, D. B. (2001). Using propensity scores to help design observational studies: application to the tobacco litigation. *Health Services and Outcomes Research Methodology* 2, 169–188.
- Rubin, D. B. (2009). Should observational studies be designed to allow lack of balance in covariate distributions across treatment groups? *Statistics in Medicine* 28, 1420–1423.
- Ruppert, D., S. J. Sheather, and M. P. Wand (1995). An effective bandwidth selector for local least squares regression. *Journal of the American Statistical Association* 90, 1257–1270.
- Sato, M., J. Singh, S. Takemori, T. Sonoda, Q. Zhang, and T. Ohkuma (2019). Uplif-based evaluation and optimization of recommenders. In *Conference on Recommender Systems*.
- Seaman, S. R. and S. Vansteelandt (2018). Introduction to double robust methods for incomplete data. *Statistical Science* 33, 184–197.
- Semenova, V. and V. Chernozhukov (2021). Debiased machine learning of conditional average treatment effects and other causal functions. *The Econometrics Journal* 24, 264–289.
- Smith, T. W. and J. Kim (2010). *Paid Sick Days: Attitudes and Experiences*. Public Welfare Foundation.
- Soiza, R. L., C. Scicluna, and E. C. Thomson (2021). Efficacy and safety of covid-19 vaccines in older people. *Age and Ageing* 50, 279–283.
- Tan, Z. (2007). Comment: understanding or, ps and dr. *Statistical Science* 22, 560–568.
- Tan, Z. (2020). Regularized calibrated estimation of propensity scores with model misspecification and high-dimensional data. *Biometrika* 107, 137–158.
- Velentgas, P., N. A. Dreyer, P. Nourjah, S. R. Smith, and M. M. Torchia (2013). *Developing a Protocol for Observational Comparative Effectiveness Research: A User's Guide*. Agency for Healthcare Research and Quality (AHRQ).
- Wager, S. and S. Athey (2018). Estimation and inference of heterogeneous treatment effects

## REFERENCES

---

- using random forests. *Journal of the American Statistical Association* 113, 1228–1242.
- Wand, M., C. Moler, and B. Ripley (2021). *KernSmooth: Functions for Kernel Smoothing Supporting Wand & Jones (1995)*. <https://CRAN.R-project.org/package=KernSmooth>.
- Wang, L. and L. Yang (2009). Spline estimation of single index model. *Statistica Sinica* 19, 765–783.
- Wei, L., Z. Yeying, and G. Debashis (2017). On estimating regression-based causal effects using sufficient dimension reduction. *Biometrika* 104, 51–65.
- Wu, P., Z. Tan, W. Hu, and X.-H. Zhou (2022). Model-assisted inference for covariate-specific treatment effects with high-dimensional data. *Statistica Sinica (To Appear)*.
- Wu, P., X. Tong, Y. Wang, J. Liang, and X.-H. Zhou (2022). Robust quasi-oracle estimation of average causal effects. *Biostatistics and Epidemiology* 6, 144–163.
- Wu, P., X. Xu, X. Tong, Q. Jiang, and B. Lu (2021). Semiparametric estimation for average causal effects using propensity score-based spline. *Journal of Statistical Planning and Inference* 212, 153–168.
- Yin, Y. (2018). Assessing the treatment effect heterogeneity with a latent variable. *Statistica Sinica* 28, 115–135.
- Zhai, Y., T. A. Santibanez, K. E. Kahn, C. L. Black, and M. A. de Perio (2018). Paid sick leave benefits, influenza vaccination, and taking sick days due to influenza-like illness among u.s. workers. *Vaccine* 36, 7316–7323.
- Zhang, G. and R. Little (2009). Extensions of the penalized spline of propensity prediction method of imputation. *Biometrics* 65, 911–918.
- Zhou, T., M. R. Elliott, and R. J. A. Little (2019). Penalized spline of propensity methods for treatment comparison. *Journal of the American Statistical Association* 114, 1–19.
- Zimmert, M. and M. Lechner (2019). Nonparametric estimation of causal heterogeneity under high-dimensional confounding. <https://arxiv.org/abs/1908.08779v1>.