# Solution group representations as quantum symmetries of graphs

David E. Roberson<sup>\*1</sup> and Simon Schmidt<sup>+2</sup>

<sup>1</sup>Department of Applied Mathematics and Computer Science, Technical University of Denmark, DK-2800 Lyngby, Denmark <sup>2</sup>QMATH, Department of Mathematical Sciences, University of Copenhagen, Universitetsparken 5, 2100 Copenhagen Ø, Denmark

November 25, 2021

#### Abstract

In 2019, Aterias et al. constructed pairs of quantum isomorphic, non-isomorphic graphs from linear constraint systems. This article deals with quantum automorphisms and quantum isomorphisms of colored versions of those graphs. We show that the quantum automorphism group of such a colored graph is the dual of the homogeneous solution group of the underlying linear constraint system. Given a vertex– and edge-colored graph with certain properties, we construct an uncolored graph that has the same quantum automorphism group as the colored graph we started with. Using those results, we obtain the first known example of a graph that has quantum symmetry and finite quantum automorphism group. Furthermore, we construct a pair of quantum isomorphic, non-isomorphic graphs that both have no quantum symmetry.

## 1 Introduction

Generalizing Mermin's magic square [12], linear system games were introduced by Cleve and Mittal in [7]. It was then shown by Cleve, Liu and Slofstra [6], that perfect quantum strategies for those games are related to the representations of the associated solution group; a finitely presented group with generators and relations reflected by the underlying linear system.

Together with his collaborators, the first author [1] constructed quantum isomorphic, but non-isomorphic graphs from linear system games having perfect quantum strategies but no such classical strategies. In this case, a quantum isomorphism corresponds to a perfect quantum strategy of the isomorphism game which was also introduced in [1].

Quantum automorphism groups of graphs were first introduced by Banica [2] and Bichon [5] to obtain examples of quantum permutation groups. They are generalizations of the automorphism group of a graph in the framework of Woronowicz' compact matrix quantum groups. An interesting question to ask is which quantum permutation groups can be realized as the quantum automorphism group of some graph. This question has for example been considered in [4].

In this work, we give a correspondence between representations of solution groups and quantum isomorphisms/quantum automorphisms of colored versions of the graphs appearing in [1]. More specifically, we will see that the quantum automorphism groups of the colored

<sup>\*</sup>davideroberson@gmail.com

<sup>&</sup>lt;sup>†</sup>sisc@math.ku.dk

versions of those graphs are given by the dual of the homogeneous solution group of the underlying linear constraint system. Furthermore, representations of the non-homogeneous solution group provide quantum isomorphisms between the colored versions of the graphs.

Additionally, we will discuss a decoloring procedure for vertex– and edge-colored graphs which does not change the quantum automorphism group for certain graphs. This procedure decolors the vertices by adding paths of different lengths to vertices of different colors. In a second step, we get rid of the edge-colors by subdividing the colored edges and then adding paths of different lengths to the newly added vertices.

The previous results allow us to construct two explicit examples of graphs that were not known before: First, we construct a graph that has quantum symmetry and finite quantum automorphism group. Second, we obtain a pair of graphs that are quantum isomorphic and non-isomorphic, where both graphs additionally do not have quantum symmetry.

The article is structured as follows. In Section 2, we briefly discuss compact quantum groups and give the definition of the quantum automorphism group of a vertex– and edgecolored graph. In Section 3, we define colored versions of the graphs appearing in [1]. Here we prove that the dual of the solution group is the quantum automorphism group of such a graph. Section 4 deals with decoloring the graphs from the previous section without changing the quantum automorphism group. In Section 5, we provide (uncolored) graphs whose quantum automorphism group is the dual of a solution group. We in particular obtain a graph with quantum symmetry and finite quantum automorphism group, see Corollary 5.6. Finally, we discuss quantum isomorphisms of the colored graphs in Section 6. Here, we present our example of a pair of graphs that are quantum isomorphic but non-isomorphic, where both graphs do not have quantum symmetry, see Corollary 6.10.

## 2 Preliminaries

We start with the definition of a compact quantum group, see [20]. Throughout this article, we write  $A \otimes B$  for the minimal tensor product of the  $C^*$ -algebras A and B.

**Definition 2.1.** A *compact quantum group*  $\mathbb{G}$  is a pair  $(C(\mathbb{G}), \Delta)$ , where  $C(\mathbb{G})$  is a unital  $C^*$ algebra and  $\Delta : C(\mathbb{G}) \to C(\mathbb{G}) \otimes C(\mathbb{G})$  is a unital \*-homomorphism with the following properties

- $(\Delta \otimes id) \circ \Delta = (id \otimes \Delta) \circ \Delta$  (coassociativity)
- $\Delta(C(\mathbb{G}))(1 \otimes C(\mathbb{G}))$  and  $\Delta(C(\mathbb{G}))(C(\mathbb{G}) \otimes 1)$  are dense in  $C(\mathbb{G}) \otimes C(\mathbb{G})$ .

**Definition 2.2.** Let  $\mathbb{G} = (C(\mathbb{G}), \Delta_{\mathbb{G}})$  and  $\mathbb{H} = (C(\mathbb{H}), \Delta_{\mathbb{H}})$  be compact quantum groups. Then  $\mathbb{G}$  and  $\mathbb{H}$  are isomorphic as compact quantum groups if there exists a \*-isomorphism  $\varphi : C(\mathbb{G}) \to C(\mathbb{H})$  such that  $\Delta_{\mathbb{H}} \circ \varphi = (\varphi \otimes \varphi) \circ \Delta_{\mathbb{G}}$ . We then write  $\mathbb{G} \cong \mathbb{H}$ .

**Definition 2.3.** Let  $\mathbb{G} = (C(\mathbb{G}), \Delta_{\mathbb{G}})$  be a compact quantum group. We say that  $\mathbb{G}$  is *finite* if the  $C^*$ -algebra  $C(\mathbb{G})$  is finite-dimensional.

The following example can for example be found in [11, Example 4.5].

**Example 2.4.** Let  $\Gamma$  be a discrete group. Note that the group  $C^*$ -algebra  $C^*(\Gamma)$  can be written as

$$C^{*}(\Gamma) = C^{*}(u_{g}, g \in \Gamma | u_{g} \text{ unitary, } u_{gh} = u_{g}u_{h}, u_{g^{-1}} = u_{g}^{*}),$$

see for example [18, A.6]. Let  $\Delta$  be the \*-homomorphism  $\Delta : C^*(\Gamma) \to C^*(\Gamma) \otimes C^*(\Gamma)$ ,  $\Delta(u_g) = u_g \otimes u_g$ . Then the pair  $(C^*(\Gamma), \Delta)$  is a compact quantum group known as the *dual* of  $\Gamma$  which we denote by  $\hat{\Gamma}$ .

We will now define compact matrix quantum groups, a subclass of compact quantum groups.

**Definition 2.5** ([19]). A compact matrix quantum group  $\mathbb{G}$  is a pair  $(C(\mathbb{G}), u)$ , where  $C(\mathbb{G})$  is a unital  $C^*$ -algebra and  $u = (u_{ij}) \in M_n(C(\mathbb{G}))$  is a matrix such that

- the elements  $u_{ij}$ ,  $1 \le i, j \le n$ , generate  $C(\mathbb{G})$ ,
- the \*-homomorphism  $\Delta : C(\mathbb{G}) \to C(\mathbb{G}) \otimes C(\mathbb{G}), u_{ij} \mapsto \sum_{k=1}^{n} u_{ik} \otimes u_{kj}$  exists,
- the matrix u and its transpose  $u^T$  are invertible.

The matrix u is usually called *fundamental representation* of  $\mathbb{G}$ .

A very important example is the quantum symmetric group, the quantum analogue of the symmetric group. It was defined by Wang in [17].

**Definition 2.6.** The quantum symmetric group  $S_n^+ = (C(S_n^+), u)$  is the compact matrix quantum group, where

$$C(S_n^+) := C^*(u_{ij}, 1 \le i, j \le n | u_{ij} = u_{ij}^* = u_{ij}^2, \sum_{k=1}^n u_{ik} = \sum_{k=1}^n u_{kj} = 1)$$

Note that a matrix  $u = (u_{ij})_{i,j \in [n]}$  with entries from a *nontrivial* unital  $C^*$ -algebra satisfying  $u_{ij} = u_{ij}^* = u_{ij}^2$  and  $\sum_{k=1}^n u_{ik} = \sum_{k=1}^n u_{kj} = 1$ , as in the definition above, is known as a *magic* unitary.

Quantum automorphism groups of finite graphs were defined in [2], [5]. We give a more general definition in the following, for vertex- and edge-colored graphs. Note that edgecolorings are similar to distances in finite quantum metric spaces, as considered in [3]. For us, graphs are undirected and do not have loops nor multiple edges. A colored graph is a graph G with vertex set V and edge set E along with a coloring function  $c: V \cup E \to S$  for some set S. We refer to c(x) as the color of the vertex/edge x. To be specific, we will sometimes refer to colored graphs as vertex- and edge-colored graphs. Furthermore, we will also consider edge-colored graphs where the coloring function is defined only on the edge set E, i.e., edges but not vertices are colored. In either case we will use  $E_c$  to refer to the set of edges of color *c*. Also,  $A_{G_c}$  will denote the adjacency matrix of the edge color *c*, i.e.,  $(A_{G_c})_{ij} = 1$  if  $(i, j) \in E_c$ and  $(A_{G_c})_{ij} = 0$  otherwise.

**Definition 2.7.** Let G be a vertex- and edge-colored graph. The quantum automorphism group Qut(G) is the compact matrix quantum group (C(Qut(G)), u), where C(Qut(G)) is the universal  $C^*$ -algebra with generators  $u_{ij}$ ,  $i, j \in V(G)$  and relations

$$u_{ij} = u_{ij}^* = u_{ij}^2, \qquad i, j \in V(G),$$
(1)

$$\sum u_{il} = 1 = \sum u_{li}, \qquad i \in V(G), \qquad (2)$$

$$u_{ij} = 0,$$
  

$$A_{G_c}u = uA_{G_c}$$
for all  $i, j \in V(G)$  with  $c(i) \neq c(j),$ 
for all edge colors  $c_i$ 
(4)

$$G_c u = u A_{G_c}$$
 for all edge colors  $c$ , (4)

where (4) is nothing but  $\sum_{k} u_{ik}(A_{G_c})_{kj} = \sum_{k} (A_{G_c})_{ik} u_{kj}$  for all  $i, j \in V(G)$  and all edge-colors c.

It is not immediately obvious that this defines a compact matrix quantum group. By Definition 2.5 we have to show that u and  $u^T$  are invertible as well as that the comultiplication  $\Delta : C(\operatorname{Qut}(G)) \to C(\operatorname{Qut}(G)) \otimes C(\operatorname{Qut}(G)), u_{ij} \mapsto \sum_k u_{ik} \otimes u_{kj}$  is a \*-homomorphism. It is enough to show  $\Delta(\sum_k u_{ik}(A_{G_c})_{kj}) = \Delta(\sum_k (A_{G_c})_{ik}u_{kj})$  for  $i, j \in V(G)$ , c edge-color and  $\Delta(u_{ij}) = 0$  for  $c(i) \neq c(j)$ , since we know that  $S_n^+$  is a compact matrix quantum group. The first equation follows for example from [15, Lemma 2.1.2]. For the second equation, we take vertices i, j with  $c(i) \neq c(j)$ . We have  $\Delta(u_{ij}) = \sum_k u_{ik} \otimes u_{kj}$ . Note that there is no  $k \in V(G)$  with c(k) = c(i) and c(k) = c(j), since we assumed  $c(i) \neq c(j)$ . We deduce  $u_{ik} \otimes u_{kj} = 0$  for all k and thus  $\Delta(u_{ij}) = 0$ . Summarizing, we see that Qut(G) is indeed a compact matrix quantum group.

The following lemma gives relations that are equivalent to Relation (4), see [15, Proposition 2.1.3].

**Lemma 2.8.** Let  $u_{ij}$ ,  $1 \le i, j \le n$ , be the generators of  $C(S_n^+)$ . Then Relation (4) is equivalent to the relations

 $u_{ij}u_{kl} = 0$  if  $(i, k) \in E_c$ ,  $(j, l) \notin E_c$  for some edge-color c,  $u_{ij}u_{kl} = 0$  if  $(i, k) \notin E_c$ ,  $(j, l) \in E_c$  for some edge-color c.

## 3 Colored graphs whose quantum automorphism group is the dual of a solution group

In the following definition we use 1 to denote the identity element of a group.

**Definition 3.1.** Let  $M \in \mathbb{F}_2^{m \times n}$  and  $b \in \mathbb{F}_2^m$  with  $b \neq 0$ . The solution group  $\Gamma(M, b)$  of the linear system Mx = b is the group generated by elements  $x_i$  for  $i \in [n]$  and an element  $\gamma$  satisfying the following relations:

1. 
$$x_i^2 = 1$$
 for all  $i \in [n]$ ;

2.  $x_i x_j = x_j x_i$  if there exists  $k \in [m]$  s.t.  $M_{ki} = M_{kj} = 1$ ;

3. 
$$\prod_{i:M_{k}=1} x_i = \gamma^{b_k}$$
 for all  $k \in [m]$ ;

4. 
$$\gamma^2 = 1$$

5.  $x_i \gamma = \gamma x_i$  for all  $i \in [n]$ .

If b = 0, then we refer to  $\Gamma(M, b)$  as the *homogeneous solution group* of the system Mx = 0, and define this the same as above except that we add the relation  $\gamma = 1$ . This is equivalent to removing  $\gamma$  from the list of generators, changing the righthand side of the equation in (3) to 1, and removing items (4) and (5). We will sometimes also use  $\Gamma_0(M)$  to denote this group.

We will denote by  $S_k(M)$  the set  $\{i \in [n] : M_{ki} = 1\}$ , often writing simply  $S_k$  when M is clear from context. We use  $\pm 1^S$  to denote the set of functions  $\alpha : S \to \{1, -1\}$ , and will typically write  $\alpha_i$  instead of  $\alpha(i)$ . We will also use  $\pm 1^S_0$  to denote the subset of such functions satisfying  $\prod_{i \in S} \alpha_i = 1$ , and similarly use  $\pm 1^S_1$  for the set of such functions satisfying  $\prod_{i \in S} \alpha_i = -1$ .

**Definition 3.2.** Let  $M \in \mathbb{F}_2^{m \times n}$  and  $b \in \mathbb{F}_2^m$ . Define the colored graph G := G(M, b) as follows. The vertex set of G is  $\{(k, \alpha) : k \in [m], \alpha \in \pm 1_{b_k}^{S_k}\}$ . The color of a vertex  $v = (k, \alpha)$ , denoted c(v), is k. We also define the subsets  $V_k = \{v \in V(G) : c(v) = k\}$  for all  $k \in [m]$ , and these partition the vertex set. For any  $l, k \in [m]$  such that  $S_l \cap S_k \neq \emptyset$ , the graph G contains all (non-loop) edges between  $V_l$  and  $V_k$  (thus each  $V_k$  induces a complete subgraph). Given an edge e between adjacent vertices  $(l, \alpha)$  and  $(k, \beta)$ , the color of e, denoted c(e), is equal to the function  $\alpha \Delta \beta \in \pm 1^{S_l \cap S_k}$  defined as  $(\alpha \Delta \beta)_i = \alpha_i \beta_i$ .

Note that  $\alpha \triangle \beta = \beta \triangle \alpha$  and so the edge colors defined above really are edge colors and not arc (directed edge) colors.

**Remark 3.3.** According to the above definition, it is possible for two edges e and e' between pairs of vertices  $(l, \alpha), (k, \beta)$  and  $(l', \alpha'), (k', \beta')$  to be colored the same color even if l is not equal to either l' nor k', i.e., the edges are between different pairs of subsets  $V_l, V_k$  and  $V_{l'}, V_{k'}$ . This can happen since it is possible that  $S_l \cap S_k = S_{l'} \cap S_{k'}$  and  $\alpha \Delta \beta = \alpha' \Delta \beta'$ . However, we do wish such edges to be distinguished by the (quantum) automorphism group of G(M, b), i.e.,  $u_{(l,\alpha),(l'\alpha')}u_{(k,\beta),(k'\beta')} = 0$  where u is the fundamental representation of  $Qut(G(M, b))^1$ . This could be done explicitly by defining the color of the edge e (for instance) to be c(e) = $(\{l,k\}, \alpha \Delta \beta)$ . However, this is redundant for our purposes since such edges e and e' as described above are already distinguished by the (quantum) automorphism group due to the vertex colors of the endpoints of the edges. In other words, the edges between  $V_l$  and  $V_k$  can be thought of as being *implicitly* colored distinctly from those between  $V_{l'}$  and  $V_{k'}$  whenever  $\{l, k\} \neq \{l', k'\}$ .

**Remark 3.4.** We also note that in order to reduce the total number of colors used, for each pair  $l, k \in [m]$ , we can choose one color  $\alpha \in \pm 1^{S_l \cap S_k}$  and replace these edges with non-edges. Moreover, instead of coloring the edges between  $V_l$  and  $V_k$  with functions from  $\pm 1^{S_l \cap S_k}$ , we can simply use  $\{1, \ldots, 2^{|S_l \cap S_k|} - 1\}$ , and this will not change Qut(G) as explained in the previous remark. In fact we will need to do this for some of our results later on.

**Example 3.5.** Consider the linear system Mx = b, where

$$M = \begin{pmatrix} 1 & 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 1 \end{pmatrix}, b = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

Then the graph G(M, b) is the one given in Figure 1.




Figure 1: The graph G(M, b) for M and b as above. The vertices on the left hand side are the solutions of  $x_1x_2x_3 = 1$ , the vertices on the right hand side are the solutions of  $x_1x_4x_5 = -1$ . We used Remarks 3.3 and 3.4 to reduce the number of colors needed in the graph.

**Lemma 3.6.** Let  $M \in \mathbb{F}_2^{m \times n}$  and  $b \in \mathbb{F}_2^m$ . Set G = G(M, b). If u is the fundamental representation of Qut(G), then u has the form

where each  $u^{(k)}$  is a magic unitary. Furthermore,  $u^{(k)}_{\alpha,\beta} := u_{(k,\alpha),(k,\beta)}$  depends only on k and the value of  $\alpha \triangle \beta$  and therefore the entries of  $u^{(k)}$  pairwise commute for each  $k \in [m]$ .

<sup>&</sup>lt;sup>1</sup>This is the quantum analog of there being no automorphism mapping the edge  $(l, \alpha)(k, \beta)$  to  $(l'\alpha')(k', \beta')$ .

*Proof.* If  $l \neq k$ , then  $c((l, \alpha)) = l \neq k = c((k, \beta))$  and thus  $u_{(l,\alpha),(k,\beta)} = 0$ . This shows that u has the block form given in the lemma statement. Moreover, each diagonal block must be a magic unitary since u is a magic unitary.

Now fix  $k \in [m]$ . Pick  $\tilde{\alpha}, \tilde{\beta} \in \pm 1_{b_k}^{S_k}$  arbitrarily. Suppose that  $\alpha, \beta \in \pm 1_{b_k}^{S_k}$  are such that  $\alpha \Delta \beta = \tilde{\alpha} \Delta \tilde{\beta}$ . Note that since all these functions are elements of  $\pm 1_{b_k}^{S_k}$ , the operation  $\Delta$  is simply the pointwise product, and therefore we have that  $\tilde{\alpha} \Delta \alpha = \tilde{\beta} \Delta \beta$ . In particular, this implies that the color of the edge between  $(k, \tilde{\alpha})$  and  $(k, \alpha)$  is equal to the color of the edge between  $(k, \tilde{\beta})$  and  $(k, \beta')$  if and only if  $\beta' = \beta$ . Therefore,  $u_{\tilde{\alpha}, \tilde{\beta}}^{(k)} u_{\alpha, \beta'}^{(k)} = 0$  unless  $\beta' = \beta$ . We similarly have that  $u_{\tilde{\alpha}, \tilde{\beta}'}^{(k)} u_{\alpha, \beta}^{(k)} = 0$  unless  $\tilde{\beta}' = \tilde{\beta}$ . It follows that

$$u_{\tilde{\alpha},\tilde{\beta}}^{(k)} = u_{\tilde{\alpha},\tilde{\beta}}^{(k)} \left( \sum_{\beta' \in \pm 1_{b_k}^{S_k}} u_{\alpha,\beta'}^{(k)} \right) = u_{\tilde{\alpha},\tilde{\beta}}^{(k)} u_{\alpha,\beta}^{(k)} = \left( \sum_{\tilde{\beta}' \in \pm 1_{b_k}^{S_k}} u_{\tilde{\alpha},\tilde{\beta}'}^{(k)} \right) u_{\alpha,\beta}^{(k)} = u_{\alpha,\beta}^{(k)}$$

Thus the operator  $u_{\alpha,\beta}^{(k)}$  depends only on the value of  $\alpha \triangle \beta$  (and k) as claimed. Let us therefore use  $u_{\delta}^{(k)}$  to denote  $u_{\alpha,\beta}^{(k)}$  where  $\delta = \alpha \triangle \beta$ . For any fixed  $\alpha \in \pm 1_{b_k}^{S_k}$ , the set  $\{\alpha \triangle \beta : \beta \in \pm 1_{b_k}^{S_k}\}$  is equal to  $\pm 1_0^{S_k}$ . Thus the set of operators appearing in any row (and similarly any column) of  $u^{(k)}$  is  $\{u_{\delta}^{(k)} : \delta \in \pm 1_0^{S_k}\}$ . Since the entries of any given row of a magic unitary commute, and every row of  $u^{(k)}$  contains the same set of entries, we have that all entries of  $u^{(k)}$  commute.  $\Box$ 

**Remark 3.7.** Let  $\Gamma = \Gamma(M, 0)$  be the homogeneous solution group of the system Mx = 0. By Example 2.4, the group  $C^*$ -algebra  $C^*(\Gamma)$  can be written as

$$C^{*}(\Gamma) = C^{*}(x_{i}, i \in [n] | x_{i} = x_{i}^{*}, x_{i}^{2} = 1, x_{i}x_{j} = x_{j}x_{i} \text{ if there exists } k \in [m] \text{ s.t. } M_{ki} = M_{kj} = 1,$$
$$\prod_{i:M_{ki}=1} x_{i} = 1 \text{ for all } k \in [m]).$$

Note that we slightly abuse the notation by writing  $x_i$  instead of  $u_{x_i}$ .

**Theorem 3.8.** Let  $M \in \mathbb{F}_2^{m \times n}$  and  $b \in \mathbb{F}_2^m$ . Set G = G(M, b) and  $\Gamma = \Gamma(M, 0) = \Gamma_0(M)$ . Let  $\Delta_{\Gamma} : C^*(\Gamma) \to C^*(\Gamma) \otimes C^*(\Gamma)$  denote the coproduct given by  $\Delta_{\Gamma}(g) = g \otimes g$  for  $g \in \Gamma$ , and let  $\Delta_G : C(\operatorname{Qut}(G)) \to C(\operatorname{Qut}(G)) \otimes C(\operatorname{Qut}(G))$  denote the coproduct given by  $\Delta_G(u_{a,c}) = \sum_{b \in V(G)} u_{a,b} \otimes u_{b,c}$  where  $u = (u_{a,c})$  is the fundamental representation of  $\operatorname{Qut}(G)$ . Then there exists an isomorphism  $\varphi : C^*(\Gamma) \to C(\operatorname{Qut}(G))$  such that  $\Delta_G \circ \varphi = (\varphi \otimes \varphi) \circ \Delta_{\Gamma}$ .

*Proof.* The proof is structured as follows: we use the universal properties of  $C^*(\Gamma)$  and C(Qut(G)) respectively to first obtain a \*-homomorphism  $\varphi_1 : C^*(\Gamma) \to C(\text{Qut}(G))$ , and then obtain a \*-homomorphism  $\varphi_2 : C(\text{Qut}(G)) \to C^*(\Gamma)$ . We will then show that  $\varphi_1$  and  $\varphi_2$  are inverses of each other thus proving that they are in fact isomorphisms. Lastly, we will prove that  $\varphi_1$  intertwines the coproducts as described in the theorem statement.

Step 1: Construction of a \*-homomorphism  $\varphi_1 : C^*(\Gamma) \to C(Qut(G))$ .

For this step, we will construct elements  $y_i$  of C(Qut(G)) that satisfy the relations of the generators  $x_i$  of  $C^*(\Gamma)$  given in Remark 3.7. This proves that there is a \*-homomorphism  $\varphi_1$  from  $C^*(\Gamma)$  to C(Qut(G)) such that  $\varphi_1(x_i) = y_i$ . Later we will see that  $\varphi_1$  is in fact an isomorphism.

Let u be the fundamental representation of  $\operatorname{Qut}(G)$ . By Lemma 3.6,  $u = \bigoplus_{k \in [m]} u^{(k)}$  where each  $u^{(k)}$  is a magic unitary indexed by  $V_k = \{(k, \alpha) : \alpha \in \pm 1_{b_k}^{S_k}\}$ . Moreover, the operator  $u_{\alpha,\beta}^{(k)} := u_{(k,\alpha),(k,\beta)}$  depends only on k and the value of  $\alpha \Delta \beta \in \pm 1_0^{S_k}$ . Thus, as in the proof of Lemma 3.6, for each  $\delta \in \pm 1_0^{S_k}$  we let  $u_{\delta}^{(k)}$  denote  $u_{\alpha,\beta}^{(k)}$  such that  $\alpha \Delta \beta = \delta$ . Note that for any  $\alpha \in \pm 1_{b_k}^{S_k}$ , we have that the set  $\{\alpha \triangle \beta : \beta \in \pm 1_{b_k}^{S_k}\}$  is equal to  $\pm 1_0^{S_k}$  and thus every row/column of  $u^{(k)}$  contains the same set of operators and

$$\sum_{\delta \in \pm 1_0^{S_k}} u_{\delta}^{(k)} = \sum_{\beta \in \pm 1_{b_k}^{S_k}} = \sum_{\alpha \in \pm 1_{b_k}^{S_k}} u_{\alpha,\beta}^{(k)} = 1.$$
(6)

Now we define  $y_i^{(k)} = \sum_{\alpha \in \pm 1_{\alpha}^{S_k}} \alpha_i u_{\alpha}^{(k)}$  for all  $k \in [m]$  and  $i \in S_k$ . We first aim to show that  $y_i^{(k)}$  does not depend on k.

Fix  $i \in [n]$  and  $l, k \in [m]$  such that  $i \in S_l \cap S_k$ . Consider the edges between the subsets  $V_l$  and  $V_k$ . For each  $\delta \in \pm 1^{S_l \cap S_k}$ , let  $A^{\delta}$  be the adjacency matrix of the graph consisting of the edges of G colored  $\delta$ . Further, let  $B^{\delta}$  be the submatrix of  $A^{\delta}$  consisting of the rows indexed by  $V_l$  and columns indexed by  $V_k$ . In other words,

$$B^{\delta}_{(l,lpha),(k,eta)} = egin{cases} 1 & ext{if } lpha riangle eta \ 0 & ext{o.w.} \end{cases}$$

Also define  $A^+ = \sum_{\delta \in \pm 1^{S_l \cap S_k}, \delta_i = +1} A^{\delta}$  and  $A^- = \sum_{\delta \in \pm 1^{S_l \cap S_k}, \delta_i = -1} A^{\delta}$ , and similarly define  $B^+$ and  $B^-$ . Then

$$B_{(l,\alpha),(k,\beta)}^{+} = \begin{cases} 1 & \text{if } \alpha_i = \beta_i \\ 0 & \text{o.w.} \end{cases} \quad \& \quad B_{(l,\alpha),(k,\beta)}^{-} = \begin{cases} 1 & \text{if } \alpha_i = -\beta_i \\ 0 & \text{o.w.} \end{cases}$$

Since u must commute with each  $A^{\delta}$  by definition of Qut(G), it must also commute with both  $A^+$  and  $A^-$ . This implies that  $u^{(l)}B^+ = B^+u^{(k)}$  and  $u^{(l)}B^- = B^-u^{(k)}$ . Considering the  $(l, \alpha), (k, \beta)$  entry of both sides of the former equation, we see that

$$\sum_{\substack{\alpha' \in \pm 1_{b_l}^{S_l} \\ \text{s.t. } \alpha'_i = \beta_i}} u_{\alpha,\alpha'}^{(l)} = \sum_{\substack{\beta' \in \pm 1_{b_k}^{S_k} \\ \text{s.t. } \beta'_i = \alpha_i}} u_{\beta',\beta}^{(k)}$$

Note that for every term in the above sums, we have that  $\alpha_i \alpha'_i = \alpha_i \beta_i = \beta'_i \beta_i$ . Thus, reexpressing  $u_{\alpha,\alpha'}^{(l)}$  as  $u_{\tilde{\alpha}}^{(l)}$  where  $\tilde{\alpha} = \alpha \triangle \alpha'$ , and similarly for  $u_{\beta',\beta}^{(k)}$  we have that

$$\sum_{\substack{\tilde{\alpha} \in \pm \mathbf{1}_{0}^{S_{l}} \\ \text{s.t. } \tilde{\alpha}_{i} = \alpha_{i}\beta_{i}}} u_{\tilde{\alpha}}^{(l)} = \sum_{\substack{\tilde{\beta} \in \pm \mathbf{1}_{0}^{S_{k}} \\ \text{s.t. } \tilde{\beta}_{i} = \alpha_{i}\beta_{i}}} u_{\tilde{\beta}}^{(k)}$$

Doing the same for  $u^{(l)}B^- = B^- u^{(k)}$  yields the same equation but with  $\alpha_i \beta_i$  replaced with  $-\alpha_i\beta_i$  and combining these proves that

$$\sum_{\tilde{\alpha} \in \pm 1_0^{S_l}, \tilde{\alpha}_i = x} u_{\tilde{\alpha}}^{(l)} = \sum_{\tilde{\beta} \in \pm 1_0^{S_k}, \tilde{\beta}_i = x} u_{\tilde{\beta}}^{(k)}$$

for any  $x \in \{+1, -1\}$ . Therefore,

$$y_{i}^{(l)} = \left(\sum_{\substack{\alpha \in \pm 1_{0}^{S_{l}} \\ \text{s.t. } \alpha_{i} = +1}} u_{\alpha}^{(l)}\right) - \left(\sum_{\substack{\alpha \in \pm 1_{0}^{S_{l}} \\ \text{s.t. } \alpha_{i} = -1}} u_{\alpha}^{(l)}\right) = \left(\sum_{\substack{\alpha \in \pm 1_{0}^{S_{k}} \\ \text{s.t. } \alpha_{i} = +1}} u_{\alpha}^{(k)}\right) - \left(\sum_{\substack{\alpha \in \pm 1_{0}^{S_{k}} \\ \text{s.t. } \alpha_{i} = -1}} u_{\alpha}^{(k)}\right) = y_{i}^{(k)}.$$

So we have shown that the value of  $y_i^{(k)}$  does not depend on k, and thus we will simply denote this operator by  $y_i$ .

Now note that since  $y_i$  is a linear combination (with real coefficients) of the operators  $u_{\alpha}^{(k)}$  for  $k \in [m]$  such that  $i \in S_k$ , and these operators are entries of the magic unitary u, we have that  $y_i^* = y_i$ . Also, by Equation (6) we have that  $u_{\alpha}^{(k)}u_{\beta}^{(k)} = 0$  for  $\alpha \neq \beta$  and thus

$$y_i^2 = \left(\sum_{\alpha \in \pm 1_0^{S_k}} \alpha_i u_{\alpha}^{(k)}\right)^2 = \sum_{\alpha \in \pm 1_0^{S_k}} \alpha_i^2 \left(u_{\alpha}^{(k)}\right)^2 = \sum_{\alpha \in \pm 1_0^{S_k}} u_{\alpha}^{(k)} = 1.$$

പ

Thus the  $y_i$  satisfy relation (1) from Definition 3.1.

Next we will show that relation (2) of Definition 3.1 holds, i.e., that  $y_i y_j = y_j y_i$  if there exists  $k \in [m]$  such that  $i, j \in S_k$ . Suppose that i, j, k are as described. Then  $y_i = y_i^{(k)}$  and  $y_j = y_j^{(k)}$  are both linear combinations of the entries of  $u^{(k)}$  which pairwise commute by Lemma 3.6. Therefore  $y_i$  and  $y_j$  commute as desired.

Lastly, we must show that relation (3) of Definition 3.1 holds. Recall that we are trying to show that  $C^*(\Gamma_0(M)) \cong C(\text{Qut}(G))$ , i.e., we are in the *homogeneous* solution group case. Thus we must show that  $\prod_{i \in S_k} y_i = 1$  for all  $k \in [m]$ . We have that

$$\prod_{i \in S_k} y_i = \prod_{i \in S_k} y_i^{(k)}$$
$$= \prod_{i \in S_k} \left( \sum_{\alpha \in \pm 1_0^{S_k}} \alpha_i u_\alpha^{(k)} \right)$$

Since the elements  $u_{\alpha}^{(k)}$  are pairwise orthogonal, when we expand the above product all cross terms disappear and we obtain

$$\prod_{i \in S_k} y_i = \sum_{\alpha \in \pm 1_0^{S_k}} \left( \prod_{i \in S_k} \alpha_i \right) u_\alpha^{(k)} = \sum_{\alpha \in \pm 1_0^{S_k}} u_\alpha^{(k)} = 1,$$

as desired. Therefore the elements  $y_i \in C(\text{Qut}(G))$  for  $i \in [n]$  satisfy the relations of the generators of  $C^*(\Gamma)$  and thus there exists a \*-homomorphism  $\varphi_1$  from  $C^*(\Gamma)$  to C(Qut(G)) such that  $\varphi_1(x_i) = y_i$  for all  $i \in [n]$ .

Step 2: Construction of a \*-homomorphism  $\varphi_2 : C(\operatorname{Qut}(G)) \to C^*(\Gamma)$ .

For this step, we will construct elements  $v_{(l,\alpha),(k,\beta)}$  of  $C^*(\Gamma)$  that satisfy all the relations of the generators  $u_{(l,\alpha),(k,\beta)}$  of C(Qut(G)). This proves that there is a \*-homomorphism  $\varphi_2$  from C(Qut(G)) to  $C^*(\Gamma)$  such that  $\varphi_2(u_{(l,\alpha),(k,\beta)}) = v_{(l,\alpha),(k,\beta)}$ . Later we will see that  $\varphi_2$  is an isomorphism.

For each  $i \in [n]$ , define  $p_i^{\pm} = \frac{1}{2}(1 \pm x_i)$ . It is straightforward to check that  $p_i^{\pm}$  is a projection, i.e.,  $(p_i^{\pm})^* = p_i^{\pm} = (p_i^{\pm})^2$ , and that  $x_i = p_i^+ - p_i^-$  and  $1 = p_i^+ + p_i^-$ . Note that the latter implies that  $p_i^+$  and  $p_i^-$  are orthogonal, i.e.,  $p_i^+ p_i^- = 0$ . We will abuse notation somewhat and write  $p_i^{\alpha_i}$  to denote  $p_i^+$  whenever  $\alpha_i = +1$  and similarly for  $p_i^-$  when  $\alpha_i = -1$ . Next, for all  $k \in [m]$ and  $\alpha \in \pm 1^{S_k}$  define  $v_{\alpha}^{(k)} = \prod_{i \in S_k} p_i^{\alpha_i}$ . Note that the  $v_{\alpha}^{(k)}$  are indeed well defined since the commutativity of the elements  $p_i^{\pm}$  for  $i \in S_k$  follows from the commutativity of the  $x_i$  for  $i \in S_k$  (i.e., relation (2) from Definition 3.1). Since it is the product of pairwise commuting projections, we have that  $v_{\alpha}^{(k)}$  is a projection. Furthermore,

$$\sum_{\alpha \in \pm 1^{S_k}} v_{\alpha}^{(k)} = \sum_{\alpha \in \pm 1^{S_k}} \prod_{i \in S_k} p_i^{\alpha_i} = \prod_{i \in S_k} (p_i^+ + p_i^-) = \prod_{i \in S_k} 1 = 1.$$
(7)

This also implies that for a fixed  $k \in [m]$  the projections  $v_{\alpha}^{(k)}$  are pairwise orthogonal.

Now suppose that  $\alpha \in \pm 1_1^{S_k}$ , i.e., that  $\prod_{i \in S_k} \alpha_i = -1$ . Then, using the easily checked fact that  $x_i p_i^{\alpha_i} = \alpha_i p_i^{\alpha_i}$ , we have that

$$v_{\alpha}^{(k)} = \left(\prod_{i \in S_k} x_i\right) v_{\alpha}^{(k)} = \prod_{i \in S_k} x_i p_i^{\alpha_i} = \prod_{i \in S_k} \alpha_i p_i^{\alpha_i} = \left(\prod_{i \in S_k} \alpha_i\right) v_{\alpha}^{(k)} = -v_{\alpha}^{(k)}.$$
(8)

Thus  $v_{\alpha}^{(k)} = 0$  for all  $\alpha \in \pm 1_1^{S_k}$ . Combining this with Equation (7), we have that

$$\sum_{\alpha \in \pm 1_0^{S_k}} v_{\alpha}^{(k)} = 1.$$
(9)

Next, for all  $k \in [m]$  and  $\alpha, \beta \in \pm 1_{b_k}^{S_k}$  we define  $v_{\alpha,\beta}^{(k)} := v_{\delta}^{(k)}$  where  $\delta = \alpha \triangle \beta$ . Lastly, define v to the the matrix indexed by V(G) such that

$$v_{(l,\alpha),(k,\beta)} = \begin{cases} v_{\alpha,\beta}^{(l)} & \text{if } l = k \\ 0 & \text{o.w.} \end{cases}$$

We aim to show that v satisfies all of the relations satisfied by the fundamental representation u of  $\operatorname{Qut}(G)$ . First, v is a magic unitary since all of its entries are projections and the sum of its  $(l, \alpha)$ -row is simply  $\sum_{\beta \in \pm 1_{b_l}^{S_l}} v_{\alpha,\beta}^{(l)} = \sum_{\delta \in \pm 1_0^{S_l}} v_{\delta}^{(l)} = 1$ . Next we must show that for  $a, b \in V(G)$ , we have  $v_{a,b} = 0$  whenever  $c(a) \neq c(b)$ , i.e., the colors of a and b are different. Recall from the definition of G(M, b) that the color of the vertex  $(l, \alpha)$  is l. Thus v satisfies this relation by definition. Lastly, we must show that for  $a, b, a', b' \in V(G)$ , we have  $v_{a,b}v_{a',b'} = 0$  whenever the edges  $\{a, a'\}$  and  $\{b, b'\}$  have different colors (or when one is an edge and the other is not). The previously shown relation already implies this one unless c(a) = c(b) and c(a') = c(b'). So we may assume that  $a = (l, \alpha), b = (l, \beta), a' = (k, \alpha')$ , and  $b' = (k, \beta')$ . Thus, letting  $\delta = \alpha \Delta \beta$  and  $\delta' = \alpha' \Delta \beta'$ , we have  $v_{a,b} = v_{\alpha,\beta}^{(l)} = v_{\delta}^{(l)}$  and  $v_{a',b'} = v_{\alpha',\beta}^{(k)} = v_{\delta'}^{(k)}$ . Suppose the edges  $\{a, a'\}$  and  $\{b, b'\}$  do have different colors. By definition of G(M, b) this is equivalent to  $\alpha \Delta \alpha' \neq \beta \Delta \beta'$ . This means there exists  $j \in S_l \cap S_k$  such that  $\alpha_j \alpha'_j \neq \beta_j \beta'_j$ . Since all these terms are in  $\{+1, -1\}$ , we have that  $\delta_j = \alpha_j \beta_j \neq \alpha'_j \beta'_j = \delta'_j$ . This implies that  $p_j^{\delta_j} p_j^{\delta_j} = 0$ . Therefore,

$$v_{a,b}v_{a',b'} = v_{\delta}^{(l)}v_{\delta'}^{(k)} = \left(\prod_{i\in S_l} p_i^{\delta_i}\right) \left(\prod_{i\in S_k} p_i^{\delta'_i}\right) = p_j^{\delta_j}p_j^{\delta'_j} \left(\prod_{i\in S_l\setminus\{j\}} p_i^{\delta_i}\right) \left(\prod_{i\in S_k\setminus\{j\}} p_i^{\delta'_i}\right) = 0.$$

So we have shown that v is a magic unitary satisfying all of the relations satisfied by u and thus by the universal property of C(Qut(G)), there exists a \*-homomorphism  $\varphi_2$  from C(Qut(G))to  $C^*(\Gamma)$  such that  $\varphi_2(u_{(l,\alpha),(k,\beta)}) = v_{(l,\alpha),(k,\beta)}$ , which is equivalent to  $\varphi_2\left(u_{\delta}^{(k)}\right) = v_{\delta}^{(k)}$  for all  $k \in [m]$  and  $\delta \in \pm 1_0^{S_k}$ .

*Step 3: Showing that*  $\varphi_1$  *and*  $\varphi_2$  *are inverses of each other.* 

We first show that  $\varphi_2 \circ \varphi_1 : C^*(\Gamma) \to C^*(\Gamma)$  is the identity. Since both  $\varphi_1$  and  $\varphi_2$  are \*-homomorphisms, it suffices to show that  $\varphi_2 \circ \varphi_1$  acts as the identity on the generators  $x_i$  of  $C^*(\Gamma)$ . From Step 1 we have that  $\varphi_1(x_i) = y_i = \sum_{\alpha \in \pm 1_0^{S_k}} \alpha_i u_{\alpha}^{(k)}$  for any  $k \in [m]$  such that  $i \in S_k$ . From Step 2 we have that  $\varphi_2(u_{\alpha}^{(k)}) = v_{\alpha}^{(k)} = \prod_{i \in S_k} p_i^{\alpha_i}$ . Therefore,

$$\varphi_2 \circ \varphi_1(x_i) = \sum_{\alpha \in \pm 1_0^{S_k}} \alpha_i \varphi_2(u_\alpha^{(k)}) = \sum_{\alpha \in \pm 1_0^{S_k}} \alpha_i v_\alpha^{(k)}.$$

To see that the final expression above is equal to  $x_i$ , we will show that multiplying it by  $x_i$  yields 1. First, note that since  $x_i p_i^{\alpha_i} = \alpha_i p_i^{\alpha_i}$ , we have that

$$x_i v_{\alpha}^{(k)} = x_i \prod_{j \in S_k} p_j^{\alpha_j} = x_i p_i^{\alpha_i} \prod_{j \in S_k \setminus \{i\}} p_j^{\alpha_j} = \alpha_i p_i^{\alpha_i} \prod_{j \in S_k \setminus \{i\}} p_j^{\alpha_j} = \alpha_i \prod_{j \in S_k} p_j^{\alpha_j} = \alpha_i v_{\alpha}^{(k)}.$$

Thus, making use of Equation (9) in the last equality, we have that

$$x_i(\varphi_2 \circ \varphi_1(x_i)) = x_i\left(\sum_{\alpha \in \pm 1_0^{S_k}} \alpha_i v_{\alpha}^{(k)}\right) = \sum_{\alpha \in \pm 1_0^{S_k}} \alpha_i x_i v_{\alpha}^{(k)} = \sum_{\alpha \in \pm 1_0^{S_k}} \alpha_i^2 v_{\alpha}^{(k)} = \sum_{\alpha \in \pm 1_0^{S_k}} v_{\alpha}^{(k)} = 1,$$

as desired. It follows that  $\varphi_2 \circ \varphi_1(x_i)$  is the inverse of  $x_i$ , which is its own inverse by definition. Therefore  $\varphi_2 \circ \varphi_1(x_i) = x_i$  and thus  $\varphi_2 \circ \varphi_1$  is the identity map on  $C^*(\Gamma)$ .

Now we will show that  $\varphi_1 \circ \varphi_2 : C(\operatorname{Qut}(G)) \to C(\operatorname{Qut}(G))$  is the identity map. As above, it suffices to show that  $\varphi_1 \circ \varphi_2$  acts as the identity on all entries of the fundamental representation u. Other than 0, the entries of u are precisely the elements  $u_{\alpha}^{(k)}$  for  $k \in [m]$  and  $\alpha \in \pm 1_0^{S_k}$ . From Step 2 we have that  $\varphi_2(u_{\alpha}^{(k)}) = v_{\alpha}^{(k)} = \prod_{i \in S_k} p_i^{\alpha_i} = \prod_{i \in S_k} \frac{1}{2}(1 + \alpha_i x_i)$ , and from Step 1 we have that  $\varphi_1(x_i) = y_i = \sum_{\alpha \in \pm 1_0^{S_k}} \alpha_i u_{\alpha}^{(k)}$ . Therefore,

$$\varphi_1 \circ \varphi_2(u_{\alpha}^{(k)}) = \varphi_1\left(\prod_{i \in S_k} \frac{1}{2}(1 + \alpha_i x_i)\right) = \prod_{i \in S_k} \frac{1}{2}(1 + \alpha_i y_i),$$
(10)

where we have used the fact that  $\varphi_1$  maps  $1 \in C^*(\Gamma)$  to  $1 \in C(\operatorname{Qut}(G))$  which follows from  $\varphi_1(1) = \varphi_1(x_i^2) = \varphi_1(x_i)^2 = y_i^2 = 1$ . Let us use  $\tilde{u}_{\alpha}^{(k)}$  to denote the final expression of Equation (10). We wish to show that  $\tilde{u}_{\alpha}^{(k)} = u_{\alpha}^{(k)}$ . Since the  $y_i$  satisfy all of the relations that the  $x_i$  satisfy, the same argument as for the  $v_{\alpha}^{(k)}$  can be used to show that, for fixed  $k \in [m]$  the  $\tilde{u}_{\alpha}^{(k)}$  are pairwise orthogonal projections satisfying  $\sum_{\alpha \in \pm 1_0^{S_k}} \tilde{u}_{\alpha}^{(k)} = 1$ . For  $i \in S_k$  and  $\beta \in \pm 1_0^{S_k}$  we have that

$$u_{\beta}^{(k)}y_{i} = u_{\beta}^{(k)}\sum_{\alpha \in \pm 1_{0}^{S_{k}}} \alpha_{i}u_{\alpha}^{(k)} = \beta_{i}u_{\beta}^{(k)}$$

Suppose that  $\alpha \neq \beta$ . Then there exists  $j \in S_k$  such that  $\alpha_j \neq \beta_j$  and therefore

$$\begin{split} u_{\beta}^{(k)} \tilde{u}_{\alpha}^{(k)} &= u_{\beta}^{(k)} \prod_{i \in S_{k}} \frac{1}{2} (1 + \alpha_{i} y_{i}) \\ &= u_{\beta}^{(k)} \frac{1}{2} (1 + \alpha_{j} y_{j}) \prod_{i \in S_{k} \setminus \{j\}} \frac{1}{2} (1 + \alpha_{i} y_{i}) \\ &= \frac{1}{2} (u_{\beta}^{(k)} + \alpha_{j} u_{\beta}^{(k)} y_{j}) \prod_{i \in S_{k} \setminus \{j\}} \frac{1}{2} (1 + \alpha_{i} y_{i}) \\ &= \frac{1}{2} (u_{\beta}^{(k)} + \alpha_{j} \beta_{j} u_{\beta}^{(k)}) \prod_{i \in S_{k} \setminus \{j\}} \frac{1}{2} (1 + \alpha_{i} y_{i}) \\ &= \frac{1}{2} (u_{\beta}^{(k)} - u_{\beta}^{(k)}) \prod_{i \in S_{k} \setminus \{j\}} \frac{1}{2} (1 + \alpha_{i} y_{i}) \\ &= \frac{1}{2} (u_{\beta}^{(k)} - u_{\beta}^{(k)}) \prod_{i \in S_{k} \setminus \{j\}} \frac{1}{2} (1 + \alpha_{i} y_{i}) \\ &= 0 \end{split}$$

where the penultimate equality comes from the fact that  $\alpha_j \neq \beta_j$  implies that  $\alpha_j \beta_j = -1$ . Using the above we have that

$$u_{\alpha}^{(k)} = u_{\alpha}^{(k)} \sum_{\beta \in \pm 1_0^{S_k}} \tilde{u}_{\beta}^{(k)} = u_{\alpha}^{(k)} \tilde{u}_{\alpha}^{(k)} = \sum_{\beta \in \pm 1_0^{S_k}} u_{\beta}^{(k)} \tilde{u}_{\alpha}^{(k)} = \tilde{u}_{\alpha}^{(k)}.$$

This completes the proof that  $\varphi_1 \circ \varphi_2(u_{\alpha}^{(k)}) = u_{\alpha}^{(k)}$  and thus that  $\varphi_1 \circ \varphi_2$  is the identity on  $C(\operatorname{Qut}(G))$ . Combining with the above, this proves that  $\varphi_1 : C^*(\Gamma) \to C(\operatorname{Qut}(G))$  and  $\varphi_2 : C(\operatorname{Qut}(G)) \to C^*(\Gamma)$  are isomorphisms of these  $C^*$ -algebras which are inverse to each other.

## Step 4: Showing that $\Delta_G \circ \varphi_1 = (\varphi_1 \otimes \varphi_1) \circ \Delta_{\Gamma}$ .

Since  $\varphi_1$  and the coproducts are \*-homomorphisms, it suffices to prove the identity on the generators, i.e., that  $\Delta_G(\varphi_1(x_i)) = (\varphi_1 \otimes \varphi_1)(\Delta_{\Gamma}(x_i))$  for all  $i \in [n]$ . From the above, we have that  $\varphi_1(x_i) = y_i = \sum_{\delta \in \pm 1_0^{S_k}} \delta_i u_{\delta}^{(k)}$  for any  $k \in [m]$  such that  $i \in S_k$ . Thus  $(\varphi_1 \otimes \varphi_1)(\Delta_{\Gamma}(x_i)) = (\varphi_1 \otimes \varphi_1)(x_i \otimes x_i) = y_i \otimes y_i$ . We aim to show that  $\Delta_G(\varphi_1(x_i)) = \Delta_G(y_i)$  is equal to  $y_i \otimes y_i$ . First, let us determine  $\Delta_G(u_{\delta}^{(k)})$  for  $\delta \in \pm 1_0^{S_k}$ . Picking  $\alpha \in \pm 1_{b_k}^{S_k}$  arbitrarily, we have that

First, let us determine  $\Delta_G(u_{\delta}^{(\kappa)})$  for  $\delta \in \pm 1_0^{S_k}$ . Picking  $\alpha \in \pm 1_{b_k}^{S_k}$  arbitrarily, we have that  $u_{\delta}^{(k)} = u_{\alpha,\delta\Delta\alpha}^{(k)}$  and thus

$$\Delta_G \left( u_{\delta}^{(k)} \right) = \Delta_G \left( u_{\alpha,\delta \Delta \alpha}^{(k)} \right)$$
  
=  $\Delta_G \left( u_{(k,\alpha),(k,\delta \Delta \alpha)} \right)$   
=  $\sum_{\substack{l \in [m] \\ \beta \in \pm 1_{b_l}^{S_l}}} u_{(k,\alpha),(l,\beta)} \otimes u_{(l,\beta),(k,\delta \Delta \alpha)}$   
=  $\sum_{\substack{\beta \in \pm 1_{b_k}^{S_k}}} u_{(k,\alpha),(k,\beta)} \otimes u_{(k,\beta),(k,\delta \Delta \alpha)}$   
=  $\sum_{\delta' \in \pm 1_{0}^{S_k}} u_{\delta'}^{(k)} \otimes u_{\delta \Delta \delta'}^{(k)}.$ 

Now we will show that  $\Delta_G(y_i) = y_i \otimes y_i$ . We have that

$$\begin{split} \Delta_{G}(y_{i}) &= \Delta_{G} \left( \sum_{\delta \in \pm 1_{0}^{S_{k}}} \delta_{i} u_{\delta}^{(k)} \right) \\ &= \sum_{\delta \in \pm 1_{0}^{S_{k}}} \delta_{i} \Delta_{G} \left( u_{\delta}^{(k)} \right) \\ &= \sum_{\delta, \delta' \in \pm 1_{0}^{S_{k}}} \delta_{i} \left( u_{\delta'}^{(k)} \otimes u_{\delta \Delta \delta'}^{(k)} \right) \\ &= \sum_{\delta, \delta' \in \pm 1_{0}^{S_{k}}} \delta_{i} \delta_{i}^{\prime 2} \left( u_{\delta'}^{(k)} \otimes u_{\delta \Delta \delta'}^{(k)} \right) \\ &= \sum_{\delta, \delta' \in \pm 1_{0}^{S_{k}}} \left( \delta_{i}^{\prime} u_{\delta'}^{(k)} \right) \otimes \left( \delta_{i} \delta_{i}^{\prime} u_{\delta \Delta \delta'}^{(k)} \right) \\ &= \sum_{\delta' \in \pm 1_{0}^{S_{k}}} \left( \delta_{i}^{\prime} u_{\delta'}^{(k)} \otimes \left( \sum_{\delta \in \pm 1_{0}^{S_{k}}} \delta_{i}^{\prime\prime} u_{\delta \Delta \delta'}^{(k)} \right) \right) \\ &= \sum_{\delta' \in \pm 1_{0}^{S_{k}}} \delta_{i}^{\prime} u_{\delta'}^{(k)} \otimes y_{i} \end{split}$$

$$= y_i \otimes y_i.$$

Since we have already shown that  $(\varphi_1 \otimes \varphi_1) \circ \Delta_{\Gamma}(x_i) = y_i \otimes y_i$ , this completes the proof.  $\Box$ 

## 4 Constructing uncolored graphs that have the same quantum automorphism group as a given colored graph

In the following, we will describe a procedure that given certain vertex– and edge-colored graphs produces a decolored graph with isomorphic quantum automorphism group. This procedure is divided into two steps: We first decolor the vertices and then the edges. We start with a lemma known from [8, Lemma 3.2.3]. The degree deg(v) of a vertex v denotes the number of neighbors of v in the graph G.

**Lemma 4.1.** Let G be a finite graph,  $A_G$  be its adjacency matrix and  $u_{vw}$ ,  $1 \le v, w \le n$  be the generators of C(Qut(G)). If  $(A_G^l)_{vv} \ne (A_G^l)_{ww}$  for some  $l \in \mathbb{N}$ , then  $u_{vw} = 0$ . Particularly, if  $\deg(v) \ne \deg(w)$ , then  $u_{vw} = 0$ .

The following lemma can be found in [14, Lemma 3.2]. The distance d(v, w) between two vertices  $v, w \in V(G)$  is the length of a shortest path connecting v and w.

**Lemma 4.2.** Let G be a finite graph and  $u_{vw}$ ,  $1 \le v, w \le n$  be the generators of C(Qut(G)). If we have  $d(v, p) \ne d(w, q)$ , then  $u_{vw}u_{pq} = 0$ .

**Lemma 4.3.** Let G be a finite graph and u the fundamental representation of Qut(G). Let  $v, w \in V(G)$ . If there exists  $p \in V(G)$  with d(v, p) = k such that  $deg(p) \neq deg(q)$  for all  $q \in V(G)$  with d(w,q) = k, then  $u_{vw} = 0$ .

*Proof.* Let  $v, w \in V(G)$  and  $p \in V(G)$ , d(v, p) = k as above. Using Relation (2) and Lemma 4.2, we get

$$u_{vw} = u_{vw} \left( \sum_{q} u_{pq} \right) = u_{vw} \left( \sum_{q;d(w,q)=k} u_{pq} \right).$$

We conclude

$$u_{vw} = u_{vw} \left( \sum_{q;d(w,q)=k} u_{pq} \right) = 0,$$

since we have  $deg(p) \neq deg(q)$  and thus  $u_{pq} = 0$  by Lemma 4.1.

**Remark 4.4.** Lemma 4.1, Lemma 4.2 and Lemma 4.3 also work for vertex– and edge-colored graphs (take the decolored adjacency matrix in Lemma 4.1), since colors just add more relations on the generators of C(Qut(G)).

We will now define a edge-colored graph G' from the vertex– and edge-colored graph G.

**Definition 4.5.** Let *G* be a vertex – and edge-colored graph. Depending on the color *c*, attach a path of length  $n_c \in \mathbb{N}_0$  to every vertex colored *c*, where  $n_{c_1} \neq n_{c_2}$  for colors  $c_1 \neq c_2$  and then decolor the vertices of the graph. We choose one of the edge-colors of *G* and let the edges in the paths all have this edge-color. We denote this new edge-colored (but not vertex-colored) graph by *G*'.

The next lemma gives a description of the fundamental representation of Qut(G'). We will see in the following theorem that the quantum automorphism groups of G and G' are isomorphic.




Figure 2: Example for the construction of the graph G' from G. We chose the edge-colors between the added paths to be black.

**Lemma 4.6.** Let G be a vertex – and edge-colored graph with  $\deg(v) \ge 3$  for all  $v \in V(G)$  and let G' be the graph as in Definition 4.5. Denote by  $v_i$ ,  $0 \le i \le n_{c(v)}$ , the vertices in the added path with  $d(v, v_i) = i$  (thus  $v = v_0$ ) and let  $V_i = \{x \in V(G'); x = v_i \text{ for some } v \in V(G)\}$ . Then the fundamental representation u of Qut(G') is of the form

where additionally  $u_{v_iw_i} = u_{v_0w_0}$  for all *i* and  $u_{v_iw_i} = 0$  for  $c(v) \neq c(w)$ , c(.) being the vertex-colors in the original graph *G*.

*Proof.* Step 1: It holds  $u_{v_iw_0} = u_{v_0w_i} = 0$  for  $i \neq 0$ . We know that it holds  $\deg(v_i) \in \{1, 2\}$  for vertices  $v_i$  with i > 0. Since we have  $\deg(w_0) \ge 3$  by assumption, we get  $u_{v_iw_0} = 0$  by Lemma 4.1. We similarly obtain  $u_{v_0w_i} = 0$ .

Step 2: We have  $u_{v_iw_j} = 0$  for  $i \neq j, i, j > 0$ .

First assume i < j. Then  $v_0$  is a vertex with  $d(v_i, v_0) = i$  and  $\deg(v_0) \ge 3$ . Since i < j, we know that the vertices q with  $d(w_j, q) = i$  are in the path added to w and thus  $\deg(q) \in \{1, 2\}$ . We deduce  $u_{v_iw_j} = 0$  by Lemma 4.3. The case i > j follows similarly.

Step 3: It holds  $u_{v_iw_i} = u_{v_0w_0}$  for all *i*. We first show  $u_{v_0w_0} = u_{v_1w_1}$ . It holds

$$u_{v_0w_0} = u_{v_0w_0} \left( \sum_{\substack{k; (k,w_0) \in E}} u_{v_1k} \right)$$
$$= u_{v_0w_0} \left( u_{v_1w_1} + \sum_{\substack{p_0; (p_0,w_0) \in E, \\ p_0 \in V_0}} u_{v_1p_0} \right)$$
$$= u_{v_0w_0} u_{v_1w_1}$$
$$= (u_{v_0w_0} + u_{v_0w_2}) u_{v_1w_1}$$

$$= \left(\sum_{\substack{k;(k,w_1)\in E}} u_{v_0k}\right) u_{v_1w_1}$$
$$= u_{v_1w_1},$$

since  $u_{v_1p_0} = 0$  and  $u_{v_0w_2} = 0$  by *Step 1*. Furthermore, for  $i \ge 1$ , we have

$$u_{v_{i}w_{i}} = u_{v_{i}w_{i}} \left( \sum_{k;(k,w_{i})\in E} u_{v_{i+1}k} \right)$$
  
=  $u_{v_{i}w_{i}} (u_{v_{i+1}w_{i+1}} + u_{v_{i+1}w_{i-1}})$   
=  $u_{v_{i}w_{i}} u_{v_{i+1}w_{i+1}}$   
=  $(u_{v_{i}w_{i}} + u_{v_{i}w_{i+2}})u_{v_{i+1}w_{i+1}}$   
=  $\left( \sum_{k;(k,w_{i+1})\in E} u_{v_{i}k} \right) u_{v_{i+1}w_{i+1}}$   
=  $u_{v_{i+1}w_{i+1}}$ 

since it holds that  $u_{v_{i+1}w_{i-1}} = 0$  and  $u_{v_iw_{i+2}} = 0$  by *Step 1* or *Step 2*. Note that if there is no  $w_{i+2}$ , we still get  $u_{v_iw_i} = u_{v_{i+1}w_{i+1}}$  by a similar calculation, since then  $w_i$  is the only neighbor of  $w_{i+1}$ .

Step 4: It holds  $u_{v_iw_i} = 0$  for  $c(v) \neq c(w)$ .

Since  $c(v) \neq c(w)$ , we know  $n_{c(v)} \neq n_{c(w)}$ . Assume  $n := n_{c(v)} < n_{c(w)}$ . Then we have  $u_{v_n w_n} = 0$  by Lemma 4.1, since  $\deg(v_n) = 1 \neq 2 = \deg(w_n)$ . We deduce  $u_{v_i w_i} = u_{v_0 w_0} = u_{v_n w_n} = 0$  for all *i* by *Step 3*. The case  $n_{c(v)} > n_{c(w)}$  follows similarly.

**Theorem 4.7.** Let G be a vertex – and edge-colored graph with  $\deg(v) \ge 3$  for all  $v \in V(G)$  and let G' be as in Definition 4.5. Then there exists a \*-isomorphism  $\varphi : C(\operatorname{Qut}(G)) \to C(\operatorname{Qut}(G'))$  such that  $\Delta_{G'} \circ \varphi = (\varphi \otimes \varphi) \circ \Delta_G$ .

*Proof.* Let u be the fundamental representation of Qut(G'). As in Lemma 4.6, we denote by  $v_i$ ,  $0 \le i \le n_{c(v)}$  the vertices in the added path with  $d(v, v_i) = i$  and let  $V_i = \{x \in V(G'); x = v_i \text{ for some } v \in V(G)\}$ . Let  $c_e$  be the color of the edges in the attached paths in G'. For colors  $c \ne c_e$ , we get that  $A_{G'_c}$  is equal to

By Lemma 4.6, we directly see that  $uA_{G'_c} = A_{G'_c}u$  implies  $u_0A_{G_c} = A_{G_c}u_0$  for  $c \neq c_e$ ,  $u_0$  being the  $(V_0 \times V_0)$ -block in u (see (11)). The adjacency matrix  $A_{G'_{c_e}}$  is of the form

where  $B_i$  is the  $(V_{i-1} \times V_i)$ -matrix with  $(B_i)_{a_{i-1}b_i} = \delta_{ab}$ . Also by Lemma 4.6, we see that  $uA_{G'_{c_e}} = A_{G'_{c_e}}u$  implies  $u_0A_{G_{c_e}} = A_{G_{c_e}}u_0$ . Furthermore, since  $(u_i)_{v_iw_i} = (u_0)_{v_0w_0}$  for all i, we see that  $C(\operatorname{Qut}(G'))$  is generated by the entries of  $u_0$ . We conclude that  $C(\operatorname{Qut}(G'))$  is generated by a magic unitary  $u_0$  that fulfills  $u_0A_{G_c} = A_{G_c}u_0$  for all edge-colors c and  $(u_0)_{v_0w_0} = 0$  for  $c(v) \neq c(w)$  (see Lemma 4.6). This yields a surjective \*-homomorphism  $\varphi : C(\operatorname{Qut}(G)) \to C(\operatorname{Qut}(G'))$ ,  $w_{ab} \mapsto (u_0)_{ab}$ , where w is the fundamental representation of  $\operatorname{Qut}(G)$ .

For the other direction, take the fundamental representation w of Qut(G) and build the matrix w' as follows

where we put  $(w_i)_{a_ib_i} = w_{a_0b_0}$ . Note that w' is a magic unitary if and only if the matrices  $w_i$  are magic unitaries. We compute

$$\sum_{b;n_c(b)\geq i} (w_i)_{a_ib_i} = \sum_{b;n_c(b)\geq i} w_{ab} = \sum_b w_{ab} = 1,$$

where we used  $w_{ab} = 0$  for  $c(a) \neq c(b)$  and the fact that w is a magic unitary. Similarly, we get  $\sum_{a;n_c(a)\geq i}(w_i)_{a_ib_i} = 1$  and thus w' is a magic unitary. It remains to show that w' commutes with  $A_{G'_c}$  for all edge colors c. We deduce from (12) and (14) that  $wA_{G_c} = A_{G_c}w$  implies  $w'A_{G'_c} = A_{G'_c}w'$  for  $c \neq c_e$ . From (13) and (14), we see that  $w'A_{G'_{ce}} = A_{G'_{ce}}w'$  is equivalent to  $wA_{G_{ce}} = A_{G_{ce}}w, w_iB_{i+1} = B_{i+1}w_{i+1}, w_{i+1}B_{i+1}^t = B_{i+1}^tw_i$ . Note that we have

$$(w_i B_{i+1})_{a_i b_{i+1}} = \sum_k (w_i)_{a_i k} (B_{i+1})_{k b_{i+1}}$$
  
=  $(w_i)_{a_i b_i}$   
=  $(w)_{a_0 b_0}$   
=  $(w_{i+1})_{a_{i+1} b_{i+1}}$   
=  $\sum_k (B_{i+1})_{a_i k} (w_{i+1})_{k b_{i+1}}$   
=  $(B_{i+1} w_{i+1})_{a_i b_{i+1}}$ 

by definition of  $w_i$  and  $B_i$ . We similarly get that  $w_{i+1}B_{i+1}^t = B_{i+1}^t w_i$  is automatically fulfilled by definition of  $w_i$  and  $B_i$ . Since we have  $wA_{G_{c_e}} = A_{G_{c_e}} w$  by assumption, we obtain  $w'A_{G'_{c_e}} = A_{G'_{c_e}}w'$ . Summarizing, we have  $w'A_{G'_c} = A_{G'_c}w'$  for all edge-colors c. This yields a surjective \*-homomorphism  $\tilde{\varphi} : C(\operatorname{Qut}(G')) \to C(\operatorname{Qut}(G)), u_{xy} \mapsto w'_{xy}$  which is inverse to  $\varphi$ . Thus,  $\varphi$  is a \*-isomorphism.

It remains to show that  $\Delta_{G'} \circ \varphi = (\varphi \otimes \varphi) \circ \Delta_G$ . But this is true because

$$\Delta_{G'} \circ \varphi(w_{ij}) = \Delta_{G'}((u_0)_{ij})$$
$$= \sum_k (u_0)_{ik} \otimes (u_0)_{kj}$$
$$= (\varphi \otimes \varphi) \left(\sum_k w_{ik} \otimes w_{kj}\right)$$

$$= (\varphi \otimes \varphi) \circ \Delta(w_{ij}).$$

We will now deal with the edge-colors of G'. First, we need the following easy lemma.

**Lemma 4.8.** Let  $w_{ij}$ ,  $1 \le i, j \le n$  be elements in a  $C^*$ -algebra such that the matrix  $w = (w_{ij})_{1 \le i, j \le n}$  is a magic unitary. Then  $w_{ij}w_{kl} + w_{il}w_{kj}$  is a projection if and only if  $w_{ij}w_{kl} = w_{kl}w_{ij}$  and  $w_{il}w_{kj} = w_{kj}w_{il}$ .

*Proof.* Let  $w_{ij}w_{kl} + w_{il}w_{kj}$  be a projection. Since it is self-adjoint, we have

$$w_{ij}w_{kl} + w_{il}w_{kj} = w_{kl}w_{ij} + w_{kj}w_{il}$$

Multiplying by  $w_{ij}$  and  $w_{il}$ , respectively, yields  $w_{ij}w_{kl} = w_{ij}w_{kl}w_{ij}$  and  $w_{il}w_{kj} = w_{il}w_{kj}w_{il}$ . But, by taking adjoints, this implies  $w_{ij}w_{kl} = w_{kl}w_{ij}$  and  $w_{il}w_{kj} = w_{kj}w_{il}$ . The other direction is clear, since  $w_{ij}w_{kl} + w_{il}w_{kj}$  is the sum of two orthogonal projections if  $w_{ij}w_{kl} = w_{kl}w_{ij}$  and  $w_{il}w_{kj} = w_{kj}w_{il}$ .

We will define a (uncolored) graph G'' from the edge-colored graph G'. In the next theorem, we will then see that for certain graphs, the quantum automorphism groups of G' and G'' are isomorphic. Let G be a graph and  $e = (u, v) \in E(G)$ . We say that we subdivide e if we delete the edge e = (u, v) from G and add a vertex w as well as edges (u, w) and (w, v) to the graph.

**Definition 4.9.** Let *G* be a vertex – and edge-colored graph. First decolor the vertices by applying the construction as in Definition 4.5 to obtain the edge-colored graph *G'*. We denote the color of the newly added edges in the graph *G'* by  $c_0$ . We subdivide each colored edge with  $c(e) \neq c_0$  and add a path of length  $m_c$  to the subdivision, where  $m_{c_1} \neq m_{c_2}$  for colors  $c_1 \neq c_2$  and then decolor the edges in the graph *G'*. We call this graph *G''*.




Figure 3: Example for the construction of the graph G'' from G'. We chose black to be the edge-color  $c_0$ .

**Lemma 4.10.** Let G be a vertex – and edge-colored graph such that  $\deg(v) \ge 3$  for all  $v \in V(G)$ and let G" be the graph as in Definition 4.9. We denote the vertex that subdivided the edge e in G by  $e_0$  and the vertices in the added path with  $d(e_0, e_i) = i$  by  $e_i$ ,  $1 \le i \le m_{c(e)}$ . Furthermore  $E'_i = \{x \in V(G''); x = e_i \text{ for some } e \in E(G)\}$ . Then, the fundamental representation u of Qut(G'') is of the form

where  $u_{e_if_i} = u_{vx}u_{wy} + u_{vy}u_{wx}$  for c(e) = c(f), e = (v, w), f = (x, y) and  $u_{e_if_i} = 0$  for  $c(e) \neq c(f)$ .

*Proof.* Step 1: It holds  $u_{e_iv_j} = u_{v_je_i} = 0$  for  $v_j \in V(G')$ ,  $e_i \in E'_i$ . First assume i > 0, j = 0. Then  $\deg(e_i) \in \{1, 2\}$  and  $\deg(v_0) \ge 3$ , therefore  $u_{e_iv_0} = 0$  by Lemma 4.1.

Now, let i = 0, j > 0. If  $m_{c(e_0)} > 0$ , then  $\deg(e_0) = 3 \neq \deg(v_j)$ , since  $\deg(v_j) \in \{1, 2\}$ . Therefore  $u_{e_0v_j} = 0$  by Lemma 4.1. Let  $m_{c(e_0)} = 0$ . If  $n_{c(v_0)} = 1$ , then  $\deg(v_1) = 1 \neq 2 = \deg(e_0)$ and thus  $u_{e_0v_1} = 0$  by Lemma 4.1. If  $n_{c(v_0)} \ge 2$ , then  $v_j$  has at least one neighbor of degree 1 or 2. Since  $e_0$  only has neighbors of degree  $\ge 3$ , we get  $u_{e_0v_j} = 0$  by Lemma 4.3.

Assume  $i \neq j$  and  $i, j \neq 0$ . Let furthermore i < j. Then  $e_0$  is a vertex with  $d(e_i, e_0) = i$  and  $deg(e_0) = 3$ . Since i < j, we know that the vertices q with  $d(v_j, q) = i$  are in the path and thus  $deg(q) \in \{1, 2\}$ . We deduce  $u_{e_iv_j} = 0$  by Lemma 4.3. The case i > j follows similarly.

It remains to show  $u_{e_iv_i} = 0$ . If  $n_{c(v_0)} > 0$ , then  $\deg(v_0) \ge 4$  and  $d(v_0, v_i) = i$ . We know  $\deg(q) \le 3$  for vertices q with  $d(q, e_i) = i$ , since either  $q = e_0$  or q is a vertex in the added path. We deduce  $u_{e_iv_i} = 0$  by Lemma 4.3. Now assume  $n_{c(v_0)} = 0$ . If  $m_{c(e_0)} = 0$ , then we know  $\deg(e_0) = 2$  and thus  $u_{e_0v_0} = 0$  by Lemma 4.1. If  $m_{c(e_0)} > 0$ , then  $e_0$  has a neighbor of degree 1 or 2. If  $v_0$  has no neighbor of this degree, then  $u_{e_0v_0} = 0$  by Lemma 4.3. The vertex  $v_0$  only has a neighbor of degree 2 if there exists a subdivision  $f_0$  of some edge  $f = (v_0, w_0)$  with  $m_{c(f)} = 0$ . Then, it holds

$$u_{e_0v_0} = u_{e_0v_0} \left( \sum_{k; (k,e_0) \in E(G'')} u_{kf_0} \right) = u_{e_0v_0} u_{e_1f_0}.$$

If  $m_{c(e_0)} = 1$ , then  $\deg(e_1) = 1 \neq 2 = \deg(f_0)$  which yields  $u_{e_1f_0} = 0$  by Lemma 4.1. If  $m_{c(e_0)} \geq 2$ , then  $e_1$  has a neighbor  $e_2$  with  $\deg(e_2) \in \{1, 2\}$ . Since the neighbors  $v_0$ ,  $w_0$  of  $f_0$  have degree  $\geq 3$ , we get  $u_{e_1f_0} = 0$  by Lemma 4.3. We deduce  $u_{e_0v_0} = 0$  in all those cases.

Step 2: It holds  $u_{e_i f_0} = u_{e_0 f_i} = 0$  for  $i \neq 0$ .

We know that it holds  $\deg(e_i) \in \{1, 2\}$  for vertices  $e_i$  with i > 0. If  $m_{c(f_0)} > 0$ , then we have  $\deg(f_0) = 3$  and get  $u_{e_i f_0} = 0$  by Lemma 4.1. Let  $m_{c(f_0)} = 0$ . If  $m_{c(e_0)} = 1$ , then  $\deg(e_1) = 1$  and thus  $u_{e_1 f_0} = 0$  by Lemma 4.1. If  $m_{c(e_0)} \ge 2$ , then  $e_i$  has a neighbor with degree 1 or 2. Since the neighbors of  $f_0$  have degree  $\ge 3$  by assumption, we get  $u_{e_i f_0} = 0$  by Lemma 4.3. We similarly obtain  $u_{e_0 f_i} = 0$ .

Step 3: It holds  $u_{e_if_j} = 0$  for  $i \neq j, i, j > 0$ .

First assume i < j. Then  $e_0$  is a vertex with  $d(e_i, e_0) = i$  and  $deg(e_0) = 3$ . Since i < j, we know that the vertices q with  $d(f_j, q) = i$  are in the path and thus  $deg(q) \in \{1, 2\}$ . We deduce  $u_{e_if_i} = 0$  by Lemma 4.3. The case i > j follows similarly.

Step 4: It holds  $u_{e_if_i} = u_{e_0f_0}$  for all *i*. We first show  $u_{e_0f_0} = u_{e_1f_1}$ . It holds

$$\begin{aligned} u_{e_0f_0} &= u_{e_0f_0} \left( \sum_{k;(k,f_0) \in E(G'')} u_{e_1k} \right) \\ &= u_{e_0f_0} \left( u_{e_1f_1} + \sum_{p_0;(p_0,f_0) \in E(G''), p_0 \in V(G')} u_{e_1p_0} \right) \\ &= u_{e_0f_0} u_{e_1f_1} \\ &= (u_{e_0f_0} + u_{e_0f_2}) u_{e_1f_1} \\ &= \left( \sum_{k;(k,f_1) \in E(G'')} u_{e_0k} \right) u_{e_1f_1} \\ &= u_{e_1f_1}, \end{aligned}$$

since  $u_{e_1p_0} = 0$  by *Step 1* and  $u_{e_0f_2} = 0$  by *Step 2*. Furthermore, for  $i \ge 1$ , we have

$$\begin{split} u_{e_if_i} &= u_{e_if_i} \left( \sum_{\substack{k; (k,f_i) \in E(G'') \\ k; (k,f_i) \in E(G'')}} u_{e_{i+1}k} \right) \\ &= u_{e_if_i} (u_{e_{i+1}f_{i+1}} + u_{e_{i+1}f_{i-1}}) \\ &= u_{e_if_i} u_{e_{i+1}f_{i+1}} \\ &= (u_{e_if_i} + u_{e_if_{i+2}}) u_{e_{i+1}f_{i+1}} \\ &= \left( \sum_{\substack{k; (k,f_{i+1}) \in E(G'') \\ k; (k,f_{i+1}) \in E(G'')}} u_{e_ik} \right) u_{e_{i+1}f_{i+1}} \\ &= u_{e_{i+1}f_{i+1}} \end{split}$$

since we have  $u_{e_{i+1}f_{i-1}} = 0$  and  $u_{e_if_{i+2}} = 0$  by *Step 2* or *Step 3*. Note that if there is no  $f_{i+2}$ , we still get  $u_{e_if_i} = u_{e_{i+1}f_{i+1}}$  by a similar calculation, since then  $f_i$  is the only neighbor of  $f_{i+1}$ .

Step 5: It holds  $u_{e_if_i} = 0$  for  $c(e) \neq c(f)$ .

Since  $c(e) \neq c(f)$ , we know  $m_{c(e)} \neq m_{c(f)}$ . First assume  $m := m_{c(e)} < m_{c(f)}$ . Then we have  $u_{e_m f_m} = 0$  by Lemma 4.1, since  $\deg(e_m) = 1 \neq 2 = \deg(f_m)$ . We deduce  $u_{e_i f_i} = u_{e_0 f_0} = u_{e_m f_m} = 0$  for all *i* by *Step 3*. The case  $m_{c(e)} > m_{c(f)}$  is similar.

Step 6: It holds  $u_{e_if_i} = u_{vx}u_{wy} + u_{vy}u_{wx}$  for e = (v, w), f = (x, y). We have

$$u_{e_0f_0} = \left(\sum_{k;(k,f_0)\in E(G'')} u_{vk}\right) u_{e_0f_0} \left(\sum_{l;(l,f_0)\in E(G'')} u_{wl}\right)$$
$$= (u_{vx} + u_{vy} + u_{vf_1}) u_{e_0f_0} (u_{wx} + u_{wy} + u_{wf_1})$$
$$= (u_{vx} + u_{vy}) u_{e_0f_0} (u_{wx} + u_{wy}),$$

where we used  $u_{vf_1} = 0$ ,  $u_{wf_1} = 0$  by *Step 1*. Note that  $e_0$  is the only vertex in  $E'_0$  that is a common neighbor of v and w. Thus, we have  $(u_{vx} + u_{vy})u_{kf_0}(u_{wx} + u_{wy}) = 0$  for all  $k \neq e_0$  by Lemma 4.2 and *Step 1*. Therefore, we get

$$u_{e_0f_0} = (u_{vx} + u_{vy})u_{e_0f_0}(u_{wx} + u_{wy})$$

$$= (u_{vx} + u_{vy}) \left( \sum_{k;(k,v) \in E(G''),(k,w) \in E(G'')} u_{kf_0} \right) (u_{wx} + u_{wy})$$
  
=  $(u_{vx} + u_{vy})(u_{wx} + u_{wy})$   
=  $u_{vx}u_{wy} + u_{vy}u_{wx}.$ 

By Step 4, we obtain  $u_{e_if_i} = u_{vx}u_{wy} + u_{vy}u_{wx}$ .

**Remark 4.11.** Note that the operator  $u_{(v,w)(x,y)} = u_{vx}u_{wy} + u_{vy}u_{wx}$  does not depend on the order of (v, w) or (x, y), since

$$\begin{split} u_{(v,w)(x,y)} &= u_{vx}u_{wy} + u_{vy}u_{wx} = u_{vy}u_{wx} + u_{vx}u_{wy} = u_{(v,w),(y,x)}, \\ u_{(v,w)(y,x)} &= u_{vy}u_{wx} + u_{vx}u_{wy} = u_{wx}u_{vy} + u_{wy}u_{vx} = u_{(w,v),(x,y)}. \end{split}$$

Before stating the theorem, we first need to define a quantum subgroup of the quantum automorphism group of the graph G'. It is straightforward to check that the comultiplication is a \*-homomorphism.

**Definition 4.12.** Let *G* be a vertex – and edge-colored graph and *G'* as in Definiton 4.5. We define  $\operatorname{Qut}_{c_0}^*(G')$  to be the compact matrix quantum group whose corresponding *C*\*-algebra is generated by a magic unitary *x* with  $xA_{G_c} = A_{G_c}x$  for every edge color *c* and  $x_{ik}x_{jl} = x_{jl}x_{ik}$  for  $c((i, j)) = c((k, l)), c \neq c_0$ , where  $c_0$  is the edge-color we choose for the newly added edges in *G'*.

**Theorem 4.13.** Let G be a vertex – and edge-colored graph such that  $\deg(v) \ge 3$  for all  $v \in V(G)$ . Let G" be the graph as in Definition 4.9 and let  $\operatorname{Qut}_{c_0}^*(G')$  be the compact matrix quantum group as in Definition 4.12. Then there exists a \*-isomorphism  $\varphi : C(\operatorname{Qut}_{c_0}^*(G')) \to C(\operatorname{Qut}(G''))$  such that  $\Delta_{G''} \circ \varphi = (\varphi \otimes \varphi) \circ \Delta_{G'}$ .

*Proof.* Let u be the fundamental representation of Qut(G''). As in Lemma 4.10, we denote the vertex that subdivided the edge e in G by  $e_0$  and the vertices in the added path with  $d(e_0, e_i) = i$  by  $e_i$ ,  $1 \le i \le m_{c(e)}$ . Moreover, we denote by  $v_i$ ,  $0 \le i \le n_{c(v)}$  the vertices in the added path with  $d(v, v_i) = i$ . The adjacency matrix of G'' is of the form

where  $A_{G_0}$  is the adjacency matrix of the edge-color that is not subdivided together with the paths from the construction of G',  $B_{G''}$  is the matrix with

$$(B_{G''})_{ve} = \begin{cases} 1 & \text{for } v \text{ incident to } e, \\ 0 & \text{otherwise,} \end{cases}$$

and  $C_i$  is the  $(E'_{i-1} \times E'_i)$ -matrix with  $(C_i)_{a_{i-1}b_i} = \delta_{ab}$ . Using Lemma 4.10, we see that  $uA_{G''} = A_{G''}u$  implies  $wA_{G'_0} = A_{G'_0}w$ ,  $wB_{G''} = B_{G''}u_0$ ,  $u_0B^t_{G''} = B^t_{G''}w$ , where w and  $u_0$  are blocks in u (see (15)). Furthermore, since  $(u_i)_{e_if_i} = w_{ak}w_{bl} + w_{al}w_{bk}$  for e = (a, b), f = (k, l), we also have that  $C(\operatorname{Qut}(G''))$  is generated by the entries of the matrix w. We will now show that w fulfills

the relations of the generators of  $C(\operatorname{Qut}_{c_0}^*(G'))$ . We already know  $wA_{G'_0} = A_{G'_0}w$ . We will now show  $wA_{G_c} = A_{G_c}w$  for every edge color  $c \neq c_0$ . By Lemma 2.8, this is equivalent to  $w_{ik}w_{jl} = 0$ for  $c((i, j)) \neq c((k, l))$  if  $(i, j) \in E'_0$  or  $(k, l) \in E'_0$ . We will show that the elements of w fulfill those relations. Let  $v \in V(G')$ ,  $(a, b) = e \in E'_0$ . It holds

$$w_{va} + w_{vb} = \sum_{k;k \text{ inc. to } e} w_{vk}$$
  

$$= \sum_{k} w_{vk} (B_{G''})_{ke}$$
  

$$= (wB_{G''})_{ve}$$
  

$$= (B_{G''}u_0)_{ve}$$
  

$$= \sum_{k} (B_{G''})_{vk} (u_0)_{ke}$$
  

$$= \sum_{k;k \text{ inc. to } v} (u_0)_{ke}$$
  

$$= \sum_{s;(v,s)\in E_0} (w_{va}w_{sb} + w_{vb}w_{sa}).$$
(17)

Multiplying  $w_{va}$  from left and right yields

$$w_{va} = \sum_{s;(v,s)\in E_0} w_{va} w_{sb} w_{va}$$

We deduce  $\sum_{s;(v,s)\notin E'_0} w_{va}w_{sb}w_{va} = 0$  and since all elements in the sum are positive, we get  $w_{va}w_{sb}w_{va} = 0$  for all  $v, s \in V(G')$  with  $(v, s) \notin E'_0$ . We deduce  $w_{va}w_{sb} = 0$  for all  $v, s \in V$  with  $(v, s) \notin E'_0$ . Similarly, by using  $u_0B^t_{G''} = B^t_{G''}w$ , we get  $w_{av}w_{bs} = 0$  for all  $v, s \in V(G')$  with  $(v, s) \notin E'_0$ . It remains to show  $w_{ik}w_{jl} = 0$  for  $c(e) \neq c(f)$ ,  $e = (i, j) \in E'_0$ ,  $f = (k, l) \in E'_0$ . We know  $(u_0)_{ef} = 0$  for  $c(e) \neq c(f)$  from Lemma 4.10. Then

$$w_{ik}w_{jl} + w_{il}w_{jk} = (u_0)_{ef} = 0$$

and by multiplying  $w_{ik}$  and  $w_{il}$  from the left, respectively, we get  $w_{ik}w_{jl} = 0$  and  $w_{il}w_{jk} = 0$ . Thus, we have shown  $w_{ik}w_{jl} = 0$  for  $c((i, j)) \neq c((k, l))$  if  $(i, j) \in E'_0$  or  $(k, l) \in E'_0$  which is equivalent to  $wA_{G_c} = A_{G_c}w$  for every edge color  $c \neq c_0$ .

Summarizing, we get  $wA_{G_c} = A_{G_c}w$  for every edge color c. By Lemma 4.10, we furthermore know that  $(u_0)_{ab} = w_{ik}w_{jl} + w_{il}w_{jk}$  for a = (i, j), b = (k, l) and c(a) = c(b),  $c(a) \neq c_0$ and thus  $w_{ik}w_{jl} + w_{il}w_{jk}$  is a projection. Then Lemma 4.8 yields  $w_{ij}w_{kl} = w_{kl}w_{ij}$  and  $w_{il}w_{kj} = w_{kj}w_{il}$ . Therefore, we get a surjective \*-homomorphism  $\varphi : C(\operatorname{Qut}^*_{c_0}(G')) \to C(\operatorname{Qut}(G''), w'_{ij} \mapsto w_{ij})$ , where w' is the fundamental representation of  $\operatorname{Qut}^*_{c_0}(G')$ .

For the other direction, take the fundamental representation w' of  $\operatorname{Qut}_{c_0}^*(G')$  and build the matrix w'' as follows

|        | V(G') | $E'_0$ |        |   | $E'_m$   |
|--------|-------|--------|--------|---|----------|
| V(G')  | (w')  | 0      | 0      |   | 0 \      |
| $E'_0$ | 0     | $u'_0$ | 0      |   | 0        |
| ÷      | 0     | 0      | $u_1'$ |   | 0        |
| ÷      |       | ÷      | ÷      | · | 0        |
| $E'_m$ | 0     | 0      | 0      | 0 | $u'_m$ / |

where we put  $(u'_i)_{e_i f_i} = w'_{ak}w'_{bl} + w'_{al}w'_{bk}$  for e = (a, b), f = (k, l). Those elements are projections by Lemma 4.8, because we have  $w'_{ak}w'_{bl} = w'_{bl}w'_{ak}$  for  $(a, b), (k, l) \in E'_0$  (for the same color by assumption, for different colors because the product is 0). Note that w'' is a magic unitary if and only if the matrices  $u'_i$  are magic unitaries. Let  $e = (a, b) \in E'_0$ . We compute

$$\sum_{f;n_c(f)\geq i} (u'_i)_{e_if_i} = \sum_{k,l;n_c((k,l))\geq i} (w'_{ak}w'_{bl} + w'_{al}w'_{bk}) = \sum_{k,l;(k,l)\in E'_0} (w'_{ak}w'_{bl} + w'_{al}w'_{bk}) = 1,$$

where we used  $w'_{ak}w'_{bl} = 0$  for  $c((a, b)) \neq c((k, l))$  and the fact that w' is a magic unitary. We get  $\sum_{f;n_c(f)\geq i} (u'_i)_{f_ie_i} = 1$  similarly and therefore w'' is a magic unitary. It remains to show that w'' commutes with  $A_{G''}$ . Similar to equation (17), we see that

$$(w'B_{G''})_{i(a,b)} = w'_{ia} + w'_{ib}$$
 and  $(B_{G''}u'_0)_{i(a,b)} = \sum_{x;(i,x)\in E'_0} (w'_{ia}w'_{xb} + w'_{ib}w'_{xa}).$ 

Using  $w'_{ia}w'_{xb} = 0$  for  $(i, x) \notin E'_0, (a, b) \in E'_0$ , we get

$$\sum_{v;(i,x)\in E'_0} (w'_{ia}w'_{xb} + w'_{ib}w'_{xa}) = \sum_x (w'_{ia}w'_{xb} + w'_{ib}w'_{xa}) = w'_{ia} + w'_{ib},$$

and therefore  $w'B_{G''} = B_{G''}u'_0$ . We obtain  $u'_0B^t_{G''} = B^t_{G''}w'$  similarly. From (16) and the form of w'', we see that  $w''A_{G''} = A_{G''}w''$  is equivalent to  $w'A_{G'_0} = A_{G'_0}w'$ ,  $w'B_{G''} = B_{G''}u'_0$ ,  $u'_0B^t_{G''} = B^t_{G''}w'$ ,  $u'_iC_{i+1} = C_{i+1}u'_{i+1}$ ,  $u'_{i+1}C^t_{i+1} = C^t_{i+1}u'_i$ . Note that  $u'_iC_{i+1} = C_{i+1}u'_{i+1}$ ,  $u'_{i+1}C^t_{i+1} = C^t_{i+1}u'_i$  are automatically fulfilled by definition of  $u'_i$  and  $C_i$ . Therefore, we get  $w''A_{G''} = A_{G''}w''$ . This yields a surjective \*-homomorphism  $\tilde{\varphi} : C(\operatorname{Qut}(G'')) \to C(\operatorname{Qut}^*(G')_{c_0})$ ,  $u_{ij} \mapsto w''_{ij}$  which is inverse to  $\varphi$ . Thus,  $\varphi$  is a \*-isomorphism.

It remains to show that  $\Delta_{G''} \circ \varphi = (\varphi \otimes \varphi) \circ \Delta_{G'}$ , which is true because of the following:

$$\Delta_{G''} \circ \varphi(w'_{ij}) = \Delta_{G'}(w_{ij})$$
  
=  $\sum_k w_{ik} \otimes w_{kj}$   
=  $(\varphi \otimes \varphi) \left( \sum_k w'_{ik} \otimes w'_{kj} \right)$   
=  $(\varphi \otimes \varphi) \circ \Delta_{G'}(w'_{ij}).$ 

| 1 | - |  |  |
|---|---|--|--|
|   |   |  |  |
|   |   |  |  |

We get the following corollary, which we will use in the next section to construct a graph with quantum symmetry and finite quantum automorphism group.

**Corollary 4.14.** Let G be a vertex – and edge-colored graph such that  $deg(v) \ge 3$  for all  $v \in V(G)$ . Denote by G' and G'' the graphs constructed in Definitions 4.5 and 4.9. If  $Qut_{c_0}^*(G') \cong Qut(G')$ , then  $Qut(G'') \cong Qut(G)$ .

## 5 (Uncolored) graphs whose quantum automorphism group is the dual of a solution group

In this section, we will look at certain graphs from Definition 3.2, where we replace one of the edge colors by non-edges. Furthermore, we restrict to graphs coming from linear constraint systems as in the following definition. By using Theorem 3.8 and the decoloring procedure in Section 4, we will obtain (decolored) graphs whose quantum automorphism group is the dual of the corresponding solution group.

**Definition 5.1.** Let *H* be a connected graph with vertex set [m] and label the edges  $1, \ldots, n := |E(H)|$ . Let  $M_H \in \mathbb{F}_2^{m \times n}$  be the matrix, where

$$(M_H)_{ki} = \begin{cases} 1 & \text{if } k \in V(H) \text{ is incident to } i \in E(H); \\ 0 & \text{o.w.} \end{cases}$$

**Definition 5.2.** Let *H* be a connected graph, let  $M_H$  be as in Definition 5.1 and let  $G(M_H, b)$  be the colored graph as in Definition 3.2. Recall  $S_k(M_H) = \{i \in [n] : (M_H)_{ki} = 1\}$ . Note that it holds  $|S_l(M_H) \cap S_k(M_H)| \le 1$  for all  $l \ne k$ , since otherwise there would be multiple edges between *l* and *k* in the graph *H* by definition of  $M_H$ . By Remarks 3.3 and 3.4, we know that we can color the edges between any  $V_l$  and  $V_k$ ,  $k \ne l$ , by the same  $(2^{|S_k \cap S_l|} - 1)$  numbers. In our case, this means that for every  $|S_l \cap S_k| \ne 0$ , we can use the same two numbers for any  $V_l$ ,  $V_k$ . We color the edge between  $(l, \alpha)$ ,  $(k, \beta)$  by  $\alpha_i \beta_i \in \{-1, 1\}$ , where  $i \in [n]$  is the number for which  $(M_H)_{ki} = (M_H)_{li} = 1$ . We then define the graph  $G_*(M_H, b)$  to be the graph obtained from the previous graph by replacing the edges of color 1 by non-edges.

**Remark 5.3.** As already mentioned in Remark 3.4, Theorem 3.8 is true if we replace  $G(M_H, 0)$  by  $G_*(M_H, 0)$ .

For the next lemma, recall the definition of  $\text{Qut}_{c_0}^*(G')$  from Definition 4.12. Note that a matching in a graph is a set of non-adjacent edges, i.e. no two edges share a common vertex.

**Lemma 5.4.** Let H be a graph and  $G_*(M_H, 0)$  as in Definition 5.2. We choose the color of the added paths in  $G'_*(M_H, 0)$  to be -1, i.e. the same color as any edge between  $V_l$  and  $V_k$  for  $k \neq l$ . Choosing  $c_0 = -1$ , we get  $\operatorname{Qut}_{-1}^*(G'_*(M_H, 0)) \cong \operatorname{Qut}(G'_*(M_H, 0))$ .

*Proof.* We start by showing that every edge-color class except  $c_0$  in  $G'_*(M_H, 0)$  is a matching. For this, first note that every edge between  $V_k$  and  $V_l$ ,  $k \neq l$  as well as every edge in an added path has color  $c_0$ . Thus, the remaining edges are between vertices  $(k, \alpha)$ ,  $(k, \beta)$  in  $V_k$  for some k. Thus, the color of the edge is the function  $\alpha \Delta \beta : S_k \to {\pm 1}^{S_k}$ ,  $(\alpha \Delta \beta)_i = \alpha_i \beta_i$ . Fix a vertex  $(k, \alpha)$ . For every  $\beta \neq \beta'$  it holds  $\alpha \Delta \beta \neq \alpha \Delta \beta'$ , as otherwise  $\beta_i = \beta'_i$  for all i. Thus, no two edges of the same color share a common vertex, which means that the color classes are matchings.

Let  $c((i, k)) = c((j, l)) \neq c_0$ . Then

$$u_{ij}u_{kl} = u_{ij}\left(\sum_{a;c((j,a))=c((i,k))} u_{ka}\right) = u_{ij}\left(\sum_{a} u_{ka}\right) = u_{ij},$$

because *l* is the only neighbor of *j* such that c((i,k)) = c((j,l)), since the color classes are matchings. Using the adjoint then yields  $u_{ij}u_{kl} = u_{kl}u_{ij}$ . We deduce  $\operatorname{Qut}_{-1}^*(G'_*(M_H, 0)) \cong \operatorname{Qut}(G'_*(M_H, 0))$ .

We get the following theorem.

**Theorem 5.5.** Let H be a graph with  $\deg(v) \ge 2$  for  $v \in V(H)$  and  $G_*(M_H, 0)$  be as in Definition 5.2. Let  $G''_*(M_H, 0)$  be the graph as in Lemma 5.4. Then  $\operatorname{Qut}(G''_*(M_H, 0)) \cong \widehat{\Gamma_H}$ .

*Proof.* First note that we have  $\deg(v) \ge 3$  for  $v \in V(G_*(M_H, 0))$  (since  $\deg(w) \ge 2$  for  $w \in V(H)$ , we see that every  $v \in V(G_*(M_H, 0))$  has at least one neighbor that is associated to the same equation and at least two neighbors that come from different equations but share one of the at least two variables in the equation). We also know  $\operatorname{Qut}_{-1}^*(G'_*(M_H, 0)) \cong \operatorname{Qut}(G'_*(M_H, 0))$  by Lemma 5.4. Thus, Corollary 4.14 yields  $\operatorname{Qut}(G''_*(M_H, 0)) \cong \operatorname{Qut}(G_*(M_H, 0))$ . Using Theorem 3.8, we deduce  $\operatorname{Qut}(G''_*(M_H, 0)) \cong \widehat{\Gamma_H}$ .

Using the theorem above, we are able to obtain a graph with quantum symmetry and finite quantum automorphism group. This graph is constructed from the constraint system associated to  $K_{3,4}$ .

**Corollary 5.6.** The graph  $G''_*(M_{K_{3,4}}, 0)$  has quantum symmetry and finite quantum automorphism group.

*Proof.* It is easy to see that  $\deg(v) \geq 2$  for all  $v \in V(K_{3,4})$ . Therefore  $C(\operatorname{Qut}(G''_*(M_{K_{3,4}}, 0)) \cong C^*(\Gamma_{K_{3,4}}))$  by the previous theorem. Furthermore, it was shown in [13] (by using Sage [16]) that  $\Gamma_{K_{3,4}}$  is finite and non-abelian, which yields that  $C^*(\Gamma_{K_{3,4}})$  is finite-dimensional and non-commutative. By the isomorphism above, we get that  $C(\operatorname{Qut}(G''_*(M_{K_{3,4}}, 0)))$  is finite-dimensional and non-commutative which yields the assertion.

## 6 Quantum Isomorphisms

In this section we consider quantum isomorphisms between the colored graphs G(M, b) and G(M, b') for  $b \neq b'$ . In particular we give analogs of Lemma 3.6 and Theorem 3.8 for this case. From this we are able to obtain non-isomorphic colored graphs G(M, b) and G(M, b') that are quantum isomorphic but neither G(M, b) nor G(M, b') has quantum symmetry. Further, applying decoloring techniques as in Section 4 allows us to produce non-isomorphic uncolored graphs G and G' without quantum symmetry that are nevertheless quantum isomorphic. This appears to be the first known such example.

To define quantum isomorphism of colored graphs, we must provide a suitable generalization of the isomorphism game. The way to do this is quite natural.

**Definition 6.1** (Isomorphism game for colored graphs). Given colored graphs G and G', with respective color functions c and c', the (G, G')-isomorphism game has as inputs and outputs for both Alice and Bob the set  $V(G) \cup V(G')^2$ . To win, upon receiving  $x \in V(G)$  Alice (respectively Bob) must respond with  $y \in V(G')$  and vice versa. If this condition is met, then there is a vertex  $g_a \in V(G)$  that is either Alice's input or output, and there is similarly  $g_b \in V(G)$  and  $g'_a, g'_b \in V(G')$ . Alice and Bob then win if the following conditions are met:

- 1.  $c(g_a) = c'(g'_a)$  and  $c(g_b) = c'(g'_b)$ ;
- 2.  $g_a = g_b$  if and only if  $g'_a = g'_b$ ;
- 3.  $(g_a, g_b)$  is an edge of color *c* if and only if  $(g'_a, g'_b)$  is an edge of color *c*.

We then say that two colored graphs G and G' are *quantum isomorphic* if there is a quantum strategy<sup>3</sup> that wins the (G, G')-isomorphism game with probability 1. In [10] it was shown that (uncolored) graphs G and G' are quantum isomorphic if and only if there exists a magic unitary u such that  $A_G u = uA_{G'}$ . Precisely the same proof applied to colored graphs gives the following:

**Proposition 6.2.** Colored graphs G and G' with coloring functions c and c' are quantum isomorphic if and only if there exists a  $V(G) \times V(G')$  magic unitary u such that  $c(g) \neq c'(g')$  implies  $u_{gg'} = 0$  for  $g \in V(G)$ ,  $g' \in V(G')$ , and  $A_{G_c}u = uA_{G'_c}$  for all edge colors c.

Based on the above, we define the following<sup>4</sup>:

<sup>&</sup>lt;sup>2</sup>We are implicitly assuming here that the vertex sets of G and G' are disjoint. Alternatively we can assume that as part of their input they are told which graph their input should be associated to, and similarly for their output.

<sup>&</sup>lt;sup>3</sup>There are a few different models of quantum strategies that can be considered. In this work we are always referring to the commuting operator framework when we mention quantum strategies.

<sup>&</sup>lt;sup>4</sup>This  $C^*$ -algebra is analogous to the \*-algebra of a nonlocal game defined in [9], applied to the isomorphism game.

**Definition 6.3** (Isomorphism algebra). Given colored graphs G and G' with coloring functions c and c', we define the *isomorphism algebra*, denoted Iso(G, G') as the universal  $C^*$ -algebra generated by elements  $u_{gg'}$  for  $g \in V(G)$ ,  $g' \in V(G')$  such that

$$u_{gg'}^2 = u_{gg'}^* = u_{gg'} \text{ for all } g \in V(G), g' \in V(G');$$
(18)

$$\sum_{h \in V(G)} u_{hg'} = 1 = \sum_{h' \in V(G')} u_{gh'} \text{ for all } g \in V(G), g' \in V(G');$$
(19)

$$u_{ag'} = 0 \text{ if } c(g) \neq c'(g');$$
(20)

$$A_{G_c}u = uA_{G'_c} \text{ for all edge colors } c.$$
(21)

It is clear from Proposition 6.2 and the above definition that the following holds:

**Proposition 6.4.** Let G and G' be colored graphs. Then  $G \cong_q G'$  if and only if Iso(G, G') is nontrivial.

We will also need to define the following algebra for our results in this section:

**Definition 6.5.** For  $M \in \mathbb{F}_2^{m \times n}$  and  $b \in \mathbb{F}_2^m$ , we define  $\mathcal{A}(M, b)$  to be the universal  $C^*$  algebra generated by  $x_1, \ldots, x_n$  such that

- 1.  $x_i^2 = 1$  for all  $i \in [n]$ ;
- 2.  $x_i x_j = x_j x_i$  if there exists  $k \in [m]$  such that  $M_{ki} = M_{kj} = 1$ ;
- 3.  $\prod_{i:M_{ki}=1} x_i = (-1)^{b_k}$  for all  $k \in [m]$ .

Note that  $\mathcal{A}(M, b)$  is not  $C^*(\Gamma)$  for  $\Gamma = \Gamma(M, b)$ , unless b = 0. Rather, for  $b \neq 0$ , the algebra  $\mathcal{A}(M, b)$  is isomorphic to the algebra  $pC^*(\Gamma(M, b)) := \{px : x \in C^*(\Gamma(M, b))\}$  where  $p = \frac{1}{2}(1 - \gamma)$ , which may be trivial for some M and b. We will show that Iso(G(M, b), G(M, b')) is isomorphic to  $\mathcal{A}(M, b + b')$ , but first we need an analog of Lemma 3.6. We omit the proof because it is almost identical to the proof of Lemma 3.6. The only difference is that one index of the magic unitary u is a vertex of G(M, b) and the other is a vertex of G(M, b') instead of both being from the same graph.

**Lemma 6.6.** Let  $M \in \mathbb{F}_2^{m \times n}$  and  $b, b' \in \mathbb{F}_2^m$ . Set G = G(M, b) and G' = G(M, b'). If u is a  $V(G) \times V(G')$  magic unitary that gives a quantum isomorphism from G to G' then u has the form

|          | $V_1(G')$     | $V_2(G')$ |           |   | $V_m(G')$   |      |
|----------|---------------|-----------|-----------|---|---|------|
| $V_1(G)$ | $( u^{(1)} )$ | 0         | 0         |   | 0   |      |
| $V_2(G)$ | 0             | $u^{(2)}$ | 0         |   | 0   |      |
| :        | 0             | 0         | $u^{(3)}$ |   | $\begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ u^{(m)} \end{pmatrix}$ | (22) |
| ÷        |               | ÷         | ÷         | · | 0   |      |
| $V_m(G)$ | $\setminus 0$ | 0         | 0         | 0 | $u^{(m)}$ /   |      |

where each  $u^{(k)}$  is a magic unitary. Furthermore,  $u^{(k)}_{\alpha,\beta} := u_{(k,\alpha),(k,\beta)}$  depends only on k and the value of  $\alpha \triangle \beta$  and therefore the entries of  $u^{(k)}$  pairwise commute for each  $k \in [m]$ .

We are now able to prove the following partial analog of Theorem 3.8:

**Theorem 6.7.** Let  $M \in \mathbb{F}_2^{m \times n}$  and  $b \neq b' \in \mathbb{F}_2^m$ . Set G = G(M, b), G' = G(M, b'), and  $\mathcal{A} = \mathcal{A}(M, b + b')$ . Then there exists an isomorphism  $\varphi : \mathcal{A} \to Iso(G, G')$ .

*Proof.* The proof is very similar to that of Theorem 3.8. We will first give a homomorphism  $\varphi_1$ from A to Iso(G,G') and then a homomorphism  $\varphi_2$  in the other direction, and finally show that these are inverses of each other and therefore isomorphisms.

### Step 1: Construction of a \*-homomorphism $\varphi_1 : \mathcal{A} \to Iso(G, G')$ .

For this step, we will construct elements  $y_i$  of Iso(G, G') that satisfy the relations of the generators  $x_i$  of  $\mathcal{A}$ . This proves that there is a \*-homomorphism  $\varphi_1$  from  $\mathcal{A}$  to Iso(G, G') such that  $\varphi_1(x_i) = y_i$ . Later we will see that  $\varphi_1$  is in fact an isomorphism.

Let u be magic unitary of generators for Iso(G, G'). By Lemma 6.6,  $u = \bigoplus_{k \in [m]} u^{(k)}$  where each  $u^{(k)}$  is a  $V_k(G) \times V_k(G')$  magic unitary. Moreover, the operator  $u^{(k)}_{\alpha,\beta} := u_{(k,\alpha),(k,\beta)}$  depends only on k and the value of  $\alpha \Delta \beta \in \pm 1^{S_k}_{b_k+b'_k}$ . Thus, as in the proof of Lemma 3.6, for each  $\delta \in \pm 1_{b_k+b'_k}^{S_k}$  we let  $u_{\delta}^{(k)}$  denote  $u_{\alpha,\beta}^{(k)}$  such that  $\alpha \triangle \beta = \delta$ . Note that for any  $\alpha \in \pm 1_{b_k}^{S_k}$ , we have that

$$\{u_{\delta}^{(k)}: \delta \in \pm 1_{b_k + b'_k}^{S_k}\} = \{u_{\alpha,\beta}^{(k)}: \beta \in \pm 1_{b'_k}^{S_k}\},\tag{23}$$

and thus every row (and similarly every column) of  $u^{(k)}$  contains the same set of operators. Now we define  $y_i^{(k)} = \sum_{\alpha \in \pm 1_{b_k+b'_k}} \alpha_i u_{\alpha}^{(k)}$  for all  $k \in [m]$  and  $i \in S_k$ . We first aim to show

that  $y_i^{(k)}$  does not depend on k.

Fix  $i \in [n]$  and  $l, k \in [m]$  such that  $i \in S_l \cap S_k$ . For each  $\delta \in \pm 1^{S_l \cap S_k}$ , let  $A_{\delta}$  (resp.  $A'_{\delta}$ ) be the adjacency matrices of the graph consisting of the edges of G (resp. G') colored  $\delta$ . Further, let  $B_{\delta}$  (resp.  $B'_{\delta}$ ) be the submatrix of  $A_{\delta}$  (resp.  $A'_{\delta}$ ) consisting of the rows indexed by  $V_l(G)$  and columns indexed by  $V_k(G)$  (resp.  $V_l(G')$  and  $V_k(G')$ ). In other words,

$$(B_{\delta})_{(l,\alpha),(k,\beta)} = \begin{cases} 1 & \text{if } \alpha \triangle \beta = \delta \\ 0 & \text{o.w.} \end{cases}$$

and similarly for  $B'_{\delta}$ . Also define  $A_+ = \sum_{\delta \in \pm 1^{S_l \cap S_k}, \delta_i = +1} A_{\delta}$  and  $A_- = \sum_{\delta \in \pm 1^{S_l \cap S_k}, \delta_i = -1} A_{\delta}$ , and similarly define  $A'_+$ ,  $A'_-$ ,  $B_+$ ,  $B_-$ ,  $B'_+$ , and  $B'_-$ . Then

$$(B_{+})_{(l,\alpha),(k,\beta)} = \begin{cases} 1 & \text{if } \alpha_{i} = \beta_{i} \\ 0 & \text{o.w.} \end{cases} & \& \quad (B_{-})_{(l,\alpha),(k,\beta)} = \begin{cases} 1 & \text{if } \alpha_{i} = -\beta_{i} \\ 0 & \text{o.w.} \end{cases}$$

and similarly for  $B'_+$  and  $B'_-$ . Since u must satisfy  $A_{\delta}u = uA'_{\delta}$  by definition, it must also satisfy  $A_{\pm}u = uA'_{\pm}$ . This implies that  $B_+u^{(k)} = u^{(l)}B'_+$  and  $B_-u^{(k)} = u^{(l)}B'_-$ . Considering the  $(l, \alpha), (k, \beta)$  entry of both sides of the former equation, we see that

$$\sum_{\substack{\beta' \in \pm 1_{b_k}^{S_k} \\ \text{s.t. } \beta'_i = \alpha_i}} u_{\beta',\beta}^{(k)} = \sum_{\substack{\alpha' \in \pm 1_{b'_l}^{S_l} \\ \text{s.t. } \alpha'_i = \beta_i}} u_{\alpha,\alpha'}^{(l)}$$

Note that for every term in the above sums, we have that  $\alpha_i \alpha'_i = \alpha_i \beta_i = \beta'_i \beta_i$ . Thus, reexpressing  $u_{\alpha,\alpha'}^{(l)}$  as  $u_{\tilde{\alpha}}^{(l)}$  where  $\tilde{\alpha} = \alpha \triangle \alpha'$ , and similarly for  $u_{\beta',\beta}^{(k)}$  we have that

$$\sum_{\substack{\tilde{\beta} \in \pm 1_{b_k+b'_k}^{S_k} \\ \text{s.t. } \tilde{\beta}_i = \alpha_i \beta_i}} u_{\tilde{\beta}}^{(k)} = \sum_{\substack{\tilde{\alpha} \in \pm 1_{b_l+b'_l}^{S_l} \\ \text{s.t. } \tilde{\alpha}_i = \alpha_i \beta_i}} u_{\tilde{\alpha}}^{(l)}$$

Doing the same for  $B_{-}u^{(k)} = u^{(l)}B'_{-}$  yields the same equation but with  $\alpha_{i}\beta_{i}$  replaced with  $-\alpha_i\beta_i$  and combining these proves that

$$\sum_{\tilde{\beta} \in \pm 1_{b_k + b'_k}^{S_k}, \tilde{\beta}_i = x} u_{\tilde{\beta}}^{(k)} = \sum_{\tilde{\alpha} \in \pm 1_{b_l + b'_l}^{S_l}, \tilde{\alpha}_i = x} u_{\tilde{\alpha}}^{(l)}$$

for any  $x \in \{+1, -1\}$ . Therefore,

$$y_{i}^{(l)} = \left(\sum_{\substack{\alpha \in \pm 1_{b_{l}+b_{l}'}^{S_{l}} \\ \text{s.t. } \alpha_{i} = +1}} u_{\alpha}^{(l)}\right) - \left(\sum_{\substack{\alpha \in \pm 1_{b_{l}+b_{l}'}^{S_{l}} \\ \text{s.t. } \alpha_{i} = -1}} u_{\alpha}^{(l)}\right) = \left(\sum_{\substack{\alpha \in \pm 1_{b_{k}+b_{k}'}^{S_{k}} \\ \text{s.t. } \alpha_{i} = +1}} u_{\alpha}^{(k)}\right) - \left(\sum_{\substack{\alpha \in \pm 1_{b_{k}+b_{k}'}^{S_{k}} \\ \text{s.t. } \alpha_{i} = -1}} u_{\alpha}^{(k)}\right) = y_{i}^{(k)}.$$

So we have shown that the value of  $y_i^{(k)}$  does not depend on k, and thus we will simply denote this operator by  $y_i$ .

Now note that since  $y_i$  is a linear combination (with real coefficients) of the operators  $u_{\alpha}^{(k)}$  for  $k \in [m]$  such that  $i \in S_k$ , and these operators are entries of the magic unitary u, we have that  $y_i^* = y_i$ . Also, by (23) we have that  $u_{\alpha}^{(k)}u_{\beta}^{(k)} = 0$  for  $\alpha \neq \beta$  and thus

$$y_i^2 = \left(\sum_{\alpha \in \pm 1_0^{S_k}} \alpha_i u_{\alpha}^{(k)}\right)^2 = \sum_{\alpha \in \pm 1_0^{S_k}} \alpha_i^2 \left(u_{\alpha}^{(k)}\right)^2 = \sum_{\alpha \in \pm 1_0^{S_k}} u_{\alpha}^{(k)} = 1.$$

Thus the  $y_i$  satisfy relation (1) from Definition 6.5.

Next we will show that relation (2) of Definition 6.5 holds, i.e., that  $y_i y_j = y_j y_i$  if there exists  $k \in [m]$  such that  $i, j \in S_k$ . Suppose that i, j, k are as described. Then  $y_i = y_i^{(k)}$  and  $y_j = y_j^{(k)}$  are both linear combinations of the entries of  $u^{(k)}$  which pairwise commute by Lemma 6.6. Therefore  $y_i$  and  $y_j$  commute as desired.

Lastly, we must show that relation (3) of Definition 6.5 holds, i.e., that  $\prod_{i \in S_k} y_i = (-1)^{b_k + b'_k}$  for all  $k \in [m]$ . We have that

$$\prod_{i \in S_k} y_i = \prod_{i \in S_k} y_i^{(k)}$$
$$= \prod_{i \in S_k} \left( \sum_{\alpha \in \pm 1_{b_k + b'_k}^{S_k}} \alpha_i u_\alpha^{(k)} \right)$$

Since the elements  $u_{\alpha}^{(k)}$  are pairwise orthogonal, when we expand the above product all cross terms disappear and we obtain

$$\prod_{i \in S_k} y_i = \sum_{\alpha \in \pm 1_{b_k + b'_k}^{S_k}} \left( \prod_{i \in S_k} \alpha_i \right) u_{\alpha}^{(k)} = \sum_{\alpha \in \pm 1_{b_k + b'_k}^{S_k}} (-1)^{b_k + b'_k} u_{\alpha}^{(k)} = (-1)^{b_k + b'_k},$$

since  $\prod_{i \in S_k} \alpha_i = (-1)^{b_k + b'_k}$  for all  $\alpha \in \pm 1^{S_k}_{b_k + b'_k}$  by definition. Therefore the elements  $y_i \in Iso(G, G')$  for  $i \in [n]$  satisfy the relations of the generators of  $\mathcal{A}$  and thus there exists a \*-homomorphism  $\varphi_1$  from  $\mathcal{A}$  to Iso(G, G') such that  $\varphi_1(x_i) = y_i$  for all  $i \in [n]$ .

Step 2: Construction of a \*-homomorphism  $\varphi_2 : Iso(G, G') \to \mathcal{A}$ .

This step is almost identical to Step 2 of the proof of Theorem 3.8, and so we omit it. We only remark that the biggest change is that when showing that  $v_{\alpha}^{(k)} = 0$  for  $\alpha \notin \pm 1_{b_k+b'_k}^{S_k}$ , the analog of Equation (8) gets an additional factor of  $(-1)^{b_k+b'_k}$  in the first and last expressions.

*Step 3: Showing that*  $\varphi_1$  *and*  $\varphi_2$  *are inverses of each other.* 

This step is nearly identical to Step 3 of the proof of Theorem 3.8 and so we omit it.

**Corollary 6.8.** Let  $M \in \mathbb{F}_2^{m \times n}$  and  $b \neq b' \in \mathbb{F}_2^m$ . Then the following are equivalent:

- 1. there is a quantum isomorphism from G(M, b) to G(M, b');
- 2. the algebra  $\mathcal{A}(M, b + b')$  is nontrivial;
- 3. the element  $\gamma$  of the solution group  $\Gamma(M, b + b')$  is not equal to the identity.

*Proof.* Proposition 6.2 states that (1) is equivalent to Iso(G(M, b), G(M, b')) being nontrivial, which is equivalent to (2) by Theorem 6.7. To see that (2) and (3) are equivalent, recall that  $\mathcal{A}(M, b + b')$  is isomorphic to  $pC^*(\Gamma(M, b + b')) = \{px : x \in nC^*(\Gamma(M, b + b'))\}$  where  $p = \frac{1}{2}(1 - \gamma)$ , and so these algebras are nontrivial if and only if  $\gamma \neq 1$  in  $C^*(\Gamma(M, b + b'))$  which is equivalent to  $\gamma$  not being equal to the identity in  $\Gamma(M, b + b')$ .

Note that the equivalence of (1) and (3) in the corollary above differs from earlier results (e.g., from [1]) because the graphs G(M, b) and G(M, b') here are *colored* graphs, rather than uncolored ones.

We are now able to give two non-isomorphic colored graphs without quantum symmetry that are quantum isomorphic. Recall the definition of  $G_*(M, 0)$  from Definition 5.2.

**Corollary 6.9.** Let M be the incidence matrix of  $K_{3,3}$ , and let  $b \in \mathbb{F}_2^{V(K_{3,3})}$  be any vector with exactly one entry equal to 1. Then  $G_*(M, 0)$  and  $G_*(M, b)$  are non-isomorphic colored graphs with no quantum symmetry which are nevertheless quantum isomorphic.

*Proof.* In this case it is well known that the element  $\gamma$  of the solution group  $\Gamma(M, b + 0) = \Gamma(M, b)$  is not equal to the identity [7], since this is the solution group of the Mermin-Peres magic square game. It then follows from Corollary 6.8 that  $G_*(M, 0)$  and  $G_*(M, b)$  are quantum isomorphic. On the other hand, it is known that even the uncolored versions of these graphs are non-isomorphic [1], and thus the colored versions are as well. Lastly, it was shown in [13] that  $\Gamma(M, 0)$  is abelian, and therefore by Theorem 3.8 the algebras  $C(\text{Qut}(G_*(M, 0)))$  and  $C(\text{Qut}(G_*(M, b)))$  are commutative, i.e., the graphs  $G_*(M, 0)$  and  $G_*(M, b)$  have no quantum symmetry.

**Corollary 6.10.** Let M be the incidence matrix of  $K_{3,3}$ , and let  $b \in \mathbb{F}_2^{V(K_{3,3})}$  be any vector with exactly one entry equal to 1. Let  $G''_*(M, 0)$  and  $G''_*(M, b)$  the graphs constructed from  $G_*(M, 0)$  and  $G_*(M, b)$  as in Lemma 5.4, where we choose the added paths in both graphs to have the same length for vertices of the same color and edges of the same color. Then  $G''_*(M, 0)$  and  $G''_*(M, b)$  are non-isomorphic graphs with no quantum symmetry which are nevertheless quantum isomorphic.

*Proof.* We start by showing that  $G''_*(M, 0)$  and  $G''_*(M, b)$  have no quantum symmetry. By the previous corollary, we know that  $G_*(M, 0)$  and  $G_*(M, b)$  have no quantum symmetry. We have  $C(\operatorname{Qut}^*_{-1}(G'_*(M, 0))) \cong C(\operatorname{Qut}(G'_*(M, 0)))$  and  $C(\operatorname{Qut}^*_{-1}(G'_*(M, b))) \cong C(\operatorname{Qut}(G'_*(M, b)))$  by Lemma 5.4. Therefore, Corollary 4.14 yields that  $G''_*(M, 0)$  and  $G''_*(M, b)$  have no quantum symmetry.

It remains to show that  $G''_*(M, 0)$  and  $G''_*(M, b)$  are quantum isomorphic. By Corollary 6.9, we know that  $G_*(M, 0)$  and  $G_*(M, b)$  are quantum isomorphic. Thus, there exists a  $C^*$ -algebra A and a matrix  $U \in M_n(A)$  such that Relations (18)–(21) hold. In  $G''_*(M, 0)$ , we denote the vertex that subdivided the edge e in  $G_*(M, 0)$  by  $e_0$  and the vertices in the added path with  $d(e_0, e_i) = i$  by  $e_i$ ,  $1 \le i \le m_{c(e)}$ . Moreover, we denote by  $v_i$ ,  $0 \le i \le n_{c(v)}$  the vertices in the added path with  $d(v, v_i) = i$ . We similarly label the vertices of  $G''_*(M, b)$ .

Note that by a similar reasoning as in Lemma 5.4, we get  $u_{ij}u_{kl} = u_{kl}u_{ij}$  for all e = (i, k), f = (j, l) with  $c(e) \neq c_0 \neq c'(f)$ . Define the magic unitary *W* as follows:

$$\begin{split} w_{i_k j_l} &= \delta_{kl} u_{ij} \text{ for } i \in V(G(M,0)), j \in V(G(M,b)), \\ w_{e_k f_l} &= \delta_{kl} (u_{ac} u_{bd} + u_{bd} u_{ac}) \text{ for } e = (a,b) \in E(G(M,0)), f = (c,d) \in E(G(M,b)), \end{split}$$

$$\begin{split} w_{i_k f_l} &= 0 \text{ for } i \in V(G(M,0)), f \in E(G(M,b)), \\ w_{e_k j_l} &= 0 \text{ for } e \in E(G(M,0)), j \in V(G(M,b)), \end{split}$$

for all occurring k, l. Similar to the proofs of Theorem 4.7 and Theorem 4.13, we see that  $A_{G''_*(M,0)}W = WA_{G''_*(M,b)}$ . We conclude that  $G''_*(M,0)$  and  $G''_*(M,b)$  are quantum isomorphic.

### Acknowledgments.

**DR** was partially supported by the VILLUM Centre of Excellence for the Mathematics of Quantum Theory (QMATH) project no. 35544.

**SSch** would like to thank Matthew Daws and Christian Voigt for helpful discussions. This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie grant agreement No. 101030346.

## References

- Albert Atserias, Laura Mančinska, David E. Roberson, Robert Šámal, Simone Severini, and Antonios Varvitsiotis. Quantum and non-signalling graph isomorphisms. J. Combin. Theory Ser. B, 136:289–328, 2019.
- [2] Teodor Banica. Quantum automorphism groups of homogeneous graphs. J. Funct. Anal., 224(2):243–280, 2005.
- [3] Teodor Banica. Quantum automorphism groups of small metric spaces. *Pacific J. Math.*, 219(1):27–51, 2005.
- [4] Teodor Banica and J.P. MacCarthy. The Frucht property in the quantum group setting. *arXiv:2106.04999*, 2021.
- [5] Julien Bichon. Quantum automorphism groups of finite graphs. *Proc. Amer. Math. Soc.*, 131(3):665–673, 2003.
- [6] Richard Cleve, Li Liu, and William Slofstra. Perfect commuting-operator strategies for linear system games. J. Math. Phys., 58(1):012202, 7, 2017.
- [7] Richard Cleve and Rajat Mittal. Characterization of binary constraint system games. In Automata, languages, and programming. Part I, volume 8572 of Lecture Notes in Comput. Sci., pages 320–331. Springer, Heidelberg, 2014.
- [8] Melanie Fulton. The quantum automorphism group and undirected trees. *PhD Thesis, Virginia*, 2006.
- [9] J William Helton, Kyle P Meyer, Vern I Paulsen, and Matthew Satriano. Algebras, synchronous games, and chromatic numbers of graphs. *New York Journal of Mathematics*, 2019.
- [10] Martino Lupini, Laura Mančinska, and David E. Roberson. Nonlocal games and quantum permutation groups. *Journal of Functional Analysis*, 279(5):108592, 2020. doi:https://doi.org/10.1016/j.jfa.2020.108592.
- [11] Ann Maes and Alfons Van Daele. Notes on compact quantum groups. *Nieuw Arch. Wisk.* (4), 16(1-2):73–112, 1998.
- [12] N. David Mermin. Simple unified form for the major no-hidden-variables theorems. *Phys. Rev. Lett.*, 65, 1990.

- [13] Connor Paul-Paddock. Algebraic and combinatorial aspects of incidence groups and linear system non-local games arising from graphs. *Master's thesis*, 2019.
- [14] Simon Schmidt. On the quantum symmetry of distance-transitive graphs. *Advances in Mathematics*, 368:107150, 2020.
- [15] Simon Schmidt. Quantum automorphism groups of finite graphs. *PhD Thesis, Saarland University*, 2020.
- [16] The Sage Developers. SageMath, the Sage Mathematics Software System (Version 8.9), 2019. https://www.sagemath.org.
- [17] Shuzhou Wang. Quantum symmetry groups of finite spaces. *Comm. Math. Phys.*, 195(1):195–211, 1998.
- [18] Moritz Weber. On *C*\*-algebras generated by isometries with twisted commutation relations. *J. Funct. Anal.*, 264(8):1975–2004, 2013.
- [19] S. L. Woronowicz. Compact matrix pseudogroups. Comm. Math. Phys., 111(4):613–665, 1987.
- [20] S. L. Woronowicz. Compact quantum groups. In *Symétries quantiques (Les Houches, 1995)*, pages 845–884. North-Holland, Amsterdam, 1998.