

# COMPLETE INTERPOLATING SEQUENCES FOR THE GAUSSIAN SHIFT-INVARIANT SPACE

ANTON BARANOV, YURII BELOV, AND KARLHEINZ GRÖCHENIG

ABSTRACT. We give a full description of complete interpolating sequences for the shift-invariant space generated by the Gaussian. As a consequence, we rederive the known density conditions for sampling and interpolation.

## 1. MAIN RESULTS

Consider the shift-invariant space of functions on  $\mathbb{R}$  with Gaussian generator  $g(z) = e^{-az^2}$  for  $a > 0$  defined as

$$V^2 = \left\{ f(z) = \sum_{n \in \mathbb{Z}} c_n e^{-a(z-n)^2} : (c_n) \in \ell^2(\mathbb{Z}) \right\}.$$

We consider the space  $V^2$  as a subspace of  $L^2(\mathbb{R})$  with the usual  $L^2$ -norm.

The space  $V^2$  belongs to the general family of shift-invariant spaces. Given a generator  $g \in L^2(\mathbb{R})$ , such a space is defined as

$$V^2(g) = \left\{ f(z) = \sum_{n \in \mathbb{Z}} c_n g(z-n) : (c_n) \in \ell^2(\mathbb{Z}) \right\} \subseteq L^2(\mathbb{R}).$$

The primary example is the classical Paley–Wiener space  $PW = \{f \in L^2(\mathbb{R}) : \text{supp } \hat{f} \subseteq [-1/2, 1/2]\}$ , which, by the sampling theorem of Shannon–Whittaker–Kotelnikov, can be identified with the shift-invariant space  $V^2(\frac{\sin \pi x}{\pi x})$ . In signal processing [1] shift-invariant spaces are often taken as a substitute for the Paley–Wiener space. A unifying feature of both  $V^2$  (Gaussian generator) and  $PW$  (sinc-generator) is the fact that both spaces can be viewed as spaces of entire functions by interpreting the variable  $z$  to be in  $\mathbb{C}$ .

It is easy to see that the norm equivalence  $\|f\|_{L^2(\mathbb{R})} \asymp \|(c_n)\|_{\ell^2(\mathbb{Z})}$  holds on  $V^2$  with some absolute constants. In what follows it will be convenient for us to work with the second quantity and so we put  $\|f\|_{V^2} := \|(c_n)\|_{\ell^2(\mathbb{Z})}$ , and we often identify  $V^2$  with  $\ell^2(\mathbb{Z})$  via the mapping  $(c_n) \mapsto f(z) = \sum_{n \in \mathbb{Z}} c_n e^{-(z-n)^2}$ .

A sequence  $\Lambda = \{\lambda_n\}_{n \in \mathbb{Z}} \subset \mathbb{R}$  is said to be *sampling for  $V^2$* , if

$$\sum_{n \in \mathbb{Z}} |f(\lambda_n)|^2 \asymp \|f\|_{V^2}^2, \quad f \in V^2,$$

---

2000 *Mathematics Subject Classification.* Primary 30H20; Secondary 30D10, 30E05, 42C15, 94A20.

*Key words and phrases.* Sampling, interpolation, Riesz bases, small Fock spaces, shift-invariant space, Avdonin-type condition.

A. B. and Yu. B. were supported by the Ministry of Science and Higher Education of the Russian Federation, agreement No 075-15-2021-602, and by the Russian Foundation for Basic Research grant 20-51-14001-ANF-a. K. G. was supported in part by the project P31887-N32 of the Austrian Science Fund (FWF).

and we say that  $\Lambda$  is *interpolating for  $V^2$*  if for every  $(a_n) \in \ell^2(\mathbb{Z})$  there exists  $f \in V^2$  such that  $f(\lambda_n) = a_n$ . If  $\Lambda$  is both sampling and interpolating (or, equivalently, the solution of the interpolation problem is unique), we say that  $\Lambda$  is a *complete interpolating sequence*.

Sampling and interpolation in the space  $V^2$  and in more general shift-invariant spaces were studied in [8, 9]. The main result can be formulated as follows. Recall that  $\Lambda$  is said to be separated if  $\inf_{\lambda, \lambda' \in \Lambda, \lambda \neq \lambda'} |\lambda - \lambda'| > 0$ , and denote by  $D^+(\Lambda)$  and  $D^-(\Lambda)$  the usual upper and lower Beurling densities

$$D^+(\Lambda) = \lim_{r \rightarrow \infty} \frac{\sup_{x \in \mathbb{R}} \text{card}(\Lambda \cap [x, x+r])}{r}, \quad D^-(\Lambda) = \lim_{r \rightarrow \infty} \frac{\inf_{x \in \mathbb{R}} \text{card}(\Lambda \cap [x, x+r])}{r}.$$

**Theorem 1.1.** (i) *Sufficiency: Every separated sequence  $\Lambda \subset \mathbb{R}$  with  $D^-(\Lambda) > 1$  is sampling for  $V^2$ . Moreover, it contains a complete interpolating sequence.*  
(ii) *Necessity: If  $\Lambda$  is sampling for  $V^2$ , then  $\Lambda$  is a finite union of separated subsequences and  $\Lambda$  contains a separated sequence  $\tilde{\Lambda}$  such that  $D^-(\tilde{\Lambda}) \geq 1$ .*

**Theorem 1.2.** (i) *Every separated sequence  $\Lambda$  with  $D^+(\Lambda) < 1$  is a set of interpolation for  $V^2$ . Moreover, it can be enlarged to a complete interpolating sequence.*  
(ii) *If  $\Lambda$  is a set of interpolation for  $V^2$ , then  $\Lambda$  is separated and  $D^+(\Lambda) \leq 1$ .*

One of the main insights obtained in [8, 9] is the similarity of  $V^2$  with the Paley–Wiener space  $PW$  with respect to sampling and interpolation. In particular, for both  $V^2$  and  $PW$  the same density conditions hold. For  $PW$  these go back to Beurling, Kahane, and Landau [6, 14, 15, 18].

Theorems 1.1 and 1.2 provide an almost characterization of sampling sets and of interpolating sets, but they leave open the case of critical density. If a set is simultaneously sampling and interpolating, then  $D(\Lambda) = 1$ . The case of the critical density is much more subtle, because anything may happen. For  $PW$  the problem of complete interpolating sequences was solved in [12] and [16], for  $V^2$  it was open so far.

Our main result is a complete and explicit description of complete interpolating sequences for  $V^2$ . Moreover, this characterization can be extended to the larger class of complex Gaussians

$$g_c(z) = e^{-cz^2}, \quad c = a + ib, \quad a > 0, \quad b \in \mathbb{R}.$$

We denote the space  $V^2(g_c)$  by  $V_c^2$ .

It is easy to see that for  $f(z) = \sum_{n \in \mathbb{Z}} c_n e^{-c(z-n)^2}$  we still have  $\|f\|_{L^2(\mathbb{R})} \asymp \|(c_n)\|_{\ell^2(\mathbb{Z})}$  and so we can define the norm in  $V_c^2$  by  $\|f\|_{V_c^2} := \|(c_n)\|_{\ell^2(\mathbb{Z})}$ .

**Theorem 1.3.** *Given  $c \in \mathbb{C}$  with  $a = \text{Re } c > 0$ , an increasing sequence  $\Lambda \subset \mathbb{R}$  is a complete interpolating sequence for  $V_c^2$ , if and only if  $\Lambda$  is separated and there exists an enumeration  $\Lambda = \{\lambda_n\}_{n \in \mathbb{Z}}$ ,  $\lambda_n = n + \delta_n$ ,  $n \in \mathbb{Z}$ , such that*

- (a)  $(\delta_n)_{n \in \mathbb{Z}} \in \ell^\infty$ ;
- (b) *there exists  $N \geq 1$  and  $\delta > 0$  such that*

$$\sup_{n \in \mathbb{Z}} \frac{1}{N} \left| \sum_{k=n+1}^{n+N} \delta_k \right| \leq \delta < \frac{1}{2}.$$

Theorem 1.3 is stronger than the results in [8]. Indeed, as a corollary, we will deduce the density results for sampling or interpolation obtained previously in [8] and extend them to the spaces generated by complex Gaussians.

At least to us, Theorem 1.3 is quite surprising, as it shows a marked difference between  $V_c^2$  and  $PW$ . The characterizing conditions go back to Avdonin who showed that conditions (a) and (b) are sufficient for complete interpolating sequences in  $PW$ . However, they are far from necessary in  $PW$ . By contrast, in the shift-invariant spaces  $V_c^2$  these conditions provide a complete characterization.

To prove Theorem 1.3, we reduce the problem to the study of similar problems in some Fock-type space of entire functions. After this reduction is done, one can apply the description of complete interpolating sequences in this space obtained in [4].

## 2. UNITARY EQUIVALENCE WITH A FOCK-TYPE SPACE

For a parameter  $a > 0$ , we consider the Fock-type space of entire functions, sometimes referred to as a *small Fock space*, defined by

$$\mathcal{F} = \mathcal{F}_a = \left\{ F \text{ entire} : \|F\|_{\mathcal{F}_a}^2 = \frac{1}{2\sqrt{2\pi a}} \int_{\mathbb{C}} |F(w)|^2 e^{-\frac{1}{2a}(\log|w|)^2} dm_2(w) < \infty \right\},$$

where  $m_2$  is the Lebesgue measure  $\frac{1}{\pi} dx dy$ . A simple computation shows that for a function  $F(w) = \sum_{n \geq 0} b_n w^n$  one has

$$\|F\|_{\mathcal{F}_a}^2 = \sum_{n \geq 0} |b_n|^2 e^{2a(n+1)^2}.$$

In what follows we assume  $c \in \mathbb{C}$  to be fixed. To see the connection between  $\mathcal{F}_a$  and  $V_c^2$ , we note that

$$\sum_{n \in \mathbb{Z}} c_n e^{-c(z-n)^2} = e^{-cz^2} \sum_{n \in \mathbb{Z}} c_n e^{-cn^2} e^{2cnz}.$$

Introducing a new variable  $w = e^{2cz}$ ,  $V_c^2$  becomes unitarily equivalent to the space of functions representable as

$$\left\{ g(w) = \sum_{n \in \mathbb{Z}} d_n w^n : \|g\|^2 := \sum_{n \in \mathbb{Z}} |d_n|^2 e^{2an^2} < \infty \right\}.$$

This space consists of functions analytic in  $\mathbb{C} \setminus \{0\}$ . It is more convenient, however, to split this space into three parts, corresponding to positive and negative powers of  $w$ , and the constant term. Namely, for  $f \in V_c^2$  we write

$$\begin{aligned} f(z) &= \sum_{n \in \mathbb{Z}} c_n e^{-c(z-n)^2} = \sum_{n < 0} c_n e^{-c(z-n)^2} + c_0 e^{-cz^2} + \sum_{n > 0} c_n e^{-c(z-n)^2} \\ (1) \quad &= f_-(z) + c_0 e^{-cz^2} + f_+(z). \end{aligned}$$

With respect to this decomposition  $V_c^2$  splits into a direct sum  $V_c^2 = V_-^2 \oplus e^{-cz^2} \mathbb{C} \oplus V_+^2$ , which is, in terms of the coefficients, simply  $\ell^2(\mathbb{Z}) = \ell^2(\mathbb{N}_-) \oplus \mathbb{C} \oplus \ell^2(\mathbb{N})$ .

Next, for  $w = e^{2cz}$ , set

$$F_+(w) = \sum_{n=1}^{\infty} c_n e^{-cn^2} w^{n-1}, \quad F_-(w) = \sum_{n=1}^{\infty} c_{-n} e^{-cn^2} w^{n-1},$$

then in view of (1) we can write

$$f(z) = e^{-cz^2} (w^{-1} F_-(w^{-1}) + c_0 + w F_+(w)).$$

It follows from the discussion above that  $F_+, F_- \in \mathcal{F}_a$  and

$$\|F_+\|_{\mathcal{F}_a} = \left( \sum_{n>0} |c_n e^{-cn^2}|^2 e^{2an^2} \right)^{1/2} = \|c\|_2 = \|f_+\|_{V_c^2},$$

and  $\|F_-\|_{\mathcal{F}_a} = \|f_-\|_{V_c^2}$ . In addition every function  $F_+, F_- \in \mathcal{F}_a$  can appear in this representation and

$$wF_+(w) = e^{cz^2} f_+(z) \Big|_{e^{2cz}=w}.$$

This representation of elements in  $V_c^2$  makes it possible to reduce the study of complete interpolating sequences in  $V_c^2$  to the same problem in the Fock-type space  $\mathcal{F}_a$ .

Let us introduce the notions of sampling and interpolation for the space  $\mathcal{F}_a$ . Let  $\varphi(z) = \frac{1}{4a} \log^2 |z|$ . In the following we drop the reference to the parameter  $a$ , when it is not needed. We say that the sequence  $\{w_n\}$  is sampling for  $\mathcal{F}$ , if

$$(2) \quad \sum_n (1 + |w_n|^2) e^{-2\varphi(w_n)} |F(w_n)|^2 \asymp \|F\|_{\mathcal{F}}^2, \quad F \in \mathcal{F}.$$

Analogously,  $\{w_n\}$  is interpolating for  $\mathcal{F}$  if for every sequence  $\{a_n\}$  satisfying  $\sum_n (1 + |w_n|^2) e^{-2\varphi(w_n)} |a_n|^2 < \infty$  there exists  $F \in \mathcal{F}$  with  $F(w_n) = a_n$ . Recall from [5, Lemma 2.7] that the reproducing kernel  $k_w$  of  $\mathcal{F}$  has the norm

$$(3) \quad \|k_w\|_{\mathcal{F}}^2 \asymp \frac{e^{2\varphi(w)}}{1 + |w|^2}.$$

Thus the weights in the sampling inequality are nothing but  $\|k_{w_n}\|_{\mathcal{F}}^{-2}$ . So the norm equivalence (2) reads as  $\sum_n |\langle f, k_{w_n} \rangle_{\mathcal{F}}|^2 \|k_{w_n}\|_{\mathcal{F}}^{-2} \asymp \|f\|_{\mathcal{F}}^2$ . We may thus formulate the questions about sampling and interpolation in the equivalent language of frames and Riesz sequences of normalized reproducing kernels  $\tilde{k}_{w_n} = k_{w_n} / \|k_{w_n}\|_{\mathcal{F}}$  in  $\mathcal{F}$ . Namely,  $\{w_n\}$  is sampling for  $\mathcal{F}$  if and only if  $\{\tilde{k}_{w_n}\}$  is a frame in  $\mathcal{F}$ , while  $\{w_n\}$  is interpolating if and only if  $\{\tilde{k}_{w_n}\}$  is a Riesz sequence. Finally,  $\{w_n\}$  is a complete interpolating sequence if and only if  $\{\tilde{k}_{w_n}\}$  is a Riesz basis in  $\mathcal{F}$ .

We now formulate the description of complete interpolating sequences for  $\mathcal{F}_a$ . First it was shown in [5] that the sequence  $\{e^{2an}\}_{n \geq 1}$  is a complete interpolating sequence for  $\mathcal{F}_a$ . Then a full characterization of complete interpolating sequences was obtained in [4]. Remarkably, this characterization is formulated in terms of perturbations of the complete interpolating sequence  $\{e^{2an}\}_{n \geq 1}$ , somewhat in the spirit of the theorems of Kadets and Avdonin for complex exponentials (although this condition is only sufficient in  $PW$ ). Let  $\{w_n\}_{n \geq 1}$  be a sequence such that  $0 < |w_n| \leq |w_{n+1}|$  and let

$$w_n = e^{2an} e^{\delta_n} e^{i\theta_n}$$

with  $\delta_n, \theta_n \in \mathbb{R}$ . In [4] the following result was proved.

**Theorem 2.1.** [4] *A sequence  $\{w_n\}_{n \geq 1}$  is a complete interpolating sequence for  $\mathcal{F}_a^1$ , if and only if*

- (i)  $\{w_n\}$  is  $\mathcal{F}_a$ -separated, i.e., there exists  $\gamma > 0$  such that  $|w_m - w_n| \geq \gamma |w_n|$ ,  $m \neq n$ ;
- (ii)  $(\delta_n)_{n \geq 1} \in \ell^\infty$ ;

---

<sup>1</sup>Note that in [4] the normalization for the small Fock space is different; to formulate the results from [4] one needs to set  $\alpha = 1/(4a)$ .

(iii) there exists  $N \geq 1$  and  $\delta > 0$  such that

$$\sup_{n \geq 1} \frac{1}{N} \left| \sum_{k=n+1}^{n+N} \delta_k \right| \leq \delta < a.$$

Note that the result does not depend on the arguments of  $w_n$ , but only on their moduli. For the description of complete interpolating sequences for a more general class of Hilbert spaces of entire functions see [3].

### 3. PROOF OF THEOREM 1.3: SUFFICIENCY

Assume that  $\Lambda \subset \mathbb{R}$  is separated and that there exists an enumeration  $\Lambda = \{\lambda_m\}_{m \in \mathbb{Z}}$ ,  $\lambda_m = m + \delta_m$ ,  $m \in \mathbb{Z}$ , such that conditions (a) and (b) of Theorem 1.3 are satisfied. We need to show that the mapping

$$(4) \quad J : (c_n)_{n \in \mathbb{Z}} \mapsto (f(\lambda_m))_{m \in \mathbb{Z}}$$

is an isomorphism of  $\ell^2(\mathbb{Z})$  onto itself. Again we identify the sequence  $(c_n)$  and the corresponding function  $f \in V_c^2$ .

Recall that  $\varphi(w) = \frac{1}{4a}(\log |w|)^2$ , and put

$$w_m = e^{2c\lambda_m} = e^{2am} e^{2a\delta_m} e^{2ib(m+\delta_m)}.$$

The Avdonin-type condition (iii) of Theorem 2.1 reads as  $\frac{1}{N} \left| \sum_{k=n+1}^{n+N} (2a\delta_k) \right| \leq \delta < a$ , which amounts precisely to the assumption (b) of Theorem 1.3. Thus each of the sequences  $\{w_m\}_{m \geq 1}$  and  $\{w_m^{-1}\}_{m \geq 1}$  is a complete interpolating sequence for  $\mathcal{F}_a$ .

*Claim 1.* The mapping  $U_+ : (c_n)_{n \geq 1} \mapsto (f_+(\lambda_m))_{m \geq 1}$  is an isomorphism from  $\ell^2(\mathbb{N})$  onto itself. Indeed, since  $\varphi(w) = (\log |w|)^2 / (4a) = \lambda^2$  whenever  $w = e^{2c\lambda}$ , we have

$$(5) \quad f_+(\lambda_m) = e^{-c\lambda_m^2} w_m F_+(w_m) = e^{-\varphi(w_m)} e^{-ib\lambda_m^2} w_m F_+(w_m),$$

and we conclude that  $(f_+(\lambda_m))_{m \geq 1} \in \ell^2(\mathbb{N})$ , because  $\{w_m\}$  is sampling, and that every sequence in  $\ell^2(\mathbb{N})$  can be obtained in this way from some function in  $F_+$ , because  $\{w_m\}$  is interpolating. Thus  $U_+$  is one-to-one and onto. Similarly the mapping  $U_- : (c_n)_{n \leq -1} \mapsto (f_-(\lambda_m))_{m \leq -1}$  is an isomorphism on  $\ell^2(\mathbb{N}_-)$ .

*Claim 2.* The mapping  $K_+ : (c_n)_{n \geq 1} \mapsto (f_+(\lambda_m))_{m \leq -1}$  is compact from  $\ell^2(\mathbb{N})$  to  $\ell^2(\mathbb{N}_-)$ . Since

$$f_+(\lambda_m) = \sum_{n \geq 1} c_n e^{-c(m+\delta_m-n)^2},$$

the matrix of  $K_+$  has the entries  $e^{-c(m+\delta_m-n)^2}$ ,  $m < 0, n > 0$ . The boundedness of  $\delta_m$  yields that  $\sum_{m < 0, n > 0} e^{-a(|m|+n-\delta_m)^2} < \infty$ , thus  $K_+$  is a Hilbert–Schmidt operator. Similarly, the operator  $K_- : (c_n)_{n \leq -1} \mapsto (f_-(\lambda_m))_{m \geq 1}$  is compact from  $\ell^2(\mathbb{N}_-)$  to  $\ell^2(\mathbb{N})$ .

Now we define the operator  $U$  on  $\ell^2(\mathbb{Z}) = \ell^2(\mathbb{N}_-) \oplus \mathbb{C} \oplus \ell^2(\mathbb{N})$  by the formula

$$(6) \quad U : ((c_n)_{n \leq -1}; c_0; (c_n)_{n \geq 1}) \mapsto ((f_-(\lambda_m))_{m \leq -1}; f(\lambda_0); (f_+(\lambda_m))_{m \geq 1})$$

By Claim 1  $U$  is an isomorphism from  $\ell^2(\mathbb{Z})$  onto itself. At the same time, by Claim 2, the map  $J : (c_n)_{n \in \mathbb{Z}} \mapsto (f(\lambda_m))_{m \in \mathbb{Z}}$  differs from  $U$  by a compact operator, since  $f(\lambda_m) = f_+(\lambda_m) + c_0 e^{-c\lambda_m^2} + f_-(\lambda_m)$ . Thus, to prove that  $J$  is invertible it is sufficient to show that  $J$  is one-to-one, which is usually easier. In our context it means that  $\Lambda$  is a uniqueness set for the space  $V_c^2$ .

*Claim 3.*  $\Lambda$  is a uniqueness set for the space  $V_c^2$ . Assume that  $f \in V_c^2$  and  $f(\lambda_m) = 0$  for all  $m$ . Then the function

$$F(w) = w^{-1}F_-(w^{-1}) + c_0 + wF_+(w)$$

vanishes on the set  $w_m = e^{2c\lambda_m}$ .

Since  $W_+ = \{w_m\}_{m \geq 1}$  is a complete interpolating sequence for  $\mathcal{F}$ , there exists a so-called *generating function*  $G_+$  for this sequence. This is a function with simple zeros exactly at  $w_m$ ,  $m \geq 1$ , and no other zeros, such that  $G_+ \notin \mathcal{F}$ , but  $\frac{G_+}{w-w_m} \in \mathcal{F}$  for all  $m$ . In fact, the system  $\left\{\frac{G_+}{G'_+(w_m)(w-w_m)}\right\}_{m \geq 1}$  is the biorthogonal system in  $\mathcal{F}$  to the system of reproducing kernels  $\{k_{w_m}\}_{m \geq 1}$ .

We will need several estimates for the generating functions. First, let  $G_0$  be the generating function for the sequence  $W_0 = \{e^{2am}\}_{m \geq 1}$ . It was shown in [5, Lemma 2.6] that

$$|G_0(w)| \asymp e^{\varphi(w)} \frac{\text{dist}(w, W_0)}{1 + |w|^{3/2}}, \quad w \in \mathbb{C}.$$

Using standard estimates of lacunary canonical products it is easy to show (see [4, Section 3, p. 1373]) that for the generating function  $G_+$  of the perturbed sequence  $W_+ = \{w_m\}_{m \geq 1}$  with  $|w_m| = e^{2am+2a\delta m}$  one has

$$\frac{|G_+(w)|}{|G_0(w)|} \asymp \exp\left(-2a \sum_{k=1}^m \delta_k\right) \frac{\text{dist}(w, W_+)}{\text{dist}(w, W_0)}$$

whenever  $e^{a(2m-1)} \leq |w| \leq e^{a(2m+1)}$ . By condition (b) of Theorem 1.3, for sufficiently large  $m$  we have  $\exp(-2a \sum_{k=1}^m \delta_k) \geq \exp(-2a\delta m) \asymp |w|^{-\delta}$ , where  $\delta < \frac{1}{2}$ . Hence,

$$(7) \quad |G_+(w)| \gtrsim \frac{\text{dist}(w, W_+)}{(1 + |w|)^{\frac{3}{2} + \delta}} e^{\varphi(w)}.$$

Analogously, we define the function  $G_-$ , the generating function of the sequence  $W_- = \{w_n^{-1}\}_{n \leq -1}$ . Consequently, the zero set of the function  $(w-w_0)G_+(w)G_-(w^{-1})$  is precisely  $\tilde{\Lambda} = \{w_n : n \in \mathbb{Z}\}$ . Since by assumption  $F$  vanishes also on  $\tilde{\Lambda}$ ,  $F$  possesses the factorization

$$F(w) = (w-w_0)G_+(w)G_-(w^{-1})H(w)$$

for some function  $H$  analytic in  $\mathbb{C} \setminus \{0\}$ . We will show that  $H$  is an entire function which tends to 0 at infinity, and thus must be identically zero. Clearly,  $w^{-1}F_-(w^{-1}) \rightarrow 0$ , as  $|w| \rightarrow \infty$  and

$$|F_+(w)| = |\langle F_+, k_w \rangle| \leq \|F_+\|_{\mathcal{F}} \|k_w\|_{\mathcal{F}} \lesssim \frac{e^{\varphi(w)}}{1 + |w|}.$$

by (3). Consequently,

$$(8) \quad |F(w)| \lesssim 1 + |wF_+(w)| \lesssim |w| \frac{e^{\varphi(w)}}{1 + |w|} \leq e^{\varphi(w)}, \quad |w| \rightarrow \infty.$$

Thus, using (7), (8), and the fact that  $G_-(w^{-1}) \rightarrow G_-(0) \neq 0$ ,  $|w| \rightarrow \infty$ , we get

$$\begin{aligned} |H(w)| &= \frac{|F(w)|}{|(w-w_0)G_+(w)G_-(w)|} \\ &\lesssim e^{\varphi(w)} \cdot \frac{(1 + |w|)^{\frac{3}{2} + \delta}}{|w| \text{dist}(w, W_+)} e^{-\varphi(w)} \lesssim \frac{(1 + |w|)^{\frac{1}{2} + \delta}}{\text{dist}(w, W_+)}, \quad |w| \rightarrow \infty. \end{aligned}$$

It follows from the  $\mathcal{F}$ -separation of  $W_+$  and the inclusion  $W_+ \subset \mathbb{R}$  that there exists a sequence of radii  $r_k \rightarrow \infty$  such that  $\text{dist}(w, W_+) \geq \gamma|w|$  when  $|w| = r_k$ , with  $\gamma > 0$  independent on  $k$ . Since  $\delta < \frac{1}{2}$ , we conclude that  $\max_{|w|=r_k} |H(w)| \rightarrow 0$ , as  $k \rightarrow \infty$ . The maximum principle implies that also  $H(w) \rightarrow 0$ , as  $|w| \rightarrow \infty$ .

Analogously, replacing  $w$  by  $w^{-1}$  and using the estimate (7) for  $G_-$ , we conclude that  $H(w) \rightarrow 0$  as  $w \rightarrow 0$ . Consequently the singularity of  $H$  at 0 is removable, and  $H$  is thus entire and bounded. Thus  $H \equiv 0$  and so  $F \equiv 0$ , as was to be shown.  $\square$

To conclude we formulate a simple, sufficient condition for complete interpolating sets in the style of Kadets, see e.g. [19].

**Corollary 3.1.** *Assume that  $|\delta_n| \leq \delta < \frac{1}{2}$  for all  $n \in \mathbb{Z}$ . Then  $\{n + \delta_n\}$  is a complete interpolating sequence for  $V_c^2$ .*

Note that in the Paley–Wiener space  $PW$  the corresponding result holds with  $\delta < 1/4$ . The constant  $1/2$  is sharp, as the set  $\{n + 1/2 : n \in \mathbb{Z}\}$  is not a sampling sequence for  $V_a^2$  with  $a > 0$  [13].

#### 4. PROOF OF THEOREM 1.3: NECESSITY

For the proof of necessity we need to reverse the argument of the sufficiency part. Assume that  $\Lambda$  is a complete interpolating sequence for  $V_c^2$ . This means that the operator  $J$  given by (4) is an isomorphism of  $\ell^2(\mathbb{Z})$  onto itself.

Since a complete interpolating sequence for  $V_c^2$  is always separated, the operators  $K_+$  and  $K_-$  defined in Claim 2 are compact, whence  $U$  is a compact perturbation of the isomorphism  $J$ .

It follows that the kernel of  $U$  has finite dimension and that the range of  $U$  has finite codimension:

$$(9) \quad \dim \text{Ker } U < \infty \quad \text{and} \quad \text{codim Range } U < \infty.$$

In the language of frame theory [10, Sec. 8.7] one speaks of the deficit and the excess of the set of reproducing kernels. The condition  $\text{Ker } U \neq \{0\}$  means that the reproducing kernels associated to the sampling points  $\lambda_n, n \in \mathbb{Z}$ , are not complete, but span a proper subspace (the *deficit*). The finite codimension of the Range of  $U$  means that there are linear dependencies of the reproducing kernels, in other words, too many functions for a Riesz basis (the *excess*). The finiteness condition of (9) implies that we can construct a Riesz basis of reproducing kernels by adding and/or removing finitely many points.

In our context we will use an important property of the Fock space  $\mathcal{F}$ . Assume that  $F \in \mathcal{F}$  satisfies  $F(w_1) = 0$  and  $w_2 \in \mathbb{C}, w_2 \neq w_1$ . Then the function  $\tilde{F}(z) = \frac{z-w_2}{z-w_1}F(z)$  is again in  $\mathcal{F}$ . This property is called the *division property* of  $\mathcal{F}$ .

**Step 1.** To start with the proof of necessity, assume that (9) holds. Recall that  $V_+^2 = \{f \in V_c^2 : f = f_+\} \cong \ell^2(\mathbb{N})$ , and consider the restriction  $U_+ = U|_{V_+^2}$  of  $U$  to the subspace  $V_+^2$ , given by  $U_+(c_n)_{n \geq 1} = (f_+(\lambda_m))_{m \geq 1}$ . From the properties of  $U$  we conclude that the kernel  $\mathcal{N} = \text{Ker } U_+$  in  $V_+^2$  also is finite-dimensional and that  $\text{Range } U_+$  is a closed subspace of  $\ell^2(\mathbb{N})$  of finite codimension. Put  $w_m = e^{2e\lambda_m}$ , as before. If  $U_+$  is an isomorphism between  $V_+^2$  and  $\ell^2(\mathbb{N})$ , then, by (5),  $\{w_m\}_{m \geq 1}$  is a complete interpolating sequence for  $\mathcal{F}$ . If not, we distinguish two cases.

**Case 1.** Assume that  $U_+$  has a nontrivial kernel  $\mathcal{N}$ . Then  $U_+|_{V_+^2 \ominus \mathcal{N}}$  is an isomorphism on its image. Thus,

$$\sum_{m \geq 1} |f_+(\lambda_m)|^2 \asymp \|f_+\|_{V_+^2}^2, \quad f_+ \in V_+^2 \ominus \mathcal{N}.$$

Note that  $f \in \mathcal{N}$  (i.e.  $f_+(\lambda_m) = 0$ ,  $m \geq 1$ ) if and only if  $F_+(w_m) = 0$ ,  $m \geq 1$ . By (5), we have

$$\sum_{m \geq 1} (1 + |w_m|^2) e^{-2\varphi(w_m)} |F_+(w_m)|^2 \asymp \|F_+\|_{\mathcal{F}}^2, \quad F_+ \in \mathcal{F} \ominus \mathcal{M},$$

where  $\mathcal{M}$  is the finite-dimensional subspace in  $\mathcal{F}$  which consists of all functions vanishing on  $\{w_m\}_{m \geq 1}$ .

We will need the following simple lemma.

**Lemma 4.1.** *Let  $\mathcal{M} = \{F \in \mathcal{F} : F(w_m) = 0, m \geq 1\}$  and  $M = \dim \mathcal{M} < \infty$ . Then there exists a set of points  $\{\mu_k\}_{1 \leq k \leq M}$  which is a uniqueness set for  $\mathcal{M}$ , while  $\{\mu_k\}_{1 \leq k \leq M-1}$  is a non-uniqueness set for  $\mathcal{M}$ .*

*Proof.* We argue by induction on  $M$ . The base  $M = 1$  (i.e.  $\mathcal{M} = \text{Lin}\{F_1\}$ ) is trivial since one can take as  $\mu_1$  any point such that  $F_1(\mu_1) \neq 0$ . Assume that we can prove the statement for  $M - 1$ . Take any point  $\mu_1 \in \mathbb{C}$  such that there is a function  $F_0 \in \mathcal{M}$  with  $F_0(\mu_1) \neq 0$  and consider  $\mathcal{M}_1 = \{F \in \mathcal{M} : F(\mu_1) = 0\}$ . It is obvious that  $\dim \mathcal{M} = M - 1$ . By the induction hypothesis there exist points  $\mu_2, \dots, \mu_M$  which form a uniqueness set for  $\mathcal{M}_1$ , while  $\{\mu_2, \dots, \mu_{M-1}\}$  is a non-uniqueness set for  $\mathcal{M}_1$ . Then the points  $\{\mu_k\}_{1 \leq k \leq M}$  have the required property.  $\square$

Let  $\{\mu_k\}_{1 \leq k \leq M}$  be the set constructed in the lemma. Since  $\mathcal{M}$  is finite-dimensional, it follows that

$$\sum_{k=1}^M (1 + |\mu_k|^2) e^{-2\varphi(\mu_k)} |F_+(\mu_k)|^2 \asymp \|F_+\|_{\mathcal{F}}^2, \quad F_+ \in \mathcal{M}.$$

Let  $\tilde{W} = \{w_m\}_{m \geq 1} \cup \{\mu_k\}_{1 \leq k \leq M}$  the enlarged set. By construction, the sampling operator  $F_+ \mapsto (F_+(\tilde{w}_k))_{w_k \in \tilde{W}}$  is one-to-one and it has closed range of finite codimension in  $\ell^2(\tilde{W})$ . Consequently, by [10, Thm. 8.29, (c)],  $\tilde{W}$  is a sampling set for  $\mathcal{F}$ .

By Lemma 4.1 the set  $\tilde{W} \setminus \{\mu_M\}$  is a non-uniqueness set for  $\mathcal{F}$  and therefore there exists a function  $H \in \mathcal{F}$ , such that  $H(w_m) = H(\mu_k) = 0$  for  $m \in \mathbb{N}$  and  $k = 1, \dots, M - 1$ , but  $H(\mu_M) \neq 0$ . We now use the division property of  $\mathcal{F}$  and show that every set  $\tilde{W} \setminus \{v\}$ , where  $v \in \tilde{W} \setminus \{\mu_M\}$ , is also a non-uniqueness set in  $\mathcal{F}$ . Consider  $\tilde{H}(w) = \frac{w - \mu_M}{w - v} H(w)$ . Then  $\tilde{H}$  is non-zero in  $\mathcal{F}$ , and  $\tilde{H}$  vanishes on  $\tilde{W} \setminus \{v\}$ , but  $\tilde{H}$  cannot vanish at  $v$ , because  $\tilde{W}$  is sampling on  $\mathcal{F}_+$ .

It follows that the frame of reproducing kernels  $\{k_w\}_{w \in \tilde{W}}$  fails to be complete after removal of *any* of its members. We conclude that the normalized reproducing kernels corresponding to the sampling set  $\tilde{W}$  form an *exact frame*, i.e., a frame which fails to remain a frame after removal of any of its vectors. By well-known results (see, e.g., [10, 19]) the normalized reproducing kernels at  $\tilde{W}$  form a Riesz basis in  $\mathcal{F}$ , and, equivalently,  $\tilde{W}$  is a complete interpolating sequence for  $\mathcal{F}$ .

**Case 2.** Assume now that  $\text{Ker } U_+ = 0$ , whence  $U_+$  is an isomorphism onto its image and so  $\{w_m\}_{m \geq 1}$  is already sampling for  $\mathcal{F}$ . Let  $\text{codim Range } U_+ = M$ . This means that



the frame of reproducing kernels  $\{k_{w_m}\}_{m \geq 1}$  has finite excess, and therefore it becomes a Riesz basis after the removal of finitely many elements, see [11, Thm. 2.4] and [10].

Repeating the argument for the negative part of  $V_c^2$ , we conclude that the set  $\{w_m^{-1}\}_{m \leq -1}$  either can be enlarged to a complete interpolating set for  $\mathcal{F}$  or becomes a complete interpolating set after the removal of a finite set of points.

**Step 2.** It remains to show that after an appropriate enumeration the sequence  $\{\lambda_m\}$  will satisfy conditions (a) and (b) of Theorem 1.3. We note that both these conditions and also the property of being a complete interpolating sequence for  $\mathcal{F}$  are stable with respect to moving a finite number of points (without gluing them). Formally, we remove a finite subset  $F_1 \subseteq \Lambda$  and add a finite set  $F_2$ , such that  $\text{card } F_1 = \text{card } F_2$  and  $F_2 \cap \Lambda = \emptyset$ . After relabeling, the new set  $\Lambda' = (\Lambda \setminus F_1) \cup F_2$  satisfies conditions (a) and (b), if and only if  $\Lambda$  satisfies conditions (a) and (b).

In Step 1 we have shown that either (i) there exists a finite set  $F_+ = \{\mu_j\}_{j=1, \dots, M} \subset \mathbb{R}$  such that  $\tilde{W}_+ = \{w_m\}_{m \geq 1} \cup \{e^{2c\mu_j}\}_{j=1, \dots, M}$  is a complete interpolating set for  $\mathcal{F}$  or (ii) we can remove a finite set  $J_+ \subseteq \{w_m\}_{m \geq 1}$ , such that  $\tilde{W}_+ = \{w_m\}_{m \geq 1} \setminus J_+$  is a complete interpolating set for  $\mathcal{F}$ . Using the original points  $\lambda_m$  instead of  $e^{2c\lambda_m}$  and calling the removed points also  $F_+$ , the relabeled sequence  $\{\lambda_m\}_{m \geq 0} \cup F_+$  or  $\{\lambda_m\}_{m \geq 0} \setminus F_+$  then satisfies a one-sided version of (a) and (b) by Theorem 2.1.

We apply the same procedure to the sequence  $\{w_m^{-1}\}_{m < 0}$  and after adding a finite set  $F_- = \{\nu_j\}$ , we obtain a complete interpolating sequence  $\tilde{W}_- = \{w_m^{-1}\}_{m < 0} \cup \{e^{-2c\nu_j}\}$  or  $\tilde{W}_- = \{w_m^{-1}\}_{m < 0} \setminus J_-$ . Again a relabeling the sequence  $\{\lambda_m\}_{m < 0} \cup F_-$  or  $\{\lambda_m\}_{m < 0} \setminus F_-$  satisfies conditions (a) and (b).

By taking the union of both sequences and returning to  $\lambda_m, \mu_j, \nu_j$  (instead of  $w_m$ ), we see that there exist finite sets  $F_1$  and  $F_2$ , satisfying  $F_1 \subseteq \Lambda$  and  $F_2 \cap \Lambda = \emptyset$ , such that  $\Lambda' = (\Lambda \setminus F_1) \cup F_2$  can be enumerated to satisfy (a) and (b). It is possible that  $F_1 = \emptyset$  (we add points to both  $\{w_m\}_{m \geq 0}$  and to  $\{w_m\}_{m < 0}$ ) or  $F_2 = \emptyset$  (we remove points from both sets).

Hence, by sufficiency part of Theorem 1.3,  $\Lambda'$  is a complete interpolating sequence for  $V_c^2$ . Note that  $\Lambda = (\Lambda' \setminus F_2) \cup F_1$ . Let  $\text{card } F_1 = \text{card } F_2 + \ell$ . If  $\ell > 0$ , we have moved  $\text{card } F_2$  points and added  $\ell$  additional points. Then  $\Lambda$  fails to be a complete interpolating sequence for  $V_c^2$  in contradiction to the assumption. If  $\ell < 0$ , we have moved  $\text{card } F_1$  points and removed  $|\ell|$  additional points. Again  $\Lambda$  fails to be a complete interpolating sequence.

We conclude that  $\text{card } F_1 = \text{card } F_2$  and  $\Lambda$  is obtained from  $\Lambda'$  by moving finitely many points. As observed above, since  $\Lambda'$  satisfies conditions (a) and (b), so must the original sequence  $\Lambda$ , as was to be proved.  $\square$

## 5. PROOF OF THE DENSITY RESULTS AND REMARKS

Finally we indicate how Theorems 1.1 and 1.2 follow from the characterization of complete interpolating sequences in  $V_c^2$ . We use a simple ‘‘combinatorial’’ argument from [17] in order to show that any separated sequence  $\Lambda$  with  $D^-(\Lambda) > 1$  contains a subsequence  $\Lambda' = \{m + \delta_m\}_{m \in \mathbb{Z}}$  with  $\delta_m$  satisfying Avdonin-type conditions (a) and (b) of Theorem 1.3. Analogously, any separated  $\Lambda$  with  $D^+(\Lambda) < 1$  can be enhanced to such a set  $\Lambda'$ . We refer to [4] for the precise details, as they are almost identical.

The necessary density conditions for sampling and interpolation in a shift-invariant space are well-known, see, e.g., [1]. Let us show that they can also be deduced from the results for  $\mathcal{F}$  in [4].

Let  $\Lambda$  be an interpolating set for  $V_c^2$ . Then the mapping  $f \mapsto \{f(\lambda_m)\}_{m \geq 1}$  acts from  $V_c^2$  onto  $\ell^2(\mathbb{N})$ . By Claim 2 of the proof of sufficiency part in Theorem 1.3 the map  $f \mapsto \{f_-(\lambda_m)\}_{m \geq 1}$  is a compact operator from  $V_c^2$  to  $\ell^2(\mathbb{N})$ , whence the operator  $f \mapsto \{f_+(\lambda_m)\}_{m \geq 1}$  has the closed range of finite codimension. Then it is easy to see that for some  $m_0 \geq 1$  the map  $f \mapsto \{f_+(\lambda_m)\}_{m \geq m_0}$  will be onto. Hence, the set  $\{\lambda_m\}_{m \geq m_0}$  will be an interpolating set for  $V_+^2$  and so  $\{w_m\}_{m \geq m_0}$  will be interpolating for  $\mathcal{F}$ . We conclude that  $D^+(\Lambda) \leq 1$ .

Now assume that  $\Lambda$  is a sampling set for  $V_c^2$  and, in particular, for  $V_+^2$ . Since  $\Lambda$  is a finite union of separated sets, the sequence  $\{f_+(\lambda_m)\}_{m \leq -1}$  decays very fast and it is easy to show that the set  $\{\lambda_m\}_{m \geq m_0}$  will be sampling for  $V_+^2$ .

**Further remarks.** 1. Let  $V_c^p = \{f(z) = \sum_{n \in \mathbb{Z}} c_n e^{-c(z-n)^2} : (c_n) \in \ell^p(\mathbb{Z})\}$  be the subspace generated by shifts of the Gaussian with  $\ell^p$  coefficients. This is a closed subspace of  $L^p(\mathbb{R})$ . Again, a sequence  $\Lambda = \{\lambda_n : n \in \mathbb{Z}\}$  is called a sampling set for  $V_c^p$ , if

$$\sum_{n \in \mathbb{Z}} |f(\lambda_n)|^p \asymp \|f\|_{V_c^p}^p \asymp \|c\|_p^p, \quad f \in V_c^p,$$

and *interpolating* for  $V_c^p$  if any  $(a_n) \in \ell^p(\mathbb{Z})$  can be interpolated by a function  $f \in V_c^p$ . Based on the general theory of sampling in shift-invariant spaces with a nice generator, we can assert that *a set  $\Lambda \subseteq \mathbb{R}$  is sampling for  $V_c^2$ , if and only if  $\Lambda$  is sampling for some  $V_c^{p_0}$ , if and only if  $\Lambda$  is sampling for  $V_c^p$  for all  $p, 1 \leq p \leq \infty$*  [8, Thm. 3.1]. A similar statement holds for the interpolation property. Theorem 1.3 therefore provides a characterization for complete interpolating sequences for all spaces  $V_c^p$ .

2. *Sign retrieval.* The sign retrieval problem asks whether a real-valued function  $f$  in some function space is uniquely determined by its unsigned values  $|f(\lambda)|$  on some set  $\Lambda$ . For  $V_a^2$  with a real-valued Gaussian generator, Theorem 1.3 implies the following result on sign retrieval.

**Corollary 5.1.** *Let  $\Lambda = \{n/2 + \delta_n : n \in \mathbb{Z}\}$  be uniformly separated such that  $\{\delta_n\}$  satisfies the Avdonin-type conditions (a) and (b) with  $\delta < 1/4$ . Then every  $f \in V_a^2$  is uniquely determined by its unsigned values  $\{|f(n/2 + \delta_n)| : n \in \mathbb{Z}\}$ .*

The proof is the same as in [7], we simply replace sampling sets of density  $D^-(\Lambda) > 2$  by a complete interpolating sequence for a dilated version of  $V_a^2$ . Thus sign retrieval is possible even at the critical density.

3. It would be interesting to obtain similar results for complete interpolating sequences in shift-invariant spaces with other generators. Even a sharp version of Corollary 3.1 would have important consequences for Gabor frames.

## REFERENCES

- [1] A. Aldroubi, K. Gröchenig, Nonuniform sampling and reconstruction in shift-invariant spaces. *SIAM Rev.* **43** (2001), 4, 585–620.
- [2] S.A. Avdonin, On the question of Riesz bases of exponential functions in  $L^2$ , *Vestnik Leningrad. Univ.* No. **13** Mat. Meh. Astronom. Vyp. 3 (1974), 5–12 (in Russian).

- [3] Yu. Belov, T. Mengestie, K. Seip, Discrete Hilbert transforms on sparse sequences, *Proc. London Math. Soc.* **103** (2011), 73–105.
- [4] A. Baranov, A. Dumont, A. Hartmann, K. Kellay, Sampling, interpolation and Riesz bases in small Fock spaces, *J. Math. Pures Appl.* **103** (2015), 6, 1358–1389.
- [5] A. Borichev, Yu. Lyubarskii, Riesz bases of reproducing kernels in Fock type spaces, *J. Inst. Math. Jussieu* **9** (2010), 449–461.
- [6] A. Beurling, *The Collected Works of Arne Beurling. Vol. 2.* Contemporary Mathematicians. Birkhäuser Boston Inc., Boston, MA, 1989. Harmonic analysis, Edited by L. Carleson, P. Malliavin, J. Neuberger and J. Wermer.
- [7] K. Gröchenig, Phase-retrieval in shift-invariant spaces with Gaussian generator. *J. Fourier Anal. Appl.* **26** (2020), 3, Paper No. 52.
- [8] K. Gröchenig, J.L. Romero, J. Stöckler, Sampling theorems for shift-invariant spaces, Gabor frames, and totally positive functions, *Invent. Math.* **211** (2018), 3, 1119–1148.
- [9] K. Gröchenig, J.L. Romero, J. Stöckler, Sharp results on sampling with derivatives in shift-invariant spaces and multi-window Gabor frames, *Constr. Approx.* **51** (2020), 1, 1–25.
- [10] C. Heil, *A Basis Theory Primer, Expanded Edition*, Birkhäuser, Boston, 2011.
- [11] J. R. Holub, Pre-frame operators, Besselian frames, and near-Riesz bases in Hilbert spaces. *Proc. Amer. Math. Soc.*, **122** (1994), 1, 779–785.
- [12] S. Hruscev, N. Nikolski, B. Pavlov, Unconditional bases of exponentials and of reproducing kernels. Complex Analysis and Spectral Theory, 214–335, Lecture Notes in Math. **864**, Springer-Verlag, Berlin, 1981.
- [13] A. Janssen, The Zak transform and sampling theorems for wavelet subspaces. *IEEE Trans. Signal Process.* **41** (1993) 3360–3365.
- [14] J.-P. Kahane, Pseudo-périodicité et séries de Fourier lacunaires. *Ann. Sci. École Norm. Sup.* **79** (1962), 93–150.
- [15] H. J. Landau, Necessary density conditions for sampling and interpolation of certain entire functions. *Acta Math.* **117** (1967), 37–52.
- [16] J. Ortega-Cerdà, K. Seip. Fourier frames. *Ann. of Math. (2)* **155** (2002), 3, 789–806.
- [17] K. Seip, A simple construction of exponential bases in  $L^2$  of the union of several intervals, *Proc. Edinburgh Math. Soc.* **38** (1995), 1, 171–177.
- [18] K. Seip. *Interpolation and Sampling in Spaces of Analytic Functions*, University Lecture Series, Vol. 33, American Mathematical Society, Providence, RI, 2004.
- [19] R.M. Young, *An Introduction to Nonharmonic Fourier Series*, Academic Press, New York, 1980.

DEPARTMENT OF MATHEMATICS AND MECHANICS, ST. PETERSBURG STATE UNIVERSITY, ST. PETERSBURG, RUSSIA

*Email address:* anton.d.baranov@gmail.com

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, ST. PETERSBURG STATE UNIVERSITY, ST. PETERSBURG, RUSSIA

*Email address:* j\_b\_juri\_belov@mail.ru

FACULTY OF MATHEMATICS, UNIVERSITY OF VIENNA, OSKAR-MORGENSTERN-PLATZ 1, A-1090 VIENNA, AUSTRIA

*Email address:* karlheinz.groechenig@univie.ac.at