

# GURARIĬ OPERATORS ARE GENERIC

TARAS BANAKH<sup>1,2</sup> AND JOANNA GARBULIŃSKA-WĘGRZYN<sup>2</sup>

ABSTRACT. Answering a question of Garbulińska-Węgrzyn and Kubiś, we prove that Gurariĭ operators form a dense  $G_\delta$ -set in the space  $\mathcal{B}(\mathbb{G})$  of all nonexpansive operators on the Gurariĭ space  $\mathbb{G}$ , endowed with the strong operator topology. This implies that universal operators on  $\mathbb{G}$  form a residual set in  $\mathcal{B}(\mathbb{G})$ .

## 1. INTRODUCTION

All *Banach spaces* considered in this paper are real and separable, all *operators* are linear bounded operators between separable Banach spaces. An *isometric embedding* (resp.  $\varepsilon$ -*isometric embedding* for some  $\varepsilon > 0$ ) of Banach spaces is any injective operator  $T : X \rightarrow Y$  such that  $\|T(x)\| = \|x\|$  (resp.  $(1 - \varepsilon)\|x\| < \|T(x)\| < (1 + \varepsilon)\|x\|$ ) for any nonzero  $x \in X$ . An operator  $T : X \rightarrow Y$  is called *nonexpansive* if  $\|T\| \leq 1$ .

A Banach space  $X$  is called *Gurariĭ* if for any finite-dimensional Banach spaces  $A \subseteq B$  and any  $\varepsilon > 0$ , any isometric embedding  $i : B \rightarrow X$  extends to an  $\varepsilon$ -isometric embedding  $\bar{i} : A \rightarrow X$ . The first example of a Gurariĭ space  $\mathbb{G}$  was constructed by Gurariĭ in [4]. In [9] Lusky proved that any two Gurariĭ spaces are isometrically isomorphic, so up to an isometry there exists a unique Gurariĭ space  $\mathbb{G}$ . A simple proof of the uniqueness of the Gurariĭ space was found by Kubiś and Solecki [8]. In [7], Kubiś suggested an elementary game-theoretic construction of the Gurariĭ space. It is known [3] that the Gurariĭ space is *universal* in the sense that it contains an isometric copy of any separable Banach space.

An operator  $U : V \rightarrow W$  between Banach spaces is defined to be *universal* if for every operator  $T : X \rightarrow Y$  with  $\|T\| \leq \|U\|$ , there exist linear isometric embeddings  $i : X \rightarrow V$ ,  $j : Y \rightarrow W$  such that the diagram

$$\begin{array}{ccc} V & \xrightarrow{U} & W \\ \uparrow i & & \uparrow j \\ X & \xrightarrow{T} & Y \end{array}$$

is commutative, that is,  $U \circ i = j \circ T$ .

In [1], [2] Garbulińska-Węgrzyn and Kubiś constructed a universal operator  $\Omega : \mathbb{G} \rightarrow \mathbb{G}$  using the technique of Fraïssé limits. More precisely, they defined the notion of a Gurariĭ operator (which is an operator counterpart of the notion of a Gurariĭ space), constructed a Gurariĭ operator (as the Fraïssé limit in a suitable category) and proved that every Gurariĭ operator is universal.

An operator  $G : X \rightarrow Y$  between Banach spaces is called *Gurariĭ* if  $G$  is nonexpansive and for any  $\varepsilon > 0$ , any nonexpansive operator  $T : A \rightarrow B$  between finite-dimensional Banach spaces, any Banach subspaces  $A_0 \subseteq A$ ,  $B_0 \subseteq B$  with  $T[A_0] \subseteq B_0$ , and any isometric embeddings  $i_0 : A_0 \subseteq X$ ,  $j_0 : B_0 \subseteq Y$  with  $G \circ i_0 = j_0 \circ T|_{A_0}$ , there exist  $\varepsilon$ -isometric embeddings  $i : A \rightarrow X$  and  $j : B \rightarrow Y$  such that  $i|_{A_0} = i_0$ ,  $j|_{B_0} = j_0$  and  $G \circ i = j \circ T$ .

2020 *Mathematics Subject Classification.* 47A05; 47A65; 46B04; 46B28; 54B52; 54H05; 54H15.

*Key words and phrases.* Gurariĭ space, universal operator.

An example of a Gurariĭ operator  $\Omega : \mathbb{G} \rightarrow \mathbb{G}$  was constructed in [1]. By [1, Theorem 3.5], any Gurariĭ operator  $G : X \rightarrow Y$  is isometric to the Gurariĭ operator  $\Omega : \mathbb{G} \rightarrow \mathbb{G}$  in the sense that there exist bijective isometries  $i : X \rightarrow \mathbb{G}$  and  $j : Y \rightarrow \mathbb{G}$  such that  $\Omega \circ i = j \circ G$ . Therefore, a Gurariĭ operator is unique up to an isometry (like the Gurariĭ space). By [1, Theorem 3.3], every Gurariĭ operator is universal. Therefore, the set  $\mathcal{G}(\mathbb{G})$  of Gurariĭ operators from a Gurariĭ space  $\mathbb{G}$  to itself is a subset of the set  $\mathcal{U}(\mathbb{G})$  of nonexpansive universal operators from  $\mathbb{G}$  to  $\mathbb{G}$ . In its turn,  $\mathcal{U}(\mathbb{G})$  is a subset of the space  $\mathcal{B}(\mathbb{G})$  of all nonexpansive operators from  $\mathbb{G}$  to  $\mathbb{G}$ . The space  $\mathcal{B}(\mathbb{G})$  is endowed with the strong operator topology. Let us recall that the *strong operator topology* on  $\mathcal{B}(\mathbb{G})$  is the smallest topology such that for every  $x \in \mathbb{G}$  the map  $\delta_x : \mathcal{B}(\mathbb{G}) \rightarrow \mathbb{G}$ ,  $\delta_x : T \mapsto T(x)$ , is continuous.

In [2, Question 1] Garbulińska-Węgrzyn and Kubiś asked whether the set  $\mathcal{G}(\mathbb{G})$  of Gurariĭ operators is residual in the space  $\mathcal{B}(\mathbb{G})$ . Let us recall that a subset  $R$  of a topological space  $X$  is *residual* if it contains a dense  $G_\delta$ -subset of  $X$ .

In this paper we answer Question 1 of [2] affirmatively proving the following main result.

**Theorem 1.** *The set  $\mathcal{G}(\mathbb{G})$  of Gurariĭ operators is dense  $G_\delta$  in the space  $\mathcal{B}(\mathbb{G})$  of all nonexpansive operators on  $\mathbb{G}$ .*

Since  $\mathcal{G}(\mathbb{G}) \subseteq \mathcal{U}(\mathbb{G})$ , Theorem 1 implies the following corollary.

**Corollary 1.** *The set  $\mathcal{U}(\mathbb{G})$  of universal nonexpansive operators on  $\mathbb{G}$  is residual in the space  $\mathcal{B}(\mathbb{G})$  of all nonexpansive operators on  $\mathbb{G}$ .*

**Remark 1.** Observe that the space  $\mathcal{B}(\mathbb{G})$  is a topological semigroup with respect to the operation of composition of operators. The group  $\text{Iso}(\mathbb{G})$  of linear isometries of  $\mathbb{G}$  is Polish [6, 9.3] and hence is a  $G_\delta$ -subgroup of the topological semigroup  $\mathcal{B}(\mathbb{G})$ . By [1, 3.5], any Gurariĭ operators are isometric, which implies the set  $\mathcal{G}(\mathbb{G})$  coincides with the orbit  $\{I \circ \Omega \circ J : I, J \in \text{Iso}(\mathbb{G})\}$  of any Gurariĭ operator  $\Omega : \mathbb{G} \rightarrow \mathbb{G}$  under the two-sided action of the Polish group  $\text{Iso}(\mathbb{G}) \times \text{Iso}(\mathbb{G})$  on  $\mathcal{B}(\mathbb{G})$ , and this orbit is a unique dense  $G_\delta$ -orbit of this action. Theorem 1 also shows that Gurariĭ operators are “generic” elements of the space  $\mathcal{B}(\mathbb{G})$  in the Baire category sense.

## 2. PRELIMINARIES

By  $\mathbb{R}_+$  we denote the half-line  $[0, \infty)$ . Each natural number  $n$  is identified with the set  $\{0, \dots, n-1\}$ . For a function  $f : X \rightarrow Y$  between sets and a subset  $A \subseteq X$  we denote by  $f[A]$  the image  $\{f(x) : x \in A\}$  of the set  $A$  under the map  $f$ .

For a Banach space  $X$ , point  $x \in X$  and positive real number  $r$ , let

$$B_X(x, r) = \{y \in X : \|x - y\| < r\}$$

be the open ball of radius  $r$  with center  $x$  in the Banach space  $X$ . If  $x = 0$ , then we shall write  $B_X(r)$  instead of  $B_X(0, r)$ .

For an operator  $T : X \rightarrow Y$  between Banach spaces, let

$$\begin{aligned} \|T\| &= \inf\{r \in \mathbb{R}_+ : T[B_X(1)] \subseteq B_Y(r)\} \quad \text{and} \\ \|T^\leftarrow\| &= \inf(\{r \in \mathbb{R}_+ : T[X] \cap B_Y(1) \subseteq T[B_X(r)]\} \cup \{\infty\}). \end{aligned}$$

The (finite or infinite) number  $\|T^\leftarrow\|$  is called the *openness norm* of  $T$ . Observe that an operator  $T : X \rightarrow Y$  between Banach spaces has finite openness norm  $\|T^\leftarrow\|$  if and only if the operator  $T : X \rightarrow T[X]$  is open if and only if  $T[X]$  is a closed subspace of  $Y$ . In particular, for any operator  $T : X \rightarrow Y$  from a finite-dimensional Banach space  $X$ , the openness norm  $\|T^\leftarrow\|$  is finite. For a bijective operator  $T : X \rightarrow Y$  the openness norm  $\|T^\leftarrow\|$  is equal to the norm  $\|T^{-1}\|$  of the inverse operator  $T^{-1} : Y \rightarrow X$ .

For a non-negative integer  $n$  and a real number  $p \in [1, \infty)$ , let  $\ell_p^n$  be the Banach space  $\mathbb{R}^n$  endowed with the  $\ell_p$ -norm

$$\|x\| = \left( \sum_{i \in n} |x_i|^p \right)^{\frac{1}{p}} \quad \text{for any } x = (x_i)_{i \in n} \in \mathbb{R}^n.$$

A Banach space is *Hilbert* if its norm is generated by an inner product  $\langle \cdot, \cdot \rangle$  in the sense that  $\|x\| = \sqrt{\langle x, x \rangle}$  for all  $x \in X$ . In particular, for every  $n \in \mathbb{N}$  the Banach space  $\ell_2^n$  is Hilbert: its norm is generated by the inner product

$$\langle x, y \rangle = \sum_{i \in n} x_i y_i.$$

It is well-known that the inner product  $\langle \cdot, \cdot \rangle$  of a Hilbert space can be uniquely recovered from the norm by the formula

$$\langle x, y \rangle = \frac{1}{4}(\|x + y\|^2 - \|x - y\|^2).$$

For a subset  $A \subseteq X$  of a Hilbert space  $X$  we denote by

$$A^\perp = \{x \in X : \forall a \in A \ \langle x, a \rangle = 0\}$$

the *orthogonal complement* of  $A$  in  $X$ .

For an operator  $T : X \rightarrow Y$  between Banach spaces,  $\ker(T)$  stands for the *kernel*  $T^{-1}(0)$  of  $T$ .

**Lemma 1.** *Let  $T : X \rightarrow Y$  be an operator from a Hilbert space  $X$  to a Banach space  $Y$  and  $K = \ker(T)$  be the kernel of  $T$ . Then  $T[B_X(1)] = T[B_X(1) \cap K^\perp]$ .*

*Proof.* The inclusion  $T[B_X(1) \cap K^\perp] \subseteq T[B_X(1)]$  is trivial. To prove that  $T[B_X(1)] \subseteq T[B_X(1) \cap K^\perp]$ , fix any element  $y \in T[B_X(1)]$  and find  $x \in B_X(1)$  with  $T(x) = y$ . Let  $x'$  be the orthogonal projection of  $x$  onto the subspace  $K$ . Then  $x - x' \in K^\perp$  and  $T(x - x') = T(x) - T(x') = T(x) - 0 = y$ . Since  $x = x' + (x - x')$  and  $\langle x', x - x' \rangle = 0$  in the Hilbert space  $X$ , we have  $\|x - x'\|^2 = \|x\|^2 - \|x'\|^2 \leq \|x\|^2 < 1$  by the Pitagoras Theorem. Therefore,  $y = T(x - x') \in T[B_X(1) \cap K^\perp]$ .  $\square$

**Lemma 2.** *Let  $X$  be a Hilbert space of finite dimension  $n > 0$ , and  $T : X \rightarrow Y$  be an operator to a Banach space  $Y$ . Then*

$$\|T\| \leq \sqrt{n} \max_{1 \leq i \leq n} \|T(e_i)\|$$

for any orthonormal basis  $e_1, \dots, e_n$  in  $X$ .

*Proof.* Consider the isomorphism  $I : \ell_1^n \rightarrow X$  defined by  $I : (x_1, \dots, x_n) \mapsto \sum_{i=1}^n x_i e_i$ . It is easy to see that  $\|I\| = 1$ . Using Cauchy–Bunyakovsky–Schwartz inequality, one can show that  $\|I^{-1}\| = \sqrt{n}$ . It follows that

$$\|T\| = \|T \circ I \circ I^{-1}\| \leq \|T \circ I\| \cdot \|I^{-1}\| = \max_{1 \leq i \leq n} \|T(e_i)\| \cdot \sqrt{n}.$$

$\square$

The *Banach-Mazur distance* between two isomorphic Banach spaces  $X, Y$  is defined as

$$d_{BM}(X, Y) := \inf\{\|I\| \cdot \|I^{-1}\| : I \text{ is an isomorphism between } X \text{ and } Y\}.$$

The following proposition is a classical result of John [5] (see also [10, Theorem 2.3.2]).

**Proposition 1 (John).** *For any  $n$ -dimensional Banach space  $X$ , we have  $d_{BM}(X, \ell_2^n) \leq \sqrt{n}$ .*

**Lemma 3.** *Let  $\varepsilon, \delta$  be positive real numbers. Let  $i, j : X \rightarrow Y$  be two operators between Banach spaces. If  $\|i - j\| \leq \delta$  and  $i$  is an  $\varepsilon$ -isometric embedding, then  $j$  is an  $(\varepsilon + \delta)$ -isometric embedding.*

*Proof.* Given any nonzero  $x \in X$ , observe that

$$\|j(x)\| \leq \|i(x)\| + \|i(x) - j(x)\| < (1 + \varepsilon)\|x\| + \|i - j\| \cdot \|x\| \leq (1 + \varepsilon)\|x\| + \delta\|x\| = (1 + \varepsilon + \delta)\|x\|$$

and

$$\|j(x)\| \geq \|i(x)\| - \|i(x) - j(x)\| > (1 - \varepsilon)\|x\| - \delta\|x\| = (1 - \varepsilon - \delta)\|x\|,$$

which means that  $j$  is an  $(\varepsilon + \delta)$ -isometric embedding.  $\square$

**Lemma 4.** *If  $G : \mathbb{G} \rightarrow \mathbb{G}$  is a Gurariĭ operator, then  $G[B_{\mathbb{G}}(1)] = B_{\mathbb{G}}(1)$  and hence  $\|G\| = 1 = \|G^{\leftarrow}\|$ .*

*Proof.* The nonexpansive property of  $G$  ensures that  $G[B_{\mathbb{G}}(1)] \subseteq B_{\mathbb{G}}(1)$ . To show that  $B_{\mathbb{G}}(1) \subseteq G[B_{\mathbb{G}}(1)]$ , fix any element  $y \in B_{\mathbb{G}}(1)$ . If  $\|y\| = 0$ , then  $y = 0 = G(0) \in G[B_{\mathbb{G}}(1)]$ . So, we assume that  $\|y\| > 0$ .

Let  $X = Y = \mathbb{R}$  and  $T : X \rightarrow Y$  be the identity operator. Let  $X_0 = \{0\} \subset X$ ,  $Y_0 = Y$ ,  $i_0 : X_0 \rightarrow \{0\} \subseteq \mathbb{G}$  be the trivial isometric embedding and  $j : Y_0 \rightarrow \mathbb{G}$  be the isometric embedding defined by  $j(t) = \frac{ty}{\|y\|}$ . Since  $\|y\| < 1$ , there exists  $\varepsilon > 0$  such that  $(1 + \varepsilon)\|y\| < 1$ . Since the operator  $G$  is Gurariĭ, there exists an  $\varepsilon$ -isometric embedding  $i : X \rightarrow \mathbb{G}$  such that  $G \circ i = j \circ T$ . Then the element  $x = i(\|y\|)$  has  $G(x) = G \circ i(\|y\|) = j \circ T(\|y\|) = j(\|y\|) = y$  and  $\|x\| = \|i(\|y\|)\| < (1 + \varepsilon)\|y\| < 1$ . Therefore,  $y = G(x) \in G[B_{\mathbb{G}}(1)]$  and hence  $B_{\mathbb{G}}(1) = G[B_{\mathbb{G}}(1)]$ . The latter equality implies that  $\|G\| = 1 = \|G^{\leftarrow}\|$ .  $\square$

In the proof of Theorem 1 we shall apply the following homogeneity property of the Gurariĭ space, proved by Kubiś and Solecki in [8, 1.1].

**Lemma 5.** *Let  $X$  be a finite-dimensional linear subspace of the Gurariĭ space  $\mathbb{G}$ . For every  $\varepsilon > 0$  and every  $\varepsilon$ -isometric embedding  $f : X \rightarrow \mathbb{G}$  there exists a bijective isometry  $I : \mathbb{G} \rightarrow \mathbb{G}$  such that  $\|f - I|_X\| < \varepsilon$ .*

### 3. CHARACTERIZING GURARIĬ OPERATORS

In this section we shall present several characterizations of Gurariĭ operators. The following characterization was proved in [1].

**Theorem 2.** *An operator  $G : \mathbb{G} \rightarrow \mathbb{G}$  is Gurariĭ if and only if it satisfies the following condition:*

(GA) *for any  $\varepsilon > 0$ , any nonexpansive operator  $T : X \rightarrow Y$  between finite-dimensional Banach spaces, any Banach subspaces  $X_0 \subseteq X$ ,  $Y_0 \subseteq Y$  with  $T[X_0] \subseteq Y_0$ , and any isometric embeddings  $i_0 : X_0 \rightarrow \mathbb{G}$ ,  $j_0 : Y_0 \rightarrow \mathbb{G}$  with  $G \circ i_0 = j_0 \circ T|_{X_0}$ , there exist  $\varepsilon$ -isometric embeddings  $i : X \rightarrow \mathbb{G}$ ,  $j : Y \rightarrow \mathbb{G}$  such that*

$$\|i|_{X_0} - i_0\| < \varepsilon, \quad \|j|_{Y_0} - j_0\| < \varepsilon, \quad \text{and} \quad \|G \circ i - j \circ T\| < \varepsilon.$$

Next, we show that the isometric embeddings  $i_0, j_0$  in the above characterization can be replaced by  $\delta$ -isometric embeddings for a sufficiently small  $\delta$ . For non-negative real numbers  $\varepsilon, n, m, t$  let  $\delta(\varepsilon, n, m, t)$  be the largest number  $\delta$  such that

$$\delta(1 + (1 + \delta)(n + 1) \max\{1, t\sqrt{nm}\}) \leq \varepsilon.$$

**Theorem 3.** *A nonexpansive operator  $G : \mathbb{G} \rightarrow \mathbb{G}$  is Gurariĭ if and only if it satisfies the condition:*

(GB) for any  $\varepsilon \in (0, 1]$ , any nonexpansive operator  $T : X \rightarrow Y$  between finite-dimensional Banach spaces, any Banach subspaces  $X_0 \subseteq X$ ,  $Y_0 \subseteq Y$  with  $T_0[X_0] \subseteq Y_0$  where  $T_0 = T|_{X_0} : X_0 \rightarrow Y_0$ , and any  $\delta$ -isometric embeddings  $i_0 : X_0 \rightarrow \mathbb{G}$ ,  $j_0 : Y_0 \rightarrow \mathbb{G}$  with

$$\|G \circ i_0 - j_0 \circ T_0\| < \delta := \delta(\varepsilon, \dim(X_0), \dim(Y_0), \|T_0^{\leftarrow}\|)$$

there exist  $\varepsilon$ -isometric embeddings  $i : X \rightarrow \mathbb{G}$ ,  $j : Y \rightarrow \mathbb{G}$  such that

$$\|i|_{X_0} - i_0\| < \varepsilon, \quad \|j|_{Y_0} - j_0\| < \varepsilon, \quad \text{and} \quad \|G \circ i - j \circ T\| < \varepsilon.$$

*Proof.* The ‘‘if’’ part follows immediately from Theorem 2 and the trivial implication (GB) $\Rightarrow$ (GA).

To prove the ‘‘only if’’ part, we need to show that every Gurariĭ operator  $G : \mathbb{G} \rightarrow \mathbb{G}$  has the property (GB). Fix any  $\varepsilon \in (0, 1]$ , any nonexpansive operator  $T : X \rightarrow Y$  between finite-dimensional Banach spaces, any Banach subspaces  $X_0 \subseteq X$  and  $Y_0 \subseteq Y$  with  $T[X_0] \subseteq Y_0$ , and any  $\delta$ -isometric embeddings  $i_0 : X_0 \rightarrow \mathbb{G}$ ,  $j_0 : Y_0 \rightarrow \mathbb{G}$  such that  $\|G \circ i_0 - j_0 \circ T_0\| < \delta$  where  $\delta = \delta(\varepsilon, \dim(X_0), \dim(Y_0), \|T_0^{\leftarrow}\|)$  and  $T_0 = T|_{X_0} : X_0 \rightarrow Y_0$ .

Let  $n = \dim X_0$  and  $m = \dim Y_0$ . By Proposition 1, there exist isomorphisms  $I : X_0 \rightarrow \ell_2^n$  and  $J : Y_0 \rightarrow \ell_2^m$  such that  $\|I\| \cdot \|I^{-1}\| \leq \sqrt{n}$  and  $\|J\| \cdot \|J^{-1}\| \leq \sqrt{m}$ .

Fix a basis  $e_1, \dots, e_k, e_{k+1}, \dots, e_n$  in  $X_0$  and a basis  $e'_1, \dots, e'_{n-k}, e'_{n-k+1}, \dots, e'_m$  in  $Y_0$  such that:

- (1)  $I(e_1), \dots, I(e_n)$  is an orthonormal basis for the Hilbert space  $\ell_2^n$ ;
- (2)  $e_1, \dots, e_k$  is a basis for the subspace  $\ker(T_0)$ ;
- (3)  $e'_l = T(e_{l+k})$  for  $l \in \{1, \dots, n-k\}$ ;
- (4)  $J(e'_{n-k+1}), \dots, J(e'_m)$  is an orthonormal basis for the orthogonal complement of the space  $J[T[X_0]]$  in the Hilbert space  $\ell_2^m$ .

Let  $K = I[\ker(T_0)] \subseteq \ell_2^n$  and  $K^\perp$  be the orthogonal complement of  $K$  in the Hilbert space  $\ell_2^n$ . The choice of the basis  $e_1, \dots, e_n$  guarantees that  $I(e_1), \dots, I(e_k)$  and  $I(e_{k+1}), \dots, I(e_n)$  are orthonormal bases for the spaces  $K$  and  $K^\perp$ , respectively. Observe that the operator  $T_\perp = T \circ I^{-1}|_{K^\perp} : K^\perp \rightarrow T[X_0]$  is an isomorphism. By Lemma 1,

$$T_\perp[B_{K^\perp}(1)] = T \circ I^{-1}[B_{K^\perp}(1)] = T \circ I^{-1}[B_{\ell_2^n}(1)].$$

Also  $I[B_{X_0}(1)] \subseteq B_{\ell_2^n}(\|I\|)$  and hence  $B_{X_0}(r) \subseteq I^{-1}[B_{\ell_2^n}(\|I\|r)]$  for every  $r \in \mathbb{R}_+$ . Then

$$\begin{aligned} \{r \in \mathbb{R}_+ : B_{T[X_0]}(1) \subseteq T[B_{X_0}(r)]\} &\subseteq \\ \{r \in \mathbb{R}_+ : B_{T[X_0]}(1) \subseteq T \circ I^{-1}[B_{\ell_2^n}(\|I\|r)]\} &= \\ \{r \in \mathbb{R}_+ : B_{T[X_0]}(1) \subseteq T_\perp[B_{K^\perp}(\|I\|r)]\} & \end{aligned}$$

and thus

$$\begin{aligned} \|T_0^{\leftarrow}\| &= \inf\{r \in \mathbb{R}_+ : B_{T[X_0]}(1) \subseteq T[B_{X_0}(r)]\} \geq \\ &\geq \inf\{r \in \mathbb{R}_+ : B_{T[X_0]}(1) \subseteq T_\perp[B_{K^\perp}(\|I\|r)]\} = \\ &= \|I\|^{-1} \inf\{\|I\|r : B_{T[X_0]}(1) \subseteq T_\perp[B_{K^\perp}(\|I\|r)]\} = \|I\|^{-1} \|T_\perp^{-1}\|. \end{aligned}$$

Therefore,

$$\|T_\perp^{-1}\| \leq \|T_0^{\leftarrow}\| \cdot \|I\|.$$

By condition (2), for any  $l \in \{1, \dots, k\}$  we have  $e_l \in \ker(T_0)$  and hence

$$\begin{aligned} \|G(i_0(e_l))\| &= \|G \circ i_0(e_l) - j_0 \circ T_0(e_l)\| \leq \|G \circ i_0 - j_0 \circ T_0\| \cdot \|e_l\| < \\ &< \delta \cdot \|I^{-1}\| \cdot \|I(e_l)\| = \delta \cdot \|I^{-1}\|. \end{aligned}$$

Lemma 4 implies that

$$G[i_0(e_l) + B_{\mathbb{G}}(\delta\|I^{-1}\|)] = G(i_0(e_l)) + B_{\mathbb{G}}(\delta\|I^{-1}\|) \ni 0.$$

Then there exist  $y_l \in i_0(e_l) + B_{\mathbb{G}}(\delta\|I^{-1}\|)$  such that  $y_l \in \ker G$ .

Consider the operator  $i'_0 : X_0 \rightarrow \mathbb{G}$  such that  $i'_0(e_l) = y_l$  for all  $l \in \{1, \dots, k\}$  and  $i'_0(e_l) = i_0(e_l)$  for all  $l \in \{k+1, \dots, n\}$ .

**Claim 1.**  $\|i'_0 - i_0\| < \delta n$  and  $i'_0$  is a  $\delta(n+1)$ -isometric embedding.

*Proof.* By Lemma 2,

$$\begin{aligned} \|i'_0 - i_0\| &= \|(i'_0 - i_0) \circ I^{-1} \circ I\| \leq \|(i'_0 - i_0) \circ I^{-1}\| \cdot \|I\| \leq \\ &\leq \|I\| \sqrt{n} \max_{1 \leq l \leq n} \|(i'_0 - i_0) \circ I^{-1}(I(e_l))\| = \|I\| \sqrt{n} \max_{1 \leq l \leq n} \|i'_0(e_l) - i_0(e_l)\| = \\ &= \|I\| \sqrt{n} \max_{1 \leq l \leq k} \|y_l - i_0(e_l)\| < \|I\| \sqrt{n} \delta \|I^{-1}\| \leq \delta n. \end{aligned}$$

Since  $i_0$  is a  $\delta$ -isometric embedding and  $\|i_0 - i'_0\| < \delta n$ , the operator  $i'_0 : X_0 \rightarrow \mathbb{G}$  is a  $\delta(n+1)$ -isometric embedding by Lemma 3.  $\square$

Let  $j'_0 : Y_0 \rightarrow \mathbb{G}$  be a unique operator such that  $j'_0(e'_l) = G \circ i'_0(e_{l+k}) = G \circ i_0(e_{l+k})$  for all  $l \in \{1, \dots, n-k\}$  and  $j'_0(e'_l) = j_0(e'_l)$  for all  $l \in \{n-k+1, \dots, m\}$ . Observe that for every  $l \in \{1, \dots, k\}$  we have

$$j'_0 \circ T_0(e_l) = j'_0(0) = 0 = G(y_l) = G \circ i'_0(e_l),$$

and for every  $l \in \{k+1, \dots, n\}$  we have

$$j'_0 \circ T_0(e_l) = j'_0(e'_{l-k}) = G \circ i'_0(e_{l-k+k}) = G \circ i'_0(e_l).$$

Therefore,  $j'_0 \circ T_0(e_l) = G \circ i'_0(e_l)$  for every  $l \in \{1, \dots, n\}$ . Taking into account that  $e_1, \dots, e_n$  is a basis of the space  $X_0$ , we conclude that

$$(1) \quad j'_0 \circ T_0 = G \circ i'_0.$$

**Claim 2.**  $\|j'_0 - j_0\| \leq \delta(n+1)\sqrt{nm}\|T_0^{\leftarrow}\|$  and  $j'_0$  is a  $\delta(1 + (n+1)\sqrt{nm}\|T_0^{\leftarrow}\|)$ -isometric embedding.

*Proof.* Since  $j'_0 \circ T_0 = G \circ i'_0$ , we have

$$\begin{aligned} \|(j'_0 - j_0) \circ T_0\| &= \|G \circ i'_0 - j_0 \circ T_0\| \leq \|G \circ i'_0 - G \circ i_0\| + \|G \circ i_0 - j_0 \circ T_0\| \\ &< \|G\| \cdot \|i'_0 - i_0\| + \delta \leq \delta n + \delta = \delta(n+1) \end{aligned}$$

and

$$\begin{aligned} \|(j'_0 - j_0) \upharpoonright_{T[X_0]}\| &= \|(j'_0 - j_0) \circ T_{\perp} \circ T_{\perp}^{-1}\| = \\ &= \|(j'_0 - j_0) \circ T_0 \circ I^{-1} \circ T_{\perp}^{-1}\| \leq \|(j'_0 - j_0) \circ T_0\| \cdot \|I^{-1}\| \cdot \|T_{\perp}^{-1}\| \leq \\ &\leq \delta(n+1)\|I^{-1}\| \cdot \|T_0^{\leftarrow}\| \cdot \|I\| \leq \delta(n+1)\sqrt{n}\|T_0^{\leftarrow}\|. \end{aligned}$$

Given any element  $z \in Y_0$ , write  $J(z)$  as  $J(z) = J(x) + J(y)$  for some  $x \in T[X_0]$  and  $y \in Y_0$  such that  $J(y) \in (J[T[X_0]])^{\perp}$ . Since the vectors  $J(x)$  and  $J(y)$  are orthogonal in the Hilbert space  $\ell_2^m$ , the Pitagoras Theorem ensures that  $\|J(x)\| \leq \|J(z)\|$ . Taking into account that  $y$  belongs to the linear hull of the vectors  $e'_{n-k+1}, \dots, e'_m$  and  $j'_0(e'_l) = j_0(e'_l)$  for all  $l \in \{n-k+1, \dots, m\}$ , we conclude that  $j'_0(y) = j_0(y)$ . Then

$$\begin{aligned} \|j'_0(z) - j_0(z)\| &= \|j'_0(x+y) - j_0(x+y)\| = \|j'_0(x) - j_0(x)\| = \|(j'_0 - j_0) \circ J^{-1} \circ J(x)\| \leq \\ &\leq \|(j'_0 - j_0) \upharpoonright_{T[X_0]}\| \cdot \|J^{-1}\| \cdot \|J(x)\| \leq \|(j'_0 - j_0) \upharpoonright_{T[X_0]}\| \cdot \|J^{-1}\| \cdot \|J(z)\| \leq \\ &\leq \delta(n+1)\sqrt{n}\|T_0^{\leftarrow}\| \cdot \|J^{-1}\| \cdot \|J\| \cdot \|z\| \leq \delta(n+1)\sqrt{n}\|T_0^{\leftarrow}\| \sqrt{m} \cdot \|z\|. \end{aligned}$$

and hence

$$\|j'_0 - j_0\| \leq \delta(n+1)\sqrt{nm}\|T_0^{\leftarrow}\|.$$

Since  $j_0$  is a  $\delta$ -isometric embedding, the operator  $j'_0$  is a  $\delta(1 + (n+1)\sqrt{nm}\|T_0^{\leftarrow}\|)$ -isometric embedding by Lemma 3.  $\square$

Let

$$r = \frac{1}{1 + \delta(n+1) \max\{1, \sqrt{nm}\|T_0^{\leftarrow}\|\}},$$

$D_X$  be the convex hull of the set  $(i'_0)^{-1}[B_{\mathbb{G}}(1)] \cup B_X(r)$  in  $X$  and  $D_Y$  be the convex hull of the set  $(j'_0)^{-1}[B_{\mathbb{G}}(1)] \cup B_Y(r)$  in  $Y$ .

**Claim 3.**  $B_X(r) \subseteq D_X \subseteq B_X(\frac{1}{1-\delta(n+1)})$  and  $D_X \cap X_0 = (i'_0)^{-1}[B_{\mathbb{G}}(1)]$ .

*Proof.* The inclusion  $B_X(r) \subseteq D_X$  follows from the definition of  $D_X$ . To see that  $D_X \subseteq B_X(\frac{1}{1-\delta(n+1)})$ , take any  $x \in (i'_0)^{-1}[B_{\mathbb{G}}(1)]$ . By Claim 1,  $i'_0$  is a  $\delta(n+1)$ -isometric embedding and hence  $(1 - \delta(n+1))\|x\| \leq \|i'_0(x)\| < 1$  and  $\|x\| < \frac{1}{1-\delta(n+1)}$ . Then

$$D_X \subseteq \text{conv}((i'_0)^{-1}[B_{\mathbb{G}}(1)] \cup B_X(r)) \subseteq B_X(\frac{1}{1-\delta(n+1)}).$$

By Claim 1,  $i'_0[B_{X_0}(r)] \subseteq B_{\mathbb{G}}(r\|i'_0\|) \subseteq B_{\mathbb{G}}(r(1+\delta(n+1))) \subseteq B_{\mathbb{G}}(1)$  and  $i'_0[(i'_0)^{-1}[B_{\mathbb{G}}(1)]] \subseteq B_{\mathbb{G}}(1)$ , which implies that  $i'_0[D_X \cap X_0] \subseteq B_{\mathbb{G}}(1)$  and  $D_X \cap X_0 = (i'_0)^{-1}[B_{\mathbb{G}}(1)]$  by the injectivity of  $i'_0$ .  $\square$

By analogy, we can apply Claim 2 and prove

**Claim 4.**  $B_Y(r) \subseteq D_Y \subseteq B_Y(\frac{1}{1-\delta(1+(n+1)\sqrt{nm}\|T_0^{\leftarrow}\|)})$  and  $D_Y \cap Y_0 = (j'_0)^{-1}[B_{\mathbb{G}}(1)]$ .

**Claim 5.**  $T[D_X] \subseteq D_Y$ .

*Proof.* Observe that for every  $x \in (i'_0)^{-1}[B_{\mathbb{G}}(1)]$ , the equality  $j'_0 \circ T_0 = G \circ i'_0$  proved in (1) and the inequality  $\|G\| \leq 1$  imply

$$j'_0 \circ T_0(x) = G \circ i'_0(x) \in G[B_{\mathbb{G}}(1)] \subseteq B_{\mathbb{G}}(1)$$

and hence  $T(x) = T_0(x) \in (j'_0)^{-1}[B_{\mathbb{G}}(1)] \subseteq D_Y$ . On the other hand, for every  $x \in B_X(r)$  the inequality  $\|T\| \leq 1$  implies  $T(x) \in T[B_X(r)] \subseteq B_Y(r) \subseteq D_Y$ . Then

$$\begin{aligned} T[D_X] &= T[\text{conv}((i'_0)^{-1}[B_{\mathbb{G}}(1)] \cup B_X(r))] = \\ &\quad \text{conv}(T[(i'_0)^{-1}[B_{\mathbb{G}}(1)] \cup B_X(r)]) \subseteq \text{conv}(D_Y) = D_Y. \end{aligned}$$

$\square$

Let  $X'$  be the space  $X$  endowed with the norm  $\|\cdot\|_{X'}$  defined by

$$\|x\|_{X'} = \inf\{t \in \mathbb{R}_+ : x \in t \cdot D_X\},$$

and let  $Y'$  be the space  $Y$  endowed with the norm  $\|\cdot\|_{Y'}$

$$\|y\|_{Y'} = \inf\{t \in \mathbb{R}_+ : y \in t \cdot D_Y\}.$$

It follows that  $B_{X'}(1) = D_X$  and  $B_{Y'}(1) = D_Y$ . Let  $\text{Id}_{X,X'} : X \rightarrow X'$ ,  $\text{Id}_{X',X} : X' \rightarrow X$ ,  $\text{Id}_{Y,Y'} : Y \rightarrow Y'$ , and  $\text{Id}_{Y',Y} : Y' \rightarrow Y$  be the identity operators.

In the Banach spaces  $X'$  and  $Y'$  consider the subspaces  $X'_0 = \text{Id}_{X,X'}[X_0]$  and  $Y'_0 = \text{Id}_{Y,Y'}[Y_0]$ . Let  $\text{Id}_{X_0,X'_0} : X_0 \rightarrow X'_0$ ,  $\text{Id}_{X'_0,X_0} : X'_0 \rightarrow X_0$ ,  $\text{Id}_{Y_0,Y'_0} : Y_0 \rightarrow Y'_0$ , and  $\text{Id}_{Y'_0,Y_0} : Y'_0 \rightarrow Y_0$  be the identity operators.

Claims 3 and 4 imply that  $i'_0 \circ \text{Id}_{X'_0,X_0} : X'_0 \rightarrow \mathbb{G}$  and  $j'_0 \circ \text{Id}_{Y'_0,Y_0} : Y'_0 \rightarrow \mathbb{G}$  are isometric embeddings, and Claim 5 implies that the operator  $T' = \text{Id}_{Y,Y'} \circ T \circ \text{Id}_{X',X} : X' \rightarrow Y'$  is nonexpansive. Let  $T'_0 = T'|_{X'_0}$ . Since  $G \circ i'_0 \circ \text{Id}_{X'_0,X_0} = j'_0 \circ T'_0$ , we can apply the definition of the GurariĬ space and find  $\delta$ -isometric embeddings  $i' : X' \rightarrow \mathbb{G}$  and  $j' : Y' \rightarrow \mathbb{G}$  such that

$\|i' \upharpoonright_{X'_0} - i'_0 \circ \text{Id}_{X'_0, X_0}\| < \delta$ ,  $\|j' \upharpoonright_{Y'_0} - j'_0 \circ \text{Id}_{Y'_0, Y_0}\| < \delta$  and  $\|G \circ i' - j \circ T'\| < \delta$ . Consider the operators  $i = i' \circ \text{Id}_{X, X'} : X \rightarrow \mathbb{G}$  and  $j = j' \circ \text{Id}_{Y, Y'} : Y \rightarrow \mathbb{G}$ . It remains to prove that  $i, j$  are  $\varepsilon$ -isometric embeddings and

$$\max\{\|i \upharpoonright_{X_0} - i_0\|, \|j \upharpoonright_{Y_0} - j_0\|, \|G \circ i - j \circ T\|\} < \varepsilon.$$

To see that  $i$  is an  $\varepsilon$ -isometric embedding, take any nonzero  $x \in X$  and applying Claim 1, conclude that

$$\begin{aligned} \|i(x)\|_{\mathbb{G}} &= \|i' \circ \text{Id}_{X, X'}(x)\|_{\mathbb{G}} \leq \|i'\| \cdot \|\text{Id}_{X, X'}(x)\|_{X'} < (1 + \delta) \frac{1}{r} \|x\|_X = \\ &= (1 + \delta) \cdot (1 + \delta(n + 1) \max\{1, \sqrt{nm}\|T_0^{\leftarrow}\|\}) \|x\|_X \leq (1 + \varepsilon) \|x\|_X, \end{aligned}$$

by the choice of  $\delta = \delta(\varepsilon, n, m, \|T_0^{\leftarrow}\|)$ . On the other hand the inclusion  $D_X \subseteq B_X(\frac{1}{1 - \delta(n + 1)})$  proved in Claim 3 implies

$$\|i(x)\|_{\mathbb{G}} = \|i' \circ \text{Id}_{X, X'}(x)\|_{\mathbb{G}} > (1 - \delta) \|\text{Id}_{X, X'}(x)\|_{X'} \geq (1 - \delta)(1 - \delta(n + 1)) \|x\|_X \geq (1 - \varepsilon) \|x\|_X$$

by the choice of  $\delta = \delta(\varepsilon, n, m, \|T_0^{\leftarrow}\|)$ . This means that  $i : X \rightarrow \mathbb{G}$  is an  $\varepsilon$ -isometric embedding.

By analogy we can prove that  $j$  is an  $\varepsilon$ -isometric embedding.

The inclusion  $B_X(r) \subseteq D_X$  implies  $\|\text{Id}_{X_0, X'_0}\| \leq \|\text{Id}_{X, X'}\| \leq \frac{1}{r}$  and hence

$$\|i \upharpoonright_{X_0} - i_0\| = \|i' \circ \text{Id}_{X_0, X'_0} - i'_0 \circ \text{Id}_{X_0, X'_0}\| \leq \|i' - i'_0\| \cdot \|\text{Id}_{X_0, X'_0}\| < \frac{\delta}{r} \leq \varepsilon.$$

By analogy we can prove that  $\|j \upharpoonright_{Y_0} - j_0\| < \varepsilon$ . Finally, observe that

$$\begin{aligned} \|G \circ i - j \circ T\| &= \|G \circ i' \circ \text{Id}_{X, X'} - j' \circ \text{Id}_{Y, Y'} \circ T'\| = \|G \circ i' \circ \text{Id}_{X, X'} - j' \circ T' \circ \text{Id}_{X, X'}\| \leq \\ &\leq \|G \circ i' - j' \circ T'\| \cdot \|\text{Id}_{X, X'}\| < \delta \cdot \|\text{Id}_{X, X'}\| \leq \frac{\delta}{r} \leq \varepsilon. \end{aligned}$$

□

Next, we show that the spaces  $X, Y$ , operator  $T$ , subspaces  $X_0, Y_0$  and  $\delta$ -isometries  $i_0, j_0$  in Theorem 3 can be chosen from a suitable countable family of standard objects, defined below.

A *standard linear space* is the linear space  $\mathbb{R}^n$  for some  $n \in \omega$ . For a standard linear space  $X = \mathbb{R}^n$  by  $E_X = \{x \in \{0, 1\}^n : |x^{-1}(1)| = 1\}$  we denote the standard unit basis in  $X = \mathbb{R}^n$ . A linear subspace  $X_0$  of a standard linear space  $X$  is called *standard* if  $X_0$  coincides with the linear hull of some subset  $E \subseteq E_X$ . It is clear that each standard linear space  $X$  has exactly  $2^{\dim(X)}$  standard linear subspaces.

An operator  $T : X \rightarrow Y$  between linear spaces is called *standard* if there exist numbers  $k, n, m$  such that  $X = \mathbb{R}^n$ ,  $Y = \mathbb{R}^m$ ,  $0 \leq n - k \leq m$  and  $T$  assigns to each  $x \in \mathbb{R}^n$  the vector  $y = T(x)$  defined by

$$y(i) = \begin{cases} x(i + k) & \text{if } i < n - k \\ 0 & \text{otherwise.} \end{cases}$$

In this case  $\ker(T) = \mathbb{R}^k \times \{0\}^{n-k}$  and  $T[X] = \mathbb{R}^{n-k} \times \{0\}^{m-n+k}$ .

A norm  $\|\cdot\|$  on a standard linear space  $X = \mathbb{R}^n$  is called *rational* if its closed unit ball  $\{x \in X : \|x\| \leq 1\}$  coincides with the convex hull of some finite set  $F \subseteq \mathbb{Q}^n \subseteq \mathbb{R}^n$ . It is clear that the family of rational norms on  $\mathbb{R}^n$  is countable.

A *rational Banach space* is a standard linear space endowed with a rational norm.

For a countable dense set  $D$  in the Gurarii space  $\mathbb{G}$ , let  $\mathcal{F}_D$  be the family of all 9-tuples  $(\varepsilon, X, Y, T, X_0, Y_0, T_0, i_0, j_0)$  consisting of

- a rational number  $\varepsilon \in (0, 1]$ ;



- rational Banach spaces  $X, Y$ ;
- a nonexpansive standard operator  $T : X \rightarrow Y$ ;
- standard linear subspaces  $X_0 \subseteq X$  and  $Y_0 \subseteq Y$  such that  $T[X_0] \subseteq Y_0$ ;
- the operator  $T_0 = T|_{X_0} : X_0 \rightarrow Y_0$ ;
- $\delta(\varepsilon, \dim(X_0), \dim(Y_0), \|T_0^{\leftarrow}\|)$ -isometric embeddings  $i_0 : X_0 \rightarrow \mathbb{G}$  and  $j_0 : Y_0 \rightarrow \mathbb{G}$  such that  $i_0(E_X \cap X_0) \cup j_0(E_Y \cap Y_0) \subseteq D$ .

It is easy to see that the family  $\mathcal{F}_D$  is countable.

**Theorem 4.** *Let  $D$  be a countable dense subset of the Gurariĭ space  $\mathbb{G}$ . A nonexpansive operator  $G : \mathbb{G} \rightarrow \mathbb{G}$  is Gurariĭ if and only if it satisfies the condition:*

(GC) *for any 9-tuple  $(\varepsilon, X, Y, T, X_0, Y_0, T_0, i_0, j_0) \in \mathcal{F}_D$  with*

$$\|G \circ i_0 - j_0 \circ T|_{X_0}\| < \delta(\varepsilon, \dim(X_0), \dim(Y_0), \|T_0^{\leftarrow}\|),$$

*there exist  $\varepsilon$ -isometric embeddings  $i : X \rightarrow \mathbb{G}$ ,  $j : Y \rightarrow \mathbb{G}$  such that*

$$\|i|_{X_0} - i_0\| < \varepsilon, \quad \|j|_{Y_0} - j_0\| < \varepsilon, \quad \text{and} \quad \|G \circ i - j \circ T\| < \varepsilon.$$

*Proof.* The ‘‘only if’’ part follows immediately from the ‘‘only if’’ part of Theorem 3. To prove the ‘‘if’’ part, assume that the condition (GC) is satisfied. By Theorem 3, the Gurariĭ property of the operator  $G$  will follow as soon as we show that  $G$  satisfies the condition (GB). To this end, fix any  $\varepsilon \in (0, 1]$ , any nonexpansive operator  $T : X \rightarrow Y$  between finite-dimensional Banach spaces, any linear subspaces  $X_0 \subseteq X$  and  $Y_0 \subseteq Y$  with  $T[X_0] \subseteq Y_0$ , and  $\delta$ -isometric embeddings  $i_0 : X_0 \rightarrow \mathbb{G}$  and  $j_0 : Y_0 \rightarrow \mathbb{G}$  such that  $\|G \circ i_0 - j_0 \circ T_0\| < \delta := \delta(\varepsilon, \dim(X_0), \dim(Y_0), \|T_0^{\leftarrow}\|)$ , where  $T_0 = T|_{X_0} : X_0 \rightarrow Y_0$ .

Choose a basis  $E_X = \{e_1, \dots, e_{\dim(X)}\}$  in the space  $X$  such that

- $\{e_1, \dots, e_{\dim(\ker(T_0))}\}$  is a basis for the linear space  $\ker(T_0)$ ;
- $\{e_1, \dots, e_{\dim(\ker(T))}\}$  is a basis for the linear space  $\ker(T)$ ;
- $\{e_1, \dots, e_{\dim(\ker(T_0))}\} \cup \{e_{\dim(\ker(T))+1}, \dots, e_{\dim(\ker(T))+\dim(X_0)-\dim(\ker(T_0))}\}$  is a basis for the space  $X_0$ ;
- $\{e_1, \dots, e_{\dim(T^{-1}[Y_0])}\}$  is a basis for the linear space  $T^{-1}[Y_0]$ .

Let  $E_Y = \{e'_1, \dots, e'_{\dim(Y)}\}$  be a basis for the linear space  $Y$  such that  $e'_i = T(e_{\dim(\ker(T))+i})$  for all  $i \in \{1, \dots, \dim(X) - \dim(\ker(T))\}$ . The bases  $E_X$  and  $E_Y$  allow us to identify the spaces  $X, Y$  with the standard linear spaces  $\mathbb{R}^{\dim(X)}$  and  $\mathbb{R}^{\dim(Y)}$ . The choice of the bases  $E_X, E_Y$  ensures that the operator  $T : X \rightarrow Y$  is standard and the subspaces  $X_0 \subseteq X$  and  $Y_0 \subseteq Y$  are standard.

By the compactness of the unit spheres in the Banach spaces  $X_0, Y_0$ , there exists  $\delta' < \delta$  such that the  $\delta$ -isometric embeddings  $i_0 : X_0 \rightarrow \mathbb{G}$  and  $j_0 : Y_0 \rightarrow \mathbb{G}$  are  $\delta'$ -isometric embeddings. Since  $\|G \circ i_0 - j_0 \circ T_0\| < \delta$ , we can additionally assume that  $\|G \circ i_0 - j_0 \circ T_0\| < \delta'$ .

By the continuity of the function  $\delta(\cdot, \dim(X_0), \dim(Y_0), \cdot)$ , there exists a positive real number  $\nu$  such that

$$\delta' + 2\nu \leq \delta(\varepsilon', \dim(X_0), \dim(Y_0), t)$$

for any  $\varepsilon', t \in \mathbb{R}$  with

$$\frac{\varepsilon - 2\nu}{1 + \nu} \leq \varepsilon' \leq \varepsilon \quad \text{and} \quad \|T_0^{\leftarrow}\| \leq t \leq (1 + \nu)^2 \|T_0^{\leftarrow}\|.$$

By the density of  $D$  in  $\mathbb{G}$ , the  $\delta'$ -isometric embeddings  $i_0 : X_0 \rightarrow \mathbb{G}$  and  $j_0 : Y_0 \rightarrow \mathbb{G}$  can be approximated by  $\delta'$ -isometric embeddings  $i''_0 : X_0 \rightarrow \mathbb{G}$  and  $j''_0 : Y_0 \rightarrow \mathbb{G}$  such that  $i''_0(E_X \cap X_0) \cup j''_0(E_Y \cap Y_0) \subseteq D$ ,  $\max\{\|i''_0 - i_0\|, \|j''_0 - j_0\|\} < \nu$ , and  $\|G \circ i''_0 - j''_0 \circ T_0\| < \delta'$ . The norms  $\|\cdot\|_X$  and  $\|\cdot\|_Y$  on the standard linear spaces  $X, Y$  can be approximated by rational

norms  $\|\cdot\|_{X'} : X \rightarrow [0, \infty)$  and  $\|\cdot\|_{Y'} : Y \rightarrow [0, \infty)$  such that for every nonzero elements  $x \in X$  and  $y \in Y$  the following conditions are satisfied:

- (a)  $\|x\|_X \leq \|x\|_{X'} < (1 + \nu)\|x\|_X$  and  $\frac{1}{1+\nu}\|y\|_Y \leq \|y\|_{Y'} \leq \|y\|_Y$ ;
- (b)  $(1 - \delta')\|x\|_{X'} < \|i'_0(x)\|_{\mathbb{G}} < (1 + \delta')\|x\|_{X'}$  and  $(1 - \delta')\|y\|_{Y'} < \|j'_0(y)\|_{\mathbb{G}} < (1 + \delta')\|y\|_{X'}$ .

Let  $X'$  (resp.  $Y'$ ) be the rational Banach space  $X$  (resp.  $Y$ ) endowed with the rational norm  $\|\cdot\|_{X'}$  (resp.  $\|\cdot\|_{Y'}$ ). The condition (a) implies

$$(2) \quad B_{X'}(1) \subseteq B_X(1) \subseteq B_{X'}(1 + \nu) \quad \text{and} \quad B_Y(1) \subseteq B_{Y'}(1) \subseteq B_Y(1 + \nu).$$

Let

$$\text{Id}_{X, X'} : X \rightarrow X', \quad \text{Id}_{X', X} : X' \rightarrow X, \quad \text{Id}_{Y, Y'} : Y \rightarrow Y', \quad \text{and} \quad \text{Id}_{Y', Y} : Y' \rightarrow Y$$

be the identity operators. Let  $X'_0 = \text{Id}_{X, X'}[X_0]$ ,  $Y'_0 = \text{Id}_{Y, Y'}[Y_0]$ , and

$$\text{Id}_{X_0, X'_0} : X_0 \rightarrow X'_0, \quad \text{Id}_{X'_0, X_0} : X'_0 \rightarrow X_0, \quad \text{Id}_{Y_0, Y'_0} : Y_0 \rightarrow Y'_0 \quad \text{and} \quad \text{Id}_{Y'_0, Y_0} : Y'_0 \rightarrow Y_0$$

be the identity operators. Let  $i'_0 = i''_0 \circ \text{Id}_{X'_0, X_0}$  and  $j'_0 = j''_0 \circ \text{Id}_{Y'_0, Y_0}$ .

The condition (a) implies that  $\max\{\|\text{Id}_{X', X}\|, \|\text{Id}_{Y, Y'}\|\} \leq 1$  and hence the operator  $T' = \text{Id}_{Y, Y'} \circ T \circ \text{Id}_{X', X} : X' \rightarrow Y'$  is nonexpansive. Let  $T'_0 = T' \upharpoonright_{X'_0}$ . The inclusions (2) imply

$$\|T_0^{\leftarrow}\| \leq \|(T'_0)^{\leftarrow}\| \leq (1 + \nu)^2 \|T_0^{\leftarrow}\|.$$

Choose any rational  $\varepsilon'$  such that  $\frac{\varepsilon - 2\nu}{1 + \nu} \leq \varepsilon' \leq \frac{\varepsilon - \nu}{1 + \nu}$ . The choice of  $\nu$  ensures that

$$\delta' + 2\nu \leq \delta(\varepsilon', \dim(X_0), \dim(Y_0), \|(T'_0)^{\leftarrow}\|).$$

Then the 9-tuple  $(\varepsilon', X', Y', T', X'_0, Y'_0, T'_0, i'_0, j'_0)$  belongs to the family  $\mathcal{F}_D$ . Since

$$\begin{aligned} \|G \circ i'_0 - j'_0 \circ T'_0\| &\leq \|G \circ i''_0 - j''_0 \circ T_0\| \cdot \|\text{Id}_{X'_0, X_0}\| \leq \|G \circ i''_0 - j''_0 \circ T_0\| \leq \\ &\leq \|G \circ i''_0 - G \circ i_0\| + \|G \circ i_0 - j_0 \circ T_0\| + \|j_0 \circ T_0 - j''_0 \circ T_0\| \leq \\ &\leq \|G\| \cdot \|i''_0 - i_0\| + \delta' + \|j_0 - j''_0\| \cdot \|T_0\| \leq \nu + \delta' + \nu \leq \delta(\varepsilon', \dim(X_0), \dim(Y_0), \|(T'_0)^{\leftarrow}\|). \end{aligned}$$

By the condition (GC), there exist  $\varepsilon'$ -isometric embeddings  $i' : X' \rightarrow \mathbb{G}$  and  $j' : Y' \rightarrow \mathbb{G}$  such that

$$\max\{\|i' \upharpoonright_{X'_0} - i'_0\|, \|j' \upharpoonright_{Y'_0} - j'_0\|, \|G \circ i' - j' \circ T'\|\} < \varepsilon'.$$

The inequality  $\varepsilon'(1 + \nu) \leq \varepsilon$  and the condition (a) imply that the maps  $i = i' \circ \text{Id}_{X, X'}$  and  $j = j' \circ \text{Id}_{Y, Y'}$  are  $\varepsilon$ -isometric embeddings. It remains to show that

$$\max\{\|i \upharpoonright_{X_0} - i_0\|, \|j \upharpoonright_{Y_0} - j_0\|, \|G \circ i - j \circ T\|\} < \varepsilon.$$

Observe that

$$\begin{aligned} \|i \upharpoonright_{X_0} - i_0\| &= \|i' \circ \text{Id}_{X_0, X'_0} - i''_0\| + \|i''_0 - i_0\| \leq \\ &\leq \|i' \upharpoonright_{X'_0} - i''_0 \circ \text{Id}_{X'_0, X_0}\| \cdot \|\text{Id}_{X_0, X'_0}\| + \|i''_0 - i_0\| < \varepsilon'(1 + \nu) + \nu \leq \varepsilon. \end{aligned}$$

By analogy we can prove that  $\|j \upharpoonright_{Y_0} - j_0\| < \varepsilon$ . Finally,

$$\|G \circ i - j \circ T\| = \|G \circ i' \circ \text{Id}_{X, X'} - j' \circ T' \circ \text{Id}_{X, X'}\| \leq \|G \circ i' - j' \circ T'\| \cdot \|\text{Id}_{X, X'}\| \leq \varepsilon'(1 + \nu) < \varepsilon.$$

□

## 4. PROOF OF THEOREM 1

First we show that the set  $\mathcal{G}(\mathbb{G})$  of Gurariĭ operators is dense in the space  $\mathcal{B}(\mathbb{G})$ .

Fix any nonexpansive operator  $T \in \mathcal{B}(\mathbb{G})$  and any neighborhood  $O_T$  of  $T$  in  $\mathcal{B}(\mathbb{G})$ . By the definition of the strong operator topology, there exist  $\varepsilon > 0$  and a finite-dimensional subspace  $E \subseteq \mathbb{G}$  such that  $\{S \in \mathcal{B}(\mathbb{G}) : \|(S - T)|_E\| \leq 2\varepsilon\} \subseteq O_T$ . Consider the finite-dimensional Banach space  $F = T[E] \subseteq \mathbb{G}$ . Let  $\Omega : \mathbb{G} \rightarrow \mathbb{G}$  be the Gurariĭ operator, constructed in [1]. By [1, 3.3], the operator  $\Omega$  is universal. So, there exist isometric embeddings  $i : E \rightarrow \mathbb{G}$  and  $j : F \rightarrow \mathbb{G}$  such that  $\Omega \circ i = j \circ T|_E$ . By Lemma 5, for the isometric embeddings  $i : E \rightarrow \mathbb{G}$  and  $j : F \rightarrow \mathbb{G}$  there exist bijective isometries  $I, J \in \text{Iso}(\mathbb{G})$  such that  $\|i - I|_E\| < \varepsilon$  and  $\|j - J|_F\| < \varepsilon$ . Then  $S = J^{-1} \circ \Omega \circ I$  is a Gurariĭ operator such that for every  $x \in E$  we have

$$\begin{aligned} \|S(x) - T(x)\| &= \|J^{-1} \circ \Omega \circ I(x) - T(x)\| \leq \\ &\leq \|J^{-1} \circ \Omega \circ I(x) - J^{-1} \circ \Omega \circ i(x)\| + \|J^{-1} \circ \Omega \circ i(x) - T(x)\| \leq \\ &\leq \|J^{-1}\| \cdot \|\Omega\| \cdot \|I(x) - i(x)\| + \|J^{-1} \circ j \circ T(x) - T(x)\| = \\ &= \|I(x) - i(x)\| + \|j \circ T(x) - J \circ T(x)\| \leq \|I|_E - i\| \cdot \|x\| + \|j - J|_F\| \cdot \|T\| \cdot \|x\| \leq \\ &\leq \varepsilon\|x\| + \varepsilon\|x\| = 2\varepsilon\|x\| \end{aligned}$$

and hence  $\|(S - T)|_E\| \leq 2\varepsilon$ . The choice of  $\varepsilon$  ensures that  $S \in O_T \cap \mathcal{G}(\mathbb{G})$ , witnessing that the subspace  $\mathcal{G}(\mathbb{G})$  is dense in  $\mathcal{B}(\mathbb{G})$ .

Now we shall prove that the dense set  $\mathcal{G}(\mathbb{G})$  is of type  $G_\delta$  in  $\mathcal{B}(\mathbb{G})$ . Fix any countable dense set  $D$  in the Gurariĭ space  $\mathbb{G}$ . For every 9-tuple

$$t = (\varepsilon, X, Y, T, X_0, Y_0, T_0, i_0, j_0) \in \mathcal{F}_D$$

and the number  $\delta_t = \delta(\varepsilon, \dim(X_0), \dim(Y_0), \|T_0^{\leftarrow}\|)$ , consider the set

$$\mathcal{F}_t = \{G \in \mathcal{B}(\mathbb{G}) : \|G \circ i_0 - j_0 \circ T_0\| \geq \delta_t\}$$

and the set  $\mathcal{U}_t$  of all operators  $G \in \mathcal{B}(\mathbb{G})$  such that  $\|G \circ i_0 - j_0 \circ T_0\| < \delta_t$  and there exist  $\varepsilon$ -isometric embeddings  $i : X \rightarrow \mathbb{G}$  and  $j : Y \rightarrow \mathbb{G}$  such that

$$\max\{\|i|_{X_0} - i_0\|, \|j|_{Y_0} - j_0\|, \|G \circ i - j \circ T\|\} < \varepsilon.$$

It is easy to see that the set  $\mathcal{F}_t$  is closed in  $\mathcal{B}(\mathbb{G})$  and the set  $\mathcal{U}_t$  is open in  $\mathcal{B}(\mathbb{G})$ . Then their union  $(\mathcal{F}_t \cup \mathcal{U}_t)$  is a  $G_\delta$ -set in  $\mathcal{B}(\mathbb{G})$ . Theorem 4 ensures that  $\mathcal{G}(\mathbb{G}) = \bigcap_{t \in \mathcal{F}_D} (\mathcal{F}_t \cup \mathcal{U}_t)$  and hence  $\mathcal{G}(\mathbb{G})$  is a dense  $G_\delta$ -set in  $\mathcal{B}(\mathbb{G})$ .

## REFERENCES

- [1] J. Garbulińska-Węgrzyn, W. Kubiś, *A universal operator on the Gurariĭ space*, Journal of Operator Theory **73** (2015) 143–158.
- [2] J. Garbulińska-Węgrzyn, W. Kubiś, *A note on universal operators between separable Banach spaces*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM **114**:3 (2020), Paper No. 148, 11 pp.
- [3] Ju L. Gevorkjan, *Universality of the spaces of almost universal placement*, Funkcional. Anal. i Priložen **8** (1974) 72 (in Russian).
- [4] V.I. Gurariĭ, *Space of universal disposition, isotropic spaces and the Mazur problem on rotations of Banach spaces*, Sibirsk Math. Ž. **7** (1966) 1002–1013 (in Russian).
- [5] F. John, *Extremum problems with inequalities as subsidiary conditions*, Courant Anniversary volume, (1948), 187–204, Interscience, New York, 1948.
- [6] A.S. Kechris, *Classical Descriptive Set Theory*, Springer, 1995.
- [7] W. Kubiś, *Game-theoretic characterization of the Gurariĭ space*, Archiv der Mathematik **110** (2018) 53–59.
- [8] W. Kubiś, S. Solecki, *A proof of uniqueness of the Gurariĭ space*, Israel J. Math. **195** (2013) 449–456.
- [9] W. Lusky, *The Gurariĭ spaces are unique*, Arch. Math. (Basel) **27** (1976) 627–635.
- [10] W. B. Johnson, J. Lindenstrauss, *Handbook of the geometry of the Banach spaces. Volum 1*, North-Holland, 2001.

<sup>1</sup>IVAN FRANKO NATIONAL UNIVERSITY OF LVIV, UKRAINE  
*Email address:* `t.o.banakh@gmail.com`

<sup>2</sup>JAN KOCHANOWSKI UNIVERSITY, KIELCE, POLAND  
*Email address:* `jgarbulinska@ujk.edu.pl`