

**THE MATHIEU GROUP M_{23} AS ADDITIVE FUNCTIONS ON
THE FINITE FIELD OF SIZE 2^{11}**

YIMING BING*, BRIGHT HU, RONNI HU, RHIANNA LI, STEFAN LU, FINN MCDONALD,
MICHAEL SUN, NICHOLAS WOLFE, JOSHUA YAO, LEON ZHOU, NATHAN ZHOU

ABSTRACT. We explicitly extend the standard permutation action of the Mathieu group M_{23} on a 23 element set $C = C_{23}$ contained in a finite field of 2^{11} elements $\mathbb{F}_{2^{11}}$ to additive functions on this finite field. That is we represent M_{23} as functions $\varphi : \mathbb{F}_{2^{11}} \rightarrow \mathbb{F}_{2^{11}}$ such that $\varphi(x+y) = \varphi(x) + \varphi(y)$ and $\varphi|_C$ is the standard permutation action. We give explicit 11×11 matrices for the pair of standard generators of order 23 and order 5, as well as many tables to help facilitate future calculations.

INTRODUCTION

The Mathieu group M_{23} was first introduced in 1873 in [3]. (We translated this paper to English [1]). Since then, there has been much interest surrounding this group; for example, it is one of the sporadic simple finite groups in the classification program of finite simple groups and the only sporadic group for which the Inverse Galois Problem remains open.

In 1957 [4] proved the existence of an irreducible faithful representation of M_{23} as 11×11 matrices over $\mathbb{Z}/2\mathbb{Z}$ using the characterisation of M_{23} as the automorphism group of a Steiner system, where certain key lemmas were left to computer calculations and omitted out of necessity. There were also many mentions of such a definition which restricts to the standard permutation action that can be found online such as Wikipedia or Mathoverflow but we were not able to find an explicit construction of this elegant characterisation in the literature.

That is, let $C := C_{23}$ be the 23 element subgroup of the multiplicative group of the finite field of 2^{11} elements $\mathbb{F}_{2^{11}}$. We want to represent M_{23} as invertible functions

$$\varphi : \mathbb{F}_{2^{11}} \rightarrow \mathbb{F}_{2^{11}}$$

such that $\varphi(C) \subseteq C$ and $\varphi(x+y) = \varphi(x) + \varphi(y)$ for all $x, y \in \mathbb{F}_{2^{11}}$ and the restriction

$$\varphi|_C : C \rightarrow C$$

is a corresponding permutation from M_{23} acting transitively on a 23 element set.

In this article we proceed to construct a representation satisfying these criteria very explicitly in the hopes that we and other young mathematicians can gain a better understanding and make an easier transition into the subject matter. We do not mention Steiner systems nor Golay codes to achieve this representation, which we believe will add much greater accessibility for young readers. All parts of the proof are directly verifiable and carried out by hand.

We also include extensive tables detailing many of the explicit relationships between the elements of the finite structures involved for completeness in the hopes that they

also might be useful in future studies. We finally conclude with stating matrices for the generators of M_{23} .

1. THE FIELD $F_{2^{11}}$

We construct $F_{2^{11}}$ using the irreducible polynomial $x^{11} + x^2 + 1 \pmod{2}$

$$\mathbb{F}_{2^{11}} = F_{2^{11}}[x]/(x^{11} + x^2 + 1)F_{2^{11}}[x] \cong \{aX^2 + bX + c \mid a, b, c \in F_{2^{11}}, X^{11} + X^2 + 1 = 0\}$$

and with $x \mapsto X$ we express everything as powers of X in the first table. This polynomial appears on a list of irreducible polynomials in [2] and our table also serves as an explicit proof that both the polynomial is irreducible and that the quotient by this polynomial is a field with 2^{11} elements. The polynomial was chosen because it has the fewest terms with the lowest power of x . It is clear that every degree 11 polynomial will have the x^{11} term and if irreducible will have constant term 1 and at least 3 terms ($x^{11} + 1$ is reducible, for example substitute $x = 1$). It suffices to check that $x^{11} + x + 1$ is reducible.

While it is known the multiplicative group is cyclic (can be expressed as powers of a single element), our table below also proves it explicitly here using the X as the generator. It suffices to check that $X^{23} \neq 1$ and $X^{89} \neq 1$. The initials track who participated in the calculations. Many of these entries are not needed in our proof but are included for completeness.

Power	Expression	Binary String	Initials
X^{11}	$X^2 + 1$	00000000101	MS
X^{12}	$X^3 + X$	00000001010	BH
X^{13}	$X^4 + X^2$	00000010100	BH
X^{14}	$X^5 + X^3$	00000101000	BH
X^{15}	$X^6 + X^4$	00001010000	BH
X^{16}	$X^7 + X^5$	00010100000	FM, BH
X^{17}	$X^8 + X^6$	00101000000	BH
X^{18}	$X^9 + X^7$	01010000000	BH
X^{19}	$X^{10} + X^8$	10100000000	BH
X^{20}	$X^9 + X^2 + 1$	01000000101	MS, BH
X^{21}	$X^{10} + X^3 + X$	10000001010	BH
X^{22}	$X^4 + 1$	00000010001	FM
X^{23}	$X^5 + X$	00000100010	FM
X^{24}	$X^6 + X^2$	00001000100	FM
X^{25}	$X^7 + X^3$	00010001000	FM
X^{26}	$X^8 + X^4$	00100010000	FM
X^{27}	$X^9 + X^5$	01000100000	FM
X^{28}	$X^{10} + X^6$	10001000000	FM
X^{29}	$X^7 + X^2 + 1$	00010000101	FM
X^{30}	$X^8 + X^3 + X$	00100001010	FM
X^{31}	$X^9 + X^4 + X^2$	01000010100	FM
X^{32}	$X^{10} + X^5 + X^3$	10000101000	FM
X^{33}	$X^6 + X^4 + X^2 + 1$	00001010101	FM
X^{34}	$X^7 + X^5 + X^3 + X$	00010101010	BH
X^{35}	$X^8 + X^6 + X^4 + X^2$	00101010100	BH
X^{36}	$X^9 + X^7 + X^5 + X^3$	01010101000	BH
X^{37}	$X^{10} + X^8 + X^6 + X^4$	10101010000	BH
X^{38}	$X^9 + X^7 + X^5 + X^2 + 1$	01010100101	BH
X^{39}	$X^{10} + X^8 + X^6 + X^3 + X$	10101001010	BH

X^{40}	$X^9 + X^7 + X^4 + 1$	01010010001	MS, BH
X^{41}	$X^{10} + X^8 + X^5 + X$	10100100010	BH
X^{42}	$X^9 + X^6 + 1$	01001000001	BH
X^{43}	$X^{10} + X^7 + X$	10010000010	BH
X^{44}	$X^8 + 1$	00100000001	BH
X^{45}	$X^9 + X$	01000000010	BH
X^{46}	$X^{10} + X^2$	10000000100	BH
X^{47}	$X^3 + X^2 + 1$	00000001101	BH
X^{48}	$X^4 + X^3 + X$	00000011010	BH
X^{49}	$X^5 + X^4 + X^2$	00000110100	BH
X^{50}	$X^6 + X^5 + X^3$	00001101000	BH
X^{51}	$X^7 + X^6 + X^4$	00011010000	BH
X^{52}	$X^8 + X^7 + X^5$	00110100000	BH
X^{53}	$X^9 + X^8 + X^6$	01101000000	BH
X^{54}	$X^{10} + X^9 + X^7$	11010000000	BH
X^{55}	$X^{10} + X^8 + X^2 + 1$	10100000101	RK
X^{56}	$X^9 + X^3 + X^2 + X + 1$	01000001111	BH
X^{57}	$X^{10} + X^4 + X^3 + X^2 + X$	10000011110	BH
X^{58}	$X^5 + X^4 + X^3 + 1$	00000111001	BH
X^{59}	$X^6 + X^5 + X^4 + X$	00001110010	BH
X^{60}	$X^7 + X^6 + X^5 + X^2$	00011100100	BH
X^{61}	$X^8 + X^7 + X^6 + X^3$	00111001000	BH
X^{62}	$X^9 + X^8 + X^7 + X^4$	01110010000	BH
X^{63}	$X^{10} + X^9 + X^8 + X^5$	11100100000	BH
X^{64}	$X^{10} + X^9 + X^6 + X^2 + 1$	11001000101	MS, BH
X^{65}	$X^{10} + X^7 + X^3 + X^2 + X + 1$	10010001111	BH
X^{66}	$X^8 + X^4 + X^3 + X + 1$	00100011011	FM
X^{67}	$X^9 + X^5 + X^4 + X^2 + X$	01000110110	BH
X^{68}	$X^{10} + X^6 + X^5 + X^3 + X^2$	10001101100	BH
X^{69}	$X^7 + X^6 + X^4 + X^3 + X^2 + 1$	00011011101	BH
X^{70}	$X^8 + X^7 + X^5 + X^4 + X^3 + X$	00110111010	BH
X^{71}	$X^9 + X^8 + X^6 + X^5 + X^4 + X^2$	01101110100	BH
X^{72}	$X^{10} + X^9 + X^7 + X^6 + X^5 + X^3$	11011101000	BH
X^{73}	$X^{10} + X^8 + X^7 + X^6 + X^4 + X^2 + 1$	10111010101	BH
X^{74}	$X^9 + X^8 + X^7 + X^5 + X^3 + X^2 + X + 1$	01110101111	BH
X^{75}	$X^{10} + X^9 + X^8 + X^6 + X^4 + X^3 + X^2 + X$	11101011110	BH
X^{76}	$X^{10} + X^9 + X^7 + X^5 + X^4 + X^3 + 1$	11010111001	BH
X^{77}	$X^{10} + X^8 + X^6 + X^5 + X^4 + X^2 + X + 1$	10101110111	FM
X^{78}	$X^9 + X^7 + X^6 + X^5 + X^3 + X + 1$	01011101011	BH
X^{79}	$X^{10} + X^8 + X^7 + X^6 + X^4 + X^2 + X$	10111010110	BH
X^{80}	$X^9 + X^8 + X^7 + X^5 + X^3 + 1$	01110101001	BH
X^{81}	$X^{10} + X^9 + X^8 + X^6 + X^4 + X$	11101010010	BH
X^{82}	$X^{10} + X^9 + X^7 + X^5 + 1$	11010100001	BH
X^{83}	$X^{10} + X^8 + X^6 + X^2 + X + 1$	10101000111	BH
X^{84}	$X^9 + X^7 + X^3 + X + 1$	01010001011	BH
X^{85}	$X^{10} + X^8 + X^4 + X^2 + X$	10100010110	BH
X^{86}	$X^9 + X^5 + X^3 + 1$	01000101001	BH
X^{87}	$X^{10} + X^6 + X^4 + X$	10001010010	BH
X^{88}	$X^7 + X^5 + 1$	00010100001	FM
X^{89}	$X^8 + X^6 + X$	00101000010	BH
X^{90}	$X^9 + X^7 + X^2$	01010000100	BH
X^{91}	$X^{10} + X^8 + X^3$	10100001000	BH
X^{92}	$X^9 + X^4 + X^2 + 1$	01000010101	BH

X^{93}	$X^{10} + X^5 + X^3 + X$	10000101010	BH
X^{94}	$X^6 + X^4 + 1$	00001010001	BH
X^{95}	$X^7 + X^5 + X$	00010100010	BH
X^{96}	$X^8 + X^6 + X^2$	00101000100	RH, MS, BH
X^{97}	$X^9 + X^7 + X^3$	01010001000	BH
X^{98}	$X^{10} + X^8 + X^4$	10100010000	BH
X^{99}	$X^9 + X^5 + X^2 + 1$	01000100101	BH
X^{100}	$X^{10} + X^6 + X^3 + X$	10001001010	BH

1.1. **The Powers of X^{89} .** Let $\alpha = X^{89}$, here are the elements of C .

Power	Expression	Binary	Initials
α	$X^8 + X^6 + X$	00101000010	MS
α^2	$X^7 + X^5 + X^3 + X^2 + X$	00010101110	MS
α^3	$X^{10} + X^7 + X^3 + X^2$	10010001100	MS
α^4	$X^{10} + X^6 + X^5 + X^4 + X^3 + X^2$	10001111100	MS
α^5	$X^8 + X^7 + X^6 + X^5 + 1$	00111100001	SL, BH, YB
α^6	$X^9 + X^6 + X^5 + X^4 + X^3 + X^2 + 1$	01001111101	ML, RK
α^7	$X^{10} + X^9 + X^8 + X^7 + X^4 + X^2 + X$	11110010110	RH, MS
α^8	$X^{10} + X^9 + X^8 + X^6 + X^4 + X^3 + X^2 + X + 1$	11101011111	RH, MS
α^9	$X^{10} + X^8 + X^7 + X^6 + X^5 + X^3 + X^2 + X$	10111101110	RL
α^{10}	$X^{10} + X^7 + X + 1$	10010000011	JY, LZ
α^{11}	$X^7 + X^5 + X^2 + X + 1$	00010100111	MS, RH
α^{12}	$X^{10} + X^9 + X^8 + X^7 + X^6 + X^4 + X^3 + X + 1$	11111011011	RH,MS,JY,YB
α^{13}	$X^8 + X^7 + X^5 + X$	00110100010	RL, MS, JY
α^{14}	$X^8 + X^4 + X^3 + 1$	00100011001	MS,LZ,RL
α^{15}	$X^{10} + X^8 + X^7 + X^6 + X^5 + X^4 + X^2 + 1$	11010011110	MS,RL,JY
α^{16}	$X^8 + X^6 + X^5 + X^4 + X^3 + X$	00101111010	MS, BH, RL
α^{17}	$X^{10} + X^9 + X^7 + X^6$	11011000000	RH,MS,LZ,YB
α^{18}	$X^{10} + X^9 + X^7 + X^6 + X^4 + X + 1$	11011010011	YB, BH
α^{19}	$X^8 + X^5 + X^4 + X^3 + X^2 + X + 1$	00100111111	NZ, MS
α^{20}	$X^9 + X^5 + X^3$	01000101000	BH
α^{21}	$X^{10} + X^9 + X^8 + X^6 + X^4 + X^2$	11101010100	NW
α^{22}	$X^{10} + X^5 + X^4 + X^3 + X^2 + 1$	10000111101	YB
α^{23}	1	00000000001	MS,BH
α^{24}	$X^8 + X^6 + X$	00101000010	YB

Notes: 1,2,3,4,5,10,20 powers are used to calculate the 23rd power of alpha as one so they are very likely correct. Similarly with 2,5,16,21 (NW). Also checked 11,12,22, 17,18,19,20 all checked.

1.2. **Powers of X As Powers of α .** Let $A = \{1, \alpha, \alpha^2, \dots, \alpha^{10}\}$, we see from the table below that this is a additively independent set. This is because we know the set $\chi = \{1, X, X^2, \dots, X^{10}\}$ is additively independent and we write each $x \in \chi$ as combinations of $a \in A$.

Power	Expression	Initials
X^0	1	BH, YB
X^1	$\alpha^{10} + \alpha^5 + \alpha^3 + \alpha^2 + \alpha$	MS, LS
X^2	$\alpha^9 + \alpha^7 + \alpha^6 + 1$	MS, JG, TC, LS, JT
X^3	$\alpha^9 + \alpha^7 + \alpha^6 + \alpha^5 + \alpha^2 + \alpha$	MS, LZ, NZ
X^4	$\alpha^8 + \alpha^6 + \alpha^5 + \alpha^4 + \alpha^2 + 1$	LZ, NW, MS
X^5	$\alpha^9 + \alpha^3 + \alpha$	RH, MS
X^6	$\alpha^{10} + \alpha^8 + \alpha^6 + \alpha^5$	LZ, FM, TC
X^7	$\alpha^{10} + \alpha^9 + \alpha^2 + \alpha + 1$	MS
X^8	$\alpha^8 + \alpha^6 + \alpha^3 + \alpha^2$	RH, MS, LZ, NW
X^9	$\alpha^{10} + \alpha^9 + \alpha^6 + \alpha^5 + \alpha^4 + \alpha^3 + 1$	YB
X^{10}	$\alpha^{10} + \alpha^9 + \alpha^5 + \alpha^3$	MS, JG

We list some examples of additive functions on $\mathbb{F}_{2^{11}}$ that preserve C .

Example 1.1.

$$x \mapsto \alpha x.$$

Example 1.2.

$$\text{Fr}(x) = x^2$$

1.3. Powers of α as the Sum of Lower Powers of α .

Power	Expression	Initial
α^{11}	$\alpha^9 + \alpha^7 + \alpha^6 + \alpha^5 + \alpha + 1$	MS, RL
α^{12}	$\alpha^{10} + \alpha^8 + \alpha^7 + \alpha^6 + \alpha^2 + \alpha$	MS
α^{13}	$\alpha^8 + \alpha^6 + \alpha^5 + \alpha^3 + \alpha^2 + \alpha + 1$	MS, RL, LZ
α^{14}	$\alpha^9 + \alpha^7 + \alpha^6 + \alpha^4 + \alpha^3 + \alpha^2 + \alpha$	MS, RH
α^{15}	$\alpha^{10} + \alpha^8 + \alpha^7 + \alpha^5 + \alpha^4 + \alpha^3 + \alpha^2$	BH
α^{16}	$\alpha^8 + \alpha^7 + \alpha^4 + \alpha^3 + \alpha + 1$	MS
α^{17}	$\alpha^9 + \alpha^8 + \alpha^5 + \alpha^4 + \alpha^2 + \alpha$	MS
α^{18}	$\alpha^{10} + \alpha^9 + \alpha^6 + \alpha^5 + \alpha^3 + \alpha^2$	MS
α^{19}	$\alpha^{10} + \alpha^9 + \alpha^5 + \alpha^4 + \alpha^3 + \alpha + 1$	MS, BH
α^{20}	$\alpha^{10} + \alpha^9 + \alpha^7 + \alpha^4 + \alpha^2 + 1$	LZ, BH
α^{21}	$\alpha^{10} + \alpha^9 + \alpha^8 + \alpha^7 + \alpha^6 + \alpha^3 + 1$	MS, FM
α^{22}	$\alpha^{10} + \alpha^8 + \alpha^6 + \alpha^5 + \alpha^4 + 1$	LZ, BH
α^{23}	1	BH, MS, YB

Here is an example which does not preserve C .

Example 1.3. Example of function defined on 11 elements of C : $f(1) = 1$, $f(\alpha) = \alpha^2$, $f(\alpha^2) = \alpha$, $f(\alpha^i) = \alpha^i$ for $i \neq 1, 2$ $i < 11$. no adjacent transposition can be together with a full cycle as otherwise we get S_{23} .

Explicity, $f(\alpha^{11}) = \alpha^9 + \alpha^7 + \alpha^6 + \alpha^5 + \alpha^2 + 1 \notin C$

2. EXTENDING FUNCTIONS

2.1. Powers of β as Powers of α and vice versa.

Power of β	Cor. Power of α	Power of α	Cor. Power of β
β	α^5	α	β^{14}
β^2	α^{10}	α^2	β^5
β^3	α^{15}	α^3	β^{19}
β^4	α^{20}	α^4	β^{10}
β^5	α^2	α^5	β
β^6	α^7	α^6	β^{15}
β^7	α^{12}	α^7	β^6
β^8	α^{17}	α^8	β^{20}
β^9	α^{22}	α^9	β^{11}
β^{10}	α^4	α^{10}	β^2
β^{11}	α^9	α^{11}	β^{16}
β^{12}	α^{14}	α^{12}	β^7
β^{13}	α^{19}	α^{13}	β^{21}
β^{14}	α	α^{14}	β^{12}
β^{15}	α^6	α^{15}	β^3
β^{16}	α^{11}	α^{16}	β^{17}
β^{17}	α^{16}	α^{17}	β^8
β^{18}	α^{21}	α^{18}	β^{22}
β^{19}	α^3	α^{19}	β^{13}
β^{20}	α^8	α^{20}	β^4
β^{21}	α^{13}	α^{21}	β^{18}
β^{22}	α^{18}	α^{22}	β^9
β^{23}	α^{23}	α^{23}	β^{23}

Checked by YB

Let $A = \{1, \alpha, \alpha^2, \dots, \alpha^{10}\}$ and $\chi = \{1, X, X^2, \dots, X^{10}\}$ be the additively independent sets from previous sections.

Definition 2.1. Let $f : C \rightarrow C$ by $f(\beta^j) = \beta^{j+1}$. Under the identification

$$\beta^j \mapsto j$$

f can be given by the cycle

$$(1, 2, 3, \dots, 22, 23).$$

Definition 2.2. Let $g : C \rightarrow C$ be defined as

$$(3, 17, 10, 7, 9)(4, 13, 14, 19, 5)(8, 18, 11, 12, 23)(15, 20, 22, 21, 16)$$

under the identification

$$\beta^j \mapsto j.$$

Proposition 2.1 (Mathieu [3]). *The functions f and g generate M_{23} inside the symmetric group S_{23} .*

Proposition 2.2. *The function f extends uniquely to an additive function on $\mathbb{F}_{2^{11}}$. Moreover we have*

$$f(x) = \beta x$$

for all $x \in \mathbb{F}_{2^{11}}$.

Proof. f is uniquely determined as an additive function by its values on the first 11 powers of alpha, while we note that multiplication by β is clearly additive and agrees with f on C . Hence f extends uniquely to multiplication by β and is additive. \square

Theorem 2.1. *The function g extends uniquely to an additive function on $\mathbb{F}_{2^{11}}$.*

Proof. It suffices to show that outputs of α^{11} to α^{22} agree with those determined by additivity on the additively independent set $1, \alpha, \alpha^2, \dots, \alpha^{10}$. The remaining outputs are then defined uniquely by additivity and the inputs in A .

$$\begin{aligned}
 g(\alpha^{11}) &= g(\alpha^9 + \alpha^7 + \alpha^6 + \alpha^5 + \alpha^1 + 1) \\
 &= g(\beta^{11} + \beta^6 + \beta^{15} + \beta + \beta^{14} + \beta^{23}) \\
 &= g(\beta^{11}) + g(\beta^6) + g(\beta^{15}) + g(\beta) + g(\beta^{14}) + g(\beta^{23}) \\
 &= \beta^{12} + \beta^6 + \beta^{20} + \beta + \beta^{19} + \beta^8 \\
 &= \alpha^{14} + \alpha^7 + \alpha^8 + \alpha^5 + \alpha^3 + \alpha^{17} \\
 &= (\alpha^9 + \alpha^7 + \alpha^6 + \alpha^4 + \alpha^3 + \alpha^2 + \alpha) + \alpha^7 + \alpha^8 + \alpha^5 + \alpha^3 + \alpha^{17} \\
 &= \alpha^9 + \alpha^8 + \alpha^6 + \alpha^5 + \alpha^4 + \alpha^2 + \alpha + \alpha^{17} \\
 &= \alpha^9 + \alpha^8 + \alpha^6 + \alpha^5 + \alpha^4 + \alpha^2 + \alpha + (\alpha^9 + \alpha^8 + \alpha^5 + \alpha^4 + \alpha^2 + \alpha) \\
 &= \alpha^6, \quad g(\beta^{16}) = \beta^{15}, \quad \text{YB, RL.}
 \end{aligned}$$

$$\begin{aligned}
 g(\alpha^{12}) &= g(\alpha^{10} + \alpha^8 + \alpha^7 + \alpha^6 + \alpha^2 + \alpha) \\
 &= g(\beta^2 + \beta^{20} + \beta^6 + \beta^{15} + \beta^5 + \beta^{14}) \\
 &= g(\beta^2) + g(\beta^{20}) + g(\beta^6) + g(\beta^{15}) + g(\beta^5) + g(\beta^{14}) \\
 &= \beta^2 + \beta^{22} + \beta^6 + \beta^{20} + \beta^4 + \beta^{19} \\
 &= \alpha^{10} + \alpha^{18} + \alpha^7 + \alpha^8 + \alpha^{20} + \alpha^3 \\
 &= \alpha^{10} + (\alpha^{10} + \alpha^9 + \alpha^6 + \alpha^5 + \alpha^3 + \alpha^2) + \alpha^7 + \alpha^8 + \alpha^{20} + \alpha^3 \\
 &= \alpha^9 + \alpha^6 + \alpha^5 + \alpha^2 + \alpha^7 + \alpha^8 + \alpha^{20} \\
 &= \alpha^9 + \alpha^6 + \alpha^5 + \alpha^2 + \alpha^7 + \alpha^8 + (\alpha^{10} + \alpha^9 + \alpha^7 + \alpha^4 + \alpha^2 + 1) \\
 &= \alpha^6 + \alpha^5 + \alpha^8 + (\alpha^{10} + \alpha^4 + 1) \\
 &= \alpha^{10} + \alpha^8 + \alpha^6 + \alpha^5 + \alpha^4 + 1 \\
 &= \alpha^{22}, \quad g(\beta^7) = \beta^9, \quad \text{YB, RL}
 \end{aligned}$$

$$\begin{aligned}
 g(\alpha^{13}) &= g(\alpha^8 + \alpha^6 + \alpha^5 + \alpha^3 + \alpha^2 + \alpha + 1) \\
 &= g(\beta^{20} + \beta^{15} + \beta + \beta^{19} + \beta^5 + \beta^{14} + \beta^{23}) \\
 &= \beta^{22} + \beta^{20} + \beta + \beta^5 + \beta^4 + \beta^{19} + \beta^8 \\
 &= \alpha^{18} + \alpha^8 + \alpha^5 + \alpha^2 + \alpha^{20} + \alpha^3 + \alpha^{17} \\
 &= (\alpha^{10} + \alpha^9 + \alpha^6 + \alpha^5 + \alpha^3 + \alpha^2) + \alpha^8 + \alpha^5 + \alpha^2 + \alpha^3 + \alpha^{20} + \alpha^{17} \\
 &= \alpha^{10} + \alpha^9 + \alpha^8 + \alpha^6 + (\alpha^{10} + \alpha^9 + \alpha^7 + \alpha^4 + \alpha^2 + 1) + \alpha^{17} \\
 &= \alpha^8 + \alpha^7 + \alpha^6 + \alpha^4 + \alpha^2 + 1 + (\alpha^9 + \alpha^8 + \alpha^5 + \alpha^4 + \alpha^2 + \alpha) \\
 &= \alpha^9 + \alpha^7 + \alpha^6 + \alpha^5 + \alpha + 1 \\
 &= \alpha^{11}, \quad g(\beta^{21}) = \beta^{16}, \quad \text{YB, RL, LZ}
 \end{aligned}$$

$$\begin{aligned}
g(\alpha^{14}) &= g(\alpha^9 + \alpha^7 + \alpha^6 + \alpha^4 + \alpha^3 + \alpha^2 + \alpha) \\
&= g(\beta^{11} + \beta^6 + \beta^{15} + \beta^{10} + \beta^{19} + \beta^5 + \beta^{14}) \\
&= g(\beta^{11}) + g(\beta^6) + g(\beta^{15}) + g(\beta^{10}) + g(\beta^{19}) + g(\beta^5) + g(\beta^{14}) \\
&= \beta^{12} + \beta^6 + \beta^{20} + \beta^7 + \beta^5 + \beta^4 + \beta^{19} \\
&= \alpha^{14} + \alpha^7 + \alpha^8 + \alpha^{12} + \alpha^2 + \alpha^{20} + \alpha^3 \\
&= (\alpha^9 + \alpha^7 + \alpha^6 + \alpha^4 + \alpha^3 + \alpha^2 + \alpha) + \alpha^7 + \alpha^8 + \alpha^2 + \alpha^3 + \alpha^{12} + \alpha^{20} \\
&= \alpha^9 + \alpha^8 + \alpha^6 + \alpha^4 + \alpha + (\alpha^{10} + \alpha^8 + \alpha^7 + \alpha^6 + \alpha^2 + \alpha) + \alpha^{20} \\
&= \alpha^{10} + \alpha^9 + \alpha^7 + \alpha^4 + \alpha^2 + (\alpha^{10} + \alpha^9 + \alpha^7 + \alpha^4 + \alpha^2 + 1) \\
&= 1 \\
&= \alpha^{23}, \quad g(\beta^{12}) = \beta^{23}, \quad \text{YB, RL, NZ}
\end{aligned}$$

$$\begin{aligned}
g(\alpha^{15}) &= g(\alpha^{10} + \alpha^8 + \alpha^7 + \alpha^5 + \alpha^4 + \alpha^3 + \alpha^2) \\
&= g(\beta^2 + \beta^{20} + \beta^6 + \beta + \beta^{10} + \beta^{19} + \beta^5) \\
&= g(\beta^2) + g(\beta^{20}) + g(\beta^6) + g(\beta) + g(\beta^{10}) + g(\beta^{19}) + g(\beta^5) \\
&= \beta^2 + \beta^{22} + \beta^6 + \beta + \beta^7 + \beta^5 + \beta^4 \\
&= \alpha^{10} + \alpha^{18} + \alpha^7 + \alpha^5 + \alpha^{12} + \alpha^2 + \alpha^{20} \\
&= \alpha^{10} + (\alpha^{10} + \alpha^9 + \alpha^6 + \alpha^5 + \alpha^3 + \alpha^2) + \alpha^7 + \alpha^5 + \alpha^{12} + \alpha^2 + \alpha^{20} \\
&= \alpha^9 + \alpha^7 + \alpha^6 + \alpha^3 + \alpha^{12} + \alpha^{20} \\
&= \alpha^9 + \alpha^7 + \alpha^6 + \alpha^3 + (\alpha^{10} + \alpha^8 + \alpha^7 + \alpha^6 + \alpha^2 + \alpha) + \alpha^{20} \\
&= \alpha^{10} + \alpha^9 + \alpha^8 + \alpha^3 + \alpha^2 + \alpha + \alpha^{20} \\
&= \alpha^{10} + \alpha^9 + \alpha^8 + \alpha^3 + \alpha^2 + \alpha + (\alpha^{10} + \alpha^9 + \alpha^7 + \alpha^4 + \alpha^2 + 1) \\
&= \alpha^8 + \alpha^7 + \alpha^4 + \alpha^3 + \alpha + 1 \\
&= \alpha^{16}, \quad g(\beta^3) = \beta^{17}, \quad \text{YB, RL, NZ}
\end{aligned}$$

$$\begin{aligned}
 g(\alpha^{16}) &= g(\alpha^8 + \alpha^7 + \alpha^4 + \alpha^3 + \alpha + 1) \\
 &= g(\beta^{20} + \beta^6 + \beta^{10} + \beta^{19} + \beta^{14} + \beta^{23}) \\
 &= g(\beta^{20}) + g(\beta^6) + g(\beta^{10}) + g(\beta^{19}) + g(\beta^{14}) + g(\beta^{23}) \\
 &= \beta^{22} + \beta^6 + \beta^7 + \beta^5 + \beta^{19} + \beta^8 \\
 &= \alpha^{18} + \alpha^7 + \alpha^{12} + \alpha^2 + \alpha^3 + \alpha^{17} \\
 &= (\alpha^{10} + \alpha^9 + \alpha^6 + \alpha^5 + \alpha^3 + \alpha^2) + \alpha^7 + \alpha^{12} + \alpha^2 + \alpha^3 + \alpha^{17} \\
 &= \alpha^{10} + \alpha^9 + \alpha^7 + \alpha^6 + \alpha^5 + \alpha^{12} + \alpha^{17} \\
 &= \alpha^{10} + \alpha^9 + \alpha^7 + \alpha^6 + \alpha^5 + (\alpha^{10} + \alpha^8 + \alpha^7 + \alpha^6 + \alpha^2 + \alpha) + \alpha^{17} \\
 &= \alpha^9 + \alpha^8 + \alpha^5 + \alpha^2 + \alpha + \alpha^{17} \\
 &= \alpha^9 + \alpha^8 + \alpha^5 + \alpha^2 + \alpha + (\alpha^9 + \alpha^8 + \alpha^5 + \alpha^4 + \alpha^2 + \alpha) \\
 &= \alpha^4 \\
 &= \alpha^4, \quad g(\beta^{17}) = \beta^{10}, \quad \text{YB, RL, NZ}
 \end{aligned}$$

$$\begin{aligned}
 g(\alpha^{17}) &= g(\alpha^9 + \alpha^8 + \alpha^5 + \alpha^4 + \alpha^2 + \alpha) \\
 &= g(\beta^{11} + \beta^{20} + \beta + \beta^{10} + \beta^5 + \beta^{14}) \\
 &= g(\beta^{11}) + g(\beta^{20}) + g(\beta) + g(\beta^{10}) + g(\beta^5) + g(\beta^{14}) \\
 &= \beta^{12} + \beta^{22} + \beta + \beta^7 + \beta^4 + \beta^{19} \\
 &= \alpha^{14} + \alpha^{18} + \alpha^5 + \alpha^{12} + \alpha^{20} + \alpha^3 \\
 &= (\alpha^9 + \alpha^7 + \alpha^6 + \alpha^4 + \alpha^3 + \alpha^2 + \alpha) + \alpha^{18} + \alpha^5 + \alpha^{12} + \alpha^{20} + \alpha^3 \\
 &= \alpha^9 + \alpha^7 + \alpha^6 + \alpha^5 + \alpha^4 + \alpha^2 + \alpha + \alpha^{18} + \alpha^{12} + \alpha^{20} \\
 &= \alpha^9 + \alpha^7 + \alpha^6 + \alpha^5 + \alpha^4 + \alpha^2 + \alpha + (\alpha^{10} + \alpha^9 + \alpha^6 + \alpha^5 + \alpha^3 + \alpha^2) + \alpha^{12} + \alpha^{20} \\
 &= \alpha^{10} + \alpha^7 + \alpha^4 + \alpha^3 + \alpha + \alpha^{12} + \alpha^{20} \\
 &= \alpha^{10} + \alpha^7 + \alpha^4 + \alpha^3 + \alpha + (\alpha^{10} + \alpha^8 + \alpha^7 + \alpha^6 + \alpha^2 + \alpha) + \alpha^{20} \\
 &= \alpha^8 + \alpha^6 + \alpha^4 + \alpha^3 + \alpha^2 + \alpha^{20} \\
 &= \alpha^8 + \alpha^6 + \alpha^4 + \alpha^3 + \alpha^2 + (\alpha^{10} + \alpha^9 + \alpha^7 + \alpha^4 + \alpha^2 + 1) \\
 &= \alpha^{10} + \alpha^9 + \alpha^8 + \alpha^7 + \alpha^6 + \alpha^3 + 1 \\
 &= \alpha^{21}, \quad g(\beta^8) = \beta^{18}, \quad \text{YB, NZ, RL}
 \end{aligned}$$

$$\begin{aligned}
g(\alpha^{18}) &= g(\alpha^{10} + \alpha^9 + \alpha^6 + \alpha^5 + \alpha^3 + \alpha^2) \\
&= g(\beta^2 + \beta^{11} + \beta^{15} + \beta + \beta^{19} + \beta^5) \\
&= g(\beta^2) + g(\beta^{11}) + g(\beta^{15}) + g(\beta) + g(\beta^{19}) + g(\beta^5) \\
&= \beta^2 + \beta^{12} + \beta^{20} + \beta + \beta^5 + \beta^4 \\
&= \alpha^{10} + \alpha^{14} + \alpha^8 + \alpha^5 + \alpha^2 + \alpha^{20} \\
&= \alpha^{10} + (\alpha^9 + \alpha^7 + \alpha^6 + \alpha^4 + \alpha^3 + \alpha^2 + \alpha) + \alpha^8 + \alpha^5 + \alpha^2 + \alpha^{20} \\
&= \alpha^{10} + \alpha^9 + \alpha^8 + \alpha^7 + \alpha^6 + \alpha^5 + \alpha^4 + \alpha^3 + \alpha + \alpha^{20} \\
&= \alpha^{10} + \alpha^9 + \alpha^8 + \alpha^7 + \alpha^6 + \alpha^5 + \alpha^4 + \alpha^3 + \alpha + (\alpha^{10} + \alpha^9 + \alpha^7 + \alpha^4 + \alpha^2 + 1) \\
&= \alpha^8 + \alpha^6 + \alpha^5 + \alpha^3 + \alpha^2 + \alpha + 1 \\
&= \alpha^{13}, \quad g(\beta^{22}) = \beta^{21}, \quad \text{YB, NZ}
\end{aligned}$$

$$\begin{aligned}
g(\alpha^{19}) &= g(\alpha^{10} + \alpha^9 + \alpha^5 + \alpha^4 + \alpha^3 + \alpha + 1) \\
&= g(\beta^2 + \beta^{11} + \beta + \beta^{10} + \beta^{19} + \beta^{14} + \beta^{23}) \\
&= g(\beta^2) + g(\beta^{11}) + g(\beta) + g(\beta^{10}) + g(\beta^{19}) + g(\beta^{14}) + g(\beta^{23}) \\
&= \beta^2 + \beta^{12} + \beta + \beta^7 + \beta^5 + \beta^{19} + \beta^8 \\
&= \alpha^{10} + \alpha^{14} + \alpha^5 + \alpha^{12} + \alpha^2 + \alpha^3 + \alpha^{17} \\
&= \alpha^{10} + (\alpha^9 + \alpha^7 + \alpha^6 + \alpha^4 + \alpha^3 + \alpha^2 + \alpha) + \alpha^5 + \alpha^{12} + \alpha^2 + \alpha^3 + \alpha^{17} \\
&= \alpha^{10} + \alpha^9 + \alpha^7 + \alpha^6 + \alpha^5 + \alpha^4 + \alpha + \alpha^{12} + \alpha^{17} \\
&= \alpha^{10} + \alpha^9 + \alpha^7 + \alpha^6 + \alpha^5 + \alpha^4 + \alpha + (\alpha^{10} + \alpha^8 + \alpha^7 + \alpha^6 + \alpha^2 + \alpha) + \alpha^{17} \\
&= \alpha^9 + \alpha^8 + \alpha^5 + \alpha^4 + \alpha^2 + \alpha^{17} \\
&= \alpha^9 + \alpha^8 + \alpha^5 + \alpha^4 + \alpha^2 + (\alpha^9 + \alpha^8 + \alpha^5 + \alpha^4 + \alpha^2 + \alpha) \\
&= \alpha, \quad g(\beta^{13}) = \beta^{14}, \quad \text{YB, NZ}
\end{aligned}$$

$$\begin{aligned}
 g(\alpha^{20}) &= g(\alpha^{10} + \alpha^9 + \alpha^7 + \alpha^4 + \alpha^2 + 1) \\
 &= g(\beta^2 + \beta^{11} + \beta^6 + \beta^{10} + \beta^5 + \beta^{23}) \\
 &= g(\beta^2) + g(\beta^{11}) + g(\beta^6) + g(\beta^{10}) + g(\beta^5) + g(\beta^{23}) \\
 &= \beta^2 + \beta^{12} + \beta^6 + \beta^7 + \beta^4 + \beta^8 \\
 &= \alpha^{10} + \alpha^{14} + \alpha^7 + \alpha^{12} + \alpha^{20} + \alpha^{17} \\
 &= \alpha^{10} + (\alpha^9 + \alpha^7 + \alpha^6 + \alpha^4 + \alpha^3 + \alpha^2 + \alpha) + \alpha^7 + \alpha^{12} + \alpha^{20} + \alpha^{17} \\
 &= \alpha^{10} + \alpha^9 + \alpha^6 + \alpha^4 + \alpha^3 + \alpha^2 + \alpha + \alpha^{12} + \alpha^{20} + \alpha^{17} \\
 &= \alpha^{10} + \alpha^9 + \alpha^6 + \alpha^4 + \alpha^3 + \alpha^2 + \alpha + (\alpha^{10} + \alpha^8 + \alpha^7 + \alpha^6 + \alpha^2 + \alpha) + \alpha^{20} + \alpha^{17} \\
 &= \alpha^9 + \alpha^8 + \alpha^7 + \alpha^4 + \alpha^3 + \alpha^{20} + \alpha^{17} \\
 &= \alpha^9 + \alpha^8 + \alpha^7 + \alpha^4 + \alpha^3 + (\alpha^{10} + \alpha^9 + \alpha^7 + \alpha^4 + \alpha^2 + 1) + \alpha^{17} \\
 &= \alpha^{10} + \alpha^8 + \alpha^3 + \alpha^2 + 1 + \alpha^{17} \\
 &= \alpha^{10} + \alpha^8 + \alpha^3 + \alpha^2 + 1 + (\alpha^9 + \alpha^8 + \alpha^5 + \alpha^4 + \alpha^2 + \alpha) \\
 &= \alpha^{10} + \alpha^9 + \alpha^5 + \alpha^4 + \alpha^3 + \alpha + 1 \\
 &= \alpha^{19}, \quad g(\beta^4) = \beta^{13}, \quad \text{YB, NZ}
 \end{aligned}$$

$$\begin{aligned}
 g(\alpha^{21}) &= g(\alpha^{10} + \alpha^9 + \alpha^8 + \alpha^7 + \alpha^6 + \alpha^3 + 1) \\
 &= g(\beta^2 + \beta^{11} + \beta^{20} + \beta^6 + \beta^{15} + \beta^{19} + \beta^{23}) \\
 &= g(\beta^2) + g(\beta^{11}) + g(\beta^{20}) + g(\beta^6) + g(\beta^{15}) + g(\beta^{19}) + g(\beta^{23}) \\
 &= \beta^2 + \beta^{12} + \beta^{22} + \beta^6 + \beta^{20} + \beta^5 + \beta^8 \\
 &= \alpha^{10} + \alpha^{14} + \alpha^{18} + \alpha^7 + \alpha^8 + \alpha^2 + \alpha^{17} \\
 &= \alpha^{10} + (\alpha^9 + \alpha^7 + \alpha^6 + \alpha^4 + \alpha^3 + \alpha^2 + \alpha) + \alpha^{18} + \alpha^7 + \alpha^8 + \alpha^2 + \alpha^{17} \\
 &= \alpha^{10} + \alpha^9 + \alpha^8 + \alpha^6 + \alpha^4 + \alpha^3 + \alpha + \alpha^{18} + \alpha^{17} \\
 &= \alpha^{10} + \alpha^9 + \alpha^8 + \alpha^6 + \alpha^4 + \alpha^3 + \alpha + (\alpha^{10} + \alpha^9 + \alpha^6 + \alpha^5 + \alpha^3 + \alpha^2) + \alpha^{17} \\
 &= \alpha^8 + \alpha^5 + \alpha^4 + \alpha^2 + \alpha + \alpha^{17} \\
 &= \alpha^8 + \alpha^5 + \alpha^4 + \alpha^2 + \alpha + (\alpha^9 + \alpha^8 + \alpha^5 + \alpha^4 + \alpha^2 + \alpha) \\
 &= \alpha^9 \\
 &= \alpha^9, \quad g(\beta^{18}) = \beta^{11}, \quad \text{YB, NZ}
 \end{aligned}$$

$$\begin{aligned}
g(\alpha^{22}) &= g(\alpha^{10} + \alpha^8 + \alpha^6 + \alpha^5 + \alpha^4 + 1) \\
&= g(\beta^2 + \beta^{20} + \beta^{15} + \beta + \beta^{10} + \beta^{23}) \\
&= g(\beta^2) + g(\beta^{20}) + g(\beta^{15}) + g(\beta) + g(\beta^{10}) + g(\beta^{23}) \\
&= \beta^2 + \beta^{22} + \beta^{20} + \beta + \beta^7 + \beta^8 \\
&= \alpha^{10} + \alpha^{18} + \alpha^8 + \alpha^5 + \alpha^{12} + \alpha^{17} \\
&= \alpha^{10} + (\alpha^{10} + \alpha^9 + \alpha^6 + \alpha^5 + \alpha^3 + \alpha^2) + \alpha^8 + \alpha^5 + \alpha^{12} + \alpha^{17} \\
&= \alpha^9 + \alpha^8 + \alpha^6 + \alpha^3 + \alpha^2 + \alpha^{12} + \alpha^{17} \\
&= \alpha^9 + \alpha^8 + \alpha^6 + \alpha^3 + \alpha^2 + (\alpha^{10} + \alpha^8 + \alpha^7 + \alpha^6 + \alpha^2 + \alpha) + \alpha^{17} \\
&= \alpha^{10} + \alpha^9 + \alpha^7 + \alpha^3 + \alpha + \alpha^{17} \\
&= \alpha^{10} + \alpha^9 + \alpha^7 + \alpha^3 + \alpha + (\alpha^9 + \alpha^8 + \alpha^5 + \alpha^4 + \alpha^2 + \alpha) \\
&= \alpha^{10} + \alpha^8 + \alpha^7 + \alpha^5 + \alpha^4 + \alpha^3 + \alpha^2 \\
&= \alpha^{15}, \quad g(\beta^9) = \beta^3, \quad \text{YB, NZ}
\end{aligned}$$

□

Remark 2.1. The additivity of g was first tested for $\beta := \alpha$ and found to be not additive. We then tried $\beta := \alpha^5$ because it was unrelated to α under the squaring function and suspect that those obtained from successive squaring of α^5 should also produce an additive g while those which aren't do not.

2.2. Matrices. Since the functions f and g can be extended to additive functions on $\mathbb{F}_{2^{11}}$ we can express them as matrices with respect to the bases A and χ ordered as they were written in their definitions.

$$[g]_A^A = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 & 0 & 1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 \end{pmatrix}$$

$$[f]_A^A = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 1 \end{pmatrix}$$

$$[f]_\chi^\chi = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 & 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 1 & 0 & 0 & 0 & 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 & 1 & 0 & 0 & 1 & 0 & 1 \end{pmatrix}$$

$$[g]_\chi^\chi = \begin{pmatrix} 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 1 & 0 & 0 & 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 1 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 1 & 0 & 1 \end{pmatrix}$$

Theorem 2.2. *There is a faithful 11 dimensional irreducible representation of M_{23} over \mathbb{F}_2 such that the generators f and g are given by the matrices $[f]_\chi^\chi$ and $[g]_\chi^\chi$ with respect to χ , while it is given by $[f]_A^A$ and $[g]_A^A$ with respect to A . The 23 element subgroup of the multiplicative group of $\mathbb{F}_{2^{11}}$ is preserved by M_{23} under this representation and the restrictions of f to C and g to C under the identification*

$$j \mapsto \alpha^{5j}$$

is given by

$$f|_C = (1, 2, 3, \dots, 22, 23)$$

and

$$g|_C = (3, 17, 10, 7, 9)(4, 13, 14, 19, 5)(8, 18, 11, 12, 23)(15, 20, 22, 21, 16).$$

Furthermore, 11 is the minimal dimension of a faithful \mathbb{F}_2 representation.

Proof. Since f and g correspond to generators of M_{23} as permutations in S_{23} , they will also generate M_{23} as functions on $\mathbb{F}_{2^{11}}$. (Faithful). Moreover, since f and g are additive (\mathbb{F}_2 linear), the functions they generate will also be additive and hence we have a representation of M_{23} .

Now since this representation is faithful, it suffices to show that there are no smaller faithful representations and this will imply irreducibility as well. To do this it suffices to observe that for the size of M_{23} to divide the size of GL , we need $23|(2^i-1)$ for some i but this forces $i \geq 11$. □

Remark 2.2. Since M_{23} is simple, every non-trivial representation is faithful.

Remark 2.3. We also tested to see if this representation extends to M_{24} using the generators f, g and

$$h = (1, 24)(2, 23)(3, 12)(4, 16)(5, 18)(6, 10)(7, 20)(8, 14)(9, 21)(11, 17)(13, 22)(15, 19),$$

which appeared in [3] with $B = 21A$, $C = A$ and $D = -A$, but h does not seem to be additive for any choice of β obtained from α^5 by successive squaring. In each case we tested an additivity relation that does not involve β and obtained a contradiction of the form $\beta = 0$.

ACKNOWLEDGEMENTS

We are very grateful for the encouragement and support of the parents of our talented young researchers many of whom are not yet even in middle school. Honourable mention to Gavin Tran, Rhianna Kho and Max Lau for participating. We also acknowledge free resources such as Wikipedia, ATLAS, groupprops, MathStackExchange, MathOverflow and open access journals for their assistance. This project was completed without access to journal subscriptions.

REFERENCES

- [1] Y. Bing et al. “French to English Mathieu 1873 Translation”. In: *placeholder journal* placeholder volume (0), placeholder pages. URL: <https://placeholder>.
- [2] R. Church. “Tables of Irreducible Polynomials for the First Four Prime Moduli”. In: *Annals of Mathematics* 36 (1935), pp. 198–209.
- [3] Emile Mathieu. “Sur la fonction cinq fois transitive de 24 quantités.[On the Five times transitive function of 24 quantities]”. In: *Journal de mathématiques pures et appliquées 2e série* 18 (1873), pp. 25–46.
- [4] L. J. Paige. “A Note on Mathieu Groups”. In: *Canadian Journal of Mathematics* 9 (1957), pp. 15–18.

DR MICHAEL SUN’S SCHOOL OF MATHS, NSW, AUSTRALIA
Corresponding author: humanyimingbing@gmail.com