

Updated determinations of $|V_{us}|$ with tau decays using new recent estimates of radiative corrections for light-meson leptonic decay rates

A. Lusiani^{1*},

¹ Scuola Normale Superiore and INFN sezione di Pisa, Italy

* alberto.lusiani@pi.infn.it

January 13, 2022

*16th International Workshop on Tau Lepton Physics (TAU2021),
September 27 – October 1, 2021
doi:[10.21468/SciPostPhysProc.7](https://doi.org/10.21468/SciPostPhysProc.7)*

Abstract

We update the $|V_{us}|$ determinations using the HFLAV 2018 report tau branching fraction results with recent new estimates of the $\pi\ell 2$ and $K\ell 2$ radiative corrections. There are minor changes of the central values and uncertainties.

Contents

1	Introduction	1
2	$ V_{us} $ from $B(\tau \rightarrow K \nu)/B(\tau \rightarrow \pi \nu)$ and $B(\tau \rightarrow K \nu)$	2
3	Conclusions	3
	References	3

1 Introduction

Recent measurements of $|V_{ud}|$, $|V_{us}|$ and $|V_{ub}|$ are not consistent with the unitarity condition on the first row of the CKM matrix [1, 2]. Tau decay measurements are used to determine $|V_{us}|$ [3, 4], supplementing the more precise determinations that are obtained using kaon decays. All these estimates also rely on lattice QCD estimates of form factors and decay constants [5, 6], with the exception of the $|V_{us}|$ determinations based on the total branching fraction of the tau lepton into strange final states. Two of the $|V_{us}|$ determinations using tau measurements rely on estimates of the radiative corrections for the branching fractions $B(\tau \rightarrow \pi/K \nu)$, which are computed using

also the radiative corrections for $B(\pi \rightarrow \ell \nu)$ and $B(K \rightarrow \ell \nu)$ [7]. New estimates of these radiative corrections have been computed with a novel approach using lattice QCD+QED [8]. We evaluate in the following the impact of these new estimates on the $|V_{us}|$ determinations using tau decay measurements.

2 $|V_{us}|$ from $B(\tau \rightarrow K \nu)/B(\tau \rightarrow \pi \nu)$ and $B(\tau \rightarrow K \nu)$

We use the tau branching fractions of the HFLAV 2018 report fit [3] and we compute $|V_{us}|$ using the updated external inputs provided by the Review of Particle Physics [9], by the FLAG review of lattice QCD calculations [5, 6], by CODATA 2018 [10]. In updating the CODATA constants from the values used in the HFLAV 2018 report, a numerical transcription error that slightly affected this $|V_{us}|$ determination in the HFLAV 2018 report has been corrected. Finally, we obtain $|V_{us}|$ using both the original [7] and the recently published [8] radiative corrections.

We compute $|V_{us}|$ as in the HFLAV 2018 report from the ratio of branching fractions $B(\tau^- \rightarrow K^- \nu_\tau)/B(\tau^- \rightarrow \pi^- \nu_\tau)$ and from the branching fraction $B(\tau^- \rightarrow K^- \nu_\tau)$ using the equations

$$\frac{B(\tau^- \rightarrow K^- \nu_\tau)}{B(\tau^- \rightarrow \pi^- \nu_\tau)} = \frac{f_{K^\pm}^2 |V_{us}|^2 (m_\tau^2 - m_K^2)^2}{f_{\pi^\pm}^2 |V_{ud}|^2 (m_\tau^2 - m_\pi^2)^2} \frac{1 + \delta R_{\tau/K}}{1 + \delta R_{\tau/\pi}} (1 + \delta R_{K\mu 2/\pi\mu 2}),$$

$$B(\tau^- \rightarrow K^- \nu_\tau) = \frac{G_F^2}{16\pi\hbar} f_{K^\pm}^2 |V_{us}|^2 \tau_\tau m_\tau^3 \left(1 - \frac{m_K^2}{m_\tau^2}\right)^2 (1 + \delta R_{\tau/K})(1 + \delta R_{K\mu 2}),$$

respectively. We use Refs. [11–14] to compute the radiative-correction factor

$$\frac{1 + \delta R_{\tau/K}}{1 + \delta R_{\tau/\pi}} = \frac{1 + (0.90 \pm 0.22)\%}{1 + (0.16 \pm 0.14)\%}.$$

$|V_{ud}| = 0.97373 \pm 0.00031$ is taken from a 2020 updated determination [1].

The radiation correction terms $\delta R_{K\mu 2/\pi\mu 2} = -0.69 \pm 0.17$ and $\delta R_{K\mu 2} = 1.07 \pm 0.21$ are provided without the isospin-breaking corrections by Refs. [7, 15–18]. The same sources report also $\delta R_{\pi\mu 2} = 1.76 \pm 0.21$. The three estimates are consistent with a correlation of 0.67 between $\delta R_{\pi\mu 2}$ and $\delta R_{K\mu 2}$, which is used when computing $|V_{us}|$. These radiative correction terms are used with the lattice QCD decay constants that include isospin-breaking corrections from the FLAG 2019 lattice QCD averages with $N_f = 2 + 1 + 1$: $f_{K^\pm}/f_{\pi^\pm} = 1.1932 \pm 0.0021$ [5, 19–22] and $f_{K^\pm} = 155.7 \pm 0.3$ MeV [5, 20, 21, 23].

New estimates of the radiation correction terms inclusive of isospin-breaking corrections $\delta R'_{K\mu 2/\pi\mu 2} = -1.26 \pm 0.14$ and $\delta R'_{K\mu 2} = 0.24 \pm 0.10$ are provided by Ref. [8]. The same source reports also $\delta R'_{\pi\mu 2} = 1.53 \pm 0.19$. The three estimates are consistent with a correlation of 0.63 between $\delta R'_{\pi\mu 2}$ and $\delta R'_{K\mu 2}$, which originates from a correlation of 0.794 between $\delta R'_{\pi\mu 2}$ the non-isospin-breaking component of $\delta R'_{K\mu 2}$ [24]. These radiative correction terms are used with the isospin-symmetric lattice QCD decay constants with $N_f = 2 + 1 + 1$ provided by Ref. [8]: $f_K/f_\pi = 1.1966 \pm 0.0018$ and $f_K = 156.1 \pm 0.2$ MeV.

Table 1 reports the values of $|V_{us}|_{\tau K/\pi}$ from $B(\tau \rightarrow K \nu)/B(\tau \rightarrow \pi \nu)$ and $|V_{us}|_{\tau K}$ from $B(\tau \rightarrow K \nu)$ using both the original and the new $\pi\ell 2$ and $K\ell 2$ radiative corrections. The improvements in the precision of the radiative corrections results in minor improvements on the $|V_{us}|$

Table 1: $|V_{us}|$ determinations. $|V_{us}|_{\tau s}$ denotes $|V_{us}|$ obtained with the $\tau \rightarrow X_s \nu$ inclusive method using the HFLAV 2018 report [3], complemented with the updated external inputs described in the text. $|V_{us}|_{\text{uni}}$ denotes the value of $|V_{us}|$ assuming the CKM matrix unitarity, $|V_{ud}|$ from Ref. [1] and $|V_{us}|$ from Ref. [9]. The determinations of $|V_{us}|_{\tau K/\pi}$ and $|V_{us}|_{\tau K}$ using the exclusive tau branching fractions is reported using both the original and the new estimates of the $\pi\ell 2$ and $K\ell 2$ radiative corrections. $|V_{us}|_{\tau}$ denotes the combination of all three $|V_{us}|$ determinations with tau decay measurements. The signed difference with respect to $|V_{us}|_{\text{uni}}$ in standard deviations is reported for all $|V_{us}|$ determinations.

	$ V_{us} $			
$ V_{us} _{\text{uni}}$	0.2277 ± 0.0013		0.0σ	
$ V_{us} _{\tau s}$	0.2192 ± 0.0019		-3.6σ	
	Cirigliano & Neufeld 2011		Di Carlo <i>et al.</i> 2019	
$ V_{us} _{\tau K/\pi}$	0.2234 ± 0.0015	-2.0σ	0.2235 ± 0.0015	-2.0σ
$ V_{us} _{\tau K}$	0.2226 ± 0.0015	-2.6σ	0.2229 ± 0.0014	-2.6σ
$ V_{us} _{\tau}$	0.2217 ± 0.0013	-3.2σ	0.2219 ± 0.0013	-3.1σ

determinations because of other larger contributing uncertainties. The changes of the central values are also minor compared with the total uncertainties. Figure 1 reports the $|V_{us}|$ determinations in this document compared with $|V_{us}|$ from the CKM matrix unitarity and $|V_{us}|$ from kaon decay measurements.

3 Conclusions

Recent new slightly more precise estimates of the $\pi\ell 2$ and $K\ell 2$ radiative corrections have been used to update the $|V_{us}|$ determination using the tau branching fractions of the HFLAV 2018 report, resulting in minor changes of the central values and uncertainties.

References

- [1] J. C. Hardy and I. S. Towner, *Superaligned $0^+ \rightarrow 0^+$ nuclear β decays: 2020 critical survey, with implications for V_{ud} and CKM unitarity*, Phys. Rev. C **102**(4), 045501 (2020), doi:[10.1103/PhysRevC.102.045501](https://doi.org/10.1103/PhysRevC.102.045501).
- [2] C.-Y. Seng, D. Galviz, W. J. Marciano and U.-G. Meißner, *An update on $|V_{us}|$ and $|V_{us}/V_{ud}|$ from semileptonic kaon and pion decays* (2021), [2107.14708](https://arxiv.org/abs/2107.14708).
- [3] Y. S. Amhis *et al.*, *Averages of b -hadron, c -hadron, and τ -lepton properties as of 2018*, Eur. Phys. J. **C81**, 226 (2021), doi:[10.1140/epjc/s10052-020-8156-7](https://doi.org/10.1140/epjc/s10052-020-8156-7), updated results and plots available at <https://hflav.web.cern.ch/>, 1909.12524.




Figure 1: $|V_{us}|$ determinations using tau decay measurements with the new radiative corrections in Ref. [8], compared with $|V_{us}|$ from the CKM matrix unitarity and $|V_{us}|$ from kaon measurements in the Review of Particle Physics [9].

- [4] A. Lusiani, *Improved determination of $|V_{us}|$ with τ decays*, PoS ICHEP2020, 447 (2021), doi:[10.22323/1.390.0447](https://doi.org/10.22323/1.390.0447).
- [5] S. Aoki *et al.*, *FLAG Review 2019* (2019), [1902.08191](https://arxiv.org/abs/1902.08191).
- [6] S. Aoki *et al.*, *FLAG Review 2019 December 2020 web update*, FLAG web site (2020).
- [7] V. Cirigliano and H. Neufeld, *A note on isospin violation in $Pl_2(\text{gamma})$ decays*, Phys. Lett. **B700**, 7 (2011), doi:[10.1016/j.physletb.2011.04.038](https://doi.org/10.1016/j.physletb.2011.04.038), [1102.0563](https://arxiv.org/abs/1102.0563).
- [8] M. Di Carlo, D. Giusti, V. Lubicz, G. Martinelli, C. T. Sachrajda, F. Sanfilippo, S. Simula and N. Tantalo, *Light-meson leptonic decay rates in lattice QCD+QED*, Phys. Rev. D **100**(3), 034514 (2019), doi:[10.1103/PhysRevD.100.034514](https://doi.org/10.1103/PhysRevD.100.034514), [1904.08731](https://arxiv.org/abs/1904.08731).
- [9] P. Zyla *et al.*, *Review of Particle Physics*, PTEP **2020**(8), 083C01 (2020), doi:[10.1093/ptep/ptaa104](https://doi.org/10.1093/ptep/ptaa104).
- [10] E. Tiesinga, P. J. Mohr, D. B. Newell and B. N. Taylor, *CODATA recommended values of the fundamental physical constants: 2018**, Rev. Mod. Phys. **93**(2), 025010 (2021), doi:[10.1103/RevModPhys.93.025010](https://doi.org/10.1103/RevModPhys.93.025010), Database developed by J. Baker, M. Douma, and S. Kotochigova. Available at <http://physics.nist.gov/constants>, National Institute of Standards and Technology, Gaithersburg, MD 20899.
- [11] W. J. Marciano and A. Sirlin, *Radiative corrections to π_{l2} decays*, Phys. Rev. Lett. **71**, 3629 (1993), doi:[10.1103/PhysRevLett.71.3629](https://doi.org/10.1103/PhysRevLett.71.3629).

- [12] R. Decker and M. Finkemeier, *Radiative corrections to the decay $\tau \rightarrow \pi(K)\nu_\tau$* , Phys. Lett. **B334**, 199 (1994), doi:[10.1016/0370-2693\(94\)90611-4](https://doi.org/10.1016/0370-2693(94)90611-4).
- [13] R. Decker and M. Finkemeier, *Short and long distance effects in the decay $\tau \rightarrow \pi\nu_\tau(\gamma)$* , Nucl. Phys. **B438**, 17 (1995), doi:[10.1016/0550-3213\(95\)00597-L](https://doi.org/10.1016/0550-3213(95)00597-L), [hep-ph/9403385](https://arxiv.org/abs/hep-ph/9403385).
- [14] R. Decker and M. Finkemeier, *Radiative corrections to the decay $\tau \rightarrow \pi\nu_\tau$* , Nucl. Phys. Proc. Suppl. **40**, 453 (1995), doi:[10.1016/0920-5632\(95\)00170-E](https://doi.org/10.1016/0920-5632(95)00170-E), [hep-ph/9411316](https://arxiv.org/abs/hep-ph/9411316).
- [15] A. Pich, *Precision Tau Physics*, Prog. Part. Nucl. Phys. **75**, 41 (2014), doi:[10.1016/j.pnpnp.2013.11.002](https://doi.org/10.1016/j.pnpnp.2013.11.002), [1310.7922](https://arxiv.org/abs/1310.7922).
- [16] W. J. Marciano, *Precise determination of $|V(us)|$ from lattice calculations of pseudoscalar decay constants*, Phys. Rev. Lett. **93**, 231803 (2004), doi:[10.1103/PhysRevLett.93.231803](https://doi.org/10.1103/PhysRevLett.93.231803), [hep-ph/0402299](https://arxiv.org/abs/hep-ph/0402299).
- [17] M. Tanabashi *et al.*, *Review of Particle Physics*, Phys. Rev. **D98**(3), 030001 (2018), doi:[10.1103/PhysRevD.98.030001](https://doi.org/10.1103/PhysRevD.98.030001).
- [18] J. L. Rosner, S. Stone and R. S. Van de Water, *Leptonic Decays of Charged Pseudoscalar Mesons - 2015* (2015), [1509.02220](https://arxiv.org/abs/1509.02220).
- [19] N. Miller *et al.*, *F_K/F_π from Möbius Domain-Wall fermions solved on gradient-flowed HISQ ensembles*, Phys. Rev. D **102**(3), 034507 (2020), doi:[10.1103/PhysRevD.102.034507](https://doi.org/10.1103/PhysRevD.102.034507), [2005.04795](https://arxiv.org/abs/2005.04795).
- [20] R. J. Dowdall, C. T. H. Davies, G. P. Lepage and C. McNeile, *V_{us} from pi and K decay constants in full lattice QCD with physical u, d, s and c quarks*, Phys. Rev. **D88**, 074504 (2013), doi:[10.1103/PhysRevD.88.074504](https://doi.org/10.1103/PhysRevD.88.074504), [1303.1670](https://arxiv.org/abs/1303.1670).
- [21] N. Carrasco *et al.*, *Leptonic decay constants f_K, f_D , and f_{D_s} with $N_f = 2 + 1 + 1$ twisted-mass lattice QCD*, Phys. Rev. **D91**, 054507 (2015), doi:[10.1103/PhysRevD.91.054507](https://doi.org/10.1103/PhysRevD.91.054507), [1411.7908](https://arxiv.org/abs/1411.7908).
- [22] A. Bazavov *et al.*, *B- and D-meson leptonic decay constants from four-flavor lattice QCD*, Phys. Rev. **D98**, 074512 (2018), doi:[10.1103/PhysRevD.98.074512](https://doi.org/10.1103/PhysRevD.98.074512), [1712.09262](https://arxiv.org/abs/1712.09262).
- [23] A. Bazavov *et al.*, *Charmed and light pseudoscalar meson decay constants from four-flavor lattice QCD with physical light quarks*, Phys. Rev. **D90**, 074509 (2014), doi:[10.1103/PhysRevD.90.074509](https://doi.org/10.1103/PhysRevD.90.074509), [1407.3772](https://arxiv.org/abs/1407.3772).
- [24] S. Simula, private communication (2020).