

On Some Applications of the Sagnac Effect

¹A. H. Ziaie*, ¹H. Moradpour†, ²V. B. Bezerra‡, ³A. Jawad§

¹*Research Institute for Astronomy and Astrophysics of Maragha (RIAAM),
University of Maragheh, P. O. Box 55136-553, Maragheh, Iran*

²*Departamento de Física, CCEN - Universidade Federal da
Paraíba; C.P. 5008, CEP 58.051-970, João Pessoa, PB, Brasil*

³*Department of Mathematics, COMSATS University Islamabad, Lahore-Campus, Lahore, 54000, Pakistan*

Considering exact spacetimes representing rotating black holes and naked singularities, we study the possibility that the Sagnac effect detects *i*) higher dimensions, *ii*) rotation of black holes in higher dimension, and also, *iii*) distinguishes black holes from naked singularities. The results indicate that the Sagnac time delay gets affected by the presence of extra-dimension or its associated angular momentum. This time delay is also different in the spacetime of a naked singularity compared to that of a black hole. Hence, the Sagnac effect may be used as an experiment for better understanding of spacetimes with higher dimensions or those that admit naked singularities.

I. INTRODUCTION

In order to study the so-called Aether, various experiments such as the well-known Michelson-Morley experiment, and the Sagnac experiment have been proposed, and employed [1–3]. Circular relative motion is the backbone of the Sagnac effect which gets along well with the ideas of testing modified gravities and detecting a variety of phenomena predicted in general relativity (GR) [4–12]. Indeed, this experiment is a gravitational counterpart of the Aharonov-Bohm effect [13], an achievement which helps us find a simple way to calculate the Sagnac time delay [5, 6, 13].

The problem of the number of spacetime dimensions has attracted much attention during the past decades and is still a notable challenge in modern physics [14] and specially in black hole physics [15]; the dimensions number and also the probable rotation of black hole in other dimensions affect the spacetime geometry. In this regard, the Kerr metric or Kerr geometry describing the geometry of empty spacetime around a rotating uncharged axially-symmetric black hole is a widely used metric in the literature [16]. This rich solution has inspired physicists to seek for different types of rotating black holes like BTZ [17], and Myers-Perry ones [18]. These solutions clearly prove the implications of dimensions other than four on the properties of black hole, motivating us to study the Sagnac experiment as a tool to examine physical properties of such solutions. This indeed helps us finding a way to verify the existence of other dimensions.

Our present understanding of black holes in the universe implies that these compact objects in the universe are formed as the end state of a typical gravitational collapse of matter. From a theoretical perspective, a direct consequence of the use of GR reveals that, a continual

gravitational collapse always leads to spacetime singularities necessarily hidden by an event horizon, according to cosmic censorship hypothesis (CCH) [19]. Hence, no signal emanated from the neighborhood of singular region can reach distant observers. However, neither a conclusive proof or disproof nor a concrete mathematical formulation of CCH has been provided up until now leaving this hypothesis as one of the most outstanding unresolved problems in GR. On the other hand, there are great debates on possible inevitability of the spacetime singularities [30]. In this regard, recent studies of collapse scenarios show that GR field equations also admit naked singularities, depending on the initial conditions and different matter configurations [20, 21]. In this line, the possibility of distinguishing black holes from naked singularities is an attractive challenge [23–30].

Here, we are initially going to address the possibility that the Sagnac experiment detects the number of spacetime dimensions as well as the rotation of black holes in other dimensions. This is the task of the subsequent section. The propose of the third section is to show that the Sagnac experiment also can distinguish black holes from naked singularities. We set the units so that $c = G = 1$.

II. SAGNAC TIME DELAY OF ROTATING BLACK HOLES

Our aim in the present section is to calculate the Sagnac time delay for two well-known black hole solutions. Considering the results obtained in [6], we proceed with the following steps

- Without loss of generality we work within equatorial plane by setting $\theta = \pi/2$. Then, we make coordinate transformation on azimuthal coordinate: $\phi \rightarrow \phi + \omega_0 t$, where ω_0 is the rotational velocity of the source of signals and t is the proper time as measured by a clock rotating with constant angular velocity ω_0 [4, 6, 7].
- Define four-vector gravitomagnetic potential as: $A_\mu = -g_{\mu 0}/g_{00}$, and

*ah.ziaie@maragheh.ac.ir

†hn.moradpour@maragheh.ac.ir

‡valdir@fisica.ufpb.br

§abduljawad@cuilahore.edu.pk

- Perform the integration

$$\Delta t = 2\sqrt{-g_{00}} \int_0^{2\pi} A_\phi d\phi, \quad (1)$$

in order to evaluate the Sagnac time delay.

Now, we consider a satellite orbiting in a circular orbit of radius $R = \text{constant}$ with constant angular velocity ω_0 around the gravitating body [6, 31]. This satellite acts as a source/receiver of light beams propagating along circular orbits of the same radius around a source of gravitational field which here is taken as a Kerr black hole, see Fig. (1). To calculate the Sagnac time delay for this black hole, we begin with its line element given by

$$\begin{aligned} ds^2 = & - \left[1 - \frac{2Mr}{r^2 + a^2 \cos^2 \theta} \right] dt^2 + \frac{r^2 + a^2 \cos^2 \theta}{a^2 - 2Mr + r^2} dr^2 \\ & + (r^2 + a^2 \cos^2 \theta) d\theta^2 - \frac{4Mra \sin^2 \theta}{r^2 + a^2 \cos^2 \theta} dt d\phi \\ & + \left[r^2 + a^2 + \frac{2Mra^2 \sin^2 \theta}{r^2 + a^2 \cos^2 \theta} \right] d\phi^2, \end{aligned} \quad (2)$$

where $a = J/M$ is the angular momentum per black hole mass. A straightforward calculation then reveals that the Sagnac time delay for a Kerr black hole is obtained as

$$\Delta t_K = \frac{4\pi [\omega_0 (Ra^2 + 2Ma^2 + R^3) - 2aM]}{R \left[1 - \frac{2M}{R} + \frac{4aM\omega_0}{R} - \omega_0^2 \left(\frac{2Ma^2}{R} + a^2 + R^2 \right) \right]^{\frac{1}{2}}}. \quad (3)$$

This time difference, as measured by a sensitive clock which is carried by the satellite corresponds to the round trip time for a co-propagating light beam traveling along a circular orbit. For a counter-propagating beam we must set $a \rightarrow -a$. It can be also verified that in the limit $M \rightarrow 0$, $a \rightarrow 0$, the Sagnac time delay reduces to its counterpart in 4-dimensional Minkowski spacetime. We note that the condition on Δt_K to be a real and positive number requires that the numerator of expression (3) be positive and its denominator assumes real numbers. We further note that the denominator of expression (3) has two roots given by

$$\begin{aligned} a_- &= \frac{2m\omega_0 - \sqrt{R^2\omega_0^2 - 2mR^3\omega_0^4 - R^4\omega_0^4}}{2m\omega_0^2 + R\omega_0^2}, \\ a_+ &= \frac{2m\omega_0 + \sqrt{R^2\omega_0^2 - 2mR^3\omega_0^4 - R^4\omega_0^4}}{2m\omega_0^2 + R\omega_0^2}. \end{aligned} \quad (4)$$

Hence for Δt_K to be finite we require that the black hole angular momentum satisfies the condition $a_- < a < a_+$. In Fig. (2) we have sketched the allowed regions for mass and angular momentum satisfying the above conditions. The upper and lower panels present the allowed regions for counter and co-propagating beams, respectively. Figure (3) presents the behavior of Δt_K against black hole mass. We observe that for co-propagating beams, as the black hole angular momentum increases the Sagnac time




FIG. 1: The time difference as measured by a clock rotating around the black hole with constant angular velocity ω_0 between the emission and absorption of the co-propagating (+) and counter-propagating (-) light beams (red circular orbit). The black point represents a satellite carrying the source/receiver.

delay decreases and such a decrement depends on black hole mass. In fact, the rotation of black hole assists shortening of the light path as a result of frame dragging effect. For counter-propagating beams, the faster the black hole rotates, the longer it takes for the beam to complete its orbit. In the limit $a \rightarrow 0$ we still have non-vanishing time delay which is due to nonzero rotational velocity of the source, however, the more the black hole weight, the greater the Sagnac time delay.

Next, we proceed to calculate the Sagnac time delay for a 5-dimensional black hole known as Myers-Perry black hole whose line element is given by

$$\begin{aligned} ds^2 = & -dt^2 + \frac{M (dt + a \sin^2 \theta d\phi + b \cos^2 \theta d\psi)^2}{r^2 + a^2 \cos^2 \theta + b^2 \sin^2 \theta} \\ & + \frac{r^2 (r^2 + a^2 \cos^2 \theta + b^2 \sin^2 \theta)}{(r^2 + a^2)(r^2 + b^2) - Mr^2} dr^2 \\ & + (r^2 + a^2) \sin^2 \theta d\phi^2 + (r^2 + b^2) \cos^2 \theta d\psi^2 \\ & + (r^2 + a^2 \cos^2 \theta + b^2 \sin^2 \theta) d\theta^2, \end{aligned} \quad (5)$$

where b is the angular momentum of fifth-dimension. Similarly to the above procedure we get the Sagnac time delay as

$$\begin{aligned} \Delta t_{MP} = & \frac{4\pi}{(R^2 + b^2)^{\frac{1}{2}}} \times \\ & \frac{[\omega_0 (2Ma^2 + a^2b^2 + R^2b^2 + R^2a^2 + R^4) - 2aM]}{[R^2 + b^2 - 2M + \mathcal{G}]^{\frac{1}{2}}}, \end{aligned} \quad (6)$$




FIG. 2: The allowed regions of mass and angular momentum for $\omega_0 = 0.15$ and $R = 2.0$ (black region), $\omega_0 = 0.18$ and $R = 2.1$ (red region), $\omega_0 = 0.14$ and $R = 1.5$ (blue region) and $\omega_0 = 0.12$ and $R = 1.3$ (gray region). The upper (lower) panel stands for counter (co)-propagating beams.

where

$$\mathcal{G} = \omega_0 [4aM - \omega_0(2Ma^2 + a^2b^2 + R^2b^2 + R^2a^2 + R^4)]. \quad (7)$$

We note that in the limit $a = b \rightarrow 0$, $M \rightarrow 0$, the above expression reduces to Sagnac time delay in 5-dimensional Minkowski spacetime. At first glance we observe that even if we set $b = 0$, the Sagnac time delay of a Kerr black hole is different to that of Myers-Perry whose angular momentum in fifth dimension is zero. This means that even if there is no rotation in the fifth dimension, the round trip time for a 4-dimensional black hole, as given by Eq. (3) is different from a 5-dimensional black hole, due to the presence of extra dimension. This has been shown in Fig. (4) where we have plotted the two




FIG. 3: Behavior of Sagnac time delay versus angular momentum for ω_0 and R chosen from regions of Fig. (2) and different values of Kerr black hole mass. The family of solid curves represent the counter-propagating and those of dashed ones represent the co-propagating beams. For the model parameters we have set $\omega_0 = 0.15$, $R = 2.0$ and $M = 0.7$ (black curves), $\omega_0 = 0.14$, $R = 1.5$ and $M = 0.6$ (blue curves) and $\omega_0 = 0.12$, $R = 1.3$ and $M = 0.35$ (red curves).




FIG. 4: Behavior of Sagnac time delay for Kerr (dashed curve) and Myers-Perry (solid curve) black holes versus angular momentum of fourth-dimension. For the model parameters we have set $b = 0$, $\omega_0 = 0.15$, $R = 2.0$ and $M = 0.7$. The critical value of angular momentum is found as $a_{cr} = 0.4428$.

time delays against the angular momentum of the fourth-dimension. As we observe, both time delays are decreasing functions of angular momentum, however, for the case of Kerr black hole, the Sagnac time delay decreases at a higher rate, see dashed curve. There is a critical value for the angular momentum of fourth-dimension (a_{cr}) at which $\Delta t_K = \Delta t_{MP}$. For $a < a_{cr}$ we have $\Delta t_K > \Delta t_{MP}$ and vice versa. Also in the limit $a \rightarrow 0$ both time delays are non zero due to the angular velocity of the source/receiver but $\Delta t_{MP}(a = 0) < \Delta t_K(a = 0)$. Figure




FIG. 5: Behavior of Sagnac time delay for Kerr (dashed curve) and Myers-Perry (solid curves) black holes versus angular momentum of fourth-dimension. For the model parameters we have set $b = 0$ (black curve), $b = 1$ (blue curve), $b = 2$ (red curve), $b = 10$ (gray curve), $\omega_0 = 0.15$, $R = 2.0$ and $M = 0.7$.

(5) shows a comparison of Sagnac time delay for Kerr and Myers-Perry black holes for different values of angular momentum of fifth-dimension. Interestingly, we observe that increasing the b parameter shifts Δt_{MP} from decreasing to increasing function of angular momentum of the fourth-dimension. This means that the faster the black hole rotates in fifth dimension the more chance we may have to distinguish it, through the Sagnac effect, from a purely 4-dimensional Kerr black hole.

III. DISCRIMINATING BETWEEN BLACK HOLE AND NAKED SINGULARITIES

Our aim in the present section is to employ the Sagnac effect as a tool to distinguish between a black hole and naked singularity. Let us begin with the line element for a rotating naked singularity which in Boyer-Lindquist coordinates is given by [32]

$$\begin{aligned}
 ds^2 = & -\frac{\Delta - a^2 \sin^2 \theta}{\Sigma} dt^2 + \frac{\Sigma}{\Delta} dr^2 + \Sigma d\theta^2 \\
 & + \sin^2 \theta \left[\Sigma + \left(2 - \frac{\Delta - a^2 \sin^2 \theta}{\Sigma} \right) a^2 \sin^2 \theta \right] d\phi^2 \\
 & - 2 \left(1 - \frac{\Delta - a^2 \sin^2 \theta}{\Sigma} \right) a \sin^2 \theta dt d\phi, \quad (8)
 \end{aligned}$$

where

$$\begin{aligned}
 \Sigma = h^{1-\mu} \rho, \quad \Delta = r^2 + a^2 - \frac{2Mr}{\mu}, \quad \rho = r^2 + a^2 \cos^2 \theta, \\
 \mu = \frac{M}{\sqrt{M^2 + q^2}}, \quad h = 1 - \frac{2Mr}{\mu\rho}, \quad (9)
 \end{aligned}$$

and M and q are the Arnowitt-Deser-Misner (ADM) mass and scalar charge and are constant real param-

eters. The above metric is the rotating version of Janis-Newman-Winicour solution [33] and represents the space-time of a rotating globally naked singularity (for $0 \leq \mu < 1$) [34] in the presence of a massless scalar field whose radial dependence is given by

$$\Phi(r) = \frac{\sqrt{1-\mu^2}}{4} \ln \left(1 - \frac{2Mr}{\mu\rho} \right). \quad (10)$$

It can be easily seen that for $\mu = 1$ (or equivalently $q = 0$) the Kerr metric is recovered.

Now, assume that the source/receiver orbits with angular velocity ω_0 at constant radius $r = R$ in the plane $\sin \theta = k$. Simple calculations then give

$$ds^2 = \frac{k^2}{B} (Dd\phi - a dt)^2 - \frac{A}{B} (dt - ak^2 d\phi)^2, \quad (11)$$

where

$$A = \Delta \Big|_{r=R}, \quad B = \Sigma \Big|_{r=R}^{\sin \theta = k}, \quad D = (\Sigma + a^2 k^2) \Big|_{r=R}^{\theta=0}. \quad (12)$$

If we denote the angular velocity of light beam as Ω , then $\phi = \Omega t$. The angular velocity of light rays is then obtained as

$$\Omega_{\pm} = \frac{a \pm \frac{\sqrt{A}}{k}}{D \pm ak\sqrt{A}}, \quad (13)$$

where use has been made of the condition $ds^2 = 0$ for null geodesics. We note that the above result is valid within the planes for which $D \pm ak\sqrt{A} \neq 0$. During the orbital rotation of the light beams, the source/receiver rotates at angular distance ϕ_0 and consequently, when the two light beams reach the source/receiver they experience phase difference, depending on the relative motion of the light beams with respect to the source/receiver. Indeed, the light beam which is co(counter)-rotating with the source/receiver, will travel at angular distance $\phi_0 + 2\pi$ ($\phi_0 - 2\pi$). Thus, for the distances that the light beams travel we have

$$\frac{\Omega_{\pm}}{\omega_0} \phi_{0\pm} = \phi_{0\pm} \pm 2\pi, \quad (14)$$

whence we get

$$\phi_{0\pm} = \pm \frac{2\pi\omega_0}{\Omega_{\pm} - \omega_0}. \quad (15)$$

We note that $\phi_{0\pm}$ is the angular distance travelled with the light beam with angular velocity Ω_{\pm} . From Eq.(11) we get the following time difference for the source/receiver as

$$dt = \frac{d\phi}{\omega_0 \sqrt{B}} \left[k^2 (D\omega_0 - a)^2 - A (1 - ak^2 \omega_0)^2 \right]^{\frac{1}{2}}. \quad (16)$$

Finally, integrating from ϕ_{0-} to ϕ_{0+} along with utilizing expression (14) we can find the Sagnac time difference




FIG. 6: Behavior of Sagnac time delay for Kerr black hole (red curve) and a rotating naked singularity (black curves) versus angular momentum. For the model parameters we have set $\omega_0 = 0.15$, $R = 2.0$, $M = 0.7$ and $\mu = 0.95$ (dashed curve), $\mu = 0.85$ (dotted curve), $\mu = 0.75$ (long-dashed curve).

between the two light signals. For $k = 1$, the Sagnac time delay reads

$$\Delta t_{NS} = \frac{4\pi [\omega_0 (D^2 - a^2 A) - a(D - A)]}{Rh^{\frac{1-\mu}{2}} \sqrt{A(1 - a\omega_0)^2 - (D\omega_0 - a)^2}}. \quad (17)$$

It can be checked that for $\mu = 1$ the above relation reduces to the Sagnac time delay for a Kerr spacetime. In Fig. (6) we have compared the Sagnac time delay for a Kerr black hole and its naked singularity counterpart. As we observe, both time delays are decreasing functions of angular momentum, however, the time delay for naked singularity decreases more rapidly for smaller values of μ parameter. Thus, for a specific angular momentum, the lesser the value of μ parameter, the more chance we have to distinguish a naked singularity from a black hole with same angular momentum.

IV. SUMMARY AND CONCLUDING REMARKS

Motivated by the idea that if spacetime admits extra-dimensions, then its implications should appear in gravi-

tational measurements around black holes, in the present work, we employed the Kerr and the Myers-Perry metrics and studied the capability of Sagnac effect in recognizing possible existence of the fifth dimension as well as the rotation of black holes in this dimension. To this aim, we studied the interference process of light beams in a rotating frame in curved spacetime. The corresponding set up is based on an interferometer carried by a satellite that uniformly rotates in a circular orbit around the black hole. This satellite also carries a clock that measures the difference between propagation time of each of the co-propagating and counter-propagating beams [6, 31].

We then observed that even if black hole has no rotation in fifth dimension, the Sagnac time delay of a Kerr black hole will be different from that of Myers-Perry due to the mere presence of the fifth dimension. We also found that the change in angular momentum of the fifth dimension affects the Sagnac time delay so that it could be distinguished from a four-dimensional Kerr black hole. Although, our focus was on four and five dimensional cases, the present research can legally be extended to higher dimensions.

Moreover, using the Janis-Newman-Winicour metric which represents the spacetime of a rotating naked singularity, it is shown that the Sagnac effect may also be used to distinguish naked rotating singularities from a Kerr black hole. The outcomes of the present article confirm the power of Sagnac effect in studying possible existence of extra-dimensions and naked singularities in the Universe.

Acknowledgments

V.B.B. is partially supported by the National Council for Scientific and Technological Development – CNPq (Brazil), through the Research Project No. 307211/2020-7. This paper is published as part of a research project supported by the University of Maragheh Research Affairs Office.

[1] G. Sagnac, *Comptes Rendus* **129**, 818 (1899).
[2] G. Sagnac, *Comptes Rendus* **157**, 708 (1913).
[3] H. Fizeau, *Comptes Rendus* **33**, 349 (1851).
[4] A. Tartaglia, *Phys. Rev. D* **58**, 064009 (1998).
[5] G. Rizzi and M. L. Ruggiero, *Relativity in Rotating Frames. Fundamental Theories of Physics*, vol 135. Springer, Dordrecht.

[6] M. L. Ruggiero, *Nuovo Cim. B* **119**, 893 (2004).
[7] M. L. Ruggiero, *Galaxies* **3**, no.2, 84-102 (2015).
[8] I. Zeylikovich and J. Malinsky, *Class. Quant. Grav.* **28**, 155011 (2011).
[9] R. K. Karimov, R. N. Izmailov, A. A. Potapov and K. K. Nandi, *Gen. Rel. Grav.* **50**, 44 (2018).
[10] E. Benedetto, F. Feleppa, I. Licata, H. Moradpour and

- C. Corda, Eur. Phys. J. C **79**, 187 (2019).
- [11] F. Feleppa, H. Moradpour, C. Corda and S. Aghababaei, EPL (2021). doi:10.1209/0295-5075/ac1240,
- [12] E. Benedetto, F. Feleppa, G. Iovane and E. Laserra, Int. J. Geo. Meth. Mod. Phys. **17**, 2050128 (2020).
- [13] J. J. Sakurai, Phys. Rev. D **21**, 2993 (1980).
- [14] M. Corman et al., arXiv:2109.08748.
- [15] R. Emparan and H. S. Reall, Living Rev. Relativ. **11**, 6 (2008).
- [16] S. A. Teukolsky, Class. Quantum Grav. **32**, 124006 (2015).
- [17] S. Carlip, Class. Quantum Grav. **12**, 2853 (1995).
- [18] R. C. Myers and M. J. Perry, Annals Phys. **172**, 304 (1986).
- [19] R. Penrose, Riv. Nuovo Cimento **1**, 252 (1969); [Gen.Rel. Grav. **34**, 1141 (2002)];
R. Penrose, in *General relativity, An Einstein Centenary Survey*, edited by S. W. Hawking and W. Israel (Cambridge University Press, Cambridge, England, 1973).
- [20] P. S. Joshi, *Gravitational Collapse and Spacetime Singularities*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, 2012).
- [21] S. L. Shapiro and S. A. Teukolsky, Phys. Rev. L **66**, 994 (1991)
- [22] S. W. Hawking, R. Penrose, *The Nature of Space and Time*. Princeton University Press, (2010).
- [23] P. S. Joshi, I. H. Dwivedi, Phys. Rev. D **47**, 5357 (1993).
- [24] K. S. Virbhadra, G. F. R. Ellis, Phys. Rev. D **65**, 103004 (2002).
- [25] G. N. Gyulchev, S. S. Yazadjiev, Phys. Rev. D. **78**, 083004 (2008).
- [26] P. S. Joshi, Scientific American, **300**, 36 (2009).
- [27] P. S. Joshi, D. Malafarina, R. Narayan, Class. Quant. Grav. **31**, 015002 (2014).
- [28] H. Liu, M. Zhou, C. Bambi, JCAP **08**, 044 (2018).
- [29] R. Shaikh, P. S. Joshi, JCAP **10**, 064 (2019).
- [30] H. Zhang, Sci Rep **7**, 4000 (2017).
- [31] A. Tartaglia, Advances in Space Research, **66**, 2757 (2020).
- [32] G. N. Gyulchev, S. S. Yazadjiev, Phys. Rev. D. **78**, (2008) 083004.
- [33] K. S. Virbhadra, Int. J. Mod. Phys. A **12**, 4831 (1997).
- [34] K. S. Virbhadra, S. Jhingan and P. S. Joshi, Int. J. Mod. Phys. D **6**, 357 (1997).