

The X-ray corona turns into the relativistic jet in the micro quasar GRS 1915+105

Mariano Méndez^{*,1}, Konstantinos Karpouzas^{1,2}, Federico García^{1,3}, Liang Zhang², Yuexin Zhang¹,
Tomaso M. Belloni⁴ & Diego Altamirano²

¹*Kapteyn Astronomical Institute, University of Groningen, Postbus 800, 9700 AV Groningen, The Netherlands*

²*School of Physics and Astronomy, University of Southampton, Southampton, SO17 1BJ, UK*

³*Instituto Argentino de Radioastronomía (CCT La Plata, CONICET; CICPBA; UNLP), C.C.5, (1894) Villa Elisa, Buenos Aires, Argentina*

⁴*INAF - Osservatorio Astronomico di Brera, Via E. Bianchi 46, I-23807 Merate, Italy*

1 **GRS 1915+105¹ was the first stellar-mass black-hole in our Galaxy to display a superluminal**
2 **radio jet², similar to those observed in active galactic nuclei with a supermassive black hole**
3 **at the centre³. It has been proposed that the radio emission in GRS 1915+105 is fed by insta-**
4 **bilities in the accretion disc⁴ by which the inner parts of the accretion flow is ejected in the**
5 **jet⁵⁻⁷. Here we show that there is a significant correlation between: (i) the radio flux, coming**
6 **from the jet, and the flux of the iron emission line, coming from the disc and, (ii) the tem-**
7 **perature of the corona that produces the high-energy part of the X-ray spectrum via inverse**
8 **Compton scattering and the amplitude of a high-frequency variability component coming**
9 **from the innermost part of the accretion flow. At the same time, the radio flux and the flux of**
10 **the iron line are strongly anti-correlated with the temperature of the X-ray corona and the**

11 **amplitude of the high-frequency variability component. These correlations persist over ~ 10**
12 **years, despite the highly variable X-ray and radio properties of the source in that period^{8,9}.**
13 **Our findings provide, for the first time, incontrovertible evidence that the energy that powers**
14 **this black-hole system can be directed either to the X-ray corona or the jet. When this energy**
15 **is used to power the corona, raising its temperature, there is less energy left to fuel the jet**
16 **and the radio flux drops, and vice versa. These facts, plus the modelling of the variability in**
17 **this source show conclusively that in GRS 1915+105 the X-ray corona morphs into the jet.**

18

19 The X-ray spectrum of black-hole binaries can be decomposed into three main radiation
20 components: (i) A thermal component that dominates the emission at low (soft) energies¹⁰, below
21 ~ 5 keV, due to an accretion disc through which mass flows from the binary companion to the black
22 hole. (ii) A power-law like component that dominates the spectrum at energies above ~ 5 – 10
23 keV, caused by inverse-Compton scattering of the disc photons in a corona of highly-energetic
24 electrons¹¹. If the energy distribution of the electrons is Maxwellian, this component features a
25 high-energy cut off at an energy that depends upon the electron temperature of the corona, kT_e .
26 (iii) A broad emission line at ~ 6.5 – 7 keV due to iron, produced when photons from the corona
27 reflect off the disc, with the line profile being set by special- and general-relativistic effects^{12,13}.
28 The X-ray emission of accreting black-hole binaries exhibits high-amplitude variability from tens
29 of milliseconds to decades^{8,14}. Depending on the relative importance of these spectral components
30 and the strength of the variability, accreting black holes display different states¹⁵: In the hard states
31 the X-ray spectrum (~ 1 – 20 keV range) is dominated by emission from the corona, and the Fourier

32 power spectrum shows variability of up to 50% of the average luminosity over a broad range of time
33 scales plus narrow quasi-periodic oscillations, QPOs. In the soft states the emission is dominated
34 by the accretion disc, and the variability drops to less than 5% of the average luminosity.

35 Black-hole binaries in the hard states emit in radio with a spectrum that is consistent with
36 self-absorbed synchrotron radiation from an optically thick and compact jet¹⁶. During the transition
37 from the hard to the soft states, some black-hole binaries show radio emission from individual,
38 spatially-resolved, plasma clouds that are ejected in a jet at speeds close to the speed of light. The
39 radio spectrum of these discrete ejections is consistent with synchrotron emission from optically-
40 thin material².

41 GRS 1915+105 harbours a $12^{+2.0}_{-1.8}$ solar-mass black hole¹⁷ and is very variable both in X-rays
42 and radio wavelengths^{8,9}. In X-rays, the emission switches from times in which a bright accre-
43 tion disc with a temperature $kT_{\text{bb}} \approx 2$ keV and small inner-radius dominates the spectrum, to times
44 in which the corona dominates the spectrum, the disc is relatively cool, $kT_{\text{bb}} \approx 0.5-1$ keV and is
45 inferred to have a large inner radius⁴. In several observations, mostly those with a cool disc, in ad-
46 dition to a band-limited noise component, a narrow and strong QPO, called type-C QPO¹⁸, appears
47 in the Fourier power spectrum at frequencies between ~ 0.4 Hz and 6.5 Hz¹⁹. The frequency of
48 the QPO increases as the temperature of the disc increases and the spectrum of the source softens.
49 Besides this QPO (and harmonics and sub-harmonics of the fundamental frequency), the power
50 spectrum of GRS 1915+105 sometimes shows a broad variability component at $\sim 60-80$ Hz²⁰ that
51 we will call the high-frequency bump. The high-frequency bump appears when the spectrum of the

52 source is dominated by the corona, but observations in which the corona dominates the emission
53 do not always show this high-frequency bump. At the same time, observations of GRS 1915+105
54 in which the spectrum is dominated by the corona are sometimes, but not always, accompanied by
55 high radio fluxes^{21,22}. (We show the power-density spectra of two observations of GRS 1915+105
56 with the QPOs and the bump indicated in the Extended Data Figure 5.)

57 We studied a large dataset of 1800+ X-ray observations of GRS 1915+105 obtained with
58 the Rossi X-ray Timing Explorer (RXTE) between 1996 and 2012, combined with almost daily
59 observations of the source at 15 GHz with the Ryle telescope. Our final sample consists of 410
60 observations for which we have simultaneous data both in X-rays and radio. For each of these ob-
61 servations we have a measurement of (i) the radio flux density at 15 GHz, (ii) the X-ray hardness
62 ratio calculated as the ratio of the intensity in the 13–60 keV to the 2–7 keV band, (iii) the fre-
63 quency of the fundamental component of the type-C QPO, (iv) the phase lag at the QPO frequency
64 for photons in the 5.7–15 keV band with respect to those in the 2–5.7 keV band, (v) the 2–60
65 keV fractional rms amplitude of the high-frequency bump, and (vi) the best-fitting parameters to
66 the X-ray energy spectra of the source. (See the section Methods for details of the analysis and an
67 explanation of some of these quantities). These vastly different types of data, consisting of X-ray
68 and radio fluxes and spectral and timing properties of a single source, come from wavelengths that
69 are more than eight orders of magnitude apart and sample time scales that are more than eleven
70 orders of magnitude different, from ten milliseconds to a decade.

71 In Figure 1 we plot the X-ray hardness ratio as a function of the frequency of the QPO for

72 these 410 observations of GRS 1915+105. The x and y axes of the four panels in this Figure are
73 the same, while the colours of the points in each panel represent, respectively, the simultaneous 15-
74 GHz radio-flux measurements from the jet (Fig. 1a), the electron temperature of the X-ray corona
75 (Fig. 1b), the flux of the iron emission line in the X-ray spectrum (Fig. 1c), and the fractional rms
76 amplitude of the high-frequency bump (Fig. 1d). In all four panels blue (red) indicates high (low)
77 values of the quantity represented by the colours.

78 The QPO frequency generally increases as the hardness ratio decreases and the source spec-
79 trum softens, consistent with a decreasing inner radius of the accretion disc that leads to an increase
80 of both the disc flux¹⁰ and QPO frequency^{23,24}. The relation, however, is significantly broader than
81 the spread expected from the errors in each quantity. (The errors are smaller than the size of the
82 points.)

83 The colours of the points in the four panels of this Figure show that the radio flux, the flux of
84 the iron line, the temperature of the corona and the fractional rms amplitude of the high-frequency
85 bump depend upon both QPO frequency and hardness ratio. This double dependence is the reason
86 for the striking match between the two panels on the left and the two on the right, and the almost
87 perfect mirror match between the two top and the two bottom panels. In other words, the breadth
88 of the hardness-ratio/QPO-frequency relation is consistently set by either of the four quantities,
89 such that if we plotted the data in a diagram with the QPO frequency along the x axis, the hardness
90 ratio along the y axis and any of these four quantities along the z axis, the points would lie on a
91 2-dimensional surface in 3D.

92 The two panels of Figure 2 show the flux of the iron line as a function of the X-ray flux
 93 in the 2–25-keV band for all observations of GRS 1915+105. In the left panel the colour of the
 94 points represents the temperature of the corona, while in the right panel the colours represent the
 95 phase lags between the 2–5.7 keV and 5.7–15 keV bands at the frequency of the QPO. While the
 96 flux of the iron line increases as the total flux of the source increases, when the temperature of the
 97 corona is low, the radio flux is high and the lags at the QPO frequency are positive the correlation²⁵
 98 is steeper than when the temperature is high, the radio flux is low and the lags are negative. To
 99 confirm that our measurements of the line flux are not biased because of the relatively low spectral
 100 resolution of the RXTE/PCA instrument, we also include in this Figure independent measurements
 101 of the flux of the line with Chandra and Suzaku taken from the literature^{25,26} (black plus and cross
 102 symbols).

103 These results provide a unique clue to understand the nature of the corona and the jet in this
 104 object, and what powers these components. Inverse Compton scattering cools down the corona by
 105 transferring energy from the electrons to the soft disc photons. The temperature of the corona in
 106 black-hole binaries, however, increases during periods in which the photon flux of the disc drops
 107 and Compton cooling is less effective. This means that a source of power balances the inverse
 108 Compton cooling and sets the temperature of the corona. Our findings show that in GRS 1915+105
 109 the energy provided by this mechanism is split to either power the jet or heat the corona.

110 The thermal energy stored in a spherical corona of optical depth τ and size L around a black
 111 hole of mass M is $E_{th} \approx 3.4 \times 10^{26} \tau \left(\frac{kT_e}{1 \text{ keV}} \right) \left(\frac{M}{M_\odot} \right)^2 \left(\frac{L}{R_g} \right)^2$ erg, where $R_g = GM/c^2$ is the

112 gravitational radius of the black hole and G and c are, respectively, the gravitational constant and
113 the speed of light²⁷.

114 As discussed in the section Extended Results and Discussion, the change of sign of the lags
115 is explained if a fraction of the disc photons that are Comptonised in the corona impinges back
116 onto the disc before reaching the observer²⁸. When this feedback fraction is low, time delays due
117 to Comptonisation dominate and the lags are positive; when this fraction is high, reprocessed disc
118 photons reach the observer later than those from the corona and the lags become negative. Taking
119 the corona sizes from fits with this model²⁹, $L \approx 10-1200R_g$, plus $\tau \approx 1-6$ and $kT_e \approx 5-40$ keV
120 from the spectral fits, we find that $E_{th} \approx 10^{31}-10^{35}$ erg. If this energy is released over the time
121 scale of the high-frequency bump¹, the thermal luminosity is 2 to 5 orders of magnitude lower
122 than the observed luminosity of the corona in GRS 1915+105. The alternative is that the corona
123 is powered by magnetic energy^{27,30}, e.g., shear energy due to differential rotation of the magnetic-
124 field lines that thread the accretion disc. This magnetic energy would also be responsible for the
125 synchrotron radio emission and the jet ejection mechanism in black-hole binaries.

126 We propose that in GRS 1915+105 the corona turns into the jet and that both are, at different
127 times, the same physical component. Based on the results shown here and fits with the model of
128 the lags that we present in the section Extended Results and Discussion, the process proceeds as
129 follows: (1) When the QPO frequency is ~ 6 Hz the corona is relatively large and enshrouds the
130 inner parts of the accretion disc (Fig. 3a); as seen from the corona, the disc covers half of the sky

¹Note that this is the shortest variability time scale in the data; using any other time scale longer than this one to estimate the thermal luminosity would make the discrepancy even bigger.

131 and hence there is a high probability that photons from the corona return to the disc leading to a
132 high feedback fraction and negative QPO lags. As the QPO frequency decreases from ~ 6 Hz to
133 ~ 2 Hz, the inner edge of the disc moves outwards and the magnitude of the lags and the size of
134 the corona decrease. In this phase the magnetic field that threads the disc and powers the corona
135 is relatively disorganised, the magnetic energy is dissipated stochastically leading to a high corona
136 temperature. (2) At a QPO frequency² of ~ 2 Hz the size of the corona becomes equal to the
137 truncation radius of the accretion disc, the feedback fraction decreases abruptly and the QPO lags
138 become zero. From this point on the corona moves inside the accretion disc (Fig. 3b), the magnetic
139 field starts to become coherent on longer spatial scales, the stochastic energy dissipation decreases
140 and the temperature of the corona drops. At the same time the magnetic-field lines start to channel
141 material from the corona into the direction perpendicular to the accretion disc, and low-level radio
142 emission from the jet starts to appear. (3) As the QPO frequency continues decreasing below ~ 2
143 Hz the inner disc radius moves further out and the size of the corona increases again but, given
144 that the lags are positive, this time the feedback fraction remains low. From this it follows that
145 the corona does not cover the inner parts of the disc, which can only be the case if the geometry
146 of the corona changes such that it becomes more prominent in the direction perpendicular to the
147 disc (Fig. 3c); as seen from the corona, the disc now covers a much smaller area of the sky. As the
148 magnetic-field lines become more spatially coherent³¹ there is less stochastic energy dissipation,
149 the corona temperature drops further and the material from the corona that was channeled off the
150 originally extended corona is expelled away from the accretion-disc plane and becomes the radio

²The transition occurs when the lags for the QPO change from negative to positive; although the exact value of the QPO frequency¹⁹ at which this transition occurs is between 1.8 and 2 Hz, here we will always write 2 Hz for simplicity.

151 jet (Fig. 3d).

152 Our multi-wavelength correlations match the proposal that, during the initial parts of an
153 outburst, the X-ray corona of the black-hole binary MAXI J1820+070 contracts³² and then re-
154 expands³³. Here we show conclusively that, as previously speculated^{7,34,35}, in the case of GRS
155 1915+105 the contracting corona becomes the radio jet and that, at least part of the time, the
156 corona and the jet are actually one and the same physical component^{36–38}.

157 The appearance of radio flares when the QPO frequency is below 2 Hz reinforces our inter-
158 pretation. Figure 4 shows that in the periods that the frequency of the QPO moves more or less
159 stochastically between ~ 2 Hz and ~ 8 Hz (red circles) the radio flux (light blue curve) is low. Oc-
160 casionally, the QPO frequency evolves in a more systematic way: it starts to decrease, crosses the
161 value of 2 Hz and moves up again; at the same time the radio flux increases sharply and a radio
162 flare lasting a few tens of days is observed. Figure 4 shows this behaviour over a period of about
163 500 days in which the source shows two radio flares. Extended Data Figure 9 shows that the same
164 behaviour repeats consistently over a period of 10 years and about a dozen radio flares.

165 The relation between the iron-line flux and the total flux is consistent with the above scenario.
166 The flux of the iron line depends more strongly upon X-ray flux when the temperature of the corona
167 is low, that is when the corona has turned into the jet²⁵, than when the corona is more extended and
168 covers the inner parts of the accretion disc (Fig. 2) . This points to a lamp-post geometry³⁹ with
169 the corona, which is now the jet, illuminating the disc anisotropically^{40,41}. Because the jet does not
170 cover the accretion disc, the flux of the iron line is not (or mildly) attenuated by the corona^{42,43}.

171 The behaviour we observe in GRS 1915+105 could explain the deviations from a single
172 track in the radio/X-ray correlation of other accreting galactic black holes^{44–49} If, as in the case of
173 GRS 1915+105, the energy powering these systems is used to either accelerate the jet or heat the
174 corona, different sources, or the same source at different times, will show lower radio fluxes at a
175 given X-ray flux (or, equivalently, higher X-ray fluxes at a given radio flux), depending on how
176 much energy is directed towards, respectively, launching the radio jet or heating the X-ray corona.

References

1. Fender, R. & Belloni, T. GRS 1915+105 and the Disc-Jet Coupling in Accreting Black Hole Systems. *Annu. Rev. Astron. Astrophys.* **42**, 317–364 (2004).
2. Mirabel, I. F. & Rodríguez, L. F. A superluminal source in the Galaxy. *Nature* **371**, 46–48 (1994).
3. Pearson, T. J. *et al.* Superluminal expansion of quasar 3C273. *Nature* **290**, 365–368 (1981).
4. Belloni, T., Méndez, M., King, A. R., van der Klis, M. & van Paradijs, J. A Unified Model for the Spectral Variability in GRS 1915+105. *Astrophys. J.* **488**, L109–L112 (1997).
5. Mirabel, I. F. *et al.* Accretion instabilities and jet formation in GRS 1915+105. *Astron. Astrophys.* **330**, L9–L12 (1998).
6. Eikenberry, S. S., Matthews, K., Morgan, E. H., Remillard, R. A. & Nelson, R. W. Evidence for a Disk-Jet Interaction in the Microquasar GRS 1915+105. *Astrophys. J.* **494**, L61–L64 (1998).

7. Fender, R. P., Belloni, T. M. & Gallo, E. Towards a unified model for black hole X-ray binary jets. *Mon. Not. R. Astron. Soc.* **355**, 1105–1118 (2004).
8. Belloni, T., Klein-Wolt, M., Méndez, M., van der Klis, M. & van Paradijs, J. A model-independent analysis of the variability of GRS 1915+105. *Astron. Astrophys.* **355**, 271–290 (2000).
9. Pooley, G. G. & Fender, R. P. The variable radio emission from GRS 1915+105. *Mon. Not. R. Astron. Soc.* **292**, 925–933 (1997).
10. Shakura, N. I. & Sunyaev, R. A. Reprint of 1973A&A....24..337S. Black holes in binary systems. Observational appearance. *Astron. Astrophys.* **500**, 33–51 (1973).
11. Sunyaev, R. A. & Titarchuk, L. G. Comptonization of X-rays in plasma clouds. Typical radiation spectra. *Astron. Astrophys.* **500**, 167–184 (1980).
12. Fabian, A. C., Rees, M. J., Stella, L. & White, N. E. X-ray fluorescence from the inner disc in Cygnus X-1. *Mon. Not. R. Astron. Soc.* **238**, 729–736 (1989).
13. Fabian, A. C. *et al.* Broad line emission from iron K- and L-shell transitions in the active galaxy 1H0707-495. *Nature* **459**, 540–542 (2009).
14. Greiner, J., Morgan, E. H. & Remillard, R. A. Rossi X-Ray Timing Explorer Observations of GRS 1915+105. *Astrophys. J.* **473**, L107 (1996).
15. Méndez, M. & van der Klis, M. The EXOSAT Data on GX 339-4: Further Evidence for an “Intermediate” State. *Astrophys. J.* **479**, 926–932 (1997).

16. Fender, R. P., Spencer, R. E., Newell, S. J. & Tzioumis, A. K. High-resolution radio observations of the black hole candidate GX 339-4. *Mon. Not. R. Astron. Soc.* **286**, L29–L32 (1997).
17. Reid, M. J. *et al.* A Parallax Distance to the Microquasar GRS 1915+105 and a Revised Estimate of its Black Hole Mass. *Astrophys. J.* **796**, 2 (2014).
18. Casella, P., Belloni, T. & Stella, L. The ABC of Low-Frequency Quasi-periodic Oscillations in Black Hole Candidates: Analogies with Z Sources. *Astrophys. J.* **629**, 403–407 (2005).
19. Zhang, L. *et al.* A systematic analysis of the phase lags associated with the type-C quasi-periodic oscillation in GRS 1915+105. *Mon. Not. R. Astron. Soc.* **494**, 1375–1386 (2020).
20. Trudolyubov, S. P. On the Two Types of Steady Hard X-Ray States of GRS 1915+105. *Astrophys. J.* **558**, 276–282 (2001).
21. Munro, M. P., Morgan, E. H. & Remillard, R. A. Quasi-periodic Oscillations and Spectral States in GRS 1915+105. *Astrophys. J.* **527**, 321–340 (1999).
22. Fender, R. P. *et al.* MERLIN observations of relativistic ejections from GRS 1915+105. *Mon. Not. R. Astron. Soc.* **304**, 865–876 (1999).
23. Stella, L. & Vietri, M. Lense-Thirring Precession and Quasi-periodic Oscillations in Low-Mass X-Ray Binaries. *Astrophys. J.* **492**, L59–L62 (1998).
24. Ingram, A., Done, C. & Fragile, P. C. Low-frequency quasi-periodic oscillations spectra and Lense-Thirring precession. *Mon. Not. R. Astron. Soc.* **397**, L101–L105 (2009).

25. Neilsen, J. & Lee, J. C. Accretion disk winds as the jet suppression mechanism in the microquasar GRS 1915+105. *Nature* **458**, 481–484 (2009).
26. Mizumoto, M., Ebisawa, K., Tsujimoto, M. & Inoue, H. Origin of the X-ray broad iron spectral feature in GRS 1915+105. *Publ. astr. Soc. Japan* **68**, S16 (June 2016).
27. Merloni, A. & Fabian, A. C. Accretion disc coronae as magnetic reservoirs. *Mon. Not. R. Astron. Soc.* **321**, 549–552 (2001).
28. Karpouzas, K. *et al.* The Comptonizing medium of the neutron star in 4U 1636 - 53 through its lower kilohertz quasi-periodic oscillations. *Mon. Not. R. Astron. Soc.* **492**, 1399–1415 (2020).
29. Karpouzas, K. *et al.* A variable corona for GRS 1915+105. *Mon. Not. R. Astron. Soc.* **503**, 5522–5533 (2021).
30. Malzac, J., Merloni, A. & Fabian, A. C. Jet-disc coupling through a common energy reservoir in the black hole XTE J1118+480. *Mon. Not. R. Astron. Soc.* **351**, 253–264 (2004).
31. Meier, D. L. Magnetically Dominated Accretion Flows (MDAFS) and Jet Production in the Low hard State. *Ap&SS* **300**, 55–65 (2005).
32. Kara, E. *et al.* The corona contracts in a black-hole transient. *Nature* **565**, 198–201 (2019).
33. Wang, J. *et al.* Disk, Corona, Jet Connection in the Intermediate State of MAXI J1820+070 Revealed by NICER Spectral-timing Analysis. *Astrophys. J.* **910**, L3 (2021).
34. Vadawale, S. V. *et al.* On the Origin of the Various Types of Radio Emission in GRS 1915+105. *Astrophys. J.* **597**, 1023–1035 (2003).

35. Rodriguez, J., Corbel, S. & Tomsick, J. A. Spectral Evolution of the Microquasar XTE J1550-564 over Its Entire 2000 Outburst. *Astrophys. J.* **595**, 1032–1038 (2003).
36. Levinson, A. & Blandford, R. On the Jets Associated with Galactic Superluminal Sources. *Astrophys. J.* **456**, L29 (1996).
37. Giannios, D., Kylafis, N. D. & Psaltis, D. Spectra and time variability of Galactic black-hole X-ray sources in the low/hard state. *Astron. Astrophys.* **425**, 163–169 (2004).
38. Markoff, S., Nowak, M. A. & Wilms, J. Going with the Flow: Can the Base of Jets Subsume the Role of Compact Accretion Disk Coronae? *Astrophys. J.* **635**, 1203–1216 (2005).
39. Miniutti, G. & Fabian, A. C. A light bending model for the X-ray temporal and spectral properties of accreting black holes. *Mon. Not. R. Astron. Soc.* **349**, 1435–1448 (2004).
40. Kylafis, N. D., Reig, P. & Papadakis, I. A quantitative explanation of the type-B QPOs in GX 339-4. *Astron. Astrophys.* **640**, L16 (2020).
41. Reig, P. & Kylafis, N. D. Illumination of the accretion disk in black hole binaries: An extended jet as the primary source of hard X-rays. *Astron. Astrophys.* **646**, A112 (2021).
42. Matt, G., Fabian, A. C. & Reynolds, C. S. Geometrical and chemical dependence of K-shell X-ray features. *Mon. Not. R. Astron. Soc.* **289**, 175–184 (1997).
43. Petrucci, P. O., Merloni, A., Fabian, A., Haardt, F. & Gallo, E. The effects of a Comptonizing corona on the appearance of the reflection components in accreting black hole spectra. *Mon. Not. R. Astron. Soc.* **328**, 501–510 (2001).

44. Hannikainen, D. C., Hunstead, R. W., Campbell-Wilson, D. & Sood, R. K. MOST radio monitoring of GX 339-4. *Astron. Astrophys.* **337**, 460–464 (1998).
45. Corbel, S., Nowak, M. A., Fender, R. P., Tzioumis, A. K. & Markoff, S. Radio/X-ray correlation in the low/hard state of GX 339-4. *Astron. Astrophys.* **400**, 1007–1012 (2003).
46. Gallo, E., Fender, R. P. & Pooley, G. G. A universal radio-X-ray correlation in low/hard state black hole binaries. *Mon. Not. R. Astron. Soc.* **344**, 60–72 (2003).
47. Coriat, M. *et al.* Radiatively efficient accreting black holes in the hard state: the case study of H1743-322. *Mon. Not. R. Astron. Soc.* **414**, 677–690 (2011).
48. Gallo, E., Miller, B. P. & Fender, R. Assessing luminosity correlations via cluster analysis: evidence for dual tracks in the radio/X-ray domain of black hole X-ray binaries. *Mon. Not. R. Astron. Soc.* **423**, 590–599 (2012).
49. Gallo, E., Degenaar, N. & van den Eijnden, J. Hard state neutron star and black hole X-ray binaries in the radio:X-ray luminosity plane. *Mon. Not. R. Astron. Soc.* **478**, L132–L136 (2018).


Acknowledgements This work is part of the research programme Athena with project number 184.034.002, which is (partly) financed by the Dutch Research Council (NWO). Y.Z. acknowledges the support from China Scholarship Council Scholarship (201906100030). TMB acknowledges financial contribution from the agreement ASI-INAF n.2017-14-H.0, PRIN-INAF 2019 N.15, and thanks the Team Meeting at the International Space Science Institute (Bern) for fruitful discussions. D.A acknowledges support from the Royal Society. O.B. and D.A. acknowledge support of a Diamond Jubilee International Visiting Fellowship by the University of Southampton. We thank G. Pooley for making the radio data available. This research has made use of data and/or software provided by the High Energy Astrophysics Science Archive Research Center (HEASARC), which is a service of the Astrophysics Science Division at NASA/GSFC. This research made use of NASA's Astrophysics Data System.

Author Contributions All authors contributed to interpretation of results and edited the manuscript. M.M. led the interpretation, obtained spectral parameters and wrote the manuscript. K.K. wrote the model that triggered this research, produced initial radio/timing plots, fitted rms and lag spectra and co-led the interpretation. F.G. produced initial 3D radio/timing/spectral plots, fitted rms and lag spectra of the QPO and co-led the interpretation. M.M., K.K. and F.G. measured extra QPO frequencies. L.Z. obtained parameters of the QPO. Y.Z. obtained parameters of the high-frequency bump. T.M.B. had the idea to study the high-frequency bump in connection with the radio flux. D.A. and O.B. discussed the results and contributed to the interpretation.


Competing Interests The authors declare that they have no competing financial interests.

Supplementary Information is available for this paper.


Correspondence Correspondence and requests for materials should be addressed to MM (email: mariano@astro.rug.nl). Reprints and permissions information is available at www.nature.com/reprints




1a



1b





1c



1d


1 Hardness ratio vs. QPO frequency of GRS 1915+105

Each point corresponds to one of the 410 observations of GRS 1915+105. The hardness ratio is the ratio of the observed intensity from 13–60 keV to 2–7 keV in units of the Crab nebula. Average errors of the QPO frequency and hardness ratio are, respectively, ± 0.05 Hz and ± 0.001 Crab units, and are both smaller than the size of the points. The colour of the points in each panel represent: **a**, The 15-GHz radio-flux with average error of $\pm(1-5)$ mJy. **b**, The electron temperature of the X-ray corona with average error of ± 0.75 keV. **c**, The flux of the iron emission line with average error of $\pm 0.7 \times 10^{-10}$ erg cm⁻² s⁻¹. **d**, The fractional rms amplitude of the high-frequency bump with average error of $\pm 0.7\%$. In panel **d** the circles indicate measured values and the triangles upper limits. (All errors and upper limits in this work represent $1-\sigma$ and 95% confidence intervals, respectively.)




2 Flux of the iron line vs. total flux in the 2–25 keV range for GRS 1915+105

Each coloured point corresponds to one of the 410 observations of GRS 1915+105. The colour scale represents the temperature of the corona (left panel) and the phase lags of the QPO in radians (right panel). Negative (positive) lags indicate that, at the QPO frequency, the soft photons in the band 2–5.7 keV arrive after (before) the hard ones in the band 5.7–15 keV. The average error of the plotted quantities is shown at the bottom right of the plots. The lines indicate the two separate correlations. The plus and cross symbols correspond to measurements of the the flux of the line using observations of, respectively, Chandra²⁵ and Suzaku²⁶.



3 Schematic picture of the corona turning into the jet in GRS 1915+105

When the QPO frequency is in the 2–6 Hz range the lags are negative, the corona is hot and extended, and covers the inner parts of the disc (panel **a**). The magnitude of the lags and the size of the corona decrease and the inner disc radius increases as the QPO frequency decreases. When the QPO frequency is ~ 2 Hz the lags are zero and the size of the corona is equal to the inner radius of the accretion disc (panel **b**). As the frequency of the QPO decreases further below ~ 2 Hz the lags turn positive and the corona becomes the jet (panels **c** and **d**).



4 Time evolution of QPO frequency and radio flux for GRS 1915+105

The points connected by a line show the time evolution of the frequency of the QPO in the X-ray power spectrum of GRS 1915+105. The light-blue curve (smoothed with a Gaussian kernel) shows the simultaneous measurements of the radio flux density at 15 GHz (y axis rescaled). A strong radio flare appears on the two occasions in which the lags of the QPO turn from soft (red points) to hard (red/blue points), corresponding to the QPO frequency crossing from above to below ~ 2 Hz. The horizontal band shows the range of QPO frequencies over which the transition occurs. Here we show the first two radio flares in our data, but the same behaviour occurs consistently for all radio flares during our observations (see Supplementary Material).

177 **Methods**

178 **Power spectra:** We examined all the RXTE archival observations of GRS1915+105 from 1996 to
179 2012 obtained with the Proportional Counter Array (PCA). The observations that we used for our
180 analysis belong to the class⁸ χ , state C, equivalent to one of the hard states in other (transient) black-
181 hole sources. For each observation we computed the Fourier power spectrum in the full energy
182 band (absolute PCA channel 0–249) every 128 s with a time resolution of 1/512 s, corresponding
183 to a Nyquist frequency of 256 Hz. We averaged all the 128-s power spectra within an observation,
184 subtracted the contribution due to Poisson noise⁵⁰ and normalised⁵¹ these averaged power spectra to
185 units of fractional rms squared per Hz. (We ignored the background count rate for this calculation
186 since it was always negligible compared to the source count rate.) We subsequently applied a
187 logarithmic frequency rebin to the data such that the size of a bin increases by $\exp(1/100)$ with
188 respect to the size of the previous one, and we used XSPEC version 12.9 to fit the resulting power
189 spectra with a sum of Lorentzian functions⁵² that represent the broad-band noise component and
190 a number of QPOs¹⁹. As in previous studies²⁰ we included a Gaussian component centred at zero
191 frequency in the model to fit a high-frequency bump at 60–80 Hz in the power spectra. (We show
192 two power spectra with the best-fitting model in Extended Data Figure 5.) For the rest of the
193 analysis we selected only observations where at least one narrow QPO peak is present on top of
194 the broad-band noise component in the power spectra, which is typical for the type-C QPOs¹⁸.
195 Based on the fitting results we only retained features that were detected at a significance greater
196 than $3\text{-}\sigma$ and had a Q factor, defined as the ratio of the QPO frequency to its width, of 2 or more.
197 We further checked the spectrogram of each observation, which shows visually the Fourier power

198 spectrum as it varies with time, and excluded ~ 130 observations in which the QPO frequency
199 changed significantly within a full RXTE observation. Our final sample contains a total of 620
200 observations.

201 **Lag spectra:** Following the method described in^{53,54} we produced a frequency-dependent phase-
202 lag spectrum (lag-frequency spectrum) between the 2–5.7 and 5.7–15 keV energy bands for each
203 observation. Since our sample includes observations during the PCA calibration epochs 3–5,
204 to account for changes in the PCA channel-to-energy gain factor we selected the closest absolute
205 channels that matched these energy bands, but the exact boundaries of each band still differ slightly
206 between epochs. To calculate the phase lags of the QPO we averaged the lag-frequency spectra
207 around the centroid frequency of the QPO, $\nu_0 \pm FWHM/2$, where $FWHM$ is the full-width at
208 half-maximum of the Lorentzian that we used to fit the QPO profile. In principle, the phase lags
209 in the range of frequencies of the QPO can be affected by the lags of the underlying broad-band
210 noise component. However, in GRS 1915+105 the rms amplitude of the QPO is much higher than
211 that of the broad-band noise and the phase lags at the frequency of the QPO are dominated by
212 the QPO itself, with the contribution of the noise component being negligible⁵⁵. In this work, a
213 positive (negative) lag means that the hard photons lag (lead) the soft photons. No correction for
214 the dead-time-driven cross-talk⁵⁶ was done because this effect was found to be negligible. (See¹⁹
215 for other details of the timing analysis.) In Extended Data Figure 6 we show nine representative
216 power and lag-frequency spectra covering the range of frequencies of the QPO in GRS 1915+105.

217 **Energy Spectra:** We used the RXTE/PCA³ Standard 2 data to extract energy spectra separately
 218 for each observation in our sample. We corrected the energy spectra for dead time and used the
 219 FTOOLS PCABACKEST and PCARSP in HEADAS v.6.27 to, respectively, extract background spec-
 220 tra and produce response files for each observation. We fitted the energy spectra of all observa-
 221 tions jointly with the model VPHABS*(DISKBB+GAUSS+NTHCOMP). The component VPHABS
 222 accounts for the interstellar absorption along the line of sight to the source; we used the abundance
 223 and cross-section tables given by⁵⁷ and⁵⁸, respectively, with the column density of hydrogen, N_{H} ,
 224 linked across observations and free to vary. Because N_{H} in the direction of GRS 1915+105 is quite
 225 high⁵⁹, $N_{\text{H}} \gtrsim 6 \times 10^{22}$ atoms cm^{-2} , we also let the abundance of iron in the interstellar material free
 226 to vary, since neutral iron produces an absorption edge at $E \approx 7.1$ keV that was apparent in the fit
 227 residuals. The component DISKBB⁶⁰ represents the emission from an optically thick and geometri-
 228 cally thin accretion disc, and has parameters kT_{bb} and N_{dbb} that are, respectively, the temperature
 229 at the inner disc radius and the normalisation of the component defined as the ratio of the inner
 230 disc radius to the distance to the source squared multiplied by the cosine of the inclination of the
 231 disc with respect to the line of sight. The component NTHCOMP⁶¹ represents the emission due to
 232 inverse-Compton scattering from the corona. The parameters of this component are the power-law
 233 index, Γ , the electron temperature of the corona, kT_{e} , the temperature of the source of soft pho-
 234 tons that enter into, and are up scattered in, the corona, kT_{seed} , and a normalisation that gives the
 235 flux density at 1 keV. We assumed that the source of seed soft photons is the accretion disc, and

³We also fitted the RXTE/HEXTE (High Energy X-ray Timing Experiment) data of those observations in which
 the instrument was operational; the results were consistent with the ones of the RXTE/PCA, but since data of this
 instrument were not available for all observations, we did not use the RXTE/HEXTE data for the rest of the analysis.

236 linked the temperature kT_{seed} to the temperature of the DISKBB component for each observation
237 separately during the fits. In this model, the optical depth of the corona, τ , assumed to be ho-
238 mogeneous, is a function¹¹ of Γ and kT_e . Finally, the component GAUSS represents a broad iron
239 emission line at 6.5–7 keV due to reflection of corona photons off the accretion disc^{12,13}. While
240 a full reflection model^{62,63} would be more appropriate to describe this effect, the PCA instrument
241 does not have enough spectral resolution to distinguish between the profile of a line in a full reflec-
242 tion model and a Gaussian component. The parameters of the GAUSS component are the energy,
243 width and total photon flux of the line, E_g , σ , and N_g , respectively. Finally, we calculated the
244 total observed flux of the source and the intrinsic (i.e., not affected by interstellar absorption) flux
245 of the full model and of all the individual model components in the 2–25 keV energy range. In
246 Extended Data Figure 7 we show the energy spectra, with the best-fitting model, of the same two
247 observations in Extended Data Figure 5.

248 If the iron emission line is due to reflection of corona photons off the accretion disc, and
249 the width of the line is a consequence of Doppler boosting, Doppler beaming and gravitational
250 redshift, it would be more appropriate to use a full reflection component that includes relativistic
251 broadening instead of a Gaussian to fit the spectrum. To explore this we fitted all the spectra
252 with the model `VPHABS*(DISKBB+RELXILLCP+NTHCOMP)`, where the component `RELXILLCP`⁶³
253 represents the reflection off the accretion disc of the emission from the `NTHCOMP` component.
254 Compared to the fits with a Gaussian, this model has six extra free parameters. Given that the
255 RXTE/PCA instrument has limited spectral resolution and does not extend below ~ 3 keV, the
256 model is insensitive to some of the parameters of `RELXILLCP`, and some of these parameters become

257 highly correlated with those of the DISC component. We therefore fixed the spin of the black hole
258 to $a_* = 0.95$ (ref. [59]) and the inclination angle of the accretion disc to the line of sight to
259 $i = 65^\circ$ (ref. [2]). We further linked the two emissivity indices during the fits, which eliminates
260 the parameter that gives the radius in the disc at which the power-law index of the emissivity profile
261 changes, and fixed the inner radius of the disc at the radius of the innermost stable circular orbit⁵⁹.
262 Finally we linked the power-law index, Γ , and the electron temperature, kT_e , of RELXILLCP to
263 the corresponding parameters of NTHCOMP. The extra free parameters of the fit are one emissivity
264 index, the reflection fraction, the ionisation parameter and the iron abundance of the disc. We
265 obtain equally good fits as with the model above, with the reflection fraction ranging from 1%
266 to 20%. Because the reflected component is much-lower than the Comptonised component, the
267 parameters of the NTHCOMP in these fits are consistent with those from the fits with a Gaussian
268 line. The biggest impact of fitting the spectra with a reflection component is that the parameters of
269 the disc are less well constrained in the model with RELXILLCP than in the model with a Gaussian.
270 This is understandable given that the emission of the reflection component at energies below $\sim 4-5$
271 keV, where the disc emission peaks, is higher than that of a Gaussian.

272 The RELXILLCP model does not provide the flux or the equivalent width of the Gaussian, how-
273 ever the reflection fraction of RELXILLCP is generally correlated with the parameters of the line⁶⁴.
274 We plotted the flux of the iron line from the fits with a Gaussian against the reflection fraction from
275 the fits with RELXILLCP and confirmed that this is indeed the case in our fits. We therefore obtain
276 similar plots to those in Figure 2 if instead of the flux of the Gaussian line we plot the reflection
277 fraction of RELXILLCP vs. total flux. We prefer to use the former because, as we explained above,

278 the model with a Gaussian line has less free parameters than the full reflection model which, given
279 the limited spectral resolution of the RXTE/PCA instrument, leads to degeneracies of the parame-
280 ters in the fits with the full reflection model, and because by using a Gaussian we can easily include
281 in the plot the measurements of the flux of the iron line obtained with Chandra and Suzaku, as we
282 did in Figure 2.

283 **Hardness ratio:** For each RXTE observation of GRS 1915+105 we computed a hardness ratio
284 value defined as the ratio of the background-subtracted count rate of the source in the 13–60 keV
285 band to that in the 2–7 keV band. As with the lag-frequency spectra, we selected the closest
286 absolute channels that matched these energy bands in each PCA gain epoch. Before calculating
287 the ratios, we corrected each observed count rate for instrumental dead time and normalised them
288 by the count rate in the same band from the Crab nebula to account for possible changes of the
289 effective area of the instrument with time.

290 **Radio fluxes:** For the flux-density data of GRS 1915+105 in radio we used measurements⁹ from
291 the Ryle Telescope at 15 GHz with the four mobile antennas set in a compact array configuration
292 within 100 m of the nearest fixed antenna yielding a resolution of ~ 30 arcsec at that frequency.
293 The flux-density scale was calibrated with observations of the nearby quasars 3C 48 or 3C 286.
294 The observations consist mostly of 32-s samples with an rms noise of 6 mJy that decreases as the
295 square root of the integration time; flux-density values below about 1 mJy may be unreliable. See⁹
296 for other details of the analysis of the radio data.

297 Finally, we cross-correlated all the X-ray and radio data based on the date of the observations,

298 which left us with a sample of 410 observations with simultaneous radio flux densities at 15 GHz
299 and X-ray energy, power-density and lag-frequency spectra and hardness ratios.

300 Having described the observations and analysis we used, it is worth noting that the mea-
301 surements presented in this paper come from very different types of data and totally independent
302 analysis techniques: The hardness ratio, iron-line flux and kT_e come from X-ray light curves and
303 time-averaged energy spectra, the frequency and lags of the QPO and the rms amplitude of the
304 high-frequency bump come from Fourier power spectra of high-time resolution data, and the radio
305 flux was measured independently in a totally different frequency band than the X-ray data.

306 **Extended results and discussion**

307 **Fits to the power spectra:** In Extended Data Figure 5 we plot two representative Fourier power
308 spectra of GRS 1915+105 to show both the type-C QPO and the high-frequency bump. The power
309 spectrum plotted in black corresponds to the observation from September 21 1998 (ObsID 30703-
310 01-34-00), while the power spectrum plotted in blue corresponds to the observation from April
311 10 1998 (ObsID 30402-01-09-01). Both power spectra were calculated using photons from the
312 full PCA band, nominally 2–60 keV. The QPO is the narrow feature that appears in both power
313 spectra at ~ 2 Hz, with a second harmonic and subharmonic at, respectively, ~ 4 Hz and ~ 1 Hz.
314 In the case of GRS 1915+105 it is relatively straightforward to distinguish the QPO fundamental
315 from the subharmonic and harmonic components because the QPO fundamental, subharmonic and
316 harmonic components show very different rms amplitude and lags dependence when plotted vs.
317 their own frequency¹⁹, and the fundamental is always the strongest peak in the power spectrum.

318 The model includes also two broad Lorentzians centred, respectively, at 0 Hz and 2.5 Hz to
319 fit the the broad-band noise component in the power spectrum. The high-frequency bump is the
320 broad feature extending up to ~ 100 Hz in the black power spectrum^{52,65–70}. We fitted this bump
321 with a zero-centred Gaussian and let the width and normalisation free to vary during the fits. It
322 is apparent that the QPO at 2 Hz and its second harmonic and subharmonic are present in both
323 power spectra, but the high-frequency bump is only present in the observation of September 1998
324 and significantly absent in the one of April 1998. We computed the rms amplitude of the high-
325 frequency bump by integrating the power of the best-fitting Gaussian from 0 Hz to ∞ . The rms
326 amplitude of the high-frequency bump in the observation of September 1998 is $8.0 \pm 0.3\%$ of the

327 total luminosity of the source in the 2–60 keV band, whereas the 95% confidence upper limit of
328 the rms amplitude in the observation of April 1998 is only 1% of the total luminosity. Since the
329 high-frequency bump was not significantly detected in the observation of April 1998, to calculate
330 the upper limit we fixed the width of the Gaussian to the value we obtained from the fit to the
331 observation of September 1998. We followed the same procedure to compute the rms amplitude
332 or, when appropriate, the upper limits of the high-frequency bump in the rest of the observations.
333 In the observations in which the high-frequency bump is significantly detected the width of this
334 feature ranges from ~ 30 Hz to ~ 100 Hz. In those observations in which the high-frequency bump
335 was not significantly detected, to compute the upper limits we fixed the width of the Gaussian to
336 70 Hz, which is the average value that we obtained from the fits of the observations in which the
337 high-frequency bump was significantly detected. We report a detailed analysis of the properties of
338 the high-frequency bump in a separate paper⁷¹.

339 **Fits to the energy spectra:** In Extended Data Figure 7 we show the energy spectra of the same
340 two observations for which we show the power spectra in Extended Data Figure 5. The spectrum
341 plotted in black corresponds to the observation from September 21 1998 (ObsID 30703-01-34-
342 00), while the spectrum plotted in blue corresponds to the observation from April 10 1998 (ObsID
343 30402-01-09-01). This Figure shows that the flux of the source was lower and, at the same time,
344 the spectrum was harder (extending to higher energies) in September 1998 than in April 1998.
345 This difference is reflected in the values of the best-fitting temperature of the corona in these two
346 observations, which are $kT_e=10.9\pm 0.8$ keV and $kT_e=6.3\pm 0.5$ in the observations of September
347 1998 and April 1998, respectively. From the comparison of this Figure and Extended Data Figure 5

348 it is apparent that the high-frequency bump is present when the temperature of the corona is high.
349 Indeed, when we fit all the observations we find that the rms amplitude of the high-frequency bump
350 is higher when the temperature of the corona is higher, regardless of the values of the hardness ratio
351 and QPO frequency in each observation (see Figure 1).

352 **Energy-dependent rms amplitude and phase lags of the QPO. The corona becomes the jet:**
353 In Extended Data Figure 6 we plot the power- and lag-frequency spectra of nine observations of
354 GRS 1915+105, representative of the range of frequencies covered by the type-C QPO. The phase
355 lags of the QPO change from positive when the frequency of the QPO is lower than ~ 2 Hz to
356 negative when the frequency of the QPO is higher than ~ 2 Hz. (See¹⁹ for details of this result and a
357 discussion of the implication of this behaviour.) At the same time, as the QPO frequency increases
358 the slope of the lag spectrum decreases²⁹, going from positive when the frequency increases from
359 from ~ 0.5 Hz to ~ 2 Hz, to 0 when the frequency is ~ 2 Hz, and finally to negative when the QPO
360 frequency increases further from ~ 2 Hz to ~ 6 Hz.

361 The energy-dependent rms amplitude and phase lags of the QPO in GRS 1915+105 are well
362 explained by a time-dependent model²⁸ in which photons from the accretion disc are inverse Compton
363 scattered in a corona of characteristic size L , and a fraction of those photons subsequently
364 impinges back onto the disc before reaching the observer. The time-averaged version of this variability
365 model is equivalent to the NTHCOMP model in XSPEC. In the current version of the model
366 we assume that the corona is spherically symmetric and has constant optical depth and electron
367 temperature. Most likely the geometry of the corona is more complex than that, and the optical
368 depth and electron temperature change as a function of the distance to the black hole. In a non-

369 spherical corona the rms amplitude and lags of the QPO may change with the inclination of the
 370 binary system with respect to the line of sight^{72,73}. We note, however, that both the rms amplitude
 371 and lags of the QPO in this and other sources change in a systematic way with QPO frequency,
 372 which in turn changes over time scales of minutes to hours, much faster than potential changes
 373 of the binary inclination that are expected to happen over periods of a few tens of days to a few
 374 thousand years⁷⁴. This evinces that, although inclination may play a role, the thermodynamical
 375 properties of the corona have a much larger effect than inclination in setting the rms amplitude and
 376 lags of the QPO.

377 Fits with this time-dependent model to the data of GRS 1915+105 show^{29,75} that, as the QPO
 378 frequency decreases from ~ 6 Hz to ~ 2 Hz, the feedback fraction is close to 100% and remains
 379 more or less constant. At the same time the size⁴ of the corona decreases from $L \approx 1000 - 2000$ km
 380 ($\sim 60 - 120 R_g$ for a 12 solar-mass black hole) to $L \approx 100$ km ($\sim 6 - 10 R_g$), the lags are negative and
 381 their magnitude decreases. At a QPO frequency of ~ 2 Hz the corona size is minimum, the feedback
 382 fraction drops abruptly to zero and the lags become zero. If the QPO frequency is produced by
 383 Lense-Thirring precession^{23,24} of the inner edge of the accretion disc, at a QPO frequency of 2 Hz
 384 the size of the corona is more or less equal to the inner disc radius (dashed line in Extended Data
 385 Figure 8). When the QPO frequency decreases below ~ 2 Hz, the corona size increases again up
 386 to $L \approx 2 \times 10^4$ km ($\sim 1200 R_g$) at the lowest QPO frequency, the feedback fraction remains constant
 387 close to zero and the lags become positive and increase with decreasing QPO frequency. Although

⁴In the model we solve the Kompaneets equation⁷⁶ for a spherical corona; since the actual geometry of the corona is likely different, the values given here should be considered as a characteristic size of the corona.

388 the size of the corona increases again when the QPO frequency is below ~ 2 Hz, the fact that the
389 feedback fraction is close to zero whereas when the QPO frequency is above ~ 2 Hz the feedback
390 fraction is high, implies that the corona does not cover the inner parts of the accretion disc. This in
391 turn demonstrates that the geometry of the corona has changed (see Fig. 3).

392 Extended Data Figure 9 shows the time evolution of the QPO frequency superimposed on 10
393 years of the radio flux measurements of GRS 1915+105 with the Ryle telescope. To produce this
394 Figure we reanalysed all the RXTE observations in the period of the radio monitoring campaign,
395 and added 357 observations with QPOs that were not in the sample of 410 observations discussed
396 so far. These 357 observations were not included in the original analysis¹⁹ because the QPO
397 frequency drifts by a few tenths of Hz in the observation period, which could bias the measurement
398 of the rms and lags of the QPO. But since the frequency drift is only slightly larger than the error
399 of the average QPO frequency in each observation, we decided to use these data for this Figure
400 because this allows us to have a denser coverage of the time evolution of the QPO frequency during
401 the radio monitoring campaign. Extended Data Figure 9 shows that the intervals of strong radio
402 flares⁵ coincide with the times when the QPO frequency is below 2 Hz. This behaviour repeats
403 consistently for all the radio flares in these observations; there is no radio flare in which the QPO
404 frequency is not below 2 Hz, and there is no case of a QPO frequency below 2 Hz without a radio
405 flare. Given the data, we estimated that the probability of having a QPO frequency below 2 Hz and
406 a radio flare at the same time if the two phenomena were uncorrelated is less than 2×10^{-10} .

⁵While it is apparent to the eye from the Figure, it is difficult to quantify what a strong radio flare is. Here we classify a flare as strong if its flux density exceeds 50 mJy for more than 10 days and the flare lasts more than 100 days.

407 Taken together, the results shown in Extended Data Figures 8 and 9 and the sudden change of
408 the feedback fraction^{29,75} from ~ 1 to ~ 0 when the QPO frequency is, respectively, above or below
409 2 Hz are consistent with the scenario described earlier, in which the energy that is initially stored
410 in the X-ray corona is gradually released into the radio jet and, quite possibly, the X-ray corona
411 itself becomes the jet.


412 The morphing of the extended corona into the jet is strengthened by the relation between
413 the actual luminosity of the corona and that expected if the emission is due to inverse-Compton
414 scattering of seed photons from the accretion disc. In Extended Figure 10 we show the ratio of
415 the expected to the observed corona flux vs. the observed flux of the corona. We calculated the
416 expected flux as the product of the bolometric disc flux times e^y , where y is the Compton parameter.
417 (This is the expression for the case that the optical depth is $\tau > 1$.) Since the Compton y parameter
418 gives the number of scatterings times the relative energy that a photon gains per scattering, if the
419 seed-photon source is the disc, the expected and observed corona fluxes should be roughly the
420 same. This Figure shows that the observed and expected fluxes are the same (within a factor 3)
421 only when the lags of the QPO are negative (yellow, orange and red points), i.e., when the QPO
422 frequency is higher than ~ 2 Hz and the disc is enshrouded by the corona. When the lags are
423 positive (light blue and blue points) and the corona morphed into the jet, if the seed photons were
424 from the disc the expected flux of the corona is much lower than the one observed. In this phase a
425 different source must be providing the seed photons for Comptonisation. Since in this phase the jet
426 is active, the most likely source of seed photons is synchrotron emission⁷⁷ from the same electrons
427 that produce the jet. Indeed, if we calculate the expected corona flux by multiplying the observed

428 synchrotron flux coming from the jet extrapolated to the X-ray band times e^y , the expected corona
429 flux is much higher when the lags of the QPO are positive than when they are negative.

430 **Data availability** All the X-ray data used in this study are available NASA's High Energy Astro-
431 physics Science Archive Research Center <https://heasarc.gsfc.nasa.gov/>. The radio
432 data used in this study are available through the website [http://www.mrao.cam.ac.uk/](http://www.mrao.cam.ac.uk/~guy/)
433 [~guy/](http://www.mrao.cam.ac.uk/~guy/).


434 **Code availability** The data reduction was done using HEADAS v6.27, while the model fitting
435 of energy, power and lag-energy spectra was done with XSPEC; both packages are available at
436 the HEASARC website (<https://heasarc.gsfc.nasa.gov/>). The timing analysis was
437 performed with the GHATS package developed by T.M.B. and is available on request (http://astrosat.iucaa.in/~astrosat/GHATS_Package/Home.html). All figures were
438 made in TOPCAT, a JAVA-based scientific plotting package developed by Mike Taylor (<http://www.star.bris.ac.uk/~mbt/topcat/>).
439
440

Extended data figures and tables




Extended Data Figure 5 Representative power spectra of GRS 1915+105

X-ray Fourier power spectra of two observations of GRS 1915+105 (data with errors) fitted with a model consisting of several Lorentzians (solid lines with the individual components indicated with dotted lines) for the QPO (plus harmonic and sub-harmonic components) and a zero-centred Gaussian for the high-frequency bump. In both power spectra the strong QPO fundamental component is at ~ 2 Hz; the high-frequency bump peaking at ~ 70 Hz is significantly detected, with an rms amplitude of $8.0 \pm 0.3\%$, in the power spectrum plotted in black (ObsID 30703-01-34-00 of September 21 1998), and is significantly absent, with a 95% confidence upper limit of the rms amplitude of 1%, in the power spectrum plotted in blue (ObsID 30402-01-09-01 of April 10 1998).




Extended Data Figure 6 Power-density (upper half of each panel) and phase-lag (lower half of each panel) spectra of nine observations of GRS 1915+105 with RXTE. Each panel corresponds to an observation in which the QPO was at the frequency given, together with the ObsID, in the legend of the panel. We calculated the power spectra using the photons from the full PCA band, and the phase-lag spectrum using photons in the 5.7–15 keV band with respect to the photons in the 2–5.7 keV band. The lags are given in units of $\text{rad}/2\pi$ and indicate whether, at a given frequency, the variability of the low-energy photons lag (negative lags) or lead (positive lags) that of the high-energy photons. The black vertical lines indicate the range of frequencies that we used to measure the lags of the QPO.




Extended Data Figure 7 Representative energy spectra of GRS 1915+105

X-ray energy spectra (data points with errors) of the same two observations of GRS 1915+105 shown in Extended Data Figure 5, fitted with a model consisting of the emission from the accretion disc, the corona and a broad iron line. The dashed lines represent the best-fitting model of the corona, while the solid lines through the data correspond to the full model. In the spectrum plotted in black (ObsID 30703-01-34-00 of September 21 1998) the temperature of the corona is $kT_e=10.9\pm 0.8$ keV, while in the one plotted in blue (ObsID 30402-01-09-01 of April 10 1998) $kT_e=6.3\pm 0.5$ keV. The difference in temperature between the two spectra is apparent by the fact that, at high energies, the flux drops quicker in the spectrum plotted in blue than in the one plotted in black. The bottom panel shows the residuals, $\Delta\chi=(\text{data}-\text{model})/\text{error}$, of the best-fitting model.




Extended Data Figure 8 Size of the corona as a function of QPO frequency for GRS 1915+105

Each point corresponds to one of the 410 observations of GRS 1915+105. The size of the corona comes from the fits of a variable-Comptonisation model^{28,29} to the rms and lag spectra of the QPO⁷⁵. The colour of the points indicates the feedback fraction, the fraction of photons from the corona that returns to the disc. The black solid line is the weighted average of the points. The grey dashed line shows the inner radius of the accretion disc in km assuming that the QPO frequency is the Lense-Thirring precession frequency for a 12 solar-mass black hole.



Extended Data Figure 9 Time evolution of QPO frequency and radio flux for GRS 1915+105

The points connected by a line show the time evolution of the frequency of the QPO in the X-ray power spectrum of GRS 1915+105. The light-blue curve (smoothed with a Gaussian kernel) shows the simultaneous measurements of the radio flux density at 15 GHz (y axis rescaled). A strong radio flare appears whenever the lags of the QPO turn from soft (red points) to hard (blue points), corresponding to the QPO frequency crossing from above to below ~ 2 Hz. The horizontal band shows the range of QPO frequencies over which the transition occurs. The probability of having a positive QPO lag *always* and *only* during a radio flare if the two phenomena were uncorrelated is less than 2×10^{-10} . (Fig. 4 in the main text shows a zoom-in of the first two flares.)



Extended Data Figure 10 Expected corona flux in GRS 1915+105

Ratio of the expected to the observed corona flux vs. observed corona flux. The expected corona flux is calculated as the bolometric disc flux times e^y , where y is the Compton parameter. Each point corresponds to one of the 410 observations of GRS 1915+105. The average error of the plotted quantities is shown at the upper right of the plot. The colour scale represents the phase lags, in units of $(\text{rad}/2\pi)$, of the QPO (see Fig. 2 for an explanation). The triangles indicate upper limits of the ratio of the expected to the observed corona flux. The expected and observed fluxes are roughly the same (within a factor of ~ 3) when the lags are negative, the corona is extended and hot, and it covers the inner parts of the accretion disc (colours towards red), whereas the expected flux drops significantly when the lags are positive, the geometry of the corona changes and its temperature drops as it turns into the jet (colours towards blue).

References

50. Zhang, W., Jahoda, K., Swank, J. H., Morgan, E. H. & Giles, A. B. Dead-Time Modifications to Fast Fourier Transform Power Spectra. *Astrophys. J.* **449**, 930 (1995).
51. Belloni, T. & Hasinger, G. An atlas of aperiodic variability in HMXB. *Astron. Astrophys.* **230**, 103–119 (1990).
52. Nowak, M. A. Are there three peaks in the power spectra of GX 339-4 and Cyg X-1? *Mon. Not. R. Astron. Soc.* **318**, 361–367 (2000).
53. Vaughan, B. A. & Nowak, M. A. X-Ray Variability Coherence: How to Compute It, What It Means, and How It Constrains Models of GX 339-4 and Cygnus X-1. *Astrophys. J.* **474**, L43–L46 (1997).
54. Nowak, M. A., Vaughan, B. A., Wilms, J., Dove, J. B. & Begelman, M. C. Rossi X-Ray Timing Explorer Observation of Cygnus X-1. II. Timing Analysis. *Astrophys. J.* **510**, 874–891 (1999).
55. van den Eijnden, J., Ingram, A. & Uttley, P. Probing the origin of quasi-periodic oscillations: the short-time-scale evolution of phase lags in GRS 1915+105. *Mon. Not. R. Astron. Soc.* **458**, 3655–3666 (2016).
56. van der Klis, M. *et al.* The Complex Cross-Spectra of Cygnus X-2 and GX 5-1. *Astrophys. J.* **319**, L13 (1987).
57. Wilms, J., Allen, A. & McCray, R. On the Absorption of X-Rays in the Interstellar Medium. *Astrophys. J.* **542**, 914–924 (2000).

58. Verner, D. A., Ferland, G. J., Korista, K. T. & Yakovlev, D. G. Atomic Data for Astrophysics. II. New Analytic FITS for Photoionization Cross Sections of Atoms and Ions. *Astrophys. J.* **465**, 487 (1996).
59. Miller, J. M. *et al.* NuSTAR Spectroscopy of GRS 1915+105: Disk Reflection, Spin, and Connections to Jets. *Astrophys. J.* **775**, L45 (2013).
60. Mitsuda, K. *et al.* Energy spectra of low-mass binary X-ray sources observed from Tenma. *Publ. astr. Soc. Japan* **36**, 741–759 (1984).
61. Zdziarski, A. A., Johnson, W. N. & Magdziarz, P. Broad-band γ -ray and X-ray spectra of NGC 4151 and their implications for physical processes and geometry. *Mon. Not. R. Astron. Soc.* **283**, 193–206 (1996).
62. García, J. *et al.* Improved Reflection Models of Black Hole Accretion Disks: Treating the Angular Distribution of X-Rays. *Astrophys. J.* **782**, 76 (2014).
63. Dauser, T., Garcia, J., Parker, M. L., Fabian, A. C. & Wilms, J. The role of the reflection fraction in constraining black hole spin. *Mon. Not. R. Astron. Soc.* **444**, L100–L104 (2014).
64. Dunn, R. J. H., Fender, R. P., Körding, E. G., Cabanac, C. & Belloni, T. Studying the X-ray hysteresis in GX 339-4: the disc and iron line over one decade. *Mon. Not. R. Astron. Soc.* **387**, 545–563 (June 2008).
65. Belloni, T., Méndez, M., van der Klis, M., Lewin, W. H. G. & Dieters, S. A State Transition of GX 339-4 Observed with the Rossi X-Ray Timing Explorer. *Astrophys. J.* **519**, L159–L163 (1999).

66. Revnivtsev, M., Gilfanov, M. & Churazov, E. High frequencies in the power spectrum of Cyg X-1 in the hard and soft spectral states. *Astron. Astrophys.* **363**, 1013–1018 (2000).
67. Belloni, T., Psaltis, D. & van der Klis, M. A Unified Description of the Timing Features of Accreting X-Ray Binaries. *Astrophys. J.* **572**, 392–406 (2002).
68. Pottschmidt, K. *et al.* Long term variability of Cygnus X-1. I. X-ray spectral-temporal correlations in the hard state. *Astron. Astrophys.* **407**, 1039–1058 (2003).
69. Motta, S. E. *et al.* Black hole spin measurements through the relativistic precession model: XTE J1550-564. *Mon. Not. R. Astron. Soc.* **439**, L65–L69 (2014).
70. Motta, S. E., Belloni, T. M., Stella, L., Muñoz-Darias, T. & Fender, R. Precise mass and spin measurements for a stellar-mass black hole through X-ray timing: the case of GRO J1655-40. *Mon. Not. R. Astron. Soc.* **437**, 2554–2565 (2014).
71. Zhang, Y. *et al.* *The evolution of the high-frequency variability in the black hole candidate GRS 1915+105 as seen by RXTE* *Mon. Not. R. Astron. Soc.* submitted. 2021.
72. Motta, S. E. *et al.* Geometrical constraints on the origin of timing signals from black holes. *Mon. Not. R. Astron. Soc.* **447**, 2059–2072 (2015).
73. van den Eijnden, J. *et al.* Inclination dependence of QPO phase lags in black hole X-ray binaries. *Mon. Not. R. Astron. Soc.* **464**, 2643–2659 (2017).
74. Wijers, R. A. M. J. & Pringle, J. E. Warped accretion discs and the long periods in X-ray binaries. *Mon. Not. R. Astron. Soc.* **308**, 207–220 (Sept. 1999).

75. García, F. *et al.* *The evolving properties of the corona of GRS 1915+105: A spectral-timing perspective through Comptonisation modelling* *Mon. Not. R. Astron. Soc.* submitted. 2021.
76. Kompaneets, A. S. The Establishment of Thermal Equilibrium between Quanta and Electrons. *Soviet Journal of Experimental and Theoretical Physics* **4**, 730–737 (1957).
77. Veledina, A. Interplay of spectral components in timing properties of accreting compact objects. *Mon. Not. R. Astron. Soc.* **481**, 4236–4249 (2018).