

Dark Matter Physics from the CMB-S4 Experiment

Cora Dvorkin^{*1}, Renée Hlozek^{2,3}, Rui An⁴, Kimberly K. Boddy⁵, Francis-Yan Cyr-Racine⁶, Gerrit S. Farren⁷, Vera Gluscevic⁴, Daniel Grin⁸, David J. E. Marsh⁹, Joel Meyers¹⁰, Keir K. Rogers², Katelin Schutz¹¹, and Weishuang Linda Xu¹²

¹Department of Physics, Harvard University, 17 Oxford Street, Cambridge, MA 02138, USA

²Dunlap Institute for Astronomy and Astrophysics, University of Toronto, 50 St. George Street, Toronto, ON M5S 3H4, Canada

³Department of Astronomy and Astrophysics, University of Toronto, 50 St. George Street, Toronto, ON M5S 3H4, Canada

⁴Department of Physics and Astronomy, University of Southern California, Los Angeles, CA, 90089, USA

⁵Department of Physics, The University of Texas at Austin, Austin, TX 78712, USA

⁶Department of Physics and Astronomy, University of New Mexico, Albuquerque, NM 87106, USA

⁷Department of Applied Mathematics and Theoretical Physics, University of Cambridge, Cambridge CB3 0WA, United Kingdom

⁸Department of Physics and Astronomy, Haverford College, 370 Lancaster Ave, Haverford, PA 19041, United States

⁹Department of Physics, King's College, London WC2R 2LS, United Kingdom

¹⁰Department of Physics, Southern Methodist University, Dallas, TX 75275, USA

¹¹Department of Physics, Ernest Rutherford Physics Building, 3600 Rue University, Montréal, QC H3A 2T8

¹²Berkeley Center for Theoretical Physics, South Hall Rd, Berkeley, CA 94720, United States

Endorsers: Kevork N. Abazajian, Peter Adshead, Mustafa Amin, Denis Barkats, Darcy Barron, Amy N. Bender, Bradford A. Benson, Colin Bischoff, Julian Borrill, Thejs Brinckmann, John E. Carlstrom, Clarence Chang, Jacques Delabrouille, Eleonora Di Valentino, Thomas Essinger-Hileman, Simone Ferraro, Jon E. Gudmundsson, Selim C Hotinli, Kevin M. Huffenberger, Bradley R. Johnson, Louis Legrand, Marilena Loverde, Yin-Zhe Ma, Tony Mroczkowski, Moritz Münchmeyer, Michael D. Niemack, Giorgio Orlando, Alexandra Rahlin, Christian L. Reichardt, Mathieu Remazeilles, John E. Ruhl, Emmanuel Schaan, Radek Stompor, Cynthia Trendafilova, Matthieu Tristram, Gensheng Wang, Scott Watson, Edward J. Wollack, W.L. Kimmy Wu, Zhilei Xu, Andrea Zonca

Abstract

The nature of dark matter is one of the major puzzles of fundamental physics, integral to the understanding of our universe across almost every epoch. The search for dark matter takes place at different energy scales, and use data ranging from particle colliders to astrophysical surveys. We focus here on CMB-S4, a future ground-based Cosmic Microwave Background (CMB) experiment, which is expected to provide exquisite measurements of the CMB temperature and polarization anisotropies. These measurements (on their own and in combination with other surveys) will allow for new means to shed light on the nature of dark matter.

*cdvorkin@g.harvard.edu

1 Introduction

Cosmological and astrophysical observations have shown us that approximately 84% of all matter in the universe is some form of non-baryonic dark matter (DM). The nature of dark matter is still unknown, though a combination of laboratory experiments and astronomical observations have placed broad limits on the physical properties of dark matter candidates. We expect to see a huge improvement in the sensitivity of astrophysical measurements in the next decade, which will provide very deep insights for a wide range of dark matter scenarios. Of particular interest are observations of the Cosmic Microwave Background (CMB), primordial light from a time when the universe was roughly 380 000 years old. The CMB encodes information about the earliest moments of the history of the universe and its evolution over cosmic times. The statistics of the anisotropies in the temperature and polarization of the CMB are sensitive to the densities of the various constituents of the universe and their interactions, the cosmic expansion history, and the formation of cosmic structure. This white paper focuses on how observations with CMB-S4 will provide further insights into the nature of dark matter, enabling the possibility to distinguish among several proposed models of dark matter and the ‘standard’ cold dark matter (CDM) picture.

The observational future of CMB science is bright. Current state-of-the-art telescopes and detectors are being replaced with ever more sensitive instruments. The next-generation CMB-S4 [1] experiment will measure the CMB in multiple frequencies to noise levels of $\approx 1 \mu\text{K-arcmin}$ with a resolution of $< 1.5 \text{ arcmin}$ over 70% of the sky (in addition to measuring about 3% of the sky to a greater depth of $\sim 0.5 \mu\text{K-arcmin}$ with both $\sim 25 \text{ arcmin}$ and $\sim 2.5 \text{ arcmin}$ resolution). CMB-S4 will be able to probe departures from the Standard Model (SM) of particle physics through deep, high-resolution maps of the sky in CMB temperature and polarization fluctuations.

These measurements will provide a snapshot of the universe as it was around the time of recombination, and they will also reveal the imprints of structure growth at much later times. Gravitational lensing of the CMB [2] leads to characteristic distortions of the primary CMB maps, allowing us to statistically reconstruct maps of the integrated line-of-sight density. Scattering of CMB photons in galaxy clusters (the Sunyaev-Zel’dovich effect) [3, 4] allows for the identification of the most massive bound structures in the universe out to very high redshifts.

Cosmological measurements in general and CMB measurements in particular provide insights into dark matter physics that are complementary to direct, indirect, and collider searches for dark matter. Cosmological observables are impacted by the influence of dark matter on the entire cosmic history. Dark matter constraints derived from cosmology do not rely on assumptions about the dark matter density in the neighborhood of the Earth or of any astrophysical object. Furthermore, CMB observations are sensitive to regions of parameter space that are out of reach of current direct searches. Figure 1 provides an overview of the various means by which CMB-S4 will search for the effects of dark matter physics across a wide range of mass scales.

In the sections below, we contextualize a range of dark matter scenarios and describe the expected sensitivity of CMB-S4 to a variety of DM candidates.




Figure 1: CMB-S4 will be able to probe light relics, axions, warm dark matter, and different dark matter scenarios. The relevant mass to which the CMB is sensitive to is shown for each case. The observable that drives the constraint is shown in different colors. Figure from Ref. [1].

2 Dark matter annihilation

Weakly interacting massive particles (WIMPs) are some of the most compelling candidates for DM, where the amplitude of the self-annihilation cross section can give rise to the observed DM density today. When dark matter annihilates, there is an injection of energy that heats and ionizes the baryonic gas, and this energy is inversely proportional to the DM mass. If this happens during and after recombination, it will affect the CMB temperature and polarization fluctuations [5, 6]. A higher ionization fraction will increase the optical depth of the photons, and therefore it will produce a suppression in the CMB acoustic peaks. Furthermore, the extra scattering of photons will affect the polarization fluctuations at large scales. Measurements by the *Planck* satellite at degree angular scales provided strong bounds on annihilation and decay of DM at the sub-GeV scale [7, 8], complementing direct detection searches that probe heavier masses. Future measurements from next-generation ground-based CMB experiments are expected to improve the current sensitivity by a factor of 2-3 [9–11].

3 Sensitivity to thermal-relic dark matter mass

Current null results from direct detection searches for heavy WIMPs have inspired WIMP-like and other models for light thermal-relic DM, with sub-GeV masses [12–18]. If the light thermal-relic DM particles are in thermal contact with the rest of the plasma during Big Bang Nucleosynthesis (BBN), which is the case for majority of the popular WIMP models

in the current literature [19–29], and if their masses are around $\sim 0.01\text{--}20$ MeV, they become non-relativistic right around the time of BBN. The resulting DM annihilation into Standard Model particles can affect the early expansion history, altering the time at which proton-to-neutron conversion, and various other nuclear processes happen, and therefore impact the production of light elements [30–32], including the primordial abundance of helium-4, Y_p . Annihilation products can additionally alter the radiation content in the universe, changing the effective number of light species N_{eff} [23]. Both of these effects are imprinted on the CMB anisotropies.

The current CMB constraints on the mass of light DM were derived using *Planck* data [22, 24, 25, 27, 33, 34], as well as the small-scale measurements from the Atacama Cosmology Telescope and South Pole Telescope [35], together with the primordial abundance measurements of helium and deuterium. The CMB bound on DM mass is primarily driven by the measurement of N_{eff} , and is stronger than the bounds derived from primordial abundance measurements from pristine Lyman- α systems, for most DM models. CMB-S4 will deliver a measurement of N_{eff} with percent-level precision, $\sigma(N_{\text{eff}}) = 0.03$, promising to put very stringent lower bounds on the mass of thermal-relic DM (or detect it), regardless of the details of its coupling to the Standard Model. Previous studies [35] found that the lower bounds on the mass from CMB-S4 will be able to exclude most of the mass range that could affect the process of BBN [29, 35].

4 Light (but massive) relics - LiMRs

There have been many proposed extensions of the Standard Model that introduce the existence of light, weakly interacting particles. A general category is the one of light (but massive) relics (or LiMRs), which are particles that were in thermal contact with the SM in the early universe and relativistic when they decoupled. The decoupling of these relics while relativistic gives these particles significant streaming motion, which sets a scale below which they cannot cluster, altering the large-scale structure (LSS) of our universe.

One of the key signatures of these new light species is their contribution to the early universe energy density as radiation, which alters the expansion history of the universe and the epoch of radiation-matter equality, and additionally manifests in the CMB by modifying the Silk damping tail. This signature of new dark radiation is commonly parametrized as ΔN_{eff} , and is quartically sensitive to the temperature of the light relic. In turn, measurements of the aforementioned scale-dependent suppression of large-scale structure are sensitive to both the mass and temperature of the LiMR [36], and thus our reach for new physics in this arena is greatly strengthened by joint analysis of CMB and LSS data.

A previous analysis of CMB temperature and polarization anisotropy measurements from the *Planck* satellite and full-shape LSS information from BOSS DR12 [37] sets limits of $m_X < 2.3$ eV (2σ) for fermionic relics decoupling at TeV-scale temperatures. A notable application for this constraint is for light gravitinos in gauge-mediated SUSY scenarios, where the scale of SUSY breaking is bounded from above at 70 TeV. The considerable improvement of CMB anisotropy measurements expected with CMB-S4, with a notably sharpened sensitivity to N_{eff} contributions, would provide significant additional power towards detecting cosmic LiMRs (or the absence thereof). For the illustrative example of the




Figure 2: Dark matter (DM) – proton cross section parameter space that can be probed with CMB-S4, for velocity-independent interactions. CMB-S4 is forecast [38] to improve existing *Planck* CMB bounds [39–41] by over an order of magnitude. CMB-S4 is complementary to direct-detection searches [42–44] and from the X-ray Quantum Calorimeter (XQC) [45], which have limited sensitivity to light (sub-GeV) and strongly-interacting DM. Large-scale structure limits come from the Lyman-alpha forest ($\text{Ly}\alpha\text{f}$) from SDSS-I [40] and from Keck-HIRES & VLT-UVES spectra [46]; and from the Milky Way (MW) satellite mass function [47–49].

gravitino, analysis of CMB-S4 and improved LSS data (such as from the forthcoming Dark Energy Spectroscopic Instrument - DESI) in tandem will yield unprecedented sensitivities: the ability to probe the entire parameter space at the 2σ level, and 3σ detection reach for all masses eV-scale and above.

5 Dark matter-baryon/electron interactions

A generic and well-studied feature of dark matter models is some form of elastic interaction with Standard Model particles, in particular electrons [49, 50] or nuclei [39–41, 46, 51–54]. These interactions are well constrained by terrestrial direct detection experiments,

but cosmological data can serve as a powerful complement in several ways: firstly, to extend experimental sensitivity to smaller ($m_\chi \ll \text{GeV}$) DM masses, where direct detection technologies are still developing; secondly, to increase our access to higher cross sections (such as would be expected for composite models) that would be lost to overburden and atmospheric scattering in underground experiments; and finally, in the case that a detection is made, to provide insight on how DM-SM interaction strength scales with energy and momenta transfer.

In the early universe, scattering with baryons or electrons may thermalize the dark matter distribution, washing out density fluctuations and impeding the growth of structure. The decreased clustering of matter imprints on the Cosmic Microwave Background as a suppression of temperature and polarization anisotropies at small scales, and the modified relative velocity between DM and the SM plasma further induces a shift in the scale of acoustic oscillations. As such, the increased precision of CMB-S4 temperature and polarization data, in addition to gravitational lensing at $\sim \text{arcmin}$ angular scales [38], will drastically improve the competitiveness of cosmological data in constraining dark matter physics, as well as access significant regions of parameter space that have been heretofore unprobed. For instance, for a velocity-independent cross section and 1 GeV dark matter mass, CMB-S4 will set a bound on the scattering with baryons of $\sigma < 6 \times 10^{-27} \text{cm}^2$ at 95% C.L. [38] (see Fig. 2).

6 Dark matter freeze-in

Dark matter could achieve its observed abundance via annihilation or decay of Standard Model particles in the early universe without thermalizing with the plasma. This is a generic mechanism known as DM “freeze-in” (see e.g. Refs. [55, 56]). The parameter space for freeze-in represents the smallest relevant coupling the DM can have with the SM plasma that affects early Universe observables. If freeze-in DM couples to the Standard Model via a light mediator, most DM is produced at late times, and so in this sense freeze-in is insensitive to initial conditions. If the light mediator is the Standard Model photon or a kinetically-mixed dark photon (which is likely given strong constraints on other light mediators [57, 58]), then DM becomes effectively charged. The freeze-in scenario is one of the few effective ways of making charged DM given the exclusion of the freeze-out scenario [59]. Freeze-in of effectively charged DM is a key target benchmark for many proposed and ongoing direct detection experiments [60–73]. Additionally, since the DM is non-thermal in this scenario, freeze-in is one of the few allowed mechanisms for making sub-MeV DM from the SM plasma while still remaining consistent with bounds from BBN and N_{eff} on thermal dark sectors (see the discussion in Section 3).

The portal responsible for DM production necessarily implies a DM-baryon interaction, creating effects that are similar to those described in Section 5 that are highly complementary to direct detection experiments. However, freeze-in requires a different analysis because scattering via a light mediator scales steeply with velocity as v^{-4} and therefore the highly non-thermal phase space distribution must be taken into account to derive accurate limits [74]. Previous studies showed that current constraints from CMB measurements from the *Planck* satellite set a freeze-in DM mass bound with $m_\chi \lesssim 18 \text{keV}$ being excluded

at the 95% C.L., while CMB-S4 will be sensitive to masses up to $m_\chi \approx 29$ keV [75].

7 Dark matter-neutrino interactions

Dark matter interactions with Standard Model particles have received plenty of attention in the literature. One particularly intriguing scenario is the scattering of DM with Standard-Model neutrinos in the early universe [76–83]. In this case, DM interactions lead to a suppression of the primordial density fluctuations through collisional damping [76, 78], which leaves noticeable signatures in the CMB anisotropy power spectra and the matter power spectrum, and ultimately affects the large-scale structure of the universe we observe today [84–86].

DM-neutrino interactions can affect CMB temperature and polarization power spectra by increasing the amplitude of the CMB acoustic peaks, and shifting them to higher multipoles ℓ , with respect to the non-interaction scenario [83, 84, 87]. Previous CMB analyses suggest that the constraints on cosmological parameters (in particular, DM–neutrino scattering cross section, and the sum of the neutrino masses), which rely primarily on measurements from *Planck* at this time, could improve significantly with more accurate temperature and polarization data from the next-generation ground-based experiments. We expect the high-resolution measurements with CMB-S4 would increase the sensitivity to DM-neutrino interactions.

8 Dark matter-dark radiation interaction

In many theories (see e.g. Refs. [88–95]), DM is part of an extended dark sector in which new interactions, such as a dark $U(1)$ force [96–98], can couple the DM to relativistic degrees of freedom at early times. DM interacting with such dark radiation affects the CMB in two distinct ways. First, the presence of this extra dark radiation affects the expansion history of the universe, affects the Silk damping tail of the CMB (similarly to the impact of N_{eff}), generates an early Integrated Sachs-Wolfe (ISW) effect, and modifies the redshift of matter-radiation equality. However, unlike the standard free-streaming neutrinos, this dark radiation is tightly-coupled (fluid-like) at early cosmological epochs, hence resulting in different phase and amplitude shift to the CMB acoustic peaks [99–102] for modes entering the causal horizon before the onset of dark radiation free-streaming.

Second, the large dark radiation pressure prohibits the growth of dark matter fluctuations on scales entering the horizon before kinematic decoupling of this DM-dark radiation fluid. On these characteristic length scales, the presence of dark acoustic oscillations (DAOs) and Silk-like damping decreases the depth of the gravitational potential near recombination, reducing the source term of CMB temperature fluctuations. In addition to these impacts on the primary CMB, the presence of DAOs and the small-scale damping of matter fluctuation will affect the lensing of the CMB photons as they travel towards us from the last scattering surface.

The DM-dark radiation models with the greatest deviations from the standard picture have already been constrained with data from the *Planck* satellite [103, 104]. In general,




Figure 3: Fractional difference of the CMB lensing spectrum $C_L^{\phi\phi}$ between a standard Λ CDM model (with massless neutrinos) and four different interacting DM models (“ETHOS” models), each characterized by a different interacting fraction. For comparison, we also display a standard massive neutrino model with $\sum m_\nu = 0.06$ eV. The parameter $a_4 = 3 \times 10^2 \text{ Mpc}^{-1}$ characterizes the strength of the interaction between DM and dark radiation [105].

models in which all of the DM couples to dark radiation at the time when Fourier modes probed by the CMB enter the horizon are ruled out. This leaves scenarios in which only a small fraction of the DM couples to dark radiation as viable candidate to be probed by CMB-S4, which will strongly constrain any models that are currently still allowed given *Planck* data.

As an illustration of the reach of CMB-S4, we show in Fig. 3 the impact of different interacting DM models on the CMB lensing power spectrum. We can see here the fractional difference between interacting DM models and the standard Λ CDM case with the sum of neutrino masses $\sum m_\nu = 0$, for models where the fraction of interacting DM f_{int} is varied from 2% to 5%, highlighting the fact that CMB-S4 will rule out models with fractions as low as 2%. Importantly, the interacting DM-dark radiation model predicts a different shape to the lensing power spectrum than that of massive neutrinos, highlighting that CMB-S4 will be able to distinguish between these two possibilities.

9 Axions and other ultra-light bosons

Ultra-light axions (ULAs) are an attractive candidate for dark matter [106–112] due to the fact that they behave like Λ CDM on large scales but also alleviate challenges to Λ CDM on small scales [108, 113]. Furthermore, they can also behave as an early dark energy component to mitigate differences between high-redshift and local measurements of the Hubble constant H_0 [114].

The key feature of ultra-light axions dark matter is that they have de Broglie wavelengths that are macroscopic, which would generate cored galactic halo density profiles

on scales of $\sim 0.7 (m_\phi/10^{-22} \text{ eV})^{-1/2} \text{ kpc}$, where m_ϕ is the mass of the axion. This feature has gone some way to resolving the cusp-core conundrum of Milky-Way dwarf spheroidal galaxies [108, 115–122]. As discussed in Ref. [118], a dynamical analysis of Fornax and Sculptor suggests that $0.3 \times 10^{-22} \text{ eV} \lesssim m_\phi \lesssim 10^{-22} \text{ eV}$ can fit the profile of these dwarf spheroidal galaxies, however some tensions remain in solving both the ultra-faint dwarf satellites and the dwarf spheroidal population [123, 124].

While baryonic effects have been suggested to mitigate challenges to Λ CDM at small scales, ULAs have been claimed to partially resolve these issues by suppressing the number of low-mass subhalos around Milky-Way scale halos and by the depression of satellite galaxy masses in halos relative to Λ CDM expectations [117, 125–127].

Depending on their assumed mass, ULAs affect the CMB power spectrum on large scales through the ISW effect, and on small scales in the damping tail. The most sensitive probe of large-scale structure from measurements of the CMB is through the measurement of the lensing deflection on small scales. It is this regime that will see the largest improvement with future CMB experiments such as CMB-S4, as shown in Figure 4. In addition, CMB photons may be Compton re-scattered by electrons along the line of sight. When this occurs in galaxy clusters, the resulting kinetic Sunyaev-Zel’dovich (kSZ) induces additional CMB temperature anisotropies that probe the real-time growth of structure, and can be detected via cross-correlation with galaxy surveys. ULAs will boost this signal (due to increased bias in a structure-suppressing cosmology), yielding a potential improvement in sensitivity [128]. Even better sensitivity is possible using the Ostriker-Vishniac (OV) CMB anisotropies, induced by gas inhomogeneities during the mildly non-linear reionization epoch.

In contrast to other DM candidates, the ULA field could be excited prior to thermalization of the cosmic plasma. In particular, if the $U(1)$ symmetry of ULAs is broken before the end of inflation, isocurvature fluctuations would be excited [129–135], with an amplitude determined by the Hubble parameter H_I during inflation and m_ϕ [136–140]. If H_I can be independently determined using CMB B-mode polarization anisotropies (e.g., by future experiments like CMB-S4 itself [1], Simons Observatory [141], or LiteBIRD [142]) via the production of tensor modes [143, 144], a ULA energy density contribution of $\Omega_\phi \sim 0.01$ could be detected by CMB-S4 [1, 145]. Current constraints on isocurvature and tensor modes allow a fractional ULA contribution of $\sim 10\%$ to the total DM budget [145, 146].

Alternatively, if the ULA $U(1)$ symmetry is broken after the end of inflation, a white-noise power spectrum of isocurvature would be produced. Depending on details of ULA production (such as thermal corrections to the ULA mass), CMB-S4 sensitivity to isocurvature fluctuations (when combined with CMB lensing, and independent probes of galaxy clustering and lensing shear) could test ULA masses as high as $m_\phi \simeq 10^{-17} \text{ eV}$ [147].

10 Conclusion

CMB measurements provide a means to assess the impact of dark matter across the whole of cosmic history. The sensitivity of the CMB measurements to the dark sector do not rely on assumptions about the local dark matter distribution and are insensitive to the details of astrophysical modeling that are necessary for some indirect searches. We summarize




Figure 4: Ultra-light boson dark matter (UDM) parameter space that can be probed with CMB-S4, compared to other current (*solid*) and projected (*lines*) cosmological bounds. *In turquoise*, UDM parameter space probed by kinetic Sunyaev-Zel’dovich measurements combining CMB-S4 data (including primary anisotropies) with large-scale structure information from the Dark Energy Spectroscopic Instrument (kSZ-S4) [128]; *in blue*, forecast CMB-S4 UDM sensitivity using temperature, polarization and lensing data [146]; *in green*, forecast Ostriker-Vishniac effect bounds from CMB-S4 (OV-S4) [128]. Existing CMB bounds come from *Planck* [114, 148], while LSS bounds come from galaxy clustering combined with *Planck* (+BOSS) [149], galaxy weak lensing combined with *Planck* (+DES) [150], the UV luminosity function and optical depth to reionization (Rei.) [151], and the Lyman-alpha forest (Lyaf) [152–156].

Model	Limit
DM annihilation	$m_\chi > 30 - 50$ GeV (thermal cross section, 20% of energy deposition)
Thermal DM mass	$m_\chi > 10 - 15$ MeV (depending on DM models)
LiMRs	3σ (non-)detection for $m_\chi > \mathcal{O}(1$ eV), in combination with DESI
DM-baryon scattering	$\sigma < 6 \times 10^{-27}$ cm ² (for 1 GeV DM)
Sub-MeV DM freeze-in	$m_\chi > 29$ keV
DM-DR interaction	$f_{\text{int}} < 2\%$
ULA: 10^{-31} eV $< m_\phi < 10^{-24.5}$	$\Omega_\phi < 3 \times 10^{-3}$

Table 1: Summary of sensitivity for different dark matter scenarios using CMB-S4.

the key results for the dark matter scenarios discussed in this paper in Table 1.

CMB-S4 will probe deep into the damping tail of the CMB fluctuations, where much of the imprints from different dark matter scenarios deviates most strongly from the stan-

standard Λ CDM picture. Additionally, the improved measurements of the CMB lensing power spectrum will provide insights into the clustering of matter across a wide range of scales. Measurements of matter clustering are valuable for deriving constraints on dark matter physics, since clustering is expected to be suppressed at various scales for many dark matter scenarios.

CMB-S4 will dramatically improve the precision with which we measure CMB anisotropies. This leap in sensitivity will enable new insights into the nature of dark matter, complementary to what is expected from laboratory experiments and other astrophysical searches.

Acknowledgments

CD is partially supported by Department of Energy (DOE) grant DE-SC0020223. RH is a CIFAR Azrieli Global in the Gravity and the Extreme Universe program, and a Sloan Fellow. The Dunlap Institute is funded through an endowment established by the David Dunlap family and the University of Toronto. Canadian co-authors acknowledge support from the Natural Sciences and Engineering Research Council of Canada. Computations were performed on the SciNet supercomputer at the SciNet HPC Consortium. SciNet is funded by: the Canada Foundation for Innovation; the Government of Ontario; Ontario Research Fund - Research Excellence; and the University of Toronto. KB acknowledges support from the NSF Grant No. PHY-2112884. FYCR was supported in part by the NSF grant No. AST-2008696. GSF acknowledges support through the Isaac Newton Studentship and the Helen Stone Scholarship at the University of Cambridge. VG and RA acknowledge the support from NASA through the Astrophysics Theory Program, Award Number 21-ATP21-0135. DJEM is supported by an STFC Ernest Rutherford Fellowship. JM is supported by the US Department of Energy under Grant DE-SC0010129.

References

- [1] K. Abazajian et al. CMB-S4 Science Case, Reference Design, and Project Plan. *arXiv e-prints*, pp. arXiv:1907.04473, July 2019.
- [2] A. Lewis and A. Challinor. Weak gravitational lensing of the cmb. *Phys. Rept.*, 429:1–65, 2006.
- [3] R. A. Sunyaev and Y. B. Zeldovich. The Velocity of clusters of galaxies relative to the microwave background. The Possibility of its measurement. *Mon. Not. Roy. Astron. Soc.*, 190:413–420, 1980.
- [4] R. A. Sunyaev and Y. B. Zeldovich. Microwave background radiation as a probe of the contemporary structure and history of the universe. *Ann. Rev. Astron. Astrophys.*, 18:537–560, 1980.
- [5] X.-L. Chen and M. Kamionkowski. Particle decays during the cosmic dark ages. *Phys. Rev.*, D70:043502, 2004.

- [6] N. Padmanabhan and D. P. Finkbeiner. Detecting dark matter annihilation with CMB polarization: Signatures and experimental prospects. *Phys. Rev.*, D72:023508, 2005.
- [7] N. Aghanim et al. Planck 2018 results. VI. Cosmological parameters. *Astron. Astrophys.*, 641:A6, 2020. [Erratum: *Astron. Astrophys.* 652, C4 (2021)].
- [8] T. R. Slatyer and C.-L. Wu. General Constraints on Dark Matter Decay from the Cosmic Microwave Background. *Phys. Rev. D*, 95(2):023010, 2017.
- [9] M. S. Madhavacheril et al. Current Dark Matter Annihilation Constraints from CMB and Low-Redshift Data. *Phys. Rev.*, D89:103508, 2014.
- [10] W. L. K. Wu et al. A Guide to Designing Future Ground-based Cosmic Microwave Background Experiments. *Astrophys. J.*, 788:138, 2014.
- [11] D. Green et al. Aspects of Dark Matter Annihilation in Cosmology. *JCAP*, 04:025, 2019.
- [12] D. Hooper and K. M. Zurek. Natural supersymmetric model with MeV dark matter. *PhRvD*, 77(8):087302, April 2008.
- [13] J. L. Feng and J. Kumar. Dark-Matter Particles without Weak-Scale Masses or Weak Interactions. *PhRvL*, 101(23):231301, December 2008.
- [14] D. Hooper et al. MeV dark matter and small scale structure. *PhRvD*, 76(10):103515, November 2007.
- [15] C. Boehm and P. Fayet. Scalar dark matter candidates. *Nuclear Physics B*, 683(1):219–263, April 2004.
- [16] C. Boehm et al. MeV Dark Matter: Has It Been Detected? *PhRvL*, 92(10):101301, March 2004.
- [17] D. Hooper et al. Possible Evidence for MeV Dark Matter in Dwarf Spheroidals. *PhRvL*, 93(16):161302, October 2004.
- [18] C. Boehm et al. Light and heavy dark matter particles. *PhRvD*, 69(10):101302, May 2004.
- [19] G. Jungman et al. Supersymmetric dark matter. *PhR*, 267:195–373, March 1996.
- [20] C. Boehm et al. Increasing N_{eff} with particles in thermal equilibrium with neutrinos. *JCAP*, 2012(12):027, December 2012.
- [21] L. Bergström. Dark matter evidence, particle physics candidates and detection methods. *Annalen der Physik*, 524(9-10):479–496, October 2012.
- [22] C. Boehm et al. A lower bound on the mass of cold thermal dark matter from Planck. *JCAP*, 2013(8):041, August 2013.

- [23] G. Steigman. Equivalent neutrinos, light WIMPs, and the chimera of dark radiation. *PhRvD*, 87(10):103517, May 2013.
- [24] K. M. Nollett and G. Steigman. BBN and the CMB constrain light, electromagnetically coupled WIMPs. *PhRvD*, 89(8):083508, April 2014.
- [25] K. M. Nollett and G. Steigman. BBN and the CMB constrain neutrino coupled light WIMPs. *PhRvD*, 91(8):083505, April 2015.
- [26] R. J. Wilkinson et al. Ruling out the light weakly interacting massive particle explanation of the Galactic 511 keV line. *PhRvD*, 94(10):103525, November 2016.
- [27] M. Escudero. Neutrino decoupling beyond the Standard Model: CMB constraints on the Dark Matter mass with a fast and precise N_{eff} evaluation. *JCAP*, 2019(2):007, February 2019.
- [28] P. F. Depta et al. BBN constraints on the annihilation of MeV-scale dark matter. *JCAP*, 2019(4):029, April 2019.
- [29] N. Sabti et al. Refined bounds on MeV-scale thermal dark sectors from BBN and the CMB. *JCAP*, 2020(1):004, January 2020.
- [30] E. W. Kolb et al. Effect of interacting particles on primordial nucleosynthesis. *PhRvD*, 34(8):2197–2205, October 1986.
- [31] P. D. Serpico and G. G. Raffelt. MeV-mass dark matter and primordial nucleosynthesis. *PhRvD*, 70(4):043526, August 2004.
- [32] C. Boehm et al. Can annihilating dark matter be lighter than a few GeVs? *Journal of Physics G Nuclear Physics*, 30(3):279–285, March 2004.
- [33] G. Steigman and K. M. Nollett. Light WIMPs, equivalent neutrinos, BBN, and the CMB. *MmSAI*, 85:175, January 2014.
- [34] C. Giovanetti et al. Joint CMB and BBN Constraints on Light Dark Sectors with Dark Radiation. *arXiv e-prints*, pp. arXiv:2109.03246, September 2021.
- [35] R. An et al. What does cosmology tell us about the mass of thermal-relic dark matter? *arXiv e-prints*, pp. arXiv:2202.03515, February 2022.
- [36] N. DePorzio et al. Finding eV-scale light relics with cosmological observables. *Phys. Rev. D*, 103(2):023504, 2021.
- [37] W. L. Xu et al. Cosmological Constraints on Light (but Massive) Relics. 7 2021.
- [38] Z. Li et al. Disentangling Dark Physics with Cosmic Microwave Background Experiments. *Phys. Rev. D*, 98(12):123524, 2018.
- [39] K. K. Boddy and V. Gluscevic. First cosmological constraint on the effective theory of dark matter-proton interactions. *Phys. Rev. D*, 98:083510, Oct 2018.

- [40] W. L. Xu et al. Probing sub-GeV dark matter-baryon scattering with cosmological observables. *PhRvD*, 97:103530, May 2018.
- [41] C. Dvorkin et al. Constraining Dark Matter-Baryon Scattering with Linear Cosmology. *Phys. Rev. D*, 89(2):023519, 2014.
- [42] T. Emken and C. Kouvaris. How blind are underground and surface detectors to strongly interacting dark matter? *Phys. Rev. D*, 97:115047, Jun 2018.
- [43] G. Angloher et al. Results on MeV-scale dark matter from a gram-scale cryogenic calorimeter operated above ground. *European Physical Journal C*, 77(9):637, September 2017.
- [44] E. Armengaud et al. Searching for low-mass dark matter particles with a massive Ge bolometer operated above ground. *PhRvD*, 99(8):082003, April 2019.
- [45] M. S. Mahdawi and G. R. Farrar. Constraints on Dark Matter with a moderately large and velocity-dependent DM-nucleon cross-section. *JCAP*, 2018(10):007, October 2018.
- [46] K. K. Rogers et al. New limits on light dark matter - proton cross section from the cosmic large-scale structure. 11 2021.
- [47] K. Maamari et al. Bounds on Velocity-dependent Dark Matter-Proton Scattering from Milky Way Satellite Abundance. *ApJL*, 907(2):L46, February 2021.
- [48] E. O. Nadler et al. Constraints on Dark Matter Properties from Observations of Milky Way Satellite Galaxies. *PhRvL*, 126(9):091101, March 2021.
- [49] M. A. Buen-Abad et al. Cosmological Constraints on Dark Matter Interactions with Ordinary Matter. 7 2021.
- [50] D. V. Nguyen et al. Observational constraints on dark matter scattering with electrons. *Phys. Rev. D*, 104(10):103521, 2021.
- [51] X.-l. Chen et al. Cosmic microwave background and large scale structure limits on the interaction between dark matter and baryons. *Phys. Rev. D*, 65:123515, 2002.
- [52] V. Gluscevic and K. K. Boddy. Constraints on Scattering of keV–TeV Dark Matter with Protons in the Early Universe. *Phys. Rev. Lett.*, 121(8):081301, 2018.
- [53] T. R. Slatyer and C.-L. Wu. Early-Universe constraints on dark matter-baryon scattering and their implications for a global 21 cm signal. *Phys. Rev. D*, 98(2):023013, 2018.
- [54] K. K. Boddy et al. Critical assessment of CMB limits on dark matter-baryon scattering: New treatment of the relative bulk velocity. *Phys. Rev. D*, 98(12):123506, 2018.
- [55] L. J. Hall et al. Freeze-In Production of FIMP Dark Matter. *JHEP*, 03:080, 2010.

- [56] N. Bernal et al. The Dawn of FIMP Dark Matter: A Review of Models and Constraints. *Int. J. Mod. Phys. A*, 32(27):1730023, 2017.
- [57] S. Knapen et al. Light Dark Matter: Models and Constraints. *Phys. Rev. D*, 96(11):115021, 2017.
- [58] D. Green and S. Rajendran. The Cosmology of Sub-MeV Dark Matter. *JHEP*, 10:013, 2017.
- [59] S. D. McDermott et al. Turning off the Lights: How Dark is Dark Matter? *Phys. Rev. D*, 83:063509, 2011.
- [60] R. Essig et al. Direct Detection of Sub-GeV Dark Matter. *Phys. Rev.*, D85:076007, 2012.
- [61] R. Essig et al. First Direct Detection Limits on sub-GeV Dark Matter from XENON10. *Phys. Rev. Lett.*, 109:021301, 2012.
- [62] R. Essig et al. Direct Detection of sub-GeV Dark Matter with Semiconductor Targets. *JHEP*, 05:046, 2016.
- [63] A. Aguilar-Arevalo et al. Constraints on Light Dark Matter Particles Interacting with Electrons from DAMIC at SNOLAB. *Phys. Rev. Lett.*, 123(18):181802, 2019.
- [64] L. Barak et al. SENSEI: Direct-Detection Results on sub-GeV Dark Matter from a New Skipper-CCD. 4 2020.
- [65] D. Amaral et al. Constraints on low-mass, relic dark matter candidates from a surface-operated SuperCDMS single-charge sensitive detector. 5 2020.
- [66] P. Agnes et al. Constraints on Sub-GeV Dark-Matter–Electron Scattering from the DarkSide-50 Experiment. *Phys. Rev. Lett.*, 121(11):111303, 2018.
- [67] Y. Hochberg et al. Detection of sub-MeV Dark Matter with Three-Dimensional Dirac Materials. *Phys. Rev.*, D97(1):015004, 2018.
- [68] R. M. Geilhufe et al. Dirac Materials for Sub-MeV Dark Matter Detection: New Targets and Improved Formalism. *Phys. Rev. D*, 101(5):055005, 2020.
- [69] A. Coskuner et al. Directional Dark Matter Detection in Anisotropic Dirac Materials. 9 2019.
- [70] S. Knapen et al. Detection of Light Dark Matter With Optical Phonons in Polar Materials. 2017.
- [71] S. Griffin et al. Directional Detection of Light Dark Matter with Polar Materials. 2018.
- [72] S. M. Griffin et al. Multichannel direct detection of light dark matter: Target comparison. *Phys. Rev. D*, 101(5):055004, 2020.

- [73] A. Berlin et al. Directly Deflecting Particle Dark Matter. *Phys. Rev. Lett.*, 124(1):011801, 2020.
- [74] C. Dvorkin et al. Making dark matter out of light: freeze-in from plasma effects. *Phys. Rev. D*, 99(11):115009, 2019.
- [75] C. Dvorkin et al. Cosmology of Sub-MeV Dark Matter Freeze-In. *Phys. Rev. Lett.*, 127(11):111301, 2021.
- [76] C. Boehm et al. Constraining dark matter candidates from structure formation. *Physics Letters B*, 518(1-2):8–14, October 2001.
- [77] C. Boehm et al. WIMP matter power spectra and small scale power generation. *arXiv e-prints*, pp. astro-ph/0309652, September 2003.
- [78] C. Boehm and R. Schaeffer. Constraints on Dark Matter interactions from structure formation: damping lengths. *A&A*, 438(2):419–442, August 2005.
- [79] P. Serra et al. Constraints on neutrino-dark matter interactions from cosmic microwave background and large scale structure data. *PhRvD*, 81(4):043507, February 2010.
- [80] G. Mangano et al. Cosmological bounds on dark-matter-neutrino interactions. *PhRvD*, 74(4):043517, August 2006.
- [81] Y. Ali-Haïmoud et al. Constraints on Dark Matter Interactions with Standard Model Particles from Cosmic Microwave Background Spectral Distortions. *PhRvL*, 115(7):071304, August 2015.
- [82] E. Di Valentino et al. Reducing the H_0 and σ_8 tensions with dark matter-neutrino interactions. *PhRvD*, 97(4):043513, February 2018.
- [83] M. R. Mosbech et al. The full Boltzmann hierarchy for dark matter-massive neutrino interactions. *JCAP*, 2021(3):066, March 2021.
- [84] R. J. Wilkinson et al. Constraining dark matter-neutrino interactions using the CMB and large-scale structure. *JCAP*, 2014(5):011, May 2014.
- [85] A. Olivares-Del Campo et al. Dark matter-neutrino interactions through the lens of their cosmological implications. *PhRvD*, 97(7):075039, April 2018.
- [86] D. C. Hooper and M. Lucca. Hints of dark matter-neutrino interactions in Lyman- α data. *arXiv e-prints*, pp. arXiv:2110.04024, October 2021.
- [87] A. Paul et al. Shedding light on dark matter and neutrino interactions from cosmology. *JCAP*, 2021(10):017, October 2021.
- [88] R. Foot. Mirror matter-type dark matter. *Int. J. Mod. Phys.*, D13:2161–2192, 2004.
- [89] D. E. Kaplan et al. Atomic Dark Matter. *icap*, 1005:021, 2010.

- [90] D. E. Kaplan et al. Dark Atoms: Asymmetry and Direct Detection. *JCAP*, 1110:011, 2011.
- [91] F.-Y. Cyr-Racine and K. Sigurdson. Cosmology of atomic dark matter. *Phys. Rev.*, D87(10):103515, 2013.
- [92] L. G. van den Aarssen et al. Is dark matter with long-range interactions a solution to all small-scale problems of Λ CDM cosmology? *Phys. Rev. Lett.*, 109:231301, 2012.
- [93] J. M. Cline et al. Scattering properties of dark atoms and molecules. *Phys. Rev. D*, 89(4):043514, 2014.
- [94] M. A. Buen-Abad et al. Non-Abelian dark matter and dark radiation. *Phys. Rev.*, D92(2):023531, 2015.
- [95] J. Lesgourgues et al. Evidence for dark matter interactions in cosmological precision data? *JCAP*, 1602(02):037, 2016.
- [96] L. Ackerman et al. Dark Matter and Dark Radiation. *Phys. Rev. D*, 79:023519, 2009.
- [97] J. L. Feng et al. Hidden Charged Dark Matter. *jcap*, 0907:004, 2009.
- [98] P. Agrawal et al. Make Dark Matter Charged Again. *JCAP*, 1705(05):022, 2017.
- [99] S. Bashinsky and U. Seljak. Neutrino perturbations in CMB anisotropy and matter clustering. *Phys. Rev.*, D69:083002, 2004.
- [100] F.-Y. Cyr-Racine and K. Sigurdson. Limits on Neutrino-Neutrino Scattering in the Early Universe. *Phys. Rev.*, D90(12):123533, 2014.
- [101] B. Follin et al. First Detection of the Acoustic Oscillation Phase Shift Expected from the Cosmic Neutrino Background. *Phys. Rev. Lett.*, 115(9):091301, 2015.
- [102] D. Baumann et al. Phases of New Physics in the CMB. *JCAP*, 1601:007, 2016.
- [103] F.-Y. Cyr-Racine et al. Constraints on Large-Scale Dark Acoustic Oscillations from Cosmology. *Phys. Rev. D*, 89(6):063517, 2014.
- [104] M. Archidiacono et al. Constraining Dark Matter-Dark Radiation interactions with CMB, BAO, and Lyman- α . *JCAP*, 10:055, 2019.
- [105] F.-Y. Cyr-Racine et al. ETHOS—an effective theory of structure formation: From dark particle physics to the matter distribution of the Universe. *Phys. Rev.*, D93(12):123527, 2016.
- [106] T. Matos and J. A. Nieto. Topics on Kaluza-Klein theory. *Revista Mexicana de Fisica*, 39:81–131, August 1993.
- [107] T. Matos et al. Scalar field as dark matter in the universe. *Classical and Quantum Gravity*, 17(7):1707–1712, April 2000.

- [108] W. Hu et al. Cold and fuzzy dark matter. *Phys. Rev. Lett.*, 85:1158–1161, 2000.
- [109] A. Suárez and T. Matos. Structure formation with scalar-field dark matter: the fluid approach. *MNRAS*, 416(1):87–93, September 2011.
- [110] J.-C. Hwang and H. Noh. Axion as a cold dark matter candidate. *Physics Letters B*, 680(1):1–3, September 2009.
- [111] C.-G. Park et al. Axion as a cold dark matter candidate: Low-mass case. *PhRvD*, 86(8):083535, October 2012.
- [112] J. Magaña et al. Structure formation with scalar field dark matter: the field approach. *JCAP*, 2012(10):003, October 2012.
- [113] J. S. Bullock and M. Boylan-Kolchin. Small-Scale Challenges to the Λ CDM Paradigm. *ARA&A*, 55(1):343–387, August 2017.
- [114] V. Poulin et al. Cosmological implications of ultralight axionlike fields. *PhRvD*, 98(8):083525, October 2018.
- [115] C. H. Lee and S. J. Sin. Quantum aspect of dark matter and star formation in elliptical galaxies. *Journal of Korean Physical Society*, 28(1):16–19, February 1995.
- [116] A. Arbey et al. Galactic halos of fluid dark matter. *PhRvD*, 68(2):023511, July 2003.
- [117] D. J. E. Marsh and J. Silk. A Model For Halo Formation With Axion Mixed Dark Matter. *Mon. Not. Roy. Astron. Soc.*, 437(3):2652–2663, 2014.
- [118] D. J. E. Marsh and A.-R. Pop. Axion dark matter, solitons and the cusp-core problem. *MNRAS*, 451(3):2479–2492, August 2015.
- [119] L. A. Ureña-López and A. X. Gonzalez-Morales. Towards accurate cosmological predictions for rapidly oscillating scalar fields as dark matter. *JCAP*, 2016(7):048, July 2016.
- [120] T. Bernal et al. Scalar field dark matter in clusters of galaxies. *MNRAS*, 468(3):3135–3149, July 2017.
- [121] H. Deng et al. Can light dark matter solve the core-cusp problem? *PhRvD*, 98(2):023513, July 2018.
- [122] T. Broadhurst et al. Ghostly galaxies as solitons of Bose-Einstein dark matter. *PhRvD*, 101(8):083012, April 2020.
- [123] M. Safarzadeh and D. N. Spergel. Ultra-light Dark Matter Is Incompatible with the Milky Way’s Dwarf Satellites. *ApJ*, 893(1):21, April 2020.
- [124] E. Calabrese and D. N. Spergel. Ultra-light dark matter in ultra-faint dwarf galaxies. *MNRAS*, 460(4):4397–4402, August 2016.

- [125] D. J. E. Marsh. Axion Cosmology. *Phys. Rept.*, 643:1–79, 2016.
- [126] D. J. E. Marsh. WarmAndFuzzy: the halo model beyond CDM. 2016.
- [127] L. Hui et al. Ultralight scalars as cosmological dark matter. *PhRvD*, 95(4):043541, February 2017.
- [128] G. S. Farren et al. Ultra-light axions and the kinetic Sunyaev-Zel’dovich Effect. *arXiv e-prints*, pp. arXiv:2109.13268, September 2021.
- [129] M. Axenides et al. Development of Axion Perturbations in an Axion Dominated Universe. *Phys. Lett.*, B126:178, 1983.
- [130] M. S. Turner et al. Formation of Structure in an Axion Dominated Universe. *Phys. Lett. B*, 125:35, 1983. [Erratum: *Phys.Lett.B* 125, 519 (1983)].
- [131] D. H. Lyth. Axions and inflation: Sitting in the vacuum. *Phys. Rev. D*, 45:3394–3404, 1992.
- [132] P. Fox et al. Probing a QCD string axion with precision cosmological measurements. 2004.
- [133] M. P. Hertzberg et al. Axion Cosmology and the Energy Scale of Inflation. *Phys.Rev.*, D78:083507, 2008.
- [134] E. Komatsu et al. Five-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Interpretation. *Astrophys. J. Suppl.*, 180:330–376, 2009.
- [135] D. Langlois. Lectures on inflation and cosmological perturbations. *Lect. Notes Phys.*, 800:1–57, 2010.
- [136] L. Visinelli and P. Gondolo. Dark Matter Axions Revisited. *Phys. Rev. D*, 80:035024, 2009.
- [137] D. J. E. Marsh et al. Axiverse cosmology and the energy scale of inflation. *Phys. Rev.*, D87:121701, 2013.
- [138] D. J. E. Marsh et al. Tensor Interpretation of BICEP2 Results Severely Constrains Axion Dark Matter. *Phys. Rev. Lett.*, 113(1):011801, 2014.
- [139] L. Visinelli and P. Gondolo. Axion cold dark matter in view of BICEP2 results. *Phys. Rev. Lett.*, 113:011802, 2014.
- [140] L. Visinelli. Light axion-like dark matter must be present during inflation. *Phys. Rev. D*, 96(2):023013, 2017.
- [141] N. Galitzki. The Simons Observatory: Project Overview. 2018.
- [142] A. Suzuki et al. The LiteBIRD Satellite Mission: Sub-Kelvin Instrument. *Journal of Low Temperature Physics*, 193(5-6):1048–1056, December 2018.

- [143] U. Seljak and M. Zaldarriaga. Signature of gravity waves in polarization of the microwave background. *Phys.Rev.Lett.*, 78:2054–2057, 1997.
- [144] M. Kamionkowski et al. Statistics of cosmic microwave background polarization. *Phys. Rev. D*, 55:7368–7388, 1997.
- [145] R. Hlozek et al. Using the Full Power of the Cosmic Microwave Background to Probe Axion Dark Matter. *Mon. Not. Roy. Astron. Soc.*, 476(3):3063–3085, 2018.
- [146] R. Hložek et al. Future CMB tests of dark matter: ultra-light axions and massive neutrinos. 2016.
- [147] M. Feix et al. Post-inflationary axion isocurvature perturbations facing CMB and large-scale structure. *JCAP*, 11:046, 2020.
- [148] R. Hložek et al. A search for ultralight axions using precision cosmological data. *Phys. Rev.*, D91(10):103512, 2015.
- [149] A. Laguë et al. Constraining ultralight axions with galaxy surveys. *JCAP*, 2022(1):049, January 2022.
- [150] M. Dentler et al. Fuzzy Dark Matter and the Dark Energy Survey Year 1 Data. *arXiv e-prints*, pp. arXiv:2111.01199, November 2021.
- [151] B. Bozek et al. Galaxy UV-luminosity function and reionization constraints on axion dark matter. *Mon. Not. Roy. Astron. Soc.*, 450(1):209–222, 2015.
- [152] K. K. Rogers and H. V. Peiris. Strong Bound on Canonical Ultralight Axion Dark Matter from the Lyman-Alpha Forest. *PhRvL*, 126(7):071302, February 2021.
- [153] K. K. Rogers and H. V. Peiris. General framework for cosmological dark matter bounds using N -body simulations. *PhRvD*, 103(4):043526, February 2021.
- [154] T. Kobayashi et al. Lyman- α constraints on ultralight scalar dark matter: Implications for the early and late universe. *Phys. Rev. D*, 96(12):123514, 2017.
- [155] V. Iršič et al. First Constraints on Fuzzy Dark Matter from Lyman- α Forest Data and Hydrodynamical Simulations. *PhRvL*, 119(3):031302, July 2017.
- [156] E. Armengaud et al. Constraining the mass of light bosonic dark matter using SDSS Lyman- α forest. *MNRAS*, 471(4):4606–4614, November 2017.