

# NON-LOCAL HEAT EQUATION WITH MOVING BOUNDARY AND CURVE-CROSSING OF DELAYED BROWNIAN MOTION

GIACOME ASCIONE, PIERRE PATIE, AND BRUNO TOALDO

ABSTRACT. In this paper we consider a non-local (in time) heat equation on a time-increasing parabolic set whose boundary is determined by a suitable curve. We provide a notion of solution for this equation and we study well-posedness under Dirichlet conditions outside the domain. A maximum principle is proved and used to derive uniqueness and continuity with respect to the initial datum of the solutions of the Dirichlet problem. Existence is proved by showing a stochastic representation based on the delayed Brownian motion killed on the boundary. Related properties on crossing probabilities of the delayed Brownian motion are obtained. The asymptotic behaviour of the mean square displacement of the process is determined, showing that the diffusive behaviour is anomalous.

## CONTENTS

1. Introduction	1
1.1. The non-local heat equation with moving boundary	3
2. Weak maximum principle	6
2.1. Applications of the maximum principle	11
3. Stochastic representation for the non-local problem	12
3.1. Preliminary results	12
3.2. The non-local problem	18
4. Anomalous diffusive behaviour	33
5. Examples	36
Appendix A. Proofs of some technical results	39
A.1. Proof of Proposition 3.4	39
A.2. Proof of Proposition 3.6	43
A.3. Proof of Proposition 3.8	45
A.4. Proof of Proposition 3.14	46
Acknowledgements	46
References	46

## 1. INTRODUCTION

Given a one-dimensional Brownian motion  $B(t)$  and a function  $t \in \mathbb{R}_0^+ := [0, +\infty) \mapsto \varphi(t) \in \mathbb{R}$ , representing a boundary, one can define the corresponding

---

*Date:* March 21, 2022.

*2020 Mathematics Subject Classification.* 35R37, 60K50.

*Key words and phrases.* Semi-Markov process; anomalous diffusion; non-local heat equation; subordinator; inverse subordinator.

killed process as

$$B^\dagger(t) = \begin{cases} B(t), & t < \mathcal{T}, \\ \infty, & t \geq \mathcal{T}, \end{cases} \quad (1.1)$$

where  $\mathcal{T} = \inf \{t \geq 0 : B(t) \geq \varphi(t)\}$ , i.e., when the Brownian motion hits the boundary it is absorbed in a cemetery point  $\infty \notin \mathbb{R}$ . The interplay between the process  $B^\dagger$  and the problem of the form

$$\begin{cases} \partial_t u(t, x) = \frac{1}{2} \partial_x^2 u(t, x), & x < \varphi(t), t > 0, \\ u(t, \varphi(t)) = 0, & t > 0, \\ u(0, x) = f(x), \end{cases} \quad (1.2)$$

where  $f(x)$  is a suitable initial datum, is nowadays an understood topic (see, e.g., [34] and references therein): the one-dimensional density of  $B^\dagger$  upon killing is the fundamental solution to (1.2).

In this paper we consider a non-local (in time) counterpart of (1.2). We study the heat equation

$$\partial_t \int_0^t (u(x, s) - u(x, 0)) k(t-s) ds = \frac{1}{2} \partial_x^2 u(x, t), \quad (x, t) \in \mathcal{O}, \quad (1.3)$$

under suitable assumptions on  $k(t)$ , for a time increasing parabolic domain  $\mathcal{O}$ , with *exterior* Dirichlet condition. The fact that Dirichlet condition should be given outside  $\mathcal{O}$  and not only on the boundary is an intrinsic peculiarity of the non-local problems, i.e., the integral on the left-hand side of (1.3) involves  $(x, t) \notin \mathcal{O}$  and far from the boundary. The assumptions on  $k$  will be given so that the corresponding equation on the whole domain  $\mathbb{R} \times \mathbb{R}^+$  (the non-local counterpart of the classical heat equation) is solved by the one-dimensional density of the so-called delayed Brownian motion, i.e., a Brownian motion time-changed with the inverse of a (independent) strictly increasing Lévy process. As we will discuss later, it will be sufficient to assume that the kernel  $k : (0, \infty) \mapsto [0, \infty)$ , is non-increasing, vanishing at infinity and singular at zero but integrable on  $(0, 1]$ . The prototype is the choice

$$k(t) = t^{-\beta} / \Gamma(1 - \beta), \quad \beta \in (0, 1), \quad (1.4)$$

i.e., the case when the operator on the left-hand side of (1.3) is a  $\beta$ -fractional-time derivative. The corresponding process is called *delayed* Brownian motion (see, e.g., [36]) since it has continuous paths with interval of constancies (induced by the time-change) acting as a trapping effect for the original Brownian motion.

In addition to the purely theoretical interests arising from the stochastic and analytical point of view, there are strong motivations behind the study of this kind of problems coming from mathematical modeling of anomalous diffusion. With this in mind, non-local (fractional-type) equations on (fixed) domains, have been intensively studied in the last decades (for example, the reader can consult [44] and references therein for information on anomalous diffusion and how they interact with fractional equations, [40] for a stochastic approach to fractional diffusion on bounded domains, [56] for trapping models, [13] for the asymptotic behaviour). Non-local diffusion equations with moving boundaries have been considered, for example, in [54; 55; 62; 63; 64], mainly with analytical methods and motivated by mathematical modeling of anomalous diffusion. Precisely, fractional diffusion problems with moving boundaries arise, for example, when considering memory

effects in the latent heat accumulation on the melting interface of a solid body, together with a suitable condition on it (see [63; 64] and further generalizations in [19]). It is clear from these works that the moving boundaries make the classical methodologies of integro-differential (fractional-type) equations ineffective. The moving boundary problem studied in [55] is the most similar to our situation: the author considered a time-fractional problem (which means, in our framework, to assume (1.4)) with two moving boundaries, he provided a notion of solution and, for some explicit boundaries, he proved an existence result by resorting to special functions and superposition principles.

Our main results establish well-posedness for the general problem associated with (1.3): we provide a suitable notion of solution (and fundamental solution) for (1.3), we establish a weak maximum principle from which uniqueness (for elliptic operators more general than  $\partial_x^2$ ) and continuity with respect to the initial datum follow. We prove existence of the solution by providing a stochastic representation based on the killed delayed Brownian motion, i.e., a Brownian motion time-changed via the inverse of an independent strictly increasing Lévy process and then killed on the curve representing the moving boundary. With this at hand, other useful results on the curve-crossing probabilities of the delayed Brownian motion, such as a reflection principle and the asymptotic behaviour of the mean square displacement of the process, are obtained.

Here is a rigorous description of the problem we are going to address.

**1.1. The non-local heat equation with moving boundary.** A function  $\varpi : \mathbb{R}^+ \mapsto \mathbb{R}$ , where  $\mathbb{R}^+ := (0, \infty)$ , is said to be a Bernstein function if it is of class  $C^\infty$ , it is non-negative and  $(-1)^n \partial_\lambda^n \varpi(\lambda) \geq 0$ , for all  $n \in \mathbb{N}$  and  $\lambda > 0$ . It can be proved [58, Definition 3.1] that  $\varpi$  is a Bernstein function if, and only if, it has the representation

$$\varpi(\lambda) = a + b\lambda + \int_0^{+\infty} (1 - e^{-\lambda s}) \nu(ds) \quad (1.5)$$

for  $a, b \geq 0$  and where  $\nu$  is a positive measure on  $\mathbb{R}^+$  such that

$$\int_0^{+\infty} (s \wedge 1) \nu(ds) < \infty. \quad (1.6)$$

See more on Bernstein functions on [25; 58]. The measure  $\nu$  is said to be a Lévy measure. In what follows we will always assume that  $a = 0$ . Furthermore, we will consider only the case  $b = 0$ , for the sake of simplicity. We emphasize that the results proved in the paper could be generalized to include the case  $b > 0$ , but this would yield to some cumbersome technical computations.

From now on we will use the symbol  $\Phi(\lambda)$  for a Bernstein function having the representation

$$\Phi(\lambda) := \int_0^{+\infty} (1 - e^{-\lambda s}) \nu(ds). \quad (1.7)$$

Let us also denote by  $\bar{\nu}$  the tail of the Lévy measure, i.e., for any  $s > 0$ ,

$$\bar{\nu}(s) := \nu(s, +\infty) \quad (1.8)$$

and define, for a suitable class of functions  $f : \mathbb{R}_0^+ \rightarrow \mathbb{R}$ ,

$$\partial_t^\Phi f(t) := \partial_t \int_0^t \bar{\nu}(t-s) (f(s) - f(0)) ds. \quad (1.9)$$

Note that  $\bar{\nu}$  is non-increasing and using (1.6) it is easy to see that

$$\int_0^t \bar{\nu}(s) ds < \infty, \text{ for all } t > 0, \quad (1.10)$$

i.e.,  $\bar{\nu} \in L^1_{\text{loc}}(\mathbb{R}^+)$ . By the discussion above it is clear that the class of functions being the tail of a Lévy measure (in the sense of (1.8)) is quite large. If  $k : (0, \infty) \mapsto [0, \infty)$  is a non-increasing, vanishing at infinity, right-continuous function integrable on  $(0, 1]$  then its distributional derivative  $k(dt)$  (that is a negative Radon measure) satisfies (by [24, Theorem 3.3.1])

$$-\int_0^\infty (t \wedge 1) k(dt) < +\infty$$

hence it can be viewed as the tail of the Lévy measure  $-k(dt)$ . We have the following useful representation of  $\partial_t^\Phi$  acting on absolutely continuous functions.

**Lemma 1.1.** *Fix  $T > 0$ . For  $f \in AC([0, T])$  it is true that  $\partial_t^\Phi f$  exists a.e. in  $[0, T]$ ,*

$$\partial_t^\Phi f(t) = \int_0^t \bar{\nu}(t-s) \partial_t f(s) ds, \quad (1.11)$$

for almost any  $t \in [0, T]$  and  $t \in [0, T] \mapsto \partial_t^\Phi f(t)$  is in  $L^1([0, T])$ . Moreover, if  $f$  is Lipschitz, then equation (1.11) holds for any  $t \in [0, T]$  and  $\partial_t^\Phi f(t)$  is continuous on  $[0, T]$ .

*Proof.* Let us first observe that  $\partial_t f(t)$  belongs to  $L^1(0, T)$ . Thus, the function  $g(t) = \int_0^t \bar{\nu}(t-s) \partial_t f(s) ds$  belongs to  $L^1(0, T)$  (being the convolution product of two  $L^1$  functions) and  $G(t) = \int_0^t g(s) ds$  is absolutely continuous. Next, let us observe that

$$\begin{aligned} \int_0^t \int_0^s \bar{\nu}(\tau) |\partial_t f(s-\tau)| d\tau ds &= \int_0^t \bar{\nu}(\tau) \int_\tau^t |\partial_t f(s-\tau)| ds d\tau \\ &= \int_0^t \bar{\nu}(\tau) \int_0^{t-\tau} |\partial_t f(z)| dz d\tau \\ &\leq \left( \int_0^t \bar{\nu}(\tau) d\tau \right) \left( \int_0^t |\partial_t f(z)| dz \right) < +\infty. \end{aligned} \quad (1.12)$$

Let us evaluate explicitly  $G(t)$ . We have

$$\begin{aligned} G(t) &= \int_0^t \int_0^s \bar{\nu}(s-\tau) \partial_t f(\tau) d\tau ds \\ &= \int_0^t \int_0^s \bar{\nu}(\tau) \partial_t f(s-\tau) d\tau ds \\ &= \int_0^t \bar{\nu}(\tau) \int_\tau^t \partial_t f(s-\tau) ds d\tau, \end{aligned}$$

where we could use Fubini's theorem by (1.12). Now let us consider the change of variables  $s - \tau = z$  and let us use [65, Theorem 7.29] to achieve

$$\begin{aligned} G(t) &= \int_0^t \bar{\nu}(\tau) \int_0^{t-\tau} \partial_t f(z) dz d\tau \\ &= \int_0^t \bar{\nu}(\tau) (f(t-\tau) - f(0)) d\tau. \end{aligned}$$

Hence, in particular, the right-hand side is also absolutely continuous and we can differentiate both sides to conclude the first part of the proof. Concerning the second part of the proof, it is a direct consequence of [1, Proposition 1.3.2].  $\square$

Let  $\varphi : \mathbb{R}_0^+ \rightarrow \mathbb{R}$  be a non-decreasing function and define  $\mathbb{R}^{+,N} = \mathbb{R}_0^+ \times \mathbb{R}^N$ . Let

$$\mathcal{O} = \{(t, x) \in \mathbb{R}^{+,1} : t > 0, x < \varphi(t)\}, \quad \partial_0 \mathcal{O} := \{(0, x) : x < \varphi(0)\}. \quad (1.13)$$

We are interested in the  $\Phi$ -fractional heat equation

$$\partial_t^\Phi u(t, x) = \frac{1}{2} \partial_x^2 u(t, x), \quad (t, x) \in \mathcal{O}, \quad (1.14)$$

subject to

$$\begin{cases} u(t, x) = 0, & (t, x) \in \mathcal{O}^c \setminus \partial_0 \mathcal{O}, \\ u(t, x) = f(x), & (t, x) \in \partial_0 \mathcal{O}, \\ \lim_{x \rightarrow -\infty} u(t, x) = 0, & \text{locally uniformly for } t > 0, \end{cases} \quad (1.15)$$

where  $f$  is a suitable initial datum with compact support contained in  $(-\infty, \varphi(0))$ . Here, with *locally uniformly* we mean *uniformly in any compact subset*. Upon giving a suitable notion of classical solution to the problem, we show a weak maximum principle (from which uniqueness and continuity with respect to the initial datum follow) and a stochastic representation of the solution. It turns out that this stochastic representation is based on the delayed Brownian motion killed on the boundary  $\varphi(t)$ . Here is the rigorous construction of this process.

Let  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, B, \sigma, \mathbb{P})$  be a Markov additive process in the sense of [16, Definition 1.2.]. Suppose further that  $\sigma = \{\sigma(t), t \geq 0\}$ , is a subordinator, i.e., a non-decreasing Lévy process started at zero and having Lévy symbol

$$\mathbb{E} e^{-\lambda \sigma(t)} = e^{-t \Phi(\lambda)}, \quad (1.16)$$

and that  $B = \{B(t), t \geq 0\}$  is a Brownian motion on  $\mathbb{R}$ , independent of  $\sigma$ . Define, for any  $t \geq 0$ ,

$$L(t) := \inf \{s \geq 0 : \sigma(s) > t\}. \quad (1.17)$$

The process  $L = \{L(t), t \geq 0\}$  is (in general) right-continuous and non-decreasing. However, when considering the existence result with the stochastic representation, we will always assume that  $\nu(0, \infty) = +\infty$  so that  $\sigma$  is a strictly increasing subordinator: in this case the process  $L$  has continuous sample paths. Let us denote, for any probability distribution  $\mathbf{m}$ , by  $\mathbb{P}_{\mathbf{m}}$  the probability measure obtained by conditioning to the fact that  $B(0)$  admits distribution  $\mathbf{m}$ . In particular, if  $\mathbf{m} = \delta_y$  for some  $y \in \mathbb{R}$ , let us denote  $\mathbb{P}_{\delta_y} = \mathbb{P}_y$  and  $\mathbb{P} := \mathbb{P}_0$ . Consider then the process  $X_\Phi = \{X_\Phi(t), t \geq 0\}$ , where, for any  $t \geq 0$ ,

$$X_\Phi(t) := B(L(t)) \quad (1.18)$$

which is called *delayed* Brownian motion (e.g., [14; 36]). This process has continuous trajectories, as it is the composition of continuous processes. Further, it has interval of constancy induced by the time-change, since the jumps of  $\sigma$  produce intervals of constancy in the path of  $L$ . There is a wide recent literature concerning processes of this form, in particular on their governing equations [5; 6; 9; 15; 28; 32; 36; 40; 43; 47; 48; 49; 51; 60; 61] their interpretation as scale limit of CTRWs [7; 8; 31; 38; 37; 39; 42; 53; 59], distributional and path properties [4; 14; 50; 56] and applications in different fields [21; 23; 44; 57]. As we will see in details in the paper, the fact

that  $X_\Phi$  is semi-Markovian will be crucial for our purposes (we suggest [26; 27; 42] for a discussion on the semi-Markov property which can be applied to  $X_\Phi$ ).

Now let

$$X_\Phi^\dagger(t) := \begin{cases} X_\Phi(t), & t < \mathcal{T}, \\ \infty, & t \geq \mathcal{T}, \end{cases} \quad (1.19)$$

where

$$\mathcal{T} := \inf \{t \geq 0 : X_\Phi(t) \geq \varphi(t)\}, \quad (1.20)$$

and the state  $\infty \notin \mathbb{R}$  is a *cemetery* point and is absorbing. Then define, for any  $A \in \mathcal{B}$ , where  $\mathcal{B}$  is the Borel  $\sigma$ -algebra on  $\mathbb{R}$ , the sub-probability measure on  $\mathcal{B}$

$$\mathcal{B} \ni A \mapsto \mathbb{P}_y \left( X_\Phi^\dagger(t) \in A \right). \quad (1.21)$$

We show that, for any  $y \in (-\infty, \varphi(0))$ ,

$$\mathbb{P}_y \left( X_\Phi^\dagger(t) \in A \right) = \int_A q(t, x; y) dx \quad (1.22)$$

for a suitable density  $q(t, x; y)$  and that

$$\int_{(-\infty, \varphi(0))} f(y) q(t, x; y) dy \quad (1.23)$$

is the unique classical solution (in the appropriate sense, see Definition 2.1 below) to (1.14) subject to (1.15) for  $f \in C_c((-\infty, \varphi(0)))$ , i.e., the space of functions which are continuous and compactly supported on  $(-\infty, \varphi(0))$ . We will prove this result under appropriate technical conditions (see **A1**, **A2(a) – A2(d)**, **A3** and **A4** below).

The paper is organized as follows. We deal with the uniqueness problem on more general (than  $\mathcal{O}$ ) increasing parabolic sets and with a more general (than  $\partial_x^2$ ) advection-diffusion operator and therefore we discuss separately, in the general framework, the concept of solution, its uniqueness and the continuity with respect to the initial datum by proving a maximum principle (see Section 2). Existence is proved with stochastic methods in Section 3. The anomalous diffusive behaviour is studied in Section 4.

## 2. WEAK MAXIMUM PRINCIPLE

First of all, let us introduce some notation. Let  $\mathbb{R}_T^{+,N} := [0, T] \times \mathbb{R}^N$  and consider  $\mathfrak{D}$  to be a subset of  $\mathbb{R}_T^{+,N}$ , possibly  $T = +\infty$  with  $\mathbb{R}_{+\infty}^{+,N} := \mathbb{R}_0^+ \times \mathbb{R}^N =: \mathbb{R}^{+,N}$ . With  $\overline{\mathfrak{D}}$  we denote the closure of  $\mathfrak{D}$ . Let us define the parabolic interior of  $\mathfrak{D}$  as the set  $\mathfrak{D}^*$  such that

$$(t_0, x_0) \in \mathfrak{D}^* \Leftrightarrow \exists r > 0 : B_r(t_0, x_0) \cap \{(t, x) \in \mathbb{R}_T^{+,N} : t \leq t_0\} \subset \mathfrak{D} \quad (2.1)$$

where with  $B_r(t_0, x_0)$  we denote the ball of radius  $r > 0$  and center  $(t_0, x_0)$  in  $\mathbb{R}^{N+1}$ . Generally one has  $\mathfrak{D}^* \subset \mathfrak{D} \subset \overline{\mathfrak{D}}$ . We say that a set  $\mathfrak{D}$  is a parabolic open set if  $\mathfrak{D} = \mathfrak{D}^*$ . From now on, let us suppose that  $\mathfrak{D}$  is a parabolic open set. Let us also stress that  $\mathcal{O}$  defined in (1.13) is a parabolic open subset of  $\mathbb{R}^{+,1}$ . Therefore the notation related to  $\mathfrak{D}$  that we introduce below is extended directly to  $\mathcal{O}$ .

We define the parabolic boundary of  $\mathfrak{D}$  as  $\partial_p \mathfrak{D} = \overline{\mathfrak{D}} \setminus \mathfrak{D}^*$ . Next let us define the following set-valued functions:

$$\mathfrak{D}_1 : x \in \mathbb{R}^N \mapsto \mathfrak{D}_1(x) := \{t \in [0, T] : (t, x) \in \mathfrak{D}\}$$

$$\mathfrak{D}_2 : t \in [0, T] \mapsto \mathfrak{D}_2(t) := \{x \in \mathbb{R}^N : (t, x) \in \mathfrak{D}\}.$$

For any  $A \subset \mathbb{R}^N$  we denote  $\mathfrak{D}_1(A) = \bigcup_{x \in A} \mathfrak{D}_1(x)$  and for any  $A \subset [0, T]$  we denote  $\mathfrak{D}_2(A) = \bigcup_{t \in A} \mathfrak{D}_2(t)$ . In particular, let us set  $\mathfrak{D}_1 := \mathfrak{D}_1(\mathbb{R}^N)$  and  $\mathfrak{D}_2 := \mathfrak{D}_2([0, T])$ . All this notation is extended to  $\overline{\mathfrak{D}}$ . Finally, let us assume that  $\mathfrak{D}$  is non-decreasing in  $t$ , i.e., that the function  $[0, T] \ni t \mapsto \mathfrak{D}_2(t)$  is non-decreasing with respect to the inclusion.

We investigate time-nonlocal advection-diffusion equations on the parabolic domain  $\mathfrak{D}$ . In particular, let us consider the advection-diffusion operator  $L$  on  $\mathfrak{D}_2$  given by

$$Lu(x) = p_2(x)\Delta u(x) + \langle p_1(x), \nabla u(x) \rangle \quad (2.2)$$

where  $p_2 : \mathfrak{D}_2 \rightarrow \mathbb{R}^+$  and  $p_1 : \mathfrak{D}_2 \rightarrow \mathbb{R}^N$ , for any function  $u : \mathfrak{D}_2 \rightarrow \mathbb{R}$  in  $C^2(\mathbb{R}^N \mathfrak{D})$ .

**Definition 2.1.** *We say that a function  $u(t, x) : [0, T] \times \mathfrak{D}_2 \mapsto \mathbb{R}$  (possibly  $\mathbb{R}_0^+$  in place of  $[0, T]$ ) is a classical solution of*

$$\partial_t^\Phi u(t, x) = Lu(t, x), \quad (t, x) \in \mathfrak{D}, \quad (2.3)$$

if

- (1)  $u \in C^0(\overline{\mathfrak{D}})$ ;
- (2)  $\forall t \in \mathfrak{D}_1$  with  $t > 0$  it holds  $u(t, \cdot) \in C^2(\mathfrak{D}_2(t))$ ;
- (3)  $\partial_t^\Phi u(t, x)$  exists for any  $(t, x) \in \mathfrak{D}$ ;
- (4) for any  $x \in \mathfrak{D}_2$ ,  $\partial_t^\Phi u(\cdot, x) \in C^0(\mathfrak{D}_1(x))$ ;
- (5) it satisfies (2.3).

**Remark 2.2.** *Our notion of solution is similar to the one given in [55, Definition 2.7] for the fractional case. Precisely we ask for regularity of the solution inside the domain in place that in the whole cylinder  $[0, T] \times \mathfrak{D}_2$ . Continuity of the non-local operator (in place of continuity of the time-derivative) is needed to extend, via a density argument, the extremal point property. Finally, since we prove general well-posedness, to guarantee both existence and uniqueness of the solution we consider exterior Dirichlet conditions in place of just boundary conditions.*

Our aim in this section is to show that the solutions of the non-local advection-diffusion equation (2.3) admits maximum on the parabolic boundary of  $\mathfrak{D}$ . We divide the case when  $\mathfrak{D}$  is bounded (and, without loss of generality, connected) to the case where it is unbounded. The latter is needed since we want to apply the maximum principle to the problem (1.14).

**Theorem 2.3.** *Let  $\mathfrak{D}$  be bounded and connected. Suppose that  $u$  is a classical solution (in the sense above) of*

$$\partial_t^\Phi u(t, x) = Lu(t, x), \quad (t, x) \in \mathfrak{D}, \quad (2.4)$$

such that  $u(\cdot, x)$  is constant whenever  $(t, x) \notin \mathfrak{D}$ . Then it is true that

$$u(t, x) \leq \max_{(\tau, y) \in \partial_p \mathfrak{D}} u(\tau, y), \quad \text{for all } (t, x) \in \overline{\mathfrak{D}}. \quad (2.5)$$

**Theorem 2.4.** *Let  $N = 1$  and  $\mathfrak{D} = \{(t, x) : x < \varphi(t), t \in [0, T]\}$  for  $t \mapsto \varphi(t)$  is non-decreasing. Let  $u(t, x)$  be a classical solution to (2.3) in the sense above. Further suppose that  $u(t, x)$  is such that  $u(\cdot, x)$  is constant when  $(t, x) \notin \mathfrak{D}$  and that*

there exists a function  $u_\infty(t)$  continuous in  $[0, T]$  so that  $u(t, x) \rightarrow u_\infty(t)$  uniformly as  $x \rightarrow -\infty$  for  $t \in [0, T]$ . Then

$$u(t, x) \leq \max \left\{ \max_{(y, \tau) \in \partial_p \mathfrak{D}} u(y, \tau), \max_{\tau \in [0, T]} u_\infty(\tau) \right\}, \text{ for all } (t, x) \in \overline{\mathfrak{D}}. \quad (2.6)$$

**Remark 2.5.** Note that we will show the stochastic representation (1.23) for  $u(t, x)$  subject to Dirichlet conditions on the time-dependent boundary  $x = \varphi(t)$ . It follows that if, for example, the function  $f(x)$  is negative then the maximum of  $u(t, x)$  is on the set  $x = \varphi(t)$  where  $u(t, x) = 0$ . If, instead,  $f(x)$  is positive, the maximum is on the set  $(0, x)$ ,  $x \leq \varphi(0)$ .

Moreover the same result applies to  $-u(t, x)$ . In particular this implies that if  $f(x)$  is of prescribed sign then  $u$  will preserve it.

To prove Theorem 2.3, We first need to show a sort of compatibility between maximum points of a function and its non-local derivative. This is done in the forthcoming two results.

**Lemma 2.6.** Let  $T > 0$  be arbitrary and let  $f : I \subset [0, T] \rightarrow \mathbb{R}$  where  $I$  is an interval containing 0. If  $t_0 \in (0, T]$  is a maximum point of  $f$  and  $f \in W^{1,1}(0, t_0) \cap C^1((0, t_0))$ , then  $\partial_t^\Phi f(t_0) \geq 0$ .

*Proof.* Let us consider the function  $g(\tau) = f(t_0) - f(\tau)$  for  $\tau \in [0, T]$ , so that, being  $t_0$  the maximum point of  $f$ , it holds  $g(\tau) \geq 0$  for any  $\tau \in [0, T]$  and  $\partial_t^\Phi g(\tau) = -\partial_t^\Phi f(\tau)$  for any  $\tau \in (0, T]$ . Let us observe that the assertion is trivial if  $g$  is constant, thus let us only consider the case in which  $g$  is not constant. Being  $f \in C^1((0, t_0))$  by assumption,  $g \in C^1([\varepsilon, t_0])$  for any  $\varepsilon \in (0, t_0)$ , so that  $g$  is Lipschitz on  $[\varepsilon, t_0]$  and thus

$$\forall \varepsilon \in (0, T) \exists C_\varepsilon > 0 : g(\tau) \leq C_\varepsilon |t_0 - \tau| \quad \forall \tau \in [\varepsilon, T]. \quad (2.7)$$

Note that by hypotheses  $g$  is absolutely continuous on  $[0, t_0]$ , thus it follows that we can use (1.11) to say that

$$\partial_t^\Phi g(t_0) = \int_0^{t_0} \bar{\nu}(t_0 - \tau) g'(\tau) d\tau, \quad (2.8)$$

where  $g'$  is the derivative of  $g$ . Let us split the integral in  $\varepsilon \in (0, t_0)$ , where  $\varepsilon$  will be chosen in what follows, to obtain

$$\partial_t^\Phi g(t_0) = \int_0^\varepsilon \bar{\nu}(t_0 - \tau) g'(\tau) d\tau + \int_\varepsilon^{t_0} \bar{\nu}(t_0 - \tau) g'(\tau) d\tau \quad (2.9)$$

$$= I_1(\varepsilon) + I_2(\varepsilon). \quad (2.10)$$

Now let us consider  $I_2(\varepsilon)$ . Being  $g$  Lipschitz in  $[\varepsilon, t_0]$  and  $\bar{\nu} \in L^1(0, t_0 - \varepsilon)$ , we can use dominated convergence theorem to obtain

$$I_2(\varepsilon) = \lim_{a \rightarrow 0^+} \int_\varepsilon^{t_0-a} \bar{\nu}(t_0 - \tau) g'(\tau) d\tau = \lim_{a \rightarrow 0^+} \int_a^{t_0-\varepsilon} \bar{\nu}(z) g'(t-z) dz. \quad (2.11)$$

Now let us recall that the function  $\bar{\nu}$  is non-increasing and finite in  $[a, t_0 - \varepsilon]$ , hence it is of bounded variation with distributional derivative given by  $-\nu$ . Thus we can apply integration by parts (see [24, Theorem 3.3.1]) to obtain

$$I_2(\varepsilon) = \lim_{a \rightarrow 0} \left[ \bar{\nu}(a) g(t_0 - a) - \bar{\nu}(t_0 - \varepsilon) g(\varepsilon) - \int_a^{t_0-\varepsilon} g(t_0 - z) \nu(dz) \right]. \quad (2.12)$$

Combining inequality (2.7) with the fact that  $t\bar{\nu}(t) \rightarrow 0$  as  $t \rightarrow 0$  (see, e.g., [58, page 17]) and also using monotone convergence theorem, we get

$$I_2(\varepsilon) = -\bar{\nu}(t_0 - \varepsilon)g(\varepsilon) - \int_0^{t_0 - \varepsilon} g(t_0 - \tau)\nu(d\tau). \quad (2.13)$$

Since  $g$  is not constant on  $(0, t_0)$ , we know there exists  $\varepsilon_0 \in (0, t_0)$  such that  $g(\varepsilon_0) > 0$  and we can suppose  $\varepsilon < \varepsilon_0$ . We have, by continuity of  $g$  and the fact that  $-\bar{\nu}(t_0 - \varepsilon)g(\varepsilon) \leq 0$ ,

$$I_2(\varepsilon) \leq - \int_0^{t_0 - \varepsilon_0} g(t_0 - \tau)\nu(d\tau) =: -C < 0, \quad (2.14)$$

where  $C$  is finite since, by (2.7),

$$\int_0^{t_0 - \varepsilon_0} g(t_0 - \tau)\nu(d\tau) \leq C_{\varepsilon_0} \int_0^{t_0 - \varepsilon_0} \tau \nu(d\tau) < +\infty.$$

Now let us consider  $I_1(\varepsilon)$ . By absolute continuity of  $g$  and monotonicity of  $\bar{\nu}$  we clearly have

$$I_1(\varepsilon) \leq \bar{\nu}(t_0 - \varepsilon) \int_0^\varepsilon \frac{dg}{dt}(\tau) d\tau \leq \bar{\nu}(t_0 - \varepsilon_0) \int_0^\varepsilon \left| \frac{dg}{dt}(\tau) \right| d\tau. \quad (2.15)$$

By absolute continuity of the Lebesgue integral we have that there exists a  $\varepsilon \in (0, \varepsilon_0)$  such that  $I_1(\varepsilon) \leq C/2$ . Let us fix this  $\varepsilon$  to achieve

$$\partial_t^\Phi g(t_0) = I_1(\varepsilon) + I_2(\varepsilon) \leq -\frac{C}{2} < 0 \quad (2.16)$$

concluding the proof.  $\square$

It turns out that the assumption  $f \in C^1((0, t_0])$  in the previous result is unnecessary and it can be extended and can be replaced by existence and continuity of  $\partial_t^\Phi f(t)$ .

**Lemma 2.7.** *Let  $T > 0$  be arbitrary. Let  $f$  be a continuous function in  $[0, T]$  and set  $t_0 = \min \operatorname{argmax}(f)$ . Suppose  $t_0 > 0$  and  $\partial_t^\Phi f(t)$  is well-defined and continuous in an open interval  $I \subset (0, T)$  containing  $t_0$ . Then  $\partial_t^\Phi f(t_0) \geq 0$ .*

*Proof.* Denote  $f_0(t) := f(t) - f(0)$ , let

$$g_f(s) := \int_0^s \bar{\nu}(s - v)f_0(v)dv \quad (2.17)$$

and observe that since  $\partial_t^\Phi f(t)$  is continuous on  $[\varepsilon, T]$  then  $g_f \in C^0([0, T]) \cap C^1(I)$ . Consider now a sequence  $\varphi_n$ ,  $n \in \mathbb{N}$ , of Friedrich mollifiers. Since the support of  $\varphi_n$  is compact, it is easy to see that for any open interval  $I' \subset \bar{I} \subset I$  containing  $t_0$  it holds

$$\varphi_n * \frac{dg_f}{dt} = \frac{d}{dt}(\varphi_n * g_f), \quad (2.18)$$

on  $\bar{I}$  for  $n$  sufficiently large. However, we can rewrite

$$g_f = (f_0 \mathbf{1}_{[0, +\infty)}) * (\bar{\nu} \mathbf{1}_{[0, +\infty)}) \quad (2.19)$$

thus by associativity and commutativity of the convolution operation we get

$$\varphi_n * \frac{dg_f}{dt} = \frac{d}{dt}(\bar{\nu} \mathbf{1}_{[0, +\infty)} * (\varphi_n * f_0 \mathbf{1}_{[0, +\infty)})) = \partial_t^\Phi (\varphi_n * f_0 \mathbf{1}_{[0, +\infty)}). \quad (2.20)$$

Note now that if  $t_0$  is a maximum point for  $f$ , so it is for  $f_0$  and vice versa. Let  $f_n := \varphi_n * f_0 \mathbf{1}_{[0,+\infty)}$  and observe that the sequence uniformly converges towards  $f_0$  in  $[0, T]$  (by [1, Lemma 1.3.3] since  $f_0$  is continuous). In particular it uniformly converges to  $f_0$  in  $[0, t_0]$  and, being  $t_0$  the unique maximum point in  $[0, t_0]$  of  $f_0$ , it follows that the maximum points  $t_n^0$  of  $f_n$  in  $[0, t_0]$  converge towards  $t_0$ . Moreover  $f_n$  is  $C^1((0, T])$  and thus it holds, by Lemma 2.6,

$$\partial_t^\Phi f_n(t_n^0) \geq 0. \quad (2.21)$$

However, by [1, Lemma 1.3.3] and (2.20),  $\partial_t^\Phi f_n = \varphi_n * \partial_t^\Phi f$  uniformly converges towards  $\partial_t^\Phi f$  in  $\bar{I}$ . Hence, taking the limit in (2.21) we conclude the proof.  $\square$

Now we proceed with the proof of Theorem 2.3, following a line of reasoning similar to [35, Theorem 2]. We adapt the main arguments of the proof to our case by extending on the sections  $\mathfrak{D}_1(x)$  the function  $u$  by continuity with a constant function: this stems on the idea that the non-local derivative does not take into account intervals in which the function is constant, thus we are not changing the nature of such derivative. Moreover, in the proof we will need to take into account all the difficulties related to the parabolic topology of  $\mathfrak{D}$ .

*Proof of Theorem 2.3.* Let us set  $M_1 = \max_{(\tau, y) \in \partial_p \mathfrak{D}} u(\tau, y)$  and let us argue by contradiction. Suppose that there exists a point  $(t_0, x_0) \in \mathfrak{D}$  such that  $u(t_0, x_0) = M_2 > M_1$ . Let us define  $\varepsilon = M_2 - M_1 > 0$  and consider the auxiliary function

$$w(t, x) = u(t, x) + \frac{\varepsilon}{2} \frac{T-t}{T} \quad \forall (t, x) \in \bar{\mathfrak{D}} \quad (2.22)$$

For any  $x \in \mathfrak{D}_2$  define  $\mathfrak{T}(x) = \min \bar{\mathfrak{D}}(x)$  and observe that for  $(\mathfrak{T}(x), x) \in \partial_p \mathfrak{D}$  it holds

$$w(\mathfrak{T}(x), x) = u(\mathfrak{T}(x), x) + \frac{\varepsilon}{2} \frac{T - \mathfrak{T}(x)}{T}. \quad (2.23)$$

Thus, if we set  $w(t, x) = w(\mathfrak{T}(x), x)$  and  $u(t, x) = u(\mathfrak{T}(x), x)$  for any  $x \in \mathfrak{D}_2$  and  $t \in [0, \mathfrak{T}(x)]$  whenever  $\mathfrak{T}(x) \neq 0$  the functions  $w(\cdot, x)$  and  $u(\cdot, x)$  are continuous on  $[0, T]$  for any fixed  $x \in \mathfrak{D}_2$ . Since  $\frac{T-t}{T} < 1$  we have that

$$w(t, x) \leq u(t, x) + \frac{\varepsilon}{2} \quad \forall (t, x) \in [0, T] \times \mathfrak{D}_2. \quad (2.24)$$

In the point  $(t_0, x_0)$  we obtain

$$w(t_0, x_0) = u(t_0, x_0) + \frac{\varepsilon}{2} \frac{T-t_0}{T} \geq u(t_0, x_0) = \varepsilon + M_1 \quad (2.25)$$

Now let us consider  $(t, x) \in \partial_p \mathfrak{D}$ . We have  $M_1 \geq u(t, x)$  and then, by also using equation (2.24),

$$w(t_0, x_0) \geq \varepsilon + M_1 \geq \varepsilon + u(t, x) \geq \frac{\varepsilon}{2} + w(t, x), \quad \forall (t, x) \in \partial_p \mathfrak{D}. \quad (2.26)$$

Now, since  $w \in C^0(\bar{\mathfrak{D}})$ , we know there exists a point  $(t_1, x_1) \in \bar{\mathfrak{D}}$  such that  $w(t_1, x_1) = \max_{(t, x) \in \bar{\mathfrak{D}}} w(t, x)$ . Without loss of generality, we can suppose  $t_1 = \min\{t \in [0, T] : w(t, x_1) = \max_{(t, x) \in \bar{\mathfrak{D}}} w(t, x)\}$ . However, since  $w(t_0, x_0) > w(t, x)$  for any  $(t, x) \in \partial_p \mathfrak{D}$ , one should have  $(t_1, x_1) \in \mathfrak{D}^*$ . Moreover, if  $\mathfrak{T}(x_1) \neq 0$ , we have

$$w(t_1, x_1) > w(\mathfrak{T}(x_1), x_1) = w(t, x_1), \quad \forall t \in (0, \mathfrak{T}(x_1)), \quad (2.27)$$

hence  $t_1$  is the first maximum point of  $w(\cdot, x_1)$ . Finally observe that  $w(t_1, x_1) > w(t_0, x_0) > \varepsilon > 0$ .

Since by construction we know that  $w(\cdot, x_1)$  is continuous in  $[0, T]$  and admits a non-local derivative which is continuous on  $(\mathfrak{I}(x_1), T]$ , we have that

$$\begin{cases} \nabla w(t_1, x_1) = 0, \\ \Delta w(t_1, x_1) \leq 0, \\ \partial_t^\Phi w(t_1, x_1) \geq 0, \end{cases} \quad (2.28)$$

where we used Lemma 2.7. Moreover by linearity of  $\partial_t^\Phi$  we get that

$$\partial_t^\Phi u(t, x) = \partial_t^\Phi w(t, x) + \frac{\varepsilon}{2T} I(t), \quad \forall (t, x) \in \mathfrak{D}, \quad (2.29)$$

where

$$I(t) = \int_0^t \bar{v}(\tau) d\tau > 0. \quad (2.30)$$

Thus we have

$$\partial_t^\Phi u(t_1, x_1) - Lu(t_1, x_1) = \partial_t^\Phi w(t_1, x_1) + \frac{\varepsilon}{2T} I(t_1) - p_2(x_1) \Delta w(t_1, x_1) > 0 \quad (2.31)$$

which is absurd since  $u$  is solution of Equation (2.4).  $\square$

*Proof of Theorem 2.4.* The proof can be conducted in the very same way as the previous one by using Weierstrass' Theorem on the set  $\bar{\mathfrak{D}} \cup (\{-\infty\} \times [0, T])$ .  $\square$

**2.1. Applications of the maximum principle.** As a direct consequence of the weak maximum principle (Theorems 2.3 and 2.4) we obtain uniqueness of the solution of the problem (1.14). The notation used so far for the set  $\mathfrak{D}$  are extended to  $\mathcal{O}$  in the obvious way.

**Corollary 2.8.** *Let  $\varphi : \mathbb{R}_0^+ \rightarrow \mathbb{R}$  be a non-decreasing function. Let  $u_1, u_2$  be two classical solutions of (1.14) subject to (1.15). Then  $u_1 \equiv u_2$ .*

*Proof.* Fix  $T > 0$  and consider  $\mathcal{O}^T = \mathcal{O} \cap ([0, T] \times \mathbb{R})$ . Let us observe that the function  $w = u_1 - u_2$  solves

$$\partial_t^\Phi w(t, x) = \frac{1}{2} \partial_x^2 w(t, x), \quad (t, x) \in \mathcal{O}, \quad (2.32)$$

$w(t, x) = 0$  for any  $(t, x) \in ([0, T] \times \mathbb{R}) \setminus \mathcal{O}$  and  $\lim_{x \rightarrow -\infty} w(t, x) = 0$  uniformly for  $t \in [0, T]$ . Thus, by Theorem 2.4, we have that for any  $(t, x) \in \bar{\mathcal{O}}^T$  it holds

$$w(t, x) \leq \max \left\{ \max_{(\tau, y) \in \partial_p \mathcal{O}^T} w(\tau, y), \max_{\tau \in [0, T]} u_\infty(\tau) \right\} = 0. \quad (2.33)$$

Arguing in the same way with  $-w(t, x)$ , we have that  $w \equiv 0$  in  $\bar{\mathcal{O}}^T$  and then  $u_1 \equiv u_2$  in  $\bar{\mathcal{O}}^T$ . Since  $T > 0$  is arbitrary we conclude the proof.  $\square$

Another consequence of the weak maximum principle is the continuity with respect to the initial data.

**Corollary 2.9.** *Let  $\varphi : \mathbb{R}_0^+ \rightarrow \mathbb{R}$  be a non-decreasing function. Let  $u_1, u_2$  be two classical solutions of (1.14) subject to (1.15) with  $u_i(0, x) = f_i(x)$ ,  $i = 1, 2$ , for  $f_i(x)$  continuous and bounded on  $(-\infty, \varphi(0))$  with  $\lim_{x \rightarrow -\infty} f_i(x) = \lim_{x \rightarrow \varphi(0)} f_i(x) = 0$ . Then*

$$\sup_{(t, x) \in \mathcal{O}} |u_1(t, x) - u_2(t, x)| \leq \sup_{x \leq \varphi(0)} |f_1(x) - f_2(x)|. \quad (2.34)$$

*Proof.* Arguing as in Corollary 2.8 on  $u_1 - u_2$ , by Theorem 2.4 it follows that

$$\begin{aligned}
& \sup_{(t,x) \in \mathcal{O}^T} |u_1(t,x) - u_2(t,x)| \\
&= \max \left\{ \sup_{(t,x) \in \mathcal{O}^T} u_1(t,x) - u_2(t,x), \sup_{(t,x) \in \mathcal{O}^T} u_2(t,x) - u_1(t,x) \right\} \\
&= \max \left\{ \max \left\{ 0, \sup_{x \leq \varphi(0)} f_1(x) - f_2(x) \right\}, \max \left\{ 0, \sup_{x \leq \varphi(0)} f_2(x) - f_1(x) \right\} \right\} \\
&= \max \left\{ \sup_{x \leq \varphi(0)} f_1(x) - f_2(x), \sup_{x \leq \varphi(0)} f_2(x) - f_1(x) \right\} \\
&= \sup_{x \leq \varphi(0)} |f_1(x) - f_2(x)|. \tag{2.35}
\end{aligned}$$

Taking the supremum on  $T$  we conclude the proof.  $\square$

**Remark 2.10.** *It is clear that Corollaries 2.8 and 2.9 can be proved also in the case when  $\mathcal{O}$  is bounded as in Theorem 2.3, with Dirichlet exterior conditions.*

Note that we have proved uniqueness and continuity with respect to the initial data of the classical solution. We conclude the well-posedness argument by proving existence of the classical solution for the problem (1.14) by means of a stochastic representation.

### 3. STOCHASTIC REPRESENTATION FOR THE NON-LOCAL PROBLEM

In this section we show the stochastic representation (1.23) of the (unique) classical solution to (1.14). Therefore we first need to provide a representation for the distribution

$$\mathcal{B} \ni A \mapsto \mathbb{P}_y \left( X_{\Phi}^{\dagger}(t) \in A \right). \tag{3.1}$$

**3.1. Preliminary results.** Under suitable assumptions on the Lévy measure  $\nu$  and the moving boundary  $\varphi$ , it turns out that (3.1) is absolutely continuous with respect to the Lebesgue measure and can be expressed using the one-dimensional density of the free (unkilled) process  $X_{\Phi}$ . To show this, we have to exploit the semi-Markov property of the aforementioned process.

Using the independence of  $B$  and  $\sigma$ , it is easy to see that, under  $\mathbb{P}_y$ , for any  $t > 0$ , the law of  $X_{\Phi}(t)$  is absolutely continuous with a density given by

$$p_{\Phi}(t, x; y) = \int_0^{+\infty} p(s, x; y) f_L(s; t) ds \tag{3.2}$$

where  $f_L(\cdot; t)$  is the density of  $L(t)$  with  $f_L(s; 0) = 0$  for  $s > 0$ , and  $p(s, x; y)$  is the transition density of  $B$ , i.e.,

$$p(s, x; y) = \frac{1}{\sqrt{2\pi s}} e^{-\frac{(x-y)^2}{2s}}, \quad (s, x, y) \in (0, \infty) \times \mathbb{R}^2. \tag{3.3}$$

The existence of the density of  $L(t)$  is guaranteed by [38, Theorem 3.1] under the condition

- The Lévy measure satisfies  $\nu(0, \infty) = +\infty$ . (A1)

The fact that  $X_\Phi$  is semi-Markovian means that it can be embedded in a Markov process in the following sense: if we let

$$G(t) := \sigma(L(t)-), \quad H(t) := \sigma(L(t)), \quad (3.4)$$

then the processes  $\{(X_\Phi(t), t - G(t)), t \geq 0\}$  and  $\{(X_\Phi(t), H(t) - t), t \geq 0\}$  are time homogeneous strong Markov processes with respect to filtrations  $(\mathcal{F}_{L(t)-})_{t \geq 0}$  and  $(\mathcal{F}_{L(t)})_{t \geq 0}$  respectively (see [42, Section 4]).

In particular, the evolution of  $X_\Phi$  on the time interval  $[H(t), \infty)$  depends only on the position of  $X_\Phi$  at time  $H(t)$ , meaning that  $H(t)$  is a random time with respect to  $X_\Phi$  exhibits the Markov property (see the discussion at [42, page 1704]). This can be equivalently seen by observing that, since  $(B, \sigma)$  is Markov additive in the sense of [16, Definition 1.2],  $(\Omega, \mathcal{F}, (\mathcal{F}_{L(t)})_{t \geq 0}, \overline{\mathcal{M}}, X_\Phi, \mathbb{P}_x)$  is regenerative in the sense of [27, Definition 2.2 (and Example 2.13)], where the regenerative set  $\overline{\mathcal{M}}$  is given by the closure of

$$\mathcal{M}(\omega) = \{t \in \mathbb{R}_0^+ : t = \sigma(u, \omega) \text{ for some } u \geq 0\}, \quad \omega \in \Omega. \quad (3.5)$$

Furthermore,  $X_\Phi$  exhibits the strong regeneration property in the sense of [27, eq. (2.14)], so that the strong Markov property holds with respect to any  $(\mathcal{F}_{L(t)})_{t \geq 0}$ -Markov time. In the following we will need to ensure that  $X_\Phi$  overleaps for the first time the boundary  $\varphi$  in a point that does not belong to an interval of constancy of the process, in other words  $\mathcal{T} \in \overline{\mathcal{M}}$  almost surely. In such a case, since

$$H(t, \omega) = \inf \{s > t : s \in \overline{\mathcal{M}}(\omega)\}, \quad \omega \in \Omega, \quad (3.6)$$

it holds  $H(\mathcal{T}) = \mathcal{T}$  almost surely and then  $X_\Phi$  exhibits the strong Markov property at  $\mathcal{T}$ . A sufficient condition for such a property is given by asking that  $\varphi$  is non-decreasing and continuous. Indeed, while the discussion is trivial if  $X_\Phi(0) \geq \varphi(0)$  since  $\mathcal{T} = 0$ , if  $X_\Phi(0) < \varphi(0)$  it is clear that either  $\mathcal{T} = \infty$  or  $X_\Phi(\mathcal{T}) \geq \varphi(\mathcal{T})$ . We only need to study the latter case. For the sake of the reader, we omit the dependence on  $\omega \in \Omega$ . Being both  $X_\Phi$  and  $\varphi$  continuous, the set  $\{t \in [0, +\infty) : X_\Phi(t) - \varphi(t) = 0\}$  is closed and

$$0 < \mathcal{T} = \min\{t \in [0, +\infty) : X_\Phi(t) - \varphi(t) = 0\}.$$

Now suppose that  $\mathcal{T} \notin \overline{\mathcal{M}}$ . Then we know that  $\mathcal{T} \in (\sigma(w-), \sigma(w))$  for some  $w > 0$  and

$$X_\Phi(\sigma(w-)) = X_\Phi(\mathcal{T}) = \varphi(\mathcal{T}) \geq \varphi(\sigma(w-)).$$

Combining this information with the fact that  $X_\Phi(0) < \varphi(0)$ , we get that there exist  $t_0 \in (0, \sigma(w-)]$ , and then in particular  $t_0 < \mathcal{T}$ , such that  $X_\Phi(t_0) - \varphi(t_0) = 0$ , which is a contradiction.

In the previous argument we used the continuity of  $X_\Phi$  and  $\varphi$ , which also implies another simple consequence. Indeed, for any  $w_0 \geq 0$  and  $y < \varphi(0)$ ,

$$\mathbb{P}_y(\mathcal{T} \in \{w_0\}) \leq \mathbb{P}_y(X_\Phi(w_0) = \varphi(w_0)) = 0, \quad (3.7)$$

i.e.,  $\mathcal{T}$  is a continuous r.v. while if  $y \geq \varphi(0)$  it is clear that  $\mathcal{T} \equiv 0$  almost surely. Let us stress that all the previous arguments also hold if  $\varphi$  is just non-decreasing and right-continuous, but continuity will be used in further results. Therefore, we assume that  $\varphi$  is a non-decreasing continuous boundary. However, as we will see in the following, we need some additional regularity to handle more subtle properties of the fundamental solution of (1.14) under the conditions (1.15). Despite we did

not justify some of them up to now, let us write down all the assumptions we need on the moving boundary  $\varphi$ .

- $t \mapsto \varphi(t)$  is non-decreasing and continuous; (A2(a))
- if  $\varphi(t_0) = \varphi(t_1)$  for some  $t_0 < t_1$ , then  $\varphi(t) = \varphi(t_0)$  for any  $t \geq t_0$ ; (A2(b))
- $t \mapsto \varphi(t)$  is bounded by a constant  $l > 0$ ; (A2(c))
- $t \mapsto \varphi(t)$  is locally Lipschitz. (A2(d))

**Lemma 3.1.** *Under A1 and A2(a), it is true that, for  $t > 0$  and for any  $A \in \mathcal{B}$ ,*

$$\mathbb{P}_y \left( X_{\Phi}^{\dagger}(t) \in A \right) = \int_A q(t, x; y) dx \quad (3.8)$$

where, for  $(t, x, y) \in \mathbb{R}^{+,2}$ ,

$$q(t, x; y) = p_{\Phi}(t, x; y) - \int_0^t p_{\Phi}(t-w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw). \quad (3.9)$$

*Proof.* The statement is clear if  $y \geq \varphi(0)$  as in this case  $q(t, x; y) = 0$  for any  $t \geq 0$  and  $x \in \mathbb{R}$ . Thus let us focus on the case  $y < \varphi(0)$ . Let us first observe that for any Borel set  $A \in \mathcal{B}$  and any  $t \geq 0$  it holds

$$\begin{aligned} & \mathbb{P}_y \left( X_{\Phi}^{\dagger}(t) \in A \right) \\ &= \mathbb{P}_y(\mathcal{T} > t, X_{\Phi}(t) \in A) \\ &= \mathbb{P}_y(X_{\Phi}(t) \in A) - \mathbb{P}_y(\mathcal{T} \leq t, X_{\Phi}(t) \in A) \end{aligned} \quad (3.10)$$

$$= \mathbb{P}_y(X_{\Phi}(t) \in A) - \int_0^t \mathbb{P}_y(X_{\Phi}(t) \in A | \mathcal{T} = w) \mathbb{P}_y(\mathcal{T} \in dw), \quad (3.11)$$

where we used (3.7). Since  $\varphi$  is non-decreasing and continuous we have that  $\mathcal{T} \in \overline{\mathcal{M}}$  and thus  $X_{\Phi}$  is Markov at  $\mathcal{T}$  (see the discussion above). Hence

$$\begin{aligned} & \mathbb{P}_y \left( X_{\Phi}^{\dagger}(t) \in A \right) \\ &= \mathbb{P}_y(X_{\Phi}(t) \in A) - \int_0^t \mathbb{P}_y(X_{\Phi}(t) \in A | X_{\Phi}(H(w)) = \varphi(w), \mathcal{T} = w) \mathbb{P}_y(\mathcal{T} \in dw) \\ &= \mathbb{P}_y(X_{\Phi}(t) \in A) - \int_0^t \mathbb{P}_{\varphi(w)}(X_{\Phi}(t-w) \in A) \mathbb{P}_y(\mathcal{T} \in dw) \end{aligned} \quad (3.12)$$

and then, by using Fubini's theorem

$$\begin{aligned} & \mathbb{P}_y \left( X_{\Phi}^{\dagger}(t) \in A \right) \\ &= \int_A p_{\Phi}(t, x; y) dx - \int_A \int_0^t p_{\Phi}(t-w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) dx. \end{aligned} \quad (3.13)$$

Being  $A$  any Borel set and  $t \geq 0$  arbitrary, we conclude the proof.  $\square$

**Remark 3.2.** *Equation (3.9) is already well-known in the Markov context and it is called the Dynkin-Hunt formula. A precise statement of such a formula is given in [20, Corollary 3.14]. Here we proved a semi-Markov generalization of it.*

It is clear that the smoothness properties of  $q(t, x; y)$  depend on  $p_{\Phi}(t, x; y)$ . Furthermore, smoothness properties of  $p_{\Phi}(t, x; y)$  depend on  $f_L(s, t)$  which can be expressed by assuming smoothness of the density of the corresponding subordinator.

Such a density is, in turn, smooth enough under the following assumption (which is called Orey's condition).

- There exist  $\gamma \in (0, 2)$ ,  $C_\gamma > 0$  and  $r_0 > 0$  such that

$$\int_{(0,r)} s^2 \nu(ds) > C_\gamma r^\gamma, \text{ for any } r < r_0. \quad (\mathbf{A3})$$

**Remark 3.3.** *It is not difficult to see that, due to the integrability condition  $\int_{(0,1)} s \nu(ds) < \infty$  for the Lévy measure of a subordinator, we must have  $\gamma \in (1, 2)$ .*

The name of such condition follows from [46], in which it is shown that it implies

$$|\mathbb{E}[e^{i\xi\sigma(1)}]| \leq e^{-\frac{C_\gamma}{4}\xi^{2-\gamma}} \quad (3.14)$$

for  $|\xi|$  big enough. If we write  $\Psi(\xi) := \Phi(-i\xi)$ , it holds

$$\mathbb{E}[e^{i\xi\sigma(t)}] = e^{-t\Psi(\xi)} \quad (3.15)$$

and thus Orey's condition implies

$$|\mathbb{E}[e^{i\xi\sigma(t)}]| \leq e^{-\frac{C_\gamma t}{4}\xi^{2-\gamma}}, \text{ for } |\xi| \text{ big enough.} \quad (3.16)$$

We remark that this also implies that the corresponding subordinator admits, for any fixed  $t > 0$ , a one-dimensional density  $s \mapsto g_\sigma(s, t)$  which is infinitely differentiable on  $(0, \infty)$ .

Using the previous assumptions, we can prove suitable regularity properties for  $p_\Phi$ . In particular, we show that  $t \mapsto p_\Phi(t, x, y)$  is differentiable on  $(0, \infty)$  and this partial derivative is continuous as a function of three variables whenever  $x \neq y$  and further, it can be extended as  $t \rightarrow 0$  in such a way to be locally dominated. Being the result quite technical, the proof of it is postponed to Appendix A.1.

**Proposition 3.4.** *Under A1 and A3 we have that:*

- (1) *For any  $x \neq y$  the function  $p_\Phi(\cdot, x; y)$  is differentiable on  $(0, \infty)$  and*

$$\partial_t p_\Phi(t, x; y) = \int_0^{+\infty} p(s, x; y) \partial_t f_L(s, t) ds. \quad (3.17)$$

- (2) *The derivative  $\partial_t p_\Phi(t, x; y)$  is continuous on  $(0, \infty) \times (\mathbb{R}^2 \setminus \{(x, x), x \in \mathbb{R}\})$  and further  $\lim_{t \rightarrow 0^+} \partial_t p_\Phi(t, x; y) \rightarrow 0$  locally uniformly with respect to  $(x, y) \in \mathbb{R}^2 \setminus \{(x, x), x \in \mathbb{R}\}$ .*
- (3) *For any  $a_1 < a_0 < a < b < b_0 < b_1$  and  $T > 0$ , let  $I_T = [0, T]$ ,  $I_1 = [a_1, b_1] \setminus [a_0, b_0]$  and  $I_2 = [a, b]$  and define  $\mathcal{I} := I_T \times I_1 \times I_2$ . Then there exists a function  $f_{\mathcal{I}} \in L^1([a_1, b_1])$  such that*

$$|\partial_t p_\Phi(t, x, y)| \leq f_{\mathcal{I}}(x) \text{ for all } (t, x; y) \in \mathcal{I}. \quad (3.18)$$

**Remark 3.5.** *Let us stress out that Theorem 3.4 implies, via Lemma 1.1, that  $\partial_t^\Phi p(\cdot, x; y)$  is continuous as  $x \neq y$ .*

We also need to prove some regularity properties of the function  $\partial_t^\Phi p_\Phi(t, x, y)$ . It turns out that the latter follow if we ensure the existence of  $\partial_s f_L(\cdot, t)$  and its integrability against  $p(\cdot, x; y)$ . This is partially done in the following Proposition, whose proof is given in Appendix A.2.

**Proposition 3.6.** Under **A1** and **A3** the derivative  $\partial_s f_L(\cdot, t)$  exists on  $(0, \infty)$  and

$$(0, \infty) \ni s \mapsto \partial_s f_L(s, t) = \int_0^t \bar{\nu}(\tau) \partial_s g_\sigma(t - \tau, s) d\tau. \quad (3.19)$$

Further, it holds  $s f_L(s, t) \rightarrow 0$  and  $s^2 |\partial_s f_L(s, t)| \rightarrow 0$  as  $s \rightarrow +\infty$ , locally uniformly with respect to  $t \in \mathbb{R}^+$ .

Once  $\partial_s f_L(s, t)$  exists and behaves at infinity as described by Proposition 3.6, we only need to ensure its integrability against  $p(s, x; y)$  near 0. Such property will be explicitly requested via the following assumption, as it will be also used to bound  $\partial_x^2 p_\Phi(t, x; y)$  near the diagonal  $x = y$ .

- For any  $0 < a \leq b$  there exists a function  $h(s)$  and a  $\delta > 0$  such that  $|\partial_s f_L(s, t)| \leq h(s)$  for any  $(s, t) \in (0, \delta) \times [a, b]$  (A4)  
and  $s \mapsto h(s)/\sqrt{s}$  is integrable in  $(0, \delta)$ .

**Remark 3.7.** We observe that **A4** holds in several cases, as we will see in Section 5 below where we will compute some examples. We conjecture that the validity of **A4** could be checked, for more general subordinators, by using the same strategy that is adopted in the examples. In practice, the technique implies that a power series representation can be obtained for the density  $f_L(s, t)$ , allowing to differentiate under the sum. We suppose that such a representation can be obtained with Laplace inversion, exactly as it has been done in the references cited in the examples. However, it should rely on some cumbersome computation and we will take care of these technical details in a subsequent paper.

The previous results refer to the regularity of  $p_\Phi(t, x; y)$  with respect to the time variable  $t$ . Concerning the space variable  $x$  we will use the following result, whose proof is left in Appendix A.3.

**Proposition 3.8.** Under **A1**,  $\partial_x p_\Phi(t, x, y)$  and  $\partial_x^2 p_\Phi(t, x, y)$  exist and are continuous functions in  $\mathbb{R}_0^+ \times (\mathbb{R}^2 \setminus \{(x, x), x \in \mathbb{R}\})$ . Furthermore, it holds

$$\partial_x p_\Phi(t, x; y) = -\sqrt{\frac{2}{\pi}} \int_0^{+\infty} \frac{(x-y)e^{-\frac{(x-y)^2}{2s}}}{s^{\frac{3}{2}}} f_L(s; t) ds, \quad (3.20)$$

$$\partial_x^2 p_\Phi(t, x; y) = \int_0^{+\infty} \sqrt{\frac{2}{\pi s^3}} \left( \frac{(x-y)^2}{s} - 1 \right) e^{-\frac{(x-y)^2}{2s}} f_L(s; t) ds. \quad (3.21)$$

Finally, in order to use Equation (3.9), together with the regularity results we gave in this Section, we also need to prove that  $p_\Phi$  is the fundamental solution of the non-local heat equation in the whole domain. Let us stress out that this result can be easily achieved via semigroup theory appealing to the abstract Cauchy problem satisfied by the mapping  $t \mapsto \mathbb{E}^x u(X_\Phi(t))$ . Another possibility would be resorting to the theory in [28, Section 5] which, however, rests on slightly different assumptions on the non-local time operator. We are interested in the fact that  $p_\Phi$  is a pointwise (almost everywhere) solution of the aforementioned equation.

**Theorem 3.9.** Suppose Assumptions **A1**, **A3** hold. Then, for  $t > 0$  and  $s > 0$ , it is true that

$$\partial_t^\Phi f_L(s, t) = -\partial_s f_L(s, t). \quad (3.22)$$

Furthermore, for  $t > 0$ ,  $y \in \mathbb{R}$  and  $x \neq y$ , it is true that

$$\partial_t^\Phi p_\Phi(t, x; y) = \frac{1}{2} \partial_x^2 p_\Phi(t, x; y). \quad (3.23)$$

*Proof.* We first prove (3.23), then the proof of (3.22) can be conducted analogously. We prove before that, for  $x \neq y$ , and every  $t > 0$ ,

$$\begin{aligned} I_t^\Phi p_\Phi(t, x; y) &:= \int_0^t p_\Phi(s, x; y) \bar{\nu}(t-s) ds \\ &= \frac{1}{2} \int_0^t \partial_x^2 p_\Phi(s, x; y) ds =: I_t \frac{1}{2} \partial_x^2 p_\Phi(t, x; y). \end{aligned} \quad (3.24)$$

Once this is done, Proposition 3.8 guarantees that the right-hand side of (3.24) is in  $C^1$  and thus we can differentiate both sides of the equation to get (3.23). Denoting  $\tilde{u}(\lambda)$  the Laplace transform ( $t \mapsto \lambda$ ) of a function  $u(t)$  we have that, for  $x \neq y$ , using Tonelli's theorem and (3.2),

$$\begin{aligned} \tilde{p}_\Phi(\lambda, x; y) &= \int_0^{+\infty} e^{-\lambda t} \int_0^{+\infty} p(s, x; y) f_L(s, t) ds dt \\ &= \frac{\Phi(\lambda)}{\lambda} \tilde{p}(\Phi(\lambda), x; y) \end{aligned} \quad (3.25)$$

where we used also [38, eq. (3.13)] to say that

$$\tilde{f}_L(s, \lambda) = \frac{\Phi(\lambda)}{\lambda} e^{-s\Phi(\lambda)}. \quad (3.26)$$

Let us recall that

$$\int_0^{+\infty} e^{-\lambda t} |\partial_x p(t, x; y)| dt < \infty \quad (3.27)$$

as well as

$$\int_0^{+\infty} e^{-\lambda t} |\partial_x^2 p(t, x; y)| dt < \infty \quad (3.28)$$

and

$$\partial_t p(t, x; y) = \frac{1}{2} \partial_x^2 p(t, x; y). \quad (3.29)$$

Now since

$$\int_0^{+\infty} e^{-\lambda t} \bar{\nu}(t) dt = \frac{\Phi(\lambda)}{\lambda}, \quad (3.30)$$

we obtain that

$$\begin{aligned} &\widetilde{I_t^\Phi p_\Phi}(\lambda, x; y) \\ &= \frac{\Phi(\lambda)}{\lambda} \frac{\Phi(\lambda)}{\lambda} \tilde{p}(\Phi(\lambda), x; y) \quad \text{by (3.25), (3.30) and [1, Proposition 1.6.4]} \\ &= \frac{\Phi(\lambda)}{\lambda^2} \widetilde{\partial_t p}(\Phi(\lambda), x; y) \quad \text{by [1, Corollary 1.6.6]} \\ &= \frac{\Phi(\lambda)}{2\lambda^2} \widetilde{\partial_x^2 p}(\Phi(\lambda), x; y) \quad \text{by (3.29)} \\ &= \frac{1}{\lambda} \frac{1}{2} \partial_x^2 \frac{\Phi(\lambda)}{\lambda} \tilde{p}(\Phi(\lambda), x; y) \quad \text{by (3.27) and (3.28)} \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{\lambda} \frac{1}{2} \partial_x^2 \widetilde{p}_\Phi(\lambda, x; y) && \text{by (3.25)} \\
&= \frac{1}{2} \widetilde{I_t \partial_x^2 p_\Phi}(\lambda, x; y) && \text{by [1, Corollary 1.6.5].} \quad (3.31)
\end{aligned}$$

Since both-sides of (3.24) are continuous functions, then (3.31) guarantees that (3.24) is true for any  $t > 0$ , by injectivity of Laplace transform [1, Theorem 1.7.3]. Analogously, to prove (3.22) we just need to show that

$$I_t^\Phi f_L(s, t) := \int_0^t f_L(s, \tau) \bar{v}(t - \tau) d\tau = - \int_0^t \partial_s f_L(s, \tau) d\tau, \quad (3.32)$$

arguing as above with a double Laplace transform and observing that the existence of both sides of (3.32) is guaranteed by Proposition 3.6.  $\square$

**Remark 3.10.** *Let us observe that  $p_\Phi(t, x; y)$  is actually the fundamental solution of the non-local heat equation. Indeed, for any initial datum  $f \in C_c(\mathbb{R})$ , the function*

$$u(t, x) = \int_{\mathbb{R}} p_\Phi(t, x; y) f(y) dy$$

*is  $C^2$  in the  $x$  variable for any fixed  $t > 0$ , admits continuous non-local derivative in the  $t$  variable (for  $t > 0$ ) for any fixed  $x \in \mathbb{R}$  and solves*

$$\begin{cases} \partial_t^\Phi u(t, x) = \frac{1}{2} \partial_x^2 u(t, x) & (t, x) \in \mathbb{R}^+ \times \mathbb{R} \\ u(0, x) = f(x) & x \in \mathbb{R}. \end{cases}$$

*Moreover, it is the unique solution such that  $\lim_{x \rightarrow \pm\infty} u(t, x) = 0$  locally uniformly in  $t \geq 0$ . Finally, let us observe that if  $\Phi$  is a complete Bernstein function, then, by [2, Theorem 3], it is analytic (as a function  $u : t \in \mathbb{R}_+ \mapsto C^0(\mathbb{R})$ ) for  $t > 0$ .*

**3.2. The non-local problem.** Here is the precise statement on the existence of the solution to (1.14), by means of a the stochastic representation based on  $q(t, x; y)$ , i.e., the density of the killed time-changed Brownian motion.

**Theorem 3.11.** *Let  $\mathcal{O}$  be defined as in (1.13). Under **A1**, **A2(a)**, **A2(b)**, **A3** and **A4**, for any function  $f \in C_c(-\infty, \varphi(0))$ , the unique classical solution of*

$$\partial_t^\Phi u(x, t) = \frac{1}{2} \partial_x^2 u(x, t), \quad (t, x) \in \mathcal{O}, \quad (3.33)$$

*in the sense of Definition 2.1, subject to the boundary conditions*

$$\begin{cases} u(t, x) = 0, & (t, x) \in \mathcal{O}^c \setminus \partial_0 \mathcal{O}, \\ \lim_{x \rightarrow -\infty} u(t, x) = 0, & \text{locally uniformly on } t > 0, \end{cases} \quad (3.34)$$

*and initial condition*

$$u(t, x) = f(x), \quad (t, x) \in \partial_0 \mathcal{O}, \quad (3.35)$$

*is given by*

$$u(t, x) = \int_{-\infty}^{\varphi(0)} q(t, x; y) f(y) dy, \quad (3.36)$$

*where  $(t, x; y) \mapsto q(t, x; y)$  is defined in Lemma 3.1. Furthermore  $q(t, x; y)$  weakly converges to  $\delta_y(dx)$  as  $t \rightarrow 0$ . As a consequence,  $q(t, x; y)$  is a fundamental solution of (3.33) under the conditions (3.34) and (3.35).*

*Moreover, suppose  $q_1(t, x; y)$  is another fundamental solution of (3.33) under the conditions (3.34) and (3.35). Then there exists a set  $\mathcal{N} \subset (-\infty, \varphi(0))$ , with*

zero Lebesgue measure, such that  $q(t, x; y) = q_1(t, x; y)$  for any  $(t, x) \in \mathcal{O}$  and  $y \in (-\infty, \varphi(0)) \setminus \mathcal{N}$ .

*Proof.* Let us first show that  $q(t, x; y)$  satisfies eq. (3.33), for  $x \neq y$ , subject to (3.34) and weakly converges to  $\delta_y(dx)$ , for fixed  $y \in (-\infty, \varphi(0))$ . To do this we resort to the representation given in Lemma 3.1. Fix  $y \in (-\infty, \varphi(0))$  and observe that continuity of  $\mathbb{R}^{+,1} \ni (t, x) \mapsto q(t, x; y)$  follows from equation (3.9) and the fact that  $\mathcal{T}$  is a continuous r.v (see (3.7)), since  $q(\cdot, \cdot; y)$  is the sum of two continuous functions in  $\mathbb{R}^{+,1}$ .

Now we show that the boundary conditions hold. It is easy to see that if  $x > \varphi(t)$  for some  $t \geq 0$  then  $q(t, x; y) = 0$ . Indeed, for any  $\varepsilon > 0$  small enough, we have

$$\mathbb{P}_y(\mathcal{T} > t, X_\Phi(t) \in [x - \varepsilon, x + \varepsilon]) = 0 \quad (3.37)$$

and then

$$\frac{1}{2\varepsilon} \int_{x-\varepsilon}^{x+\varepsilon} q(t, z; y) dz = \frac{1}{2\varepsilon} \mathbb{P}_y(\mathcal{T} > t, X_\Phi(t) \in [x - \varepsilon, x + \varepsilon]) = 0. \quad (3.38)$$

Thus, being  $q$  continuous, taking the limit as  $\varepsilon \rightarrow 0$ , we achieve  $q(t, x; y) = 0$ . By continuity, the same holds as  $x = \varphi(t)$ , in particular  $q(t, \varphi(t); y) = 0$ . Let us also observe that, for fixed  $y \in (-\infty, \varphi(0))$ ,

$$\lim_{x \rightarrow -\infty} q(t, x; y) = 0, \quad \text{locally uniformly for } t > 0, \quad (3.39)$$

as a direct consequence of the fact that

$$\lim_{x \rightarrow -\infty} p_\Phi(t, x; y) = 0$$

locally uniformly for  $(t, y) \in \mathbb{R}^{+,1}$ . Indeed, if we define  $\mathcal{K} := [0, T] \times [a, +\infty)$  for  $T > 0$  and  $a \in \mathbb{R}$  arbitrary,

$$\begin{aligned} \sup_{(t,y) \in \mathcal{K}} p_\Phi(t, x; y) &\leq \sup_{t \in [0, T]} \int_0^\infty \frac{1}{\sqrt{2\pi s}} e^{-\frac{(a-x)^2}{2s}} f_L(s, t) ds \\ &\leq \frac{1}{\sqrt{2\pi e}(a-x)}. \end{aligned} \quad (3.40)$$

Now we show that the initial condition is satisfied. Let  $f \in C_b(\mathbb{R})$  and observe that, by Lemma 3.1,

$$\begin{aligned} &\int_{\mathbb{R}} q(t, x; y) f(x) dx \\ &= \int_{\mathbb{R}} p_\Phi(t, x; y) f(x) dx - \int_0^t \int_{\mathbb{R}} p_\Phi(t-w, x; \varphi(w)) f(x) dx \mathbb{P}_y(\mathcal{T} \in dw). \end{aligned} \quad (3.41)$$

Clearly, we only need to show that the second term in (3.41) tends to zero, since the first one converges towards  $f(y)$ . To do this note that

$$\left| \int_0^t \int_{\mathbb{R}} p_\Phi(t-w, x; \varphi(w)) f(x) dx \mathbb{P}_y(\mathcal{T} \in dw) \right| \leq \|f\|_{L^\infty(\mathbb{R})} \mathbb{P}(\mathcal{T} \leq t) \rightarrow 0, \quad (3.42)$$

as  $t \rightarrow 0$ . This proves the weak convergence of  $q(t, x; y)$  to  $\delta_y(dx)$  as  $t \rightarrow 0$ . We focus now on the equation. First recall that  $\partial_t^\Phi p(t, x; y)$  and  $\partial_x^2 p_\Phi(t, x; y)$  exist

respectively by Propositions 3.4 and 3.8. By (3.23), we could write, for  $x \neq y$ ,

$$\begin{aligned}
\partial_t^\Phi q(t, x; y) &= \partial_t^\Phi p_\Phi(t, x; y) - \partial_t^\Phi \int_0^t p_\Phi(t-w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) \\
&= \partial_t^\Phi p_\Phi(t, x; y) - \int_0^t \partial_t^\Phi p_\Phi(t-w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) \\
&= \frac{1}{2} \partial_x^2 p_\Phi(t, x; y) - \frac{1}{2} \int_0^t \partial_x^2 p_\Phi(t-w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) \\
&= \frac{1}{2} \left( \partial_x^2 p_\Phi(t, x; y) - \partial_x^2 \int_0^t p_\Phi(t-w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) \right) \\
&= \frac{1}{2} \partial_x^2 q(t, x; y). \tag{3.43}
\end{aligned}$$

provided that

$$\partial_x \int_0^t p_\Phi(t-w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) = \int_0^t \partial_x p_\Phi(t-w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) \tag{3.44}$$

$$\partial_x^2 \int_0^t p_\Phi(t-w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) = \int_0^t \partial_x^2 p_\Phi(t-w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) \tag{3.45}$$

$$\partial_t^\Phi \int_0^t p_\Phi(t-w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) = \int_0^t \partial_t^\Phi p_\Phi(t-w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw). \tag{3.46}$$

Let us first prove (3.44). Let us suppose, without loss of generality, that  $\varphi(0) > 0$  and  $y = 0$ . By (3.20) we get

$$\partial_x p_\Phi(t, x; 0) = -\sqrt{\frac{2}{\pi}} \int_0^{+\infty} \frac{x e^{-\frac{x^2}{2s}}}{s^{\frac{3}{2}}} f_L(s; t) ds. \tag{3.47}$$

that, by using the change of variables  $z = \frac{x}{\sqrt{2s}}$  for  $x > 0$ , becomes

$$\partial_x p_\Phi(t, x; 0) = -\frac{4}{\sqrt{\pi}} \int_0^{+\infty} e^{-z^2} f_L\left(\frac{x}{2z^2}; t\right) dz. \tag{3.48}$$

Since, by Proposition 3.6,  $\lim_{s \rightarrow +\infty} f_L(s; t) = 0$  locally uniformly with respect to  $t \in \mathbb{R}^+$ , then  $f_L(s; t)$  is bounded in  $[0, +\infty) \times [a, b]$  for any  $0 < a < b$ . Thus, by dominated convergence theorem, we have

$$\lim_{x \rightarrow 0^+} \partial_x p_\Phi(t, x; 0) = -\frac{\bar{\nu}(t)}{2}, \tag{3.49}$$

and the limit is locally uniform with respect to  $t \in \mathbb{R}^+$ , where we used the fact that  $f_L(0, t) = \bar{\nu}(t)$ . With the same argument we have, in general,

$$\lim_{x \rightarrow 0^\pm} \partial_x p_\Phi(t, x; 0) = \mp \frac{\bar{\nu}(t)}{2} \tag{3.50}$$

and both limits hold locally uniformly with respect to  $t \in \mathbb{R}^+$ . Now let us consider the integral of  $p_\Phi$  against  $\mathbb{P}_y(\mathcal{T} \in dw)$ . Since  $(t, x) \in \mathcal{O}$ , we have that  $x < \varphi(t)$ . By **A2(a)** we know that the cardinality of  $\{t \geq 0 : x = \varphi(t)\}$  is at most 1 and thus, by continuity of  $\mathcal{T}$ ,  $\partial_x p_\Phi(t-w, x; \varphi(w))$  is defined  $\mathbb{P}_y(\mathcal{T} \in dw)$ -almost everywhere. If  $x < \varphi(0)$ , then, for any  $w \in [0, t]$ , there exist  $a, b \in \mathbb{R}$  such that

$a < x - \varphi(t) < x - \varphi(w) < x - \varphi(0) < b < 0$  and then we know, from (3.20), that  $|\partial_x p_\Phi(t - w, z; \varphi(w))|$  is bounded as  $w \leq t$  and  $z$  belongs to a sufficiently small neighbourhood of  $x$  (so that  $z - \varphi(0)$  is separated from 0). This is enough to prove (3.44) as  $x < \varphi(0)$ . Now let us suppose  $x \in [\varphi(0), \varphi(t)]$ . Then, by A2(a) we know that there exists a unique  $0 \leq w_0 < t$  such that  $x = \varphi(w_0)$ . Let us consider  $\varepsilon > 0$  such that  $w_0 < w_0 + \varepsilon < t$  and let us write

$$\begin{aligned} & \partial_x \int_0^t p_\Phi(t - w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) \\ &= \partial_x \left( \int_0^{w_0 + \varepsilon} p_\Phi(t - w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) \right. \\ & \quad \left. + \int_{w_0 + \varepsilon}^t p_\Phi(t - w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) \right). \end{aligned}$$

We can take the derivative inside the second integral with the same argument as in the case  $x < \varphi(0)$ . Concerning the first integral, since  $0 < t - w_0 - \varepsilon \leq t - w$ , by the locally uniform limits (3.50) we know that  $|\partial_x p_\Phi(t - w, z; \varphi(w))|$  is bounded as  $z$  belongs to a small neighbourhood of  $x$ . This concludes the proof of (3.44).

Now we show (3.45). Again, in case  $x < \varphi(0)$ , equation (3.21) guarantees that  $|\partial_x^2 p_\Phi(t - w, z; \varphi(w))|$  is bounded as  $w \leq t$  and  $z$  belongs to a small neighbourhood of  $x$ , thus we only have to study the case  $x \in [\varphi(0), \varphi(t)]$ . To deal with it we proceed as follows. Note that, if  $x \neq 0$ ,

$$\begin{aligned} & \partial_t^\Phi p_\Phi(t, x; 0) \\ &= \int_0^t \bar{v}(t - \tau) \partial_t p_\Phi(\tau, x; 0) d\tau && \text{by Lemma 1.1 and Theorem 3.4} \\ &= \int_0^t \bar{v}(t - \tau) \int_0^\infty p(s, x; 0) \partial_t f_L(s, \tau) ds d\tau && \text{by (3.17)} \\ &= \int_0^{+\infty} p(s, x; 0) \partial_t^\Phi f_L(s; t) ds, && \text{by Fubini's theorem} \\ &= - \int_0^{+\infty} p(s, x; 0) \partial_s f_L(s; t) ds && \text{by Lemma 3.9.} \end{aligned} \tag{3.51}$$

Now fix  $[a, b]$  with  $0 < a < t < b$ , let  $\delta$  be fixed as in A4 and observe that

$$\begin{aligned} \int_0^{+\infty} p(s, x; 0) |\partial_s f_L(s; t)| ds &= \int_0^\delta p(s, x; 0) |\partial_s f_L(s; t)| ds \\ & \quad + \int_\delta^{+\infty} p(s, x; 0) |\partial_s f_L(s; t)| ds \\ &\leq \frac{1}{\sqrt{2\pi}} \int_0^\delta s^{-\frac{1}{2}} |\partial_s f_L(s; t)| ds \\ & \quad + \frac{1}{\sqrt{2\pi}} \int_\delta^{+\infty} s^{-\frac{1}{2}} |\partial_s f_L(s; t)| ds \\ &=: \frac{1}{\sqrt{2\pi}} (I_1 + I_2). \end{aligned} \tag{3.52}$$

Working with  $I_1$ , by **A4** we know that there exists a function  $h(s)$  such that  $|\partial_s f_L(s; t)| \leq h(s)$  and

$$I_1 \leq \int_0^\delta s^{-1/2} h(s) ds < +\infty.$$

Concerning  $I_2$  we have, by Proposition 3.6, that  $s^2 |\partial_s f(s; t)| \leq C_1(\delta)$  where

$$C_1(\delta) := \max_{(s,t) \in K_\delta} s^2 |\partial_s f(s; t)|. \quad (3.53)$$

This leads to

$$I_2 \leq C_1(\delta) \int_\delta^{+\infty} s^{-5/2} ds < +\infty.$$

Thus we can use dominated convergence to say that

$$\lim_{x \rightarrow 0} \partial_t^\Phi p_\Phi(t, x; 0) = -\frac{1}{\sqrt{2\pi}} \int_0^{+\infty} s^{-\frac{1}{2}} \partial_s f_L(s; t) ds \quad (3.54)$$

and thus, by (3.23),

$$\lim_{x \rightarrow 0} \partial_x^2 p_\Phi(t, x; 0) = -\frac{2}{\sqrt{2\pi}} \int_0^{+\infty} s^{-\frac{1}{2}} \partial_s f_L(s; t) ds.$$

Note that both limits are locally uniform with respect to  $t \in \mathbb{R}^+$ . Arguing as we did with the first derivative, we obtain that (3.45) is true even if  $x \in [\varphi(0), \varphi(t)]$ .

Now we prove (3.46). By Fubini's theorem we easily get that

$$\begin{aligned} & \partial_t^\Phi \left( \int_0^t p_\Phi(t-w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) \right) \\ &= \partial_t \int_0^t \int_0^{t-s} p_\Phi(t-s-w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) \bar{\nu}(s) ds \\ &= \partial_t \int_0^t \int_0^{t-w} p_\Phi(t-s-w, x; \varphi(w)) \nu(s) ds \mathbb{P}_y(\mathcal{T} \in dw) \\ &= \partial_t \int_0^{+\infty} k(t, w, x) \mathbb{P}_y(\mathcal{T} \in dw), \end{aligned} \quad (3.55)$$

where we set

$$k(t, w, x) := \mathbb{1}_{\mathbb{R}^+}(t-w) \int_0^{t-w} p_\Phi(t-s-w, x; \varphi(w)) \bar{\nu}(s) ds. \quad (3.56)$$

Note that, for  $w \neq t$ ,

$$\partial_t k(t, w, x) = \mathbb{1}_{\mathbb{R}^+}(t-w) \partial_t^\Phi p_\Phi(t-w, x; \varphi(w)). \quad (3.57)$$

Moreover, let us remark that if  $x - \varphi(w)$  is separated from 0 as  $|w - t| \rightarrow 0$ , when  $t \in [a, b]$  for some constants  $0 < a < b$ , then, by item (2) of Proposition 3.4,

$$\lim_{\substack{|t-w| \rightarrow 0 \\ t \in [a, b]}} \partial_t k(t, w, x) = 0. \quad (3.58)$$

The latter limit is clearly true for any  $x < \varphi(0)$ . In this case, rewriting (3.21) as

$$\partial_x^2 p_\Phi(t, x; y) = \mathbb{E} \left[ \sqrt{\frac{2}{\pi(L(t))^3}} \left( \frac{(x-y)^2}{L(t)} - 1 \right) e^{\frac{(x-y)^2}{L(t)}} \right] \quad (3.59)$$

it is clear that  $|\partial_x^2 p_\Phi(s-w, x; \varphi(w))|$  is bounded as  $s$  belongs to a small neighbourhood of  $t > 0$  and  $w \leq s$  and then Equation (3.23) guarantees that this is also true

for  $|\partial_t^\Phi p_\Phi(s-w, x; \varphi(w))|$ . Combining the latter information with (3.58), we know that  $|\partial_t k(s, w, x)|$  is bounded as  $s$  belongs to a small neighbourhood of  $t > 0$  and  $w \in \mathbb{R}_0^+$ . Thus, for  $x < \varphi(0)$ , we conclude that

$$\begin{aligned} \partial_t^\Phi \left( \int_0^t p_\Phi(t-w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) \right) &= \partial_t \int_0^\infty k(t, w, x) \mathbb{P}_y(\mathcal{T} \in dw) \\ &= \int_0^\infty \partial_t k(t, w, x) \mathbb{P}_y(\mathcal{T} \in dw) \\ &= \int_0^t \partial_t^\Phi p_\Phi(t-w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw), \end{aligned} \quad (3.60)$$

where we used dominated convergence theorem and (3.57).

Now let us consider the case  $x \in [\varphi(0), \varphi(t)]$ . In this case we know there exists a unique  $w_0$  such that  $\varphi(w_0) = x$ . Note that  $t - w_0 > 0$  and then there exists  $a_0$  such that  $t - w_0 > a_0$ . If  $t < b$  for some  $b > 0$ , we get  $w_0 < b - a_0$  and  $a_0 < t - w_0 < b$ . Arguing as in equations (3.51) and (3.52) we have

$$|\partial_t^\Phi p_\Phi(t - w_0, x - \varphi(w_0); 0)| \leq \frac{1}{\sqrt{2\pi}}(I_1 + I_2). \quad (3.61)$$

Now observe that by (3.54), as soon as  $s-w$  belongs to a compact set separated from zero, we can extend  $(s, w) \mapsto \partial_t^\Phi p_\Phi(s-w, x - \varphi(w); 0)$  by continuity in  $(s, w_0)$ . Let us now consider  $0 < a < b$  such that  $b-a < a_0$  and  $w_0 < a < t < b$ . Observe that for any  $s \in [a, b]$ , it holds  $s-b+a_0 \geq a-b+a_0 > 0$ . Moreover, for any  $0 \leq w < b-a_0$  it holds  $0 < a-b+a_0 \leq s-b+a_0 < s-w \leq b$ . Thus, if  $s \in [a, b]$  and  $w \in [0, b_1]$  for some  $b_1 < b-a_0$  such that  $w_0 < b_1$ ,  $s-w$  belongs to a compact set separated from zero,  $w_0 \in [0, b_1]$  and the function  $(s, w) \in [a, b] \times [0, b_1] \rightarrow |\partial_t^\Phi p_\Phi(s-w, x - \varphi(w); 0)|$  is continuous. In particular, by continuity and (3.61), there exists a compact set  $K' \subset [a, b] \times [0, b_1]$  such that  $(t, w_0)$  belongs to the interior of  $K'$  and

$$|\partial_t^\Phi p_\Phi(s-w, x - \varphi(w); 0)| \leq \frac{2}{\sqrt{2\pi}}(I_1 + I_2), \quad \forall (s, w) \in K'. \quad (3.62)$$

Without loss of generality we can suppose that  $K' = [a_2, b_2] \times [w_0 - \delta_w, w_0 + \delta_w]$  for some  $a < a_2 < t < b_2 < b$  and  $\delta_w > 0$  such that  $w_0 + \delta_w < a$ , therefore if  $|w - w_0| > \delta_w$  then there exists  $\varepsilon > 0$  such that  $|x - \varphi(w)| > \varepsilon$ . Furthermore, we have that there exists  $M > 0$  such that, for any  $w \leq b$ ,  $|x - \varphi(w)| < M$ . Then, for  $s \in [a_2, b_2]$ ,  $|w - w_0| > \delta_w$  and  $w \leq s$ , by (3.23) and (3.59), we have that

$$|\partial_t^\Phi p_\Phi(s-w, x - \varphi(w); 0)| \leq C_2(\varepsilon, M), \quad (3.63)$$

where  $C_2(\varepsilon, M)$  is a suitable constant. Moreover, since, as  $s \in [a_2, b_2]$ ,  $x - \varphi(w)$  is separated from 0 if  $|s - w|$  is small, (3.58) holds and (3.63) then implies

$$|\partial_t k(s, w, x)| \leq C_2(\varepsilon, M), \quad \forall s \in [a_2, b_2], \quad |w - w_0| > \delta_w. \quad (3.64)$$

Combining (3.62) and (3.64) yields

$$|\partial_t k(s, w, x)| \leq \max \left\{ C_2(\varepsilon, M), \frac{2}{\sqrt{2\pi}}(I_1 + I_2) \right\},$$

for all  $w \in \mathbb{R}_0^+$  and  $s \in [a_2, b_2]$ . The same argument as in (3.60) implies, via dominated convergence theorem, that equation (3.46) holds true even if  $x \in [\varphi(0), \varphi(t)]$ .

Now we consider the problem with initial data  $f \in C_c((-\infty, \varphi(0)))$ . To do this, let us define

$$u(t, x) = \int_{-\infty}^{\varphi(0)} q(t, x; y) f(y) dy.$$

Since  $q(t, x; y)$  is continuous in  $(t, x) \in \mathbb{R}^{+,1}$  with  $t > 0$  for any fixed  $y < \varphi(0)$ , if we show that  $q(t, x; y)$  is bounded in a neighbourhood of  $(t, x)$  by a constant independent of  $y \in \text{supp}(f)$ , where  $\text{supp}(f)$  denotes the support of  $f$ , we obtain, by dominated convergence theorem, that  $u$  is continuous in  $(t, x)$ . This is clearly true if  $x > \varphi(t)$ . If  $x < \varphi(t)$ , then there exist  $0 < a_t < t < b_t$  and  $a_x < x < b_x < \varphi(0)$  such that for  $(s, z) \in [a_t, b_t] \times [a_x, b_x]$  and  $w \in [0, t]$  it holds  $(t - w, x; \varphi(w)) \in \mathcal{K}$  where  $\mathcal{K} \subset \mathbb{R}^{+,3}$  is a compact set such that  $(0, 0, 0) \notin \mathcal{K}$ . Hence, for  $(s, z) \in [a_t, b_t] \times [a_x, b_x]$  it holds

$$\int_0^s |p_{\Phi}(s - w, z; \varphi(w))| \mathbb{P}_y(\mathcal{T} \in dw) \leq C_3(\mathcal{K}) \mathbb{P}_y(\mathcal{T} \leq s) \leq C_3(\mathcal{K}). \quad (3.65)$$

Moreover, we have, by (3.2) and the fact that  $L$  is increasing

$$p_{\Phi}(s, z; y) = \mathbb{E} \left[ \frac{1}{\sqrt{2\pi L(s)}} e^{-\frac{(z-y)^2}{2L(s)}} \right] \leq \mathbb{E} \left[ \frac{1}{\sqrt{2\pi L(a_t)}} \right] < +\infty, \quad (3.66)$$

where the latter inequality is implied by [3, Lemma 2.3]. These two inequalities prove that  $q(s, z; y)$  is bounded by a constant independent of  $y$  if  $(s, z)$  belongs to a neighbourhood of  $(t, x)$  with  $x < \varphi(t)$ .

We only need to prove the same statement in a point of the form  $(t, \varphi(t))$  for  $t > 0$ . To do this, let us first consider the function

$$g(t) = \mathbb{E}[(L(t))^{-\frac{1}{2}}]$$

that is well-defined by [3, Lemma 2.3]. Let us observe that we can evaluate the Laplace transform, by Fubini's theorem,

$$\tilde{g}(\lambda) = \frac{\Phi(\lambda)}{\lambda} \int_0^{+\infty} s^{-\frac{1}{2}} e^{-s\Phi(\lambda)} ds = \frac{\sqrt{\pi\Phi(\lambda)}}{\lambda}, \quad \lambda > 0.$$

In particular, this means that  $g \in L_{\text{loc}}^1(\mathbb{R}_0^+)$ . Now let us observe that, by (3.66), we get

$$\begin{aligned} \int_0^s |p_{\Phi}(s - w, z; \varphi(w))| \mathbb{P}_y(\mathcal{T} \in dw) &\leq \frac{1}{\sqrt{2\pi}} \int_0^s g(s - w) \mathbb{P}_y(\mathcal{T} \in dw) \\ &= \frac{\mathbb{E}_y[g(s - \mathcal{T}) \mathbf{1}_{(0,s)}(\mathcal{T})]}{\sqrt{2\pi}}. \end{aligned} \quad (3.67)$$

Let us stress that, since  $L$  is almost surely non-decreasing, the function  $w \in (0, s) \mapsto g(s - w)$  is non-decreasing. Let  $M_{\varphi} = \max_{w \in [0, b_t]} \varphi(w)$  and  $\mathcal{T}_M := \inf\{t \geq 0 : X_{\Phi}(t) \geq M_{\varphi}\}$ . It is clear that  $\mathcal{T} \mathbf{1}_{(0,s)}(\mathcal{T}) \leq \mathcal{T}_M \mathbf{1}_{(0,s)}(\mathcal{T}_M)$  almost surely. Hence we have,

$$\mathbb{E}_y[g(s - \mathcal{T}) \mathbf{1}_{(0,s)}(\mathcal{T})] \leq \mathbb{E}_y[g(s - \mathcal{T}_M) \mathbf{1}_{(0,s)}(\mathcal{T}_M)].$$

Now, since  $M_{\varphi}$  is meant for  $t$  fixed, we have that  $\mathcal{T}_M \stackrel{d}{=} \sigma(T_M)$  where  $T_M := \inf\{t \geq 0 : B(t) \geq M_{\varphi}\}$  (see [4]). With this in mind, we have that  $\mathcal{T}_M$  is an absolutely

continuous random variable with density

$$p_{M,\Phi}(t; y) = \int_0^{+\infty} \frac{(M_\varphi - y)}{\sqrt{2\pi s^3}} e^{-\frac{(M_\varphi - y)^2}{2s}} f_\Phi(s; t) ds = \mathbb{E} \left[ \frac{(M_\varphi - y)}{\sqrt{2\pi(L(t))^3}} e^{-\frac{(M_\varphi - y)^2}{2L(t)}} \right].$$

Hence we have

$$\mathbb{E}_y[g(s - \mathcal{T}_M)\mathbf{1}_{(0,s)}(\mathcal{T}_M)] = \int_0^s g(s - w)p_{M,\Phi}(w; y)dw. \quad (3.68)$$

If we denote  $m = \min \text{supp}(f)$ , we have

$$\begin{aligned} p_{M,\Phi}(w; y) \mathbb{E} \left[ \frac{(M_\varphi - y)}{\sqrt{2\pi(L(w))^3}} e^{-\frac{(M_\varphi - y)^2}{2L(w)}} \right] &\leq \mathbb{E} \left[ \frac{(M_\varphi - m)}{\sqrt{2\pi(L(w))^3}} e^{-\frac{(M_\varphi - \varphi(0))^2}{2L(w)}} \right] \\ &\leq C_4(M_\varphi, m) \end{aligned}$$

where  $C_4(M_\varphi, m) = \sup_{\tau \in \mathbb{R}^+} \frac{(M_\varphi - m)}{\sqrt{2\pi\tau^3}} e^{-\frac{(M_\varphi - \varphi(0))^2}{2\tau}}$ , and then (3.68) implies

$$\begin{aligned} \mathbb{E}_y[g(s - \mathcal{T}_M)\mathbf{1}_{(0,s)}(\mathcal{T}_M)] &\leq C_4(M_\varphi, m) \int_0^s g(s - w)dw \\ &\leq C_4(M_\varphi, m) \int_0^{b_t} g(w)dw =: C_5(M_\varphi, m, b_t). \end{aligned} \quad (3.69)$$

Tracing back to (3.67) we get

$$\int_0^s |p_\Phi(s - w, z; \varphi(w))| \mathbb{P}_y(\mathcal{T} \in dw) \leq \frac{C_5(m, M_\varphi, b_t)}{\sqrt{2\pi}}$$

Combining the previous inequality with (3.66) we conclude that  $q(s, z; y)$  is bounded as  $(s, z)$  belong to a suitable neighbourhood of  $(t, \varphi(t))$  by a constant independent of  $y \in \text{supp}(f)$ . Thus we conclude that  $u$  is continuous in any  $(t, x) \in \mathbb{R}^{+,1}$  such that  $t > 0$ .

We are left to prove the continuity of  $u$  on points of the form  $(0, x)$  as  $x \leq \varphi(0)$ . Writing explicitly the integral representation of  $q$  we have

$$u(t, x) = \int_{-\infty}^{\varphi(0)} p_\Phi(t, x; y) f(y) dy - \int_{-\infty}^{\varphi(0)} \int_0^t p_\Phi(t - w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) f(y) dy. \quad (3.70)$$

In the same spirit of (3.69) it is possible to see that

$$\int_{-\infty}^{\varphi(0)} \int_0^t p_\Phi(t - w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) f(y) dy \rightarrow 0 \quad (3.71)$$

locally uniformly to 0 as  $t \rightarrow 0^+$ , thus we only have to work with the first integral in (3.70).

Let us fix any  $(0, x_0) \in \partial_0 \mathcal{O}$  and consider a sequence  $(t_n, x_n) \in \mathcal{O}$  such that  $(t_n, x_n) \rightarrow (0, x_0)$ . If  $x_0 \notin \text{supp}(f)$ , then we may assume that

$$d = \min_{n \in \mathbb{N}} \text{dist}(x_n, \text{supp}(f)) > 0. \quad (3.72)$$

Thus we have

$$\begin{aligned} \left| \int_{-\infty}^{\varphi(0)} p_\Phi(t_n, x_n; y) f(y) dy \right| &\leq \|f\|_{L^\infty(\text{supp}(f))} \int_{\text{supp}(f)} p_\Phi(t_n, x_n - y; 0) dy \\ &\leq \|f\|_{L^\infty(\text{supp}(f))} |\text{supp}(f)| p_\Phi(t_n, d; 0) \rightarrow 0. \end{aligned}$$

Now let us consider  $x_0 \in \text{supp}(f)$ . Being  $\int_{\mathbb{R}} p_{\Phi}(t, x - y; 0) dy = 1$ , we have

$$\left| \int_{-\infty}^{\varphi(0)} p_{\Phi}(t, x; y) f(y) dy - f(x_0) \right| \leq \int_{\mathbb{R}} p_{\Phi}(t, x - y; 0) |f(y) - f(x_0)| dy.$$

Being  $f$  continuous, fix  $\varepsilon > 0$  and consider  $\delta > 0$  such that if  $|x_0 - y| < \delta$  then  $|f(y) - f(x_0)| < \varepsilon$ . Thus we have

$$\begin{aligned} & \left| \int_{-\infty}^{\varphi(0)} p_{\Phi}(t_n, x_n; y) f(y) dy - f(x_0) \right| \\ & \leq \int_{|x_0 - y| < \delta} p_{\Phi}(t_n, x_n - y; 0) |f(y) - f(x_0)| dy \\ & \quad + \int_{|x_0 - y| \geq \delta} p_{\Phi}(t_n, x_n - y; 0) |f(y) - f(x_0)| dy \\ & =: I_3^n + I_4^n. \end{aligned} \tag{3.73}$$

Concerning  $I_3^n$ , we have

$$I_3^n < \varepsilon \int_{|x_0 - y| < \delta} p_{\Phi}(t_n, x_n - y; 0) dy \leq \varepsilon \int_{\mathbb{R}} p_{\Phi}(t_n, x_n - y; 0) dy = \varepsilon. \tag{3.74}$$

Concerning  $I_4^n$ , we may assume  $|x_n - x_0| \leq \frac{\delta}{2}$ . Then we have

$$\delta \leq |x_0 - y| \leq |x_n - y| + |x_n - x_0| \leq |x_n - y| + \frac{\delta}{2},$$

that is to say  $|x_n - y| \geq \frac{\delta}{2}$ . Thus it holds

$$\begin{aligned} I_4^n & \leq \int_{|x_n - y| \geq \frac{\delta}{2}} p_{\Phi}(t_n, x_n - y; 0) |f(y) - f(x_0)| dy \\ & \leq 2 \|f\|_{L^\infty(\text{supp}(f))} \int_{\frac{\delta}{2}}^{+\infty} p_{\Phi}(t_n, z; 0) dz \\ & = 2 \|f\|_{L^\infty(\text{supp}(f))} \mathbb{P} \left( X_{\Phi}(t_n) > \frac{\delta}{2} \right). \end{aligned} \tag{3.75}$$

Combining equations (3.74) and (3.75) with (3.73) we get

$$\left| \int_{-\infty}^{\varphi(0)} p_{\Phi}(t_n, x_n; y) f(y) dy - f(x_0) \right| \leq \varepsilon + 2 \|f\|_{L^\infty(\text{supp}(f))} \mathbb{P} \left( X_{\Phi}(t_n) > \frac{\delta}{2} \right).$$

Taking the limit superior as  $n \rightarrow +\infty$  we have

$$\limsup_{n \rightarrow +\infty} \left| \int_{-\infty}^{\varphi(0)} p_{\Phi}(t_n, x_n; y) f(y) dy - f(x_0) \right| \leq \varepsilon,$$

and therefore, since  $\varepsilon > 0$  is arbitrary, we have

$$\lim_{n \rightarrow +\infty} \left| \int_{-\infty}^{\varphi(0)} p_{\Phi}(t_n, x_n; y) f(y) dy - f(x_0) \right| = 0.$$

Thus we have that  $u \in C^0(\overline{\mathcal{O}})$ , that is item (1) of Definition 2.1, together with the initial condition (3.35).

We prove now Item (2) of Def. 2.1. Now let us fix any  $(t, x) \in \mathcal{O}$ . Recall that the arguments we used to prove equation (3.44) actually show that as  $t > 0$ ,  $\partial_x p_\Phi(t, x - y; 0)$  is locally bounded as  $y \in \text{supp}(f)$  and that

$$\int_0^t |\partial_x p_\Phi(t - w, x; \varphi(w))| \mathbb{P}_y(\mathcal{T} \in dw) \leq C_9$$

for some constant  $C_9 > 0$ , as  $y < \varphi(0)$ . In particular by dominated convergence theorem we conclude that

$$\partial_x u(t, x) = \int_{-\infty}^{\varphi(0)} \partial_x q(t, x; y) f(y) dy.$$

In the same spirit we obtain that

$$\partial_x^2 u(t, x) = \int_{-\infty}^{\varphi(0)} \partial_x^2 q(t, x; y) f(y) dy.$$

and also that  $\partial_x q(t, x; y)$  and  $\partial_x^2 q(t, x; y)$  are locally bounded with bounds independent of  $y \in \text{supp}(f)$ . Furthermore since we have shown that, for any fixed  $t > 0$ , the functions  $\mathcal{O}_2(t) \ni x \mapsto \partial_x q(t, x; y)$  and  $\mathcal{O}_2(t) \ni x \mapsto \partial_x^2 q(t, x; y)$  are continuous for  $x \neq y$ , a simple application of the dominated convergence theorem implies that, for fixed  $t > 0$ ,  $u(t, \cdot) \in C^2(\mathcal{O}_2(t))$ .

In the same spirit we can deal with Items (3) and (4) of Definition 2.1. We have that

$$\begin{aligned} & \int_0^t \int_{-\infty}^{\varphi(0)} q(t - \tau, x; y) f(y) dy \bar{\nu}(\tau) d\tau \\ &= \int_{-\infty}^{\varphi(0)} \int_0^t q(t - \tau, x; y) f(y) \bar{\nu}(\tau) d\tau dy \\ &= \int_{-\infty}^{\varphi(0)} \int_0^t p_\Phi(t - \tau, x; y) f(y) \bar{\nu}(\tau) d\tau dy \end{aligned} \quad (3.76)$$

$$+ \int_{-\infty}^{\varphi(0)} \int_0^t \int_0^{t-\tau} p_\Phi(t - \tau - w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) \bar{\nu}(\tau) d\tau f(y) dy. \quad (3.77)$$

by a simple application of Fubini's theorem. Taking the derivative, we should have

$$\begin{aligned} \partial_t^\Phi u(x, t) &= \partial_t \int_0^{+\infty} \int_0^t p_\Phi(t - \tau, x; y) f(y) \bar{\nu}(\tau) d\tau dy \\ &\quad + \partial_t \int_0^{+\infty} \int_0^t \int_0^{t-\tau} p_\Phi(t - \tau - w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) \bar{\nu}(\tau) d\tau f(y) dy, \end{aligned}$$

provided that the derivatives exist. To ascertain this, we show that we can move them inside the integrals. Indeed we have shown that  $|\partial_t^\Phi p_\Phi(t, x - y; 0)|$  is locally bounded with respect to  $t > 0$  and  $x - y$  and

$$\partial_t^\Phi \int_0^t p_\Phi(t - w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) = \int_0^t \partial_t^\Phi p_\Phi(t - w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) \leq C_{10},$$

for some  $C_{10} > 0$ , as  $y \in \text{supp}(f)$  that is compact. Hence we have

$$\partial_t^\Phi u(x, t) = \int_{-\infty}^{\varphi(0)} \partial_t^\Phi p_\Phi(t, x; y) f(y) dy$$

$$\begin{aligned}
& + \int_{-\infty}^{\varphi(0)} \left( \partial_t^\Phi \int_0^t p_\Phi(t-w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) \right) f(y) dy \\
& = \int_{-\infty}^{\varphi(0)} \partial_t^\Phi q(t, x; y) f(y) dy.
\end{aligned}$$

This obviously implies point (3) of Definition 2.1. Moreover, as it was for the second derivative, being  $\partial_t^\Phi q(t, x; y)$  locally bounded as  $(t, x) \in \mathcal{O}$  and, for fixed  $x \in \mathcal{O}_2$ , continuous in  $\mathcal{O}_1(x)$  for almost all  $y \in (-\infty, \varphi(0))$ , a simple application of the dominated convergence theorem then yields  $\partial_t^\Phi u(\cdot, x) \in C^0(\mathcal{O}_1(x))$  for any fixed  $x \in \mathcal{O}_2$ .

Item (5) of Definition 2.1 therefore follows since

$$\begin{aligned}
\partial_t^\Phi u(x, t) & = \int_{-\infty}^{\varphi(0)} \partial_t^\Phi q(t, x; y) f(y) dy \\
& = \frac{1}{2} \int_{-\infty}^{\varphi(0)} \partial_x^2 q(t, x; y) f(y) dy = \frac{1}{2} \partial_x^2 u(x, t).
\end{aligned}$$

Concerning the boundary and the limit conditions, they easily follow from the respective ones of  $q$ . Therefore, uniqueness follows from the weak maximum principle, i.e., Corollary 2.8.

Now we show uniqueness of  $q(t, x; y)$  for almost all  $y$ . To do this, suppose that  $q_1(t, x; y)$  is another fundamental solution of Equation (3.33). Define, for any  $\psi \in C_c^\infty(-\infty, \varphi(0))$

$$u^\psi(x, t) := \int_{-\infty}^{\varphi(0)} q(t, x; y) \psi(y) dy \quad u_1^\psi(x, t) := \int_{-\infty}^{\varphi(0)} q_1(t, x; y) \psi(y) dy.$$

By uniqueness of the solution for compactly supported continuous initial data we have  $u^\psi(x, t) = u_1^\psi(x, t)$  and thus

$$\int_{-\infty}^{\varphi(0)} (q(t, x; y) - q_1(t, x; y)) \psi(y) dy = 0. \quad (3.78)$$

Since  $\psi(y)$  is arbitrary, we conclude that  $q(t, x; y) = q_1(t, x; y)$  for almost any  $y \in (-\infty, \varphi(0))$  by Fundamental Lemma of Calculus of Variations (e.g. [17, Theorem 1.24]).  $\square$

We remark that the fundamental solution is unique for all  $y \in (-\infty, \varphi(0))$  under the further assumptions **A2(c)** and **A2(d)**. This is made rigorous in the forthcoming result.

**Theorem 3.12.** *Assume **A1**, **A2(a)** – **A2(d)**, **A3**, **A4** and suppose further that  $\limsup_{\lambda \rightarrow \infty} \frac{\Phi(2\lambda)}{\Phi(\lambda)} < 2$ . Then we have that, for any fixed  $(t, x) \in \mathcal{O}$ , the function*

$$(-\infty, \varphi(0)) \ni y \mapsto q(t, x; y) \quad (3.79)$$

*is continuous.*

*Proof.* Using (3.9), it is clear that the continuity of  $q(t, x; \cdot)$  only depends on the continuity of

$$(-\infty, \varphi(0)) \ni y \mapsto \int_0^t p_\Phi(t-w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw). \quad (3.80)$$

In order to have continuity of (3.80), by Helly's Theorem (e.g., [30, Theorem 4, page 370]) it will be sufficient to prove that  $\mathbb{P}_{y_n}(\mathcal{T} \leq t) \rightarrow \mathbb{P}_y(\mathcal{T} \leq t)$  for every  $t$  as  $y_n \rightarrow y$  in  $(-\infty, \varphi(0))$  (we recall that  $t \mapsto \mathbb{P}_y(\mathcal{T} \leq t)$  is continuous by (3.7)). Note that, by space homogeneity of Brownian motion (and therefore of  $X_\Phi(t)$ ) and paths' continuity, we have that

$$\mathbb{P}_y(\mathcal{T} \leq t) = \mathbb{P}(\mathcal{T}^y \leq t) \quad (3.81)$$

where

$$\mathcal{T}^y := \inf \{t \geq 0 : y + X_\Phi(t) - \varphi(t) > 0\}. \quad (3.82)$$

Therefore we have to prove that

$$\mathbb{P}(\mathcal{T}^{y_n} \leq t) \rightarrow \mathbb{P}(\mathcal{T}^y \leq t) \quad (3.83)$$

whenever  $y_n \rightarrow y$ . For this we resort to [66, Theorem 13.6.4], as follows. Let  $D \equiv D[0, \infty)$  be the space of real valued right-continuous functions with finite left-limit and let  $E \equiv E[0, \infty) \subset D$  the set containing elements  $d \in D$  such that  $\lim_{t \rightarrow +\infty} \sup_{0 \leq s \leq t} d(s) = \infty$ . Let us first observe that for any  $n \in \mathbb{N}$  it holds  $y_n + X_\Phi(\cdot) - \varphi(\cdot) \in E$  almost surely. Indeed, Assumption **A2(c)** guarantees that  $\varphi(t) \leq l$  for some constant  $l < \infty$  while, being  $\lim_{t \rightarrow +\infty} L(t) = +\infty$ , it holds  $\limsup_{t \rightarrow \infty} X_\Phi(t) = +\infty$  almost surely. Setting  $C \equiv C^0(\mathbb{R}_0^+) \subset D$ , it is also clear that  $y + X_\Phi(\cdot) - \varphi(\cdot) \in C$  almost surely and that  $y_n + X_\Phi(\cdot) - \varphi(\cdot) \rightarrow y + X_\Phi(\cdot) - \varphi(\cdot)$  almost surely in the uniform topology of  $C$ , thus implying that the convergence also holds in  $E$  with Skorokhod's  $M$  topology. To use [66, Theorem 13.6.4], let us observe that  $\mathcal{T}^{y_n}$  are the first passage times of the process  $y_n + X_\Phi(t) - \varphi(t)$  through 0, thus (3.83) follows if we prove that for any  $\varepsilon > 0$  the process  $y + X_\Phi(t) - \varphi(t)$  does not assume the value 0 in the interval  $(\mathcal{T}^y - \varepsilon, \mathcal{T}^y)$  almost surely. This is implied by the fact that the random variable  $\max_{0 \leq s \leq t} y + X_\Phi(s) - \varphi(s)$  does not admit any atom at 0 for any  $t > 0$ . Indeed, if for  $\omega \in \Omega$  there exists  $t < \mathcal{T}^y$  such that  $y + X_\Phi(t, \omega) - \varphi(t) = 0$ , then  $\max_{0 \leq s \leq z} y + X_\Phi(s, \omega) - \varphi(s) = 0$  for any  $t \leq z < \mathcal{T}^y$  and, in particular, we can suppose  $z \in \mathbb{Q}$ . Thus, if  $\max_{0 \leq s \leq t} y + X_\Phi(s) - \varphi(s)$  does not admit any atom at 0 for any  $t > 0$ , we have

$$\begin{aligned} & \mathbb{P}(\exists t < \mathcal{T}^y : y + X_\Phi(t) - \varphi(t) = 0) \\ & \leq \mathbb{P}(\exists z < \mathcal{T}^y, z \in \mathbb{Q} : \max_{0 \leq s \leq z} y + X_\Phi(s) - \varphi(s) = 0) \\ & \leq \mathbb{P}(\exists z \in \mathbb{Q} : \max_{0 \leq s \leq z} y + X_\Phi(s) - \varphi(s) = 0) \\ & \leq \sum_{z \in \mathbb{Q}} \mathbb{P}(\max_{0 \leq s \leq z} y + X_\Phi(s) - \varphi(s) = 0) = 0. \end{aligned}$$

The fact that the summands above are zero can be ascertained as follows. Denote  $Y(t) := y + X_\Phi(t)$ . Then

$$\begin{aligned} & \mathbb{P}\left(\max_{0 \leq w \leq t} (Y(w) - \varphi(w)) = 0\right) \\ & \leq \mathbb{P}(Y(w) - \varphi(w) \leq 0, \forall w \in [\mathcal{T}^y, t], \mathcal{T}^y \leq t) \\ & = \int_0^t \mathbb{P}(Y(w) - \varphi(w) \leq 0, \forall w \in [z, t] \mid \mathcal{T}^y = z) \mathbb{P}(\mathcal{T}^y \in dz). \quad (3.84) \end{aligned}$$

Since the boundary is increasing, we can apply the Markov property at the renewal time  $\mathcal{T}^y = z$  as in Lemma 3.1 to say that

$$\begin{aligned} & \mathbb{P}(Y(w) - \varphi(w) \leq 0, \forall w \in [z, t] \mid \mathcal{T}^y = z) \\ &= \mathbb{P}_{\varphi(z)-y}(Y(w) - \varphi(w+z) \leq 0, \forall w \in [0, t-z]) \\ &\leq \mathbb{P}_{\varphi(z)-y}(\exists \bar{\epsilon} > 0 : \forall t \in [0, \bar{\epsilon}], Y(t) - \varphi(z+t) \leq 0). \end{aligned} \quad (3.85)$$

Use now **A2(c)** and **A2(d)** to say that

$$\int_0^{+\infty} (\varphi(z+t) - \varphi(z)) \nu(dt) < \infty. \quad (3.86)$$

A simple restatement of [10, Proposition III.10] tells us that

$$\lim_{t \rightarrow 0} \frac{L(t)}{\varphi(z+t) - \varphi(z)} = +\infty, \text{ a.s.}, \quad (3.87)$$

from which it follows that, for almost all  $\omega \in \Omega$ , there exists  $\bar{\epsilon}_0(\omega)$  such that  $\varphi(t+z) - \varphi(z) \leq L(t, \omega)$ , for all  $t \in [0, \bar{\epsilon}_0(\omega)]$ , that implies that, a.s.,  $Y(t, \omega) - \varphi(t+z) \geq Y(t, \omega) - \varphi(z) - L(t, \omega)$ , for all  $t \in [0, \bar{\epsilon}_0(\omega)]$ . Therefore, using (3.85), we have that

$$\begin{aligned} & \mathbb{P}(Y(w) - \varphi(w) \leq 0, \forall w \in [z, t] \mid \mathcal{T}^y = z) \\ &\leq \mathbb{P}_{\varphi(z)-y}(\exists \bar{\epsilon} < \bar{\epsilon}_0 : \forall t \in [0, \bar{\epsilon}], Y(t) - \varphi(z+t) \leq 0) \\ &\leq \mathbb{P}_{\varphi(z)-y}(\exists \bar{\epsilon} < \bar{\epsilon}_0 : \forall t \in [0, \bar{\epsilon}], Y(t) - L(t) - \varphi(z) \leq 0) \\ &= \mathbb{P}_{\varphi(z)}(\exists \bar{\epsilon} < \bar{\epsilon}_0 : \forall t \in [0, \bar{\epsilon}], B(L(t)) - L(t) \leq \varphi(z)). \end{aligned} \quad (3.88)$$

The fact that (3.88) is zero can be ascertained by observing that  $B(L(t)) - L(t)$  is the (independent) time-change of a Brownian motion with drift  $B(s) - s$ . The corresponding event for the drifted Brownian motion is clearly zero (e.g., [45, Theorem 1.38]). Moreover, by the continuity of the paths of  $L(t)$ ,

$$\{B(L(t, \omega), \omega) - L(t, \omega) : t \in [0, \epsilon_0(\omega)]\} = \{B(t, \omega) - t : t \in [0, L(\epsilon_0(\omega), \omega)]\}. \quad (3.89)$$

Since 0 is a renewal point and by definition of  $\epsilon_0$ , it follows that  $L(\epsilon_0) > 0$ , a.s. and thus the result follows.  $\square$

**Remark 3.13.** Note that **A2(c)** is not crucial for Theorem 3.12. In practice we only need that, a.s.,

$$\limsup_{t \rightarrow +\infty} X_{\Phi}(t) - \varphi(t) = \infty \quad (3.90)$$

and that (3.86) is true. A possible way to determine a boundary  $t \mapsto \varphi(t)$  which is not bounded but so that (3.90) is true is to use the law of the iterated logarithm for  $X_{\Phi}(t)$ , i.e., [36, Theorem 4].

Let us also observe that the condition  $\limsup_{\lambda \rightarrow \infty} \frac{\Phi(2\lambda)}{\Phi(\lambda)} < 2$  is, for instance, implied by asking that  $\Phi$  is regularly varying at infinity of order  $\alpha \in [0, 1)$ . Furthermore, it is also equivalent to  $\limsup_{\lambda \rightarrow \infty} \frac{\lambda \Phi'(\lambda)}{\Phi(\lambda)} < 1$  (see [10, Exercise III.6]).

Finally, let us recall that the hypothesis  $\limsup_{\lambda \rightarrow \infty} \frac{\Phi(2\lambda)}{\Phi(\lambda)} < 2$  is not needed if, additionally,  $\frac{\varphi(z+t) - \varphi(z)}{t}$  is decreasing for any  $z > 0$ . Indeed, in such case we refer to a restatement of [10, Theorem III.9]. Such a restatement is provided in [11, Proposition 4.4].

Theorem 3.12 can be actually improved, as we will show that the function  $q(t, x; y)$  is continuous as a function of three variables. For this we need the following technical statement, whose proof is left in Appendix A.4

**Proposition 3.14.** *Let  $F_n, F : \mathbb{R}_0^+ \mapsto \mathbb{R}$  for any  $n \in \mathbb{N}$  such that  $F_n(t) \rightarrow F(t)$  as  $n \rightarrow \infty$ , for any  $t \in \mathbb{R}_0^+$ . Suppose further that  $F_n, F$  are continuous, non-decreasing and  $\lim_{t \rightarrow \infty} F_n(t) = \lim_{t \rightarrow \infty} F(t) = l \in \mathbb{R}$ . Then  $F_n \rightarrow F$  uniformly.*

Now we can prove the aforementioned property.

**Theorem 3.15.** *Assume A1, A2(a) – A2(d), A3, A4 and suppose further that  $\limsup_{\lambda \rightarrow \infty} \frac{\Phi(2\lambda)}{\Phi(\lambda)} < 2$ . Then the function*

$$\mathcal{O} \times \partial_0 \mathcal{O} \ni (t, x; y) \mapsto q(t, x; y) \quad (3.91)$$

is continuous.

*Proof.* Fix  $(t, x; y) \in \mathcal{O} \times \partial_0 \mathcal{O}$  and an arbitrary sequence  $(t_n, x_n; y_n) \rightarrow (t, x; y)$  in  $\mathcal{O} \times \partial_0 \mathcal{O}$ . Observe that by the integral representation of  $q(t, x; y)$  given in (3.9) we only have to study continuity of the second summand since  $(t, x; y) \mapsto p_\Phi(t, x; y)$  is clearly continuous. Therefore, considering the second summand, we obtain

$$\begin{aligned} & \left| \int_0^{t_n} p_\Phi(t_n - w, x_n; \varphi(w)) \mathbb{P}_{y_n}(\mathcal{T} \in dw) - \int_0^t p_\Phi(t - w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) \right| \\ & \leq \left| \int_0^{t_n} p_\Phi(t_n - w, x_n; \varphi(w)) \mathbb{P}_{y_n}(\mathcal{T} \in dw) - \int_0^t p_\Phi(t - w, x; \varphi(w)) \mathbb{P}_{y_n}(\mathcal{T} \in dw) \right| \\ & \quad + \left| \int_0^t p_\Phi(t - w, x; \varphi(w)) \mathbb{P}_{y_n}(\mathcal{T} \in dw) - \int_0^t p_\Phi(t - w, x; \varphi(w)) \mathbb{P}_y(\mathcal{T} \in dw) \right|. \end{aligned} \quad (3.92)$$

The second term in (3.92) tends to zero by Theorem 3.12, hence we proceed only with the first one. Taking  $t_n \uparrow t$ , we have that

$$\begin{aligned} & \left| \int_0^{t_n} p_\Phi(t_n - w, x_n; \varphi(w)) \mathbb{P}_{y_n}(\mathcal{T} \in dw) - \int_0^t p_\Phi(t - w, x; \varphi(w)) \mathbb{P}_{y_n}(\mathcal{T} \in dw) \right| \\ & \leq \int_0^{t_n} |p_\Phi(t_n - w, x_n, \varphi(w)) - p_\Phi(t - w, x, \varphi(w))| \mathbb{P}_{y_n}(\mathcal{T} \in dw) \end{aligned} \quad (3.93)$$

$$+ \int_{t_n}^t p_\Phi(t - w, x, \varphi(w)) \mathbb{P}_{y_n}(\mathcal{T} \in dw). \quad (3.94)$$

First we deal with (3.93). We need to distinguish two cases.

Consider first the case  $x \in [\varphi(0), \varphi(t)]$ . Then, by A2(a) and A2(b), there exists a unique  $w_0$  such that  $x = \varphi(w_0)$ . Since  $w_0 \neq t$ , then there exists  $\eta > 0$  such that  $|t - w_0| > 4\eta$ . We can suppose, without loss of generality, that  $|t_n - t| < 2\eta$  and then  $|t_n - w_0| > 2\eta$ . Now let us split (3.93) as

$$\int_{|w - w_0| > \eta, w < t_n} |p_\Phi(t_n - w, x_n, \varphi(w)) - p_\Phi(t - w, x, \varphi(w))| \mathbb{P}_{y_n}(\mathcal{T} \in dw) \quad (3.95)$$

$$+ \int_{|w - w_0| \leq \eta, w < t_n} |p_\Phi(t_n - w, x_n, \varphi(w)) - p_\Phi(t - w, x, \varphi(w))| \mathbb{P}_{y_n}(\mathcal{T} \in dw). \quad (3.96)$$

Let us work out with (3.95). Observe that for  $\tau \in [t_1, t]$  and  $w \in \Theta_1 := \{w \in [0, t] : |w - w_0| \geq \eta\}$ , it holds  $\tau - w \in [0, t - w_0 - \eta] \cup [t_1 - w_0 + \eta, t]$  (note that  $t - w_0 - \eta < t_1 - w_0 + \eta$  since  $t - t_1 < 2\eta$ ). Furthermore,  $\tau - w \leq t - w_0 - \eta$  implies  $w_0 - w \leq t - \tau - \eta < \eta$ . Since  $|w - w_0| \geq \eta$ , it must hold  $w \geq w_0 + \eta$ , so that  $\varphi(w) \geq \varphi(w_0 + \eta) > x$ . Now consider  $\delta > 0$  such that  $x + 3\delta \leq \varphi(w_0 + \eta)$  and suppose that, without loss of generality,  $|x_n - x| < \delta$  for any  $n$ . Define now the compact set

$$K_1 := [t_1, t] \times \Theta_1 \times [x - \delta, x + \delta] \quad (3.97)$$

so that the function  $K_1 \ni (\tau, w, z) \mapsto p_\Phi(\tau - w, z; \varphi(w))$  is continuous (note that if  $\tau - w$  approaches zero then  $\varphi(w) \geq x + 3\delta > x + 2\delta > x + \delta \geq z$ ). Fix  $\mu > 0$ : by uniform continuity we know that there exists  $n$  big enough such that

$$|p_\Phi(t_n - w, x_n, \varphi(w)) - p_\Phi(t - w, x, \varphi(w))| < \mu, \quad (3.98)$$

for  $w \in \Theta_1$ .

Now we consider (3.96). We can suppose, without loss of generality, that  $t_1 > w_0$ . In particular this means that  $t_1 > w_0 + \eta$ . Observe that for  $|w - w_0| \leq \eta$  we have  $t_1 - w \geq \eta$ . Consider again  $\tau \in [t_1, t]$ . Then  $\tau - w \geq t_1 - w \geq \eta$ . Let  $\Theta_2 := \{w \in [0, t] : |w - w_0| \leq \eta\}$  and define the compact set

$$K_2 = [t_1, t] \times \Theta_2 \times [x - \delta, x + \delta]. \quad (3.99)$$

Since for any  $(\tau, w, z) \in K_2$  we have  $\tau - w \geq \eta > 0$ , then  $K_2 \ni (\tau, z; w) \mapsto p_\Phi(\tau - w, z; \varphi(w))$  is continuous. Hence, we can suppose that  $n$  is big enough so that (3.98) holds even if  $w \in \Theta_2$ . In conclusion we get

$$\int_0^{t_n} |p_\Phi(t_n - w, x_n, \varphi(w)) - p_\Phi(t - w, x, \varphi(w))| \mathbb{P}_{y_n}(\mathcal{T} \in dw) \leq \mu. \quad (3.100)$$

Since  $\mu > 0$  is arbitrary we conclude that

$$\int_0^{t_n} |p_\Phi(t_n - w, x_n, \varphi(w)) - p_\Phi(t - w, x, \varphi(w))| \mathbb{P}_{y_n}(\mathcal{T} \in dw) \rightarrow 0. \quad (3.101)$$

If, instead,  $x < \varphi(0)$  then we consider directly the compact set

$$K_3 = [t_1, t] \times [0, t] \times [x - \delta, x + \delta], \quad (3.102)$$

where now  $\delta > 0$  is chosen so that  $x + \delta < \varphi(0)$  and we argue as before.

Now we deal with (3.94). Without loss of generality, we can suppose  $x < \varphi(w)$  for any  $w \in [t_n, t]$ . Then  $p_\Phi(t - w, x; \varphi(w)) \leq C(t, x)$  so that,

$$\int_{t_n}^t p_\Phi(t - w, x, \varphi(w)) \mathbb{P}_{y_n}(\mathcal{T} \in dw) \leq C(t, x) (\mathbb{P}_{y_n}(\mathcal{T} \leq t) - \mathbb{P}_{y_n}(\mathcal{T} \leq t_n)). \quad (3.103)$$

Now observe that we have shown in the previous theorem that  $\mathbb{P}_{y_n}(\mathcal{T} \leq t) \rightarrow \mathbb{P}_y(\mathcal{T} \leq t)$ . However, the functions  $t \mapsto \mathbb{P}_y(\mathcal{T} \leq t)$ ,  $t \mapsto \mathbb{P}_{y_n}(\mathcal{T} \leq t)$  are continuous, monotone, bounded and they have the same limit. Hence  $\mathbb{P}_{y_n}(\mathcal{T} \leq t) \rightarrow \mathbb{P}_y(\mathcal{T} \leq t)$  uniformly in  $t$  as it follows from Proposition 3.14. Thus we achieve that (3.103) tends to zero as  $n \rightarrow \infty$ . The same arguments apply to the case  $t_n \downarrow t$ , concluding the proof.  $\square$

As a consequence of the previous result we get uniqueness of a continuous fundamental solution.

**Corollary 3.16.** *Under the hypotheses of Theorem 3.15, the function  $(t, x; y) \mapsto q(t, x; y)$  is the unique fundamental solution of the non-local problem introduced in Theorem 3.11 which is continuous as a function of  $y \in \partial_0 \mathcal{O}$  for fixed  $(t, x) \in \mathcal{O}$ .*

*Proof.* Let  $q_1(t, x; y)$  be another fundamental solution continuous in  $y$  for fixed  $(t, x)$ . Fix  $(t, x) \in \mathcal{O}$  arbitrary. Use (3.78) to conclude that  $q(t, x; y) = q_1(t, x; y)$  a.e. By continuity we conclude the proof.  $\square$

**Remark 3.17.** *If, for fixed  $(t, x) \in \mathcal{O}$ ,  $y \mapsto q_1(t, x; y)$  is not continuous,  $q_1(t, x; y) = q(t, x; y)$  in any continuity point of  $q_1$ .*

#### 4. ANOMALOUS DIFFUSIVE BEHAVIOUR

In this section we obtain an estimate of the asymptotic behaviour of the mean square displacement of the (fundamental) solution of the non-local problem, i.e., we estimate the quantity

$$(\overline{\Delta x})^2(t) := \int_{-\infty}^{+\infty} x^2 \mathbb{P}_0 \left( X_{\Phi}^{\dagger}(t) \in dx \right). \quad (4.1)$$

The asymptotic behaviour of time-changed processes related with non-local equations is crucial for applications in the context of anomalous diffusion, see the recent contribution [29] and references therein. It turns out that (4.1) can be expressed by means of the potential measure of the corresponding subordinator, i.e.,

$$U(t) := \mathbb{E} \int_0^{+\infty} \mathbb{1}_{\{\sigma(s) \leq t\}} ds. \quad (4.2)$$

In our framework this measure admits the density

$$u_{\Phi}(t) := \int_0^{+\infty} g_{\sigma}(t, s) ds, \quad (4.3)$$

where  $t \mapsto g_{\sigma}(t, s)$  is the density of the subordinator  $\sigma(s)$ . To ascertain this, note that the density  $g_{\sigma}$  exists by **A3** and therefore, by applying Tonelli repeatedly, we obtain for any  $t > 0$

$$\begin{aligned} U(t) &= \int_0^{+\infty} \mathbb{P}(\sigma(s) \leq t) ds \\ &= \int_0^t \int_0^{+\infty} g_{\sigma}(w, s) ds dw \\ &= \int_0^t u_{\Phi}(w) dw. \end{aligned} \quad (4.4)$$

From the representation above it is easy to see that

$$\int_0^{+\infty} e^{-\lambda t} u_{\Phi}(t) dt = \frac{1}{\Phi(\lambda)}, \quad (4.5)$$

and this will be used below. To obtain the result we will make use of the reflection principle for the time-changed Brownian motion that we prove in the following proposition.

**Proposition 4.1.** *We have that, for any  $c > 0$ ,*

$$\mathbb{P} \left( \sup_{0 \leq s \leq t} X_{\Phi}(s) > c \right) = 2 \mathbb{P} (X_{\Phi}(t) > c). \quad (4.6)$$

*Proof.* By conditioning on  $L(t)$  and using independence we have that

$$\begin{aligned}
\mathbb{P}\left(\sup_{0 \leq s \leq t} X_\Phi(s) > c\right) &= \mathbb{P}\left(\sup_{0 \leq w \leq L(t)} B(w) > c\right) \\
&= \mathbb{E} \mathbb{P}\left(\sup_{0 \leq w \leq L(t)} B(w) > c \mid L(t)\right) \\
&= \int_0^\infty \mathbb{P}\left(\sup_{0 \leq w \leq s} B(w) > c\right) f_L(s, t) ds \\
&= 2 \int_0^\infty \mathbb{P}(B(s) > c) f_L(s, t) ds \\
&= 2 \mathbb{P}(X_\Phi(t) > c). \tag{4.7}
\end{aligned}$$

□

The asymptotic behaviour of the mean square displacement can be now obtained using the reflection principle. We have the following result.

**Theorem 4.2.** *Assume **A1** and **A2(a) – A2(d)** and suppose further that  $\Phi(\lambda)$  varies regularly at  $0+$  with index  $\gamma \in (0, 1]$ . Then*

$$\frac{\gamma}{1+\gamma} \frac{1}{\Gamma(1+\gamma)} \leq \liminf_{t \rightarrow \infty} \Phi(1/t) (\Delta x)^2(t) \leq \limsup_{t \rightarrow \infty} \Phi(1/t) (\Delta x)^2(t) \leq \frac{1}{\Gamma(1+\gamma)}. \tag{4.8}$$

*Proof.* By Lemma 3.1 we obtain

$$\begin{aligned}
(\overline{\Delta x})^2(t) &= \int_{-\infty}^{+\infty} x^2 q(t, x; 0) dx \\
&= \mathbb{E}[X_\phi(t)]^2 - \int_0^t \mathbb{E}^{\varphi(w)} [X_\Phi(t-w)]^2 \mathbb{P}(\mathcal{T} \in dw) \\
&= \mathbb{E}[X_\phi(t)]^2 - \int_0^t \mathbb{E}[\varphi(w) + X_\Phi(t-w)]^2 \mathbb{P}(\mathcal{T} \in dw), \tag{4.9}
\end{aligned}$$

where the second last step is justified by Tonelli's theorem while the last step by space homogeneity. Let us first show the upper bound. Note that

$$\begin{aligned}
\mathbb{E}[X_\Phi(t)]^2 &= \int_0^{+\infty} \int_{-\infty}^{+\infty} x^2 \frac{1}{\sqrt{2\pi s}} e^{-\frac{(x-y)^2}{2s}} dx f_L(s, t) ds \\
&= \mathbb{E}L(t) \\
&= \int_0^{+\infty} \mathbb{P}(L(t) > s) ds \\
&= \mathbb{E} \int_0^{+\infty} \mathbf{1}_{\{\sigma(s) \leq t\}} ds. \tag{4.10}
\end{aligned}$$

The last quantity is the potential measure of the subordinator. Hence we easily obtain the upper bound

$$(\Delta x)^2(t) \leq U(t)$$

where the asymptotic behaviour is given by [11, Proposition 1.5] as follows

$$U(t) \sim \frac{1}{\Gamma(1+\gamma)} \frac{1}{\Phi(1/t)}. \tag{4.11}$$

Concerning the lower bound it is easy to see that  $\mathbb{E}X_\Phi(t) = 0$ , therefore we obtain

$$\begin{aligned} (\overline{\Delta x})^2(t) &= U(t) - \int_0^t (\varphi^2(w) + U(t-w)) \mathbb{P}(\mathcal{T} \in dw) \\ &\geq U(t) - \int_0^t U(t-w) \mathbb{P}(\mathcal{T} \in dw) - l^2. \end{aligned}$$

By denoting

$$\mathcal{T}_0 := \inf \{s \geq 0 : X_\Phi(s) = \varphi(0)\} \quad (4.12)$$

we understand that, a.s.,  $\mathcal{T} \geq \mathcal{T}_0$ . Thus, from (4.12), it follows that

$$\begin{aligned} (\overline{\Delta x})^2(t) &\geq U(t) - \int_0^t U(t-w) \mathbb{P}(\mathcal{T} \in dw) - l^2 \\ &= U(t) - \mathbb{E}U(t-\mathcal{T})\mathbb{1}_{\{\mathcal{T} \leq t\}} - l^2 \\ &\geq U(t) - \mathbb{E}U(t-\mathcal{T}_0)\mathbb{1}_{\{\mathcal{T}_0 \leq t\}} - l^2 \\ &= U(t) - \int_0^t U(t-w) \mathbb{P}(\mathcal{T}_0 \in dw) - l^2 \end{aligned} \quad (4.13)$$

where in the second last step we used that  $t \mapsto U(t)$  is non-decreasing. Observe that, by the reflection principle,

$$\begin{aligned} \int_0^t U(t-w) \mathbb{P}(\mathcal{T}_0 \in dw) &= 2 \int_0^t U(t-w) \partial_w \mathbb{P}(X_\Phi(w) > \varphi(0)) dw \\ &= 2 \int_0^t u_\Phi(t-w) \mathbb{P}(X_\Phi(w) > \varphi(0)) dw. \end{aligned} \quad (4.14)$$

Denote now

$$\mathcal{U}(t) := \int_0^t \int_0^s U(s-w) \mathbb{P}(\mathcal{T}_0 \in dw) ds \quad (4.15)$$

and the corresponding Laplace-Stieltjes transform

$$\tilde{\mathcal{U}}(\lambda) := \int_0^{+\infty} e^{-\lambda t} d\mathcal{U}(t). \quad (4.16)$$

Let us recall that the Tauberian theorem [18, Theorem XIII.5.2] says that

$$\mathcal{U}(t) \sim \frac{t^\rho \mathcal{L}(t)}{\Gamma(1+\rho)} \text{ as } t \rightarrow +\infty \quad \iff \quad \tilde{\mathcal{U}}(\lambda) \sim \lambda^{-\rho} \mathcal{L}(1/\lambda) \text{ as } \lambda \rightarrow 0+ \quad (4.17)$$

where  $\mathcal{L}(t)$  is slowly varying at infinity and  $\rho > 0$ . Observe that, denoting with  $K_\nu(z)$  the modified Bessel function of the second kind, we have

$$\begin{aligned} &\tilde{\mathcal{U}}(\lambda) \\ &= 2 \int_0^{+\infty} e^{-\lambda t} \int_0^t u_\Phi(t-w) \mathbb{P}(X_\Phi(w) > \varphi(0)) dw dt \quad \text{by (4.14)} \\ &= 2 \frac{1}{\Phi(\lambda)} \int_0^{+\infty} e^{-\lambda w} \int_{\varphi(0)}^\infty \int_0^{+\infty} \frac{e^{-\frac{x^2}{2s}}}{\sqrt{2\pi s}} f_L(s, w) ds dx dw \quad \text{by [1, Proposition 1.6.4] and (4.5)} \\ &= 2 \frac{1}{\Phi(\lambda)} \frac{\Phi(\lambda)}{\lambda} \int_{\varphi(0)}^\infty \int_0^{+\infty} e^{-s\Phi(\lambda)} \frac{e^{-\frac{x^2}{2s}}}{\sqrt{2\pi s}} ds dx \quad \text{by Tonelli's theorem and [38, eq. (3.13)]} \end{aligned}$$

$$\begin{aligned}
&= \frac{4}{\lambda} \frac{1}{\sqrt{2\pi}} \int_{\varphi(0)}^{\infty} \left( \frac{x^2}{2\Phi(\lambda)} \right)^{1/4} K_{1/2} \left( 2\sqrt{\frac{x^2}{2}\Phi(\lambda)} \right) dx && \text{by [22, formula 3.471.9]} \\
&= \frac{2}{\sqrt{2\lambda}} \frac{1}{\sqrt{\Phi(\lambda)}} \int_{\varphi(0)}^{\infty} e^{-\sqrt{2}x\sqrt{\Phi(\lambda)}} dx && \text{by [22, formula 8.469.3]} \\
&= \frac{1}{\lambda} \frac{1}{\Phi(\lambda)} e^{-\sqrt{2}\varphi(0)\sqrt{\Phi(\lambda)}} \\
&\sim \frac{1}{\lambda^{1+\gamma}} \frac{1}{\mathcal{L}(1/\lambda)} \quad \text{as } \lambda \rightarrow 0+, && (4.18)
\end{aligned}$$

where the asymptotic behaviour follows since  $\Phi(\lambda)$  varies regularly at  $0+$ , i.e.  $\Phi(\lambda) = \lambda^\gamma \mathcal{L}(1/\lambda)$ , with index  $\gamma \in (0, 1]$ . Since  $\mathcal{L}(\lambda)$  is slowly varying at infinity, then  $1/\mathcal{L}(\lambda)$  is slowly varying at infinity and thus it follows by (4.17) that, as  $t \rightarrow +\infty$ ,

$$U(t) \sim \frac{1}{\Gamma(2+\gamma)} \frac{t}{t^{-\gamma} \mathcal{L}(t)} = \frac{1}{\Gamma(2+\gamma)} \frac{t}{\Phi(1/t)} \quad (4.19)$$

so that, by the monotone density theorem [12, Theorem 1.7.2]

$$\int_0^t U(t-w) \mathbb{P}(\mathcal{T}_0 \in dw) \sim \frac{1}{\Gamma(2+\gamma)} \frac{1}{\Phi(1/t)} \text{ as } t \rightarrow +\infty. \quad (4.20)$$

The monotone density theorem can be applied since the function

$$t \mapsto \int_0^t U(t-w) \mathbb{P}(\mathcal{T} \in dw) \quad (4.21)$$

is non-decreasing, as can be ascertained by observing that for arbitrary  $h, t > 0$ ,

$$\begin{aligned}
&\int_0^{t+h} U(t+h-w) \mathbb{P}(\mathcal{T} \in dw) - \int_0^t U(t-w) \mathbb{P}(\mathcal{T} \in dw) \\
&= \int_0^t (U(t+h-w) - U(t-w)) \mathbb{P}(\mathcal{T} \in dw) + \int_t^{t+h} U(t+h-w) \mathbb{P}(\mathcal{T} \in dw)
\end{aligned} \quad (4.22)$$

which is positive since  $t \mapsto U(t)$  is non-decreasing. Putting pieces together we have, for the lower bound, that, as  $t \rightarrow +\infty$ ,

$$U(t) - \int_0^t U(t-w) \mathbb{P}(\mathcal{T}_0 \in dw) - t^2 \sim \left(1 - \frac{1}{1+\gamma}\right) \frac{1}{\Gamma(1+\gamma)} \frac{1}{\Phi(1/t)} \quad (4.23)$$

and this yields the desired result.  $\square$

## 5. EXAMPLES

In this section we provide some examples of our theory where some computations can be performed explicitly. We discuss, in particular, some specific kernels and subordinators satisfying the assumptions. For this, one has to specify the Bernstein function from which the kernel  $\bar{\nu}(t)$  appearing in the operator  $\partial_t^\Phi$  and the related subordinator  $\sigma(t)$  follow.

**Example 5.1** (The fractional problem and the  $\beta$ -stable subordinator). *The prototype of the whole theory is given by the Bernstein function  $\Phi(\lambda) = \lambda^\beta$  for  $\beta \in (0, 1)$ .*

Then the corresponding Lévy measure has a Lebesgue density given by

$$\nu(s) = \frac{\beta s^{-\beta-1}}{\Gamma(1-\beta)} \quad (5.1)$$

so that

$$\bar{\nu}(s) = \frac{s^{-\beta}}{\Gamma(1-\beta)}. \quad (5.2)$$

It is clear that, in this framework, **A1** and **A3** are verified. The corresponding operator  $\partial_t^\Phi$ , obtained using the kernel (5.2), becomes a  $\beta$ -fractional derivative. The subordinator corresponding to  $\Phi(\lambda) = \lambda^\beta$  is called a  $\beta$ -stable subordinator, say  $\sigma_\beta(t)$ , and its inverse is called the inverse  $\beta$ -stable subordinator  $L_\beta(t)$ . It turns out that the assumptions **A4** is also satisfied. Indeed, it is known (see [41]) that, denoting by  $g_\beta(s)$  a density of the random variable  $\sigma_\beta(1)$  and  $f_\beta(s; t)$  a density of  $L_\beta(t)$ ,

$$f_\beta(s; t) = \frac{t}{\beta} s^{-1-\frac{1}{\beta}} g_\beta(ts^{-\frac{1}{\beta}}). \quad (5.3)$$

Using (5.3) we can check **A4**. Let us recall that the function  $g_\beta(x)$  admits the following series representation (see [52, Eq. (7)])

$$g_\beta(x) = \frac{1}{\pi} \sum_{j=1}^{+\infty} (-1)^{j+1} \frac{\Gamma(1+j\beta)}{j!} \sin(\pi\beta j) x^{-1-\beta j}. \quad (5.4)$$

This leads to the power series (with respect to  $s$ ) representation

$$\begin{aligned} f_\beta(s; t) &= \sum_{j=1}^{\infty} (-1)^{j+1} \frac{\Gamma(1+j\beta)}{\beta j!} \frac{\sin(\pi\beta j)}{\pi} t^{-\beta j} s^{j-1} \\ &= \sum_{j=0}^{\infty} (-1)^j \frac{\Gamma(1+(j+1)\beta)}{\beta(j+1)!} \frac{\sin(\pi\beta(j+1))}{\pi} t^{-\beta(j+1)} s^j \end{aligned} \quad (5.5)$$

which is uniformly convergent for  $s$  near zero and can be differentiated term by term, leading to

$$\begin{aligned} \partial_s f_\beta(s, t) &= \sum_{j=1}^{\infty} (-1)^j j \frac{\Gamma(1+(j+1)\beta)}{\beta(j+1)!} \frac{\sin(\pi\beta(j+1))}{\pi} t^{-\beta(j+1)} s^{j-1} \\ &= \sum_{j=0}^{\infty} (-1)^{j+1} (j+1) \frac{\Gamma(1+(j+2)\beta)}{\beta(j+2)!} \frac{\sin(\pi\beta(j+2))}{\pi} t^{-\beta(j+2)} s^j \\ &= \sum_{j=1}^{\infty} (-1)^{j+1} (j+1) \frac{\Gamma(1+(j+2)\beta)}{\beta(j+2)!} \frac{\sin(\pi\beta(j+2))}{\pi} t^{-\beta(j+2)} s^j \\ &\quad - \frac{\Gamma(1+2\beta)}{\beta} \frac{\sin(2\pi\beta)}{\pi} t^{-2\beta}. \end{aligned} \quad (5.6)$$

This proves **A4**. Note that, if  $\beta \neq 1/2$ , using Euler's reflection formula for the Gamma function, we obtain

$$\partial_s f_\beta(0+, t) = -\frac{\Gamma(1+2\beta)}{\beta\Gamma(2\beta)\Gamma(1-2\beta)} t^{-2\beta} = -\frac{2}{\Gamma(1-2\beta)} t^{-2\beta}, \quad (5.7)$$

while for  $\beta = 1/2$ , it is easy to see that  $\partial_s f_\beta(0+, t) = 0$ .

Observe further that  $\lambda^\beta$  varies regularly at  $0+$  with index  $\beta$ , so that the mean square displacement behaviour determined in Theorem 4.2 is, in this case,

$$\frac{\beta}{1+\beta} \frac{1}{\Gamma(1+\beta)} \leq \liminf_{t \rightarrow \infty} \frac{(\Delta x)^2(t)}{t^\beta} \leq \limsup_{t \rightarrow \infty} \frac{(\Delta x)^2(t)}{t^\beta} \leq \frac{1}{\Gamma(1+\beta)}. \quad (5.8)$$

**Example 5.2** (The relativistic stable subordinator). Suppose that the Bernstein function is, for  $\beta \in (0, 1)$ ,  $m > 0$ ,

$$\Phi(\lambda) = (\lambda + m)^\beta - m^\beta \quad (5.9)$$

which corresponds to the choice of a Lévy measure with Lévy density

$$\nu(s) = \frac{\beta s^{-\beta-1}}{\Gamma(1-\beta)} e^{-ms} \quad (5.10)$$

so that

$$\bar{\nu}(s) = \frac{\beta m^\beta}{\Gamma(1-\beta)} \Gamma(-\beta, ms) \quad (5.11)$$

where  $\Gamma(\cdot, \cdot)$  denotes the upper incomplete Gamma function. The corresponding subordinator is called, in this case, the relativistic stable subordinator. In this situation it is clear that **A1** and **A3** are satisfied. Furthermore, we can use [33, Proposition 2.1] to say that the corresponding inverse subordinator has a density  $f_{\beta,m}(s, t)$  with the series representation

$$f_{\beta,m}(s, t) = \frac{e^{m^\beta s}}{\pi} \sum_{k=0}^{\infty} (-1)^k \frac{s^k m^{\beta(k+1)}}{k!} C_k(t, m, \beta) =: e^{m^\beta s} \Sigma_{\beta,m}(s, t) \quad (5.12)$$

where

$$C_k(t, m, \beta) := \Gamma(1 + \beta(k+1)) \Gamma(-\beta(k+1), mt) \sin((k+1)\beta\pi) - \Gamma(1 + \beta k) \Gamma(-\beta k, mt) \sin(k\pi\beta). \quad (5.13)$$

Let us observe that

$$\begin{aligned} \partial_s \Sigma_{\beta,m}(s, t) &= \sum_{k=1}^{\infty} (-1)^k \frac{1}{\pi} \frac{s^{k-1} m^{\beta(k+1)}}{(k-1)!} C_k(t, m, \beta) \\ &= - \sum_{k=0}^{\infty} (-1)^k \frac{1}{\pi} \frac{s^k m^{\beta(k+2)}}{k!} C_{k+1}(t, m, \beta) \end{aligned} \quad (5.14)$$

and thus

$$\partial_s f_{\beta,m}(s, t) = \sum_{k=0}^{\infty} (-1)^k e^{m^\beta s} \frac{1}{\pi} \frac{s^k m^{\beta(k+2)}}{k!} (C_k(t, m, \beta) - C_{k+1}(t, m, \beta)). \quad (5.15)$$

It follows that

$$\partial_s f_{\beta,m}(0+, t) = \begin{cases} 2\beta m^{2\beta} \left( \frac{\Gamma(-\beta, mt)}{\Gamma(1-\beta)} - \frac{\Gamma(-2\beta, mt)}{\Gamma(1-2\beta)} \right), & \beta \neq 1/2, \\ m\Gamma(-1/2, mt)/\sqrt{\pi}, & \beta = 1/2 \end{cases} \quad (5.16)$$

and then **A4** is also satisfied.

In this case the function  $(\lambda + m)^\beta - m^\beta$  varies regularly at  $0+$  with index  $\gamma = 1$ , so the asymptotic behaviour of the mean square displacement determined in Theorem 4.2 becomes

$$\begin{aligned} \frac{1}{2} &\leq \liminf_{t \rightarrow +\infty} \left( \left( \frac{1}{t} + m \right)^\beta - m^\beta \right) (\Delta x)^2(t) \\ &\leq \limsup_{t \rightarrow +\infty} \left( \left( \frac{1}{t} + m \right)^\beta - m^\beta \right) (\Delta x)^2(t) \leq 1 \end{aligned} \quad (5.17)$$

and thus

$$\frac{1}{2\beta m^\beta} \leq \liminf_{t \rightarrow +\infty} \frac{(\Delta x)^2(t)}{t} \leq \limsup_{t \rightarrow +\infty} \frac{(\Delta x)^2(t)}{t} \leq \frac{1}{\beta m^\beta}. \quad (5.18)$$

#### APPENDIX A. PROOFS OF SOME TECHNICAL RESULTS

**A.1. Proof of Proposition 3.4.** In this section we will always assume both **A1** and **A3**. Before providing the proof of Proposition 3.4, we need to prove the following technical lemmas.

**Lemma A.1.** *For any fixed  $s \in \mathbb{R}^+$ , the function  $t \mapsto f_L(s; t)$  is differentiable on  $\mathbb{R}^+$  and*

$$\partial_t f_L(s; t) = \int_0^t \bar{\nu}(\tau) \partial_t g_\sigma(t - \tau; s) d\tau. \quad (A.1)$$

Moreover, denoting  $I_\Phi(t) = \int_0^t \bar{\nu}(\tau) d\tau$ , it holds

$$|\partial_t f_L(s; t)| \leq \frac{1}{\pi} I_\Phi(t) \int_0^{+\infty} \xi e^{-s\Re(\Psi(\xi))} d\xi. \quad (A.2)$$

*Proof.* By [38, Theorem 3.1] we have that

$$f_L(s, t) = \int_0^t \bar{\nu}(\tau) g_\sigma(t - \tau, s) d\tau. \quad (A.3)$$

Consider  $h > 0$  and observe that

$$\frac{f_L(s; t+h) - f_L(s; t)}{h} = \int_0^t \bar{\nu}(\tau) \frac{g_\sigma(t+h-\tau; s) - g_\sigma(t-\tau; s)}{h} d\tau \quad (A.4)$$

$$+ \int_t^{t+h} \bar{\nu}(\tau) \frac{g_\sigma(t+h-\tau; s)}{h} d\tau \quad (A.5)$$

$$=: I_1(h) + I_2(h). \quad (A.6)$$

We want to take the limit as  $h \rightarrow 0^+$ . Concerning  $I_2(h)$ , we have

$$I_2(h) = \int_0^h \bar{\nu}(t+h-\tau) \frac{g_\sigma(\tau; s)}{h} d\tau \quad (A.7)$$

$$= \int_0^h (\bar{\nu}(t+h-\tau) - \bar{\nu}(t-\tau)) \frac{g_\sigma(\tau; s)}{h} d\tau \quad (A.8)$$

$$+ \int_0^h \bar{\nu}(t-\tau) \frac{g_\sigma(\tau; s)}{h} d\tau \quad (A.9)$$

$$=: I_3(h) + I_4(h). \quad (A.10)$$

It is clear that

$$\lim_{h \rightarrow 0^+} I_4(h) = \bar{\nu}(t) g_\sigma(0^+; s) = 0. \quad (A.11)$$

To work with  $I_3(h)$ , let us observe that, by continuity of  $\bar{\nu}$ , for any  $\varepsilon > 0$  there exists  $\delta > 0$  such that

$$|\bar{\nu}(t+h-\tau) - \bar{\nu}(t-\tau)| \leq 2\pi\varepsilon, \quad \forall h \in (0, \delta). \quad (\text{A.12})$$

Moreover, since

$$g_\sigma(t; s) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-i\xi t} e^{-s\Psi(\xi)} d\xi \quad (\text{A.13})$$

and, in particular,

$$|g_\sigma(t; s)| \leq \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-s\Re(\Psi(\xi))} d\xi, \quad (\text{A.14})$$

we get

$$|I_3(h)| \leq 2\varepsilon \int_0^{+\infty} e^{-s\Re(\Psi(\xi))} d\xi, \quad h \in (0, \delta), \quad (\text{A.15})$$

that implies

$$\limsup_{h \rightarrow 0^+} |I_3(h)| \leq 2\varepsilon \int_0^{+\infty} e^{-s\Re(\Psi(\xi))} d\xi. \quad (\text{A.16})$$

Sending  $\varepsilon \rightarrow 0^+$  we obtain  $\lim_{h \rightarrow 0^+} I_3(h) = 0$ .

Finally let us consider  $I_1(h)$ , whose integrand is dominated via the inequality

$$\bar{\nu}(\tau) \frac{|g_\sigma(t+h-\tau; s) - g_\sigma(t-\tau; s)|}{h} \leq \frac{1}{\pi} \bar{\nu}(\tau) \int_0^{+\infty} \xi e^{-s\Re(\Psi(\xi))} d\xi \quad (\text{A.17})$$

that is integrable on  $(0, t)$ . Hence we can take the limit inside the integral sign to get

$$\lim_{h \rightarrow 0^+} \frac{f_L(s; t+h) - f_L(s; t)}{h} = \int_0^t \bar{\nu}(\tau) \partial_t g_\sigma(t-\tau; s) d\tau. \quad (\text{A.18})$$

The argument for  $h < 0$  is exactly the same. Finally, the upper bound follows from (A.17).  $\square$

**Lemma A.2.** *Let  $x \neq y$  and  $z > 0$ . Then it holds*

$$\int_0^{+\infty} s^{-\frac{1}{2}} e^{-\frac{(x-y)^2}{2s} - zs} ds = \sqrt{\frac{\pi}{z}} e^{-|x-y|\sqrt{2z}}. \quad (\text{A.19})$$

*Proof.* First of all let us rewrite the integral as

$$\begin{aligned} & \int_0^{+\infty} s^{-\frac{1}{2}} e^{-\frac{(x-y)^2}{2s} - zs} ds \\ &= 2 \left( \frac{|x-y|}{\sqrt{2z}} \right)^{\frac{1}{2}} \frac{1}{2} \left( \frac{|x-y|}{\sqrt{2z}} \right)^{-\frac{1}{2}} \int_0^{+\infty} s^{-(-\frac{1}{2})-1} e^{-\frac{2z}{2} \left( \frac{(x-y)^2}{2zs} - s \right)} ds. \end{aligned}$$

Now, by [22, Formula 3.471.9], we obtain

$$\int_0^{+\infty} s^{-\frac{1}{2}} e^{-\frac{(x-y)^2}{2s} - zs} ds = 2 \left( \frac{|x-y|}{\sqrt{2z}} \right)^{\frac{1}{2}} K_{-\frac{1}{2}}(\sqrt{2z}|x-y|), \quad (\text{A.20})$$

where  $K_\nu(z)$  is the modified Bessel function of the third kind (see [22]). In particular, it holds, by [22, Formula 8.469.3]

$$K_{-\frac{1}{2}}(\sqrt{2z}|x-y|) = \sqrt{\frac{\pi}{2\sqrt{2z}|x-y|}} e^{-|x-y|\sqrt{2z}}, \quad (\text{A.21})$$

concluding the proof.  $\square$

**Lemma A.3.** *There exist  $\delta_0 > 0$  and  $C > 0$  such that*

$$\Re(\Psi(\xi)) \geq C\xi^2, \quad 0 < \xi < \delta_0. \quad (\text{A.22})$$

*Proof of Lemma A.3.* Let us recall, by [58, Proposition 3.5], that  $\Phi$  admits a unique continuous extension  $\Phi : \{z \in \mathbb{C} : \Re(z) \geq 0\} \rightarrow \{z \in \mathbb{C} : \Re(z) \geq 0\}$ . Thus, in particular, we have that  $\Re(\Psi(\xi)) \geq 0$  for any  $\xi \in \mathbb{R}$ . Moreover, if  $\xi \neq 0$ , let us observe that

$$\Re(\Psi(\xi)) = \int_0^{+\infty} (1 - \cos(\xi t))\nu(t)dt > 0. \quad (\text{A.23})$$

Recalling that there exists  $\delta_0 > 0$  such that, for any  $x < \delta_0$ , it holds  $1 - \cos x \geq x^2/4$ , we get

$$\Re(\Psi(\xi)) \geq \int_0^{\frac{\delta_0}{\xi}} \frac{\xi^2 t^2}{4} \nu(dt) \geq \frac{\xi^2}{4} \int_0^1 t^2 \nu(dt), \quad 0 < \xi < \delta_0. \quad (\text{A.24})$$

□

Now we are ready to prove the aforementioned Proposition.

*Proof of Proposition 3.4.* (1) We resort to Lemma A.1. Let us consider the integrand in Equation (3.17). By Equation (A.2) we have that

$$\frac{e^{-\frac{(x-y)^2}{2s}}}{\sqrt{2\pi s}} |\partial_t f_L(s; t)| \leq \frac{e^{-\frac{(x-y)^2}{2s}}}{\pi\sqrt{2\pi s}} I_\Phi(t) \int_0^{+\infty} \xi e^{-s\Re(\Psi(\xi))} d\xi. \quad (\text{A.25})$$

Since  $I_\Phi(t)$  is locally bounded on  $(0, +\infty)$ , we only have to show that  $G(s) = \frac{e^{-\frac{(x-y)^2}{2s}}}{\pi\sqrt{2\pi s}} \int_0^{+\infty} \xi e^{-s\Re(\Psi(\xi))} d\xi$  is integrable in  $(0, +\infty)$ . By using Fubini's theorem we have

$$\int_0^{+\infty} G(s)ds = \frac{1}{\pi\sqrt{2\pi}} \int_0^{+\infty} \xi \int_0^{+\infty} s^{-\frac{1}{2}} e^{-\frac{(x-y)^2}{2s} - s\Re(\Psi(\xi))} ds d\xi \quad (\text{A.26})$$

$$= \frac{1}{\sqrt{2\pi}} \int_0^{+\infty} \frac{\xi}{\sqrt{\Re(\Psi(\xi))}} e^{-|x-y|\sqrt{2\Re(\Psi(\xi))}} d\xi, \quad (\text{A.27})$$

where we also used Lemma A.2. Let us consider  $M > 0$  big enough such that, by Orey's condition (3.14), it holds

$$e^{-s\Re(\Psi(\xi))} \leq \exp\left\{-\frac{C_\gamma s}{4}\xi^\alpha\right\}, \quad \xi > M, \quad (\text{A.28})$$

for  $\alpha = 2 - \gamma$ . This implies in particular

$$\Re(\Psi(\xi)) \geq \frac{C_\gamma}{4}\xi^\alpha, \quad \xi > M. \quad (\text{A.29})$$

Let us also consider  $0 < \delta < \min\{M, \delta_0\}$ , where  $\delta_0 > 0$  is defined in Lemma A.3. By splitting the integral in  $\delta$  and  $M$ , we obtain

$$\int_0^{+\infty} \frac{\xi}{\sqrt{\Re(\Psi(\xi))}} e^{-|x-y|\sqrt{2\Re(\Psi(\xi))}} d\xi = \int_0^\delta \frac{\xi}{\sqrt{\Re(\Psi(\xi))}} e^{-|x-y|\sqrt{2\Re(\Psi(\xi))}} d\xi \quad (\text{A.30})$$

$$+ \int_\delta^M \frac{\xi}{\sqrt{\Re(\Psi(\xi))}} e^{-|x-y|\sqrt{2\Re(\Psi(\xi))}} d\xi \quad (\text{A.31})$$

$$+ \int_M^{+\infty} \frac{\xi}{\sqrt{\Re(\Psi(\xi))}} e^{-|x-y|\sqrt{2\Re(\Psi(\xi))}} d\xi \quad (\text{A.32})$$

$$=: I_1 + I_2 + I_3. \quad (\text{A.33})$$

Now we estimate the quantities in (A.33). By Lemma A.3 we already know that  $I_1 < \infty$ . Also  $I_2 < \infty$  clearly, thanks to the continuity of  $\frac{\xi}{\sqrt{\Re(\Psi(\xi))}}$  in  $[\delta, M]$ . Finally, concerning  $I_3$ , (A.29) implies

$$I_3 \leq C \int_M^{+\infty} \xi^{1-\frac{\alpha}{2}} e^{-|x-y|\sqrt{2\xi^{\frac{\alpha}{2}}}} d\xi < +\infty, \quad (\text{A.34})$$

concluding the proof.

- (2) To show the first assertion, let us consider any compact set  $\mathcal{K}_{(x,y)} \subseteq \mathbb{R}^2$  such that  $\mathcal{K}_{(x,y)} \cap \{(x, x), x \in \mathbb{R}\} = \emptyset$  and let  $\mathcal{K}_T = [0, T]$  for some  $T > 0$ . Let us show that  $\partial_t p_\Phi(t, x; y)$  is continuous in  $\mathcal{K} := \mathcal{K}_T \times \mathcal{K}_{(x,y)}$ . To do this, let us first observe that  $\partial_t f_L(s; t)$  is continuous in  $t$  for  $t > 0$  and fixed  $s > 0$ : this can be ascertained by using the representation in Lemma A.1 and the fact that by Orey's condition  $\partial_t g_\sigma(t - \tau; s)$  is continuous in  $t$ . Moreover, (A.2) guarantees that we can extend  $\partial_t f_L(s; t)$  by right-continuity in  $t = 0$ , setting  $\partial_t f_L(s; 0) = 0$ . Denoting  $m = \min_{(x,y) \in \mathcal{K}_{(x,y)}} |x-y| > 0$  and combining (3.3) and (A.2), it is clear that the integrand of (3.17) is dominated by

$$\frac{e^{-\frac{m^2}{2s}}}{\pi\sqrt{s}} I_\Phi(T) \int_0^{+\infty} \xi e^{-s\Re(\Psi(\xi))} d\xi, \quad (\text{A.35})$$

that is integrable in  $(0, +\infty)$  with respect to the variable  $s$ . This concludes the proof.

- (3) Let us define the functions

$$g_{\mathcal{I}}(x, s) := \begin{cases} \frac{1}{\pi\sqrt{2\pi s}} e^{-\frac{(a-x)^2}{2s}} I_\Phi(\tau) \int_0^{+\infty} \xi e^{-s\Re(\Psi(\xi))} d\xi, & x < a_0, \\ \frac{1}{\pi\sqrt{2\pi s}} e^{-\frac{(x-b)^2}{2s}} I_\Phi(\tau) \int_0^{+\infty} \xi e^{-s\Re(\Psi(\xi))} d\xi, & x > b_0, \\ 0, & x \in [a_0, b_0], \end{cases} \quad (\text{A.36})$$

and

$$f_{\mathcal{I}}(x) := \begin{cases} C(a_0, b_0, a, b, \tau), & x \in [a_0, b_0], \\ \int_0^{+\infty} g_{\mathcal{I}}(x, s) ds, & \text{elsewhere,} \end{cases} \quad (\text{A.37})$$

where  $C(a_0, b_0, a, b, \tau)$  is a suitable constant such that  $|\partial_t p_\Phi(t, x; y)| \leq f_{\mathcal{I}}(x)$  as  $(t, x, y) \in \mathcal{I}$ , that exists by item (2). We only have to show that  $f_{\mathcal{I}} \in L^1([a_1, b_1])$ . Clearly

$$\int_{a_0}^{b_0} f_{\mathcal{I}}(x) dx = (b_0 - a_0) C(a_0, b_0, a, b, \tau). \quad (\text{A.38})$$

On the interval  $(b_0, b_1)$  we have

$$\int_{b_0}^{b_1} f_{\mathcal{I}}(x) dx = \int_{b_0}^{b_1} \int_0^{+\infty} \frac{e^{-\frac{(x-b)^2}{2s}}}{\pi\sqrt{2\pi s}} I_\Phi(\tau) \int_0^{+\infty} \xi e^{-s\Re(\Psi(\xi))} d\xi ds dx. \quad (\text{A.39})$$

By Lemma A.2 and repeatedly using Fubini's Theorem we have that

$$\int_{b_0}^{b_1} f_{\mathcal{I}}(x) dx$$

$$\begin{aligned}
&= I_{\Phi}(\tau) \int_{b_0}^{b_1} \int_0^{+\infty} \frac{e^{-\frac{(x-b)^2}{2s}}}{\pi\sqrt{2\pi s}} \int_0^{+\infty} \xi e^{-s\Re(\Psi(\xi))} d\xi ds dx \\
&= I_{\Phi}(\tau) \int_{b_0}^{b_1} \int_0^{+\infty} \frac{\xi}{\pi\sqrt{\Re(\Psi(\xi))}} e^{-(x-b)\sqrt{2\Re(\Psi(\xi))}} d\xi dx \\
&= I_{\Phi}(\tau) \int_{b_0}^{b_1} \int_0^{\delta_0} \frac{\xi}{\pi\sqrt{\Re(\Psi(\xi))}} e^{-(x-b)\sqrt{2\Re(\Psi(\xi))}} d\xi \\
&\quad + I_{\Phi}(\tau) \int_{b_0}^{b_1} \int_{\delta_0}^M \frac{\xi}{\pi\sqrt{\Re(\Psi(\xi))}} e^{-(x-b)\sqrt{2\Re(\Psi(\xi))}} d\xi \\
&\quad + I_{\Phi}(\tau) \int_{b_0}^{b_1} \int_M^{\infty} \frac{\xi}{\pi\sqrt{\Re(\Psi(\xi))}} e^{-(x-b)\sqrt{2\Re(\Psi(\xi))}} d\xi \\
&=: I_1 + I_2 + I_3, \tag{A.40}
\end{aligned}$$

where  $\delta_0$  is defined in Lemma A.3 and  $M$  in (A.29). The first integral  $I_1$  is finite since

$$I_1 \leq \frac{1}{\pi} I_{\Phi}(T)(b_1 - b_0) \int_0^{x_0} \frac{\xi}{\sqrt{\Re(\Psi(\xi))}} d\xi < \infty, \tag{A.41}$$

by Lemma A.3. The term  $I_2$  is finite since the integrand is continuous. For  $I_3$  note that, for  $\alpha = 2 - \gamma$ ,

$$\begin{aligned}
I_3 &\leq I_{\Phi}(\tau) \int_{b_0}^{\infty} \int_M^{\infty} \frac{2}{\sqrt{C_{\gamma}}} \xi^{1-\frac{\alpha}{2}} e^{-(x-b)\sqrt{\frac{C_{\gamma}}{2}\xi^{\alpha}}} d\xi dx \\
&\leq \frac{1}{C_{\gamma}} I_{\Phi}(\tau) \int_M^{\infty} \xi^{1-\alpha} e^{-(b_0-b)\sqrt{C_0/2}\xi^{\alpha/2}} d\xi < \infty, \tag{A.42}
\end{aligned}$$

where we used (A.29) and A.3. Thus we have that  $\int_{b_0}^{b_1} f_{\mathcal{I}}(x)dx < +\infty$ . Arguing in a similar way in  $(a_1, a_0)$ , we conclude the proof.  $\square$

**A.2. Proof of Proposition 3.6.** Let us recall that

$$g_{\sigma}(\tau; s) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-i\xi\tau - s\Psi(\xi)} d\xi. \tag{A.43}$$

Using (A.43) we have

$$\left| \frac{g_{\sigma}(\tau; s) - g_{\sigma}(\tau; s')}{s - s'} \right| \leq \frac{1}{2\pi} \int_{-\infty}^{+\infty} \left| \frac{e^{-s\Psi(\xi)} - e^{-s'\Psi(\xi)}}{s - s'} \right| d\xi. \tag{A.44}$$

Suppose  $s > s'$ . By (A.44) and the fact that  $x^{-1}(1 - e^{-x}) \leq 1$ , we find

$$\begin{aligned}
&\left| \frac{g_{\sigma}(\tau; s) - g_{\sigma}(\tau; s')}{s - s'} \right| \\
&\leq \frac{1}{2\pi} \int_{-\infty}^{+\infty} \left[ e^{-s\Re(\Psi(\xi))} \left| \frac{e^{-is\Im(\Psi(\xi))} - e^{-is'\Im(\Psi(\xi))}}{\Im(\Psi(\xi))(s - s')} \right| |\Im(\Psi(\xi))| \right. \\
&\quad \left. + e^{-s\Re(\Psi(\xi))} \left| \frac{e^{-(s-s')\Re(\Psi(\xi))} - 1}{\Re(\Psi(\xi))(s - s')} \right| \Re(\Psi(\xi)) \right] d\xi
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-s\Re(\Psi(\xi))} |\Im(\Psi(\xi))| d\xi + \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-s\Re(\Psi(\xi))} \Re(\Psi(\xi)) d\xi \\
&= \frac{1}{2\pi} \int_{-M}^M e^{-s\Re(\Psi(\xi))} (|\Im(\Psi(\xi))| + \Re(\Psi(\xi))) d\xi \\
&\quad + \frac{1}{2\pi} \int_{|\xi|>M} e^{-s\Re(\Psi(\xi))} |\Im(\Psi(\xi))| d\xi \\
&\quad + \frac{1}{2\pi} \int_{|\xi|>M} e^{-s\Re(\Psi(\xi))} \Re(\Psi(\xi)) d\xi \\
&=: I_1 + I_2 + I_3
\end{aligned} \tag{A.45}$$

where  $M$  is defined in (A.29). Note now that

$$\Im(\Psi(\xi)) = - \int_0^{+\infty} \sin(\xi t) \nu(dt) \tag{A.46}$$

is odd and therefore  $|\Im(\Psi(\xi))|$  is even. For  $\xi > M$  and  $t < 1$ , it is clear that  $|\sin(\xi t)| \leq \xi t$  and then

$$|\Im(\Psi(\xi))| \leq \xi \int_0^1 t \nu(dt) + \bar{\nu}(1). \tag{A.47}$$

It follows by (A.47) and (A.29) that

$$I_2 \leq \frac{1}{\pi} \int_{\xi>M} \left( \xi \int_0^1 t \nu(dt) + \bar{\nu}(1) \right) e^{-s\xi^\alpha} d\xi < \infty, \tag{A.48}$$

where  $\alpha = 2 - \gamma$ . Observe now that  $\Re(\Psi(\xi))$  is even and that  $1 - \cos(\xi t) \leq \frac{\xi^2 t^2}{2}$ , thus

$$\Re(\Psi(\xi)) = \int_0^{+\infty} (1 - \cos(\xi t)) \nu(dt) \leq \frac{\xi^2}{2} \int_0^1 t^2 \nu(dt) + 2\bar{\nu}(1). \tag{A.49}$$

It follows by (A.49) and (A.29) that

$$I_3 \leq \frac{1}{\pi} \int_{\xi>M} \left( \frac{\xi^2}{2} \int_0^1 t^2 \nu(dt) + 2\bar{\nu}(1) \right) e^{-s\xi^\alpha} d\xi < \infty. \tag{A.50}$$

It also clear that  $I_1$  is finite. Repeating the same argument for  $s' > s$  and combining the results we obtained, for fixed  $s > 0$  and  $s' \in (s/2, 3s/2)$ ,

$$\left| \frac{g(\tau, s) - g(\tau, s')}{s - s'} \right| \leq \mathcal{C}_1 \tag{A.51}$$

where  $\mathcal{C}_1 > 0$  does not depend on  $\tau$ . Therefore, using (A.3) we obtain

$$\left| \frac{f_L(s, t) - f_L(s', t)}{s - s'} \right| \leq \mathcal{C}_1 \int_0^t \bar{\nu}(\tau) d\tau. \tag{A.52}$$

Hence the first assertion follows by dominated convergence. Now we prove the claimed behaviour at infinity. By (A.3) and (A.43), it holds

$$\sup_{t \in [a, b]} s f_L(s, t) \leq \underbrace{\int_0^b \bar{\nu}(\tau) d\tau}_{=: \mathcal{C}_2} s \int_{-\infty}^{+\infty} e^{-s\Re(\Psi(\xi))} d\xi$$

$$\begin{aligned}
&\leq 2C_2 \left[ \int_0^M s e^{-s\Re(\Psi(\xi))} d\xi + s \int_M^{+\infty} e^{-s\xi^\alpha} d\xi \right] \\
&\leq 2C_2 \left[ \int_0^M s e^{-s\Re(\Psi(\xi))} d\xi + \frac{1}{\alpha} s^{1-\frac{1}{\alpha}} \Gamma(1/\alpha) \right]. \tag{A.53}
\end{aligned}$$

It is clear that (A.53) tends to zero as  $s \rightarrow \infty$  by using dominated convergence on the first integral and the fact that  $\alpha < 1$  by Remark 3.3. The last assertion follows by similar computation. Indeed, in the same spirit of (A.53), we obtain

$$\sup_{t \in [a, b]} s^2 |\partial_s f_L(s, t)| \leq 2C_2 \left[ \int_0^M |\Psi(\xi)| s^2 e^{-s\Re(\Psi(\xi))} d\xi + s^2 \int_M^\infty |\Psi(\xi)| e^{-s\xi^\alpha} d\xi \right]. \tag{A.54}$$

Using the estimates on  $\Re(\Psi(\xi))$  and  $\Im(\Psi(\xi))$  determined above we obtain, for  $\xi > M$  sufficiently big,

$$|\Psi(\xi)| \leq C_3 \xi^2 \tag{A.55}$$

for a suitable  $C_3 > 0$ . It follows that

$$\sup_{t \in [a, b]} s^2 |\partial_s f_L(s, t)| \leq 2C_2 \left[ \int_0^M |\Psi(\xi)| s^2 e^{-s\Re(\Psi(\xi))} d\xi + \frac{1}{\alpha} C_3 s^{2-\frac{3}{\alpha}} \Gamma(3/\alpha) \right]. \tag{A.56}$$

It is clear that (A.56) tends to zero as  $s \rightarrow \infty$  by using dominated convergence on the first integral and, again, the fact that  $\alpha < 1$  by Remark 3.3.  $\square$

**A.3. Proof of Proposition 3.8.** Since it holds

$$\partial_x p(s, x; y) = -\sqrt{\frac{2}{\pi s^3}} (x - y) e^{-\frac{(x-y)^2}{2s}}, \tag{A.57}$$

it is clear that (3.20) follows from (3.2) once we prove that we can take the derivative inside the integral sign. To do this, let us rewrite (3.2) as

$$p_\Phi(t, x; y) = \mathbb{E}[p(L(t), x; y)] \tag{A.58}$$

and let us observe that, for any compact set  $\mathcal{K}_{(x, y)} \subset \mathbb{R}^2 \setminus \{(x, x) : x \in \mathbb{R}\}$  it holds

$$|\partial_x p(L(t), x; y)| \leq \sqrt{\frac{2}{\pi(L(t))^3}} M e^{-\frac{m^2}{2L(t)}} \leq C,$$

where  $M = \max_{(x, y) \in \mathcal{K}_{(x, y)}} |x - y|$ ,  $m = \min_{(x, y) \in \mathcal{K}_{(x, y)}} |x - y| > 0$  and  $C > 0$  is a constant such that  $\sqrt{\frac{2}{\pi s^3}} M e^{-\frac{m^2}{2Ls}} \leq C$  for any  $s \in \mathbb{R}_0^+$ . This is enough, together with (A.58) and the continuity of the paths of  $L(t)$ , to guarantee both (3.20) and the continuity of  $\partial_x p_\Phi(t, x; y)$  on  $\mathbb{R}_0^+ \times (\mathbb{R}^2 \setminus \{(x, x) : x \in \mathbb{R}\})$ . The arguments to prove (3.21) and the continuity of  $\partial_x^2 p_\Phi(t, x; y)$  are exactly the same.  $\square$

**A.4. Proof of Proposition 3.14.** Without loss of generality we can suppose that  $l = 1$ . Fix  $\epsilon > 0$ . Then there exists  $T > 0$  such that for any  $t \geq T$  it is true that  $1 - F(t) < \epsilon/2$ . Note that  $F(t)$  is continuous in  $[0, T]$  and thus uniformly continuous. It follows that there exists  $\delta > 0$  such that for any choice of  $t_1$  and  $t_2$  in  $[0, T]$  it holds

$$|t_2 - t_1| < \delta \implies |F(t_2) - F(t_1)| < \epsilon/2. \quad (\text{A.59})$$

Take now a finite partition  $\Pi$  of  $[0, T]$  with  $\text{diam}(\Pi) < \delta$ . Since  $F_n(t) \rightarrow F(t)$  for any  $t$  we have by pointwise convergence that there exists  $\nu > 0$  such that, for any  $n > \nu$  one has

$$|F_n(t_i) - F(t_i)| < \epsilon/2 \text{ for all } t_i \in \Pi. \quad (\text{A.60})$$

Now consider  $t > T$ . We have that

$$F_n(t) \leq 1 \leq F(t) + \epsilon/2 < F(t) + \epsilon \quad (\text{A.61})$$

where we used the fact that since  $t > T$  then  $1 < F(t) + \epsilon/2$ . On the other hand, for  $n > \nu$

$$\begin{aligned} F_n(t) &\geq F_n(T) \\ &\geq F(T) - \epsilon/2 && \text{by (A.60) since } T \in \Pi \\ &\geq F(t) - (F(t) - F(T)) - \epsilon/2 \\ &\geq F(t) - \epsilon && \text{since } F(t) - F(T) \leq 1 - F(T) < \epsilon/2. \end{aligned} \quad (\text{A.62})$$

For  $t \in [0, T]$  instead, we have that there exists  $t_i, t_{i+1} \in \Pi$  such that  $t \in [t_i, t_{i+1}]$ . It follows that, for  $n > \nu$ ,

$$\begin{aligned} F_n(t) &\leq F_n(t_{i+1}) \\ &\leq F(t_{i+1}) + \epsilon/2 && \text{by (A.60) since } t_{i+1} \in \Pi \\ &\leq F(t) + (F(t_{i+1}) - F(t)) + \epsilon/2 \\ &\leq F(t) + \epsilon && \text{by (A.59) since } \text{diam}(\Pi) < \delta. \end{aligned} \quad (\text{A.63})$$

With the same argument it is possible to see that, for any  $n > \nu$

$$F_n(t) \geq F(t) - \epsilon. \quad (\text{A.64})$$

□

#### ACKNOWLEDGEMENTS

This work has been partially supported by the MIUR PRIN 2017 project ‘‘Stochastic Models for Complex Systems’’, no. 2017JFFHSH.

#### REFERENCES

- [1] W. Arendt, C.J.K. Batty, M. Hieber and F. Neubrander. Vector valued Laplace transform and Cauchy problem. Second Edition. *Birkhuser*, Berlin, 2010.
- [2] G. Ascione, N. Leonenko and E. Pirozzi. Time-Non-Local Pearson Diffusions. *Journal of Statistical Physics* 183(3): 1 – 42, 2021.
- [3] G. Ascione, Y. Mishura and E. Pirozzi. Convergence results for the time-changed fractional Ornstein–Uhlenbeck processes. *Theory of Probability and Mathematical Statistics* 104: 23 – 47, 2021.

- [4] G. Ascione, E. Pirozzi and B. Toaldo. On the exit time from open sets of some semi-Markov processes. *Annals of Applied Probability*, 30(3): 1130 – 1163, 2020.
- [5] G. Ascione. Abstract Cauchy problems for the generalized fractional calculus. *Nonlinear Analysis, Theory, Methods and Applications*, 209: 112339, 2021.
- [6] B. Baeumer and M.M. Meerschaert. Stochastic solutions for fractional Cauchy problems. *Fractional Calculus and Applied Analysis* 4: 481–500, 2001.
- [7] B. Baeumer and P. Straka. Fokker-Planck and Kolmogorov backward equations for continuous time random walk limits. *Proceedings of the American Mathematical Society*, 145: 399 – 412, 2017.
- [8] P. Becker Kern, M.M. Meerschaert and H. P. Scheffler. Limit theorems for coupled continuous time random walks, *Ann. Probab.*, 32: 730 – 756, 2004.
- [9] L. Beghin, C. Macci and C. Ricciuti. Random time-change with inverses of multivariate subordinators: Governing equations and fractional dynamics. *Stoch. Proc. Appl.*, 130(10): 6364 – 6387, 2020.
- [10] J. Bertoin. Lévy processes. *Cambridge University Press*, Cambridge, 1996.
- [11] J. Bertoin. Subordinators: examples and applications. *Lectures on Probability Theory and Statistics (Saint-Flour, 1997)*, 1 – 91. *Lectures Notes in Math.*, 1717, Springer, Berlin, 1999.
- [12] N.H. Bingham, C.M. Goldie and J.F. Teugels. Regular variation. *Cambridge University Press*, Cambridge, 1987.
- [13] R. Capitanelli and M. D’Ovidio. Asymptotics for time-changed diffusion. *Theory of Probability and Mathematical Statistics*, special volume 95 in honor of Prof. N. Leonenko: 37 – 54, 2017.
- [14] R. Capitanelli and M. D’Ovidio. Delayed and rushed motions through time change. *Alea (Rio de Janeiro)*, 17: 183 – 204, 2020.
- [15] Z.-Q. Chen. Time fractional equations and probabilistic representation. *Chaos, Solitons and Fractals*, 102: 168 – 174, 2017.
- [16] E. Cinlar. Markov additive processes. II. *Z. Wahrscheinlichkeitstheorie verw. Geb.*, 24: 95 – 121, 1972.
- [17] B. Dacorogna. Introduction to Calculus of Variations. *World Scientific Publishing*, 2014.
- [18] W. Feller. *An Introduction to Probability Theory and Its Applications*. Vol. II, 2nd Ed., Wiley, New York, 1971.
- [19] R. Garra, F. Falcini, V. R. Voller, F. Pagnini A generalized Stefan model accounting for system memory and non-locality, *International Communications in Heat and Mass Transfer* 114 (2020): 104584.
- [20] R. K. Gettoor and M. J. Sharpe. Excursions of dual processes. *Advances in Mathematics*, 45.3, 259-309, 1982.
- [21] N. Georgiou, I.Z. Kiss and E. Scalas. Solvable non-Markovian dynamic network. *Physical Review E*, 92, 042801, 2015.
- [22] I.S. Gradshteyn and I.M. Ryzhik. *Table of integrals, series, and products*. Academic press, 2014
- [23] M. Hairer, G. Iyer, L. Korolov, A. Novikov, and Z. Pajor-Gyulai. A fractional kinetic process describing the intermediate time behaviour of cellular flows. *The Annals of Probability*, 46(2): 897 – 955, 2018.
- [24] E. Hille, R. S. Phillips. Functional analysis and semi-groups. *American Mathematical Soc.*, Vol. 31, 1996.

- [25] N. Jacob. Pseudo-differential operators and Markov processes. Vol I. *Imperial College Press*, 2002.
- [26] J. Jacod. Systèmes régénératifs and processus semi-Markoviens. *Z. Wahrscheinlichkeitstheorie verw. Gebiete*, 31: 1 – 23, 1974.
- [27] H. Kaspı and B. Maisonneuve. Regenerative systems on the real line. *Ann. Probab.*, 16: 1306 – 1332, 1988.
- [28] A.N. Kochubei. General fractional calculus, evolution equations and renewal processes. *Integral Equations and Operator Theory*, 71: 583 – 600, 2011.
- [29] A.N. Kochubei, Y.G. Kondratiev and J.L. da Silva. Random time change and related evolution equations. Time asymptotic behavior. *Stochastics and Dynamics*, 20(5): 2050034, 2020.
- [30] A.N. Kolmogorov and S.V. Fomin. Introductory real analysis. *Dover Publications, Inc.*, New York, 1975.
- [31] V.N. Kolokoltsov. Generalized Continuous-Time Random Walks, subordination by hitting times, and fractional dynamics. *Theory Probab. Appl.* 53:, 594–609, 2009.
- [32] M. E. Hernández-Hernández, V.N. Kolokoltsov and L. Toniazzi. Generalised Fractional Evolution Equations of Caputo Type. *Chaos, Solitons & Fractals*, 102: 184 – 196, 2017.
- [33] A. Kumar and P. Vellaisamy. Inverse tempered stable subordinators. *Statistics and Probability Letters*, 103: 134 – 141, 2015.
- [34] H.R. Lerche. Boundary Crossing of Brownian Motion. *Springer-Verlag*, 1980.
- [35] Y. Luchko. Maximum principle for the generalized time-fractional diffusion equation. *Journal of Mathematical Analysis and Applications* 351.1: 218–223, 2009.
- [36] M. Magdziarz and R.L. Schilling. Asymptotic properties of Brownian motion delayed by inverse subordinators. *Proc. Amer. Math. Soc.*, 143: 4485 – 4501, 2015.
- [37] M. M. Meerschaert, D. A. Benson, H. P. Scheffler and P. Becker-Kern. *Governing equations and solutions of anomalous random walk limits*, Phys. Rev. E (66) 060102(R), 2002.
- [38] M.M. Meerschaert and H.P. Scheffler. Triangular array limits for continuous time random walks. *Stochastic Processes and their Applications*, 118(9): 1606 – 1633, 2008.
- [39] M.M. Meerschaert, E. Nane and P. Vellaisamy. The fractional Poisson process and the inverse stable subordinator. *Electronic Journal of Probability*, 16(59): 1600–1620, 2011.
- [40] M.M. Meerschaert, E. Nane and P. Vellaisamy. Fractional Cauchy problems on bounded domains. *The Annals of Probability*, 37(3): 979 – 1007, 2009.
- [41] M. M. Meerschaert, and P. Straka. Inverse stable subordinators. *Mathematical modelling of natural phenomena*, 8.2: 1 – 16, 2013.
- [42] M.M. Meerschaert and P. Straka. Semi-Markov approach to continuous time random walk limit processes. *The Annals of Probability*, 42(4) : 1699 – 1723, 2014.
- [43] M.M. Meerschaert and B. Toaldo. Relaxation patterns and semi-Markov dynamics. *Stoch. Proc. Appl.*,29(8): 2850 – 2879, 2019.
- [44] R. Metzler and J. Klafter. The random walk’s guide to anomalous diffusion: a fractional dynamics approach. *Physics Reports*, 339: 1 – 77, 2000.

- [45] P. Mörters and Y. Peres. Brownian motion. *Cambridge University Press*, Cambridge, 2010.
- [46] S. Orey. On continuity properties of infinitely divisible distribution functions. *The Annals of Mathematical Statistics*, 39(3): 936–937, 1968.
- [47] E. Orsingher, C. Ricciuti and B. Toaldo. Time-inhomogeneous jump processes and variable order operators. *Potential Analysis*, 45(3): 435 – 461, 2016.
- [48] E. Orsingher, C. Ricciuti and B. Toaldo. On semi-Markov processes and their Kolmogorov integro-differential equations. *Journal of Functional Analysis*, 275(4): 830 – 868, 2018.
- [49] A. Pachon, F. Polito, C. Ricciuti. On discrete-time semi-Markov processes. *Discr. Cont. Dyn. Syst. B*, 26(3): 1499 – 1529, 2020.
- [50] P. Patie, M. Savov and R.L. Loeffen. Extinction time of non-Markovian self-similar processes, persistence, annihilation of jumps and the Fréchet distribution. *J. Stat. Phys.*, 175(5): 1022 – 1041, 2019.
- [51] P. Patie and A. Srapionyan. Self-similar Cauchy problems and generalized Mittag-Leffler functions. *Frac. Calc. Appl. Anal.*, 24(2): 447 - 482, 2021.
- [52] K. A. Penson and K. Górska. Exact and explicit probability densities for one-sided Lévy stable distributions. *Physical review letters* 105.21: 210604, 2010.
- [53] C. Ricciuti and B. Toaldo. Semi-Markov models and motion in heterogeneous media. *Journal of Statistical Physics*, 169(2): 340 - 361, 2017.
- [54] S.D. Roscani. Hopf lemma for the fractional diffusion operator and its application to a fractional free-boundary problem. *Journal of Mathematical Analysis and Applications*, 434(1): 125 – 135, 2016.
- [55] S.D. Roscani. Moving-boundary problems for the time-fractional diffusion equation. *Electronic Journal of Differential Equations*, 2017(44): 1 – 12, 2017.
- [56] M. Savov and B. Toaldo. Semi-Markov processes, integro-differential equations and anomalous diffusion-aggregation. *Annales de l'Institut Henri Poincaré (B) Probability and Statistics*, 56(4): 2640 – 2671, 2020.
- [57] E. Scalas. Five years of continuous-time random walks in econophysics. The Complex Networks of Economic Interactions, *Lecture Notes in Economics and Mathematical Systems*, 567: 3–16, Springer, Berlin, 2006.
- [58] R.L. Schilling, R. Song and Z. Vondraček. Bernstein functions: theory and applications. *Walter de Gruyter GmbH & Company KG*, Vol 37 of De Gruyter Studies in Mathematics Series, 2010.
- [59] P. Straka. Variable Order Fractional Fokker-Planck Equations derived from Continuous Time Random Walks. *Physica A: Statistical Mechanics and its Applications*, 503: 451 – 463, 2018.
- [60] B. Toaldo. Convolution-type derivatives, hitting-times of subordinators and time-changed  $C_0$ -semigroups. *Potential Analysis*, 42(1): 115 – 140, 2015.
- [61] B. Toaldo. Lévy mixing related to distributed order calculus, subordinators and slow diffusions. *Journal of Mathematical Analysis and Applications*, 430(2): 1009 – 1036, 2015.
- [62] C.J. Vogl, M.J. Miksis and S.H. Davis. Moving boundary problems governed by anomalous diffusion. *Proc. R. Soc. A*, 468: 3348 – 3369, 2012.
- [63] V.R. Voller Fractional stefan problems. *International Journal of Heat and Mass Transfer*, 74: 269 – 277, 2014
- [64] V.R. Voller, F. Falcini and R. Garra. Fractional Stefan problems exhibiting lumped and distributed latent-heat memory effects. *Physical Review E*, 87:

042401, 2013.

- [65] R. L. Wheeden, A. Zygmund. Measure and integral. *Dekker*, New York, 1977.
- [66] W. Whitt. Stochastic-Process Limits. *Springer-Verlag*, Berlin, 2002.

SCUOLA SUPERIORE MERIDIONALE - UNIVERSITÀ DEGLI STUDI DI NAPOLI "FEDERICO II", LARGO  
S. MARCELLINO 10 - 80138, NAPOLI (ITALY)

*Email address:* [giacomo.ascione@unina.it](mailto:giacomo.ascione@unina.it)

CORNELL UNIVERSITY, 219 RHODES HALL - 14852, ITHACA, NY (USA)

*Email address:* [ppatie@cornell.edu](mailto:ppatie@cornell.edu)

DIPARTIMENTO DI MATEMATICA "GIUSEPPE PEANO" - UNIVERSITÀ DEGLI STUDI DI TORINO,  
VIA CARLO ALBERTO 10 - 10123, TORINO (ITALY)

*Email address:* [bruno.toaldo@unito.it](mailto:bruno.toaldo@unito.it)