An extended Rayleigh model: Properties, regression and COVID-19 application

Gauss M. Cordeiro Universidade Federal de Pernambuco gausscordeiro@gmail.com

> Gabriela M. Rodrigues Universidade de São Paulo gabrielar@usp.br

> Edwin M. M. Ortega Universidade de São Paulo edwin@usp.br

Luís H. de Santana Universidade Federal de Pernambuco desantanalh@gmail.com

> Roberto Vila Universidade de Brasília rovig1610gmail.com

Abstract

We define a four-parameter extended Rayleigh distribution, and obtain several mathematical properties including a stochastic representation. We construct a regression from the new distribution. The estimation is done by maximum likelihood. The utility of the new models is proved in two real applications.

Keywords: Censored data, generalized Rayleigh; maximum likelihood estimation; regression model, stochastic representation.

1 Introduction

The Rayleigh distribution has been employed in many areas (Johnson *et al.*, 1994). Several of its extensions have been published so far.

The cumulative distribution function (cdf) of the generalized Rayleigh (GR) distribution (Vodă, 1976) is

$$G_{\rm GR}(x;\delta,\theta) = \gamma_1(\delta+1,\theta x^2), \quad x > 0, \tag{1}$$

where $\delta > -1$ and $\theta > 0$, $\Gamma(p) = \int_0^\infty w^{p-1} e^{-w} dw$ and $\gamma_1(p, x) = \Gamma(p)^{-1} \int_0^x w^{p-1} e^{-w} dw$ are the gamma function and lower incomplete gamma function ratio, respectively.

The GR distribution includes well-known sub-models. The classical Rayleigh distribution follows when $\delta = 0$ and $\theta = \lambda^{-2}$. If $\delta = 2^{-1}$ and $\theta = (2\lambda^2)^{-1}$, it gives the Maxwell distribution. The chi-square refers to $\theta = (2\tau^2)^{-1}$, $\tau > 0$, and $\delta = n/2 - 1$, $n \in \mathbb{N}$, and the half-normal to $\delta = -2^{-1}$ and $\theta = 2\sigma^{-2}$.

The probability density function (pdf) corresponding to (1) is

$$g_{\rm GR}(x;\delta,\theta) = \frac{2\theta^{\delta+1}}{\Gamma(\delta+1)} x^{2\delta+1} e^{-\theta x^2}.$$
 (2)

Let $Z \sim \text{GR}(\delta, \theta)$ be a random variable with density (2). The moments of Z are easily obtained from the integral given in Section 3.478 of Gradshteyn and Ryzhik (2000) $\int_0^\infty x^{\nu-1} e^{-\mu x^p} dx = \frac{\Gamma(\nu/p)}{p} \mu^{\nu/p}$, where $p, \nu, \mu > 0$. Indeed, the sth ordinary moment of Z (for a positive real number s) is

$$\mu'_{s}(\delta,\theta) = \frac{\Gamma(s/2 + \delta + 1)}{\theta^{s/2} \Gamma(\delta + 1)}.$$
(3)

The cdf of the generalized odd log-logistic-G (GOLL-G) family follows from Cordeiro *et al.* (2017)

$$F(x;\alpha,\beta,\boldsymbol{\xi}) = \frac{G(x;\boldsymbol{\xi})^{\alpha\beta}}{G(x;\boldsymbol{\xi})^{\alpha\beta} + \left[1 - G(x;\boldsymbol{\xi})^{\beta}\right]^{\alpha}},\tag{4}$$

where $\alpha > 0$ and $\beta > 0$ are two extra parameters.

The generalized log-logistic (Gleaton and Lynch, 2006) and proportional reversed hazard rate (Gupta and Gupta, 2007) are special models of Equation (4). Further, the parent comes when $\alpha = \beta = 1$. Prataviera *et al.* (2018) discussed the generalized odd log-logistic flexible Weibull distribution.

If $g(x; \boldsymbol{\xi})$ is the parent density, the pdf corresponding to (4) can be written as

$$f(x;\alpha,\beta,\boldsymbol{\xi}) = \frac{\alpha\beta g(x;\boldsymbol{\xi}) G(x;\boldsymbol{\xi})^{\alpha\beta-1} [1 - G(x;\boldsymbol{\xi})^{\beta}]^{\alpha-1}}{\left\{G(x;\boldsymbol{\xi})^{\alpha\beta} + [1 - G(x;\boldsymbol{\xi})^{\beta}]^{\alpha}\right\}^{2}}.$$
(5)

The paper is structured as follows. Section 2 defines the generalized odd log-logistic generalized Rayleigh (GOLLGR) distribution, addresses some asymptotes and quantile function (qf), gives a linear representation for the family density, reports moments and generating function, and addresses maximum likelihood estimation. A new regression is constructed in Section 3, and some simulation studies are carried out as well as residual analysis. Two lifetime data sets in Section 4 show the utility of the new models. Some conclusions are reported in Section 5.

2 The GOLLGR model, properties and estimation

The GOLLGR density is defined (for x > 0) by inserting (1) and (2) in Equation (5)

$$f(x) = f(x;\alpha,\beta,\delta,\theta) = \frac{2\alpha\beta\theta^{\delta+1} x^{2\delta+1} e^{-\theta x^2} \gamma_1 \left(\delta+1,\theta x^2\right)^{\alpha\beta-1} \left[1-\gamma_1 \left(\delta+1,\theta x^2\right)^{\beta}\right]^{\alpha-1}}{\Gamma(\delta+1) \left\{\gamma_1 \left(\delta+1,\theta x^2\right)^{\alpha\beta}+\left[1-\gamma_1 \left(\delta+1,\theta x^2\right)^{\beta}\right]^{\alpha}\right\}^2}, \quad (6)$$

and its hazard rate function (hrf) is

$$h(x) = \frac{2\alpha\beta\theta^{\delta+1} x^{2\delta+1} e^{-\theta x^2} \gamma_1 (\delta+1, \theta x^2)^{\alpha\beta-1}}{\Gamma(\delta+1) \left[1 - \gamma_1 (\delta+1, \theta x^2)^{\beta}\right] \left\{\gamma_1 (\delta+1, \theta x^2)^{\alpha\beta} + \left[1 - \gamma_1 (\delta+1, \theta x^2)^{\beta}\right]^{\alpha}\right\}}$$
(7)

We have $\lim_{x\to\infty} f(x) = 0$, and

$$\lim_{x \to 0^{+}} f(x) = \begin{cases} \frac{2\sqrt{\theta}}{\sqrt{\pi}}, & \alpha\beta - 1 = 0, \ 2\delta + 1 = 0, \\ 0, & \alpha\beta - 1 > 0, \ 2\delta + 1 > 0, \\ \infty, & \alpha\beta - 1 > 0, \ 2\delta + 1 < 0, \ 0 < \delta + 1 < \frac{2\delta + 1}{2(1 - \alpha\beta)}, \\ \frac{2\alpha\beta\sqrt{\theta}}{\Gamma(\delta + 1)} \left[\frac{2(1 - \alpha\beta)}{\Gamma(\delta + 1)(2\delta + 1)}\right]^{\alpha\beta - 1}, & \alpha\beta - 1 > 0, \ 2\delta + 1 < 0, \ \delta + 1 = \frac{2\delta + 1}{2(1 - \alpha\beta)}, \\ 0, & \alpha\beta - 1 > 0, \ 2\delta + 1 < 0, \ \delta + 1 > \frac{2\delta + 1}{2(1 - \alpha\beta)}, \\ \infty, & \alpha\beta - 1 < 0, \ 2\delta + 1 > 0, \ \delta + 1 > \frac{2\delta + 1}{2(1 - \alpha\beta)}, \\ \infty, & \alpha\beta - 1 < 0, \ 2\delta + 1 > 0, \ \delta + 1 > \frac{2\delta + 1}{2(1 - \alpha\beta)}, \\ 0, & \alpha\beta - 1 < 0, \ 2\delta + 1 > 0, \ \delta + 1 = \frac{2\delta + 1}{2(1 - \alpha\beta)}, \\ 0, & \alpha\beta - 1 < 0, \ 2\delta + 1 > 0, \ \delta + 1 = \frac{2\delta + 1}{2(1 - \alpha\beta)}, \\ 0, & \alpha\beta - 1 < 0, \ 2\delta + 1 > 0, \ \delta + 1 = \frac{2\delta + 1}{2(1 - \alpha\beta)}, \\ 0, & \alpha\beta - 1 < 0, \ 2\delta + 1 > 0, \ 0 < \delta + 1 < \frac{2\delta + 1}{2(1 - \alpha\beta)}, \\ \infty, & \alpha\beta - 1 < 0, \ 2\delta + 1 > 0, \ 0 < \delta + 1 < \frac{2\delta + 1}{2(1 - \alpha\beta)}, \\ \infty, & \alpha\beta - 1 < 0, \ 2\delta + 1 > 0, \ 0 < \delta + 1 < \frac{2\delta + 1}{2(1 - \alpha\beta)}, \\ \infty, & \alpha\beta - 1 < 0, \ 2\delta + 1 > 0, \ 0 < \delta + 1 < \frac{2\delta + 1}{2(1 - \alpha\beta)}, \\ \infty, & \alpha\beta - 1 < 0, \ 2\delta + 1 < 0, \ \alpha\beta - 1 < 0 \ 2\delta + 1 < 0. \end{cases}$$

Further, $\lim_{x\to\infty} h(x) = \infty$ and $\lim_{x\to 0^+} h(x) = \lim_{x\to 0^+} f(x)$.

Hereafter, we omit the arguments of the functions, and let $X \sim \text{GOLLGR}(\alpha, \beta, \delta, \theta)$ be a random variable with pdf (6), The odd log-logistic GR follows when $\beta = 1$, and the proportional reversed hazard rate GR when $\alpha = 1$. Plots of the pdf and hrf of X are displayed in Figures1 and 2, respectively. They reveal the bimodality and asymmetry of the pdf, and different shapes of the hrf.

2.1 Asymptotes and quantile function

The asymptotes below follow from Equations (4) and (5)

$$F(x) \sim G(x)^{\alpha\beta}, \quad f(x), h(x) \sim \alpha\beta g(x)G(x)^{\alpha\beta-1} \quad \text{as} \quad x \to 0^+,$$



Figure 1: The GOLLGR pdf. (a) $\delta = 1.5$ and $\theta = 15$. (b) $\alpha = 0.3$, $\beta = 2$ and $\theta = 15$. (c) $\alpha = 0.3$, $\beta = 1.5$ and $\delta = 1.5$.

and

$$f(x) \sim \alpha \beta g(x) \left[1 - G(x)^{\beta}\right]^{\alpha - 1}, \quad h(x) \sim \frac{\alpha \beta g(x)}{\left[1 - G(x)^{\beta}\right]} \quad \text{as} \quad x \to \infty.$$

For the GOLLGR distribution, we obtain

$$F(x) \sim \gamma_1 (\delta + 1, \theta x^2)^{\alpha \beta},$$

$$f(x), h(x) \sim \frac{2\alpha \beta \theta^{\delta + 1}}{\Gamma(\delta + 1)} x^{2\delta + 1} e^{-\theta x^2} \gamma_1 (\delta + 1, \theta x^2)^{\alpha \beta - 1} \quad \text{as} \quad x \to 0^+,$$

and

$$f(x) \sim \frac{2\alpha\beta\theta^{\delta+1} x^{2\delta+1} e^{-\theta x^2} [1 - \gamma_1 (\delta + 1, \theta x^2)^{\beta}]^{\alpha-1}}{\Gamma(\delta + 1)},$$

$$h(x) \sim \frac{2\alpha\beta\theta^{\delta+1} x^{2\delta+1} e^{-\theta x^2}}{\Gamma(\delta + 1) [1 - \gamma_1 (\delta + 1, \theta x^2)^{\beta}]} \quad \text{as} \quad x \to \infty.$$

By combining the inverse functions of (1) and (4), the qf of X can be expressed as

$$x = Q(u) = Q_{\rm GR} \left(\left[\frac{\left(\frac{u}{1-u}\right)^{\frac{1}{\alpha}}}{1 + \left(\frac{u}{1-u}\right)^{\frac{1}{\alpha}}} \right]^{1/\beta} \right), \tag{9}$$



Figure 2: The GOLLGR hrf. (a) $\alpha = 0.1$ and $\delta = 1.5$. (b) $\alpha = 0.3$ and $\beta = 2.5$. (c) $\alpha = 0.1$.

where the GR qf $Q_{\rm GR}(z) = G_{\rm GR}^{-1}(z; \delta, \theta)$ comes as

$$Q_{\rm GR}(z) = \left[\theta^{-1} \gamma^{-1} (\delta + 1; z)\right]^{1/2}.$$
 (10)

Here, $\gamma^{-1}(\delta + 1; z)$ is the gamma qf with shape $\delta + 1$ and unity scale. So, the GOLLGR variates follow easily from (9).

2.2 Stochastic representation

The pdf of a log-logistic random variable $Y \sim LL(\nu, \alpha)$ is

$$f_Y(y;\nu,\alpha) = \frac{(\alpha/\nu)(y/\nu)^{\alpha-1}}{\left[1 + (y/\nu)^{\alpha}\right]^2},$$
(11)

where $\nu > 0$ and $\alpha > 0$ are scale and shape, respectively.

If $X \sim \text{GOLLGR}(\alpha, \beta, \delta, \theta)$ and $Y \sim \text{LL}(1, \alpha)$, we can write from (4) and (10)

$$F(x;\alpha,\beta,\delta,\theta) = \mathbb{P}\left(Y \leqslant \frac{G_{\mathrm{GR}}(x;\delta,\theta)^{\beta}}{1 - G_{\mathrm{GR}}(x;\delta,\theta)^{\beta}}\right)$$
$$= \mathbb{P}\left(\left(\frac{Y}{1+Y}\right)^{1/\beta} \leqslant G_{\mathrm{GR}}(x;\delta,\theta)\right) = \mathbb{P}\left(\left[\theta^{-1}\gamma^{-1}\left(\delta+1;\left(\frac{Y}{1+Y}\right)^{1/\beta}\right)\right]^{1/2} \leqslant x\right),$$
(12)

where $G_{\rm GR}(x; \delta, \theta) = \gamma_1(\delta + 1, \theta x^2), x > 0$. So, we have proved the following:

Proposition 1. The GOLLGR random variable X admits the stochastic representation:

$$X = \left[\theta^{-1} \gamma^{-1} \left(\delta + 1; \left(\frac{Y}{1+Y}\right)^{1/\beta}\right)\right]^{1/2} \quad for \quad Y \sim LL(1, \alpha).$$

2.3 Critical points and modality

For brevity, we define

$$T(x) = T(x; \beta, \delta, \theta) = \frac{G_{\rm GR}(x; \delta, \theta)^{\beta}}{1 - G_{\rm GR}(x; \delta, \theta)^{\beta}}.$$
(13)

Since $T(x/k; \beta, \delta, \theta) = T(x; \beta, \delta, \theta/k^2), k > 0$, the next result follows from (12):

Proposition 2. If $X \sim \text{GOLLGR}(\alpha, \beta, \delta, \theta)$, then $k X \sim \text{GOLLGR}(\alpha, \beta, \delta, \theta/k^2)$.

By differentiating (12) with respect to x, we obtain

$$f(x;\alpha,\beta,\delta,\theta) = f_Y(T(x);1,\alpha)T'(x), \quad Y \sim \mathrm{LL}(1,\alpha).$$
(14)

Then, the derivative of $f(x; \alpha, \beta, \delta, \theta)$ is

$$f'(x;\alpha,\beta,\delta,\theta) = f'_Y(T(x);1,\alpha)[T'(x)]^2 + f_Y(T(x);1,\alpha)T''(x).$$
(15)

Since

$$f'_Y(t;1,\alpha) = -f_Y(t;1,\alpha)r[t]$$
 with $r[t] = \frac{t^{\alpha} + \alpha(t^{\alpha} - 1) + 1}{t(t^{\alpha} + 1)}$

Equation (15) can be expressed as

$$f'(x; \alpha, \beta, \delta, \theta) = f_Y(T(x); 1, \alpha) \{ T''(x) - r[T(x)][T'(x)]^2 \},\$$

where

$$T'(x) = \frac{\beta g_{\rm GR}(x;\delta,\theta)T(x)}{G_{\rm GR}(x;\delta,\theta)[1 - G_{\rm GR}(x;\delta,\theta)^{\beta}]},$$

$$T''(x) = T'(x) \left\{ \frac{g'_{\rm GR}(x;\delta,\theta)}{g_{\rm GR}(x;\delta,\theta)} + g_{\rm GR}(x;\delta,\theta) \frac{(\beta+1)G_{\rm GR}(x;\delta,\theta)^{\beta} + \beta - 1}{G_{\rm GR}(x;\delta,\theta)[1 - G_{\rm GR}(x;\delta,\theta)^{\beta}]} \right\},$$

and

$$\frac{g'_{\rm GR}(x;\delta,\theta)}{g_{\rm GR}(x;\delta,\theta)} = \frac{2\delta + 2\theta x^2 - 1}{x}.$$
(16)

Then,

$$f'(x;\alpha,\beta,\delta,\theta) = f_Y(T(x);1,\alpha)T'(x) \left\{ \frac{g'_{\rm GR}(x;\delta,\theta)}{g_{\rm GR}(x;\delta,\theta)} + \frac{g_{\rm GR}(x;\delta,\theta)}{G_{\rm GR}(x;\delta,\theta)} \left[T(x)+1\right] \left\{ (\beta+1)G_{\rm GR}(x;\delta,\theta)^{\beta} - \alpha\beta \left[\frac{T(x)^{\alpha}-1}{T(x)^{\alpha}+1}\right] - 1 \right\} \right\}$$

Equation (14) gives $f_Y(T(x); 1, \alpha)T'(x) = f(x; \alpha, \beta, \delta, \theta)$, which is a positive function. Hence, any critical point of the pdf of X satisfies the non-linear equation:

$$\frac{g_{\rm GR}'(x;\delta,\theta)}{g_{\rm GR}(x;\delta,\theta)} + \frac{g_{\rm GR}(x;\delta,\theta)}{G_{\rm GR}(x;\delta,\theta)} \left[T(x)+1\right] \left\{ (\beta+1)G_{\rm GR}(x;\delta,\theta)^{\beta} - \alpha\beta \left[\frac{T(x)^{\alpha}-1}{T(x)^{\alpha}+1}\right] - 1 \right\} = 0.$$

The previous result can be written equivalently as:

Proposition 3. A critical point of the pdf of X satisfies

$$\frac{y''}{(y')^2} + \frac{(\beta+1)y^{\beta}[y^{\alpha\beta} + (1-y^{\beta})^{\alpha}] - (\alpha\beta+1)y^{\alpha\beta} - 2(1-y^{\beta})^{\alpha}}{y(1-y^{\beta})[y^{\alpha\beta} + (1-y^{\beta})^{\alpha}]} = 0,$$
(17)

where $y = y(x) = G_{GR}(x; \delta, \theta) = \gamma_1(\delta + 1, \theta x^2).$

In the remainder of this section, we use Equation (17) and the limit in (8) to analyze the modality of the pdf of X when $\alpha = 1$.

For $\alpha = 1$, Equation (17) reduces to

$$\frac{g_{\rm GR}(x;\delta,\theta)}{G_{\rm GR}(x;\delta,\theta)} = \frac{\delta + \theta x^2 - (1/2)}{x}$$

Equivalently,

$$G_{\rm GR}(x;\delta,\theta) = g(x),\tag{18}$$

where

$$g(x) = x g_{\text{GR}}(x; \delta, \theta) = \frac{2\theta^{\delta+1}}{\Gamma(\delta+1)} \frac{x^{2(\delta+1)} e^{-\theta x^2}}{\delta + \theta x^2 - (1/2)}$$

A careful analysis shows that, on $(0, \infty)$, g(x) is decreasing/decreasing-increasing-decreasing when $\delta < 1/2$ or unimodal when $\delta \ge 1/2$, and $\lim_{x\to 0^+} g(x) = \lim_{x\to\infty} g(x) = 0$. Moreover, g(x) has a vertical asymptotic at $x_{va} = \sqrt{[(1/2) - \delta]/\theta}$ when $\delta < 1/2$, with g(x) < 0 for all $x < x_{va}$ and g(x) > 0 for all $x > x_{va}$. Since $G_{\text{GR}}(x; \delta, \theta)$ is increasing on $(0, \infty)$, because this one is a cdf, it is plausible to expect that, by varying the parameters, Equation (18) has at most three roots on $(0, \infty)$. So, the pdf of X has at most three critical points on $(0, \infty)$. In the following, we analyze some possible scenarios:

- If the GOLLGR pdf has a single critical point, say x_0 , and $\beta > 1$ and $\delta \ge 1/2$, we have by the second limit in (8) $\lim_{x\to 0^+} f(x; \alpha, \beta, \delta, \theta) = \lim_{x\to\infty} f(x; \alpha, \beta, \delta, \theta) = 0$. Then, x_0 is really a maximum point, and the pdf of X is unimodal.
- If the GOLLGR pdf has three single critical points, say $x_1 < x_2 < x_3$, and $\beta > 1$ and $\delta \ge 1/2$, again, by the second limit in (8), we have $\lim_{x\to 0^+} f(x; \alpha, \beta, \delta, \theta) = \lim_{x\to\infty} f(x; \alpha, \beta, \delta, \theta) = 0$. Hence, x_1 and x_3 are maximum points, and x_2 is a minimum point, and the GOLLGR pdf is bimodal.

In general, as done previously, we can use the limit in (8) and the number of critical points of the pdf of X to obtain the result:

Theorem 4. The GOLLGR pdf is decreasing/ decreasing-increasing-decreasing/unimodal or bimodal.

2.4 Tail behavior

Here, we prove that, under certain constraints in the parameter space, the distribution of X has thinner tails than an exponential distribution. More precisely, we prove the following two results:

Proposition 5. For any $\alpha \ge 1$ and any t > 0,

$$\lim_{x \to \infty} \frac{\mathrm{e}^{-tx}}{1 - F(x; \alpha, \beta, \delta, \theta)} = \infty.$$
(19)

Proof. If $Y \sim LL(1, \alpha)$, by (12), $F(x; \alpha, \beta, \delta, \theta) = \mathbb{P}(Y \leq T(x))$. Moreover, it is well-known that $\mathbb{P}(Y \leq y) = y^{\alpha}/(1+y^{\alpha})$. Then, from the definition (13) of T, we have (for any t > 0),

$$\frac{\mathrm{e}^{-tx}}{1 - F(x;\alpha,\beta,\delta,\theta)} = \frac{\mathrm{e}^{-tx}}{1 - \mathbb{P}(Y \leqslant T(x))} = \mathrm{e}^{-tx}[1 + T(x)^{\alpha}]$$
$$\geqslant \mathrm{e}^{-tx}T(x)^{\alpha} = \frac{\mathrm{e}^{-tx}G_{\mathrm{GR}}(x;\delta,\theta)^{\alpha\beta}}{\left[1 - G_{\mathrm{GR}}(x;\delta,\theta)^{\beta}\right]^{\alpha}}.$$
(20)

The L'Hôpital's rule yields

$$\lim_{x \to \infty} \frac{\mathrm{e}^{-tx}}{\left[1 - G_{\mathrm{GR}}(x;\delta,\theta)^{\beta}\right]^{\alpha}} = \lim_{x \to \infty} \frac{t\left[\frac{\mathrm{e}^{-tx}}{g_{\mathrm{GR}}(x;\delta,\theta)}\right]}{\alpha\beta\left[1 - G_{\mathrm{GR}}(x;\delta,\theta)^{\beta}\right]^{\alpha-1}G_{\mathrm{GR}}(x;\delta,\theta)^{\beta-1}}.$$
 (21)

Since (for $\alpha \ge 1$),

$$\lim_{x \to \infty} \frac{\mathrm{e}^{-tx}}{g_{\mathrm{GR}}(x;\delta,\theta)} = \left[\frac{2\theta^{\delta+1}}{\Gamma(\delta+1)} x^{2\delta+1} \,\mathrm{e}^{-\theta x^2 + tx}\right]^{-1} = \infty$$

and $\lim_{x\to\infty} G_{\rm GR}(x;\delta,\theta) = 1$. We have from (21)

$$\lim_{x \to \infty} \frac{\mathrm{e}^{-tx}}{\left[1 - G_{\mathrm{GR}}(x; \delta, \theta)^{\beta}\right]^{\alpha}} = \infty$$

By taking $x \to \infty$ for both sides of inequality (20) and by using the above limit, it follows (19). **Proposition 6.** For any $0 < \beta \leq 1$ and $\delta > 0$, the limit (for t > 0) (19) holds.

Proof. By inequality in (20), it is enough to prove

$$\lim_{x \to \infty} \frac{\mathrm{e}^{-tx}}{\left[1 - G_{\mathrm{GR}}(x;\delta,\theta)^{\beta}\right]^{\alpha}} = \lim_{x \to \infty} \frac{1}{\mathrm{e}^{tx} \left[1 - \gamma_1 (\delta + 1, \theta x^2)^{\beta}\right]^{\alpha}} = \infty.$$
(22)

Indeed, by using the C_p inequality:

 $\forall x,y \geq 0; \ (x+y)^p \leqslant C_p(x^p+y^p), \quad \text{where } p > 0 \text{ and } C_p = \max\{1, 2^{p-1}\};$

we have (for $0 < \beta \leq 1$)

$$1 - \gamma_1 (\delta + 1, \theta x^2)^{\beta} \leq [1 - \gamma_1 (\delta + 1, \theta x^2)]^{\beta} = \Gamma_1 (\delta + 1, \theta x^2)^{\beta},$$
(23)

where $\Gamma_1(p,x) = \Gamma(p)^{-1} \int_x^\infty w^{p-1} e^{-w} dw$ is the upper incomplete gamma function ratio.

By using the inequality of Natalini and Palumbo (2000): for a > 1, B > 1 and x > B(a - 1)/(B - 1),

$$\Gamma(a, x) < B x^{a-1} e^{-x}$$

we have (for $x > \sqrt{B\delta/[\theta(B-1)]}$ and $\delta > 0$)

$$\Gamma_1(\delta+1,\theta x^2)^{\beta} < B^{\beta} \,\theta^{\beta\delta} \,\Gamma(\delta+1)^{-\beta} x^{2\beta\delta} \,\mathrm{e}^{-\beta\theta x^2}.$$
(24)

By combining (23) and (24), we obtain (for any $x > \sqrt{B\delta/[\theta(B-1)]}$)

$$e^{tx} \left[1 - \gamma_1 (\delta + 1, \theta x^2)^{\beta} \right]^{\alpha} < B^{\alpha\beta} \theta^{\alpha\beta\delta} \Gamma(\delta + 1)^{-\alpha\beta} x^{2\alpha\beta\delta} e^{-\alpha\beta\theta x^2 + tx}.$$

Letting $x \to \infty$ in the above inequality, we have $e^{tx} [1 - \gamma_1 (\delta + 1, \theta x^2)^{\beta}]^{\alpha}$ tends to zero, proving the limit in (22). Thus, we complete the proof.

2.5 Linear Representation

First, the exponentiated-G ("Exp-G") random variable $W \sim \text{Exp}^c G$ for a continuous cdf G(x)and c > 0, has cdf $H_c(x) = G(x)^c$ and pdf $h_c(x) = c g(x) G(x)^{c-1}$. Many Exp-G properties were published in the last three decades.

We obtain after some algebra the power series

$$G(x)^{\alpha\beta} + \left[1 - G(x)^{\beta}\right]^{\alpha} = \sum_{k=0}^{\infty} c_k G(x)^k , \qquad (25)$$

where $a_k = a_k(\alpha\beta) = \sum_{j=k}^{\infty} (-1)^{j+k} {\alpha\beta \choose j} {j \choose k}$, and

$$c_k = c_k(\alpha, \beta) = a_k(\alpha\beta) + \sum_{i=0}^{\infty} \sum_{j=k}^{\infty} (-1)^{i+j+k} {\alpha \choose i} {i\beta \choose j} {j \choose k}$$

Equation (4) can be rewritten from the ratio of two power series as

$$F(x) = \sum_{k=0}^{\infty} d_k G(x)^k,$$
(26)

where $d_k = d_k(\alpha, \beta)$'s are found found recursively (for k > 0, $d_0 = a_0/c_0$)

$$d_k = c_0^{-1} \left(a_k + \sum_{r=1}^k c_r \, d_{k-r} \right).$$

By differentiating (26) and changing indices

$$f(x) = \sum_{l=0}^{\infty} (l+1) \, d_{l+1} \, g(x) \, G(x)^l$$

For the GR model, we obtain

$$f(x) = \sum_{l=0}^{\infty} (l+1) d_{l+1}(\alpha,\beta) \frac{2\theta^{\delta+1}}{\Gamma(\delta+1)} x^{2\delta+1} e^{-\theta x^2} \gamma_1(\delta+1,\theta x^2)^l,$$
(27)

,

The power series for the incomplete gamma function ratio holds

$$\gamma_1(\delta + 1, \theta x^2) = \frac{(\theta x^2)^{\delta + 1}}{\Gamma(\delta + 1)} \sum_{m=0}^{\infty} \frac{(-1)^m (\theta x^2)^m}{m! (\alpha + 1 + m)}$$

Equation 0.314 in Gradshteyn and Ryzhik (2000) gives (for a natural number $l \ge 1$)

$$\left(\sum_{m=0}^{\infty} q_m x^m\right)^l = \sum_{m=0}^{\infty} e_m^{(l)} x^m$$

where $e_0^{(l)} = q_0^l$, and $e_m^{(l)}$ (for $l \ge 1$) can be found from

$$e_m^{(l)} = \frac{1}{m q_0} \sum_{i=1}^m [(l+1) \, i - m] \, q_i \, e_{m-i}^{(l)}.$$
⁽²⁸⁾

Then,

$$\gamma_1(\delta+1,\theta x^2)^l = \frac{(\theta x^2)^{l(\delta+1)}}{\Gamma(\delta+1)^l} \sum_{m=0}^{\infty} e_m^{(l)} x^{2m} , \qquad (29)$$

where the quantities $e_m^{(l)}$ follow from (28) with the constants

$$q_m = \frac{(-1)^m \theta^m}{(\delta + 1 + m)m!}$$

for m = 0, 1, ...Further, we set the conditions $e_0^{(0)} = 1$ and $e_m^{(0)} = 0$ for $m \ge 1$. Hence, inserting (29) in Equation (27) (under these conditions) yields

$$f(x) = \sum_{l,m=0}^{\infty} \frac{2(l+1)d_{l+1}(\alpha,\beta)e_m^{(l)}}{\Gamma(\delta+1)^{l+1}\theta^m} \theta^{[l(\delta+1)+m+\delta]+1} x^{2[l(\delta+1)+m+\delta]+1} e^{-\theta x^2}$$

and then

$$f(x) = \sum_{l,m=0}^{\infty} w_{l,m} g_{\text{GR}}(x;\theta,\delta_{l,m}^*) , \qquad (30)$$

where $\delta^*_{l,m} = l(\delta + 1) + m + \delta$, and the coefficients are

$$w_{l,m} = w_{l,m}(\theta, \delta, \alpha, \beta) = \frac{(l+1)\Gamma(\delta_{l,m}^* + 1) d_{l+1}(\alpha, \beta) e_m^{(l)}}{\theta^m \Gamma(\delta+1)^{l+1}}$$

Equation (30) is useful to obtain some properties of X from those of the GR model.

2.6 Properties

The sth ordinary moment of X comes from (3) and (30) as

$$E(X^s) = \sum_{l,m=0}^{\infty} w_{l,m}(\theta, \delta, \alpha, \beta) \frac{\Gamma(s/2 + \delta_{l,m}^* + 1)}{\theta^{s/2} \Gamma(\delta_{l,m}^* + 1)}$$

The sth incomplete moment of X follows from (30) as

$$\begin{split} m_{s}(x) &= \sum_{l,m=0}^{\infty} \frac{w_{l,m}(\theta, \delta, \alpha, \beta)}{\Gamma(\delta_{l,m}^{*} + 1)} \int_{0}^{x} 2\theta^{\delta_{l,m}^{*} + 1} t^{s} t^{2\delta_{l,m}^{*} + 1} e^{-\theta t^{2}} dt \\ &= \sum_{l,m=0}^{\infty} w_{l,m}(\theta, \delta, \alpha, \beta) \frac{\Gamma(\delta_{l,m}^{*} + s/2 + 1)}{\Gamma(\delta_{l,m}^{*} + 1)\theta^{s/2}} \int_{0}^{x} \frac{2\theta^{\delta_{l,m}^{*} + s/2 + 1}}{\Gamma(\delta_{l,m}^{*} + s/2 + 1)} t^{2(\delta_{l,m}^{*} + s/2) + 1} e^{-\theta t^{2}} dt \\ &= \sum_{l,m=0}^{\infty} w_{l,m}(\theta, \delta, \alpha, \beta) \frac{\Gamma(s/2 + \delta_{l,m}^{*} + 1)}{\theta^{s/2} \Gamma(\delta_{l,m}^{*} + 1)} \gamma_{1}(\delta_{l,m}^{*} + s/2 + 1, \theta x^{2}) \,. \end{split}$$

The mean deviations and Bonferroni and Lorenz curves of X are obtained from $m_1(x)$. The generating function (gf) of X can be expressed as

$$M(t) = \int_0^\infty e^{tx} \sum_{l,m=0}^\infty w_{l,m}(\theta,\delta,\alpha,\beta) g_{\rm GR}(x;\theta,\delta_{l,m}^*) dx,$$

that is,

$$M(t) = \sum_{l,m=0}^{\infty} \frac{2 w_{l,m}(\theta, \delta, \alpha, \beta) \, \theta^{\delta_{l,m}^* + 1}}{\Gamma(\delta_{l,m}^* + 1)} \int_0^\infty x^{2\delta_{l,m}^* + 1} \, \mathrm{e}^{-\theta x^2 + tx} \, dx \, .$$

From Equation 2.3.15.3 in Prudnikov et al. (1986), we can write

$$\int_0^\infty x^{\alpha-1} \mathrm{e}^{-px^2-qx} dx = \frac{\Gamma(\alpha)}{(2p)^{\alpha/2}} \exp\left(\frac{q^2}{8p}\right) D_{-\alpha}\left(\frac{q}{\sqrt{2p}}\right) \,,$$

where $\operatorname{Re}(\alpha)$, $\operatorname{Re}(p) > 0$, and

$$D_p(y) = \frac{\exp(-y^2/4)}{\Gamma(-p)} \int_0^\infty \exp\{-(wy + w^2/2)\} w^{-(p+1)} dw.$$

Thus,

$$M(t) = \sum_{l,m=0}^{\infty} \frac{2 w_{l,m} \theta^{\delta_{l,m}^*+1}}{\Gamma(\delta_{l,m}^*+1)} \frac{\Gamma(\tilde{\delta}_{l,m})}{(2\theta)^{\tilde{\delta}_{l,m/2}}} \exp\left(\frac{t^2}{8\theta}\right) D_{-\tilde{\delta}_{l,m}}\left(-\frac{t}{\sqrt{2\theta}}\right),$$

where $\tilde{\delta}_{l,m} = 2 \left(\delta^*_{l,m} + 1 \right)$.

2.7 Estimation

We obtain the maximum likelihood estimate (MLE) of $\boldsymbol{\eta} = (\alpha, \beta, \delta, \theta)^{\top}$ given the data x_1, \ldots, x_n from the GOLLGR distribution.

The total log-likelihood function for η is

$$l_{n}(\boldsymbol{\eta}) = n \left[\log(\alpha\beta) + (\delta+1)\log\theta \right] + (2\delta+1) \sum_{i=1}^{n} \log x_{i} - \theta \sum_{i=1}^{n} x_{i}^{2} + (\alpha\beta-1) \sum_{i=1}^{n} \log \gamma_{1}(\delta+1, \theta x_{i}^{2}) + (\alpha-1) \sum_{i=1}^{n} \log[1 - \gamma_{1}(\delta+1, \theta x_{i}^{2})^{\beta}] - 2 \sum_{i=1}^{n} \log \left\{ \gamma_{1}(\delta+1, \theta x_{i}^{2})^{\alpha\beta} + [1 - \gamma_{1}(\delta+1, \theta x_{i}^{2})^{\beta}]^{\alpha} \right\}.$$
(31)

The maximization of (31) can be done using the R software or SAS (PROC NLMIXED), among others.

3 The GOLLGR regression

Recently some papers on regression models have been published, for example, see, Hashimoto *et al.* (2019), Prataviera *et al.* (2020), Silva *et al.* (2020) and Vasconcelos *et al.* (2021). Based on these papers we introduced the regression model based on the GOLLGR distribution.

The systematic components of the GOLLGR regression for X are defined by

$$\theta_i = \exp(\mathbf{v}_i^{\top} \boldsymbol{\lambda}_1)$$
 and $\delta_i = \exp(\mathbf{v}_i^{\top} \boldsymbol{\lambda}_2) - 1$ $i = 1, \dots, n,$ (32)

where $\boldsymbol{\lambda}_1 = (\lambda_{11}, \dots, \lambda_{1p})^{\top}$ and $\boldsymbol{\lambda}_2 = (\lambda_{21}, \dots, \lambda_{2p})^{\top}$ are vectors of unknown coefficients, and $\mathbf{v}_i^{\top} = (v_{i1}, \dots, v_{ip})$ is a vector of known explanatory variables.

The survival function of X comes from (4) as

$$S(x_i|\mathbf{v}) = \frac{[1 - \gamma_1(\delta_i + 1, \theta_i \, x_i^2)^{\beta}]^{\alpha}}{\gamma_1(\delta_i + 1, \theta_i \, x_i^2)^{\alpha\beta} + [1 - \gamma_1(\delta_i + 1, \theta_i \, x_i^2)^{\beta}]^{\alpha}}.$$
(33)

Equation (33) opens new possibilities for fitting different types of regressions. The odd loglogistic GR (OLLGR) regression follows when $\beta = 1$, the exponentiated GR (EGR) regression when $\alpha = 1$, and the GR regression when $\beta = \alpha = 1$.

Let X_i be the lifetime and C_i be the non-informative censoring time (assuming independent),

and $x_i = \min\{X_i, C_i\}$. The total log-likelihood function for $\boldsymbol{\eta} = (a, b, \boldsymbol{\lambda}_1^{\top})^{\top}$ from regression (32) is

$$l(\boldsymbol{\eta}) = r \log\left(\frac{\alpha \beta 2}{\Gamma(\delta_{i}+1)}\right) + (\delta_{i}+1) \sum_{i \in F} \log(\theta_{i}) + (2 \,\delta_{i}+1) \sum_{i \in F} \log(x_{i}) - \sum_{i \in F} \theta_{i} x_{i}^{2} \\ + (\alpha \beta - 1) \sum_{i \in F} \log[\gamma_{1}(\delta_{i}+1,\theta_{i} \, x_{i}^{2})] + (\alpha - 1) \sum_{i \in F} \log[1 - \gamma_{1}(\delta_{i}+1,\theta_{i} \, x_{i}^{2})^{\beta}] \\ - 2 \sum_{i \in F} \log\left\{\gamma_{1}(\delta_{i}+1,\theta_{i} \, x_{i}^{2})^{\alpha \beta} + [1 - \gamma_{1}(\delta_{i}+1,\theta_{i} \, x_{i}^{2})^{\beta}]^{\alpha}\right\} \\ + \sum_{i \in C} \log\left\{\frac{[1 - \gamma_{1}(\delta_{i}+1,\theta_{i} \, x_{i}^{2})^{\alpha \beta} + [1 - \gamma_{1}(\delta_{i}+1,\theta_{i} \, x_{i}^{2})^{\beta}]^{\alpha}}{\gamma_{1}(\delta_{i}+1,\theta_{i} \, x_{i}^{2})^{\alpha \beta} + [1 - \gamma_{1}(\delta_{i}+1,\theta_{i} \, x_{i}^{2})^{\beta}]^{\alpha}}\right\},$$
(34)

where F and C are the sets of observed lifetimes and censoring times, r is the number of uncensored observations (failures). The MLE $\hat{\eta}$ of the vector of unknown parameters can be found by maximizing (34).

3.1 Residual analysis

The quantile residuals (qrs) (Dunn and Smith, 1996) have been adopted frequently in regression applications to verify possible deviations from the model assumptions. For the proposed regression, they are

$$qr_{i} = \Phi^{-1} \left\{ \frac{\gamma_{1}(\hat{\delta}_{i}+1,\hat{\theta}_{i}x_{i}^{2})^{\hat{\alpha}\hat{\beta}}}{\gamma_{1}(\hat{\delta}_{i}+1,\hat{\theta}_{i}x_{i}^{2})^{\hat{\alpha}\hat{\beta}} + [1-\gamma_{1}(\hat{\delta}_{i}+1,\hat{\theta}_{i}x_{i}^{2})^{\hat{\beta}}]^{\hat{\alpha}}} \right\},$$
(35)

where $\Phi^{-1}(\cdot)$ is the standard normal qf.

3.2 Simulations

In this section, two simulation studies are presented by using gamlss package in R software.

The GOLLGR distribution

First, we generate 1,000 samples from Equation (9) with $\alpha = 0.35$, $\beta = 0.55$, $\delta = -0.55$ and $\theta = 0.11$, and sample sizes n = 50, 150 and 500, and calculate the MLEs. The average estimates (AEs), biases and mean squared errors (MSEs) in Table 1 show that the AEs converge to the true values and the biases and MSEs decrease when n increases.

The GOLLGR regression

	n = 50			n = 150			n = 500		
Parameter	AE	Bias	MSE	AE	Bias	MSE	AE	Bias	MSE
α	0.3533	0.0033	0.0205	0.3487	-0.0013	0.0050	0.3488	-0.0012	0.0027
β	0.7140	0.1640	0.3857	0.5861	0.0361	0.0197	0.5688	0.0188	0.0095
δ	-0.5217	0.0283	0.1054	-0.5512	-0.0012	0.0044	-0.5514	-0.0014	0.0018
σ	0.1484	0.0384	0.0115	0.1188	0.0088	0.0015	0.1148	0.0048	0.0008

Table 1: Simulation results from the GOLLGR distribution.

The second study examines the accuracy of the MLEs in the proposed regression. The observations are generated from $X_i \sim \text{GOLLGR}(\alpha, \beta, \delta_i, \theta_i)$ and $v_{1i} \sim \text{Binomial}(1, 0.5)$, where v_{1i} is taken in two groups (0 and 1). We consider 1,000 samples for $\alpha = 0.37$, $\beta = 0.61$, $\lambda_{10} = 0.55$, $\lambda_{11} = 1.75$, $\lambda_{20} = 0.65$ and $\lambda_{21} = 2.75$ and n = 150, 350 and 650, and the simulation process follows as:

(i) Generate $v_{1i} \sim \text{Binomial}(1, 0.5);$

(ii) Calculate δ_i and θ_i from the systematic components: $\delta_i = \exp(\lambda_{10} + \lambda_{11}v_{1i}) - 1$ and $\theta_i = \exp(\lambda_{20} + \lambda_{21}v_{1i})$, respectively;

(iii) Generate $u_i \sim U(0, 1)$;

(iv) The previous steps yield x_i 's from (9).

The numbers in Table 2 reveal that the AEs converge to the true values, and the biases and MSEs decrease when n increases, thus indicating the consistency of the estimators.

	n = 150			n = 350			n = 650		
Parameters	AE	Bias	MSE	AE	Bias	MSE	AE	Bias	MSE
α	0.3736	0.0036	0.0146	0.3700	0.0000	0.0055	0.3722	0.0022	0.0028
β	0.7324	0.1224	0.2991	0.6615	0.0515	0.0897	0.6317	0.0217	0.0332
λ_{10}	0.5917	0.0417	0.2022	0.5787	0.0287	0.0973	0.5618	0.0118	0.0467
λ_{11}	1.7715	0.0215	0.0535	1.7586	0.0086	0.0219	1.7553	0.0053	0.0114
λ_{20}	0.7308	0.0808	0.1444	0.6955	0.0455	0.0606	0.6640	0.0140	0.0307
λ_{21}	2.7400	-0.0100	0.0582	2.7450	-0.0050	0.0249	2.7518	0.0018	0.0127

Table 2: Simulation results from the GOLLGR regression.

4 Applications

Here, we compare the GOLLGR model and its GR, EGR, OLLGR sub-models in two applications. We determine the MLEs, and the criteria: Akaike Information Criterion (AIC), Consistent Akaike, Information Criterion (CAIC), and Bayesian Information Criterion (BIC).

4.1 Application 1: Voltage data

We consider the times of failure and running times for a field-tracking study (Meeker and Escobar, 1998). We fit the Rayleigh distribution ($\alpha = \beta = 1$) to find initial values for θ and δ . All computations are done through the NLMIXED subroutine in SAS. Table 3 lists the MLEs (their standard errors in parentheses), and the previous measures, which reveal that the GOLLG distribution can be chosen as the best model.

Model	α	β	δ	θ	AIC	CAIC	BIC
GOLLGR	0.0437	0.4611	34.6605	0.0011	362.0	363.6	367.6
	(0.0111)	(01515)	(0.00002)	(3.958E-6)			
OLLGR	0.0246	1	37.0100	0.0010	383.2	384.1	387.4
	(0.0041)		(0.0070)	(0.00006)			
EGR	1	0.0133	35.3427	0.00019	366.5	367.4	370.7
		(0.0014)	(0.0051)	(0.00003)			
GR	1	1	-0.5079	0.000011	368.4	368.8	371.2
			(0.1147)	(5.1E-6)			

Table 3: Findings for voltage data.

The likelihood ratio (LR) statistics in Table 4 indicate that the GOLLGR distribution is the best model among the others. The histogram and the plots of the estimated densities in Figure 3(a), and those of the empirical and estimated survival functions in Figure 3(b), support the previous conclusion.

Table 4: LR statistics for voltage data.

Model	Hypotheses	LR statistic	<i>p</i> -value
GOLLGR vs OLLGR	$H_0: \beta = 1 \text{ vs } H_1: H_0 \text{ is false}$	23.2	< 0.00001
GOLLGR vs EGR	$H_0: \alpha = 1$ vs $H_1: H_0$ is false	6.5	0.01079
GOLLGR vs GR	$H_0: \beta = \alpha = 1$ vs $H_1: H_0$ is false	10.4	0.0055

4.2 Application 2: COVID-19 data

The second application refers to lifetimes of individuals diagnosed with COVID-19 (Coronavirus Disease 1999) (Galvão and Roncalli, 2021). Since it was declared an international health emergency, many studies have been conducted to obtain information about the clinical, epidemiological and prognostic aspects of the disease; see, for example, Cordeiro *et al.* (2021a), Cordeiro *et al.* (2021b) and Marinho *et al.* (2021).



Figure 3: Voltage data: (a) Estimated densities. (b) Estimated survival functions and the empirical survival.

In Brazil, the epidemiological data are disclosed by the Health Information System (available in: https://opendatasus.saude.gov.br/en/dataset/srag-2021-e-2022. In this analysis, we work with the gamlss package of R.

In this study, 881 patients infected by the virus are considered, confirmed by the RT-PCR test method. The participants consisted of hospitalized patients and outpatients living in the city of Campinas (Brazil) in January and February 2021. The survival consisted of the interval between the first symptoms until the date of death due to COVID-19 (failure). Deaths due to other causes or after the is 73.6%. Equation (32) is considered with factors associated with the highest risk of death. The results are compared with the OLLGR, EGR and GR sub-regressions.

The following variables were considered for each patient (i = 1, ..., 881):

- x_i : time until death due to COVID-19 (in days);
- cens_i : $\operatorname{censoring indicator} (0 = \operatorname{censored}, 1 = \operatorname{observed lifetime});$
- v_{i1} : age (in years);
- v_{i2} : diabetes mellitus (0= no or not reported, 1= yes).

The total number of patients suffering from the comorbidity diabetes was 264 (29.97%), among whom 104 (39.39%) died. In turn, of the 617 patients (70.03%) without the disease or who did

not report it, 128 (20.75%) died. Figure 4 presents the Kaplan-Meier survival curve, showing the greater risk of death among patients suffering from diabetes.



Figure 4: Kaplan-Meier survival curve for the variable diabetes mellitus (1 = yes, 0 = no or not informed).

The statistics in Table 5 support that the GOLLGR regression can be chosen as the best model. Further, the LR statistics in Table 6 indicate that the wider regression yields the best fit. Table 7 reports the MLEs (SEs in parentheses) from the fitted GOLLGR regression.

Model	AIC	BIC	CAIC
GOLLGR	2222.54	2260.78	2237.66
OLLGR	2237.73	2271.19	2262.63
EGR	2240.17	2273.63	2265.07
GR	2238.66	2267.34	2273.34

Table 5: Findings from the fitted regressions to COVID-19 data.

Figure 5 provides the graphs of the quantile residuals (qrs) (35). The residual index plot (Figure 5a) reveals that the qrs have a random behavior and that only four observations are outside the [-3,3] range. The normal probability plot for the qrs (Figure 5b) indicates that the residuals

Model	Hypotheses	LR statistic	<i>p</i> -value
GOLLGR vs OLLGR	$H_0: \beta = 1 \text{ vs } H_1: H_0 \text{ is false}$	17.1915	< 0.00001
GOLLGR vs EGR	$H_0: \alpha = 1$ vs $H_1: H_0$ is false	19.6298	< 0.00001
GOLLGR vs GR	$H_0: \beta = \alpha = 1$ vs $H_1: H_0$ is false	20.1202	< 0.00001

Table 6: LR statistics for COVID-19 data.

Table 7: Results from the fitted GOLLGR regressions to COVID-19 data.

	MLE	SE	p-value
λ_{10}	-0.3599	0.0289	< 0.0001
λ_{11}	-0.0028	0.0006	< 0.0001
λ_{12}	0.0468	0.0359	0.1920
λ_{20}	-10.1148	0.1176	< 0.0001
λ_{21}	0.0480	0.0022	< 0.0001
λ_{22}	0.3331	0.0774	< 0.0001
$\log(\alpha)$	-0.9748	0.0130	
$\log(\beta)$	2.0127	0.0130	

follow approximately a normal distribution, which support the fitted regression. Thus, there is no evidence against the GOLLGR regression assumptions.



Figure 5: COVID-19 data: (a) Index plot of the qrs. (b) Normal probability plot of the qrs.

Some interpretations are in order. Table 7 shows that the covariable age is significant, meaning that older individuals tend to have a progressively shorter period until death due to this coronavirus. It is noted a significant difference between individuals with and without diabetes mellitus in relation to the time until death by COVID-19.

5 Conclusions

There is a clear need for extended well-known distributions and their successful applications in several areas. The Rayleigh distribution plays a crucial role in modelling and analyzing lifetime data, and several extensions of this distribution have been published in recent years. We constructed a new regression based on the four-parameter extended Rayleigh distribution, and showed its utility in the analysis of lifetime data.

References

- Cordeiro, G.M., Alizadeh, M., Ozel, G., Hosseini, B., Ortega, E.M.M. and Altun, E. (2017) The generalized odd log-logistic family of distributions: properties, regression models and applications. *Journal of Statistical Computation and Simulation*, 87, 908-932.
- [2] Cordeiro, G.M., Figueiredo, D., Silva, L., Ortega, E.M.M., and Prataviera, F. (2021a) Explaining COVID-19 mortality rates in the first wave in Europe. *Model Assisted Statistics and Applications*, 16, 211-221.
- [3] Cordeiro, G.M., de Azevedo Cysneiro, F.J., and Cabral, P.C. (2021b) Estatísticas Básicas e Modelagem de Regressão das taxas de mortalidade por COVID-19 nos Estados Brasileiros. Brazilian Journal of Development, 72, 117735-117749.
- [4] Dunn, P., and Smyth, G. (1996) Randomized quantile residuals. Journal of Computational and Graphical Statistics, 5, 236–44.
- [5] Galvão, M.H.R., and Roncalli, A.G. (2021) Factors associated with increased risk of death from COVID-19: survival analysis based on confirmed cases. *Brazilian Journal of Epidemiology*, 23, 1-10.
- [6] Gleaton, J.U. and Lynch, J.D. (2006) Properties of generalized log-logistic families of lifetime distributions. *Journal of Probability and Statistical Science*, 4, 51-64.
- [7] Gradshteyn, I.S. and Ryzhik, I.M. (2000) Table of integrals, series, and products. San Diego, Academic Press.
- [8] Gupta, R.C. and Gupta, R.D. (2007) Proportional reversed hazard rate model and its applications. Journal of Statistical Planning and Inference, 137, 3525-3536.

- [9] Hashimoto, E.M., Silva, G.O., Ortega, E.M.M. and Cordeiro, G.M. (2019). Log-Burr XII gamma-Weibull regression model with andom effects and censored data. Journal of Statisyical Theory and Practice, 13, 1-18.
- [10] Johnson, N.L., Kotz, S. and Balakrishnan, N. (1994) Continuous univariate distributions. New York, John Wiley and Sons.
- [11] Marinho, P.R.D., Cordeiro, G.M., Coelho, H.F. and Brandão, S.C.S. (2021) Covid-19 in Brazil: A sad scenario. Cytokine & growth factor reviews, 58, 51-54.
- [12] Meeker, W.Q. and Escobar, L.A. (1998) Statistical Methods for Reliability Data. Wiley, New York.
- [13] Natalini, P. and Palumbo, B. (2000) Inequalities for the incomplete gamma function. Mathematical Inequalities & Applications, 3, 69-77.
- [14] Prataviera, F., Ortega, E.M.M., Gauss Cordeiro, G.M., Pescim, R.R and Verssani, B.A.W. (2018) A new generalized odd log-logistic flexible Weibull regression model with applications in repairable systems. *Reliability Engineering & System Safety*, **176**: 13-26.
- [15] Prataviera, F., Loibel, S.M.C., Grego, K.F., Ortega, E.M.M. and Cordeiro, G.M. (2020). Modelling non-proportional hazard for survival data with different systematic components. *Environmental and Ecological Statistics*, 27, 467-489.
- [16] Prudnikov, A.P., Brychkov, Y.A. and Marichev, O.I. (1986) Integrals and series. New York, Gordon and Breach Science Publishers.
- [17] Silva, G.O., Cordeiro, G.M. and Ortega, E.M.M. (2020). Surviving and non surviving fraction regression models based on the beta modified Weibull distribution. *Model Assisted Statistics* and Applications, 15, 111-126.
- [18] Vasconcelos, J.C.S., Cordeiro, G.M., Ortega, E.M.M. and Ribeiro, J.G. (2021). A regression model for extreme events and the presence of bimodality with application in energy generation data. *IET Renewable Power Generation*, 48, p. 452-461, 2021.
- [19] Vodă, V.G. (1976) Inferential procedures on a generalized Rayleigh variate, I. Applications of Mathematics, 21, 395-412.