

# Simulating dynamical phases of chiral $p + ip$ superconductors with a trapped ion magnet

Athreya Shankar,<sup>1,2,\*</sup> Emil A. Yuzbashyan,<sup>3</sup> Victor Gurarie,<sup>4,5</sup>  
Peter Zoller,<sup>1,2</sup> John J. Bollinger,<sup>6</sup> and Ana Maria Rey<sup>7,5,†</sup>

<sup>1</sup>*Institute for Theoretical Physics, University of Innsbruck, Innsbruck, Austria*

<sup>2</sup>*Institute for Quantum Optics and Quantum Information of the Austrian Academy of Sciences, Innsbruck, Austria*

<sup>3</sup>*Center for Materials Theory, Rutgers University, Piscataway, New Jersey 08854, USA*

<sup>4</sup>*Department of Physics, University of Colorado, Boulder, CO 80309*

<sup>5</sup>*Center for Theory of Quantum Matter, University of Colorado, Boulder, CO 80309*

<sup>6</sup>*National Institute of Standards and Technology, Boulder, CO 80309*

<sup>7</sup>*JILA, National Institute of Standards and Technology,  
and Department of Physics, University of Colorado, Boulder, CO 80309*

(Dated: April 13, 2022)

Two-dimensional  $p + ip$  superconductors and superfluids are systems that feature chiral behavior emerging from the Cooper pairing of electrons or neutral fermionic atoms with non-zero angular momentum. Their realization has been a longstanding goal because they offer great potential utility for quantum computation and memory. However, they have so far eluded experimental observation both in solid state systems as well as in ultracold quantum gases. Here, we propose to leverage the tremendous control offered by rotating two-dimensional trapped-ion crystals in a Penning trap to simulate the dynamical phases of two-dimensional  $p + ip$  superfluids. This is accomplished by mapping the presence or absence of a Cooper pair into an effective spin-1/2 system encoded in the ions' electronic levels. We show how to infer the topological properties of the dynamical phases, and discuss the role of beyond mean-field corrections. More broadly, our work opens the door to use trapped ion systems to explore exotic models of topological superconductivity and also paves the way to generate and manipulate skyrmionic spin textures in these platforms.

## I. INTRODUCTION

The observation and classification of dynamical behaviors in quantum many-body systems constitute a core milestone in quantum science. One fascinating and promising paradigm comprises the dynamical phases predicted to emerge from quenches of superconductors and superfluids [1–3], systems that feature Cooper pairing of electrons or neutral fermionic atoms. In particular, topological  $p + ip$  Bardeen–Cooper–Schrieffer superconductors (in charged electrons) or superfluids (in neutral atoms)—systems that feature non-trivial topological properties [4] and gapless, chiral edge states that circulate around the boundary—are especially exciting given their potential use for topological quantum computation. Despite intensive theoretical efforts,  $p + ip$  superfluids have eluded experimental observation, with the only confirmed realization being the A-phase of  $^3\text{He}$ , which ironically is one of the oldest-known superfluids but is also hard to control and manipulate. Ultracold fermionic quantum gases are currently the leading platform for quantum simulation of strongly correlated matter. However, in spite of all their attractive features, the control and manipulation of  $p$ -wave interactions in these systems has remained a challenge since  $p$ -wave interactions are weak under standard conditions and require Feshbach resonances to enhance

them. The latter unfortunately introduce strong three-body processes which make the gas unstable and destroy the desired pairing processes [5–9], although schemes circumventing this problem have been proposed [10–12]. Theory proposals have also suggested the observation of topological superfluids by suddenly bringing weakly interacting atoms close to a Feshbach resonance [13], but to date experimental efforts remain unsuccessful.

In the present work, we propose a pathway towards the observation of non-equilibrium dynamical phases of topological  $p + ip$  superfluids by using a 2D crystal of ions in a Penning trap. This platform offers a high degree of control and flexibility in state initialization, interaction control and readout, that have been previously leveraged for the sensing of weak electric fields and for the simulation of quantum magnets [14, 15]. In this system, we propose to encode a spin-1/2 degree of freedom in two electronic states of the ions, which, via the Anderson pseudospin mapping [16], are used to simulate the presence or the absence of a Cooper pair.

Our proposal takes advantage of the fact that the ion crystal in a Penning trap is rotating in the lab frame [17]. This feature has never before been exploited in the context of quantum simulation and, in fact, it is often viewed as an impediment, e.g., to perform single site addressing. We show that by tuning the orientation and parameters of the laser beams that are typically used to couple the electronic and motional degrees of freedom of the crystal, we can engineer controllable effective interactions that simulate the Hamiltonian of a  $p + ip$  superfluid. Tuning the laser parameters also allows us to i) prepare initial

\* athreya.shankar@uibk.ac.at

† arey@jilaui1.colorado.edu




FIG. 1. **Probing dynamical phases of  $p + ip$  superfluids using ion crystals in Penning traps.** (a) The fermionic model is mapped on to spins encoded in the internal states of the ions, where spin up (down) represents the absence (presence) of a Cooper pair (inverted Anderson pseudospin mapping). Here we show representative spin textures that can be engineered for the topologically nontrivial BCS and trivial BEC phases. (b) The dynamical phases are classified according to the long time behavior of the magnitude of an order parameter— Phase I:  $|\Psi(t)| \rightarrow 0$ ; Phase II:  $|\Psi(t)| \rightarrow$  non-zero constant; Phase III:  $|\Psi(t)|$  displays persistent oscillations. (c) Schematic of our experimental proposal. State initialization,  $p$ -wave interactions and readout are all achieved using appropriate parameters for a pair of optical dipole force (ODF) lasers and a pair of co-propagating Raman lasers. In contrast to prior implementations, the ODF difference wavevector  $\delta\mathbf{k}$  has both an out-of-plane and in-plane component (see side view). The result is a tilted traveling wave lattice that crosses the crystal plane slightly obliquely and thus couples the ions' electronic degrees of freedom, the out-of-plane center-of-mass mode and the in-plane crystal rotation.

states that resemble the low energy conditions of a  $p$ -wave superfluid, ii) control the relative strength between the kinetic energy and pairing interaction terms required to observe the three different mean-field dynamical phases predicted to exist in  $p \pm ip$  superfluids [4] and iii) measure a superconducting order parameter for classifying the dynamical phases. Moreover, since state-of-the-art ion crystals are not in the thermodynamic limit but are instead limited to  $\lesssim 500$  ions, they naturally open a path to explore modifications to the non-equilibrium dynamics arising from beyond-mean-field effects.

A key appeal of  $p + ip$  superfluids compared to ordinary superfluids is the possibility of featuring states with nontrivial topological order. In an ordinary superfluid or superconductor, the BCS and the BEC regimes—which respectively favor weakly bound Cooper pairs and a Bose-Einstein Condensate of tightly bound molecules made of two fermions—are continuously connected and are only distinguished by the strength of the pairing. In contrast, the two regimes exhibit different topological behaviors in 2D  $p + ip$  superfluids, with a genuine quantum phase transition separating the topologically nontrivial BCS phase from the topologically trivial BEC phase in the equilibrium situation. This feature extends into the non-equilibrium regime, where the dynamical phases exhibit a dynamical topological quantum phase transition [4]. Here, we show how to engineer both topologically trivial and nontrivial dynamical phases in our system and demonstrate how their topological character can be distinguished by inferring an appropriate winding number and additionally confirmed by measurements of the effective Cooper pair distribution function.

## II. 2D $p + ip$ SUPERFLUIDS

The Hamiltonian that governs the low energy sector of  $p + ip$  superconductors is given (setting  $\hbar = 1$ ) by

$$\hat{H} = \sum_{\mathbf{p}} \frac{p^2}{2m} \hat{c}_{\mathbf{p}}^\dagger \hat{c}_{\mathbf{p}} - \frac{G}{2m} \sum_{\mathbf{p}, \mathbf{q}} \mathbf{p} \cdot \mathbf{q} \hat{c}_{\mathbf{p}}^\dagger \hat{c}_{-\mathbf{p}}^\dagger \hat{c}_{-\mathbf{q}} \hat{c}_{\mathbf{q}}. \quad (1)$$

Here,  $m$  is the electron mass and  $\hat{c}_{\mathbf{p}}^\dagger, \hat{c}_{\mathbf{p}}$  are fermionic creation and annihilation operators for a fermion with momentum  $\mathbf{p}$ . The first term describes the single particle dispersion and the second term the attractive ( $G > 0$ )  $p$ -wave interactions that lead to the formation of Cooper pairs. For the case of 2D  $p + ip$  superconductors, the momentum  $\mathbf{p} \equiv p_x \hat{\mathbf{e}}_x + p_y \hat{\mathbf{e}}_y$  is restricted to the  $x - y$  plane. This Hamiltonian assumes that Cooper pairs are only created and destroyed with zero center-of-mass momentum and neglects pair-breaking processes.

Under these conditions, the low-energy physics can be mapped on to the dynamics of a collection of interacting spin-1/2 systems via the Anderson pseudospin mapping that introduces spin-1/2 operators at each momentum  $\mathbf{p}$ :

$$\begin{aligned} 2\hat{s}_{\mathbf{p}}^Z &= \hat{c}_{\mathbf{p}}^\dagger \hat{c}_{\mathbf{p}} + \hat{c}_{-\mathbf{p}}^\dagger \hat{c}_{-\mathbf{p}} - 1, \\ \hat{s}_{\mathbf{p}}^+ &= \hat{c}_{\mathbf{p}}^\dagger \hat{c}_{-\mathbf{p}}^\dagger, \quad \hat{s}_{\mathbf{p}}^- = \hat{c}_{-\mathbf{p}} \hat{c}_{\mathbf{p}}. \end{aligned} \quad (2)$$

Here, the presence or absence of a Cooper pair at momentum  $\mathbf{p}$  corresponds to the eigenstates  $|\uparrow\rangle_{\mathbf{p}}, |\downarrow\rangle_{\mathbf{p}}$  of  $\hat{s}_{\mathbf{p}}^Z$  with eigenvalues  $\pm 1/2$  respectively, and the raising and lowering operators  $\hat{s}_{\mathbf{p}}^\pm$  describe the creation and annihilation of this Cooper pair. In terms of the Anderson pseudospin operators, Hamiltonian (1) can be expressed

as

$$\hat{H} = \sum_{\mathbf{p}}' \frac{p^2}{m} \hat{s}_{\mathbf{p}}^Z - \frac{2G}{m} \sum_{\mathbf{p}, \mathbf{q}}' \mathbf{p} \cdot \mathbf{q} \hat{s}_{\mathbf{p}}^+ \hat{s}_{\mathbf{q}}^- . \quad (3)$$

The primed sums indicate that in the spin model, the summation is only over  $\mathbf{p}, \mathbf{q}$  in one half-plane, say  $p_y, q_y \geq 0$ , so that all the spin operators are independent.

The ground state of the spin model (3) possesses the property that the spin orientation is correlated with the azimuthal angle  $\phi_{\mathbf{p}}$  in momentum space, giving rise to chiral spin textures as depicted in Fig. 1(a). A winding number  $Q$  can be ascribed to the spin texture, based on which the state can be classified as belonging to a topologically non-trivial BCS phase ( $Q = 1$ ) or a topologically trivial BEC phase ( $Q = 0$ ) phase. Physically, assuming spins at large momenta are always held fixed in  $|\downarrow\rangle$ , the spin texture is topologically non-trivial if the central spin is in  $|\uparrow\rangle$  whereas it is trivial if the central spin is in  $|\downarrow\rangle$ .

The quench dynamics of  $p + ip$  superconductors were theoretically studied [4] by considering a chiral variant of the spin model (3) given by

$$\hat{H} = \sum_{\mathbf{p}}' \frac{p^2}{m} \hat{s}_{\mathbf{p}}^Z - \frac{G}{m} \sum_{\mathbf{p}, \mathbf{q}}' p q e^{-i(\phi_{\mathbf{p}} - \phi_{\mathbf{q}})} \hat{s}_{\mathbf{p}}^+ \hat{s}_{\mathbf{q}}^- , \quad (4)$$

where  $p$  and  $\phi_{\mathbf{p}}$  are the magnitude and azimuthal angle for the 2D momentum  $\mathbf{p}$ . This Hamiltonian breaks time-reversal symmetry explicitly, and preferentially selects  $p_x - ip_y$  over  $p_x + ip_y$  pairing, which are degenerate in the time-reversal-invariant Hamiltonian (1). Nevertheless both Hamiltonians possess the same  $p + ip$  ground state and dynamical phases in the thermodynamic limit. Therefore, in this work, we focus on the quantum simulation of the chiral model (4).

In the thermodynamic limit, where mean-field theory is exact, the dynamics can be pictured as each spin precessing about a local magnetic field:

$$\frac{d}{dt} \langle \hat{s}_{\mathbf{p}} \rangle = \langle \hat{s}_{\mathbf{p}} \rangle \times \mathbf{B}_{\mathbf{p}} . \quad (5)$$

Here,  $\langle \hat{s}_{\mathbf{p}} \rangle$  is the expectation value of the spin vector at momentum  $\mathbf{p}$  and  $\mathbf{B}_{\mathbf{p}}$  is the local magnetic field with components

$$\begin{aligned} B_{\mathbf{p}}^X &= -p \cos(\phi_{\mathbf{p}}) \text{Re}[\Psi] - p \sin(\phi_{\mathbf{p}}) \text{Im}[\Psi] , \\ B_{\mathbf{p}}^Y &= p \cos(\phi_{\mathbf{p}}) \text{Im}[\Psi] - p \sin(\phi_{\mathbf{p}}) \text{Re}[\Psi] , \\ B_{\mathbf{p}}^Z &= -\frac{p^2}{m} \end{aligned} \quad (6)$$

written in terms of an order parameter  $\Psi$  given by

$$\Psi(t) = -\frac{2G}{m} \sum_{\mathbf{p}}' p e^{i\phi_{\mathbf{p}}} \langle \hat{s}_{\mathbf{p}}^-(t) \rangle . \quad (7)$$

Mean-field theory predicts the emergence of three dynamical phases when the system is initialized in its ground state and the pairing strength  $G$  is quenched.

They have been classified according to the long-time behavior of  $|\Psi(t)|$  as illustrated in Fig. 1(b), where we plot a dimensionless and normalized version of the order parameter, [see Eq. (12)]. In Phase I, the single-particle kinetic energy term ( $B_{\mathbf{p}}^Z$ ) dominates and  $|\Psi(t)| \rightarrow 0$ . In phases II and III interactions instead stabilize a finite order parameter. In Phase II,  $|\Psi(t)|$  tends to a non-zero constant value, while in Phase III, also known as a self-generated Floquet phase [18]),  $|\Psi(t)|$  features persistent oscillations.

The topological properties of the dynamical phases are best understood in terms of a second winding number  $W$ . Although this quantity is formally defined in terms of retarded single-particle Green functions[4], it can be physically interpreted in Phase II as the winding of the magnetic field texture in an appropriate rotating frame. In the thermodynamic limit, mean-field theory predicts that the long-time order parameter in phase II can be written as

$$\Psi(t) = \Psi_{\infty} e^{-2i\mu_{\infty} t} , \quad (8)$$

where  $\Psi_{\infty}$  is the magnitude in the limit  $t \rightarrow \infty$  and  $\mu_{\infty}$  is a dynamical chemical potential. In a frame rotating at  $2\mu_{\infty}$ , the spins precess under a static effective magnetic field  $\bar{\mathbf{B}}_{\mathbf{p}} \equiv \mathbf{B}_{\mathbf{p}} + 2\mu_{\infty} \hat{\mathbf{e}}_Z$  whose texture can be analogous to the spin texture in a  $p + ip$  ground state. The winding number  $W$  is computed as

$$W = \frac{1}{4\pi} \int dp_x dp_y \hat{\mathbf{B}}_{\mathbf{p}} \cdot \left( \frac{d\hat{\mathbf{B}}_{\mathbf{p}}}{dp_x} \times \frac{d\hat{\mathbf{B}}_{\mathbf{p}}}{dp_y} \right) , \quad (9)$$

where  $\hat{\mathbf{B}}_{\mathbf{p}}$  denotes the corresponding unit vector in the rotating frame. While the spins at large momenta  $p \rightarrow \infty$  experience a field  $\bar{\mathbf{B}}_{\mathbf{p}} \approx -(p^2/m)\hat{\mathbf{e}}_Z$  that points down, the central spin at  $\mathbf{p} = \mathbf{0}$  is isolated from the other spins and experiences an effective magnetic field  $\bar{\mathbf{B}}_{\mathbf{0}} = 2\mu_{\infty} \hat{\mathbf{e}}_Z$ . Therefore, the magnetic field texture is BCS-like and topologically nontrivial ( $W = 1$ ) for  $\mu_{\infty} > 0$  ( $\bar{\mathbf{B}}_{\mathbf{0}}$  pointing up) while it is BEC-like and topologically trivial ( $W = 0$ ) for  $\mu_{\infty} < 0$  ( $\bar{\mathbf{B}}_{\mathbf{0}}$  pointing down).

### III. IMPLEMENTATION WITH PENNING TRAPS

We now discuss how the spin model (3) can be simulated with ion crystals in a Penning trap, where the pseudospin-1/2 system is encoded in two long-lived hyperfine states of each trapped ion. In this trap, ions self-organize into a planar crystal with an approximate triangular lattice structure under the influence of static trapping fields. An electric quadrupole field  $\mathcal{E}$  accomplishes axial trapping and confines the ions to a single plane. The addition of a strong axial magnetic field  $\mathcal{B}$  leads to an  $\mathcal{E} \times \mathcal{B}$  drift of the ions in this plane. This rotation provides radial confinement and the corresponding rotation frequency  $\omega_r$  can be precisely controlled by additional electrodes. The out-of-plane motion of a crystal

of  $N$  ions is described using  $N$  normal modes of vibration, called the drumhead modes. The highest frequency drumhead mode is the center-of-mass (c.m.) mode, which is well separated from the rest of the modes and hence can be well resolved.

In our modeling, the drumhead c.m. mode is treated quantum mechanically and described by bosonic creation and annihilation operators  $\hat{a}_1^\dagger, \hat{a}_1$ . On the other hand, the planar motion is dominated by the crystal rotation, and is hence treated classically with the  $x_j$  and  $y_j$  coordinates of ion  $j$  undergoing uniform circular motion at radius  $r_j$  from the trap center, with frequency  $\omega_r$  and azimuthal phase offset  $\phi_j$ .

As a first step to realize Hamiltonian (4), we engineer a Jaynes-Cummings type interaction between each spin and the drumhead c.m. mode, with the coupling depending on the planar position of the ion as viewed in the crystal rotating frame. As we explain shortly, the Hamiltonian we engineer is given by

$$\hat{H}_{2\text{ch}} = \sum_{j=1}^N B_1 \tilde{r}_j^2 \hat{\mathbf{s}}_j^Z + \delta_1 \hat{a}_1^\dagger \hat{a}_1 - \sum_{j=1}^N \frac{G}{i\sqrt{N}} \tilde{r}_j \left( \hat{\mathbf{s}}_j^- \hat{a}_1^\dagger e^{i\phi_j} - \hat{a}_1 \hat{\mathbf{s}}_j^+ e^{-i\phi_j} \right). \quad (10)$$

This Hamiltonian is written in a rotated spin space  $\mathcal{Z} \equiv -X$ ,  $\mathcal{X} \equiv Z$ ,  $\mathcal{Y} \equiv Y$ , and  $\hat{\mathbf{s}}_j$  denotes spin operators in this rotated space. Here,  $\tilde{r}_j = r_j/R$  is a normalized radial coordinate,  $B_1$  is a frequency controlling the dispersion of the spins,  $\delta_1$  is an effective detuning of the c.m. mode from the spins and  $G$  is the spin-mode coupling strength. In particular, the amplitude and phase of the coupling of spin  $j$  to the c.m. mode respectively depend on  $\tilde{r}_j$  and  $\phi_j$ .

Equation (10) describes the Hamiltonian for the so-called two-channel model of a  $p$ -wave superconductor. The c.m. mode plays the role of the bosonic molecular channel, while each ion encodes an Anderson pseudospin in its electronic states. Here, spin up (down) indicates the presence (absence) of a Cooper pair. While the Anderson pseudospins live in a lattice in momentum space where the coordinates are  $(p_x, p_y)$ , the role of momentum is instead played here by the position  $(x_j, y_j)$  of each ion  $j$  in the crystal plane. ‘Momentum’-dependent rates appear in the single-particle and interaction terms through the radius  $r_j$  and the phase factors  $e^{\pm i\phi_j}$ .

Subsequently, an effective spin model can be derived in the situation when  $\delta_1 \gg G, B_1$ . Using effective Hamiltonian theory [19] and assuming that the c.m. mode is in the motional ground state, we obtain the one-channel model given by

$$\hat{H}_{1\text{ch}} = K \sum_j \tilde{r}_j^2 \hat{\mathbf{s}}_j^Z - \frac{J}{N} \sum_{j \neq k} \tilde{r}_j \tilde{r}_k \hat{\mathbf{s}}_j^+ \hat{\mathbf{s}}_k^- e^{-i(\phi_j - \phi_k)}, \quad (11)$$

where  $J = G^2/\delta_1$  and  $K = B_1 - J/N$ . Equation (11) is essentially the one-channel  $p$ -wave Hamiltonian (4) that we wish to simulate.

We now briefly outline how Hamiltonian (10) can be engineered while presenting the detailed derivation in Appendix A. Coupling between the spins and the crystal motion is enabled by the application of an optical dipole force (ODF) that gives rise to spatially dependent AC Stark shifts on the spin states. The ODF is generated by two traveling-wave lasers with difference wavevector  $\Delta\mathbf{k}$  and beatnote frequency  $\mu_r$ . In typical applications, only the axial motion is coupled to the spin and hence  $\Delta\mathbf{k} \parallel \hat{z}$ . However, in this work, we consider the  $\Delta\mathbf{k}$  to have non-zero components both along the  $\hat{z}$  and  $\hat{x}$  directions [Fig. 1(c)]. The result is a spatially varying AC Stark shift that depends on both the in-plane and out-of-plane motions of the ions, thereby coupling the spins to the motion along both the directions.

A second ingredient in our proposal consists of a pair of co-propagating Raman lasers that drives spin flips without coupling to the motion. We assume that the two Raman lasers have an identical but tunable beam waist  $w$ , leading to an effective two-photon Rabi frequency that is radially varying as  $B(r) = B_0 e^{-r^2/w^2}$  and corresponding Hamiltonian  $\hat{H}_{\text{Raman},j} = B(r_j) \hat{\mathbf{s}}_j^X$ . For  $w \gg R$ , where  $R$  is the crystal radius, we can approximate  $B(r) \approx B_0 - B_0 r^2/w^2$ .

The role of the Raman drive is twofold and becomes apparent in the rotated spin space (see Appendix A). First, the spatially homogeneous drive with strength  $B_0$  serves to break the symmetry between a Jaynes-Cummings and an anti-Jaynes-Cummings type interaction of the spins and the drumhead c.m. mode that arise due to the ODF. The Jaynes-Cummings term can then be selectively brought near resonance by an appropriate choice of the ODF beatnote frequency  $\mu_r$ . Second, the beam waist  $w$  serves as a control knob for tuning the single particle dispersion, i.e.  $B_1 = B_0 R^2/w^2$ .

We present potential experimental parameters for realizing our proposal in Appendix B and study the adverse impact of off-resonant terms in Appendices C and D. Our study suggests that it is possible to operate in parameter regimes where the off-resonant terms have only a small effect. Although our study of off-resonant terms is extensive up to second order in small parameters, we have not considered possible accidental resonances or near-resonances that may occur at higher order because of the sheer number of higher-order terms. The impact of these terms and the parameter regimes where they are negligible could potentially be explored directly on the quantum simulator. In addition, for typical operating conditions, we estimate that decoherence from off-resonant light scattering may limit the simulation time. However, we note that the relative strength of coherent interaction to decoherence can be increased, for instance, by the choice of ion species and transition, by enhancing coherent coupling by parametric amplification [20] or by working at a different operating point for the optical dipole force.

### A. Initialization and readout

To observe dynamical phases generated by Hamiltonian (11), the spins must be initialized in a state possessing a chiral spin texture with a nonzero order parameter, such as the ones shown in Fig. 1(a). In the case of the trapped ion crystal, it is convenient to work with a normalized order parameter  $\Psi$  defined as

$$\Psi(t) = \frac{2}{N} \sum_{j=1}^N \tilde{r}_j e^{i\phi_j} \langle \hat{\mathbf{s}}_j^-(t) \rangle. \quad (12)$$

For engineering interactions, the ODF beatnote frequency  $\mu_r$  was tuned to couple the spins, the drumhead c.m. mode and the planar rotation. For preparing chiral initial states, we instead tune  $\mu_r$  to only couple the spins to the planar rotation without involving the drumhead c.m. mode. By additionally tuning the beam waist  $w_{\text{ODF}}$  of the ODF lasers, both BCS and BEC-like initial spin textures can be prepared. The initialization Hamiltonian is of the form

$$\hat{H}_{\text{init}} = \frac{\Omega_0}{2} \sum_j e^{-r_j^2/w_{\text{ODF}}^2} \tilde{r}_j (\hat{\mathbf{s}}_j^+ e^{-i\phi_j} + \hat{\mathbf{s}}_j^- e^{i\phi_j}), \quad (13)$$

where  $\Omega_0$  is an effective drive strength. This Hamiltonian drives single-spin rotations where the axis of rotation for ion  $j$  depends on the azimuthal angle  $\phi_j$  in the rotating frame. Starting with all spins initialized in  $|\uparrow\rangle_{\mathcal{Z}}$ , setting  $w_{\text{ODF}} \gg R$ , and using a pulse area  $\Omega_0 T = \pi$  results in a BCS-like spin texture [e.g. Fig. 1(a)], that can be used to observe phases I and II. Here, we have exploited the fact that the magnitude of the Rabi frequency increases with the radius so that the central spin is unaffected while the outermost spins are rotated to  $|\downarrow\rangle_{\mathcal{Z}}$ . A BEC-like spin texture [e.g. Fig. 1(a)] can be engineered by setting  $w_{\text{ODF}} < R$  and starting with all spins in  $|\downarrow\rangle_{\mathcal{Z}}$ . In this way, ions at the center, where  $\tilde{r}_j \sim 0$ , are unaffected whereas ions at the crystal boundary are also left unchanged since the beam intensity tapers to zero. On the other hand, ions in the intermediate region experience some degree of spin rotation and thereby give rise to a BEC-like texture. For observing phase III dynamics, a BCS-like initial state with a sharp domain wall between  $|\downarrow\rangle_{\mathcal{Z}}$  and  $|\uparrow\rangle_{\mathcal{Z}}$  spins and a small value of  $|\Psi(0)|$  is suitable. To prepare the domain wall, an optical pumping beam with a narrow waist, followed by a global  $\pi/2$  pulse, can be used to selectively prepare ions in the central region in  $|\uparrow\rangle_{\mathcal{Z}}$  those outside it in  $|\downarrow\rangle_{\mathcal{Z}}$ . A small initial  $|\Psi(0)|$  can then be induced by a short-time application of Hamiltonian (13). A detailed description of state initialization is presented in Appendix (G).

Measurement of the real and imaginary parts of the order parameter is also enabled by the Hamiltonian (13). To demonstrate this, we first introduce site-dependent orthogonal axes in the crystal plane and in the rotating frame,

$$\begin{aligned} \hat{\mathbf{e}}_{\mathcal{X}'_j} &= \sin \phi_j \hat{\mathbf{e}}_{\mathcal{X}} - \cos \phi_j \hat{\mathbf{e}}_{\mathcal{Y}}, \\ \hat{\mathbf{e}}_{\mathcal{Y}'_j} &= \cos \phi_j \hat{\mathbf{e}}_{\mathcal{X}} + \sin \phi_j \hat{\mathbf{e}}_{\mathcal{Y}}, \end{aligned} \quad (14)$$

such that  $\hat{\mathbf{e}}_{\mathcal{X}'_j} \times \hat{\mathbf{e}}_{\mathcal{Y}'_j} = \hat{\mathbf{e}}_{\mathcal{Z}}$ . In terms of these axes, we can write

$$\text{Re}[\Psi] = \frac{2}{N} \sum_j \tilde{r}_j \langle \hat{\mathbf{s}}_j^{\mathcal{Y}'_j} \rangle, \quad \text{Im}[\Psi] = \frac{2}{N} \sum_j \tilde{r}_j \langle \hat{\mathbf{s}}_j^{\mathcal{X}'_j} \rangle, \quad (15)$$

and with  $w_{\text{ODF}} \gg R$ , Hamiltonian (13) can be expressed as

$$\hat{H}_{\text{init}} = \Omega_0 \sum_j \tilde{r}_j \hat{\mathbf{s}}_j^{\mathcal{Y}'_j}. \quad (16)$$

After running the quantum simulation up to some time  $T$ , we evolve the system under (16) for a further time  $t$ . This leads to

$$\begin{aligned} \langle \hat{\mathbf{s}}_j^{\mathcal{Z}}(T+t) \rangle &= \langle \hat{\mathbf{s}}_j^{\mathcal{Z}}(T) \rangle \cos(\Omega_0 \tilde{r}_j t) - \langle \hat{\mathbf{s}}_j^{\mathcal{X}'_j}(T) \rangle \sin(\Omega_0 \tilde{r}_j t) \\ &\approx \langle \hat{\mathbf{s}}_j^{\mathcal{Z}}(T) \rangle - \left( \Omega_0 \tilde{r}_j \langle \hat{\mathbf{s}}_j^{\mathcal{X}'_j}(T) \rangle \right) t, \end{aligned} \quad (17)$$

where the approximation holds true for short rotation times. Summing over all the ions, we get

$$\langle \hat{\mathcal{J}}^{\mathcal{Z}}(t) \rangle - \langle \hat{\mathcal{J}}^{\mathcal{Z}}(0) \rangle \approx -\frac{\Omega_0}{2} \text{Im}[\Psi(T)] t, \quad (18)$$

where  $\hat{\mathcal{J}}^{\mathcal{Z}} = \sum_j \hat{\mathbf{s}}_j^{\mathcal{Z}}$ . The imaginary part of the order parameter determines the rate of change of  $\langle \hat{\mathcal{J}}^{\mathcal{Z}}(t) \rangle$  as the rotation time  $t$  is increased. This quantity is accessible by a global fluorescence measurement after applying a global  $\pi/2$  pulse to rotate  $\mathcal{Z} \rightarrow \mathcal{Z}$ . Furthermore, a phase offset of  $\pi/2$  can be introduced in Hamiltonian (13) by shifting the phase of the ODF beatnote, which can be used to measure  $\text{Re}[\Psi(T)]$  in a similar manner.

## IV. PROBING THE DYNAMICAL PHASES

Having established protocols for initializing BCS-like and BEC-like initial states, for engineering interactions, and for measuring the order parameter, we now proceed to discuss how the dynamical phases can be observed in a crystal stored in a Penning trap. The purple lines in Fig. 2 show the mean-field predictions for the time evolution of  $|\Psi(t)|$  under different initial conditions and interaction strengths, the latter characterized by the ratio  $K/J$ . The magnitude  $|\Psi(t)|$  decays to 0 in phase I, saturates to a non-zero constant in phase II and displays persistent oscillations in phase III. However, given the finite number of ions ( $N = 200$ ), we are motivated to investigate the validity of mean-field theory in describing our system. Towards this end, we simulate the dynamics under  $\hat{H}_{\text{1ch}}$  via the discrete truncated Wigner approximation (dTWA) method that accounts for the quantum noise of the initial state [21] (see Appendix E). The results of the dTWA simulations are shown by the red lines in Fig. 2. The dTWA and mean-field results agree well in Phase I where the single-particle dephasing dominates




FIG. 2. **Manifestation of dynamical phases in a 200 ion crystal.** Time evolution of  $|\Psi(t)|$  in phases I, II and III computed using mean-field theory and the discrete truncated Wigner approximation (dTWA) method. The dTWA results show that the finite size of the crystal leads to a decay of  $|\Psi(t)|$  even in phases II and III. However, the buildup of quantum correlations in these phases is captured by a second order parameter  $\tilde{\Psi}$  [Eq. (19)]. In all cases, the decay of  $|\Psi(t)|$  in the absence of interactions is plotted for reference. Crystal parameters are detailed in Appendix B and chiral spin states are initialized according to Appendix G. In (a)  $K/J = 10$ , while in (b) and (c)  $K/J = 1$ .

the interactions. In contrast, the dTWA results deviate significantly from the mean-field predictions in Phases II and III. In both cases, quantum noise causes the order parameter to eventually decay to zero in the long time limit.

Importantly, the decay of the mean-field order parameter  $|\Psi(t)|$  in phases II and III is accompanied by the development of quantum correlations, which is captured in a more appropriate order parameter  $\tilde{\Psi}$  defined as

$$\tilde{\Psi} = \frac{2}{N} \left| \sum_{j \neq k} \tilde{r}_j \tilde{r}_k \langle \hat{\mathbf{s}}_j^+ \hat{\mathbf{s}}_k^- \rangle e^{-i(\phi_j - \phi_k)} \right|^{1/2}. \quad (19)$$

We note that  $\tilde{\Psi}$  is just a measure of the interaction part of the Hamiltonian  $\hat{H}_{1\text{ch}}$  [Eq. (11)]. Figure 2 shows that  $\tilde{\Psi}$  stabilizes to a non-zero constant in phases II and III indicating the persistence of dynamical p-wave superfluidity in these phases.

Even though the mean-field order parameter  $|\Psi(t)|$  disappears at long times, the three phases can be distinguished in the short time dynamics of this observable. Figure 2 shows that for  $Jt \lesssim 20$ , the mean-field and dTWA results are in approximate agreement in all three phases. The magnitude  $|\Psi(t)|$  does not decay in Phase II whereas it exhibits strong amplification in Phase III. In an experiment, the stabilization of superfluidity by interactions can be sharply demonstrated by comparing the dynamics of  $|\Psi(t)|$  in the presence and absence of the ODF drive; in the absence of interactions,  $|\Psi(t)|$  decays towards zero even on short time scales.

## V. INFERRING TOPOLOGY

In contrast to *s*-wave superconductors, ground states and dynamical phases of *p*-wave superconductors can

possess nontrivial topological properties. We now discuss how the topology of the dynamical phases can be probed in the ion simulator. Figure 3(a) shows the winding number  $W$  computed in mean-field theory as the ratio  $K/J$  is tuned for a crystal of  $N = 200$  ions in the Penning trap (see Appendix F for details of this calculation). Representative examples of the magnetic field texture are also shown, which demonstrate the transition from a topologically trivial BEC-like texture ( $K/J < 0.6$ ,  $W \approx 0$ ) to a topologically non-trivial BCS-like texture ( $K/J > 0.6$ ,  $W \approx 1$ ). Remarkably, the topologically trivial and non-trivial phases can be distinguished by measurements of the real and imaginary parts of the order parameter. From Eq. (8), the sense of rotation of the order parameter in the complex plane—clockwise ( $\mu_\infty > 0$ ) or counterclockwise ( $\mu_\infty < 0$ )—is a direct measurement of the sign of  $\mu_\infty$ , and consequently, enables us to infer the BCS-like or BEC-like nature of the effective magnetic field texture. Figures 3(b) and (c) show that the sense of rotation of the order parameter is different for  $K/J = 0.35$  and  $K/J = 0.85$ , clearly indicating the transition from a topologically trivial to a topologically nontrivial dynamical phase as the ratio  $K/J$  is tuned. While mean-field theory predicts the order parameter to precess with an approximately constant radius in the complex plane, the build-up of quantum correlations cause the order parameter to spiral in towards the origin at long times, consistent with Fig. 2(b). Nevertheless, the decay does not modify the topological nature of the dynamical phases.

The preservation of the topology in the regime where the order parameter is decaying can further be confirmed by measuring the so-called Cooper pair distribution function (CPDF)  $\gamma(\mathbf{p})$  [4]. This function characterizes the nonequilibrium distribution of the quasiparticle spectrum in the asymptotic steady state and provides information about the topology of the dynamical phases: The topology of the dynamical phase is trivial (non-trivial) if the




FIG. 3. **Inferring topological properties of the dynamical phases.** (a) Mean-field, long-time winding number of the effective magnetic field texture as the relative strength of the single-particle ( $K$ ) and interaction terms ( $J$ ) are varied, starting from a BEC-like initial state. In the strong interaction case ( $K/J < 0.6$ ), the texture is BEC-like and topologically trivial, while in the weak interaction case ( $K/J > 0.6$ ), it is BCS-like and topologically nontrivial. Representative field textures are shown, where each arrow now indicates the unit vector of the effective magnetic field acting at that site, and the color code indicates the normalized  $B_z$  component. (b-c) The sense of rotation of the order parameter in the complex plane reveals the winding number of the underlying magnetic field texture. The topologically trivial (nontrivial) dynamical phase is associated with a counterclockwise (clockwise) rotation of  $\Psi(t)$ . While mean-field predicts a stable limit cycle (gray dashed line), dTWA calculations (color dots) show the order parameter spiralling in towards the origin at long times (plotted here until  $Jt = 100$ ). The color gradient indicates the arrow of time. (d) The Cooper pair distribution function (CPDF) displays an even (odd) number of zero-crossings in the topologically trivial (non-trivial) case. These features are preserved even at times  $Jt \sim 100$ , when the order parameter has decayed considerably due to quantum fluctuations. For comparison, the dashed lines show the CPDF computed using mean-field theory. Crystal parameters are detailed in Appendix B and chiral spin states are initialized according to Appendix G.

number of zero-crossings of this function is even (odd) [4]. In superconductors the CPDF can be measured via time-resolved ARPES [22]. In the spin model,  $\gamma(\mathbf{p})$ , maps to  $\gamma(\tilde{r}_j)$  and corresponds to the projection of the local spin vector  $\hat{\mathbf{s}}_j$  onto the local effective magnetic field  $\bar{\mathbf{B}}_j$  at a given radius  $\tilde{r}_j$ :

$$\gamma(\tilde{r}_j) = \frac{\langle \overline{\hat{\mathbf{s}}_j} \rangle \cdot \langle \overline{\mathbf{B}}_j \rangle}{\sqrt{\langle \overline{\hat{\mathbf{s}}_j} \rangle \cdot \langle \overline{\hat{\mathbf{s}}_j} \rangle} \sqrt{\langle \overline{\mathbf{B}}_j \rangle \cdot \langle \overline{\mathbf{B}}_j \rangle}}. \quad (20)$$

Here, the overbar denotes that the spin and effective magnetic field are measured in a frame rotating at  $2\mu_\infty$  and the denominator ensures that the quantity is the direction cosine of the spin vector with respect to the local

magnetic field. Figure 3(d) shows the CPDF as a function of ion radius from the trap center for  $K/J = 0.35$  and  $K/J = 0.85$ . The curves in Fig. 3(d) have been computed accounting for quantum fluctuations and by running the simulation up to a time  $Jt = 100$ , for which the decay of the order parameter is clearly visible in Fig. 3(b-c). We find that the number of zero-crossings of the CPDF enable an inference of the topology even after significant decay of the order parameter. We note that site-resolved measurement of the spin components is sufficient to measure the effective local magnetic field (see Eq. (6)), while the value of  $\mu_\infty$  can be obtained from a Fourier transform of the time series of  $\Psi(t)$ .




FIG. 4. **Realizing the two-channel model.** The drum-head c.m. mode plays the role of the molecular channel in the two-channel  $p$ -wave model. As the coupling to the c.m. mode is tuned from off-resonant ( $\delta_1^2/G^2 \gg 1$ ), to near-resonance ( $\delta_1^2/G^2 \sim 0$ ), the role of the bosonic channel is no longer negligible and its effects can be clearly observed in the time evolution of  $|\Psi(t)|$ . Here, we fix  $B_1 = G/\sqrt{10}$  as  $\delta_1$  is varied to obtain the different curves. Crystal parameters are detailed in Appendix B and chiral spin states are initialized according to Appendix G.

## VI. REALIZING A TWO-CHANNEL MODEL

So far, we have focused on a regime where the c.m. mode is coupled off-resonantly to the spins and can hence be adiabatically eliminated, giving rise to an effective one-channel model description in terms of the Anderson spins alone. The c.m. mode plays the role of the bosonic molecular channel in the two-channel  $p$ -wave model. By suitably tuning the ODF difference frequency  $\mu_r$ , a near-resonant coupling with the c.m. mode can be engineered, thereby enabling simulation of the more general two-channel Hamiltonian (10). Thus, our trapped ion simulator allows the exploration of the distinct physical behaviors featured by both the one-channel and two-channel models in the same experimental setting. Figure 4 shows the time evolution of  $|\Psi(t)|$  as the coupling to the c.m. mode is tuned from a far off-resonant regime ( $\delta_1^2/G^2 \gg 1$ ) to the resonant regime ( $\delta_1^2/G^2 \sim 0$ ). The curves are computed accounting for the initial quantum noise of the spins, which are in a BCS-like state, as well as the c.m. mode, which is assumed to be in the ground state. In the off-resonant regime, the behavior is similar to the one-channel model as the occupation of the c.m. mode remains negligible at all times. However,  $|\Psi(t)|$  exhibits pronounced oscillatory behavior in the near-resonant and resonant regimes where excitations are exchanged between the spins and the c.m. mode. Interestingly, these results indicate that for fixed  $B_1$ , there is an optimal  $\delta_1$  in order to stabilize  $|\Psi(t)|$ . In addition to probing the role of the molecular channel, near-resonant coupling also enhances the interaction strength and may hence exhibit stronger signatures of the interaction dynamics in the time before sources of decoherence such as light scatter-

ing from the ODF beams kick in (Appendix B).

## VII. CONCLUSION AND OUTLOOK

We have proposed a protocol to simulate the quench dynamics of  $p \pm ip$  superfluids using ion crystals stored in a Penning trap. By utilizing the Anderson pseudospin mapping, we make a fermionic Hamiltonian amenable to simulation using spins encoded in the internal states of ions. In this way, we not only leverage the versatile toolbox of the trapped ion platform but we also circumvent limitations that arise with direct simulations of the fermionic model using ultracold quantum gases, where  $p$ -wave superfluids have been highly elusive till date.

We demonstrated how signatures of all three dynamical phases can be observed using crystals with  $\sim 200$  ions. In the thermodynamic limit, the 2D  $p + ip$  model becomes integrable and the dynamics is well described by mean field theory. However, beyond mean field effects kick in at finite number of fermions; the relatively small number of spins in the trapped ion crystal naturally opens the avenue to observe beyond mean-field effects in the quench dynamics of  $p + ip$  superfluids and also facilitates the emulation of small superconducting grains [23].

We showed how the topological character of the dynamical phases can be inferred via the sense of rotation of the order parameter in the complex plane. Since our proposal maps the fermionic momentum on to the real-space position of ions, the addition of site-resolved detection of spins in the trap can provide time and momentum-resolved measurements of the simulated system, analogous to techniques used in solid state materials such as ARPES [24]. Along these lines, we showed how the Cooper pair distribution function can be measured with site-resolved detection and can be used to verify the topology of the dynamical phase.

By tuning closer to resonance with the c.m. mode, a two-channel  $p$ -wave model can be realized and the nonzero occupation of a molecular channel can be additionally simulated and investigated. It will be interesting to explore conditions where a dynamically active molecular channel can enhance superfluidity [25], in a way reminiscent to analogous phenomena recently studied in solid state quantum optics[26–36]. For example, our system can be used to study the system response when the c.m. mode (i.e. the molecular channel) is driven or even squeezed via parametric amplification [20, 37]. Furthermore, although we have considered a regime where the other drumhead modes are off-resonant, strong spin-mode coupling can lead to weak excitations of more than a single mode, leading to the emergence of spatial inhomogeneities than can emulate the phenomenon of Cooper pair turbulence [38].

Beyond the simulation of  $p + ip$  superconductors, trapped ions could also be used to study further exotic superconducting systems such as chiral  $d_{x^2-y^2} + id_{xy}$  superfluids [39]. Such systems can be simulated in the Pen-



ning trap by engineering the phase of the ODF beams to have spatial variations, which is possible with the use of deformable mirrors.

Finally, we note that the fermionic statistics of the original  $p$ -wave model is not present in the corresponding spin mapping, and therefore our proposal cannot be used to produce Majorana fermions [40, 41]. However, the protocols we have developed can be used to both produce skyrmionic spin textures as well as stabilize them using interactions, which could find applications in demonstrating skyrmion qubits for quantum computing [42].

#### ACKNOWLEDGEMENTS

We thank Allison Carter for providing estimates for the decoherence rates from light scattering. We thank

Diego Barberena and Bryce Bullock for a careful reading and comments on the manuscript. This work is supported by the European Union's Horizon 2020 research and innovation program under Grant Agreements No. 817482 (Pasquans), by the Simons Collaboration on Ultra-Quantum Matter, which is a grant from the Simons Foundation (651440, P. Z.), and by LASCEM via AFOSR No. 64896-PH-QC. Support is also acknowledged from the AFOSR grants FA9550-18-1-0319 and FA9550-19-1-0275, by the NSF JILA-PFC PHY-1734006, QLCI-OMA -2016244, by the U.S. Department of Energy, Office of Science, National Quantum Information Science Research Centers, Quantum Systems Accelerator, and by NIST. JJB acknowledges support from the DARPA ON-ISQ program and AFOSR grant FA9550-201-0019.

## Appendix A: Derivation of effective Hamiltonians

We show how the combination of optical dipole force (ODF) and Raman beams with suitable geometries leads to effective one-channel and two-channel models for  $p$ -wave interactions in ion crystals stored in a Penning trap.

### 1. Optical dipole force with tilted wavevectors

The Hamiltonian corresponding to only the ODF interaction is

$$\hat{H} = \sum_{j=1}^N \omega_s \hat{s}_j^Z + \sum_{n=1}^N \omega_n \hat{a}_n^\dagger \hat{a}_n + 2 \sum_{j=1}^N \delta_{\text{AC}} \sin(\Delta k_x x_j + \Delta k_z \hat{z}_j - \mu_r t) \hat{s}_j^Z, \quad (\text{A1})$$

where  $\omega_s$  is the spin transition frequency,  $\delta_{\text{AC}}$  is the strength of the optical dipole force,  $\Delta \mathbf{k} = \Delta k_x \hat{\mathbf{e}}_x + \Delta k_z \hat{\mathbf{e}}_z$  is the difference wavevector of the ODF beams and  $\mu_r$  is their difference frequency. The out-of-plane motion is treated quantum mechanically and is represented by the operator  $\hat{z}_j$ . In terms of the drumhead modes, it can be expressed as

$$\Delta k_z \hat{z}_j = \sum_{n=1}^N \eta_n \mathcal{M}_{jn} (\hat{a}_n + \hat{a}_n^\dagger), \quad (\text{A2})$$

where  $\hat{a}_n, \hat{a}_n^\dagger$  are annihilation and creation operators for mode  $n$  with frequency  $\omega_n$ ,  $\eta_n = \Delta k_z \sqrt{\hbar/(2m\omega_n)}$  is the Lamb-Dicke parameter,  $m$  is the ion mass, and  $\mathcal{M}_{jn}$  is the displacement of ion  $j$  under the influence of mode  $n$ . In contrast, the in-plane motion is dominated by the crystal rotation and is represented by the classical coordinate  $x_j(t) = r_j \cos(\omega_r t + \phi_j)$ , where  $r_j$  is the distance of the ion from the trap center and  $\phi_j$  is the azimuthal angle in the rotating frame.

### 2. Co-propagating Raman beams

We now introduce a pair of Raman beams that drive resonant two-photon transitions between the spin states. The beams are assumed to be copropagating so that their difference wavevector approximately vanishes [43] and hence does not induce any spin-motion coupling. In a frame rotating at  $\omega_s$ , the total Hamiltonian including the ODF and the Raman beams is

$$\hat{H} = \sum_{j=1}^N B_j \hat{s}_j^X + \sum_{n=1}^N \omega_n \hat{a}_n^\dagger \hat{a}_n + 2 \sum_{j=1}^N \delta_{\text{AC}} \sin(\Delta k_x x_j + \Delta k_z \hat{z}_j - \mu_r t) \hat{s}_j^Z. \quad (\text{A3})$$

Here,  $B_j$  is the effective two-photon Rabi frequency at the site of ion  $j$ . Assuming a beam waist  $w \gg R$  for the Raman lasers, where  $R$  is the crystal radius, we can approximate  $B_j \approx B_0 - B_0 r_j^2/w^2$ .

We now analyze the spin dynamics in a rotated spin space such that  $\hat{\mathbf{s}}_Z \equiv -\hat{s}_X$  and  $\hat{\mathbf{s}}_X \equiv \hat{s}_Z$ . Further transforming to a frame rotating at  $B_0$ , the Hamiltonian in the rotated spin space is

$$\hat{H} = \sum_{j=1}^N B_0 \frac{r_j^2}{w^2} \hat{s}_j^Z + \sum_{n=1}^N \omega_n \hat{a}_n^\dagger \hat{a}_n + \sum_{j=1}^N \delta_{\text{AC}} \sin(\Delta k_x x_j + \Delta k_z \hat{z}_j - \mu_r t) (\hat{\mathbf{s}}_j^+ e^{-iB_0 t} + \hat{\mathbf{s}}_j^- e^{iB_0 t}). \quad (\text{A4})$$

We can now see the twofold role played by the Raman drive: By providing a splitting  $2B_0$  between  $\hat{\mathbf{s}}_j^+$  and  $\hat{\mathbf{s}}_j^-$ , it will enable retention of only desired interactions and enable ‘‘rotating out’’ unwanted interactions. Second, the beam waist  $w$  acts as a knob to tune the single particle dispersion.

### 3. Small angle approximation

We now consider the sine function appearing in Eq. (A4). For three arguments  $A = \Delta k_x x_j, B = \Delta k_z \hat{z}_j, C = -\mu_r t$ , we can expand

$$\sin(A + B + C) = \sin(A + B) \cos(C) + \cos(A + B) \sin(C) \quad (\text{A5})$$

We assume that  $A, B \ll 1$  and expand the relevant trigonometric functions in the small angle limit. The result, correct to second order in  $A, B$ , is

$$\sin(A + B + C) \approx (A + B) \cos C + \left(1 - \frac{A^2 + B^2}{2} - AB\right) \sin C. \quad (\text{A6})$$

#### 4. Obtaining the two-channel model

We express the resonance requirements as a sum of frequencies appearing in the argument of complex exponentials multiplying each interaction term. To do so, we first note that the motion along  $x$  can be written as  $x_j = r_j \cos(\omega_r t + \phi_j) \equiv (r_j/2)(e^{i(\omega_r t + \phi_j)} + e^{-i(\omega_r t + \phi_j)})$ , while the motion along  $z$  can be expanded as

$$\Delta k_z \hat{z}_j(t) = \sum_{n=1}^N \eta_n \mathcal{M}_{jn} (\hat{a}_n e^{-i\omega_n t} + \hat{a}_n^\dagger e^{i\omega_n t}), \quad (\text{A7})$$

This expansion assumes we have moved to an interaction picture with respect to the free phonon frequencies  $\omega_n$ . The homogeneous Raman drive sets a frequency  $B_0$  for the spins as seen from Eq. (A4). Finally, the terms  $\cos(-\mu_r t), \sin(-\mu_r t)$  can be expanded with complex exponentials of the form  $e^{\pm i\mu_r t}$ . For discussing the rotating wave approximations, we neglect the small contribution arising from the spatially inhomogeneous component of the Raman beams, i.e. we assume the beams waist  $w \rightarrow \infty$ .

We tune  $\mu_r$  to selectively induce a coupling between the spins ( $B_0$ ), the drumhead c.m. mode ( $\omega_1$ ) and the planar rotation ( $\omega_r$ ). In particular, we adjust  $\mu_r$  such that  $B_0 - \mu_r + \omega_1 + \omega_r = \delta_1$ . We will later show using realistic experimental parameters that all unwanted terms are far-off resonant and can be neglected in a rotating-wave approximation. Then, the only near-resonant term stems from the  $AB \sin C$  type term in Eq. (A6), and is of the form

$$\hat{H}_1 = - \sum_{j=1}^N \frac{G}{i\sqrt{N}} \tilde{r}_j \left( \hat{\mathbf{s}}_j^- \hat{a}_1^\dagger e^{i(\delta_1 t + \phi_j)} - \hat{\mathbf{s}}_j^+ \hat{a}_1 e^{-i(\delta_1 t + \phi_j)} \right), \quad (\text{A8})$$

where  $\tilde{r}_j = r_j/R$  is the ion radius normalized to the crystal radius  $R$  (assuming a nearly circular crystal) and  $G$  is given by

$$G = \frac{\delta_{\text{AC}}(\eta_1) (\Delta k_x R)}{4}. \quad (\text{A9})$$

Here,  $\eta_1 = \Delta k_z \sqrt{\hbar/(2m\omega_1)}$  is the Lamb-Dicke parameter for the c.m. mode. The quantity  $\Delta k_x R$  can be thought of as an effective Lamb-Dicke parameter for the in-plane motion.

The above analysis was carried out assuming that the beam waist of the Raman lasers  $w \rightarrow \infty$ . Restoring a finite  $w$  and performing a frame transformation for the c.m. mode, we arrive at the effective Hamiltonian

$$\hat{H}_{2\text{ch}} = \sum_{j=1}^N B_1 \tilde{r}_j^2 \hat{\mathbf{s}}_j^z + \delta_1 \hat{a}_1^\dagger \hat{a}_1 - \sum_{j=1}^N \frac{G}{i\sqrt{N}} \tilde{r}_j \left( \hat{\mathbf{s}}_j^- \hat{a}_1^\dagger e^{i\phi_j} - \hat{\mathbf{s}}_j^+ \hat{a}_1 e^{-i\phi_j} \right), \quad (\text{A10})$$

where  $B_1 = B_0 R^2/w^2$ .

#### 5. Effective spin-exchange interaction

We now eliminate the c.m. mode from Eq. (A10) using effective Hamiltonian theory [19]. We express energy resonance requirements once again as complex exponentials by defining  $\delta_1^j = \delta_1 - B_1 \tilde{r}_j^2$ . Using a frame transformation for the spins and the c.m. mode, Eq. (A10) can be written as

$$\hat{H}_1 = - \sum_{j=1}^N \frac{G}{i\sqrt{N}} \tilde{r}_j \left( \hat{\mathbf{s}}_j^- \hat{a}_1^\dagger e^{i(\delta_1^j t + \phi_j)} - \hat{\mathbf{s}}_j^+ \hat{a}_1 e^{-i(\delta_1^j t + \phi_j)} \right). \quad (\text{A11})$$

By assuming that  $\delta_1^j$  are large compared to the maximum interaction strength  $G$ , a spin-spin Hamiltonian can be derived using effective Hamiltonian theory. The result is

$$\hat{H}_{\text{eff}} = \sum_{j,k=1}^N \frac{G^2}{Nh(\delta_1^j, \delta_1^k)} \tilde{r}_j \tilde{r}_k [\hat{a}_1^\dagger \hat{\mathbf{s}}_j^- e^{i(\delta_1^j t + \phi_j)}, \hat{a}_1 \hat{\mathbf{s}}_k^+ e^{-i(\delta_1^k t + \phi_k)}], \quad (\text{A12})$$

where  $h(a, b)$  is the harmonic mean of  $a, b$ . The commutator evaluates to

$$[\hat{a}_1^\dagger \hat{\mathbf{s}}_j^-, \hat{\mathbf{s}}_k^+ \hat{a}_1] = \begin{cases} -\hat{\mathbf{s}}_j^- \hat{\mathbf{s}}_k^+, & j \neq k, \\ -2\hat{a}_1^\dagger \hat{a}_1 \hat{\mathbf{s}}_j^z - \hat{\mathbf{s}}_j^+ \hat{\mathbf{s}}_j^-, & j = k. \end{cases} \quad (\text{A13})$$

The term containing  $\hat{a}_1^\dagger \hat{a}_1$  can be neglected if the c.m. mode is initially in the ground state such that  $\langle \hat{a}_1^\dagger \hat{a}_1 \rangle = 0$ . The effective Hamiltonian is therefore

$$\hat{H}_{\text{eff}} = - \sum_{j,k=1}^N \frac{G^2}{Nh(\delta_1^j, \delta_1^k)} \tilde{r}_j \tilde{r}_k \hat{\mathbf{s}}_j^+ \hat{\mathbf{s}}_k^- e^{-i[(\delta_1^j - \delta_1^k)t + (\phi_j - \phi_k)]}. \quad (\text{A14})$$

Transforming back to the non-rotating frame gives

$$\hat{H}_{\text{eff}} = \sum_{j=1}^N \left( B_1 \tilde{r}_j^2 - \frac{G^2}{N\delta_1^j} \right) \hat{\mathbf{s}}_j^z - \sum_{j=1}^N \sum_{k \neq j}^N \frac{G^2}{Nh(\delta_1^j, \delta_1^k)} \tilde{r}_j \tilde{r}_k \hat{\mathbf{s}}_j^+ \hat{\mathbf{s}}_k^- e^{-i(\phi_j - \phi_k)}, \quad (\text{A15})$$

Since  $\delta_1 \gg G^2/\delta_1$  and  $B_1$  is comparable to the latter frequency, as a first approximation we can assume  $\delta_1^j \approx \delta_1^k = \delta_1$  and neglect site dependency in the denominators of the effective frequencies. This leads to the Hamiltonian for the one-channel model:

$$\hat{H}_{1\text{ch}} = K \sum_{j=1}^N \tilde{r}_j^2 \hat{\mathbf{s}}_j^z - \frac{J}{N} \sum_{j=1}^N \sum_{k \neq j}^N \tilde{r}_j \tilde{r}_k \hat{\mathbf{s}}_j^+ \hat{\mathbf{s}}_k^- e^{-i(\phi_j - \phi_k)}, \quad (\text{A16})$$

where  $J = G^2/\delta_1$  and  $K = B_1 - J/N$ . We note that the total magnetization  $\hat{\mathcal{J}}^z = \sum_j \hat{\mathbf{s}}_j^z$  is conserved by this Hamiltonian.

## Appendix B: Experimental parameters for implementation

In this section, we provide experimental parameters for implementing our proposal. These parameters are based on settings used in the NIST Penning trap, where 2D crystals of tens to hundreds of  ${}^9\text{Be}^+$  ions are routinely prepared for quantum simulation and sensing.

### 1. Trapping parameters

Two-dimensional crystals of  ${}^9\text{Be}^+$  ions are formed in the Penning trap by a combination of an electric quadrupole field providing axial confinement and a strong axial magnetic field  $\mathcal{B} \approx 4.46$  T (cyclotron frequency  $\omega_c/(2\pi) \approx 7.6$  MHz) that aids in radial confinement. The spin-1/2 degree of freedom is encoded in two long-lived hyperfine levels of each  ${}^9\text{Be}^+$  ion, i.e.  $|\uparrow\rangle \equiv |2S_{1/2}, 1/2\rangle$  and  $|\downarrow\rangle \equiv |2S_{1/2}, -1/2\rangle$ . For a crystal with  $N = 200$  ions, the crystal radius is  $R \approx 100$   $\mu\text{m}$ . Here we consider two sets of trapping parameters. In case A, we will set the rotation frequency to  $\omega_r/(2\pi) = 180$  kHz and choose an axial trapping frequency, which is also the drumhead c.m. frequency, to be  $\omega_1/(2\pi) = 1.59$  MHz. In case B, we choose a faster rotating crystal with a higher axial trapping frequency, viz.  $\omega_r/(2\pi) = 900$  kHz and  $\omega_1/(2\pi) = 3.42$  MHz. In the following, we explicitly refer to the different cases when specifying parameters that are not the same in the two cases. While case A represents the commonly used trapping parameters, the reason we consider two sets of parameters will become clear in Appendix D, where we show that certain off-resonant terms are only negligible for the faster rotating crystal, i.e. case B.

In general, both the ODF beams and the Raman beams intersect the crystal plane at a non-zero angle. However, for the quantum simulation, we require that these beams have a radially varying intensity profile in the crystal plane. This can be achieved by using laser beams with elliptical beams waists, whose cross-section in the crystal plane will have a radial intensity profile.

### 2. ODF interaction

The optical dipole force is realized using a pair of lasers that intersect the crystal at approximately equal and opposite angles relative to the crystal plane, see Fig. 1(c). These lasers induce spatially varying AC Stark shifts on

the pseudospin states by coupling these states to the  $|2P_{3/2}\rangle$  manifold and give rise to a Hamiltonian of the form described in Eq. (A1).

From Fig. 1(c), the wavevectors of the ODF lasers are given by

$$\begin{aligned}\mathbf{k}_u &= (k \cos \theta) \hat{\mathbf{e}}_x - (k \sin \theta) \hat{\mathbf{e}}_z, \\ \mathbf{k}_l &= (k \cos(\theta + \delta\theta)) \hat{\mathbf{e}}_x + (k \sin(\theta + \delta\theta)) \hat{\mathbf{e}}_z,\end{aligned}\tag{B1}$$

where  $u, l$  denote the upper and lower ODF beams respectively and  $k = |\mathbf{k}_u| = |\mathbf{k}_l|$ . Denoting  $k_x = k \cos \theta, k_z = k \sin \theta$ , we can write

$$\begin{aligned}\mathbf{k}_u &= k_x \hat{\mathbf{e}}_x - k_z \hat{\mathbf{e}}_z, \\ \mathbf{k}_l &\approx (k_x - k_z \delta\theta) \hat{\mathbf{e}}_x + (k_z + k_x \delta\theta) \hat{\mathbf{e}}_z.\end{aligned}\tag{B2}$$

The difference wavevector can then be expressed as

$$\Delta\mathbf{k} = k_z \delta\theta \hat{\mathbf{e}}_x - (2k_z - k_x \delta\theta) \hat{\mathbf{e}}_z \equiv \Delta k_x \hat{\mathbf{e}}_x + \Delta k_z \hat{\mathbf{e}}_z.\tag{B3}$$

The wavevector magnitude of each ODF laser is  $k \approx 2 \times 10^7 \text{m}^{-1}$ . The Lamb-Dicke parameter along the axial direction is  $\eta_1 = \Delta k_z l_1^{\text{ZP}}$ , where  $\Delta k_z \approx 2k \sin(\theta)$  is the difference wavevector along the  $z$ -direction and  $l_1^{\text{ZP}} = \sqrt{\hbar/(2m\omega_1)}$  is the zero-point motion of the c.m. mode. Here, we have neglected the small correction to  $\Delta k_z$  that arises from a non-zero misalignment  $\delta\theta$ . To obtain a value  $\eta_1$ , the ODF lasers must be oriented at angles  $\pm\theta$  with respect to the crystal plane, with  $\theta = \sin^{-1}(\eta_1/(2kl_1^{\text{ZP}}))$ . For  $\eta_1 \approx 0.3$ , we find  $\theta \approx 23.4^\circ$  for case A and  $\theta \approx 35.7^\circ$  for case B.

An analogous small parameter along the  $x$ -direction is given by  $\eta_x = \Delta k_x R$ , where  $\Delta k_x = k \delta\theta \sin \theta$  is the difference wavevector in the crystal plane. To obtain a value  $\eta_x$ , the required misalignment is given by  $\delta\theta = \eta_x/(kR \sin \theta)$ . For  $\eta_x \approx 0.3$ , we find  $\delta\theta \approx 0.017^\circ$  for case A and  $\delta\theta \approx 0.016^\circ$  for case B.

We assume that  $\delta_{\text{AC}}/(2\pi) \approx 40$  kHz. Then, the interaction strength  $G$  for the two-channel model is given by

$$\frac{G}{2\pi} = \frac{\delta_{\text{AC}} \eta_1 \eta_x}{4(2\pi)} = 900 \text{ Hz}.\tag{B4}$$

By assuming a detuning of  $\delta_1/(2\pi) = 2$  kHz from the c.m. mode, we can arrive at the effective one-channel model coupling strength  $J$  as

$$\frac{J}{2\pi} = \frac{G^2}{\delta_1} = 405 \text{ Hz}.\tag{B5}$$

In this parameter regime, the ratio  $\delta_1^2/G^2 \sim 5$ , and therefore the simulation will have a small two-channel character to it in addition to the dominant one-channel model (see Fig. 4).

### 3. Raman beams

Two-photon Raman transitions between the pseudospin states can be engineered by introducing a pair of copropagating Raman lasers that couple these states to the  $|2P_{3/2}\rangle$  manifold in a far detuned regime. The purpose of the Raman beams is twofold. First, the spatially homogeneous two-photon Raman frequency  $B_0$  breaks the symmetry between the  $\hat{\mathbf{s}}^+$  and  $\hat{\mathbf{s}}^-$  terms, see Eq. (A4). We take this value to be  $B_0/(2\pi) = 10$  kHz. Second, the radially varying intensity of the Raman beams tunes the dispersion  $K$  of the spins. We estimate the scale of the required beam waist  $w$  as the value at which  $K \approx J$ , i.e.

$$K \approx \frac{B_0 R^2}{w^2} = J \implies w \approx 497 \text{ } \mu\text{m}.\tag{B6}$$

An alternative mechanism to generate a Raman beam intensity gradient is by utilizing the Doppler shifts arising from the crystal rotation. Although the Raman beams are copropagating, their difference wavevector  $|\Delta\mathbf{k}_R| \neq 0$  because of the frequency splitting  $\omega_s$  of the spin states, i.e.  $|\Delta\mathbf{k}_R| = \omega_s/c$ , where  $c$  is the speed of light in vacuum. Assuming the Raman beams are propagating in the  $x-z$  plane and make an angle  $\theta_R$  with the crystal plane, the Hamiltonian for a single ion interacting with the Raman beams is given by

$$\hat{H}_{R,j} = \omega_s \hat{\mathbf{s}}_j^Z + \frac{B_0}{2} \left( \hat{\mathbf{s}}_j^+ e^{i[|\Delta\mathbf{k}_R| \cos \theta_R x_j(t) - \Delta\omega_R t]} + \hat{\mathbf{s}}_j^- e^{-i[|\Delta\mathbf{k}_R| \cos \theta_R x_j(t) - \Delta\omega_R t]} \right).\tag{B7}$$

In the Doppler free case, two-photon resonance is satisfied by setting  $\Delta\omega_R = \omega_s$ . In this case, transforming to a frame rotating at  $\omega_s$  leads to the interaction Hamiltonian

$$\hat{H}_{R,j}^I = \frac{B_0}{2} \left( \hat{\mathbf{s}}_j^+ e^{i|\Delta\mathbf{k}_R| \cos\theta_R x_j(t)} + \hat{\mathbf{s}}_j^- e^{-i|\Delta\mathbf{k}_R| \cos\theta_R x_j(t)} \right). \quad (\text{B8})$$

The Doppler shift is time modulated because  $x_j(t) = r_j \cos(\omega_r t + \phi_j)$ . Defining  $\eta_j^R = |\Delta\mathbf{k}_R| \cos\theta_R r_j$ , we can use the Jacobi-Anger expansion to write

$$e^{-i\eta_j^R \cos(\omega_r t + \phi_j)} = \sum_{n=-\infty}^{\infty} (-i)^n J_n(\eta_j^R) e^{-in(\omega_r t + \phi_j)}, \quad (\text{B9})$$

where  $J_n(x)$  is the  $n^{\text{th}}$  Bessel function of the first kind. The  $n = 0$  term is then given by

$$\hat{H}_{R,j}^{I,(0)} = \frac{B_0 J_0(\eta_j^R)}{2} \hat{\mathbf{s}}_j^x \approx \frac{B_0}{2} \left( 1 - \frac{(\eta_j^R)^2}{4} \right) \hat{\mathbf{s}}_j^x = \frac{1}{2} \left( B_0 - \frac{B_0 \omega_s^2 R^2}{4c^2} \tilde{r}_j^2 \right) \hat{\mathbf{s}}_j^x. \quad (\text{B10})$$

The approximation is valid for  $(\eta_j^R)^2 \ll 1$ . To verify this, we consider a worst-case situation when  $\theta_R = 0$  and  $r_j = R = 100 \mu\text{m}$ , where  $R$  is the crystal radius. Using  $\omega_s/(2\pi) = 124 \text{ GHz}$ , we estimate  $\eta_j^R \approx 0.26$  and  $(\eta_j^R)^2 \approx 0.067 \ll 1$ . The achievable value of  $B_1$  in this case is given by

$$B_1 = \frac{(\eta_j^R)^2}{4} B_0 \approx 0.017 B_0 \approx 2\pi \times 170 \text{ Hz}, \quad (\text{B11})$$

where we have used  $B_0/(2\pi) = 10 \text{ kHz}$ . Therefore, it appears that  $B_1$  can be partially realized even without a beam waist by simply exploiting the crystal rotation.

#### 4. Decoherence from off-resonant light scattering

We separately estimate the contributions from the ODF beams and the Raman beams and find them to be [44, 45]

$$\frac{\Gamma_{\text{ODF}}}{2\pi} \approx 38 \text{ Hz}, \quad \frac{\Gamma_{\text{Raman}}}{2\pi} \approx 6 \text{ Hz}. \quad (\text{B12})$$

The total decoherence rate is then  $\Gamma_{\text{tot}}/(2\pi) = (\Gamma_{\text{ODF}} + \Gamma_{\text{Raman}})/(2\pi) \approx 44 \text{ Hz}$ .

With the chosen parameters, we estimate the typical time up to which the simulation can be run as  $Jt \approx J(1/\Gamma_{\text{tot}}) \sim 9$ . As we mention in the Main Text, the ratio of the coherent interaction to the decoherence rate can be enhanced by a number of means, including choosing a different ion species and transition, enhancing coherent coupling by parametric amplification and by working at a different ODF operating point.

#### Appendix C: Rotating-wave approximations

In deriving the two-channel model, we assumed that all other terms in the expansion Eq. (A6) could be considered rapidly oscillating. We now check whether this is true by first listing the various resonance conditions and estimating their values using real experimental parameters.

We first consider the term of the form  $AB \sin C \hat{\mathbf{s}}_j^-$ , for which the possible arguments for the complex exponentials are (up to an overall sign, which gives the hermitian conjugate process with  $\hat{\mathbf{s}}_j^+$ )

$$\begin{aligned} T_{11} &\equiv B_0 - \mu_r + \omega_n + \omega_r = \delta_n \\ T_{12} &\equiv B_0 - \mu_r + \omega_n - \omega_r = \delta_n - 2\omega_r \\ T_{13} &\equiv B_0 - \mu_r - \omega_n + \omega_r = \delta_n - 2\omega_n \\ T_{14} &\equiv B_0 - \mu_r - \omega_n - \omega_r = \delta_n - 2\omega_n - 2\omega_r \\ T_{15} &= B_0 + \mu_r + \omega_n + \omega_r = \delta_n + 2\mu_r \\ T_{16} &\equiv B_0 + \mu_r + \omega_n - \omega_r = \delta_n + 2\mu_r - 2\omega_r \\ T_{17} &\equiv B_0 + \mu_r - \omega_n + \omega_r = \delta_n + 2\mu_r - 2\omega_n \\ T_{18} &\equiv B_0 + \mu_r - \omega_n - \omega_r = -\delta_n + 2B_0. \end{aligned} \quad (\text{C1})$$

We next consider the term of the form  $B^2 \sin C \hat{\mathbf{s}}_j^-$ . Here, we have

$$\begin{aligned}
T_{21} &\equiv B_0 - \mu_r + \omega_n + \omega_k = \delta_1 + (\omega_n + \omega_k) - (\omega_1 + \omega_r) \\
T_{22} &\equiv B_0 - \mu_r + \omega_n - \omega_k = \delta_1 + (\omega_n - \omega_k) - (\omega_1 + \omega_r) \\
T_{23} &\equiv B_0 - \mu_r - \omega_n + \omega_k = \delta_1 + (\omega_k - \omega_n) - (\omega_1 + \omega_r) \\
T_{24} &\equiv B_0 - \mu_r - \omega_n - \omega_k = \delta_1 - (\omega_n + \omega_k) - (\omega_1 + \omega_r) \\
T_{25} &\equiv B_0 + \mu_r + \omega_n + \omega_k = \delta_1 + 2\mu_r + (\omega_n + \omega_k) - (\omega_1 + \omega_r) \\
T_{26} &\equiv B_0 + \mu_r + \omega_n - \omega_k = \delta_1 + 2\mu_r + (\omega_n - \omega_k) - (\omega_1 + \omega_r) \\
T_{27} &\equiv B_0 + \mu_r - \omega_n + \omega_k = \delta_1 + 2\mu_r + (\omega_k - \omega_n) - (\omega_1 + \omega_r) \\
T_{28} &\equiv B_0 + \mu_r - \omega_n - \omega_k = \delta_1 + 2\mu_r - (\omega_n + \omega_k) - (\omega_1 + \omega_r).
\end{aligned} \tag{C2}$$

Next, we consider terms of the form  $A^2 \sin C \hat{\mathbf{s}}_j^-$ . For these terms, we get

$$\begin{aligned}
T_{31} &\equiv B_0 - \mu_r + \omega_r + \omega_r = \delta_n + \omega_r - \omega_n \\
T_{32} &\equiv B_0 - \mu_r + \omega_r - \omega_r = \delta_n - \omega_r - \omega_n \\
T_{33} &\equiv B_0 - \mu_r - \omega_r + \omega_r = \delta_n - \omega_r - \omega_n \\
T_{34} &\equiv B_0 - \mu_r - \omega_r - \omega_r = \delta_n - 3\omega_r - \omega_n \\
T_{35} &\equiv B_0 + \mu_r + \omega_r + \omega_r = \delta_n + 2\mu_r + \omega_r - \omega_n \\
T_{36} &\equiv B_0 + \mu_r + \omega_r - \omega_r = \delta_n + 2\mu_r - \omega_r - \omega_n \\
T_{37} &\equiv B_0 + \mu_r - \omega_r + \omega_r = \delta_n + 2\mu_r - \omega_r - \omega_n \\
T_{38} &\equiv B_0 + \mu_r - \omega_r - \omega_r = \delta_n + 2\mu_r - 3\omega_r - \omega_n.
\end{aligned} \tag{C3}$$

Now, we consider terms of the form  $\sin C \hat{\mathbf{s}}_j^-$ . Here, the complex exponentials are simply  $T_{41} \equiv B_0 + \mu_r$  and  $T_{42} \equiv B_0 - \mu_r$ , which evaluate to  $\delta_n - \omega_n - \omega_r$  and  $\delta_n + 2\mu_r - \omega_n - \omega_r$ .

We now turn to the term of the form  $B \cos C \hat{\mathbf{s}}_j^-$ . Here, we have the following detunings

$$\begin{aligned}
T_{51} &\equiv B_0 - \mu_r + \omega_n = \delta_n - \omega_r \\
T_{52} &\equiv B_0 - \mu_r - \omega_n = \delta_n - 2\omega_n - \omega_r \\
T_{53} &\equiv B_0 + \mu_r + \omega_n = \delta_n + 2\mu_r - \omega_r \\
T_{54} &\equiv B_0 + \mu_r - \omega_n = \delta_n + 2\mu_r - 2\omega_n - \omega_r.
\end{aligned} \tag{C4}$$

Finally, we consider the term of the form  $A \cos C \hat{\mathbf{s}}_j^-$ . Here, we get

$$\begin{aligned}
T_{61} &\equiv B_0 - \mu_r + \omega_r = \delta_n - \omega_n \\
T_{62} &\equiv B_0 - \mu_r - \omega_r = \delta_n - 2\omega_r - \omega_n \\
T_{63} &\equiv B_0 + \mu_r + \omega_r = \delta_n + 2\mu_r - \omega_n \\
T_{64} &\equiv B_0 + \mu_r - \omega_r = \delta_n + 2\mu_r - 2\omega_r - \omega_n.
\end{aligned} \tag{C5}$$

In order to estimate these expressions, we compute the equilibrium crystal structure for cases A and B, and obtain the drumhead mode spectrum to obtain the frequencies  $\omega_n$ . In Table I, we provide the maximum and minimum values possible for each of these terms and for cases A and B. If the maximum and minimum values are large compared to  $|\delta_1|$  and have the same sign, then there are no accidental resonances and the terms can be safely neglected. We have excluded the c.m. term when evaluating the range of  $T_{11}$  as this is precisely the term of interest with  $\delta_1/(2\pi) \leq 2$  kHz. All these terms are far off-resonant and preliminary numerical studies suggest that they can be neglected as a first approximation for simulating the two-channel model if the c.m. mode is tuned close to resonance, i.e.  $\delta_1/(2\pi) \lesssim 1$  kHz.

#### Appendix D: One-channel model and off-resonant terms

For simulating the one-channel model, we assumed that  $\delta_1/(2\pi) = 2$  kHz constitutes a large detuning based on which the c.m. mode can be adiabatically eliminated. The resulting interaction is fourth order in the small parameters  $A, B$  of the small angle expansion Eq. (A6). This raises the question whether some of the other terms in Eq. (A6), although off-resonant, could potentially compete with the effective one-channel interactions because they are of lower order in the small parameters. In this section, we consider the effect of terms that are zeroth and first order in the small parameters on the effective dynamics of the one-channel model. We also estimate off-resonant effects that arise from the second-order term  $AB \sin C$  since some terms in this group are not very far off-resonant.

term	case A	case B
$T_{11}$	(-527, -19)	(-505, -21)
$T_{12}$	(-887, -358)	(-2305, -1798)
$T_{13}$	(-3178, -2649)	(-6849, -6343)
$T_{14}$	(-3538, -3009)	(-8649, -8143)
$T_{15}$	(3029, 3558)	(8163, 8669)
$T_{16}$	(2669, 3198)	(6363, 6869)
$T_{17}$	(378, 907)	(1818, 2325)
$T_{18}$	(18, 547)	(18, 525)
$T_{21}$	(354, 1412)	(1514, 2528)
$T_{22}$	(-2297, -1239)	(-4830, -3817)
$T_{23}$	(-2297, -1239)	(-4830, -3817)
$T_{24}$	(-4948, -3890)	(-11175, -10162)
$T_{25}$	(3910, 4968)	(10182, 11195)
$T_{26}$	(1259, 2317)	(3837, 4850)
$T_{27}$	(1259, 2317)	(3837, 4850)
$T_{28}$	(-1392, -334)	(-2508, -1494)
$T_{31}$	(-1408, -1408)	(-2524, -2524)
$T_{32}$	(-1768, -1768)	(-4324, -4324)
$T_{33}$	(-1768, -1768)	(-4324, -4324)
$T_{34}$	(-2128, -2128)	(-6124, -6124)
$T_{35}$	(2148, 2148)	(6144, 6144)
$T_{36}$	(1788, 1788)	(4344, 4344)
$T_{37}$	(1788, 1788)	(4344, 4344)
$T_{38}$	(1428, 1428)	(2544, 2544)
$T_{41}$	(1788, 1788)	(4344, 4344)
$T_{42}$	(-1768, -1768)	(-4324, -4324)
$T_{51}$	(-707, -178)	(-1405, -898)
$T_{52}$	(-3358, -2829)	(-7749, -7243)
$T_{53}$	(2849, 3378)	(7263, 7769)
$T_{54}$	(198, 727)	(918, 1425)
$T_{61}$	(-1588, -1588)	(-3424, -3424)
$T_{62}$	(-1948, -1948)	(-5224, -5224)
$T_{63}$	(1968, 1968)	(5244, 5244)
$T_{64}$	(1608, 1608)	(3444, 3444)

TABLE I. Minimum and maximum values of various resonance conditions, showing that all unwanted terms are far off-resonant for both trapping parameters, viz. case A and case B. The frequencies are reported in units of kilohertz.

### 1. $\sin C$ term

This term is of the form

$$\hat{H}_4 = \sum_j \frac{\delta_{AC}}{2i} \left( \hat{\mathbf{s}}_j^+ e^{-i(\mu_r + B_j)t} + \hat{\mathbf{s}}_j^- e^{-i(\mu_r - B_j)t} \right) + \text{h.c.} \quad (\text{D1})$$



Using effective Hamiltonian theory (EHT) and assuming  $B_j \ll \mu_r$  and  $B_j \approx B_0$  leads to

$$\hat{H}_{4,\text{eff}} = \sum_j \frac{\delta_{\text{AC}}^2 B_0}{\mu_r^2} \hat{\mathbf{s}}_j^z. \quad (\text{D2})$$

This term represents a small AC Stark shift that leads to collective spin precession at a frequency  $\sim 5$  Hz for case A and  $\sim 0.9$  Hz for case B. Hence, its effect can be considered negligible.

## 2. $A \cos C$ term

This term is of the form

$$\begin{aligned} \hat{H}_6 = \sum_j \frac{\delta_{\text{AC}} \eta_x \tilde{r}_j}{4} & \left( \hat{\mathbf{s}}_j^+ e^{-i[(\mu_r + \omega_r + B_j)t + \phi_j]} + \hat{\mathbf{s}}_j^+ e^{-i[(\mu_r - \omega_r + B_j)t - \phi_j]} \right. \\ & \left. + \hat{\mathbf{s}}_j^- e^{-i[(\mu_r + \omega_r - B_j)t + \phi_j]} + \hat{\mathbf{s}}_j^- e^{-i[(\mu_r - \omega_r - B_j)t - \phi_j]} \right) + \text{h.c.} \end{aligned} \quad (\text{D3})$$

Using EHT leads to

$$\hat{H}_{6,\text{eff}} = - \sum_j \frac{\delta_{\text{AC}}^2 \eta_x^2 \tilde{r}_j^2}{8} \left[ \frac{1}{\mu_r + \omega_r + B_j} + \frac{1}{\mu_r - \omega_r + B_j} - \frac{1}{\mu_r + \omega_r - B_j} - \frac{1}{\mu_r - \omega_r - B_j} \right] \hat{\mathbf{s}}_j^z. \quad (\text{D4})$$

Assuming  $B_j \sim B_0 \ll \omega_r, \mu_r$ , we get

$$\hat{H}_{6,\text{eff}} = \sum_j \frac{\delta_{\text{AC}}^2 B_0 \eta_x^2 \tilde{r}_j^2}{4} \left[ \frac{1}{(\mu_r + \omega_r)^2} + \frac{1}{(\mu_r - \omega_r)^2} \right] \hat{\mathbf{s}}_j^z. \quad (\text{D5})$$

This term results in a radius-dependent AC Stark shift. We estimate its maximal value by setting  $\tilde{r}_j = 1$ , for which we find a precession frequency  $\sim 0.5$  Hz for case A and  $\sim 0.04$  Hz for case B. These are very small compared to the dispersion generated by the Raman beam waist and hence we neglect these terms.

## 3. $B \cos C$ term

This term is of the form

$$\begin{aligned} \hat{H}_5 = \sum_{j,n} \frac{\delta_{\text{AC}} \eta_n \mathcal{M}_{jn}}{2} & \left( \hat{\mathbf{s}}_j^+ \hat{a}_n e^{-i(\mu_r + B_0 + \omega_n)t} + \hat{\mathbf{s}}_j^+ \hat{a}_n^\dagger e^{-i(\mu_r + B_0 - \omega_n)t} \right. \\ & \left. + \hat{\mathbf{s}}_j^- \hat{a}_n e^{-i(\mu_r - B_0 + \omega_n)t} + \hat{\mathbf{s}}_j^- \hat{a}_n^\dagger e^{-i(\mu_r - B_0 - \omega_n)t} \right) + \text{h.c.} \end{aligned} \quad (\text{D6})$$

For applying EHT, we evaluate the following commutators

$$\begin{aligned} [\hat{\mathbf{s}}_j^+ \hat{a}_n, \hat{\mathbf{s}}_k^- \hat{a}_m^\dagger] &= 2\hat{a}_m^\dagger \hat{a}_n \hat{\mathbf{s}}_j^z \delta_{jk} + \delta_{nm} \hat{\mathbf{s}}_j^+ \hat{\mathbf{s}}_k^-, \\ [\hat{\mathbf{s}}_j^+ \hat{a}_n^\dagger, \hat{\mathbf{s}}_k^- \hat{a}_m] &= 2\hat{a}_m \hat{a}_n^\dagger \hat{\mathbf{s}}_j^z \delta_{jk} - \delta_{nm} \hat{\mathbf{s}}_j^+ \hat{\mathbf{s}}_k^-, \\ [\hat{\mathbf{s}}_j^- \hat{a}_n, \hat{\mathbf{s}}_k^+ \hat{a}_m^\dagger] &= -2\hat{a}_m^\dagger \hat{a}_n \hat{\mathbf{s}}_j^z \delta_{jk} + \delta_{nm} \hat{\mathbf{s}}_j^- \hat{\mathbf{s}}_k^+, \\ [\hat{\mathbf{s}}_j^- \hat{a}_n^\dagger, \hat{\mathbf{s}}_k^+ \hat{a}_m] &= -2\hat{a}_m \hat{a}_n^\dagger \hat{\mathbf{s}}_j^z \delta_{jk} - \delta_{nm} \hat{\mathbf{s}}_j^- \hat{\mathbf{s}}_k^+. \end{aligned} \quad (\text{D7})$$

We only consider the effective role of each term in the parenthesis independently. We briefly comment on cross-resonances between different terms in the parentheses as well as between different groups of terms (in particular  $B \cos C$  and  $AB \sin C$ ) at the end of this section. With the assumption of ground state cooling, we make the replacement  $\hat{a}_m^\dagger \hat{a}_n \rightarrow 0$  and  $\hat{a}_m \hat{a}_n^\dagger \rightarrow \delta_{nm}$ . We then have the effective Hamiltonian

$$\begin{aligned} \hat{H}_{5,\text{eff}} = & - \sum_j \sum_n \frac{\delta_{\text{AC}}^2 \eta_n^2 \mathcal{M}_{jn}^2}{4} \left[ \frac{1}{\mu_r + B_0 + \omega_n} + \frac{1}{\mu_r + B_0 - \omega_n} - \frac{1}{\mu_r - B_0 + \omega_n} - \frac{1}{\mu_r - B_0 - \omega_n} \right] \hat{\mathbf{s}}_j^z \\ & - \sum_{j \neq k} \sum_n \frac{\delta_{\text{AC}}^2 \eta_n^2 \mathcal{M}_{jn} \mathcal{M}_{kn}}{4} \left[ \frac{1}{\mu_r + B_0 + \omega_n} - \frac{1}{\mu_r + B_0 - \omega_n} + \frac{1}{\mu_r - B_0 + \omega_n} - \frac{1}{\mu_r - B_0 - \omega_n} \right] \hat{\mathbf{s}}_j^+ \hat{\mathbf{s}}_k^-. \end{aligned} \quad (\text{D8})$$

As a first approximation, we neglect terms with  $\mu_r + \omega_n$  in the denominator since they are small compared with terms that have  $\mu_r - \omega_n$  in the denominator. Using the fact that  $B_0 \ll \mu_r - \omega_n$ , we arrive at the approximate effective Hamiltonian

$$\hat{H}_{5,\text{eff}} = \sum_{j,k} J_{5,jk} \hat{\mathbf{s}}_j^+ \hat{\mathbf{s}}_k^-, \quad (\text{D9})$$

where the interaction matrix  $J_5$  has elements given by

$$J_{5,jk} = \begin{cases} \sum_n \frac{\delta_{\text{AC}}^2 B_0 \eta_n^2 \mathcal{M}_{jn}^2}{2(\mu_r - \omega_n)^2}, & j = k \\ \sum_n \frac{\delta_{\text{AC}}^2 \eta_n^2 \mathcal{M}_{jn} \mathcal{M}_{kn}}{2(\mu_r - \omega_n)}, & j \neq k. \end{cases} \quad (\text{D10})$$

The effective Hamiltonian  $\hat{H}_{5,\text{eff}}$  mediates achiral spin-exchange type interactions that directly compete with the chiral spin-exchange that we wish to engineer. We study the impact of this term numerically in Fig. 5 for case A and case B using mean-field theory. We find that in case A, this interaction causes  $|\psi(t)|$  to rapidly decay toward zero on short timescales whereas in case B, the impact of this term is rather small. From the form of the elements in the coupling matrix  $J_5$ , a larger rotation frequency  $\omega_r$  increases the denominator, i.e. makes these terms smaller and hence their impact is smaller in case B.

#### 4. $AB \sin C$ term

The interaction giving rise to the two-channel model is present in this second-order term and was discussed previously. Here, we estimate the contribution of other terms present in this interaction since some of them are not very far off-resonant.

This term is of the form

$$\hat{H}_1 = - \sum_{j,n} \frac{\delta_{\text{AC}} \eta_x \tilde{r}_j \eta_n \mathcal{M}_{jn}}{4i} \left( \hat{\mathbf{s}}_j^+ \hat{a}_n^\dagger e^{-i[(\mu_r - \omega_n - \omega_r + B_0)t - \phi_j]} + \hat{\mathbf{s}}_j^+ \hat{a}_n^\dagger e^{-i[(\mu_r - \omega_n + \omega_r + B_0)t + \phi_j]} \right. \\ \left. + \hat{\mathbf{s}}_j^- \hat{a}_n^\dagger e^{-i[(\mu_r - \omega_n - \omega_r - B_0)t - \phi_j]} + \hat{\mathbf{s}}_j^- \hat{a}_n^\dagger e^{-i[(\mu_r - \omega_n + \omega_r - B_0)t + \phi_j]} \right). \quad (\text{D11})$$

In writing the above equation, we have already ignored the terms that have exponentials containing the combination  $\mu_r + \omega_n$ . Following a calculation similar to the  $B \cos C$  term, we find an effective Hamiltonian given by

$$\hat{H}_{1,\text{eff}} = \sum_{j,k} J_{11,jk} \hat{\mathbf{s}}_j^+ \hat{\mathbf{s}}_k^- + \sum_{j,k} J_{12,jk} \hat{\mathbf{s}}_j^+ \hat{\mathbf{s}}_k^-, \quad (\text{D12})$$

where the interaction matrices  $J_{11}, J_{1,2}$  have elements given by

$$J_{11,jk} = \begin{cases} \sum_n \frac{\delta_{\text{AC}}^2 \eta_x^2 \tilde{r}_j^2 \eta_n^2 \mathcal{M}_{jn}^2}{16(\delta_n - 2B_0)}, & j = k \\ - \sum_n \frac{\delta_{\text{AC}}^2 \eta_x^2 \tilde{r}_j \tilde{r}_k \eta_n^2 \mathcal{M}_{jn} \mathcal{M}_{kn}}{16} \left[ \frac{1}{\delta_n - 2B_0} + \frac{1}{\delta_n - 2\omega_r} \right] e^{i(\phi_j - \phi_k)}, & j \neq k, \end{cases} \\ J_{12,jk} = \begin{cases} - \sum_n \frac{\delta_{\text{AC}}^2 \eta_x^2 \tilde{r}_j^2 \eta_n^2 \mathcal{M}_{jn}^2}{16\delta_n}, & j = k \\ - \sum_n \frac{\delta_{\text{AC}}^2 \eta_x^2 \tilde{r}_j \tilde{r}_k \eta_n^2 \mathcal{M}_{jn} \mathcal{M}_{kn}}{16} \left[ \frac{1}{\delta_n - 2\omega_r} + \frac{1}{\delta_n} \right] e^{-i(\phi_j - \phi_k)}, & j \neq k. \end{cases} \quad (\text{D13})$$

The  $J_{12}$  matrix now represents the chiral spin-exchange interactions arising from all drumhead modes. On the other hand, the  $J_{11}$  matrix describes anti-chiral interactions mediated by these modes. We estimate the impact of these terms numerically in Fig. 5 and find that they do not significantly impact the one-channel model dynamics in both cases A and B.

Our study of off-resonant interactions shows that trapping parameters can be found where the impact of these terms can be neglected for realizing the one-channel model. We have also briefly looked at the possibility of cross-resonances between the individual terms in the  $B \cos C$  and  $AB \sin C$  subgroups as well as between these groups. While accidental near-resonances may occur between terms involving a drumhead mode  $n$  at frequency  $\omega_n$  and, say, a mode  $m$  at  $\omega_m \approx \omega_n \pm 2B_0$  or  $\omega_m \approx \omega_n \pm \omega_r$ , such terms typically come with a prefactor  $\hat{a}_n^\dagger \hat{a}_m$  and only lead to




FIG. 5. Impact of off-resonant interactions on the one-channel model dynamics for crystals formed under two different trapping parameters, (a) case A and (b) case B (see Appendix B 1). In both cases, we progressively add terms to the pure one-channel model and study their impact. We add the following terms in the specified order: Chiral spin-exchange by all modes ( $J_{12}$ ), anti-chiral spin-exchange by all modes ( $J_{11}$ ) and achiral spin-exchange by all modes ( $J_5$ ).

single-spin terms, and hence they may possibly be neglected if the drumhead modes are near-ground-state cooled. A brief inspection of other possible cross-resonances between terms up to second order indicates that the resulting effective terms are small single-spin terms that don't affect the spin-spin interaction term.

We end this section by noting that we have studied terms in the small angle expansion (A6) up to second order in the small parameters and explored the effective impact of off-resonant terms that appear important on the dynamics of interest. However, we have not considered possible accidental resonances or near-resonances that may occur at third and higher order in the small angle expansion because of the sheer number of terms that appear at higher orders. Their impact and the parameter regimes where they are negligible could potentially be explored directly on the quantum simulator.

## Appendix E: Numerical methods

While the classification of dynamical phases is based on a mean-field study in the thermodynamic limit, the finite size of our system has motivated us to consider beyond-mean-field techniques for numerical solution. Here, we present the mean-field equations for the one-channel and two-channel models, discuss how we include the quantum noise of the initial state using the discrete truncated Wigner approximation (dTWA) method, and benchmark the performance of dTWA using special crystals where the geometry allows for exact numerical solution of the Schrodinger equation.

### 1. Mean-field equations of motion for one-channel and two-channel models

In the mean-field theory for the one-channel model, we replace the spin vector operator  $\hat{\mathbf{s}}_j$  at each site  $j$  by a vector of  $c$ -numbers  $\mathbf{s}_j \equiv (\mathbf{s}_j^x, \mathbf{s}_j^y, \mathbf{s}_j^z)$ . The commutation relations are replaced by Poisson brackets,  $\{\mathbf{s}_j^a, \mathbf{s}_k^b\} = i\epsilon_{abc}\mathbf{s}_j^c\delta_{jk}$ . Defining  $\mathbf{s}_j^\pm = \mathbf{s}_j^x \pm i\mathbf{s}_j^y$ , the dynamics under Hamiltonian (A16) are given by

$$\begin{aligned} \frac{d}{dt}\mathbf{s}_j^+ &= iK\tilde{r}_j^2\mathbf{s}_j^+ + iJ\tilde{r}_j\mathbf{s}_j^z e^{i\phi_j}\Psi_j^*, \\ \frac{d}{dt}\mathbf{s}_j^z &= -J\tilde{r}_j\text{Im}[\mathbf{s}_j^+ e^{-i\phi_j}\Psi_j] \text{,} \end{aligned} \quad (\text{E1})$$

where  $\Psi_j = (2/N)\sum_{k\neq j}\tilde{r}_k\mathbf{s}_k^- e^{i\phi_k}$ .

To obtain the mean-field equations for the two-channel model governed by Hamiltonian (A10), we additionally

replace the annihilation operator  $\hat{a}_1$  of the c.m. mode by a complex amplitude  $\alpha_1$ . The resulting equations are

$$\begin{aligned}\frac{d}{dt}\mathfrak{s}_j^+ &= iB_1\tilde{r}_j^2\mathfrak{s}_j^+ + \frac{2G}{\sqrt{N}}\alpha_1^*\mathfrak{s}_j^Z\tilde{r}_je^{i\phi_j}, \\ \frac{d}{dt}\mathfrak{s}_j^Z &= -\frac{2G}{\sqrt{N}}\tilde{r}_j\text{Re}[s_j^+\alpha_1e^{-i\phi_j}], \\ \frac{d}{dt}\alpha_1 &= -i\delta_0\alpha_1 + \frac{\sqrt{N}G}{2}\Psi,\end{aligned}\tag{E2}$$

with  $\Psi = (2/N)\sum_j\tilde{r}_j\mathfrak{s}_j^-e^{i\phi_j}$ .

For mean-field simulations, the above equations are numerically evolved starting from initial conditions where the  $c$ -numbers are assigned the expectation values of the corresponding quantum operators in the initial state.

## 2. Accounting for initial quantum noise

For finite size systems, quantum corrections to the mean-field dynamics become important. To explore their effects, we simulate the effects of the quantum noise of the initial state by evolving several trajectories under the mean-field equations starting from randomly drawn initial conditions. For the spin degrees of freedom, we first find the mean spin direction  $\hat{\mathbf{e}}_j^{\parallel}$  in the initial state. Next, we identify two mutually orthogonal spin directions,  $\hat{\mathbf{e}}_j^{\perp,1}, \hat{\mathbf{e}}_j^{\perp,2}$ , in the plane perpendicular to the mean spin. The initial spin vector can then be written as

$$\mathfrak{s}_j = \mathfrak{s}_j^{\parallel}\hat{\mathbf{e}}_j^{\parallel} + \mathfrak{s}_j^{\perp,1}\hat{\mathbf{e}}_j^{\perp,1} + \mathfrak{s}_j^{\perp,2}\hat{\mathbf{e}}_j^{\perp,2}.\tag{E3}$$

For mean-field simulations, we set  $\mathfrak{s}_j^{\parallel} = 1/2, \mathfrak{s}_j^{\perp,1} = \mathfrak{s}_j^{\perp,2} = 0$ . To go beyond mean-field, we use the prescription of the discrete truncated Wigner approximation [21], according to which  $\mathfrak{s}_j^{\perp,1}$  and  $\mathfrak{s}_j^{\perp,2}$  are independently and randomly chosen to be  $\pm 1/2$  with equal probability.

In the case of the two-channel model, we additionally draw the complex amplitude  $\alpha_1$  from the Wigner distribution of the initial state of the c.m. mode, which we always assume is the motional ground state in this work. Therefore, the real and imaginary parts of  $\alpha_1$  are independently drawn from a Gaussian distribution with zero mean and a variance of  $1/4$ .

## 3. Benchmarking the dTWA results

In the case of a real crystal in a Penning trap, the triangular lattice is only approximate and hence every ion is typically at a slightly different radius from the trap center. This makes an exact solution of the Schrodinger equation subject to Hamiltonian (A16) exponentially hard. Therefore, in order to test the reliability of the dTWA results, we test this technique on a hypothetical ideal crystal made of  $M$  concentric rings of ions, for which an exact numerical solution to the Schrodinger equation is feasible. We assume that the number of ions  $N_m$  in ring  $m = 1, \dots, M$  is given by  $6(m-1) + \delta_{m,1}$ , which mimics the hexagonal lattice structure of Penning trap crystals. Here, the first ring  $m = 1$  is taken to be the single ion at the crystal center. We can then define total angular momentum operators for each ring,  $\hat{\mathfrak{J}}_m^{\pm}, \hat{\mathfrak{J}}_m^Z$  as

$$\hat{\mathfrak{J}}_m^{\pm} = \sum_{j \in \text{ring } m} \hat{\mathfrak{s}}_j^{\pm} e^{\mp i\phi_j}, \quad \hat{\mathfrak{J}}_m^Z = \sum_{j \in \text{ring } m} \hat{\mathfrak{s}}_j^Z.\tag{E4}$$

These operators are readily seen to obey the usual angular momentum commutation relations. In terms of these operators, the one-channel model (A16) can be expressed as

$$\hat{H}_{\text{eff}} = B_1 \sum_{m=1}^M \tilde{r}_m^2 \hat{\mathfrak{J}}_m^Z - \frac{J}{N} \sum_{m,m'=1}^M \tilde{r}_m \tilde{r}_{m'} \hat{\mathfrak{J}}_m^+ \hat{\mathfrak{J}}_{m'}^-\tag{E5}$$

For a crystal with  $M$  rings, the total number of ions is  $N(M) = 1 + 3M(M-1)$ . The computational complexity is significantly reduced in the total angular momentum picture, because, for the initial states we consider, we only need to track the fully symmetric subspace of each ring. Therefore, the number of basis states in each ring is reduced from  $2^{N_m}$  to  $N_m + 1$ , thereby enabling the rapid simulation of exact dynamics for crystals with up to  $M = 5$  rings ( $N(5) = 61$  ions). Figure 6 shows the excellent agreement of the dTWA calculation with the exact solution for crystals with  $M = 4$  and  $M = 5$  rings, confirming the validity of the dTWA technique for beyond-mean-field calculations in this work.




FIG. 6. Benchmarking the discrete truncated Wigner approximation (dTWA) method. Solid lines are computed by numerical propagation of the Schrodinger equation while the dotted lines are obtained using the dTWA method. Here, we assume that the crystal is made of perfectly concentric rings of ions and the spin state is initialized in a BCS-like state according to Appendix G 1. Here, we have used the ratio  $K/J = 1$ .

### Appendix F: Computation of winding numbers

In continuous 2D space, the winding number of a unit vector field  $\hat{\mathbf{u}}(x, y)$  is defined as the surface integral

$$W = \frac{1}{4\pi} \int dx dy \hat{\mathbf{u}} \cdot \left( \frac{d\hat{\mathbf{u}}}{dx} \times \frac{d\hat{\mathbf{u}}}{dy} \right). \quad (\text{F1})$$

In the crystal, the winding number calculation must be carried out on a lattice with discrete sites. Here, the prescription is to identify triplets of neighbors by introducing a triangulation of the crystal lattice as shown in Fig. 7 [46]. We use the Delaunay triangulation, wherein triangles are formed between neighboring triplets in such a way that no vertex of the crystal lies inside the circumcircle of each triangle. For each triangle, we label the vertices  $A, B, C$  such that the directed edges give rise to a face normal pointing upward from the crystal plane, i.e.  $\overrightarrow{AB} \times \overrightarrow{BC} \parallel \hat{\mathbf{e}}_z$ . Having identified such ordered triplets of vertices, a solid angle  $\Omega_{ABC}$  is introduced for each triangle, defined as

$$\tan \left( \frac{\Omega_{ABC}}{2} \right) = \frac{\hat{\mathbf{u}}_A \cdot (\hat{\mathbf{u}}_B \times \hat{\mathbf{u}}_C)}{1 + \hat{\mathbf{u}}_A \cdot \hat{\mathbf{u}}_B + \hat{\mathbf{u}}_B \cdot \hat{\mathbf{u}}_C + \hat{\mathbf{u}}_C \cdot \hat{\mathbf{u}}_A}. \quad (\text{F2})$$

The winding number on the discrete lattice is obtained by summing the solid angle  $\Omega_{ABC}$  over all triangles  $\Delta_{ABC}$  of the triangulation:

$$W = \frac{1}{4\pi} \sum_{\Delta_{ABC}} \Omega_{ABC}. \quad (\text{F3})$$

In the present work, we call the winding number  $Q$  if the vector field is taken as the spin texture, while we label it as  $W$  if the vector field is the effective magnetic field texture in the frame rotating at  $2\mu_\infty$ .

### Appendix G: State initialization

To generate chiral BCS-like and BEC-like initial states, we take advantage of the term of the form  $A \cos C \hat{\mathbf{s}}_j^-$  that is available in the small-angle expansion of the ODF interaction [Eq. (A6)]. This term describes a coupling of the spins with the planar rotation without involving the drumhead c.m. mode. We assume that the ODF lasers have a tunable beam waist  $w_{\text{ODF}}$ . By tuning  $\mu_r = B_0 + \omega_r$  and ignoring rapidly oscillating terms, the effective interaction is given by

$$\hat{H}_{\text{init}} = \sum_j \frac{\Omega_j}{2} \tilde{r}_j (\hat{\mathbf{s}}_j^+ e^{-i\phi_j} + \hat{\mathbf{s}}_j^- e^{i\phi_j}), \quad (\text{G1})$$

where  $\Omega_j = \Omega_0 e^{-r_j^2/w_{\text{ODF}}^2}$  with  $\Omega_0 = \delta_{AC}(\Delta k_x R)/2$  and  $\tilde{r}_j = r_j/R$ . The Hamiltonian  $\hat{H}_{\text{init}}$  describes non-interacting spins each undergoing rotation under a local magnetic field. Using the local axes introduced in Eq. (14), we can




FIG. 7. Demonstration of Delaunay triangulation for a crystal with 200 ions. The triangulation is used to identify triplets of neighbors on the discrete lattice for computing the winding number of the spin texture ( $Q$ ) or that of the effective magnetic field texture ( $W$ ).




FIG. 8. Initial spin textures: (a) BCS texture for phase I and II, (b) BEC texture for winding number studies and (c) BCS initialization for phase III.

compactly write

$$\hat{H}_{\text{init}} = \sum_j \Omega_j \hat{\mathbf{s}}_j^y. \quad (\text{G2})$$

For the initialization, we assume that the beam waist of the Raman beams is much larger than the crystal radius, i.e.  $w \gg R$  so that the dispersion arising from the spatial variation of the Raman beams can be neglected.

We now describe the initialization protocols for various cases that we discuss in the Main Text. A plot of these spin textures is shown in Fig. 8.

### 1. BCS initialization for phases I and II

We assume that the beam waist of the ODF lasers are much larger than the crystal radius, i.e.,  $w_{\text{ODF}} \gg R$ . Then,  $\hat{H}_{\text{init}}$  reduces to

$$\hat{H}_{\text{init}} = \Omega_0 \sum_j \tilde{r}_j \hat{\mathbf{s}}_j^y. \quad (\text{G3})$$

We initialize all spins in  $|\uparrow\rangle_{\mathcal{Z}}$  (i.e. in the rotated spin space) by an appropriate global  $\pi/2$  pulse. Setting the maximum pulse area  $\Omega_0 T = \pi$ , the outermost spin are then rotated all the way to  $|\downarrow\rangle_{\mathcal{Z}}$  whereas the central spin is unaffected by virtue of the dependence of the Rabi frequency on  $\tilde{r}_j$ . Data shown in Fig. 2(a-b) and Fig. 4 are obtained using this initial state.

## 2. BEC initialization for winding number studies

We assume that  $w_{\text{ODF}} < \sqrt{2}R$ . The radial modulation of the Rabi frequency results in a maximum Rabi frequency  $\Omega_{\text{max}}$  at radius  $r_{\text{max}}$  that are respectively given by

$$\Omega_{\text{max}} = \Omega_0 \frac{w_{\text{ODF}}}{R\sqrt{2e}}, \quad r_{\text{max}} = \frac{w_{\text{ODF}}}{\sqrt{2}}. \quad (\text{G4})$$

Therefore, for  $w_{\text{ODF}} < \sqrt{2}R$ , the maximum pulse area for a fixed rotation time  $T$  is experienced by a spin somewhere in the middle of the crystal. We initialize all spins in  $|\downarrow\rangle_{\mathcal{Z}}$ . For the winding number study, we set the maximum pulse area  $\Omega_{\text{max}}T = \pi$  and use a beam waist  $w_{\text{ODF}} = 0.3R$ , which ensures that a large number of the outer spins are negligibly rotated. This ensures that the winding number  $W$  is reasonably quantized. Data shown in Fig. 3 are obtained using this initial state.

## 3. BCS initialization for phase III

Preparing the initial state for phase III requires the presence of a sharp domain wall and an order parameter of small magnitude. To obtain the domain wall, first spins are initialized in  $|\downarrow\rangle_{\mathcal{Z}}$  (i.e. in the unrotated spin space). An optical pumping beam selectively excites spins in a central region of radius  $r_d$  (chosen to be  $r_d = R/2$ ) to  $|\uparrow\rangle_{\mathcal{Z}}$ . A  $\pi/2$  pulse about the  $Y$ -axis then respectively converts the central and outer regions to domains of  $|\uparrow\rangle_{\mathcal{Z}}$  and  $|\downarrow\rangle_{\mathcal{Z}}$  spins (i.e. in the rotated spin space). To initialize a small order parameter, we assume  $w_{\text{ODF}} = R/2$ . We set the maximum pulse area to be  $\Omega_{\text{max}}T = 0.1\pi$ , i.e. the spin rotation is through rather small angles. Furthermore, since the spins in the different domains start in opposite orientations, a cancellation occurs resulting in a very small magnitude for the initial order parameter. Data shown in Fig. 2(c) are obtained using this initial state.

- 
- [1] R. A. Barankov and L. S. Levitov, “Synchronization in the bcs pairing dynamics as a critical phenomenon,” *Phys. Rev. Lett.* **96**, 230403 (2006).
  - [2] E. A. Yuzbashyan, M. Dzero, V. Gurarie, and M. S. Foster, “Quantum quench phase diagrams of an  $s$ -wave bcs-bec condensate,” *Phys. Rev. A* **91**, 033628 (2015).
  - [3] Robert J. Lewis-Swan, Diego Barberena, Julia R. K. Cline, Dylan J. Young, James K. Thompson, and Ana Maria Rey, “Cavity-qed quantum simulator of dynamical phases of a bardeen-cooper-schrieffer superconductor,” *Phys. Rev. Lett.* **126**, 173601 (2021).
  - [4] Matthew S. Foster, Maxim Dzero, Victor Gurarie, and Emil A. Yuzbashyan, “Quantum quench in a  $p+ip$  superfluid: Winding numbers and topological states far from equilibrium,” *Phys. Rev. B* **88**, 104511 (2013).
  - [5] C A Regal, C Ticknor, J L Bohn, and D S Jin, “Tuning  $p$ -wave interactions in an ultracold Fermi gas of atoms,” *Phys. Rev. Lett.* **90**, 053201 (2003).
  - [6] C. A. Regal, C. Ticknor, J. L. Bohn, and D. S. Jin, “Tuning  $p$ -wave interactions in an ultracold fermi gas of atoms,” *Phys. Rev. Lett.* **90**, 053201 (2003).
  - [7] C. H. Schunck, M. W. Zwierlein, C. A. Stan, S. M. F. Raupach, W. Ketterle, A. Simoni, E. Tiesinga, C. J. Williams, and P. S. Julienne, “Feshbach resonances in fermionic  ${}^6\text{Li}$ ,” *Phys. Rev. A* **71**, 045601 (2005).
  - [8] J P Gaebler, J T Stewart, J L Bohn, and D S Jin, “ $p$ -wave Feshbach molecules,” *Phys. Rev. Lett.* **98**, 200403 (2007).
  - [9] Kenneth Günter, Thilo Stöferle, Henning Moritz, Michael Köhl, and Tilman Esslinger, “ $p$ -wave interactions in low-dimensional fermionic gases,” *Phys. Rev. Lett.* **95**, 230401 (2005).
  - [10] Y.-J. Han, Y.-H. Chan, W. Yi, A. J. Daley, S. Diehl, P. Zoller, and L.-M. Duan, “Stabilization of the  $p$ -wave superfluid state in an optical lattice,” *Phys. Rev. Lett.* **103**, 070404 (2009).
  - [11] N. R. Cooper and G. V. Shlyapnikov, “Stable topological superfluid phase of ultracold polar fermionic molecules,” *Phys. Rev. Lett.* **103**, 155302 (2009).
  - [12] A. K. Fedorov, S. I. Matveenko, V. I. Yudson, and G. V. Shlyapnikov, “Novel  $p$ -wave superfluids of fermionic polar molecules,” *Scientific Reports* **6**, 27448 (2016).
  - [13] Matthew S. Foster, Victor Gurarie, Maxim Dzero, and Emil A. Yuzbashyan, “Quench-induced floquet topological  $p$ -wave superfluids,” *Phys. Rev. Lett.* **113**, 076403 (2014).
  - [14] Kevin A. Gilmore, Matthew Affolter, Robert J. Lewis-Swan, Diego Barberena, Elena Jordan, Ana Maria Rey, and John J. Bollinger, “Quantum-enhanced sensing of displacements and electric fields with two-dimensional trapped-ion crystals,” *Science* **373**, 673–678 (2021), <https://www.science.org/doi/pdf/10.1126/science.abi5226>.
  - [15] Martin Gärttner, Justin G Bohnet, Arghavan Safavi-Naini, Michael L Wall, John J Bollinger, and Ana Maria Rey, “Measuring out-of-time-order correlations and multiple quantum spectra in a trapped-ion quantum magnet,” *Nature Physics* **13**, 781–786 (2017).
  - [16] P. W. Anderson, “Random-phase approximation in the theory of superconductivity,” *Phys. Rev.* **112**, 1900–1916 (1958).

- [17] Athreya Shankar, Chen Tang, Matthew Affolter, Kevin Gilmore, Daniel H. E. Dubin, Scott Parker, Murray J. Holland, and John J. Bollinger, “Broadening of the drumhead-mode spectrum due to in-plane thermal fluctuations of two-dimensional trapped ion crystals in a penning trap,” *Phys. Rev. A* **102**, 053106 (2020).
- [18] R. A. Barankov and L. S. Levitov, “Synchronization in the bcs pairing dynamics as a critical phenomenon,” *Phys. Rev. Lett.* **96**, 230403 (2006).
- [19] D F James and J Jerke, “Effective hamiltonian theory and its applications in quantum information,” *Canadian Journal of Physics* **85**, 625–632 (2007), <https://doi.org/10.1139/p07-060>.
- [20] Wenchao Ge, Brian C. Sawyer, Joseph W. Britton, Kurt Jacobs, John J. Bollinger, and Michael Foss-Feig, “Trapped ion quantum information processing with squeezed phonons,” *Phys. Rev. Lett.* **122**, 030501 (2019).
- [21] J. Schachenmayer, A. Pikovski, and A. M. Rey, “Many-body quantum spin dynamics with monte carlo trajectories on a discrete phase space,” *Phys. Rev. X* **5**, 011022 (2015).
- [22] Lukas Schwarz, Benedikt Fauseweh, and Dirk Manske, “Momentum-resolved analysis of condensate dynamic and higgs oscillations in quenched superconductors with time-resolved arpes,” *Phys. Rev. B* **101**, 224510 (2020).
- [23] Jan van Delft, “Superconductivity in ultrasmall metallic grains,” *Annalen der Physik* **10**, 219–276 (2001).
- [24] Yunxiang Liao and Matthew S. Foster, “Spectroscopic probes of isolated nonequilibrium quantum matter: Quantum quenches, floquet states, and distribution functions,” *Phys. Rev. A* **92**, 053620 (2015).
- [25] Shane P Kelly, James K Thompson, Ana Maria Rey, and Jamir Marino, “Resonant light enhances phase coherence in a cavity qed simulator of fermionic superfluidity,” arXiv preprint arXiv:2202.05851 (2022).
- [26] Ryusuke Matsunaga, Naoto Tsuji, Hiroyuki Fujita, Arata Sugioka, Kazumasa Makise, Yoshinori Uzawa, Hirotaka Terai, Zhen Wang, Hideo Aoki, and Ryo Shimano, “Light-induced collective pseudospin precession resonating with Higgs mode in a superconductor,” *Science* **345**, 1145–1149 (2014).
- [27] Ryusuke Matsunaga, Yuki I. Hamada, Kazumasa Makise, Yoshinori Uzawa, Hirotaka Terai, Zhen Wang, and Ryo Shimano, “Higgs amplitude mode in the bcs superconductors  $\text{Nb}_{1-x}\text{Ti}_x\text{N}$  induced by terahertz pulse excitation,” *Phys. Rev. Lett.* **111**, 057002 (2013).
- [28] Ryusuke Matsunaga, Naoto Tsuji, Kazumasa Makise, Hirotaka Terai, Hideo Aoki, and Ryo Shimano, “Polarization-resolved terahertz third-harmonic generation in a single-crystal superconductor nbn: Dominance of the higgs mode beyond the bcs approximation,” *Phys. Rev. B* **96**, 020505 (2017).
- [29] M. Mitrano, A. Cantaluppi, D. Nicoletti, S. Kaiser, A. Perucchi, S. Lupi, P. Di Pietro, D. Pontiroli, M. Riccò, S. R. Clark, D. Jaksch, and A. Cavalleri, “Possible light-induced superconductivity in  $\text{k}_3\text{c}_{60}$  at high temperature,” *Nature* **530**, 461–464 (2016).
- [30] R. Mankowsky, A. Subedi, M. Först, S. O. Mariager, M. Chollet, H. T. Lemke, J. S. Robinson, J. M. Glowina, M. P. Minitti, A. Frano, M. Fechner, N. A. Spaldin, T. Loew, B. Keimer, A. Georges, and A. Cavalleri, “Non-linear lattice dynamics as a basis for enhanced superconductivity in  $\text{YBa}_2\text{Cu}_3\text{O}_{6.5}$ ,” *Nature* **516**, 71–73 (2014).
- [31] Kazuki Isoyama, Naotaka Yoshikawa, Kota Katsumi, Jeremy Wong, Naoki Shikama, Yuki Sakishita, Fuyuki Nabeshima, Atsutaka Maeda, and Ryo Shimano, “Light-induced enhancement of superconductivity in iron-based superconductor  $\text{FeSe}_{0.5}\text{Te}_{0.5}$ ,” *Communications Physics* **4**, 160 (2021).
- [32] M. A. Sentef, M. Ruggenthaler, and A. Rubio, “Cavity quantum-electrodynamical polaritonically enhanced electron-phonon coupling and its influence on superconductivity,” *Science Advances* **4**, eaau6969 (2018), <https://www.science.org/doi/pdf/10.1126/sciadv.aau6969>.
- [33] Frank Schlawin and Dieter Jaksch, “Cavity-mediated unconventional pairing in ultracold fermionic atoms,” *Phys. Rev. Lett.* **123**, 133601 (2019).
- [34] Jonathan B. Curtis, Zachary M. Raines, Andrew A. Allocca, Mohammad Hafezi, and Victor M. Galitski, “Cavity quantum eliasberg enhancement of superconductivity,” *Phys. Rev. Lett.* **122**, 167002 (2019).
- [35] Anoop Thomas, Eloïse Devaux, Kalaivanan Nagarajan, Thibault Chervy, Marcus Seidel, David Hagemüller, Stefan Schütz, Johannes Schachenmayer, Cyrilaque Genet, Guido Pupillo, and Thomas W. Ebbesen, “Exploring superconductivity under strong coupling with the vacuum electromagnetic field,” (2019), arXiv:1911.01459 [cond-mat.supr-con].
- [36] Ahana Chakraborty and Francesco Piazza, “Long-range photon fluctuations enhance photon-mediated electron pairing and superconductivity,” *Phys. Rev. Lett.* **127**, 177002 (2021).
- [37] S. C. Burd, R. Srinivas, H. M. Knaack, W. Ge, A. C. Wilson, D. J. Wineland, D. Leibfried, J. J. Bollinger, D. T. C. Allcock, and D. H. Slichter, “Quantum amplification of boson-mediated interactions,” *Nature Physics* **17**, 898–902 (2021).
- [38] M. Dzero, E. A. Yuzbashyan, and B. L. Altshuler, “Cooper pair turbulence in atomic fermi gases,” *EPL (Europhysics Letters)* **85**, 20004 (2009).
- [39] Mark H. Fischer, Titus Neupert, Christian Platt, Andreas P. Schnyder, Werner Hanke, Jun Goryo, Ronny Thomale, and Manfred Sigrist, “Chiral  $d$ -wave superconductivity in srptas,” *Phys. Rev. B* **89**, 020509 (2014).
- [40] Liang Jiang, Takuya Kitagawa, Jason Alicea, A. R. Akhmerov, David Pekker, Gil Refael, J. Ignacio Cirac, Eugene Demler, Mikhail D. Lukin, and Peter Zoller, “Majorana fermions in equilibrium and in driven cold-atom quantum wires,” *Phys. Rev. Lett.* **106**, 220402 (2011).
- [41] Christina V. Kraus, Marcello Dalmonte, Mikhail A. Baranov, Andreas M. Läuchli, and P. Zoller, “Majorana edge states in atomic wires coupled by pair hopping,” *Phys. Rev. Lett.* **111**, 173004 (2013).
- [42] Christina Psaroudaki and Christos Panagopoulos, “Skyrmion qubits: A new class of quantum logic elements based on nanoscale magnetization,” *Phys. Rev. Lett.* **127**, 067201 (2021).
- [43] We discuss this approximation in more detail in Appendix B.
- [44] H. Uys, M. J. Biercuk, A. P. VanDevender, C. Ospelkaus, D. Meiser, R. Ozeri, and J. J. Bollinger, “Decoherence due to elastic rayleigh scattering,” *Phys. Rev. Lett.* **105**, 200401 (2010).
- [45] Allison Carter, private communication (to be published).
- [46] Jan Müller, *Magnetic Skyrmions and Topological Domain Walls*, Ph.D. thesis, Universität zu Köln (2018).