

ALMA Lensing Cluster Survey:
ALMA-Herschel Joint Study of Lensed Dusty Star-Forming Galaxies across $z \simeq 0.5 - 6$

FENGWU SUN ¹, EIICHI EGAMI ¹, SEIJI FUJIMOTO ^{2,3}, TIMOTHY RAWLE ⁴, FRANZ E. BAUER ^{5,6},
KOTARO KOHNO ^{7,8}, IAN SMAIL ⁹, PABLO G. PÉREZ-GONZÁLEZ ¹⁰, YIPING AO ^{11,12}, SCOTT C. CHAPMAN,^{13,14}
FRANCOISE COMBES ^{15,16}, MIROSLAVA DESSAUGES-ZAVADSKY ¹⁷, DANIEL ESPADA ^{18,19},
JORGE GONZÁLEZ-LÓPEZ ^{20,21}, ANTON M. KOEKEMOER ²², VASILY KOKOREV ^{2,3}, MINJU M. LEE ²³,
KANA MOROKUMA-MATSUI ^{7,24,25}, ALEJANDRA M. MUÑOZ ARANCIBIA ^{6,26}, MASAMUNE OGURI ^{27,28,29},
ROSER PELLÓ ³⁰, YOSHIHIRO UEDA ³¹, RYOSUKE UEMATSU ³¹, FRANCESCO VALENTINO ^{2,3},
PAUL VAN DER WERF ³², GREGORY L. WALTH ³³, MICHAEL ZEMCOV ^{34,35} AND ADI ZITRIN ³⁶

¹Steward Observatory, University of Arizona, 933 N. Cherry Avenue, Tucson, 85721, USA

²Cosmic Dawn Center (DAWN), Jagtvej 128, DK2200 Copenhagen N, Denmark

³Niels Bohr Institute, University of Copenhagen, Lyngbyvej 2, DK2100 Copenhagen Ø, Denmark

⁴European Space Agency (ESA), ESA Office, Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

⁵Instituto de Astrofísica, Facultad de Física, Pontificia Universidad Católica de Chile Av. Vicuña Mackenna 4860, 782-0436 Macul, Santiago, Chile

⁶Millennium Institute of Astrophysics, Nuncio Monseñor Sótero Sanz 100, Providencia, Santiago, Chile

⁷Institute of Astronomy, Graduate School of Science, The University of Tokyo, 2-21-1 Osawa, Mitaka, Tokyo 181-0015, Japan

⁸Research Center for the Early Universe, School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

⁹Centre for Extragalactic Astronomy, Department of Physics, Durham University, South Road, Durham, DH1 3LE, UK

¹⁰Centro de Astrobiología, Departamento de Astrofísica, CSIC-INTA, Cra. de Ajalvir km.4 E-28850-Torrejón de Ardoz, Madrid, Spain

¹¹Purple Mountain Observatory & Key Laboratory for Radio Astronomy, Chinese Academy of Sciences, Nanjing, China

¹²School of Astronomy and Space Science, University of Science and Technology of China, Hefei, Anhui, China

¹³Eureka Scientific, Inc. 2452 Delmer Street Suite 100, Oakland, CA 94602-3017, USA

¹⁴Dalhousie University Dept. of Physics and Atmospheric Science Coburg Road Halifax, B3H1A6, Canada

¹⁵Sorbonne Université, Observatoire de Paris, Université PSL, CNRS, LERMA, 75014 Paris, France

¹⁶Collège de France, 11 Place Marcelin Berthelot, 75231 Paris, France

¹⁷Observatoire de Genève, Université de Genève, 51, Ch. des Maillettes, 1290 Versoix, Switzerland

¹⁸SKA Organisation, Lower Withington, Macclesfield, Cheshire SK11 9DL, UK

¹⁹Departamento de Física Teórica y del Cosmos, Campus de Fuentenueva, Universidad de Granada, E18071-Granada, Spain

²⁰Las Campanas Observatory, Carnegie Institution of Washington, Casilla 601, La Serena, Chile

²¹Núcleo de Astronomía de la Facultad de Ingeniería y Ciencias, Universidad Diego Portales, Av. Ejército Libertador 441, Santiago, Chile

²²Space Telescope Science Institute, 3700 San Martin Dr., Baltimore, MD 21218, USA

²³Max-Planck-Institut für Extraterrestrische Physik (MPE), Giessenbachstr. 1, D-85748 Garching, Germany

²⁴Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 3-1-1 Yoshinodai, Chuo-ku, Sagami-hara, Kanagawa 252-5210, Japan

²⁵Chile Observatory, National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka-shi, Tokyo 181-8588, Japan

²⁶Center for Mathematical Modeling, University of Chile, AFB170001, Chile

²⁷Research Center for the Early Universe, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

²⁸Center for Frontier Science, Chiba University, 1-33 Yayoi-cho, Inage-ku, Chiba 263-8522, Japan

²⁹Kavli Institute for the Physics and Mathematics of the Universe (WPI), The University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa-shi, Chiba, 277-8583, Japan

³⁰Aix Marseille Université, CNRS, CNES, LAM (Laboratoire d'Astrophysique de Marseille), UMR 7326, 13388, Marseille, France

³¹Department of Astronomy, Kyoto University, Kyoto 606-8502, Japan

³²Leiden Observatory, Leiden University, P.O. Box 9513, NL-2300 RA Leiden, The Netherlands

³³The Observatories of the Carnegie Institution for Science, 813 Santa Barbara Street, Pasadena, CA 91101, USA

³⁴Rochester Institute of Technology, 1 Lomb Memorial Drive, Rochester, NY 14623, USA

³⁵*Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109, USA*

³⁶*Physics Department, Ben-Gurion University of the Negev, P.O. Box 653, Be'er-Sheva 84105, Israel*

(Received October 29, 2021; Revised April 22, 2022; Accepted May 3, 2022)

Submitted to ApJ

ABSTRACT

We present an ALMA-*Herschel* joint analysis of sources detected by the ALMA Lensing Cluster Survey (ALCS) at 1.15 mm. *Herschel*/PACS and SPIRE data at 100–500 μm are deblended for 180 ALMA sources in 33 lensing cluster fields that are either detected securely (141 sources; in our main sample) or tentatively at $S/N \geq 4$ with cross-matched *HST/Spitzer* counterparts, down to a deblended 1.15 mm flux density of ~ 0.02 mJy. We performed far-infrared spectral energy distribution modeling and derived the physical properties of dusty star formation for 125 sources (109 independently) that are detected at $> 2\sigma$ in at least one *Herschel* band. 27 secure ALCS sources are not detected in any *Herschel* bands, including 17 optical/near-IR-dark sources that likely reside at $z = 4.2 \pm 1.2$. The 16–50–84 percentiles of the redshift distribution are 1.15–2.08–3.59 for ALCS sources in the main sample, suggesting an increasing fraction of $z \simeq 1 - 2$ galaxies among fainter millimeter sources ($f_{1150} \sim 0.1$ mJy). With a median lensing magnification factor of $\mu = 2.6_{-0.8}^{+2.6}$, ALCS sources in the main sample exhibit a median intrinsic star-formation rate of $94_{-54}^{+84} M_{\odot} \text{ yr}^{-1}$, lower than that of conventional submillimeter galaxies at similar redshifts by a factor of ~ 3 . Our study suggests weak or no redshift evolution of dust temperature with $L_{\text{IR}} < 10^{12} L_{\odot}$ galaxies within our sample at $z \simeq 0 - 2$. At $L_{\text{IR}} > 10^{12} L_{\odot}$, the dust temperatures show no evolution across $z \simeq 1 - 4$ while being lower than those in the local Universe. For the highest-redshift source in our sample ($z = 6.07$), we can rule out an extreme dust temperature (> 80 K) that was reported for MACS0416 Y1 at $z = 8.31$.

Keywords: submillimeter: galaxies – galaxies: evolution – galaxies: high-redshift

1. INTRODUCTION

In massive star-forming galaxies at cosmological distances, a large fraction of star formation is found to be obscured by dust (e.g., Ivison et al. 1998; Heinis et al. 2014; Whitaker et al. 2017; Fudamoto et al. 2020). Observations at far-infrared (far-IR) wavelengths directly sample the thermal continuum emission from dust grains in the interstellar medium (ISM), a reliable tracer of recent star formation activity (e.g., Kennicutt & Evans 2012). With the high sensitivity and spatial resolution of Atacama Large Millimeter/submillimeter Array (ALMA), submillimeter galaxies (SMGs; or dusty star-forming galaxies, DSFGs as they are often called) have been studied up to a redshift of 6.9 (Strandet et al. 2017; Marrone et al. 2018), and dust continuum emission from Lyman-break galaxies (LBGs) have also been revealed up to a redshift of 8.3 (e.g., Tamura et al. 2019; Bakx et al. 2020). ALMA studies of dust obscured star formation, combined with observations obtained at rest-frame UV/optical wavelengths (e.g., with *Hubble Space Telescope*, *HST*), provide a comprehensive picture on galaxy formation and evolution across the past 13 Gyrs (e.g., Bouwens et al. 2020; see a recent review by Hodge & da Cunha 2020).

In order to discover and study the physical properties of SMGs that are intrinsically faint (0.01–1 mJy around 1 mm wavelength), the effect of gravitational lensing has been widely utilized, which allowed the first detection of SMGs (Smail et al. 1997). The ALMA Lensing Cluster Survey (ALCS) is an ALMA Cycle-6 large programs (PI: Kohno; Kohno 2019; Kohno et al. in preparation) dedicated to survey intrinsically faint continuum sources and line emitters with the assistance of gravitational lensing. By surveying a total image-plane sky area of ~ 134 arcmin² (primary beam response greater than 0.3) down to a depth of 0.07 mJy beam⁻¹ (1σ), ALCS aimed to detect > 100 continuum sources at $\geq 5\sigma$ significance at 1.15 mm. ALCS has an effective survey area of ~ 10 arcmin² for sources brighter than 0.1 mJy at 1.15 mm ($> 5\sigma$; lensing corrected). This is the largest survey obtained with ALMA Band 6 at comparable depth so far. The detected continuum sources can then be used to examine the origin of cosmic infrared background (CIB), measure the [C II] luminosity function in the Epoch of Reionization (EoR; Fujimoto et al. 2021), and constrain the evolution of gas and dust content of galaxies around the peak of cosmic star-formation history (CSFH).

All the 33 cluster fields were selected from the best studied clusters primarily from the Abell (Abell 1958; Abell et al. 1989) and MACS (Ebeling et al. 2001) catalogs that have been observed with *HST* Treasury Programs including Cluster Lensing And Supernova survey with *Hubble* (CLASH, 12 clusters; PI: Postman; Postman et al. 2012), *Hubble* Frontier Fields (HFF, 5 clusters; PI: Lotz; Lotz et al. 2017) and Reionization Lensing Cluster Survey (RELICS, 16 clusters; PI: Coe; Coe et al. 2019). These survey programs utilized both the Advanced Camera for Surveys (ACS) and Wide Field Camera 3 (WFC3) to obtain deep and high-resolution images of massive galaxy clusters at $z = 0.2 - 0.9$ from the optical to near-infrared (near-IR). Combined with *Spitzer*/IRAC coverage of at least at medium depth (the median 5σ depth at $4.5\ \mu\text{m}$ is 23.3 ± 0.1 AB mag; Sun et al. 2021a), these *HST* data provide direct constraints on unobscured or mildly obscured stellar components in both the environment of the massive galaxy cluster as well as in gravitationally lensed galaxies in the distant Universe.

The ALCS fields were also observed by the *Herschel* Space Observatory (Pilbratt et al. 2010) in the wavelength range from 100 to 500 μm . Launched in 2009 and retired in 2013, *Herschel* was designed to study the dust-obscured universe at submillimeter wavelengths. *Herschel* data are critical for the interpretation of the ALCS fields due to its unique wavelength coverage of thermal dust continuum of high-redshift ($z \gtrsim 1$) galaxies. *Herschel* and ALMA observations both detect thermal emissions from dust heated by the UV radiation from young stars. With a good sampling of far-IR spectral energy distributions (SEDs) with 4–6 bands in total, ALMA and *Herschel* data provide critical constraints on photometric redshifts, thermal dust temperature and dust mass of ALCS-selected galaxies.

Herschel data of the ALCS clusters were obtained through various programs. Among them, the *Herschel* Lensing Survey (HLS; Egami et al. 2010; Sun et al. 2021b) is the largest program imaging the fields of massive galaxy clusters to study cluster-lensed high-redshift galaxies. Deep *Herschel* observations in blank fields are often subject to confusion noise (Nguyen et al. 2010). This restricts the detection of typical ultra luminous infrared galaxies (ULIRGs; $L_{\text{IR}} \geq 10^{12} L_{\odot}$) beyond $z \sim 2$ (c.f. Rawle et al. 2016). However, with the lensing magnification by massive clusters, which typically do not contain far-IR-bright galaxies in the cluster core (e.g., Rawle et al. 2012), we are able to break the confusion limit and discover intrinsically faint sources (Smail et al. 1997). This has been demonstrated by the *Herschel* detection of the $z = 2.8$ LIRG ($L_{\text{IR}} = 10^{11} - 10^{12} L_{\odot}$)

behind the Bullet cluster with a lensing magnification of $\mu \sim 75$ (Rex et al. 2010). With cluster lensing, Sklias et al. (2014) and Dessauges-Zavadsky et al. (2015) explored the star formation history, dust extinction and molecular gas content of LIRGs at $z \simeq 1.5 - 3$, and recent ALMA observations of cluster-lensed *Herschel* sources revealed the existence of low-surface-brightness SMGs with extended dust continua (Sun et al. 2021b).

In this work, we present the ALMA-*Herschel* joint analysis of the dusty star-forming galaxies detected by ALCS. Similar work has been presented in Rawle et al. (2016, hereafter R16) for the six HFF clusters, and here we expand the sample to 28 more cluster fields and use high-resolution ($\sim 1''$) ALMA continuum maps as priors for source extraction, in contrast to the mid-IR priors (*Spitzer* and *WISE*) used in R16. Because of the well known negative K -correction, the selection function of SMGs at millimeter wavelengths is nearly constant in terms of cold dust mass across $z \simeq 1 - 6$. Therefore, compared with R16, the use of ALMA priors allows a more extensive and accurate measurements of *Herschel* flux densities of sources at higher redshifts ($z > 2$), constraining the redshift distribution, dust temperatures and star-formation rates (SFRs) of millimeter sources towards the faint end ($f_{1150} \sim 0.02$ mJy).

This paper is arranged as follows: In Section 2, we introduce the sample discovered by the ALCS (will be described in greater detail by Fujimoto et al. in preparation) and the obtained *Herschel* data together with the data reduction techniques. Section 3 presents the procedure for source extraction using the *Herschel* data. Section 4 presents the far-IR SED fitting and photometric redshift estimate. Section 5 presents the analysis of lensing magnification and uncertainty. In Section 6, we discuss the statistical results of galaxy properties and their implications. The summary can be found in Section 7. Throughout this work, we assume a flat Λ CDM cosmology with $h = 0.7$ and $\Omega_m = 0.3$. We define the IR luminosity (L_{IR}) as the integrated luminosity over a rest-frame wavelength range from 8 to 1000 μm .

2. OBSERVATIONS AND DATA REDUCTION

2.1. ALMA Data and the Sample

All of the sources in this work are selected with the ALCS, which will be detailed in Fujimoto et al. (in preparation). ALMA Band-6 observations for the 33 clusters were conducted through Program 2018.1.00035.L (ALCS; PI: Kohno), and we also combined archival data from Programs 2013.1.00999.S and 2015.1.01425.S (ALMA Frontier Fields; PI: Bauer; González-López et al. 2017). The list of the ALCS clusters with their coordinates, short names (e.g., M0553

for MACSJ0553.4-3342) and *HST* program names is presented in Table 1. The observations were obtained at a central wavelength of 1.15 mm with a 15 GHz total bandwidth (i.e., two tunings of dual polarization; 250.0–257.5 GHz and 265.0–272.5 GHz). The use of two tunings instead of one allows us to search for line-emitting galaxies over a larger volume, which is another important science goal of ALCS. This, for example, led to the serendipitous discovery of a [C II] emitter at $z = 6.072$ (Fujimoto et al. 2021). All the ALMA data were reduced with CASA (McMullin et al. 2007) with different pipelines versions for observations obtained in different cycles (e.g., v5.4.0 for 26 clusters observed in Cycle 6 and v5.6.1 for the remaining clusters in Cycle 7). Natural-weighting continuum imaging was performed at both the native (full width at half maximum, FWHM $\sim 1''$) and *uv*-tapered ($\sim 2''$) resolutions with the CASA TCLEAN algorithm.

Through a peak pixel identification routine of SEXTRACTOR (Bertin & Arnouts 1996) with the ALMA maps at both the native and $2''$ -tapered resolutions (before primary beam correction), we securely detected 141 sources that are either at (i) signal-to-noise ratio $S/N_{\text{nat}} \geq 5$ in the native-resolution maps, or (ii) $S/N_{\text{tap}} \geq 4.5$ in the $2''$ -tapered maps, over an area of $\sim 134 \text{ arcmin}^2$ with primary beam response greater than 0.3 (Fujimoto et al. in preparation). Based on the number count of negative peaks, the number of spurious source above these S/N cuts is expected to be around one. We further refer to these 141 secure ALCS sources as the *Main Sample*.

258 sources were tentatively detected at (i) $S/N_{\text{nat}} = 4-5$ in the native-resolution maps and (ii) $S/N_{\text{tap}} < 4.5$ in the $2''$ -tapered maps, down to a minimal flux density of $\sim 0.2 \text{ mJy}$ at 1.15 mm. Based on *HST* and *Spitzer*/IRAC images, we identified 39 of these sources with near/mid-IR counterparts within a separation of $1''$. Given the high source densities in cluster fields ($\sim 0.06 \text{ arcsec}^{-2}$ in the *HST*/F160W band; Sun et al. 2021a), we expect 7 ± 3 pairs of random associations among these tentative ALCS sources and cross-matched near/mid-IR counterparts. Assuming that the majority of these *HST*/*Spitzer*-matched sources are real, we note that the number found is consistent with the number difference between positive and negative peaks in the ALMA maps in this S/N range (37 in total). We further refer to these 39 tentative ALCS sources as the *Secondary Sample*, but warn that $18 \pm 8\%$ of these sources are likely to be false matched.

We note that the continuum S/N of ALCS sources could be boosted by serendipitous emission line detections, for example, CO(5-4) line at $z = 1.11 - 1.17$ and

1.24 – 1.31 (e.g., M0553-ID133/190/249 at $z = 1.142$, Ebeling et al. 2017; Sun et al. 2021b). All the line emitters will be reported by a future paper of the collaboration. However, because of a large bandwidth (15 GHz), the CO line contamination to continuum S/N and flux density is limited to $\lesssim 1-10\%$. In addition, ALCS can only sample faint high- J CO lines (upper J number at ≥ 7) for sources at $z \gtrsim 2$. According to the CO spectral line energy distribution (SLED) of high-redshift SMGs reported in the literature (e.g., Greve et al. 2014; Béthermin et al. 2016; Yang et al. 2017; Birkin et al. 2021), even the high- J CO SLED is as flat as those reported for local active-galactic-nucleus-host (AGN-host) galaxies (e.g., Rosenberg et al. 2015), the CO contamination will be $\sim 1 - 10\%$ at most. Only one [C II] emitter was found among all continuum sources (Fujimoto et al. 2021; Laporte et al. 2021), and the continuum flux density of this source was measured on line-subtracted continuum image.

2.2. *Herschel*/PACS

The Photodetector Array Camera and Spectrometer (PACS; Poglitsch et al. 2010) on *Herschel* enabled simultaneous observations at $160 \mu\text{m}$ (red channel) with the long-wavelength camera and either at 70 or $100 \mu\text{m}$ (blue/green channel) with the short-wavelength camera. Eighteen out of the total 33 clusters were imaged with PACS at both 100 and $160 \mu\text{m}$, and two clusters were also observed with PACS at $70 \mu\text{m}$ (M1149 and AS1063). The analysis of PACS $70 \mu\text{m}$ data in these cluster fields has been presented by R16, and the only two matched sources are AS1063-ID17 ($z = 1.44$, $f_{70} = 7.3 \pm 0.9 \text{ mJy}$) and AS1063-ID147 ($z = 0.610$, $f_{70} = 28.8 \pm 2.3 \text{ mJy}$; analyzed in detail by Walth et al. 2019).

16 out of the 18 clusters were observed by PACS as part of the HLS (Egami et al. 2010; Sun et al. 2021b), which combines an Open-Time Key Program (KPOT; program ID: KPOT_eegami_1; nine clusters) and an Open-Time Cycle 2 (OT2; program ID: OT2_eegami_5; seven clusters) Program (both PI: Egami). The remaining two clusters, namely Abell370 and RXJ1347-1145, were observed as part of the PACS Evolutionary Probe (PEP; program ID: KPGT_dlutz_1, PI: Lutz; Lutz et al. 2011). All of the PACS 100 and $160 \mu\text{m}$ observations consist of two orthogonal scan maps, each comprising 18–22 repetitions of 13 parallel 4-arcmin scan legs. The summary of the PACS observations, including the observation IDs and total scan time for each cluster, is presented in Table 1.

We followed the same data reduction procedure as detailed in R16 for the HFF clusters. The PACS images were generated with UNIMAP (Piazzo et al. 2015) with a




Figure 1. Median RMS noise of SPIRE pixels (RMS_{map} ; measured as the noise of sky background after sources being filtered out) at $250\ \mu\text{m}$ within the footprints of 33 ALCS clusters. 18 clusters were observed with *Herschel* in the “deep” mode (in blue), and they have deep SPIRE images ($\text{RMS}_{\text{map}} < 9\ \text{mJy beam}^{-1}$) as well as PACS ones. The remaining 15 clusters were observed in the “snapshot” mode (in orange), and they have shallower SPIRE data ($\text{RMS}_{\text{map}} > 9\ \text{mJy beam}^{-1}$; see Section 2.3) with no PACS coverage. The number of main-sample ALCS sources in each cluster field is noted at the bottom of each bar.

pixel scale of $1''.0$ at $100\ \mu\text{m}$ and $2''.0$ at $160\ \mu\text{m}$. The final PACS image products have a typical field of view (FoV) with a radius of $\sim 4'$, covering the full ALMA footprints of the 18 clusters. The typical beam sizes are $7''.4$ and $11''.4$ at 100 and $160\ \mu\text{m}$, and the depths of the PACS data at the cluster center are presented in Table 2.

2.3. *Herschel*/SPIRE

The Spectral and Photometric Imaging Receiver (SPIRE; Griffin et al. 2010) on *Herschel* worked simultaneously at 250 , 350 and $500\ \mu\text{m}$. All of the 33 clusters were imaged with SPIRE in two observing modes with different depths. The 18 clusters also observed with PACS were scanned with SPIRE in the “deep” mode down to confusion-limited depths ($\text{RMS}_{\text{map}} \sim 6\ \text{mJy beam}^{-1}$ at $250\ \mu\text{m}$; measured as the noise of sky background after sources being filtered out), and the remaining 15 clusters were observed in the “snapshot” mode with a shorter scan duration and thus at shallower depths ($\text{RMS}_{\text{map}} \sim 11\ \text{mJy beam}^{-1}$; as visualized in Figure 1).

Among the total 18 clusters in the “deep” mode, 16 of them were observed as part of the HLS. Observations of the nine clusters through KPOT_eegami_1 consisted of 20 repetitions in the large scan map mode, each with two $4'$ scans and cross-scans (total scan time as $t_{\text{scan}} \sim 1.7\ \text{h}$ per cluster). The other seven clusters observed through Open-Time Cycle 1/2 Programs (OT1_eegami_4 and OT2_eegami_5; both PI: Egami) were imaged through 11-repetition small scan maps (one in OT1 and ten in OT2), and each repetition consisted of one scan and

one cross-scan of $4'$ length ($t_{\text{scan}} \sim 0.4\ \text{h}$ per cluster). The remaining two clusters, A370 and R1347, were observed as part of the *Herschel* Multi-tiered Extragalactic Survey (HerMES; program ID: KPGT_soliver_1; PI: Oliver; Oliver et al. 2012). Both of these clusters were observed with eight small scan maps (six repetitions per each covering the cluster core) and three large scan maps (one repetition per each with $38'$ length covering a wider area), and the total scan time is $3.5\ \text{h}$ per cluster. The final SPIRE map products of these two clusters have a wider FoV, but the central depths are comparable to those of the HLS data.

All of the 15 clusters in the “snapshot” mode were observed as part of the HLS through single-repetition small scan maps, and each repetition consisted of one scan and one cross-scan of a $4'$ length ($t_{\text{scan}} \sim 3\ \text{min}$ per cluster). Fourteen of them were observed by the OT1 program OT1_eegami_4, and the remaining one was observed by the OT2 program OT2_eegami_6 (PI: Egami).

Table 1 summarizes the IDs and total scan times of all the SPIRE observations. All of the SPIRE data were processed by the standard reduction pipeline in HIPE v12.2 (Ott 2010) which is also detailed in R16. The output pixel sizes of the final image products are $6''$, $9''$ and $12''$ at 250 , 350 and $500\ \mu\text{m}$. The typical radii of the SPIRE FoVs are $7'$ for the 15 clusters observed in the “snapshot” mode, $8'$ for the seven clusters observed in OT1/2, $11'$ for the nine clusters observed in KPOT, and $33'$ for A370 and R1347. The full survey area of the ALCS was covered by these SPIRE images. The typical

beam sizes are $18''$, $24''$ and $35''$ in these three bands, and the depths of the SPIRE data at the cluster center are presented in Table 2.

2.4. Ancillary *HST* and *Spitzer* Data

For the purpose of enhancing the astrometric accuracy of *Herschel* data, we include the *Spitzer*/IRAC data of these 33 cluster fields in our analysis obtained from the NASA/IPAC Infrared Science Archive (IRSA¹). We also include the *HST* data of all the cluster fields but only for comparing the positions of dust continuum sources with the stellar components. We defer the study of optical/near-IR counterparts and panchromatic SED modeling of ALCS sources to another paper from the collaboration.

2.5. Redshift Catalogs

To supply accurate redshifts for far-IR SED modeling (Section 4), we cross-matched the ALCS source sample with the spectroscopic redshift (z_{spec}) catalogs made available by the CLASH-VLT spectroscopic survey (Biviano et al. 2013), Grism Lens-Amplified Survey from Space (GLASS; Schmidt et al. 2014; Treu et al. 2015; Wang et al. 2015) and recent VLT/MUSE surveys of massive cluster fields by Caminha et al. (2019) and Richard et al. (2021). A maximum separation of $1''.5$ is allowed for cross-matching, which is comparable to the FWHM of IRAC PSF. We also include redshifts for a few sources reported by various studies in the literature (e.g., M0553 triply lensed system at $z = 1.14$; Ebeling et al. 2017) or private communication (e.g., M0417-ID46/58/121, an *HST* *H*-faint triply lensed system at $z = 3.65$; Kohno et al. in preparation). In addition, we also include two ALMA-HFF sources reported by Laporte et al. (2017) with their z_{spec} 's derived from the GLASS detection of the 4000 \AA break, and a triply lensed ALCS source system that belongs to a MUSE-confirmed galaxy group at $z = 4.32$ (Caputi et al. 2021). Spectroscopic redshifts are available for 60 ALCS sources in both the main and secondary samples.

We also utilized the *HST* photometric redshift (z_{phot}) catalogs of optical/near-IR sources tabulated by CLASH (Molino et al. 2017), HFF (Shipley et al. 2018) and RELICS (Coe et al. 2019) groups. Sources are cross-matched by their coordinates and a maximum separation of $1''.5$ is allowed. Fujimoto et al. (2016) reported a median offset of $0''.25$ between the *HST* and ALMA centroids of ALMA sources, and such an observed offset could be larger in cluster fields because of the lensing magnification. We also apply visual inspections of the

HST F814W, F105W and F160W images to remove any conspicuous mismatch. We identified catalogued *HST* z_{phot} measurements for 125 ALCS sources in both the main and secondary samples, including 49 sources with additional spectroscopic redshifts.

3. HERSCHEL SOURCE EXTRACTION

3.1. Preparation

3.1.1. Image Alignment

Following R16, we first aligned all the *Herschel* images to the IRAC Channel 1 ($3.6 \mu\text{m}$) images before the actual source extraction at $100\text{--}500 \mu\text{m}$. We cross-matched the IRAC $3.6 \mu\text{m}$ source catalog in each field with the $\sim 10\text{--}20$ brightest sources detected in *Herschel* bands using DAOFIND (Stetson 1987). We then computed the median RA and Dec offsets of the matched sources in the *Herschel* and IRAC bands, and corrected these for the *Herschel* data. We only calculated the offsets independently for the PACS $100 \mu\text{m}$ and SPIRE $250 \mu\text{m}$ data and applied the same astrometric shift to other bands of the same instrument. This is because the offsets between different bands of the same *Herschel* instrument have been well calibrated. The median corrected offsets are $0''.9$ and $1''.3$ for PACS and SPIRE images, consistent with those reported in R16.

3.1.2. Input ALCS Source Catalog

We constructed the input catalog for *Herschel* source extraction using 180 ALCS sources at $S/N \geq 4$ (Section 2.1). Among them, 141 secure sources (main sample) were extracted in the first two iterations, and 39 tentative sources with matched *HST*/*Spitzer* counterparts (secondary sample) were then extracted on the residual images. We note that 85% of tentative ALCS sources at $S/N_{\text{nat}} = 4\text{--}5$ and $S/N_{\text{tap}} < 4.5$ do not show any near/mid-IR counterpart. These sources are expected to mostly be spurious and not included for *Herschel* photometry. However, it is possible that a few of them represent highly obscured high-redshift galaxies ($A_V \gtrsim 5$, $z \gtrsim 4$), which are missed by this study. The coordinates of ALMA sources were used as positional priors, and the S/N 's of ALMA detections were later used to rank the priority of extraction in Section 3.2.

3.1.3. Background Subtraction

We estimated and subtracted the 2D sky background of *Herschel* images using sigma-clipped statistics in each mesh of a grid that covers the whole input data frame. This is the same algorithm used by SEXTRACTOR (Bertin & Arnouts 1996). The typical box size of the mesh was $21 \times 21 / 16 \times 16$ pixels for PACS/SPIRE bands, corresponding to an area of $10/20$ beams. The

¹ <https://irsa.ipac.caltech.edu/>




Figure 2. *Herschel* images of Abell2744 (A2744) observed in the “deep” mode. PACS 100/160 μm images are shown on the left and SPIRE 250/350/500 μm images are on the right. Each panel consists of the scientific image (left) and residual image after PSF source extraction (right). In each panel, the red patch represents the ALCS footprint (primary beam response cut at 0.25), and the magenta circles (blue squares) represent the secure ALCS sources (main sample) extracted at $S/N > 2$ ($S/N < 2$) in each *Herschel* band. The cyan diamonds are tentative ALCS sources (secondary sample) extracted at $S/N > 2$ in the *Herschel* bands. The green triangles are ALMA-faint *Herschel* sources (see Section 3.2), which we also included for source extraction. One-arcmin scale bar is shown at the lower-right corner of each panel.

size of median filter, which was applied to suppress possible overestimate of background due to bright sources, was $7 \times 7 / 5 \times 5$ pixels for PACS/SPIRE bands (corresponding to an area of $1/2$ beams). In the cluster fields with strong extended emissions seen in the SPIRE bands (R0032, M2129, P171 and A2163), we slightly reduced the sizes of mesh and filter for a better removal of foreground large-scale emissions. The 2D RMS map was then created as a by-product of this background subtraction process.

3.1.4. Neighborhood Examination

Our *Herschel* source extraction process started from the bluest band for each cluster, i.e., 100 μm for the 18 cluster observed with PACS in the “deep” mode, and

250 μm for the other 15 clusters observed in the “snapshot” mode. Therefore, in the bluest band, we first extracted a source catalog using the DAOFIND algorithm, which contains sources detected or undetected by the ALCS. We then matched and removed the ALCS sources in this catalog within a maximum separation of $1/3$ the point-spread function (PSF) FWHM (i.e., $2''$ at 100 μm and $6''$ at 250 μm). Therefore, this catalog represents the 1.15 mm-undetected *Herschel* sources that may blend with ALMA-detected sources in *Herschel* data at longer wavelengths (e.g., 500 μm). Compared with ALCS sources, these sources are likely at a lower redshift, and thus their SEDs peak at shorter wavelength and drop rapidly at longer wavelength. We only included ALMA-faint *Herschel* sources within a separa-

tion of $30''$ ($100\ \mu\text{m}$) or $36''$ ($250\ \mu\text{m}$) from the ALCS sources. This is because a larger searching area would not further increase the quality of source deblending for the ALCS sources.

To enhance the accuracy of flux extraction, we also manually added or adjusted the positions of several ALMA-faint *Herschel* sources according to the coordinates of their IRAC counterparts. This affects 5% of all ALMA-faint *Herschel* sources. For *Herschel* bands at longer wavelengths, we directly used this list of ALMA-faint *Herschel* sources in the bluest band. However, we find that if an ALMA-faint *Herschel* source cannot be extracted at $>10\ \text{mJy}$ for “deep”-mode clusters or $>18\ \text{mJy}$ for “snapshot”-mode clusters in a given SPIRE band, this source will be very unlikely to be detected above 2σ at any redder band. Therefore, such sources will be removed from fitting at longer wavelengths.

3.2. Iterative *Herschel* Photometry

Herschel source extraction was performed with an iterative PSF photometry approach in an increasing order of wavelength using PHOTUTILS (Bradley et al. 2019). We adopted the PSF models of PACS and SPIRE from the *Herschel* Science Archive (HSA)². We also applied the spacecraft orientation angle to calculate the realistic PSF for the data taken in each cluster field.

3.2.1. Initial Guess of Flux Densities

To provide initial guesses of flux densities for PSF photometry, we performed circular aperture photometry for all the ALCS sources and ALMA-faint *Herschel* sources. The radii of the apertures were $5''$, $8''$, $12''$, $15''$, $18''$ from 100 to $500\ \mu\text{m}$, and the aperture correction factors (1.92, 1.90, 1.66, 1.79, 2.24) were computed based on the PSF models. In the PACS bands, these initial guesses of flux densities were used for all the sources in the main sample and ALMA-faint *Herschel* sources. In the SPIRE bands, these initial guesses were only used for all the $S/N_{\text{ALMA}} \equiv \max(S/N_{\text{nat}}, S/N_{\text{tap}}) \geq 10$ sources and ALMA-faint *Herschel* sources due to a stronger source blending issue.

3.2.2. Iterations of PSF Photometry

The PSF photometry was performed in three rounds of iterations in an decreasing order of the significance of detection. Because most of the ALCS sources are compact in spatial extent ($\text{FWHM} \lesssim 1''$), we assumed point-like profile for all of the sources to be extracted. In each iteration, we only kept the results of those sources with




Figure 3. *Herschel*/SPIRE images of MACSJ0417.5-1154 (M0417) observed in the “snapshot” mode. The layout and symbols are the same as those in Figure 2.

positive extracted flux densities. The uncertainties of extracted flux densities were estimated from the covariance matrix of least squares fitting.

In the first iteration, we tentatively extract sources that were (i) ALCS sources at $S/N_{\text{ALMA}} \geq 10$, or (ii) ALMA-faint *Herschel* sources described above. These two types of objects should be the brightest sources seen in a given *Herschel* map. Therefore, an accurate flux density modeling of these sources will provide helpful guesses for the final combined source models in the whole data frame. We applied the rotated PSFs and modeled the flux densities at given source positions. The DAOGROUP algorithm (Stetson 1987) was adopted to group sources within a separation of one beam FWHM. The best-fit model was then stored and used as prior information for the next iteration.

In the second iteration, we extracted sources that were (i) ALCS sources in the main sample, or (ii) ALMA-faint *Herschel* sources. These two types of objects should

² <https://www.cosmos.esa.int/web/herschel/ancillary-data-products>

be secure sources and thus their flux densities should be positive in any *Herschel* band. The flux priors were given by the first iteration or aperture photometry if the sources were not modeled previously. In SPIRE bands, the initial flux guesses of sources at $S/N_{\text{ALMA}} < 10$ were assumed as RMS_{map} . With a similar PSF photometry routine, we modeled and subtracted sources with positive best-fit flux densities. In this intermediate step, most of the real sources in the current *Herschel* band have been extracted. The best-fit model and residual maps were recorded for the next iteration.

In the last iteration, we extracted tentative ALCS sources in the secondary sample. Source extraction was performed on the residual map of the second iteration, and the initial guesses of the flux densities were set to zero. Only sources modeled with positive flux densities were kept. The best-fit model and residual was then recorded as part of the final products.

Figure 2 and 3 display the *Herschel* scientific and residual images (i.e., before and after the source extraction) of two cluster fields observed in both the “deep” and “snapshot” modes, namely A2744 (Figure 2; “deep”) and M0417 (Figure 3; “snapshot”). Secure, tentative ALCS sources and ALMA-faint *Herschel* sources extracted at $S/N > 2$ are shown as open magenta, cyan and green circles, respectively. No significant residual in SPIRE bands can be found within the ALCS footprint shown as the region enclosed by red solid line, although weak ring-shape residuals can be identified in PACS bands for a few very bright sources ($f \gtrsim 50$ mJy). This could be caused by the invalidity of point-source assumption or mismatch of PSF models, but our examination in Appendix A suggests no gain or loss of PACS flux densities through this PSF photometry routine.

3.3. Non-detections

As a consequence of the limited depths of the *Herschel* data when compared with the deep ALCS data, a significant number of sources were not successfully extracted in the *Herschel* bands (see statistics in Section 3.5). Therefore, we only provide 3σ upper limits of their flux densities. As pointed out by R16 and other works, because the source positions are known from the ALMA data, the actual *Herschel* flux limit of non-detections are lower than the nominal confusion noise limit (e.g., Nguyen et al. 2010).

Based on the flux densities and their uncertainties modeled with PSF photometry, we calculated the median flux density uncertainty of the extracted secure sources in each band and each cluster field (main ALCS sample, *Herschel* $S/N > 2$). These uncertainties were obtained based on positional priors using the covariance




Figure 4. RMS noise of *Herschel* images measured from the prior-based catalogs ($\text{RMS}_{\text{prior}}$; see Section 3.3) versus the 2D uncertainty maps (RMS_{map}). Measurements obtained on the maps from 100 to 500 μm are shown as symbols with different colors as labeled in the upper-left legend. The dashed line indicates the case in which the two RMS noises are identical, and the solid line represents the median ratio of $\text{RMS}_{\text{prior}}$ to RMS_{map} as 0.43 ± 0.01 .

matrix of least squares PSF modeling, and thus we define them as $\text{RMS}_{\text{prior}}$. Table 2 presents all the measured $\text{RMS}_{\text{prior}}$ along with the RMS_{map} which is directly measured from the 2D uncertainty map within the ALCS footprint. We also compare the $\text{RMS}_{\text{prior}}$ with RMS_{map} measured for all our *Herschel* data in Figure 4, and we find a median ratio of $\text{RMS}_{\text{prior}}$ to RMS_{map} as 0.43 ± 0.01 in the *Herschel*/SPIRE bands. Similar value can also be found for the PACS bands. This means that with prior knowledge of source positions, the actual 3σ limit of non-detection is around 1.3 times of the local background RMS_{map} . Such an upper limit is adopted for all the *Herschel* non-detections in this work.

The median 3σ depths derived for the 18 clusters observed in the “deep” mode are 7.5, 7.6 and 8.2 mJy at 250, 350 and 500 μm . These are consistent with the 3σ RMS of deblended flux densities using the cross identification procedure reported by Roseboom et al. (2010, for HerMES fields), and only slightly lower than the reported value in Swinbank et al. (2014) at 500 μm , which also included ALMA positional priors for deblending.

3.4. Special Sources

Several ALCS sources show secure close companions (i.e., angular separation less than $6''$, which is one third of the beam FWHM at 250 μm) at $S/N \geq 5$ in the ALMA maps. Five of these seven systems have already been confirmed as lensed arcs or multiply lensed systems




Figure 5. The seven special cases of sources with close companions that we redistributed the *Herschel* flux densities based on the ALMA flux density ratios (see Section 3.4). In each postage stamp image, we show the *HST* WFC3-IR/F160W image, *Spitzer* IRAC 3.6 μm image, ALMA Band-6 image at native resolution and SPIRE 250 μm image (500 μm for ACT0102-ID215/224) from the left to right. ALCS sources are labeled with open blue circles, and cluster names, ALMA source IDs and scale bars are shown in the corners of the F814W images.

with *HST* or ALMA data. Due to the coarse resolution of the *Herschel* data, especially those of SPIRE, the flux density ratios among these source groups might be incorrectly modeled in Section 3.2.

In this step, we re-distributed the *Herschel* flux densities of these source groups according to their ALMA flux density ratios. If a source was resolved on the native-resolution ALMA map with a major-axis FWHM less than $3''$ (morphological parameters modeled with CASA IMFIT; Fujimoto et al. in preparation), we adopted the ALMA flux density measured with a circular aperture of $r = 2''$. For sources with larger FWHMs, we adopted the best-fit ALMA flux densities derived from surface brightness profile modeling (assuming 2D Gaussian profile with IMFIT). For unresolved sources, we used the peak flux densities per beam measured on the $2''$ -tapered maps. We note that the redistribution of *Herschel* flux densities assumes a fixed far-IR SED shape among blended sources in each group. Only one source from each blended group is considered for the discussions of dust temperature in Section 6.4.

We redistributed the *Herschel* fluxes for all the ALCS sources in the main sample and within a separation of $6''$. This includes: (i) ACT0102-ID215/224 (lensed galaxy pair known as “la Flaca”; Lindner et al. 2015; Wu et al. 2018; Caputi et al. 2021), (ii) ACT0102-

ID223/294 (lensed galaxy pair; Wu et al. 2018), (iii) M0417-ID221/223 (two ALMA sources at $S/N \gtrsim 6$ with a separation of $2.3''$; Kohno et al. in preparation), (iv) M0553-ID133/190/249 (triply lensed arc; Ebeling et al. 2017; Sun et al. 2021b), (v) M1206-ID55/60/61 (known as the “Cosmic Snake”; Ebeling et al. 2009; Cava et al. 2018; Dessauges-Zavadsky et al. 2019), (vi) R0032-ID53/55/57/58 (lensed arc; Dessauges-Zavadsky et al. 2017), (vii) R1347-ID145/148 (IR-bright lensed arc in R1347). In Figure 5 we display the *HST*, ALMA and SPIRE images of all these source groups. Note that we only redistributed the 500 μm flux densities of ACT0102-ID215/224 because of a moderate separation ($9''$).

3.5. Statistics of *Herschel* Detections

By comparing our PSF photometric results with those derived with aperture photometry, PSF photometry using mid-IR priors (R16) and different software (XID+; Hurley et al. 2017) as presented in Appendix A, we confirmed the quality of our *Herschel* flux density measurements. In Table C1 we present the *Herschel* photometric catalog of 180 ALCS sources in both the main and secondary sample (Section 2.1). The definition of these samples is also illustrated in Figure 6.

Figure 7 shows the *Herschel* detection rates as functions of ALMA S/N cut from 100 to 500 μm . The rates of 100 μm and 250 μm detection ($> 2\sigma$) are almost constant




Figure 6. Summary of ALMA and *Herschel* detections and sample definition. The *Main Sample* includes all 141 secure ALCS sources detected at either $S/N_{\text{nat}} \geq 5$ in the native-resolution maps or $S/N_{\text{tap}} \geq 4.5$ in the $2''$ -tapered maps. The *Secondary Sample* includes 39 tentative ALCS sources ($S/N_{\text{nat}} = 4 - 5$ and $S/N_{\text{tap}} < 4.5$) showing *HST/Spitzer* counterparts within $1''$ offset. Table C1 presents the *Herschel* photometric catalog of these 180 sources (shown as blue circles). Among 141 sources in the main sample, 105 of them are star-forming galaxies which are detected at $S/N > 2$ in at least one *Herschel* band. Together with 20 star-forming galaxies in the secondary sample above the same *Herschel* detection threshold, these galaxies are presented in Table C2 for their physical properties (shown as red filled regions). 43 sources that are undetected in all *Herschel* bands (including 27 “*Herschel*-faint” galaxies in the main sample; see Section 3.5 and Section 4.3), 11 BCGs and one mismatched source (A2537-ID06) are not included in Table C2 (shown as shallow filled regions).




Figure 7. Detection ($> 2\sigma$) rates of ALCS sources as functions of ALMA S/N cut in all five *Herschel* bands. Line color and style of each band are shown in the lower-right legend, and values in parentheses indicate numbers of $> 2\sigma$ *Herschel* detections among secure ALCS sources ($S/N_{\text{ALMA}} \geq 5$).

at $S/N_{\text{ALMA}} \geq 5$. However, the detection rates at long

wavelength (350 and 500 μm) are clearly correlated with the significance of ALMA sources. This is likely the consequence of (i) a larger beam size and stronger blending effect towards longer wavelength, and (ii) a decreasing fraction of high-redshift sources ($z > 3$) towards lower 1.15 mm flux density (e.g., Béthermin et al. 2015; Casey et al. 2018; Popping et al. 2020). We also note one caveat that certain extragalactic ALMA surveys of rest-frame UV/optical-selected galaxies may have a selection bias against highly dust-obscured sources at $z \gtrsim 3$. For such surveys, the most accurate measurement of the redshift distribution can be obtained after the sample is spectroscopically complete (cf., Reuter et al. 2020; Chen et al. 2021).

Among the total of 141 secure ALCS sources at $S/N_{\text{nat}} \geq 5$ or $S/N_{\text{tap}} \geq 4.5$ (58 of which fall in the PACS coverage), we successfully extracted 40, 44, 94, 77 and 55 sources at 100, 160, 250, 350 and 500 μm at above a 2σ significance, respectively. The SPIRE detection rate in the “deep”-mode clusters is higher than that in the “snapshot”-mode clusters by $\sim 10\%$. 113

(99) ALCS sources were detected at $> 2\sigma$ ($> 3\sigma$) in one *Herschel* band at least (including eight BCGs), and 91 sources were detected at $> 2\sigma$ in two *Herschel* bands at least.

Only 28 secure ALCS sources (20% of the total 141 sources) can not be extracted at $S/N > 2$ in any *Herschel* band. The 16-50-84th percentiles of 1.15 mm flux densities of these sources (0.45–0.92–1.41 mJy) are smaller than those of *Herschel*-detected sources (0.66–1.22–3.00 mJy). Except for M1206-ID58 as a brightest cluster galaxy, BCG, at $z = 0.440$, we refer to the remaining 27 sources as *Herschel*-faint galaxies in later analysis (Section 4.3; also called as *Herschel*-dropout galaxies in Boone et al. 2013).

Among the 39 tentative ALCS sources at $S/N_{\text{nat}} = 4-5$ and $S/N_{\text{tap}} < 4.5$, 22 of them can be extracted above a 2σ significance in at least one *Herschel* band, including one BCG and one mismatched source. The remaining 17 sources are undetected in any *Herschel* band including one BCG (R0032-ID162). These sources are excluded for further analysis because of a higher false ID rate.

We further justify such a 2σ detection threshold by calculating the joint probability of spurious sources through a χ^2 statistic of the detection significance in all available *Herschel* bands. For all ALCS sources extracted at $S/N > 2$ in any *Herschel* band, only five sources ($4 \pm 2\%$) exhibit p -values of spurious detection at above 0.01, including three secure ALCS sources (ACT0102-ID118, P171-ID69, P171-ID177) and two tentative sources (AS295-ID269, M1115-ID33). The largest p -value = 0.03 is seen for source P171-ID69 with $S/N_{\text{ALMA}} = 19.7$. Therefore, we conclude that with the single-band 2σ detection threshold, the number of spurious *Herschel* sources will be on the order of unity.

4. SED FITTING

4.1. Methodology

We perform far-IR SED modeling of our sample with the best available redshifts (z_{best}) using MAGPHYS (da Cunha et al. 2008, 2015). Here the z_{best} is either spectroscopic redshift (z_{spec}), published *HST*-derived z_{phot} (Section 2.5) or far-IR z_{phot} (priority from high to low). Redshift uncertainty is propagated into the uncertainties of derived physical properties through a Monte-Carlo sampling of z_{phot} likelihood when z_{spec} is not available. In order to derive and validate far-IR z_{phot} for sources without z_{spec} or *HST* z_{phot} , we also perform simultaneous far-IR SED fitting and photometric redshift estimate of our sample using MAGPHYS+PHOTO-Z (Battisti et al. 2019), the photo-z extension of MAGPHYS.

MAGPHYS assumes a Chabrier (2003) initial mass function (IMF), a continuous delayed exponential star-

formation history (SFH) with random starburst, and energy balance between dust absorption in the UV and re-emission in the IR. At far-IR wavelengths, the dust model assumed by MAGPHYS includes a warm (30–80 K) and a cold (20–40 K) component, and the prior distribution of luminosity-weighted dust temperature (T_{dust}) peaks around 37 K with a 1σ dispersion of $\sim 20\%$. Such a dust temperature is comparable to those of widely adopted spectral templates at around median $L_{\text{IR}} \sim 10^{12} L_{\odot}$ including Chary & Elbaz (2001) and Magdis et al. (2012, which is based on Draine & Li 2007 model). For a full description of the models and parameters assumed by MAGPHYS, see da Cunha et al. (2008, 2015) and Battisti et al. (2019). Here, we only include five or three bands of *Herschel* data and ALMA 1.15 mm flux densities for our SED modeling. Further optical/near-IR counterpart matching, uniform *HST* and *Spitzer* photometry and panchromatic SED fitting will be presented by another paper of the ALCS collaboration, and certain conclusions on T_{dust} and redshift distribution depending on the far-IR z_{phot} 's may be further revised.

Here, we highlight several caveats of our SED modeling obtained with MAGPHYS. First of all, the accuracy of far-IR z_{phot} is subject to a well-known degeneracy between redshift and dust temperature, typically showing an error around $\Delta z \sim 0.2(1+z)$. In addition, MAGPHYS+PHOTO-Z assumes a prior redshift distribution peaking at $z \sim 1.7$, and in practice we find that such a prior will lead to an artificial shift of z_{phot} estimate towards such a redshift. To address this issue, we adopt a uniform redshift prior instead. Furthermore, the non-thermal emission of BCGs seen at 1.15 mm cannot be properly modeled, and thus their boosted ALMA flux densities (e.g., M1931-ID41, Fogarty et al. 2019) will lead to a wrong estimate of IR luminosity and SFR. Therefore, we do not perform SED modeling for all the known BCGs. We also note that only three lensed ALCS sources are detected in X-ray among 31 clusters fields with publicly available *Chandra* data (A370-ID110, M0416-ID117, M0329-ID11; these sources will be discussed by Uematsu et al. from the collaboration). We also estimate the upper limit of AGN contribution to the derived IR luminosities. We assume the SKIRTOR model (Stalevski et al. 2012, 2016) for a type-2 AGN SED with an inclination angle of 70° . To estimate the upper limit of X-ray luminosity, we used a simple absorbed power-law model with a photon index of 1.9 and an intrinsic absorption of $\log(N_{\text{H}}/\text{cm}^2) = 23$. The median X-ray luminosity is $L_X < 2 \times 10^{43} \text{ erg s}^{-1}$ for X-ray undetected sources, corresponding to an IR luminosity of $\lesssim 6 \times 10^9 L_{\odot}$. Therefore, the AGN contamination will not be a concern for the majority ($\gtrsim 95\%$)




Figure 8. *Left:* Comparison of far-IR photometric redshifts and spectroscopic redshifts (see the references of z_{spec} 's in Table C2). Dashed black line indicates the cases where far-IR z_{phot} 's are identical to z_{spec} 's. *Right:* Comparison of photometric redshifts obtained through far-IR (*Herschel* and ALMA) and optical/near-IR (*HST*) SED fitting. Squares denote ALCS sources in the main sample, and downward triangles denote those in the secondary sample. All the sources are color-coded with their observed offsets between the *HST* and ALMA counterparts. Blue density map and contours represent the overall distribution of the cross-matched sample, except for 22 sources that are either out of *HST*/WFC3-IR coverage or detected at above 2σ in only one *Herschel* band (shown with transparent symbols). Dashed black line indicates the cases where z_{phot} 's obtained through far-IR and *HST* SED fitting are identical.

of ALCS sources, but we also note that in the case of a Compton thick AGN ($\log(N_{\text{H}}/\text{cm}^2) > 24$), the upper limit on L_X can be larger by more than an order of magnitude. Finally, MAGPHYS can only provide weak constraints on the physical properties of *Herschel*-faint galaxies individually, which are specifically discussed in Section 4.3.

Table C2 presents a summary of the best-fit galaxy properties of 125 ALCS sources detected at $S/N \geq 4$, including 47 sources that are spectroscopically confirmed and additional 42 sources with catalogued *HST* z_{phot} . This sample, further referred to as the ALCS-*Herschel* joint sample, includes 105 secure (i.e., the main sample) and 20 tentative ALCS sources (the secondary sample) detected above 2σ in one *Herschel* band at least, except for 11 BCGs and one special source (A2537-ID06) due to the poorness of SED fitting. A2537-ID06 is only detected with ALMA at $S/N=4.2$ and offset from a passive cluster dwarf galaxy by $0''.9$, and therefore it is likely a false detection with *Herschel*/SPIRE fluxes coming from an ALMA-faint *Herschel* source. The definition of this sample is also visualized as the red filled regions in Figure 6. The postage stamp images and best-fit far-IR SEDs of these 125 sources are shown in Appendix B.

We further model the dust temperature of sources in Table C2 with the *Herschel* and ALMA data. We fit the dust continuum emissions of all sources with mod-

ified blackbody (MBB) using the best available redshifts. The dust absorption coefficient is assumed to be $\kappa = 0.40 \times (\nu/250)^\beta$ in unit of $\text{cm}^2 \text{g}^{-1}$, where ν is the frequency in GHz in the rest frame. We assume a fixed dust emissivity of $\beta = 1.8$, which was widely adopted in previous studies (e.g., Díaz-Santos et al. 2017; Dudzevičiūtė et al. 2020; Sun et al. 2021b) and supported by a recent 2 mm study of SMGs at $z \simeq 1 - 3$ (da Cunha et al. 2021). Following previous work including Greve et al. (2012) and Sun et al. (2021b), we only fit the SED over a rest-frame wavelength of $50 \mu\text{m}$ to avoid optically thick regime and eliminate any possible contribution of warm dust component at shorter wavelength. We note that luminous SMGs ($L_{\text{IR}} \simeq 10^{12.5} - 10^{13} L_\odot$) are found to be optically thick at $\lambda_{\text{thick}} \sim 100 \mu\text{m}$ (e.g., Spilker et al. 2016; Simpson et al. 2017; Dudzevičiūtė et al. 2020), but we argue that less luminous ALCS sources have lower dust mass densities (e.g., Sun et al. 2021b) and therefore smaller optical depths at the same wavelength in the rest frame. The best-fit dust temperatures and masses (M_{dust}) are also presented in Table C2. The uncertainty of redshift is propagated into that of the dust temperature if the z_{spec} is unknown. The dust masses derived from this fitting procedure are also consistent with those from MAGPHYS.

We also note that many literature use the rest-frame wavelength of far-IR SED peak (λ_{peak}) to quantify the

luminosity-weighted dust temperature (e.g., Casey et al. 2018; Reuter et al. 2020; Burnham et al. 2021). This is because λ_{peak} is less dependent on dust opacity assumption compared with T_{dust} . Under the dust absorption coefficient assumption that we adopt, the conversion between λ_{peak} and T_{dust} is $\lambda_{\text{peak}} = 3 \times 10^3 T_{\text{dust}}^{-1} \mu\text{m K}$.

4.2. Validity of Far-IR Photometric Redshifts

To validate the far-IR photometric redshifts derived with MAGPHYS+PHOTO-Z, we first compare the far-IR z_{phot} 's of 47 spectroscopically confirmed sources with their z_{spec} 's in the left panel of Figure 8. The median redshift discrepancy is $(z_{\text{phot}} - z_{\text{spec}})/(1 + z_{\text{spec}}) = -0.01 \pm 0.04$, indicating an excellent agreement. We apply a Kolmogorov–Smirnov (K–S) test on the discrepancy between z_{phot} and z_{spec} divided by the uncertainty of far-IR z_{phot} , in comparison with the standard normal distribution. We conclude that the standard deviation of $(z_{\text{phot}} - z_{\text{spec}})/(1 + z_{\text{spec}})$ is well predicted by the uncertainty of far-IR z_{phot} (p -value=0.59).

The comparison of far-IR and *HST* z_{phot} 's is shown in the right panel of Figure 8. We find a general agreement of z_{phot} estimate between *HST* and far-IR SED modeling albeit with a significant dispersion. We note that the best-fit linear slope of $z_{\text{phot,FIR}}(z_{\text{phot,HST}})$ is only 0.31 ± 0.08 . However, this is mainly contributed by ~ 4 tentative sources in the secondary sample with $z_{\text{phot,HST}} \sim 4$ but $z_{\text{phot,FIR}} \sim 2$, which may be subject to a high false-ID rate ($\sim 18\%$). The discrepancy between the two z_{phot} 's, defined as $(z_{\text{phot,HST}} - z_{\text{phot,FIR}})/[1 + (z_{\text{phot,HST}} + z_{\text{phot,FIR}})/2]$, is observed to be $0.01^{+0.27}_{-0.23}$ (16–50–84th percentile) with a typical uncertainty of 0.21. We perform a K–S test on the photometric redshift discrepancies divided by their uncertainties of the cross-matched sources. The null hypothesis that the far-IR-*HST* z_{phot} discrepancies relative to their uncertainties are drawn from a standard normal distribution cannot be ruled out (p -value=0.33), reinforcing the agreement between these two photometric redshift estimates.

4.3. Herschel-Faint Galaxies

Among the 27 *Herschel*-faint galaxies reported in Section 3.5, two of the sources have been spectroscopically confirmed. They are R0600-ID164 as a [C II]-emitting lensed arc at $z = 6.072$ (Fujimoto et al. 2021; Laporte et al. 2021) and R0032-ID32 as the faintest component of a multiply lensed arc at $z = 3.631$ (Dessauges-Zavadsky et al. 2017). In addition to this, eight sources exhibit *HST* counterparts with tabulated photometric redshifts (median $z_{\text{phot}} = 2.0 \pm 1.0$; Molino et al. 2017; Coe et al. 2019). Postage stamp images of these ten sources are shown in Appendix B (Figure B3). Far-IR SEDs of




Figure 9. *Top:* Stacked *Herschel*/SPIRE images of 12 *Herschel*-faint galaxies ($S/N_{\text{ALMA}} > 5.5$) without z_{spec} or *HST* z_{phot} at 250, 350 and 500 μm . The white circles represent the apertures used for photometry. Flux density of the stacked source in each band is noted in the lower-left corner of each plot. Scale bars of 20'' are shown in the lower-right corners. *Bottom:* Far-IR SED of stacked sources (open black squares). The probability distribution of redshift is shown as the inset plot. The median far-IR SEDs of sources in the ALCS-*Herschel* joint sample are shown as open gray circles. Composite SEDs of AS2UDS SMGs (stack of $z > 3$ sources; Dudzevičiūtė et al. 2020) at $z = 2 - 6$, normalized to the median 1.15 mm flux density of *Herschel*-faint galaxies, are shown for comparison.

these galaxies are also modeled with MAGPHYS, and we note that the adopted IR spectral templates are essentially MBB spectra at around $T_{\text{dust}} = 35 \pm 6 \text{ K}$ without exceeding the *Herschel* non-detection limits except for SM0723-ID93 ($S/N_{\text{ALMA}} = 4.6$ in 2''-tapered map), which is likely a random association between a cluster dwarf galaxy with spurious ALMA source.

The remaining 17 sources do not have cross-matched *HST* z_{phot} because they are intrinsically faint shortward of 1.6 μm and/or out of *HST*/WFC3-IR coverage (Figure B4 in Appendix B). Such a near-IR-dark (also often called as “*H*-dropout/faint”) nature suggests that they are likely dust-obscured star-forming galaxies at $z \sim 4$ that have raised general interest in recent studies (e.g., Simpson et al. 2014; Franco et al. 2018; Yamaguchi et al. 2019; Alcalde Pampliega et al. 2019; Wang et al. 2019; Williams et al. 2019; Dudzevičiūtė et al. 2020;

Smail et al. 2021; Gómez-Guijarro et al. 2022; Sun et al. 2021a). For each individual sources, the non-detections in the *HST* and *Herschel* bands prevent us from deriving useful constraints of their redshifts and physical properties.

To address this issue, we stack the *Herschel* residual images of *Herschel*-faint galaxies without spectroscopic or *HST* photometric redshifts. We note that four out of five sources at $S/N_{\text{ALMA}} < 5.5$ do not show any counterpart in *HST* or *Spitzer* bands (ACT0102-ID11, M2129-ID24, ACT0102-ID251 and R1347-ID51). Therefore, these sources could be spurious detections, or highly obscured galaxies at very high redshift (i.e., similar to R0600-ID67 and R0949-ID19, the brightest *Herschel*-faint galaxies in Figure B4 that do not show any *HST* or *Spitzer* counterpart; and also the ALMA-only [C II]-emitters at $z > 6$ reported recently by Fudamoto et al. 2021). Our stacking analysis suggests that including these sources will lead to a lower S/N in SPIRE 350 and 500 μm bands, and therefore we only present the stacked SEDs of 12 sources at $S/N_{\text{ALMA}} > 5.5$.

We first normalize the *Herschel*/SPIRE residual and uncertainty images of all sources by their ALMA flux densities. Here the residual images are the scientific images with all the other *Herschel* sources subtracted assuming point-source models as described in Section 3.2. PACS images are not stacked because of the unavailability for most sources. We stack all the images in each SPIRE band using an inverse-variance-weighting method. The stacked SPIRE images are presented in Figure 9. We measure the flux densities of stacked sources using an aperture of $r_{\text{aper}} = 18''$ with appropriate aperture-correction factors. The sky background is subtracted using the median of sigma-clipped local annulus, and photometric uncertainty is estimated from the RMS of that.

The stacked source can be detected at $\sim 3\sigma$ in SPIRE 350 and 500 μm band while remains undetected at 250 μm ($< 2\sigma$). The stacked far-IR SED is shown below the stacked SPIRE images in Figure 9. We also show the median far-IR SEDs of ALCS sources that are detected with *Herschel* to visualize the clear difference in the continuum strength at below 500 μm . Here all the photometric data are normalized to the median ALMA flux density (1.09 mJy). With MAGPHYS+PHOTO-Z, we derive a median z_{phot} of 4.2 ± 1.2 (uncertainties denote the 16-84th percentiles of the likelihood distribution), IR luminosity of $10^{12.1 \pm 0.3} \mu^{-1} L_{\odot}$ and SFR of $100_{-50}^{+100} \mu^{-1} M_{\odot} \text{yr}^{-1}$ before lensing magnification correction. The derived redshift is consistent with those of *HST* *H*-faint galaxies in previous studies (e.g., Simp-




Figure 10. Stacked histogram of lensing magnification factor (μ) of 105 *Herschel*-detected ALCS sources in the main sample (filled light red bars), 27 *Herschel*-faint sources (hatched light red bars) and 20 *Herschel*-detected sources in the secondary sample (solid red steps). The dashed black line denotes the median magnification factor ($\mu_{\text{med}} = 2.6$) of the whole sample. No significant difference in μ_{med} can be identified for all three subsamples. The bin size is $\Delta \log(\mu) = 0.1$.

son et al. 2014; Wang et al. 2019; Alcalde Pampliega et al. 2019; Dudzevičiūtė et al. 2020; Sun et al. 2021a).

Using the composite SED templates of AS2UDS SMGs at $z > 3$ (Dudzevičiūtė et al. 2020) and ALESS SMGs at $z > 3.5$ (da Cunha et al. 2015), we derive best-fit z_{phot} of 3.8 ± 0.4 and 5.2 ± 0.6 , respectively. Given the large scattering of T_{dust} for sources at given luminosity and redshift (e.g., Schreiber et al. 2018; Dudzevičiūtė et al. 2020), the redshift uncertainty can be significantly underestimated with single template matching techniques. In addition to this, the median IR luminosity of SMGs in the ALESS $z > 3.5$ sample is ~ 10 times higher than that of stacked *Herschel*-faint sources and therefore likely exhibits a higher T_{dust} . This likely leads to an overestimated z_{phot} through template matching for the ALESS $z > 3.5$ sample.

5. LENS MODELING

5.1. Models, Magnifications and Multiple Images

We calculate the lensing magnification factor (μ) based on the best available redshifts. We adopt two sets of parametric lens models, the so-called Zitrin-NFW lens models (Zitrin et al. 2013, 2015) for all the HFF and CLASH clusters, and GLAFIC lens models (Oguri 2010; Okabe et al. 2020) for the RELICS clusters. The lensing magnification is derived using the

maps³ of projected cluster mass surface density (κ) and weak lensing shear (γ) at the centroid of ALCS source as $\mu = [(1 - \kappa \cdot D_{ls}/D_s)^2 - (\gamma \cdot D_{ls}/D_s)^2]^{-1}$, where D_{ls} is the angular diameter distance between the lens and the source, and D_s is the angular diameter distance to the source. We assume no magnification ($\mu = 1$) for sources within or in front of the cluster fields ($z_s < z_{cl} + 0.1$), following R16. For 16 *Herschel*-faint galaxies without catalogued redshifts (Section 4.3), we calculate their magnification at $z_s = 4.2$ uniformly.

The distribution of magnification factors of the ALCS-*Herschel* joint sample and *Herschel*-faint sample is shown in Figure 10. The 16–50–84th percentiles of the distribution of μ are 1.8–2.6–5.2, and seven sources exhibit a strong magnification with $\mu > 10$. The median lensing magnification suggests that the ALCS has reached a great depth that typical surveys in blank fields would require a $\sim 7\times$ longer observing time to achieve.

Among the 125 sources in our ALCS-*Herschel* joint sample, six groups of sources have been spectroscopically confirmed as multiply imaged systems. These include (i) ACT0102-ID118/215/224 (Caputi et al. 2021), (ii) M0417-ID46/58/121 (Kohno et al. in preparation), (iii) M0553-ID133/190/249 (Ebeling et al. 2017), (iv) M1206-ID27/55/60/61 (Ebeling et al. 2009), (v) R0032-ID53/55/57/58 (Dessauges-Zavadsky et al. 2017) and (vi) R1347-ID145/148 (Richard et al. 2021). Additionally, M1931-ID47/55/61, R0032-ID208/281/304, and R0032-ID127/131/198 are also multiply lensed candidates yet to be spectroscopically confirmed, including *HST* *H*-faint sources M1931-ID47 and R0032-ID208/281/304 that are part of the *Herschel*-faint sample. This reduces the number of independent sources in Table C2 to 109. Multiply imaged sources of the same system are shown separately in diagrams for individual sources (e.g., Figure 8), but only counted once in all statistics of physical properties (e.g., redshifts, SFRs and dust temperatures) in Section 6.

5.2. Uncertainties

The detailed characterization of the lensing magnification uncertainty (σ_μ) will be presented by a forthcoming source count paper of the ALCS collaboration (Fujimoto et al. in preparation), and here we only present qualitative estimate with a few quick methods. First of all, we estimate the σ_μ based on the uncertainty of photometric redshift for sources without spectroscopic confirmation. The typical lensing magnification error propagated from

z_{phot} uncertainty is $\sigma_\mu/\mu \sim 5\%$. In addition to this, we also compare the derived magnification factors using ALMA and *HST* source centroids, and the typical difference is found to be less than 1%. In order to quantify the σ_μ caused by extended source profile, we also measure the average magnification factor within a radius of $0''.6$ from the ALMA source centroid. Such an effect is negligible ($\sigma_\mu/\mu \lesssim 2\%$) in most cases except for M0159-ID24 and R0032-ID57 (Figure 5) because of the galaxy-galaxy lensing effect, which provides a stronger magnification gradient over a smaller angular scale.

As an alternative method, we also analyze the uncertainty map of magnification presented by Zitrin et al. (2013, 2015) in the CLASH cluster fields, obtained through a Monte Carlo Markov Chain (MCMC) fitting routines when the cluster mass models were constructed. We find that the σ_μ/μ is around $\sim 5\%$ at $\mu = 3$ assuming a fixed sky position and redshift. However, this uncertainty could be as large as $\sim 10\%$ at $\mu = 10$ and $\sim 50\%$ at $\mu = 100$, indicating that sources with larger magnifications are subject to a larger relative uncertainty and therefore their intrinsic physical properties (e.g., L_{IR} and SFR) are more uncertain.

Finally, based on the standard deviation of magnifications predicted by different lens models of the same clusters produced by different methods and groups (i.e., GLAFIC, CATS, Zitrin-NFW and Zitrin Light-Traces-Mass (LTM) for Frontier Field clusters; Oguri 2010; Richard et al. 2014; Zitrin et al. 2015; Kawamata et al. 2016, 2018), we find a typical magnification uncertainty of $\sigma_\mu/\mu \sim 20\%$. This is comparable to the σ_μ reported in R16 ($\sigma_\mu = 0.5$, $\sim 20\%$ of the median magnification for sources in this work).

6. DISCUSSION

6.1. Far-IR Colors

We further study the far-IR colors of *Herschel*-detected ALCS sources. Figure 11 displays four color-color diagrams of our sample, covering the full wavelength range from $100\ \mu\text{m}$ to $1.15\ \text{mm}$. Based on the average IR spectral templates in Rieke et al. (2009), we superimpose the redshift-evolution tracks of a typical star-forming galaxy (SFG; $L_{\text{IR}} = 10^{10.25} L_\odot$, corresponding to a SFR_{IR} of $\sim 2 M_\odot \text{yr}^{-1}$ assuming the conversion factor in Kennicutt & Evans 2012) and a ULIRG ($L_{\text{IR}} = 10^{12} L_\odot$, corresponding to a SFR_{IR} of $\gtrsim 100 M_\odot \text{yr}^{-1}$) on all of the four color-color diagrams. The dust temperatures of the SFG and ULIRG templates are around 20 K and 40 K. We also compare the distribution of our ALMA-selected sample with the R16 sample in two of the color-color diagrams where the $1.15\ \text{mm}$ flux density is not invoked.

³ These maps are made available on Mikulski Archive for Space Telescopes (MAST) as high-level science products (HLSP; <https://archive.stsci.edu/hlsp/>) and scaled to $D_{ls}/D_s = 1$.




Figure 11. Far-IR colors of *Herschel*-detected ALCS sources, color-coded with the redshifts. (a) PACS–SPIRE color (f_{100}/f_{250}) versus PACS color (f_{100}/f_{160}). (b) PACS–ALMA color (f_{100}/f_{1150}) versus PACS color (f_{100}/f_{160}). (c) SPIRE color (f_{250}/f_{350}) versus SPIRE color (f_{250}/f_{500}). (d) SPIRE–ALMA color (f_{350}/f_{1150}) versus SPIRE color (f_{250}/f_{500}). Sources shown as squares are galaxies in the main sample ($S/N \geq 5$), and downward triangles denote those in the secondary sample ($S/N = 4 - 5$). BCGs are shown in diamonds. We also compute and plot the redshift tracks of star-forming galaxies ($L_{\text{IR}} = 10^{10.25} L_{\odot}$, $T_{\text{dust}} = 20$ K; blue tracks) and ULIRGs ($L_{\text{IR}} = 10^{12} L_{\odot}$, $T_{\text{dust}} = 40$ K; red tracks) based on the spectral templates of Rieke et al. (2009), and far-IR colors at redshifts of 0, 1, 2, 3, 4 and 5 are labeled out with pentagrams. In panel (a) and (c), the green contours enclose 68% and 95% of the far-IR color distribution of *Herschel* sources detected in *Hubble* Frontier Fields (R16), and the background color maps denote the median redshift of these sources in each bin.

6.1.1. Selection bias

Through a comparison with the redshift-evolution tracks computed from the SED templates in Rieke et al. (2009), the color-color distributions of our sample match those of SFGs at $0 < z \lesssim 2.5$ and ULIRGs at $1 \lesssim z \lesssim 5$. At $z \gtrsim 2.5$, the depth of *Herschel* data is not deep enough shallow to select low- T_{dust} sources (i.e., ~ 20 K for typical SFGs) even with a lensing magnification factor of a few tens.

The selection function of ALCS is nearly flat in terms of cold dust mass across $z \simeq 1 - 6$ ($M_{\text{dust}} \gtrsim 10^8 \mu^{-1} M_{\odot}$). However, with regard to a fixed L_{IR} , the ALMA-*Herschel* joint selection do bias against low-redshift

($z \lesssim 1$) galaxies with high dust temperature (~ 40 K, ULIRG-like; see panel a and b of Figure 11). This could be interpreted as a combined effect of survey volume and 1.15 mm selection-limit. At $z < 1$, the volume density of ULIRGs is ~ 100 times lower than that of SFGs (e.g., Gruppioni et al. 2013), and therefore the expected number of ULIRG detections with ALCS in this redshift range is only on the order of unity. In addition, if we compare two galaxies with $T_{\text{dust}} = 20$ K and 40 K at the same redshift (e.g., $z = 0.5$) and the same intrinsic IR luminosity, the warmer galaxy should be fainter at 1.15 mm and thus less likely to be selected by ALCS. These galaxies with high T_{dust} are likely to be selected as ALMA-faint

Herschel sources, which are not included in our ALCS-*Herschel* joint sample. This further suggests that the observed-frame 1.15 mm selection has more bias in L_{IR} than M_{dust} (e.g., Scoville et al. 2014; Dudzevičiūtė et al. 2021) because the $M_{\text{dust}} - f_{1150}$ relation has less dependence on the dust temperature.

6.1.2. Comparison with Rawle et al. (2016)

We find that the majority of the *Herschel*-detected ALCS sources follow the similar distribution of far-IR colors as the *Herschel*-selected sources in R16 (Figure 11, panel a and c). However, only 4% of the sources in R16 are at $z > 2$, in contrast to a large fraction of 43% in our ALCS-*Herschel* joint sample. Sources in R16 were selected based on *Spitzer* and *WISE* mid-IR priors, and $> 4\sigma$ detections in at least two *Herschel* bands were required. Such a selection is biased against high-redshift sources which are faint in bluer *Herschel* bands (e.g., PACS 100 μm) but likely detectable with SPIRE 500 μm and ALMA Band 6. As shown in the panel (a) of Figure 11, PACS-selected sources in R16 are mostly at $z \lesssim 1$, which are populated by ALMA-faint *Herschel* sources that are not included in the ALCS sample. Meanwhile, as discussed earlier in this subsection, the ALCS-*Herschel* joint sample is biased against low-redshift sources with moderately low IR luminosity ($L_{\text{IR}} < 10^{11} L_{\odot}$) but high dust temperature. The combined selection effects lead to the significant difference in the redshift distributions of *Herschel* source samples of these two works.

6.2. Redshift versus 1.15 mm Flux Density

The top-left panel of Figure 12 displays the redshift distribution of sources in our ALCS-*Herschel* joint sample. The 16-50-84th percentiles of redshifts are 1.11–1.90–2.95 (1.05–1.81–2.96) for the main (full) joint sample, and no obvious difference in redshift distribution can be found among sources in clusters observed with “deep” and “snapshot” mode. The uncertainty of photometric redshifts are accounted for by the derivation of percentiles through a Monte-Carlo sampling of the probability distributions. Among the joint sample, spectroscopically confirmed sources are generally at slightly lower redshifts. The 16-50-84th percentiles of z_{spec} ’s are 1.06–1.55–2.90 (1.03–1.49–2.70) for 26 (31) independent sources in the main (full) sample. However, we note that this ALCS-*Herschel* joint sample does not include 27 *Herschel*-faint galaxies that likely reside at higher redshifts (Section 4.3).

To estimate the median redshift of the full ALCS sample, we must include the *Herschel*-faint sources. Based on the probability distribution of 17 optical/near-IR-dark *Herschel*-faint galaxies (intrinsically 15 sources

after removing multiply lensed images) derived with MAGPHYS+PHOTO-Z ($z_{\text{phot}} = 4.2 \pm 1.2$), the 16–50–84th percentiles of redshifts are 1.15–2.08–3.59 for secure ALCS sources in the main sample (middle-left panel of Figure 12; note that cluster member galaxies are not included).

The median redshift of secure ALCS sources is higher than that of the main ASPECS sample ($z_{\text{med}} = 1.80 \pm 0.15$) in the *Hubble* Ultra Deep Field (HUDF; Aravena et al. 2020), but smaller than those of slightly shallower surveys including AS2UDS ($z_{\text{med}} = 2.61 \pm 0.08$ down to ~ 1 mJy at 850 μm ; Dudzevičiūtė et al. 2020), ASAGAO ($z_{\text{med}} = 2.38 \pm 0.14$ down to ~ 0.2 mJy at 1.2 mm, note that near-IR-dark ALMA sources at $z \sim 4$ are not included; Yamaguchi et al. 2019, 2020) and GOODS-ALMA ($z_{\text{med}} = 2.56 \pm 0.13$ down to ~ 0.5 mJy at 1.1 mm in their main sample; Franco et al. 2018; Gómez-Guijarro et al. 2022). The median redshift of ALCS sources is also smaller than those of millimeter-selected strongly lensed SMGs with wider but shallower surveys (e.g., $z_{\text{med}} = 2.9 \pm 0.1$ with the *Planck*’s dusty GEMS sample, Cañameras et al. 2015; $z_{\text{med}} = 3.9 \pm 0.2$ with the SPT sample, Reuter et al. 2020), as well as that of unlensed sources selected with the MORA survey at 2 mm ($z_{\text{med}} = 3.6 \pm 0.3$ down to ~ 0.3 mJy, Casey et al. 2021).

The median redshift of millimeter and submillimeter sources as a function of flux density limit is a key test for the evolution model of dust-obscured star-formation history of the Universe (see a review by Hodge & da Cunha 2020). The right panel of Figure 12 shows the best available redshifts versus intrinsic 1.15 mm flux densities of all ALCS sources in the main sample (excluding cluster member galaxies; filled squares) and *Herschel*-detected ALCS sources in the secondary sample (open downward triangles). Assuming the probability distribution of redshifts for optical/near-IR-dark *Herschel*-faint galaxies derived with MAGPHYS+PHOTO-Z, we are able to derive the median redshift of secure ALCS sources above given 1.15 mm flux density cuts (i.e., $z_{\text{med}}(> f_{1150})$; magenta circles). The uncertainty of photometric redshift is propagated into the uncertainty of $z_{\text{med}}(> f)$ through a Monte-Carlo sampling of z_{phot} likelihood.

The median redshifts of secure ALCS sources decrease from $z_{\text{med}} = 2.40 \pm 0.29$ at $f_{1150} > 1$ mJy to $z_{\text{med}} = 2.04 \pm 0.12$ at $f_{1150} > 0.1$ mJy, where our survey is $\sim 50\%$ complete given the depth and median lensing magnification. We also perform K-S test for the redshift distributions of ALCS and ASPECS sources above 0.1 mJy, and no obvious difference can be found (p-value=0.64). For spectroscopically confirmed sample, the decrease of median redshift is not conspicuous




Figure 12. *Left:* Distribution of best available redshifts of ALCS sources. ALCS-*Herschel* sources in the main (filled light red bars) and secondary sample (solid red steps), catalogued in Table C2 (i.e., excluding *Herschel*-faint galaxies), are shown in the top panel. After assuming the probability distribution for the redshifts of optical/near-IR-dark *Herschel*-faint galaxies derived with magphys+photo- z ($z_{\text{phot}} = 4.2 \pm 1.2$, Section 4.3), the redshift distribution of sources in the main sample ($S/N_{\text{nat}} \geq 5$ or $S/N_{\text{tap}} \geq 4.5$) is shown in middle panel. Note that cluster member galaxies are not included. The redshift distributions of reference samples, including AS2UDS (Dudzevičiūtė et al. 2020, orange steps; scaled by a factor of 1/7 for the purpose of displaying), ASPECS (Aravena et al. 2020, purple steps) and GOODS-ALMA (Gómez-Guijarro et al. 2022, blue steps) are shown in the lower panel. Vertical dashed line indicates median redshift for each sample. The bin sizes are $\Delta z = 0.4$. *Right:* Best available redshifts versus ALMA 1.15 mm flux densities corrected for lensing magnification. Sources in the main (secondary) sample are shown as filled gray squares (open downward triangles). The median z_{best} and f_{1150} of 17 optical/near-IR-dark *Herschel*-faint galaxies are visualized as the filled red square. Magenta circles denote the median redshifts of secure ALCS sources at above given 1.15 mm flux density thresholds. Blue diamonds denote z_{med} at various 1.1–1.2 mm flux density thresholds in literature (Michałowski et al. 2012; Yun et al. 2012; Miettinen et al. 2017; Brisbin et al. 2017; Dunlop et al. 2017; Aravena et al. 2020; Gómez-Guijarro et al. 2022). The median redshifts as functions of 1.1 mm flux density cut, modeled by Casey et al. (2018, both dust-rich and dust-poor scenarios for $z > 4$ Universe) and Lagos et al. (2020), are plotted for comparison.

because of a smaller sample size ($z_{\text{med}} = 1.71 \pm 0.87$ at > 1 mJy to $z_{\text{med}} = 1.60 \pm 0.21$ at > 0.1 mJy). We note that a higher redshift assumption (e.g., $z_{\text{med}} \sim 6$) of optical/near-IR-dark *Herschel*-faint galaxies will not change our measurements of $z_{\text{med}}(> f)$ despite a larger standard error. We also compute the median redshifts of ALCS sources in logarithmic flux density bins from 0.1 to 2 mJy (bin size is 0.1 dex), and the null hypothesis that there is no monotonic relation between z_{med} and f_{1150} can be ruled out (p -value = 0.03, computed from Spearman’s rank correlation coefficient $\rho = 0.56$).

Further compared with previous 1.1–1.2 mm surveys of SMGs (Michałowski et al. 2012; Yun et al. 2012; Miettinen et al. 2017; Brisbin et al. 2017; Dunlop et al. 2017; Aravena et al. 2020; Gómez-Guijarro et al. 2022), the derived $z_{\text{med}}(> f_{1150})$ function suggests an increasing fraction of $z \simeq 1 - 2$ galaxies at $f_{1150} < 1$ mJy (also shown

with ASPECS and semi-empirical model presented in Popping et al. 2020). A linear fitting to the $z_{\text{med}}(> \log(f_{1150}))$ measurements suggests that the positive correlation is significant ($> 5\sigma$). We note that $z_{\text{med}}(> f_{1150})$ below 0.5 mJy was poorly probed with previous surveys because of either limited volume (ASPECS-like) or relatively shallower depth (e.g. GOODS-ALMA and single-dish surveys). Similar $z_{\text{med}}(> f)$ trends were also reported at 870 μm (Ivison et al. 2007; Stach et al. 2019; Simpson et al. 2020; Birkin et al. 2021; Chen et al. 2021).

The overall $z_{\text{med}}(> f_{1150})$ function obtained with the main ALCS sample is lower than that predicted by the SHARK semi-analytic model presented in Lagos et al. (2020, the offset is $\Delta z \sim -0.4$), but very close to that empirically modeled by Casey et al. (2018, $\Delta z \sim 0.1$) assuming a negligible contribution ($< 10\%$) of SMGs to the cosmic SFR density at $z > 4$, also known as

the “dust-poor” scenario. Nevertheless, the observed $z_{\text{med}}(> f_{1150})$ of our sample is much lower than that of “dust-rich” scenario modeled by Casey et al. (2018, $\Delta z \sim -0.7$) in which SMGs dominate the star formation ($\sim 90\%$) at $z > 4$. This may suggest that the majority of cosmic star formation at $z > 4$ will be hosted in unobscured environment, similar to the conclusions drawn in Dudzevičiūtė et al. (2020), Bouwens et al. (2020), Casey et al. (2021) and Zavala et al. (2021). However, we note that the decreasing obscured fraction of cosmic SFR density at $z > 4$ cannot be considered as the unique cause of positive $z_{\text{med}}(> f_{1150})$ relation. Other scenarios, e.g., a steep faint-end slope of dust mass function at $z = 1 \sim 2$, can lead to a similar $z_{\text{med}}(> f_{1150})$ function as observed. Detailed characterization of IR luminosity function and cosmic SFR density evolution will be presented in a future work.

6.3. Statistics of Intrinsic (Source-Plane) Properties

We study the distribution of the intrinsic SFRs (total SFR derived by MAGPHYS and corrected for lensing magnification) for ALCS sources in Figure 13. For all ALCS sources in the main sample (excluding cluster member galaxies and including *Herschel*-faint galaxies), the 16–50–84 percentiles of the distribution are 40, 94 and $178 M_{\odot} \text{ yr}^{-1}$, slightly larger than those of the GOALS sample (16–50–84 percentiles of SFR as 25, 45 and $175 M_{\odot} \text{ yr}^{-1}$; Armus et al. 2009; Howell et al. 2010), which is mostly comprised of LIRGs in the local Universe. *Herschel*-detected sources in the secondary sample exhibit LIRG-like SFRs with 16–50–84th percentiles of the distribution as 10, 25 and $97 M_{\odot} \text{ yr}^{-1}$.

We also compare the intrinsic SFRs with those of ALMA sources reported by AS2UDS (707 SMGs at $z_{\text{med}} = 2.6$; Stach et al. 2019; Dudzevičiūtė et al. 2020), GOODS-ALMA (35 sources at $z_{\text{med}} = 2.7$ studied in Franco et al. 2020; note that the SFRs are recomputed assuming Chabrier 2003 IMF) and ASPECS survey (32 sources in the main sample at $z_{\text{med}} = 1.8$; González-López et al. 2020; Aravena et al. 2020). Although the SFR distribution is similar in the image plane (lensing uncorrected), the median source-plane SFR (intrinsic) of ALCS sources is lower than that of conventional SMGs (i.e., $f > 1 \text{ mJy}$ at $850 \mu\text{m}$) in the AS2UDS sample by a factor of ~ 3 (median SFR = $236 \pm 8 M_{\odot} \text{ yr}^{-1}$; although 92 sources in AS2UDS sample are at $< 100 M_{\odot} \text{ yr}^{-1}$ with a median redshift of 2.0 ± 0.1). Below $30 M_{\odot} \text{ yr}^{-1}$, which is the median SFR of 32 sources in the main ASPECS sample down to a fidelity of 50%, our sample contain more sources than ASPECS ($N = 16$), including 11 *Herschel*-detected and ~ 10 *Herschel*-faint sources at $S/N \geq 5$, as well as 12 tentative sources in the secondary




Figure 13. Stacked histograms of observed (top panel) and intrinsic SFRs (middle panel) of ALCS sources. In the top panel, we show the image-plane SFR distributions of sources in the ALCS-*Herschel* joint sample at $S/N_{\text{ALMA}} \geq 4$ detected in both the “deep” mode (hatched gray bars) and “snapshot” mode (solid black steps). In the middle panel, we show the source-plane (intrinsic) SFR distributions of *Herschel*-detected sources in the main sample (shallow red filled bars), *Herschel*-faint sources (hatched bars) and *Herschel*-detected sources in the secondary sample (solid red steps). Cluster member galaxies are not included. We also show the SFR distributions of ALMA sources reported by AS2UDS (Dudzevičiūtė et al. 2020, orange steps; scaled by a factor of 1/7 for the purpose of displaying), GOODS-ALMA (Franco et al. 2020, blue steps) and ASPECS (Aravena et al. 2020, purple steps) in the bottom panel. Vertical dashed line in each panel indicates the median SFR of each sample. The bin size is $\Delta \log(\text{SFR}) = 0.2$.

sample. Because of the same software (MAGPHYS) and SFR tracer (far-IR) being used, these comparisons are fair. The median ratio between SFR and L_{IR} is found to be $10^{-10.1} M_{\odot} \text{ yr}^{-1} L_{\odot}^{-1}$ for both the main samples of ALCS and ASPECS, which is only offset from the conversion factor in Kennicutt & Evans (2012) by 0.1 dex (assuming Chabrier IMF). We also note that the ASPECS sample presented by Aravena et al. (2020) did not enforce any *Herschel* detection, and no SPIRE flux density information was given because of blending issues, making it difficult to directly and reliably constrain the dust temperature from the far-IR SED. This highlights the uniqueness of our ALCS-*Herschel* joint sample as the probe of (sub-)LIRG population at $z \lesssim 2$.




Figure 14. Intrinsic IR luminosity (left) and SFR (right) of ALCS sources versus redshift, including all sources in the main sample (*Herschel*-detected sources and *Herschel*-faint galaxies with z_{spec} or *HST*-derived z_{phot} ; shown as squares) and *Herschel*-detected sources in the secondary sample (downward triangles). Spectroscopically confirmed sources are shown with filled black symbols and the open symbols denote z_{phot} -only ones. The stacked optical/near-IR-dark *Herschel*-faint sources (Section 4.3) are shown as the filled red squares. Vertical bars indicate the logarithmic magnification ($\log \mu$) corrected for each source. In the left panel, we plot the detection limit of ALMA ($f > 0.3$ mJy) and *Herschel* ($S/N > 2$, $f \gtrsim 5$ mJy) in the “deep” mode without lensing magnification, assuming a ULIRG spectral template ($L_{\text{IR}} = 10^{12} L_{\odot}$ and $T_{\text{dust}} \sim 40$ K; Rieke et al. 2009). ALMA-faint *Herschel* sources and *Herschel*-faint galaxies that are not included in Table C2 (ALCS-*Herschel* joint sample) are mostly detected in the shaded green and red regions, respectively. In the right panel, we plot the SFRs of “main sequence” (MS) galaxies as a function of redshift at fixed stellar masses of 10^{11} and $10^{10.6} M_{\odot}$ (dotted and dashed line; Speagle et al. 2014). The 1σ dispersion of SFR_{MS} (~ 0.3 dex) at $M_{\text{star}} = 10^{10.6} M_{\odot}$ is shown as the filled grey region. ALMA sources reported by AS2UDS (Dudzevičiūtė et al. 2020, orange crosses) and ASPECS (Aravena et al. 2020, purple diamonds) are also plotted for comparison. The typical uncertainties of redshifts, L_{IR} and SFR are shown to the left of the lower-right legends.

Figure 14 shows the distribution of intrinsic IR luminosity and SFR as functions of redshift, highlighting the less vigorous star formation among the ALCS sources compared with unlensed SMGs (Dudzevičiūtė et al. 2020). We calculate the nominal detection limits of infrared galaxies in the ALMA (> 0.3 mJy at 1.15 mm) and *Herschel* ($> 2\sigma$ in the “deep” mode; $\gtrsim 5$ mJy) bands assuming a ULIRG template at $L_{\text{IR}} = 10^{12} L_{\odot}$ (Rieke et al. 2009). We find that the intrinsic luminosities of 29% (37%) sources in the ALCS-*Herschel* joint sample are lower than the nominal detection limit of *Herschel* (ALMA). This demonstrates the benefit of sky survey in lensing cluster fields especially for *Herschel* because the lensing magnification will allow us to extract and study sources below the nominal confusion-noise limit.

Sources whose observed flux densities are below either of the ALMA or *Herschel* detection thresholds are excluded from the ALCS-*Herschel* joint sample (Table C2). These sources are referred to as ALMA-faint *Herschel* sources (Section 3.2) or *Herschel*-faint galaxies (Section 4.3; also included in Figure 14) in our study. Based on the left panel of Figure 14, we find out that the ALMA-faint *Herschel* sources are likely distributed at $z \simeq 0 - 2$, consistent with the typical redshifts of *Herschel* sources detected in the Frontier Fields (R16).

ALCS sources without secure *Herschel* detection primarily reside at $z \gtrsim 3$, consistent with the redshifts of *Herschel*-faint galaxies presented in Section 4.3.

We also find that $\sim 77\%$ of ALCS sources (excluding cluster member galaxies) host a lower SFR than the star-formation “main sequence” (MS) at a fixed stellar mass of $M_{\text{star}} = 10^{11} M_{\odot}$ (Speagle et al. 2014). Assuming that the majority of ALCS sources are on the MS (e.g., Aravena et al. 2020; Sun et al. 2021b), this suggests a lower intrinsic stellar mass than unlensed SMG samples in literature (e.g., median $M_{\text{star}} = 10^{11.1} M_{\odot}$ in Dudzevičiūtė et al. 2020). The median stellar mass is likely around $10^{10.6 \pm 0.2} M_{\odot}$ based on a comparison with the MS across $z \simeq 1 - 4$ (Speagle et al. 2014), which will be further analyzed and constrained by a future work from the collaboration. The 1σ scattering of measured SFRs around the $\text{SFR}_{\text{MS}}(M_{\text{star}} = 10^{10.6} M_{\odot})$ is 0.4 dex.

6.4. Dust Temperature versus IR Luminosity

We study the relation between the dust temperature and intrinsic IR luminosity of sources in the ALCS-*Herschel* joint sample as shown in the left panel of Figure 15. Sources with redistributed *Herschel* fluxes (Section 3.3) are considered only once among those in each blended group. We also compare our sample with a variety of galaxies, including nearby galax-




Figure 15. *Left:* Dust temperature versus intrinsic IR luminosity of sources in the ALCS-*Herschel* joint sample. Symbols of ALCS sources are the same as those in Figure 14, and the typical uncertainty is shown in the lower-right corner. We also compare our sample with 850- μm -selected SMGs (AS2UDS sample at $z_{\text{med}} = 2.6$; Dudzevičiūtė et al. 2020), 450- μm -selected SMGs (STUDIES sample at $z_{\text{med}} = 1.8$; Lim et al. 2020), lensed SMGs (SPT sample at $z_{\text{med}} = 3.9$; Spilker et al. 2016; Strandet et al. 2016; Reuter et al. 2020; *Planck*’s dusty GEMS sample at $z_{\text{med}} = 2.9$; Cañameras et al. 2015, 2018), low-redshift ($z \lesssim 1$) LIRGs (HerMES/PEP sample; Symeonidis et al. 2013), local (U)LIRGs (GOALS sample; Díaz-Santos et al. 2017) and nearby galaxies (KINGFISH sample; Skibba et al. 2011; Hunt et al. 2015). Symbols of reference sample are labeled in the legend in the upper-left corner. *Right:* Dust temperature versus redshift of ALCS-*Herschel* sources in the intrinsic IR luminosity bins of $10^{11.5} - 10^{12} L_{\odot}$ (i.e., LIRGs; top) and $10^{12} - 10^{12.5} L_{\odot}$ (i.e., ULIRGs; bottom). Note that we only consider sources either with z_{spec} or accurate *HST* z_{phot} , and therefore the uncertainty of T_{dust} for each source is less than 20%. The p -value of Spearman’s rank correlation is shown in the lower-right corner of each panel, suggesting no conspicuous evolution of T_{dust} for the ULIRG bin but a tentative redshift dependence for the LIRG bin. Shaded gray regions indicate the $T_{\text{dust}} - z$ space that the samples are incomplete assuming 16th percentile lensing magnification ($\mu = 1.8$), where the sources are not included for p -value calculation. The gradation in gray scale indicates the incompleteness from the highest to lowest L_{IR} in each luminosity bin. 450- μm -selected SMGs (STUDIES sample, Lim et al. 2020; green pluses) and 850- μm -selected SMGs (AS2UDS sample, Dudzevičiūtė et al. 2020; orange crosses) with relatively accurate T_{dust} measurements (uncertainty $< 15\%$) are also shown for comparison. The median T_{dust} of local galaxies in the GOALS sample (Díaz-Santos et al. 2017) is shown as the horizontal short dashed line in each bin.

ies (KINGFISH sample; Skibba et al. 2011; Hunt et al. 2015), local (U)LIRGs (GOALS sample; Díaz-Santos et al. 2017), low-redshift LIRGs (HerMES/PEP sample at $z \lesssim 1$; Symeonidis et al. 2013), 850- μm -selected SMGs at $z_{\text{med}} = 2.6$ (AS2UDS sample; Dudzevičiūtė et al. 2020), 450- μm -selected SMGs at $z_{\text{med}} = 1.8$ (STUDIES sample; Lim et al. 2020) and galaxy-lensed SMGs at $z_{\text{med}} = 3.9$ (SPT sample; Spilker et al. 2016; Strandet et al. 2016; Reuter et al. 2020).

The median dust temperature is 32.0 ± 0.5 K for the ALCS-*Herschel* joint sample (with or without far-IR- z_{phot} galaxies included). Spectroscopically confirmed sources have a slightly higher median dust temperature (33.6 ± 0.9 K) than that of sources without z_{spec} ’s (31.9 ± 0.5 K). We identify a weak positive $L_{\text{IR}} - T_{\text{dust}}$ relation as $T_{\text{dust}} = (2.2 \pm 1.7) \log(L_{\text{IR}}/10^{12}) + (32.6 \pm 0.7)$ where the units of L_{IR} and T_{dust} are L_{\odot} and K, respectively. Such a weak $L_{\text{IR}} - T_{\text{dust}}$ relation (Spearman’s $\rho = 0.22$, p -value=0.03; uncertainties of T_{dust} are considered) is not consistent with the strong relations drawn from most of the high-redshift reference relations

(see also Burnham et al. 2021). It is, however, consistent with some cosmological simulation results including Liang et al. (2019).

We argue that a weak observed $L_{\text{IR}} - T_{\text{dust}}$ relation is a consequence of inhomogeneous $L_{\text{IR}}(T_{\text{dust}})$ selection threshold caused by lensing effect. The $L_{\text{IR}} - T_{\text{dust}}$ relation is a joint effect of both physics (the Stefan-Boltzmann law) and selection. If lensing magnification is not applied, the detection threshold of L_{IR} at each given redshift and millimeter flux density (e.g., $z = 2$ and $f_{1150} = 0.3$ mJy) will be exactly a monotonic function of T_{dust} (e.g., Lim et al. 2020). Combined with the selection bias towards high- L_{IR} and high- T_{dust} source at higher redshifts in the *Herschel* bands, a positive $L_{\text{IR}} - T_{\text{dust}}$ relation could be identified. In the image plane, we do find a positive $\mu L_{\text{IR}} - T_{\text{dust}}$ correlation at 4σ significance. However, with the lensing magnification, we are able to detect sources with lower L_{IR} at given T_{dust} inhomogeneously, leading to a weaker $L_{\text{IR}} - T_{\text{dust}}$ relation in the source plane. This is also seen with the strongly lensed SPT sources (Spilker et al. 2016; Reuter

et al. 2020) where the significance of $L_{\text{IR}} - T_{\text{dust}}$ relation is also around 2σ , despite that SPT sources are far more luminous. We also note that a strong $L_{\text{IR}} - T_{\text{dust}}$ relation can be identified for the joint sample of ALCS and SPT. This is because the sources of two surveys are selected in distinct ranges of IR luminosities, and the increase of T_{dust} over a wider range of L_{IR} becomes significant enough ($T \propto L^{0.25} R^{-0.5}$ according to the Stefan-Boltzmann law).

We find that at an intrinsic IR luminosity between $10^{11} - 10^{12} L_{\odot}$ (i.e., LIRGs; $z = 1.5^{+0.5}_{-0.4}$ for 16-50-84th percentiles of the redshift distribution), the dust temperatures of ALCS sources resemble those of local analogs (Díaz-Santos et al. 2017) and low-redshift LIRGs selected by *Herschel*/SPIRE in cosmological deep fields (Symeonidis et al. 2013). This indicates no or weak evolution of the dust temperature of LIRG-like galaxies from $z \sim 2$ to the local Universe, consistent with the conclusion made based on lensed HLS sources at $z_{\text{med}} = 1.9$ on $\Sigma_{\text{IR}} - T_{\text{dust}}$ plane (Sun et al. 2021b).

At $L_{\text{IR}} \gtrsim 10^{12} L_{\odot}$, ALCS sources ($z = 2.3^{+0.8}_{-0.9}$ for 16-50-84th percentiles of the redshift distribution) show lower dust temperatures than those of local ULIRGs, resembling 850- μm -selected SMGs at $z_{\text{med}} = 2.6$ (AS2UDS sample; Dudzevičiūtė et al. 2020) except for one warm outlying system (R0032 lensed arc at $z = 3.63$, $T_{\text{dust}} \sim 50$ K; Figure 5 and Dessauges-Zavadsky et al. 2017). As a result, the difference in T_{dust} between LIRGs and ULIRGs in the ALCS-*Herschel* joint sample at $z \gtrsim 1$ is not significant, which is reflected by the weak slope of $L_{\text{IR}} - T_{\text{dust}}$ relation. Previous works have reported a lower dust temperature or larger optical depth in $z \simeq 1 - 3$ SMGs compared to their local analogs (e.g., Symeonidis et al. 2009, 2013; Hwang et al. 2010; Swinbank et al. 2014). As also pointed out by a few of these studies, the high IR luminosity with relatively low dust temperature is likely caused by a larger size of star-forming region in SMGs ($R_e = 1 \sim 2$ kpc, e.g., Simpson et al. 2015; Ikarashi et al. 2015; Hodge et al. 2016; Rupopakarn et al. 2016; Fujimoto et al. 2017; Elbaz et al. 2018; Lang et al. 2019; Gullberg et al. 2019; Sun et al. 2021b; Gómez-Guijarro et al. 2022) in contrast to the compact size often seen in local ULIRGs (e.g., ~ 0.1 kpc in Arp 220; Soifer et al. 2000; Barcos-Muñoz et al. 2017; Sakamoto et al. 2017).

We further study the redshift evolution of dust temperature in the IR luminosity bins of $10^{11.5} - 10^{12} L_{\odot}$ (LIRGs) and $10^{12} - 10^{12.5} L_{\odot}$ (ULIRGs). Because 84% of ALCS sources have lensing magnification factors greater than 1.8, our joint sample is $\gtrsim 80\%$ complete for galaxies in the LIRG bin at $z \lesssim 2.5$ and $T_{\text{dust}} \lesssim 35$ K, and galaxies in the ULIRG bin at $z \lesssim 4$ and $T_{\text{dust}} \lesssim 45$ K.

The right panel of Figure 15 shows the dust temperatures versus redshifts of galaxies in the two L_{IR} bins, with the incomplete region on $T_{\text{dust}} - z$ plane visualized with gray shade. Note that we only include galaxies with uncertainties of $\Delta T/T$ less than 20%, which only consist of sources with z_{spec} or accurate *HST* z_{phot} .

To test the existence of any monotonic redshift evolution of T_{dust} , we compute the Spearman's rank correlation coefficient and p -value for sources within the redshift ranges of high completeness. A weak positive $T_{\text{dust}}(z)$ relation can be tentatively drawn for sources in the $L_{\text{IR}} = 10^{11.5} - 10^{12} L_{\odot}$ bin (p -value = 0.06), with sources at $z \lesssim 2$ showing lower T_{dust} than that of local LIRGs (Díaz-Santos et al. 2017). We argue that the selection bias against low-redshift sources with warm dust temperatures ($T_{\text{dust}} \gtrsim 35$ K; discussed in Section 6.1) is likely the cause of this weak $T_{\text{dust}}(z)$ relation.

We note that similar trend is also seen with Symeonidis et al. (2013) for galaxies with $L_{\text{IR}} = 10^{11.6} - 10^{11.8} L_{\odot}$ at $z < 1$, and positive $T_{\text{dust}}(z)$ trend has been suggested by certain stacking analyses (e.g., Magnelli et al. 2014; Schreiber et al. 2018; Simpson et al. 2019) and simulations (e.g., Liang et al. 2019; also Lagos et al. 2020 but a weak $T_{\text{dust}}(z)$ evolution). However, the samples used by most of the stacking analyses are stellar-mass-selected, which are different in total SFR from L_{IR} - or M_{dust} -selected samples obtained with submillimeter/millimeter surveys. This means that the evolution of $T_{\text{dust}}(z)$ at fixed stellar mass can be a combined effect of the weak evolution of $L_{\text{IR}} - T_{\text{dust}}$ relation and strong evolution of star-forming main sequence (i.e., increasing SFR/ M_{star} and thus $L_{\text{IR}}/M_{\text{star}}$ towards higher redshifts; see also a recent study by Drew & Casey 2022). The positive $T_{\text{dust}}(z)$ trend is not seen in 450/850- μm -selected SMGs in the AS2UDS and STUDIES sample (Dudzevičiūtė et al. 2020, 2021; Lim et al. 2020).

For ULIRGs in the bin of $L_{\text{IR}} = 10^{12} - 10^{12.5} L_{\odot}$, no redshift evolution of T_{dust} can be identified (p -value = 0.71). Similar conclusion can also be drawn with the STUDIES and AS2UDS SMGs with relatively accurate T_{dust} measurements (uncertainty $< 15\%$; typically requires ≥ 1 -band SPIRE detection) in this L_{IR} bin.

6.5. Gas Depletion Time Scale

Assuming a canonical gas-to-dust mass ratio (GDR) of $\delta_{\text{GDR}} = 100$, we can estimate a median gas depletion time scale for sources in the ALCS-*Herschel* joint sample as $t_{\text{dep}} = \delta_{\text{GDR}} M_{\text{dust}}/\text{SFR} = 190^{+266}_{-95}$ Myr (error bar denotes 1σ dispersion). Sources with higher T_{dust} generally show shorter t_{dep} (equivalently higher star-forming efficiency, $\text{SFE} = \text{SFR}/M_{\text{gas}}$). For sources only with far-IR z_{phot} , their dust temperatures are close to the peak




Figure 16. Molecular gas depletion time scale ($t_{\text{dep}} = \delta_{\text{GDR}} M_{\text{dust}} / \text{SFR}$) of sources in the ALCS-*Herschel* joint sample as a function of redshift. The typical uncertainty is shown in the lower-right corner. Symbols are the same as those in Figure 14, but we de-emphasize far-IR- z_{phot} sources with semi-transparent symbols to address the degeneracy between T_{dust} priors and t_{dep} . Sources in the L_{IR} -limited ULIRG subsample at $z \simeq 1 - 4$ where ALMA and SPIRE detections are $\gtrsim 80\%$ complete (see Section 6.4 and the lower-right panel of Figure 15) are highlighted with red open circles, and no redshift dependence of t_{dep} can be identified. We also plot sources in ASPECS main sample (Aravena et al. 2020, purple diamonds), AS2UDS (Dudzevičiūtė et al. 2020, orange crosses), GOODS-ALMA (Franco et al. 2020, blue pluses), PHIBSS (Tacconi et al. 2018, cyan dots), *Planck*’s dusty GEMs (Cañameras et al. 2018, pink diamonds) and SPT sample (Reuter et al. 2020, shallow red circles) for comparisons. $t_{\text{dep}}(z)$ relations based on the prescriptions of Scoville et al. (2017), Tacconi et al. (2018) and Liu et al. (2019) are shown in magenta, cyan and dashed black lines, respectively, assuming $M_{\text{star}} = 10^{10.6} M_{\odot}$ and main-sequence SFR.

of prior T_{dust} distribution as assumed by MAGPHYS, and therefore the t_{dep} is nearly constant. Despite such a degeneracy, the derived median t_{dep} does not change significantly if we exclude sources without z_{spec} (166 ± 25 Myr) or *HST* z_{phot} (219 ± 30 Myr) from the statistics. We further note that t_{dep} is a conserved quantity independent of lensing magnification effect.

The derived t_{dep} of our sample is broadly consistent with those derived with AS2UDS sample (295_{-185}^{+307} Myr; Dudzevičiūtė et al. 2020), ASPECS main sample (299_{-167}^{+431} Myr; Aravena et al. 2020) and HLS bright lensed source sample (226_{-73}^{+196} Myr; Sun et al. 2021b) using the same method (i.e., $\delta_{\text{GDR}} = 100$ with M_{dust} derived from MAGPHYS SED modeling). Although the requirement of *Herschel* detections will result in a selection bias towards sources with higher T_{dust} and shorter

t_{dep} , the use of *Herschel* data provides characterization of obscured SFR and t_{dep} in higher precision.

Figure 16 shows the gas depletion time scale as a function of redshift for sources in the ALCS-*Herschel* joint sample. We also compare ALCS sources with ASPECS main sample (Aravena et al. 2020), AS2UDS (Dudzevičiūtė et al. 2020), GOODS-ALMA (Franco et al. 2020) and PHIBSS sample (Tacconi et al. 2018) across comparable redshift ranges. The best-fit models of $t_{\text{dep}}(z)$ at $M_{\text{star}} = 10^{10.6} M_{\odot}$ and main-sequence SFR (Speagle et al. 2014), based on the prescriptions of Scoville et al. (2017), Tacconi et al. (2018) and Liu et al. (2019), are also plotted for comparison. Sources in our sample exhibit a shorter depletion time scale than the model predictions. This is caused by the IR-selection nature (i.e., favoring sources above the star-forming MS with higher L_{IR} and T_{dust}) and potentially low T_{dust} assumptions for single-band millimeter continuum sources in previous studies (e.g., 25 K in Scoville et al. 2017). Additionally, δ_{GDR} is observed to be a function of metallicity (e.g., Leroy et al. 2011; $\delta_{\text{GDR}} \sim 200$ at half solar metallicity), which could introduce further uncertainty to M_{gas} and t_{dep} .

Although a declining trend of t_{dep} towards higher redshift can be identified, we argue that this is a selection effect of sources with higher L_{IR} and T_{dust} towards higher redshift. If we restrict the sample to an intrinsic IR luminosity of $10^{12} - 10^{12.5} L_{\odot}$ at $z \simeq 1 - 4$ where the ALMA and *Herschel*/SPIRE detections are $\gtrsim 80\%$ complete (Section 6.4 and the lower-right panel of Figure 15), we find no significant redshift dependence of t_{dep} (the p -value of Spearman’s rank correlation is 0.21). This is consistent with the conclusion of Dudzevičiūtė et al. (2020) drawn upon L_{IR} -limited sample with both ALMA and SPIRE detections.

6.6. Dust Temperature of a [C II]-Emitter at $z = 6.07$

R0600-ID164 is a [C II]-emitting lensed arc at $z = 6.072$ discovered blindly by the ALCS (named as RXCJ0600–2007 z6.1/z6.2 in Fujimoto et al. 2021). Laporte et al. (2021) reported the ALMA dust continuum detection of this source ($S/N = 4.84$ on the $2''$ -tapered map). Although this source is not detected in any *Herschel* band, the upper limits of flux densities at 250–500 μm can be used to constrain its dust temperature.

Figure 17 displays the observed far-IR SED of R0600-ID164 before lensing magnification correction (red open circles; upper limits are at 3σ). We compare the SED with modified blackbody spectral templates at intrinsic $T_{\text{dust}} = 50 - 125$ K with the CMB effects considered following the prescription of da Cunha et al. (2013). Based on the dust mass, dust continuum size reported




Figure 17. Rest-frame far-IR SED of R0600-ID164 (red open circles), a [C II]-emitting lensed arc at $z = 6.07$ blindly discovered with the ALCS (Fujimoto et al. 2021; Laporte et al. 2021). Modified blackbody spectra at $T_{\text{dust}} = 40 - 120$ K (with dust emissivity $\beta = 1.6$ and 2.0 as dashed and solid lines) are plotted for comparison. Compared with the MACS0416 Y1 (open black squares; Bakx et al. 2020) with unusually warm dust temperature ($T_{\text{dust}} > 80$ K) and/or steep dust emissivity index ($\beta > 2$), we are able to rule out such a scenario with *Herschel*/SPIRE non-detections. SEDs of galaxies at $z \simeq 6 - 7$ with normal dust temperature (40–50 K; Hashimoto et al. 2019; Harikane et al. 2020; Bakx et al. 2021) are shown as green dots. Note: all SEDs of galaxies in literature are scaled to the same flux density of R0600-ID184 at $160 \mu\text{m}$ in the rest frame. Upper limits are at 3σ .

in Laporte et al. (2021) and dust absorption coefficient adopted in Section 4.1, we find a low dust mass surface density of $(6 \pm 3) \times 10^6 M_{\odot} \text{kpc}^{-2}$. Therefore, the optical depth of dust continuum is on the order of unity at $\lambda_{\text{thick}} \lesssim 30 \mu\text{m}$ in the rest frame. The dust emissivities are assumed as $\beta = 1.6$ and 2.0 . We also compare R0600-ID164 with galaxies at $z \simeq 6 - 7$ with normal dust temperatures (40–50 K; shown as green dots) confirmed with ALMA continuum detections in two bands at least, including J1211–0118, J0217–0208 ($z = 6.03$ and 6.20 , respectively; Harikane et al. 2020), A1689-zD1 ($z = 7.13$; Watson et al. 2015; Knudsen et al. 2017; Inoue et al. 2020; Bakx et al. 2021) and B14–65666 ($z = 7.15$; Hashimoto et al. 2019; Sugahara et al. 2021).

Bakx et al. (2020) showed that MACS0416 Y1, a lensed galaxy at $z = 8.31$ (black open squares in Figure 17; also Tamura et al. 2019), exhibits abnormally warm dust temperature ($T_{\text{dust}} > 80$ K, 90% confidence) and/or steep dust emissivity index ($\beta > 2$). With the three-band SPIRE flux density upper limits, we calculate the χ^2 of non-detections assuming a $T_{\text{dust}} = 80$ K, $\beta = 2.0$ and $\lambda_{\text{thick}} = 30 \mu\text{m}$ dust continuum model. The derived χ^2 is 6.6 (reduced $\chi^2_{\nu} = 2.2$), indicating

that such a model can be ruled out at $> 90\%$ confidence level. Although assuming lower dust emissivity and longer λ_{thick} will lead to a smaller χ^2 , we can draw the conclusion that the T_{dust} and β of R0600-ID164 are not as extreme as those observed for MACS0416 Y1. However, we also note that the differential lensing effect seen around the caustic line ($\mu = 30 - 160$; Fujimoto et al. 2021) may introduce further uncertainty on the dust temperature constraint.

7. SUMMARY

We present an ALMA-*Herschel* joint study of 1.15 mm continuum sources detected by the ALMA Lensing Cluster Survey (ALCS), an ALMA Cycle-6 large program (PI: Kohno) dedicated for surveying intrinsically faint millimeter sources with the assistance of gravitational lensing. All of the 33 ALCS cluster fields have been observed by *Herschel*/SPIRE at 250, 350 and $500 \mu\text{m}$; 18 of them were observed down to confusion-limited noise levels, and have high-quality *Herschel*/PACS 100 and $160 \mu\text{m}$ coverages (i.e., in the “deep” mode). We conducted PSF flux extraction for all 141 secure ALCS sources ($S/N_{\text{nat}} \geq 5$ in the native-resolution maps or $S/N_{\text{tap}} \geq 4.5$ in the $2''$ -tapered maps; the main sample) and 39 tentative ALCS sources with near/mid-IR counterparts ($S/N_{\text{nat}} = 4 - 5$ and $S/N_{\text{tap}} < 4.5$; the secondary sample). We then performed far-infrared SED modeling for 125 of them, which are detected in the *Herschel* bands at $S/N \geq 2$ (the ALCS-*Herschel* joint sample). The main results are the following:

1. 113 out of the 141 secure ALCS sources were detected at $> 2\sigma$ in at least one *Herschel* band, and 22 out of 39 tentative ALCS sources were detected above the same threshold (Table C1). The single-band *Herschel* detection rate remains nearly constant as a function of S/N_{ALMA} at 100 and $250 \mu\text{m}$, but clearly correlated with S/N_{ALMA} at longer wavelength (350 and $500 \mu\text{m}$).
2. We conducted far-IR SED modeling and derived photometric redshifts for 125 *Herschel*-detected ALCS sources (109 independently) excluding BCGs. Among them, 47 sources are spectroscopically confirmed, and additional 42 sources have catalogued *HST* photometric redshifts. Physical properties (L_{IR} , SFR, M_{dust} , T_{dust}) of sources in this ALCS-*Herschel* joint sample are presented in Table C2, derived based on the best available redshifts (z_{best}).
3. 27 lensed ALCS sources in the main sample are not detected in any *Herschel* band ($< 2\sigma$). Among these *Herschel*-faint galaxies, ten sources have catalogued z_{spec} ($z = 3.631, 6.072$; two sources) or

- HST* z_{phot} ($z_{\text{med}} = 2.0 \pm 1.0$; eight sources). The remaining 17 optical/near-IR-dark sources likely reside at $z_{\text{phot}} = 4.2 \pm 1.2$, hosting a typical IR luminosity of $10^{11.7 \pm 0.3} (2.6/\mu) L_{\odot}$ and obscured SFR of $40_{-20}^{+40} (2.6/\mu) M_{\odot} \text{ yr}^{-1}$ (median lensing magnification factor $\mu_{\text{med}} = 2.6$).
4. ALCS sources are cold-dust-mass-selected ($M_{\text{dust}} \gtrsim 10^8 \mu^{-1} M_{\odot}$) across $z \simeq 1 - 6$. However, at a fixed L_{IR} , the ALMA-*Herschel* joint selection is biased against galaxies with local-ULIRG-like dust temperature ($T_{\text{dust}} \sim 40 \text{ K}$) at $z \lesssim 1$. This is because (i) the effective survey volume of ALCS at $z < 1$ is limited for the selection of ULIRGs ($L_{\text{IR}} > 10^{12} L_{\odot}$), and (ii) at given redshift and intrinsic L_{IR} , galaxies with higher dust temperatures (i.e., ULIRG-like) will appear fainter in ALMA Band 6, and thus it is more challenging to detect them with the ALCS.
 5. The 16–50–84th percentiles of the redshift distribution of the ALCS sources in the main sample (excluding cluster member galaxies) are 1.15–2.08–3.59. The median redshift of secure ALCS sources is higher than that of sources selected with the deep ASPECS survey (Aravena et al. 2020), but lower than those of sources in shallower ALMA Band-6/7 surveys (e.g., Dudzevičiūtė et al. 2020; Yamaguchi et al. 2020; Gómez-Guijarro et al. 2022). Together with the median redshift as a function of 1.15 mm flux density cut ($z_{\text{med}}(> f_{1150})$), this suggests an increasing fraction of $z \simeq 1 - 2$ galaxies among fainter millimeter sources ($\sim 0.1 \text{ mJy}$) and potentially decreasing obscured fraction of cosmic star formation at $z > 4$.
 6. With a median lensing magnification of $\mu = 2.6_{-0.8}^{+2.6}$, we derive an intrinsic SFR distribution of $94_{-54}^{+84} M_{\odot} \text{ yr}^{-1}$ for sources at $S/N \geq 5$ (errorbar denotes 1σ dispersion). The intrinsic SFRs and IR luminosities of our sample are slightly higher than those of local LIRGs (e.g., GOALS sample; Howell et al. 2010) but lower than those of conventional unlensed SMGs selected at $z \sim 2 - 3$ by a factor of ~ 3 (Dudzevičiūtė et al. 2020).
 7. We compare the dust temperatures (modeled by a modified blackbody spectrum with $\beta = 1.8$) versus IR luminosities with various galaxy samples from the literature. The median T_{dust} is $32.0 \pm 0.5 \text{ K}$ for the ALCS-*Herschel* joint sample. Our result suggests no or weak evolution of the T_{dust} of LIRGs ($L_{\text{IR}} < 10^{12} L_{\odot}$) from $z \sim 2$ to the local Universe at a given L_{IR} . At $L_{\text{IR}} \gtrsim 10^{12} L_{\odot}$, ALCS sources exhibit cooler dust temperature compared with local ULIRGs, and no evolution of T_{dust} can

be found at $z \simeq 1 - 4$, as has been reported by previous SMG surveys (Dudzevičiūtė et al. 2020).

8. Assuming a canonical gas-to-dust ratio of 100, the gas depletion time scales for sources in the ALCS-*Herschel* sample are found to be $190_{-95}^{+265} \text{ Myr}$. For sources in the IR luminosity range of $10^{12} - 10^{12.5} L_{\odot}$, no redshift evolution of t_{dep} can be identified across $z \simeq 1 - 4$.
9. The $z = 6.072$ [C II]-emitting lensed arc, R0600-ID164, is the highest-redshift source in our ALCS continuum source sample ($S/N \geq 4$) confirmed so far (Fujimoto et al. 2021; Laporte et al. 2021). With the *Herschel*/SPIRE non-detections, we can rule out a MACS0416 Y1-like warm dust temperature ($T_{\text{dust}} > 80 \text{ K}$; Bakx et al. 2020) at $> 90\%$ confidence level.

With the lensing magnification provided by massive galaxy clusters, our joint analysis based on ALMA and *Herschel* observations reveal the population of galaxies at $z \simeq 1 - 3$ with moderate star formation rate at a few tens of solar masses per year. ALMA surveys in blank fields would require ~ 7 times longer observation times to reach the same depth. In the high-redshift regime, we discover 17 optical/near-IR-dark *Herschel*-faint sources that are likely dust-obscured star-forming galaxies at $z \sim 4$. These sources are excellent targets for multi-wavelength follow-up. Future ALMA spectral line-scan and *JWST*/NIRSpec observations will provide key insights into the nature of these distant dusty star-forming galaxies, revealing their contribution to the obscured cosmic SFR density at $z > 4$.

ACKNOWLEDGMENTS

We thank the anonymous referee for helpful comments. FS acknowledges support from the NRAO Student Observing Support (SOS) award SOSPA7-022. FS and EE acknowledge funding from JWST/NIRCam contract to the University of Arizona, NAS5-02105. KK acknowledges support from JSPS KAKENHI Grant Number JP17H06130 and the NAOJ ALMA Scientific Research Grant Number 2017-06B. IRS acknowledges support from STFC (ST/T000244/1). P.G.P.-G. acknowledges support from Spanish Government grant PGC2018-093499-B-I00. MO acknowledges support from JSPS KAKENHI Grant Numbers JP18K03693, JP20H00181, JP20H05856, JP22H01260. AZ acknowledges support from the Ministry of Science and Technology, Israel. We thank Ugnė Dudzevičiūtė for sharing the

composite SEDs of AS2UDS SMGs. We thank Claudia Lagos for helpful discussion.

This paper makes use of the following ALMA data: ADS/JAO.ALMA#2018.1.00035.L, 2013.1.00999.S and 2015.1.01425.S. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada), MOST and ASIAA (Taiwan), and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. This work is based on observations made with *Herschel*. *Herschel* is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA. This work is based (in part) on observa-

tions made with the *Spitzer Space Telescope*, which was operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA. This research is based on observations made with the NASA/ESA *Hubble Space Telescope* obtained from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5–26555. Some of the data presented in this paper were obtained from the Mikulski Archive for Space Telescopes (MAST) at the Space Telescope Science Institute. The specific observations analyzed can be accessed via [Frontier Fields](#), [CLASH](#) and [RELICS](#).

Facilities: HST (ACS and WFC3); Spitzer (IRAC); Herschel (PACS and SPIRE); ALMA.

Software: astropy (Astropy Collaboration et al. 2013), Photutils (Bradley et al. 2019), MAGPHYS (da Cunha et al. 2008, 2015; Battisti et al. 2019).

Table 1. Summary of *Herschel* observations for 33 ALCS galaxy cluster fields

Cluster Name	Group ^a	Coordinates		Short Name	<i>Herschel</i> /PACS 100/160 μm		<i>Herschel</i> /SPIRE 250/350/500 μm	
		RA	Dec		Observation ID	t_{obs} (h) ^b	Observation ID	t_{obs} (s) ^b
18 HLS-“deep” clusters:								
Abell209	CLASH	01:31:52.5	−13:36:38	A209	134218841[8,9]	4.28	1342188581	5803
Abell383	CLASH	02:48:03.3	−03:31:44	A383	134218915[1,2]	4.28	1342189503	5803
MACS0329.7–0211	CLASH	03:29:41.6	−02:11:47	M0329	134224928[0,1]	4.28	13422[14564,39844]	1580
MACS0429.6–0253	CLASH	04:29:36.1	−02:53:08	M0429	1342250[641,836]	4.28	13422[39932,41124]	1580
MACS1115.9+0129	CLASH	11:15:52.0	01:29:56	M1115	13422476[72,91]	4.28	13422[23226,56866]	1580
MACS1206.2–0847	CLASH	12:06:12.2	−08:48:02	M1206	134225745[5,6]	4.28	13422[34856,47273]	1580
MACS1311.0–0310	CLASH	13:11:01.6	−03:10:39	M1311	13422486[26,56]	4.28	13422[34800,59416]	1580
RXJ1347–1145	CLASH	13:47:30.5	−11:45:10	R1347	134221383[6,7]	5.23	13422[01256–63,47859–61]	12728
MACS1423.8+2404	CLASH	14:23:47.7	24:04:40	M1423	134218821[5,6]	5.47	1342188159	6636
MACS1931.8–2635	CLASH	19:31:49.6	−26:34:34	M1931	13422416[19,81]	4.28	13422[15993,54639]	1580
MACS2129.4–0741	CLASH	21:29:26.2	−07:41:26	M2129	134218780[1,2]	5.47	1342195710	5786
RXJ2129.7+0005	CLASH	21:29:39.9	00:05:18	R2129	134218725[6,7]	5.32	1342188167	6636
Abell2744	HFF	00:14:21.2	−30:23:50	A2744	134218825[1,2]	5.47	1342188584	5803
Abell370	HFF	02:39:52.9	−01:34:36	A370	13422333[2,3]	5.23	13422[01311–18,48002–04]	12728
MACSJ0416.1–2403	HFF	04:16:08.9	−24:04:28	M0416	134225029[1,2]	4.28	13422[39858,41122]	1580
MACSJ1149.5+2223	HFF	11:49:36.3	22:23:58	M1149	134221179[7,8]	4.28	1342222841	5786
AbellS1063	HFF	22:48:44.4	−44:31:48	AS1063	134218822[2,3]	5.47	1342188165	6636
Abell2537	RELICS	23:08:22.2	−02:11:32	A2537	1342187[799,800]	5.47	1342188179	6636
15 HLS-“snapshot” clusters:								
RXCJ0032.1+1808	RELICS	00:32:11.0	18:07:49	R0032	1342234685	169
MACSJ0035.4–2015	RELICS	00:35:26.9	−20:15:40	M0035	1342234697	169
ACTCLJ0102–49151	RELICS	01:03:00.0	−49:16:22	ACT0102	1342258408	169
MACSJ0159.8–0849	RELICS	01:59:49.4	−08:50:00	M0159	1342237535	169
AbellS295	RELICS	02:45:31.3	−53:02:24	AS295	1342236215	169
MACSJ0257.1–2325	RELICS	02:57:10.2	−23:26:11	M0257	1342214559	169
PLCKG171.9–40.7	RELICS	03:12:56.9	08:22:19	P171	1342239833	169
Abell3192	RELICS	03:58:53.0	−29:55:44	A3192	1342239861	169
MACSJ0417.5–1154	RELICS	04:17:33.7	−11:54:22	M0417	1342239855	169
MACSJ0553.4–3342	RELICS	05:53:23.0	−33:42:29	M0553	1342227700	169

Table 1 continued

Table 1 (*continued*)

Cluster Name	Group ^a	Coordinates		Short Name	<i>Herschel</i> /PACS 100/160 μm		<i>Herschel</i> /SPIRE 250/350/500 μm	
		RA	Dec		Observation ID	t_{obs} (h) ^b	Observation ID	t_{obs} (s) ^b
RXCJ0600.1–2007	RELICS	06:00:09.7	–20:08:08	R0600	1342230801	169
SMACSJ0723.3–7327	RELICS	07:23:19.4	–73:27:15	SM0723	1342229668	169
RXCJ0949.8+1707	RELICS	09:49:50.8	17:07:15	R0949	1342246604	169
Abell2163	RELICS	16:15:48.3	–06:07:36	A2163	1342229566	169
RXCJ2211.7–0350	RELICS	22:11:45.9	–03:49:44	R2211	1342211362	169

NOTE—Clusters with *Herschel*/PACS data are considered as observed in the “deep” mode, and the remaining ones are considered as observed in the “snapshot” mode (see Section 2.3). *Herschel* observation IDs in brackets indicate the difference in the last a few digits, for example, the SPIRE data of Abell370 was taken with observation IDs 1342201311–1342201318 and 1342248002–1342248004.

^a Group name of *HST* Program.

^b Total scan time of all observations.

Table 2. Summary of median 1σ depth of prior-based *Herschel* catalogues within the ALCS footprints

Cluster Name	$N(\text{main})^a$	PACS 100 μm	PACS 160 μm	SPIRE 250 μm	SPIRE 350 μm	SPIRE 500 μm
		(mJy)	(mJy)	(mJy)	(mJy)	(mJy)
Abell209	1	... (1.5)	... (3.0)	... (5.0)	... (5.1)	... (5.7)
Abell383	1	0.6 (1.5)	1.8 (2.9)	2.8 (5.5)	2.7 (5.1)	... (5.9)
MACS0329.7–0211	1	0.5 (1.4)	... (2.8)	... (7.1)	... (6.8)	... (7.5)
MACS0429.6–0253	3	0.6 (1.4)	1.0 (2.9)	3.3 (7.5)	... (6.9)	... (8.0)
MACS1115.9+0129	4	0.8 (1.4)	0.8 (2.9)	3.2 (7.8)	4.3 (6.5)	3.9 (7.1)
MACS1206.2–0847	6	0.5 (1.4)	1.7 (3.1)	15.5 (8.6)	2.2 (7.5)	4.4 (8.2)
MACS1311.0–0310	2	0.5 (1.4)	1.3 (2.8)	3.2 (7.4)	2.7 (6.5)	2.6 (7.2)
RXJ1347–1145	6	0.5 (1.3)	0.7 (2.4)	2.4 (4.3)	6.7 (4.3)	2.0 (5.2)
MACS1423.8+2404	2	0.6 (1.5)	1.3 (2.9)	3.4 (5.0)	3.0 (5.0)	2.3 (5.6)
MACS1931.8–2635	4	1.0 (1.4)	1.2 (2.8)	2.8 (7.0)	3.0 (6.5)	3.0 (7.2)
MACS2129.4–0741	2	0.5 (1.5)	0.8 (3.4)	2.1 (6.8)	2.9 (6.9)	... (6.9)
RXJ2129.7+0005	2	... (1.3)	1.1 (3.1)	3.2 (5.5)	... (5.7)	4.1 (6.1)
Abell2744	6	0.9 (1.5)	1.5 (3.0)	2.4 (5.1)	3.3 (5.4)	2.8 (5.8)
Abell370	5	0.5 (1.3)	1.3 (2.8)	2.8 (5.0)	3.2 (5.3)	2.2 (6.2)
MACSJ0416.1–2403	4	0.5 (1.4)	0.8 (2.7)	2.7 (7.0)	1.9 (6.0)	1.7 (7.3)
MACSJ1149.5+2223	1	0.9 (1.5)	3.2 (3.0)	2.1 (5.4)	2.2 (4.9)	3.5 (5.6)
AbellS1063	4	0.9 (1.4)	0.8 (2.3)	3.3 (5.5)	5.0 (5.6)	3.2 (6.4)
Abell2537	2	1.5 (1.5)	2.2 (3.1)	3.2 (5.9)	4.0 (6.1)	2.6 (6.0)
RXCJ0032.1+1808	17	4.5 (14.4)	6.4 (16.6)	4.8 (16.9)
MACSJ0035.4–2015	2	3.6 (10.7)	... (11.6)	... (12.9)
ACTCLJ0102–49151	12	4.7 (11.8)	4.8 (11.7)	5.0 (15.7)
MACSJ0159.8–0849	4	4.5 (11.1)	4.1 (12.0)	5.2 (13.7)
AbellS295	0 (11.7)	... (12.4)	... (15.2)
MACSJ0257.1–2325	1	4.4 (11.8)	4.5 (11.8)	4.7 (15.3)
PLCKG171.9–40.7	3	5.1 (12.3)	... (12.8)	... (13.7)
Abell3192	4	4.4 (11.1)	4.7 (12.1)	4.6 (14.1)
MACSJ0417.5–1154	7	4.7 (10.9)	5.7 (13.0)	5.2 (15.2)
MACSJ0553.4–3342	13	4.5 (10.6)	4.6 (11.7)	6.1 (14.5)
RXCJ0600.1–2007	4	4.4 (11.5)	5.6 (12.4)	... (13.8)
SMACSJ0723.3–7327	2 (12.2)	... (13.2)	... (15.5)
RXCJ0949.8+1707	4	5.0 (12.0)	5.3 (12.1)	5.7 (14.6)
Abell2163	0 (11.1)	... (10.9)	... (13.8)
RXCJ2211.7–0350	3	5.0 (12.5)	6.3 (12.6)	6.9 (14.4)

Table 2 *continued*

Table 2 (continued)

Cluster Name	$N(\text{main})^a$	PACS 100 μm	PACS 160 μm	SPIRE 250 μm	SPIRE 350 μm	SPIRE 500 μm
		(mJy)	(mJy)	(mJy)	(mJy)	(mJy)

NOTE—Values in parenthesis are RMS noises directly measured from 2D *Herschel* uncertainty maps within the ALCS footprints (i.e., without any dedicated positional prior; Section 3.3).

^aNumber of ALMA sources in the main sample ($S/N_{\text{nat}} \geq 5$ in the native-resolution maps, or $S/N_{\text{nat}} \geq 4.5$ in the $2''$ -tapered maps).

APPENDIX

A. QUALITY OF THE PHOTOMETRIC RESULTS




Figure A1. *Left:* Ratios of flux densities measured with aperture (f_{aper}) and PSF (f_{psf}) photometry in the five *Herschel* bands (100–500 μm from the top to the bottom). All the sources detected in the *Herschel* images at $S/N > 5$ are shown as the empty hatched histograms, and the ALCS sources are shown as the solid filled ones. Dashed black lines indicate the cases where the flux densities obtained through PSF and aperture photometry are identical. The number of sources and the median $f_{\text{aper}}/f_{\text{psf}}$ ratio (with 1σ standard error) are also shown in each panel for each *Herschel* band. *Right:* *Herschel* flux density ratios of submillimeter sources in *Hubble* Frontier Fields measured by Rawle et al. (2016) and this work (i.e., both detected and undetected by the ALCS), plotted as a function of S/N in the corresponding *Herschel* band (S/N_{Herschel}). Symbols are the same as those in Figure 4 and also labeled in the upper-right corner. The solid gray line indicates the case in which the two measurements are identical, and the dashed lines indicate the cases of $\pm 10\%$ deviation. The shaded region denotes the 1σ dispersion range of flux ratios for sources detected at any given S/N_{Herschel} .

In order to examine the quality of our PSF photometric results, we first compared our *Herschel* flux densities (f_{psf}) with those obtained with aperture photometry (f_{aper}) in the left panel of Figure A1. This comparison includes both the ALCS sources and ALMA-undetected *Herschel* sources extracted at $S/N > 5$ in each *Herschel* band. We find that the majority of *Herschel* sources exhibit comparable flux densities obtained through both methods, i.e., $f_{\text{aper}}/f_{\text{psf}} \sim 1$, up to a wavelength of 350 μm (see statistics for each band in the left panel of Figure A1). Because of source blending, a number excess of sources can be found at $f_{\text{aper}}/f_{\text{psf}} > 1$ especially at longer wavelength (i.e., 500 μm). Such an effect is inevitable for the majority of ALCS sources given the large beam size of $35''$ at 500 μm . We then conclude that our flux densities obtained through PSF photometry are not subject to any obvious systematic offset from those by aperture photometry.




Figure A2. Comparisons of *Herschel*/SPIRE flux densities, including both submillimeter sources detected and undetected by the ALCS, measured with XID+ (Hurley et al. 2017) and our iterative extraction procedure (Section 3). Comparison in the SPIRE 250, 350 and 500 μm band are shown in the left, central and right panel, respectively. In each panel, the diagonal solid black line indicates identical flux densities measured with the two procedures, and the shaded gray region indicates the expected 1σ dispersion range of flux densities given the depth of data taken in the “snapshot” mode.

We also compare our flux density measurements of sources in Frontier Fields with R16. R16 reported 263 secure *Herschel* detections within the *HST*/ACS footprints of the six HFF clusters (both central regions and parallel footprints). We cross-match our photometric measurements (f_{This} ; including both ALCS sources and ALMA-undetected *Herschel* sources) with those of R16 (f_{R16}) allowing a maximum angular separation of $1''.5$. The flux density ratios are plotted as a function of *Herschel*-band S/N’s in the right panel of Figure A1. We find a general consistency between the flux densities of these two works (median flux ratio is $f_{\text{R16}}/f_{\text{This}} = 0.98 \pm 0.02$) without any conspicuous systematic offset.

To evaluate the accuracy of SPIRE photometry without PACS information, we also experimented extraction from 250 μm (instead of 100 μm) for the “deep”-mode clusters. The resultant SPIRE flux densities are generally consistent with those extracted using PACS information at the 1σ confidence level.

We also extracted the fluxes of ALCS sources using XID+ (Hurley et al. 2017). XID+ is a prior-based *Herschel*/SPIRE flux extraction software built upon a probabilistic Bayesian framework. The positional prior that we used was the same as described in Section 3, i.e., including both ALCS-detected sources at $S/N_{\text{ALMA}} \geq 4$ and ALMA-undetected *Herschel* sources. Compared with our iterative extraction procedure, XID+ does not distinguish the priority of input sources.

Figure A2 shows the comparisons of SPIRE flux densities measured by both de-blending routines. We observe a general consistency between the flux density estimate using both methods. For ALCS sources in the main sample, the median difference of flux densities estimated with two routines is around 1 mJy in all the three bands, and the standard deviation of flux density differences are well predicted by the joint uncertainty of both measurements. We further analyze ALCS sources whose XID+-derived flux densities are different from those extracted iteratively by a factor of > 2 . The median separation of these sources to their nearest neighbors is found to be $7'' \pm 6''$, significantly smaller than the typical separation of $14'' \pm 10''$ for the full sample. Therefore, we conclude that the large photometric discrepancy among these sources is mainly caused by strong blending effect. With prior ALMA flux density information, iterative extraction procedure has the potential to enhance the photometric accuracy in crowded lensing cluster fields.

B. POSTAGE STAMP IMAGES AND FAR-IR SEDS OF ALCS SOURCES

In the online journal, we show the figure sets of *HST*-*Spitzer*-ALMA postage stamp images and best-fit far-IR SED plots of all 125 sources in the ALCS-*Herschel* joint sample (Table C2). These include:

Fig. Set B1. Postage stamp images and far-IR SEDs of 105 sources in the main ALCS-*Herschel* sample.

Fig. Set B2. Postage stamp images and far-IR SEDs of 20 sources in the secondary ALCS-*Herschel* sample.




Figure B1. *Top:* Postage stamp images of A2537-ID42 ($S/N_{\text{ALMA}} = 6.8$), one of the 105 sources in the main ALCS-*Herschel* joint sample. From the left to the right we show the ALMA Band-6 continuum image, *HST* F814W/F105W/F160W true color image and *Spitzer*/IRAC 3.6/4.5 μm true color image. The galaxy is in the center of each image labeled by green or magenta circles. Other ALCS sources in the field, if exist, are also labeled out with their S/N and ID noted. A two-arcsec scale bar is plotted in the lower right corner of ALMA continuum image. *Bottom:* Far-IR SED of A2537-ID42. *Herschel* and ALMA photometric measurements are shown as open red squares (upper limits are at 3σ). Best-fit SED model derived with MAGPHYS at z_{best} is shown as solid black line. For sources without spectroscopic confirmation (z_{spec}), we also plot the spectral templates of (U)LIRG ($L_{\text{IR}} = 10^{12} L_{\odot}$; Rieke et al. 2009) at $z = 2 - 6$ (normalized to the 1.15 mm flux density) for comparison. The complete figure set (105 images) is available in the online journal.

In addition to these, we also show the ALMA-*HST*-*Spitzer* postage stamps images of ten *Herschel*-faint galaxies that have z_{spec} or catalogued *HST* z_{phot} (Figure B3), and the remaining 17 *Herschel*-faint galaxies without z_{spec} or catalogued *HST* z_{phot} (i.e., optical/near-IR-dark; Figure B4).

C. CATALOGS OF HERSCHEL PHOTOMETRY AND DERIVED PHYSICAL PROPERTIES

Table C1 presents the *Herschel* flux density measurements of (i) 141 secure ALCS sources at $S/N_{\text{nat}} \geq 5$ or $S/N_{\text{tap}} \geq 4.5$ (the main sample), and (ii) 39 tentative ALCS sources at $S/N_{\text{nat}} = 4 - 5$ and $S/N_{\text{tap}} < 4.5$ with cross-matched *HST* or *Spitzer*/IRAC counterparts within $1''$ separation (the secondary sample). Table C2 presents the galaxy properties of 125 ALCS sources that are detected at $> 2\sigma$ in ≥ 1 *Herschel* band (i.e., ALCS-*Herschel* joint sample). Note that BCGs and *Herschel*-faint galaxies are not included in this table. The definition of the samples in these two tables are also visualized in Figure 6.





Figure B2. Same as Figure B1 but for A383-ID24 ($S/N_{\text{ALMA}} = 4.0$), one of the 20 sources in the secondary ALCS-*Herschel* joint sample. The complete figure set (20 images) is available in the online journal.

Table C1. Summary of *Herschel* photometry

#	Name	Coordinates		S/N_{ALMA}	<i>Herschel</i> /PACS		<i>Herschel</i> /SPIRE		
		RA	Dec		100 μm	160 μm	250 μm	350 μm	500 μm
(mJy)									
<i>Main Sample</i> ($S/N_{\text{nat}} \geq 5$ or $S/N_{\text{tap}} \geq 4.5$)									
1	A209-ID38 [↓]	01:31:53.38	-13:37:17.4	5.2	<1.5	<2.3	<6.4	<6.6	<7.4
2	A2163-ID11	16:15:49.06	-06:08:13.9	4.6 [†]	37.7 \pm 5.3	24.1 \pm 5.2	<17.2
3	A2537-ID42	23:08:24.43	-02:11:05.5	6.8	19.5 \pm 0.5	44.0 \pm 0.9	49.7 \pm 1.9	39.8 \pm 2.5	20.2 \pm 2.6
4	A2537-ID49	23:08:17.92	-02:11:12.7	5.2	33.0 \pm 2.5	55.9 \pm 3.4	36.1 \pm 4.5	30.9 \pm 5.5	8.2 \pm 4.3
5	A2537-ID66	23:08:19.68	-02:11:25.3	4.9	15.5 \pm 0.7	20.5 \pm 2.6	20.3 \pm 5.2	<22.9	<7.8
6	A2744-ID07	00:14:19.12	-30:22:42.2	8.0	<0.8	<2.3	15.4 \pm 6.6	18.1 \pm 3.7	7.7 \pm 3.0
7	A2744-ID21	00:14:22.09	-30:22:49.7	6.8	4.4 \pm 1.2	13.0 \pm 2.1	7.1 \pm 2.9	<13.6	<7.7
8	A2744-ID33	00:14:20.40	-30:22:54.5	15.0	0.9 \pm 0.5	2.1 \pm 1.4	5.5 \pm 5.3	5.2 \pm 4.7	15.6 \pm 4.4
9	A2744-ID56	00:14:17.58	-30:23:00.6	11.9	28.3 \pm 0.6	61.7 \pm 1.2	64.3 \pm 1.9	46.6 \pm 2.4	24.0 \pm 2.6
10	A2744-ID81	00:14:19.80	-30:23:07.7	26.7	<1.6	4.3 \pm 1.8	9.8 \pm 5.4	15.8 \pm 4.6	<15.0
11	A2744-ID319	00:14:18.26	-30:24:47.5	14.2	<1.9	7.8 \pm 1.3	11.7 \pm 1.9	10.5 \pm 3.0	9.5 \pm 2.6
12	A3192-ID31	03:58:50.02	-29:56:30.8	32.6	124.9 \pm 4.4	88.4 \pm 4.7	40.3 \pm 4.6
13	A3192-ID40	03:58:51.13	-29:55:10.9	26.0	20.0 \pm 4.4	15.8 \pm 5.1	<20.5

Table C1 *continued*

Herschel-faint galaxies with spectroscopic redshifts:



Herschel-faint galaxies with HST photometric redshifts:




Figure B3. Postage stamp images of ten *Herschel*-faint galaxies with either spectroscopic redshifts (R0032-ID32 and R0600-ID164) or *HST* photometric redshifts (the remaining eight sources). In each category, sources are shown in descending order of ALMA S/N. The layout of these postage stamp images is the same as that in Figure B1. The best available redshift is noted in the lower right corner of *Spitzer*/*IRAC* image.

Table C1 (*continued*)

#	Name	Coordinates		S/N _{ALMA}	<i>Herschel</i> /PACS		<i>Herschel</i> /SPIRE		
		RA	Dec		100 μm	160 μm	250 μm	350 μm	500 μm
					(mJy)	(mJy)	(mJy)	(mJy)	(mJy)
14	A3192-ID131	03:58:47.63	-29:56:33.5	11.5	12.8 \pm 4.0	10.0 \pm 4.7	<13.8
15	A3192-ID138	03:58:57.13	-29:54:53.5	8.1	20.8 \pm 4.5	12.2 \pm 4.4	<15.4
16	A370-ID18	02:39:50.20	-01:35:42.1	12.9	1.2 \pm 0.5	6.1 \pm 1.3	9.8 \pm 6.2	17.9 \pm 9.2	<11.5
17	A370-ID31	02:39:54.43	-01:33:36.5	5.9	34.8 \pm 0.6	49.1 \pm 1.0	32.3 \pm 1.8	17.3 \pm 5.4	<20.9
18	A370-ID103	02:39:58.15	-01:34:24.5	6.8	8.1 \pm 0.5	22.7 \pm 3.2	33.6 \pm 10.1	27.3 \pm 2.8	13.0 \pm 2.2
19	A370-ID110	02:39:56.57	-01:34:26.3	24.9	48.8 \pm 0.6	105.5 \pm 0.9	108.8 \pm 2.8	76.7 \pm 2.8	30.9 \pm 2.2
20	A370-ID146	02:39:51.44	-01:34:41.5	5.4	0.6 \pm 0.5	5.1 \pm 1.4	9.5 \pm 4.9	8.3 \pm 3.5	<5.8
21	A383-ID40 [•]	02:48:03.39	-03:31:44.8	34.2	14.1 \pm 0.6	24.0 \pm 1.8	16.6 \pm 2.8	7.3 \pm 2.7	<7.7

Table C1 *continued*

Herschel-faint galaxies without optical/near-IR counterparts/redshifts:


Figure B4. Same as Figure B3 but for 17 *Herschel*-faint sources without optical/near-IR counterparts or catalogued redshifts.

Table C1 (continued)

#	Name	Coordinates		S/N _{ALMA}	<i>Herschel</i> /PACS		<i>Herschel</i> /SPIRE		
		RA	Dec		100 μm	160 μm	250 μm	350 μm	500 μm
					(mJy)	(mJy)	(mJy)	(mJy)	(mJy)
22	ACT0102-ID11 [↓]	01:02:59.47	-49:17:25.4	5.3	<13.6	<15.0	<20.5
23	ACT0102-ID22	01:03:03.04	-49:17:11.2	7.9	6.1 \pm 5.3	10.2 \pm 4.8	12.2 \pm 5.6
24	ACT0102-ID50	01:02:59.40	-49:16:55.6	5.8	18.8 \pm 4.1	12.2 \pm 4.4	11.0 \pm 5.0
25	ACT0102-ID52 [↓]	01:03:01.47	-49:16:56.6	5.8	6.3 \pm 4.1	<15.3	<20.5
26	ACT0102-ID118	01:02:51.09	-49:15:38.9	34.3	9.0 \pm 5.2	14.6 \pm 7.3	9.0 \pm 5.7
27	ACT0102-ID160	01:03:00.20	-49:16:03.7	4.9 [†]	14.3 \pm 3.6	<12.0	<20.4
28	ACT0102-ID215	01:02:54.92	-49:15:14.7	42.4	<16.1	<22.6	11.4 \pm 3.8
29	ACT0102-ID223	01:02:49.24	-49:15:08.9	7.9	11.7 \pm 3.9	14.0 \pm 4.7	10.6 \pm 3.5
30	ACT0102-ID224	01:02:55.68	-49:15:09.1	91.8	18.4 \pm 5.4	28.7 \pm 7.5	28.2 \pm 9.4
31	ACT0102-ID241	01:02:58.13	-49:14:56.2	5.4	6.2 \pm 3.4	12.0 \pm 4.3	12.2 \pm 4.3
32	ACT0102-ID251 [↓]	01:02:55.07	-49:14:19.2	5.0	<14.8	<15.5	<20.1
33	ACT0102-ID276 [↓]	01:02:49.29	-49:14:38.2	9.9	<14.6	6.3 \pm 4.0	<20.1
34	ACT0102-ID294	01:02:49.37	-49:15:05.3	19.1 [†]	36.0 \pm 12.0	42.9 \pm 14.3	32.6 \pm 10.9
35	AS1063-ID15	22:48:46.58	-44:30:46.9	7.7 [†]	6.3 \pm 1.0	10.4 \pm 0.8	5.3 \pm 3.6	<25.4	<27.3
36	AS1063-ID17	22:48:47.29	-44:30:48.2	15.5	24.0 \pm 0.5	61.6 \pm 0.8	62.7 \pm 3.6	48.4 \pm 8.5	22.5 \pm 9.2
37	AS1063-ID147	22:48:41.82	-44:31:57.5	6.0 [†]	66.9 \pm 0.9	98.4 \pm 0.9	67.8 \pm 3.3	30.8 \pm 5.0	7.3 \pm 3.2

Table C1 continued

Table C1 (continued)

#	Name	Coordinates		S/N _{ALMA}	<i>Herschel</i> /PACS		<i>Herschel</i> /SPIRE		
		RA	Dec		100 μm	160 μm	250 μm	350 μm	500 μm
					(mJy)	(mJy)	(mJy)	(mJy)	(mJy)
38	AS1063-ID222	22:48:49.06	-44:32:25.4	85.7	<1.4	3.0 \pm 0.9	10.6 \pm 2.6	21.8 \pm 3.2	23.3 \pm 2.6
39	M0035-ID41	00:35:20.46	-20:15:35.2	7.2	8.3 \pm 3.6	<13.4	<13.7
40	M0035-ID68 \downarrow	00:35:30.76	-20:15:54.9	11.1	<14.3	<15.1	<14.5
41	M0035-ID94	00:35:27.24	-20:16:17.4	4.6 \uparrow	23.6 \pm 3.4	<12.8	<16.7
42	M0159-ID05	01:59:51.69	-08:49:10.1	16.0	24.0 \pm 4.0	20.4 \pm 4.1	10.2 \pm 5.4
43	M0159-ID24	01:59:47.33	-08:49:33.2	6.3	24.1 \pm 5.2	<15.7	9.0 \pm 6.6
44	M0159-ID46 \bullet	01:59:49.35	-08:49:59.0	32.8	11.7 \pm 4.4	<15.6	14.2 \pm 5.2
45	M0159-ID61	01:59:51.56	-08:50:13.3	5.8	14.1 \pm 4.7	<15.6	<17.8
46	M0257-ID13	02:57:11.25	-23:25:43.6	12.2 \uparrow	11.7 \pm 4.4	13.4 \pm 4.5	15.1 \pm 4.7
47	M0329-ID11	03:29:41.76	-02:10:56.8	11.1	1.7 \pm 0.5	1.6 \pm 0.8	<9.2	<8.6	<9.7
48	M0416-ID51	04:16:06.97	-24:04:00.0	7.1	1.8 \pm 0.5	4.0 \pm 0.7	<9.0	<7.8	<9.4
49	M0416-ID79	04:16:11.67	-24:04:19.5	5.3	1.4 \pm 0.5	1.1 \pm 0.9	6.3 \pm 2.7	<7.8	<9.4
50	M0416-ID117	04:16:10.80	-24:04:47.6	15.7	3.9 \pm 0.5	10.4 \pm 0.8	12.4 \pm 2.2	9.3 \pm 1.9	9.8 \pm 1.7
51	M0416-ID160	04:16:08.83	-24:05:22.5	6.0	3.5 \pm 0.5	5.8 \pm 0.8	6.0 \pm 2.7	3.5 \pm 2.7	<9.4
52	M0417-ID46	04:17:33.32	-11:55:03.3	48.0	45.4 \pm 5.8	39.0 \pm 6.2	30.7 \pm 5.1
53	M0417-ID49	04:17:40.15	-11:55:00.8	36.1	43.4 \pm 4.7	60.6 \pm 7.7	35.8 \pm 5.6
54	M0417-ID58	04:17:35.65	-11:54:53.3	30.4	11.1 \pm 7.3	14.8 \pm 5.3	11.1 \pm 5.7
55	M0417-ID121	04:17:37.02	-11:54:19.8	20.7	10.9 \pm 6.3	6.9 \pm 4.7	17.6 \pm 5.2
56	M0417-ID204	04:17:32.23	-11:52:39.1	4.5 \uparrow	13.8 \pm 4.4	<18.2	<34.2
57	M0417-ID218	04:17:32.01	-11:52:57.4	14.5	37.7 \pm 5.1	44.0 \pm 8.9	14.9 \pm 7.4
58	M0417-ID221	04:17:32.48	-11:53:02.8	7.7	20.9 \pm 1.8	14.3 \pm 2.9	7.3 \pm 3.7
59	M0417-ID223	04:17:32.55	-11:53:05.0	6.0	38.6 \pm 3.2	26.4 \pm 5.3	13.6 \pm 6.8
60	M0429-ID04 \downarrow	04:29:35.01	-02:52:39.3	8.4	<0.7	<2.3	<9.7	<9.0	5.7 \pm 4.1
61	M0429-ID19 \bullet	04:29:36.03	-02:53:06.8	27.6	31.2 \pm 0.6	44.4 \pm 1.1	25.5 \pm 3.9	7.3 \pm 5.1	<10.4
62	M0429-ID27	04:29:37.10	-02:53:35.2	6.9	4.2 \pm 0.5	8.8 \pm 0.9	12.5 \pm 2.6	5.2 \pm 3.5	6.3 \pm 3.4
63	M0553-ID17	05:53:15.40	-33:41:35.4	8.5	19.5 \pm 4.9	12.4 \pm 5.7	<16.4
64	M0553-ID18 \downarrow	05:53:16.75	-33:41:35.6	4.5 \uparrow	<14.6	5.7 \pm 5.7	<18.7
65	M0553-ID58 \downarrow	05:53:30.18	-33:41:52.1	7.5	<13.3	<15.3	<18.9
66	M0553-ID61	05:53:21.49	-33:41:54.7	10.7	17.9 \pm 4.5	15.5 \pm 4.6	<22.6
67	M0553-ID67 \downarrow	05:53:29.30	-33:41:56.3	10.0	<59.2	27.3 \pm 22.9	<18.9
68	M0553-ID133	05:53:27.79	-33:42:16.0	39.0	119.3 \pm 3.1	83.7 \pm 4.2	43.5 \pm 6.1
69	M0553-ID190	05:53:27.85	-33:42:30.6	67.5	179.1 \pm 4.7	125.6 \pm 6.3	65.2 \pm 9.2
70	M0553-ID200	05:53:33.40	-33:42:31.7	5.1 \uparrow	20.6 \pm 7.5	<18.8	<18.9
71	M0553-ID249	05:53:27.63	-33:42:43.9	44.0	126.1 \pm 3.3	88.4 \pm 4.4	45.9 \pm 6.5
72	M0553-ID275	05:53:26.05	-33:42:51.6	6.6	47.0 \pm 4.2	38.1 \pm 6.0	9.6 \pm 6.0
73	M0553-ID303	05:53:31.98	-33:43:00.5	8.2 \uparrow	79.2 \pm 4.5	48.7 \pm 4.7	12.6 \pm 4.3
74	M0553-ID355	05:53:23.28	-33:43:16.0	12.4	12.4 \pm 3.8	24.1 \pm 4.5	17.3 \pm 4.3
75	M0553-ID375	05:53:13.29	-33:43:21.3	8.3	18.9 \pm 4.7	9.9 \pm 4.3	<16.2
76	M0553-ID398	05:53:29.87	-33:43:26.0	13.5	9.2 \pm 4.3	10.9 \pm 4.4	5.0 \pm 4.3
77	M1115-ID02	11:15:50.69	01:30:35.5	11.3	24.8 \pm 3.0	38.2 \pm 1.8	72.8 \pm 6.6	72.0 \pm 4.6	49.2 \pm 4.6
78	M1115-ID04	11:15:52.03	01:30:28.2	7.6	16.5 \pm 0.8	29.1 \pm 0.9	35.5 \pm 3.2	27.7 \pm 3.9	12.8 \pm 3.2
79	M1115-ID34	11:15:54.14	01:29:56.3	16.1	0.7 \pm 0.5	2.5 \pm 0.8	7.7 \pm 3.2	<8.3	<9.2
80	M1115-ID36 \bullet	11:15:51.90	01:29:54.9	7.5	2.8 \pm 0.5	3.4 \pm 0.8	<10.2	<8.4	<7.6
81	M1149-ID77	11:49:36.10	22:24:24.5	5.5 \uparrow	2.2 \pm 0.9	9.4 \pm 3.2	15.4 \pm 2.1	11.5 \pm 2.2	7.3 \pm 3.5
82	M1149-ID229	11:49:34.04	22:23:16.7	4.7 \uparrow	2.3 \pm 0.8	4.9 \pm 1.5	<15.3	<6.4	<7.3
83	M1206-ID27	12:06:11.26	-08:47:43.6	7.7 \uparrow	14.2 \pm 0.6	33.3 \pm 6.0	26.9 \pm 11.1	42.6 \pm 6.0	13.0 \pm 4.7
84	M1206-ID54 \downarrow	12:06:15.94	-08:47:59.4	7.4 \uparrow	0.7 \pm 0.5	<2.4	<11.4	<9.4	3.6 \pm 2.6
85	M1206-ID55	12:06:10.74	-08:48:00.6	9.2 \uparrow	24.8 \pm 0.3	44.9 \pm 1.2	42.2 \pm 14.1	26.4 \pm 1.6	11.1 \pm 3.5
86	M1206-ID58 \bullet	12:06:12.15	-08:48:03.5	6.5 \uparrow	<0.7	<2.4	<11.1	<9.8	<10.7
87	M1206-ID60	12:06:10.75	-08:48:05.3	10.4 \uparrow	29.7 \pm 0.4	53.7 \pm 1.5	50.5 \pm 16.8	31.6 \pm 1.9	13.3 \pm 4.1
88	M1206-ID61	12:06:10.80	-08:48:08.9	8.0 \uparrow	40.6 \pm 0.6	73.4 \pm 2.0	69.1 \pm 23.0	43.3 \pm 2.6	18.1 \pm 5.6

Table C1 continued

Table C1 (continued)

#	Name	Coordinates		S/N _{ALMA}	Herschel/PACS		Herschel/SPIRE		
		RA	Dec		100 μ m	160 μ m	250 μ m	350 μ m	500 μ m
					(mJy)	(mJy)	(mJy)	(mJy)	(mJy)
89	M1311-ID27	13:11:01.32	-03:10:33.6	9.2 [†]	5.8±0.5	19.0±1.5	28.5±3.0	23.3±2.4	13.6±2.6
90	M1311-ID33	13:11:00.14	-03:10:42.4	8.3	1.4±0.5	13.8±1.2	9.4±3.4	10.4±3.0	6.8±3.9
91	M1423-ID38	14:23:47.01	24:04:54.2	5.5	5.3±0.6	9.4±1.2	14.4±3.4	8.7±3.0	<7.2
92	M1423-ID50 [•]	14:23:47.88	24:04:42.4	23.1	11.3±0.6	19.7±1.3	22.5±3.4	13.8±3.1	14.1±2.3
93	M1931-ID41 [•]	19:31:49.63	-26:34:33.2	83.3	233.6±1.4	243.3±1.3	121.0±2.8	44.7±3.0	18.5±3.0
94	M1931-ID47 [↓]	19:31:52.08	-26:34:36.4	7.7	<0.7	<2.9	<9.2	<8.5	<6.9
95	M1931-ID55	19:31:48.10	-26:34:43.4	11.6 [†]	7.6±0.6	13.2±1.0	9.3±2.8	12.5±3.0	4.4±3.0
96	M1931-ID61	19:31:49.28	-26:34:50.9	10.5 [†]	0.6±0.6	2.2±1.2	5.3±2.8	7.2±3.3	6.4±3.4
97	M2129-ID24 [↓]	21:29:25.28	-07:41:07.7	5.2 [†]	<0.7	<2.7	<9.7	<9.2	<8.9
98	M2129-ID46	21:29:29.45	-07:41:31.2	6.2	4.8±0.5	11.1±0.8	12.9±2.1	8.6±2.9	<9.1
99	P171-ID69	03:12:52.76	08:22:45.0	19.7	11.6±5.4	9.8±7.7	<17.4
100	P171-ID162 [↓]	03:12:58.70	08:23:29.2	9.0	<16.6	<17.4	<23.6
101	P171-ID177	03:12:55.48	08:23:44.0	7.3 [†]	11.9±4.8	<21.2	<21.6
102	R0032-ID32 [↓]	00:32:09.77	18:06:24.5	9.6	9.0±7.5	14.3±8.7	<23.3
103	R0032-ID53	00:32:08.22	18:06:40.3	11.3	21.8±1.9	20.4±2.8	12.3±3.2
104	R0032-ID55	00:32:07.84	18:06:46.1	18.9 [†]	41.6±3.6	38.9±5.4	23.5±6.1
105	R0032-ID57	00:32:07.71	18:06:48.8	7.5	7.4±0.6	7.0±1.0	4.2±1.1
106	R0032-ID58	00:32:07.56	18:06:51.4	12.2	22.7±2.0	21.2±2.9	12.9±3.3
107	R0032-ID81 [↓]	00:32:12.86	18:07:02.0	5.2	<17.6	<19.6	<21.3
108	R0032-ID127	00:32:13.16	18:08:14.3	53.6	13.2±5.3	30.2±7.4	<38.9
109	R0032-ID131	00:32:12.19	18:08:13.3	46.8	12.7±5.3	15.6±7.4	32.7±11.4
110	R0032-ID198	00:32:08.54	18:08:26.7	18.3	33.8±7.7	24.9±8.4	<23.0
111	R0032-ID208 [↓]	00:32:13.97	18:08:33.7	11.2	<15.6	<21.7	10.6±8.8
112	R0032-ID220	00:32:08.61	18:09:08.9	6.8 [†]	33.1±9.5	18.5±18.3	<23.1
113	R0032-ID238	00:32:07.83	18:09:33.8	6.6	8.6±3.7	7.2±6.5	19.4±4.8
114	R0032-ID250 [↓]	00:32:08.76	18:09:20.8	8.0	<17.1	11.4±11.3	<23.1
115	R0032-ID276	00:32:08.61	18:08:59.2	14.9	13.0±10.3	20.0±17.1	36.7±9.1
116	R0032-ID281 [↓]	00:32:12.25	18:08:49.0	9.7	<15.7	<21.8	<21.8
117	R0032-ID287	00:32:07.70	18:08:52.9	11.4	46.3±7.8	30.8±7.9	<36.2
118	R0032-ID304 [↓]	00:32:08.95	18:08:41.8	11.2	13.2±7.7	<27.0	<22.7
119	R0600-ID12	06:00:06.32	-20:10:04.6	5.3	19.9±5.8	22.0±5.6	<19.7
120	R0600-ID13	06:00:05.05	-20:09:51.1	9.9	41.4±3.9	27.3±5.7	9.5±6.6
121	R0600-ID67 [↓]	06:00:05.01	-20:09:24.3	38.1	<15.5	<16.3	<18.0
122	R0600-ID111	06:00:08.91	-20:08:55.9	5.2	15.2±4.4	9.5±4.4	<14.7
123	R0600-ID164 [↓]	06:00:09.10	-20:08:26.5	4.8 [†]	<15.1	<16.1	<18.0
124	R0949-ID10	09:49:52.13	17:05:47.2	61.9	8.5±4.5	14.1±5.3	24.7±5.9
125	R0949-ID19 [↓]	09:49:53.85	17:05:57.7	13.1	<16.2	<15.9	<16.5
126	R0949-ID122	09:49:51.16	17:08:12.8	5.3	12.8±4.6	5.6±5.4	17.4±5.6
127	R0949-ID124	09:49:54.62	17:08:10.5	12.1	19.0±5.4	9.9±5.0	11.3±7.3
128	R1347-ID41	13:47:31.11	-11:44:38.8	7.2	6.5±0.4	17.3±0.7	13.2±1.6	5.5±1.8	6.1±4.4
129	R1347-ID51 [↓]	13:47:33.74	-11:44:48.9	5.0	<0.6	<1.9	<5.6	<5.5	<13.2
130	R1347-ID75 [•]	13:47:30.63	-11:45:09.5	57.0	5.8±0.4	6.4±0.7	<5.6	<5.5	9.4±4.3
131	R1347-ID145	13:47:27.65	-11:45:51.1	22.0	12.0±0.5	29.6±0.7	42.8±2.6	37.1±7.9	21.6±1.1
132	R1347-ID148	13:47:27.85	-11:45:55.8	21.4 [†]	16.5±0.7	40.7±0.9	58.8±3.5	51.0±10.8	29.7±1.4
133	R1347-ID166	13:47:28.00	-11:46:12.2	5.0	<1.3	1.0±0.8	10.3±2.3	15.5±5.5	16.1±2.6
134	R2129-ID20 [•]	21:29:39.96	00:05:21.0	17.8	1.1±0.6	<2.4	6.2±2.4	5.9±3.8	<12.3
135	R2129-ID37	21:29:38.24	00:04:52.6	5.8	1.1±1.0	9.3±1.1	9.8±4.0	10.3±7.5	8.7±4.1
136	R2211-ID19	22:11:47.70	-03:51:03.6	6.5	19.5±6.4	8.9±5.0	<18.7
137	R2211-ID35	22:11:42.91	-03:50:41.7	17.8 [†]	131.1±5.0	94.8±6.3	46.0±6.9
138	R2211-ID171	22:11:42.20	-03:50:23.8	10.1	24.4±4.7	11.1±6.3	<20.8
139	SM0723-ID61 [↓]	07:23:03.86	-73:27:06.2	5.8	<16.2	<17.4	<20.2

Table C1 continued

Table C1 (*continued*)

#	Name	Coordinates		S/N _{ALMA}	<i>Herschel</i> /PACS		<i>Herschel</i> /SPIRE		
		RA	Dec		100 μm	160 μm	250 μm	350 μm	500 μm
					(mJy)	(mJy)	(mJy)	(mJy)	(mJy)
140	SM0723-ID93 [↓]	07:23:03.15	-73:27:21.0	4.7 [†]	<16.2	<17.4	<20.2
141	SM0723-ID124 [↓]	07:23:25.08	-73:27:38.9	5.7	<15.7	<17.1	<20.2
<i>Secondary Sample</i> (S/N _{nat} = 4 - 5 and S/N _{tap} < 4.5)									
142	A2537-ID06	23:08:20.81	-02:10:34.6	4.2	<0.8	<2.4	6.8±2.8	6.3±2.5	4.9±2.2
143	A2537-ID24	23:08:21.17	-02:10:54.5	4.5	18.4±0.5	30.9±0.8	24.7±1.5	18.0±2.5	9.9±2.2
144	A2744-ID17	00:14:19.51	-30:22:48.6	4.4	<0.8	<2.3	<6.6	<6.9	<7.7
145	A2744-ID47	00:14:17.29	-30:22:58.6	4.7	3.4±0.6	2.8±1.0	3.7±1.8	<5.9	<7.7
146	A2744-ID176	00:14:17.36	-30:23:45.4	4.2 [†]	0.7±0.5	<2.3	<6.8	<7.0	2.3±1.5
147	A2744-ID178	00:14:24.12	-30:23:46.0	4.7	4.9±0.5	9.3±0.9	11.8±1.5	3.6±1.8	<7.5
148	A2744-ID227	00:14:16.56	-30:24:10.0	4.6	7.6±0.5	9.2±0.8	<6.8	<7.1	1.8±1.5
149	A3192-ID83	03:58:53.92	-29:55:55.4	4.2	<14.4	<13.2	<18.4
150	A3192-ID154	03:58:56.69	-29:54:20.1	4.2	<11.7	<12.8	<13.1
151	A370-ID27	02:39:53.54	-01:33:35.3	4.3	1.4±0.5	2.3±0.8	5.4±1.4	3.0±1.6	2.0±1.7
152	A370-ID172	02:39:47.38	-01:34:53.1	4.3	<0.7	<2.3	<6.7	<6.9	<8.1
153	A383-ID24	02:48:02.83	-03:31:26.5	4.0	17.5±0.6	27.6±1.0	21.2±1.8	8.1±1.5	<7.7
154	A383-ID50	02:48:02.82	-03:31:57.5	4.2	5.9±0.5	5.1±0.8	3.0±1.7	<6.6	<7.7
155	ACT0102-ID128	01:02:50.47	-49:15:41.7	4.5	<14.6	<14.8	<20.2
156	AS295-ID09	02:45:36.63	-53:03:27.4	4.3	<15.8	<13.6	8.2±6.1
157	AS295-ID269	02:45:22.33	-53:00:48.9	4.3	12.3±5.3	7.5±5.0	<15.9
158	M0035-ID33	00:35:25.50	-20:15:29.5	4.1	<12.7	<15.1	<16.7
159	M0416-ID120	04:16:10.51	-24:04:49.0	4.4	<1.5	<2.3	<6.5	<7.8	<9.4
160	M0416-ID138	04:16:10.52	-24:05:04.8	4.9	3.2±0.5	2.7±0.8	6.0±2.2	4.1±2.0	<9.4
161	M0416-ID156	04:16:08.79	-24:05:17.6	4.5	3.9±0.5	0.9±0.7	<9.3	<7.9	<9.4
162	M1115-ID33	11:15:54.34	01:29:57.1	4.3	0.8±0.5	1.9±0.8	<10.2	<8.4	<9.2
163	M1149-ID27	11:49:40.00	22:24:57.1	4.6	<0.7	4.1±0.8	6.2±1.4	3.6±1.6	4.1±1.8
164	M1149-ID95	11:49:41.05	22:24:16.9	4.1	1.3±0.5	3.5±0.8	4.6±1.8	7.6±1.8	4.1±1.4
165	M1206-ID38	12:06:08.25	-08:47:51.1	4.4	3.2±0.5	6.9±0.9	<11.1	9.0±2.4	4.5±2.2
166	M1206-ID84	12:06:13.13	-08:48:27.2	4.4	14.9±0.5	20.2±1.5	9.8±4.1	<12.6	<10.2
167	M1423-ID52	14:23:47.57	24:04:37.6	4.4	3.9±0.5	<2.7	<6.5	<6.5	<7.2
168	M1423-ID76	14:23:48.03	24:04:12.2	5.0	3.3±0.6	3.7±0.8	3.8±1.9	<6.3	<7.3
169	M1931-ID42 [•]	19:31:49.48	-26:34:31.3	4.2	13.5±1.3	4.3±1.2	<9.1	<8.5	<9.3
170	M1931-ID69	19:31:47.63	-26:35:02.2	4.3	<0.7	4.4±0.7	<9.2	<8.5	<6.2
171	M2129-ID62	21:29:29.63	-07:41:37.7	4.3	4.5±0.5	6.2±0.8	<5.9	<10.4	<9.1
172	P171-ID161	03:12:55.35	08:23:43.7	4.3	<16.6	<21.2	<21.5
173	R0032-ID63	00:32:11.39	18:06:52.2	4.5	<18.1	<19.6	<22.1
174	R0032-ID162 [•]	00:32:11.53	18:07:52.3	4.8	8.2±4.5	<21.9	<21.4
175	R0032-ID245	00:32:09.46	18:09:28.9	4.6	8.2±4.3	<20.9	<12.9
176	R0949-ID14	09:49:52.36	17:05:46.6	4.7	<16.7	<15.9	<14.0
177	R0949-ID113	09:49:53.87	17:08:17.0	4.0	<15.4	<15.7	<14.9
178	R0949-ID119	09:49:52.12	17:08:09.7	4.2	18.8±4.1	<13.8	<18.9
179	R2211-ID164	22:11:44.72	-03:50:13.7	4.6	<16.3	<16.3	<18.7
180	SM0723-ID98	07:23:02.63	-73:27:24.9	4.1	<16.2	<17.4	<20.2

NOTE—Here we only present *Herschel* photometry for all secure ALMA sources (141 sources in total; main sample) and tentative ALMA sources with cross-matched *HST* or *Spitzer* counterparts (39 sources in total; secondary sample).

[↓] *Herschel*-faint sources with low significance of detection (S/N<2) in all *Herschel* bands but secure detection (S/N≥5) at 1.15 mm (see Section 4.3).

[•] BCGs in corresponding clusters. M1931-ID42 was identified as the extended tail of the BCG M1931-ID41 (Fogarty et al. 2019).

[†] Measured from 2''-tapered ALMA maps instead of native-resolution maps due to a higher significance of detection.

Table C2. Summary of MAGPHYS SED fitting results

#	Name	z_{best}^a	z_{FIR}^b	Ref. ^a	μ^c	Model ^d	$\log(L_{\text{IR}})^e$	$\log(\text{SFR})^e$	$\log(M_{\text{dust}})^e$	T_{dust}^f
							($\mu^{-1}L_{\odot}$)	($\mu^{-1}M_{\odot} \text{ yr}^{-1}$)	($\mu^{-1}M_{\odot}$)	(K)
Main Sample ($S/N_{\text{ALMA}} \geq 5$)										
1	A2163-ID11	$1.23^{+0.68}_{-0.10}$	$1.08^{+0.44}_{-0.43}$	HST	3.2	(1)	12.48 ± 0.33	2.31 ± 0.44	8.39 ± 0.11	35.5 ± 7.7
2	A2537-ID42	$1.13^{+0.07}_{-0.07}$	$1.16^{+0.48}_{-0.54}$	HST	3.6	(1)	12.43 ± 0.06	2.27 ± 0.15	8.77 ± 0.06	30.2 ± 1.1
3	A2537-ID49	$1.07^{+0.43}_{-0.38}$...	FIR	2.9	(1)	12.44 ± 0.43	2.33 ± 0.49	8.36 ± 0.10	35.7 ± 7.2
4	A2537-ID66	$3.09^{+0.06}_{-0.10}$	$1.16^{+0.40}_{-0.43}$	HST	15.5	(1)	13.25 ± 0.04	3.39 ± 0.21	7.72 ± 0.16	63.7 ± 9.8
5	A2744-ID07	2.41	$1.61^{+0.59}_{-0.53}$	(1)	1.9	(2)	12.64 ± 0.16	2.52 ± 0.23	8.30 ± 0.09	41.4 ± 4.8
6	A2744-ID21	2.64	$1.78^{+0.68}_{-0.60}$	(1)	1.9	(2)	12.67 ± 0.04	2.61 ± 0.11	8.18 ± 0.15	33.9 ± 4.0
7	A2744-ID33	3.06	$3.05^{+1.08}_{-0.92}$	(1)	2.2	(2)	12.43 ± 0.11	2.32 ± 0.16	8.70 ± 0.10	31.4 ± 3.4
8	A2744-ID56	1.50	$0.99^{+0.36}_{-0.41}$	(1)	3.2	(2)	12.83 ± 0.02	2.61 ± 0.18	8.61 ± 0.04	37.5 ± 0.5
9	A2744-ID81	2.90	$2.46^{+0.82}_{-0.76}$	(2)	3.3	(2)	12.56 ± 0.10	2.45 ± 0.15	8.57 ± 0.07	34.9 ± 3.0
10	A2744-ID319	2.58	$3.45^{+1.14}_{-0.70}$	(1)	1.9	(2)	12.64 ± 0.06	2.57 ± 0.10	9.29 ± 0.07	26.0 ± 0.9
11	A3192-ID31	$1.22^{+0.06}_{-0.05}$	$1.20^{+0.39}_{-0.47}$	HST	2.4	(1)	12.90 ± 0.12	2.68 ± 0.31	9.02 ± 0.06	33.1 ± 1.1
12	A3192-ID40	$3.01^{+0.96}_{-0.87}$...	FIR	2.7	(1)	12.85 ± 0.25	2.77 ± 0.28	9.04 ± 0.20	31.6 ± 7.4
13	A3192-ID131	$2.35^{+0.80}_{-0.73}$...	FIR	2.9	(1)	12.41 ± 0.33	2.30 ± 0.40	8.58 ± 0.22	31.9 ± 7.7
14	A3192-ID138	$0.92^{+0.39}_{-0.07}$	$1.54^{+0.53}_{-0.50}$	HST	3.0	(1)	11.80 ± 0.34	1.64 ± 0.43	8.53 ± 0.15	24.5 ± 3.7
15	A370-ID18	$2.48^{+0.29}_{-0.15}$	$2.51^{+0.80}_{-0.82}$	HST	2.7	(2)	12.49 ± 0.13	2.36 ± 0.18	8.64 ± 0.11	27.5 ± 4.6
16	A370-ID31	0.38	$1.32^{+0.42}_{-0.52}$	(3)	1.0	(-1)	11.38 ± 0.09	1.31 ± 0.16	8.13 ± 0.09	28.2 ± 0.2
17	A370-ID103	$1.22^{+0.02}_{-0.04}$	$1.51^{+0.55}_{-0.57}$	HST	1.0	(-1)	12.23 ± 0.07	2.03 ± 0.18	8.73 ± 0.08	28.5 ± 1.0
18	A370-ID110	1.06	$1.07^{+0.40}_{-0.46}$	(4)	2.3	(2)	12.73 ± 0.04	2.50 ± 0.18	8.94 ± 0.04	31.1 ± 0.2
19	A370-ID146	1.07	$1.74^{+0.53}_{-0.54}$	(5)	8.9	(2)	11.34 ± 0.05	1.02 ± 0.25	8.25 ± 0.12	26.1 ± 1.7
20	ACT0102-ID22	$2.74^{+1.10}_{-0.86}$...	FIR	1.9	(1)	12.41 ± 0.28	2.30 ± 0.36	8.61 ± 0.17	32.0 ± 9.1
21	ACT0102-ID50	$2.24^{+0.16}_{-0.12}$	$1.54^{+0.53}_{-0.50}$	HST	2.3	(1)	12.58 ± 0.16	2.47 ± 0.23	8.23 ± 0.09	41.1 ± 4.2
22	ACT0102-ID118	4.32	$3.76^{+1.41}_{-1.12}$	(6)	5.2	(1)	12.89 ± 0.17	2.81 ± 0.20	8.89 ± 0.13	34.8 ± 4.5
23	ACT0102-ID160	$2.20^{+0.18}_{-0.15}$	$1.82^{+0.65}_{-0.59}$	HST	2.5	(1)	12.43 ± 0.17	2.31 ± 0.25	8.31 ± 0.11	33.9 ± 3.8
24	ACT0102-ID215	4.32	$4.26^{+1.48}_{-1.27}$	(6)	8.8	(1)	12.83 ± 0.19	2.76 ± 0.23	9.03 ± 0.15	30.1 ± 3.6
25	ACT0102-ID223	$1.82^{+0.09}_{-0.11}$	$2.14^{+0.68}_{-0.70}$	HST	2.0	(1)	12.21 ± 0.17	2.08 ± 0.27	8.65 ± 0.11	28.4 ± 2.4
26	ACT0102-ID224	4.32	$3.90^{+1.38}_{-1.10}$	(6)	9.2	(1)	13.30 ± 0.09	3.22 ± 0.12	9.37 ± 0.08	33.4 ± 2.0
27	ACT0102-ID241	$2.46^{+0.29}_{-0.21}$	$2.50^{+0.91}_{-0.78}$	HST	1.9	(1)	12.32 ± 0.17	2.21 ± 0.24	8.59 ± 0.15	30.6 ± 3.6
28	ACT0102-ID294	$2.14^{+0.68}_{-0.70}$...	FIR	2.4	(1)	12.84 ± 0.33	2.72 ± 0.41	9.04 ± 0.17	31.8 ± 7.3
29	AS1063-ID15	1.43	$2.24^{+0.50}_{-0.34}$	(5)	2.1	(2)	12.17 ± 0.06	2.08 ± 0.10	9.02 ± 0.05	26.4 ± 0.6
30	AS1063-ID17	1.44	$1.12^{+0.39}_{-0.44}$	(5)	2.4	(2)	12.75 ± 0.03	2.52 ± 0.19	8.61 ± 0.04	37.1 ± 0.6
31	AS1063-ID147	0.61	$0.78^{+0.25}_{-0.28}$	(7)	5.7	(2)	12.12 ± 0.05	1.92 ± 0.23	8.41 ± 0.07	32.9 ± 0.2
32	AS1063-ID222	$2.48^{+0.14}_{-0.08}$	$4.30^{+1.35}_{-1.28}$	HST	1.8	(2)	12.57 ± 0.07	2.39 ± 0.16	9.73 ± 0.06	21.6 ± 0.9
33	M0035-ID41	$2.67^{+1.28}_{-0.86}$...	FIR	2.1	(1)	12.37 ± 0.33	2.25 ± 0.40	8.49 ± 0.24	29.9 ± 9.6
34	M0035-ID94	0.36	$1.06^{+0.41}_{-0.41}$	(8)	1.0	(-1)	11.17 ± 0.22	1.00 ± 0.39	8.01 ± 0.16	21.9 ± 2.4
35	M0159-ID05	$1.81^{+0.09}_{-0.39}$	$2.05^{+0.63}_{-0.67}$	HST	2.4	(1)	12.46 ± 0.18	2.35 ± 0.26	8.81 ± 0.10	29.5 ± 2.8
36	M0159-ID24	$1.48^{+0.51}_{-0.50}$...	FIR	0.7	(1)	12.36 ± 0.29	2.16 ± 0.40	8.40 ± 0.14	33.2 ± 8.7
37	M0159-ID61	$1.08^{+0.06}_{-0.05}$	$1.61^{+0.70}_{-0.55}$	HST	2.9	(1)	11.84 ± 0.22	1.67 ± 0.34	8.33 ± 0.16	26.3 ± 3.5
38	M0257-ID13	$1.96^{+0.44}_{-0.41}$	$2.71^{+0.94}_{-0.82}$	HST	1.1	(1)	12.27 ± 0.25	2.14 ± 0.36	9.05 ± 0.13	24.9 ± 3.9
39	M0329-ID11	$3.08^{+0.03}_{-0.06}$	$2.26^{+0.88}_{-0.56}$	HST	2.2	(2)	12.38 ± 0.12	2.29 ± 0.14	8.63 ± 0.16	...
40	M0416-ID51	1.96	$2.06^{+0.67}_{-0.77}$	(5)	2.2	(2)	12.04 ± 0.05	1.93 ± 0.10	8.30 ± 0.11	32.6 ± 1.3
41	M0416-ID79	$2.20^{+0.03}_{-0.02}$	$2.10^{+0.80}_{-0.72}$	HST	1.9	(2)	11.90 ± 0.09	1.79 ± 0.15	8.13 ± 0.17	31.9 ± 2.9
42	M0416-ID117	2.09	$2.71^{+0.60}_{-0.55}$	(5)	1.8	(2)	12.44 ± 0.04	2.39 ± 0.09	8.76 ± 0.09	34.6 ± 0.7
43	M0416-ID160	$1.89^{+0.24}_{-0.03}$	$1.58^{+0.54}_{-0.47}$	HST	1.9	(2)	12.14 ± 0.10	2.06 ± 0.14	8.00 ± 0.23	37.4 ± 2.9
44	M0417-ID46	3.65	$2.34^{+0.72}_{-0.66}$	(9)	5.2	(1)	13.43 ± 0.09	3.37 ± 0.13	8.85 ± 0.04	44.0 ± 1.2
45	M0417-ID49	$2.20^{+0.69}_{-0.71}$...	FIR	1.9	(1)	12.92 ± 0.38	2.79 ± 0.48	9.24 ± 0.27	30.5 ± 6.7
46	M0417-ID58	3.65	$3.46^{+1.20}_{-1.01}$	(9)	4.0	(1)	12.82 ± 0.15	2.73 ± 0.19	8.95 ± 0.11	33.3 ± 2.6
47	M0417-ID121	3.65	$3.20^{+1.17}_{-0.95}$	(9)	3.6	(1)	12.71 ± 0.17	2.62 ± 0.20	8.74 ± 0.10	34.9 ± 3.2
48	M0417-ID204	$2.05^{+0.89}_{-0.76}$...	FIR	3.0	(1)	12.37 ± 0.27	2.25 ± 0.34	8.58 ± 0.27	30.4 ± 9.4
49	M0417-ID218	$1.63^{+0.52}_{-0.51}$...	FIR	4.0	(1)	12.61 ± 0.36	2.47 ± 0.46	8.77 ± 0.17	32.0 ± 6.4

Table C2 continued

Table C2 (continued)

#	Name	z_{best}^a	z_{FIR}^b	Ref. a	μ^c	Model d	$\log(L_{\text{IR}})^e$	$\log(\text{SFR})^e$	$\log(M_{\text{dust}})^e$	T_{dust}^f
							($\mu^{-1}L_{\odot}$)	($\mu^{-1}M_{\odot} \text{ yr}^{-1}$)	($\mu^{-1}M_{\odot}$)	(K)
50	M0417-ID221	$1.15^{+0.07}_{-0.64}$	$1.80^{+0.57}_{-0.55}$	HST	2.9	(1)	11.97 ± 0.20	1.85 ± 0.36	8.73 ± 0.12	24.8 ± 4.2
51	M0417-ID223	$1.10^{+0.09}_{-0.08}$	$1.64^{+0.58}_{-0.53}$	HST	2.8	(1)	12.24 ± 0.19	2.10 ± 0.31	8.84 ± 0.21	25.5 ± 2.4
52	M0429-ID27	$2.72^{+0.77}_{-0.40}$	$2.42^{+0.57}_{-0.52}$	HST	4.0	(2)	12.69 ± 0.23	2.63 ± 0.27	8.52 ± 0.19	34.7 ± 5.8
53	M0553-ID17	$2.49^{+0.84}_{-0.74}$...	FIR	1.6	(1)	12.63 ± 0.30	2.53 ± 0.36	8.80 ± 0.22	31.7 ± 7.5
54	M0553-ID61	$1.49^{+0.07}_{-0.09}$	$2.08^{+0.65}_{-0.68}$	HST	2.0	(1)	12.14 ± 0.18	2.01 ± 0.30	8.81 ± 0.08	25.8 ± 1.9
55	M0553-ID133	1.14	$1.30^{+0.40}_{-0.46}$	(10)	4.4	(1)	12.80 ± 0.12	2.62 ± 0.30	9.10 ± 0.04	30.4 ± 0.5
56	M0553-ID190	1.14	$1.30^{+0.40}_{-0.46}$	(10)	6.6	(1)	12.98 ± 0.12	2.79 ± 0.30	9.28 ± 0.04	30.6 ± 0.4
57	M0553-ID200	$1.71^{+0.82}_{-0.58}$...	FIR	2.2	(1)	12.36 ± 0.31	2.21 ± 0.39	8.41 ± 0.15	31.3 ± 9.1
58	M0553-ID249	1.14	$1.30^{+0.40}_{-0.46}$	(10)	5.8	(1)	12.82 ± 0.12	2.64 ± 0.30	9.12 ± 0.04	30.5 ± 0.5
59	M0553-ID275	$0.94^{+0.06}_{-0.05}$	$0.87^{+0.33}_{-0.34}$	HST	2.5	(1)	12.41 ± 0.17	2.22 ± 0.34	8.36 ± 0.11	34.7 ± 3.0
60	M0553-ID303	0.84	$0.98^{+0.39}_{-0.39}$	(8)	1.5	(1)	12.48 ± 0.16	2.31 ± 0.31	8.66 ± 0.12	30.8 ± 2.7
61	M0553-ID355	$1.82^{+0.11}_{-0.06}$	$1.91^{+0.65}_{-0.66}$	HST	1.8	(1)	12.32 ± 0.14	2.17 ± 0.25	8.64 ± 0.09	29.9 ± 2.0
62	M0553-ID375	$2.42^{+0.80}_{-0.72}$...	FIR	1.0	(-1)	12.54 ± 0.29	2.44 ± 0.34	8.69 ± 0.20	31.9 ± 7.4
63	M0553-ID398	$3.39^{+1.22}_{-0.99}$...	FIR	1.7	(1)	12.62 ± 0.27	2.53 ± 0.32	8.79 ± 0.22	32.8 ± 8.7
64	M1115-ID02	$1.92^{+0.11}_{-0.10}$	$1.14^{+0.68}_{-0.25}$	HST	3.1	(2)	13.04 ± 0.09	2.91 ± 0.14	9.11 ± 0.07	33.9 ± 1.3
65	M1115-ID04	1.60	$1.38^{+0.39}_{-0.50}$	(11)	2.7	(2)	12.66 ± 0.03	2.55 ± 0.09	8.51 ± 0.06	36.2 ± 1.1
66	M1115-ID34	$3.21^{+1.05}_{-0.94}$...	FIR	2.9	(2)	12.53 ± 0.28	2.43 ± 0.31	8.73 ± 0.23	25.6 ± 6.9
67	M1149-ID77	1.46	$1.61^{+0.52}_{-0.51}$	(2)	3.0	(2)	12.01 ± 0.05	1.81 ± 0.22	8.38 ± 0.09	30.6 ± 1.6
68	M1149-ID229	$1.38^{+0.28}_{-0.06}$	$2.05^{+0.67}_{-0.72}$	HST	2.0	(2)	11.79 ± 0.13	1.63 ± 0.20	8.54 ± 0.14	25.5 ± 2.5
69	M1206-ID27	1.04	$1.16^{+0.45}_{-0.49}$	(12)	5.2	(2)	12.20 ± 0.04	1.98 ± 0.21	8.56 ± 0.06	29.9 ± 1.2
70	M1206-ID55	1.04	$1.23^{+0.31}_{-0.53}$	(12)	7.7	(2)	12.37 ± 0.05	2.24 ± 0.11	8.54 ± 0.05	31.5 ± 0.5
71	M1206-ID60	1.04	$1.23^{+0.31}_{-0.53}$	(12)	25.5	(2)	12.45 ± 0.05	2.32 ± 0.11	8.62 ± 0.05	31.6 ± 0.5
72	M1206-ID61	1.04	$1.22^{+0.33}_{-0.51}$	(12)	73.1	(2)	12.58 ± 0.05	2.46 ± 0.11	8.76 ± 0.07	31.5 ± 0.6
73	M1311-ID27	2.19	$1.55^{+0.61}_{-0.59}$	(11)	15.8	(2)	12.74 ± 0.03	2.61 ± 0.10	8.45 ± 0.05	39.9 ± 0.8
74	M1311-ID33	$2.32^{+0.77}_{-0.20}$	$1.78^{+0.46}_{-0.44}$	HST	2.6	(2)	12.53 ± 0.21	2.42 ± 0.33	8.30 ± 0.12	32.0 ± 5.4
75	M1423-ID38	$1.90^{+0.04}_{-0.14}$	$1.60^{+0.49}_{-0.47}$	HST	4.3	(2)	12.36 ± 0.08	2.27 ± 0.12	8.18 ± 0.13	37.2 ± 1.9
76	M1931-ID55	$2.15^{+0.65}_{-0.34}$...	FIR	2.9	(2)	12.60 ± 0.25	2.56 ± 0.29	8.80 ± 0.15	35.6 ± 5.6
77	M1931-ID61	$3.02^{+1.03}_{-0.89}$...	FIR	2.4	(2)	12.34 ± 0.34	2.24 ± 0.42	8.56 ± 0.25	32.0 ± 8.1
78	M2129-ID46	1.48	$1.20^{+0.43}_{-0.47}$	(5)	3.7	(2)	12.06 ± 0.04	1.91 ± 0.16	7.91 ± 0.12	36.7 ± 1.8
79	P171-ID69	$3.57^{+1.42}_{-1.08}$...	FIR	2.7	(1)	12.77 ± 0.27	2.69 ± 0.31	8.95 ± 0.25	30.1 ± 9.0
80	P171-ID177	$2.55^{+1.11}_{-0.83}$...	FIR	2.6	(1)	12.47 ± 0.27	2.35 ± 0.33	8.58 ± 0.18	31.5 ± 9.3
81	R0032-ID53	3.63	$1.95^{+0.62}_{-0.64}$	(13)	9.2	(1)	13.18 ± 0.08	3.16 ± 0.15	8.34 ± 0.04	49.6 ± 1.5
82	R0032-ID55	3.63	$1.95^{+0.62}_{-0.64}$	(13)	18.8	(1)	13.46 ± 0.08	3.44 ± 0.16	8.61 ± 0.04	49.6 ± 1.2
83	R0032-ID57	3.63	$1.95^{+0.64}_{-0.65}$	(13)	7.5	(1)	12.70 ± 0.08	2.66 ± 0.14	7.95 ± 0.07	49.6 ± 2.7
84	R0032-ID58	3.63	$1.95^{+0.62}_{-0.64}$	(13)	7.7	(1)	13.20 ± 0.07	3.18 ± 0.15	8.36 ± 0.04	49.6 ± 1.5
85	R0032-ID127	$3.14^{+1.03}_{-0.95}$...	FIR	9.7	(1)	12.89 ± 0.23	2.79 ± 0.29	9.16 ± 0.18	31.4 ± 7.7
86	R0032-ID131	$3.27^{+1.15}_{-0.95}$...	FIR	3.3	(1)	12.83 ± 0.32	2.74 ± 0.36	9.09 ± 0.28	31.9 ± 8.1
87	R0032-ID198	$1.63^{+0.55}_{-0.51}$...	FIR	2.0	(1)	12.55 ± 0.54	2.41 ± 0.63	8.67 ± 0.16	32.3 ± 6.9
88	R0032-ID220	$1.47^{+0.55}_{-0.50}$...	FIR	2.1	(1)	12.47 ± 0.34	2.31 ± 0.42	8.52 ± 0.14	33.6 ± 8.1
89	R0032-ID238	$2.95^{+1.04}_{-0.89}$...	FIR	1.7	(1)	12.57 ± 0.29	2.47 ± 0.35	8.84 ± 0.25	31.3 ± 8.0
90	R0032-ID276	$1.99^{+0.83}_{-0.68}$...	FIR	2.6	(1)	12.46 ± 0.37	2.31 ± 0.45	8.56 ± 0.20	33.1 ± 9.6
91	R0032-ID287	$1.32^{+0.42}_{-0.45}$...	FIR	1.6	(1)	12.56 ± 0.29	2.41 ± 0.38	8.62 ± 0.11	33.0 ± 6.7
92	R0600-ID12	$1.09^{+0.47}_{-0.43}$...	FIR	1.2	(1)	12.14 ± 0.36	1.96 ± 0.46	8.17 ± 0.12	33.6 ± 8.7
93	R0600-ID13	1.27	$1.23^{+0.40}_{-0.45}$	(8)	1.3	(1)	12.49 ± 0.13	2.34 ± 0.26	8.52 ± 0.07	33.6 ± 1.8
94	R0600-ID111	$1.92^{+0.07}_{-0.10}$	$1.15^{+0.52}_{-0.47}$	HST	2.7	(1)	12.31 ± 0.19	2.18 ± 0.29	7.97 ± 0.09	46.1 ± 8.1
95	R0949-ID10	$3.87^{+1.28}_{-1.14}$...	FIR	3.0	(1)	12.94 ± 0.35	2.85 ± 0.41	9.23 ± 0.31	31.5 ± 8.0
96	R0949-ID122	$1.77^{+0.72}_{-0.60}$...	FIR	2.8	(1)	12.19 ± 0.42	2.05 ± 0.53	8.31 ± 0.22	32.2 ± 8.7
97	R0949-ID124	$2.89^{+1.01}_{-0.85}$...	FIR	3.3	(1)	12.69 ± 0.31	2.60 ± 0.35	8.86 ± 0.24	32.4 ± 8.1
98	R1347-ID41	0.85	$1.51^{+0.72}_{-0.51}$	(11)	4.4	(2)	11.60 ± 0.04	1.50 ± 0.15	8.16 ± 0.08	30.7 ± 0.8
99	R1347-ID145	1.77	$1.68^{+0.61}_{-0.57}$	(14)	3.3	(2)	12.77 ± 0.03	2.62 ± 0.10	8.84 ± 0.05	34.0 ± 0.3
100	R1347-ID148	1.77	$1.68^{+0.61}_{-0.57}$	(14)	5.8	(2)	12.91 ± 0.04	2.76 ± 0.10	8.98 ± 0.04	34.0 ± 0.2

Table C2 continued

Table C2 (continued)

#	Name	z_{best}^a	z_{FIR}^b	Ref. a	μ^c	Model d	$\log(L_{\text{IR}})^e$	$\log(\text{SFR})^e$	$\log(M_{\text{dust}})^e$	T_{dust}^f
							$(\mu^{-1}L_{\odot})$	$(\mu^{-1}M_{\odot} \text{ yr}^{-1})$	$(\mu^{-1}M_{\odot})$	(K)
101	R1347-ID166	$2.43^{+0.69}_{-0.78}$...	FIR	3.2	(2)	12.19 ± 0.37	2.00 ± 0.54	8.59 ± 0.20	34.0 ± 7.6
102	R2129-ID37	$2.48^{+0.65}_{-0.08}$	$1.85^{+0.56}_{-0.55}$	HST	1.9	(2)	12.52 ± 0.13	2.41 ± 0.19	8.24 ± 0.08	36.0 ± 5.3
103	R2211-ID19	$1.58^{+0.66}_{-0.54}$...	FIR	2.0	(1)	12.24 ± 0.31	2.10 ± 0.39	8.31 ± 0.16	33.1 ± 8.9
104	R2211-ID35	$1.10^{+0.36}_{-0.44}$...	FIR	2.4	(1)	12.86 ± 0.48	2.62 ± 0.62	8.99 ± 0.12	32.9 ± 6.3
105	R2211-ID171	$2.31^{+0.74}_{-0.69}$...	FIR	2.5	(1)	12.65 ± 0.43	2.56 ± 0.50	8.80 ± 0.23	32.1 ± 7.2
Secondary Sample ($S/N_{\text{ALMA}} = 4 - 5$)										
106	A2537-ID24	$3.11^{+0.10}_{-0.06}$	$1.04^{+0.34}_{-0.38}$	HST	19.8	(1)	13.27 ± 0.08	3.44 ± 0.22	7.83 ± 0.06	65.9 ± 5.3
107	A2744-ID47	$1.65^{+0.02}_{-0.03}$	$1.60^{+0.52}_{-0.70}$	HST	2.8	(2)	11.93 ± 0.07	1.87 ± 0.11	8.48 ± 0.33	32.9 ± 5.1
108	A2744-ID178	0.94	$1.85^{+0.69}_{-0.52}$	(15)	2.0	(2)	11.61 ± 0.06	1.50 ± 0.12	8.36 ± 0.16	32.4 ± 1.1
109	A2744-ID227	2.58	$1.04^{+0.54}_{-0.46}$	(5)	1.8	(2)	12.71 ± 0.05	2.77 ± 0.19	7.88 ± 0.27	...
110	A370-ID27	$3.14^{+0.04}_{-0.04}$	$2.58^{+0.92}_{-0.79}$	HST	5.2	(2)	12.35 ± 0.06	2.28 ± 0.10	8.39 ± 0.26	34.5 ± 3.4
111	A383-ID24	$0.67^{+0.04}_{-0.03}$	$0.84^{+0.29}_{-0.32}$	HST	2.6	(2)	11.65 ± 0.08	1.48 ± 0.23	7.95 ± 0.10	32.9 ± 0.8
112	A383-ID50	1.01	$0.62^{+0.45}_{-0.32}$	(16)	13.2	(2)	11.62 ± 0.07	1.57 ± 0.16	7.12 ± 0.34	33.7 ± 6.4
113	AS295-ID269	$2.34^{+1.19}_{-0.90}$...	FIR	2.0	(1)	12.37 ± 0.31	2.26 ± 0.38	8.56 ± 0.28	...
114	M0416-ID138	$1.84^{+0.02}_{-0.02}$	$1.44^{+0.44}_{-0.41}$	HST	1.5	(2)	11.99 ± 0.07	1.93 ± 0.11	8.07 ± 0.22	32.6 ± 2.3
115	M0416-ID156	$1.57^{+0.05}_{-0.03}$	$0.43^{+0.83}_{-0.39}$	HST	1.8	(2)	11.84 ± 0.06	1.81 ± 0.15	8.21 ± 0.23	24.2 ± 3.2
116	M1115-ID33	$0.74^{+0.02}_{-0.05}$	$2.84^{+1.12}_{-0.87}$	HST	1.6	(2)	10.69 ± 0.14	0.44 ± 0.27	7.69 ± 0.59	15.8 ± 1.1
117	M1149-ID27	$5.05^{+0.13}_{-0.26}$	$2.15^{+0.81}_{-0.70}$	HST	2.4	(2)	12.88 ± 0.08	2.89 ± 0.19	7.96 ± 0.17	46.6 ± 7.0
118	M1149-ID95	$1.12^{+0.19}_{-0.53}$	$2.22^{+0.87}_{-0.76}$	HST	1.3	(2)	11.40 ± 0.43	1.20 ± 0.56	8.42 ± 0.36	23.0 ± 4.2
119	M1206-ID38	1.49	$1.57^{+0.52}_{-0.54}$	(17)	4.2	(2)	11.94 ± 0.05	1.81 ± 0.12	8.09 ± 0.17	32.1 ± 1.6
120	M1206-ID84	0.48	$1.31^{+0.46}_{-0.46}$	(17)	1.0	(-1)	11.33 ± 0.11	1.24 ± 0.16	8.03 ± 0.20	30.6 ± 0.9
121	M1423-ID52	$1.23^{+0.05}_{-0.05}$	$1.28^{+0.43}_{-0.47}$	HST	4.0	(2)	11.79 ± 0.08	1.64 ± 0.18	7.95 ± 0.21	...
122	M1423-ID76	$0.67^{+0.14}_{-0.23}$	$1.89^{+0.36}_{-0.47}$	HST	1.2	(2)	11.05 ± 0.41	0.95 ± 0.45	7.96 ± 0.54	33.7 ± 4.9
123	M1931-ID69	$1.60^{+0.60}_{-0.54}$...	FIR	1.8	(2)	11.77 ± 0.41	1.65 ± 0.49	7.95 ± 0.27	33.1 ± 7.6
124	M2129-ID62	1.48	$1.46^{+0.42}_{-0.43}$	(5)	2.7	(2)	11.98 ± 0.05	1.90 ± 0.10	8.09 ± 0.17	34.8 ± 2.7
125	R0949-ID119	$0.56^{+0.09}_{-0.05}$	$1.43^{+0.50}_{-0.49}$	HST	1.5	(1)	11.39 ± 0.22	1.21 ± 0.38	8.32 ± 0.15	20.3 ± 2.4

^a Best redshifts. *HST* or far-IR photometric redshifts are presented as the median of their likelihood distributions with 1σ confidence range. References of spectroscopic redshifts: (1) CO spectroscopy (private communication from F. Bauer), (2) GLASS and Laporte et al. (2017), (3) Wold et al. (2012), (4) Soucail et al. (1999), (5) GLASS (Treu et al. 2015; Schmidt et al. 2014), (6) Caputi et al. (2021), (7) Gómez et al. (2012), GLASS (Treu et al. 2015; Schmidt et al. 2014) and Walth et al. (2019). (8) CO spectroscopy (ALCS; Fujimoto et al. in preparation), (9) CO spectroscopy (Kohno et al. in preparation), (10) Ebeling et al. (2017), (11) Caminha et al. (2019), (12) “Cosmic Snake” (e.g., Ebeling et al. 2009; Biviano et al. 2013; Cava et al. 2018), (13) Dessauges-Zavadsky et al. (2017), (14) Richard et al. (2021), (15) GLASS (Wang et al. 2015), (16) Sand et al. (2004), (17) Biviano et al. (2013). *HST* photometric redshifts are taken from Molino et al. (2017), Shipley et al. (2018) and Coe et al. (2019) for sources detected in CLASH, HFF and RELICS cluster fields, respectively.

^b Far-IR photometric redshifts modeled with MAGPHYS+PHOTO-Z (Battisti et al. 2019; see Section 4.1). This column is left blank if $z_{\text{best}} = z_{\text{FIR}}$.

^c Lensing magnification factors are calculated based on the cluster mass models and the best redshifts of sources (see Section 5.1, also for the evaluation of uncertainty).

^d Adopted models of lensing magnifications. (1): GLAFIC (Oguri 2010), (2): Zitrin NFW (Zitrin et al. 2013, 2015), (-1): lensing models are not available (e.g., the sources are cluster member galaxies, or out of *HST* footprint).

^e These quantities, derived with MAGPHYS (da Cunha et al. 2008, 2015; Battisti et al. 2019), are not corrected for lensing magnification.

^f Modeled with MBB spectrum with fixed dust emissivity at $\beta = 1.8$ (Section 4.1).

REFERENCES

- Abell, G. O. 1958, *ApJS*, 3, 211, doi: [10.1086/190036](https://doi.org/10.1086/190036)
- Abell, G. O., Corwin, Harold G., J., & Olowin, R. P. 1989, *ApJS*, 70, 1, doi: [10.1086/191333](https://doi.org/10.1086/191333)
- Alcalde Pampliega, B., Pérez-González, P. G., Barro, G., et al. 2019, *ApJ*, 876, 135, doi: [10.3847/1538-4357/ab14f2](https://doi.org/10.3847/1538-4357/ab14f2)
- Aravena, M., Boogaard, L., González-López, J., et al. 2020, *ApJ*, 901, 79, doi: [10.3847/1538-4357/ab99a2](https://doi.org/10.3847/1538-4357/ab99a2)
- Armus, L., Mazzarella, J. M., Evans, A. S., et al. 2009, *PASP*, 121, 559, doi: [10.1086/600092](https://doi.org/10.1086/600092)
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33, doi: [10.1051/0004-6361/201322068](https://doi.org/10.1051/0004-6361/201322068)
- Bakx, T. J. L. C., Tamura, Y., Hashimoto, T., et al. 2020, *MNRAS*, 493, 4294, doi: [10.1093/mnras/staa509](https://doi.org/10.1093/mnras/staa509)

- Bakx, T. J. L. C., Sommovigo, L., Carniani, S., et al. 2021, MNRAS, 508, L58, doi: [10.1093/mnrasl/slab104](https://doi.org/10.1093/mnrasl/slab104)
- Barcos-Muñoz, L., Leroy, A. K., Evans, A. S., et al. 2017, ApJ, 843, 117, doi: [10.3847/1538-4357/aa789a](https://doi.org/10.3847/1538-4357/aa789a)
- Battisti, A. J., da Cunha, E., Grasha, K., et al. 2019, ApJ, 882, 61, doi: [10.3847/1538-4357/ab345d](https://doi.org/10.3847/1538-4357/ab345d)
- Bertin, E., & Arnouts, S. 1996, A&AS, 117, 393, doi: [10.1051/aas:1996164](https://doi.org/10.1051/aas:1996164)
- Béthermin, M., De Breuck, C., Sargent, M., & Daddi, E. 2015, A&A, 576, L9, doi: [10.1051/0004-6361/201525718](https://doi.org/10.1051/0004-6361/201525718)
- Béthermin, M., De Breuck, C., Gullberg, B., et al. 2016, A&A, 586, L7, doi: [10.1051/0004-6361/201527739](https://doi.org/10.1051/0004-6361/201527739)
- Birkin, J. E., Weiss, A., Wardlow, J. L., et al. 2021, MNRAS, 501, 3926, doi: [10.1093/mnras/staa3862](https://doi.org/10.1093/mnras/staa3862)
- Biviano, A., Rosati, P., Balestra, I., et al. 2013, A&A, 558, A1, doi: [10.1051/0004-6361/201321955](https://doi.org/10.1051/0004-6361/201321955)
- Boone, F., Clément, B., Richard, J., et al. 2013, A&A, 559, L1, doi: [10.1051/0004-6361/201322552](https://doi.org/10.1051/0004-6361/201322552)
- Bouwens, R., González-López, J., Aravena, M., et al. 2020, ApJ, 902, 112, doi: [10.3847/1538-4357/abb830](https://doi.org/10.3847/1538-4357/abb830)
- Bradley, L., Sipőcz, B., Robitaille, T., et al. 2019, astropy/photutils: v0.7.2, v0.7.2, Zenodo, doi: [10.5281/zenodo.3568287](https://doi.org/10.5281/zenodo.3568287)
- Brisbin, D., Miettinen, O., Aravena, M., et al. 2017, A&A, 608, A15, doi: [10.1051/0004-6361/201730558](https://doi.org/10.1051/0004-6361/201730558)
- Burnham, A. D., Casey, C. M., Zavala, J. A., et al. 2021, ApJ, 910, 89, doi: [10.3847/1538-4357/abe401](https://doi.org/10.3847/1538-4357/abe401)
- Cañameras, R., Nesvadba, N. P. H., Guery, D., et al. 2015, A&A, 581, A105, doi: [10.1051/0004-6361/201425128](https://doi.org/10.1051/0004-6361/201425128)
- Cañameras, R., Yang, C., Nesvadba, N. P. H., et al. 2018, A&A, 620, A61, doi: [10.1051/0004-6361/201833625](https://doi.org/10.1051/0004-6361/201833625)
- Caminha, G. B., Rosati, P., Grillo, C., et al. 2019, A&A, 632, A36, doi: [10.1051/0004-6361/201935454](https://doi.org/10.1051/0004-6361/201935454)
- Caputi, K. I., Caminha, G. B., Fujimoto, S., et al. 2021, ApJ, 908, 146, doi: [10.3847/1538-4357/abd4d0](https://doi.org/10.3847/1538-4357/abd4d0)
- Casey, C. M., Zavala, J. A., Spilker, J., et al. 2018, ApJ, 862, 77, doi: [10.3847/1538-4357/aac82d](https://doi.org/10.3847/1538-4357/aac82d)
- Casey, C. M., Zavala, J. A., Manning, S. M., et al. 2021, ApJ, 923, 215, doi: [10.3847/1538-4357/ac2eb4](https://doi.org/10.3847/1538-4357/ac2eb4)
- Cava, A., Schaerer, D., Richard, J., et al. 2018, Nature Astronomy, 2, 76, doi: [10.1038/s41550-017-0295-x](https://doi.org/10.1038/s41550-017-0295-x)
- Chabrier, G. 2003, PASP, 115, 763, doi: [10.1086/376392](https://doi.org/10.1086/376392)
- Chary, R., & Elbaz, D. 2001, ApJ, 556, 562, doi: [10.1086/321609](https://doi.org/10.1086/321609)
- Chen, C.-C., Liao, C.-L., Smail, I., et al. 2021, arXiv e-prints, arXiv:2112.07430, <https://arxiv.org/abs/2112.07430>
- Coe, D., Salmon, B., Bradač, M., et al. 2019, ApJ, 884, 85, doi: [10.3847/1538-4357/ab412b](https://doi.org/10.3847/1538-4357/ab412b)
- da Cunha, E., Charlot, S., & Elbaz, D. 2008, MNRAS, 388, 1595, doi: [10.1111/j.1365-2966.2008.13535.x](https://doi.org/10.1111/j.1365-2966.2008.13535.x)
- da Cunha, E., Groves, B., Walter, F., et al. 2013, ApJ, 766, 13, doi: [10.1088/0004-637X/766/1/13](https://doi.org/10.1088/0004-637X/766/1/13)
- da Cunha, E., Walter, F., Smail, I. R., et al. 2015, ApJ, 806, 110, doi: [10.1088/0004-637X/806/1/110](https://doi.org/10.1088/0004-637X/806/1/110)
- da Cunha, E., Hodge, J. A., Casey, C. M., et al. 2021, ApJ, 919, 30, doi: [10.3847/1538-4357/ac0ae0](https://doi.org/10.3847/1538-4357/ac0ae0)
- Dessauges-Zavadsky, M., Zamojski, M., Schaerer, D., et al. 2015, A&A, 577, A50, doi: [10.1051/0004-6361/201424661](https://doi.org/10.1051/0004-6361/201424661)
- Dessauges-Zavadsky, M., Zamojski, M., Rujopakarn, W., et al. 2017, A&A, 605, A81, doi: [10.1051/0004-6361/201628513](https://doi.org/10.1051/0004-6361/201628513)
- Dessauges-Zavadsky, M., Richard, J., Combes, F., et al. 2019, Nature Astronomy, 3, 1115, doi: [10.1038/s41550-019-0874-0](https://doi.org/10.1038/s41550-019-0874-0)
- Díaz-Santos, T., Armus, L., Charmandaris, V., et al. 2017, ApJ, 846, 32, doi: [10.3847/1538-4357/aa81d7](https://doi.org/10.3847/1538-4357/aa81d7)
- Draine, B. T., & Li, A. 2007, ApJ, 657, 810, doi: [10.1086/511055](https://doi.org/10.1086/511055)
- Drew, P. M., & Casey, C. M. 2022, arXiv e-prints, arXiv:2203.16655, <https://arxiv.org/abs/2203.16655>
- Dudzevičiūtė, U., Smail, I., Swinbank, A. M., et al. 2020, MNRAS, 494, 3828, doi: [10.1093/mnras/staa769](https://doi.org/10.1093/mnras/staa769)
- . 2021, MNRAS, 500, 942, doi: [10.1093/mnras/staa3285](https://doi.org/10.1093/mnras/staa3285)
- Dunlop, J. S., McLure, R. J., Biggs, A. D., et al. 2017, MNRAS, 466, 861, doi: [10.1093/mnras/stw3088](https://doi.org/10.1093/mnras/stw3088)
- Ebeling, H., Edge, A. C., & Henry, J. P. 2001, ApJ, 553, 668, doi: [10.1086/320958](https://doi.org/10.1086/320958)
- Ebeling, H., Ma, C. J., Kneib, J. P., et al. 2009, MNRAS, 395, 1213, doi: [10.1111/j.1365-2966.2009.14502.x](https://doi.org/10.1111/j.1365-2966.2009.14502.x)
- Ebeling, H., Qi, J., & Richard, J. 2017, MNRAS, 471, 3305, doi: [10.1093/mnras/stx1636](https://doi.org/10.1093/mnras/stx1636)
- Egami, E., Rex, M., Rawle, T. D., et al. 2010, A&A, 518, L12, doi: [10.1051/0004-6361/201014696](https://doi.org/10.1051/0004-6361/201014696)
- Elbaz, D., Leiton, R., Nagar, N., et al. 2018, A&A, 616, A110, doi: [10.1051/0004-6361/201732370](https://doi.org/10.1051/0004-6361/201732370)
- Fogarty, K., Postman, M., Li, Y., et al. 2019, ApJ, 879, 103, doi: [10.3847/1538-4357/ab22a4](https://doi.org/10.3847/1538-4357/ab22a4)
- Franco, M., Elbaz, D., Béthermin, M., et al. 2018, A&A, 620, A152, doi: [10.1051/0004-6361/201832928](https://doi.org/10.1051/0004-6361/201832928)
- Franco, M., Elbaz, D., Zhou, L., et al. 2020, A&A, 643, A30, doi: [10.1051/0004-6361/202038312](https://doi.org/10.1051/0004-6361/202038312)
- Fudamoto, Y., Oesch, P. A., Faisst, A., et al. 2020, A&A, 643, A4, doi: [10.1051/0004-6361/202038163](https://doi.org/10.1051/0004-6361/202038163)
- Fudamoto, Y., Oesch, P. A., Schouws, S., et al. 2021, Nature, 597, 489–492, doi: [10.1038/s41586-021-03846-z](https://doi.org/10.1038/s41586-021-03846-z)
- Fujimoto, S., Ouchi, M., Ono, Y., et al. 2016, ApJS, 222, 1, doi: [10.3847/0067-0049/222/1/1](https://doi.org/10.3847/0067-0049/222/1/1)

- Fujimoto, S., Ouchi, M., Shibuya, T., & Nagai, H. 2017, *ApJ*, 850, 83, doi: [10.3847/1538-4357/aa93e6](https://doi.org/10.3847/1538-4357/aa93e6)
- Fujimoto, S., Oguri, M., Brammer, G., et al. 2021, *ApJ*, 911, 99, doi: [10.3847/1538-4357/abd7ec](https://doi.org/10.3847/1538-4357/abd7ec)
- Fujimoto, S., Hatsukade, K., Yoshimura, Y., et al. in preparation
- Gómez, P. L., Valkonen, L. E., Romer, A. K., et al. 2012, *AJ*, 144, 79, doi: [10.1088/0004-6256/144/3/79](https://doi.org/10.1088/0004-6256/144/3/79)
- Gómez-Guijarro, C., Elbaz, D., Xiao, M., et al. 2022, *A&A*, 658, A43, doi: [10.1051/0004-6361/202141615](https://doi.org/10.1051/0004-6361/202141615)
- González-López, J., Bauer, F. E., Romero-Cañizales, C., et al. 2017, *A&A*, 597, A41, doi: [10.1051/0004-6361/201628806](https://doi.org/10.1051/0004-6361/201628806)
- González-López, J., Novak, M., Decarli, R., et al. 2020, *ApJ*, 897, 91, doi: [10.3847/1538-4357/ab765b](https://doi.org/10.3847/1538-4357/ab765b)
- Greve, T. R., Vieira, J. D., Weiß, A., et al. 2012, *ApJ*, 756, 101, doi: [10.1088/0004-637X/756/1/101](https://doi.org/10.1088/0004-637X/756/1/101)
- Greve, T. R., Leonidaki, I., Xilouris, E. M., et al. 2014, *ApJ*, 794, 142, doi: [10.1088/0004-637X/794/2/142](https://doi.org/10.1088/0004-637X/794/2/142)
- Griffin, M. J., Abergel, A., Abreu, A., et al. 2010, *A&A*, 518, L3, doi: [10.1051/0004-6361/201014519](https://doi.org/10.1051/0004-6361/201014519)
- Gruppioni, C., Pozzi, F., Rodighiero, G., et al. 2013, *MNRAS*, 432, 23, doi: [10.1093/mnras/stt308](https://doi.org/10.1093/mnras/stt308)
- Gullberg, B., Smail, I., Swinbank, A. M., et al. 2019, *MNRAS*, 490, 4956, doi: [10.1093/mnras/stz2835](https://doi.org/10.1093/mnras/stz2835)
- Harikane, Y., Ouchi, M., Inoue, A. K., et al. 2020, *ApJ*, 896, 93, doi: [10.3847/1538-4357/ab94bd](https://doi.org/10.3847/1538-4357/ab94bd)
- Hashimoto, T., Inoue, A. K., Mawatari, K., et al. 2019, *PASJ*, 71, 71, doi: [10.1093/pasj/psz049](https://doi.org/10.1093/pasj/psz049)
- Heinis, S., Buat, V., Béthermin, M., et al. 2014, *MNRAS*, 437, 1268, doi: [10.1093/mnras/stt1960](https://doi.org/10.1093/mnras/stt1960)
- Hodge, J. A., & da Cunha, E. 2020, *Royal Society Open Science*, 7, 200556, doi: [10.1098/rsos.200556](https://doi.org/10.1098/rsos.200556)
- Hodge, J. A., Swinbank, A. M., Simpson, J. M., et al. 2016, *ApJ*, 833, 103, doi: [10.3847/1538-4357/833/1/103](https://doi.org/10.3847/1538-4357/833/1/103)
- Howell, J. H., Armus, L., Mazzarella, J. M., et al. 2010, *ApJ*, 715, 572, doi: [10.1088/0004-637X/715/1/572](https://doi.org/10.1088/0004-637X/715/1/572)
- Hunt, L. K., Draine, B. T., Bianchi, S., et al. 2015, *A&A*, 576, A33, doi: [10.1051/0004-6361/201424734](https://doi.org/10.1051/0004-6361/201424734)
- Hurley, P. D., Oliver, S., Betancourt, M., et al. 2017, *MNRAS*, 464, 885, doi: [10.1093/mnras/stw2375](https://doi.org/10.1093/mnras/stw2375)
- Hwang, H. S., Elbaz, D., Magdis, G., et al. 2010, *MNRAS*, 409, 75, doi: [10.1111/j.1365-2966.2010.17645.x](https://doi.org/10.1111/j.1365-2966.2010.17645.x)
- Ikarashi, S., Ivison, R. J., Caputi, K. I., et al. 2015, *ApJ*, 810, 133, doi: [10.1088/0004-637X/810/2/133](https://doi.org/10.1088/0004-637X/810/2/133)
- Inoue, A. K., Hashimoto, T., Chihara, H., & Koike, C. 2020, *MNRAS*, 495, 1577, doi: [10.1093/mnras/staa1203](https://doi.org/10.1093/mnras/staa1203)
- Ivison, R. J., Smail, I., Le Borgne, J. F., et al. 1998, *MNRAS*, 298, 583, doi: [10.1046/j.1365-8711.1998.01677.x](https://doi.org/10.1046/j.1365-8711.1998.01677.x)
- Ivison, R. J., Greve, T. R., Dunlop, J. S., et al. 2007, *MNRAS*, 380, 199, doi: [10.1111/j.1365-2966.2007.12044.x](https://doi.org/10.1111/j.1365-2966.2007.12044.x)
- Kawamata, R., Ishigaki, M., Shimasaku, K., et al. 2018, *ApJ*, 855, 4, doi: [10.3847/1538-4357/aaa6cf](https://doi.org/10.3847/1538-4357/aaa6cf)
- Kawamata, R., Oguri, M., Ishigaki, M., Shimasaku, K., & Ouchi, M. 2016, *ApJ*, 819, 114, doi: [10.3847/0004-637X/819/2/114](https://doi.org/10.3847/0004-637X/819/2/114)
- Kennicutt, R. C., & Evans, N. J. 2012, *ARA&A*, 50, 531, doi: [10.1146/annurev-astro-081811-125610](https://doi.org/10.1146/annurev-astro-081811-125610)
- Knudsen, K. K., Watson, D., Frayer, D., et al. 2017, *MNRAS*, 466, 138, doi: [10.1093/mnras/stw3066](https://doi.org/10.1093/mnras/stw3066)
- Kohno, K. 2019, in *ALMA2019: Science Results and Cross-Facility Synergies*, 64, doi: [10.5281/zenodo.3585294](https://doi.org/10.5281/zenodo.3585294)
- Kohno, K., Fujimoto, S., Hatsukade, K., et al. in preparation
- Lagos, C. d. P., da Cunha, E., Robotham, A. S. G., et al. 2020, *MNRAS*, 499, 1948, doi: [10.1093/mnras/staa2861](https://doi.org/10.1093/mnras/staa2861)
- Lang, P., Schinnerer, E., Smail, I., et al. 2019, *ApJ*, 879, 54, doi: [10.3847/1538-4357/ab1f77](https://doi.org/10.3847/1538-4357/ab1f77)
- Laporte, N., Ellis, R. S., Boone, F., et al. 2017, *ApJL*, 837, L21, doi: [10.3847/2041-8213/aa62aa](https://doi.org/10.3847/2041-8213/aa62aa)
- Laporte, N., Zitrin, A., Ellis, R. S., et al. 2021, *MNRAS*, 505, 4838, doi: [10.1093/mnras/stab191](https://doi.org/10.1093/mnras/stab191)
- Leroy, A. K., Bolatto, A., Gordon, K., et al. 2011, *ApJ*, 737, 12, doi: [10.1088/0004-637X/737/1/12](https://doi.org/10.1088/0004-637X/737/1/12)
- Liang, L., Feldmann, R., Kereš, D., et al. 2019, *MNRAS*, 489, 1397, doi: [10.1093/mnras/stz2134](https://doi.org/10.1093/mnras/stz2134)
- Lim, C.-F., Wang, W.-H., Smail, I., et al. 2020, *ApJ*, 889, 80, doi: [10.3847/1538-4357/ab607f](https://doi.org/10.3847/1538-4357/ab607f)
- Lindner, R. R., Aguirre, P., Baker, A. J., et al. 2015, *ApJ*, 803, 79, doi: [10.1088/0004-637X/803/2/79](https://doi.org/10.1088/0004-637X/803/2/79)
- Liu, D., Lang, P., Magnelli, B., et al. 2019, *ApJS*, 244, 40, doi: [10.3847/1538-4365/ab42da](https://doi.org/10.3847/1538-4365/ab42da)
- Lotz, J. M., Koekemoer, A., Coe, D., et al. 2017, *ApJ*, 837, 97, doi: [10.3847/1538-4357/837/1/97](https://doi.org/10.3847/1538-4357/837/1/97)
- Lutz, D., Poglitsch, A., Altieri, B., et al. 2011, *A&A*, 532, A90, doi: [10.1051/0004-6361/201117107](https://doi.org/10.1051/0004-6361/201117107)
- Magdis, G. E., Daddi, E., Béthermin, M., et al. 2012, *ApJ*, 760, 6, doi: [10.1088/0004-637X/760/1/6](https://doi.org/10.1088/0004-637X/760/1/6)
- Magnelli, B., Lutz, D., Saintonge, A., et al. 2014, *A&A*, 561, A86, doi: [10.1051/0004-6361/201322217](https://doi.org/10.1051/0004-6361/201322217)
- Marrone, D. P., Spilker, J. S., Hayward, C. C., et al. 2018, *Nature*, 553, 51, doi: [10.1038/nature24629](https://doi.org/10.1038/nature24629)
- McMullin, J. P., Waters, B., Schiebel, D., Young, W., & Golap, K. 2007, *Astronomical Society of the Pacific Conference Series*, Vol. 376, *CASA Architecture and Applications*, ed. R. A. Shaw, F. Hill, & D. J. Bell, 127
- Michałowski, M. J., Dunlop, J. S., Ivison, R. J., et al. 2012, *MNRAS*, 426, 1845, doi: [10.1111/j.1365-2966.2012.21828.x](https://doi.org/10.1111/j.1365-2966.2012.21828.x)

- Miettinen, O., Delvecchio, I., Smolčić, V., et al. 2017, *A&A*, 606, A17, doi: [10.1051/0004-6361/201730762](https://doi.org/10.1051/0004-6361/201730762)
- Molino, A., Benítez, N., Ascaso, B., et al. 2017, *MNRAS*, 470, 95, doi: [10.1093/mnras/stx1243](https://doi.org/10.1093/mnras/stx1243)
- Nguyen, H. T., Schulz, B., Levenson, L., et al. 2010, *A&A*, 518, L5, doi: [10.1051/0004-6361/201014680](https://doi.org/10.1051/0004-6361/201014680)
- Oguri, M. 2010, *PASJ*, 62, 1017, doi: [10.1093/pasj/62.4.1017](https://doi.org/10.1093/pasj/62.4.1017)
- Okabe, T., Oguri, M., Peirani, S., et al. 2020, *MNRAS*, 496, 2591, doi: [10.1093/mnras/staa1479](https://doi.org/10.1093/mnras/staa1479)
- Oliver, S. J., Bock, J., Altieri, B., et al. 2012, *MNRAS*, 424, 1614, doi: [10.1111/j.1365-2966.2012.20912.x](https://doi.org/10.1111/j.1365-2966.2012.20912.x)
- Ott, S. 2010, *Astronomical Society of the Pacific Conference Series*, Vol. 434, *The Herschel Data Processing System — HIPE and Pipelines — Up and Running Since the Start of the Mission*, ed. Y. Mizumoto, K. I. Morita, & M. Ohishi, 139
- Piazzo, L., Calzoletti, L., Faustini, F., et al. 2015, *MNRAS*, 447, 1471, doi: [10.1093/mnras/stu2453](https://doi.org/10.1093/mnras/stu2453)
- Pilbratt, G. L., Riedinger, J. R., Passvogel, T., et al. 2010, *A&A*, 518, L1, doi: [10.1051/0004-6361/201014759](https://doi.org/10.1051/0004-6361/201014759)
- Poglitsch, A., Waelkens, C., Geis, N., et al. 2010, *A&A*, 518, L2, doi: [10.1051/0004-6361/201014535](https://doi.org/10.1051/0004-6361/201014535)
- Popping, G., Walter, F., Behroozi, P., et al. 2020, *ApJ*, 891, 135, doi: [10.3847/1538-4357/ab76c0](https://doi.org/10.3847/1538-4357/ab76c0)
- Postman, M., Coe, D., Benítez, N., et al. 2012, *ApJS*, 199, 25, doi: [10.1088/0067-0049/199/2/25](https://doi.org/10.1088/0067-0049/199/2/25)
- Rawle, T. D., Edge, A. C., Egami, E., et al. 2012, *ApJ*, 747, 29, doi: [10.1088/0004-637X/747/1/29](https://doi.org/10.1088/0004-637X/747/1/29)
- Rawle, T. D., Altieri, B., Egami, E., et al. 2016, *MNRAS*, 459, 1626, doi: [10.1093/mnras/stw712](https://doi.org/10.1093/mnras/stw712)
- Reuter, C., Vieira, J. D., Spilker, J. S., et al. 2020, *ApJ*, 902, 78, doi: [10.3847/1538-4357/abb599](https://doi.org/10.3847/1538-4357/abb599)
- Rex, M., Rawle, T. D., Egami, E., et al. 2010, *A&A*, 518, L13, doi: [10.1051/0004-6361/201014693](https://doi.org/10.1051/0004-6361/201014693)
- Richard, J., Jauzac, M., Limousin, M., et al. 2014, *MNRAS*, 444, 268, doi: [10.1093/mnras/stu1395](https://doi.org/10.1093/mnras/stu1395)
- Richard, J., Claeysens, A., Lagattuta, D., et al. 2021, *A&A*, 646, A83, doi: [10.1051/0004-6361/202039462](https://doi.org/10.1051/0004-6361/202039462)
- Rieke, G. H., Alonso-Herrero, A., Weiner, B. J., et al. 2009, *ApJ*, 692, 556, doi: [10.1088/0004-637X/692/1/556](https://doi.org/10.1088/0004-637X/692/1/556)
- Roseboom, I. G., Oliver, S. J., Kunz, M., et al. 2010, *MNRAS*, 409, 48, doi: [10.1111/j.1365-2966.2010.17634.x](https://doi.org/10.1111/j.1365-2966.2010.17634.x)
- Rosenberg, M. J. F., van der Werf, P. P., Aalto, S., et al. 2015, *ApJ*, 801, 72, doi: [10.1088/0004-637X/801/2/72](https://doi.org/10.1088/0004-637X/801/2/72)
- Rujopakarn, W., Dunlop, J. S., Rieke, G. H., et al. 2016, *ApJ*, 833, 12, doi: [10.3847/0004-637X/833/1/12](https://doi.org/10.3847/0004-637X/833/1/12)
- Sakamoto, K., Aalto, S., Barcos-Muñoz, L., et al. 2017, *ApJ*, 849, 14, doi: [10.3847/1538-4357/aa8f4b](https://doi.org/10.3847/1538-4357/aa8f4b)
- Sand, D. J., Treu, T., Smith, G. P., & Ellis, R. S. 2004, *ApJ*, 604, 88, doi: [10.1086/382146](https://doi.org/10.1086/382146)
- Schmidt, K. B., Treu, T., Brammer, G. B., et al. 2014, *ApJL*, 782, L36, doi: [10.1088/2041-8205/782/2/L36](https://doi.org/10.1088/2041-8205/782/2/L36)
- Schreiber, C., Elbaz, D., Pannella, M., et al. 2018, *A&A*, 609, A30, doi: [10.1051/0004-6361/201731506](https://doi.org/10.1051/0004-6361/201731506)
- Scoville, N., Aussel, H., Sheth, K., et al. 2014, *ApJ*, 783, 84, doi: [10.1088/0004-637X/783/2/84](https://doi.org/10.1088/0004-637X/783/2/84)
- Scoville, N., Lee, N., Vanden Bout, P., et al. 2017, *ApJ*, 837, 150, doi: [10.3847/1538-4357/aa61a0](https://doi.org/10.3847/1538-4357/aa61a0)
- Shipley, H. V., Lange-Vagle, D., Marchesini, D., et al. 2018, *ApJS*, 235, 14, doi: [10.3847/1538-4365/aaacce](https://doi.org/10.3847/1538-4365/aaacce)
- Simpson, J. M., Swinbank, A. M., Smail, I., et al. 2014, *ApJ*, 788, 125, doi: [10.1088/0004-637X/788/2/125](https://doi.org/10.1088/0004-637X/788/2/125)
- Simpson, J. M., Smail, I., Swinbank, A. M., et al. 2015, *ApJ*, 799, 81, doi: [10.1088/0004-637X/799/1/81](https://doi.org/10.1088/0004-637X/799/1/81)
- . 2017, *ApJ*, 839, 58, doi: [10.3847/1538-4357/aa65d0](https://doi.org/10.3847/1538-4357/aa65d0)
- . 2019, *ApJ*, 880, 43, doi: [10.3847/1538-4357/ab23ff](https://doi.org/10.3847/1538-4357/ab23ff)
- Simpson, J. M., Smail, I., Dudzevičiūtė, U., et al. 2020, *MNRAS*, 495, 3409, doi: [10.1093/mnras/staa1345](https://doi.org/10.1093/mnras/staa1345)
- Skibba, R. A., Engelbracht, C. W., Dale, D., et al. 2011, *ApJ*, 738, 89, doi: [10.1088/0004-637X/738/1/89](https://doi.org/10.1088/0004-637X/738/1/89)
- Sklias, P., Zamojski, M., Schaerer, D., et al. 2014, *A&A*, 561, A149, doi: [10.1051/0004-6361/201322424](https://doi.org/10.1051/0004-6361/201322424)
- Smail, I., Ivison, R. J., & Blain, A. W. 1997, *ApJL*, 490, L5, doi: [10.1086/311017](https://doi.org/10.1086/311017)
- Smail, I., Dudzevičiūtė, U., Stach, S. M., et al. 2021, *MNRAS*, 502, 3426, doi: [10.1093/mnras/stab283](https://doi.org/10.1093/mnras/stab283)
- Soifer, B. T., Neugebauer, G., Matthews, K., et al. 2000, *AJ*, 119, 509, doi: [10.1086/301233](https://doi.org/10.1086/301233)
- Soucail, G., Kneib, J. P., Bézécourt, J., et al. 1999, *A&A*, 343, L70. <https://arxiv.org/abs/astro-ph/9901123>
- Speagle, J. S., Steinhardt, C. L., Capak, P. L., & Silverman, J. D. 2014, *ApJS*, 214, 15, doi: [10.1088/0067-0049/214/2/15](https://doi.org/10.1088/0067-0049/214/2/15)
- Spilker, J. S., Marrone, D. P., Aravena, M., et al. 2016, *ApJ*, 826, 112, doi: [10.3847/0004-637X/826/2/112](https://doi.org/10.3847/0004-637X/826/2/112)
- Stach, S. M., Dudzevičiūtė, U., Smail, I., et al. 2019, *MNRAS*, 487, 4648, doi: [10.1093/mnras/stz1536](https://doi.org/10.1093/mnras/stz1536)
- Stalevski, M., Fritz, J., Baes, M., Nakos, T., & Popović, L. Č. 2012, *MNRAS*, 420, 2756, doi: [10.1111/j.1365-2966.2011.19775.x](https://doi.org/10.1111/j.1365-2966.2011.19775.x)
- Stalevski, M., Ricci, C., Ueda, Y., et al. 2016, *MNRAS*, 458, 2288, doi: [10.1093/mnras/stw444](https://doi.org/10.1093/mnras/stw444)
- Stetson, P. B. 1987, *PASP*, 99, 191, doi: [10.1086/131977](https://doi.org/10.1086/131977)
- Strandet, M. L., Weiss, A., Vieira, J. D., et al. 2016, *ApJ*, 822, 80, doi: [10.3847/0004-637X/822/2/80](https://doi.org/10.3847/0004-637X/822/2/80)
- Strandet, M. L., Weiss, A., De Breuck, C., et al. 2017, *ApJL*, 842, L15, doi: [10.3847/2041-8213/aa74b0](https://doi.org/10.3847/2041-8213/aa74b0)

- Sugahara, Y., Inoue, A. K., Hashimoto, T., et al. 2021, *ApJ*, 923, 5, doi: [10.3847/1538-4357/ac2a36](https://doi.org/10.3847/1538-4357/ac2a36)
- Sun, F., Egami, E., Pérez-González, P. G., et al. 2021a, *ApJ*, 922, 114, doi: [10.3847/1538-4357/ac2578](https://doi.org/10.3847/1538-4357/ac2578)
- Sun, F., Egami, E., Rawle, T. D., et al. 2021b, *ApJ*, 908, 192, doi: [10.3847/1538-4357/abd6e4](https://doi.org/10.3847/1538-4357/abd6e4)
- Swinbank, A. M., Simpson, J. M., Smail, I., et al. 2014, *MNRAS*, 438, 1267, doi: [10.1093/mnras/stt2273](https://doi.org/10.1093/mnras/stt2273)
- Symeonidis, M., Page, M. J., Seymour, N., et al. 2009, *MNRAS*, 397, 1728, doi: [10.1111/j.1365-2966.2009.15040.x](https://doi.org/10.1111/j.1365-2966.2009.15040.x)
- Symeonidis, M., Vaccari, M., Berta, S., et al. 2013, *MNRAS*, 431, 2317, doi: [10.1093/mnras/stt330](https://doi.org/10.1093/mnras/stt330)
- Tacconi, L. J., Genzel, R., Saintonge, A., et al. 2018, *ApJ*, 853, 179, doi: [10.3847/1538-4357/aaa4b4](https://doi.org/10.3847/1538-4357/aaa4b4)
- Tamura, Y., Mawatari, K., Hashimoto, T., et al. 2019, *ApJ*, 874, 27, doi: [10.3847/1538-4357/ab0374](https://doi.org/10.3847/1538-4357/ab0374)
- Treu, T., Schmidt, K. B., Brammer, G. B., et al. 2015, *ApJ*, 812, 114, doi: [10.1088/0004-637X/812/2/114](https://doi.org/10.1088/0004-637X/812/2/114)
- Walth, G. L., Egami, E., Clément, B., et al. 2019, *ApJ*, 877, 7, doi: [10.3847/1538-4357/ab16d8](https://doi.org/10.3847/1538-4357/ab16d8)
- Wang, T., Schreiber, C., Elbaz, D., et al. 2019, *Nature*, 572, 211, doi: [10.1038/s41586-019-1452-4](https://doi.org/10.1038/s41586-019-1452-4)
- Wang, X., Hoag, A., Huang, K. H., et al. 2015, *ApJ*, 811, 29, doi: [10.1088/0004-637X/811/1/29](https://doi.org/10.1088/0004-637X/811/1/29)
- Watson, D., Christensen, L., Knudsen, K. K., et al. 2015, *Nature*, 519, 327, doi: [10.1038/nature14164](https://doi.org/10.1038/nature14164)
- Whitaker, K. E., Pope, A., Cybulski, R., et al. 2017, *ApJ*, 850, 208, doi: [10.3847/1538-4357/aa94ce](https://doi.org/10.3847/1538-4357/aa94ce)
- Williams, C. C., Labbe, I., Spilker, J., et al. 2019, *ApJ*, 884, 154, doi: [10.3847/1538-4357/ab44aa](https://doi.org/10.3847/1538-4357/ab44aa)
- Wold, I. G. B., Owen, F. N., Wang, W.-H., Barger, A. J., & Keenan, R. C. 2012, *ApJS*, 202, 2, doi: [10.1088/0067-0049/202/1/2](https://doi.org/10.1088/0067-0049/202/1/2)
- Wu, J. F., Aguirre, P., Baker, A. J., et al. 2018, *ApJ*, 853, 195, doi: [10.3847/1538-4357/aaa0dc](https://doi.org/10.3847/1538-4357/aaa0dc)
- Yamaguchi, Y., Kohno, K., Hatsukade, B., et al. 2019, *ApJ*, 878, 73, doi: [10.3847/1538-4357/ab0d22](https://doi.org/10.3847/1538-4357/ab0d22)
- . 2020, *PASJ*, 72, 69, doi: [10.1093/pasj/psaa057](https://doi.org/10.1093/pasj/psaa057)
- Yang, C., Omont, A., Beelen, A., et al. 2017, *A&A*, 608, A144, doi: [10.1051/0004-6361/201731391](https://doi.org/10.1051/0004-6361/201731391)
- Yun, M. S., Scott, K. S., Guo, Y., et al. 2012, *MNRAS*, 420, 957, doi: [10.1111/j.1365-2966.2011.19898.x](https://doi.org/10.1111/j.1365-2966.2011.19898.x)
- Zavala, J. A., Casey, C. M., Manning, S. M., et al. 2021, *ApJ*, 909, 165, doi: [10.3847/1538-4357/abdb27](https://doi.org/10.3847/1538-4357/abdb27)
- Zitrin, A., Meneghetti, M., Umetsu, K., et al. 2013, *ApJL*, 762, L30, doi: [10.1088/2041-8205/762/2/L30](https://doi.org/10.1088/2041-8205/762/2/L30)
- Zitrin, A., Fabris, A., Merten, J., et al. 2015, *ApJ*, 801, 44, doi: [10.1088/0004-637X/801/1/44](https://doi.org/10.1088/0004-637X/801/1/44)