

# Decomposition of enveloping algebras of simple Lie algebras and their related polynomial algebras

Rutwig Campoamor-Stursberg<sup>1\*</sup>; Ian Marquette<sup>2 †</sup>

<sup>1</sup> Instituto de Matemática Interdisciplinar and Dpto. Geometría y Topología, UCM, E-28040 Madrid, Spain

<sup>2</sup> School of Mathematics and Physics, The University of Queensland Brisbane, QLD 4072, Australia

April 26, 2022

## Abstract

The decomposition problem of the enveloping algebra of a simple Lie algebra is reconsidered combining both the analytical and the algebraic approach, showing its relation with the internal labelling problem with respect to a nilpotent subalgebra. A lower bound for the number of generators of the commutant as well as the maximal Abelian subalgebra are obtained. The decomposition problem for the exceptional Lie algebra  $G_2$  is completely solved.

## 1 Introduction

Decomposition theorems for enveloping algebras of (simple) Lie algebras constitute a classical result within the structure theory, and have found extensive application in representation theory [1]. For physical problems, these techniques have passed unnoticed for a long time, until the systematic study of superintegrable and quasi-solvable systems has shown how the detailed analysis of enveloping algebras, as well as its analytical counterpart in terms of differential operators can be of much use in solving these problems [2–5]. This has motivated various approaches to polynomial algebras, both from the perspective of rings of functions as well as from the purely algebraic formalism (see [6–10] and references therein). Polynomial algebras have attracted attention on their own right, in particular, the quadratic case [8], showing to what extent different properties of Lie algebras, such as the Jacobi identity or the Poincaré–Birkhoff–Witt bases are preserved, providing a new scope for building other types of algebraic structures. Polynomial algebras of this type have also found applications within the context of Gröbner bases and adjacent areas [11], the generalization of space-time conformal algebras [12], as well as the description of symmetries and orthogonal polynomials related to quantum models [4, 5]. This shows the diversity of problems in which polynomial algebras emerge naturally. Here, we intend to point out the connection between polynomial algebras and some classical problems of simple Lie algebras and their enveloping algebra.

In this work we reconsider the decomposition problem of enveloping algebras of simple Lie algebras combining the purely algebraic approach with an analytical interpretation. More precisely, we show that finding the commutant  $C_{\mathcal{U}(\mathfrak{s})}(P)$  of the highest weight vector for the adjoint representation is computationally equivalent to determine the subgroup scalars for a distinguished nilpotent Lie algebra, namely the nilradical  $\mathfrak{n}$  of the Borel subalgebra of  $\mathfrak{s}$ . This allows us to estimate a lower bound for the number of polynomials in the enveloping algebra required to span the commutant, as well as to determine the dimension of its maximal Abelian subalgebra. The case of the three classical rank-two simple Lie algebras, known from the literature (see [13, 14] and references therein) are revisited from this perspective. Further, we solve the decomposition problem for the exceptional Lie algebra  $G_2$ , which is considerably involved from the computational point of view. It is shown that the polynomials giving rise to the decomposition of the enveloping algebra determine a 17-dimensional polynomial algebra admitting several non-Abelian subalgebras. In the general context, it is shown that the commutant  $C_{\mathcal{U}(\mathfrak{s})}(P)$  always admits a non-Abelian polynomial subalgebra  $\mathcal{A}$  spanned by polynomials forming an integrity

---

\*rutwig@ucm.es

†i.marquette@uq.edu.au

basis for the system of partial differential equations (PDEs) associated to the reduction chain  $\mathfrak{n} \subset \mathfrak{s}$ . We finish with some comments on how the procedure can be extended to reduction chains of semisimple Lie algebras not related to the decomposition of enveloping algebras.

## 2 Presentation of semisimple Lie algebras

One of the most convenient descriptions of complex semisimple Lie algebras is given in terms of the so-called Serre relations, that allow to reconstruct the Lie algebra from its root system and the Dynkin diagram [15].

Let  $\mathfrak{s}$  be a complex semisimple Lie algebra of rank  $\ell$ ,  $\{H_1, \dots, H_\ell\}$  a basis of a Cartan subalgebra  $\mathfrak{h}$ ,  $\mathcal{R}$  the corresponding root system and  $\Delta = \{\alpha_1, \dots, \alpha_\ell\}$  a basis of simple roots. Recall that the Cartan integers are thus given by

$$n(i, j) = \langle \alpha_j, \alpha_i \rangle = \frac{2(\alpha_j, \alpha_i)}{(\alpha_i, \alpha_i)},$$

where  $(\alpha_i, \alpha_j) = \kappa(H_i, H_j)$  is defined in terms of the nondegenerate Killing form  $\kappa$ . Then we can always find  $3\ell$  generators  $H_i, X_i, Y_i$  satisfying the relations

$$\begin{aligned} [H_i, H_j] &= 0, & [H_i, X_j] &= n(i, j)X_j, & [H_i, Y_j] &= -n(i, j)Y_j, \\ [X_i, Y_j] &= \delta_i^j H_i, & \text{ad}(X_i)^{1-n(i, j)} X_j &= 0, & \text{ad}(Y_i)^{1-n(i, j)} Y_j &= 0 \quad (i \neq j). \end{aligned} \quad (1)$$

Clearly, the generators  $X_i$  are associated to the simple roots  $\alpha_i$ , while the  $Y_i$  correspond to the opposite roots (see e.g. [15]).

To any linear form  $\alpha \in \mathfrak{h}^*$  we associate the weight space  $\mathfrak{s}_\alpha$  defined by

$$\mathfrak{s}_\alpha = \{X \in \mathfrak{s} \mid [H, X] = \alpha(H)X, \quad H \in \mathfrak{h}\},$$

Clearly  $\mathfrak{h} = \mathfrak{s}^0$  and the  $\alpha$  such that  $\mathfrak{s}^\alpha \neq 0$  correspond to the roots in  $\mathcal{R}$ . In terms of the weight spaces, the Lie algebra  $\mathfrak{s}$  can be decomposed as

$$\mathfrak{s} = \mathfrak{h} \oplus \sum_{\alpha \in \mathcal{R}} \mathfrak{s}_\alpha. \quad (2)$$

This actually induces a grading in  $\mathfrak{s}$ , as for any  $\alpha_i, \alpha_j \in \mathcal{R}$  the relations (1) imply that  $[\mathfrak{s}^{\alpha_i}, \mathfrak{s}^{\alpha_j}] \subset \mathfrak{s}^{\alpha_i + \alpha_j}$ . Let thus  $\mathcal{U}(\mathfrak{s})$  be the universal enveloping algebra of  $\mathfrak{s}$ . For any fixed positive integer  $p$ , we denote by  $\mathcal{U}_{(p)}(\mathfrak{g})$  the subspace generated by the monomials  $X_1^{a_1} \dots X_n^{a_n}$  satisfying the constraint  $a_1 + a_2 + \dots + a_n \leq p$ , where  $n = \dim \mathfrak{s}$ . Using this relation, we say that an element  $P \in \mathcal{U}(\mathfrak{s})$  is of degree  $d$  if  $d = \inf \{k \mid P \in \mathcal{U}_{(k)}(\mathfrak{s})\}$ . As the enveloping algebra is naturally filtered, we have the relation

$$\mathcal{U}_{(0)}(\mathfrak{s}) = \mathbb{C}, \quad \mathcal{U}_{(p)}(\mathfrak{s})\mathcal{U}_{(q)}(\mathfrak{s}) \subset \mathcal{U}_{(p+q)}(\mathfrak{s}), \quad p, q \geq 0. \quad (3)$$

A direct consequence of the filtration is that each  $\mathcal{U}_{(p)}(\mathfrak{s})$  is a finite-dimensional representation of  $\mathfrak{s}$ , hence the enveloping algebra  $\mathcal{U}(\mathfrak{s})$  is a sum of finite-dimensional representations of the simple Lie algebra  $\mathfrak{s}$ .

From the general structure theory (see e.g. [16, 17]) it is easily seen that the adjoint action of  $\mathfrak{s}$  on  $\mathcal{U}(\mathfrak{s})$  (respectively, the symmetric algebra  $S(\mathfrak{s})$ ) is given by

$$\begin{aligned} P \in \mathcal{U}(\mathfrak{s}) &\mapsto P.X_i := [X_i, P] = X_i P - P X_i \in \mathcal{U}(\mathfrak{s}), \\ P(x_1, \dots, x_n) \in S(\mathfrak{g}) &\mapsto \widehat{X}_i(P) = C_{ij}^k x_k \frac{\partial}{\partial x_j} \in S(\mathfrak{g}), \end{aligned} \quad (4)$$

where  $\{X_1, \dots, X_n\}$  is a basis of  $\mathfrak{s}$ . A linear isomorphism  $\Lambda : S(\mathfrak{s}) \rightarrow \mathcal{U}(\mathfrak{s})$  that commutes with the adjoint action is easily obtained through the symmetrization map

$$\Lambda(x_{j_1} \dots x_{j_p}) = \frac{1}{p!} \sum_{\sigma \in \mathbf{S}_p} X_{j_{\sigma(1)}} \dots X_{j_{\sigma(p)}}, \quad (5)$$

where  $\mathbf{S}(p)$  denotes the symmetric group of order  $p!$ . For the space  $S^{(p)}(\mathfrak{s})$  of homogeneous polynomials having degree  $p$ , the relation  $\mathcal{U}^{(p)}(\mathfrak{s}) = \Lambda(S^{(p)}(\mathfrak{s}))$  further allows us to decompose the subspace  $\mathcal{U}_{(p)}(\mathfrak{g})$  as a direct sum of homogeneous elements:  $\mathcal{U}_{(p)}(\mathfrak{s}) = \sum_{k=0}^p \mathcal{U}^{(k)}(\mathfrak{s})$ . Thus for any  $P \in \mathcal{U}_{(p)}(\mathfrak{s}), Q \in \mathcal{U}_{(q)}(\mathfrak{s})$  the commutator satisfies

$$[P, Q] \in \mathcal{U}_{(p+q-1)}(\mathfrak{s}).$$

Further, as a direct consequence of the Poincaré–Birkhoff–Witt theorem, we have the dimension formula

$$\dim \mathcal{U}^{(p)}(\mathfrak{s}) = \dim \frac{\mathcal{U}_{(p)}(\mathfrak{s})}{\mathcal{U}_{(p-1)}(\mathfrak{s})} = \dim S^{(p)}(\mathfrak{s}) = \binom{\dim \mathfrak{s} + p - 1}{p}. \quad (6)$$

Invariant polynomials of  $\mathfrak{s}$  are defined as the centre of  $\mathcal{U}(\mathfrak{s})$ :

$$Z(\mathcal{U}(\mathfrak{s})) = \{P \in \mathcal{U}(\mathfrak{s}) \mid [\mathfrak{s}, P] = 0\}. \quad (7)$$

These elements can be identified with the (polynomial) solutions of the differential operators in (4) (see [16, 17] for details).

The commutant  $\mathcal{C}_{\mathcal{U}(\mathfrak{s})}(P)$  of an element  $P \in \mathcal{U}(\mathfrak{s})$  is defined as the centralizer of  $P$  in the enveloping algebra, that is, the set of elements in  $\mathcal{U}(\mathfrak{g})$  that commute with  $P$

$$\mathcal{C}_{\mathcal{U}(\mathfrak{s})}(P) = \{Q \in \mathcal{U}(\mathfrak{s}) \mid [P, Q] = 0\}. \quad (8)$$

As  $\mathfrak{s}$  is semisimple, the commutant is finitely generated, a property that allows us to find an integrity basis for it.

## 2.1 Decomposition of enveloping algebras and commutants

As mentioned before, the adjoint action (4) implies that the universal enveloping algebra  $\mathcal{U}(\mathfrak{s})$  can be seen as a representation of  $\mathfrak{s}$ , which is moreover completely reducible whenever the Lie algebra is semisimple [16]. We therefore have a decomposition

$$\mathcal{U}(\mathfrak{s}) = \bigoplus_{k=0}^{\infty} V_k, \quad (9)$$

where  $V_k = \mathcal{U}(\mathfrak{s}).v_{\lambda_k}$  is an irreducible component spanned by a highest vector  $v_{\lambda_k}$  with weight  $\lambda_k \in \mathfrak{h}^*$  for the extended adjoint action, i.e.,

$$[H_i, v_{\lambda_k}] = \lambda_k(H_i)v_{\lambda_k}, \quad H_i \in \mathfrak{h}, \quad [X_i, v_{\lambda_k}] = 0, \quad X_i \in \mathfrak{s}_{\alpha_i}, \quad \alpha_i \in \Delta.$$

Therefore, finding the components in the decomposition (9) amounts to compute the elements in the enveloping algebra  $\mathcal{U}(\mathfrak{s})$  that commute with the generators associated to the positive roots of  $\mathfrak{s}$ . Due to the Serre relations (1), it suffices to consider the elements in  $\mathcal{U}(\mathfrak{s})$  that commute with the generators  $X_i$  associated to simple roots  $\alpha_i \in \Delta$ , as any other elements of  $\mathfrak{n}$  are obtained by commutators. In this context, we observe that the generators  $X_i$  associated to simple roots generate a nilpotent Lie algebra  $\mathfrak{n}$  (see equation (1)).<sup>1</sup>, so that the decomposition problem corresponds to find the centralizer of  $\mathfrak{n}$  in  $\mathcal{U}(\mathfrak{s})$ :

$$\mathcal{C}_{\mathcal{U}(\mathfrak{s})}(\mathfrak{n}) = \{P \in \mathcal{U}(\mathfrak{s}) \mid [X, P] = 0, \quad X \in \mathfrak{n}\} \quad (10)$$

By the algebraic properties of enveloping algebras, we can always find a finite number of polynomials  $\{P_1, \dots, P_s\}$  such that, as vector space,  $\mathcal{C}_{\mathcal{U}(\mathfrak{s})}(\mathfrak{n})$  is spanned by

$$P_1^{a_1} P_2^{a_2} \dots P_s^{a_s}, \quad a_i \in \mathbb{N} \cup 0, \quad (11)$$

and where the coefficients  $a_i$  are possibly constrained by some relation. It should be observed that these polynomials are not necessarily algebraically independent, but merely linearly independent, and that they generate a non-Abelian polynomial algebra. This implies in practice that suitable dependent polynomials must be added to a functionally independent set of polynomials in order to obtain a basis of the type (11). With respect to a given Cartan subalgebra  $\mathfrak{h}$ , if the weight of each  $P_i$  is given by  $(\lambda_1^i, \dots, \lambda_\ell^i)$ , the weight of an element (11) is given by

$$\omega_i = \sum_{i=1}^s a_i (\lambda_1^i, \dots, \lambda_\ell^i). \quad (12)$$

Any element  $P \in \mathcal{C}_{\mathcal{U}(\mathfrak{s})}(\mathfrak{n})$  can thus be labeled using its degree  $d$  as polynomial and its weight  $\mu = (\lambda_1, \dots, \lambda_\ell)$  with respect to the (fixed) Cartan subalgebra. For given values of  $a_i$ , the element in (11) generates an irreducible representation of  $\mathfrak{s}$  of dimension  $d_{a_1, \dots, a_s}$ . In order to fulfill the decomposition (9), the identity

$$\dim \mathcal{U}^{(p)} = \sum_{(a_1, \dots, a_s) \in \Phi_p} d_{a_1, \dots, a_s}, \quad \Phi_p = \left\{ (a_1, \dots, a_s) \mid \sum_{k=1}^s a_k \deg P_k = p \right\} \quad \text{mod } \mathcal{S}, \quad p \geq 1 \quad (13)$$

<sup>1</sup>This algebra actually corresponds to the nilradical  $\mathfrak{n}$  of the Borel subalgebra  $\mathfrak{b}(\Delta)$  of  $\mathfrak{s}$  (the maximal solvable subalgebra of  $\mathfrak{s}$ )

must be satisfied, where  $\mathcal{S}$  denotes the set of relations that constrain the coefficients  $a_i$ .

As the commutant  $\mathcal{C}_{\mathcal{U}(\mathfrak{s})}(\mathfrak{n})$  involves generators that do not belong to the subalgebra  $\mathfrak{n}$ , in addition to the invariants of  $\mathfrak{n}$  there will be additional polynomials depending on the variables of  $\mathfrak{s}$ , in particular the Casimir operators of the simple Lie algebra  $\mathfrak{s}$ . We can suppose without loss of generalization that for the generic element (11), the (primitive) Casimir operators  $I_1, \dots, I_\ell$  of  $\mathfrak{s}$  satisfy

$$P_{s+1-\ell} = I_1, \dots, P_s = I_\ell. \quad (14)$$

Defining the set

$$\mathcal{B} = \{P_1^{a_1} P_2^{a_2} \dots P_{s-\ell}^{a_{s-\ell}}, \quad a_i \in \mathbb{N} \cup 0\}, \quad (15)$$

it follows easily that  $\mathcal{B}$  forms a basis of the commutant  $\mathcal{C}_{\mathcal{U}(\mathfrak{s})}(\mathfrak{n})$  seen as a free module over  $\mathbb{C}[I_1, \dots, I_\ell]$ . This is an immediate consequence of the Schur lemma, as the Casimir operators act as scalar matrices on each irreducible  $\mathfrak{s}$ -representation. As the invariants of  $\mathfrak{s}$  have zero highest weight (with respect to a Cartan subalgebra), the highest weight of the vectors in (11) are determined by the polynomials which have a nonvanishing commutator with at least one generator of  $\mathfrak{s}$  not belonging to the nilpotent subalgebra  $\mathfrak{n}$ . In particular, the generator  $X_\beta$  corresponding to the highest root  $\beta \in \mathcal{R}^+$  always has a nonzero weight, as it lies in the closure of the Weyl chamber. The Casimir operators of  $\mathfrak{n}$ , as they do not commute with all generators of  $\mathfrak{s}$ , also have a nonvanishing weight. In terms of representation theory, this implies that for any finite-dimensional irreducible complex representation  $\varphi : \mathfrak{s} \rightarrow \text{End}(V)$ , the elements  $\varphi(P)$  with  $P \in \mathcal{B}$  span the space of highest weight vectors in the representation [13].

### 3 Analytic properties of the commutant

Alternatively to the analysis of the universal enveloping algebra  $\mathcal{U}(\mathfrak{s})$ , we can use the differential operators in (4) to compute the elements in the commutant analytically. As  $\mathfrak{n}$  is a subalgebra, this correspond to the so-called labelling problem with respect to the chain  $\mathfrak{n} \subset \mathfrak{s}$  (see [18–20]).

Starting from the embedding  $\mathfrak{n} \subset \mathfrak{s}$ , we extend an arbitrary basis  $\{X_1, \dots, X_m\}$  of the subalgebra  $\mathfrak{n}$  to a basis  $\{X_1, \dots, X_m, Y_1, \dots, Y_{n-m}\}$  of  $\mathfrak{s}$ , such that the commutators of the subalgebra generators are given by

$$[X_i, X_j] = C_{ij}^k X_k, \quad [X_i, Y_p] = D_{ip}^k X_k + E_{ip}^q Y_q \quad (16)$$

where  $i, j, k \in \{1, \dots, m\}$  and  $p, q \in \{1, \dots, n-m\}$ . The operators that commute with the elements in  $\mathfrak{n}$  hence correspond to the solutions of the system of PDEs

$$\widehat{X}_i = -C_{ij}^k x_k \frac{\partial}{\partial x_j} - (D_{ip}^k x_k + E_{ip}^q y_q) \frac{\partial}{\partial y_p}, \quad 1 \leq i \leq m. \quad (17)$$

where  $\{x_1, \dots, x_n, y_1, \dots, y_{n-m}\}$  are the coordinates in a dual basis of  $\mathfrak{B}$ . We observe that solutions  $F$  to the system (17) such that  $\frac{\partial F}{\partial y_p} = 0$  for all  $p \in \{1, \dots, n-m\}$  correspond to the Casimir invariants of the subalgebra, while a genuine subgroup scalar must explicitly depend on the variables  $\{y_1, \dots, y_{n-m}\}$  [20]. Now the system (17) has exactly  $n - r'$  independent solutions, where  $r'$  denotes the rank of the  $m \times n$  coefficient matrix of  $\mathfrak{n}$  as subalgebra of  $\mathfrak{s}$ . Although an arbitrary integrity basis for the system (17) does not necessarily generate a polynomial algebra, so that the analytical approach provides at best a lower bound  $d_0$  on the number of required polynomials to obtain the decomposition (9), it may be asked whether for any simple Lie algebra there exists at least an integrity basis for the system (17) such that forms a polynomial algebra. We observe that starting from such a subalgebra, the commutant  $\mathcal{C}_{\mathcal{U}(\mathfrak{s})}(\mathfrak{n})$  can formally be constructed, adding those elements that are algebraically dependent (but linearly independent) required to satisfy the dimension condition (13).

The following table gives the lower bound  $d_0$  obtained analytically for the complex simple Lie algebras:

Type	$A_\ell$	$B_\ell, C_\ell$	$D_\ell$	$G_2$	$F_4$	$E_6$	$E_7$	$E_8$	(18)
$d_0$	$\frac{\ell(\ell+3)}{2}$	$\ell^2 + \ell$	$\ell^2$	8	28	42	70	128	

Among these polynomials, the invariants of  $\mathfrak{n}$  are included. As the subalgebra is nilpotent, we can always find a maximal set of functionally independent solutions formed by polynomials, i.e., an integrity basis [21]. In particular, its centre is generated by the generator associated to the highest root in  $\mathcal{R}$ , and the commutant contains exactly one linear polynomial. From these solutions,  $\ell + \mathcal{N}(\mathfrak{n})$  correspond to the Casimir invariants of either  $\mathfrak{s}$  or  $\mathfrak{n}$ ,<sup>2</sup> so that the number of available operators is given by  $\chi = n - r' - \ell - \mathcal{N}(\mathfrak{n})$ . It can be easily

<sup>2</sup>Here we use the fact that the nilradical of the Borel subalgebra and  $\mathfrak{s}$  can never have common Casimir invariants.

shown (see e.g. [19]) that  $m = r'$ , which implies that  $\chi = 2n_0$ . It should however be noted that among these  $2n_0$  solutions, at most  $n_0$  correspond to operators that commute with each other [19]. We conclude from this that the maximal number of operators that commute with the subalgebra  $\mathfrak{n}$  and with each other is given by

$$\xi = n_0 + \ell + \mathcal{N}(\mathfrak{n}) = \frac{\dim \mathfrak{s} - \dim \mathfrak{n} + \ell + \mathcal{N}(\mathfrak{n})}{2}. \quad (19)$$

The remaining independent solutions of system (17) necessarily have some nontrivial commutator. Expressed in other words, we conclude that the commutant  $\mathcal{C}_{\mathcal{U}(\mathfrak{s})}(\mathfrak{n})$  always contains a maximal Abelian subalgebra of dimension  $\xi$ .

As the generators of  $\mathfrak{n}$  correspond to the positive roots of  $\mathfrak{s}$ , we can easily deduce the number of Casimir invariants of  $\mathfrak{n}$  making use of the so-called Maurer–Cartan equations of the subalgebra [22]. From (2) we know that  $\mathfrak{n} = \sum_{\alpha \in \mathcal{R}^+} s_\alpha$ . Let  $\omega_\alpha$  denote the invariant 1-form corresponding to the generator of the weight space  $s_\alpha$ . For any positive root, the Maurer–Cartan equations can be written formally as

$$d\omega_\alpha = \sum_{\beta, \gamma \in \mathcal{R}^+} \lambda^{\beta, \gamma} \omega_\beta \wedge \omega_\gamma, \quad (20)$$

where  $\lambda^{\beta, \gamma} = 0$  whenever  $\beta + \gamma \neq \alpha$ . In particular, if  $\alpha_i \in \Delta$  is a simple root, it is obvious that the form is closed, i.e.  $d\omega_{\alpha_i} = 0$ , while for non-simple roots the relations  $d\omega_\alpha \neq 0$  holds. We now define a generic 2-form  $\theta = \sum_{\alpha \in \mathcal{R}^+} d\omega_\alpha$  and the index  $j_0(\mathfrak{n})$  as the lowest natural number such that

$$\bigwedge_{i=1}^{j_0(\mathfrak{n})} \theta \neq 0, \quad \bigwedge_{i=1}^{j_0(\mathfrak{n})+1} \theta \equiv 0. \quad (21)$$

Then the number  $\mathcal{N}(\mathfrak{n})$  of invariants is given by  $\mathcal{N}(\mathfrak{n}) = \dim \mathfrak{n} - 2j_0(\mathfrak{n})$  [22].

**Proposition 1** *Let  $\mathfrak{n}$  be the nilradical of the Borel subalgebra of a complex simple Lie algebra  $\mathfrak{s}$ .*

1. *If  $\mathfrak{s} \simeq A_\ell$ ,  $\dim \mathfrak{n} = \frac{\ell(\ell+1)}{2}$  and  $\mathcal{N}(\mathfrak{n}) = \left\lfloor \frac{\ell+1}{2} \right\rfloor$  for  $\ell \geq 1$ .*
2. *If  $\mathfrak{s} \simeq B_\ell$ ,  $\dim \mathfrak{n} = \ell^2$  and  $\mathcal{N}(\mathfrak{n}) = \ell$  for  $\ell \geq 2$ .*
3. *If  $\mathfrak{s} \simeq C_\ell$ ,  $\dim \mathfrak{n} = \ell^2$  and  $\mathcal{N}(\mathfrak{n}) = \ell$  for  $\ell \geq 3$ .*
4. *If  $\mathfrak{s} \simeq D_\ell$ ,  $\dim \mathfrak{n} = \ell^2 - \ell$  and*

$$\mathcal{N}(\mathfrak{n}) = \begin{cases} \ell & \ell = 2p, p \geq 2 \\ \ell - 1 & \ell = 2p + 1, p \geq 2 \end{cases}$$

5. *If  $\mathfrak{s}$  is an exceptional Lie algebra of rank  $\ell$ , then  $\mathcal{N}(\mathfrak{n}) = \ell$ .*

We prove the assertion for the type  $A_\ell$ , the argument being the same for the remaining types of simple classical algebras. The positive roots  $\mathcal{R}^+$  of  $A_\ell$  are given by

$$\beta_{j,k} = \sum_{p=j}^k \alpha_p, \quad j, k = 1, \dots, \ell, \quad j \leq k \quad (22)$$

Let  $\omega_\alpha$  denote the 1-forms. It is clear from the structure of the root system of  $A_\ell$  that the Maurer–Cartan forms have the generic form

$$d\omega_{\beta_{j,k}} = \sum_{s=j}^k \lambda_{j,k}^s \omega_{\beta_{j,s}} \wedge \omega_{\beta_{s+1,k}}, \quad (23)$$

with coefficients  $\lambda_{j,k}^s$  such that  $d(d\omega_{\beta_{j,k}}) = 0$  holds. We separate the analysis according to the parity of  $\ell$ . Let  $\ell = 2p$  and consider the 2-form

$$\theta = d\omega_{\beta_{1,2p}} + \sum_{q=1}^{p-1} \sum_{s=0}^{p-q-\frac{1}{2}} d\omega_{\beta_{2q,2q+2s+1}} + \sum_{q=1}^{p-1} \sum_{s=0}^{p-q-2} d\omega_{\beta_{2q+1,2q+2s+2}} \quad (24)$$

A routine computation shows that the form  $\theta$  satisfies

$$\bigwedge_{i=1}^{p^2} \theta \neq 0, \quad \bigwedge_{i=1}^{p^2+1} \theta \equiv 0, \quad (25)$$

and that adding additional 2-forms  $d\beta_{j,k}$  does not increase the degree of  $\theta$ , thus  $j_0(\mathfrak{n}) = p^2$ . Therefore the number of Casimir invariants of  $\setminus$  is given by

$$\mathcal{N}(\mathfrak{n}) = 2p^2 + p - 2p^2 = p = \left\lceil \frac{2p+1}{2} \right\rceil \quad (26)$$

For odd  $\ell = 2p + 1$ , we consider the 2-form

$$\theta = d\omega_{\beta_{1,2p}} + \sum_{q=1}^{p-1} \sum_{s=0}^{p-q-\frac{1}{2}} d\omega_{\beta_{2q,2q+2s+1}} + \sum_{q=1}^{p-1} \sum_{s=0}^{p-q-1} d\omega_{\beta_{2q+1,2q+2s+2}} \quad (27)$$

satisfying  $\bigwedge^{p^2+p} \theta \neq 0$ ,  $\bigwedge^{p^2+p+1} \theta \equiv 0$ , and the addition of other 2-form does not increase the degree  $j_0(\mathfrak{n}) = p^2 + p$ , from which we deduce that  $\mathcal{N}(\mathfrak{n}) = 2p^2 + 3p + 1 - 2p^2 - 2p = p + 1 = \left\lceil \frac{2p+2}{2} \right\rceil$ .

## 4 Analysis of rank 2 semisimple Lie algebras

### 4.1 $D_2 = \mathfrak{so}(4)$

Let us first consider the semisimple but not simple Lie algebra  $D_2 = A_1^2$ . In this case, the root system is reducible and indeed the disjoint union of the root systems corresponding to  $A_1$ . Choosing a basis adapted to the root system, the commutators are

$$\begin{aligned} [X_1, X_2] &= 2X_2, & [X_1, X_3] &= -2X_3, & [X_2, X_3] &= X_1, \\ [X_4, X_5] &= 2X_5, & [X_4, X_6] &= -2X_6, & [X_5, X_6] &= X_4. \end{aligned} \quad (28)$$

As expected from the fact that  $\mathfrak{n} = \mathbb{R}\langle X_2, X_4 \rangle$  is a decomposable algebra, the commutant is also decomposable. There are four algebraically independent polynomials in  $\mathcal{C}_{\mathcal{U}(\mathfrak{s})}(\mathfrak{n})$ , given by  $X_1, X_4$  themselves, as well as the two Casimir operators  $I_{21} = X_1^2 + 2(X_2X_3 + X_3X_2)$ ,  $I_{22} = X_4^2 + 2(X_5X_6 + X_6X_5)$  of  $D_2$ . These elements form an integrity basis for the solutions of the system (17), and generate the (Abelian) commutant. Hence any element in  $\mathcal{C}_{\mathcal{U}(\mathfrak{s})}(\mathfrak{n})$  can be written as

$$P = X_1^{a_1} X_4^{a_2} I_{21}^{a_3} I_{22}^{a_4}, \quad a_i \in \mathbb{N} \cup 0. \quad (29)$$

The decomposition (9) is straightforward.

### 4.2 $A_2 = \mathfrak{sl}(3, \mathbb{C})$

The decomposition of the enveloping algebra of  $A_2$  has already been considered by various authors, using both analytical and algebraic tools ([13, 14] and references therein). Here we proceed somewhat differently, looking for the algebraic structure of commutants.

Starting from the defining representation  $[1, 0]$ , the Lie algebra  $\mathfrak{sl}(3, \mathbb{C})$  is best given in terms of the elementary matrices  $E_{ij}$  defined by

$$(E_{ij})_{kl} = \delta_i^k \delta_j^l, \quad 1 \leq i, j, k, l \leq 3.$$

and subjected to the constraint  $\text{Tr}(X) = 0$  for each  $X \in \mathfrak{sl}(3, \mathbb{C})$ . Table 1 gives the commutation relations in terms of these generators.

Without loss of generality, we consider the Cartan subalgebra  $\mathfrak{h}$  generated by  $H_1 = E_{11} - E_{22}$  and  $H_2 = E_{22} - E_{33}$ . We observe that, due to this choice,  $E_{12}$  and  $E_{23}$  are the generators associated to the simple roots  $\alpha_1, \alpha_2$  of  $\Delta$ , while  $E_{13}$  is associated to the highest weight  $\alpha_1 + \alpha_2$  of the adjoint representation. Similarly,  $E_{21}, E_{32}$  and  $E_{31}$  correspond to the negative roots  $-\alpha_1, -\alpha_2$  and  $-(\alpha_1 + \alpha_2)$ , respectively. Now  $E_{12}$  and  $E_{23}$  generate a three-dimensional nilpotent algebra  $\mathfrak{n}$ <sup>3</sup>, the centre  $E_{13}$  of which, as an invariant, commutes with all generators of  $\mathfrak{n}$ . There are two other types of elements in the commutant  $\mathcal{C}_{\mathcal{U}(\mathfrak{s})}(\mathfrak{n})$ , namely the (generalized Casimir) invariants of  $\mathfrak{s}$ , that commute with all generators of the Lie algebra, and those polynomials that commute with the subalgebra  $\mathfrak{n}$ , but not with all generators of  $\mathfrak{s}$ , corresponding to subgroup scalars for the coadjoint representation. Using the analytical method (see [19, 20]) it can be easily shown that an integrity basis of functions that commute with  $\mathfrak{n}$  is formed by five independent elements. A direct computation of the

<sup>3</sup>It is indeed isomorphic to the Heisenberg algebra  $\mathfrak{h}_1$ .

Table 1: Commutators of the elementary matrices

$[\circ, \circ]$	$E_{11}$	$E_{22}$	$E_{33}$	$E_{12}$	$E_{13}$	$E_{23}$	$E_{21}$	$E_{31}$	$E_{32}$
$E_{11}$	0	0	0	$E_{12}$	$E_{13}$	0	$-E_{21}$	$-E_{31}$	0
$E_{22}$	0	0	0	$-E_{12}$	0	$E_{23}$	$E_{21}$	0	$-E_{32}$
$E_{33}$	0	0	0	0	$-E_{13}$	$-E_{23}$	0	$E_{31}$	$E_{32}$
$E_{12}$	$-E_{12}$	$E_{12}$	0	0	0	$E_{13}$	$E_{11} - E_{22}$	$-E_{32}$	0
$E_{13}$	$-E_{13}$	0	$E_{13}$	0	0	0	$-E_{23}$	$E_{11} - E_{33}$	$E_{12}$
$E_{23}$	0	$-E_{23}$	$E_{23}$	$-E_{13}$	0	0	0	$E_{21}$	$E_{22} - E_{33}$
$E_{21}$	$E_{21}$	$-E_{21}$	0	$E_{22} - E_{11}$	$E_{23}$	0	0	0	$-E_{31}$
$E_{31}$	$E_{31}$	0	$-E_{31}$	$E_{32}$	$E_{33} - E_{11}$	$-E_{21}$	0	0	0
$E_{32}$	0	$E_{32}$	$-E_{31}$	0	$-E_{12}$	$E_{33} - E_{22}$	$E_{31}$	0	0

polynomials of order at most three in the generators of  $\mathfrak{g}$  shows that there are exactly six linearly independent elements, given respectively by

$$\begin{aligned}
 B_1 &= E_{13}, & B_2 &= 3E_{12}E_{23} + (H_1 - H_2)E_{13}, \\
 C_1 &= E_{32}E_{13}^2 - E_{23}E_{12}^2 + H_2E_{12}E_{13}, & C_2 &= -E_{21}E_{13}^2 + E_{12}E_{23}^2 + H_1E_{23}E_{13}, \\
 I_2 &= \frac{1}{3}(H_1^2 + H_1H_2 + H_2^2) + H_1 + H_2 + E_{21}E_{12} + E_{31}E_{13} + E_{32}E_{23}, \\
 I_3 &= \frac{2}{27}H_1^3 + \frac{1}{9}H_1^2H_2 - \frac{1}{9}H_1H_2^2 - \frac{2}{27}H_2^3 + E_{32}E_{21}E_{13} + E_{31}E_{12}E_{23} - E_{31}E_{13} - E_{32}E_{23} \\
 &\quad + \frac{1}{3}E_{31}(H_1 - H_2)E_{13} + \frac{1}{3}E_{21}(H_1 + 2H_2)E_{12} - \frac{1}{3}E_{32}(2H_1 + H_2)E_{23} + E_{21}E_{12} \\
 &\quad + \frac{1}{3}(H_1^2 - H_2^2) + \frac{1}{3}(H_1 - H_2).
 \end{aligned} \tag{30}$$

We observe that, with respect to the Cartan subalgebra  $\mathfrak{h}$ , the polynomials  $B_1$  and  $B_2$  have weight  $(1,1)$ , while  $C_1$  is of weight  $(3,0)$  and  $C_2$  of weight  $(0,3)$  respectively. The Casimir operators  $I_2$  and  $I_3$  are obviously of weight  $(0,0)$ . As observed, at most five among these six operators are functionally independent, as follows from the algebraic dependence relation

$$C_1C_2 + B_1^3(I_2 + I_3) - \frac{1}{3}B_1^2B_2(I_2 - 2) - \frac{1}{3}B_1B_2^2 + \frac{1}{27}B_2^3 = 0. \tag{31}$$

The set (30) contains an integrity basis. As Casimir operators of  $\mathfrak{g}$ ,  $I_2$  and  $I_3$  generate an Abelian ideal, to which  $B_1, B_2$  can be added, thus exhausting the maximal number of commuting elements within the commutant (see equation (19)). Therefore, the pair  $C_1, C_2$  must lead to nonvanishing commutators. We obtain that  $B_1, B_2, C_1, C_2$  generate a five-dimensional quartic polynomial algebra with nontrivial commutators

$$\begin{aligned}
 [B_2, C_1] &= 3B_1C_1, & [B_2, C_2] &= -3B_1C_2, \\
 [C_1, C_2] &= -B_1^3I_2 - B_1^2B_2 + \frac{1}{3}B_1B_2^2
 \end{aligned} \tag{32}$$

It can be verified (see [13]) that any other polynomial of order  $d \geq 4$  is obtained as an algebraic expression in terms of  $B_1, B_2, C_1, C_2, I_2, I_3$ , showing that any element of the commutant  $\mathcal{C}_{\mathcal{U}(\mathfrak{g})}(\mathfrak{n})$  can be written as

$$P = B_1^{a_1} B_2^{a_2} C_1^{a_3} C_2^{a_4} I_2^{a_5} I_3^{a_6}, \quad a_i \in \mathbb{N} \cup 0 \tag{33}$$

subjected to the constraint  $a_3a_4 = 0$ , as a consequence of equation (31). We also observe that the five elements  $B_1, B_2, C_1, C_2, I_2$  are functionally independent and thus can be chosen as an integrity basis. As follows from the commutators, they also form a polynomial subalgebra  $\mathcal{A}$  of the commutant.

### 4.3 $B_2 = \mathfrak{so}(5, \mathbb{C})$

For the orthogonal Lie algebra  $\mathfrak{so}(5, \mathbb{C})$ , the root system  $R$  has rank two with positive roots

$$R_+ = \{\alpha_1, \alpha_2, \alpha_1 + \alpha_2, 2\alpha_1 + \alpha_2\} \tag{34}$$

Identifying the generators of the positive roots spaces with the elements  $E_1, E_2, E_3, E_4$  and the negative roots with  $F_1, F_2, F_3, F_4$ , as well as the generators  $H_1, H_2$  of the Cartan subalgebra  $\mathfrak{h}$ , the commutators are given in the following table:



Table 2: Commutators of  $\mathfrak{so}(5, \mathbb{C})$ 

$[\circ, \circ]$	$H_1$	$H_2$	$E_1$	$E_2$	$E_3$	$E_4$	$F_1$	$F_2$	$F_3$	$F_4$
$H_1$	0	0	0	$E_2$	$E_3$	$E_4$	0	$-F_2$	$-F_3$	$-F_4$
$H_2$	0	0	$E_1$	$-E_2$	0	$E_4$	$-F_1$	$F_2$	0	$-F_4$
$E_1$	0	0	0	$E_3$	$E_4$	0	$H_2$	0	$-F_2$	$-F_3$
$E_2$	$-E_2$	$E_2$	$-E_3$	0	0	0	0	$H_1 - H_2$	$F_1$	0
$E_3$	$-E_3$	0	$-E_4$	0	0	0	$-E_2$	$E_1$	$H_1$	$F_1$
$E_4$	$-E_4$	$-E_4$	0	0	0	0	$-E_3$	0	$E_1$	$H_1 + H_2$
$F_1$	0	$F_1$	$-H_2$	0	$E_2$	$E_3$	0	$-F_3$	$-F_4$	0
$F_2$	$F_2$	$-F_2$	0	$H_2 - H_1$	$-E_1$	0	$F_3$	0	0	0
$F_3$	$F_3$	0	$F_2$	$-F_1$	$-H_1$	$-E_1$	$F_4$	0	0	0
$F_4$	$F_4$	$F_4$	$F_3$	0	$-F_1$	$-H_1 - H_2$	0	0	0	0

The nilradical  $\mathfrak{n}$  of the Borel subalgebra, generated by  $E_1, E_2, E_3, E_4$ , is isomorphic to the only indecomposable nilpotent Lie algebra of dimension four  $\mathfrak{n}_{4,1}$ , and possesses two Casimir operators that are linear and quadratic in the generators, given by  $P_1 = E_4$  and  $P_2 = E_2 E_4 - \frac{1}{2} E_3^3$  respectively (see [23]). The weight of these polynomials are (1,1) and (2,0) respectively. Using formula (17), it can be easily verified that there are six algebraically independent solutions to the system. In addition to the two previous solutions  $P_1$  and  $P_2$ , up to order four there are three additional polynomials that commute with the subalgebra  $\mathfrak{n}$  but not with the whole orthogonal algebra:

$$\begin{aligned}
 P_3 &= E_3 - E_1 E_2 + E_4 F_1 - H_2 E_3, \\
 P_4 &= -E_4 H_1 + E_4 H_2 - 2E_2 E_1^2 + 2E_3 E_1 H_1 - 2E_3 E_1 H_2 - E_4 H_1^2 + 2E_4 H_2 H_1 \\
 &\quad - E_4 H_2^2 + 2F_2 E_3^2 - 4F_2 E_4 E_2, \\
 P_5 &= -2E_4 E_3 - \frac{1}{2} E_3^2 E_1 + \frac{5}{2} E_4 E_2 E_1 - E_4 E_3 H_1 + F_1 E_4^2 + \frac{1}{2} E_3 E_2 E_1^2 - \frac{1}{2} E_3^2 E_1 H_1 \\
 &\quad + \frac{1}{2} E_3^2 E_1 H_2 + \frac{1}{2} E_4 E_2 E_1 H_1 + \frac{1}{2} E_4 E_2 E_1 H_2 - \frac{1}{2} E_4 E_3 H_2 H_1 + \frac{1}{2} E_4 E_3 H_2^2 - \frac{1}{2} F_1 E_4 E_3 E_1 \\
 &\quad + \frac{1}{2} F_1 E_4^2 H_1 - \frac{1}{2} F_1 E_4^2 H_2 - \frac{1}{2} F_2 E_3^3 + F_2 E_4 E_3 E_2 - \frac{1}{2} F_3 E_4 E_3^2 + F_3 E_4^2 E_2
 \end{aligned} \tag{35}$$

Finally, the two Casimir operators  $C_2$  and  $C_4$  are given by

$$\begin{aligned}
 C_2 &= H_1 + \frac{1}{3} H_2 - \frac{2}{3} E_1 F_1 - \frac{2}{3} E_2 F_2 - \frac{2}{3} E_3 F_3 - \frac{2}{3} E_4 F_4 - \frac{1}{3} H_1^2 - \frac{1}{3} H_2^2, \\
 C_4 &= -H_2 - 2F_1 E_1 - 2F_4 E_4 - \frac{3}{2} H_2 H_1 - H_2^2 - 3F_1 E_1 H_1 + 2F_2 E_2 H_2 - 3F_2 F_1 E_3 \\
 &\quad - 2F_3 E_2 E_1 - F_3 E_3 H_2 - 2F_3 F_1 E_4 - F_4 E_4 H_1 - 2F_4 E_4 H_2 - \frac{1}{2} H_2 H_1^2 - \frac{3}{2} H_1 H_2^2 \\
 &\quad - F_1 E_1 H_1^2 + F_2 E_2 H_2 H_1 - F_2 F_1 E_2 E_1 - F_2 F_1 E_3 H_1 - F_2 F_1 E_3 H_2 + F_2 F_1^2 E_4 - \frac{1}{2} F_2^2 E_2^2 \\
 &\quad - F_3 E_2 E_1 H_1 - F_3 E_2 E_1 H_2 - F_3 E_3 H_2^2 - F_3 F_1 E_4 H_1 + F_3 F_1 E_4 H_2 - F_3 F_2 E_3 E_2 - F_3^2 E_4 E_2 \\
 &\quad + F_4 E_2 E_1^2 - F_4 E_3 E_1 H_1 + F_4 E_3 E_1 H_2 - F_4 E_4 H_2 H_1 - F_4 F_1 E_4 E_1 - F_4 F_2 E_3^2 + F_4 F_2 E_4 E_2 \\
 &\quad - F_4 F_3 E_4 E_3 - \frac{1}{2} F_4^2 E_4^2 - \frac{1}{2} H_1^2 H_2^2.
 \end{aligned} \tag{36}$$

These operators exhaust the number of linearly independent elements in the commutant, and thus generate  $\mathcal{C}_{\mathcal{U}(\mathfrak{s})}(\mathfrak{n})$  (see also [24] for a derivation using the Poisson bracket formalism). Among these seven operators, only six are algebraically independent, as follows from the relation (see [24])

$$P_5^2 - P_2 P_4 + 2P_1 P_3^2 + P_3^3 P_4 - 4P_1 P_2 + P_1 P_2 (6 - P_4) - P_2 P_4 + P_1^2 P_2 C_4 = 0. \tag{37}$$

The only nonvanishing commutators among these operators are given by

$$\begin{aligned}
 [P_3, P_4] &= -2P_1 P_3 + 4P_5, \quad [P_4, P_5] = 2P_1^2 P_3 + P_1 P_3 P_4 - 4P_1 P_5, \\
 [P_3, P_5] &= -P_1 P_2 + \frac{3}{2} P_1 P_2 C_2 - \frac{1}{2} P_1 P_3^2 - \frac{1}{2} P_2 P_4.
 \end{aligned} \tag{38}$$

It follows that any element  $P$  in the commutant can be written as

$$P = P_1^{a_1} P_2^{a_2} P_3^{a_3} P_4^{a_4} P_5^{a_5} C_2^{a_6} C_4^{a_7}, \quad a_i \in \mathbb{N} \cup 0 \tag{39}$$



subjected to the constraint  $a_5 = 0, 1$  due to relation (37).

We observe that, according to formula (19), at most five of the commutant generators commute with each other. In particular, the polynomials  $\{P_1, \dots, P_5, C_2\}$ , which can be taken as an integrity basis for the system (17), further generate a polynomial (cubic) subalgebra of  $\mathcal{C}_{\mathcal{U}(\mathfrak{s})}(\mathfrak{n})$ .

## 5 The exceptional Lie algebra $G_2$

The last of the (reduced) root systems of rank two corresponds to the 14-dimensional exceptional Lie algebra  $G_2$ , with the set of positive roots given by

$$R_+ = \{\alpha_1, \alpha_2, \alpha_1 + \alpha_2, 2\alpha_1 + \alpha_2, 3\alpha_1 + \alpha_2, 3\alpha_1 + 2\alpha_2\} \quad (40)$$

Let  $X_1, X_2$  denote the generators of the Cartan subalgebra and  $X_3, X_5, X_7, X_9, X_{11}, X_{13}$  be the generators associated to the positive roots, with  $X_4, X_6, X_8, X_{10}, X_{12}, X_{14}$  being those corresponding to the negative roots. Over this basis, the commutators are given by

Table 3: Commutators of  $G_2$

[,]	$X_3$	$X_4$	$X_5$	$X_6$	$X_7$	$X_8$	$X_9$	$X_{10}$	$X_{11}$	$X_{12}$	$X_{13}$	$X_{14}$
$X_1$	$2X_3$	$-2X_4$	$-3X_5$	$3X_6$	$-X_7$	$X_8$	$X_9$	$-X_{10}$	$3X_{11}$	$-3X_{12}$	0	0
$X_2$	$-X_3$	$X_4$	$2X_5$	$-2X_6$	$X_7$	$-X_8$	0	0	$-X_{11}$	$X_{12}$	$X_{13}$	$-X_{14}$
$X_3$	0	$X_1$	$X_7$	0	$2X_9$	$-3X_6$	$-3X_{11}$	$-2X_8$	0	$X_{10}$	0	0
$X_4$		0	0	$-X_8$	$3X_5$	$-2X_{10}$	$2X_7$	$3X_{12}$	$-X_9$	0	0	0
$X_5$			0	$X_2$	0	$X_4$	0	0	$-X_{13}$	0	0	$X_{12}$
$X_6$				0	$-X_3$	0	0	0	0	$X_{14}$	$-X_{11}$	0
$X_7$					0	$X_1 + 3X_2$	$-3X_{13}$	$2X_4$	0	0	0	$X_{10}$
$X_8$						0	$-2X_3$	$3X_{14}$	0	0	$-X_9$	0
$X_9$							0	$2X_1 + 3X_2$	0	$-X_4$	0	$-X_8$
$X_{10}$								0	$X_3$	0	$X_7$	0
$X_{11}$									0	$X_1 + X_2$	0	$-X_6$
$X_{12}$										0	$X_5$	0
$X_{13}$											0	$X_1 + 2X_2$
$X_{14}$												0

The nilradical  $\mathfrak{n}$  of the Borel subalgebra of  $G_2$ , generated by  $X_3, X_5$ , is of dimension six and isomorphic to the nilpotent Lie algebra  $\mathfrak{n}_{6,19}$  listed in [23]. It possesses two Casimir operators of degrees one and two given by  $Q_1 = X_{13}$  and  $Q_2 = X_9^2 - 3X_3X_{13} + 3X_7X_{11}$ , respectively, and their weight with respect to the Cartan subalgebra  $\mathfrak{h} = \mathbb{R}\langle X_1, X_2 \rangle$  is given by  $(0,1)$  and  $(2,0)$ . The Casimir operators  $C_2 = 5X_1 + 15X_2 + X_1^2 + X_{10}X_9 + 3X_{12}X_{11} + 3X_{14}X_{13} + 3X_2X_1 + 3X_2^2 + X_4X_3 + 3X_6X_5 + X_8X_7$  and  $C_6$  of  $G_2$  have degrees 2 and 6 (the explicit expression of the latter operator is skipped because of its length) and both have weight  $(0,0)$ . These four polynomials commute with each other. As an integrity basis for the system (17) is formed by eight polynomials, four additional independent polynomials must be found. From these eight operators, at most six can commute, as follows from equation (19). We will see that, in this case, the commutant  $\mathcal{C}_{\mathcal{U}(\mathfrak{s})}(\mathfrak{n})$  requires more than twice this number of operators, making the decomposition of the enveloping algebra of  $G_2$  a computationally demanding problem.

For computational simplicity, and due to the dimension and the length of the polynomials commuting with  $X_3, X_5$ , it is convenient to proceed searching for polynomials with a given degree  $d$  and weight  $(\lambda, \mu)$ , that will be denoted by  $O_d^{[\lambda, \mu]}$ . The strategy is to find an integrity basis formed by polynomials of lowest possible order, so that it contains a set of six independent commuting polynomials, and to complete it to a set of linearly independent polynomials.

Up to degree four, the following seven polynomials in the commutant are linearly independent:

- $d = 1$ :  $O_1^{[0,1]} = Q_1$
- $d = 2$ :  $O_2^{[2,0]} = Q_2, O_2^{[0,0]} = C_2,$
- $d = 3$ :

$$O_3^{[3,0]} = 2X_9^3 + 27X_{11}^2X_5 - 27X_{13}X_{11}X_2 - 27X_{13}^2X_6 + 9X_{11}X_9X_7 - 9X_{13}X_9X_3,$$

$$O_3^{[1,0]} = X_{10}X_9^2 + 3(X_{11}X_{10}X_7 + X_{11}X_4X_1 - X_{13}X_{10}X_3) + 9(X_{11}X_4X_2 + X_{11}X_8X_5 + X_{13}X_6X_4) \\ + 3(X_{13}X_8X_1 + X_7X_3X_2 - X_5X_3^2 + X_7^2X_6 + X_9X_2X_1) + X_9X_4X_3 + X_9X_8X_7 + X_9X_1^2 \\ - 9X_9X_6X_5 + 2X_9 + 3X_1X_4 - 6X_{13}X_8 - 6X_7X_3 + X_9X_1.$$

- $d = 4$ :

$$\begin{aligned}
O_4^{[2,0]} = & 2X_9^2 X_{13} X_{14} + 2X_9^2 X_{11} X_{12} - 3X_8^2 X_{13} X_{13} - 2X_7^2 X_8 X_{11} - X_7 X_9 X_{10} X_{11} + 6X_7 X_{11} X_{13} X_{14} \\
& + 6X_5 X_8 X_9 X_{11} + 9X_6 X_{10} X_{13}^2 + X_6 X_7^2 X_9 + 6X_7 X_{11}^2 X_{12} - 4X_5 X_6 X_9^2 - 3X_5 X_6 X_7 X_{11} - 3X_4^2 X_{11}^2 \\
& + X_3 X_9 X_{10} X_{13} + 6X_4 X_6 X_9 X_{13} - 6X_4 X_8 X_{11} X_{13} - 9X_5 X_{10} X_{11} X_{11} + 2X_3 X_7 X_8 X_{13} - X_2^2 X_9^2 \\
& + 2X_3^2 X_4 X_{13} - X_3^2 X_5 X_9 - 2X_3 X_4 X_7 X_{11} + 3X_3 X_5 X_6 X_{13} - 6X_3 X_{11} X_{12} X_{13} - 6X_3 X_{13}^2 X_{14} \\
& + 9X_2 X_3 X_5 X_{11} + X_2 X_3 X_7 X_9 - 9X_2 X_6 X_7 X_{13} - 6X_2 X_8 X_9 X_{13} - 3X_2^2 X_7 X_{11} - 2X_1 X_8 X_9 X_{13} \\
& - X_1^2 X_3 X_{13} + X_1^2 X_7 X_{11} - 6X_1 X_2 X_3 X_{13} + 6X_1 X_3 X_5 X_{11} - 2X_1 X_4 X_9 X_{11} - 6X_1 X_6 X_7 X_{13} \\
& + (9X_2 X_{10} X_{11} - 6X_2^2 X_3 + 5X_1 X_3) X_{13} - 11X_1 X_7 X_{11} - 6X_{10} X_{11} X_{13} + 6X_2 X_3 X_{13} - 4X_1 X_9^2 \\
& + (12X_3 X_5 - 15X_2 X_7) X_{11} + 2X_3 X_7 X_9 - 2X_4 X_9 X_{11} - 3X_6 X_7 X_{13} - 5X_2 X_9^2 - 5X_7 X_{11} + \frac{4}{3} X_9^2.
\end{aligned}$$

$$\begin{aligned}
O_4^{[0,2]} = & X_7^2 X_9^2 + 4X_7^3 X_{11} - 18X_5 X_7 X_9 X_{11} - 18X_1 X_5 X_{11} X_{13} - 2X_1 X_7 X_9 X_{13} - 12X_3 X_4 X_{13}^2 - 4X_5 X_9^3 \\
& + X_3 (12X_5 X_9 - 4X_7^2) X_{13} + 12X_4 X_7 X_{11} X_{13} + 4X_4 X_9^2 X_{13} - 3X_1^2 X_{13}^2 - 27X_5^2 X_{11}^2 - 6X_1 X_{13}^2 \\
& - 81X_5 X_{11} X_{13} - 3X_7 X_9 X_{13}.
\end{aligned}$$

These polynomials, jointly with the Casimir operator  $O_6^{[0,0]} = C_6$  of  $G_2$ , are algebraically independent and can thus be considered as an integrity basis  $\mathcal{I}$  of the system (17) associated to the nilpotent Lie algebra  $\mathfrak{n}$ . Starting from this set  $\mathcal{I}$ , we analyze the commutators in order to obtain a basis for the commutant. To this extent, we relabel  $C_2$  as  $Q_3$  and the cubic and quartic polynomials as

$$Q_4 = O_3^{[3,0]}, \quad Q_5 = O_3^{[1,0]}, \quad Q_6 = O_4^{[2,0]}, \quad Q_7 = O_4^{[0,2]}.$$

It is immediate to verify that  $Q_1, \dots, Q_5$  form an Abelian algebra. Adding the sixth-order Casimir operator of  $G_2$ , we obtain the maximal number of operators in  $\mathcal{C}_{\mathcal{U}(s)}(\mathfrak{n})$  that commute with each other (see equation (19)). Considering now the commutator of  $Q_4$  and  $Q_7$ , we observe that it decomposes as

$$[Q_4, Q_7] = -54Q_1 O_5^{[3,1]}, \quad (41)$$

where

$$\begin{aligned}
O_5^{[3,1]} = & 2X_8 X_9^2 X_{13}^2 + 6X_7 X_8 X_{11} X_{13}^2 + X_7^2 X_9^2 X_{11} + 4X_7^3 X_{11}^2 + 3X_6 X_7 X_9 X_{13}^2 - 4X_5 X_9^3 X_{11} - 18X_5 X_7 X_9 X_{11}^2 \\
& + 27X_5 X_6 X_{11} X_{13}^2 - 27X_5^2 X_{11}^3 + 2X_4 X_9^2 X_{11} X_{13} + 6X_4 X_7 X_{11}^2 X_{13} - 6X_3 X_8 X_{13}^3 - X_3 X_7 X_9^2 X_{13} \\
& - 8X_3 X_7^2 X_{11} X_{13} + 15X_3 X_5 X_9 X_{11} X_{13} - 6X_3 X_4 X_{11} X_{13}^2 + 4X_3^2 X_7 X_{13}^2 + X_2 (2X_9^3 + 9X_7 X_9 X_{11}) X_{13} \\
& + 27X_2 X_5 X_{11}^2 X_{13} - 6X_2 X_3 X_9 X_{13}^2 - X_1 X_7 X_9 X_{11} X_{13} + (9X_1 X_6 X_{13}^2 - 9X_1 X_5 X_{11}^2) X_{13} + \frac{2}{3} X_9^3 X_{13} \\
& + (X_1 X_3 X_9 X_{13} + 9X_1 X_2 X_{11} X_{13} - 7X_3 X_9 X_{13} + 24X_{11} X_{13} - 72X_5 X_{11}^2) X_{13}.
\end{aligned}$$

This fifth-order polynomial is linearly independent of  $\mathcal{I}$ , hence must be added to the set. Two further fifth-order operators are obtained by successive commutators:<sup>4</sup>

$$[Q_5, Q_7] = 27Q_1 O_5^{[1,1]}, \quad [Q_5, O_5^{[1,1]}] = Q_1 Q_2 Q_3^2 - Q_1 Q_5^2 - 6Q_2 O_5^{[0,1]}, \quad (42)$$

The operators in  $\mathcal{J}_1 = \mathcal{I} \cup \{O_5^{[0,1]}, O_5^{[1,1]}, O_5^{[3,1]}\}$  are linearly independent, and exhaust the polynomials of order  $d \leq 5$  having this property. We label them as  $Q_8 = O_5^{[0,1]}$ ,  $Q_9 = O_5^{[1,1]}$ ,  $Q_{10} = O_5^{[3,1]}$ . The commutator  $[Q_5, Q_6] = 27O_6^{[3,0]}$  provides a linearly independent operator of order six that we label as  $Q_{11}$ . The commutator  $[Q_6, Q_7] = 12Q_1 O_6^{[2,1]}$  gives rise to an operator  $Q_{12} = O_6^{[2,1]}$  that is also independent on the previous eleven polynomials. A routine computation shows that  $Q_{11}, Q_{12}$  and  $Q_{13} = C_6$  are the only sixth-order polynomials linearly independent of the elements in  $\mathcal{J}_1$ . An operator of seventh-order not expressible in terms of the previous elements is obtained from the commutator

$$[Q_5, Q_8] = 6O_7^{[1,1]}. \quad (43)$$

We label this polynomial as  $Q_{14}$ . Two additional linearly independent operators are obtained from the commutators of the latter as

$$[Q_8, Q_{14}] = -\frac{4}{9} Q_3^2 Q_5 Q_7 - \frac{1}{6} Q_1^2 Q_5 Q_{13} - \frac{4}{3} Q_5 O_8^{[0,2]} - \frac{2}{9} Q_3 O_9^{[1,2]} \quad (44)$$

<sup>4</sup>For these as well as the following operators, we omit their explicit expression due to their length.

We denote them as  $Q_{15} = O_8^{[0,2]}$  and  $Q_{16} = O_9^{[1,2]}$  respectively. Finally, the commutator  $[Q_8, Q_{15}] = 4O_{12}^{[0,3]}$  provides a polynomial of order twelve that is linearly independent of  $\{Q_1, \dots, Q_{16}\}$ , and that we denote as  $Q_{17}$ . Analyzing higher orders does not lead to new polynomials that are linearly independent of the former. Any other commutator among these operators can hence be expressed in terms of  $\{Q_1, \dots, Q_{17}\}$ . Excluding the commutators above, the remaining non-vanishing commutators are given by<sup>5</sup>

$$\begin{aligned}
[Q_4, Q_8] &= -27Q_1Q_{11}, & [Q_4, Q_9] &= 3Q_1(2Q_2Q_6 - Q_4Q_5 - 6Q_2^2Q_3), & [Q_4, Q_{10}] &= Q_1(Q_4^2 - 4Q_2^3), \\
[Q_4, Q_{12}] &= 3Q_1(Q_4Q_6 - 2Q_2^2Q_5 - 3Q_2Q_3Q_4), & [Q_4, Q_{14}] &= Q_1(Q_2Q_5^2 - 3Q_2^2Q_3^2 + Q_3Q_4Q_5 + 4Q_2Q_3Q_6), \\
[Q_4, Q_{15}] &= 3Q_1(Q_6Q_9 - 3Q_2Q_3Q_9 + Q_5Q_{17}), & [Q_4, Q_{16}] &= 18Q_1(Q_6Q_{12} - Q_2Q_3Q_9) - 243Q_1^2Q_5Q_{11} \\
&\quad - 27Q_1(Q_3Q_4Q_9 - 3Q_3Q_5Q_{10}), \\
[Q_4, Q_{17}] &= \frac{9}{2}Q_1^3(2Q_3^2Q_5Q_6 - Q_3^4Q_4 + Q_2Q_5(Q_{13} - 9Q_3^3) - Q_3Q_5^3) - \frac{3}{4}Q_1\left(Q_5^3Q_7 + 3Q_4Q_8^2 - \frac{1}{2}Q_4Q_7Q_{13}\right) \\
&\quad + \frac{27}{4}Q_1\left(Q_5Q_9^2 - Q_2Q_3^2Q_5 + Q_3Q_9Q_{12} + \frac{3}{2}Q_4Q_8^2\right) + \frac{27}{2}Q_1^2\left(Q_3^2Q_4Q_8 - Q_5Q_6Q_8 + \frac{9}{2}Q_2Q_3Q_5Q_8\right), \\
[Q_5, Q_{10}] &= \frac{1}{3}Q_1(Q_4Q_5 - 2Q_2Q_6 + 6Q_2^2Q_3), & [Q_5, Q_{11}] &= \frac{2}{9}(Q_6^2 - Q_3Q_4Q_5 + 2Q_2^2Q_3^2) - \frac{2}{9}Q_2(Q_5^2 - 4Q_3Q_6), \\
[Q_5, Q_{12}] &= 2Q_1(Q_3^2Q_4 + 3Q_2Q_3Q_5 - Q_5Q_6) - 3Q_4Q_8, \\
[Q_5, Q_{14}] &= \frac{1}{3}Q_1Q_2(8Q_3^3 - Q_{13}) + \frac{4}{3}Q_1Q_3(Q_5^2 - Q_3Q_6) + 2Q_6Q_8 - 4Q_2Q_3Q_8, & [Q_5, Q_{15}] &= (2Q_1Q_3^2 - 3Q_8)Q_9, \\
[Q_5, Q_{16}] &= 12Q_1Q_3(Q_3Q_{12} - Q_5Q_9) + 72Q_1Q_5Q_{14} - 18Q_8Q_{12}, \\
[Q_5, Q_{17}] &= -\frac{9}{2}Q_1^3Q_3^4Q_5 + \frac{27}{2}Q_1^2Q_3^2Q_5Q_8 - Q_1\left(Q_3^2Q_{16} + \frac{9}{2}Q_3Q_9Q_{15} - \frac{1}{8}Q_5Q_7Q_{13} + \frac{81}{8}Q_5Q_8^2 + 3Q_3^3Q_5Q_7\right) \\
&\quad + \frac{9}{2}Q_3Q_5Q_7Q_8 + \frac{3}{2}Q_8Q_{16}, & [Q_6, Q_8] &= -2Q_5Q_9 - 2Q_3Q_{12}, \\
[Q_6, Q_9] &= 2Q_1(Q_3^2Q_4 + 9Q_2Q_3Q_5 - 2Q_5Q_6) - 3Q_4Q_8, & [Q_6, Q_{10}] &= \frac{2}{3}Q_1(Q_4Q_6 - 2Q_2^2Q_5 - 3Q_2Q_3Q_4), \\
[Q_6, Q_{11}] &= \frac{4}{9}Q_2\left(Q_5Q_6 - 4Q_2Q_3Q_5 - \frac{2}{3}Q_3^2Q_4\right) + \frac{2}{9}Q_4(Q_3Q_6 - Q_5^2), & [Q_6, Q_{12}] &= 2Q_1Q_6^2 - 6Q_2^2Q_8 + 2Q_1 \times \\
&\quad (2Q_2^2Q_3^2 - 3Q_2Q_5^2 - 3Q_2Q_3Q_6), & [Q_6, Q_{15}] &= 2Q_1(Q_3^2Q_{12} + 3Q_3Q_5Q_9 + 9Q_5Q_{14}) - 3Q_8Q_{12}, \\
[Q_6, Q_{14}] &= \frac{5}{3}Q_1\left(Q_2Q_3^2Q_5 + Q_5^3 - Q_3Q_5Q_6 - \frac{2}{5}Q_3^3Q_4 + \frac{1}{10}Q_4Q_{13}\right) + Q_3Q_4Q_8 + 2Q_2Q_5Q_8, \\
[Q_6, Q_{16}] &= 6Q_1\left((2Q_2Q_3^2 - Q_5^2 + 2Q_3Q_6)Q_9 - 9Q_3^3Q_{10} - 81Q_8Q_{11}\right) + 9Q_1(10Q_3Q_5Q_{12} - Q_{10}Q_{13}) + 324Q_1^2 \times \\
&\quad Q_3^2Q_{11} + 81Q_3Q_8Q_{10} - 18Q_2Q_8Q_9, & [Q_6, Q_{17}] &= 3Q_1^3\left(Q_3^4Q_6 - 3Q_2Q_3^5 + 6Q_3^3Q_5^2 - \frac{1}{2}Q_5^2Q_{13}\right) \\
&\quad + 9Q_1^2(Q_3^2Q_6 + 3Q_2Q_3^3Q_8 - 3Q_3Q_5^2Q_8) - \frac{3}{4}Q_5^2Q_7Q_8 + \frac{9}{4}Q_8Q_9^2 + \frac{27}{4}Q_1(3Q_2Q_3Q_8^2 - Q_6Q_8^2) \\
&\quad + 5Q_1Q_3^2Q_5Q_7 + 3Q_1(Q_3^2Q_9^2 - 3Q_5^2Q_{15} - 9Q_3Q_9Q_{14}) + \frac{1}{4}Q_1(Q_6Q_7Q_{13} - 3Q_2Q_3Q_7Q_{13}), \\
[Q_7, Q_9] &= 2Q_1Q_5Q_7, & [Q_7, Q_{10}] &= 2Q_1(8Q_1^2Q_2Q_5 - Q_4Q_7), & [Q_7, Q_{11}] &= -2Q_1Q_4Q_8 + \frac{4}{3}Q_1^2 \times \\
&\quad (Q_3^2Q_4 + 4Q_2Q_3Q_5 - Q_5Q_6), & [Q_7, Q_{12}] &= 12Q_1^2Q_2(2Q_1Q_3^2 - 3Q_8) + 4Q_1(3Q_2Q_3 - Q_6)Q_7, \\
[Q_7, Q_{14}] &= -\frac{2}{3}Q_1(Q_{16} + 4Q_3Q_5Q_7) - 6Q_1^3Q_3^2Q_5, & [Q_7, Q_{16}] &= 3Q_1(8Q_3Q_7 + 18Q_1^2Q_3^2)Q_9 + 18Q_1Q_7Q_{14} \\
&\quad - 81Q_1^2Q_8, & [Q_8, Q_9] &= 9Q_1Q_5Q_8 - \frac{8}{3}Q_3Q_5Q_7 - 6Q_1^2Q_3^2Q_5 - \frac{2}{3}Q_{16}, \\
[Q_8, Q_9] &= \frac{2}{3}Q_1^2(Q_3^2Q_4 - Q_5Q_6 + 4Q_2Q_3Q_5) - Q_1Q_4Q_8, & [Q_8, Q_{11}] &= \frac{8}{9}Q_1\left(Q_2Q_3^2 + \frac{1}{3}Q_5^2\right)Q_5 + \frac{1}{27}Q_1 \times \\
&\quad Q_4Q_{13} - \frac{4}{9}Q_1Q_3Q_5Q_6, & [Q_8, Q_{12}] &= 4Q_1^2Q_2Q_3^3 - 6Q_1Q_2Q_3Q_8 - Q_9^2 + (2Q_2Q_3^2 + \frac{1}{3}Q_5^2 - \frac{2}{3}Q_2Q_6)Q_7, \\
[Q_8, Q_{16}] &= 18Q_1^2Q_3^2(Q_3Q_9 + 3Q_{14}) + 4Q_3^2Q_7Q_9 - 27Q_1(Q_3Q_8Q_9 - 3Q_8Q_{14}) + 30Q_3Q_7Q_{14} - \frac{3}{2}Q_1^2Q_9Q_{13},
\end{aligned}$$

<sup>5</sup>Due to simplicity, we give the commutators and relations in its symmetric form.

$$\begin{aligned}
[Q_8, Q_{17}] &= 2Q_1^4 Q_3^6 - 9Q_1^3 Q_3^4 Q_8 + Q_1^2 \left( Q_3^5 Q_7 + \frac{27}{2} Q_3^2 Q_8^2 + \frac{1}{6} Q_3^2 Q_7 - 12Q_3^3 Q_{15} + Q_{13} Q_{15} \right) - \frac{27}{4} Q_1 Q_8^3 \\
&\quad - 3Q_1 Q_3 (Q_3^2 Q_7 - 6Q_8 Q_{15}) + \frac{1}{12} Q_3 Q_7 (Q_7 Q_{13} + 27Q_8^2 - 36Q_3 Q_{15}) - \frac{1}{4} Q_1 Q_4 Q_7 Q_{13} + 6Q_{15}^2, \\
[Q_9, Q_{10}] &= -Q_1 (Q_5 Q_{10} + Q_4 Q_9), \quad [Q_9, Q_{11}] = 3Q_1 Q_5 Q_{11} + \frac{1}{3} Q_1 (Q_4 Q_9 + 5Q_5 Q_{10}) + \frac{2}{9} (Q_2 Q_5 Q_9 - Q_6 Q_{12}), \\
[Q_9, Q_{12}] &= 3Q_1 ((3Q_2 Q_3 - Q_6) Q_9 - 2Q_3^2 Q_{10}) + 9Q_8 Q_{10}, \quad [Q_9, Q_{14}] = \frac{4}{3} Q_1 Q_3 (Q_5 Q_9 + Q_3 Q_{12}) + Q_1 Q_5 Q_{14} \\
&\quad - 2Q_8 Q_{12}, \quad [Q_9, Q_{15}] = Q_5 Q_7 Q_8 - \frac{2}{3} Q_1 Q_3^2 Q_5 Q_7, \\
[Q_9, Q_{16}] &= 9Q_1^3 (4Q_2 Q_3^4 - Q_3^2 Q_5^2) + Q_1 \left( 108Q_2 Q_3^2 + \frac{27}{2} Q_5^2 \right) Q_8 + 81Q_1 (Q_2 Q_8^2 + Q_9 Q_{14}) - 6Q_6 Q_7 Q_8 \\
&\quad + Q_1 (4Q_3^2 Q_6 Q_7 + 3Q_2 Q_7 Q_{13}), \\
[Q_9, Q_{17}] &= \frac{9}{2} Q_1^2 Q_3^2 (Q_1 Q_3^2 - 3Q_8) Q_9 + Q_1 Q_3 \left( Q_3^2 Q_7 Q_9 + 3Q_3 Q_7 Q_{14} - \frac{9}{2} Q_9 Q_{15} \right) - \frac{3}{2} Q_3 Q_7 Q_8 Q_9 - \frac{9}{2} Q_7 Q_8 Q_{14} \\
&\quad + \frac{1}{8} Q_1 (Q_7 Q_9 Q_{13} + 81Q_8^2 Q_9), \quad [Q_{10}, Q_{11}] = -Q_1 Q_4 Q_{11}, \quad [Q_{10}, Q_{12}] = Q_1 (Q_6 - 3Q_2 Q_3) Q_{10} \\
&\quad + 2Q_1 Q_3^2 Q_9, \quad [Q_{10}, Q_{14}] = \frac{3}{2} Q_1^2 Q_5 Q_{11} - \frac{2}{9} Q_1 (Q_6 Q_{12} + 2Q_2 Q_5 Q_9 - 3Q_3 Q_5 Q_{10}), \\
[Q_{10}, Q_{15}] &= -\frac{4}{3} Q_1^3 Q_2 Q_3^2 Q_5 + 2Q_1^2 Q_2 Q_5 Q_8 + Q_1 \left( Q_9 Q_{12} - Q_2 Q_3 Q_5 Q_7 + \frac{1}{3} Q_5 Q_6 Q_7 \right), \\
[Q_{10}, Q_{16}] &= Q_1^3 (7Q_2^2 Q_6 - 4Q_3^2 Q_4 Q_5 + 3Q_2^2 Q_{13} + 3Q_2 Q_3 (4Q_3 Q_6 + 6Q_5^2 - 12Q_2 Q_3^2)) + 6Q_1^2 (Q_4 Q_5 - 3Q_2 Q_6) Q_8 \\
&\quad + 54Q_1^2 Q_2^2 Q_3 Q_8 + Q_1 Q_2 Q_3 (8Q_6 - 9Q_2 Q_3) Q_7 - \frac{1}{3} Q_1 (Q_2 Q_5^2 + 5Q_6^2) Q_7 + Q_1 (6Q_2 Q_9^2 - 5Q_3 Q_4 Q_5 Q_7), \\
[Q_{10}, Q_{17}] &= \frac{3}{2} Q_1^3 Q_3 \left( Q_5^2 + 5Q_2 Q_3^2 - \frac{2}{3} Q_3 Q_6 \right) Q_9 + 2Q_1^2 Q_3^2 \left( Q_5 Q_{12} - \frac{9}{4} Q_3^2 Q_{10} \right) - \frac{1}{2} Q_1^3 Q_2 Q_9 Q_{13} - \frac{3}{4} Q_1 Q_9^3 \\
&\quad + \frac{3}{2} Q_1^2 \left( Q_6 - \frac{15}{2} Q_2 Q_3 \right) Q_8 Q_9 - 3Q_1^2 \left( Q_5 Q_8 Q_{12} - \frac{9}{2} Q_3^2 Q_8 Q_{10} \right) + \frac{3}{4} Q_1 Q_3 (Q_3 Q_5 Q_7 Q_{12} - Q_6 Q_7 Q_9) \\
&\quad + \frac{3}{4} Q_1 (Q_5^2 + 3Q_2 Q_3^2) Q_7 Q_9 - \frac{3}{8} Q_1 (Q_7 Q_9 - 13 + 27Q_8 Q_{10}^2), \\
[Q_{11}, Q_{12}] &= 3Q_1 (Q_6 - 3Q_2 Q_3) Q_{11} + \frac{2}{9} Q_2 (Q_6 - 3Q_2 Q_3) Q_9 - \frac{1}{9} Q_4 Q_5 Q_9 + \frac{1}{3} (Q_5^2 + 6Q_2 Q_3^2 - 2Q_3 Q_6) Q_{10}, \\
[Q_{11}, Q_{14}] &= \frac{1}{9} Q_3^2 (Q_5 Q_{10} - Q_4 Q_9) + \frac{2}{27} (Q_3 Q_6 - Q_5^2) Q_{12} + \frac{2}{27} (Q_5 Q_6 - 4Q_2 Q_3 Q_5) Q_9, \\
[Q_{11}, Q_{15}] &= \frac{2}{9} Q_1^2 (Q_3^4 Q_4 + 13Q_2 Q_3^3 Q_5 + 6Q_3 Q_5^3 - 4Q_3^2 Q_5 Q_6) - \frac{1}{3} Q_1^2 Q_2 Q_5 Q_{13} - \frac{13}{3} Q_1 Q_2 Q_3 Q_5 Q_8 - \frac{1}{3} Q_5 Q_9^2 \\
&\quad + \frac{2}{3} Q_1 (2Q_5 Q_6 - Q_3^2 Q_4) Q_8 - \frac{1}{9} Q_3 (Q_5 Q_6 Q_7 + 3Q_9 Q_{12}) + \frac{1}{54} (Q_4 Q_{13} + 2Q_5^3) Q_7 + \frac{1}{2} Q_4 Q_8^2 \\
&\quad + \frac{1}{3} Q_2 Q_3^2 Q_5 Q_7, \\
[Q_{11}, Q_{16}] &= \frac{1}{3} Q_1^2 (68Q_2^2 Q_3^4 + 8Q_3^3 Q_4 + 26Q_2 Q_3^2 Q_5^2 + 2(Q_2 Q_6 - Q_4 Q_5) Q_{13} + 10Q_5^4 - 2Q_3 (Q_5^2 + 24Q_2 Q_3^2) Q_6) \\
&\quad + \frac{8}{3} Q_1^2 Q_3^2 Q_6^2 - 2Q_1^2 Q_2^2 Q_3 Q_{13} - 4Q_1 (Q_6^2 + Q_2 Q_5^2 + Q_3 Q_4 Q_5 - 6Q_2 Q_3 Q_6 - 8Q_2^2 Q_3^2) Q_8 + Q_6 Q_9^2 \\
&\quad + 3Q_2^2 \left( 2Q_3^3 Q_7 - Q_8^2 - \frac{1}{27} Q_7 Q_{13} \right) - \frac{1}{9} (Q_5^2 Q_6 - 10Q_3 Q_6^2 + 4Q_2 Q_3 Q_5^2 - 6Q_3^2 Q_4 Q_5 + 48Q_2 Q_3^2 Q_6) Q_7 \\
&\quad - 2Q_9 - 2Q_3 Q_9^2, \\
[Q_{11}, Q_{17}] &= \frac{1}{2} Q_1^2 \left( Q_3^3 Q_6 - \frac{4}{3} Q_3^2 Q_5^2 - \frac{22}{3} Q_2 Q_3^4 + \frac{1}{2} Q_2 Q_3 Q_{13} - \frac{1}{6} Q_6 Q_{13} \right) Q_9 + \frac{27}{2} Q_1^2 Q_3^2 \left( Q_8 Q_{11} - \frac{7}{81} Q_3 Q_5 Q_{12} \right) \\
&\quad + Q_1 Q_3^3 (Q_1 Q_3^2 - 3Q_8) Q_{10} + \frac{1}{6} \left( Q_3^2 Q_6 - \frac{4}{3} Q_3 Q_5^2 - 3Q_2 Q_3^3 \right) Q_7 Q_9 - \frac{1}{6} Q_3^2 Q_5 Q_7 Q_{12} - \frac{9}{2} Q_1^3 Q_3^4 Q_{11} \\
&\quad + \frac{1}{4} Q_1 (7Q_2 Q_3^2 - 3Q_9 - 3Q_6 + 8Q_5^2) Q_8 Q_9 + \frac{3}{2} Q_2 Q_8^2 Q_9 + \frac{9}{4} Q_3 Q_8^2 Q_{10} + \frac{7}{4} Q_1 Q_3 Q_5 Q_8 Q_{12} - \frac{81}{8} Q_1 \times \\
&\quad Q_8^2 Q_{11} + \frac{1}{12} (Q_3 Q_7 + 3Q_1^2 Q_3^2) Q_{10} Q_{13} + \frac{1}{18} Q_2 Q_7 Q_9 Q_{13} - \frac{3}{8} Q_1 Q_8 Q_{10} Q_{13} - Q_1^2 Q_3 Q_5^2 Q_{14} - \frac{1}{6} \times \\
&\quad (Q_5^2 Q_7 - 9Q_9^2 Q_{14}),
\end{aligned}$$

$$\begin{aligned}
[Q_{12}, Q_{14}] &= \frac{9}{2} Q_1^2 Q_3^2 Q_{11} - \frac{1}{6} Q_1 (Q_2 Q_3^2 + 8Q_5^2 - 3Q_3 Q_6) Q_9 - 2Q_2 Q_8 Q_9 + (3Q_3 Q_8 - 2Q_1 Q_3^3) Q_{10} \\
&\quad - \frac{27}{4} Q_1 Q_8 Q_{11} - \frac{1}{2} Q_1 Q_3 Q_5 Q_{12} + \frac{1}{4} Q_1 Q_{10} Q_{13}, \\
[Q_{12}, Q_{15}] &= -2Q_1^3 Q_2 Q_3^4 + 6Q_1^2 Q_2 Q_3^2 Q_8 + 2Q_1 \left( Q_2 Q_3^3 + \frac{3}{4} Q_3 Q_5^2 - \frac{1}{3} Q_3^2 Q_6 \right) Q_7 + Q_6 (Q_6 - 3Q_2 Q_3) Q_7 Q_8 \\
&\quad - \frac{1}{2} Q_1 Q_2 (Q_7 Q_{13} + 9Q_8^2) - \frac{3}{2} Q_1 Q - 3Q - 9^2 - 18Q_1 Q_9 Q_{14}, \\
[Q_{12}, Q_{16}] &= 3Q_1^3 (7Q_3^4 Q_4 + 25Q_2 Q_3^3 Q_5 - 4Q_3^2 Q_5 Q_6 - 3Q_2 Q_5 Q_{13}) + 9Q_1^2 \left( Q_5 Q_6 - 7Q_3^2 Q_4 - \frac{45}{2} Q_2 Q_3 Q_5 \right) Q_8 \\
&\quad + Q_1 \left( 6Q_3^3 Q_4 + \frac{89}{2} Q_2 Q_3^2 Q_5 - \frac{9}{2} Q_5^3 - \frac{27}{2} Q_3 Q_5 Q_6 \right) Q_7 - (9Q_3 Q_4 + 6Q_2 Q_5) Q_7 Q_8 - \frac{9}{2} Q_1 Q_5 Q_9^2 \\
&\quad + \frac{189}{4} Q_1 Q_4 Q_8^2 - \frac{27}{2} Q_1 Q_3 Q_9 Q_{12} + \frac{3}{4} Q_1 Q_4 Q_7 Q_{13}, \\
[Q_{12}, Q_{17}] &= \frac{3}{2} Q_1^3 Q_5 Q_9 Q_{13} - Q_1 Q_3^2 (Q_5 Q_7 + 9Q_1^2 Q_3 Q_5) Q_9 + 9Q_1 (Q_5 Q_9 + Q_3 Q_{12}) Q_{15} + \frac{27}{2} Q_1^2 Q_3 Q_5 Q_8 Q_9 \\
&\quad - \frac{1}{4} Q_1 (Q_7 Q_{12} Q_{13} + 27Q_8^2 Q_{12}) - 3Q_1^2 Q_3^2 (Q_1 Q_3^2 - 3Q_8) Q_{12} + \frac{3}{2} Q_5 Q_7 Q_8 Q_9, \\
[Q_{14}, Q_{15}] &= \frac{2}{3} \left( Q_8 - \frac{2}{3} Q_1 Q_3^2 \right) Q_{16} - Q_1^3 Q_3^4 Q_5 + 3Q_1^2 Q_3^2 Q_5 Q_8 - \frac{10}{9} Q_1 Q_3^2 Q_5 Q_7 + \frac{5}{3} Q_3 Q_5 Q_7 Q_8 - \frac{9}{4} Q_1 \times \\
&\quad Q_5 Q_8^2 + \frac{1}{12} Q_1 Q_5 (Q_7 Q_{13} - 36Q_3 Q_{15}), \\
[Q_{14}, Q_{16}] &= 2Q_1^3 (Q_5^2 - 2Q_2 Q_3^2) Q_{13} - 12Q_1^3 Q_3^3 (Q_5^2 - 3Q_2 Q_3^2) - 13Q_1^3 Q_3^4 Q_6 + 6Q_1^2 Q_2 (Q_{13} - 18Q_3^3) Q_8 \\
&\quad + 18(Q_1^2 Q_3 Q_5^2 - Q_2 Q_3^2 Q_7) Q_8 - \frac{1}{12} Q_1 (Q_6 Q_7 Q_{13} + 7Q_3^2 Q_5^2 Q_7) - \frac{16}{3} Q_1 Q_3^2 Q_6 Q_7 + 6Q_8 Q_9^2 \\
&\quad + 39Q_1^2 Q_3^2 Q_6 Q_8 - \frac{117}{4} Q_1 Q_6 Q_8^2 + 17Q_1 Q_5^2 Q_{15} + 2(Q_5^2 + 4Q_3 Q_6) Q_7 Q_8 + 12Q_1 Q_2 Q_3^4 Q_7 \\
&\quad - Q_1 Q_2 Q_3 (Q_7 Q_{13} - 81Q_8^2) - \frac{7}{4} Q_1 Q_3^2 Q_9^2, \\
[Q_{14}, Q_{17}] &= \frac{1}{8} Q_1^2 (2Q_1 Q_3^2 Q_9 - 3Q_8 Q_9) Q_{13} + \left( \frac{9}{2} Q_1^2 Q_3^2 Q_8 - \frac{3}{2} Q_1^3 Q_3^4 \right) (Q_3 Q_9 + 3Q_{14}) + \frac{9}{2} Q_1 Q_3 Q_{14} Q_{15} \\
&\quad + \frac{1}{2} (Q_1 Q_3^2 - 6Q_8) Q_9 Q_{15} + \frac{3}{2} \left( Q_3 Q_7 - \frac{27}{4} Q_1 Q_8 \right) Q_8 Q_{14} - \frac{1}{8} Q_1 (Q_7 Q_{13} + 8Q_3^3 Q_7) Q_{14} \\
&\quad + \frac{1}{24} Q_1 Q_3 Q_7 Q_9 Q_{13} - \frac{1}{3} Q_1 Q_3^4 Q_7 Q_9 + \frac{1}{2} Q_3^2 Q_7 Q_8 Q_9 - \frac{27}{8} Q_1 Q_3 Q_8^2 Q_9, \\
[Q_{15}, Q_{16}] &= 18Q_1^2 Q_3^2 (Q_1 Q_3^2 - 3Q_8) Q_9 + \left( 2Q_1 Q_3^3 Q_7 - 3Q_3 Q_7 Q_8 + \frac{81}{2} Q_1 Q_8^2 \right) Q_9 - 12Q_1 Q_3^2 Q_7 Q_{14} \\
&\quad + \frac{3}{2} Q_1 Q_7 Q_9 Q_{13} + 18Q_7 Q_8 Q_{14} + 162Q_1 Q_{14} Q_{15}, \\
[Q_{15}, Q_{17}] &= Q_1^3 Q_3^4 (9Q_{15} - Q_3^2 Q_7) + \frac{27}{8} (Q_8 - 2Q_1 Q_3^2) Q_7 Q_8^2 + \frac{1}{4} Q_1 (Q_7 Q_{13} + 81Q_8^2 - 36Q_3 Q - 15) Q_{15} \\
&\quad + \frac{9}{2} Q_1^2 Q_3^4 Q_7 Q_8 + \frac{1}{8} \left( Q_8 - \frac{2}{3} Q_1 Q_3^2 \right) Q_7^2 Q_{13} + 3Q_1 Q_3^2 (Q_3 Q_7 - 9Q_1 Q_8) Q_{15} - \frac{3}{2} Q_3 Q_7 Q_8 Q_{15}, \\
[Q_{16}, Q_{17}] &= 27Q_1^5 Q_3^6 Q_5 - \frac{243}{2} Q_1^4 Q_3^4 Q_5 Q_8 + \frac{729}{4} Q_1^2 \left( Q_1 Q_3^2 Q_5 - \frac{1}{2} Q_5 Q_8 \right) Q_8^2 + \frac{243}{2} Q_1^2 Q_3 Q_5 Q_8 Q_{15} \\
&\quad + 9Q_1 Q_3^4 (Q_5 Q_7 + 4Q_1^2 Q_3 Q_5) Q_7 + 81Q_1 Q_5 (Q_{15}^2 + Q_3 Q_7 Q_8^2) + Q_1^3 Q_3^3 \left( \frac{9}{2} Q_3 Q_{16} - 81Q_5 Q_{15} \right) \\
&\quad - 108Q_1^2 Q_3^3 Q_5 Q_7 Q_8 + \frac{9}{8} Q_1^2 Q_5 \left( Q_7 Q_8 - \frac{2}{3} Q_1 Q_3^2 Q_7 \right) Q_{13} + \frac{27}{2} Q_5 (Q_1^3 Q_{13} Q_{15} - Q_3^2 Q_7^2 Q_8) \\
&\quad + 9Q_5 Q_7 \left( Q_8 - \frac{2}{3} Q_1 Q_3^2 \right) Q_{15} + \frac{9}{2} Q_3 (Q_1 Q_{15} - Q_7 Q_8) Q_{16} - \frac{1}{8} Q_1 (Q_7 Q_{13} - 81Q_8^2) Q_{16} \\
&\quad + 3Q_1 Q_3^2 \left( Q_3 Q_7 - \frac{9}{2} Q_1 Q_8 \right) Q_{16}.
\end{aligned}$$

As only eight among these seventeen polynomials are functionally independent, a certain number of algebraic dependence relations are expected. In contrast to the previous examples, the computation of a complete set of relations for  $G_2$  is rather long and cumbersome, ultimately leading to 57 constraints.

$$\begin{aligned}
& Q_4Q_9 + 3Q_5Q_{10} - 2Q_2Q_{12} = 0, \quad 6Q_2Q_{14} - Q_2Q_3Q_9 + Q_6Q_9 + Q_5Q_{12} = 0, \\
& Q_5^2Q_7 + 3Q_9^2 - 12Q_2Q_{15} = 0, \quad 3Q_1^2Q_5^3 + Q_5Q_6Q_7 - 3Q_4Q_{15} + Q_2Q_{16} = 0, \\
& 6Q_1^2(Q_5^3 - Q_2Q_3^2Q_5) - 3Q_2Q_3Q_5Q_7 + Q_5Q_6Q_7 + 9Q_1Q_2Q_5Q_8 + 3Q_9Q_{12} - 6Q_4Q_{15} = 0, \\
& 6Q_1^2Q_3^2Q_5Q_9 + 2Q_3Q_5Q_7Q_9 - 9Q_1Q_5Q_8Q_9 + 6Q_5Q_7Q_{14} + 6Q_{12}Q_{15} + 12Q_2Q_{17} = 0, \\
& 6Q_1^2Q_3^2Q_5Q_9 + 4Q_3Q_5Q_7Q_9 - 9Q_1Q_5Q_8Q_9 - 6Q_5Q_7Q_{14} - 12Q_{12}Q_{15} + 2Q_9Q_{16} = 0, \\
& 2Q_2^2Q_9 - 9Q_2Q_3Q_{10} + 3Q_6Q_{10} + 27Q_1Q_2Q_{11} - Q_4Q_{12} = 0, \\
& 2Q_2Q_5Q_9 - Q_3Q_4Q_9 - 9Q_3Q_5Q_{10} + 27Q_1Q_5Q_{11} + 2Q_6Q_{12} + 6Q_4Q_{14} = 0, \\
& 3Q_{12}^2 - 6Q_1^2Q_3^2Q_4Q_5 - 3Q_1^2Q_5^2Q_6 - 3Q_3Q_4Q_5Q_7 - Q_2Q_5^2Q_7 + 9Q_1Q_4Q_5Q_8 - Q_4Q_{16} = 0, \\
& 36Q_1^2Q_2^2Q_3^3 - 12Q_1^2Q_2Q_3^2Q_6 + (9Q_2^2Q_3^2 - Q_2Q_5^2 - 6Q_2Q_3Q_6 + Q_6^2)Q_7 + 18Q_1Q_2(Q_6 - 3Q_2Q_3)Q_8 \\
& - 3Q_2Q_9^2 + 3Q_{12}^2 - 3Q_1^2Q_2^2Q_{13} = 0, \\
& 6Q_1^2Q_3^2Q_4Q_5 - 12Q_1^2Q_2^2Q_3^3 + 12Q_1^2Q_2Q_3Q_5^2 + (12Q_1^2Q_2Q_3^2 - 6Q_1^2Q_5^2)Q_6 - 6Q_2^2Q_3^2Q_7 - 2Q_6^2Q_7 + 2Q_2Q_5^2Q_7 \\
& + 2(Q_3Q_4Q_5 + 4Q_2Q_3Q_6)Q_7 + 18Q_1\left(Q_2^2Q_3 - \frac{1}{2}Q_4Q_5 - Q_2Q_6\right)Q_8 + 54Q_{10}Q_{14} = 0, \\
& 12Q_1^2Q_2^2Q_3^2 - 12Q_1^2Q_2Q_5^2 + 6Q_2^2Q_3Q_7 + Q_4Q_5Q_7 - 2Q_2Q_6Q_7 - 18Q_1Q_2^2Q_8 - 9Q_9Q_{10} = 0, \\
& 12Q_1Q_2^2Q_3^3 + 2Q_1Q_3^2Q_4Q_5 + 12Q_1Q_2Q_3Q_5^2 - 4Q_1Q_2Q_3^2Q_6 - 2Q_1Q_5^2Q_6 - 18Q_2^2Q_3Q_8 - 3Q_4Q_5Q_8 \\
& + 6Q_2Q_6Q_8 - 2Q_1Q_2^2Q_{13} + 27Q_9Q_{11} = 0, \\
& 3(Q_3Q_5Q_7 - 3Q_1Q_5Q_8)Q_{12} + 3Q_1^2(2Q_3^2Q_5Q_{12} - Q_3Q_5^2Q_9 + 6Q_5^2Q_{14}) - Q_5^2Q_7Q_9 + Q_{12}Q_{16} + 6Q_4Q_{17} = 0, \\
& 36Q_1^2Q_2Q_3^3Q_9 - 12Q_1^2Q_3^2Q_6Q_9 + 9Q_2Q_3^2Q_7Q_9 - Q_5^2Q_7Q_9 - 5Q_3Q_6Q_7Q_9 - 54Q_1Q_2Q_3Q_8Q_9 - 6Q_6Q_7Q_{14} \\
& + 18Q_1Q_6Q_8Q_9 - 3Q_9^3 + 6Q_1^2Q_3^2Q_5Q_{12} + 3Q_3Q_5Q_7Q_{12} - 9Q_1Q_5Q_8Q_{12} - 3Q_1^2Q_2Q_9Q_{13} + 2Q_{12}Q_{16} = 0, \\
& 6Q_1Q_3^2Q_{10} - 2Q_1Q_6Q_9 - 9Q_8Q_{10} + 9Q_7Q_{11} - 4Q_1Q_5Q_{12} = 0, \\
& 6Q_1^2Q_2Q_3^2Q_9 - 6Q_1^2Q_5^2Q_9 + 3Q_2Q_3Q_7Q_9 - Q_6Q_7Q_9 - 9Q_1Q_2Q_8Q_9 + Q_5Q_7Q_{12} - 18Q_{10}Q_{15} = 0, \\
& 4Q_5^2Q_9 + 4Q_3Q_6Q_9 + 54Q_1Q_3^2Q_{11} - 81Q_8Q_{11} + 12Q_3Q_5Q_{12} - 3Q_{10}Q_{13} + 12Q_6Q_{14} = 0, \\
& 6Q_1Q_2Q_3^3Q_9 + 4Q_1Q_3Q_5^2Q_9 - 2Q_1Q_3^2Q_6Q_9 - 9Q_2Q_3Q_8Q_9 + 3Q_6Q_8Q_9 + 2Q_1Q_3^2Q_5Q_{12} - 3Q_5Q_8Q_{12} \\
& - Q_1Q_2Q_9Q_{13} + 12Q_1Q_5^2Q_{14} + 54Q_{11}Q_{15} = 0, \\
& 12Q_1^2Q_2Q_3^4 - 18Q_1^2Q_3^2Q_5^2 - 9Q_3Q_5^2Q_7 - 36Q_1Q_2Q_3^2Q_8 + 27Q_1Q_5^2Q_8 + 27Q_2Q_8^2 - 3Q_3Q_9^2 + Q_2Q_7Q_{13} \\
& + 18Q_9Q_{14} - 2Q_5Q_{16} = 0, \\
& 12Q_1^2Q_2Q_3^4 - 12Q_1^2Q_3^2Q_5^2 - 6Q_3Q_5^2Q_7 - 36Q_1Q_2Q_3^2Q_8 + 18Q_1Q_5^2Q_8 + 27Q_2Q_8^2 - 6Q_3Q_9^2 + Q_2Q_7Q_{13} \\
& + 36Q_9Q_{14} + 12Q_6Q_{15} = 0, \\
& 12Q_1^2Q_3^4Q_9 - 36Q_1Q_3^2Q_8Q_9 + 27Q_8^2Q_9 + Q_7Q_9Q_{13} - 12Q_3Q_9Q_{15} + 72Q_{14}Q_{15} + 24Q_5Q_{17} = 0, \\
& 36Q_1Q_3^2Q_4Q_8 - 12Q_1^2Q_3^4Q_4 - 144Q_1^2Q_2Q_3^3Q_5 + 48Q_1^2Q_3^2Q_5Q_6 - 36Q_2Q_3^2Q_5Q_7 + 24Q_3Q_5Q_6Q_7 + 12Q_5Q_9^2 \\
& + 216Q_1Q_2Q_3Q_5Q_8 - 72Q_1Q_5Q_6Q_8 - 27Q_4Q_8^2 + 12Q_1^2Q_2Q_5Q_{13} - Q_4Q_7Q_{13} + 4Q_6Q_{16} = 0, \\
& 12Q_1^2Q_3^4Q_4 + 72Q_1^2Q_2Q_3^3Q_5 - 12Q_1^2Q_3^2Q_5Q_6 + 2(Q_5^3 + 9Q_2Q_3^2Q_5 - 3Q_3Q_5Q_6)Q_7 - 36Q_1Q_3^2Q_4Q_8 - 6Q_5Q_9^2 \\
& - 108Q_1Q_2Q_3Q_5Q_8 + 18Q_1Q_5Q_6Q_8 + 27Q_4Q_8^2 - 6Q_3Q_9Q_{12} - 6Q_1^2Q_2Q_5Q_{13} + Q_4Q_7Q_{13} + 36Q_{12}Q_{14} = 0, \\
& 12Q_3Q_5Q_7Q_{14} - (36Q_1^2Q_3^3Q_5 - 8Q_3^2Q_5Q_7 + 54Q_1Q_3Q_5Q_8)Q_9 + 3Q_1^2Q_5Q_9Q_{13} + 36Q_1^2Q_3^2Q_5Q_{14} + 12Q_{14}Q_{16} \\
& - 54Q_1Q_5Q_8Q_{14} + 12Q_5Q_9Q_{15} - 12Q_3Q_{12}Q_{15} - 24Q_6Q_{17} = 0, \\
& 24Q_1^2Q_3^3Q_5Q_9 + 4Q_3^2Q_5Q_7Q_9 - 36Q_1Q_3Q_5Q_8Q_9 + 12Q_1^2Q_3^4Q_{12} - 36Q_1Q_3^2Q_8Q_{12} + (Q_7Q_{12} - 3Q_1^2Q_5Q_9)Q_{13} \\
& + 27Q_8^2Q_{12} + (108Q_1^2Q_3^3Q_5 + 48Q_3Q_5Q_7 - 162Q_1Q_5Q_8)Q_{14} - 12Q_5Q_9Q_{15} - 12Q_3Q_{12}Q_{15} + 12Q_{14}Q_{16} = 0, \\
& 108Q_3Q_5Q_7Q_{15} - 12Q_1^2Q_3^4Q_5Q_7 + 36Q_1Q_3^2Q_5Q_7Q_8 - 27Q_5Q_7Q_8^2 - Q_5Q_7^2Q_{13} + 216Q_1^2Q_3^2Q_5Q_{15} + 24Q_{15}Q_{16} \\
& - 324Q_1Q_5Q_8Q_{15} + 72Q_9Q_{17} = 0, \\
& Q_4^2Q_7 - 12Q_1^2Q_2Q_4Q_5 - 12Q_1^2Q_2^2Q_6 - 4Q_2^3Q_7 + 27Q_{10}^2 = 0, \\
& 8Q_1Q_2^3Q_3^2 - 2Q_1Q_3^2Q_4^2 - 12Q_1Q_2Q_3Q_4Q_5 - 8Q_1Q_2^2Q_5^2 - 12Q_1Q_2^2Q_3Q_6 + 2Q_1Q_4Q_5Q_6 + 4Q_1Q_2Q_6^2 - 12Q_2^3Q_8 \\
& + 3Q_4^2Q_8 + 81Q_{10}Q_{11} = 0, \\
& Q_4Q_6Q_7 - 6Q_1^2Q_2Q_3^2Q_4 - 6Q_1^2Q_2Q_5Q_6 - 3Q_2Q_3Q_4Q_7 - 2Q_2^2Q_5Q_7 + 9Q_1Q_2Q_4Q_8 + 9Q_{10}Q_{12} = 0, \\
& 9Q_3Q_5Q_7Q_{10} - 6Q_1^2Q_3^2Q_4Q_9 - 6Q_1^2Q_5Q_6Q_9 - 3Q_3Q_4Q_7Q_9 - 2Q_2Q_5Q_7Q_9 + 9Q_1Q_4Q_8Q_9 + 18Q_1^2Q_3^2Q_5Q_{10} \\
& - 27Q_1Q_5Q_8Q_{10} + 2Q_6Q_7Q_{12} + 6Q_{10}Q_{16} = 0,
\end{aligned}$$

$$\begin{aligned}
& 12Q_1^2(Q_3^2Q_5Q_6 - Q_3^4Q_4 - 6Q_2Q_3^3Q_5)Q_7 + -18Q_2Q_3^2Q_5Q_7^2 - 2Q_5^3Q_7^2 + 6Q_3Q_5Q_6Q_7^2 + 36Q_1Q_3^2Q_4Q_7Q_8 \\
& + 108Q_1Q_2Q_3Q_5Q_7Q_8 - 18Q_1Q_5Q_6Q_7Q_8 - 27Q_4Q_7Q_8^2 + 36Q_1^2Q_3Q_5Q_9^2 + 18Q_5Q_7Q_9^2 + 36Q_1^2Q_3^2Q_9Q_{12} \\
& + 18Q_3Q_7Q_9Q_{12} - 54Q_1Q_8Q_9Q_{12} + 6Q_1^2Q_2Q_5Q_7Q_{13} - Q_4Q_7^2Q_{13} - 216Q_1^2Q_5Q_9Q_{14} + 216Q_{10}Q_{17} = 0, \\
& 12Q_3Q_4Q_5Q_6 - 36Q_2Q_3^2Q_4Q_5 - 48Q_2^2Q_3Q_5^2 - 4Q_4Q_5^3 - (36Q_2^2Q_3^2 + 12Q_2Q_5^2)Q_6 + 24Q_2Q_3Q_6^2 - 4Q_6^3 + 4Q_2^3Q_{13} \\
& - Q_4^2Q_{13} + 729Q_{11}^2 = 0, \\
& 6Q_1Q_2Q_3^3Q_4 + 4Q_1Q_2^2Q_3^2Q_5 - 4Q_1Q_2Q_5^3 - 2Q_1(Q_3^2Q_4 + 3Q_2Q_3Q_5)Q_6 + 2Q_1Q_5Q_6^2 - 9Q_2Q_3Q_4Q_8 - 6Q_2^2Q_5Q_8 \\
& + 3Q_4Q_6Q_8 - Q_1Q_2Q_4Q_{13} + 27Q_{11}Q_{12} = 0, \\
& 12Q_1Q_2^2Q_3^4 - 4Q_1Q_3^3Q_4Q_5 + 8Q_1Q_2Q_3^2Q_5^2 + 4Q_1Q_5^4 - 16Q_1Q_2Q_3^3Q_6 - 8Q_1Q_3Q_5^2Q_6 + 4Q_1Q_3^2Q_6^2 - 18Q_2^2Q_3^2Q_8 \\
& + 6(Q_3Q_4Q_5 + Q_2Q_5^2 + 4Q_2Q_3Q_6 - 6Q_6^2)Q_8 - 2Q_1Q_2^2Q_3Q_{13} + Q_1(Q_4Q_5 + 2Q_2Q_6)Q_{13} + 162Q_{11}Q_{14} = 0, \\
& 6Q_1Q_3^3Q_4Q_9 + 4Q_1Q_2Q_3^2Q_5Q_9 + 4Q_1Q_5^3Q_9 + 12Q_1Q_3Q_5Q_6Q_9 - 9Q_3Q_4Q_8Q_9 - 6Q_2Q_5Q_8Q_9 + 6Q_6Q_8Q_{12} \\
& - 18Q_1Q_3^3Q_5Q_{10} + 27Q_3Q_5Q_8Q_{10} + 162Q_1^2Q_3^2Q_5Q_{11} - 243Q_1Q_5Q_8Q_{11} + 36Q_1Q_3Q_5^2Q_{12} - 4Q_1Q_3^2Q_6Q_{12} \\
& - Q_1Q_4Q_9Q_{13} - 6Q_1Q_5Q_{10}Q_{13} + 18Q_{11}Q_{16} = 0, \\
& Q_6Q_7Q_{13} - 36Q_1^2Q_3^3Q_5^2 + 12Q_1^2Q_3^4Q_6 - 9Q_3^2Q_5^2Q_7 + Q_1(54Q_3Q_5^2 - 36Q_3^2Q_6)Q_8 + 27Q_6Q_8^2 - 3Q_3^2Q_9^2 + 3Q_1^2Q_5^2Q_{13} \\
& + 36Q_3Q_9Q_{14} + 12Q_5^2Q_{15} - 108Q_{14}^2 = 0, \\
& 24Q_1^3(Q_3^6Q_4 + 6Q_2Q_3^5Q_5 + 6Q_3^3Q_5^3 - 3Q_3^4Q_5Q_6) + Q_1(36Q_2Q_3^4Q_5 + 40Q_3^2Q_5^3 - 12Q_3^3Q_5Q_6)Q_7 - 48Q_1Q_5^3Q_{15} \\
& - 108Q_1^2(Q_3^4Q_4 + 4Q_2Q_3^3Q_5^2Q_3Q_5^3 - 2Q_3^2Q_5^2Q_6)Q_8 - 6(Q_5^3 - 3Q_3Q_5Q_6 + 9Q_2Q_3^2Q_5)Q_7Q_8 + 162Q_1Q_3^2Q_4Q_8^2 \\
& + (324Q_1Q_2Q_3Q_5 - 162Q_1Q_5Q_6)Q_8^2 - 81Q_4Q_8^3 + 54Q_5Q_8Q_9^2 - (36Q_1Q_3^3Q_9 + 54Q_3Q_8Q_9)Q_{12} - 12Q_1^2Q_2Q_3^2Q_5Q_{13} \\
& - 12Q_1^3Q_3^2Q_{13} + 2Q_1Q_3^2Q_4Q_7Q_{13} - 4Q_1Q_5Q_6Q_7Q_{13} + 18Q_1^2Q_2Q_5Q_8Q_{13} - 3Q_4Q_7Q_8Q_{13} + 6Q_1Q_9Q_{12}Q_{13} \\
& - 216Q_1Q_3Q_5Q_9Q_{14} + 648Q_{11}Q_{17} = 0, \\
& 144Q_1^2Q_2(Q_3^5Q_7 + 2Q_1^2Q_3^6) - 324Q_1^2Q_3^3Q_5^2(Q_7 + Q_1^2Q_3) + 324(Q_3Q_7Q_9 - 3Q_1Q_8Q_9)Q_{14} - 36Q_3^2Q_{15} - 9Q_1^2Q_9^2Q_{13} \\
& + 12Q_1^2Q_3^4Q_6Q_7 - 81Q_3^2Q_5^2Q_7^2 + 324Q_1^3(3Q_3^2Q_5^2 - 4Q_2Q_3^4)Q_8 + (486Q_1Q_3Q_5^2 - 432Q_1Q_2Q_3^3)Q_7Q_8 - 729Q_1^2Q_5^2Q_8^2 \\
& - 36Q_1Q_3^2Q_6Q_7Q_8 + 1944Q_1^2Q_2Q_3^2Q_8^2 + 27(Q_6 + 12Q_2Q_3)Q_7Q_8^2 - 972Q_1Q_2Q_8^3 - 27Q_3^2Q_7Q_9^2 + 24Q_1^2Q_2Q_3^2Q_7Q_{13} \\
& + 12Q_2Q_3Q_7^2Q_{13} + Q_6Q_7^2Q_{13} - 36Q_1Q_2Q_7Q_8Q_{13} + 648Q(1^2)Q(3^2)Q_9Q_{14} + 4Q_{16}^2 = 0, \\
& 12Q_1^2Q_3^3(3Q_5^2 - 12Q_3Q_6)Q_7 + 9Q_3^2Q_5^2Q_7^2 - 54Q_1Q_3Q_5^2Q_7Q_8 + 36Q_1Q_3^2Q_6Q_7Q_8 - 27Q_6Q_7Q_8^2 - 9(Q_3^2Q_7 + 8Q_1^2Q_3^3)Q_9^2 \\
& + 108Q_1Q_3Q_8Q_9^2 - 3Q_1^2Q_5^2Q_7Q_{13} - Q_6Q_7^2Q_{13} + 9Q_1^2Q_9^2Q_{13} - 216Q_1^2Q_3^2Q_9Q_{14} - 108Q_3Q_7Q_9Q_{14} + 324Q_1Q_8Q_9Q_{14} \\
& - 12Q_5^2Q_7Q_{15} + 36Q_9^2Q_{15} + 72Q_{12}Q_{17} = 0, \\
& 144Q_1Q_3^3Q_5Q_7Q_8 - 486Q_1^2Q_3^2Q_5Q_8^2 - 108Q_3Q_5Q_7Q_8^2 + 243Q_1Q_5Q_8^3 - 72Q_1^4Q_3^6Q_5 - 48Q_1^2Q_3^5Q_5Q_7 + 324Q_1^3Q_3^4Q_5Q_8 \\
& - 6Q_1^2Q_3^2Q_5Q_7Q_{13} - 4Q_3Q_5Q_7^2Q_{13} + 9Q_1Q_5Q_7Q_8Q_{13} - 216Q_1^2Q_3^3Q_5Q_{15} + 324Q_1Q_3Q_5Q_8Q_{15} + 36Q_1^2Q_5Q_{13}Q_{15} \\
& + 144Q_5Q_{15}^2 - 24Q_1^2Q_3^4Q_{16} + 72Q_1Q_3^2Q_8Q_{16} - 54Q_8^2Q_{16} - 2Q_7Q_{13}Q_{16} + 24Q_3Q_{15}Q_{16} + 432Q_{14}Q_{17} = 0, \\
& 9Q_1Q_7Q_8Q_9Q_{13} + 243Q_1Q_8^3Q_9 - 6Q_1^2Q_3^2Q_7Q_9Q_{13} - 4Q_3Q_7^2Q_9Q_{13} + 72Q_1^2Q_3^4Q_7Q_{14} - 216Q_1Q_3^2Q_7Q_8Q_{14} \\
& - 72Q_1^4Q_3^6Q_9 - 48Q_1^2Q_3^5Q_7Q_9 + (324Q_1^3Q_3^4 + 144Q_1Q_3^3Q_7)Q_8Q_9 - 486Q_1^2Q_3^2Q_8^2Q_9 - 108Q_3Q_7Q_8^2Q_9 + 162Q_7Q_8^2Q_{14} \\
& + 6Q_7^2Q_{13}Q_{14} - 216Q_1^2Q_3^3Q_9Q_{15} + 324Q_1Q_3Q_8Q_9Q_{15} + 36Q_1^2Q_9Q_{13}Q_{15} - 1296Q_1^2Q_3^2Q_{14}Q_{15} - 648Q_3Q_7Q_{14}Q_{15} \\
& + 1944Q_1Q_8Q_{14}Q_{15} + 144Q_9Q_{15}^2 + 48Q_{16}Q_{17} = 0, \\
& 144Q_1^4Q_3^8Q_7 - 864Q_1^3Q_3^6Q_7Q_8 + 1944Q_1^2Q_3^4Q_7Q_8^2 - 1944Q_1Q_3^2Q_7Q_8^3 + 729Q_7Q_8^4 + 24Q_1^2Q_3^4Q_7^2Q_{13} - 72Q_1Q_3^2Q_7^2Q_8Q_{13} \\
& + 54Q_7^2Q_8^2Q_{13} + Q_7^3Q_{13}^2 - 1728Q_1^4Q_3^6Q_{15} - 864Q_1^2Q_3^5Q_7Q_{15} + 7776Q_1^3Q_3^4Q_8Q_{15} + 2592Q_1Q_3^3Q_7Q_8Q_{15} - 1728Q_{15}^3 \\
& - 11664Q_1^2Q_3^2Q_8^2Q_{15} - 1944Q_3Q_7Q_8^2Q_{15} + 5832Q_1Q_8^3Q_{15} - 72(Q_3Q_7^2 + 2Q_1^2Q_3^2Q_7 - 3216Q_1Q_7Q_8)Q_{13}Q_{15} \\
& + 5184Q_1^2Q_3^3Q_{15}^2 + 1296Q_3^2Q_7Q_{15}^2 - 7776Q_1Q_3Q_8Q_{15}^2 - 432Q_1^2Q_{13}Q_{15}^2 + 1728Q_{17}^2 = 0, \\
& - 4Q_3^3Q_9 + Q_4^2Q_9 + 18Q_2^2Q_3Q_{10} + 3Q_4Q_5Q_{10} - 6Q_2Q_6Q_{10} - 54Q_1Q_2^2Q_{11} = 0, \\
& - 6Q_3Q_4Q_5Q_9 - 4Q_2Q_5^2Q_9 - 6Q_2Q_3Q_6Q_9 + 2Q_6^2Q_9 - 18Q_3Q_5^2Q_{10} - 54Q_1Q_2Q_3^2Q_{11} + 81Q_2Q_8Q_{11} \\
& + 2Q_5Q_6Q_{12} + 3Q_2Q_{10}Q_{13} = 0, \\
& 4Q_2Q_5Q_6Q_9 + 108Q_2Q_3^2Q_5Q_{10} - 54Q_3Q_5Q_6Q_{10} - 54Q_1Q_3^2Q_4Q_{11} - 24Q_2^2Q_3Q_5Q_9 - 4Q_4Q_5^2Q_9 - 6Q_3Q_4Q_6Q_9 \\
& - 324Q_1Q_2Q_3Q_5Q_{11} + 54Q_1Q_5Q_6Q_{11} + 81Q_4Q_8Q_{11} + 4Q_6^2Q_{12} + 3Q_4Q_{10}Q_{13} = 0, \\
& 36Q_1Q_2^2Q_3^2Q_8 - 18Q_1Q_2Q_5^2Q_8 - 27Q_2^2Q_8^2 - 12Q_1^2Q_2^2Q_3^4 + 12Q_1^2Q_2Q_3^2Q_5^2 + 6Q_2Q_3Q_5^2Q_7 - Q_5^2Q_6Q_7 + 3Q_6Q_9^2 \\
& - Q_2^2Q_7Q_{13} + 6Q_5Q_9Q_{12} = 0,
\end{aligned}$$



$$\begin{aligned}
& 18Q_2Q_3Q_5Q_6Q_7 - 2Q_5Q_6^2Q_7 + 36Q_1Q_2Q_3^2Q_4Q_8 - 12Q_1^2Q_2Q_3^4Q_4 - 144Q_1^2Q_2^2Q_3^3Q_5 + 36Q_1^2Q_2Q_3^2Q_5Q_6 \\
& - 36Q_1(2^2)Q_1(3^2)Q_5Q_7 + 216Q_1Q_2^2Q_3Q_5Q_8 - 54Q_1Q_2Q_5Q_6Q_8 - 27Q_2Q_4Q_8^2 + 12Q_2Q_5Q_9^2 + 12Q_1^2Q_2^2Q_5Q_{13} \\
& - Q_2Q_4Q_7Q_{13} + 6Q_6Q_9Q_{12} = 0, \\
& 12Q_1^2(Q_3^4Q_4^2 - 4Q_2^3Q_3^4 + 12Q_2Q_3^3Q_4Q_5 + 8Q_2^2Q_3^2Q_5^2 - 4Q_2Q_5^4 + 12Q_2^2Q_3^3Q_6 - 2Q_3^2Q_4Q_5Q_6 - 4Q_2Q_3^2Q_6^2 + Q_5^2Q_6^2) \\
& + 36Q_2Q_3^2Q_4Q_5Q_7 + 48Q_2^2Q_3Q_5^2Q_7 + 4Q_4Q_5^3Q_7 + 36Q_2^2Q_3^2Q_6Q_7 - 12Q_2Q_5^2Q_6Q_7 - 24Q_2Q_3Q_6^2Q_7 + 4Q_6^3Q_7 \\
& + 144Q_1Q_2^3Q_3^2Q_8 - 36Q_1Q_3^3Q_4^2Q_8 - 216Q_1Q_2Q_3Q_4Q_5Q_8 - 144Q_1Q_2^2Q_5^2Q_8 - 216Q_1Q_2^2Q_3Q_6Q_8 + 27Q_4^2Q_8^2 \\
& + 36Q_1Q_4Q_5Q_6Q_8 + 72Q_1Q_2Q_6^2Q_8 - 108Q_2^3Q_8^2 - 12Q_1^2Q_2Q_4Q_5Q_{13} - 12Q_1^2Q_2^2Q_6Q_{13} - 4Q_2^3Q_7Q_{13} + Q_4^2Q_7Q_{13} \\
& - 12Q_3Q_4Q_5Q_6Q_7 = 0, \\
& 12Q_1^2(Q_2^2Q_5^2Q_6 + 12Q_2^2Q_3^3 + 2Q_2^2Q_3^2Q_4Q_5 - Q_2Q_4Q_5^3 - 4Q_2^3Q_3^2Q_6) + 36Q_2^4Q_3^2Q_7 + 12Q_2^2Q_3Q_4Q_5Q_7 + Q_4^2Q_5^2Q_7 \\
& - 24Q_2^3Q_3Q_6Q_7 + 4Q_2^2Q_6^2Q_7 - 216Q_1Q_2^4Q_3Q_8 - 36Q_1Q_2^2Q_4Q_5Q_8 + 72Q_1Q_3^2Q_6Q_8 - 12Q_2^3Q_9^2 - 12Q_1^2Q_2^4Q_{13} \\
& + 3Q_4^2Q_9^2 - 4Q_2Q_4Q_5Q_6Q_7 = 0, \\
& 36Q_2Q_3^2Q_4Q_5Q_9 + 48Q_2^2Q_3Q_5^2Q_9 + 4Q_4Q_5^3Q_9 + 36Q_2^2Q_3^2Q_6Q_9 - 12Q_2Q_5^2Q_6Q_9 - 24Q_2Q_3Q_6^2Q_9 + 324Q_1Q_2^2Q_3^3Q_{11} \\
& + 4Q_6^3Q_9 + 54Q_1Q_3^2Q_4Q_5Q_{11} + 324Q_1Q_2Q_3Q_5^2Q_{11} - 108Q_1Q_2Q_3^2Q_6Q_{11} - 54Q_1Q_5^2Q_6Q_{11} - 486Q_2^2Q_3Q_8Q_{11} \\
& - 81Q_4Q_5Q_8Q_{11} + 162Q_2Q_6Q_8Q_{11} - 4Q_2^3Q_9Q_{13} + Q_4^2Q_9Q_{13} - 54Q_1Q_2^2Q_{11}Q_{13} - 12Q_3Q_4Q_5Q_6Q_9 = 0, \\
& 12Q_1^2Q_2Q_3^4Q_9 - 12Q_1^2Q_3^2Q_5^2Q_9 - 5Q_3Q_5^2Q_7Q_9 - 36Q_1Q_2Q_3^2Q_8Q_9 + 18Q_1Q_5^2Q_8Q_9 + 27Q_2Q_8^2Q_9 + Q_2Q_7Q_9Q_{13} \\
& - 3Q_3Q_9^3 - 6Q_5^2Q_7Q_{14} + 18Q_9^2Q_{14} - 12Q_5Q_{12}Q_{15} = 0, \\
& 12Q_1^2(Q_2^3Q_3^4 - 2Q_2^2Q_3^2Q_5^2 + Q_2Q_5^4) - 12Q_2^2Q_3Q_5^2Q_7 - Q_4Q_5^3Q_7 + 3Q_2Q_5^2Q_6Q_7 - 36Q_1Q_3^2Q_3^2Q_8 + 36Q_1Q_2^2Q_5^2Q_8 \\
& + 27Q_2^3Q_8^2 - 3Q_4Q_5Q_9^2 - 3Q_2Q_6Q_9^2 + Q_2^3Q_7Q_{13} = 0, \\
& 12Q_1^2(Q_2^2Q_3^4Q_4 + 12Q_2^3Q_3^3Q_5 - 2Q_2^2Q_12Q_1^2Q_2Q_3^3Q_6) + (36Q_2^3Q_3^2Q_5 - 24Q_2^2Q_3Q_5Q_6 - Q_4Q_5^2Q_6 + 4Q_2Q_5Q_6^2)Q_7 \\
& + 72Q_1 \left( Q_2^2Q_5Q_6 - \frac{1}{2}Q_2^2Q_3^2Q_4 - 3Q_2^3Q_3Q_5 \right) Q_8 + 27Q_2^2Q_4Q_8^2 - 12Q_2^2Q_5Q_9^2 - 12Q_1^2Q_2^3Q_5Q_{13} - 3Q_4Q_6Q_9^2 \\
& + Q_2^2Q_4Q_7Q_{13} = 0, \\
& 12Q_1^2Q_2Q_3^4Q_4Q_5 + 144Q_1^2Q_2^2Q_3^3Q_5^2 - 12Q_1^2(Q_2^2Q_3^42Q_2Q_3^3Q_5^2)Q_6 + 36Q_2^2Q_3^2Q_5^2Q_7 - 12Q_2Q_3Q_5^2Q_6Q_7 - 27Q_2^2Q_6Q_8^2 \\
& + Q_5^2Q_6^2Q_7 + 36Q_1(Q_2^2Q_3^2Q_6 - Q_2Q_3^2Q_4Q_5 + Q_2Q_5^2Q_6)Q_8 - 216Q_1Q_2^2Q_3Q_5^2Q_8 + 27Q_2Q_4Q_5Q_8^2 - 12Q_2Q_5^2Q_9^2 \\
& + 3Q_6^2Q_9^2 - 12Q_1^2Q_2^2Q_3^2Q_{13} + Q_2Q_4Q_5Q_7Q_{13} - Q_2^2Q_6Q_7Q_{13} = 0, \\
& 48Q_1^2Q_2^2(Q_3^2Q_6^2 + 10Q_2^2Q_3^4) - 12Q_1^2Q_2Q_3^4Q_4^2 - 72Q_1^2Q_2^2Q_3^3Q_4Q_5 - 96Q_1^2Q_2^3Q_3^2Q_5^2 - 36Q_1^2Q_2Q_3Q_4Q_5^3 + 48Q_1^2Q_2^2Q_5^4 \\
& - 288Q_1^2Q_2^3Q_3^3Q_6 + 24Q_1^2Q_2Q_3^2Q_4Q_5Q_6 + 36Q_1^2Q_2^2Q_3Q_5^2Q_6 - 12Q_2^2Q_2Q_5^2Q_6^2 + 108Q_2^4Q_3^3Q_7 - 48Q_2^3Q_3Q_5^2Q_7 \\
& + 3Q_3Q_4^2Q_5^2Q_7 - 4Q_2Q_4Q_5^3Q_7 - 108Q_2^3Q_3^2Q_6Q_7 + 12Q_2^2Q_5^2Q_6Q_7 + 36Q_2^2Q_3Q_6^2Q_7 - 4Q_2Q_6^3Q_7 - 792Q_1Q_4^2Q_3^2Q_8 \\
& + 36Q_1(Q_2Q_3^2Q_4^2 + 3Q_2^2Q_3Q_4Q_5 + 4Q_2^3Q_5^2 + 12Q_2^3Q_3Q_6Q - Q_2Q_4Q_5Q_6 - 2Q_2^2Q_6^2)Q_8 + (108Q_2^4 - 27Q_2Q_4^2)Q_8^2 \\
& - 36Q_2^3Q_3Q_9^2 + 9Q_3Q_4^2Q_9^2 - (36Q_1^2Q_2^4Q_3 + 12Q_1^2Q_2^2Q_4Q_5 + 12Q_1^2Q_2^3Q_6 + 4Q_2^4Q_7 - Q_2Q_4^2Q_7)Q_{13} = 0.
\end{aligned}$$

Any element in the commutant  $\mathcal{C}_{U(\mathfrak{s})}(\mathfrak{n})$  can thus be expressed as the product

$$P = \prod_{i=1}^{17} Q_i^{a_i}, \quad (a_1, \dots, a_{17}) \in \mathcal{S}, \quad (45)$$

where the set  $\mathcal{S}$  of constraints consists of the 53 conditions

$$\begin{aligned}
& a_{10} = 0, 1, \quad a_{11} = 0, 1, \quad a_{12} = 0, 1, \quad a_{14} = 0, 1, \quad a_{16} = 0, 1, \quad a_{17} = 0, 1, \\
& a_2a_{12} = 0, \quad a_5a_{12} = 0, \quad a_2a_{15} = 0, \quad a_2a_{16} = 0, \quad a_2a_{17} = 0, \quad a_4a_{12} = 0, \quad a_4a_{14} = 0, \\
& a_4a_{15} = 0, \quad a_4a_{16} = 0, \quad a_4a_{17} = 0, \quad a_5a_{16} = 0, \quad a_5a_{17} = 0, \quad a_6a_{14} = 0, \quad a_6a_{15} = 0, \\
& a_6a_{16} = 0, \quad a_6a_{17} = 0, \quad a_7a_{11} = 0, \quad a_9a_{10} = 0, \quad a_9a_{11} = 0, \quad a_9a_{16} = 0, \quad a_9a_{17} = 0, \\
& a_{10}a_{11} = 0, \quad a_{10}a_{12} = 0, \quad a_{10}a_{14} = 0, \quad a_{10}a_{15} = 0, \quad a_{10}a_{16} = 0, \quad a_{10}a_{17} = 0, \quad a_{11}a_{12} = 0, \\
& a_{11}a_{14} = 0, \quad a_{11}a_{15} = 0, \quad a_{11}a_{16} = 0, \quad a_{11}a_{17} = 0, \quad a_{12}a_{14} = 0, \quad a_{12}a_{16} = 0, \quad a_{12}a_{17} = 0, \\
& a_{14}a_{16} = 0, \quad a_{14}a_{17} = 0, \quad a_{16}a_{17} = 0, \quad a_2a_6a_{10} = 0, \quad a_2a_{10}a_{13} = 0, \quad a_4a_{10}a_{13} = 0, \\
& a_5a_9a_{12} = 0, \quad a_6a_9a_{12} = 0, \quad a_5a_{12}a_{15} = 0, \quad a_2a_4a_5a_6a_7 = 0, \quad a_3a_4a_5a_6a_7 = 0, \quad a_3a_4a_5a_6a_9 = 0,
\end{aligned}$$

as well as the requirement  $(a_1, \dots, a_{17}) \notin \mathbb{Z}[v_1, v_2, v_3, v_4]$ , where

$$v_1 = (2, 3, 4, 0^{14}), \quad v_2 = (2, 4, 4, 0^{14}), \quad v_3 = (2, 2, 4, 1, 0^{13}), \quad v_4 = (2, 2, 3, 0, 2, 0^{12}). \quad (46)$$

For any admissible  $(a_1, \dots, a_{17}) \in \mathcal{S}$ , the weight  $\Lambda = [\lambda, \mu]$  of the corresponding irreducible representation of  $G_2$  is given by

$$\Lambda = [2a_2 + 3a_4 + a_5 + 2a_6 + a_9 + 3a_{10} + 3a_{11} + 2a_{12} + a_{14} + a_{16}, \\ a_1 + 2a_7 + a_8 + a_9 + a_{10} + a_{12} + a_{14} + 2a_{15} + 2a_{16} + 3a_{17}] \quad (47)$$

with dimension

$$d_{a_1, \dots, a_{17}} = \frac{1}{120} (\lambda + 1) (\mu + 1) (\lambda + \mu + 2) (\lambda + 2\mu + 3) (\lambda + 3\mu + 4) (2\lambda + 3\mu + 5) \quad (48)$$

A routine computation shows that the identity (13) is satisfied for any  $p \geq 2$ . For low values of  $p$ , the terms in the decomposition (9) are explicitly given by:<sup>6</sup>

$$\begin{aligned} p = 2 : \quad \mathcal{U}^{(2)} &= [0, 0] \oplus [1, 0] \oplus [0, 1], \\ p = 3 : \quad \mathcal{U}^{(3)} &= [1, 0] \oplus [0, 1] \oplus [3, 0] \oplus [0, 3] \oplus [2, 1], \\ p = 4 : \quad \mathcal{U}^{(4)} &= [0, 0] \oplus [2, 0]^2 \oplus [0, 2]^2 \oplus [1, 1] \oplus [4, 0] \oplus [0, 4] \oplus [3, 1] \oplus [2, 2], \\ p = 5 : \quad \mathcal{U}^{(5)} &= [1, 0] \oplus [0, 1]^2 \oplus [1, 1] \oplus [1, 2] \oplus [2, 1]^2 \oplus [3, 0]^2 \oplus [0, 3]^2 \oplus [3, 1] \oplus [3, 2] \oplus [2, 3] \\ &\quad \oplus [4, 1] \oplus [5, 0] \oplus [0, 5], \\ p = 6 : \quad \mathcal{U}^{(6)} &= [0, 0]^2 \oplus [2, 0]^3 \oplus [0, 2]^3 \oplus [1, 1] \oplus [3, 0]^2 \oplus [2, 1] \oplus [1, 2] \oplus [4, 0]^3 \oplus [0, 4]^2 \oplus [2, 2]^2 \\ &\quad \oplus [3, 1]^2 \oplus [1, 3] \oplus [3, 2] \oplus [6, 0] \oplus [5, 1] \oplus [4, 2] \oplus [3, 3] \oplus [2, 4] \oplus [0, 6]. \end{aligned}$$

The total number of terms in the decomposition for the values  $1 \leq p \leq 16$  is

$$1, 3, 5, 10, 17, 31, 48, 80, 122, 187, 274, 404, 569, 805, 1106, 1512,$$

while the number of nonequivalent irreducible representations intervening is

$$1, 3, 5, 8, 13, 19, 25, 35, 44, 55, 66, 80, 93, 109, 124, 142.$$

An interesting property of the commutant  $\mathcal{C}_{\mathcal{U}^{(s)}}(\mathfrak{n})$  of  $G_2$  that distinguishes it from previous cases is the existence of several non-Abelian polynomial subalgebras. As commented before, the polynomials  $\{Q_1, \dots, Q_5, Q_{13}\}$  generate a six-dimensional Abelian subalgebra  $\mathcal{A}_0$ . To comment only on some possibilities, we observe that if we extend  $\mathcal{A}_0$  adjoining  $Q_6$ , the commutators show that  $\mathcal{A}_1 = \{Q_1, \dots, Q_6, Q_{11}, Q_{13}\}$  determines an eight-dimensional polynomial algebra. As these eight polynomials are functionally independent, they form an integrity basis for the system (17). The subalgebra  $\mathcal{A}_1$  suffices to describe the decomposition (9) up to  $p = 3$ , and fails for  $p = 4$  due to absence of  $Q_7$ . If we adjoin the latter element, the resulting polynomial algebra is easily seen to be isomorphic to the whole commutant. However, adjoining  $Q_{10}$  to  $\mathcal{A}_1$  leads to a nine-dimensional subalgebra  $\mathcal{A}_2$  that is maximal, as the adjunction of any other element again generates the commutant  $\mathcal{C}_{\mathcal{U}^{(s)}}(\mathfrak{n})$ . On the other hand, the elements  $\{Q_1, Q_2, Q_4, Q_{10}\}$  themselves generate a minimal non-Abelian polynomial algebra.

## 6 Conclusions

The problem of determining the decomposition of the universal enveloping algebra of a simple complex Lie algebra  $\mathfrak{s}$  has been revisited combining both the purely algebraic and the analytical approach. It has been observed that determining the polynomials that commute with the elements associated to the simple roots of  $\mathfrak{s}$  is formally equivalent to computing the subgroup scalars for the reduction chain  $\mathfrak{n} \subset \mathfrak{s}$ , where  $\mathfrak{n}$  is the maximal nilpotent ideal of the Borel subalgebra of  $\mathfrak{s}$ . This fact allows us to deduce that the commutant  $\mathcal{C}_{\mathcal{U}^{(s)}}(\mathfrak{n})$  always possesses a maximal Abelian subalgebra, the dimension of which is completely determined by  $\mathfrak{n}$ . With the analytical approach, the commutant can be constructed successively starting from an integrity basis for the system of PDEs associated to the reduction chain. In this context, the question whether any integrity basis forms a non-Abelian polynomial algebra has been shown to be false in general, although for any of the cases considered, a polynomial subalgebra  $\mathcal{A}$  of the commutant  $\mathcal{C}_{\mathcal{U}^{(s)}}(\mathfrak{n})$  and whose elements form an integrity basis has been shown to exist.

The case of rank two simple Lie algebra has been analyzed in detail. For the classical algebras  $A_2$  and  $B_2$ , that have already been considered in the literature (see [13, 14, 24]), additional properties of the commutant have been described. The case of the exceptional Lie algebra  $G_2$ , which has been solved for the first time, is much more complicated. The first notable difference with the classical algebras is that number of linearly

<sup>6</sup>Conventions on representations are adapted to those used in [25].

polynomials required for the decomposition (9) is more than twice the cardinal of an integrity basis. Besides the computational difficulties due to the degree and length of the polynomials involved, for  $G_2$  the number of algebraic dependence relations is unexpectedly high, namely 57, in contrast with the classical algebras, where the number of relations is low. It has been further shown that, among the three classical Lie algebras of rank two, for  $A_2$  and  $B_2$  the polynomial subalgebra  $\mathcal{A}$  is actually maximal of codimension one, for  $D_2$  it does not exist as the commutant is itself Abelian, while for  $G_2$  the subalgebra is of codimension nine and not maximal, as it can be extended nontrivially.

In view of the analogies with the so-called internal label problem [26], we may ask whether the procedure can be adapted to other reduction chains  $\mathfrak{s}' \subset \mathfrak{s}$  of semisimple Lie algebras in order to construct polynomial algebras. It shall however taken into account that polynomials  $P$  in the enveloping algebra of  $\mathfrak{s}$  commuting with the generators of  $\mathfrak{s}'$  are not suitable to describe the decomposition (9), as the eigenvalue of  $P$  with respect to the Cartan subalgebra of  $\mathfrak{s}'$  is always zero. Nonetheless, the ansatz can be of use for the analysis of integrable systems [9, 27] endowed with some internal symmetry, as well as for (quasi-)exactly solvable (QES) systems [10, 28]. In [10], the case of  $\mathfrak{gl}(3)$  was exploited to provide new algebraic (QES) systems, while in [27],  $\mathfrak{su}(3)$  was used to provide an algebraic setting for a superintegrable Hamiltonian on the sphere. This points out how the results of this paper may be as well of interest for different areas of mathematical physics, as well as it may also reveal to be of interest in the description of wider types of representations of simple Lie algebras [29].

As an illustrating example, let us consider the simple Lie algebra  $A_2$  in the basis (1) and the regular subalgebra  $\mathfrak{sl}(2, \mathbb{C})$  generated by  $H_1, E_{12}, E_{21}$ . In this case, the generators are no more related to highest weight vectors of  $A_2$ , but formally, the commutant of the subalgebra  $\mathfrak{sl}(2, \mathbb{C})$  in  $\mathcal{U}(\mathfrak{sl}(3, \mathbb{C}))$  has also a polynomial structure. An integrity basis for the system (17) associated to the subalgebra generators contains five elements, as can be easily verified. A short computation shows that there exist six linearly independent elements that span the commutant, given by the polynomials

$$\begin{aligned} P_1 &= H_1 + 2H_2, & P_2 &= E_{13}E_{31} + E_{23}E_{32} - H_2 - \frac{1}{2}H_1, & P_3 &= H_1^2 + 4E_{12}E_{21} - 2H_1, \\ P_4 &= E_{12}E_{31}^2 - E_{21}E_{32}^2 - H_1E_{31}E_{32}, & P_5 &= E_{12}E_{23}E_{31} + E_{13}E_{21}E_{32} + H_1E_{13}E_{31} - H_2E_{23}E_{32} \\ &+ E_{23}E_{32} - E_{12}E_{21} - H_2^2 - H_1H_2 - \frac{1}{2}H_1^2 - H_2, & P_6 &= E_{12}E_{23}^2 - E_{13}E_{21}^2 + H_1E_{13}E_{23} - 2E_{13}E_{23}. \end{aligned} \quad (49)$$

The six operators generate a polynomial algebra with nonvanishing commutators

$$\begin{aligned} [P_1, P_4] &= -6P_4, & [P_1, P_6] &= 6P_6, & [P_2, P_3] &= 3P_4 + P_1P_4, & [P_2, P_6] &= 3P_6 - P_1P_6, \\ [P_4, P_5] &= \left(-\frac{1}{2}P_1^2 - 5P_1 + 4P_2 - \frac{1}{2}P_3 - 12\right)P_4, & [P_4, P_6] &= P_1P_2P_3 - P_1(P_2^2 - P_2) + (2P_2 - P_3)P_5 \\ &- \frac{1}{4}P_1P_3, & [P_5, P_6] &= 4P_2P_6 - \frac{1}{2}(P_1^2 + P_3)P_6 + 5P_1P_6 - 12P_6. \end{aligned} \quad (50)$$

The polynomials are further algebraically dependent through the relation

$$\begin{aligned} P_1^2P_2^2 - P_2^2P_3 - 4P_1P_2P_5 + 2P_1P_2P_3 - 2P_1P_2^2 - P_1^2P_2 - \frac{1}{4}P_1^2P_3 - 4P_4P_6 + 4P_5^2 \\ + 2(P_1 + 2P_2 - P_3)P_5 + 2P_1P_2 - P_2P_3 + \frac{1}{4}(P_3 - 2P_1)P_3 = 0, \end{aligned} \quad (51)$$

showing that any element in the commutant can be written as

$$P = P_1^{a_1}P_2^{a_2}P_3^{a_3}P_4^{a_4}P_5^{a_5}P_6^{a_6}, \quad a_i \in \mathbb{N} \cup 0, \quad a_5 = 0, 1. \quad (52)$$

We observe that the weight of the polynomials with respect to the Cartan subalgebra of  $\mathfrak{sl}(3, \mathbb{C})$  is  $(0, 0)$  for  $P_1, P_2, P_3$  and  $P_5$ , while  $P_4$  has weight  $(0, 3)$  and  $P_6$  has weight  $(0, -3)$ .

## Acknowledgement

RCS was supported by the research grant PID2019-106802GB-I00/AEI/10.13039/501100011033 (AEI/ FEDER, UE). IM was supported by by Australian Research Council Future Fellowship FT180100099.

## References

- [1] B. Kostant, *Amer. J. Math. Phys.* **85** (1963) 327–404.

- [2] V. Ovsienko, A. V. Turbiner, *Comptes Rendus Acad. Sci. A* **314** (1992), 13–16.
- [3] A. V. Turbiner, *J. Math. Phys.* **33** (1992) 3389–3393.
- [4] W. Miller, S. Post and P. Winternitz *J. of Phys. A: Math. and Theor.* **46** (2013) 423001
- [5] D. Latini, I. Marquette and Y.-Z. Zhang, Polynomial algebras of superintegrable systems separating in Cartesian coordinates from higher order ladder operators. arXiv.2202.13281.
- [6] L. Freidel, J. M. Maillet, *Physics Letters B* **262** (1991), 278–284.
- [7] P. Letourneau, L. Vinet, *Ann. Physics* **243** (1995) 144–168.
- [8] A. Polishchuk and L. Positselski, *Quadratic Algebras*, Amer. Math. Society, Rhode Island, 2005.
- [9] D. Latini, I. Marquette, Y.-Z. Zhang, *Ann. Physics* **426** (2021) 168397.
- [10] R. Campoamor-Stursberg, I. Marquette, *Ann. Physics* **437** (2022) 168694.
- [11] H. Li, *Noncommutative Polynomial Algebras of Solvable Type and Their Modules*, CRC, Boca Raton, 2021.
- [12] L. A. Yates and P. D. Jarvis, *J. of Phys. A: Math. and Theor.* **51** (2018) 145203
- [13] D. E. Flath, *J. Math. Phys.* **31** (1990) 1076–1077.
- [14] Č. Burdík, O. Navrátil and S. Pošta, *Acta Polytech.* **47** (2007) 25–31.
- [15] J. P. Serre *Algèbres de Lie semi-simples complexes*, New York, Benjamin Inc., 1966.
- [16] J. Dixmier, *Algèbres enveloppantes*, Gauthier-Villars, Paris, 1974.
- [17] L. Abellanas, L. Martínez Alonso, *J. Math. Phys.* **16** (1975) 1580–1584.
- [18] G. Racah, *Rendiconti R. Acad. Nazionale Lincei* **8** 108–112.
- [19] A. Peccia, R. T. Sharp, *J. Math. Phys.* **17** (1976) 1313–1316.
- [20] R. Campoamor-Stursberg, *J. Phys. A: Math. Theor.* **44** (2011) 025234.
- [21] J. Dixmier, *Comptes Rendus Acad. Sci. A* **265** (1967) 408–410.
- [22] R. Campoamor-Stursberg, *Phys. Lett. A* **327** (2004) 138–145.
- [23] L. Šnobl and P. Winternitz, *Classification and identification of Lie algebras*, Amer. Math. Society, Rhode Island, 2014.
- [24] Č. Burdík, O. Navrátil in: S. Silvestrov et al. (Eds.) *Generalized Lie Theory in Mathematics, Physics and beyond*, Springer Verlag, Berlin, pp. 297–302.
- [25] W. G. McKay, J. Patera, *Tables of Dimensions, Indices, and Branching Rules for Representations of Simple Lie Algebras*, Marcel Dekker, New York, 1981.
- [26] R. T. Sharp, *Proc. Camb. Phil. Soc.* **68** (1970) 571–578
- [27] F. Correa, M. A. del Olmo, I. Marquette, and J. Negro, *J. of Phys. A: Math. and Theor.* **54** (2020) 015205.
- [28] P. Gaboriaud, L. Vinet, S. Vinet, A. Zhedanov, *J. Phys. Conf. Series* **1194** (2019) 012034.
- [29] J.E. Humphreys, *Representation of Semisimple Lie Algebras in the BGG Category  $\mathcal{O}$* , Springer Verlag, Berlin, 2008.