

Typed Non-determinism in Functional and Concurrent Calculi

Bas van den Heuvel¹[0000-0002-8264-7371], Joseph W. N.
Paulus¹[0000-0002-1711-9361],
Daniele Nantes-Sobrinho^{2,3}[0000-0002-1959-8730], and Jorge A.
Pérez¹[0000-0002-1452-6180]

¹ University of Groningen, Groningen, The Netherlands

² University of Brasília, Brasília, Brazil

³ Imperial College London, London, UK

Abstract. We study functional and concurrent calculi with non-determinism, along with type systems to control resources based on linearity. The interplay between non-determinism and linearity is delicate: careless handling of branches can discard resources meant to be used exactly once. Here we go beyond prior work by considering non-determinism in its standard sense: once a branch is selected, the rest are discarded. Our technical contributions are three-fold. First, we introduce a π -calculus with non-deterministic choice, governed by session types. Second, we introduce a resource λ -calculus, governed by intersection types, in which non-determinism concerns fetching of resources from bags. Finally, we connect our two typed non-deterministic calculi via a correct translation.

1 Introduction

In this paper, we present new formulations of typed programming calculi with *non-determinism*. A classical ingredient of models of computation, non-determinism brings flexibility and generality in specifications. In process calculi such as CCS and the π -calculus, one source of non-determinism is choice, which is typically *non-confluent*: that is, given $P + Q$, we have either $P + Q \longrightarrow P$ or $P + Q \longrightarrow Q$. Thus, committing to a branch entails discarding the rest.

We study non-determinism as a way of increasing the expressivity of typed calculi in which resource control is based on *linearity*. The interplay between non-determinism and linearity is delicate: a careless discarding of branches can jeopardize resources meant to be used exactly once. On the concurrent side, we consider the π -calculus, the paradigmatic model of concurrency [26]. We focus on π -calculi with *session types* [13,14], in which linear logic principles ensure communication correctness: here the resources are names that perform session protocols; they can be *unrestricted* (used multiple times) and *linear* (used exactly once). To properly control resources, non-confluent non-determinism is confined to unrestricted names; linear names can only perform deterministic choices.

In this context, considering *confluent* forms of non-determinism can be appealing. Intuitively, such formulations allow all branches to proceed independently: given $P_1 \longrightarrow Q_1$ and $P_2 \longrightarrow Q_2$, then $P_1 + P_2 \longrightarrow Q_1 + P_2$ and

$P_1 + P_2 \longrightarrow P_1 + Q_2$. Because confluent non-determinism does not discard branches, it is compatible with a resource-conscious view of computation.

Confluent non-determinism has been studied mostly in the functional setting; it is present, e.g., in Pagani and Ronchi della Rocca’s resource λ -calculus [19] and in Ehrhard and Regnier’s differential λ -calculus [10]. In [19], non-determinism resides in the application of a term M to a *bag* of available resources C ; a β -reduction applies M to a resource *non-deterministically fetched* from C . Confluent non-deterministic choice is also present in the session-typed π -calculus by Caires and Pérez [5], where it expresses a choice between different implementations of the same session protocols, which are all *non-deterministically available*—they may be available but may also *fail*. In their work, a Curry-Howard correspondence between linear logic and session types (*‘propositions-as-sessions’* [6,30]) ensures confluence, protocol fidelity, and deadlock-freedom. Paulus et al. [20] relate functional and concurrent calculi with confluent non-determinism: they give a translation of a resource λ -calculus into the session π -calculus from [5], in the style of Milner’s *‘functions-as-processes’* [16].

Although results involving confluent non-determinism are most significant, usual (non-confluent) non-determinism remains of undiscussed convenience in formal modeling; consider, e.g., specifications of distributed protocols [2,18] in which commitment is essential. Indeed, non-confluent non-deterministic choice is commonplace in verification frameworks such as mCRL2 [12]. It is also relevant in functional calculi; a well-known framework is De’Liguoro and Piperno’s (untyped) non-deterministic λ -calculus [8] (see also [9] and references therein).

To further illustrate the difference between confluent and non-confluent non-determinism, we consider an example adapted from [5]: a movie server that offers a choice between buying a movie or watching its trailer. In $s\pi^+$, the typed π -calculus that we present in this paper, this server can be specified as follows:

$$\text{Server}_s = s.\text{case} \left\{ \begin{array}{l} \text{buy} : s(\text{title}); s(\text{paym}); \bar{s}[\text{movie}]; \bar{s}[] \\ \text{peek} : s(\text{title}); \bar{s}[\text{trailer}]; \bar{s}[] \end{array} \right\} \begin{array}{l} -\text{Server}_s^{\text{buy}} \\ -\text{Server}_s^{\text{peek}} \end{array}$$

where $s(-)$ and $\bar{s}[-]$ denote input and output prefixes on a name/channel s , respectively, and ‘movie’ and ‘trailer’ denote references to primitive data. Also, the free names of a process are denoted with subscripts. Process Server_s offers a choice on name s ($s.\text{case}\{-\}$) between labels **buy** and **peek**. If **buy** is received, process $\text{Server}_s^{\text{buy}}$ is launched: it receives the movie’s title and a payment method, sends the movie, and closes the session on s ($\bar{s}[]$). If **peek** is received, it proceeds as $\text{Server}_s^{\text{peek}}$: the server receives the title, sends the trailer, and closes the session.

Using the non-deterministic choice operator of $s\pi^+$, denoted ‘ \sharp ’, we can specify a process for a client Alice who is interested in the movie ‘Jaws’ but is undecided about buying the film or just watching its trailer for free:

$$\text{Alice}_s := \bar{s}.\text{buy}; \bar{s}[\text{Jaws}]; \bar{s}[\text{mcard}]; s(\text{movie}); s(); \mathbf{0} \quad \sharp \bar{s}.\text{peek}; \bar{s}[\text{Jaws}]; s(\text{trailer}); s(); \mathbf{0} \quad \begin{array}{l} -\text{Alice}_s^{\text{buy}} \\ -\text{Alice}_s^{\text{peek}} \end{array}$$

If Alice_s selects the label **buy** ($\bar{s}.\text{buy}$), process $\text{Alice}_s^{\text{buy}}$ is launched: it sends title and payment method, receives the movie, waits for the session to close ($s()$),

and then terminates (0) . If Alice_s selects `peek`, process $\text{Alice}_s^{\text{peek}}$ is launched: it sends a title, receives the trailer, waits for the session to close, and terminates. Then, process $\text{Sys} := (\nu s)(\text{Server}_s \mid \text{Alice}_s)$ denotes the composition of client and server, connected along s (using (νs)). Our semantics for $s\pi^+$, denoted \rightsquigarrow , enforces non-confluent non-determinism, as Sys can reduce to separate processes, as expected:

$$\text{Sys} \rightsquigarrow (\nu s)(\text{Server}_s^{\text{buy}} \mid \text{Alice}_s^{\text{buy}}) \quad \text{and} \quad \text{Sys} \rightsquigarrow (\nu s)(\text{Server}_s^{\text{peek}} \mid \text{Alice}_s^{\text{peek}})$$

In contrast, the confluent non-deterministic choice from [5], denoted ‘ \oplus ’, behaves differently: in their confluent semantics, Sys reduces to a *single* process including *both alternatives*, i.e., $\text{Sys} \longrightarrow (\nu s)(\text{Server}_s^{\text{buy}} \mid \text{Alice}_s^{\text{buy}}) \oplus (\nu s)(\text{Server}_s^{\text{peek}} \mid \text{Alice}_s^{\text{peek}})$.

Contributions. We study new concurrent and functional calculi with usual (non-confluent) forms of non-determinism. Framed in the typed (resource-conscious) setting, we strive for definitions that do not exert a too drastic discarding of branches (as in the non-confluent case) but also that do not exhibit a lack of commitment (as in the confluent case). Concretely, we present:

(§ 2) $s\pi^+$, a variant of the session-typed π -calculus in [5], now with non-confluent non-deterministic choice. Its semantics adapts to the typed setting the usual semantics of non-deterministic choice in the untyped π -calculus [26]. Well-typed processes enjoy type preservation and deadlock-freedom (Theorems 1 and 2).

(§ 3) $\lambda_{\mathcal{C}}^{\ell}$, a resource λ -calculus with non-determinism, enhanced with constructs for expressing resource usage and failure. Its non-idempotent intersection type system provides a quantitative measure of the need/usage of resources. Well-typed terms enjoy subject reduction and subject expansion (Theorems 3 and 4).

(§ 4) A typed translation of $\lambda_{\mathcal{C}}^{\ell}$ into $s\pi^+$, which provides further validation for our non-deterministic calculi, and casts them in the context of ‘functions-as-processes’. We prove that our translation is *correct*, i.e., it preserves types and satisfies tight operational correspondences (Theorems 5 and 6).

Moreover, § 5 closes by discussing related works. Due to space limitations, appendices contain (i) omitted material (in particular, proofs of technical results); (ii) an alternative *eager* semantics for $s\pi^+$, which we compare against the lazy semantics; and (iii) extensions of $s\pi^+$ and $\lambda_{\mathcal{C}}^{\ell}$ with *unrestricted* resources.

2 A Typed π -calculus with Non-deterministic Choice

We introduce $s\pi^+$, a session-typed π -calculus with non-deterministic choice. Following [5], session types express protocols to be executed along channels. These protocols can be *non-deterministic*: sessions may succeed but also fail. The novelty in $s\pi^+$ is the non-deterministic choice operator ‘ $P \sharp Q$ ’, whose *lazily committing semantics* is compatible with linearity. We prove that well-typed processes satisfy two key properties: *type preservation* and *deadlock-freedom*.

$P, Q ::= \mathbf{0}$	inaction	$[x \leftrightarrow y]$	forwarder
$ (\nu x)(P Q)$	connect	$P \dagger Q$	non-determinism
$ \bar{x}[y]; (P Q)$	output	$x(y); P$	input
$ \bar{x}.\ell; P$	select	$x.\mathbf{case}\{i : P\}_{i \in I}$	branch
$ \bar{x}[]$	close	$x(); P$	wait
$ x.\mathbf{some}_{w_1, \dots, w_n}; P$	expect	$\bar{x}.\mathbf{some}; P$	available
$ P Q$	parallel	$\bar{x}.\mathbf{none}$	unavailable
.....			
$P \equiv P' [P \equiv_\alpha P']$	$[x \leftrightarrow y] \equiv [y \leftrightarrow x]$	$P \mathbf{0} \equiv P$	
$(P Q) R \equiv P (Q R)$	$P Q \equiv Q P$	$(\nu x)(P Q) \equiv (\nu x)(Q P)$	
$P \dagger P \equiv P$	$P \dagger Q \equiv Q \dagger P$	$(P \dagger Q) \dagger R \equiv P \dagger (Q \dagger R)$	
$(\nu x)((P Q) R) \equiv (\nu x)(P R) Q$	$[x \notin \mathit{fn}(Q)]$		
$(\nu x)((\nu y)(P Q) R) \equiv (\nu y)((\nu x)(P R) Q)$	$[x \notin \mathit{fn}(Q), y \notin \mathit{fn}(R)]$		

Fig. 1. $s\pi^+$: syntax (top) and structural congruence (bottom).

2.1 Syntax and Semantics

We use P, Q, \dots to denote processes, and x, y, z, \dots to denote *names* representing channels. Figure 1 (top) gives the syntax of processes. $P\{y/z\}$ denotes the capture-avoiding substitution of y for z in P . Process $\mathbf{0}$ denotes inaction, and $[x \leftrightarrow y]$ is a forwarder: a bidirectional link between x and y . Parallel composition appears in two forms: while the process $P | Q$ denotes communication-free concurrency, process $(\nu x)(P | Q)$ uses restriction (νx) to express that P and Q implement complementary behaviors on x and do not share any other names.

Process $P \dagger Q$ denotes the non-deterministic choice between P and Q : intuitively, if one choice can perform a synchronization, the other option may be discarded if it cannot. Since \dagger is associative, we often omit parentheses. Also, we write $\prod_{i \in I} P_i$ for the non-deterministic choice between each P_i for $i \in I$.

Our output construct integrates parallel composition and restriction: process $\bar{x}[y]; (P | Q)$ sends a fresh name y along x and then continues as $P | Q$. The type system will ensure that behaviors on y and x are implemented by P and Q , respectively, which do not share any names—this separation defines communication-free concurrency and is key to ensuring deadlock-freedom. The input process $x(y); P$ receives a name z along x and continues as $P\{z/y\}$, which does not require the separation present in the output case. Process $x.\mathbf{case}\{i : P_i\}_{i \in I}$ denotes a branch with labeled choices indexed by the finite set I : it awaits a choice on x with continuation P_j for each $j \in I$. The process $\bar{x}.\ell; P$ selects on x the choice labeled ℓ before continuing as P . Processes $\bar{x}[]$ and $x(); P$ are dual actions for closing the session on x . We omit replicated servers $!x(y); P$ and corresponding client requests $?x[y]; P$, but they can be easily added (cf. § A).

The remaining constructs define non-deterministic sessions which may provide a protocol or fail, following [5]. Process $\bar{x}.\text{some}; P$ confirms the availability of a session on x and continues as P . Process $\bar{x}.\text{none}$ signals the failure to provide the session on x . Process $x.\text{some}_{w_1, \dots, w_n}; P$ specifies a dependency on a non-deterministic session on x (names w_1, \dots, w_n implement sessions in P). This process can either (i) synchronize with a ' $\bar{x}.\text{some}$ ' and continue as P , or (ii) synchronize with a ' $\bar{x}.\text{none}$ ', discard P , and propagate the failure to w_1, \dots, w_n . To reduce eye strain, in writing $x.\text{some}$ we freely combine names and sets of names. This way, e.g., we write $x.\text{some}_{y, fn(P), fn(Q)}$ rather than $x.\text{some}_{\{y\} \cup fn(P) \cup fn(Q)}$.

Name y is bound in $(\nu y)(P|Q)$, $\bar{x}[y]; (P|Q)$, and $x(y); P$. We write $fn(P)$ and $bn(P)$ for the free and bound names of P , respectively. We adopt Barendregt's convention.

Structural Congruence. Reduction defines the steps that a process performs on its own. It relies on *structural congruence* (\equiv), the least congruence relation on processes induced by the rules in Figure 1 (bottom). Like the syntax of processes, the definition of \equiv is aligned with the type system (defined next), such that \equiv preserves typing (subject congruence, cf. Theorem 1). Differently from [5], we do not allow distributing non-deterministic choice over parallel and restriction. As shown in § B, the position of a non-deterministic choice in a process determines how it may commit, so changing its position affects commitment.

Reduction: Intuitions and Prerequisites. Barring non-deterministic choice, our reduction rules arise as directed interpretations of proof transformations in the underlying linear logic. We follow Caires and Pfenning [6] and Wadler [30] in interpreting cut-elimination in linear logic as synchronization in $\mathfrak{s}\pi^+$.

Before delving into our reduction rules (Figure 2), it may be helpful to consider the usual reduction axiom for the (untyped) π -calculus (e.g., [17, 26]):

$$(\bar{x}[z]; P_1 + M_1) \mid (x(y); P_2 + M_2) \longrightarrow P_1 \mid P_2\{z/y\} \quad (1)$$

This axiom captures the interaction of two (binary) choices: it integrates the commitment of choice in synchronization; after the reduction step, the two branches not involved in the synchronization, M_1 and M_2 , are discarded. Our semantics of $\mathfrak{s}\pi^+$ is defined similarly: when a prefix within a branch of a choice synchronizes with its dual, that branch reduces and the entire process commits to it.

The key question at this point is: when and to which branches should we commit? In (1), a communication commits to a single branch. For $\mathfrak{s}\pi^+$, we define a *lazy semantics* that minimizes commitment as much as possible.

The intuitive idea is that multiple branches of a choice may contain the same prefix, and so all these branches represent possibilities for synchronization ("possible branches"). Other branches with different prefixes denote different possibilities ("impossible branches"). When one synchronization is chosen, the possible branches are maintained while the impossible ones are discarded.

Example 1. To distinguish possible and impossible branches, consider:

$$P := (\nu s)(s.\text{case}\{\text{buy} : \dots, \text{peek} : \dots\} \mid (\bar{s}.\text{buy}; \dots \# \bar{s}.\text{buy}; \dots \# \bar{s}.\text{peek}; \dots))$$

The branch construct (case) provides the context for the non-deterministic choice. When the case synchronizes on the ‘buy’ label, the two branches prefixed by ‘ \bar{s} .buy’ are possible, whereas the branch prefixed by ‘ \bar{s} .peek’ becomes impossible, and can be discarded. The converse occurs when the ‘peek’ label is selected. ∇

To formalize these intuitions, our reduction semantics (Figure 2) relies on some auxiliary definitions. First, we define contexts.

Definition 1. We define ND-contexts (\mathbf{N}, \mathbf{M}) as follows:

$$\mathbf{N}, \mathbf{M} ::= [\cdot] \mid \mathbf{N} \mid P \mid (\nu x)(\mathbf{N} \mid P) \mid \mathbf{N} \# P$$

The process obtained by replacing $[\cdot]$ in \mathbf{N} with P is denoted $\mathbf{N}[P]$. We refer to ND-contexts that do not use the clause ‘ $\mathbf{N} \# P$ ’ as D-contexts, denoted \mathbf{C}, \mathbf{D} .

Using D-contexts, we can express that, e.g., $\prod_{i \in I} \mathbf{C}_i[\bar{x}]$ and $\prod_{j \in J} \mathbf{D}_j[x(); Q_j]$ should match. To account for reductions with impossible branches, we define a pre-congruence on processes, denoted \succeq_S , where the parameter S denotes the subject(s) of the prefix in the possible branches. Our semantics is closed under \succeq_S . Hence, e.g., anticipating a reduction on x , the possible branch $\mathbf{C}_1[x(y); P]$ can be extended with an impossible branch to form $\mathbf{C}_1[x(y); P] \# \mathbf{C}_2[z(); Q]$.

Before defining \succeq_S (Definition 3), we first define prefixes (and their subjects). Below, we write \bar{x} to denote a finite tuple of names x_1, \dots, x_k .

Definition 2. Prefixes are defined as follows:

$$\alpha, \beta ::= \bar{x}[y] \mid x(y) \mid \bar{x}.l \mid x.\text{case} \mid \bar{x}[] \mid x() \mid \bar{x}.\text{some} \mid \bar{x}.\text{none} \mid x.\text{some}_{\bar{w}} \mid [x \leftrightarrow y]$$

The subjects of α , denoted $\text{subj}\{\alpha\}$, are $\{x, y\}$ in case of $[x \leftrightarrow y]$, or $\{x\}$. By abuse of notation, we write $\alpha; P$ even when α takes no continuation (as in $\bar{x}[]$, $\bar{x}.\text{none}$, and $[x \leftrightarrow y]$) and for $\bar{x}[y]$ which takes a parallel composition as continuation.

Definition 3. Let \bowtie denote the least relation on prefixes (Def. 2) defined by:

(i) $\bar{x}[y] \bowtie \bar{x}[z]$, (ii) $x(y) \bowtie x(z)$, and (iii) $\alpha \bowtie \alpha$ otherwise.

Given a non-empty set $S \subseteq \{x, y\}$, the pre-congruence $P \succeq_S Q$ holds when both following conditions hold:

1. $S = \{x\}$ implies
 - $P = \left(\prod_{i \in I} \mathbf{C}_i[\alpha_i; P_i] \right) \# \left(\prod_{j \in J} \mathbf{C}_j[\beta_j; Q_j] \right)$ and $Q = \prod_{i \in I} \mathbf{C}_i[\alpha_i; P_i]$, where
 - (i) $\forall i, i' \in I. \alpha_i \bowtie \alpha_{i'}$ and $\text{subj}\{\alpha_i\} = \{x\}$, and
 - (ii) $\forall i \in I. \forall j \in J. \alpha_i \not\bowtie \beta_j \wedge x \in \text{fn}(\beta_j; Q_j)$;
2. $S = \{x, y\}$ implies
 - $P = \left(\prod_{i \in I} \mathbf{C}_i[[x \leftrightarrow y]] \right) \# \left(\prod_{j \in J} \mathbf{C}_j[[x \leftrightarrow z_j]] \right) \# \left(\prod_{k \in K} \mathbf{C}_k[\alpha_k; P_k] \right)$
 - and $Q = \prod_{i \in I} \mathbf{C}_i[[x \leftrightarrow y]]$, where
 - (i) $\forall j \in J. z_j \neq y$, and (ii) $\forall k \in K. x \in \text{fn}(\alpha_k; P_k) \wedge \forall z. \alpha_k \not\bowtie [x \leftrightarrow z]$.

$$\begin{array}{c}
 [\rightsquigarrow_{\text{Id}}] (\nu x) \left(\prod_{i \in I} \mathbf{C}_i [x \leftrightarrow y] \mid Q \right) \rightsquigarrow_{x,y} \prod_{i \in I} \mathbf{C}_i [Q\{y/x\}] \\
 [\rightsquigarrow_{\otimes \otimes}] (\nu x) \left(\prod_{i \in I} \mathbf{C}_i [\bar{x}y_i]; (P_i \mid Q_i) \mid \prod_{j \in J} \mathbf{D}_j [x(z); R_j] \right) \\
 \rightsquigarrow_x \prod_{i \in I} \mathbf{C}_i \left[(\nu x) (Q_i \mid (\nu w) (P_i\{w/y_i\} \mid \prod_{j \in J} \mathbf{D}_j [R_j\{w/z\}])) \right] \\
 [\rightsquigarrow_{\otimes \&}] (\nu x) \left(\prod_{i \in I} \mathbf{C}_i [\bar{x}.k'; P_i] \mid \prod_{j \in J} \mathbf{D}_j [x.\text{case}\{k : Q_j^k\}_{k \in K}] \right) \\
 \rightsquigarrow_x (\nu x) \left(\prod_{i \in I} \mathbf{C}_i [P_i] \mid \prod_{j \in J} \mathbf{D}_j [Q_j^{k'_j}] \right) \quad [k' \in K] \\
 [\rightsquigarrow_{\mathbf{1} \perp}] (\nu x) \left(\prod_{i \in I} \mathbf{C}_i [\bar{x}[]] \mid \prod_{j \in J} \mathbf{D}_j [x(); Q_j] \right) \rightsquigarrow_x \prod_{i \in I} \mathbf{C}_i [\mathbf{0}] \mid \prod_{j \in J} \mathbf{D}_j [Q_j] \\
 [\rightsquigarrow_{\text{some}}] (\nu x) \left(\prod_{i \in I} \mathbf{C}_i [\bar{x}.\text{some}; P_i] \mid \prod_{j \in J} \mathbf{D}_j [x.\text{some}_{w_1, \dots, w_n}; Q_j] \right) \\
 \rightsquigarrow_x (\nu x) \left(\prod_{i \in I} \mathbf{C}_i [P_i] \mid \prod_{j \in J} \mathbf{D}_j [Q_j] \right) \\
 [\rightsquigarrow_{\text{none}}] (\nu x) \left(\prod_{i \in I} \mathbf{C}_i [\bar{x}.\text{none}] \mid \prod_{j \in J} \mathbf{D}_j [x.\text{some}_{w_1, \dots, w_n}; Q_j] \right) \\
 \rightsquigarrow_x \prod_{i \in I} \mathbf{C}_i [\mathbf{0}] \mid \prod_{j \in J} \mathbf{D}_j [\bar{w}_1.\text{none} \mid \dots \mid \bar{w}_n.\text{none}] \\
 [\rightsquigarrow_{\succeq_S}] \frac{x \in S \quad P \succeq_S P' \quad Q \succeq_S Q' \quad (\nu x)(P' \mid Q') \rightsquigarrow_S R}{(\nu x)(P \mid Q) \rightsquigarrow_S R} \\
 [\rightsquigarrow_{\nu \dagger}] \frac{(\nu x)(P \mid \mathbf{N}[\mathbf{C}[Q_1] \dagger \mathbf{C}[Q_2]]) \rightsquigarrow_S R}{(\nu x)(P \mid \mathbf{N}[\mathbf{C}[Q_1 \dagger Q_2]]) \rightsquigarrow_S R} \\
 [\rightsquigarrow_{\equiv}] \frac{P \equiv P' \quad P' \rightsquigarrow_S Q' \quad Q' \equiv Q}{P \rightsquigarrow_S Q} \quad [\rightsquigarrow_{\nu}] \frac{P \rightsquigarrow_S P'}{(\nu x)(P \mid Q) \rightsquigarrow_S (\nu x)(P' \mid Q)} \\
 [\rightsquigarrow_{\mid}] \frac{P \rightsquigarrow_S P'}{P \mid Q \rightsquigarrow_S P' \mid Q} \quad [\rightsquigarrow_{\dagger}] \frac{P \rightsquigarrow_S P'}{P \dagger Q \rightsquigarrow_S P' \dagger Q}
 \end{array}$$

Fig. 2. Reduction semantics for $\mathfrak{s}\pi^+$.

Intuitively, \bowtie allows us to equate output/input prefixes with the same subject (but different object). The rest of Definition 3 accounts for two kinds of reduction, using S to discard “impossible” branches. In case S is $\{x\}$ (Item 1), it concerns a synchronization on x ; in case S is $\{x, y\}$, it concerns forwarding on x and y (Item 2). In both cases, P and Q contain matching prefixes on x , while P may contain additional branches with different or blocked prefixes on x ; x must appear in the hole of the contexts in the additional branches in P (enforced with $x \in \text{fn}(\dots)$), to ensure that no matching prefixes are discarded.

Example 2. Recall process P from Example 1. To derive a synchronization with the ‘buy’ alternative of the case, we can use \succeq_S to discard the ‘peek’ alternative, as follows: $\bar{s}.\text{buy}; \dots \dagger \bar{s}.\text{buy}; \dots \dagger \bar{s}.\text{peek}; \dots \succeq_s \bar{s}.\text{buy}; \dots \dagger \bar{s}.\text{buy}; \dots$ ∇

Reduction Rules. Figure 2 gives the rules for the (lazy) reduction semantics, denoted \rightsquigarrow_S , where the set S contains the names involved in the interaction. We

omit the curly braces in this annotation; this way, e.g., we write ‘ $\rightsquigarrow_{x,y}$ ’ instead of ‘ $\rightsquigarrow_{\{x,y\}}$ ’. Also, we write \rightsquigarrow_S^k to denote a sequence of $k \geq 0$ reductions.

The first six rules in Figure 2 formalize forwarding and communication: they are defined on choices containing different D-contexts (cf. Definition 1), each with the same prefix but possibly different continuations; these rules preserve the non-deterministic choices. Rule $[\rightsquigarrow_{ID}]$ fixes S to the forwarder’s two names, and the other rules fix S to the one involved name. In particular, Rule $[\rightsquigarrow_{\otimes \mathcal{P}}]$ formalizes name communication: it involves multiple senders and multiple receivers (grouped in choices indexed by I and J , respectively). Because they proceed in lock-step, reduction leads to substitutions involving the same (fresh) name w ; also, the scopes of the choice and the contexts enclosing the senders is extended.

Rule $[\rightsquigarrow_{\succeq_S}]$ is useful to derive a synchronization that discards groups of choices. Rule $[\rightsquigarrow_{\nu \#}]$ allows inferring reductions when non-deterministic choices are not top-level: e.g., $(\nu x)(\bar{x}[] \mid (\nu y)((x(); Q_1 \# x(); Q_2) \mid R)) \rightsquigarrow_x (\nu y)(Q_1 \mid R) \# (\nu y)(Q_2 \mid R)$. The last four rules formalize that reduction is closed under structural congruence, restriction, parallel composition, and non-deterministic choice.

As mentioned earlier, a key motivation for our work is to have non-deterministic choices that effectively enforce commitment, without a too drastic discarding of alternatives. Next we illustrate this intended form of *gradual commitment*.

Example 3 (A Modified Movie Server). Consider the following variant of the movie server from the introduction, where the handling of the payment is now modeled as a branch:

$$\text{NewServer}_s := s(\text{title}); s.\text{case} \left\{ \begin{array}{l} \text{buy} : s.\text{case} \left\{ \begin{array}{l} \text{card} : s(\text{info}); \bar{s}[\text{movie}]; \bar{s}[], \\ \text{cash} : \bar{s}[\text{movie}]; \bar{s}[] \end{array} \right\}, \\ \text{peek} : \bar{s}[\text{trailer}]; \bar{s}[] \end{array} \right\}$$

Consider a client, Eve, who cannot decide between buying ‘Oppenheimer’ or watching its trailer. In the former case, she has two options for payment method:

$$\text{Eve}_s := \bar{s}[\text{Oppenheimer}]; \left(\begin{array}{l} \bar{s}.\text{buy}; \bar{s}.\text{card}; \bar{s}[\text{visa}]; s(\text{movie}); s(); \mathbf{0} \\ \# \bar{s}.\text{buy}; \bar{s}.\text{cash}; s(\text{movie}); s(); \mathbf{0} \\ \# \bar{s}.\text{peek}; s(\text{link}); s(); \mathbf{0} \end{array} \right)$$

Let $\text{Sys}^* := (\nu s)(\text{NewServer}_s \mid \text{Eve}_s)$. After sending the movie’s title, Eve’s choice (buying or watching the trailer) enables gradual commitment. We have:

$$\text{Sys}^* \rightsquigarrow_s^2 (\nu s)(s.\text{case}\{\text{card} : \dots, \text{cash} : \dots\} \mid (\bar{s}.\text{card}; \dots \# \bar{s}.\text{cash}; \dots)) =: \text{Sys}_1^*$$

$$\text{and} \quad \text{Sys}^* \rightsquigarrow_s^2 (\nu s)(\bar{s}[\text{trailer}]; \dots \mid s(\text{trailer}); \dots) =: \text{Sys}_2^*$$

Process Sys_1^* represents the situation for Eve after selecting buy, in which case the third alternative ($\bar{s}.\text{peek}; \dots$) can be discarded as an impossible branch. Process Sys_2^* represents the dual situation. From Sys_1^* , the selection of payment method completes the commitment to one alternative; we have: $\text{Sys}_1^* \rightsquigarrow_s (\nu s)(s(\text{info}); \dots \mid \bar{s}[\text{visa}]; \dots)$ and $\text{Sys}_1^* \rightsquigarrow_s (\nu s)(\bar{s}[\text{movie}]; \dots \mid s(\text{movie}); \dots)$. ∇

In Appendix B we discuss an alternative *eager* semantics that commits to a single branch upon communication, as in (1).

$[T_{\text{CUT}}] \frac{P \vdash \Gamma, x:A \quad Q \vdash \Delta, x:\bar{A}}{(\nu x)(P \mid Q) \vdash \Gamma, \Delta}$	$[T_{\text{MIX}}] \frac{P \vdash \Gamma \quad Q \vdash \Delta}{P \mid Q \vdash \Gamma, \Delta}$	
$[T_{\#}] \frac{P \vdash \Gamma \quad Q \vdash \Gamma}{P \# Q \vdash \Gamma}$	$[T_{\text{EMPTY}}] \overline{\mathbf{0} \vdash \emptyset}$	$[T_{\text{ID}}] \overline{[x \leftrightarrow y] \vdash x:A, y:\bar{A}}$
$[T_{\mathbf{1}}] \overline{\bar{x}[] \vdash x:\mathbf{1}}$	$[T_{\perp}] \frac{P \vdash \Gamma}{x(); P \vdash \Gamma, x:\perp}$	$[T_{\otimes}] \frac{P \vdash \Gamma, y:A \quad Q \vdash \Delta, x:B}{\bar{x}[y]; (P \mid Q) \vdash \Gamma, \Delta, x:A \otimes B}$
$[T_{\wp}] \frac{P \vdash \Gamma, y:A, x:B}{x(y); P \vdash \Gamma, x:A \wp B}$	$[T_{\oplus}] \frac{P \vdash \Gamma, x:A_j \quad j \in I}{\bar{x}.j; P \vdash \Gamma, x:\oplus\{i : A_i\}_{i \in I}}$	
$[T_{\&}] \frac{\forall i \in I. P_i \vdash \Gamma, x:A_i}{x.\text{case}\{i : P_i\}_{i \in I} \vdash \Gamma, x:\&\{i : A_i\}_{i \in I}}$	$[T_{\&\text{some}}] \frac{P \vdash \Gamma, x:A}{\bar{x}.\text{some}; P \vdash \Gamma, x:\&A}$	
$[T_{\&\text{none}}] \overline{\bar{x}.\text{none} \vdash x:\&A}$	$[T_{\oplus\text{some}}] \frac{P \vdash \&\Gamma, x:A}{x.\text{some}_{\text{dom}(\Gamma)}; P \vdash \&\Gamma, x:\oplus A}$	

 Fig. 3. Typing rules for $\mathfrak{s}\pi^+$.

2.2 Resource Control for $\mathfrak{s}\pi^+$ via Session Types

We define a session type system for $\mathfrak{s}\pi^+$, following ‘propositions-as-sessions’ [6,30]. As already mentioned, in a session type system, resources are names that perform protocols: the *type assignment* $x : A$ says that x should conform to the protocol specified by the session type A . We give the syntax of types:

$$A, B ::= \mathbf{1} \mid \perp \mid A \otimes B \mid A \wp B \mid \oplus\{i : A_i\}_{i \in I} \mid \&\{i : A_i\}_{i \in I} \mid \&A \mid \oplus A$$

The units $\mathbf{1}$ and \perp type closed sessions. $A \otimes B$ types a name that first outputs a name of type A and then proceeds as B . Similarly, $A \wp B$ types a name that inputs a name of type A and then proceeds as B . Types $\oplus\{i : A_i\}_{i \in I}$ and $\&\{i : A_i\}_{i \in I}$ are given to names that can select and offer a labeled choice, respectively. Then, $\&A$ is the type of a name that *may produce* a behavior of type A , or fail; dually, $\oplus A$ types a name that *may consume* a behavior of type A .

For any type A we denote its *dual* as \bar{A} . Intuitively, dual types serve to avoid communication errors: the type at one end of a channel is the dual of the type at the opposite end. Duality is an involution, defined as follows:

$$\begin{array}{llll} \bar{\bar{\mathbf{1}}} = \perp & \overline{A \otimes B} = \bar{A} \wp \bar{B} & \overline{\oplus\{i : A_i\}_{i \in I}} = \&\{i : \bar{A}_i\}_{i \in I} & \overline{\&A} = \oplus \bar{A} \\ \bar{\bar{\perp}} = \mathbf{1} & \overline{A \wp B} = \bar{A} \otimes \bar{B} & \overline{\&\{i : A_i\}_{i \in I}} = \oplus\{i : \bar{A}_i\}_{i \in I} & \overline{\oplus A} = \&\bar{A} \end{array}$$

Judgments are of the form $P \vdash \Gamma$, where P is a process and Γ is a context, a collection of type assignments. In writing $\Gamma, x : A$, we assume $x \notin \text{dom}(\Gamma)$. We write $\text{dom}(\Gamma)$ to denote the set of names appearing in Γ . We write $\&\Gamma$ to denote that $\forall x : A \in \Gamma. \exists A'. A = \&A'$.

Figure 3 gives the typing rules: they correspond to the rules in Curry-Howard interpretations of classical linear logic as session types (cf. Wadler [30]), with the rules for $\&A$ and $\oplus A$ extracted from [5], and the additional Rule $[T\#]$ for non-confluent non-deterministic choice, which modifies the confluent rule in [5].

Most rules follow [30], so we focus on those related to non-determinism. Rule $[T\&\text{some}]$ types a process with a name whose behavior can be provided, while Rule $[T\&\text{none}]$ types a name whose behavior cannot. Rule $[T\oplus\text{some}]$ types a process with a name x whose behavior may not be available. If the behavior is not available, all the sessions in the process must be canceled; hence, the rule requires all names to be typed under the $\&A$ monad.

Rule $[T\#]$ types our new non-deterministic choice operator; the branches must be typable under the same typing context. Hence, all branches denote the same sessions, which may be implemented differently. In context of a synchronization, branches that are kept are able to synchronize, whereas the discarded branches are not; nonetheless, the remaining branches still represent different implementations of the same sessions. Compared to the rule for non-determinism in [5], we do not require processes to be typable under the $\&A$ monad.

Example 4. Consider again process Eve_s from Example 3. The three branches of the non-deterministic choice give *different implementations of the same session*: assuming primitive, self-dual data types \mathbf{C} , \mathbf{M} , and \mathbf{L} , all three branches on s are typable by $\oplus\{\text{buy} : \oplus\{\text{card} : \mathbf{C} \otimes \mathbf{M} \wp \perp, \text{cash} : \mathbf{M} \wp \perp\}, \text{peek} : \mathbf{L} \wp \perp\}$. ∇

Example 5 (Unavailable Movies). Consider now a modified movie server, which offers movies that may not be yet available. We specify this server using non-deterministic choice and non-deterministically available sessions:

$$\text{BuyServ}_s := s(\text{title}); (\bar{s}.\text{none} \# \bar{s}.\text{some}; s(\text{paym}); \bar{s}[\text{movie}]; \bar{s}[]) \vdash s : T \wp (\&(\mathbf{P} \wp \mathbf{M} \otimes \mathbf{1})),$$

where $T, \mathbf{P}, \mathbf{M}$ denote primitive, self-dual data-types. While the branch ‘ $\bar{s}.\text{none}$ ’ signals that the movie is not available, the branch ‘ $\bar{s}.\text{some}; \dots$ ’ performs the expected protocol. We now define a client Ada who buys a movie for Tim, using session s ; Ada only forwards it to him (using session u) if it is actually available:

$$\begin{aligned} \text{Ada}_{s,u} &:= \bar{s}[\text{Barbie}]; s.\text{some}_u; \bar{s}[\text{visa}]; s(\text{movie}); s(); \bar{u}.\text{some}; \bar{u}[\text{movie}]; \bar{u}[] \\ &\vdash s : T \otimes (\oplus(\mathbf{P} \otimes \mathbf{M} \wp \perp)), u : \&(\mathbf{M} \otimes \mathbf{1}) \\ \text{Tim}_u &:= u.\text{some}; u(\text{movie}); u(); \mathbf{0} \vdash u : \oplus(\mathbf{M} \wp \mathbf{1}) \end{aligned}$$

Let $\text{BuySys} := (\nu s)(\text{BuyServ}_s \mid (\nu u)(\text{Ada}_{s,u} \mid \text{Tim}_u))$. Depending on whether the server has the movie ‘Barbie’ available, we have the following reductions:

$$\text{BuySys} \rightsquigarrow_s^2 (\nu u)(\bar{u}.\text{none} \mid \text{Tim}_u) \quad \text{or} \quad \text{BuySys} \rightsquigarrow_s^5 (\nu u)(\bar{u}.\text{some}; \dots \mid \text{Tim}_u) \quad \nabla$$

Our type system ensures *session fidelity* and *communication safety*, but not confluence: the former says that processes correctly follow their ascribed session protocols, and the latter that no communication errors/mismatches occur. Both properties follow from the fact that typing is consistent across structural congruence and reduction. See Appendix E.2 for details.

$M, N, L ::= x$	variable	$M \langle\langle C/x \rangle\rangle$	intermediate subst.
$ (M C)$	application	$M \langle C/\tilde{x} \rangle$	explicit subst.
$ \lambda x.M$	abstraction	$\mathbf{fail}^{\tilde{x}}$	failure
$ M[\tilde{x} \leftarrow x]$	sharing		
$C, D ::= 1$	$ \wr M \wr \cdot C$		bag
$C ::= [\cdot]$	$ C[\tilde{x} \leftarrow x]$	$ (C C)$	$ C \langle C/\tilde{x} \rangle$
			context

Fig. 4. Syntax of $\lambda_{\mathcal{C}}^{\ell}$: terms, bags, and contexts.

Theorem 1 (Type Preservation). *If $P \vdash \Gamma$, then both $P \equiv Q$ and $P \rightsquigarrow_S Q$ (for any Q and S) imply $Q \vdash \Gamma$.*

Another important, if often elusive, property in session types is *deadlock-freeness*, which ensures that processes can reduce as long as they are not inactive. Our type system satisfies deadlock-freeness for processes with fully connected names, i.e., typable under the empty context. See Appendix E.2 for details.

Theorem 2 (Deadlock-freeness). *If $P \vdash \emptyset$ and $P \neq \mathbf{0}$, then there are Q and S such that $P \rightsquigarrow_S Q$.*

3 A Non-deterministic Resource λ -calculus

We present $\lambda_{\mathcal{C}}^{\ell}$, a resource λ -calculus with non-determinism and lazy evaluation. In $\lambda_{\mathcal{C}}^{\ell}$, non-determinism is non-confluent and *implicit*, as it arises from the fetching of terms from bags of *linear* resources. This is different from $\mathfrak{s}\pi^+$, where the choice operator ‘ \sharp ’ specifies non-determinism *explicitly*. A mismatch between the number of variable occurrences and the size of the bag induces *failure*.

In $\lambda_{\mathcal{C}}^{\ell}$, the *sharing* construct $M[x_1, \dots, x_n \leftarrow x]$, expresses that x may be used in M under “aliases” x_1, \dots, x_n . Hence, it atomizes n occurrences of x in M , via an explicit pointer to n variables. This way, e.g., the λ -term $\lambda x.(x x)$ is expressed in $\lambda_{\mathcal{C}}^{\ell}$ as $\lambda x.(x_1 \wr x_2 \wr [x_1, x_2 \leftarrow x])$, where $\wr x_2 \wr$ is a bag containing x_2 .

3.1 Syntax and Reduction Semantics

Syntax. We use x, y, z, \dots for variables, and write \tilde{x} to denote a finite sequence of pairwise distinct x_i ’s, with length $|\tilde{x}|$. Figure 4 gives the syntax of terms (M, N, L) and bags (C, D). The empty bag is denoted 1 . We use C_i to denote the i -th term in C , and $\text{size}(C)$ denotes the number of elements in C . To ease readability, we often write, e.g., $\wr N_1, N_2 \wr$ as a shorthand notation for $\wr N_1 \wr \cdot \wr N_2 \wr$.

In $M[\tilde{x} \leftarrow x]$, we say that \tilde{x} are the *shared variables* and that x is the *sharing variable*. We require for each $x_i \in \tilde{x}$: (i) x_i occurs exactly once in M ; (ii) x_i is not a sharing variable. The sequence \tilde{x} can be empty: $M[\leftarrow x]$ means that x does not share any variables in M . Sharing binds the shared variables in the term.

$\frac{[\text{RS:Beta}]}{(\lambda x.M)C \longrightarrow M\langle\langle C/x \rangle\rangle}$	$\frac{[\text{RS:Ex-Sub}]}{\text{size}(C) = \tilde{x} \quad M \neq \text{fail}^{\tilde{y}}}{(M[\tilde{x} \leftarrow x])\langle\langle C/x \rangle\rangle \longrightarrow M\langle C/\tilde{x} \rangle}$	$\frac{[\text{RS : TCont}]}{M \longrightarrow N}{\mathcal{C}[M] \longrightarrow \mathcal{C}[N]}$
$\frac{[\text{RS:Fetch}^\ell]}{\text{head}(M) = x_j \quad 0 < i \leq \text{size}(C)}{M\langle C/\tilde{x}, x_j \rangle \longrightarrow M\{C_i/x_j\}\langle(C \setminus C_i)/\tilde{x}\rangle}$		
$\frac{[\text{RS:Fail}^\ell]}{\text{size}(C) \neq \tilde{x} \quad \tilde{y} = (\text{fv}(M) \setminus \{\tilde{x}\}) \cup \text{fv}(C)}{(M[\tilde{x} \leftarrow x])\langle\langle C/x \rangle\rangle \longrightarrow \text{fail}^{\tilde{y}}}$	$\frac{[\text{RS:Cons}_1]}{\tilde{y} = \text{fv}(C)}{\text{fail}^{\tilde{x}} C \longrightarrow \text{fail}^{\tilde{x} \cup \tilde{y}}}$	
$\frac{[\text{RS:Cons}_2]}{\text{size}(C) = \tilde{x} \quad \tilde{z} = \text{fv}(C)}{(\text{fail}^{\tilde{x} \cup \tilde{y}}[\tilde{x} \leftarrow x])\langle\langle C/x \rangle\rangle \longrightarrow \text{fail}^{\tilde{y} \cup \tilde{z}}}$	$\frac{[\text{RS:Cons}_3]}{\tilde{z} = \text{fv}(C)}{\text{fail}^{\tilde{y} \cup \tilde{x}}\langle C/\tilde{x} \rangle \longrightarrow \text{fail}^{\tilde{y} \cup \tilde{z}}}$	
<p>where $\text{head}(M)$ is defined as follows:</p>		
$\begin{aligned} \text{head}(x) &= x & \text{head}(\lambda x.M) &= \lambda x.M & \text{head}((M C)) &= \text{head}(M) \\ \text{head}(\text{fail}^{\tilde{x}}) &= \text{fail}^{\tilde{x}} & \text{head}(M\langle\langle C/x \rangle\rangle) &= M\langle\langle C/x \rangle\rangle & \text{head}(M\langle C/\tilde{x} \rangle) &= \text{head}(M) \\ \text{head}(M[\tilde{x} \leftarrow x]) &= \begin{cases} x & \text{head}(M) = y \text{ and } y \in \tilde{x} \\ \text{head}(M) & \text{otherwise} \end{cases} \end{aligned}$		

Fig. 5. Reduction rules for λ_c^ℓ .

An abstraction $\lambda x.M$ binds occurrences of x in M . Application $(M C)$ is as usual. The term $M\langle C/\tilde{x} \rangle$ is the *explicit substitution* of a bag C for \tilde{x} in M . We require $\text{size}(C) = |\tilde{x}|$ and for each $x_i \in \tilde{x}$: (i) x_i occurs in M ; (ii) x_i is not a sharing variable; (iii) x_i cannot occur in another explicit substitution in M . The term $M\langle\langle C/x \rangle\rangle$ denotes an intermediate explicit substitution that does not (necessarily) satisfy the conditions for explicit substitutions.

The term $\text{fail}^{\tilde{x}}$ denotes failure; the variables in \tilde{x} are “dangling” resources, which cannot be accounted for after failure. We write $\text{fv}(M)$ to denote the free variables of M , defined as expected. Term M is *closed* if $\text{fv}(M) = \emptyset$.

Semantics. Figure 5 gives the reduction semantics, denoted \longrightarrow , and the *head variable* of term M , denoted $\text{head}(M)$. Rule **[RS : Beta]** induces an intermediate substitution. Rule **[RS : Ex-Sub]** reduces an intermediate substitution to an explicit substitution, provided the size of the bag equals the number of shared variables. In case of a mismatch, the term evolves into failure via Rule **[RS : Fail]^ℓ**.

An explicit substitution $M\langle C/\tilde{x} \rangle$, where the head variable of M is $x_j \in \tilde{x}$, reduces via Rule **[R : Fetch]^ℓ**. The rule extracts a C_i from C (for some $0 < i \leq \text{size}(C)$) and substitutes it for x_j in M ; this is how fetching induces a non-deterministic choice between $\text{size}(C)$ possible reductions. Rules **[RS : Cons]_j** for

$j \in \{1, 2, 3\}$ consume terms when they meet failure. Finally, Rule [RS : TCont] closes reduction under contexts. The following example illustrates reduction.

Example 6. Consider the term $M_0 = (\lambda x.x_1 \wr x_2 \wr x_3 \mathbf{1} \wr [\tilde{x} \leftarrow x]) \wr \mathbf{fail}^\emptyset, y, I \wr$, where $I = \lambda x.(x_1[x_1 \leftarrow x])$ and $\tilde{x} = x_1, x_2, x_3$. First, M_0 evolves into an intermediate substitution (2). The bag can provide for all shared variables, so it then evolves into an explicit substitution (3):

$$M_0 \longrightarrow (x_1 \wr x_2 \wr x_3 \mathbf{1} \wr [\tilde{x} \leftarrow x]) \langle \langle \mathbf{fail}^\emptyset, y, I \rangle / x \rangle \quad (2)$$

$$\longrightarrow (x_1 \wr x_2 \wr x_3 \mathbf{1} \wr) \langle \langle \mathbf{fail}^\emptyset, y, I \rangle / \tilde{x} \rangle = M \quad (3)$$

Since $\text{head}(M) = x_1$, one of the three elements of the bag will be substituted. M represents a non-deterministic choice between the following three reductions:

$$\begin{aligned} & \nearrow (\mathbf{fail}^\emptyset \wr x_2 \wr x_3 \mathbf{1} \wr) \langle \langle y, I \rangle / x_2, x_3 \rangle = N_1 \\ M & \longrightarrow (y \wr x_2 \wr x_3 \mathbf{1} \wr) \langle \langle \mathbf{fail}^\emptyset, I \rangle / x_2, x_3 \rangle = N_2 \\ & \searrow (I \wr x_2 \wr x_3 \mathbf{1} \wr) \langle \langle \mathbf{fail}^\emptyset, y \rangle / x_2, x_3 \rangle = N_3 \quad \nabla \end{aligned}$$

3.2 Resource Control for λ_c^ℓ via Intersection Types

Our type system for λ_c^ℓ is based on non-idempotent intersection types. As in prior works [19,4], intersection types account for available resources in bags, which are unordered and have all the same type. Because we admit the term $\mathbf{fail}^{\tilde{x}}$ as typable, we say that our system enforces *well-formedness* rather than *well-typedness*. As we will see, well-typed terms form the sub-class of well-formed terms that does not include $\mathbf{fail}^{\tilde{x}}$ (see the text after Theorem 3).

Strict types (σ, τ, δ) and multiset types (π, ζ) are defined as follows:

$$\sigma, \tau, \delta ::= \mathbf{unit} \mid \pi \rightarrow \sigma \quad \pi, \zeta ::= \bigwedge_{i \in I} \sigma_i \mid \omega$$

Given a non-empty I , multiset types $\bigwedge_{i \in I} \sigma_i$ are given to bags of size $|I|$. This operator is associative, commutative, and non-idempotent (i.e., $\sigma \wedge \sigma \neq \sigma$), with identity ω . Notation σ^k stands for $\sigma \wedge \dots \wedge \sigma$ (k times, if $k > 0$) or ω (if $k = 0$).

Judgments have the form $\Gamma \vDash M : \tau$, with contexts defined as follows:

$$\Gamma, \Delta ::= - \mid \Gamma, x : \pi \mid \Gamma, x : \sigma$$

where $-$ denotes the empty context. We write $\text{dom}(\Gamma)$ for the set of variables in Γ . For $\Gamma, x : \pi$, we assume $x \notin \text{dom}(\Gamma)$. To avoid ambiguities, we write $x : \sigma^1$ to denote that the assignment involves a multiset type, rather than a strict type. Given Γ , its *core context* Γ^\downarrow concerns variables with types different from ω ; it is defined as $\Gamma^\downarrow = \{x : \pi \in \Gamma \mid \pi \neq \omega\}$.

Definition 4 (Well-formedness in λ_c^ℓ). *A term M is well-formed if there exists a context Γ and a type τ such that the rules in Figure 6 entail $\Gamma \vDash M : \tau$.*

$\frac{[\text{FS:var}^\ell]}{x : \sigma \vDash x : \sigma}$	$\frac{[\text{FS:1}^\ell]}{- \vDash \mathbf{1} : \omega}$	$\frac{[\text{FS:bag}^\ell]}{\Gamma \vDash N : \sigma \quad \Delta \vDash C : \sigma^k}{\Gamma, \Delta \vDash \{N\} \cdot C : \sigma^{k+1}}$	$\frac{[\text{FS:fail}]}{\text{dom}(\Gamma^\downarrow) = \tilde{x}}{\Gamma^\downarrow \vDash \text{fail}^{\tilde{x}} : \tau}$
$\frac{[\text{FS:weak}]}{\Gamma \vDash M : \tau}{\Gamma, x : \omega \vDash M[\leftarrow x] : \tau}$		$\frac{[\text{FS:shar}]}{\Gamma, x_1 : \sigma, \dots, x_k : \sigma \vDash M : \tau \quad k \neq 0}{\Gamma, x : \sigma^k \vDash M[x_1, \dots, x_k \leftarrow x] : \tau}$	
$\frac{[\text{FS:abs-sh}]}{\Gamma, x : \sigma^k \vDash M[\tilde{x} \leftarrow x] : \tau}{\Gamma \vDash \lambda x. (M[\tilde{x} \leftarrow x]) : \sigma^k \rightarrow \tau}$		$\frac{[\text{FS:app}]}{\Gamma \vDash M : \sigma^j \rightarrow \tau \quad \Delta \vDash C : \sigma^k}{\Gamma, \Delta \vDash M C : \tau}$	
$\frac{[\text{FS:Esub}]}{\Gamma, x : \sigma^j \vDash M[\tilde{x} \leftarrow x] : \tau \quad \Delta \vDash C : \sigma^k}{\Gamma, \Delta \vDash (M[\tilde{x} \leftarrow x]) \langle \langle C/x \rangle \rangle : \tau}$		$\frac{[\text{FS:Esub}^\ell]}{\Gamma, x_1 : \sigma, \dots, x_k : \sigma \vDash M : \tau \quad \Delta \vDash C : \sigma^k}{\Gamma, \Delta \vDash M \langle \langle C/x_1, \dots, x_k \rangle \rangle : \tau}$	

Fig. 6. Well-Formedness Rules for λ_C^ℓ .

In Figure 6, Rule $[\text{FS} : \text{var}^\ell]$ types variables. Rule $[\text{FS} : \mathbf{1}^\ell]$ types the empty bag with ω . Rule $[\text{FS} : \text{bag}^\ell]$ types the concatenation of bags. Rule $[\text{FS} : \text{fail}]$ types the term $\text{fail}^{\tilde{x}}$ with a strict type τ , provided that the domain of the core context coincides with \tilde{x} (i.e., no variable in \tilde{x} is typed with ω). Rule $[\text{FS} : \text{weak}]$ types $M[\leftarrow x]$ by weakening the context with $x : \omega$. Rule $[\text{FS} : \text{shar}]$ types $M[\tilde{x} \leftarrow x]$ with τ , provided that there are assignments to the shared variables in \tilde{x} .

Rule $[\text{FS} : \text{abs-sh}]$ types an abstraction $\lambda x. (M[\tilde{x} \leftarrow x])$ with $\sigma^k \rightarrow \tau$, provided that $M[\tilde{x} \leftarrow x] : \tau$ can be entailed from an assignment $x : \sigma^k$. Rule $[\text{FS} : \text{app}]$ types $(M C)$, provided that M has type $\sigma^j \rightarrow \tau$ and C has type σ^k . Note that, unlike usual intersection type systems, j and k may differ. Rule $[\text{FS} : \text{Esub}]$ types the intermediate substitution of a bag C of type σ^k , provided that x has type σ^j ; again, j and k may differ. Rule $[\text{FS} : \text{Esub}^\ell]$ types $M \langle \langle C/\tilde{x} \rangle \rangle$ as long as C has type $\sigma^{|\tilde{x}|}$, and each $x_i \in \tilde{x}$ is of type σ .

Well-formed terms satisfy subject reduction (SR), whereas *well-typed* terms, defined below, satisfy also subject expansion (SE). See § F.1 and § F.2 for details.

Theorem 3 (SR in λ_C^ℓ). *If $\Gamma \vDash M : \tau$ and $M \longrightarrow M'$, then $\Gamma \vDash M' : \tau$.*

From our system for well-formedness we can extract a system for *well-typed* terms, which do not include $\text{fail}^{\tilde{x}}$. Judgments for well-typedness are denoted $\Gamma \vdash M : \tau$, with rules copied from Figure 6 (the rule name prefix FS is replaced with TS), with the following modifications: (i) Rule $[\text{TS} : \text{fail}]$ is removed; (ii) Rules $[\text{TS} : \text{app}]$ and $[\text{TS} : \text{Esub}]$ are modified to disallow a mismatch between variables and resources, i.e., multiset types should match in size. Well-typed terms are also well-formed, and thus satisfy SR. Moreover, as a consequence of adopting (non-idempotent) intersection types, they also satisfy SE:

Theorem 4 (SE in λ_C^ℓ). *If $\Gamma \vdash M' : \tau$ and $M \longrightarrow M'$, then $\Gamma \vdash M : \tau$.*

$$\begin{aligned}
 \llbracket x \rrbracket_u &= \bar{x}.\mathbf{some}; [x \leftrightarrow u] \\
 \llbracket \lambda x.M \rrbracket_u &= \bar{u}.\mathbf{some}; u(x); \llbracket M \rrbracket_u \\
 \llbracket (MC) \rrbracket_u &= (\nu v)(\llbracket M \rrbracket_v \mid v.\mathbf{some}_{u, \text{fv}(C)}; \bar{v}[x]; (\llbracket C \rrbracket_x \mid [v \leftrightarrow u])) \\
 \llbracket M \langle\langle C/x \rangle\rangle \rrbracket_u &= (\nu x)(\llbracket M \rrbracket_u \mid \llbracket C \rrbracket_x) \\
 \llbracket \{ N \} \cdot C \rrbracket_x &= x.\mathbf{some}_{\text{fv}(C), \text{fv}(N)}; x(y_i); x.\mathbf{some}_{y_i, \text{fv}(C), \text{fv}(N)}; \bar{x}.\mathbf{some}; \bar{x}[z_i]; \\
 &\quad (z_i.\mathbf{some}_{\text{fv}(N)}; \llbracket N \rrbracket_{z_i} \mid \llbracket C \rrbracket_x \mid \bar{y}_i.\mathbf{none}) \\
 \llbracket \mathbf{1} \rrbracket_x &= x.\mathbf{some}_\emptyset; x(y_n); (\bar{y}_n.\mathbf{some}; \bar{y}_n[] \mid x.\mathbf{some}_\emptyset; \bar{x}.\mathbf{none}) \\
 \llbracket M \langle\{ N_1, N_2 \} / x_1, x_2 \rangle \rrbracket_u &= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(N_1)}; \llbracket N_1 \rrbracket_{z_1} \mid (\nu z_2)(z_2.\mathbf{some}_{\text{fv}(N_2)}; \llbracket N_2 \rrbracket_{z_2} \\
 &\quad \mid \amalg_{x_i \in \{x_1, x_2\}} \amalg_{x_j \in \{x_1, x_2 \setminus x_i\}} \llbracket M \rrbracket_u \{z_1/x_i\} \{z_2/x_j\})) \\
 \llbracket M[\leftarrow x] \rrbracket_u &= \bar{x}.\mathbf{some}; \bar{x}[y_i]; (y_i.\mathbf{some}_{u, \text{fv}(M)}; y_i(); \llbracket M \rrbracket_u \mid \bar{x}.\mathbf{none}) \\
 \llbracket M[\tilde{x} \leftarrow x] \rrbracket_u &= \bar{x}.\mathbf{some}; \bar{x}[y_i]; (y_i.\mathbf{some}_\emptyset; y_i(); \mathbf{0} \mid \bar{x}.\mathbf{some}; x.\mathbf{some}_{u, \text{fv}(M) \setminus \tilde{x}}; \\
 &\quad \amalg_{x_i \in \tilde{x}} x(x_i); \llbracket M[(\tilde{x} \setminus x_i) \leftarrow x] \rrbracket_u) \\
 \llbracket \mathbf{fail}^{x_1, \dots, x_k} \rrbracket_u &= \bar{u}.\mathbf{none} \mid \bar{x}_1.\mathbf{none} \mid \dots \mid \bar{x}_k.\mathbf{none}
 \end{aligned}$$

 Fig. 7. Translation of λ_c^ℓ into $s\pi^+$.

4 A Typed Translation of λ_c^ℓ into $s\pi^+$

While $s\pi^+$ features non-deterministic choice, λ_c^ℓ is a prototypical programming language in which implicit non-determinism implements fetching of resources. Resources are controlled using different type systems (session types in $s\pi^+$, intersection types in λ_c^ℓ). To reconcile these differences and illustrate the potential of $s\pi^+$ to precisely model non-determinism as found in realistic programs/protocols, we give a translation of λ_c^ℓ into $s\pi^+$. This translation preserves types (Theorem 5) and respects well-known criteria for dynamic correctness [11,22,23] (Theorem 6).

The Translation. Given a λ_c^ℓ -term M , its translation into $s\pi^+$ is denoted $\llbracket M \rrbracket_u$ and given in Figure 7. As usual, every variable x in M becomes a name x in process $\llbracket M \rrbracket_u$, where name u provides the behavior of M . A peculiarity is that, to handle failures in λ_c^ℓ , u is a non-deterministically available session: the translated term can be available or not, as signaled by prefixes $\bar{u}.\mathbf{some}$ and $\bar{u}.\mathbf{none}$, respectively. As a result, reductions from $\llbracket M \rrbracket_u$ include synchronizations that codify M 's behavior but also synchronizations that confirm a session's availability.

At its core, our translation follows Milner's. This way, e.g., the process $\llbracket (\lambda x.M) C \rrbracket_u$ enables synchronizations between $\llbracket \lambda x.M \rrbracket_v$ and $\llbracket C \rrbracket_x$ along name v , resulting in the translation of an intermediate substitution. The *key novelty* is the role and treatment of non-determinism. Accommodating non-confluent non-determinism is non-trivial, as it entails translating explicit substitutions and sharing in λ_c^ℓ using the non-deterministic choice operator \amalg in $s\pi^+$. Next we discuss these novel aspects, while highlighting differences with respect to a translation by Paulus et al. [20], which is given in the confluent setting (see § 5).

In Figure 7, non-deterministic choices occur in the translations of $M\langle C/\tilde{x}\rangle$ (explicit substitutions) and $M[\tilde{x} \leftarrow x]$ (non-empty sharing). Roughly speaking, the position of \sharp in the translation of $M\langle C/\tilde{x}\rangle$ represents the most desirable way of mimicking the fetching of terms from a bag. This use of \sharp is a central idea in our translation: as we explain below, it allows for appropriate commitment in non-deterministic choices, but also for *delayed* commitment when necessary.

For simplicity, we consider explicit substitutions $M\langle C/\tilde{x}\rangle$ where $C = \{N_1, N_2\}$ and $\tilde{x} = x_1, x_2$. The translation $\llbracket M\langle C/\tilde{x}\rangle \rrbracket_u$ uses the processes $\llbracket N_i \rrbracket_{z_i}$, where each z_i is fresh. First, each bag item confirms its behavior. Then, a variable $x_i \in \tilde{x}$ is chosen non-deterministically; we ensure that these choices consider all variables. Note that writing $\sharp_{x_i \in \{x_1, x_2\}} \sharp_{x_j \in \{x_1, x_2\} \setminus x_i}$ is equivalent to non-deterministically assigning x_i, x_j to each permutation of x_1, x_2 . The resulting choice involves $\llbracket M \rrbracket_u$ with x_i, x_j substituted by z_1, z_2 . Commitment here is triggered only via synchronizations along z_1 or z_2 ; synchronizing with $z_i.\text{some}_{fv(N_i)}$; $\llbracket N_i \rrbracket_{z_i}$ then represents fetching N_i from the bag. The size of the translated term $\llbracket M\langle C/\tilde{x}\rangle \rrbracket_u$ is exponential with respect to the size of C .

The process $\llbracket M[\tilde{x} \leftarrow x] \rrbracket_u$ proceeds as follows. First, it confirms its behavior along x . Then it sends a name y_i on x , on which a failed reduction may be handled. Next, the translation confirms again its behavior along x and non-deterministically receives a reference to an $x_i \in \tilde{x}$. Each branch consists of $\llbracket M[(\tilde{x} \setminus x_i) \leftarrow x] \rrbracket_u$. The possible choices are permuted, represented by $\sharp_{x_i \in \tilde{x}}$. Synchronizations with $\llbracket M[(\tilde{x} \setminus x_i) \leftarrow x] \rrbracket_u$ and bags delay commitment in this choice (we return to this point below). The process $\llbracket M[\leftarrow x] \rrbracket_u$ is similar but simpler: here the name x fails, as it cannot take further elements to substitute.

In case of a failure (i.e., a mismatch between the size of the bag C and the number of variables in M), our translation ensures that the confirmations of C will not succeed. This is how failure in λ_C^ℓ is correctly translated to failure in $\mathfrak{s}\pi^+$.

Translation Correctness. The translation is typed: intersection types in λ_C^ℓ are translated into session types in $\mathfrak{s}\pi^+$ (Figure 8). This translation of types abstractly describes how non-deterministic fetches are codified as non-deterministic session protocols. It is worth noting that this translation of types is the same as in [20]. This is not surprising: as we have seen, session types effectively abstract away from the behavior of processes, as all branches of a non-deterministic choice use the same typing context. Still, it is pleasant that the translation of types remains unchanged across different translations with our (non-confluent) non-determinism (in Figure 7) and with confluent non-determinism (in [20]).

To state *static* correctness, we require the following definition:

Definition 5. *Let $\Gamma = x_1 : \sigma_1, \dots, x_m : \sigma_m, v_1 : \pi_1, \dots, v_n : \pi_n$ be a context. The translation $\llbracket \cdot \rrbracket_u$ in Figure 8 extends to contexts as follows:*

$$\llbracket \Gamma \rrbracket = x_1 : \&\overline{\llbracket \sigma_1 \rrbracket}, \dots, x_m : \&\overline{\llbracket \sigma_m \rrbracket}, v_1 : \overline{\llbracket \pi_1 \rrbracket}_{(\sigma, i_1)}, \dots, v_n : \overline{\llbracket \pi_n \rrbracket}_{(\sigma, i_n)}$$

Well-formed terms translate into well-typed processes:

$$\begin{array}{l}
 \llbracket \mathbf{unit} \rrbracket = \& \mathbf{1} \qquad \llbracket \sigma^k \rightarrow \tau \rrbracket = \& (\overline{\llbracket \sigma^k \rrbracket}_{(\sigma,i)}} \wp \llbracket \tau \rrbracket) \\
 \llbracket \sigma \wedge \pi \rrbracket_{(\tau,i)} = \oplus ((\& \mathbf{1}) \wp (\oplus \& ((\oplus \llbracket \sigma \rrbracket) \otimes (\llbracket \pi \rrbracket_{(\tau,i)})))) \\
 \llbracket \omega \rrbracket_{(\sigma,i)} = \begin{cases} \oplus ((\& \mathbf{1}) \wp (\oplus \& \mathbf{1})) & \text{if } i = 0 \\ \oplus ((\& \mathbf{1}) \wp (\oplus \& ((\oplus \llbracket \sigma \rrbracket) \otimes (\llbracket \omega \rrbracket_{(\sigma,i-1)})))) & \text{if } i > 0 \end{cases}
 \end{array}$$

Fig. 8. Translation of intersection types into session types (cf. Def. 5).

Theorem 5. *If $\Gamma \vDash M : \tau$, then $\llbracket M \rrbracket_u \vdash \llbracket \Gamma \rrbracket, u : \llbracket \tau \rrbracket$.*

To state *dynamic* correctness, we rely on established notions that (abstractly) characterize *correct translations*. A language $\mathcal{L} = (L, \rightarrow)$ consists of a set of terms L and a reduction relation \rightarrow on L . Each language \mathcal{L} is assumed to contain a success constructor \checkmark . A term $T \in L$ has *success*, denoted $T \Downarrow \checkmark$, when there is a sequence of reductions (using \rightarrow) from T to a term satisfying success criteria.

Given $\mathcal{L}_1 = (L_1, \rightarrow_1)$ and $\mathcal{L}_2 = (L_2, \rightarrow_2)$, we seek translations $\llbracket \cdot \rrbracket : L_1 \rightarrow L_2$ that are correct: they satisfy well-known correctness criteria [11,22,23]. We state the set of correctness criteria that determine the correctness of a translation.

Definition 6 (Correct Translation). *Let $\mathcal{L}_1 = (\mathcal{M}, \rightarrow_1)$ and $\mathcal{L}_2 = (\mathcal{P}, \rightarrow_2)$ be two languages. Let \simeq_2 be an equivalence over \mathcal{L}_2 . We use M, M' (resp. P, P') to range over terms in \mathcal{M} (resp. \mathcal{P}). Given a translation $\llbracket \cdot \rrbracket : \mathcal{M} \rightarrow \mathcal{P}$, we define:*

Completeness: *For every M, M' such that $M \rightarrow_1^* M'$, there exists P such that $\llbracket M \rrbracket \rightarrow_2^* P \simeq_2 \llbracket M' \rrbracket$.*

Weak Soundness: *For every M and P such that $\llbracket M \rrbracket \rightarrow_2^* P$, there exist M', P' such that $M \rightarrow_1^* M'$ and $P \rightarrow_2^* P' \simeq_2 \llbracket M' \rrbracket$.*

Success Sensitivity: *For every M , we have $M \Downarrow \checkmark$ if and only if $\llbracket M \rrbracket \Downarrow \checkmark$.*

Let us write Λ to denote the set of well-formed λ_c^{ℓ} terms, and Π for the set of all well-typed π^+ processes, both including \checkmark . We have our final result:

Theorem 6 (Translation correctness under \rightsquigarrow). *The translation $\llbracket \cdot \rrbracket_- : (\Lambda, \longrightarrow) \rightarrow (\Pi, \rightsquigarrow)$ is correct (cf. Definition 6) using equivalence \equiv (Figure 1).*

The proof of Theorem 6 involves instantiating/proving each of the parts of Def. 6. Among these, *weak soundness* is the most challenging to prove. Prior work on translations of typed λ into π with confluent non-determinism [20] rely critically on confluence to match a behavior in π with a corresponding behavior in λ . Because in our setting confluence is lost, we must resort to a different proof.

As already discussed, our translation makes the implicit non-determinism in a λ_c^{ℓ} -term M explicit by adding non-deterministic choices in key points of $\llbracket M \rrbracket_u$. Our reduction \rightsquigarrow preserves those branches that simultaneously have the same prefix available (up to \bowtie). In proving weak soundness, we exploit the fact that reduction entails delayed commitment. To see this, consider the following terms:

$$(\nu x)((\alpha_1; P_1 \# \alpha_2; P_2) \mid Q) \tag{4}$$

$$(\nu x)(\alpha_1; P_1 \mid Q) \# (\nu x)(\alpha_2; P_2 \mid Q) \tag{5}$$

In (4), commitment to a choice relies on whether $\alpha_1 \bowtie \alpha_2$ holds (cf. Definition 3). If $\alpha_1 \not\bowtie \alpha_2$, a choice is made; otherwise, commitment is delayed, and depends on P_1 and P_2 . Hence, in (4) the possibility of committing to either branch is kept open. In contrast, in (5) commitment to a choice is independent of $\alpha_1 \bowtie \alpha_2$.

Our translation exploits the delayed commitment of non-determinism illustrated by (4) to mimic commitment to non-deterministic choices in $\lambda_{\mathcal{C}}^{\ell}$, which manifests in fetching resources from bags. The fact that this delayed commitment preserves information about the different branches (e.g., P_1 and P_2 in (4)) is essential to establish weak soundness, i.e., to match a behavior in $\mathfrak{s}\pi^+$ with a corresponding step in $\lambda_{\mathcal{C}}^{\ell}$. In contrast, forms of non-determinism in $\llbracket N \rrbracket_u$ that resemble (5) are useful to characterize behaviors different from fetching.

5 Summary and Related Work

We studied the interplay between resource control and non-determinism in typed calculi. We introduced $\mathfrak{s}\pi^+$ and $\lambda_{\mathcal{C}}^{\ell}$, two calculi with non-confluent non-determinism, both with type systems for resource control. Inspired by the untyped π -calculus, non-determinism in $\mathfrak{s}\pi^+$ is lazy and explicit, with session types defined following ‘propositions-as-sessions’ [5]. In $\lambda_{\mathcal{C}}^{\ell}$, non-determinism arises in the fetching of resources, and is regulated by intersection types. A correct translation of $\lambda_{\mathcal{C}}^{\ell}$ into $\mathfrak{s}\pi^+$ precisely connects their different forms of non-determinism.

Related Work Integrating (non-confluent) non-determinism within session types is non-trivial, as carelessly discarding branches would break typability. Work by Caires and Pérez [5], already mentioned, develops a confluent semantics by requiring that non-determinism is only used inside the monad $\&A$; our non-confluent semantics drops this requirement. This allows us to consider non-deterministic choices not possible in [5], such as, e.g., selections of different labels. We stress that linearity is not jeopardized: the branches of ‘ $\#$ ’ do not represent *different sessions*, but *different implementations* of the same sessions.

Atkey et al. [1] and Kokke et al. [15] extend ‘propositions-as-sessions’ with non-determinism. Their approaches are very different (conflation of the additives and bounded linear logic, respectively) and support non-determinism for unrestricted names only. Also, [1,15] do not connect with typed λ -calculi, as we do. Rocha and Caires also consider non-determinism, relying on confluence in [24] and on unrestricted names in [25]. Casal et al. [7,29] develop a type system for *mixed sessions* (sessions with mixed choices), which can express non-determinism but does not ensure deadlock-freedom. Ensuring deadlock-freedom by typing is a key feature of the ‘propositions-as-sessions’ approach that we adopt for $\mathfrak{s}\pi^+$.

Our language $\lambda_{\mathcal{C}}^{\ell}$ is most related to calculi by Boudol [3], Boudol and Lanave [4], and by Pagani and Ronchi Della Rocca [19]. Non-determinism in the calculi in [3,4] is committing and implicit; their linear resources can be consumed *at most* once, rather than *exactly* once. The work [19] considers non-committing non-determinism that is both implicit (as in $\lambda_{\mathcal{C}}^{\ell}$) and explicit (via a sum operator on terms). Both [3,19] develop (non-idempotent) intersection type systems

to regulate resources. In our type system, all terms in a bag have the same type; the system in [19] does not enforce this condition. Unlike these type systems, our system for well-formedness can type terms with a lack or an excess of resources.

Boudol and Laneve [4] and Paulus et al. [20] give translations of resource λ -calculi into π . The translation in [4] is used to study the semantics induced upon λ -terms by a translation into π ; unlike ours, it does not consider types. As already mentioned in § 4, Paulus et al. [20] relate calculi with *confluent* non-determinism: a resource λ -calculus with sums on terms, and the session π -calculus from [5]. Our translation of terms and that in [20] are very different: while here we use non-deterministic choice to mimic the sharing construct, the translation in [20] uses it to translate bags. Hence, our Theorem 6 cannot be derived from [20].

The last decade of work on ‘propositions-as-sessions’ has delivered insightful connections with typed λ -calculi—see, e.g., [30,27,28]. Excepting [20], already discussed, none of these works consider non-deterministic λ -calculi.

Acknowledgments We are grateful to the anonymous reviewers for useful comments on previous versions of this paper. We are also grateful to Mariangiola Dezani for her encouragement and suggestions. This research has been supported by the Dutch Research Council (NWO) under project No. 016.Vidi.189.046 (‘Unifying Correctness for Communicating Software’) and the EPSRC Fellowship ‘VeTSpec: Verified Trustworthy Software Specification’ (EP/R034567/1).

References

1. Atkey, R., Lindley, S., Morris, J.G.: Conflation confers concurrency. In: Lindley, S., McBride, C., Trinder, P.W., Sannella, D. (eds.) A List of Successes That Can Change the World - Essays Dedicated to Philip Wadler on the Occasion of His 60th Birthday. Lecture Notes in Computer Science, vol. 9600, pp. 32–55. Springer (2016). https://doi.org/10.1007/978-3-319-30936-1_2, https://doi.org/10.1007/978-3-319-30936-1_2
2. Berger, M., Honda, K.: The two-phase commitment protocol in an extended pi-calculus. In: Aceto, L., Victor, B. (eds.) 7th International Workshop on Expressiveness in Concurrency, EXPRESS 2000, Satellite Workshop of CONCUR 2000, State College, PA, USA, August 21, 2000. Electronic Notes in Theoretical Computer Science, vol. 39, pp. 21–46. Elsevier (2000). [https://doi.org/10.1016/S1571-0661\(05\)82502-2](https://doi.org/10.1016/S1571-0661(05)82502-2), [https://doi.org/10.1016/S1571-0661\(05\)82502-2](https://doi.org/10.1016/S1571-0661(05)82502-2)
3. Boudol, G.: The lambda-calculus with multiplicities (abstract). In: Best, E. (ed.) CONCUR ’93, 4th International Conference on Concurrency Theory, Hildesheim, Germany, August 23-26, 1993, Proceedings. Lecture Notes in Computer Science, vol. 715, pp. 1–6. Springer (1993). https://doi.org/10.1007/3-540-57208-2_1, https://doi.org/10.1007/3-540-57208-2_1
4. Boudol, G., Laneve, C.: Lambda-calculus, multiplicities, and the pi-calculus. In: Proof, Language, and Interaction, Essays in Honour of Robin Milner. pp. 659–690 (2000)
5. Caires, L., Pérez, J.A.: Linearity, control effects, and behavioral types. In: Yang, H. (ed.) Programming Languages and Systems - 26th European Sympos-

- sium on Programming, ESOP 2017, Held as Part of the European Joint Conferences on Theory and Practice of Software, ETAPS 2017, Uppsala, Sweden, April 22-29, 2017, Proceedings. Lecture Notes in Computer Science, vol. 10201, pp. 229–259. Springer (2017). https://doi.org/10.1007/978-3-662-54434-1_9, https://doi.org/10.1007/978-3-662-54434-1_9
6. Caires, L., Pfenning, F.: Session types as intuitionistic linear propositions. In: CONCUR 2010 - Concurrency Theory, 21th International Conference, CONCUR 2010, Paris, France, August 31-September 3, 2010. Proceedings. pp. 222–236 (2010). https://doi.org/10.1007/978-3-642-15375-4_16, https://doi.org/10.1007/978-3-642-15375-4_16
 7. Casal, F., Mordido, A., Vasconcelos, V.T.: Mixed sessions. Theor. Comput. Sci. **897**, 23–48 (2022). <https://doi.org/10.1016/j.tcs.2021.08.005>, <https://doi.org/10.1016/j.tcs.2021.08.005>
 8. de'Liguoro, U., Piperno, A.: Non deterministic extensions of untyped lambda-calculus. Inf. Comput. **122**(2), 149–177 (1995). <https://doi.org/10.1006/inco.1995.1145>, <https://doi.org/10.1006/inco.1995.1145>
 9. Dezani-Ciancaglini, M.: Logical Semantics for Concurrent Lambda-Calculus. Ph.D. thesis, Nijmegen University (1996), <http://www.di.unito.it/~dezani/papers/tesi.ps>
 10. Ehrhard, T., Regnier, L.: The differential lambda-calculus. Theor. Comput. Sci. **309**(1-3), 1–41 (2003). [https://doi.org/10.1016/S0304-3975\(03\)00392-X](https://doi.org/10.1016/S0304-3975(03)00392-X), [https://doi.org/10.1016/S0304-3975\(03\)00392-X](https://doi.org/10.1016/S0304-3975(03)00392-X)
 11. Gorla, D.: Towards a unified approach to encodability and separation results for process calculi. Inf. Comput. **208**(9), 1031–1053 (2010). <https://doi.org/10.1016/j.ic.2010.05.002>, <https://doi.org/10.1016/j.ic.2010.05.002>
 12. Groote, J.F., Mousavi, M.R.: Modeling and Analysis of Communicating Systems. MIT Press (2014), <https://mitpress.mit.edu/books/modeling-and-analysis-communicating-systems>
 13. Honda, K.: Types for dyadic interaction. In: Best, E. (ed.) CONCUR '93, 4th International Conference on Concurrency Theory, Hildesheim, Germany, August 23-26, 1993, Proceedings. Lecture Notes in Computer Science, vol. 715, pp. 509–523. Springer (1993). https://doi.org/10.1007/3-540-57208-2_35, https://doi.org/10.1007/3-540-57208-2_35
 14. Honda, K., Vasconcelos, V.T., Kubo, M.: Language primitives and type discipline for structured communication-based programming. In: Hankin, C. (ed.) Programming Languages and Systems - ESOP'98, 7th European Symposium on Programming, Held as Part of the European Joint Conferences on the Theory and Practice of Software, ETAPS'98, Lisbon, Portugal, March 28 - April 4, 1998, Proceedings. Lecture Notes in Computer Science, vol. 1381, pp. 122–138. Springer (1998). <https://doi.org/10.1007/BFb0053567>, <https://doi.org/10.1007/BFb0053567>
 15. Kokke, W., Morris, J.G., Wadler, P.: Towards Races in Linear Logic. Logical Methods in Computer Science **Volume 16, Issue 4** (Dec 2020). [https://doi.org/10.23638/LMCS-16\(4:15\)2020](https://doi.org/10.23638/LMCS-16(4:15)2020)
 16. Milner, R.: Functions as processes. Research Report 1154, INRIA, Sophia Antipolis (1990), final version appeared as [?]
 17. Milner, R., Parrow, J., Walker, D.: A calculus of mobile processes, I. Inf. Comput. **100**(1), 1–40 (1992). [https://doi.org/10.1016/0890-5401\(92\)90008-4](https://doi.org/10.1016/0890-5401(92)90008-4), [https://doi.org/10.1016/0890-5401\(92\)90008-4](https://doi.org/10.1016/0890-5401(92)90008-4)

18. Nestmann, U., Fuzzati, R., Merro, M.: Modeling consensus in a process calculus. In: Amadio, R.M., Lugiez, D. (eds.) CONCUR 2003 - Concurrency Theory, 14th International Conference, Marseille, France, September 3-5, 2003, Proceedings. Lecture Notes in Computer Science, vol. 2761, pp. 393–407. Springer (2003). https://doi.org/10.1007/978-3-540-45187-7_26, https://doi.org/10.1007/978-3-540-45187-7_26
19. Pagani, M., Ronchi Della Rocca, S.: Solvability in resource lambda-calculus. In: Ong, C.L. (ed.) Foundations of Software Science and Computational Structures, 13th International Conference, FOSSACS 2010, Held as Part of the Joint European Conferences on Theory and Practice of Software, ETAPS 2010, Paphos, Cyprus, March 20-28, 2010. Proceedings. Lecture Notes in Computer Science, vol. 6014, pp. 358–373. Springer (2010). https://doi.org/10.1007/978-3-642-12032-9_25, https://doi.org/10.1007/978-3-642-12032-9_25
20. Paulus, J.W.N., Nantes-Sobrinho, D., Pérez, J.A.: Non-deterministic functions as non-deterministic processes. In: Kobayashi, N. (ed.) 6th International Conference on Formal Structures for Computation and Deduction, FSCD 2021, July 17-24, 2021, Buenos Aires, Argentina (Virtual Conference). LIPIcs, vol. 195, pp. 21:1–21:22. Schloss Dagstuhl - Leibniz-Zentrum für Informatik (2021). <https://doi.org/10.4230/LIPIcs.FSCD.2021.21>, <https://doi.org/10.4230/LIPIcs.FSCD.2021.21>, Extended version on <https://arxiv.org/abs/2104.14759>
21. Paulus, J.W.N., Nantes-Sobrinho, D., Pérez, J.A.: Types and Terms Translated: Unrestricted Resources in Encoding Functions as Processes (Extended Version). CoRR abs/2112.01593 (2021), <http://arxiv.org/abs/2112.01593>
22. Peters, K.: Translational Expressiveness. Comparing Process Calculi using Encodings. Ph.D. thesis, Berlin Institute of Technology (2012), <https://doi.org/10.14279/depositonce-3416>
23. Peters, K.: Comparing process calculi using encodings. In: Pérez, J.A., Rot, J. (eds.) Proceedings Combined 26th International Workshop on Expressiveness in Concurrency and 16th Workshop on Structural Operational Semantics, EXPRESS/SOS 2019, Amsterdam, The Netherlands, 26th August 2019. EPTCS, vol. 300, pp. 19–38 (2019). <https://doi.org/10.4204/EPTCS.300.2>, <https://doi.org/10.4204/EPTCS.300.2>
24. Rocha, P., Caires, L.: Propositions-as-types and shared state. Proceedings of the ACM on Programming Languages 5(ICFP), 79:1–79:30 (Aug 2021). <https://doi.org/10.1145/3473584>
25. Rocha, P., Caires, L.: Safe Session-Based Concurrency with Shared Linear State. In: Wies, T. (ed.) Programming Languages and Systems. pp. 421–450. Lecture Notes in Computer Science, Springer Nature Switzerland, Cham (2023). https://doi.org/10.1007/978-3-031-30044-8_16
26. Sangiorgi, D., Walker, D.: The Pi-Calculus - a theory of mobile processes. Cambridge University Press (2001)
27. Toninho, B., Caires, L., Pfenning, F.: Functions as session-typed processes. In: Birkedal, L. (ed.) Foundations of Software Science and Computational Structures - 15th International Conference, FOSSACS 2012, Held as Part of the European Joint Conferences on Theory and Practice of Software, ETAPS 2012, Tallinn, Estonia, March 24 - April 1, 2012. Proceedings. Lecture Notes in Computer Science, vol. 7213, pp. 346–360. Springer (2012). https://doi.org/10.1007/978-3-642-28729-9_23, https://doi.org/10.1007/978-3-642-28729-9_23

28. Toninho, B., Yoshida, N.: On polymorphic sessions and functions - A tale of two (fully abstract) encodings. In: Ahmed, A. (ed.) Programming Languages and Systems - 27th European Symposium on Programming, ESOP 2018, Held as Part of the European Joint Conferences on Theory and Practice of Software, ETAPS 2018, Thessaloniki, Greece, April 14-20, 2018, Proceedings. Lecture Notes in Computer Science, vol. 10801, pp. 827-855. Springer (2018). https://doi.org/10.1007/978-3-319-89884-1_29, https://doi.org/10.1007/978-3-319-89884-1_29
29. Vasconcelos, V.T., Casal, F., Almeida, B., Mordido, A.: Mixed sessions. In: Müller, P. (ed.) Programming Languages and Systems - 29th European Symposium on Programming, ESOP 2020, Held as Part of the European Joint Conferences on Theory and Practice of Software, ETAPS 2020, Dublin, Ireland, April 25-30, 2020, Proceedings. Lecture Notes in Computer Science, vol. 12075, pp. 715-742. Springer (2020). https://doi.org/10.1007/978-3-030-44914-8_26, https://doi.org/10.1007/978-3-030-44914-8_26
30. Wadler, P.: Propositions as sessions. In: Thiemann, P., Findler, R.B. (eds.) ACM SIGPLAN International Conference on Functional Programming, ICFP'12, Copenhagen, Denmark, September 9-15, 2012. pp. 273-286. ACM (2012). <https://doi.org/10.1145/2364527.2364568>, <https://doi.org/10.1145/2364527.2364568>

Table of Contents

1	Introduction	1
2	A Typed π -calculus with Non-deterministic Choice	3
2.1	Syntax and Semantics	4
2.2	Resource Control for $\mathfrak{s}\pi^+$ via Session Types	9
3	A Non-deterministic Resource λ -calculus	11
3.1	Syntax and Reduction Semantics	11
3.2	Resource Control for λ_C^ℓ via Intersection Types	13
4	A Typed Translation of λ_C^ℓ into $\mathfrak{s}\pi^+$	15
5	Summary and Related Work	18
	References	19
A	Full $\mathfrak{s}\pi^+$: Replicated Servers and Clients	23
B	An Alternative Eager Semantics for $\mathfrak{s}\pi^+$	24
C	Beyond Linear Resources	27
D	Extensions in Detail: Eager Semantics and Unrestricted Resources	28
D.1	λ_C : An Extension of λ_C^ℓ with Unrestricted Bags	28
D.2	Translating λ_C into (Full) $\mathfrak{s}\pi^+$	30
E	Proofs of Type Preservation and Deadlock-freedom for (Full) $\mathfrak{s}\pi^+$	30
E.1	Eager Semantics	31
E.2	Lazy Semantics	38
F	Proofs of Subject Reduction and Subject Expansion for λ_C	43
F.1	Subject Reduction	43
F.2	Subject Expansion	47
G	Proof of Tight Correctness of the Translation under the Lazy Semantics	51
G.1	Type Preservation	52
G.2	Completeness	58
G.3	Soundness	63
G.4	Success Sensitivity	72
H	Proof of Loose Correctness of the Translation under the Eager Semantics	73
H.1	Completeness	73
H.2	Soundness	77
H.3	Success Sensitivity	86
I	Proof of Separation of Lazy and Eager Semantics	86

A Full $\mathfrak{s}\pi^+$: Replicated Servers and Clients

Full $\mathfrak{s}\pi^+$ includes unrestricted session behaviors (replicated servers and clients), not presented in Section 2. Here we discuss how to add these omitted unrestricted sessions to the system described in Section 2. The proofs of Type Preservation (Theorem 1) and Deadlock-freedom (Theorem 2) in Appendix E concern Full $\mathfrak{s}\pi^+$.

- To the syntax of processes in Figure 1 (top) we add two prefixes, $?\bar{x}[y]; P$ and $!x(y); P$, for client requests and server definitions, respectively. Both prefixes bind y in P .
- Now $fn(P)$ denotes the set of free names of P , including those used for unrestricted sessions. We write $fln(P)$ to denote the set of free linear names of P , and $fnp(P)$ for the set of free non-linear names. Note that $fnp(P) = fn(P) \setminus fln(P)$.
- To the structural congruence in Figure 1 (bottom) we add a rule that cleans up unused servers:

$$(\nu x)(!x(y); P \mid Q) \equiv Q \quad (\text{if } x \notin fn(Q)).$$

- To the lazy semantics in Figure 2 we add the following rule that initiates a session between a client and a copy of a server:

$$\begin{aligned} [\rightsquigarrow ?!] \quad & (\nu x) \left(\prod_{i \in I} C_i[?\bar{x}[y_i]; P_i] \mid \prod_{j \in J} D_j[!x(z); Q_j] \right) \\ & \rightsquigarrow_x \prod_{j \in J} D_j \left[(\nu w) \left(\prod_{i \in I} C_i[P_i\{w/z\}] \mid Q_j\{w/z\} \right) \mid !x(z); Q_j \right] \end{aligned}$$

$$\begin{array}{c}
[\rightarrow_{\text{Id}}] \quad (\nu x)(\mathbb{N}[x \leftrightarrow y] \mid Q) \longrightarrow (\mathbb{N})[Q\{y/x\}] \qquad [\rightarrow_{\text{11}}] \quad (\nu x)(\mathbb{N}[\bar{x}[]] \mid \mathbb{N}'[x(z); Q]) \longrightarrow (\mathbb{N})[\mathbf{0}] \mid (\mathbb{N}')[Q] \\
[\rightarrow_{\otimes \otimes}] \quad (\nu x)(\mathbb{N}[\bar{x}[y]; (P \mid Q)] \mid \mathbb{N}'[x(z); R]) \longrightarrow (\mathbb{N})[(\nu x)(Q \mid (\nu y)(P \mid (\mathbb{N}')[R\{y/z\}]))] \\
[\rightarrow_{\oplus \&}] \quad \forall k' \in K. (\nu x)(\mathbb{N}[\bar{x}.k'; P] \mid \mathbb{N}'[x.\text{case}\{k : Q^k\}_{k \in K}]) \longrightarrow (\nu x)((\mathbb{N})[P] \mid (\mathbb{N}')[Q^{k'}]) \\
[\rightarrow_{?!}] \quad (\nu x)(\mathbb{N}[\bar{x}[y]; P] \mid \mathbb{N}'[!x(z); Q]) \longrightarrow (\mathbb{N}')[(\nu x)((\nu y)((\mathbb{N})[P] \mid Q\{y/z\}) \mid !x(z); Q)] \\
[\rightarrow_{\text{some}}] \quad (\nu x)(\mathbb{N}[\bar{x}.\text{some}; P] \mid \mathbb{N}'[x.\text{some}_{w_1, \dots, w_n}; Q]) \longrightarrow (\nu x)((\mathbb{N})[P] \mid (\mathbb{N}')[Q]) \\
[\rightarrow_{\text{none}}] \quad (\nu x)(\mathbb{N}[\bar{x}.\text{none}] \mid \mathbb{N}'[x.\text{some}_{w_1, \dots, w_n}; Q]) \longrightarrow (\mathbb{N})[\mathbf{0}] \mid (\mathbb{N}')[\bar{w}_1.\text{none} \mid \dots \mid \bar{w}_n.\text{none}] \qquad [\rightarrow_{\equiv}] \quad \frac{P \equiv P' \quad P' \longrightarrow Q' \quad Q' \equiv Q}{P \longrightarrow Q} \\
[\rightarrow_{\nu}] \quad \frac{P \longrightarrow P'}{(\nu x)(P \mid Q) \longrightarrow (\nu x)(P' \mid Q)} \qquad [\rightarrow_{\mid}] \quad \frac{P \longrightarrow P'}{P \mid Q \longrightarrow P' \mid Q} \qquad [\rightarrow_{\ddagger}] \quad \frac{P \longrightarrow P'}{P \ddagger Q \longrightarrow P' \ddagger Q}
\end{array}$$

Fig. 9. Eager reduction semantics for $\mathfrak{s}\pi^+$.

– Writing $? \Gamma$ to denote that $\forall x \in \Gamma. \exists A. \Gamma(x) = ?A$, to the typing rules in Figure 3 we add:

$$[\text{T?}] \quad \frac{P \vdash \Gamma, y : A}{?\bar{x}[y]; P \vdash \Gamma, x : ?A} \qquad [\text{T!}] \quad \frac{P \vdash ?\Gamma, y : A}{!x(y); P \vdash ?\Gamma, x : !A} \qquad [\text{TWEAKEN}] \quad \frac{P \vdash \Gamma}{P \vdash \Gamma, x : ?A} \qquad [\text{TCONTRACT}] \quad \frac{P \vdash \Gamma, x : ?A, x' : ?A}{P\{x/x'\} \vdash \Gamma, x : ?A}$$

Moreover, we replace Rule $[\text{T}\oplus\text{some}]$ with the following:

$$[\text{T}\oplus\text{some}] \quad \frac{P \vdash \&\Gamma, ?\Delta, x : A}{x.\text{some}_{\text{dom}(\Gamma)}; P \vdash \&\Gamma, ?\Delta, x : \oplus A}$$

B An Alternative Eager Semantics for $\mathfrak{s}\pi^+$

Let us consider a variant of $\mathfrak{s}\pi^+$ in which syntax, typing, and structural congruence are as in §2, but with an *eagerly committing* semantics. The idea is simple: we fully commit to a non-deterministic choice once a prefix synchronizes.

The eager reduction semantics, denoted \longrightarrow , is given in Figure 9. This semantics implements the full commitment of non-deterministic choices by committing ND-contexts to D-contexts as follows:

Definition 7. The commitment of an ND-context \mathbb{N} , denoted (\mathbb{N}) , is defined as follows:

$$((\cdot)) := [\cdot] \qquad (\mathbb{N} \mid P) := (\mathbb{N}) \mid P \qquad ((\nu x)(\mathbb{N} \mid P)) := (\nu x)((\mathbb{N}) \mid P) \qquad (\mathbb{N} \ddagger P) := (\mathbb{N})$$

Proposition 1. For any ND-context \mathbb{N} , the context (\mathbb{N}) is a D-context.

Just as $\mathfrak{s}\pi^+$ with lazy semantics, $\mathfrak{s}\pi^+$ with \longrightarrow satisfies type preservation and deadlock-freedom. See Appendix E.1 for details.

Theorem 7 (Type Preservation: Eager Semantics). If $P \vdash \Gamma$, then both $P \equiv Q$ and $P \longrightarrow Q$ (for any S) imply $Q \vdash \Gamma$.

Proof (Sketch). If $P \equiv Q$, the thesis follows directly from Theorem 1. If $P \longrightarrow Q$, we apply induction on the derivation of the reduction. In each case, we show that the commitment of ND-contexts (Definition 7) preserves typing. \square

Theorem 8 (Deadlock-freedom: Eager Semantics). If $P \vdash \emptyset$ and $P \not\equiv \mathbf{0}$, then there is R such that $P \longrightarrow R$.

Proof (Sketch). First, we write P in such a way that we can access all its unblocked prefixes. Then we inductively show that there must be at least one pair of such prefixes that are connected by a restriction. Hence, these prefixes are duals and thus the process can reduce. \square

It is insightful to formally contrast our lazy and eager semantics. We discuss two different ways.




Fig. 10. Example 7: Reductions of M and of $\llbracket M \rrbracket_u$ under lazy and eager semantics. In $\llbracket M \rrbracket_u$, we write ‘ $\pi(X)$ ’ for the permutations of finite set X .

Lazy vs Eager, Part I: Translating λ_c^ℓ . One way of comparing \rightsquigarrow and \longrightarrow is to establish the correctness of the translation $\llbracket \cdot \rrbracket_-$ (Figure 7) but now in the eager case. It turns out that the eager semantics leads to a *strictly weaker* form of correctness, whereby completeness and weak soundness (cf. Definition 6) hold up to a precongurence \succeq_{\amalg} instead of an equivalence (as in Theorem 6).

The precongurence \succeq_{\amalg} is defined as follows:

$$\frac{}{P \succeq_{\amalg} P} \quad \frac{P_i \succeq_{\amalg} P'_i \quad i \in \{1, 2\}}{P_1 \amalg P_2 \succeq_{\amalg} P'_i} \quad \frac{P \succeq_{\amalg} P' \quad Q \succeq_{\amalg} Q'}{P \mid Q \succeq_{\amalg} P' \mid Q'} \quad \frac{P \succeq_{\amalg} P'}{(\nu x)P \succeq_{\amalg} (\nu x)P'}$$

Intuitively, $P \succeq_{\amalg} Q$ says that P has at least as many branches as Q . Translation correctness up to \succeq_{\amalg} thus means that \longrightarrow is “too eager”, as it *prematurely commits* to branches.

App. G proves that the translation under the eager semantics satisfies such criteria. Before discussing the corresponding completeness and soundness results, we present an example.

Example 7. To contrast commitment in eager and lazy semantics (and their effect on the translation’s correctness), recall from Example 6 the term M (3):

$$M = (x_1 \lambda x_2 \lambda x_3 \mathbf{1} \mathcal{J} \mathcal{J}) \langle \lambda \mathbf{fail}^0, y, I \mathcal{J} / x_1, x_2, x_3 \rangle$$

Figure 10 recalls the three branching reductions from M to N_1 , N_2 and N_3 . It also depicts a side-by-side comparison of the reductions of $\llbracket M \rrbracket_u$ under the lazy (\rightsquigarrow) and eager (\longrightarrow) semantics. In the figure, \rightsquigarrow^* and \longrightarrow^* denote the reflexive, transitive closures of \rightsquigarrow and \longrightarrow , respectively.

Under \rightsquigarrow there are three different reduction paths, each resulting directly in the translation of one of N_1, N_2, N_3 : after the first choice, the following choices are preserved. In contrast, under \longrightarrow there are six different reduction paths, each resulting in a process that relates to the translation of one of N_1, N_2, N_3 through \succeq_{\amalg} : after the first choice for an item from the bag is made, the semantics commits to choices for the other items. ∇

The correctness properties induced by \longrightarrow are *loose* (rather than *tight*, as in § 4; cf. Definition 6):

Theorem 9 (Loose Completeness (Under \longrightarrow)). *If $N \longrightarrow M$ for a well-formed closed λ_c -term N , then there exists Q such that $\llbracket N \rrbracket_u \longrightarrow^* Q$ and $\llbracket M \rrbracket_u \succeq_{\amalg} Q$.*

Proof (Sketch). By induction on reductions. See § H.1 for details. \square

Theorem 10 (Loose Weak Soundness (Under \longrightarrow)). *If $\llbracket N \rrbracket_u \longrightarrow^* Q$ for a well-formed closed λ_c -term N , then there exist N' and Q' such that (i) $N \longrightarrow^* N'$ and (ii) $Q \longrightarrow^* Q'$ with $\llbracket N' \rrbracket_u \succeq_{\amalg} Q'$.*

While \rightsquigarrow reduces multiple branches of a choice in lockstep, \longrightarrow reduces only one branch and discards the rest. Accordingly, weak soundness under \rightsquigarrow (Theorem 21) relates a sequence of lazy reductions to a sequence of reductions in $\lambda_{\mathbb{C}}^{\ell}$. In contrast, Theorem 10 is weaker: it relates a sequence of eager reductions to a subset of branches, as some branches may have been eagerly discarded. Hence, the proof of Theorem 10 has the added complexity of showing that every branch that is eagerly discarded must also be precongruent to a source reduction. This makes it difficult to apply induction directly, as we do not know which branches have been discarded in $s\pi^+$.

More in details, to prove Theorem 10, we first show that $\succeq_{\#}$ is stable under reductions. We need a way of denoting all possible reductions from a process. We define $P \longrightarrow \{P_i\}_{i \in I}$, for a fixed (maximal) finite set $I = \{i \mid P \longrightarrow P_i\}$. Similarly, we define $P \longrightarrow^* \{P_i\}_{i \in I}$ inductively: if $P \longrightarrow^* \{P_i\}_{i \in I}$ and $P_i \longrightarrow \{P_j\}_{j \in J_i}$ for each $i \in I$, then $P \longrightarrow^* \{P_j\}_{j \in J}$ with $J = \bigcup_{i \in I} J_i$. We then have the following:

Proposition 2. *If $P \succeq_{\#} Q$ and $P \longrightarrow^* \{P_i\}_{i \in I}$, then there exist J and $\{Q_j\}_{j \in J}$ such that $Q \longrightarrow^* \{Q_j\}_{j \in J}$, $J \subseteq I$, and for each $j \in J$, $P_j \succeq_{\#} Q_j$.*

Proof (Sketch). By induction on the derivation rules of the precongruence $\succeq_{\#}$. See § H.2 for details. \square

Then, using Proposition 2 we prove the following:

Lemma 1. *Let N be a well-formed closed term. If $\llbracket N \rrbracket_u \longrightarrow^* Q$, then there exist N' and $\{Q_i\}_{i \in I}$ such that (i) $N \longrightarrow^* N'$ and (ii) $Q \longrightarrow^* \{Q_i\}_{i \in I}$ where for each $j \in I$, $\llbracket N' \rrbracket_u \succeq_{\#} Q_j$.*

Lemma 1 ensures that the translation does not add behaviors not present in the source term. Theorem 10 then follows directly from Lemma 1 by taking an arbitrary Q_i .

Lazy vs Eager, Part II: Behavioral Equivalence. One may ask if the differences between \rightsquigarrow and \longrightarrow are confined to the ability to correctly translate $\lambda_{\mathbb{C}}^{\ell}$, or, relatedly, whether $\lambda_{\mathbb{C}}^{\ell}$'s formulation is responsible for these differences.

We now compare \rightsquigarrow and \longrightarrow independently from $\lambda_{\mathbb{C}}^{\ell}$ by resorting to *behavioral equivalences*. We define a simple behavioral notion of equivalence on $s\pi^+$ processes, parametric in \rightsquigarrow or \longrightarrow ; then, we prove that there are classes of processes that are equal with respect to \rightsquigarrow , but incomparable with respect to \longrightarrow (Theorem 11). A key ingredient is the following notion of observable on processes:

Definition 8. *A process P has a ready-prefix α , denoted $P \downarrow_{\alpha}$, if and only if there exist \mathbb{N}, P' such that $P \equiv \mathbb{N}[\alpha; P']$.*

We may now define:

Definition 9 (Ready-Prefix Bisimilarity). *A relation \mathbb{B} on $s\pi^+$ processes is a (strong) ready-prefix bisimulation with respect to \rightsquigarrow if and only if, for every $(P, Q) \in \mathbb{B}$,*

1. *For every P' such that $P \rightsquigarrow P'$, there exists Q' such that $Q \rightsquigarrow Q'$ and $(P', Q') \in \mathbb{B}$;*
2. *For every Q' such that $Q \rightsquigarrow Q'$, there exists P' such that $P \rightsquigarrow P'$ and $(P', Q') \in \mathbb{B}$;*
3. *For every $\alpha \bowtie \beta$, $P \downarrow_{\alpha}$ if and only if $Q \downarrow_{\beta}$.*

P and Q are ready-prefix bisimilar with respect to \rightsquigarrow , denoted $P \sim_{\mathbb{L}} Q$, if there exists a relation \mathbb{B} that is a ready-prefix bisimulation with respect to \rightsquigarrow such that $(P, Q) \in \mathbb{B}$.

A (strong) ready-prefix bisimulation with respect to \longrightarrow is defined by replacing every occurrence of ' \rightsquigarrow ' by ' \longrightarrow ' in the definition above. We write $P \sim_{\mathbb{E}} Q$ if P and Q are ready-prefix bisimilar with respect to \longrightarrow .

Ready-prefix bisimulation can highlight a subtle but significant difference between the behavior induced by our lazy and eager semantics. To demonstrate this, we consider session-typed implementations of a vending machine.

Example 8 (Two Vending Machines). Consider vending machines VM_1 and VM_2 consisting of three parts: (1) an interface, which interacts with the user to send money and choose between coffee (c) and tea (t); (2) a brewer, which produces either beverage; (3) a system, which collects the money and forwards the user's choice to the brewer. A $s\pi^+$ specification follows (below \mathbb{C} and $\mathbb{C}2$ stand for names):

$$\begin{aligned}
\text{VM}_1 &:= (\nu x)(\text{IF}_1 \mid (\nu y)(\text{Brewer} \mid \text{System})) \\
\text{VM}_2 &:= (\nu x)(\text{IF}_2 \mid (\nu y)(\text{Brewer} \mid \text{System})) \\
\text{IF}_1 &:= \bar{x}[\mathbb{C}2]; (\overline{\mathbb{C}2} \mid (\bar{x}.c; \bar{x} \mid \bar{x}.t; \bar{x})) \\
\text{IF}_2 &:= \bar{x}[\mathbb{C}2]; (\overline{\mathbb{C}2} \mid \bar{x}.c; \bar{x}) \mid \bar{x}[\mathbb{C}2]; (\overline{\mathbb{C}2} \mid \bar{x}.t; \bar{x}) \\
\text{System} &:= x(\mathbb{C}); x.\text{case} \left\{ \begin{array}{l} c : \bar{y}.c; x(); \mathbb{C}(); \bar{y}[] \\ t : \bar{y}.t; x(); \mathbb{C}(); \bar{y}[] \end{array} \right\} \\
\text{Brewer} &:= y.\text{case} \{ c : y(); \text{Brew}_c, t : y(); \text{Brew}_t \}
\end{aligned}$$

$M, N, L ::= x[*]$	variable	$M[\tilde{x} \leftarrow x]$	sharing
$\mid \lambda x.M$	abstraction	$M\langle\langle B/x \rangle\rangle$	intermediate substitution
$\mid (M B)$	application	$M\langle C/x_1, \dots, x_k \rangle$	linear substitution
$\mid \mathbf{fail}^{\tilde{x}}$	failure	$M\llbracket U/x \rrbracket$	unrestricted substitution
$[*] ::= [1] \mid [i] \quad i \in \mathbb{N}$	annotations	$A, B ::= C \star U$	bag
$U, V ::= 1^! \mid \wr M \wr^! \mid U \diamond V$	unrestricted bag	$C, D ::= 1 \mid \wr M \wr \cdot C$	linear bag
$\mathcal{C} ::= [\cdot] \mid (\mathcal{C} B) \mid \mathcal{C}\langle C/\tilde{x} \rangle \mid \mathcal{C}\llbracket U/x \rrbracket \mid \mathcal{C}[\tilde{x} \leftarrow x]$			context

Fig. 11. Syntax of $\lambda_{\mathcal{C}}$.

where $\text{Brew}_{\mathcal{C}} \vdash \emptyset$, $\text{Brew}_{\mathfrak{t}} \vdash \emptyset$, such that $\text{VM}_1 \vdash \emptyset$, $\text{VM}_2 \vdash \emptyset$.

We give two implementations of the interface: IF_1 sends the money and then chooses coffee or tea; IF_2 chooses sending the money and then requesting coffee, or sending the money and then requesting tea. Then, IF_1 and IF_2 result in two different vending machines, VM_1 and VM_2 .

We have $\text{VM}_1 \not\sim_{\mathbb{E}} \text{VM}_2$: the eager semantics distinguishes between the implementations; e.g., IF_1 has a single money slot, a button for coffee, and another button for tea, whereas IF_2 has two money slots, one for coffee, and another for tea. In contrast, under the lazy semantics, these machines are indistinguishable: $\text{VM}_1 \sim_{\mathbb{L}} \text{VM}_2$. ∇

Example 8 highlights a difference in behavior between \rightsquigarrow and \longrightarrow when a moment of choice is subtly altered. The following theorem captures this distinction (see § I):

Theorem 11. *Take $R \equiv \mathbb{N}[\alpha_1; (P \parallel Q)] \vdash \emptyset$ and $S \equiv \mathbb{N}[\alpha_2; P \parallel \alpha_3; Q] \vdash \emptyset$, where $\alpha_1 \bowtie \alpha_2 \bowtie \alpha_3$ and $\alpha_1, \alpha_2, \alpha_3$ require a continuation. Suppose that $P \not\sim_{\mathbb{L}} Q$ and $P \not\sim_{\mathbb{E}} Q$. Then (i) $R \sim_{\mathbb{L}} S$ but (ii) $R \not\sim_{\mathbb{E}} S$.*

Processes (4) and (5) from § 4 (Page 17) provide another example of a change in the moment of choice, different from the one discussed above. In (4), the choice depends on α_1 and α_2 . In (5), the choice does not depend on α_1 or α_2 , but on choices made in the context in which the process resides. Though the choice is subtly changed between (4) and (5), the impact is significant: these processes are not ready-prefix bisimilar, with respect to neither semantics. This is because, under both lazy and eager semantics, in (4) the two branches evolve in lockstep, whereas in (5) they evolve independently.

C Beyond Linear Resources

Our results extend to the language $\lambda_{\mathcal{C}}$, an extension of $\lambda_{\mathcal{C}}^{\ell}$ with a more general bag which includes unrestricted resources: resources that may be used zero or more times. Figure 11 gives the syntax of $\lambda_{\mathcal{C}}$ -terms, bags and contexts.

A key difference with $\lambda_{\mathcal{C}}^{\ell}$ is that variables x, y, z, \dots have linear and unrestricted occurrences. Notation $x[1]$ denotes a *linear* occurrence of x ; we often omit the annotation ‘[1]’, and a sequence \tilde{x} always involves linear occurrences. Notation $x[i]$ denotes an *unrestricted* occurrence of x , explicitly referencing the i -th element of an unrestricted (ordered) bag. The structure of a bag is now split into a linear and an unrestricted component: as in $\lambda_{\mathcal{C}}^{\ell}$, linear resources in bags cannot be duplicated, but unrestricted resources are always duplicated when consumed. The empty unrestricted bag is denoted $1^!$. Notation U_i denotes the singleton bag at the i -th position in U ; if there is no i -th position in U , then U_i defaults to $1^!$. We use ‘ \star ’ to combine a linear and an unrestricted bag, and unrestricted bags are joined via the non-commutative ‘ \diamond ’.

To account for the explicit distinction between linear and unrestricted occurrences of variables, we now have two forms of explicit substitution:

- $M\langle C/x_1, \dots, x_k \rangle$, a linear substitution as in $\lambda_{\mathcal{C}}^{\ell}$;
- $M\llbracket U/x^! \rrbracket$, an unrestricted substitution of an unrestricted bag U for an unrestricted variable $x^!$ in M , with the assumption that $x^!$ does not appear in another unrestricted substitution in M .

The reduction rules for $\lambda_{\mathcal{C}}$, extend the rules given in Figure 5 with some modifications to accommodate the two-component format of bags and dedicated rules for the new constructs. Here, we describe the most interesting new rules: Rule [RS:Ex-Sub] for explicit substitution, and also Rules [R:Fetch[!]] and [R:Fail[!]] for unrestricted substitution.

$$\begin{array}{c}
\text{[RS:Ex-Sub]} \\
\hline
\text{size}(C) = |\tilde{x}| \quad M \neq \mathbf{fail}^{\tilde{y}} \\
(M[\tilde{x} \leftarrow x])\langle\langle C \star U/x \rangle\rangle \longrightarrow M\langle C/\tilde{x} \rangle\llbracket U/x \rrbracket
\end{array}
\qquad
\begin{array}{c}
\text{[RS:Fetch}^! \\
\hline
\text{head}(M) = x[i] \quad U_i = \wr N \wr^! \\
M\llbracket U/x \rrbracket \longrightarrow M\{N/x[i]\}\llbracket U/x \rrbracket
\end{array}
\qquad
\begin{array}{c}
\text{[RS:Fail}^! \\
\hline
\text{head}(M) = x[i] \quad U_i = 1^! \\
M\llbracket U/x \rrbracket \longrightarrow M\{\mathbf{fail}^0/x[i]\}\llbracket U/x \rrbracket
\end{array}$$

An explicit substitution $(M[\tilde{x} \leftarrow x])\langle\langle C \star U/x \rangle\rangle$ reduces to a term in which the linear and unrestricted parts of the bag are separated into their own explicit substitutions $M\langle C/\tilde{x} \rangle\llbracket U/x \rrbracket$, if successful. The fetching of linear/unrestricted resources

from their corresponding bags is done by the appropriated fetch rules. The reduction of an unrestricted substitution $M\llbracket U/x \rrbracket$, where the head variable of M is $x[i]$, depends on U_i :

- If $U_i = \llbracket N \rrbracket^!$, then the term reduces via Rule $[\mathbf{R} : \mathbf{Fetch}^!]$ by substituting the head occurrence of $x[i]$ in M with N , denoted $M\{N/x[i]\}$; note that U_i remains available after this reduction.
- If $U_i = \mathbf{1}^!$, the head variable is instead substituted with failure via Rule $[\mathbf{R} : \mathbf{Fail}^!]$.

The definition of $\text{head}(M)$ is as in Figure 5 (bottom), extended with $\text{head}(x[i]) = x[i]$ and $\text{head}(M\llbracket U/x \rrbracket) = \text{head}(M)$. The complete set of rules is in App. D.1 .

Well-typedness and well-formedness. Types for $\lambda_{\mathbf{c}}$ extend the types for $\lambda_{\mathbf{c}}^{\ell}$ in Section 3.2 with:

$$\begin{aligned} \sigma, \tau, \delta &::= \mathbf{unit} \mid (\pi, \eta) \rightarrow \sigma \\ \eta, \epsilon &::= \sigma \mid \epsilon \diamond \eta \quad \text{list} \qquad (\pi, \eta) \quad \text{tuple} \end{aligned}$$

The list type $\epsilon \diamond \eta$ types the concatenation of unrestricted bags. It can be recursively unfolded into a finite composition of strict types $\sigma_1 \diamond \dots \diamond \sigma_n$, for some $n \geq 1$, with length n and σ_i its i -th strict type ($1 \leq i \leq n$). We write $x^! : \eta$ to denote for $x[1] : \eta_1, \dots, x[k] : \eta_k$ where η has length k . The tuple type (π, η) types concatenation of a linear bag of type π with an unrestricted bag of type η . Finally strict types are amended to allow for unrestricted functional types which go from tuple types to strict types $(\pi, \eta) \rightarrow \sigma$ rather than multiset types to strict types.

We separate contexts into two parts: linear (Γ, Δ, \dots) and unrestricted $(\Theta, \Upsilon, \dots)$:

$$\begin{aligned} \Gamma, \Delta &::= - \mid \Gamma, x : \pi \mid \Gamma, x : \sigma & \Theta, \Upsilon &::= - \mid \Theta, x^! : \eta \end{aligned}$$

Both linear and unrestricted occurrences of variables may occur at most once in a context. Judgments have the form $\Theta; \Gamma \vDash M : \tau$. We write $\vDash M : \tau$ to denote $-; - \vDash M : \tau$.

Well-formedness rules for $\lambda_{\mathbf{c}}$ extend the rules given in Figure 6 with specific rules to handle unrestricted resources, among those, we select Rules $[\mathbf{FS} : \mathbf{var}^!]$, $[\mathbf{FS} : \mathbf{abs-sh}]$, $[\mathbf{FS} : \mathbf{Esub}^!]$ and $[\mathbf{FS} : \mathbf{bag}^!]$, for typing unrestricted occurrences of a variable, abstraction of a sharing term, explicit substitution and unrestricted bags, respectively.

$$\begin{array}{c} \frac{[\mathbf{FS} : \mathbf{var}^!]}{\Theta, x^! : \eta; x : \eta_i, \Delta \vDash x : \sigma}{\Theta, x^! : \eta; \Delta \vDash x[i] : \sigma} \qquad \frac{[\mathbf{FS} : \mathbf{abs-sh}]}{\Theta, x^! : \eta; \Gamma, x : \sigma^k \vDash M[\tilde{x} \leftarrow x] : \tau \quad x \notin \text{dom}(\Gamma)}{\Theta; \Gamma \vDash \lambda x. (M[\tilde{x} \leftarrow x]) : (\sigma^k, \eta) \rightarrow \tau} \\ \\ \frac{[\mathbf{FS} : \mathbf{Esub}^!]}{\Theta, x^! : \eta; \Gamma \vDash M : \tau \quad \Theta; - \vDash U : \epsilon \quad \eta \sim \epsilon}{\Theta; \Gamma \vDash M\llbracket U/x^! \rrbracket : \tau} \qquad \frac{[\mathbf{FS} : \mathbf{bag}^!]}{\Theta; - \vDash U : \epsilon \quad \Theta; - \vDash V : \eta}{\Theta; - \vDash U \diamond V : \epsilon \diamond \eta} \end{array}$$

In Rule $[\mathbf{FS} : \mathbf{Esub}^!]$, $\eta \sim \epsilon$ denotes the fact that $\epsilon = \epsilon_1 \diamond \dots \diamond \epsilon_k \diamond \dots \diamond \epsilon_n$ and $\eta = \eta_1 \diamond \dots \diamond \eta_k$ are two list types, with $n \geq k$, such that the following hold: for all i , $1 \leq i \leq k$, $\epsilon_i = \eta_i$. The complete set of well-formedness rules for $\lambda_{\mathbf{c}}$ is in App. D.1 .

Extended translation. To extend the translation in Fig. 7 to $\lambda_{\mathbf{c}}$: The access and use of unrestricted resources in $\lambda_{\mathbf{c}}$ is codified in $s\pi^+$ by combining labeled choices and clients/servers (App. D.2).

Note: for the sake of generality the proofs in the appendices concern $\lambda_{\mathbf{c}}$.

D Extensions in Detail: Eager Semantics and Unrestricted Resources

D.1 $\lambda_{\mathbf{c}}$: An Extension of $\lambda_{\mathbf{c}}^{\ell}$ with Unrestricted Bags

Figure 12 defines the free linear variables of a term, denoted $\text{lfv}(M)$. Figure 13 gives the reduction rules for $\lambda_{\mathbf{c}}$, extending the rules given in Figure 5 with Rules $[\mathbf{R} : \mathbf{Fetch}^!]$ and $[\mathbf{R} : \mathbf{Fail}^!]$ for unrestricted substitution.

$$\begin{array}{ll}
\text{fv}(x) = \{x\} & \text{fv}((M B)) = \text{fv}(M) \cup \text{fv}(B) \\
\text{fv}(x[i]) = \emptyset & \text{fv}(M[\tilde{x} \leftarrow x]) = (\text{fv}(M) \setminus \{\tilde{x}\}) \cup \{x\} \\
\text{fv}(\mathbf{1}) = \emptyset & \text{fv}(\lambda x.(M[\tilde{x} \leftarrow x])) = \text{fv}(M[\tilde{x} \leftarrow x]) \setminus \{x\} \\
\text{fv}(\lambda M) = \text{fv}(M) & \text{fv}(M\langle C/x_1, \dots, x_k \rangle) = (\text{fv}(M) \setminus \{x_1, \dots, x_k\}) \cup \text{fv}(C) \\
\text{fv}(C \star U) = \text{fv}(C) & \text{fv}(M\langle\langle B/x \rangle\rangle) = (\text{fv}(M) \setminus \{x\}) \cup \text{fv}(B) \\
\text{fv}(\mathbf{fail}^{\tilde{x}}) = \{\tilde{x}\} & \text{fv}(\lambda M \int \cdot C) = \text{fv}(M) \cup \text{fv}(C)
\end{array}$$

In the case $\text{fv}(M) = \emptyset$, the term M is called *linearly closed*.

Fig. 12. Free Linear Variables

$$\begin{array}{c}
\begin{array}{lll}
\text{[RS:Beta]} & \text{[RS:Ex-Sub]} & \text{[RS:Fetch}^\ell] \\
\frac{}{(\lambda x.M) B \rightarrow M\langle\langle B/x \rangle\rangle} & \frac{\text{size}(C) = |\tilde{x}| \quad M \neq \mathbf{fail}^{\tilde{y}}}{(M[\tilde{x} \leftarrow x])\langle\langle C \star U/x \rangle\rangle \rightarrow M\langle\langle C/\tilde{x} \rangle\rangle\langle\langle U/x \rangle\rangle} & \frac{\text{head}(M) = x_j \quad 0 < i \leq \text{size}(C)}{M\langle\langle C/\tilde{x}, x_j \rangle\rangle \rightarrow (M\{C_i/x_j\})\langle\langle (C \setminus C_i)/\tilde{x} \rangle\rangle} \\
\text{[RS:Fail}^\ell] & \text{[RS:Fetch}^! & \text{[RS:Fail}^! \\
\frac{\text{size}(C) \neq |\tilde{x}| \quad \tilde{y} = (\text{fv}(M) \setminus \{\tilde{x}\}) \cup \text{fv}(C)}{(M[\tilde{x} \leftarrow x])\langle\langle C \star U/x \rangle\rangle \rightarrow \mathbf{fail}^{\tilde{y}}} & \frac{\text{head}(M) = x[i] \quad U_i = \lambda N \int^!}{M\langle\langle U/x \rangle\rangle \rightarrow M\{N/x[i]\}\langle\langle U/x \rangle\rangle} & \frac{\text{head}(M) = x[i] \quad U_i = \mathbf{1}^!}{M\langle\langle U/x \rangle\rangle \rightarrow M\{\mathbf{fail}^0/x[i]\}\langle\langle U/x \rangle\rangle} \\
\text{[RS:Cons}_1] & \text{[RS:Cons}_2] & \text{[RS:Cons}_3] \\
\frac{\tilde{y} = \text{fv}(C)}{\mathbf{fail}^{\tilde{x}}(C \star U) \rightarrow \mathbf{fail}^{\tilde{x} \tilde{y}}} & \frac{\text{size}(C) = |\tilde{x}| \quad \tilde{z} = \text{fv}(C)}{(\mathbf{fail}^{\tilde{x} \cup \tilde{y}}[\tilde{x} \leftarrow x])\langle\langle C \star U/x \rangle\rangle \rightarrow \mathbf{fail}^{\tilde{y} \cup \tilde{z}}} & \frac{\tilde{z} = \text{fv}(C)}{\mathbf{fail}^{\tilde{y} \cup \tilde{x}}\langle\langle C/\tilde{x} \rangle\rangle \rightarrow \mathbf{fail}^{\tilde{y} \cup \tilde{z}}} \\
\text{[RS:Cons}_4] & \text{[RS: TCont]} & \\
\frac{}{\mathbf{fail}^{\tilde{y}}\langle\langle U/x \rangle\rangle \rightarrow \mathbf{fail}^{\tilde{y}}} & \frac{}{M \rightarrow N} & \\
& \frac{}{C[M] \rightarrow C[N]} &
\end{array}
\end{array}$$

Fig. 13. Reduction rules for λ_C .

Well-typedness and well-formedness. Types for λ_C extend the types for λ_C^ℓ in Section 3.2 with:

$$\sigma, \tau, \delta ::= \mathbf{unit} \mid (\pi, \eta) \rightarrow \sigma$$

$$\eta, \epsilon ::= \sigma \mid \epsilon \diamond \eta \quad \text{list} \qquad (\pi, \eta) \quad \text{tuple}$$

The list type $\epsilon \diamond \eta$ types the concatenation of unrestricted bags. It can be recursively unfolded into a finite composition of strict types $\sigma_1 \diamond \dots \diamond \sigma_n$, for some $n \geq 1$, with length n and σ_i its i -th strict type ($1 \leq i \leq n$). We write $x^! : \eta$ to denote for $x[1] : \eta_1, \dots, x[k] : \eta_k$ where η has length k . The tuple type (π, η) types concatenation of a linear bag of type π with an unrestricted bag of type η . Finally strict types are amended to allow for unrestricted functional types which go from tuple types to strict types $(\pi, \eta) \rightarrow \sigma$ rather than multiset types to strict types.

Definition 10 ($\eta \sim \epsilon$). *Let ϵ and η be two list types, with the length of ϵ greater or equal to that of η . We say that ϵ embraces η , denoted $\eta \sim \epsilon$, whenever there exist ϵ' and ϵ'' such that: i) $\epsilon = \epsilon' \diamond \epsilon''$; ii) the size of ϵ' is that of η ; iii) for all i , $\epsilon'_i = \eta_i$.*

We separate contexts into two parts: linear (Γ, Δ, \dots) and unrestricted $(\Theta, \Upsilon, \dots)$:

$$\Gamma, \Delta ::= - \mid \Gamma, x : \pi \mid \Gamma, x : \sigma \qquad \Theta, \Upsilon ::= - \mid \Theta, x^! : \eta$$

Both linear and unrestricted occurrences of variables may occur at most once in a context.

Judgments have the form $\Theta; \Gamma \vDash M : \tau$. We write $\vDash M : \tau$ to denote $-; - \vDash M : \tau$.

Definition 11 (Well-formedness in λ_C). *A λ_C -term M is well-formed if there exists a context Θ and Γ and a type τ such that the rules in Figure 14 entail $\Theta; \Gamma \vDash M : \tau$.*

A congruence. Some terms, though syntactically different, display the same behavior. For example, assuming $x \notin \text{fv}(M)$, this holds for $M\langle\langle U/x \rangle\rangle$ and M : the former describes a substitution that “does nothing” and would result in M itself. This notion is formalized through a *congruence* (\equiv_λ) closed under syntax, given in Figure 15.

$\frac{[\text{FS:var}^\ell]}{\Theta; x : \sigma \vDash x : \sigma}$	$\frac{[\text{FS:var}^1]}{\Theta, x^1 : \eta; x : \eta_i, \Delta \vDash x : \sigma}$	$\frac{[\text{FS:1}^\ell]}{\Theta; - \vDash 1 : \omega}$	$\frac{[\text{FS:bag}^\ell]}{\Theta; \Gamma \vDash M : \sigma \quad \Theta; \Delta \vDash C : \sigma^k}$	$\frac{[\text{FS:1}^1]}{\Theta; - \vDash 1^1 : \sigma}$
$\frac{[\text{FS:bag}^1]}{\Theta; - \vDash U : \epsilon \quad \Theta; - \vDash V : \eta}$	$\frac{[\text{FS:bag}]}{\Theta; \Gamma \vDash C : \sigma^k \quad \Theta; - \vDash U : \eta}$	$\frac{[\text{FS:fail}]}{\text{dom}(\Gamma^\perp) = \tilde{x}}$	$\frac{[\text{FS:weak}]}{\Theta; \Gamma \vDash M : \tau}$	
$\frac{[\text{FS:shar}]}{\Theta; \Gamma, x_1 : \sigma, \dots, x_k : \sigma \vDash M : \tau \quad x \notin \text{dom}(\Gamma) \quad k \neq 0}$		$\frac{[\text{FS:abs-sh}]}{\Theta, x^1 : \eta; \Gamma, x : \sigma^k \vDash M[\tilde{x} \leftarrow x] : \tau \quad x \notin \text{dom}(\Gamma)}$		
$\frac{[\text{FS:app}]}{\Theta; \Gamma \vDash M : (\sigma^j, \eta) \rightarrow \tau \quad \Theta; \Delta \vDash B : (\sigma^k, \epsilon) \quad \eta \sim \epsilon}$		$\frac{[\text{FS:Esub}]}{\Theta, x^1 : \eta; \Gamma, x : \sigma^j \vDash M[\tilde{x} \leftarrow x] : \tau \quad \Theta; \Delta \vDash B : (\sigma^k, \epsilon) \quad \eta \sim \epsilon}$		
$\frac{[\text{FS:Esub}^\ell]}{\Theta; \Gamma, x_1 : \sigma, \dots, x_k : \sigma \vDash M : \tau \quad \Theta; \Delta \vDash C : \sigma^k}$		$\frac{[\text{FS:Esub}^1]}{\Theta, x^1 : \eta; \Gamma \vDash M : \tau \quad \Theta; - \vDash U : \epsilon \quad \eta \sim \epsilon}$		
$\Theta; \Gamma, \Delta \vDash (M B) : \tau$		$\Theta; \Gamma, \Delta \vDash (M[\tilde{x} \leftarrow x]) \langle\langle B/x \rangle\rangle : \tau$		
$\Theta; \Gamma, \Delta \vDash M \langle C/x_1, \dots, x_k \rangle : \tau$		$\Theta; \Gamma \vDash M \llbracket U/x^1 \rrbracket : \tau$		

Fig. 14. Well-Formedness Rules for λ_c .

D.2 Translating λ_c into (Full) $s\pi^+$

Figure 7 gives the translation of λ_c^ℓ into $s\pi^+$. Here we extend this translation to consider the extended calculus λ_c . The key differences are in the translation of unrestricted variable occurrences, intermediate substitution, abstraction, and the new structure of bags. Figure 16 gives the translation that maps terms in λ_c into processes in full $s\pi^+$, denoted $\llbracket \cdot \rrbracket_u$.

The translation of an unrestricted variable $x[j]$ first connects to a server along channel x via a request $?x^1[x_i]$ followed by a selection on $\overline{x_i}.j$.

Process $\llbracket \lambda x.(M[\tilde{x} \leftarrow x]) \rrbracket_u$ first confirms its behavior along u , followed by the reception of a channel x . The channel x provides a linear channel x^ℓ and an unrestricted channel x^1 for dedicated substitutions of the linear and unrestricted bag components. This separation is also present in the translation of $\llbracket M \langle\langle B/x \rangle\rangle \rrbracket_u$, for the same reason.

Process $\llbracket M (C \star U) \rrbracket_u$ consists of synchronizations between the translation of $\llbracket M \rrbracket_v$ and $\llbracket C \star U \rrbracket_x$: the translation of $C \star U$ evolves when M is an abstraction, say $\lambda x.(M'[\tilde{x} \leftarrow x])$. The channel x^ℓ provides the linear behavior of the bag C while x^1 provides the behavior of U ; this is done by guarding the translation of U with a server connection, such that every time a channel synchronizes with it a fresh copy of U is spawned.

Process $\llbracket M \llbracket U/x^1 \rrbracket \rrbracket_u$ consists of the composition of the translation of M and a server guarding the translation of U : in order for $\llbracket M \rrbracket_u$ to gain access to $\llbracket U \rrbracket_{x_i}$ it must first synchronize with the server channel x^1 to spawn a fresh copy of the translation of U .

To complete this section, Figure 17 gives the translation of intersection types for λ_c to session types for full $s\pi^+$.

Definition 12. *The translation $\llbracket \cdot \rrbracket_u$ in Figure 17 extends to a linear context*

$$\Gamma = x_1 : \sigma_1, \dots, x_m : \sigma_m, v_1 : \pi_1, \dots, v_n : \pi_n$$

and an unrestricted context $\Theta = x^1[1] : \eta_1, \dots, x^1[k] : \eta_k$ as follows:

$$\begin{aligned} \llbracket \Gamma \rrbracket &= x_1 : \& \llbracket \sigma_1 \rrbracket, \dots, x_m : \& \llbracket \sigma_m \rrbracket, v_1 : \llbracket \pi_1 \rrbracket_{(\sigma, i_1)}, \dots, v_n : \llbracket \pi_n \rrbracket_{(\sigma, i_n)} \\ \llbracket \Theta \rrbracket &= x^1[1] : \llbracket \eta_1 \rrbracket, \dots, x^1[k] : \llbracket \eta_k \rrbracket \end{aligned}$$

E Proofs of Type Preservation and Deadlock-freedom for (Full) $s\pi^+$

Here we prove Theorems 1 and 2 (type preservation and deadlock-freedom for the lazy semantics, respectively), as well as the analogue results for the eager semantics. In fact, deadlock-freedom for the lazy semantics follows from deadlock-freedom for the eager semantics, so we present the proofs for the eager semantics first.

$$\begin{array}{c}
\frac{\tilde{x} \# \text{fv}(B)}{(M B) \langle C / \tilde{x} \rangle \equiv_{\lambda} (M \langle C / \tilde{x} \rangle) B} \quad \frac{x \notin \text{fv}(M)}{M \llbracket U / x^{\dagger} \rrbracket \equiv_{\lambda} M} \quad \frac{x \notin \text{fv}(B)}{(M B) \llbracket U / x^{\dagger} \rrbracket \equiv_{\lambda} (M \llbracket U / x^{\dagger} \rrbracket) B} \\
\\
\frac{\tilde{x} \# \text{fv}(A)}{((M A) [\tilde{x} \leftarrow x]) \langle B / x \rangle \equiv_{\lambda} ((M [\tilde{x} \leftarrow x]) \langle B / x \rangle) A} \\
\\
\frac{\tilde{x} \# \text{fv}(A) \quad \tilde{y} \# \text{fv}(B)}{\left(((M [\tilde{y} \leftarrow y]) \langle A / y \rangle) [\tilde{x} \leftarrow x] \right) \langle B / x \rangle \equiv_{\lambda} \left((M [\tilde{y} \leftarrow y]) \langle B / y \rangle \right) [\tilde{x} \leftarrow x] \right) \langle A / x \rangle} \\
\\
\frac{\tilde{x} \# \text{fv}(B) \quad \tilde{y} \# \text{fv}(C)}{((M [\tilde{y} \leftarrow y]) \langle B / y \rangle) \langle C / \tilde{x} \rangle \equiv_{\lambda} ((M \langle C / \tilde{x} \rangle) [\tilde{y} \leftarrow y]) \langle B / y \rangle} \quad \frac{x \notin \text{fv}(B) \quad \tilde{y} \# \text{fv}(U)}{((M [\tilde{y} \leftarrow y]) \langle B / y \rangle) \llbracket U / x^{\dagger} \rrbracket \equiv_{\lambda} ((M \llbracket U / x^{\dagger} \rrbracket) [\tilde{y} \leftarrow y]) \langle B / y \rangle} \\
\\
\frac{\tilde{x} \# \text{fv}(C_2) \quad \tilde{y} \# \text{fv}(C_1)}{(M \langle C_2 / \tilde{y} \rangle) \langle C_1 / \tilde{x} \rangle \equiv_{\lambda} (M \langle C_1 / \tilde{x} \rangle) \langle C_2 / \tilde{y} \rangle} \quad \frac{\tilde{x} \# \text{fv}(U) \quad y \notin \text{fv}(C)}{(M \langle C / \tilde{x} \rangle) \llbracket U / y^{\dagger} \rrbracket \equiv_{\lambda} (M \llbracket U / y^{\dagger} \rrbracket) \langle C / \tilde{x} \rangle} \\
\\
\frac{x \notin \text{fv}(U) \quad y \notin \text{fv}(V)}{(M \llbracket V / \tilde{x} \rrbracket) \llbracket U / y^{\dagger} \rrbracket \equiv_{\lambda} (M \llbracket U / y^{\dagger} \rrbracket) \llbracket V / x^{\dagger} \rrbracket}
\end{array}$$

Fig. 15. Congruence in λ_c .

E.1 Eager Semantics

Subject Congruence

Theorem 12. *If $P \vdash \Gamma$ and $P \equiv Q$, then $Q \vdash \Gamma$.*

Proof. By induction on the derivation of the structural congruence. We first detail the base cases:

- $P \equiv_{\alpha} P' \implies P \equiv P'$. Since alpha-renaming only affects bound names, it does not affect the names in Γ , so clearly $P' \vdash \Gamma$.
- $[x \leftrightarrow y] \equiv [y \leftrightarrow x]$.

$$\overline{[x \leftrightarrow y] \vdash x:A, y:\bar{A}} \equiv \overline{[y \leftrightarrow x] \vdash x:A, y:\bar{A}}$$

- $P \mid Q \equiv Q \mid P$.

$$\frac{P \vdash \Gamma \quad Q \vdash \Delta}{P \mid Q \vdash \Gamma, \Delta} \equiv \frac{Q \vdash \Delta \quad P \vdash \Gamma}{Q \mid P \vdash \Gamma, \Delta}$$

- $(P \mid Q) \mid R \equiv P \mid (Q \mid R)$.

$$\frac{\frac{P \vdash \Gamma \quad Q \vdash \Delta}{P \mid Q \vdash \Gamma, \Delta} \quad R \vdash \Lambda}{(P \mid Q) \mid R \vdash \Gamma, \Delta, \Lambda} \equiv \frac{P \vdash \Gamma \quad \frac{Q \vdash \Delta \quad R \vdash \Lambda}{Q \mid R \vdash \Delta, \Lambda}}{P \mid (Q \mid R) \vdash \Gamma, \Delta, \Lambda}$$

- $P \mid \mathbf{0} \equiv P$

$$\frac{P \vdash \Gamma \quad \overline{\mathbf{0} \vdash \emptyset}}{P \mid \mathbf{0} \vdash \Gamma} \equiv P \vdash \Gamma$$

- $(\nu x)(P \mid Q) \equiv (\nu x)(Q \mid P)$.

$$\frac{P \vdash \Gamma, x:A \quad Q \vdash \Delta, x:\bar{A}}{(\nu x)(P \mid Q) \vdash \Gamma, \Delta} \equiv \frac{Q \vdash \Delta, x:\bar{A} \quad P \vdash \Gamma, x:A}{(\nu x)(Q \mid P) \vdash \Gamma, \Delta}$$

- $x \notin \text{fn}(Q) \implies (\nu x)((\nu y)(P \mid Q) \mid R) \equiv (\nu y)((\nu x)(P \mid R) \mid Q)$.

$$\frac{\frac{P \vdash \Gamma, y:A, x:B \quad Q \vdash \Delta, y:\bar{A}}{(\nu y)(P \mid Q) \vdash \Gamma, \Delta, x:B} \quad R \vdash \Lambda, x:\bar{B}}{(\nu x)((\nu y)(P \mid Q) \mid R) \vdash \Gamma, \Delta, \Lambda} \equiv \frac{\frac{P \vdash \Gamma, y:A, x:B \quad R \vdash \Lambda, x:\bar{B}}{(\nu x)(P \mid R) \vdash \Gamma, \Lambda, y:A} \quad Q \vdash \Delta, y:\bar{A}}{(\nu y)((\nu x)(P \mid R) \mid Q) \vdash \Gamma, \Delta, \Lambda}}$$

$$\begin{aligned}
\llbracket x \rrbracket_u &= \bar{x}.\text{some}; [x \leftrightarrow u] & \llbracket x[j] \rrbracket_u &= ?\bar{x}^1[x_i]; \bar{x}_i.j; [x_i \leftrightarrow u] \\
\llbracket \lambda x.M \rrbracket_u &= \bar{u}.\text{some}; u(x); \bar{x}.\text{some}; x(x^\ell); x(x^1); x(); \llbracket M \rrbracket_u \\
\llbracket M \langle\langle C \star U/x \rangle\rangle \rrbracket_u &= (\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M \rrbracket_u \mid \llbracket C \star U \rrbracket_x \\
\llbracket M(C \star U) \rrbracket_u &= (\nu v)(\llbracket M \rrbracket_v \mid v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x]; (\llbracket C \star U \rrbracket_x \mid [v \leftrightarrow u])) \\
\llbracket C \star U \rrbracket_x &= x.\text{some}_{\text{fv}(C)}; \bar{x}[x^\ell]; (\llbracket C \rrbracket_{x^\ell} \mid \bar{x}[x^1]; (!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid \bar{x}[])) \\
\llbracket \lambda M_j \int \cdot C \rrbracket_{x^\ell} &= \frac{x^\ell.\text{some}_{\text{fv}(C)}; x(y_i); x^\ell.\text{some}_{y_i, \text{fv}(C)}; \bar{x}^\ell.\text{some};}{\bar{x}^\ell[z_i]; (z_i.\text{some}_{\text{fv}(M_j)}; \llbracket M_j \rrbracket_{z_i} \mid (\llbracket C \setminus M_j \rrbracket_{x^\ell} \mid \bar{y}_i.\text{none}))} \\
\llbracket \mathbf{1} \rrbracket_{x^\ell} &= x^\ell.\text{some}_\emptyset; x(y_n); (\bar{y}_n.\text{some}; \bar{y}_n[] \mid x^\ell.\text{some}_\emptyset; \bar{x}^\ell.\text{none}) \\
\llbracket \mathbf{1}^1 \rrbracket_x &= \bar{x}.\text{none} & \llbracket \lambda N \int^1 \rrbracket_x &= \llbracket N \rrbracket_x & \llbracket U \rrbracket_x &= x.\text{case}\{i : \llbracket U_i \rrbracket_x\}_{U_i \in U} \\
\llbracket M \langle \lambda M_1 \int \cdot \lambda M_2 \int / x_1, x_2 \rangle \rrbracket_u &= \frac{(\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid (\nu z_2)(z_2.\text{some}_{\text{fv}(M_2)}; \llbracket M_2 \rrbracket_{z_2} \\ & \quad \mid \prod_{x_{i_1} \in \{x_1, x_2\}} \prod_{x_{i_2} \in \{x_1, x_2 \setminus x_{i_1}\}} \llbracket M \rrbracket_u \{z_1/x_{i_1}\} \{z_2/x_{i_2}\}) \dots)}{ \\
\llbracket M \llbracket U/x \rrbracket \rrbracket_u &= (\nu x^1)(\llbracket M \rrbracket_u \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
\llbracket M[\leftarrow x] \rrbracket_u &= \bar{x}^\ell.\text{some}; \bar{x}^\ell[y_i]; (y_i.\text{some}_{u, \text{fv}(M)}; y_i()); \llbracket M \rrbracket_u \mid \bar{x}^\ell.\text{none} \\
\llbracket M[\tilde{x} \leftarrow x] \rrbracket_u &= \frac{\bar{x}^\ell.\text{some}; \bar{x}^\ell[y_i]; (y_i.\text{some}_\emptyset; y_i()); \mathbf{0}}{\bar{x}^\ell.\text{some}; x^\ell.\text{some}_{u, \text{fv}(M) \setminus \bar{x}}; \prod_{x_i \in \bar{x}} x^\ell(x_i); \llbracket M[(\tilde{x} \setminus x_i) \leftarrow x] \rrbracket_u} \\
\llbracket \text{fail}^{x_1, \dots, x_k} \rrbracket_u &= \bar{u}.\text{none} \mid \bar{x}_1.\text{none} \mid \dots \mid \bar{x}_k.\text{none}
\end{aligned}$$

Fig. 16. Translation of λ_c into $\mathfrak{s}\pi^+$.

$$\begin{aligned}
\llbracket \text{unit} \rrbracket &= \& \mathbf{1} & \llbracket \eta \rrbracket &= !\&_{\eta_i \in \eta} \{i : \llbracket \eta_i \rrbracket\} \\
\llbracket (\sigma^k, \eta) \rightarrow \tau \rrbracket &= \&(\overline{\llbracket (\sigma^k, \eta) \rrbracket}_{(\sigma, i)}} \wp \llbracket \tau \rrbracket & \llbracket (\sigma^k, \eta) \rrbracket_{(\sigma, i)} &= \oplus(\overline{\llbracket \sigma^k \rrbracket}_{(\sigma, i)}} \otimes (\overline{\llbracket \eta \rrbracket} \otimes (\mathbf{1}))) \\
\llbracket \sigma \wedge \pi \rrbracket_{(\sigma, i)} &= \oplus(\overline{\llbracket \& \mathbf{1} \rrbracket} \wp (\oplus \&(\overline{\llbracket \sigma \rrbracket} \otimes (\overline{\llbracket \pi \rrbracket}_{(\sigma, i)})))) \\
\llbracket \omega \rrbracket_{(\sigma, i)} &= \begin{cases} \oplus(\overline{\llbracket \& \mathbf{1} \rrbracket} \wp (\oplus \& \mathbf{1})) & \text{if } i = 0 \\ \oplus(\overline{\llbracket \& \mathbf{1} \rrbracket} \wp (\oplus \&(\overline{\llbracket \sigma \rrbracket} \otimes (\overline{\llbracket \omega \rrbracket}_{(\sigma, i-1)})))) & \text{if } i > 0 \end{cases}
\end{aligned}$$

Fig. 17. Translation of intersection types into session types (cf. Def. 12).

$$- x \notin \text{fn}(Q) \implies (\nu x)((P \mid Q) \mid R) \equiv (\nu x)(P \mid R) \mid Q.$$

$$\frac{\frac{P \vdash \Gamma, x:A \quad Q \vdash \Delta}{P \mid Q \vdash \Gamma, \Delta, x:A} \quad R \vdash \Lambda, x:\bar{A}}{(\nu x)((P \mid Q) \mid R) \vdash \Gamma, \Delta, \Lambda} \equiv \frac{\frac{P \vdash \Gamma, x:A \quad R \vdash \Lambda, x:\bar{A}}{(\nu x)(P \mid R) \vdash \Gamma, \Lambda} \quad Q \vdash \Delta}{(\nu x)(P \mid R) \mid Q \vdash \Gamma, \Delta, \Lambda}$$

$$- x \notin \text{fn}(Q) \implies (\nu x)(!x(y); P \mid Q) \equiv Q.$$

$$\frac{\frac{P \vdash ?\Gamma, y:A}{!x(y); P \vdash ?\Gamma, x:\bar{A}} \quad \frac{Q \vdash \Delta}{Q \vdash \Delta, x:\bar{A}}}{(\nu x)(!x(y); P \mid Q) \vdash ?\Gamma, \Delta} \equiv \frac{Q \vdash \Delta}{Q \vdash ?\Gamma, \Delta}$$

$$- P \ddagger Q \equiv Q \ddagger P.$$

$$\frac{P \vdash \Gamma \quad Q \vdash \Gamma}{P \ddagger Q \vdash \Gamma} \equiv \frac{Q \vdash \Gamma \quad P \vdash \Gamma}{Q \ddagger P \vdash \Gamma}$$

$$- (P \ddagger Q) \ddagger R \equiv P \ddagger (Q \ddagger R)$$

$$\frac{\frac{P \vdash \Gamma \quad Q \vdash \Gamma}{P \ddagger Q \vdash \Gamma} \quad R \vdash \Gamma}{(P \ddagger Q) \ddagger R \vdash \Gamma} \equiv \frac{P \vdash \Gamma \quad \frac{Q \vdash \Gamma \quad R \vdash \Gamma}{Q \ddagger R \vdash \Gamma}}{P \ddagger (Q \ddagger R) \vdash \Gamma}$$

– $P \parallel P \equiv P$.

$$\frac{P \vdash \Gamma \quad P \vdash \Gamma}{P \parallel P \vdash \Gamma} \equiv P \vdash \Gamma$$

The inductive cases follow from the IH straightforwardly. Note that the rules for parallel composition do not apply directly behind the output prefix and restriction. \square

Subject Reduction

Lemma 2. *Suppose $P \vdash \Gamma, x:A$.*

1. If $P = \mathbb{N}[\bar{x}[]]$, then $A = \mathbf{1}$.
2. If $P = \mathbb{N}[x(); P']$, then $A = \perp$.
3. If $P = \mathbb{N}[\bar{x}[y]; (P' \mid P'')]$, then $A = B \otimes C$.
4. If $P = \mathbb{N}[x(y); P']$, then $A = B \wp C$.
5. If $P = \mathbb{N}[\bar{x}.j; P']$, then $A = \oplus\{i : B_i\}_{i \in I}$ where $j \in I$.
6. If $P = \mathbb{N}[x.\text{case}\{i : P'_i\}_{i \in I}]$, then $A = \&\{i : B_i\}_{i \in I}$.
7. If $P = \mathbb{N}[\bar{x}.\text{some}; P']$, then $A = \&B$.
8. If $P = \mathbb{N}[\bar{x}.\text{none}]$, then $A = \&B$.
9. If $P = \mathbb{N}[x.\text{some}_{w_1, \dots, w_n}]$, then $A = \oplus B$.
10. If $P = \mathbb{N}[?x[y]; P']$, then $A = ?B$.
11. If $P = \mathbb{N}![x(y); P']$, then $A = !B$.

Proof. Each item follows by induction on the structure of the ND-context. The base case follows by inversion of typing, and the inductive cases follow from the IH straightforwardly. \square

Lemma 3. *For each of the following items, assume $\Gamma \# \Delta$.*

1. If $\mathbb{N}[x \leftrightarrow y] \vdash \Gamma, x:A$ and $Q \vdash \Delta, x:\bar{A}$, then $\langle \mathbb{N} \rangle [Q\{y/x\}] \vdash \Gamma, \Delta$.
2. If $\mathbb{N}[\bar{x}[]] \vdash \Gamma, x:\mathbf{1}$, then $\langle \mathbb{N} \rangle [\mathbf{0}] \vdash \Gamma$.
3. If $\mathbb{N}[x(); Q] \vdash \Gamma, x:\perp$, then $\langle \mathbb{N} \rangle [Q] \vdash \Gamma$.
4. If $\text{bn}(\mathbb{N}) \# \text{fn}(\mathbb{N}')$ and $\mathbb{N}[\bar{x}[y]; (P|Q)] \vdash \Gamma, x:A \otimes B$ and $\mathbb{N}'[x(z); R] \vdash \Delta, x:\bar{A} \wp \bar{B}$, then $\langle \mathbb{N} \rangle [(\nu x)(Q | (\nu y)(P | \langle \mathbb{N}' \rangle [R\{y/z\}]))] \vdash \Gamma, \Delta$.
5. If $\mathbb{N}[\bar{x}.j; P] \vdash \Gamma, x:\oplus\{i : A_i\}_{i \in I}$ and $j \in I$, then $\langle \mathbb{N} \rangle [P] \vdash \Gamma, x:A_j$.
6. If $\mathbb{N}[x.\text{case}\{i : P_i\}_{i \in I}] \vdash \Gamma, x:\&\{i : A_i\}_{i \in I}$, then $\langle \mathbb{N} \rangle [P_i] \vdash \Gamma, x:A_i$ for every $i \in I$.
7. If $\mathbb{N}[\bar{x}.\text{some}; P] \vdash \Gamma, x:\&A$, then $\langle \mathbb{N} \rangle [P] \vdash \Gamma, x:A$.
8. If $\mathbb{N}[\bar{x}.\text{none}] \vdash \Gamma, x:\&A$, then $\langle \mathbb{N} \rangle [\mathbf{0}] \vdash \Gamma$.
9. If $\mathbb{N}[x.\text{some}_{w_1, \dots, w_n}; P] \vdash \Gamma, x:\oplus A$, then $\langle \mathbb{N} \rangle [P] \vdash \Gamma, x:A$ and $\langle \mathbb{N} \rangle [\bar{w}_1.\text{none} \mid \dots \mid \bar{w}_n.\text{none}] \vdash \Gamma$.
10. If $\text{bn}(\mathbb{N}') \# \text{fn}(\mathbb{N})$ and $\mathbb{N}[?x[y]; P] \vdash \Gamma, x:?A$ and $\mathbb{N}'![x(y); Q] \vdash \Delta, x:!\bar{A}$, then $\langle \mathbb{N}' \rangle [(\nu x)((\nu y)(\langle \mathbb{N} \rangle [P] | Q\{y/z\}) | !x(z); Q)]$.

Proof. For each item, we apply induction on the structure of the ND-contexts and detail the base cases. For simplicity, we assume no names in Γ and Δ were derived with [TWEAKEN].

1.

$$\frac{}{[x \leftrightarrow y] \vdash x:A, y:\bar{A}} \quad Q \vdash \Delta, x:\bar{A} \quad \Longrightarrow \quad Q\{y/x\} \vdash \Delta, y:\bar{A}$$

2.

$$\frac{}{[\bar{x}[]] \vdash x:\mathbf{1}} \quad \Longrightarrow \quad \frac{}{\mathbf{0} \vdash \emptyset}$$

3.

$$\frac{Q \vdash \Gamma}{x(); Q \vdash \Gamma, x:\perp} \quad \Longrightarrow \quad Q \vdash \Gamma$$

4.

$$\frac{P \vdash \Gamma, y:A \quad Q \vdash \Gamma', x:B}{[\bar{x}[y]; (P|Q)] \vdash \Gamma, \Gamma', x:A \otimes B} \quad \frac{R \vdash \Delta, z:\bar{A}, x:\bar{B}}{x(z); R \vdash \Delta, x:\bar{A} \wp \bar{B}} \quad \Longrightarrow$$

$$\frac{Q \vdash \Gamma', x:B \quad \frac{P \vdash \Gamma, y:A \quad R\{y/z\} \vdash \Delta, y:\bar{A}, x:\bar{B}}{(\nu y)(P | R\{y/z\}) \vdash \Gamma, \Delta, x:\bar{B}}}{(\nu x)(Q | (\nu y)(P | R\{y/z\})) \vdash \Gamma, \Gamma', \Delta}$$

5.

$$\frac{P \vdash \Gamma, x:A_j \quad j \in I}{\bar{x}.j; P \vdash \Gamma, x:\oplus\{i : A_i\}_{i \in I}} \Longrightarrow P \vdash \Gamma, x:A_j$$

6.

$$\frac{\forall i \in I. P_i \vdash \Gamma, x:A_i}{x.\text{case}\{i : P_i\}_{i \in I} \vdash \Gamma, x:\&\{i : A_i\}} \Longrightarrow \forall i \in I. P_i \vdash \Gamma, x:A_i$$

7.

$$\frac{P \vdash \Gamma, x:A}{\bar{x}.\text{some}; P \vdash \Gamma, x:\&A} \Longrightarrow P \vdash \Gamma, x:A$$

8.

$$\frac{}{\bar{x}.\text{none} \vdash x:\&A} \Longrightarrow \frac{}{\mathbf{0} \vdash \emptyset}$$

9.

$$\frac{P \vdash w_1:\&B_1, \dots, w_n:\&B_n, x:A}{x.\text{some}_{w_1, \dots, w_n}; P \vdash w_1:\&B_1, \dots, w_n:\&B_n, x:\oplus A} \Longrightarrow P \vdash w_1:\&B_1, \dots, w_n:\&B_n, x:A$$

$$\frac{\frac{}{\bar{w}_1.\text{none} \vdash w_1:\&B_1} \quad \dots \quad \frac{}{\bar{w}_n.\text{none} \vdash w_n:\&B_n}}{\bar{w}_1.\text{none} \mid \dots \mid \bar{w}_n.\text{none} \vdash w_1:\&B_1, \dots, w_n:\&B_n}}$$

10. This item depends on whether $x \in \text{fn}(P)$.– $x \in \text{fn}(P)$:

$$\frac{\frac{P\{x'/x\} \vdash \Gamma, y:A, x':?A}{\frac{?x[y]; (P\{x'/x\}) \vdash \Gamma, x:?A, x':?A}{?x[y]; P \vdash \Gamma, x:?A}}{\frac{Q \vdash ?\Delta, z:\bar{A}}{!x(z); Q \vdash ?\Delta, x:!\bar{A}}} \Longrightarrow$$

$$\frac{P \vdash \Gamma, y:A, x:?A \quad Q\{y/z\}\{w'/w\}_{w \in ?\Delta} \vdash ?\Delta', y:\bar{A}}{(\nu y)(P \mid Q\{y/z\}\{w'/w\}_{w \in ?\Delta}) \vdash \Gamma, ?\Delta', x:?A \quad !x(z); Q \vdash ?\Delta, x:!\bar{A}}$$

$$\frac{(\nu x)((\nu y)(P \mid Q\{y/z\}\{w'/w\}_{w \in ?\Delta}) \mid !x(z); Q) \vdash \Gamma, ?\Delta, ?\Delta'}{(\nu x)((\nu y)(P \mid Q\{y/z\}) \mid !x(z); Q) \vdash \Gamma, ?\Delta}$$

– $x \notin \text{fn}(P)$:

$$\frac{P \vdash \Gamma, y:A}{?x[y]; P \vdash \Gamma, x:?A} \quad \frac{Q \vdash ?\Delta, z:\bar{A}}{!x(z); Q \vdash ?\Delta, x:!\bar{A}} \Longrightarrow$$

$$\frac{P \vdash \Gamma, y:A \quad Q\{y/z\}\{w'/w\}_{w \in ?\Delta} \vdash ?\Delta', y:\bar{A}}{(\nu y)(P \mid Q\{y/z\}\{w'/w\}_{w \in ?\Delta}) \vdash \Gamma, ?\Delta'}$$

$$\frac{(\nu y)(P \mid Q\{y/z\}\{w'/w\}_{w \in ?\Delta}) \vdash \Gamma, ?\Delta', x:?A \quad !x(z); Q \vdash ?\Delta, x:!\bar{A}}{(\nu x)((\nu y)(P \mid Q\{y/z\}\{w'/w\}_{w \in ?\Delta}) \mid !x(z); Q) \vdash \Gamma, ?\Delta, ?\Delta'}$$

$$\frac{(\nu x)((\nu y)(P \mid Q\{y/z\}) \mid !x(z); Q) \vdash \Gamma, ?\Delta, ?\Delta'}{(\nu x)((\nu y)(P \mid Q\{y/z\}) \mid !x(z); Q) \vdash \Gamma, ?\Delta}$$

The inductive cases follow straightforwardly. Notice that the conditions on the bound and free names of the ND-contexts in items 4 and 10 make sure that no names are captured when embedding one context in the other. \square

Theorem 13 (SR for the Eager Semantics). *If $P \vdash \Gamma$ and $P \longrightarrow Q$, then $Q \vdash \Gamma$.*

Proof. By induction on the derivation of the reduction.

– Rule $[\rightarrow_{\text{ID}}]$.

$$\frac{\mathbb{N}[x \leftrightarrow y] \vdash \Gamma, x:A \quad Q \vdash \Delta, x:\bar{A}}{(\nu x)(\mathbb{N}[x \leftrightarrow y] \mid Q) \vdash \Gamma, \Delta} \quad \Longrightarrow \quad \langle \mathbb{N} \rangle [Q\{y/x\}] \vdash \Gamma, \Delta \quad (\text{Lemma 3.1})$$

– Rule $[\rightarrow_{\mathbf{1}\perp}]$.

$$\frac{\mathbb{N}[\bar{x}] \vdash \Gamma, x:\mathbf{1} \quad (\text{Lemma 2.1}) \quad \mathbb{N}'[x(); Q] \vdash \Delta, x:\perp \quad (\text{Lemma 2.2})}{\nu x(\mathbb{N}[\bar{x}] \mid \mathbb{N}'[x(); Q]) \vdash \Gamma, \Delta} \quad \Longrightarrow$$

$$\frac{\langle \mathbb{N} \rangle [\mathbf{0}] \vdash \Gamma \quad (\text{Lemma 3.2}) \quad \langle \mathbb{N}' \rangle [Q] \vdash \Delta \quad (\text{Lemma 3.3})}{\langle \mathbb{N} \rangle [\mathbf{0}] \mid \langle \mathbb{N}' \rangle [Q] \vdash \Gamma, \Delta}$$

– Rule $[\rightarrow_{\otimes \wp}]$.

$$\frac{\mathbb{N}[\bar{x}[y]; (P \mid Q)] \vdash \Gamma, x:A \otimes B \quad (\text{Lemma 2.3}) \quad \mathbb{N}'[x(z); R] \vdash \Delta, x:\bar{A} \wp \bar{B} \quad (\text{Lemma 2.4})}{(\nu x)(\mathbb{N}[\bar{x}[y]; (P \mid Q)] \mid \mathbb{N}'[x(z); R]) \vdash \Gamma, \Delta} \quad \Longrightarrow$$

$$\langle \mathbb{N} \rangle [(\nu x)(Q \mid \nu y(P \mid \langle \mathbb{N}' \rangle [R\{y/z\}])] \vdash \Gamma, \Delta \quad (\text{Lemma 3.4})$$

– Rule $[\rightarrow_{\oplus \&}]$.

$$\frac{\mathbb{N}[\bar{x}.j; P] \vdash \Gamma, x:\oplus\{i : A_i\}_{i \in I} \quad j \in I \quad (\text{Lemma 2.5}) \quad \mathbb{N}'[x.\text{case}\{i : Q_i\}_{i \in I}] \vdash \Delta, x:\&\{\bar{A}_i\}_{i \in I} \quad (\text{Lemma 2.6})}{(\nu x)(\mathbb{N}[\bar{x}.j; P] \mid \mathbb{N}'[x.\text{case}\{i : Q_i\}_{i \in I}]) \vdash \Gamma, \Delta} \quad \Longrightarrow$$

$$\frac{\langle \mathbb{N} \rangle [P] \vdash \Gamma, x:A_j \quad (\text{Lemma 3.5}) \quad \langle \mathbb{N}' \rangle [Q_j] \vdash \Delta, x:\bar{A}_j \quad (\text{Lemma 3.6})}{(\nu x)(\langle \mathbb{N} \rangle [P] \mid \langle \mathbb{N}' \rangle [Q_j]) \vdash \Gamma, \Delta}$$

– Rule $[\rightarrow_{\text{some}}]$.

$$\frac{\mathbb{N}[\bar{x}.\text{some}; P] \vdash \Gamma, x:\&A \quad (\text{Lemma 2.7}) \quad \mathbb{N}'[x()(w_1, \dots, w_n); Q] \vdash \Delta, x:\oplus\bar{A} \quad (\text{Lemma 2.9})}{(\nu x)(\mathbb{N}[\bar{x}.\text{some}; P] \mid \mathbb{N}'[x()(w_1, \dots, w_n); Q]) \vdash \Gamma, \Delta} \quad \Longrightarrow$$

$$\frac{\langle \mathbb{N} \rangle [P] \vdash \Gamma, x:A \quad (\text{Lemma 3.7}) \quad \langle \mathbb{N}' \rangle [Q] \vdash \Delta, x:\bar{A} \quad (\text{Lemma 3.9})}{(\nu x)(\langle \mathbb{N} \rangle [P] \mid \langle \mathbb{N}' \rangle [Q]) \vdash \Gamma, \Delta}$$

– Rule $[\rightarrow_{\text{none}}]$.

$$\frac{\mathbb{N}[\bar{x}.\text{none}] \vdash \Gamma, x:\&A \quad (\text{Lemma 2.8}) \quad \mathbb{N}'[x()(w_1, \dots, w_n); Q] \vdash \Delta, x:\oplus\bar{A} \quad (\text{Lemma 2.9})}{(\nu x)(\mathbb{N}[\bar{x}.\text{none}] \mid \mathbb{N}'[x()(w_1, \dots, w_n); Q]) \vdash \Gamma, \Delta} \quad \Longrightarrow$$

$$\frac{\langle \mathbb{N} \rangle [\mathbf{0}] \vdash \Gamma \quad (\text{Lemma 3.8}) \quad \langle \mathbb{N}' \rangle [\bar{w}_1.\text{none} \mid \dots \mid \bar{w}_n.\text{none}] \vdash \Delta \quad (\text{Lemma 3.9})}{\langle \mathbb{N} \rangle [\mathbf{0}] \mid \langle \mathbb{N}' \rangle [\bar{w}_1.\text{none} \mid \dots \mid \bar{w}_n.\text{none}] \vdash \Gamma, \Delta}$$

– Rule $[\rightarrow_{?!}]$.

$$\frac{\mathbb{N}[?\bar{x}[y]; P] \vdash \Gamma, x:?A \quad (\text{Lemma 2.10}) \quad \mathbb{N}'[!x(z); Q] \vdash \Delta, x:!\bar{A} \quad (\text{Lemma 2.11})}{(\nu x)(\mathbb{N}[?\bar{x}[y]; P] \mid \mathbb{N}'[!x(z); Q]) \vdash \Gamma, \Delta} \quad \Longrightarrow$$

$$\langle \mathbb{N}' \rangle [(\nu x)((\nu y)(\langle \mathbb{N} \rangle [P] \mid Q\{y/z\}) \mid !x(z); Q)] \vdash \Gamma, \Delta \quad (\text{Lemma 3.10})$$

– Rule $[\rightarrow_{\equiv}]$. Assume $P \equiv P'$ and $P' \longrightarrow Q'$ and $Q' \equiv Q$. By Theorem 12, $P' \vdash \Gamma$. By the IH, $Q' \vdash \Gamma$. By Theorem 12, $Q \vdash \Gamma$.

– Rule $[\rightarrow_{\nu}]$. Assume $P \longrightarrow P'$.

$$\frac{P \vdash \Gamma, x:A \quad Q \vdash \Delta, x:\bar{A}}{(\nu x)(P \mid Q) \vdash \Gamma, \Delta} \quad \Longrightarrow \quad \frac{P' \vdash \Gamma, x:A \quad (\text{IH}) \quad Q \vdash \Delta, x:\bar{A}}{(\nu x)(P' \mid Q) \vdash \Gamma, \Delta}$$

– Rule $[\rightarrow_{\mid}]$. Assume $P \longrightarrow P'$.

$$\frac{P \vdash \Gamma \quad Q \vdash \Delta}{P \mid Q \vdash \Gamma, \Delta} \quad \Longrightarrow \quad \frac{P' \vdash \Gamma \quad (\text{IH}) \quad Q \vdash \Delta}{P' \mid Q \vdash \Gamma, \Delta}$$

- Rule $[\rightarrow \#]$. Assume $P \rightarrow P'$.

$$\frac{P \vdash \Gamma \quad Q \vdash \Gamma}{P \# Q \vdash \Gamma} \quad \Longrightarrow \quad \frac{P' \vdash \Gamma \text{ (IH)} \quad Q \vdash \Gamma}{P' \# Q \vdash \Gamma}$$

□

Deadlock-freedom The proof uses several definitions and lemmas, which we summarize:

- Definition 13 defines single-choice multi-hole contexts, where holes may only appear on one side of non-deterministic choices. Definition 14 yields deterministic multi-hole contexts from single-choice multi-hole contexts by committing non-deterministic choices to the sides of holes. Lemma 4 ensures typing remains consistent when committing a single-choice multi-hole context.
- Lemma 5 states that any typable process not equivalent to $\mathbf{0}$ can be written as an S-context with each hole replaced by a prefixed process. Let us refer to this as the *S-context form*.
- Lemma 6 states that if a process in S-context form is typable under empty context and has a forwarder as one of its prefixes, that process contains a cut on one of the forwarder's subjects.
- Lemma 7 states that the number of prefixed processes of a process in S-context form is at least the number of cuts in the S-context. This lemma is key to the proof of Deadlock Freedom, as it is necessary to show the next lemma.
- Lemma 8 states that if a process in S-context form is typable under empty context, then there must be two of its prefixed processes that share a subject.

Definition 13 (Single-choice Multi-hole Contexts). We define single-choice multi-hole contexts (*S-contexts*, for short) as follows:

$$\mathbf{S} ::= [\cdot]_i \mid (\nu x)(\mathbf{S} \mid \mathbf{S}) \mid \mathbf{S} \mid \mathbf{S} \mid \mathbf{S} \# P$$

An *S-context* is *n-ary* if it has *n* holes $[\cdot]_1, \dots, [\cdot]_n$. We write $\mathbf{S}[P_1, \dots, P_n]$ to denote the process obtained from the *n-ary* multi-hole context \mathbf{S} by replacing each *i*-th hole in \mathbf{S} with P_i . Given an *S-context* \mathbf{S} with hole indices I and a sequence of processes $(P_i)_{i \in I}$, we write $\mathbf{S}[P_i]_{i \in I}$ to denote the process obtained from \mathbf{S} by replacing each hole with index *i* in \mathbf{S} with P_i . We say an *S-context* is a *deterministic multi-hole context* (*DM-context*, for short) if its holes do not appear inside any non-deterministic choices.

Definition 14 (Commitment of Single-choice Multi-hole Contexts). We define the commitment of *S-context* \mathbf{S} , by abuse of notation denoted $\langle \mathbf{S} \rangle$ (cf. Section 5), as follows, yielding a deterministic multi-hole context:

$$\langle [\cdot]_i \rangle := [\cdot]_i \quad \langle \mathbf{S} \mid \mathbf{S}' \rangle := \langle \mathbf{S} \rangle \mid \langle \mathbf{S}' \rangle \quad \langle (\nu x)(\mathbf{S} \mid \mathbf{S}') \rangle := (\nu x)(\langle \mathbf{S} \rangle \mid \langle \mathbf{S}' \rangle) \quad \langle \mathbf{S} \# P \rangle := \langle \mathbf{S} \rangle$$

Lemma 4. If $\mathbf{S}[P_i]_{i \in I} \vdash \Gamma$, then $\langle \mathbf{S} \rangle [P_i]_{i \in I} \vdash \Gamma$.

Proof. Straightforward, by induction on the structure of \mathbf{S} . □

Lemma 5. If $P \vdash \Gamma$ and $P \not\equiv \mathbf{0}$, then there exist *S-context* \mathbf{S} with indices I and sequence of prefixed processes $(\alpha_i; P_i)_{i \in I}$ such that $P \equiv \mathbf{S}[\alpha_i; P_i]_{i \in I}$.

Proof. Using structural congruence, we first remove all cuts with unused servers and parallel compositions with $\mathbf{0}$, obtaining $P' \equiv P$. Since $P \not\equiv \mathbf{0}$, $P' \not\equiv \mathbf{0}$. Then, we construct \mathbf{S} by induction on the typing derivation of P' . Rules [EMPTY] and [TWEAKEN] do not occur, because of how we obtained P' from P . The structural rules [TMIX], [TCUT], and [TWEAKEN] are simply copied. In case of rule [T#], we arbitrarily pick a branch to continue the construction of \mathbf{S} with, while copying the entire other branch. The other rules, which type prefixes, add a hole to \mathbf{S} ; we mark the hole with index *i* and refer to the prefixed process typed by the rule as $\alpha_i; P_i$. Clearly, $P \equiv P' = \mathbf{S}[\alpha_i; P_i]_{i \in I}$. □

Lemma 6. If $P = \mathbf{S}[\alpha_i; P_i]_{i \in I} \vdash \emptyset$ and there is $j \in I$ s.t. $\alpha_j = [x \leftrightarrow y]$, then there are $\mathbf{N}, \mathbf{N}', Q$ such that $P = \mathbf{N} \left[(\nu x)(\mathbf{N}' \left[[x \leftrightarrow y] \right] \mid Q) \right]$.

Proof. Note that there must be a restriction on x in P , because x appears free in $[x \leftrightarrow y]$ but $P \vdash \emptyset$. First, we obtain \mathbf{N} from P by replacing the restriction on x in P with a hole, referring the parallel component in which $[x \leftrightarrow y]$ appears as P' and the other parallel component as Q . Then, we obtain \mathbf{N}' from P' by replacing $[x \leftrightarrow y]$ with a hole. Clearly, $P = \mathbf{N} \left[(\nu x)(\mathbf{N}' \left[[x \leftrightarrow y] \right] \mid Q) \right]$. □

Lemma 7. If the derivation of $P = \mathbf{S}[\alpha_i; P_i]_{i \in I} \vdash \Gamma$ and \mathbf{S} is deterministic and contains *n* cuts, then $|I| \geq n + 1$.

Proof. We apply strong induction on the number n of cuts in \mathbf{S} :

- Case $n = 0$. Any \mathbf{S} -context must have at least one hole, so \mathbf{S} has at least one hole. Hence, $|I| \geq 1 = n + 1$.
- Case $n = n' + 1$. By abuse of notation, $P = P_1 \mid \dots \mid P_k$, where for each $1 \leq k' \leq k$, $P_{k'}$ is not a parallel composition. By assumption, $m \geq 1$ of the P_1, \dots, P_k are cuts. W.l.o.g., assume P_1, \dots, P_m are cuts. For each $1 \leq j \leq m$, by inversion of rule [TMIX], $P_j \vdash \Gamma_j$, and by construction, there are \mathbf{S}_j, I_j s.t. $P_j = \mathbf{S}_j[\alpha_i; P_i]_{i \in I_j}$ where \mathbf{S}_j is deterministic. We have for each $1 \leq j \leq m$ and $1 \leq j' \leq m$ where $j \neq j'$ that $I_j \# I_{j'}$, and $\bigcup_{1 \leq j \leq m} I_j \subseteq I$. Then, for each $1 \leq j \leq m$, let $1 \leq n_j \leq n$ be the number of cuts in \mathbf{S}_j . Since P_{m+1}, \dots, P_k are not cuts, we have $\sum_{1 \leq j \leq m} n_j = n$. Take any $1 \leq j \leq m$. We have $P_j = (\nu x)(P'_j \mid P''_j)$, and by inversion of rule [TCUT], $P'_j \vdash \Gamma'_j, x:A$ and $P''_j \vdash \Gamma''_j, x:\bar{A}$ where $\Gamma_j = \Gamma'_j, \Gamma''_j$. By construction, there are $\mathbf{S}'_j, \mathbf{S}''_j, I'_j, I''_j$ s.t. $P'_j = \mathbf{S}'_j[\alpha_i; P_i]_{i \in I'_j}$ and $P''_j = \mathbf{S}''_j[\alpha_i; P_i]_{i \in I''_j}$ and \mathbf{S}'_j and \mathbf{S}''_j are deterministic. We have $I'_j \# I''_j$ and $I'_j \cup I''_j = I_j$. Let n'_j and n''_j be the number of cuts in \mathbf{S}'_j and \mathbf{S}''_j , respectively. We have that $n'_j + n''_j + 1 = n_j$. Since $n_j \leq n = n' + 1$, then $n'_j, n''_j \leq n'$. Then, by the IH, $|I'_j| \geq n'_j + 1$ and $|I''_j| \geq n''_j + 1$. Therefore, $|I_j| = |I'_j \cup I''_j| = |I'_j| + |I''_j| \geq n'_j + n''_j + 1 + 1 = n_j + 1$. In conclusion,

$$|I| \geq |\bigcup_{1 \leq j \leq m} I_j| = \sum_{1 \leq j \leq m} |I_j| \geq \sum_{1 \leq j \leq m} (n_j + 1) = \sum_{1 \leq j \leq m} n_j + m = n + m \geq n + 1. \quad \square$$

Lemma 8. *If $P = \mathbf{S}[\alpha_i; P_i]_{i \in I} \vdash \emptyset$ where for each $i \in I$, $\alpha_i \neq [x \leftrightarrow y]$ for any x and y , then there are $j, k \in I$ where $j \neq k$ and $x = \text{subj}(\alpha_j) = \text{subj}(\alpha_k)$, and there are $\mathbf{N}, \mathbf{N}_j, \mathbf{N}_k$ such that $P = \mathbf{N}[(\nu x)(\mathbf{N}_j[\alpha_j] \mid \mathbf{N}_k[\alpha_k])]$.*

Proof. Let $Q = (\mathbf{S})[(\alpha_i)_{i \in I}]$. Then Q is deterministic and, by Lemma 4, $Q \vdash \emptyset$. Let n be the number of cuts in \mathbf{S} . By Lemma 7, $|I| \geq n + 1$.

Suppose, for contradiction, that for every $j, k \in I$ where $j \neq k$, we have $\text{subj}(\alpha_j) \neq \text{subj}(\alpha_k)$. Since $Q \vdash \emptyset$, for each $j \in I$, $\text{subj}(\alpha_j)$ must be bound by a cut, so \mathbf{S} must contain $|I|$ cuts. This means $|I| = n$, contradicting the fact that $|I| \geq n + 1$. Therefore, there must be $j, k \in I$ where $j \neq k$ such that $\text{subj}(\alpha_j) = \text{subj}(\alpha_k)$.

Hence, we can take $x = \text{subj}(\alpha_j) = \text{subj}(\alpha_k)$. Since $P \vdash \emptyset$ but x appears free in $\alpha_j; P_j$ and $\alpha_k; P_k$, there must be a restriction on x in \mathbf{S} containing the holes $[\cdot]_j$ and $[\cdot]_k$. We now obtain \mathbf{N} from P by replacing the restriction on x in \mathbf{S} with a hole, referring to the parallel component in which $\alpha_j; P_j$ appears as P_j and the component in which $\alpha_k; P_k$ appears as P_k . Then, we obtain \mathbf{N}_j and \mathbf{N}_k from P_j and P_k , respectively, by replacing $\alpha_j; P_j$ and $\alpha_k; P_k$ with a hole. Clearly, $P = \mathbf{N}[(\nu x)(\mathbf{N}_j[\alpha_j; P_j] \mid \mathbf{N}_k[\alpha_k; P_k])]$. \square

Theorem 8 (Deadlock-freedom: Eager Semantics). *If $P \vdash \emptyset$ and $P \neq \mathbf{0}$, then there is R such that $P \longrightarrow R$.*

Proof. By Lemma 5, there are \mathbf{S} -context \mathbf{S} with hole indices I and sequence of prefixed processes $(\alpha_i; P_i)_{i \in I}$ such that $P \equiv \mathbf{S}[\alpha_i; P_i]_{i \in I}$. The next step depends on whether there is a forwarder process among the α_i .

- If there exists $j \in I$ s.t. $\alpha_j = [x \leftrightarrow y]$ for some x and y , then by Lemma 6 there are $\mathbf{N}, \mathbf{N}', Q$ s.t. $\mathbf{S}[\alpha_i; P_i] = \mathbf{N}[(\nu x)(\mathbf{N}'[[x \leftrightarrow y]] \mid Q)]$.

$$\begin{aligned} (\nu x)(\mathbf{N}'[[x \leftrightarrow y]] \mid Q) &\longrightarrow (\mathbf{N}')[Q\{y/x\}] = R' && \text{(by rule } [\rightarrow_{\text{Id}}]) \\ \mathbf{S}[\alpha_i; P_i]_{i \in I} = \mathbf{N}[(\nu x)(\mathbf{N}'[[x \leftrightarrow y]] \mid Q)] &\longrightarrow \mathbf{N}[R'] = R && \text{(by rules } [\rightarrow_{\nu}], [\rightarrow_{\cdot}], [\rightarrow_{\perp}]) \\ P &\longrightarrow R && \text{(by rule } [\rightarrow_{\equiv}]) \end{aligned}$$

- If for each $i \in I$, $\alpha_i \neq [x \leftrightarrow y]$ for any x and y , then by Lemma 8 there are $j, k \in I$ where $j \neq k$ and $x = \text{subj}(\alpha_j) = \text{subj}(\alpha_k)$ for some x , and $\mathbf{N}, \mathbf{N}_j, \mathbf{N}_k$ such that $\mathbf{S}[\alpha_i; P_i]_{i \in I} = \mathbf{N}[(\nu x)(\mathbf{N}_j[\alpha_j; P_j] \mid \mathbf{N}_k[\alpha_k; P_k])]$.

We now show by cases on α_j that there is R' such that $(\nu x)(\mathbf{N}_j[\alpha_j; P_j] \mid \mathbf{N}_k[\alpha_k; P_k]) \longrightarrow R'$. First, note that by typability, if the type for x in $\mathbf{N}_j[\alpha_j; P_j]$ is A , then the type for x in $\mathbf{N}_k[\alpha_k; P_k]$ is \bar{A} . In the following cases, we determine more precisely the form of A by typing inversion on $\mathbf{N}_j[\alpha_j; P_j]$, and then determine the form of α_k by typing inversion using the form of \bar{A} . Note that we can exclude any cases where α_j or α_k are forwarder processes, as we assume they are not.

- If $\alpha_j; P_j = \bar{x}[]$, then $A = \mathbf{1}$ and $\bar{A} = \perp$. Hence, $\alpha_k; P_k = x(); P_k$. By rule $[\rightarrow_{\mathbf{1}\perp}]$, there is R' such that

$$(\nu x)(\mathbf{N}_j[\bar{x}[]] \mid \mathbf{N}_k[x(); P_k]) \longrightarrow R'.$$

- If $\alpha_j; P_j = \bar{x}[y]; (P'_j \mid P''_j)$ for some y , then $A = B \otimes C$ and $\bar{A} = \bar{B} \wp \bar{C}$ for some B and C . Hence, $\alpha_k; P_k = x(z); P_k$ for some z . By rule $[\rightarrow_{\otimes \wp}]$, there is R' such that

$$(\nu x)(\mathbf{N}_j[\bar{x}[y]; (P'_j \mid P''_j)] \mid \mathbf{N}_k[x(z); P_k]) \longrightarrow R'.$$

- If $\alpha_j; P_j = \bar{x}.l'; P_j$, then $A = \oplus\{l : B_l\}_{l \in L}$ and $\bar{A} = \&\{\bar{B}_l\}_{l \in L}$ for some $(B_l)_{l \in L}$ where $l' \in L$. Hence, $\alpha_k; P_k = x.\text{case}\{l : P_k^l\}_{l \in L}$. By rule $[\rightarrow_{\oplus\&}]$, there is R' such that

$$(\nu x)(N_j[\bar{x}.l'; P_j] \mid N_k[x.\text{case}\{l : P_k^l\}_{l \in L}]) \longrightarrow R'.$$

- If $\alpha_j; P_j = \bar{x}.\text{some}; P_j$, then $A = \oplus B$ and $\bar{A} = \&\bar{B}$ for some B . Hence, $\alpha_k = x.\text{some}_{w_1, \dots, w_n}; P_k$ for some w_1, \dots, w_n . By rule $[\rightarrow_{\text{some}}]$, there is R' such that

$$(\nu x)(N_j[\bar{x}.\text{some}; P_j] \mid N_k[x.\text{some}_{w_1, \dots, w_n}; P_k]) \longrightarrow R'.$$

- If $\alpha_j; P_j = \bar{x}.\text{none}$, then $A = \oplus B$ and $\bar{A} = \&\bar{B}$ for some B . Hence, $\alpha_k = x.\text{some}_{w_1, \dots, w_n}; P_k$ for some w_1, \dots, w_n . By rule $[\rightarrow_{\text{none}}]$, there is R' such that

$$(\nu x)(N_j[\bar{x}.\text{none}] \mid N_k[x.\text{some}_{w_1, \dots, w_n}; P_k]) \longrightarrow R'.$$

- If $\alpha_j; P_j = ?\bar{x}[y]; P_j$ for some y , then $A = ?B$ and $\bar{A} = !\bar{B}$ for some B . Hence, $\alpha_k; P_k = !x(z); P_k$ for some z . By rule $[\rightarrow_{?!}]$, there is R' such that

$$(\nu x)(N_j[?\bar{x}[y]; P_j] \mid N_k[!x(z); P_k]) \longrightarrow R'.$$

- Otherwise, α_j is a receiving prefix and α_k is thus a sending prefix. By cases on α_k , the proof is analogous to above.

In conclusion,

$$\begin{aligned} S[\alpha_i; P_i]_{i \in I} &= N[(\nu x)(N_j[\alpha_j; P_j] \mid N_k[\alpha_k; P_k])] \longrightarrow N[R'] = R && \text{(by rules } [\rightarrow_{\nu}], [\rightarrow_{\cdot}], [\rightarrow_{\#}] \text{)} \\ P &\longrightarrow R. && \text{(by rule } [\rightarrow_{\equiv}] \text{)} \end{aligned} \quad \square$$

E.2 Lazy Semantics

Subject Reduction

Lemma 9. *For both of the following items, assume $\Gamma \# \Delta$.*

1. *If $\forall i \in I. \forall j \in J. \text{bn}(\mathbf{C}_i) \# \text{fn}(\mathbf{D}_j)$ and $\forall i \in I. \mathbf{C}_i[\bar{x}[y_i]; (P_i \mid Q_i)] \vdash \Gamma, x:A \otimes B$ and $\forall j \in J. \mathbf{D}_j[x(z); R_j] \vdash \Delta, x:\bar{A} \wp \bar{B}$, then $\prod_{i \in I} \mathbf{C}_i[(\nu x)(Q_i \mid (\nu w)(P_i\{w/y_i\} \mid \prod_{j \in J} \mathbf{D}_j[R_j\{w/z\}]))] \vdash \Gamma, \Delta$.*
2. *If $\forall i \in I. \forall j \in J. \text{bn}(\mathbf{D}_j) \# \text{fn}(\mathbf{C}_i)$ and $\forall i \in I. \mathbf{C}_i[?\bar{x}[y_i]; P_i] \vdash \Gamma, x:?A$ and $\forall j \in J. \mathbf{D}_j[!x(z); Q_j] \vdash \Delta, x:!\bar{A}$, then $\prod_{j \in J} \mathbf{D}_j[(\nu x)((\nu w)(\prod_{i \in I} \mathbf{C}_i[P_i\{w/y_i\}] \mid Q_j\{w/z\}) \mid !x(z); Q_j)] \vdash \Gamma, \Delta$.*

Proof. Both items follow by induction on the structures of the D-contexts. For each item, we detail the base case, where $\forall i \in I. \mathbf{C}_i = [\cdot]$ and $\forall j \in J. \mathbf{D}_j = [\cdot]$. The inductive cases follow from the IH straightforwardly.

1.

$$\begin{aligned} \forall i \in I. \frac{P_i \vdash \Gamma, y_i:A \quad Q_i \vdash \Delta, x:B}{\bar{x}[y_i]; (P_i \mid Q_i) \vdash \Gamma, \Delta, x:A \otimes B} \quad \forall j \in J. \frac{R_j \vdash \Lambda, z:\bar{A}, x:\bar{B}}{x(z); R_j \vdash \Lambda, x:\bar{A} \wp \bar{B}} &\implies \\ \forall i \in I. P_i\{w/y_i\} \vdash \Gamma, w:A \quad \frac{\forall j \in J. R_j\{w/z\} \vdash \Lambda, w:\bar{A}, x:\bar{B}}{\prod_{j \in J} R_j\{w/z\} \vdash \Lambda, w:\bar{A}, x:\bar{B}} & \\ \forall i \in I. Q_i \vdash \Delta, x:B \quad \forall i \in I. (\nu w)(P_i\{w/y_i\} \mid \prod_{j \in J} R_j\{w/z\}) \vdash \Gamma, \Lambda, x:\bar{B} & \\ \frac{\forall i \in I. (\nu x)(Q_i \mid (\nu w)(P_i\{w/y_i\} \mid \prod_{j \in J} R_j\{w/z\})) \vdash \Gamma, \Delta, \Lambda}{\prod_{i \in I} (\nu x)(Q_i \mid (\nu w)(P_i\{w/y_i\} \mid \prod_{j \in J} R_j\{w/z\})) \vdash \Gamma, \Delta, \Lambda} & \end{aligned}$$

2. This item depends on whether $x \in \text{fn}(P_i)$ or not, for each $i \in I$. For simplicity, we only consider the cases where either $\forall i \in I. x \in \text{fn}(P_i)$ or $\forall i \in I. x \notin \text{fn}(P_i)$.

– $\forall i \in I. x \in \text{fn}(P_i)$.

$$\begin{array}{c}
\forall i \in I. \frac{\frac{P_i\{x'/x\} \vdash \Gamma, y_i:A, x':?A}{?x[y_i]; (P_i\{x'/x\}) \vdash \Gamma, x:?A, x':?A}}{?x[y_i]; P_i \vdash \Gamma, x:?A} \quad \forall j \in J. \frac{Q_j \vdash ?\Delta, x:\bar{A}}{!x(z); Q_j \vdash ?\Delta, x:\bar{A}} \quad \Longrightarrow \\
\\
\frac{\forall i \in I. \frac{P_i\{w/y_i\} \vdash \Gamma, w:A, x:?A}{\prod_{i \in I} P_i\{w/y_i\} \vdash \Gamma, w:A, x:?A} \quad \forall j \in J. \frac{Q_j\{w/z\}\{v'/v\}_{v \in ?\Delta} \vdash ?\Delta', w:\bar{A}}{Q_j\{w/z\}\{v'/v\}_{v \in ?\Delta} \vdash ?\Delta', w:\bar{A}}}{\prod_{i \in I} P_i\{w/y_i\} \mid Q_j\{w/z\}\{v'/v\}_{v \in ?\Delta} \vdash \Gamma, ?\Delta', x:?A} \quad \forall j \in J. \frac{Q_j \vdash ?\Delta, x:\bar{A}}{!x(z); Q_j \vdash ?\Delta, x:\bar{A}} \\
\frac{\forall j \in J. (\nu w) \left(\prod_{i \in I} P_i\{w/y_i\} \mid Q_j\{w/z\}\{v'/v\}_{v \in ?\Delta} \right) \vdash \Gamma, ?\Delta', x:?A \quad \forall j \in J. \frac{Q_j \vdash ?\Delta, x:\bar{A}}{!x(z); Q_j \vdash ?\Delta, x:\bar{A}}}{\prod_{j \in J} (\nu x) \left((\nu w) \left(\prod_{i \in I} P_i\{w/y_i\} \mid Q_j\{w/z\}\{v'/v\}_{v \in ?\Delta} \right) \mid !x(z); Q_j \right) \vdash \Gamma, ?\Delta, ?\Delta'} \\
\frac{\prod_{j \in J} (\nu x) \left((\nu w) \left(\prod_{i \in I} P_i\{w/y_i\} \mid Q_j\{w/z\}\{v'/v\}_{v \in ?\Delta} \right) \mid !x(z); Q_j \right) \vdash \Gamma, ?\Delta, ?\Delta'}{\prod_{j \in J} (\nu x) \left((\nu w) \left(\prod_{i \in I} P_i\{w/y_i\} \mid Q_j\{w/z\} \right) \mid !x(z); Q_j \right) \vdash \Gamma, ?\Delta}
\end{array}$$

– $\forall i \in I. x \notin \text{fn}(P_i)$.

$$\begin{array}{c}
\forall i \in I. \frac{P_i \vdash \Gamma, y_i:A}{?x[y_i]; P_i \vdash \Gamma, x:?A} \quad \forall j \in J. \frac{Q_j \vdash ?\Delta, x:\bar{A}}{!x(z); Q_j \vdash ?\Delta, x:\bar{A}} \quad \Longrightarrow \\
\\
\frac{\forall i \in I. \frac{P_i\{w/y_i\} \vdash \Gamma, w:A}{\prod_{i \in I} P_i\{w/y_i\} \vdash \Gamma, w:A} \quad \forall j \in J. \frac{Q_j\{w/z\}\{v'/v\}_{v \in ?\Delta} \vdash ?\Delta', w:\bar{A}}{Q_j\{w/z\}\{v'/v\}_{v \in ?\Delta} \vdash ?\Delta', w:\bar{A}}}{\prod_{i \in I} P_i\{w/y_i\} \mid Q_j\{w/z\}\{v'/v\}_{v \in ?\Delta} \vdash \Gamma, ?\Delta'} \\
\frac{\forall j \in J. (\nu w) \left(\prod_{i \in I} P_i\{w/y_i\} \mid Q_j\{w/z\}\{v'/v\}_{v \in ?\Delta} \right) \vdash \Gamma, ?\Delta'}{\prod_{i \in I} P_i\{w/y_i\} \mid Q_j\{w/z\}\{v'/v\}_{v \in ?\Delta} \vdash \Gamma, ?\Delta', x:?A} \quad \forall j \in J. \frac{Q_j \vdash ?\Delta, x:\bar{A}}{!x(z); Q_j \vdash ?\Delta, x:\bar{A}} \\
\frac{\forall j \in J. (\nu x) \left((\nu w) \left(\prod_{i \in I} P_i\{w/y_i\} \mid Q_j\{w/z\}\{v'/v\}_{v \in ?\Delta} \right) \mid !x(z); Q_j \right) \vdash \Gamma, ?\Delta, ?\Delta'}{\prod_{j \in J} (\nu x) \left((\nu w) \left(\prod_{i \in I} P_i\{w/y_i\} \mid Q_j\{w/z\}\{v'/v\}_{v \in ?\Delta} \right) \mid !x(z); Q_j \right) \vdash \Gamma, ?\Delta, ?\Delta'} \\
\frac{\prod_{j \in J} (\nu x) \left((\nu w) \left(\prod_{i \in I} P_i\{w/y_i\} \mid Q_j\{w/z\}\{v'/v\}_{v \in ?\Delta} \right) \mid !x(z); Q_j \right) \vdash \Gamma, ?\Delta, ?\Delta'}{\prod_{j \in J} (\nu x) \left((\nu w) \left(\prod_{i \in I} P_i\{w/y_i\} \mid Q_j\{w/z\} \right) \mid !x(z); Q_j \right) \vdash \Gamma, ?\Delta}
\end{array}$$

□

Theorem 14 (SR for the Lazy Semantics). *If $P \vdash \Gamma$ and $P \rightsquigarrow_S Q$, then $Q \vdash \Gamma$.*

Proof. By induction on the derivation of the reduction.

– Rule $[\rightsquigarrow_{\text{ID}}]$.

$$\frac{\forall i \in I. \mathbf{C}_i[[x \leftrightarrow y]] \vdash \Gamma, x:A}{\prod_{i \in I} \mathbf{C}_i[[x \leftrightarrow y]] \vdash \Gamma, x:A} \quad \frac{Q \vdash \Delta, x:\bar{A}}{(\nu x) \left(\prod_{i \in I} \mathbf{C}_i[[x \leftrightarrow y]] \mid Q \right) \vdash \Gamma, \Delta} \quad \Longrightarrow \quad \frac{\forall i \in I. \mathbf{C}_i[Q\{y/x\}] \vdash \Gamma, \Delta \text{ (Lemma 3.1)}}{\prod_{i \in I} \mathbf{C}_i[Q\{y/x\}] \vdash \Gamma, \Delta}$$

– Rule $[\rightsquigarrow_{\text{none}}]$.

$$\begin{array}{c}
\begin{array}{c}
\forall i \in I. \\
\mathbf{C}_i[\bar{x}.\text{none}] \vdash \Gamma, x:\&A \\
\text{(Lemma 2.8)}
\end{array}
\quad
\begin{array}{c}
\forall j \in J. \\
\mathbf{D}_j[x.\text{some}_{w_1, \dots, w_n}; Q_j] \vdash \Delta, x:\oplus\bar{A} \\
\text{(Lemma 2.9)}
\end{array} \\
\hline
\begin{array}{c}
\prod_{i \in I} \mathbf{C}_i[\bar{x}.\text{none}] \vdash \Gamma, x:\&A \quad \prod_{j \in J} \mathbf{D}_j[x.\text{some}_{w_1, \dots, w_n}; Q_j] \vdash \Delta, x:\oplus\bar{A} \\
(\nu x) \left(\prod_{i \in I} \mathbf{C}_i[\bar{x}.\text{none}] \mid \prod_{j \in J} \mathbf{D}_j[x.\text{some}_{w_1, \dots, w_n}; Q_j] \right) \vdash \Gamma, \Delta
\end{array}
\end{array}
\Rightarrow
\begin{array}{c}
\begin{array}{c}
\forall i \in I. \mathbf{C}_i[\mathbf{0}] \vdash \Gamma \quad \text{(Lemma 3.8)} \\
\prod_{i \in I} \mathbf{C}_i[P_i] \vdash \Gamma
\end{array}
\quad
\begin{array}{c}
\forall j \in J. \mathbf{D}_j[\bar{w}_1.\text{none} \mid \dots \mid \bar{w}_n.\text{none}] \vdash \Delta \quad \text{(Lemma 3.9)} \\
\prod_{j \in J} \mathbf{D}_j[\bar{w}_1.\text{none} \mid \dots \mid \bar{w}_n.\text{none}] \vdash \Delta
\end{array} \\
\hline
\prod_{i \in I} \mathbf{C}_i[\mathbf{0}] \mid \prod_{j \in J} \mathbf{D}_j[\bar{w}_1.\text{none} \mid \dots \mid \bar{w}_n.\text{none}] \vdash \Gamma, \Delta
\end{array}$$

– Rule $[\rightsquigarrow_{?!}]$.

$$\begin{array}{c}
\begin{array}{c}
\forall i \in I. \mathbf{C}_i[?\bar{x}[y_i]; P_i] \vdash \Gamma, x:?A \quad \text{(Lemma 2.10)} \\
\prod_{i \in I} \mathbf{C}_i[?\bar{x}[y_i]; P_i] \vdash \Gamma, x:?A
\end{array}
\quad
\begin{array}{c}
\forall j \in J. \mathbf{D}_j[!x(z); Q_j] \vdash \Delta, x:\bar{A} \quad \text{(Lemma 2.11)} \\
\prod_{j \in J} \mathbf{D}_j[!x(z); Q_j] \vdash \Delta, x:\bar{A}
\end{array} \\
\hline
(\nu x) \left(\prod_{i \in I} \mathbf{C}_i[?\bar{x}[y_i]; P_i] \mid \prod_{j \in J} \mathbf{D}_j[!x(z); Q_j] \right) \vdash \Gamma, \Delta
\end{array}
\Rightarrow
\begin{array}{c}
\prod_{j \in J} \mathbf{D}_j \left[(\nu x) \left((\nu w) \left(\prod_{i \in I} \mathbf{C}_i[P_i\{w/z\}] \mid Q_j\{w/z\} \right) \mid !x(z); Q_j \right) \right] \vdash \Gamma, \Delta \quad \text{(Lemma 9.2)}
\end{array}$$

□

Deadlock Freedom

Definition 15 (Multi-hole Non-deterministic Reduction Contexts).

$$\mathbf{M} ::= [\cdot] \mid (\nu x)(P \mid \mathbf{M}) \mid P \mid \mathbf{M} \mid \mathbf{M} \# \mathbf{M}$$

Lemma 10. *If $\mathbf{N}[\alpha; P] \vdash \Gamma, x:A$ and $x = \text{subj}(\alpha)$, then there are \mathbf{M} and $(\alpha_i; P_i)_{i \in I}$ such that $\mathbf{N}[\alpha; P] = \mathbf{M}[\alpha_i; P_i]_{i \in I}$ where $x \notin \text{fn}(\mathbf{M})$ and $x \in \bigcap_{i \in I} \text{fn}(\alpha_i; P_i)$ and there is $i' \in I$ such that $\alpha_{i'}; P_{i'} = \alpha; P$.*

Lemma 11. *For every multi-hole ND-context \mathbf{M} with indices I :*

- If \mathbf{M} has two or more holes, there are \mathbf{C}, \mathbf{M}_1 with indices I_1 , \mathbf{M}_2 with indices I_2 such that $\mathbf{M} = \mathbf{C}[\mathbf{M}_1 \# \mathbf{M}_2]$ where $I_1 \# I_2$ and $I = I_1 \cup I_2$.
- If \mathbf{M} has only one hole, there is \mathbf{C} such that $\mathbf{M} = \mathbf{C}$.

Definition 16.

$$\text{flat}(\mathbf{C}[\mathbf{M} \# \mathbf{M}']) := \text{flat}(\mathbf{C}[\mathbf{M}]) \# \text{flat}(\mathbf{C}[\mathbf{M}']) \quad \text{flat}(\mathbf{C}) := \mathbf{C}$$

Lemma 12. *If $\mathbf{M}[P_i]_{i \in I} \vdash \Gamma$ where $x \notin \text{fn}(\mathbf{M})$ and $\forall i \in I. x \in \text{fn}(P_i)$, then there are $(\mathbf{C}_i)_{i \in I}$ such that $\text{flat}(\mathbf{M})[P_i]_{i \in I} = \prod_{i \in I} \mathbf{C}_i[P_i]$ where $\forall i \in I. x \notin \text{fn}(\mathbf{C}_i)$.*

Lemma 13. *If*

$$(\nu x)(\mathbf{N}[\text{flat}(\mathbf{M})[\alpha_i; P_i]_{i \in I}] \mid \mathbf{N}'[\text{flat}(\mathbf{M}')[\beta_j; Q_j]_{j \in J}]) \rightsquigarrow_S R,$$

then

$$(\nu x)(\mathbf{N}[\mathbf{M}[\alpha_i; P_i]_{i \in I}] \mid \mathbf{N}'[\mathbf{M}'[\beta_j; Q_j]_{j \in J}]) \rightsquigarrow_S R.$$

Proof. By induction on the structures of \mathbf{M} and \mathbf{M}' . By Lemma 11, we only have to consider two cases for \mathbf{M} ($\mathbf{M} = \mathbf{C}[\mathbf{M}_1 \# \mathbf{M}_2]$ and $\mathbf{M} = \mathbf{C}$), and similarly for \mathbf{M}' . We only detail the base case ($\mathbf{M} = \mathbf{C}$ and $\mathbf{M}' = \mathbf{C}'$) and a representative inductive case ($\mathbf{M} = \mathbf{C}[\mathbf{M}_1 \# \mathbf{M}_2]$ and $\mathbf{M}' = \mathbf{C}'$).

- $\mathbf{M} = \mathbf{C}$ and $\mathbf{M}' = \mathbf{C}'$. Note that \mathbf{M} and \mathbf{M}' have only one hole; w.l.o.g., assume $I = J = \{1\}$.

$$\begin{aligned}
\text{flat}(\mathbf{M})[\alpha_1; P_1] &= \text{flat}(\mathbf{C})[\alpha_1; P_1] = \mathbf{C}[\alpha_1; P_1] = \mathbf{M}[\alpha_1; P_1] \\
\text{flat}(\mathbf{M}')[\beta_1; Q_1] &= \text{flat}(\mathbf{C}')[\beta_1; Q_1] = \mathbf{C}'[\beta_1; Q_1] = \mathbf{M}'[\beta_1; Q_1]
\end{aligned}$$

The thesis follows by assumption and equality.

– $\mathbf{M} = \mathbf{C}[\mathbf{M}_1 \# \mathbf{M}_2]$ and $\mathbf{M}' = \mathbf{C}'$. Note that \mathbf{M}' has only one hole; w.l.o.g., assume $J = \{1\}$.

$$\begin{aligned} & (\nu x)(\mathbf{N}[\mathit{flat}(\mathbf{M})[\alpha_i; P_i]_{i \in I} \mid \mathbf{N}'[\mathit{flat}(\mathbf{M}')[\beta_1; Q_1]]) \\ &= (\nu x)(\mathbf{N}[\mathit{flat}(\mathbf{C}[\mathbf{M}_1 \# \mathbf{M}_2])[\alpha_i; P_i]_{i \in I} \mid \mathbf{N}'[\mathit{flat}(\mathbf{C}')[\beta_1; Q_1]]) \\ &= (\nu x)(\mathbf{N}[(\mathit{flat}(\mathbf{C}[\mathbf{M}_1]) \# \mathit{flat}(\mathbf{C}[\mathbf{M}_2]))[\alpha_i; P_i]_{i \in I} \mid \mathbf{N}'[\mathbf{C}'[\beta_1; Q_1]]) \end{aligned} \quad (6)$$

There are I_1 and I_2 such that $I_1 \# I_2$ and $I = I_1 \cup I_2$ and

$$\begin{aligned} & (\nu x)(\mathbf{N}[\mathit{flat}(\mathbf{M})[\alpha_i; P_i]_{i \in I} \mid \mathbf{N}'[\mathit{flat}(\mathbf{M}')[\beta_1; Q_1]]) \\ &= (\nu x)(\mathbf{N}[\mathit{flat}(\mathbf{C}[\mathbf{M}_1])[\alpha_i; P_i]_{i \in I_1} \# \mathit{flat}(\mathbf{C}[\mathbf{M}_2])[\alpha_i; P_i]_{i \in I_2} \mid \mathbf{N}'[\mathbf{C}'[\beta_1; Q_1]]). \end{aligned} \quad (\text{by (6)}) \quad (7)$$

Let $\mathbf{N}_1 = \mathbf{N}[\cdot \# \mathit{flat}(\mathbf{C}[\mathbf{M}_2])[\alpha_i; P_i]_{i \in I_2}]$.

$$\begin{aligned} & (\nu x)(\mathbf{N}[\mathit{flat}(\mathbf{M})[\alpha_i; P_i]_{i \in I} \mid \mathbf{N}'[\mathit{flat}(\mathbf{M}')[\beta_1; Q_1]]) \\ &= (\nu x)(\mathbf{N}_1[\mathit{flat}(\mathbf{C}[\mathbf{M}_1])[\alpha_i; P_i]_{i \in I_1} \mid \mathbf{N}'[\mathbf{C}'[\beta_1; Q_1]]) \quad (\text{by (7)}) \\ &\rightsquigarrow_S R \quad (\text{by assumption}) \end{aligned} \quad (8)$$

$$\begin{aligned} & (\nu x)(\mathbf{N}_1[\mathbf{C}[\mathbf{M}_1][\alpha_i; P_i]_{i \in I_1} \mid \mathbf{N}'[\mathbf{C}'[\beta_1; Q_1]]) \\ &= (\nu x)(\mathbf{N}[\mathbf{C}[\mathbf{M}_1][\alpha_i; P_i]_{i \in I_1} \# \mathit{flat}(\mathbf{C}[\mathbf{M}_2])[\alpha_i; P_i]_{i \in I_2} \mid \mathbf{N}'[\mathbf{C}'[\beta_1; Q_1]]) \\ &\rightsquigarrow_S R \quad (\text{by IH on (8)}) \end{aligned} \quad (9)$$

Let $\mathbf{N}_2 = \mathbf{N}[\mathbf{C}[\mathbf{M}_1][\alpha_i; P_i]_{i \in I_1} \# \cdot]$.

$$\begin{aligned} & (\nu x)(\mathbf{N}[\mathbf{C}[\mathbf{M}_1][\alpha_i; P_i]_{i \in I_1} \# \mathit{flat}(\mathbf{C}[\mathbf{M}_2])[\alpha_i; P_i]_{i \in I_2} \mid \mathbf{N}'[\mathbf{C}'[\beta_1; Q_1]]) \\ &= (\nu x)(\mathbf{N}_2[\mathit{flat}(\mathbf{C}[\mathbf{M}_2])[\alpha_i; P_i]_{i \in I_2} \mid \mathbf{N}'[\mathbf{C}'[\beta_1; Q_1]]) \\ &\rightsquigarrow_S R \quad (\text{by (9)}) \end{aligned} \quad (10)$$

$$\begin{aligned} & (\nu x)(\mathbf{N}_2[\mathbf{C}[\mathbf{M}_2][\alpha_i; P_i]_{i \in I_2} \mid \mathbf{N}'[\mathbf{C}'[\beta_1; Q_1]]) \\ &= (\nu x)(\mathbf{N}[\mathbf{C}[\mathbf{M}_1][\alpha_i; P_i]_{i \in I_1} \# \mathbf{C}[\mathbf{M}_2][\alpha_i; P_i]_{i \in I_2} \mid \mathbf{N}'[\mathbf{C}'[\beta_1; Q_1]]) \\ &= (\nu x)(\mathbf{N}[\mathbf{C}[\mathbf{M}_1[\alpha_i; P_i]_{i \in I_1} \# \mathbf{C}[\mathbf{M}_2[\alpha_i; P_i]_{i \in I_2}]] \mid \mathbf{N}'[\mathbf{C}'[\beta_1; Q_1]]) \\ &\rightsquigarrow_S R \quad (\text{by IH on (10)}) \end{aligned} \quad (11)$$

$$\begin{aligned} & (\nu x)(\mathbf{N}[\mathbf{C}[\mathbf{M}_1[\alpha_i; P_i]_{i \in I_1} \# \mathbf{M}_2[\alpha_i; P_i]_{i \in I_2}]] \mid \mathbf{N}'[\mathbf{C}'[\beta_1; Q_1]]) \\ &= (\nu x)(\mathbf{N}[\mathbf{C}[(\mathbf{M}_1 \# \mathbf{M}_2)[\alpha_i; P_i]_{i \in I}]] \mid \mathbf{N}'[\mathbf{C}'[\beta_1; Q_1]]) \\ &= (\nu x)(\mathbf{N}[\mathbf{C}[\mathbf{M}_1 \# \mathbf{M}_2][\alpha_i; P_i]_{i \in I} \mid \mathbf{N}'[\mathbf{C}'[\beta_1; Q_1]]) \\ &= (\nu x)(\mathbf{N}[\mathbf{M}[\alpha_i; P_i]_{i \in I} \mid \mathbf{N}'[\mathbf{M}'[\beta_1; Q_1]]) \\ &\rightsquigarrow_S R \quad (\text{by rule } [\rightsquigarrow_{\nu \#}] \text{ on (11)}) \quad \square \end{aligned}$$

Theorem 15. *If $P \vdash \Gamma$ and $P \longrightarrow R$, then $P \rightsquigarrow_S R$.*

Proof. By induction on the derivation of the reduction. The inductive cases of rules $[\rightarrow_{\equiv}]$, $[\rightarrow_{\nu}]$, $[\rightarrow_{\perp}]$, and $[\rightarrow_{\#}]$ follow from the IH straightforwardly, using the corresponding closure rule for \rightsquigarrow . As representative base case, we consider rule $[\rightarrow_{\perp \perp}]$: $P = (\nu x)(\mathbf{N}[\overline{x}] \mid \mathbf{N}'[x(); Q]) \longrightarrow R$.

By inversion of typing, $\mathbf{N}[\overline{x}] \vdash \Gamma, x:A$, so by Lemma 10, there are \mathbf{M} and $(\alpha_i; P_i)_{i \in I}$ such that $\mathbf{N}[\overline{x}] = \mathbf{M}[\alpha_i; P_i]_{i \in I}$ where $x \notin \mathit{fn}(\mathbf{M})$ and $x \in \bigcap_{i \in I} \mathit{fn}(\alpha_i; P_i)$ and there is $i' \in I$ such that $\alpha_{i'}; P_{i'} = \overline{x}$. Similarly, there are \mathbf{M}' and $(\beta_j; Q_j)_{j \in J}$ such that $\mathbf{N}'[x(); Q] = \mathbf{M}'[\beta_j; Q_j]_{j \in J}$ where $x \notin \mathit{fn}(\mathbf{M}')$ and $x \in \bigcap_{j \in J} \mathit{fn}(\beta_j; Q_j)$ and there is $j' \in J$ such that $\beta_{j'}; Q_{j'} = x(); Q$.

By Lemma 12, $\mathit{flat}(\mathbf{M})[\alpha_i; P_i]_{i \in I} = \prod_{i \in I} \mathbf{C}_i[\alpha_i; P_i]$ and $\mathit{flat}(\mathbf{M}')[\beta_j; Q_j]_{j \in J} = \prod_{j \in J} \mathbf{C}'_j[\beta_j; Q_j]$. By typability, there is $I' \subseteq I$ such that $\forall i \in I'. \alpha_i \bowtie \overline{x}$ and $\forall i \in I \setminus I'. \alpha_i \not\bowtie \overline{x}$; hence, $i' \in I'$. Similarly, there is $J' \subseteq J$ such that $\forall j \in J'. \beta_j \bowtie x()$ and $\forall j \in J \setminus J'. \beta_j \not\bowtie x()$; hence, $j' \in J'$. Then, by Definition 3, $\mathit{flat}(\mathbf{M})[\alpha_i; P_i]_{i \in I} \succeq_x \prod_{i \in I'} \mathbf{C}_i[\overline{x}]$ and $\mathit{flat}(\mathbf{M}')[\beta_j; Q_j]_{j \in J} \succeq_x \prod_{j \in J'} \mathbf{C}'_j[x(); Q_j]$. By rule $[\rightsquigarrow_{\perp \perp}]$,

$$(\nu x)(\prod_{i \in I'} \mathbf{C}_i[\overline{x}] \mid \prod_{j \in J'} \mathbf{C}'_j[x(); Q_j]) \rightsquigarrow_x R,$$

so by rule $[\rightsquigarrow_{\succeq_x}]$,

$$(\nu x)(\mathit{flat}(\mathbf{M})[\alpha_i; P_i]_{i \in I} \mid \mathit{flat}(\mathbf{M}')[\beta_j; Q_j]_{j \in J}) \rightsquigarrow_x R.$$

Then, by Lemma 13,

$$(\nu x)(\mathbf{M}[\alpha_i; P_i]_{i \in I} \mid \mathbf{M}'[\beta_j; Q_j]_{j \in J}) \rightsquigarrow_x R.$$

As second base case, we consider rule $[\rightarrow_{\text{ID}}]$: $P = (\nu x)(\mathbf{N}[x \leftrightarrow y] \mid Q) \longrightarrow R$.

By inversion of typing, $\mathbf{N}[x \leftrightarrow y] \vdash \Gamma, x:A, y:\bar{A}$, so by Lemma 10, there are \mathbf{M} and $(\alpha_i; P_i)_{i \in I}$ such that $\mathbf{N}[x \leftrightarrow y] = \mathbf{M}[\alpha_i; P_i]_{i \in I}$ where $x \notin \text{fn}(\mathbf{M})$ and $x \in \bigcap_{i \in I} \text{fn}(\alpha_i; P_i)$ and there is $i' \in I$ such that $\alpha_{i'}; P_{i'} = [x \leftrightarrow y]$.

By Lemma 12, $\text{flat}(\mathbf{M})[\alpha_i; P_i]_{i \in I} = \prod_{i \in I} \mathbf{C}_i[\alpha_i; P_i]$. By typability, there is $I' \subseteq I$ such that $\forall i \in I'. \alpha_i = [x \leftrightarrow y]$ and $\forall i \in I \setminus I'. \alpha_i \neq [x \leftrightarrow y]$; hence, $i' \in I'$.

Then, by Definition 3, $\text{flat}(\mathbf{M})[\alpha_i; P_i]_{i \in I} \succeq_{x,y} \prod_{i \in I'} \mathbf{C}_i[[x \leftrightarrow y]]$. By rule $[\rightsquigarrow_{\text{ID}}]$,

$$(\nu x) \left(\prod_{i \in I'} \mathbf{C}_i[[x \leftrightarrow y]] \mid Q \right) \rightsquigarrow_{x,y} R,$$

so by rule $[\rightsquigarrow_{\succeq_{x,y}}]$,

$$(\nu x)(\text{flat}(\mathbf{M})[\alpha_i; P_i]_{i \in I} \mid Q) \rightsquigarrow_{x,y} R.$$

Then, by Lemma 13,

$$(\nu x)(\mathbf{M}[\alpha_i; P_i]_{i \in I} \mid Q) \rightsquigarrow_{x,y} R. \quad \square$$

Theorem 16 (DF: Lazy Semantics). *If $P \vdash \emptyset$ and $P \neq \mathbf{0}$, then $P \rightsquigarrow_S R$ for some S and R .*

Proof. As a corollary of Theorems 8 and 15. □

F Proofs of Subject Reduction and Subject Expansion for λ_c

Here we prove Theorems 3 and 4 (subject reduction and subject expansion, respectively) for λ_c .

F.1 Subject Reduction

Lemma 14 (Substitution Lemma for λ_c).

1. (Linear) If $\Theta; \Gamma, x : \sigma \vDash M : \tau$, $\text{head}(M) = x$, and $\Theta; \Delta \vDash N : \sigma$ then $\Theta; \Gamma, \Delta \vDash M\{N/x\} : \tau$.
2. (Unrestricted) If $\Theta, x^i : \eta; \Gamma \vDash M : \tau$, $\text{head}(M) = x[i]$, $\eta_i = \sigma$, and $\Theta; \cdot \vDash N : \sigma$ then $\Theta, x^i : \eta; \Gamma \vDash M\{N/x[i]\}$.

Proof.

1. By structural induction on M with $\text{head}(M) = x$. There are six cases to be analyzed:

(a) $M = x$

In this case, $\Theta; x : \sigma \vDash x : \sigma$ and $\Gamma = \emptyset$. Observe that $x\{N/x\} = N$, since $\Delta \vDash N : \sigma$, by hypothesis, the result follows.

(b) $M = M' B$.

Then $\text{head}(M' B) = \text{head}(M') = x$, and the derivation is the following:

$$[\text{FS:app}] \frac{\Theta; \Gamma_1, x : \sigma \vDash M' : (\delta^j, \eta) \rightarrow \tau \quad \Theta; \Gamma_2 \vDash B : (\delta^k, \epsilon) \quad \eta \sim \epsilon}{\Theta; \Gamma_1, \Gamma_2, x : \sigma \vDash M' B : \tau}$$

where $\Gamma = \Gamma_1, \Gamma_2$, and j, k are non-negative integers, possibly different. Since $\Delta \vdash N : \sigma$, by IH, the result holds for M' , that is,

$$\Gamma_1, \Delta \vDash M'\{N/x\} : (\delta^j, \eta) \rightarrow \tau$$

which gives the derivation:

$$[\text{FS:app}] \frac{\Theta; \Gamma_1, \Delta \vDash M'\{N/x\} : (\delta^j, \eta) \rightarrow \tau \quad \Theta; \Gamma_2 \vDash B : (\delta^k, \epsilon) \quad \eta \sim \epsilon}{\Theta; \Gamma_1, \Gamma_2, \Delta \vDash (M'\{N/x\})B : \tau}$$

From Figure 13, $(M' B)\{N/x\} = (M'\{N/x\})B$, and the result follows.

(c) $M = M'[\tilde{y} \leftarrow y]$.

Then $\text{head}(M'[\tilde{y} \leftarrow y]) = \text{head}(M') = x$, for $y \neq x$. Therefore,

$$[\text{FS:share}] \frac{\Theta; \Gamma_1, y_1 : \delta, \dots, y_k : \delta, x : \sigma \vDash M' : \tau \quad y \notin \Gamma_1 \quad k \neq 0}{\Theta; \Gamma_1, y : \delta^k, x : \sigma \vDash M'[y_1, \dots, y_k \leftarrow x] : \tau}$$

where $\Gamma = \Gamma_1, y : \delta^k$. By IH, the result follows for M' , that is,

$$\Theta; \Gamma_1, y_1 : \delta, \dots, y_k : \delta, \Delta \vDash M'\{N/x\} : \tau$$

and we have the derivation:

$$[\text{FS:shar}] \frac{\Theta; \Gamma_1, y_1 : \delta, \dots, y_k : \delta, \Delta \vDash M'\{N/x\} : \tau \quad y \notin \Gamma_1 \quad k \neq 0}{\Theta; \Gamma_1, y : \delta^k, \Delta \vDash M'\{N/x\}[\tilde{y} \leftarrow y] : \tau}$$

From Figure 13, $M'[\tilde{y} \leftarrow y]\{N/x\} = M'\{N/x\}[\tilde{y} \leftarrow y]$, and the result follows.

(d) $M = M'[\leftarrow y]$.

Then $\text{head}(M'[\leftarrow y]) = \text{head}(M') = x$ with $x \neq y$,

$$[\text{FS:weak}] \frac{\Theta; \Gamma, x : \sigma \vDash M : \tau}{\Theta; \Gamma, y : \omega, x : \sigma \vDash M[\leftarrow y] : \tau}$$

and $M'[\leftarrow y]\{N/x\} = M'\{N/x\}[\leftarrow y]$. Then by the induction hypothesis:

$$[\text{FS:weak}] \frac{\Theta; \Gamma, \Delta \vDash M\{N/x\} : \tau}{\Theta; \Gamma, y : \omega, \Delta \vDash M\{N/x\}[\leftarrow y] : \tau}$$

(e) If $M = M'\langle C/y_1, \dots, y_k \rangle$.

Then $\text{head}(M'\langle C/y_1, \dots, y_k \rangle) = \text{head}(M') = x \neq y_1, \dots, y_k$,

$$[\text{FS:Esub}^\ell] \frac{\Theta; \Gamma_1, y_1 : \delta, \dots, y_k : \delta, x : \sigma \vDash M' : \tau \quad \Theta; \Gamma_2 \vDash C : \delta^k}{\Theta; \Gamma_1, \Gamma_2, x : \sigma \vDash M'\langle C/y_1, \dots, y_k \rangle : \tau}$$

and $M'\langle C/y_1, \dots, y_k \rangle\{N/x\} = M'\{N/x\}\langle C/y_1, \dots, y_k \rangle$. Then by the induction hypothesis:

$$[\text{FS:Esub}^\ell] \frac{\Theta; \Gamma_1, \Delta, y_1 : \delta, \dots, y_k : \delta \vDash M'\{N/x\} : \tau \quad \Theta; \Gamma_2 \vDash C : \delta^k}{\Theta; \Gamma_1, \Gamma_2, \Delta \vDash M'\{N/x\}\langle C/y_1, \dots, y_k \rangle : \tau}$$

(f) If $M = M'\llbracket U/y \rrbracket$ then $\text{head}(M'\llbracket U/y \rrbracket) = \text{head}(M') = x$, and the proofs is similar to the case above.

2. By structural induction on M with $\text{head}(M) = x[i]$. There are three cases to be analyzed:

(a) $M = x[i]$.

In this case,

$$\frac{[\text{FS:var}^\ell] \frac{}{\Theta, x^i : \eta; x : \eta_i \vDash x : \sigma}}{[\text{FS:var}^!]} \frac{}{\Theta, x^i : \eta; \cdot \vDash x[i] : \sigma}$$

and $\Gamma = \emptyset$. Observe that $x[i]\{N/x[i]\} = N$, since $\Theta, x^i : \eta; \Gamma \vDash M\{N/x[i]\}$, by hypothesis, the result follows.

(b) $M = M' B$.

In this case, $\text{head}(M' B) = \text{head}(M') = x[i]$, and one has the following derivation:

$$[\text{FS:app}] \frac{\Theta, x^i : \eta; \Gamma_1 \vDash M : (\delta^j, \epsilon) \rightarrow \tau \quad \Theta, x^i : \sigma; \Gamma_2 \vDash B : (\delta^k, \epsilon') \quad \epsilon \sim \epsilon'}{\Theta, x^i : \eta; \Gamma_1, \Gamma_2 \vDash M B : \tau}$$

where $\Gamma = \Gamma_1, \Gamma_2$, δ is a strict type and j, k are non-negative integers, possibly different.

By the induction hypothesis, we get $\Theta, x^i : \eta; \Gamma_1 \vDash M'\{N/x[i]\} : (\delta^j, \epsilon) \rightarrow \tau$, which gives the derivation:

$$[\text{FS:app}] \frac{\Theta, x^i : \eta; \Gamma_1 \vDash M'\{N/x[i]\} : (\delta^j, \epsilon) \rightarrow \tau \quad \Theta, x^i : \eta; \Gamma_2 \vDash B : (\delta^k, \epsilon') \quad \epsilon \sim \epsilon'}{\Theta, x^i : \eta; \Gamma_1, \Gamma_2 \vDash (M'\{N/x[i]\})B : \tau}$$

From Figure 13, $M'\langle\langle B/y \rangle\rangle\{N/x[i]\} = M'\{N/x[i]\}\langle\langle B/y \rangle\rangle$, and the result follows.

(c) $M = M'[\tilde{y} \leftarrow y]$.

Then $\text{head}(M'[\tilde{y} \leftarrow y]) = \text{head}(M') = x[i]$, for $y \neq x$. Therefore,

$$[\text{FS:shar}] \frac{\Theta, x^i : \eta; \Gamma_1, y_1 : \delta, \dots, y_k : \delta \vDash M' : \tau \quad y \notin \Gamma_1 \quad k \neq 0}{\Theta, x^i : \eta; \Gamma_1, y : \delta^k \vDash M'[y_1, \dots, y_k \leftarrow y] : \tau}$$

where $\Gamma = \Gamma_1, y : \delta^k$. By the induction hypothesis, the result follows for M' , that is,

$$\Theta, x^i : \eta; \Gamma_1, y_1 : \delta, \dots, y_k : \delta \vDash M'\{N/x[i]\} : \tau$$

and we have the derivation:

$$[\text{FS:shar}] \frac{\Theta, x^i : \eta; \Gamma_1, y_1 : \delta, \dots, y_k : \delta \vDash M'\{N/x[i]\} : \tau \quad y \notin \Gamma_1 \quad k \neq 0}{\Theta, x^i : \eta; \Gamma_1, y : \delta^k \vDash M'\{N/x[i]\}[\tilde{y} \leftarrow y] : \tau}$$

From Figure 13 $M'[\tilde{y} \leftarrow y]\{N/x[i]\} = M'\{N/x[i]\}[\tilde{y} \leftarrow y]$, and the result follows.

(d) $M = M'[\leftarrow y]$.

Then $\text{head}(M'[\leftarrow y]) = \text{head}(M') = x[i]$ with $x \neq y$,

$$[\text{FS:weak}] \frac{\Theta, x^i : \eta; \Gamma \vDash M : \tau}{\Theta, x^i : \eta; \Gamma, y : \omega \vDash M[\leftarrow y] : \tau}$$

and $M'[\leftarrow y]\{N/x[i]\} = M'\{N/x[i]\}[\leftarrow y]$. By the induction hypothesis:

$$[\text{FS:weak}] \frac{\Theta, x^1 : \eta; \Gamma \vDash M\{N/x[i]\} : \tau}{\Theta, x^1 : \eta; \Gamma, y : \omega \vDash M\{N/x[i]\}[\leftarrow y] : \tau}$$

(e) $M = M'\langle C/y_1, \dots, y_k \rangle$.

Then $\text{head}(M'\langle C/y_1, \dots, y_k \rangle) = \text{head}(M') = x[i]$ with $x \neq y$,

$$[\text{FS:Esub}^\ell] \frac{\Theta, x^1 : \eta; \Gamma, y_1 : \delta, \dots, y_k : \delta \vDash M : \tau \quad \Theta, x^1 : \eta; \Delta \vDash C : \delta^k}{\Theta, x^1 : \eta; \Gamma, \Delta \vDash M\langle C/y_1, \dots, y_k \rangle : \tau}$$

and $M'\langle C/y_1, \dots, y_k \rangle\{N/x[i]\} = M'\{N/x[i]\}\langle C/y_1, \dots, y_k \rangle$. By the induction hypothesis:

$$[\text{FS:Esub}^\ell] \frac{\Theta, x^1 : \eta; \Gamma, y_1 : \delta, \dots, y_k : \delta \vDash M'\{N/x[i]\} : \tau \quad \Theta, x^1 : \eta; \Delta \vDash C : \delta^k}{\Theta, x^1 : \eta; \Gamma, \Delta \vDash M'\{N/x[i]\}\langle C/y_1, \dots, y_k \rangle : \tau}$$

(f) $M = M'\llbracket U/y^1 \rrbracket$.

Then $\text{head}(M'\llbracket U/y^1 \rrbracket) = \text{head}(M') = x[i]$,

$$[\text{FS:Esub}^1] \frac{\Theta, x^1 : \eta, y : \epsilon_1; \Gamma \vDash M' : \tau \quad \Theta, x^1 : \eta; - \vDash U : \epsilon_2 \quad \epsilon_1 \sim \epsilon_2}{\Theta; \Gamma \vDash M'\llbracket U/y^1 \rrbracket : \tau}$$

and $M'\llbracket U/y^1 \rrbracket\{N/x[i]\} = M'\{N/x[i]\}\llbracket U/y^1 \rrbracket$. Then by the induction hypothesis:

$$[\text{FS:Esub}^1] \frac{\Theta, x^1 : \eta, y : \epsilon_1; \Gamma \vDash M'\{N/x[i]\} : \tau \quad \Theta, x^1 : \eta; - \vDash U : \epsilon_2 \quad \epsilon_1 \sim \epsilon_2}{\Theta; \Gamma \vDash M'\{N/x[i]\}\llbracket U/y^1 \rrbracket : \tau}$$

□

Theorem 17 (SR in λ_C). *If $\Theta; \Gamma \vDash M : \tau$ and $M \longrightarrow M'$ then $\Theta; \Gamma \vDash M' : \tau$.*

Proof. By structural induction on the reduction rule from Fig. 13 applied in $M \longrightarrow M'$.

1. **Rule [RS:Beta].**

Then $M = (\lambda x. N[\tilde{x} \leftarrow x])B$ and the reduction is:

$$[\text{RS:Beta}] \frac{}{(\lambda x. N[\tilde{x} \leftarrow x])B \longrightarrow N[\tilde{x} \leftarrow x] \langle\langle B/x \rangle\rangle}$$

where $M' = N[\tilde{x} \leftarrow x] \langle\langle B/x \rangle\rangle$. Since $\Theta; \Gamma \vDash M : \tau$ we get the following derivation:

$$\begin{array}{c} [\text{FS:share}] \frac{\Theta, x^1 : \eta; \Gamma', x_1 : \sigma, \dots, x_j : \sigma \vDash N : \tau}{\Theta, x^1 : \eta; \Gamma', x : \sigma^j \vDash N[\tilde{x} \leftarrow x] : \tau} \\ [\text{FS:abs-sh}] \frac{}{\Theta; \Gamma' \vDash \lambda x. N[\tilde{x} \leftarrow x] : (\sigma^j, \eta) \rightarrow \tau} \quad \Theta; \Delta \vDash B : (\sigma^k, \epsilon) \quad \eta \sim \epsilon \\ [\text{FS:app}] \frac{}{\Theta; \Gamma', \Delta \vDash (\lambda x. N[\tilde{x} \leftarrow x])B : \tau} \end{array}$$

for $\Gamma = \Gamma', \Delta$ and $x \notin \text{dom}(\Gamma')$. Notice that:

$$[\text{FS:share}] \frac{\Theta, x^1 : \eta; \Gamma', x_1 : \sigma, \dots, x_j : \sigma \vDash N : \tau}{\Theta, x^1 : \eta; \Gamma', x : \sigma^j \vDash N[\tilde{x} \leftarrow x] : \tau} \quad \Theta; \Delta \vDash B : (\sigma^k, \epsilon) \quad \eta \sim \epsilon \\ [\text{FS:Esub}] \frac{}{\Theta; \Gamma', \Delta \vDash N[\tilde{x} \leftarrow x] \langle\langle B/x \rangle\rangle : \tau}$$

Therefore $\Theta; \Gamma', \Delta \vDash M' : \tau$ and the result follows.

2. **Rule [RS:Ex-Sub].**

Then $M = N[x_1, \dots, x_k \leftarrow x] \langle\langle C \star U/x \rangle\rangle$ where $C = \lambda N_1 \int \dots \int \lambda N_k \int$. The reduction is:

$$[\text{RS:Ex-Sub}] \frac{C = \lambda N_1 \int \dots \int \lambda N_k \int \quad N \neq \text{fail}^{\bar{y}}}{N[x_1, \dots, x_k \leftarrow x] \langle\langle C \star U/x \rangle\rangle \longrightarrow N\langle C/x_1, \dots, x_k \rangle \llbracket U/x^1 \rrbracket}$$

and $M' = N\langle C/x_1, \dots, x_k \rangle \llbracket U/x^1 \rrbracket$. To simplify the proof we take $k = 2$, as the case $k > 2$ is similar. Therefore $C = \lambda N_1 \int \cdot \lambda N_2 \int$.

Since $\Theta; \Gamma \vDash M : \tau$ we get a derivation: (we omit the labels [FS:Esub] and [FS:share])

$$\frac{\Theta, x^1 : \eta; \Gamma', x_1 : \sigma, x_2 : \sigma \vDash N : \tau \quad x \notin \text{dom}(\Gamma) \quad k \neq 0}{\Theta, x^1 : \eta; \Gamma', x : \sigma^2 \vDash N[x_1, x_2 \leftarrow x] : \tau} \quad \Theta; \Delta \vDash C \star U : (\sigma^2, \epsilon) \quad \eta \sim \epsilon \\ \Theta; \Gamma', \Delta \vDash N[x_1, x_2 \leftarrow x] \langle\langle C \star U/x \rangle\rangle : \tau$$

where $\Gamma = \Gamma', \Delta$. Consider the wf derivation for $\Pi_{1,2}$: (we omit the labels $[\text{FS} : \text{Esub}^!]$ and $[\text{FS} : \text{Esub}^\ell]$)

$$\frac{\Theta, x^! : \eta; \Gamma', x_1 : \sigma, x_2 : \sigma \Vdash N : \tau \quad \Theta; \Delta \Vdash C : \sigma^k}{\frac{\Theta, x^! : \eta; \Gamma', \Delta \Vdash N \langle C/x_1, x_2 \rangle : \tau \quad \Theta; - \Vdash U : \epsilon \quad \eta \sim \epsilon}{\Theta; \Gamma', \Delta \Vdash N \langle C/x_1, x_2 \rangle \llbracket U/x^! \rrbracket : \tau}}$$

and the result follows.

3. Rule $[\text{RS} : \text{Fetch}^\ell]$.

Then $M = N \langle C/\tilde{x}, x_j \rangle$ where $\text{head}(N) = x_j$. The reduction is:

$$[\text{RS} : \text{Fetch}_i^\ell] \frac{\text{head}(N) = x_j}{N \langle C/\tilde{x}, x_j \rangle \longrightarrow N \{C_i/x_j\} \langle (C \setminus C_i)/\tilde{x} \rangle}$$

and $M' = N \{C_i/x_j\} \langle (C \setminus C_i)/\tilde{x} \rangle$. Since $\Theta; \Gamma \Vdash M : \tau$ we get the following derivation:

$$[\text{FS} : \text{Esub}^\ell] \frac{\Theta; \Gamma', \tilde{x} : \sigma^{k-1}, x_j : \sigma \Vdash N : \tau \quad \Theta; \Delta \Vdash C : \sigma^k}{\Theta; \Gamma', \Delta \Vdash N \langle C/\tilde{x}, x_j \rangle : \tau}$$

where $\Gamma = \Gamma', \Delta$ and $\Delta = \Delta_i, \Delta'_i$ with $\Theta; \Delta_i \Vdash C_i : \sigma$. By Lemma 14, we obtain the derivation $\Theta; \Gamma', \Delta \Vdash N \{C_i/x_j\} \langle (C \setminus C_i)/\tilde{x} \rangle : \tau$ via:

$$[\text{FS} : \text{Esub}^\ell] \frac{\Theta; \Gamma', \Delta_i, \tilde{x} : \sigma^{k-1} \Vdash N \{C_i/x_j\} : \tau \quad \Theta; \Delta'_i \Vdash C \setminus C_i : \sigma^{k-1}}{\Theta; \Gamma', \Delta \Vdash N \{C_i/x_j\} \langle (C \setminus C_i)/\tilde{x} \rangle : \tau}$$

4. Rule $[\text{RS} : \text{Fetch}^!]$.

Then $M = N \llbracket U/x^! \rrbracket$ where $\text{head}(M) = x[i]$. The reduction is:

$$[\text{RS} : \text{Fetch}^!] \frac{\text{head}(M) = x[i] \quad U_i = \lambda N_i^!}{N \llbracket U/x^! \rrbracket \longrightarrow N \{N_i/x[i]\} \llbracket U/x^! \rrbracket}$$

and $M' = N \{N_i/x[i]\} \llbracket U/x^! \rrbracket$. Since $\Theta; \Gamma \Vdash M : \tau$ we get the following derivation:

$$[\text{FS} : \text{Esub}^!] \frac{\Theta, x : \eta; \Gamma \Vdash N : \tau \quad \Theta; - \Vdash U : \epsilon \quad \eta \sim \epsilon}{\Theta; \Gamma \Vdash N \llbracket U/x^! \rrbracket : \tau}$$

By Lemma 14, we obtain the derivation $\Theta, x^! : \eta; \Gamma \Vdash N \{N_i/x[i]\}$, and the result follows from:

$$[\text{FS} : \text{Esub}^!] \frac{\Theta, x^! : \eta; \Gamma \Vdash N \{N_i/x[i]\} : \tau \quad \Theta; - \Vdash U : \epsilon \quad \eta \sim \epsilon}{\Theta; \Gamma \Vdash N \{N_i/x[i]\} \llbracket U/x^! \rrbracket : \tau}$$

5. Rule $[\text{RS} : \text{TCont}]$.

Then $M = C[N]$ and the reduction is as follows:

$$[\text{RS} : \text{TCont}] \frac{N \longrightarrow N'}{C[N] \longrightarrow C[N']}$$

with $M' = C[N']$. The proof proceeds by analysing the context C . There are three cases:

(a) $C = [\cdot] B$.

In this case $M = N B$, for some B . Since $\Gamma \vdash M : \tau$ one has a derivation:

$$[\text{FS} : \text{app}] \frac{\Theta; \Gamma \Vdash N : (\sigma^j, \eta) \rightarrow \tau \quad \Theta; \Delta \Vdash B : (\sigma^k, \epsilon) \quad \eta \sim \epsilon}{\Theta; \Gamma', \Delta \Vdash N B : \tau}$$

where $\Gamma = \Gamma', \Delta$. From $\Gamma \Vdash N : \sigma^j \rightarrow \tau$ and the reduction $N \longrightarrow N'$, one has by IH that $\Gamma' \Vdash N' : \sigma^j \rightarrow \tau$.

Finally, we may type the following:

$$[\text{FS} : \text{app}] \frac{\Theta; \Gamma' \Vdash N' : (\sigma^j, \eta) \rightarrow \tau \quad \Theta; \Delta \Vdash B : (\sigma^k, \epsilon) \quad \eta \sim \epsilon}{\Theta; \Gamma', \Delta \Vdash N' B : \tau}$$

Since $M' = (C[N']) = N' B$, the result follows.

(b) Cases $C = [\cdot] \langle N/x \rangle$ and $C = [\cdot] [\tilde{x} \leftarrow x]$ are similar to the previous.

6. Rule $[\text{RS} : \text{Fail}^\ell]$.

Then $M = N[x_1, \dots, x_k \leftarrow x] \langle C \star U/x \rangle$ where $k \neq \text{size}(C)$, $\tilde{y} = (\text{lfv}(N) \setminus \{\tilde{x}\}) \cup \text{lfv}(C)$ and the reduction is:

$$[\text{RS:Fail}^\ell] \frac{k \neq \text{size}(C) \quad \tilde{y} = (\text{lfv}(N) \setminus \{\tilde{x}\}) \cup \text{lfv}(C)}{N[x_1, \dots, x_k \leftarrow x] \llbracket C \star U/x \rrbracket \longrightarrow \text{fail}^{\tilde{y}}}$$

where $M' = \text{fail}^{\tilde{y}}$. Since $\Theta, x : \eta; \Gamma', x_1 : \sigma, \dots, x_k : \sigma \Vdash M$, one has a derivation:

$$\frac{[\text{FS:Esub}] \frac{\Theta, x : \eta; \Gamma', x_1 : \sigma, \dots, x_k : \sigma \Vdash N : \tau}{\Theta, x : \eta; \Gamma', x : \sigma^k \Vdash N[x_1, \dots, x_k \leftarrow x] : \tau} \quad \Theta; \Delta \Vdash C \star U : (\sigma^j, \epsilon) \quad \eta \sim \epsilon}{[\text{FS:Esub}] \frac{\Theta, x : \eta; \Gamma', x : \sigma^k \Vdash N[x_1, \dots, x_k \leftarrow x] : \tau \quad \Theta; \Delta \Vdash C \star U : (\sigma^j, \epsilon) \quad \eta \sim \epsilon}{\Theta; \Gamma', \Delta \Vdash N[x_1, \dots, x_k \leftarrow x] \llbracket C \star U/x \rrbracket : \tau}}$$

where $\Gamma = \Gamma', \Delta$. We may type the following:

$$[\text{FS:fail}] \frac{\text{dom}(\Gamma', \Delta) = \tilde{x}}{\Theta; \Gamma', \Delta \Vdash \text{fail}^{\tilde{y}} : \tau}$$

since Γ', Δ contain assignments on the free variables in M and B . Therefore, $\Theta; \Gamma \Vdash \text{fail}^{\tilde{y}} : \tau$, by applying $[\text{FS:sum}]$ as required.

7. Rule $[\text{RS:Fail}^!]$.

Then $M = N \llbracket U/x^! \rrbracket$ where $\text{head}(M) = x[i]$ and $U_i = 1^!$ and the reduction is:

$$[\text{RS:Fail}^!] \frac{\text{head}(N) = x[i] \quad U_i = 1^!}{N \llbracket U/x^! \rrbracket \longrightarrow N\{\text{fail}^0/x[i]\} \llbracket U/x^! \rrbracket}$$

with $M' = N\{\text{fail}^0/x[i]\} \llbracket U/x^! \rrbracket$. By the induction hypothesis, one has the derivation:

$$[\text{FS:Esub}^!] \frac{\Theta, x : \eta; \Gamma \Vdash N : \tau \quad \Theta; - \Vdash U : \epsilon \quad \eta \sim \epsilon}{\Theta; \Gamma \Vdash N \llbracket U/x^! \rrbracket : \tau}$$

By Lemma 14, there exists a derivation Π_1 of $\Theta, x^! : \eta; \Gamma' \Vdash N\{\text{fail}^0/x[i]\} : \tau$. Thus,

$$[\text{FS:Esub}^!] \frac{\Theta, x^! : \eta; \Gamma \Vdash N\{\text{fail}^0/x[i]\} : \tau \quad \Theta; - \Vdash U : \epsilon \quad \eta \sim \epsilon}{\Theta; \Gamma \Vdash N\{\text{fail}^0/x[i]\} \llbracket U/x^! \rrbracket : \tau}$$

8. Rule $[\text{RS:Cons}_1]$.

Then $M = \text{fail}^{\tilde{x}} B$ and the reduction is:

$$[\text{RS:Cons}_1] \frac{\tilde{y} = \text{lfv}(C)}{\text{fail}^{\tilde{x}} C \star U \longrightarrow \text{fail}^{\tilde{x} \cup \tilde{y}}}$$

and $M' = \text{fail}^{\tilde{x} \cup \tilde{y}}$. Since $\Theta; \Gamma \Vdash M : \tau$, one has the derivation:

$$\frac{[\text{F:fail}] \frac{\Theta; \Gamma \Vdash \text{fail}^{\tilde{x}} : (\sigma^j, \eta) \rightarrow \tau}{\Theta; \Gamma' \Vdash \text{fail}^{\tilde{x}} : (\sigma^j, \eta) \rightarrow \tau} \quad [\text{FS:bag}] \frac{\Theta; \Delta \Vdash C : \sigma^k \quad \Theta; - \Vdash U : \epsilon}{\Theta; \Delta \Vdash C \star U : (\sigma^k, \epsilon) \quad \eta \sim \epsilon}}{[\text{FS:app}] \frac{\Theta; \Gamma' \Vdash \text{fail}^{\tilde{x}} : (\sigma^j, \eta) \rightarrow \tau \quad \Theta; \Delta \Vdash C \star U : (\sigma^k, \epsilon) \quad \eta \sim \epsilon}{\Theta; \Gamma', \Delta \Vdash \text{fail}^{\tilde{x}} : C \star U : \tau}}$$

Hence $\Gamma = \Gamma', \Delta$ and we may type the following:

$$[\text{F:fail}] \frac{\Theta; \Gamma \Vdash \text{fail}^{\tilde{x} \cup \tilde{y}} : \tau}{\Theta; \Gamma \Vdash \text{fail}^{\tilde{x} \cup \tilde{y}} : \tau}$$

9. The proof for the cases of $[\text{RS:Cons}_2]$, $[\text{RS:Cons}_3]$ and $[\text{RS:Cons}_4]$ proceed similarly. \square

F.2 Subject Expansion

The full well-typed rules can be seen in Figure 18.

Lemma 15 (Anti-Substitution Lemma for λ_C).

1. (Linear) If $\Theta; \Gamma \vdash M\{N/x\} : \tau$ then there exists Γ', Δ and σ such that $\Theta; \Gamma', x : \sigma \vdash M : \tau$ and $\Theta; \Delta \vdash N : \sigma$ with $\Gamma = \Gamma', \Delta$.
2. (Unrestricted) If $\Theta, x^! : \eta; \Gamma \vdash M\{N/x[i]\}$ with $\eta_i = \sigma$ then $\Theta, x^! : \eta; \Gamma \vdash M : \tau$ and $\Theta; \cdot \vdash N : \sigma$.

Proof.

1. By structural induction on M with $\text{head}(M) = x$. There are six cases to be analyzed:

$\frac{[\text{TS:var}^\ell]}{\Theta; x : \sigma \vdash x : \sigma}$	$\frac{[\text{TS:var}^1]}{\Theta, x^1 : \eta; x : \eta_i, \Delta \vdash x : \sigma \quad \Theta, x^1 : \eta; \Delta \vdash x[i] : \sigma}$	$\frac{[\text{TS:1}^\ell]}{\Theta; - \vdash 1 : \omega}$	$\frac{[\text{TS:weak}]}{\Theta; \Gamma \vdash M : \tau \quad \Theta; \Gamma, x : \omega \vdash M[\leftarrow x] : \tau}$
$\frac{[\text{TS:abs-sh}]}{\Theta, x^1 : \eta; \Gamma, x : \sigma^k \vdash M[\tilde{x} \leftarrow x] : \tau \quad x \notin \text{dom}(\Gamma) \quad \Theta; \Gamma \vdash \lambda x. (M[\tilde{x} \leftarrow x]) : (\sigma^k, \eta) \rightarrow \tau}$	$\frac{[\text{TS:bag}^\ell]}{\Theta; \Gamma \vdash M : \sigma \quad \Theta; \Delta \vdash C : \sigma^k \quad \Theta; \Gamma, \Delta \vdash \lambda M \int \cdot C : \sigma^{k+1}}$		
$\frac{[\text{TS:app}]}{\Theta; \Gamma \vdash M : (\sigma^j, \eta) \rightarrow \tau \quad \Theta; \Delta \vdash B : (\sigma^j, \eta) \quad \Theta; \Gamma, \Delta \vdash M B : \tau}$	$\frac{[\text{TS:bag}^1]}{\Theta; - \vdash U : \epsilon \quad \Theta; - \vdash V : \eta \quad \Theta; - \vdash U \diamond V : \epsilon \diamond \eta}$		
$\frac{[\text{TS:shar}]}{\Theta; \Gamma, x_1 : \sigma, \dots, x_k : \sigma \vdash M : \tau \quad x \notin \text{dom}(\Gamma) \quad k \neq 0 \quad \Theta; \Gamma, x : \sigma^k \vdash M[x_1, \dots, x_k \leftarrow x] : \tau}$	$\frac{[\text{TS:bag}]}{\Theta; \Gamma \vdash C : \sigma^k \quad \Theta; - \vdash U : \eta \quad \Theta; \Gamma \vdash C * U : (\sigma^k, \eta)}$		
$\frac{[\text{TS:Esub}^\ell]}{\Theta; \Gamma, x_1 : \sigma, \dots, x_k : \sigma \vdash M : \tau \quad \Theta; \Delta \vdash C : \sigma^k \quad \Theta; \Gamma, \Delta \vdash M \langle C/x_1, \dots, x_k \rangle : \tau}$	$\frac{[\text{TS:Esub}^1]}{\Theta, x^1 : \eta; \Gamma \vdash M : \tau \quad \Theta; - \vdash U : \eta \quad \Theta; \Gamma \vdash M \llbracket U/x^1 \rrbracket : \tau}$		
$\frac{[\text{TS:Esub}]}{\Theta, x^1 : \eta; \Gamma, x : \sigma^j \vdash M[\tilde{x} \leftarrow x] : \tau \quad \Theta; \Delta \vdash B : (\sigma^k, \eta) \quad \Theta; \Gamma, \Delta \vdash (M[\tilde{x} \leftarrow x]) \langle \langle B/x \rangle \rangle : \tau}$			

Fig. 18. Well-Typed Rules for λ_C .

- (a) $M = x$.
(b) In this case, $x\{N/x\} = N$ and so $\Gamma \vdash N : \sigma$. Take $\Gamma' = -$ and $\Delta = \Gamma$ then $\Theta; x : \sigma \vdash x : \sigma$ and $\Delta \vdash N : \sigma$ by hypothesis, the result follows.
(c) $M = M' B$.
From Figure 13, $(M' B)\{N/x\} = (M'\{N/x\})B$. Let $\Gamma = \Gamma_1, \Gamma_2$ for some Γ_1, Γ_2 and the derivation is the following:

$$\frac{[\text{TS:app}]}{\Theta; \Gamma_1 \vdash M'\{N/x\} : (\delta^j, \eta) \rightarrow \tau \quad \Theta; \Gamma_2 \vdash B : (\delta^j, \eta) \quad \Theta; \Gamma_1, \Gamma_2 \vdash (M'\{N/x\})B : \tau}$$

where $j \geq 0$. By IH there exists $\Gamma'_1, \Delta, \sigma$ such that $\Gamma_1 = \Gamma'_1, \Delta$ with $\Theta; \Gamma'_1, x : \sigma \vdash M' : (\delta^j, \eta) \rightarrow \tau$ and $\Theta; \Delta \vdash N : \sigma$. Which gives the derivation:

$$\frac{[\text{TS:app}]}{\Theta; \Gamma'_1, x : \sigma \vdash M' : (\delta^j, \eta) \rightarrow \tau \quad \Theta; \Gamma_2 \vdash B : (\delta^j, \eta) \quad \Theta; \Gamma'_1, \Gamma_2, x : \sigma \vdash M' B : \tau}$$

By taking $\Gamma' = \Gamma'_1, \Gamma_2$ the result follows.

- (d) $M = M'[\tilde{y} \leftarrow y]$ with $y \neq x$.
Then from Figure 13, $M'[\tilde{y} \leftarrow y]\{N/x\} = M'\{N/x\}[\tilde{y} \leftarrow y]$. Let $\Gamma = \Gamma_1, y : \delta^k$ with $k \neq 0$. Therefore,

$$\frac{[\text{TS:shar}]}{\Theta; \Gamma_1, y_1 : \delta, \dots, y_k : \delta \vdash M'\{N/x\} : \tau \quad y \notin \Gamma_1 \quad k \neq 0 \quad \Theta; \Gamma_1, y : \delta^k \vdash M'\{N/x\}[\tilde{y} \leftarrow y] : \tau}$$

By IH there exists $\Gamma'_1, \Delta, \sigma$ such that $\Gamma_1, y_1 : \delta, \dots, y_k : \delta = \Gamma'_1, y_1 : \delta, \dots, y_k : \delta, \Delta$ with $\Theta; \Gamma'_1, y_1 : \delta, \dots, y_k : \delta, x : \sigma \vdash M' : \tau$ and $\Theta; \Delta \vdash N : \sigma$. Which gives the derivation:

$$\frac{[\text{TS:share}]}{\Theta; \Gamma'_1, y_1 : \delta, \dots, y_k : \delta, x : \sigma \vdash M' : \tau \quad y \notin \Gamma_1 \quad k \neq 0 \quad \Theta; \Gamma'_1, y : \delta^k, x : \sigma \vdash M'[y_1, \dots, y_k \leftarrow x] : \tau}$$

By taking $\Gamma' = \Gamma'_1, y : \delta^k$ the result follows.

- (e) $M = M'[\leftarrow y]$ with $x \neq y$.
Then $M'[\leftarrow y]\{N/x\} = M'\{N/x\}[\leftarrow y]$. Therefore,

$$\frac{[\text{TS:weak}]}{\Theta; \Gamma \vdash M\{N/x\} : \tau \quad \Theta; \Gamma, y : \omega \vdash M\{N/x\}[\leftarrow y] : \tau}$$

By IH there exists Γ', Δ, σ such that $\Gamma = \Gamma', \Delta$ with $\Theta; \Gamma', x : \sigma \vdash M' : \tau$ and $\Theta; \Delta \vdash N : \sigma$. Which gives the derivation:

$$\frac{[\text{TS:weak}]}{\Theta; \Gamma', x : \sigma \vdash M : \tau \quad \Theta; \Gamma', y : \omega, x : \sigma \vdash M[\leftarrow y] : \tau}$$

By taking $\Gamma' = \Gamma', y : \omega$ the result follows.

- (f) If $M = M' \langle C/y_1, \dots, y_k \rangle$ with $x \neq y_1, \dots, y_k$.

Then $M' \langle C/y_1, \dots, y_k \rangle \{N/x\} = M' \{N/x\} \langle C/y_1, \dots, y_k \rangle$ and

$$[\text{TS:Esub}^\ell] \frac{\Theta; \Gamma_1, y_1 : \delta, \dots, y_k : \delta \vdash M' \{N/x\} : \tau \quad \Theta; \Gamma_2 \vdash C : \delta^k}{\Theta; \Gamma_1, \Gamma_2 \vdash M' \{N/x\} \langle C/y_1, \dots, y_k \rangle : \tau}$$

By IH there exists $\Gamma'_1, \Delta, \sigma$ such that $\Gamma_1, y_1 : \delta, \dots, y_k : \delta = \Gamma'_1, y_1 : \delta, \dots, y_k : \delta, \Delta$ with $\Theta; \Gamma'_1, y_1 : \delta, \dots, y_k : \delta, x : \sigma \vdash M' : \tau$ and $\Theta; \Delta \vdash N : \sigma$. Which gives the derivation:

$$[\text{TS:Esub}^\ell] \frac{\Theta; \Gamma'_1, y_1 : \delta, \dots, y_k : \delta, x : \sigma \vdash M' : \tau \quad \Theta; \Gamma_2 \vdash C : \delta^k}{\Theta; \Gamma'_1, \Gamma_2, x : \sigma \vdash M' \langle C/y_1, \dots, y_k \rangle : \tau}$$

By taking $\Gamma' = \Gamma'_1, \Gamma_2$ the result follows.

- (g) If $M = M' \llbracket U/y \rrbracket$ then $\text{head}(M' \llbracket U/y \rrbracket) = \text{head}(M') = x$, and the proofs is similar to the case above.

2. By structural induction on M with $\text{head}(M) = x[i]$. There are three cases to be analyzed:

- (a) $M = x[i]$.

Observe that $x[i] \{N/x[i]\} = N$ and let $\Theta, x^1 : \eta; \Gamma \vdash x[i] \{N/x[i]\} : \sigma$ with $\eta_i = \sigma$ In this case $\Gamma = \cdot$, and we have both

$$\frac{[\text{TS:var}^\ell] \frac{}{\Theta, x^1 : \eta; x : \eta_i \vdash x : \sigma}}{[\text{TS:var}^1] \frac{}{\Theta, x^1 : \eta; \cdot \vdash x[i] : \sigma}}$$

and $\Theta, x^1 : \eta; \Gamma \vdash N : \sigma$ and the result follows.

- (b) $M = M' B$.

From Figure 13, $M' \langle B/y \rangle \{N/x[i]\} = M' \{N/x[i]\} \langle B/y \rangle$, let $\Theta, x^1 : \eta; \Gamma_1, \Gamma_2 \vdash (M' \{N/x[i]\})B : \tau$ where $\Gamma = \Gamma_1, \Gamma_2$ and $\eta_i = \sigma$ we derive:

$$[\text{TS:app}] \frac{\Theta, x^1 : \eta; \Gamma_1 \vdash M' \{N/x[i]\} : (\delta^j, \epsilon) \rightarrow \tau \quad \Theta, x^1 : \eta; \Gamma_2 \vdash B : (\delta^j, \epsilon)}{\Theta, x^1 : \eta; \Gamma_1, \Gamma_2 \vdash (M' \{N/x[i]\})B : \tau}$$

for some strict type δ . By IH $\Theta, x^1 : \eta; \Gamma_1 \vdash M' : (\delta^j, \epsilon) \rightarrow \tau$ and $\Theta; \cdot \vdash N : \sigma$ and we derive:

$$[\text{TS:app}] \frac{\Theta, x^1 : \eta; \Gamma_1 \vdash M' : (\delta^j, \epsilon) \rightarrow \tau \quad \Theta, x^1 : \sigma; \Gamma_2 \vdash B : (\delta^j, \epsilon')}{\Theta, x^1 : \eta; \Gamma_1, \Gamma_2 \vdash M' B : \tau}$$

and the result follows.

- (c) $M = M' [\tilde{y} \leftarrow y]$.

From Figure 13 $M' [\tilde{y} \leftarrow y] \{N/x[i]\} = M' \{N/x[i]\} [\tilde{y} \leftarrow y]$, let $\Theta, x^1 : \eta; \Gamma_1, y : \delta^k \vdash M' \{N/x[i]\} [\tilde{y} \leftarrow y] : \tau$ with $\Gamma = \Gamma_1, y : \delta^k$ and $\eta_i = \sigma$, then we derive:

$$[\text{TS:shar}] \frac{\Theta, x^1 : \eta; \Gamma_1, y_1 : \delta, \dots, y_k : \delta \vdash M' \{N/x[i]\} : \tau \quad y \notin \Gamma_1 \quad k \neq 0}{\Theta, x^1 : \eta; \Gamma_1, y : \delta^k \vdash M' \{N/x[i]\} [\tilde{y} \leftarrow y] : \tau}$$

By IH $\Theta, x^1 : \eta; \Gamma_1, y_1 : \delta, \dots, y_k : \delta \vdash M' : \tau$ and $\Theta; \cdot \vdash N : \sigma$ and we derive:

$$[\text{TS:shar}] \frac{\Theta, x^1 : \eta; \Gamma_1, y_1 : \delta, \dots, y_k : \delta \vdash M' : \tau \quad y \notin \Gamma_1 \quad k \neq 0}{\Theta, x^1 : \eta; \Gamma_1, y : \delta^k \vdash M' [y_1, \dots, y_k \leftarrow y] : \tau}$$

and the result follows.

- (d) $M = M' [\leftarrow y]$.

From Figure 13 $M' [\leftarrow y] \{N/x[i]\} = M' \{N/x[i]\} [\leftarrow y]$, let $\Theta, x^1 : \eta; \Gamma, y : \omega \vdash M' \{N/x[i]\} [\leftarrow y] : \tau$ with $\Gamma = \Gamma, y : \omega$ and $\eta_i = \sigma$, then we derive:

$$[\text{TS:weak}] \frac{\Theta, x^1 : \eta; \Gamma \vdash M' \{N/x[i]\} : \tau}{\Theta, x^1 : \eta; \Gamma, y : \omega \vdash M' \{N/x[i]\} [\leftarrow y] : \tau}$$

By IH $\Theta, x^1 : \eta; \Gamma \vdash M' : \tau$ and $\Theta; \cdot \vdash N : \sigma$ and we derive:

$$[\text{TS:weak}] \frac{\Theta, x^1 : \eta; \Gamma \vdash M : \tau}{\Theta, x^1 : \eta; \Gamma, y : \omega \vdash M [\leftarrow y] : \tau}$$

and the result follows.

- (e) $M = M' \langle C/y_1, \dots, y_k \rangle$.

From Figure 13 $M' \langle C/y_1, \dots, y_k \rangle \{N/x[i]\} = M' \{N/x[i]\} \langle C/y_1, \dots, y_k \rangle$, let $\Theta, x^1 : \eta; \Gamma', \Delta \vdash M' \{N/x[i]\} \langle C/y_1, \dots, y_k \rangle : \tau$ with $\Gamma = \Gamma', \Delta$ and $\eta_i = \sigma$, then we derive:

$$[\text{TS:Esub}^\ell] \frac{\Theta, x^1 : \eta; \Gamma', y_1 : \delta, \dots, y_k : \delta \vdash M' \{N/x[i]\} : \tau \quad \Theta, x^1 : \eta; \Delta \vdash C : \delta^k}{\Theta, x^1 : \eta; \Gamma', \Delta \vdash M' \{N/x[i]\} \langle C/y_1, \dots, y_k \rangle : \tau}$$

By IH $\Theta, x^1 : \eta; \Gamma \vdash M' : \tau$ and $\Theta; \cdot \vdash N : \sigma$ and we derive:

$$[\text{TS:Esub}^\ell] \frac{\Theta, x^1 : \eta; \Gamma', y_1 : \delta, \dots, y_k : \delta \vdash M : \tau \quad \Theta, x^1 : \eta; \Delta \vdash C : \delta^k}{\Theta, x^1 : \eta; \Gamma', \Delta \vdash M \langle C/y_1, \dots, y_k \rangle : \tau}$$

and the result follows.

(f) $M = M' \llbracket U/y^1 \rrbracket$.

From Figure 13 $M' \llbracket U/y^1 \rrbracket \{N/x[i]\} = M' \{N/x[i]\} \llbracket U/y^1 \rrbracket$, let $\Theta; \Gamma \vdash M' \{N/x[i]\} \llbracket U/y^1 \rrbracket : \tau$ and $\eta_i = \sigma$, then we derive:

$$[\text{TS:Esub}^1] \frac{\Theta, x^1 : \eta, y : \epsilon; \Gamma \vdash M' \{N/x[i]\} : \tau \quad \Theta, x^1 : \eta; - \vdash U : \epsilon}{\Theta; \Gamma \vdash M' \{N/x[i]\} \llbracket U/y^1 \rrbracket : \tau}$$

By IH $\Theta, x^1 : \eta, y : \epsilon; \Gamma \vdash M' : \tau$ and $\Theta; - \vdash N : \sigma$ and we derive:

$$[\text{TS:Esub}^1] \frac{\Theta, x^1 : \eta, y : \epsilon; \Gamma \vdash M' : \tau \quad \Theta, x^1 : \eta; - \vdash U : \epsilon}{\Theta; \Gamma \vdash M' \llbracket U/y^1 \rrbracket : \tau}$$

and the result follows. \square

Theorem 18 (Subject Expansion in λ_C). *If $\Theta; \Gamma \vdash M' : \tau$ and $M \longrightarrow M'$ then $\Theta; \Gamma \vdash M : \tau$.*

Proof. By structural induction on the reduction rule from Fig. 13 applied in $M \longrightarrow M'$.

1. **Rule [RS:Beta].**

Then $M' = N[\tilde{x} \leftarrow x] \langle\langle B/x \rangle\rangle$ and the reduction is:

$$[\text{RS:Beta}] \frac{}{(\lambda x. N[\tilde{x} \leftarrow x])B \longrightarrow N[\tilde{x} \leftarrow x] \langle\langle B/x \rangle\rangle}$$

where $M = (\lambda x. N[\tilde{x} \leftarrow x])B$. Since $\Theta; \Gamma \vdash M' : \tau$ we get the following derivation:

$$[\text{TS:share}] \frac{\Theta, x^1 : \eta; \Gamma', x_1 : \sigma, \dots, x_j : \sigma \vdash N : \tau}{\Theta, x^1 : \eta; \Gamma', x : \sigma^j \vdash N[\tilde{x} \leftarrow x] : \tau} \quad \Theta; \Delta \vdash B : (\sigma^j, \eta)$$

$$[\text{TS:Esub}] \frac{}{\Theta; \Gamma', \Delta \vdash N[\tilde{x} \leftarrow x] \langle\langle B/x \rangle\rangle : \tau}$$

for $\Gamma = \Gamma', \Delta$. Notice that:

$$[\text{TS:share}] \frac{\Theta, x^1 : \eta; \Gamma', x_1 : \sigma, \dots, x_j : \sigma \vdash N : \tau}{\Theta, x^1 : \eta; \Gamma', x : \sigma^j \vdash N[\tilde{x} \leftarrow x] : \tau}$$

$$[\text{TS:abs-sh}] \frac{}{\Theta; \Gamma' \vdash \lambda x. N[\tilde{x} \leftarrow x] : (\sigma^j, \eta) \rightarrow \tau} \quad \Theta; \Delta \vdash B : (\sigma^j, \eta)$$

$$[\text{TS:app}] \frac{}{\Theta; \Gamma', \Delta \vdash (\lambda x. N[\tilde{x} \leftarrow x])B : \tau}$$

Therefore $\Theta; \Gamma', \Delta \vdash (\lambda x. N[\tilde{x} \leftarrow x])B : \tau$ and the result follows.

Then $M = (\lambda x. N[\tilde{x} \leftarrow x])B$ and the reduction is:

2. **Rule [RS:Ex-Sub].**

Then $M' = N \langle C/x_1, \dots, x_k \rangle \llbracket U/x^1 \rrbracket$ where $C = \{N_1\} \dots \{N_k\}$. The reduction is:

$$[\text{RS:Ex-Sub}] \frac{C = \{N_1\} \dots \{N_k\} \quad N \neq \text{fail}^{\tilde{y}}}{N[x_1, \dots, x_k \leftarrow x] \langle\langle C \star U/x \rangle\rangle \longrightarrow N \langle C/x_1, \dots, x_k \rangle \llbracket U/x^1 \rrbracket}$$

and $M = N[x_1, \dots, x_k \leftarrow x] \langle\langle C \star U/x \rangle\rangle$. To simplify the proof we take $k = 2$, as the case $k > 2$ is similar. Therefore $C = \{N_1\} \cdot \{N_2\}$.

Since $\Theta; \Gamma \vdash M' : \tau$ we get a derivation (we omit the labels [TS : Esub¹] and [TS : Esub^ℓ]):

$$\frac{\frac{\Theta, x^1 : \eta; \Gamma', x_1 : \sigma, x_2 : \sigma \vdash N : \tau \quad \Theta; \Delta \vdash C : \sigma^2}{\Theta, x^1 : \eta; \Gamma', \Delta \vdash N \langle C/x_1, x_2 \rangle : \tau} \quad \Theta; - \vdash U : \eta}{\Theta; \Gamma', \Delta \vdash N \langle C/x_1, x_2 \rangle \llbracket U/x^1 \rrbracket : \tau}$$

where $\Gamma = \Gamma', \Delta$. Consider the typing derivation (we omit the labels [TS : Esub] and [TS:share]):

$$\frac{\frac{\Theta, x^1 : \eta; \Gamma', x_1 : \sigma, x_2 : \sigma \vdash N : \tau}{\Theta, x^1 : \eta; \Gamma', x : \sigma^2 \vdash N[x_1, x_2 \leftarrow x] : \tau} \quad \Theta; \Delta \vdash C \star U : (\sigma^2, \eta)}{\Theta; \Gamma', \Delta \vdash N[x_1, x_2 \leftarrow x] \langle\langle C \star U/x \rangle\rangle : \tau}$$

and the result follows.

3. Rule $[\text{RS:Fetch}^\ell]$.

Then $M' = N\{C_i/x_j\}\langle(C \setminus C_i)/\tilde{x}\rangle$ where $\text{head}(N) = x_j$. The reduction is:

$$[\text{RS:Fetch}_i^\ell] \frac{\text{head}(N) = x_j}{N\langle C/\tilde{x}, x_j \rangle \longrightarrow N\{C_i/x_j\}\langle(C \setminus C_i)/\tilde{x}\rangle}$$

and $M = N\langle C/\tilde{x}, x_j \rangle$. Since $\Theta; \Gamma \vdash M' : \tau$ we get the following derivation:

$$[\text{TS:Esub}^\ell] \frac{\Theta; \Gamma', \Delta_i, \tilde{x} : \sigma^{k-1} \vdash N\{C_i/x_j\} : \tau \quad \Theta; \Delta'_i \vdash C \setminus C_i : \sigma^{k-1}}{\Theta; \Gamma', \Delta \vdash N\{C_i/x_j\}\langle(C \setminus C_i)/\tilde{x}\rangle : \tau}$$

where $\Gamma = \Gamma', \Delta$ and $\Delta = \Delta_i, \Delta'_i$. By Lemma 15, we obtain the derivation $\Theta; \Gamma', \tilde{x} : \sigma^{k-1}, x_j : \sigma \vdash N : \tau$ and $\Theta; \Delta_i \vdash C_i : \sigma$ via:

$$[\text{TS:Esub}^\ell] \frac{\Theta; \Gamma', \tilde{x} : \sigma^{k-1}, x_j : \sigma \vdash N : \tau \quad \Theta; \Delta \vdash C : \sigma^k}{\Theta; \Gamma', \Delta \vdash N\langle C/\tilde{x}, x_j \rangle : \tau}$$

4. Rule $[\text{RS:Fetch}^!]$.

Then $M' = N\{N_i/x[i]\}\llbracket U/x^! \rrbracket$ where $\text{head}(M) = x[i]$. The reduction is:

$$[\text{RS:Fetch}^!] \frac{\text{head}(N) = x[i] \quad U_i = \lambda N_i \{ \}^!}{N\llbracket U/x^! \rrbracket \longrightarrow N\{N_i/x[i]\}\llbracket U/x^! \rrbracket}$$

and $M = N\llbracket U/x^! \rrbracket$. Since $\Theta; \Gamma \vdash M' : \tau$ we get the following derivation:

$$[\text{TS:Esub}^!] \frac{\Theta, x^! : \eta; \Gamma \vdash N\{N_i/x[i]\} : \tau \quad \Theta; - \vdash U : \eta}{\Theta; \Gamma \vdash N\{N_i/x[i]\}\llbracket U/x^! \rrbracket : \tau}$$

By Lemma 15, we obtain the derivation $\Theta, x : \eta; \Gamma \vdash N : \tau$, and the result follows from:

$$[\text{TS:Esub}^!] \frac{\Theta, x : \eta; \Gamma \vdash N : \tau \quad \Theta; - \vdash U : \eta}{\Theta; \Gamma \vdash N\llbracket U/x^! \rrbracket : \tau}$$

5. Rule $[\text{RS:TCont}]$.

Then $M' = C[N']$ and the reduction is as follows:

$$[\text{RS:TCont}] \frac{N \longrightarrow N'}{C[N] \longrightarrow C[N']}$$

with $M = C[N]$. The proof proceeds by analysing the context C . There are three cases:

(a) $C = [\cdot] B$.

In this case $M' = N' B$, for some B . Since $\Gamma \vdash M' : \tau$ one has a derivation:

$$[\text{TS:app}] \frac{\Theta; \Gamma' \vdash N' : (\sigma^j, \eta) \rightarrow \tau \quad \Theta; \Delta \vdash B : (\sigma^j, \eta)}{\Theta; \Gamma', \Delta \vdash N' B : \tau}$$

where $\Gamma = \Gamma', \Delta$. From $\Gamma' \vdash N' : (\sigma^j, \eta) \rightarrow \tau$ and the reduction $N \longrightarrow N'$, one has by IH that $\Gamma' \vdash N : (\sigma^j, \eta) \rightarrow \tau$.

Finally, we may type the following:

$$[\text{TS:app}] \frac{\Theta; \Gamma' \vdash N : (\sigma^j, \eta) \rightarrow \tau \quad \Theta; \Delta \vdash B : (\sigma^j, \eta)}{\Theta; \Gamma', \Delta \vdash N B : \tau}$$

Since $M = (C[N]) = NB$, the result follows.

(b) Cases $C = [\cdot]\langle N/x \rangle$ and $C = [\cdot][\tilde{x} \leftarrow x]$ are similar to the previous.6. Rules $[\text{RS:Fail}^\ell]$, $[\text{RS:Fail}^!]$ and $[\text{RS:Cons}_1]$. These cases are trivial, since 1 , $1^!$ and $\text{fail}^{\tilde{x}\cup\tilde{y}}$ are not well-typed. \square

G Proof of Tight Correctness of the Translation under the Lazy Semantics

Here we prove Theorem 6.

Definition 17 (Success). We define $\text{head}(\check{\lambda}) = \check{\lambda}$ and $\llbracket \check{\lambda} \rrbracket_u = \check{\pi}$. We define success for terms $M \in \lambda_{\mathcal{C}}$ and $P \in \mathfrak{S}\pi^+$.

- $M \Downarrow \check{\lambda}$ if and only if, there exist $M'_1, \dots, M'_k \in \lambda_{\mathcal{C}}$ such that $M \longrightarrow^* M'$ and $\text{head}(M') = \check{\lambda}$.
- $P \Downarrow \check{\pi}$ if and only if, there exist $Q_1, Q_2 \in \mathfrak{S}\pi^+$ such that $P \longrightarrow^* (Q_1 \mid \check{\pi}) \uplus Q_2$.
- $P \Downarrow \check{\pi}$ if and only if, there exist S and $Q_1, Q_2 \in \mathfrak{S}\pi^+$ such that $P \rightsquigarrow_S^* (Q_1 \mid \check{\pi}) \uplus Q_2$.

G.1 Type Preservation

Lemma 16. $\llbracket \sigma^j \rrbracket_{(\tau_1, m)} = \llbracket \sigma^k \rrbracket_{(\tau_2, n)}$ and $\llbracket (\sigma^j, \eta) \rrbracket_{(\tau_1, m)} = \llbracket (\sigma^k, \eta) \rrbracket_{(\tau_2, n)}$ hold, provided that τ_1, τ_2, n and m are as follows:

1. If $j > k$ then take τ_1 to be an arbitrary type, $m = 0$, take τ_2 to be σ and $n = j - k$.
2. If $j < k$ then take τ_1 to be σ , $m = k - j$, take τ_2 to be an arbitrary type and $n = 0$.
3. Otherwise, if $j = k$ then take $m = n = 0$. In this case, τ_1, τ_2 are unimportant.

Proof. This proof proceeds by analyzing the conditions on types, following Paulus et al. [21]. □

Lemma 17. If $\eta \sim \epsilon$ then the following hold:

1. If $\llbracket M \rrbracket_u \vdash \llbracket \Gamma \rrbracket, \llbracket \Theta \rrbracket, x^! : \overline{\llbracket \eta \rrbracket}$ then $\llbracket M \rrbracket_u \vdash \llbracket \Gamma \rrbracket, \llbracket \Theta \rrbracket, x^! : \overline{\llbracket \epsilon \rrbracket}$.
2. If $\llbracket M \rrbracket_u \vdash \llbracket \Gamma \rrbracket, u : \llbracket (\sigma^j, \eta) \rightarrow \tau \rrbracket, \llbracket \Theta \rrbracket$ then $\llbracket M \rrbracket_u \vdash \llbracket \Gamma \rrbracket, u : \llbracket (\sigma^j, \epsilon) \rightarrow \tau \rrbracket, \llbracket \Theta \rrbracket$.

Proof. The proof is by mutual induction on the the derivations of If $\llbracket M \rrbracket_u \vdash \llbracket \Gamma \rrbracket, \llbracket \Theta \rrbracket, x^! : \overline{\llbracket \eta \rrbracket}$ and $\llbracket M \rrbracket_u \vdash \llbracket \Gamma \rrbracket, u : \llbracket (\sigma^j, \eta) \rightarrow \tau \rrbracket, \llbracket \Theta \rrbracket$ and on the structure of M .

We use Def. 12 that establishes $\llbracket \eta \rrbracket = !\&_{\eta_i \in \eta} \{i : \llbracket \eta_i \rrbracket\}$ and by duality $\overline{\llbracket \eta \rrbracket} = ? \oplus_{\eta_i \in \eta} \{i : \overline{\llbracket \eta_i \rrbracket}\}$.

1. $M = x$.

By the translation in Fig. 16: $\llbracket x \rrbracket_u = \overline{x}.\text{some}; [x \leftrightarrow u]$. We have the following derivation, for some type A :

$$\frac{\frac{\frac{[\text{TID}]}{[x \leftrightarrow u] \vdash x : \overline{A}, u : A}}{[\text{TWEAKEN}]} \frac{[x \leftrightarrow u] \vdash x : \overline{A}, u : A, x^! : \overline{\llbracket \eta \rrbracket}}{[\text{T\&}]}}{\overline{x}.\text{some}; [x \leftrightarrow u] \vdash x : \&\overline{A}, u : A, x^! : \overline{\llbracket \eta \rrbracket}}$$

The derivation is independent of $x^! : \overline{\llbracket \eta \rrbracket}$, hence the result trivially holds for $\llbracket \Gamma \rrbracket, \llbracket \Theta \rrbracket = x : \&\overline{A}, u : A$.

2. $M = x[k]$.

By the translation in Fig. 16: $\llbracket x[k] \rrbracket_u = ?x^! [x_i]; \overline{x_i}.k; [x_i \leftrightarrow u]$. We have the following derivation:

$$\frac{\frac{\frac{[\text{TID}]}{[x_i \leftrightarrow u] \vdash u : \llbracket \tau \rrbracket, x_i : \overline{\llbracket \eta_k \rrbracket}}{[\text{T\oplus}]} \frac{\overline{x_i}.k; [x_i \leftrightarrow u] \vdash u : \llbracket \tau \rrbracket, x_i : \oplus_{\eta_i \in \eta} \{i : \llbracket \eta_i \rrbracket\}}{[\text{T?}]}}{?x^! [x_i]; \overline{x_i}.k; [x_i \leftrightarrow u] \vdash u : \llbracket \tau \rrbracket, x^! : ? \oplus_{\eta_i \in \eta} \{i : \overline{\llbracket \eta_i \rrbracket}\}}{,}$$

Since $\eta \sim \epsilon$ we have that $\epsilon_k = \eta_k$ for each $k = 1, \dots, |\eta|$. Thus, the same derivation above, replacing η_i 's for ϵ_i 's entails $?x^! [x_i]; \overline{x_i}.k; [x_i \leftrightarrow u] \vdash u : \llbracket \tau \rrbracket; x^! : \overline{\llbracket \epsilon \rrbracket}$, and the result follows. For the case of $M = y[k]$ with $y \neq x$ we use the argument that the typing of y is independent on x .

3. $M = \lambda y.(M'[\tilde{y} \leftarrow y])$.

From the translation in Fig. 16:

$$\llbracket \lambda y.M'[\tilde{y} \leftarrow y] \rrbracket_u = \overline{u}.\text{some}; u(y); \underbrace{\overline{y}.\text{some}; y(y^\ell); y(y^!); y(); \llbracket M'[\tilde{y} \leftarrow y] \rrbracket_u}_P$$

$$\begin{array}{c} \Pi_1 \\ \vdots \\ [\text{T}\perp] \frac{\llbracket M'[\tilde{y} \leftarrow y] \rrbracket_u \vdash u : \llbracket \tau \rrbracket, \llbracket \Gamma' \rrbracket, y^\ell : \overline{\llbracket \sigma^k \rrbracket}_{(\sigma, i)}, \llbracket \Theta \rrbracket, y^! : \overline{\llbracket \eta \rrbracket}}{y(); \llbracket M'[\tilde{y} \leftarrow y] \rrbracket_u \vdash y : \perp, u : \llbracket \tau \rrbracket, \llbracket \Gamma' \rrbracket, y^\ell : \overline{\llbracket \sigma^k \rrbracket}_{(\sigma, i)}, \llbracket \Theta \rrbracket, y^! : \overline{\llbracket \eta \rrbracket}} \\ [\text{T}\&] \frac{y(y^!); y(); \llbracket M'[\tilde{y} \leftarrow y] \rrbracket_u \vdash y : (\overline{\llbracket \eta \rrbracket}) \wp (\perp), u : \llbracket \tau \rrbracket, \llbracket \Gamma' \rrbracket, y^\ell : \overline{\llbracket \sigma^k \rrbracket}_{(\sigma, i)}, \llbracket \Theta \rrbracket}}{y(y^\ell); y(y^!); y(); \llbracket M'[\tilde{y} \leftarrow y] \rrbracket_u \vdash y : \overline{\llbracket \sigma^k \rrbracket}_{(\sigma, i)} \wp ((\overline{\llbracket \eta \rrbracket}) \wp (\perp)), u : \llbracket \tau \rrbracket, \llbracket \Gamma' \rrbracket, \llbracket \Theta \rrbracket} \\ [\text{T}\&\text{some}] \frac{\overline{y}.\text{some}; P \vdash y : \&(\overline{\llbracket \sigma^k \rrbracket}_{(\sigma, i)}) \wp ((\overline{\llbracket \eta \rrbracket}) \wp (\perp)), u : \llbracket \tau \rrbracket, \llbracket \Gamma' \rrbracket, \llbracket \Theta \rrbracket}}{u(y); \overline{y}.\text{some}; P \vdash u : \&(\&(\overline{\llbracket \sigma^k \rrbracket}_{(\sigma, i)}) \wp ((\overline{\llbracket \eta \rrbracket}) \wp (\perp))) \wp \llbracket \tau \rrbracket, \llbracket \Gamma' \rrbracket, \llbracket \Theta \rrbracket} \\ [\text{T}\&\text{some}] \frac{\overline{u}.\text{some}; u(y); \overline{y}.\text{some}; P \vdash u : \&(\&(\overline{\llbracket \sigma^k \rrbracket}_{(\sigma, i)}) \wp ((\overline{\llbracket \eta \rrbracket}) \wp (\perp))) \wp \llbracket \tau \rrbracket, \llbracket \Gamma' \rrbracket, \llbracket \Theta \rrbracket}}{\underbrace{\&(\&(\overline{\llbracket \sigma^k \rrbracket}_{(\sigma, i)}) \wp ((\overline{\llbracket \eta \rrbracket}) \wp (\perp))) \wp \llbracket \tau \rrbracket, \llbracket \Gamma' \rrbracket, \llbracket \Theta \rrbracket}}_{\llbracket (\sigma^k, \eta) \rightarrow \tau \rrbracket} \quad \text{Def. 12}} \end{array}$$

Let us consider the following two cases:

(b) **For** $C = \lambda M' \}. C'$:

$$[\text{FS:bag}^\ell] \frac{\Theta; \Gamma' \vDash M' : \sigma \quad \Theta; \Delta \vDash C' : \sigma^k}{\Theta; \Gamma' \wedge \Delta \vDash \lambda M' \}. C' : \sigma^k}$$

Where $\Gamma = \Gamma' \wedge \Delta$. To simplify the proof, we will consider $k = 3$.

By IH we have $\llbracket M' \rrbracket_{x_i} \vdash \llbracket \Gamma' \rrbracket, x_i : \llbracket \sigma \rrbracket; \llbracket \Theta \rrbracket$ and $\llbracket C' \rrbracket_{x^\ell} \vdash \llbracket \Delta \rrbracket, x^\ell : \llbracket \sigma \wedge \sigma \rrbracket_{(\tau,j)}; \llbracket \Theta \rrbracket$.

Our translation Fig. 16 gives:

$$\begin{aligned} \llbracket C \rrbracket_{x^\ell} &= x^\ell.\text{some}_{\text{fv}(C)}; x^\ell(y_i); x^\ell.\text{some}_{y_i, \text{fv}(C)}; \overline{x^\ell}.\text{some}; \overline{x^\ell}[x_i]. \\ &\quad (x_i.\text{some}_{\text{fv}(M')}; \llbracket M' \rrbracket_{x_i} \mid \llbracket (C') \rrbracket_{x^\ell} \mid \overline{y_i}.\text{none}) \end{aligned} \quad (12)$$

Let Π_1 be the derivation:

$$\frac{\frac{\frac{\llbracket M' \rrbracket_{x_i} \vdash \llbracket \Gamma' \rrbracket, x_i : \llbracket \sigma \rrbracket, \llbracket \Theta \rrbracket}{x_i.\text{some}_{\text{fv}(M')}; \llbracket M' \rrbracket_{x_i} \vdash \llbracket \Gamma' \rrbracket, x_i : \oplus \llbracket \sigma \rrbracket, \llbracket \Theta \rrbracket}}{x_i.\text{some}_{\text{fv}(M')}; \llbracket M' \rrbracket_{x_i} \mid \overline{y_i}.\text{none} \vdash \llbracket \Gamma' \rrbracket, x_i : \oplus \llbracket \sigma \rrbracket, y_i : \& \mathbf{1}, \llbracket \Theta \rrbracket}}{\overline{y_i}.\text{none} \vdash y_i : \& \mathbf{1}}}{\overline{y_i}.\text{none} \vdash y_i : \& \mathbf{1}, \llbracket \Theta \rrbracket}}$$

Let $P = \llbracket M' \rrbracket_{x_i} \mid \llbracket C' \rrbracket_{x^\ell} \mid \overline{y_i}.\text{none}$, in the derivation Π_2 below:

$$\begin{aligned} & \frac{\Pi_1 \quad \llbracket C' \rrbracket_{x^\ell} \vdash \llbracket \Delta \rrbracket, x^\ell : \llbracket \sigma \wedge \sigma \rrbracket_{(\tau,j)}, \llbracket \Theta \rrbracket}{x^\ell[x_i].P \vdash \llbracket \Gamma' \rrbracket, \llbracket \Delta \rrbracket, y_i : \& \mathbf{1}, x^\ell : (\oplus \llbracket \sigma \rrbracket) \otimes (\llbracket \sigma \wedge \sigma \rrbracket_{(\tau,j)}), \llbracket \Theta \rrbracket}}{[\text{T}\otimes]} \\ & \frac{[\text{T}\&\text{some}]}{\underbrace{x^\ell.\text{some}; \overline{x^\ell}[x_i].P}_{P_2} \vdash \llbracket \Gamma' \rrbracket, \llbracket \Delta \rrbracket, y_i : \& \mathbf{1}, x^\ell : \&((\oplus \llbracket \sigma \rrbracket) \otimes (\llbracket \sigma \wedge \sigma \rrbracket_{(\tau,j)})), \llbracket \Theta \rrbracket}} \end{aligned}$$

Let $P_2 = \overline{x^\ell}.\text{some}; \overline{x^\ell}[x_i].P$ in the derivation below:

$$\begin{aligned} & \Pi_2 \\ & \quad \vdots \\ & \frac{P_2 \vdash \llbracket \Gamma \rrbracket, y_i : \& \mathbf{1}, x^\ell : \&((\oplus \llbracket \sigma \rrbracket) \otimes (\llbracket \sigma \wedge \sigma \rrbracket_{(\tau,j)})), \llbracket \Theta \rrbracket}{x^\ell.\text{some}_{y_i, \text{fv}(C)}; P_2 \vdash \llbracket \Gamma \rrbracket, y_i : \& \mathbf{1}, x^\ell : \oplus \&((\oplus \llbracket \sigma \rrbracket) \otimes (\llbracket \sigma \wedge \sigma \rrbracket_{(\tau,j)})), \llbracket \Theta \rrbracket}}{[\text{T}\oplus\text{some}]} \\ & \frac{[\text{T}\wp]}{x^\ell(y_i); x^\ell.\text{some}_{y_i, \text{fv}(C)}; P_2 \vdash \llbracket \Gamma \rrbracket, x^\ell : (\& \mathbf{1}) \wp (\oplus \&((\oplus \llbracket \sigma \rrbracket) \otimes (\llbracket \sigma \wedge \sigma \rrbracket_{(\tau,j)}))), \llbracket \Theta \rrbracket}}{[\text{T}\oplus\text{some}]} \\ & \frac{\llbracket C \rrbracket_{x^\ell} \vdash \llbracket \Gamma \rrbracket, x^\ell : \underbrace{\oplus((\& \mathbf{1}) \wp (\oplus \&((\oplus \llbracket \sigma \rrbracket) \otimes (\llbracket \sigma \wedge \sigma \rrbracket_{(\tau,j)}))))}_{\llbracket \sigma \wedge \sigma \wedge \sigma \rrbracket_{(\tau,j)}}}, \llbracket \Theta \rrbracket}}{\text{Def. 12}} \end{aligned}$$

Therefore, $\llbracket C \rrbracket_{x^\ell} \vdash \llbracket \Gamma \rrbracket, x^\ell : \llbracket \sigma \wedge \sigma \wedge \sigma \rrbracket_{(\tau,j)}, \llbracket \Theta \rrbracket$ and the result follows.

(c) For $\llbracket U \rrbracket_{x_i} \vdash x_i : \&_{\eta_i \in \eta} \{i : \llbracket \eta_i \rrbracket\}, \llbracket \Theta \rrbracket$.

To simplify the proof, we consider $U = 1^! \diamond \lambda M' \}^!$ with $\eta = \sigma_1 \diamond \sigma_2$ and $\&_{\eta_i \in \eta} = \&\{1 : \llbracket \sigma_1 \rrbracket, 2 : \llbracket \sigma_2 \rrbracket\}$.

Our translation gives $\llbracket U \rrbracket_{x_i} = x_i.\text{case}\{1 : \overline{x_i}.\text{none}, 2 : \llbracket M' \rrbracket_{x_i}\}$. Hence, we have:

$$[\text{FS:bag}^!] \frac{\frac{\Theta; \cdot \vDash M' : \sigma_2}{\Theta; \cdot \vDash \lambda M' \}^! : \sigma_2}}{\Theta; \cdot \vDash 1^! : \sigma_1} \quad \frac{[\text{FS:bag}^!]}{\Theta; \cdot \vDash 1^! \diamond \lambda M' \}^! : \sigma_1 \diamond \sigma_2}$$

By the induction hypothesis we have that $\Theta; \cdot \vDash M' : \sigma$ implies $\llbracket M' \rrbracket_{x_i} \vDash x_i : \llbracket \sigma \rrbracket, \llbracket \Theta \rrbracket$. Thus,

$$\begin{aligned} & \frac{[\text{T}\&\text{none}]}{\overline{x_i}.\text{none} \vdash x_i : \llbracket \sigma_1 \rrbracket}}{[\text{TWEAKEN}]} \\ & \frac{\overline{x_i}.\text{none} \vdash x_i : \llbracket \sigma_1 \rrbracket, \llbracket \Theta \rrbracket \quad \llbracket M' \rrbracket_{x_i} \vdash x_i : \llbracket \sigma_2 \rrbracket, \llbracket \Theta \rrbracket}}{x_i.\text{case}\{1 : \overline{x_i}.\text{none}, 2 : \llbracket M' \rrbracket_{x_i}\} \vdash x_i : \&\{1 : \llbracket \sigma_1 \rrbracket, 2 : \llbracket \sigma_2 \rrbracket\}, \llbracket \Theta \rrbracket}}{[\text{T}\&]} \end{aligned}$$

Therefore, $x_i.\text{case}\{1 : \overline{x_i}.\text{none}, 2 : \llbracket M' \rrbracket_{x_i}\} \vdash x_i : \&\{1 : \llbracket \sigma_1 \rrbracket, 2 : \llbracket \sigma_2 \rrbracket\}, \llbracket \Theta \rrbracket$ and the result follows.

2. The proof of type preservation for terms, relies on the analysis of ten cases:

(a) **Rule** $[\text{FS:var}^\ell]$: Then we have the following derivation:

$$[\text{FS:var}^\ell] \frac{}{\Theta; x : \tau \vDash x : \tau}$$

By Def. 12, $\llbracket x : \tau \rrbracket = x : \&\overline{\llbracket \tau \rrbracket}$, and by Fig. 16, $\llbracket x \rrbracket_u = \overline{x}.\text{some}; [x \leftrightarrow u]$. The result follows from the derivation:

$$[\text{TID}] \frac{\overline{[x \leftrightarrow u] \vdash x : \overline{[\tau]}, u : [\tau], [\Theta]}}}{[\text{T\&some}] \overline{\overline{x.\text{some}; [x \leftrightarrow u] \vdash x : \&\overline{[\tau]}, u : [\tau], [\Theta]}}}$$

(b) **Rule [FS:var[!]]**: Then we have the following derivation provided $\eta_k = \tau$:

$$[\text{FS:var}^\ell] \frac{\overline{\Theta, x^! : \eta; x : \eta_k \vDash x : \tau}}{[\text{FS:var}^!]} \overline{\Theta, x^! : \eta; - \vDash x[k] : \tau}$$

By Def. 12, $[\Theta, x^! : \eta] = [\Theta], x^! : \overline{!\&\eta_i \in \eta \{i; [\eta_i]\}}$, and by our translation (in Fig. 16), $[x[k]]_u = \overline{?x^! [x_i]; \overline{x_i}.k; [x_i \leftrightarrow u]}$. The result follows from the derivation:

$$[\text{TID}] \frac{\overline{[x_i \leftrightarrow u] \vdash u : [\tau], x_i : \overline{[\eta_k]}, [\Theta]}}{[\text{T}\oplus]} \overline{\overline{x_i}.k; [x_i \leftrightarrow u] \vdash u : [\tau], x_i : \oplus_{\eta_i \in \eta} \{i; \overline{[\eta_i]}\}, [\Theta]}}{[\text{T?}]} \overline{\overline{?x^! [x_i]; \overline{x_i}.k; [x_i \leftrightarrow u] \vdash u : [\tau], x^! : ? \oplus_{\eta_i \in \eta} \{i; \overline{[\eta_i]}\}, [\Theta]}}$$

(c) **Rule [FS:weak]**: Then we have the following derivation:

$$[\text{FS:weak}] \frac{\Theta; \Gamma \vDash M' : \tau}{\Theta; \Gamma, x : \omega \vDash M'[\leftarrow x] : \tau}$$

By Def. 12, $[\Gamma, x : \omega] = [\Gamma], x^\ell : \overline{[\omega]_{(\sigma, i_1)}}$, and by our translation Fig. 16,

$[\overline{M'[\leftarrow x]}]_u = \overline{x^\ell.\text{some}; \overline{x^\ell [y_i]}.(y_i.\text{some}_{u, \text{fv}(M')}); y_i(); [M']_u | \overline{x^\ell.\text{none}}}$.

By IH, we have $[M']_u \vdash [\Gamma], u : [\tau], [\Theta]$. The result follows from the derivation, omitting labels:

$$\frac{\frac{\frac{[M']_u \vdash [\Gamma], u : [\tau], [\Theta]}{y_i(); [M']_u \vdash y_i : \perp, [\Gamma], u : [\tau], [\Theta]}}{y_i.\text{some}_{u, \text{fv}(M')}; y_i(); [M']_u \vdash y_i : \oplus \perp, [\Gamma], u : [\tau], [\Theta]}}{\overline{x^\ell [y_i]}.(y_i.\text{some}_{u, \text{fv}(M')}); y_i(); [M']_u | \overline{x^\ell.\text{none}} \vdash x^\ell : (\oplus \perp) \otimes (\&A), [\Gamma], u : [\tau], [\Theta]}}{\frac{\overline{x^\ell.\text{none}} \vdash x^\ell : \&A}{\overline{x^\ell.\text{none}} \vdash x^\ell : \&A, [\Theta]}} \frac{\overline{x^\ell.\text{none}} \vdash x^\ell : \&A, [\Theta]}{[M'[\leftarrow x]]_u \vdash x^\ell : \&((\oplus \perp) \otimes (\&A)), [\Gamma], u : [\tau], [\Theta]}}$$

Since A is arbitrary, we can take $A = 1$ for $[\omega]_{(\sigma, 0)}$ and $A = (\overline{(\&\overline{\sigma})} \wp (\overline{[\omega]_{(\sigma, i-1)}}))$ for $[\omega]_{(\sigma, i)}$ where $i > 0$, in both cases, the result follows.

(d) **Rule [FS : abs-sh]**:

Then $M = \lambda x.(M'[\tilde{x} \leftarrow x])$, and the derivation is:

$$[\text{FS:abs-sh}] \frac{\Theta, x^! : \eta; \Gamma, x : \sigma^k \vDash M'[\tilde{x} \leftarrow x] : \tau \quad x \notin \text{dom}(\Gamma)}{\Theta; \Gamma \vDash \lambda x.(M'[\tilde{x} \leftarrow x]) : (\sigma^k, \eta) \rightarrow \tau}$$

By IH, we have $[M'[\tilde{x} \leftarrow x]]_u \vdash u : [\tau], [\Gamma], x^\ell : \overline{[\sigma^k]_{(\sigma, i)}}$, $[\Theta], x^! : \overline{[\eta]}$ and our translation (in Fig. 16) gives $[\lambda x.M'[\tilde{x} \leftarrow x]]_u = \overline{u.\text{some}; u(x); \overline{x.\text{some}}; x(x^\ell); x(x^!); x(); [M'[\tilde{x} \leftarrow x]]_u}$, whose type derivation $[M'[\tilde{x} \leftarrow x]]_u \vdash u : [(\sigma^k, \eta) \rightarrow \tau], [\Gamma], [\Theta]$, was given in the proof of Lemma 17, item 3.

(e) **Rule [FS : app]**: Then $M = M' B$, where $B = C \star U$ and the derivation is:

$$[\text{FS:app}] \frac{\Theta; \Gamma \vDash M' : (\sigma^j, \eta) \rightarrow \tau \quad \Theta; \Delta \vDash B : (\sigma^k, \epsilon) \quad \eta \sim \epsilon}{\Theta; \Gamma, \Delta \vDash M' B : \tau}$$

By IH, we have both

- $[M']_u \vdash [\Gamma], u : [(\sigma^j, \eta) \rightarrow \tau], [\Theta]$;
- $[M']_u \vdash [\Gamma], u : [(\sigma^j, \epsilon) \rightarrow \tau], [\Theta]$, by Lemma 17;
- $[B]_u \vdash [\Delta], u : \overline{[(\sigma^k, \epsilon)]_{(\tau_2, n)}}$, $[\Theta]$, for some τ_2 and some n .

Therefore, from the fact that M is well-formed and Fig. 16 and 12, we have:

- $[M'(C \star U)]_u = (\nu v)([M']_v | v.\text{some}_{u, \text{fv}(C)}; \overline{v[x].([v \leftrightarrow u] | [C \star U]_x)})$;
- $[(\sigma^j, \eta) \rightarrow \tau] = \oplus(\overline{[(\sigma^k)_{(\tau_1, m)}}] \otimes (\overline{![\eta]}))$, for some τ_1 and some m .

Also, since $[B]_u \vdash [\Delta], u : \overline{[(\sigma^k, \epsilon)]_{(\tau_2, n)}}$, $[\Theta]$, we have the following derivation II :

$$\frac{\frac{\frac{\overline{[v \leftrightarrow u] \vdash v : \overline{[\tau]}, u : [\tau]}}{[C \star U]_x \vdash [\Delta], x : \overline{[(\sigma^k, \epsilon)]_{(\tau_2, n)}}}, [\Theta]}{\overline{v[x].([v \leftrightarrow u] | [C \star U]_x)} \vdash [\Delta], v : \overline{[(\sigma^k, \epsilon)]_{(\tau_2, n)}} \otimes \overline{[\tau]}, u : [\tau], [\Theta]}}{\overline{v.\text{some}_{u, \text{fv}(C)}; \overline{v[x].([v \leftrightarrow u] | [C \star U]_x)} \vdash [\Delta], v : \oplus(\overline{[(\sigma^k, \epsilon)]_{(\tau_2, n)}} \otimes \overline{[\tau]})}, u : [\tau], [\Theta]}}{\underbrace{\overline{[(\sigma^k, \epsilon) \rightarrow \tau]}}}$$

In order to apply [TCUT], we must have that $\llbracket \sigma^j \rrbracket_{(\tau_1, m)} = \llbracket \sigma^k \rrbracket_{(\tau_2, n)}$, therefore, the choice of τ_1, τ_2, n and m , will consider the different possibilities for j and k , as in Proposition 16.

$$[\text{TCUT}] \frac{\llbracket M' \rrbracket_v \vdash \llbracket \Gamma \rrbracket, v : \llbracket (\sigma^j, \epsilon) \rightarrow \tau \rrbracket; \llbracket \Theta \rrbracket \quad \Pi}{(\nu v)(\llbracket M' \rrbracket_v \mid v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x].(\llbracket v \leftrightarrow u \rrbracket \mid \llbracket B \rrbracket_x)) \vdash \llbracket \Gamma \rrbracket, \llbracket \Delta \rrbracket, u : \llbracket \tau \rrbracket; \llbracket \Theta \rrbracket}$$

We can then conclude that $\llbracket M'B \rrbracket_u \vdash \llbracket \Gamma \rrbracket, \llbracket \Delta \rrbracket, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket$ and the result follows.

(f) **Rule [FS : share]:** Then $M = M'[x_1, \dots, x_k \leftarrow x]$ and the derivation is:

$$[\text{FS : share}] \frac{\Theta; \Delta, x_1 : \sigma, \dots, x_k : \sigma \Vdash M' : \tau \quad x \notin \Delta \quad k \neq 0}{\Theta; \Delta, x : \sigma_k \Vdash M'[x_1, \dots, x_k \leftarrow x] : \tau}$$

To simplify the proof we will consider $k = 1$ (the case in which $k > 1$ follows similarly).

By IH, we have $\llbracket M' \rrbracket_u \vdash \llbracket \Delta, x_1 : \sigma \rrbracket, u : \llbracket \tau \rrbracket; \llbracket \Theta \rrbracket$. From Fig. 16 and 12, it follows

$$\begin{aligned} & - \llbracket \Delta, x_1 : \sigma \rrbracket = \llbracket \Delta \rrbracket, \overline{x_1} : \& \llbracket \sigma \rrbracket. \\ & - \llbracket M'[x_1, \leftarrow x] \rrbracket_u = \overline{x^\ell}.\text{some}; \overline{x^\ell}[y_1].(y_1.\text{some}_\emptyset; y_1()); \mathbf{0} \mid \overline{x^\ell}.\text{some}; \overline{x^\ell}.\text{some}_{u, \text{fv}(M') \setminus x_1}; \\ & \quad \amalg_{x_i \in x_1} x(x_i); \overline{x^\ell}.\text{some}; \overline{x^\ell}[y_2].(y_2.\text{some}_{u, \text{fv}(M')}; y_2()); \llbracket M' \rrbracket_u \mid \overline{x^\ell}.\text{none} \end{aligned}$$

We shall split the expression into two parts:

$$\begin{aligned} N_1 &= \overline{x^\ell}.\text{some}; \overline{x^\ell}[y_2].(y_2.\text{some}_{u, \text{fv}(M')}; y_2()); \llbracket M' \rrbracket_u \mid \overline{x^\ell}.\text{none} \\ N_2 &= \overline{x^\ell}.\text{some}; \overline{x^\ell}[y_1].(y_1.\text{some}_\emptyset; y_1()); \mathbf{0} \mid \overline{x^\ell}.\text{some}; \overline{x^\ell}.\text{some}_{u, \text{fv}(M') \setminus x_1}; x(x_1); N_1 \end{aligned}$$

and we obtain the derivation for term N_1 as follows where we omit $\llbracket \Theta \rrbracket$ and derivation labels:

$$\begin{aligned} & \frac{\llbracket M' \rrbracket_u \vdash \llbracket \Delta, x_1 : \sigma \rrbracket, u : \llbracket \tau \rrbracket}{y_2(); \llbracket M' \rrbracket_u \vdash \llbracket \Delta, x_1 : \sigma \rrbracket, u : \llbracket \tau \rrbracket, y_2 : \perp} \\ & \frac{y_2.\text{some}_{u, \text{fv}(M')}; y_2(); \llbracket M' \rrbracket_u \vdash \llbracket \Delta, x_1 : \sigma \rrbracket, u : \llbracket \tau \rrbracket, y_2 : \oplus \perp \quad \overline{x^\ell}.\text{none} \vdash x^\ell : \& A}{\overline{x^\ell}[y_2].(y_2.\text{some}_{u, \text{fv}(M')}; y_2()); \llbracket M' \rrbracket_u \mid \overline{x^\ell}.\text{none} \vdash \llbracket \Delta, x_1 : \sigma \rrbracket, u : \llbracket \tau \rrbracket, x^\ell : (\oplus \perp) \otimes (\& A)} \\ & \frac{\overline{x^\ell}[y_2].(y_2.\text{some}_{u, \text{fv}(M')}; y_2()); \llbracket M' \rrbracket_u \mid \overline{x^\ell}.\text{none} \vdash \llbracket \Delta, x_1 : \sigma \rrbracket, u : \llbracket \tau \rrbracket, x^\ell : (\oplus \perp) \otimes (\& A)}{\underbrace{\overline{x^\ell}.\text{some}; \overline{x^\ell}[y_2].(y_2.\text{some}_{u, \text{fv}(M')}; y_2()); \llbracket M' \rrbracket_u \mid \overline{x^\ell}.\text{none}}_{N_1} \vdash \llbracket \Delta, x_1 : \sigma \rrbracket, u : \llbracket \tau \rrbracket, x^\ell : \llbracket \omega \rrbracket_{(\sigma, i)}} \end{aligned}$$

Notice that the last rule applied [T&some] assigns $x : \&((\oplus \perp) \otimes (\& A))$. Again, since A is arbitrary, we can take $A = \oplus((\& \llbracket \sigma \rrbracket) \wp (\llbracket \omega \rrbracket_{(\sigma, i-1)}))$, obtaining $x : \llbracket \omega \rrbracket_{(\sigma, i)}$.

In order to obtain a type derivation for N_2 , consider the derivation Π_1 :

$$\begin{aligned} & [\text{T}\wp] \frac{N_1 \vdash \llbracket \Delta \rrbracket, x_1 : \& \llbracket \sigma \rrbracket, u : \llbracket \tau \rrbracket, x^\ell : \llbracket \omega \rrbracket_{(\sigma, i)}, \llbracket \Theta \rrbracket}{x(x_i); N_1 \vdash \llbracket \Delta \rrbracket, u : \llbracket \tau \rrbracket, x^\ell : (\& \llbracket \sigma \rrbracket) \wp (\llbracket \omega \rrbracket_{(\sigma, i)}), \llbracket \Theta \rrbracket} \\ & [\text{T}\amalg] \frac{\amalg_{x_i \in x_1} x(x_i); N_1 \vdash \llbracket \Delta \rrbracket, u : \llbracket \tau \rrbracket, x^\ell : (\& \llbracket \sigma \rrbracket) \wp (\llbracket \omega \rrbracket_{(\sigma, i)}), \llbracket \Theta \rrbracket}{x^\ell.\text{some}_{u, \text{fv}(M') \setminus x_1}; \amalg_{x_i \in x_1} x(x_i); N_1 \vdash \llbracket \Delta \rrbracket, u : \llbracket \tau \rrbracket, x^\ell : \oplus((\& \llbracket \sigma \rrbracket) \wp (\llbracket \omega \rrbracket_{(\sigma, i)})), \llbracket \Theta \rrbracket} \\ & \frac{x^\ell.\text{some}_{u, \text{fv}(M') \setminus x_1}; \amalg_{x_i \in x_1} x(x_i); N_1 \vdash \llbracket \Delta \rrbracket, u : \llbracket \tau \rrbracket, x^\ell : \oplus((\& \llbracket \sigma \rrbracket) \wp (\llbracket \omega \rrbracket_{(\sigma, i)})), \llbracket \Theta \rrbracket}{\overline{x^\ell}.\text{some}; \overline{x^\ell}.\text{some}_{u, \text{fv}(M') \setminus x_1}; \amalg_{x_i \in x_1} x(x_i); N_1 \vdash \llbracket \Delta \rrbracket, u : \llbracket \tau \rrbracket, x^\ell : \& \oplus((\& \llbracket \sigma \rrbracket) \wp (\llbracket \omega \rrbracket_{(\sigma, i)})), \llbracket \Theta \rrbracket} \end{aligned}$$

We take $P_1 = \overline{x^\ell}.\text{some}; \overline{x^\ell}.\text{some}_{u, \text{fv}(M') \setminus x_1}; \amalg_{x_i \in x_1} x(x_i); N_1$ and $\Gamma_1 = \llbracket \Delta \rrbracket, u : \llbracket \tau \rrbracket$ and continue the derivation of N_2

$$\begin{aligned} & [\text{T}\cdot] \frac{\mathbf{0} \vdash -, \llbracket \Theta \rrbracket}{y_1(); \mathbf{0} \vdash y_1 : \perp, \llbracket \Theta \rrbracket} \quad \Pi_1 \\ & [\text{T}\perp] \frac{y_1(); \mathbf{0} \vdash y_1 : \perp, \llbracket \Theta \rrbracket}{y_1.\text{some}_\emptyset; y_1(); \mathbf{0} \vdash y_1 : \oplus \perp, \llbracket \Theta \rrbracket} \quad \vdots \\ & [\text{T}\otimes] \frac{\overline{x^\ell}[y_1].(y_1.\text{some}_\emptyset; y_1()); \mathbf{0} \mid P_1 \vdash \Gamma_1, x^\ell : \& \oplus((\& \llbracket \sigma \rrbracket) \wp (\llbracket \omega \rrbracket_{(\sigma, i)})), \llbracket \Theta \rrbracket}{\overline{x^\ell}[y_1].(y_1.\text{some}_\emptyset; y_1()); \mathbf{0} \mid P_1 \vdash \Gamma_1, x^\ell : (\oplus \perp) \otimes (\& \oplus((\& \llbracket \sigma \rrbracket) \wp (\llbracket \omega \rrbracket_{(\sigma, i)}))), \llbracket \Theta \rrbracket} \\ & \frac{\overline{x^\ell}[y_1].(y_1.\text{some}_\emptyset; y_1()); \mathbf{0} \mid P_1 \vdash \Gamma_1, x^\ell : (\oplus \perp) \otimes (\& \oplus((\& \llbracket \sigma \rrbracket) \wp (\llbracket \omega \rrbracket_{(\sigma, i)}))), \llbracket \Theta \rrbracket}{\underbrace{\overline{x^\ell}.\text{some}; \overline{x^\ell}[y_1].(y_1.\text{some}_\emptyset; y_1()); \mathbf{0} \mid P_1}_{N_2} \vdash \Gamma_1, x^\ell : \llbracket \sigma \wedge \omega \rrbracket_{(\sigma, i)}, \llbracket \Theta \rrbracket} \end{aligned}$$

Hence the theorem holds for this case.

(g) **Rule [FS : Esub]:** Then $M = (M'[\tilde{x} \leftarrow x])\langle\langle B/x \rangle\rangle$ and

$$[\text{FS:Esub}] \frac{\Theta, x' : \eta; \Gamma, x : \sigma^j \Vdash M'[\tilde{x} \leftarrow x] : \tau \quad \Theta; \Delta \Vdash B : (\sigma^k, \epsilon) \quad \eta \sim \epsilon}{\Theta; \Gamma, \Delta \Vdash (M'[\tilde{x} \leftarrow x])\langle\langle B/x \rangle\rangle : \tau}$$

By Proposition 16 and IH we have:

$$\begin{aligned} & \llbracket M'[x_1, \dots, x_k \leftarrow x] \rrbracket_u \vdash \llbracket \Gamma \rrbracket, x^\ell : \overline{\llbracket \sigma^j \rrbracket}_{(\tau, n)}, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket, x^! : \overline{\llbracket \eta \rrbracket} \\ & \llbracket M'[x_1, \dots, x_k \leftarrow x] \rrbracket_u \vdash \llbracket \Gamma \rrbracket, x^\ell : \overline{\llbracket \sigma^j \rrbracket}_{(\tau, n)}, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket, x^! : \overline{\llbracket \epsilon \rrbracket}, \text{ by Lemma 17} \\ & \llbracket B \rrbracket_x \vdash \llbracket \Delta \rrbracket, x : \llbracket (\sigma^k, \epsilon) \rrbracket_{(\tau, m)}, \llbracket \Theta \rrbracket \end{aligned}$$

From Fig. 16, we have

$$\llbracket M'[\tilde{x} \leftarrow x] \llbracket B/x \rrbracket \rrbracket_u = (\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^!); x()); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u \mid \llbracket C \star U \rrbracket_x$$

Therefore we obtain the following derivation Π :

$$\begin{aligned} & \text{[T}\perp\text{]} \frac{\llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u \vdash \llbracket \Gamma \rrbracket, x^\ell : \overline{\llbracket \sigma^j \rrbracket}_{(\tau, n)}, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket, x^! : \overline{\llbracket \epsilon \rrbracket}}{x(); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u \vdash x:\perp, \llbracket \Gamma \rrbracket, x^\ell : \overline{\llbracket \sigma^j \rrbracket}_{(\tau, n)}, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket, x^! : \overline{\llbracket \epsilon \rrbracket}} \\ & \text{[T}\wp\text{]} \frac{\text{[T}\wp\text{]} \frac{x(x^!); x(); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u \vdash x : (\overline{\llbracket \epsilon \rrbracket}) \wp \perp, \llbracket \Gamma \rrbracket, x^\ell : \overline{\llbracket \sigma^j \rrbracket}_{(\tau, n)}, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket}}{x(x^\ell); x(x^!); x(); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u \vdash x : \overline{\llbracket \sigma^j \rrbracket}_{(\tau, n)} \wp ((\overline{\llbracket \epsilon \rrbracket}) \wp \perp), \llbracket \Gamma \rrbracket, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket}} \\ & \text{[T}\&\text{some}\text{]} \frac{x(x^\ell); x(x^!); x(); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u \vdash x : \overline{\llbracket \sigma^j \rrbracket}_{(\tau, n)} \wp ((\overline{\llbracket \epsilon \rrbracket}) \wp \perp), \llbracket \Gamma \rrbracket, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket}}{\bar{x}.\text{some}; x(x^\ell); x(x^!); x(); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u \vdash x : \overline{\llbracket (\sigma^j, \epsilon) \rrbracket}_{(\tau, n)}, \llbracket \Gamma \rrbracket, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket}} \end{aligned}$$

We take $P_1 = \bar{x}.\text{some}; x(x^\ell); x(x^!); x(); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u$ and continue the derivation of Π

$$\text{[TCUT]} \frac{P_1 \vdash x : \overline{\llbracket (\sigma^j, \epsilon) \rrbracket}_{(\tau, n)}, \llbracket \Gamma \rrbracket, u : \llbracket \tau \rrbracket; \llbracket \Theta \rrbracket \quad \llbracket C \star U \rrbracket_x \vdash \llbracket \Delta \rrbracket, x : \llbracket (\sigma^k, \epsilon) \rrbracket_{(\tau, m)}, \llbracket \Theta \rrbracket}{(\nu x)(P_1 \mid \llbracket C \star U \rrbracket_x) \vdash \llbracket \Gamma \rrbracket, \llbracket \Delta \rrbracket, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket}$$

We must have that $\llbracket \sigma^j \rrbracket_{(\tau, m)} = \llbracket \sigma^k \rrbracket_{(\tau, n)}$ which by our restrictions allows. It follows that $\llbracket M'[x_1 \leftarrow x] \llbracket B/x \rrbracket \rrbracket \vdash \llbracket \Gamma, \Delta \rrbracket, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket$ and the result follows.

(h) **Rule [FS:Esub^ℓ]**: Then $M = M' \llbracket C/x_1, \dots, x_k \rrbracket$, with $C = \llbracket N_1 \rrbracket \cdot \dots \cdot \llbracket N_k \rrbracket$ and

$$\text{[FS:Esub}^\ell\text{]} \frac{\Theta; \Delta, x_1 : \sigma, \dots, x_k : \sigma \vDash M' : \tau \quad \Theta; \Gamma, \vDash C : \sigma^k}{\Theta; \Gamma, \Delta \vDash M' \llbracket C/x_1, \dots, x_k \rrbracket : \tau}$$

with $\Gamma = \Gamma_1, \dots, \Gamma_k$ and:

$$\begin{aligned} & \text{[FS:bag}^\ell\text{]} \frac{\Theta; \Gamma_k \vDash N_k : \sigma \quad \text{[FS:1}^!\text{]} \frac{}{\Theta; - \vDash 1^! : \sigma}}{\text{[FS:bag}^\ell\text{]} \frac{}{\Theta; \Gamma_1 \vDash N_1 : \sigma}} \\ & \text{[FS:bag}^\ell\text{]} \frac{\Theta; \Gamma_1 \vDash N_1 : \sigma \quad \text{[FS:bag}^\ell\text{]} \frac{}{\Theta; \Gamma_2, \dots, \Gamma_k \vDash C : \sigma^{k-1}}{\Theta; \Gamma_1, \dots, \Gamma_k \vDash \llbracket M' \rrbracket \cdot C : \sigma^k}}{\Theta; \Gamma_1, \dots, \Gamma_k \vDash \llbracket M' \rrbracket \cdot C : \sigma^k} \end{aligned}$$

By IH we have both

$$\llbracket N_1 \rrbracket_{x_1} \vdash \llbracket \Gamma_1 \rrbracket, x_1 : \llbracket \sigma \rrbracket, \llbracket \Theta \rrbracket \quad \dots \quad \llbracket N_k \rrbracket_{x_k} \vdash \llbracket \Gamma_k \rrbracket, x_k : \llbracket \sigma \rrbracket, \llbracket \Theta \rrbracket$$

$$\llbracket C \rrbracket_x \vdash \llbracket \Gamma \rrbracket, x : \llbracket \sigma^k \rrbracket_{(\sigma, i)}, \llbracket \Theta \rrbracket \quad \text{and} \quad \llbracket M' \rrbracket_u \vdash \llbracket \Gamma \rrbracket, x_1 : \&\overline{\llbracket \sigma \rrbracket}, \dots, x_k : \&\overline{\llbracket \sigma \rrbracket}, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket$$

From Fig. 16,

$$\begin{aligned} \llbracket M' \llbracket \llbracket N_1 \rrbracket \cdot \dots \cdot \llbracket N_k \rrbracket / x_1, \dots, x_k \rrbracket \rrbracket_u &= (\nu z_1)(z_1.\text{some}_{\text{fv}(N_1)}; \llbracket N_1 \rrbracket_{z_1} \mid \dots \mid (\nu z_k)(z_k.\text{some}_{\text{fv}(N_k)}; \llbracket N_k \rrbracket_{z_k} \\ & \quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \dots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \\ & \quad \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\} \dots) \end{aligned}$$

Let us take $k = 1$ and for $k > 1$ cases follow similarly omitting labels:

$$\begin{aligned} & \frac{\llbracket M' \rrbracket_u \{z_1/x_i\} \vdash \llbracket \Delta \rrbracket, z_1 : \&\overline{\llbracket \sigma \rrbracket}, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket} \quad \frac{\llbracket N_1 \rrbracket_x \vdash \llbracket \Gamma \rrbracket, x : \llbracket \sigma \rrbracket, \llbracket \Theta \rrbracket}}{\prod_{x_i \in \{x_1\}} \llbracket M' \rrbracket_u \{z_1/x_i\} \vdash \llbracket \Delta \rrbracket, z_1 : \&\overline{\llbracket \sigma \rrbracket}, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket} \quad z_1.\text{some}_{\text{fv}(N_1)}; \llbracket N_1 \rrbracket_{z_1} \vdash \llbracket \Gamma \rrbracket, z_1 : \oplus \llbracket \sigma \rrbracket, \llbracket \Theta \rrbracket}} \\ & (\nu z_1) \left(\prod_{x_i \in \{x_1\}} \llbracket M' \rrbracket_u \{z_1/x_i\} \mid z_1.\text{some}_{\text{fv}(N_1)}; \llbracket N_1 \rrbracket_{z_1} \right) \vdash \llbracket \Gamma \rrbracket, \llbracket \Delta \rrbracket, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket \end{aligned}$$

Therefore, $\llbracket M' \llbracket N_1/x_1 \rrbracket \rrbracket_u \vdash \llbracket \Gamma \rrbracket, \llbracket \Delta \rrbracket, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket$ and the result follows.

(i) **Rule [FS:Esub[!]]**: Then $M = M' \llbracket U/x^! \rrbracket$ and

$$\text{[FS:Esub}^!\text{]} \frac{\Theta, x^! : \eta; \Gamma \vDash M' : \tau \quad \Theta; - \vDash U : \eta}{\Theta; \Gamma \vDash M' \llbracket U/x^! \rrbracket : \tau}$$

By IH we have both

$$\llbracket U \rrbracket_{x_i} \vdash x_i : \&_{\eta_i \in \eta} \{i; \llbracket \eta_i \rrbracket\}, \llbracket \Theta \rrbracket \quad \text{and} \quad \llbracket M' \rrbracket_u \vdash \llbracket \Gamma \rrbracket, u : \llbracket \tau \rrbracket, x^! : \overline{\llbracket \eta \rrbracket}, \llbracket \Theta \rrbracket$$

From Definition Fig. 16, $\llbracket M' \llbracket U/x^! \rrbracket \rrbracket_u = (\nu x^!)(\llbracket M' \rrbracket_u \mid !x^!(x_i); \llbracket U \rrbracket_{x_i})$ and

$$\text{[TCUT]} \frac{\llbracket M' \rrbracket_u \vdash \llbracket \Gamma \rrbracket, u : \llbracket \tau \rrbracket, x^! : \overline{\llbracket \eta \rrbracket}, \llbracket \Theta \rrbracket} \quad \text{[T!]} \frac{\llbracket U \rrbracket_{x_i} \vdash x_i : \&_{\eta_i \in \eta} \{i; \llbracket \eta_i \rrbracket\}, \llbracket \Theta \rrbracket}{!x^!(x_i); \llbracket U \rrbracket_{x_i} \vdash x^! : \llbracket \eta \rrbracket, \llbracket \Theta \rrbracket}}{(\nu x^!)(\llbracket M' \rrbracket_u \mid !x^!(x_i); \llbracket U \rrbracket_{x_i}) \vdash \llbracket \Gamma \rrbracket, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket}$$

Therefore, $\llbracket M' \llbracket U/x^! \rrbracket \rrbracket_u \vdash \llbracket \Gamma \rrbracket, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket$ and the result follows.

(j) **Rule [FS : fail]:** Then $M = \text{fail}^{\tilde{x}}$ where $\tilde{x} = x_1, \dots, x_n$ and

$$\text{[FS:fail]} \frac{\text{dom}(\Gamma) = \tilde{x}}{\Theta; \Gamma \vDash \text{fail}^{\tilde{x}} : \tau}$$

From Fig. 16, $\llbracket \text{fail}^{x_1, \dots, x_n} \rrbracket_u = \overline{u}.\text{none} \mid \overline{x_1}.\text{none} \mid \dots \mid \overline{x_k}.\text{none}$ and

$$\text{[TMIX]} \frac{\frac{\overline{u}.\text{none} \vdash u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket} \quad \frac{\overline{x_1}.\text{none} \vdash_1 : \& \llbracket \sigma_1 \rrbracket, \llbracket \Theta \rrbracket} \quad \frac{\overline{x_n}.\text{none} \vdash x_n : \& \llbracket \sigma_n \rrbracket}{\overline{x_n}.\text{none} \vdash x_n : \& \llbracket \sigma_n \rrbracket, \llbracket \Theta \rrbracket}}{\dots} \quad \frac{\overline{u}.\text{none} \vdash u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket} \quad \overline{x_1}.\text{none} \mid \dots \mid \overline{x_k}.\text{none} \vdash x_1 : \& \llbracket \sigma_1 \rrbracket, \dots, x_n : \& \llbracket \sigma_n \rrbracket, \llbracket \Theta \rrbracket}}{\overline{u}.\text{none} \mid \overline{x_1}.\text{none} \mid \dots \mid \overline{x_k}.\text{none} \vdash x_1 : \& \llbracket \sigma_1 \rrbracket, \dots, x_n : \& \llbracket \sigma_n \rrbracket, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket}}$$

Thus, $\llbracket \text{fail}^{x_1, \dots, x_n} \rrbracket_u \vdash x_1 : \& \llbracket \sigma_1 \rrbracket, \dots, x_n : \& \llbracket \sigma_n \rrbracket, u : \llbracket \tau \rrbracket, \llbracket \Theta \rrbracket$ and the result follows. \square

G.2 Completeness

Definition 18 (Linearly Partially Open Terms). We say that a λ_C -term M is linearly partially open if $\forall x \in \text{lfv}(M)$ implies that x is not a sharing variable.

Proposition 3. Suppose N is a well-formed linearly partially open λ_C -term with $\text{head}(N) = x$ (x denoting either linear or unrestricted occurrence of x). Then,

$$\llbracket N \rrbracket_u = (\nu y_1)(\dots(\nu y_m)(\llbracket x \rrbracket_n \mid P_m) \dots \mid P_1)$$

which we shall denote as: $\llbracket N \rrbracket_u = (\nu \tilde{y})(\llbracket x \rrbracket_n \mid P)$, for some names $\tilde{y} =$ and n , and processes P .

Proof. The proof is by induction on the structure of N .

1. $N = x$: then $\llbracket x \rrbracket_u = \overline{x}.\text{some}; [x \leftrightarrow u]$. Hence $P = \mathbf{0}$ and $\tilde{y} = \emptyset$.
2. $N = x[j]$: then $\llbracket x[j] \rrbracket_u = ?x^! [x_i]; \overline{x_i}.j; [x_i \leftrightarrow u]$. Hence $P = \mathbf{0}$ and $\tilde{y} = \emptyset$.
3. $N = M (C \star U)$: then $\text{head}(M (C \star U)) = \text{head}(M) = x$ and

$$\llbracket N \rrbracket_u = \llbracket M (C \star U) \rrbracket_u = (\nu v)(\llbracket M \rrbracket_v \mid v.\text{some}_{u, \text{lfv}(C)}; \overline{v}[x].([v \leftrightarrow u] \mid \llbracket (C \star U) \rrbracket_x))$$

The result follows by induction hypothesis applied on $\llbracket M \rrbracket_u$.

4. $N = (M[\tilde{y} \leftarrow y]) \langle \langle C \star U/y \rangle \rangle$: this case does not apply since $\text{head}(N) \neq x$.
5. $N = M \langle \langle C/x_1, \dots, x_k \rangle \rangle$: then $\text{head}(M \langle \langle C/x_1, \dots, x_k \rangle \rangle) = \text{head}(M) = x$. Let $C = \lambda M_1 \dots \lambda M_k$

$$\llbracket N \rrbracket_u = (\nu z_1)(z_1.\text{some}_{\text{lfv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_k)(z_k.\text{some}_{\text{lfv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \dots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \llbracket M \rrbracket_u \{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\}) \dots)$$

The result follows by induction hypothesis applied on $\llbracket M \rrbracket_u$.

6. $N = M \llbracket U/x^! \rrbracket$: then $\text{head}(M \llbracket U/x^! \rrbracket) = \text{head}(M) = x$ and

$$\llbracket N \rrbracket_u = \llbracket M \llbracket U/x^! \rrbracket \rrbracket_u = (\nu x^!)(\llbracket M \rrbracket_u \mid !x^!(x_i); \llbracket U \rrbracket_{x_i})$$

The result follows by induction hypothesis applied on $\llbracket M \rrbracket_u$. \square

Theorem 20 (Completeness (Under \rightsquigarrow)). *If $N \longrightarrow M$ for a well-formed closed λ_C -term N , then $\llbracket N \rrbracket_u \rightsquigarrow^* \llbracket M \rrbracket_u$.*

Proof. By induction on the reduction rule applied to infer $N \longrightarrow M$. We have five cases.

1. Case [RS : Beta]:

Then $N = (\lambda x.(M'[\tilde{x} \leftarrow x]))B \longrightarrow (M'[\tilde{x} \leftarrow x])\langle\langle B/x \rangle\rangle = M$, where $B = C \star U$. The result follows from

$$\begin{aligned} \llbracket N \rrbracket_u &= (\nu v)(\llbracket \lambda x.(M'[\tilde{x} \leftarrow x]) \rrbracket_v \mid v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x]; (\llbracket C \star U \rrbracket_x \mid [v \leftrightarrow u])) \\ &= (\nu v)(\bar{v}.\text{some}; v(x); \bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_v \mid \\ &\quad v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x]; (\llbracket C \star U \rrbracket_x \mid [v \leftrightarrow u])) \\ &\rightsquigarrow (\nu v)(v(x); \bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_v \mid \bar{v}[x]; (\llbracket C \star U \rrbracket_x \mid [v \leftrightarrow u])) \\ &\rightsquigarrow (\nu v)((\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_v \mid \llbracket C \star U \rrbracket_x) \mid [v \leftrightarrow u]) \\ &\rightsquigarrow (\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u \mid \llbracket C \star U \rrbracket_x) = \llbracket M \rrbracket_u \end{aligned}$$

2. Case [RS : Ex-Sub]: Then $N = M'[x_1, \dots, x_k \leftarrow x]\langle\langle C \star U/x \rangle\rangle$, with $C = \lambda M_1 \int \dots \int M_k \int$, $k \geq 0$ and $M' \neq \text{fail}^{\bar{y}}$.

The reduction is $N = M'[x_1, \dots, x_k \leftarrow x]\langle\langle C \star U/x \rangle\rangle \longrightarrow M'\langle\langle C/x_1, \dots, x_k \rangle\rangle \llbracket U/x^1 \rrbracket = M$.

We detail the translations of $\llbracket N \rrbracket_u$ and $\llbracket M \rrbracket_u$. To simplify the proof, we will consider $k = 2$ (the case in which $k > 2$ is follows analogously. Similarly the case of $k < 2$ it contained within $k = 2$). The result follows from:

$$\begin{aligned} \llbracket N \rrbracket_u &= \llbracket M'[x_1 \leftarrow x]\langle\langle C \star U/x \rangle\rangle \rrbracket_u = (\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u \mid \llbracket C \star U \rrbracket_x) \\ &= (\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u \mid x.\text{some}_{\text{fv}(C)}; \bar{x}[x^\ell]; (\llbracket C \rrbracket_{x^\ell} \mid \bar{x}[x^1]; \\ &\quad (!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid \bar{x}[])) (:= P_{\mathbb{N}}) \\ &\rightsquigarrow^* (\nu x^1)((\nu x^\ell)(\llbracket M'[x_1, x_2 \leftarrow x] \rrbracket_u \mid \llbracket \lambda M_1 \int \cdot \int M_2 \int \rrbracket_{x^\ell} \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\ &= (\nu x^1)((\nu x^\ell)(\bar{x}^\ell.\text{some}; \bar{x}^\ell[y_1]; (y_1.\text{some}_\emptyset; y_1()); \mathbf{0} \\ &\quad \mid \bar{x}^\ell.\text{some}; x^\ell.\text{some}_{u, \text{fv}(M') \setminus \bar{x}}; \prod_{x_{i_1} \in \bar{x}} x(x_{i_1}); \llbracket M'[(\tilde{x} \setminus x_{i_1}) \leftarrow x] \rrbracket_u) \mid \\ &\quad x^\ell.\text{some}_{\text{fv}(C)}; x(y_1); x^\ell.\text{some}_{y_1, \text{fv}(C)}; \bar{x}^\ell.\text{some}; \bar{x}^\ell[z_1]; \\ &\quad (z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \llbracket \lambda M_2 \int \rrbracket_{x^\ell} \mid \bar{y}_1.\text{none})) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\ &\rightsquigarrow^* (\nu x^1)((\nu x^\ell)(\llbracket \lambda M_2 \int \rrbracket_{x^\ell} \mid (\nu z_1)(\prod_{x_{i_1} \in \bar{x}} \llbracket M'[(\tilde{x} \setminus x_{i_1}) \leftarrow x] \rrbracket_u \{z_1/x_{i_1}\} \mid \\ &\quad z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1}) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\ &= (\nu x^1)((\nu x^\ell)(x^\ell.\text{some}_{\text{fv}(C)}; x(y_2); x^\ell.\text{some}_{y_2, \text{fv}(C)}; \bar{x}^\ell.\text{some}; \bar{x}^\ell[z_2]; \\ &\quad (z_2.\text{some}_{\text{fv}(M_2)}; \llbracket M_2 \rrbracket_{z_2} \mid \llbracket \mathbf{1} \rrbracket_{x^\ell} \mid \bar{y}_2.\text{none})) \mid \\ &(\nu z_1)(\prod_{x_{i_1} \in \bar{x}} \bar{x}^\ell.\text{some}; \bar{x}^\ell[y_2]; (y_2.\text{some}_\emptyset; y_2()); \mathbf{0} \\ &\quad \mid \bar{x}^\ell.\text{some}; x^\ell.\text{some}_{u, \text{fv}(M') \setminus (\bar{x} \setminus x_{i_1})}; \prod_{x_{i_2} \in (\bar{x} \setminus x_{i_1})} x(x_{i_2}); \llbracket M'[\leftarrow x] \rrbracket_u \{z_1/x_{i_1}\} \mid \\ &\quad z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1}) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\ &\longrightarrow^* (\nu x^1)((\nu x^\ell)(\llbracket \mathbf{1} \rrbracket_{x^\ell} \mid \\ &\quad (\nu z_2)((\nu z_1)(\prod_{x_{i_1} \in \bar{x}} \prod_{x_{i_2} \in (\bar{x} \setminus x_{i_1})} x(x_{i_2}); \llbracket M'[\leftarrow x] \rrbracket_u \{z_1/x_{i_1}\} \mid \\ &\quad z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1}) \mid z_2.\text{some}_{\text{fv}(M_2)}; \llbracket M_2 \rrbracket_{z_2})) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\ &\longrightarrow^* (\nu x^1)((\nu z_2)((\nu z_1)(\prod_{x_{i_1} \in \bar{x}} \prod_{x_{i_2} \in (\bar{x} \setminus x_{i_1})} x(x_{i_2}); \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \mid \\ &\quad z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1}) \mid z_2.\text{some}_{\text{fv}(M_2)}; \llbracket M_2 \rrbracket_{z_2}) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) = \llbracket M \rrbracket_u \end{aligned}$$

3. Case [RS:Fetch^ℓ]:

Then we have $N = M'\langle\langle C/x_1, \dots, x_k \rangle\rangle$ with $\text{head}(M') = x_j$, $C = M_1, \dots, M_k$ and $N \longrightarrow M'\{M_i/x_j\}\langle\langle (C \setminus M_i)/x_1, \dots, x_k \setminus x_j \rangle\rangle = M$, for some $M_i \in C$.

On the one hand, we have:

$$\begin{aligned}
\llbracket N \rrbracket_u &= \llbracket M' \langle C/x_1, \dots, x_k, x_j \rangle \rrbracket_u = (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \dots \\
&\quad \quad \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\} \dots) \\
&= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \dots \\
&\quad \quad \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} (\nu y)(\llbracket x_j \rrbracket_y \mid P)\{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\} \\
&\quad \dots) \quad (*) \\
&= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \dots \\
&\quad \quad \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \\
&\quad \quad \quad (\nu y)(\overline{x_j}.\mathbf{some}; [x_j \leftrightarrow y] \mid P)\{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\} \dots)
\end{aligned} \tag{13}$$

where $(*)$ is inferred via Proposition 3.

Let us consider the case when $j = k$ and $M_i = M_1$ the other cases proceed similarly. Then we have the following reduction:

$$\begin{aligned}
&= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \dots \\
&\quad \quad \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \\
&\quad \quad \quad (\nu y)(\overline{x_k}.\mathbf{some}; [x_k \leftrightarrow y] \mid P)\{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\} \dots) \\
&\rightsquigarrow (\nu z_1)(\llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \prod_{x_{i_2} \in \{x_1, \dots, x_{k-1}\}} \dots \\
&\quad \quad \prod_{x_{i_k} \in \{x_1, \dots, x_{k-1} \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \\
&\quad \quad \quad (\nu y)([z_1 \leftrightarrow y] \mid P)\{z_2/x_{i_2}\} \dots \{z_k/x_{i_k}\} \dots) \\
&\rightsquigarrow (\nu z_2)(z_2.\mathbf{some}_{\text{fv}(M_2)}; \llbracket M_2 \rrbracket_{z_2} \mid \dots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \prod_{x_{i_2} \in \{x_1, \dots, x_{k-1}\}} \dots \\
&\quad \quad \prod_{x_{i_k} \in \{x_1, \dots, x_{k-1} \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \\
&\quad \quad \quad (\nu y)(\llbracket M_1 \rrbracket_y \mid P)\{z_2/x_{i_2}\} \dots \{z_k/x_{i_k}\} \dots)
\end{aligned} \tag{14}$$

On the other hand, we have:

$$\begin{aligned}
\llbracket M \rrbracket_u &= \llbracket M' \{M_1/x_k\} \langle (C \setminus M_1)/x_1, \dots, x_{k-1} \rangle \rrbracket_u \\
&= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_2)}; \llbracket M_2 \rrbracket_{z_1} \mid \dots (\nu z_{k-1})(z_{k-1}.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_{k-1}} \\
&\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_{k-1}\}} \dots \prod_{x_{i_{k-1}} \in \{x_1, \dots, x_{k-1} \setminus x_{i_1}, \dots, x_{i_{k-2}}\}} \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \dots \{z_{k-1}/x_{i_{k-1}}\} \dots)
\end{aligned} \tag{15}$$

Therefore, by (14), (15) and taking $M_{j_k} = M_i$ the result follows.

4. Case $[\mathbf{RS:Fetch}]^!$:

Then, $N = M' \llbracket U/x^! \rrbracket$ with $\text{head}(M') = x^![k]$, $U_k = \lambda N^!]$ and $N \longrightarrow M' \{N/x^! \} \llbracket U/x^! \rrbracket = M$. The result follows from

$$\begin{aligned}
\llbracket N \rrbracket_u &= \llbracket M' \llbracket U/x \rrbracket \rrbracket_u = (\nu x^1)(\llbracket M' \rrbracket_u \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&= (\nu x^1)((\nu y)(\llbracket x^1[k] \rrbracket_j \mid P) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \quad (*) \\
&= (\nu x^1)((\nu y)(?x^1[x_i]; \overline{x_i}.k; [x_i \leftrightarrow j] \mid P) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&\rightsquigarrow (\nu x^1)((\nu y)(?x^1[x_i]; \overline{x_i}.k; [x_i \leftrightarrow j] \mid P) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&\rightsquigarrow (\nu x^1)(!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid (\nu x_i)((\nu y)(\overline{x_i}.k; [x_i \leftrightarrow j] \mid P) \mid \llbracket U \rrbracket_{x_i})) \quad (16) \\
&= (\nu x^1)(!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid (\nu x_i)((\nu y)(\overline{x_i}.k; [x_i \leftrightarrow j] \mid P) \mid x_i.\mathbf{case}\{i : \llbracket U_i \rrbracket_{x_i}\}_{U_i \in U})) \\
&\rightsquigarrow (\nu x^1)(!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid (\nu x_i)((\nu y)([x_i \leftrightarrow j] \mid P) \mid \llbracket U_i \rrbracket_{x_i})) \\
&\rightsquigarrow (\nu x^1)(!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid (\nu y)(\llbracket U_i \rrbracket_j \mid P)) \\
&\rightsquigarrow (\nu x^1)(!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid (\nu y)(\llbracket N \rrbracket_j \mid P)) = \llbracket M \rrbracket_u
\end{aligned}$$

where the reductions denoted by $(*)$ are inferred via Proposition 3.

5. Case $[\mathbf{RS} : \mathbf{TCont}]$: This case follows by IH.

6. Case $[\mathbf{RS} : \mathbf{Fail}^\ell]$:

Then we have $N = M'[x_1, \dots, x_k \leftarrow x] \llbracket \langle C \star U/x \rangle \rrbracket$ with $k \neq \mathbf{size}(C)$ and $N \longrightarrow \mathbf{fail}^{\tilde{y}} = M$, where $\tilde{y} = (\mathbf{lfv}(M') \setminus \{x_1, \dots, x_k\}) \cup \mathbf{lfv}(C)$. Let $C = \llbracket M_1 \rrbracket \dots \llbracket M_l \rrbracket$ and we assume that $k > l$ and we proceed similarly for $k > l$. Hence $k = l + m$ for some $m \geq 1$

$$\begin{aligned}
\llbracket N \rrbracket_u &= \llbracket M'[x_1, \dots, x_k \leftarrow x] \llbracket \langle C \star U/x \rangle \rrbracket \rrbracket_u = (\nu x)(\overline{x}.\mathbf{some}; x(x^\ell); x(x^1); x(); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u \mid \llbracket C \star U \rrbracket_x) \\
&= (\nu x)(\overline{x}.\mathbf{some}; x(x^\ell); x(x^1); x(); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u \mid \\
&\quad x.\mathbf{some}_{\mathbf{lfv}(C)}; \overline{x}[x^\ell]; (\llbracket C \rrbracket_{x^\ell} \mid \overline{x}[x^1]; (!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid \overline{x}[]))) \\
&\rightsquigarrow^* (\nu x^1)((\nu x^\ell)(\llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u \mid \llbracket C \rrbracket_{x^\ell}) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&= (\nu x^1)((\nu x^\ell)(\overline{x}^\ell.\mathbf{some}; \overline{x}^\ell[y_1]; (y_1.\mathbf{some}_\emptyset; y_1()); \mathbf{0} \\
&\quad \mid \overline{x}^\ell.\mathbf{some}; x^\ell.\mathbf{some}_{u, \mathbf{lfv}(M') \setminus \tilde{x}}; \prod_{x_{i_1} \in \tilde{x}} x(x_{i_1}); \dots \\
&\quad \overline{x}^\ell.\mathbf{some}; \overline{x}^\ell[y_k]; (y_k.\mathbf{some}_\emptyset; y_k()); \mathbf{0} \\
&\quad \mid \overline{x}^\ell.\mathbf{some}; x^\ell.\mathbf{some}_{u, \mathbf{lfv}(M') \setminus (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{k-1}})}; \prod_{x_{i_k} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{k-1}})} x(x_{i_k}); \llbracket M'[\leftarrow x] \rrbracket_u)) \mid \\
&\quad x^\ell.\mathbf{some}_{\mathbf{lfv}(C)}; x(y_1); x^\ell.\mathbf{some}_{y_1, \mathbf{lfv}(C)}; \overline{x}^\ell.\mathbf{some}; \overline{x}^\ell[z_1]; \\
&\quad (z_1.\mathbf{some}_{\mathbf{lfv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \overline{y_1}.\mathbf{none} \mid \dots \mid x^\ell.\mathbf{some}_{\mathbf{lfv}(C)}; x(y_l); x^\ell.\mathbf{some}_{y_l, \mathbf{lfv}(M_l)}; \overline{x}^\ell.\mathbf{some}; \overline{x}^\ell[z_l]; \\
&\quad (z_l.\mathbf{some}_{\mathbf{lfv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \llbracket \mathbf{1} \rrbracket_{x^\ell} \mid \overline{y_l}.\mathbf{none}) \dots)) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \quad (:= P_{\mathbb{N}})
\end{aligned}$$

we reduce $P_{\mathbb{N}}$ arbitrarily synchronising along channels x^ℓ, y_1, \dots, y_l .

$$\begin{aligned}
P_{\mathbb{N}} &\rightsquigarrow^* (\nu x^1)((\nu x^\ell)(\llbracket 1 \rrbracket_{x^\ell} \mid (\nu z_1)(z_1.\text{some}_{\text{lfv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_l)(z_l.\text{some}_{\text{lfv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \\
&\quad \prod_{x_{i_1} \in \tilde{x}} \cdots \prod_{x_{i_l} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{l-1}})} \overline{x^\ell}.\text{some}; \overline{x^\ell}[y_{l+1}]; (y_{l+1}.\text{some}_\emptyset; y_{l+1}()); \mathbf{0} \\
&\quad \mid \overline{x^\ell}.\text{some}; x^\ell.\text{some}_{u, \text{lfv}(M') \setminus (\tilde{x} \setminus x_{i_1}, \dots, x_{i_l})}; \prod_{x_{i_{l+1}} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_l})} x(x_{i_{l+1}}); \cdots \\
&\quad \overline{x^\ell}.\text{some}; \overline{x^\ell}[y_k]; (y_k.\text{some}_\emptyset; y_k()); \mathbf{0} \\
&\quad \mid \overline{x^\ell}.\text{some}; x^\ell.\text{some}_{u, \text{lfv}(M') \setminus (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{k-1}})}; \prod_{x_{i_k} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{k-1}})} x(x_{i_k}); \\
&\quad \llbracket M'[\leftarrow x]_u \{z_1/x_{i_1}\} \cdots \{z_l/x_{i_l}\} \cdots \rrbracket \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&= (\nu x^1)((\nu x^\ell)(x^\ell.\text{some}_\emptyset; x(y_{l+1}); (\overline{y_{l+1}}.\text{some}; \overline{y_{l+1}}[] \mid x^\ell.\text{some}_\emptyset; \overline{x^\ell}.\text{none}) \\
&\quad \mid (\nu z_1)(z_1.\text{some}_{\text{lfv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_l)(z_l.\text{some}_{\text{lfv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \\
&\quad \prod_{x_{i_1} \in \tilde{x}} \cdots \prod_{x_{i_l} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{l-1}})} \overline{x^\ell}.\text{some}; \overline{x^\ell}[y_{l+1}]; (y_{l+1}.\text{some}_\emptyset; y_{l+1}()); \mathbf{0} \\
&\quad \mid \overline{x^\ell}.\text{some}; x^\ell.\text{some}_{u, \text{lfv}(M') \setminus (\tilde{x} \setminus x_{i_1}, \dots, x_{i_l})}; \prod_{x_{i_{l+1}} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_l})} x(x_{i_{l+1}}); \cdots \\
&\quad \overline{x^\ell}.\text{some}; \overline{x^\ell}[y_k]; (y_k.\text{some}_\emptyset; y_k()); \mathbf{0} \\
&\quad \mid \overline{x^\ell}.\text{some}; x^\ell.\text{some}_{u, \text{lfv}(M') \setminus (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{k-1}})}; \prod_{x_{i_k} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{k-1}})} x(x_{i_k}); \\
&\quad \llbracket M'[\leftarrow x]_u \{z_1/x_{i_1}\} \cdots \{z_l/x_{i_l}\} \cdots \rrbracket \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&\rightsquigarrow^* (\nu x^1)((\nu x^\ell)(\overline{x^\ell}.\text{none} \mid (\nu z_1)(z_1.\text{some}_{\text{lfv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_l)(z_l.\text{some}_{\text{lfv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \\
&\quad \prod_{x_{i_1} \in \tilde{x}} \cdots \prod_{x_{i_l} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{l-1}})} x^\ell.\text{some}_{u, \text{lfv}(M') \setminus (\tilde{x} \setminus x_{i_1}, \dots, x_{i_l})}; \prod_{x_{i_{l+1}} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_l})} x(x_{i_{l+1}}); \cdots \\
&\quad \overline{x^\ell}.\text{some}; \overline{x^\ell}[y_k]; (y_k.\text{some}_\emptyset; y_k()); \mathbf{0} \\
&\quad \mid \overline{x^\ell}.\text{some}; x^\ell.\text{some}_{u, \text{lfv}(M') \setminus (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{k-1}})}; \prod_{x_{i_k} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{k-1}})} x(x_{i_k}); \\
&\quad \llbracket M'[\leftarrow x]_u \{z_1/x_{i_1}\} \cdots \{z_l/x_{i_l}\} \cdots \rrbracket \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&\rightsquigarrow (\nu x^1)((\nu z_1)(z_1.\text{some}_{\text{lfv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_l)(z_l.\text{some}_{\text{lfv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \\
&\quad \prod_{x_{i_1} \in \tilde{x}} \cdots \prod_{x_{i_l} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{l-1}})} \overline{u}.\text{none} \mid \overline{(\text{lfv}(M') \setminus \tilde{x})}.\text{none} \mid \overline{(z_1, \dots, z_l)}.\text{none}) \cdots) \mid \\
&\quad !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&\rightsquigarrow^* (\nu x^1)(\prod_{x_{i_1} \in \tilde{x}} \cdots \prod_{x_{i_l} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{l-1}})} \overline{u}.\text{none} \mid \overline{(\text{lfv}(M') \setminus \tilde{x})}.\text{none} \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&\equiv \overline{u}.\text{none} \mid \overline{(\text{lfv}(M') \setminus \tilde{x})}.\text{none} = \llbracket M \rrbracket_u
\end{aligned}$$

7. Case [RS:Fail¹]:

Then, $N = M' \llbracket U/x^1 \rrbracket$ with $\text{head}(M') = x[i]$, $U_i = 1^1$ and $N \longrightarrow M' \{\text{fail}^\emptyset/x^1\} \llbracket U/x^1 \rrbracket$. The result follows from

$$\begin{aligned}
\llbracket N \rrbracket_u &= \llbracket M' \llbracket U/x^1 \rrbracket \rrbracket_u = (\nu x^1)(\llbracket M' \rrbracket_u \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&= (\nu x^1)((\nu \tilde{y})(\llbracket x[i] \rrbracket_j \mid P) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \quad (*) \\
&= (\nu x^1)((\nu \tilde{y})(\overline{x^1}[x_i]; \overline{x_i}.i; [x_i \leftrightarrow j] \mid P) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&\rightsquigarrow (\nu x^1)((\nu x_i)((\nu \tilde{y})(\overline{x_i}.i; [x_i \leftrightarrow j] \mid P) \mid \llbracket U \rrbracket_{x_i}) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&= (\nu x^1)((\nu x_i)((\nu \tilde{y})(\overline{x_i}.i; [x_i \leftrightarrow j] \mid P) \mid x_i.\text{case}\{i : \llbracket U_i \rrbracket_{x_i}\}_{U_i \in U} \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&\rightsquigarrow^* (\nu x^1)((\nu \tilde{y})(\llbracket U_i \rrbracket_j \mid P) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&= (\nu x^1)((\nu \tilde{y})(\llbracket 1^1 \rrbracket_j \mid P) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&= (\nu x^1)((\nu \tilde{y})(\overline{j}.\text{none} \mid P) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) = \llbracket M \rrbracket_u
\end{aligned}$$

8. Case [RS : Cons₁]: Then we have $N = \text{fail}^{\tilde{x}} C \star U$ and $N \longrightarrow \text{fail}^{\tilde{x} \cup \tilde{y}} M$ where $\tilde{y} = \text{lfv}(C)$. Also,

$$\begin{aligned}
\llbracket N \rrbracket_u &= \llbracket \text{fail}^{\tilde{x}} C \star U \rrbracket_u = (\nu v)(\llbracket \text{fail}^{\tilde{x}} \rrbracket_v \mid v.\text{some}_{u, \text{lfv}(C)}; \overline{v}[x]; (\llbracket C \star U \rrbracket_x \mid [v \leftrightarrow u])) \\
&= (\nu v)(\overline{v}.\text{none} \mid \overline{\tilde{x}}.\text{none} \mid v.\text{some}_{u, \text{lfv}(C)}; \overline{v}[x]; (\llbracket C \star U \rrbracket_x \mid [v \leftrightarrow u])) \\
&\rightsquigarrow \overline{\tilde{x}}.\text{none} \mid \overline{u}.\text{none} \mid \overline{\tilde{y}}.\text{none} = \llbracket M \rrbracket_u
\end{aligned}$$

9. Cases $[\text{RS} : \text{Cons}_2]$ and $[\text{RS} : \text{Cons}_3]$: These cases follow by IH similarly to Case 7.
 10. Case $[\text{RS} : \text{Cons}_4]$: Then we have $N = \text{fail}^{\tilde{y}} \llbracket U/x^! \rrbracket$ and $N \longrightarrow \text{fail}^{\tilde{y}} = M$, and

$$\begin{aligned} \llbracket N \rrbracket_u &= \llbracket \text{fail}^{\tilde{y}} \llbracket U/x^! \rrbracket \rrbracket_u = (\nu x^!)(\llbracket \text{fail}^{\tilde{y}} \rrbracket_u \mid !x^!(x_i); \llbracket U \rrbracket_{x_i}) \\ &= (\nu x^!)(\bar{u}.\text{none} \mid \bar{x}.\text{none} \mid !x^!(x_i); \llbracket U \rrbracket_{x_i}) \equiv \bar{u}.\text{none} \mid \bar{x}.\text{none} = \llbracket M \rrbracket_u \quad \square \end{aligned} \quad (17)$$

G.3 Soundness

Theorem 21 (Weak Soundness (Under \rightsquigarrow^*)). *If $\llbracket N \rrbracket_u \rightsquigarrow^* Q$ for a well-formed closed λ_C -term N , then there exist Q' and N' such that $Q \rightsquigarrow^* Q'$, $N \longrightarrow^* N'$ and $\llbracket N' \rrbracket_u \equiv Q'$.*

Proof. By induction on the structure of N and then induction on the number of reductions of $\llbracket N \rrbracket_u \rightsquigarrow^* Q$.

1. **Base case:** $N = x$, $N = x[j]$, $N = \text{fail}^0$ and $N = \lambda x.(M[\tilde{x} \leftarrow x])$.

No reductions can take place, and the result follows trivially. $Q = \llbracket N \rrbracket_u \rightsquigarrow^0 \llbracket N \rrbracket_u = Q'$ and $N \longrightarrow^0 N = N'$.

2. $N = M(C \star U)$.

Then, $\llbracket M(C \star U) \rrbracket_u = (\nu v)(\llbracket M \rrbracket_v \mid v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x]; (\llbracket C \star U \rrbracket_x \mid [v \leftrightarrow u]))$, and we are able to perform the reductions from $\llbracket M(C \star U) \rrbracket_u$.

We now proceed by induction on k , with $\llbracket N \rrbracket_u \rightsquigarrow^k Q$. There are two main cases:

- (a) When $k = 0$ the thesis follows easily:

We have $Q = \llbracket M(C \star U) \rrbracket_u \rightsquigarrow^0 \llbracket M(C \star U) \rrbracket_u = Q'$ and $M(C \star U) \longrightarrow^0 M(C \star U) = N'$.

- (b) The interesting case is when $k \geq 1$.

Then, for some process R and n, m such that $k = n + m$, we have the following:

$$\begin{aligned} \llbracket N \rrbracket_u &= (\nu v)(\llbracket M \rrbracket_v \mid v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x]; (\llbracket C \star U \rrbracket_x \mid [v \leftrightarrow u])) \\ &\rightsquigarrow^m (\nu v)(R \mid v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x]; (\llbracket C \star U \rrbracket_x \mid [v \leftrightarrow u])) \rightsquigarrow^n Q \end{aligned}$$

Thus, the first $m \geq 0$ reduction steps are internal to $\llbracket M \rrbracket_v$; type preservation in $\mathcal{S}\pi^+$ ensures that, if they occur, these reductions do not discard the possibility of synchronizing with $\bar{v}.\text{some}$. Then, the first of the $n \geq 0$ reduction steps towards Q is a synchronization between R and $v.\text{some}_{u, \text{fv}(C)}$.

We consider two sub-cases, depending on the values of m and n :

- i. $m = 0$ and $n \geq 1$:

Then $R = \llbracket M \rrbracket_v$ as $\llbracket M \rrbracket_v \rightsquigarrow^0 \llbracket M \rrbracket_v$. Notice that there are two possibilities of having an unguarded:

A. $M = (\lambda x.(M'[\tilde{x} \leftarrow x])) \langle C_1/\tilde{y}_1 \rangle \cdots \langle C_p/\tilde{y}_p \rangle \llbracket U_1/z_1^! \rrbracket \cdots \llbracket U_q/z_q^! \rrbracket$ ($p, q \geq 0$)

$$\begin{aligned} \llbracket M \rrbracket_v &= \llbracket (\lambda x.(M'[\tilde{x} \leftarrow x])) \langle C_1/\tilde{y}_1 \rangle \cdots \langle C_p/\tilde{y}_p \rangle \llbracket U_1/z_1^! \rrbracket \cdots \llbracket U_q/z_q^! \rrbracket \rrbracket_v \\ &= (\nu z_q^!)(\cdots (\nu z_1^!)(\llbracket (\lambda x.(M'[\tilde{x} \leftarrow x])) \langle C_1/\tilde{y}_1 \rangle \cdots \langle C_p/\tilde{y}_p \rangle \rrbracket_v \mid !z_1^!(z_1); \llbracket U_1 \rrbracket_{z_1} \\ &\quad \cdots \mid !z_q^!(z_q); \llbracket U_q \rrbracket_{z_q}) = Q \end{aligned}$$

Which we shall write as:

$$Q = (\nu z_q^!, \dots, z_1^!)(\llbracket (\lambda x.(M'[\tilde{x} \leftarrow x])) \langle C_1/\tilde{y}_1 \rangle \cdots \langle C_p/\tilde{y}_p \rangle \rrbracket_v \mid !z_1^!(z_1); \llbracket U_1 \rrbracket_{z_1} \cdots \mid !z_q^!(z_q); \llbracket U_q \rrbracket_{z_q})$$

for simplicity to represent the process. We also use this to simplify the translation of linear explicit substitutions of bags from:

$$\begin{aligned} \llbracket M \langle \int M_1 \int \cdots \int M_k \int /x_1, \dots, x_k \rangle \rrbracket_v &= (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\text{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\ &\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \\ &\quad \llbracket M \rrbracket_v \{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\} \cdots) \end{aligned}$$

to be represented as: $\llbracket M \langle C/\tilde{x} \rangle \rrbracket_v = (\nu \tilde{z})(\tilde{z}.\text{some}_{\text{fv}(C)}; \llbracket C \rrbracket_{\tilde{z}} \mid \prod_{\tilde{x}_i \in \text{PER}(\tilde{x})} \llbracket M \rrbracket_v \{\tilde{z}/\tilde{x}_i\})$. These representations are purely for simplicity and are not an alternative to the actual translation. We continue expanding the sub-process

$\llbracket (\lambda x.(M'[\tilde{x} \leftarrow x])) \langle C_1/\tilde{y}_1 \rangle \cdots \langle C_p/\tilde{y}_p \rangle \rrbracket_v$ using the above shortened and simplified notation:

$$\begin{aligned} \llbracket (\lambda x.(M'[\tilde{x} \leftarrow x])) \langle C_1/\tilde{y}_1 \rangle \cdots \langle C_p/\tilde{y}_p \rangle \rrbracket_v &= (\nu \tilde{w}_p)(\tilde{w}_p.\text{some}_{\text{fv}(C_p)}; \llbracket C_p \rrbracket_{\tilde{w}_p} \mid \\ &\quad \prod_{\tilde{y}_{p_i} \in \text{PER}(\tilde{y}_p)} \cdots (\nu \tilde{w}_1)(\tilde{w}_1.\text{some}_{\text{fv}(C_1)}; \llbracket C_1 \rrbracket_{\tilde{w}_1} \mid \\ &\quad \prod_{\tilde{y}_{1_i} \in \text{PER}(\tilde{y}_1)} \llbracket \lambda x.(M'[\tilde{x} \leftarrow x]) \rrbracket_v \{\tilde{w}_1/\tilde{y}_{1_i}\} \cdots \{\tilde{w}_p/\tilde{y}_{p_i}\}) \end{aligned}$$

Hence we represent $\llbracket M \rrbracket_v$ as:

$$\begin{aligned} \llbracket M \rrbracket_v &= (\nu z_q^!, \dots, z_1^!)((\nu \tilde{w}_p)(\tilde{w}_p.\text{some}_{\text{fv}(C_p)}; \llbracket C_p \rrbracket_{\tilde{w}_p} | \\ &\quad \prod_{\tilde{y}_{p_i} \in \text{PER}(\tilde{y}_p)} \dots (\nu \tilde{w}_1)(\tilde{w}_1.\text{some}_{\text{fv}(C_1)}; \llbracket C_1 \rrbracket_{\tilde{w}_1} | \\ &\quad \prod_{\tilde{y}_{1_i} \in \text{PER}(\tilde{y}_1)} \llbracket \lambda x.(M'[\tilde{x} \leftarrow x]) \rrbracket_v \{\tilde{w}_1/\tilde{y}_{1_i}\} \dots \{\tilde{w}_p/\tilde{y}_{p_i}\}) \\ &\quad | !z_1^!(z_1); \llbracket U_1 \rrbracket_{z_1} \dots | !z_q^!(z_q); \llbracket U_q \rrbracket_{z_q}) \end{aligned}$$

Finally we shall simplify the process to become:

$$\llbracket M \rrbracket_v = (\nu \tilde{z}, \tilde{w})(\prod_{i \in I} \llbracket \lambda x.(M'[\tilde{x} \leftarrow x]) \rrbracket_v \{\tilde{w}/\tilde{y}_i\} | Q'')$$

With this shape for M , we then have the following:

$$\begin{aligned} \llbracket N \rrbracket_u &= \llbracket M(C \star U) \rrbracket_u = (\nu v)(\llbracket M \rrbracket_v | v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x]; (\llbracket C \star U \rrbracket_x | [v \leftrightarrow u])) \\ &= (\nu v)((\nu \tilde{z}, \tilde{w})(\prod_{i \in I} \llbracket \lambda x.(M'[\tilde{x} \leftarrow x]) \rrbracket_v \{\tilde{w}/\tilde{y}_i\} | Q'') | v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x]; (\llbracket C \star U \rrbracket_x | [v \leftrightarrow u])) \\ &= (\nu v)((\nu \tilde{z}, \tilde{w})(\prod_{i \in I} \bar{v}.\text{some}; v(x); \bar{x}.\text{some}; x(x^\ell); x(x^1); x(); \llbracket M[\tilde{x} \leftarrow x] \rrbracket_v \{\tilde{w}/\tilde{y}_i\} | Q'')) \\ &\quad | v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x]; (\llbracket C \star U \rrbracket_x | [v \leftrightarrow u])) \\ &\rightsquigarrow (\nu v)((\nu \tilde{z}, \tilde{w})(\prod_{i \in I} v(x); \bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M[\tilde{x} \leftarrow x] \rrbracket_v \{\tilde{w}/\tilde{y}_i\} | Q'')) &= Q_1 \\ &\quad | \bar{v}[x]; (\llbracket C \star U \rrbracket_x | [v \leftrightarrow u])) \\ &\rightsquigarrow (\nu v)([v \leftrightarrow u] | (\nu x)((\nu \tilde{z}, \tilde{w})(\prod_{i \in I} \bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M[\tilde{x} \leftarrow x] \rrbracket_v \{\tilde{w}/\tilde{y}_i\} | Q'')) &= Q_2 \\ &\quad | \llbracket C \star U \rrbracket_x)) \\ &\rightsquigarrow (\nu x)((\nu \tilde{z}, \tilde{w})(\prod_{i \in I} \bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M[\tilde{x} \leftarrow x] \rrbracket_u \{\tilde{w}/\tilde{y}_i\} | Q'') | \llbracket C \star U \rrbracket_x) &= Q_3 \end{aligned}$$

We also have that

$$\begin{aligned} N &= (\lambda x.(M'[\tilde{x} \leftarrow x])) \langle C_1/\tilde{y}_1 \rangle \dots \langle C_p/\tilde{y}_p \rangle \llbracket U_1/z_1^! \rrbracket \dots \llbracket U_q/z_q^! \rrbracket (C \star U) \\ &\equiv_\lambda (\lambda x.(M'[\tilde{x} \leftarrow x])) (C \star U) \langle C_1/\tilde{y}_1 \rangle \dots \langle C_p/\tilde{y}_p \rangle \llbracket U_1/z_1^! \rrbracket \dots \llbracket U_q/z_q^! \rrbracket \\ &\longrightarrow M'[\tilde{x} \leftarrow x] \langle (C \star U)/x \rangle \langle C_1/\tilde{y}_1 \rangle \dots \langle C_p/\tilde{y}_p \rangle \llbracket U_1/z_1^! \rrbracket \dots \llbracket U_q/z_q^! \rrbracket = M \end{aligned}$$

Furthermore, we have:

$$\begin{aligned} \llbracket M \rrbracket_u &= \llbracket M'[\tilde{x} \leftarrow x] \langle (C \star U)/x \rangle \langle C_1/\tilde{y}_1 \rangle \dots \langle C_p/\tilde{y}_p \rangle \llbracket U_1/z_1^! \rrbracket \dots \llbracket U_q/z_q^! \rrbracket \rrbracket_u \\ &= (\nu x)((\nu \tilde{z}, \tilde{w})(\prod_{i \in I} \bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M[\tilde{x} \leftarrow x] \rrbracket_u \{\tilde{w}/\tilde{y}_i\} | Q'')) \\ &\quad | \llbracket C \star U \rrbracket_x \end{aligned}$$

We consider different possibilities for $n \geq 1$; in all the cases, the result follows.

When $n = 1$: We have $Q = Q_1$, $\llbracket N \rrbracket_u \rightsquigarrow^1 Q_1$. We also have that

- $Q_1 \rightsquigarrow^2 Q_3 = Q'$,
- $N \longrightarrow^1 M'[\tilde{x} \leftarrow x] \langle (C \star U)/x \rangle \langle C_1/\tilde{y}_1 \rangle \dots \langle C_p/\tilde{y}_p \rangle \llbracket U_1/z_1^! \rrbracket \dots \llbracket U_q/z_q^! \rrbracket = N'$
- and $\llbracket M'[\tilde{x} \leftarrow x] \langle (C \star U)/x \rangle \langle C_1/\tilde{y}_1 \rangle \dots \langle C_p/\tilde{y}_p \rangle \llbracket U_1/z_1^! \rrbracket \dots \llbracket U_q/z_q^! \rrbracket \rrbracket_u = Q_3$.

When $n = 2$: the analysis is similar.

When $n \geq 3$: We have $\llbracket N \rrbracket_u \rightsquigarrow^3 Q_3 \rightsquigarrow^l Q$, for $l \geq 0$. We also know that $N \longrightarrow M$, $Q_3 = \llbracket M \rrbracket_u$. By the IH, there exist Q', N' such that $Q \rightsquigarrow^i Q'$, $M \longrightarrow^j N'$ and $\llbracket N' \rrbracket_u = Q'$. Finally, $\llbracket N \rrbracket_u \rightsquigarrow^3 Q_3 \rightsquigarrow^l Q \rightsquigarrow^i Q'$ and $N \rightarrow M \longrightarrow^j N'$.

B. $M = \text{fail}^{\tilde{z}}$.

Then, $\llbracket M \rrbracket_v = \llbracket \text{fail}^{\tilde{z}} \rrbracket_v = \bar{v}.\text{none} | \tilde{z}.\text{none}$. With this shape for M , we have:

$$\begin{aligned} \llbracket N \rrbracket_u &= \llbracket (M (C \star U)) \rrbracket_u = (\nu v)(\llbracket M \rrbracket_v | v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x]; (\llbracket C \star U \rrbracket_x | [v \leftrightarrow u])) \\ &= (\nu v)(\bar{v}.\text{none} | \tilde{z}.\text{none} | v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x]; (\llbracket C \star U \rrbracket_x | [v \leftrightarrow u])) \\ &\rightsquigarrow \tilde{z}.\text{none} | \bar{u}.\text{none} | \overline{\text{fv}(C)}.\text{none} \end{aligned}$$

We also have that $N = \text{fail}^{\tilde{z}} C \star U \longrightarrow \text{fail}^{\tilde{z} \cup \text{fv}(C)} = M$. Furthermore,

$$\llbracket M \rrbracket_u = \llbracket \text{fail}^{\tilde{z} \cup \text{fv}(C)} \rrbracket_u = \tilde{z}.\text{none} | \bar{u}.\text{none} | \overline{\text{fv}(C)}.\text{none}$$

ii. When $m \geq 1$ and $n \geq 0$, we distinguish two cases:

A. When $n = 0$:

Then, $(\nu v)(R \mid v.\text{some}_{u,\text{fv}(C)}; \bar{v}[x]; (\llbracket C \star U \rrbracket_x \mid [v \leftrightarrow u])) = Q$ and $\llbracket M \rrbracket_u \rightsquigarrow^m R$ where $m \geq 1$. Then by the IH there exist R' and M' such that $R \rightsquigarrow^i R'$, $M \longrightarrow^j M'$, and $\llbracket M' \rrbracket_u = R'$. Hence we have that

$$\begin{aligned} \llbracket N \rrbracket_u &= (\nu v)(\llbracket M \rrbracket_v \mid v.\text{some}_{u,\text{fv}(C)}; \bar{v}[x]; (\llbracket C \star U \rrbracket_x \mid [v \leftrightarrow u])) \\ &\rightsquigarrow^m (\nu v)(R \mid v.\text{some}_{u,\text{fv}(C)}; \bar{v}[x]; (\llbracket C \star U \rrbracket_x \mid [v \leftrightarrow u])) = Q \end{aligned}$$

We also know that

$$Q \rightsquigarrow^i (\nu v)(R \mid v.\text{some}_{u,\text{fv}(C)}; \bar{v}[x]; (\llbracket C \star U \rrbracket_x \mid [v \leftrightarrow u])) = Q'$$

and so the λ_C term can reduce as follows: $N = (M (C \star U)) \longrightarrow^j M' (C \star U) = N'$ and $\llbracket N' \rrbracket_u = Q'$.

B. When $n \geq 1$:

Then R has an occurrence of an unguarded $\bar{v}.\text{some}$ or $\bar{v}.\text{none}$, hence it is of the form

$$\llbracket (\lambda x.(M'[\tilde{x} \leftarrow x]) \langle C_1/\tilde{y}_1 \rangle \cdots \langle C_p/\tilde{y}_p \rangle \llbracket U_1/z_1^! \rrbracket \cdots \llbracket U_q/z_q^! \rrbracket \rrbracket_v \text{ or } \llbracket \text{fail}^{\tilde{x}} \rrbracket_v. \text{ This case follows by IH.}$$

This concludes the analysis for the case $N = (M (C \star U))$.

3. $N = M[\tilde{x} \leftarrow x]$.

The sharing variable x is not free and the result follows by vacuity.

4. $N = M[\tilde{x} \leftarrow x] \langle \langle C \star U/x \rangle \rangle$. Then we have

$$\llbracket N \rrbracket_u = \llbracket M[\tilde{x} \leftarrow x] \langle \langle C \star U/x \rangle \rangle \rrbracket_u = (\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^!); x()); \llbracket M[\tilde{x} \leftarrow x] \rrbracket_u \mid \llbracket C \star U \rrbracket_x$$

Let us consider three cases.

(a) When $\text{size}(\tilde{x}) = \text{size}(C)$. Then let us consider the shape of the bag C .

i. When $C = 1$.

We have the following

$$\begin{aligned} \llbracket N \rrbracket_u &= (\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^!); x()); \llbracket M[\leftarrow x] \rrbracket_u \mid \llbracket 1 \star U \rrbracket_x \\ &= (\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^!); x()); \llbracket M[\leftarrow x] \rrbracket_u \mid \\ &\quad x.\text{some}_{\text{fv}(C)}; \bar{x}[x^\ell]; (\llbracket 1 \rrbracket_{x^\ell} \mid \bar{x}[x^!]; (!x^!(x_i); \llbracket U \rrbracket_{x_i} \mid \bar{x}[])) \\ &\rightsquigarrow (\nu x)(x(x^\ell); x(x^!); x()); \llbracket M[\leftarrow x] \rrbracket_u \mid \bar{x}[x^\ell]; (\llbracket 1 \rrbracket_{x^\ell} \mid \bar{x}[x^!]; (!x^!(x_i); \llbracket U \rrbracket_{x_i} \mid \bar{x}[])) &= Q_1 \\ &\rightsquigarrow (\nu x)(\bar{x}[x^!]; (!x^!(x_i); \llbracket U \rrbracket_{x_i} \mid \bar{x}[])) \mid (\nu x^\ell)(x(x^!); x()); \llbracket M[\leftarrow x] \rrbracket_u \mid \llbracket 1 \rrbracket_{x^\ell} &= Q_2 \\ &\rightsquigarrow (\nu x)(\bar{x}[] \mid (\nu x^!)(!x^!(x_i); \llbracket U \rrbracket_{x_i} \mid (\nu x^\ell)(x()); \llbracket M[\leftarrow x] \rrbracket_u \mid \llbracket 1 \rrbracket_{x^\ell})) &= Q_3 \\ &\rightsquigarrow (\nu x^!)(!x^!(x_i); \llbracket U \rrbracket_{x_i} \mid (\nu x^\ell)(\llbracket M[\leftarrow x] \rrbracket_u \mid \llbracket 1 \rrbracket_{x^\ell})) &= Q_4 \\ &= (\nu x^!)(!x^!(x_i); \llbracket U \rrbracket_{x_i} \mid (\nu x^\ell)(\bar{x}^\ell.\text{some}; \bar{x}^\ell[y_i]; (y_i.\text{some}_{u,\text{fv}(M)}; y_i()); \llbracket M \rrbracket_u \mid \bar{x}^\ell.\text{none}) \mid \\ &\quad x^\ell.\text{some}_\emptyset; x(y_i); (\bar{y}_i.\text{some}; \bar{y}_i[] \mid x^\ell.\text{some}_\emptyset; \bar{x}^\ell.\text{none}))) \\ &\rightsquigarrow (\nu x^!)(!x^!(x_i); \llbracket U \rrbracket_{x_i} \mid (\nu x^\ell)(\bar{x}^\ell[y_i]; (y_i.\text{some}_{u,\text{fv}(M)}; y_i()); \llbracket M \rrbracket_u \mid \bar{x}^\ell.\text{none}) \mid \\ &\quad x(y_i); (\bar{y}_i.\text{some}; \bar{y}_i[] \mid x^\ell.\text{some}_\emptyset; \bar{x}^\ell.\text{none}))) &= Q_5 \\ &\rightsquigarrow (\nu x^!)(!x^!(x_i); \llbracket U \rrbracket_{x_i} \mid (\nu x^\ell)(\bar{x}^\ell.\text{none} \mid (\nu y_i)(y_i.\text{some}_{u,\text{fv}(M)}; y_i()); \llbracket M \rrbracket_u \mid \\ &\quad \bar{y}_i.\text{some}; \bar{y}_i[] \mid x^\ell.\text{some}_\emptyset; \bar{x}^\ell.\text{none}))) &= Q_6 \\ &\rightsquigarrow (\nu x^!)(!x^!(x_i); \llbracket U \rrbracket_{x_i} \mid (\nu y_i)(y_i.\text{some}_{u,\text{fv}(M)}; y_i()); \llbracket M \rrbracket_u \mid \bar{y}_i.\text{some}; \bar{y}_i[])) &= Q_7 \\ &\rightsquigarrow (\nu x^!)(!x^!(x_i); \llbracket U \rrbracket_{x_i} \mid (\nu y_i)(y_i()); \llbracket M \rrbracket_u \mid \bar{y}_i[])) &= Q_8 \\ &\rightsquigarrow (\nu x^!)(!x^!(x_i); \llbracket U \rrbracket_{x_i} \mid \llbracket M \rrbracket_u) = \llbracket M \llbracket U/x^! \rrbracket \rrbracket_u &= Q_9 \end{aligned}$$

Notice how Q_6 has a choice however the x^ℓ name can be closed at any time so for simplicity we perform communication across this name first followed by all other communications that can take place.

Now we proceed by induction on the number of reductions $\llbracket N \rrbracket_u \rightsquigarrow^k Q$.

A. When $k = 0$, the result follows trivially. Just take $N = N'$ and $\llbracket N \rrbracket_u = Q = Q'$.

B. When $k = 1$.

We have $Q = Q_1$, $\llbracket N \rrbracket_u \rightsquigarrow^1 Q_1$ We also have that $Q_1 \rightsquigarrow^8 Q_9 = Q'$, $N \longrightarrow M \llbracket U/x^! \rrbracket = N'$ and $\llbracket N' \rrbracket_u = Q_9$

C. When $2 \leq k \leq 8$.

Proceeds similarly to the previous case

D. When $k \geq 9$.

We have $\llbracket N \rrbracket_u \rightsquigarrow^9 Q_9 \rightsquigarrow^l Q$, for $l \geq 0$. Since $Q_9 = \llbracket M \llbracket U/x^1 \rrbracket \rrbracket_u$ we apply the induction hypothesis we have that there exist Q', N' s.t. $Q \rightsquigarrow^i Q'$, $M \llbracket U/x^1 \rrbracket \longrightarrow^j N'$ and $\llbracket N' \rrbracket_u = Q'$. Then, $\llbracket N \rrbracket_u \rightsquigarrow^5 Q_5 \rightsquigarrow^l Q \rightsquigarrow^i Q'$ and by the contextual reduction rule it follows that $N = (M[\leftarrow x]) \langle \langle 1/x \rangle \rangle \longrightarrow M \llbracket U/x^1 \rrbracket \longrightarrow^j N'$ and the case holds.

ii. When $C = \wr N_1 \wr \dots \wr N_l \wr$, for $l \geq 1$. Then,

$$\begin{aligned}
\llbracket N \rrbracket_u &= \llbracket M[\tilde{x} \leftarrow x] \langle \langle C \star U/x \rangle \rangle \rrbracket_u \\
&= (\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^1); x(); \llbracket M[\tilde{x} \leftarrow x] \rrbracket_u \mid \llbracket C \star U \rrbracket_x) \\
&\rightsquigarrow^4 (\nu x^1)(!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid (\nu x^\ell)(\llbracket M[\tilde{x} \leftarrow x] \rrbracket_u \mid \llbracket C \rrbracket_{x^\ell})) \\
&= (\nu x^1)(!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid (\nu x^\ell)(\bar{x}^\ell.\text{some}; \bar{x}^\ell[y_1]; (y_1.\text{some}_\emptyset; y_1()); \mathbf{0} \\
&\quad \mid \bar{x}^\ell.\text{some}; x^\ell.\text{some}_{u, \text{fv}(M) \setminus \tilde{x}}; \prod_{x_{i_1} \in \tilde{x}} x(x_{i_1}); \dots \bar{x}^\ell.\text{some}; \bar{x}^\ell[y_1]; (y_1.\text{some}_\emptyset; y_1()); \mathbf{0} \\
&\quad \mid \bar{x}^\ell.\text{some}; x^\ell.\text{some}_{u, \text{fv}(M) \setminus (\tilde{x} \setminus x_1, \dots, x_{i_{l-1}})}; \prod_{x_{i_l} \in (\tilde{x} \setminus x_1, \dots, x_{i_{l-1}})} x(x_{i_l}); \llbracket M[\leftarrow x] \rrbracket_u \dots) \\
&\quad \mid x^\ell.\text{some}_{\text{fv}(C)}; x(y_1); x^\ell.\text{some}_{y_1, \text{fv}(C)}; \bar{x}^\ell.\text{some}; \bar{x}^\ell[z_1]; \\
&\quad (z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \bar{y}_1.\text{none} \mid \dots \\
&\quad x^\ell.\text{some}_{\text{fv}(M_l)}; x(y_l); x^\ell.\text{some}_{y_l, \text{fv}(M_l)}; \bar{x}^\ell.\text{some}; \bar{x}^\ell[z_l]; \\
&\quad (z_l.\text{some}_{\text{fv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \bar{y}_l.\text{none} \mid \llbracket \mathbf{1} \rrbracket_{x^\ell} \dots)))
\end{aligned}$$

We shall now perform multiple non-committing reductions at once. Notice that non-determinism guards the same prefixes denying the use of the reduction $[\rightsquigarrow_{\succeq x}]$ hence denying the commitment of non-determinism.

$$\begin{aligned}
&\rightsquigarrow^{6l} (\nu x^1)(!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid (\nu x^\ell)(\llbracket \mathbf{1} \rrbracket_{x^\ell} \mid \\
&\quad (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_l)(z_l.\text{some}_{\text{fv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \\
&\quad \prod_{x_{i_1} \in \tilde{x}} \prod_{x_{i_l} \in (\tilde{x} \setminus x_1, \dots, x_{i_{l-1}})} \llbracket M[\leftarrow x] \rrbracket_u \{z_1/x_{i_1}\} \dots \{z_l/x_{i_l}\} \dots)) \\
&= (\nu x^1)(!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid (\nu x^\ell)(x^\ell.\text{some}_\emptyset; x(y_{l+1}); (\bar{y}_{l+1}.\text{some}; \bar{y}_{l+1}[] \mid x^\ell.\text{some}_\emptyset; \bar{x}^\ell.\text{none}) \\
&\quad (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_l)(z_l.\text{some}_{\text{fv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \\
&\quad \prod_{x_{i_1} \in \tilde{x}} \prod_{x_{i_l} \in (\tilde{x} \setminus x_1, \dots, x_{i_{l-1}})} \bar{x}^\ell.\text{some}; \\
&\quad \bar{x}^\ell[y_{l+1}]; (y_{l+1}.\text{some}_{u, \text{fv}(M)}; y_{l+1}()); \llbracket M \rrbracket_u \{z_1/x_{i_1}\} \dots \{z_l/x_{i_l}\} \mid \bar{x}^\ell.\text{none})) \dots)) \\
&\rightsquigarrow^5 (\nu x^1)(!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid \\
&\quad (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_l)(z_l.\text{some}_{\text{fv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \\
&\quad \prod_{x_{i_1} \in \tilde{x}} \prod_{x_{i_l} \in (\tilde{x} \setminus x_1, \dots, x_{i_{l-1}})} \llbracket M \rrbracket_u \{z_1/x_{i_1}\} \dots \{z_l/x_{i_l}\} \dots)) \\
&= \llbracket M \wr \wr M_1 \wr \dots \wr M_l \wr /x_1, \dots, x_l \rrbracket \llbracket U/x^1 \rrbracket \rrbracket_u = Q_{6l+9}
\end{aligned}$$

The proof follows by induction on the number of reductions $\llbracket N \rrbracket_u \rightsquigarrow^k Q$.

A. When $k = 0$, the result follows trivially. Just take $N = N'$ and $\llbracket N \rrbracket_u = Q = Q'$.

B. When $1 \leq k \leq 6l + 9$.

Let Q_k such that $\llbracket N \rrbracket_u \rightsquigarrow^k Q_k$. We also have that $Q_k \rightsquigarrow^{6l+9-k} Q_{6l+9} = Q'$,
 $N \longrightarrow^1 M \wr \wr M_1 \wr \dots \wr M_l \wr /x_1, \dots, x_l \rrbracket \llbracket U/x^1 \rrbracket = N'$ and
 $\llbracket M \wr \wr M_1 \wr \dots \wr M_l \wr /x_1, \dots, x_l \rrbracket \llbracket U/x^1 \rrbracket \rrbracket_u = Q_{6l+9}$.

C. When $k > 6l + 9$.

Then, $\llbracket N \rrbracket_u \rightsquigarrow^{6l+9} Q_{6l+9} \rightsquigarrow^n Q$ for $n \geq 1$. Also,
 $N \longrightarrow^1 M \wr \wr M_1 \wr \dots \wr M_l \wr /x_1, \dots, x_l \rrbracket \llbracket U/x^1 \rrbracket$ and

$Q_{6l+9} = \llbracket M \wr \wr M_1 \wr \dots \wr M_l \wr /x_1, \dots, x_l \rrbracket \llbracket U/x^1 \rrbracket \rrbracket_u$.

By the induction hypothesis, there exist Q' and N' such that $Q \rightsquigarrow^i Q'$,

$M \wr \wr M_1 \wr \dots \wr M_l \wr /x_1, \dots, x_l \rrbracket \llbracket U/x^1 \rrbracket \longrightarrow^j N'$ and $\llbracket N' \rrbracket_u = Q'$.

Finally, $\llbracket N \rrbracket_u \rightsquigarrow^{6l+9} Q_{6l+9} \rightsquigarrow^n Q \rightsquigarrow^i Q'$ and

$$N \longrightarrow M \wr \wr M_1 \wr \dots \wr M_l \wr /x_1, \dots, x_l \rrbracket \llbracket U/x^1 \rrbracket \longrightarrow^j N'.$$

(b) When $\text{size}(\tilde{x}) > \text{size}(C)$.

Then we have $N = M[x_1, \dots, x_k \leftarrow x] \langle\langle C \star U/x \rangle\rangle$ with $C = \lambda N_1 \int \dots \int N_l \int \quad k > l$. $N \longrightarrow \mathbf{fail}^{\tilde{z}} = M'$ and $\tilde{z} = (\mathbf{fv}(M) \setminus \{x_1, \dots, x_k\}) \cup \mathbf{fv}(C)$. On the one hand, we have: Hence $k = l + m$ for some $m \geq 1$

$$\begin{aligned}
 \llbracket N \rrbracket_u &= \llbracket M[x_1, \dots, x_k \leftarrow x] \langle\langle C \star U/x \rangle\rangle \rrbracket_u \\
 &= (\nu x)(\bar{x}.\mathbf{some}; x(x^\ell); x(x^1); x()); \llbracket M[\tilde{x} \leftarrow x] \rrbracket_u \mid \llbracket C \star U \rrbracket_x \\
 &= (\nu x)(\bar{x}.\mathbf{some}; x(x^\ell); x(x^1); x()); \llbracket M[\tilde{x} \leftarrow x] \rrbracket_u \mid \\
 &\quad x.\mathbf{some}_{\mathbf{fv}(C)}; \bar{x}[x^\ell]; (\llbracket C \rrbracket_{x^\ell} \mid \bar{x}[x^1]; (!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid \bar{x}[])) \\
 &\rightsquigarrow^4 (\nu x^1)((\nu x^\ell)(\llbracket M[\tilde{x} \leftarrow x] \rrbracket_u \mid \llbracket C \rrbracket_{x^\ell}) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
 &= (\nu x^1)((\nu x^\ell)(\bar{x}^\ell.\mathbf{some}; \bar{x}^\ell[y_1]; (y_1.\mathbf{some}_\emptyset; y_1()); \mathbf{0} \\
 &\quad \mid \bar{x}^\ell.\mathbf{some}; x^\ell.\mathbf{some}_{u, \mathbf{fv}(M) \setminus \tilde{x}}; \prod_{x_{i_1} \in \tilde{x}} x(x_{i_1}); \dots \\
 &\quad \bar{x}^\ell.\mathbf{some}; \bar{x}^\ell[y_k]; (y_k.\mathbf{some}_\emptyset; y_k()); \mathbf{0} \\
 &\quad \mid \bar{x}^\ell.\mathbf{some}; x^\ell.\mathbf{some}_{u, \mathbf{fv}(M) \setminus (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{k-1}})}; \prod_{x_{i_k} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{k-1}})} x(x_{i_k}); \llbracket M[\leftarrow x] \rrbracket_u) \mid \\
 &\quad x^\ell.\mathbf{some}_{\mathbf{fv}(C)}; x(y_1); x^\ell.\mathbf{some}_{y_1, \mathbf{fv}(C)}; \bar{x}^\ell.\mathbf{some}; \bar{x}^\ell[z_1]; \\
 &\quad (z_1.\mathbf{some}_{\mathbf{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \bar{y}_1.\mathbf{none} \mid \dots \mid x^\ell.\mathbf{some}_{\mathbf{fv}(C)}; x(y_l); x^\ell.\mathbf{some}_{y_l, \mathbf{fv}(M_l)}; \bar{x}^\ell.\mathbf{some}; \bar{x}^\ell[z_l]; \\
 &\quad (z_l.\mathbf{some}_{\mathbf{fv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \llbracket \mathbf{1} \rrbracket_{x^\ell} \mid \bar{y}_l.\mathbf{none}) \dots)) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i} \quad (:= Q_4)
 \end{aligned} \tag{18}$$

we reduce Q_4 arbitrarily synchronising along channels x^ℓ, y_1, \dots, y_l .

$$\begin{aligned}
& Q_4 \rightsquigarrow^{6l} (\nu x^!)((\nu x^\ell)(\llbracket \mathbf{1} \rrbracket_{x^\ell} \mid (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_l)(z_l.\text{some}_{\text{fv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \\
& \quad \parallel_{x_{i_1} \in \tilde{x}} \cdots \parallel_{x_{i_l} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{l-1}})} \overline{x^\ell}.\text{some}; \overline{x^\ell}[y_{l+1}]; (y_{l+1}.\text{some}_\emptyset; y_{l+1}()); \mathbf{0} \\
& \quad \mid \overline{x^\ell}.\text{some}; x^\ell.\text{some}_{u, \text{fv}(M)}(\tilde{x} \setminus x_{i_1}, \dots, x_{i_l}); \parallel_{x_{i_{l+1}} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_l})} x(x_{i_{l+1}}); \cdots \\
& \quad \overline{x^\ell}.\text{some}; \overline{x^\ell}[y_k]; (y_k.\text{some}_\emptyset; y_k()); \mathbf{0} \\
& \quad \mid \overline{x^\ell}.\text{some}; x^\ell.\text{some}_{u, \text{fv}(M)}(\tilde{x} \setminus x_{i_1}, \dots, x_{i_{k-1}}); \parallel_{x_{i_k} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{k-1}})} x(x_{i_k}); \\
& \quad \llbracket M[\leftarrow x] \rrbracket_u \{z_1/x_{i_1}\} \cdots \{z_l/x_{i_l}\} \cdots)) \mid !x^!(x_i); \llbracket U \rrbracket_{x_i} \\
& = (\nu x^!)((\nu x^\ell)(x^\ell.\text{some}_\emptyset; x(y_{l+1}); (\overline{y_{l+1}}.\text{some}; \overline{y_{l+1}}[] \mid x^\ell.\text{some}_\emptyset; \overline{x^\ell}.\text{none}) \\
& \quad \mid (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_l)(z_l.\text{some}_{\text{fv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \\
& \quad \parallel_{x_{i_1} \in \tilde{x}} \cdots \parallel_{x_{i_l} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{l-1}})} \overline{x^\ell}.\text{some}; \overline{x^\ell}[y_{l+1}]; (y_{l+1}.\text{some}_\emptyset; y_{l+1}()); \mathbf{0} \\
& \quad \mid \overline{x^\ell}.\text{some}; x^\ell.\text{some}_{u, \text{fv}(M)}(\tilde{x} \setminus x_{i_1}, \dots, x_{i_l}); \parallel_{x_{i_{l+1}} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_l})} x(x_{i_{l+1}}); \cdots \\
& \quad \overline{x^\ell}.\text{some}; \overline{x^\ell}[y_k]; (y_k.\text{some}_\emptyset; y_k()); \mathbf{0} \\
& \quad \mid \overline{x^\ell}.\text{some}; x^\ell.\text{some}_{u, \text{fv}(M)}(\tilde{x} \setminus x_{i_1}, \dots, x_{i_{k-1}}); \parallel_{x_{i_k} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{k-1}})} x(x_{i_k}); \\
& \quad \llbracket M[\leftarrow x] \rrbracket_u \{z_1/x_{i_1}\} \cdots \{z_l/x_{i_l}\} \cdots)) \mid !x^!(x_i); \llbracket U \rrbracket_{x_i} \\
& \rightsquigarrow^5 (\nu x^!)((\nu x^\ell)(\overline{x^\ell}.\text{none} \mid (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_l)(z_l.\text{some}_{\text{fv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \\
& \quad \parallel_{x_{i_1} \in \tilde{x}} \cdots \parallel_{x_{i_l} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{l-1}})} \overline{x^\ell}.\text{some}_{u, \text{fv}(M)}(\tilde{x} \setminus x_{i_1}, \dots, x_{i_l}); \parallel_{x_{i_{l+1}} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_l})} x(x_{i_{l+1}}); \cdots \\
& \quad \overline{x^\ell}.\text{some}; \overline{x^\ell}[y_k]; (y_k.\text{some}_\emptyset; y_k()); \mathbf{0} \\
& \quad \mid \overline{x^\ell}.\text{some}; x^\ell.\text{some}_{u, \text{fv}(M)}(\tilde{x} \setminus x_{i_1}, \dots, x_{i_{k-1}}); \parallel_{x_{i_k} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{k-1}})} x(x_{i_k}); \\
& \quad \llbracket M[\leftarrow x] \rrbracket_u \{z_1/x_{i_1}\} \cdots \{z_l/x_{i_l}\} \cdots)) \mid !x^!(x_i); \llbracket U \rrbracket_{x_i} \\
& \rightsquigarrow (\nu x^!)((\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_l)(z_l.\text{some}_{\text{fv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \\
& \quad \parallel_{x_{i_1} \in \tilde{x}} \cdots \parallel_{x_{i_l} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{l-1}})} \overline{u}.\text{none} \mid (\text{fv}(M) \setminus \tilde{x}).\text{none} \mid \overline{(z_1, \dots, z_l)}.\text{none}) \cdots) \mid \\
& \quad !x^!(x_i); \llbracket U \rrbracket_{x_i} \\
& \rightsquigarrow^l (\nu x^!)(\parallel_{x_{i_1} \in \tilde{x}} \cdots \parallel_{x_{i_l} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{l-1}})} \overline{u}.\text{none} \mid \overline{(\text{fv}(M) \setminus \tilde{x})}.\text{none} \mid \overline{\text{fv}(C)}.\text{none} \mid \\
& \quad !x^!(x_i); \llbracket U \rrbracket_{x_i} \\
& \equiv \overline{u}.\text{none} \mid \overline{(\text{fv}(M) \setminus \tilde{x})}.\text{none} \mid \overline{\text{fv}(C)}.\text{none} = \llbracket \text{fail}^{\tilde{x}} \rrbracket_u = Q_{7l+10}
\end{aligned}$$

The rest of the proof is by induction on the number of reductions $\llbracket N \rrbracket_u \rightsquigarrow^j Q$.

- i. When $j = 0$, the result follows trivially. Just take $N = N'$ and $\llbracket N \rrbracket_u = Q = Q'$.
- ii. When $1 \leq j \leq 7l + 10$.

Let Q_j be such that $\llbracket N \rrbracket_u \rightsquigarrow^j Q_j$. By the steps above one has

$$Q_j \rightsquigarrow^{7l+10-j} Q_{7l+6} = Q',$$

$$N \longrightarrow^1 \text{fail}^{\tilde{x}} = N'; \text{ and } \llbracket \text{fail}^{\tilde{x}} \rrbracket_u = Q_{7l+10}.$$

- iii. When $j > 7l + 10$.

In this case, we have $\llbracket N \rrbracket_u \rightsquigarrow^{7l+10} Q_{7l+10} \rightsquigarrow^n Q$, for $n \geq 1$. We also know that $N \rightsquigarrow^1 \text{fail}^{\tilde{x}}$. However no further reductions can be performed.

(c) When $\text{size}(\tilde{x}) < \text{size}(C)$, the proof proceeds similarly to the previous case.

5. $N = M \langle C / \tilde{x} \rangle$.

In this case let us consider $C = \{M_1\} \cdots \{M_k\}$,

$$\begin{aligned}
\llbracket M \langle C / \tilde{x} \rangle \rrbracket_u & = (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\text{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
& \quad \mid \parallel_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \parallel_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \llbracket M \rrbracket_u \{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\} \cdots)
\end{aligned}$$

Therefore,

$$\begin{aligned}
\llbracket N \rrbracket_u &= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \amalg_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \amalg_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \llbracket M \rrbracket_u \{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\}) \cdots) \\
&\rightsquigarrow^m (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \amalg_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \amalg_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} R\{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\}) \cdots) \\
&\rightsquigarrow^n Q,
\end{aligned}$$

for some process R . Where \rightsquigarrow^n is a reduction that initially synchronizes with $z_i.\mathbf{some}_{\text{fv}(M_i)}$ for some $i \leq k$ when $n \geq 1$, $n + m = k \geq 1$. Type preservation in $\text{st}\pi^+$ ensures reducing $\llbracket M \rrbracket_v \rightsquigarrow^m$ does not consume possible synchronizations with $\overline{z_i}.\mathbf{some}$, if they occur. Let us consider the possible sizes of both m and n .

(a) For $m = 0$ and $n \geq 1$.

We have that $R = \llbracket M \rrbracket_u$ as $\llbracket M \rrbracket_u \rightsquigarrow^0 \llbracket M \rrbracket_u$.

Notice that there are two possibilities of having an unguarded $\overline{z_i}.\mathbf{some}$ or $\overline{z_i}.\mathbf{none}$ without internal reductions:

i. $M = \mathbf{fail}^{\tilde{x}, \tilde{y}}$.

$$\begin{aligned}
\llbracket N \rrbracket_u &= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \amalg_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \amalg_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \llbracket M \rrbracket_u \{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\}) \cdots) \\
&= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \amalg_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \amalg_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \llbracket \mathbf{fail}^{\tilde{x}, \tilde{y}} \rrbracket_u \{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\}) \cdots) \\
&= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \amalg_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \amalg_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \overline{\tilde{x}}.\mathbf{none} \mid \overline{\tilde{y}}.\mathbf{none} \mid \overline{u}.\mathbf{none} \{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\} \\
&\quad) \cdots) \\
&\equiv (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \\
&\quad \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \mid \overline{z_1}.\mathbf{none} \mid \cdots \mid \overline{z_k}.\mathbf{none} \mid \overline{\tilde{y}}.\mathbf{none} \mid \overline{u}.\mathbf{none}) \cdots) \\
&\rightsquigarrow^k \overline{\text{fv}(C)}.\mathbf{none} \mid \overline{\tilde{y}}.\mathbf{none} \mid \overline{u}.\mathbf{none} = Q',
\end{aligned}$$

and no further reductions can be performed. We also have that $N \longrightarrow \mathbf{fail}^{\tilde{y} \cup \text{fv}(C)} = N'$ and $\llbracket \mathbf{fail}^{\tilde{y} \cup \text{fv}(C)} \rrbracket_u = Q'$.

ii. $\text{head}(M) = x_i$.

Then we have the following

$$\begin{aligned}
\llbracket N \rrbracket_u &= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \amalg_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \amalg_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} (\nu \tilde{y})(\llbracket x_i \rrbracket_j \mid P)\{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\}) \cdots) \\
&= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \amalg_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \amalg_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \\
&\quad (\nu \tilde{y})(\overline{\tilde{x}_i}.\mathbf{some}; [x_i \leftrightarrow j] \mid P)\{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\}) \cdots)
\end{aligned}$$

Notice that multiple reductions can take place along any of the channels z_l . Let us consider for simplicity that $i = l = k$.

$$\begin{aligned}
\llbracket N \rrbracket_u &= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \amalg_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \amalg_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} (\nu \tilde{y})(\overline{\tilde{x}_k}.\mathbf{some}; [x_k \leftrightarrow j] \mid P)\{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\}) \cdots)
\end{aligned}$$

We show the full process of the reduction in this case for correctness. Applying multiple reductions of the form \rightsquigarrow_ν we obtain:

$$\begin{array}{c}
(\nu z_k)(z_k.\text{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
| \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \dots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \rightsquigarrow_{z_k} R_k \\
(\nu \tilde{y})(\overline{x_k}.\text{some}; [x_k \leftrightarrow j] | P)\{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\} \\
\hline
[\rightsquigarrow_{\nu}] \vdots \\
(\nu z_2)(z_2.\text{some}_{\text{fv}(M_2)}; \llbracket M_2 \rrbracket_{z_2} | \\
\dots (\nu z_k)(z_k.\text{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
| \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \dots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \rightsquigarrow_{z_k} R_1 \\
(\nu \tilde{y})(\overline{x_k}.\text{some}; [x_k \leftrightarrow j] | P)\{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\} \dots) \\
\hline
[\rightsquigarrow_{\nu}] (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} | \\
\dots (\nu z_k)(z_k.\text{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
| \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \dots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \rightsquigarrow_{z_k} (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} | R_1) \\
(\nu \tilde{y})(\overline{x_k}.\text{some}; [x_k \leftrightarrow j] | P)\{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\} \dots)
\end{array}$$

where $R_i = (\nu z_i)(z_i.\text{some}_{\text{fv}(M_i)}; \llbracket M_i \rrbracket_{z_i} | R_{i+1})$ for $i < k$
Hence we wish to show the following reduction:

$$\begin{array}{c}
(\nu z_k)(z_k.\text{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
| \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \dots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \rightsquigarrow_{z_k} R_k \\
(\nu \tilde{y})(\overline{x_k}.\text{some}; [x_k \leftrightarrow j] | P)\{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\}
\end{array}$$

We apply the reduction rule

$$\frac{P \succeq_{z_k} P' \quad Q \succeq_{z_k} Q' \quad (\nu z_k)(P' | Q') \rightsquigarrow_{z_k} R_k}{(\nu z_k)(P | Q) \rightsquigarrow_{z_k} R_k}$$

Where we take $P = z_k.\text{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \succeq_{z_k} z_k.\text{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} = P'$ and Notice that when $x_{i_k} = x_k$ we have that the substitution $\{z_k/x_{i_k}\}$ takes place

$$\begin{aligned}
Q &= \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \dots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} (\nu \tilde{y})(\overline{x_k}.\text{some}; [x_k \leftrightarrow j] | P)\{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\} \\
&= \prod_{x_{i_1} \in \{x_1, \dots, x_{k-1}\}} \dots \prod_{x_{i_{k-1}} \in \{x_1, \dots, x_{k-1} \setminus x_{i_1}, \dots, x_{i_{k-2}}\}} (\nu \tilde{y})(\overline{x_k}.\text{some}; [z_k \leftrightarrow j] | P)\{z_1/x_{i_1}\} \dots \{z_{k-1}/x_{i_{k-1}}\} \\
&\quad \# \prod_{x_{i_k} \in \{x_1, \dots, x_{k-1}\}} \prod_{x_{i_1} \in \{x_1, \dots, x_k \setminus x_{i_k}\}} \dots \prod_{x_{i_{k-1}} \in \{x_1, \dots, x_{k-1} \setminus x_{i_1}, \dots, x_{i_{k-2}}, x_{i_k}\}} \\
&\quad (\nu \tilde{y})(\overline{x_k}.\text{some}; [x_k \leftrightarrow j] | P)\{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\} \\
&\succeq_{z_k} \prod_{x_{i_1} \in \{x_1, \dots, x_{k-1}\}} \dots \prod_{x_{i_{k-1}} \in \{x_1, \dots, x_{k-1} \setminus x_{i_1}, \dots, x_{i_{k-2}}\}} (\nu \tilde{y})(\overline{x_k}.\text{some}; [z_k \leftrightarrow j] | P)\{z_1/x_{i_1}\} \dots \{z_{k-1}/x_{i_{k-1}}\} \\
&= Q'
\end{aligned}$$

Hence we wish to show the following reduction $(\nu z_k)(z_k.\text{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} | Q') \rightsquigarrow_{z_k} R_k$ We do this via the rule $[\rightsquigarrow_{\text{some}}]$ to obtain

$$R_k = (\nu z_k) \left(\llbracket M_k \rrbracket_{z_k} \mid \prod_{x_{i_1} \in \{x_1, \dots, x_{k-1}\}} \dots \prod_{x_{i_{k-1}} \in \{x_1, \dots, x_{k-1} \setminus x_{i_1}, \dots, x_{i_{k-2}}\}} (\nu \tilde{y})([z_k \leftrightarrow j] | P)\{z_1/x_{i_1}\} \dots \{z_{k-1}/x_{i_{k-1}}\} \right)$$

Hence we continue with the following reductions:

$$\begin{aligned}
&\rightsquigarrow (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_k)(\llbracket M_k \rrbracket_{z_k} \\
&\quad | \prod_{x_{i_1} \in \{x_1, \dots, x_{k-1}\}} \dots \prod_{x_{i_{k-1}} \in \{x_1, \dots, x_{k-1} \setminus x_{i_1}, \dots, x_{i_{k-2}}\}} \\
&\quad (\nu \tilde{y})([x_k \leftrightarrow j] | P)\{z_1/x_{i_1}\} \dots \{z_{k-1}/x_{i_{k-1}}\} \dots) = Q_1 \\
&\rightsquigarrow (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_{k-1})(\llbracket M_{k-1} \rrbracket_{z_{k-1}} \\
&\quad | \prod_{x_{i_1} \in \{x_1, \dots, x_{k-1}\}} \dots \prod_{x_{i_{k-1}} \in \{x_1, \dots, x_{k-1} \setminus x_{i_1}, \dots, x_{i_{k-2}}\}} \\
&\quad (\nu \tilde{y})(\llbracket M_k \rrbracket_j | P)\{z_1/x_{i_1}\} \dots \{z_{k-1}/x_{i_{k-1}}\} \dots) = Q_2
\end{aligned}$$

In addition, $N = M\langle C/\tilde{x} \rangle \longrightarrow M\{M_k/x_k\}\langle C \setminus M_k/\tilde{x} \setminus x_k \rangle = M'$. Finally,

$$\begin{aligned} \llbracket M' \rrbracket_u &= \llbracket M\{M_k/x_k\}\langle C \setminus M_k/\tilde{x} \setminus x_k \rangle \rrbracket_u \\ &= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_{k-1})(\llbracket M_{k-1} \rrbracket_{z_{k-1}} \\ &\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_{k-1}\}} \cdots \prod_{x_{i_{k-1}} \in \{x_1, \dots, x_{k-1} \setminus x_{i_1}, \dots, x_{i_{k-2}}\}} \\ &\quad (\nu \tilde{y})(\llbracket M_k \rrbracket_j \mid P)\{z_1/x_{i_1}\} \cdots \{z_{k-1}/x_{i_{k-1}}\}) \cdots) = Q_2 \end{aligned}$$

A. When $n = 1$:

Then, $Q = Q_1$ and $\llbracket N \rrbracket_u \rightsquigarrow^1 Q_1$. Also,
 $Q_1 \rightsquigarrow^1 Q_2 = Q'$, $N \longrightarrow^1 M\{M_k/x_k\}\langle C \setminus M_k/\tilde{x} \setminus x_k \rangle = N'$ and
 $\llbracket M\{M_k/x_k\}\langle C \setminus M_k/\tilde{x} \setminus x_k \rangle \rrbracket_u = Q_2$.

B. When $n \geq 2$:

Then $\llbracket N \rrbracket_u \rightsquigarrow^2 Q_2 \rightsquigarrow^l Q$, for $l \geq 0$. Also, $N \longrightarrow M'$, $Q_2 = \llbracket M' \rrbracket_u$. By the induction hypothesis, there exist Q' and N' such that $Q \rightsquigarrow^i Q'$, $M' \longrightarrow^j N'$ and $\llbracket N' \rrbracket_u = Q'$. Finally, $\llbracket N \rrbracket_u \rightsquigarrow^2 Q_2 \rightsquigarrow^l Q \rightsquigarrow^i Q'$ and $N \longrightarrow M' \longrightarrow^j N'$.

(b) For $m \geq 1$ and $n \geq 0$.

i. When $n = 0$.

Then

$$\begin{aligned} Q &= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\ &\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} R\{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\}) \cdots) \end{aligned}$$

and $\llbracket M \rrbracket_u \rightsquigarrow^m R$ where $m \geq 1$. By the IH there exist R' and M' such that $R \rightsquigarrow^i R'$, $M \longrightarrow^j M'$ and $\llbracket M' \rrbracket_u = R'$. Thus,

$$\begin{aligned} \llbracket N \rrbracket_u \rightsquigarrow^m (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\ \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} R\{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\}) \cdots) &= Q \\ \rightsquigarrow^i (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\ \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} R'\{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\}) \cdots) &= Q' \end{aligned}$$

Also, $N = M\langle C/\tilde{x} \rangle \longrightarrow^j M'\langle C/\tilde{x} \rangle = N'$ and $\llbracket N' \rrbracket_u = Q'$

ii. When $n \geq 1$. Then R has an occurrence of an unguarded $\overline{x_i}.\mathbf{some}$ or $\overline{x_i}.\mathbf{none}$, this case follows by IH.

6. $N = M\llbracket U/x^! \rrbracket$.

In this case, $\llbracket M\llbracket U/x^! \rrbracket \rrbracket_u = (\nu x^!)(\llbracket M \rrbracket_u \mid !x^!(x_i); \llbracket U \rrbracket_{x_i})$. Then,

$$\llbracket N \rrbracket_u = (\nu x^!)(\llbracket M \rrbracket_u \mid !x^!(x_i); \llbracket U \rrbracket_{x_i}) \rightsquigarrow^m (\nu x^!)(R \mid !x^!(x_i); \llbracket U \rrbracket_{x_i}) \rightsquigarrow^n Q.$$

for some process R . Where \rightsquigarrow^n is a reduction initially synchronises with $!x^!(x_i)$ when $n \geq 1$, $n + m = k \geq 1$. Type preservation in \mathfrak{S}^π ensures reducing $\llbracket M \rrbracket_v \rightsquigarrow^m$ doesn't consume possible synchronisations with $!x^!(x_i)$ if they occur. Let us consider the the possible sizes of both m and n .

(a) For $m = 0$ and $n \geq 1$.

In this case, $R = \llbracket M \rrbracket_u$ as $\llbracket M \rrbracket_u \longrightarrow^0 \llbracket M \rrbracket_u$.

Notice that the only possibility of having an unguarded $?x^![x_i]$ without internal reductions is when $\text{head}(M) = x[j]$ for some index j . Then we have the following:

$$\begin{aligned} \llbracket N \rrbracket_u &= (\nu x^!)((\nu \tilde{y})(\llbracket x[j] \rrbracket_k \mid P) \mid !x^!(x_i); \llbracket U \rrbracket_{x_i}) \\ &= (\nu x^!)((\nu \tilde{y})(?x^![x_i]; \overline{x_i}.j; [x_i \leftrightarrow k] \mid P) \mid !x^!(x_i); \llbracket U \rrbracket_{x_i}) \\ &\rightsquigarrow (\nu x^!)((\nu x_i)((\nu \tilde{y})(\overline{x_i}.j; [x_i \leftrightarrow k] \mid P) \mid \llbracket U \rrbracket_{x_i}) \mid !x^!(x_i); \llbracket U \rrbracket_{x_i}) &= Q_1 \\ &= (\nu x^!)((\nu x_i)((\nu \tilde{y})(\overline{x_i}.j; [x_i \leftrightarrow k] \mid P) \mid x_i.\mathbf{case}\{i : \llbracket U_i \rrbracket_{x_i}\}_{U_i \in U} \mid !x^!(x_i); \llbracket U \rrbracket_{x_i}) \\ &\rightsquigarrow (\nu x^!)((\nu x_i)((\nu \tilde{y})([x_i \leftrightarrow k] \mid P) \mid \llbracket U_j \rrbracket_{x_i}) \mid !x^!(x_i); \llbracket U \rrbracket_{x_i}) &= Q_2 \\ &\rightsquigarrow (\nu x^!)((\nu \tilde{y})(\llbracket U_j \rrbracket_k \mid P) \mid !x^!(x_i); \llbracket U \rrbracket_{x_i}) &= Q_3 \end{aligned}$$

We consider the two cases of the form of U_j and show that the choice of U_j is inconsequential

– When $U_j = \mathcal{N}^!$:

In this case, $N = M\llbracket U/x^! \rrbracket \longrightarrow M\{N/x^!\}\llbracket U/x^! \rrbracket = M'$. and

$$\llbracket M' \rrbracket_u = \llbracket M\{N/x^!\}\llbracket U/x^! \rrbracket \rrbracket_u = (\nu x^!)((\nu \tilde{y})(\llbracket N \rrbracket_k | P) | !x^!(x_i); \llbracket U \rrbracket_{x_i}) = Q_3$$

– When $U_i = 1^!$:

In this case, $N = M\llbracket U/x^! \rrbracket \longrightarrow M\{\mathbf{fail}^0/x^!\}\llbracket U/x^! \rrbracket = M'$.

Notice that $\llbracket 1^! \rrbracket_k = \bar{k}.\mathbf{none}$ and that $\llbracket \mathbf{fail}^0 \rrbracket_k = \bar{k}.\mathbf{none}$. In addition,

$$\begin{aligned} \llbracket M' \rrbracket_u &= \llbracket M\{\mathbf{fail}^0/x^!\}\llbracket U/x^! \rrbracket \rrbracket_u = (\nu x^!)((\nu \tilde{y})(\llbracket \mathbf{fail}^0 \rrbracket_k | P) | !x^!(x_i); \llbracket U \rrbracket_{x_i}) \\ &= (\nu x^!)((\nu \tilde{y})(\llbracket 1^! \rrbracket_k | P) | !x^!(x_i); \llbracket U \rrbracket_{x_i}) = Q_3 \end{aligned}$$

Both choices give an M that are equivalent to Q_3 .

i. When $n \leq 2$.

In this case, $Q = Q_n$ and $\llbracket N \rrbracket_u \rightsquigarrow^n Q_n$.

Also, $Q_n \rightsquigarrow^{3-n} Q_3 = Q'$, $N \longrightarrow^1 M' = N'$ and $\llbracket M' \rrbracket_u = Q_2$.

ii. When $n \geq 3$.

We have $\llbracket N \rrbracket_u \rightsquigarrow^3 Q_3 \rightsquigarrow^l Q$ for $l \geq 0$. We also know that $N \rightarrow M'$, $Q_3 = \llbracket M' \rrbracket_u$. By the IH, there exist Q and N' such that $Q \rightsquigarrow^i Q'$, $M' \longrightarrow^j N'$ and $\llbracket N' \rrbracket_u = Q'$. Finally, $\llbracket N \rrbracket_u \rightsquigarrow^3 Q_3 \rightsquigarrow^l Q \rightsquigarrow^i Q'$ and $N \rightarrow M' \longrightarrow^j N'$.

(b) For $m \geq 1$ and $n \geq 0$.

i. When $n = 0$.

Then $(\nu x^!)(R | !x^!(x_i); \llbracket U \rrbracket_{x_i}) = Q$ and $\llbracket M \rrbracket_u \rightsquigarrow^m R$ where $m \geq 1$. By the IH there exist R' and M' such that $R \rightsquigarrow^i R'$, $M \longrightarrow^j M'$ and $\llbracket M' \rrbracket_u = R'$. Hence,

$$\llbracket N \rrbracket_u = (\nu x^!)(\llbracket M \rrbracket_u | !x^!(x_i); \llbracket U \rrbracket_{x_i}) \rightsquigarrow^m (\nu x^!)(R | !x^!(x_i); \llbracket U \rrbracket_{x_i}) = Q.$$

In addition, $Q \rightsquigarrow^i (\nu x^!)(R' | !x^!(x_i); \llbracket U \rrbracket_{x_i}) = Q$, and the term can reduce as follows: $N = M\llbracket U/x^! \rrbracket \longrightarrow^j M'\llbracket U/x^! \rrbracket = N'$ and $\llbracket N' \rrbracket_u = Q'$.

ii. When $n \geq 1$.

Then R has an occurrence of an unguarded $!x^!(x_i)$, and the case follows by IH. \square

G.4 Success Sensitivity

Proposition 4 (Preservation of Success). *For all closed $M \in \lambda_c$, the following hold:*

1. $\mathbf{head}(M) = \checkmark \implies \llbracket M \rrbracket = (\nu \tilde{x})(P | \checkmark)$
2. $\llbracket M \rrbracket_u = P | \checkmark \uparrow Q \implies \mathbf{head}(M) = \checkmark$

Proof. Proof of both cases by induction on the structure of M .

1. We only need to consider terms of the following form:

(a) $M = \checkmark$:

This case is immediate.

(b) $M = N (C \star U)$:

Then, $\mathbf{head}(N (C \star U)) = \mathbf{head}(N)$. If $\mathbf{head}(N) = \checkmark$, then

$$\llbracket M(C \star U) \rrbracket_u = (\nu v)(\llbracket M \rrbracket_v | v.\mathbf{some}_{u, \text{fv}(C)}; \bar{v}[x].([v \leftrightarrow u] | \llbracket C \star U \rrbracket_x)).$$

By the IH, \checkmark is unguarded in $\llbracket N \rrbracket_u$.

(c) $M = M' \langle C/x_1, \dots, x_k \rangle$

Then we have that $\mathbf{head}(M' \langle C/x_1, \dots, x_k \rangle) = \mathbf{head}(M') = \checkmark$. Then

$$\begin{aligned} \llbracket M' \langle C/x_1, \dots, x_k \rangle \rrbracket_u &= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} | \dots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\ &\quad | \uparrow_{x_{i_1} \in \{x_1, \dots, x_k\}} \dots \uparrow_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\} \dots) \end{aligned}$$

and by the IH \checkmark is unguarded in $\llbracket M' \rrbracket_u$ hence unguarded in every summand in $\llbracket M \langle C/x_1, \dots, x_k \rangle \rrbracket_u$.

(d) $M = M' \llbracket U/x^! \rrbracket$

Then we have that $\mathbf{head}(M' \llbracket U/x^! \rrbracket) = \mathbf{head}(M') = \checkmark$. Then $\llbracket M' \llbracket U/x^! \rrbracket \rrbracket_u = (\nu x^!)(\llbracket M' \rrbracket_u | !x^!(x_i); \llbracket U \rrbracket_{x_i})$ and by the IH \checkmark is unguarded in $\llbracket M' \rrbracket_u$.

2. We only need to consider terms of the following form:

(a) **Case** $M = \checkmark$:

Then, $\llbracket \checkmark \rrbracket_u = \checkmark$ which is an unguarded occurrence of \checkmark and that $\text{head}(\checkmark) = \checkmark$.

(b) **Case** $M = N(C \star U)$:

Then, $\llbracket N(C \star U) \rrbracket_u = (\nu v)(\llbracket N \rrbracket_v \mid v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x))$. The only occurrence of an unguarded \checkmark can occur is within $\llbracket N \rrbracket_v$. By the IH, $\text{head}(N) = \checkmark$ and finally $\text{head}(N(C \star U)) = \text{head}(N)$.

(c) **Case** $M = M' \langle C/x_1, \dots, x_k \rangle$:

Then:

$$\begin{aligned} \llbracket M' \langle C/x_1, \dots, x_k \rangle \rrbracket_u &= (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots \mid (\nu z_k)(z_k.\text{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\ &\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \dots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\} \dots) \end{aligned}$$

An unguarded occurrence of \checkmark can only occur within $\llbracket M' \rrbracket_u$. By the IH, $\text{head}(M') = \checkmark$ and hence $\text{head}(M' \langle C/x_1, \dots, x_k \rangle) = \text{head}(M')$.

(d) **Case** $M = M' \llbracket U/x^1 \rrbracket$: This case is analogous to the previous. \square

Theorem 22 (Success Sensitivity (Under \rightsquigarrow)). $M \Downarrow \checkmark$ if and only if $\llbracket M \rrbracket_u \Downarrow_{\checkmark}$ for well-formed closed terms M .

Proof. We proceed with the proof in two parts.

1. Suppose that $M \Downarrow \checkmark$. We will prove that $\llbracket M \rrbracket_u \Downarrow_{\checkmark}$.

By Def. 17, there exists M' such that $M \longrightarrow^* M'$ and $\text{head}(M') = \checkmark$. By completeness, if $M \longrightarrow M'$ then there exist Q, Q' such that $\llbracket M \rrbracket_u \equiv Q \rightsquigarrow^* Q'$ and $\llbracket M' \rrbracket_u \equiv Q'$.

We wish to show that there exists Q'' such that $Q' \rightsquigarrow^* Q''$ and Q'' has an unguarded occurrence of \checkmark .

By Proposition 4 (1) we have that $\text{head}(M') = \checkmark \implies \llbracket M' \rrbracket_u = (\nu \tilde{x})(P \mid \checkmark)$. Finally $\llbracket M' \rrbracket_u = (\nu \tilde{x})(P \mid \checkmark) \equiv Q'$. Hence Q reduces to a process that has an unguarded occurrence of \checkmark .

2. Suppose that $\llbracket M \rrbracket_u \Downarrow_{\checkmark}$. We will prove that $M \Downarrow \checkmark$.

By operational soundness we have that if $\llbracket M \rrbracket_u \rightsquigarrow^* Q$ then there exist M' and Q' such that: (i) $M \longrightarrow^* M'$ and (ii) $Q \rightsquigarrow^* Q'$ with $\llbracket M' \rrbracket_u \equiv Q'$.

Since $\llbracket M \rrbracket_u \rightsquigarrow^* Q$, and $Q = Q'' \mid \checkmark$ and $Q \rightsquigarrow^* Q'$ we must have that $Q'' \rightsquigarrow^* Q'''$ with $Q' = Q''' \mid \checkmark$. As $\llbracket M' \rrbracket_u \equiv Q'$ we have that $\llbracket M' \rrbracket_u =$. Finally applying Proposition 4 (2) we have that $\llbracket M' \rrbracket_u = Q''' \mid \checkmark$ is itself a term with unguarded \checkmark , then M is itself headed with \checkmark . \square

H Proof of Loose Correctness of the Translation under the Eager Semantics

H.1 Completeness

Theorem 9 (Loose Completeness (Under \longrightarrow)). If $N \longrightarrow M$ for a well-formed closed λ_C -term N , then there exists Q such that $\llbracket N \rrbracket_u \longrightarrow^* Q$ and $\llbracket M \rrbracket_u \succeq_{\dagger} Q$.

Proof. By induction on the reduction rule applied to infer $N \longrightarrow M$. We have five cases.

1. Case [RS : Beta]: Then $N = (\lambda x.(M'[\tilde{x} \leftarrow x]))B \longrightarrow (M'[\tilde{x} \leftarrow x])\langle B/x \rangle = M$, where $B = C \star U$. The result follows easily, since

$$\begin{aligned} \llbracket N \rrbracket_u &= (\nu v)(\llbracket \lambda x.(M'[\tilde{x} \leftarrow x]) \rrbracket_v \mid v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x)) \\ &= (\nu v)(\bar{v}.\text{some}; v(x); \bar{x}.\text{some}; x(x^\ell); x(x^1); x(); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_v \mid \\ &\quad v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x)) \\ &\longrightarrow (\nu v)(v(x); \bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_v \mid \bar{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x)) \\ &\longrightarrow (\nu x)((\nu v)(\bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_v \mid [v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x) \\ &\longrightarrow (\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u \mid \llbracket C \star U \rrbracket_x) = \llbracket M \rrbracket_u \end{aligned} \tag{19}$$

2. Case [RS : Ex-Sub]: Then $N = M'[x_1, \dots, x_k \leftarrow x] \langle C \star U/x \rangle$, with $C = \lambda M_1 \dots \lambda M_k$, $k \geq 0$ and $M' \neq \text{fail}^{\tilde{v}}$.

$$\begin{aligned} \llbracket N \rrbracket_u &= (\nu v)(\llbracket \lambda x.(M'[\tilde{x} \leftarrow x]) \rrbracket_v \mid v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x)) \\ &= (\nu v)(\bar{v}.\text{some}; v(x); \bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_v \mid \\ &\quad v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x)) \\ &\longrightarrow (\nu v)(v(x); \bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_v \mid \bar{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x)) \\ &\longrightarrow (\nu x)((\nu v)(\bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_v \mid [v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x) \\ &\longrightarrow (\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u \mid \llbracket C \star U \rrbracket_x) = \llbracket M \rrbracket_u \end{aligned} \tag{20}$$

3. Case [RS : Ex-Sub]: Then $N = M'[x_1, \dots, x_k \leftarrow x] \langle\langle C \star U/x \rangle\rangle$, with $C = \lceil M_1 \rceil \dots \lceil M_k \rceil$, $k \geq 0$ and $M' \neq \text{fail}^{\bar{y}}$.

The reduction is $N = M'[x_1, \dots, x_k \leftarrow x] \langle\langle C \star U/x \rangle\rangle \longrightarrow M' \langle\langle C/x_1, \dots, x_k \rangle\rangle \llbracket U/x^1 \rrbracket = M$.

We detail the translations of $\llbracket N \rrbracket_u$ and $\llbracket M \rrbracket_u$. To simplify the proof, we consider $k = 2$ (the case in which $k > 2$ follows analogously. Similarly the case of $k = 0, 1$ it contained within $k = 2$). On the one hand, we have:

$$\begin{aligned} \llbracket N \rrbracket_u &= \llbracket M'[x_1 \leftarrow x] \langle\langle C \star U/x \rangle\rangle \rrbracket_u \\ &= (\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[x_1, x_2 \leftarrow x] \rrbracket_u \mid \llbracket C \star U \rrbracket_x \\ &= (\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[x_1, x_2 \leftarrow x] \rrbracket_u \mid x.\text{some}_{\text{fv}(C)}; \bar{x}[x^\ell].(\llbracket C \rrbracket_{x^\ell} \\ &\quad \mid \bar{x}[x^1].(!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid \bar{x}[])) \quad (:= P_{\mathbb{N}}) \end{aligned} \quad (21)$$

The reduction is $N = M'[x_1, \dots, x_k \leftarrow x] \langle\langle C \star U/x \rangle\rangle \longrightarrow M' \langle\langle C/x_1, \dots, x_k \rangle\rangle \llbracket U/x^1 \rrbracket = M$.

We detail the translations of $\llbracket N \rrbracket_u$ and $\llbracket M \rrbracket_u$. To simplify the proof, we consider $k = 2$ (the case in which $k > 2$ follows analogously. Similarly the case of $k = 0, 1$ it contained within $k = 2$). On the one hand, we have:

$$\begin{aligned} \llbracket N \rrbracket_u &= \llbracket M'[x_1 \leftarrow x] \langle\langle C \star U/x \rangle\rangle \rrbracket_u \\ &= (\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[x_1, x_2 \leftarrow x] \rrbracket_u \mid \llbracket C \star U \rrbracket_x \\ &= (\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[x_1, x_2 \leftarrow x] \rrbracket_u \mid x.\text{some}_{\text{fv}(C)}; \bar{x}[x^\ell].(\llbracket C \rrbracket_{x^\ell} \\ &\quad \mid \bar{x}[x^1].(!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid \bar{x}[])) \quad (:= P_{\mathbb{N}}) \end{aligned} \quad (22)$$

$$\begin{aligned} P_{\mathbb{N}} &\longrightarrow^* (\nu x^1)((\nu x^\ell)(\llbracket M'[x_1, x_2 \leftarrow x] \rrbracket_u \mid \llbracket \lceil M_1 \rceil \cdot \lceil M_2 \rceil \rrbracket_{x^\ell}) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\ &= (\nu x^1)((\nu x^\ell)(\bar{x}^\ell.\text{some}; \bar{x}^\ell[y_1]; (y_1.\text{some}_\emptyset; y_1()); \mathbf{0} \\ &\quad \mid \bar{x}^\ell.\text{some}; x^\ell.\text{some}_{u, \text{fv}(M') \setminus \bar{x}}; \prod_{x_{i_1} \in x_1, x_2} x(x_{i_1}); \bar{x}^\ell.\text{some}; \bar{x}^\ell[y_2]; (y_2.\text{some}_\emptyset; y_2()); \mathbf{0} \\ &\quad \mid \bar{x}^\ell.\text{some}; x^\ell.\text{some}_{u, \text{fv}(M') \setminus \bar{x}}; \prod_{x_{i_2} \in (x_1, x_2 \setminus x_{i_1})} x(x_{i_2}); \\ &\quad \bar{x}^\ell.\text{some}; \bar{x}^\ell[y_3]; (y_3.\text{some}_{u, \text{fv}(M')}; y_3()); \llbracket M' \rrbracket_u \mid \bar{x}^\ell.\text{none})) \\ &\quad \mid \\ &\quad x^\ell.\text{some}_{\text{fv}(C)}; x(y_1); x^\ell.\text{some}_{y_1, \text{fv}(C)}; \bar{x}^\ell.\text{some}; \bar{x}^\ell[z_1]; \\ &\quad (z_1.\text{some}_{\text{fv}(M_j)}; \llbracket M_j \rrbracket_{z_1} \mid \bar{y}_1.\text{none}) \\ &\quad x^\ell.\text{some}_{\text{fv}(C)}; x(y_2); x^\ell.\text{some}_{y_2, \text{fv}(C)}; \bar{x}^\ell.\text{some}; \bar{x}^\ell[z_2]; \\ &\quad (z_2.\text{some}_{\text{fv}(M_j)}; \llbracket M_j \rrbracket_{z_2} \mid \bar{y}_2.\text{none}) \\ &\quad x^\ell.\text{some}_\emptyset; x(y_3); (\bar{y}_3.\text{some}; \bar{y}_3[] \mid x^\ell.\text{some}_\emptyset; \bar{x}^\ell.\text{none}))) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\ &\longrightarrow^* (\nu x^1)((\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid (\nu z_2)(z_2.\text{some}_{\text{fv}(M_2)}; \llbracket M_2 \rrbracket_{z_2} \mid \\ &\quad \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \{z_2/x_{i_2}\})) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \end{aligned}$$

Where x_{i_1}, x_{i_2} is an arbitrary permutation of x_1, x_2 . On the other hand, we have:

$$\begin{aligned} \llbracket M \rrbracket_u &= \llbracket M' \langle\langle M_1/x_1 \rangle\rangle \llbracket U/x^1 \rrbracket \rrbracket_u \\ &= (\nu x^1)((\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid (\nu z_2)(z_2.\text{some}_{\text{fv}(M_2)}; \llbracket M_2 \rrbracket_{z_2} \\ &\quad \mid \prod_{x_{i_1} \in \{x_1, x_2\}} \prod_{x_{i_2} \in \{x_1, x_2\} \setminus \{x_{i_1}\}} \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \{z_2/x_{i_2}\})) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\ &\succeq_{\neq} (\nu x^1)((\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \\ &\quad (\nu z_2)(z_2.\text{some}_{\text{fv}(M_2)}; \llbracket M_2 \rrbracket_{z_2} \mid \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \{z_2/x_{i_2}\})) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \end{aligned} \quad (23)$$

Therefore, by (22) and (23) the result follows.

4. Case $[\text{RS:Fetch}^\ell]$: Then, we have $N = M' \langle C/x_1, \dots, x_k \rangle$ with $\text{head}(M') = x_j$, $C = M_1, \dots, M_k$ and $N \longrightarrow M' \{M_i/x_j\} \langle (C \setminus M_i)/x_1, \dots, x_k \setminus x_j \rangle = M$, for some $M_i \in C$. On the one hand, we have:

$$\begin{aligned}
\llbracket N \rrbracket_u &= (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_k)(z_k.\text{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \amalg_{x_{i_1} \in \{x_1, \dots, x_k\}} \dots \amalg_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\} \dots) \\
&= (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_k)(z_k.\text{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \amalg_{x_{i_1} \in \{x_1, \dots, x_k\}} \dots \amalg_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} (\nu \tilde{y})(\llbracket x_j \rrbracket_v \mid P) \{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\} \dots) \quad (*) \quad (24) \\
&= (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_k)(z_k.\text{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \amalg_{x_{i_1} \in \{x_1, \dots, x_k\}} \dots \amalg_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \\
&\quad (\nu \tilde{y})(\overline{x_j}.\text{some}; [x_j \leftrightarrow v] \mid P) \{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\} \dots)
\end{aligned}$$

where (*) is inferred via Proposition 3. Let us consider the case when $x_{i_k} = x_j$ the other cases proceed similarly. Then, we have reduction:

$$\begin{aligned}
&\longrightarrow (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \\
&\quad \dots (\nu z_k)(\llbracket M_k \rrbracket_{z_k} \mid (\nu \tilde{y})([x_k \leftrightarrow v] \mid P) \{z_1/x_{i_1}\} \dots \{z_{k-1}/x_{i_{k-1}}\} \dots) \\
&\longrightarrow (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_{k-1})(z_{k-1}.\text{some}_{\text{fv}(M_{k-1})}; \llbracket M_{k-1} \rrbracket_{z_{k-1}} \mid \\
&\quad (\nu \tilde{y})(\llbracket M_k \rrbracket_v \mid P) \{z_1/x_{i_1}\} \dots \{z_{k-1}/x_{i_{k-1}}\} \dots)
\end{aligned} \quad (25)$$

for some permutation $x_{i_1}, \dots, x_{i_{k-1}}$ of the bag x_1, \dots, x_{k-1} . On the other hand, we have:

$$\begin{aligned}
\llbracket M \rrbracket_u &= \llbracket M' \{M_k/x_k\} \langle (C \setminus M_k)/x_1, \dots, x_{k-1} \rangle \rrbracket_u \\
&= (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_{k-1})(z_{k-1}.\text{some}_{\text{fv}(M_{k-1})}; \llbracket M_{k-1} \rrbracket_{z_{k-1}} \\
&\quad \mid \amalg_{x_{i_1} \in \{x_1, \dots, x_{k-1}\}} \dots \amalg_{x_{i_{k-1}} \in \{x_1, \dots, x_{k-1} \setminus x_{i_1}, \dots, x_{i_{k-2}}\}} \\
&\quad (\nu \tilde{y})(\llbracket M_k \rrbracket_v \mid P) \{z_1/x_{i_1}\} \dots \{z_{k-1}/x_{i_{k-1}}\} \dots) \\
&\succeq_{\amalg} (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_{k-1})(z_{k-1}.\text{some}_{\text{fv}(M_{k-1})}; \llbracket M_{k-1} \rrbracket_{z_{k-1}} \\
&\quad \mid (\nu \tilde{y})(\llbracket M_k \rrbracket_v \mid P) \{z_1/x_{i_1}\} \dots \{z_{k-1}/x_{i_{k-1}}\} \dots)
\end{aligned} \quad (26)$$

Therefore, by (25) and (26) the result follows.

5. Case $[\text{RS:Fetch}^!]$: Then we have $N = M' \llbracket U/x^! \rrbracket$ with $\text{head}(M') = x^![k]$, $U_k = \lambda N \mathcal{J}^!$ and $N \longrightarrow M' \{N/x^!\} \llbracket U/x^! \rrbracket = M$. The result follows easily from

$$\begin{aligned}
\llbracket N \rrbracket_u &= \llbracket M' \llbracket U/x^! \rrbracket \rrbracket_u = (\nu x^!)(\llbracket M' \rrbracket_u \mid !x^!(x_k); \llbracket U \rrbracket_{x_k}) \\
&\longrightarrow^* (\nu x^!)((\nu \tilde{y})(\llbracket x^![k] \rrbracket_j \mid P) \mid !x^!(x_k); \llbracket U \rrbracket_{x_k}) \quad (*) \\
&= (\nu x^!)((\nu \tilde{y})(?x^![x_k]; \overline{x_k}.k; [x_k \leftrightarrow j] \mid P) \mid !x^!(x_k); \llbracket U \rrbracket_{x_k}) \\
&\longrightarrow (\nu x^!)((\nu \tilde{y})(\nu x_k)(\overline{x_k}.k; [x_k \leftrightarrow j] \mid \llbracket U \rrbracket_{x_k}) \mid P) \mid !x^!(x_k); \llbracket U \rrbracket_{x_k} \\
&= (\nu x^!)((\nu \tilde{y})(\nu x_k)(\overline{x_k}.k; [x_k \leftrightarrow j] \mid x_k.\text{case}\{i : \llbracket U_i \rrbracket_x\}_{U_i \in U} \mid P) \mid !x^!(x_k); \llbracket U \rrbracket_{x_k}) \\
&\longrightarrow (\nu x^!)((\nu \tilde{y})(\llbracket \lambda N \mathcal{J}^! \rrbracket_j) \mid P) \mid !x^!(x_k); \llbracket U \rrbracket_{x_k} \\
&= (\nu x^!)((\nu \tilde{y})(\llbracket N \rrbracket_j) \mid P) \mid !x^!(x_k); \llbracket U \rrbracket_{x_k} = \llbracket M \rrbracket_u
\end{aligned} \quad (27)$$

where the reductions denoted by (*) are inferred via Proposition 3.

6. Cases $[\text{RS} : \text{TCont}]$: This case follows by IH.

7. Case $[\text{RS:Fail}^\ell]$: Then we have $N = M' [x_1, \dots, x_k \leftarrow x] \langle C \star U/x \rangle$ with $k \neq \text{size}(C)$ and $N \longrightarrow \text{fail}^{\tilde{y}} = M$, where $\tilde{y} = (\text{fv}(M') \setminus \{x_1, \dots, x_k\}) \cup \text{fv}(C)$. Let $\text{size}(C) = l$ and we assume that $k > l$ and we proceed similarly for $k > l$. Hence $k = l + m$ for some $m \geq 1$

$$\begin{aligned}
\llbracket N \rrbracket_u &= \llbracket M' [x_1, \dots, x_k \leftarrow x] \llbracket C \star U/x \rrbracket \rrbracket_u = (\nu x)(\overline{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M' [\tilde{x} \leftarrow x] \rrbracket_u \mid \llbracket C \star U \rrbracket_x \\
&= (\nu x)(\overline{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M' [\tilde{x} \leftarrow x] \rrbracket_u \mid \\
&\quad x.\text{some}_{\text{fv}(C)}; \overline{x}[x^\ell].(\llbracket C \rrbracket_{x^\ell} \mid \overline{x}[x^1].(!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid \overline{x}[])) \\
&\longrightarrow^* (\nu x^1)((\nu x^\ell)(\llbracket M' [\tilde{x} \leftarrow x] \rrbracket_u \mid \llbracket C \rrbracket_{x^\ell}) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&= (\nu x^1)((\nu x^\ell)(\overline{x}^\ell.\text{some}; \overline{x}^\ell[y_1]; (y_1.\text{some}_\emptyset; y_1()); \mathbf{0} \\
&\quad \mid \overline{x}^\ell.\text{some}; x^\ell.\text{some}_{u, \text{fv}(M') \setminus \tilde{x}}; \prod_{x_{i_1} \in \tilde{x}} x(x_{i_1}); \dots \overline{x}^\ell.\text{some}; \overline{x}^\ell[y_k]; (y_k.\text{some}_\emptyset; y_k()); \mathbf{0} \\
&\quad \mid \overline{x}^\ell.\text{some}; x^\ell.\text{some}_{u, \text{fv}(M') \setminus \tilde{x}}; \prod_{x_{i_k} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{k-1}})} x(x_{i_k}); \\
&\quad \overline{x}^\ell.\text{some}; \overline{x}^\ell[y_{k+1}]; (y_{k+1}.\text{some}_{u, \text{fv}(M')}; y_{k+1}()); \llbracket M' \rrbracket_u \mid \overline{x}^\ell.\text{none})) \dots) \\
&\quad \mid x^\ell.\text{some}_{\text{fv}(C)}; x(y_1); x^\ell.\text{some}_{y_1, \text{fv}(C)}; \overline{x}^\ell.\text{some}; \overline{x}^\ell[z_1]; \\
&\quad (z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \overline{y_1}.\text{none} \mid \\
&\quad \dots x^\ell.\text{some}_{\text{fv}(M_l)}; x(y_l); x^\ell.\text{some}_{y_l, \text{fv}(C)}; \overline{x}^\ell.\text{some}; \overline{x}^\ell[z_l]; \dots \\
&\quad (z_l.\text{some}_{\text{fv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \overline{y_l}.\text{none} \mid \\
&\quad x^\ell.\text{some}_\emptyset; x(y_{l+1}); (\overline{y_{l+1}}.\text{some}; \overline{y_{l+1}}[]) \mid x^\ell.\text{some}_\emptyset; \overline{x}^\ell.\text{none})) \dots) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i} \quad (:= P_{\mathbb{N}})
\end{aligned} \tag{28}$$

we reduce $P_{\mathbb{N}}$ arbitrarily discarding non-deterministic sums.

$$\begin{aligned}
P_{\mathbb{N}} &\longrightarrow^* (\nu x^1)((\nu x^\ell)((\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_k)(z_k.\text{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \mid \\
&\quad \overline{x}^\ell.\text{some}; \overline{x}^\ell[y_{k+1}]; (y_{k+1}.\text{some}_{u, \text{fv}(M')}; y_{k+1}()); \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\} \mid \overline{x}^\ell.\text{none})) \dots) \\
&\quad \mid x^\ell.\text{some}_{\text{fv}(C) \setminus (M_1, \dots, M_k)}; x(y_{k+1}); x^\ell.\text{some}_{y_{k+1}, \text{fv}(C) \setminus (M_1, \dots, M_k)}; \overline{x}^\ell.\text{some}; \overline{x}^\ell[z_{k+1}]; \\
&\quad (z_{k+1}.\text{some}_{\text{fv}(M_{k+1})}; \llbracket M_{k+1} \rrbracket_{z_{k+1}} \mid \overline{y_1}.\text{none} \mid \\
&\quad \dots x^\ell.\text{some}_{\text{fv}(M_l)}; x(y_l); x^\ell.\text{some}_{y_l, \text{fv}(M_l)}; \overline{x}^\ell.\text{some}; \overline{x}^\ell[z_l]; \dots \\
&\quad (z_l.\text{some}_{\text{fv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \overline{y_l}.\text{none} \mid \\
&\quad x^\ell.\text{some}_\emptyset; x(y_{l+1}); (\overline{y_{l+1}}.\text{some}; \overline{y_{l+1}}[]) \mid x^\ell.\text{some}_\emptyset; \overline{x}^\ell.\text{none})) \dots) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i} \\
&\longrightarrow^* (\nu x^1)((\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_k)(z_k.\text{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \mid \\
&\quad \overline{u}.\text{none} \mid \overline{(\text{fv}(M') \setminus \tilde{x})}.\text{none} \mid \overline{z_1}.\text{none} \mid \dots \mid \overline{z_k}.\text{none} \mid \overline{(\text{fv}(C) \setminus (M_1, \dots, M_k))}.\text{none})) \dots) \\
&\quad \mid !x^1(x_i); \llbracket U \rrbracket_{x_i} \\
&\longrightarrow^* (\nu x^1)((\overline{u}.\text{none} \mid \overline{(\text{fv}(M') \setminus \tilde{x})}.\text{none} \mid \overline{(\text{fv}(C))}.\text{none}) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&\equiv \overline{u}.\text{none} \mid \overline{(\text{fv}(M') \setminus \tilde{x})}.\text{none} \mid \overline{(\text{fv}(C))}.\text{none} = \llbracket M \rrbracket_u
\end{aligned}$$

8. Case [RS:Fail¹]: Then we have $N = M' \llbracket U/x^1 \rrbracket$ with $\text{head}(M') = x[i]$, $U_i = 1^1$ and $N \longrightarrow M' \{\text{fail}^\emptyset/x^1\} \llbracket U/x^1 \rrbracket$. The result follows easily from

$$\begin{aligned}
\llbracket N \rrbracket_u &= \llbracket M' \llbracket U/x^1 \rrbracket \rrbracket_u = (\nu x^1)(\llbracket M' \rrbracket_u \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&\longrightarrow^* (\nu x^1)((\nu \tilde{y})(\llbracket x[i] \rrbracket_j \mid P) \mid !x^1(x_k); \llbracket U \rrbracket_{x_k}) \quad (*) \\
&= (\nu x^1)((\nu \tilde{y})(?x^1[x_k]; \overline{x_k}.i; [x_k \leftrightarrow j] \mid P) \mid !x^1(x_k); \llbracket U \rrbracket_{x_k}) \quad (*) \\
&\longrightarrow (\nu x^1)((\nu \tilde{y})(\nu x_k)(x_k.l_i; [x_k \leftrightarrow j] \mid \llbracket U \rrbracket_{x_k} \mid P) \mid !x^1(x_k); \llbracket U \rrbracket_{x_k}) \\
&= (\nu x^1)((\nu \tilde{y})(\nu x_k)(x_k.l_i; [x_k \leftrightarrow j] \mid x_k.\text{case}(i. \llbracket U_i \rrbracket_x)) \mid P) \mid !x^1(x_k); \llbracket U \rrbracket_{x_k} \\
&\longrightarrow (\nu x^1)((\nu \tilde{y})(\llbracket 1^1 \rrbracket_j \mid P) \mid !x^1(x_k); \llbracket U \rrbracket_{x_k}) = (\nu x^1)((\nu \tilde{y})(\overline{j}.\text{none} \mid P) \mid !x^1(x_k); \llbracket U \rrbracket_{x_k}) = \llbracket M \rrbracket_u
\end{aligned} \tag{29}$$

9. Case [RS : Cons₁]: Then we have $N = \text{fail}^{\tilde{x}} C \star U$ and $N \longrightarrow \text{fail}^{\tilde{x} \cup \tilde{y}} M$ where $\tilde{y} = \text{fv}(C)$. The result follows easily from

$$\begin{aligned}
\llbracket N \rrbracket_u &= \llbracket \text{fail}^{\tilde{x}} C \star U \rrbracket_u = (\nu v)(\llbracket \text{fail}^{\tilde{x}} \rrbracket_v \mid v.\text{some}_{u, \text{fv}(C)}; \overline{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x)) \\
&= (\nu v)(\overline{v}.\text{none} \mid \overline{\tilde{x}}.\text{none} \mid v.\text{some}_{u, \text{fv}(C)}; \overline{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x)) \\
&\longrightarrow \overline{u}.\text{none} \mid \overline{\tilde{x}}.\text{none} \mid \overline{\tilde{y}}.\text{none} = \llbracket M \rrbracket_u
\end{aligned} \tag{30}$$

10. Cases [RS : Cons₂] and [RS : Cons₃]: These cases follow by IH similarly to Case 7.

11. Case [RS:Cons₄]: Then we have $N = \mathbf{fail}^{\tilde{y}}\llbracket U/x^1 \rrbracket$ and $N \longrightarrow \mathbf{fail}^{\tilde{y}} = M$. The result follows easily from

$$\begin{aligned} \llbracket N \rrbracket_u &= \llbracket \mathbf{fail}^{\tilde{y}}\llbracket U/x^1 \rrbracket \rrbracket_u = (\nu x^1)(\llbracket \mathbf{fail}^{\tilde{y}} \rrbracket_u \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\ &= (\nu x^1)(\bar{u}.\mathbf{none} \mid \bar{x}.\mathbf{none} \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \equiv \bar{u}.\mathbf{none} \mid \bar{x}.\mathbf{none} = \llbracket M \rrbracket_u \end{aligned} \quad (31) \quad \square$$

H.2 Soundness

We define $P \longrightarrow \{P_i\}_{i \in I}$, for a fixed finite set I where $I = \{i \text{ s.t. } P \longrightarrow P_i\}$. Similarly we define $P \longrightarrow^* \{P_i\}_{i \in I}$ to be defined inductively by $P \longrightarrow^* \{P_i\}_{i \in I}$ and $P_i \longrightarrow \{P_j\}_{j \in J_i}$ for each $i \in I$ then $P \longrightarrow^* \{P_j\}_{j \in J}$ with $J = \cup_{i \in I} J_i$

Proposition 2. *If $P \succeq_{\#} Q$ and $P \longrightarrow^* \{P_i\}_{i \in I}$, then there exist J and $\{Q_j\}_{j \in J}$ such that $Q \longrightarrow^* \{Q_j\}_{j \in J}$, $J \subseteq I$, and for each $j \in J$, $P_j \succeq_{\#} Q_j$.*

Proof. Proof by induction on the precongruence rules.

- When $P = P \succeq_{\#} P = Q$ then all reductions in P are matched in Q .
- When $P = P_1 \# P_2 \succeq_{\#} Q$ with $P_i \succeq_{\#} Q \quad i \in \{1, 2\}$. Let us take $i = 1$ Then by the rule:

$$\frac{P_1 \longrightarrow P'_1}{P_1 \# P_2 \longrightarrow P'_1 \# P_2} [\rightarrow_{\#}]$$

we have $P_1 \# P_2 \longrightarrow \{P'_{1_i}\}_{i \in I_1} \# P_2$ and $P_1 \# P_2 \longrightarrow P_1 \# \{P'_{2_i}\}_{i \in I_2}$ for some I_1, I_2 Hence we have that $P_1 \# P_2 \longrightarrow \{P'_i\}_{i \in I_1 \cup I_2}$ where $P'_i = P'_{1_i} \# P_2$ if $i \in I_1$ and $P'_i = P_1 \# P'_{2_i}$ if $i \in I_2$. By the induction hypothesis $P_1 \succeq_{\#} Q$ and $P_1 \longrightarrow \{P'_{1_i}\}_{i \in I_1}$ imply $\exists J', \{Q_j\}_{j \in J'}$ s.t. $Q \longrightarrow^* \{Q_j\}_{j \in J'}$, $J' \subseteq I_1$ and $P_j \succeq_{\#} Q_j$, $\forall j \in J'$. We take $J = J'$ and hence we can deduce $P \longrightarrow \{P'_i\}_{i \in I_1 \cup I_2}$, $Q \longrightarrow^* \{Q_j\}_{j \in J}$. Finally we have that $\forall j \in J$.

$$\frac{P'_{1_j} \succeq_{\#} Q_j}{P'_{1_j} \# P_2 \succeq_{\#} Q}$$

- When $P = P_1 \mid P_2$ with $P_1 \mid P_2 \succeq_{\#} Q$. Then by the rule:

$$\frac{P_1 \longrightarrow P'_1}{P_1 \mid P_2 \longrightarrow P'_1 \mid P_2} [\rightarrow]$$

we have $P_1 \mid P_2 \longrightarrow \{P'_{1_i}\}_{i \in I_1} \mid P_2$ and $P_1 \mid P_2 \longrightarrow P_1 \mid \{P'_{2_i}\}_{i \in I_2}$ for some I_1, I_2 Hence we have that $P_1 \mid P_2 \longrightarrow \{P'_i\}_{i \in I_1 \cup I_2}$ where $P'_i = P'_{1_i} \mid P_2$ if $i \in I_1$ and $P'_i = P_1 \mid P'_{2_i}$. By the induction hypothesis $P_1 \mid P_2 \succeq_{\#} Q$ and $P_1 \longrightarrow \{P'_{1_i}\}_{i \in I_1}$ imply $\exists J'_1, \{Q_j\}_{j \in J'_1}$ s.t. $Q \longrightarrow^* \{Q_j\}_{j \in J'_1}$, $J'_1 \subseteq I_1$ and $P'_{1_j} \succeq_{\#} Q_j$, $\forall j \in J'_1$. Similarly we have that $P_2 \longrightarrow \{P'_{2_i}\}_{i \in I_2}$ imply $\exists J'_2, \{Q_j\}_{j \in J'_2}$ s.t. $Q \longrightarrow^* \{Q_j\}_{j \in J'_2}$, $J'_2 \subseteq I_2$ and $P_{2_j} \succeq_{\#} Q_j$, $\forall j \in J'_2$. We take $J = J'_1 \cup J'_2$ and hence we can deduce $P \longrightarrow \{P'_i\}_{i \in I_1 \cup I_2}$, $Q \longrightarrow^* \{Q_j\}_{j \in J}$. Finally we have that $\forall j \in J$ $P_j \succeq_{\#} Q_j$

- When $P = (\nu x)P_1$ with $(\nu x)P_1 \succeq_{\#} Q$. Then by the rule:

$$\frac{P_1 \longrightarrow P'}{(\nu x)P_1 \longrightarrow (\nu x)P'} [\rightarrow_{\nu}]$$

we have $(\nu x)P_1 \longrightarrow (\nu x)\{P'_i\}_{i \in I}$ for some I . By the induction hypothesis $(\nu x)P_1 \succeq_{\#} Q$ and $(\nu x)P_1 \longrightarrow (\nu x)\{P'_i\}_{i \in I}$ imply $\exists J', \{Q_j\}_{j \in J'}$ s.t. $Q \longrightarrow^* \{Q_j\}_{j \in J'}$, $J' \subseteq I$ and $P'_j \succeq_{\#} Q_j$, $\forall j \in J'$. We take $J = J'$ and hence we can deduce $P \longrightarrow \{P'_i\}_{i \in I}$, $Q \longrightarrow^* \{Q_j\}_{j \in J}$. Finally we have that $\forall j \in J$

$$\frac{P'_j \succeq_{\#} Q_j}{(\nu x)P'_j \succeq_{\#} (\nu x)Q_j}$$

□

Lemma 1. *Let N be a well-formed closed term. If $\llbracket N \rrbracket_u \longrightarrow^* Q$, then there exist N' and $\{Q_i\}_{i \in I}$ such that (i) $N \longrightarrow^* N'$ and (ii) $Q \longrightarrow^* \{Q_i\}_{i \in I}$ where for each $j \in I$, $\llbracket N' \rrbracket_u \succeq_{\#} Q_j$.*

Proof. By induction on the structure of N and then induction on the number of reductions of $\llbracket N \rrbracket_u \longrightarrow^* Q$.

1. **Base case:** $N = x$, $N = x[j]$, $N = \mathbf{fail}^\emptyset$ and $N = \lambda x.(M'[\tilde{x} \leftarrow x])$.

No reductions can take place, and the result follows trivially. Take $I = \{a\}$,

$$Q = \llbracket N \rrbracket_u \xrightarrow{0} \llbracket N \rrbracket_u = Q_a \text{ and } x \xrightarrow{0} x = N'.$$

2. $N = M'(C \star U)$.

Then, $\llbracket M'(C \star U) \rrbracket_u = (\nu v)(\llbracket M' \rrbracket_v \mid v.\mathbf{some}_{u, \text{fv}(C)}; \bar{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x))$, and we are able to perform the reductions from $\llbracket M'(C \star U) \rrbracket_u$.

We now proceed by induction on k , with $\llbracket N \rrbracket_u \xrightarrow{k} Q$. There are two main cases:

(a) When $k = 0$ the thesis follows easily:

$$\text{We have } i = \{a\}, Q = \llbracket M'(C \star U) \rrbracket_u \xrightarrow{0} \llbracket M'(C \star U) \rrbracket_u = Q_a \text{ and } M'(C \star U) \xrightarrow{0} M'(C \star U) = N'.$$

(b) The interesting case is when $k \geq 1$.

Then, for some process R and n, m such that $k = n + m$, we have the following:

$$\begin{aligned} \llbracket N \rrbracket_u &= (\nu v)(\llbracket M' \rrbracket_v \mid v.\mathbf{some}_{u, \text{fv}(C)}; \bar{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x)) \\ &\xrightarrow{m} (\nu v)(R \mid v.\mathbf{some}_{u, \text{fv}(C)}; \bar{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x)) \xrightarrow{n} Q \end{aligned}$$

Thus, the first $m \geq 0$ reduction steps are internal to $\llbracket M' \rrbracket_v$; type preservation in $\mathbf{s}\pi^+$ ensures that, if they occur, these reductions do not discard the possibility of synchronizing with $\bar{v}.\mathbf{some}$. Then, the first of the $n \geq 0$ reduction steps towards Q is a synchronization between R and $v.\mathbf{some}_{u, \text{fv}(C)}$.

We consider two sub-cases, depending on the values of m and n :

(b.1) Case $m = 0$ and $n \geq 1$:

Then $R = \llbracket M \rrbracket_v$ as $\llbracket M \rrbracket_v \xrightarrow{0} \llbracket M \rrbracket_v$. Notice that there are two possibilities of having an unguarded:

$$\text{i. } M' = (\lambda x.(M''[\tilde{x} \leftarrow x])) \langle C_1/\tilde{y}_1 \rangle \cdots \langle C_p/\tilde{y}_p \rangle \llbracket U_1/z_1^! \rrbracket \cdots \llbracket U_q/z_q^! \rrbracket \quad (p, q \geq 0)$$

$$\llbracket M' \rrbracket_v = \llbracket (\lambda x.(M''[\tilde{x} \leftarrow x])) \langle C_1/\tilde{y}_1 \rangle \cdots \langle C_p/\tilde{y}_p \rangle \llbracket U_1/z_1^! \rrbracket \cdots \llbracket U_q/z_q^! \rrbracket \rrbracket_v$$

For simplicity we shall denote

$$\llbracket (\lambda x.(M''[\tilde{x} \leftarrow x])) \langle C_1/\tilde{y}_1 \rangle \cdots \langle C_p/\tilde{y}_p \rangle \llbracket U_1/z_1^! \rrbracket \cdots \llbracket U_q/z_q^! \rrbracket \rrbracket_v = (\nu \tilde{y}, \tilde{z})(\llbracket \lambda x.(M''[\tilde{x} \leftarrow x]) \rrbracket_v \mid Q'')$$

where $\tilde{y} = \tilde{y}_1, \dots, \tilde{y}_p$, $\tilde{z} = z_1^!, \dots, z_q^!$ and we continue the evaluation as:

$$\begin{aligned} &= (\nu \tilde{y}, \tilde{z})(\llbracket \lambda x.(M''[\tilde{x} \leftarrow x]) \rrbracket_v \mid Q'') \\ &= (\nu \tilde{y}, \tilde{z})(\bar{v}.\mathbf{some}; v(x); \bar{x}.\mathbf{some}; x(x^\ell); x(x^1); x(); \llbracket M''[\tilde{x} \leftarrow x] \rrbracket_v \mid Q'') \end{aligned}$$

With this shape for M , we then have the following:

$$\begin{aligned} \llbracket N \rrbracket_u &= \llbracket (M' B) \rrbracket_u = (\nu v)(\llbracket M' \rrbracket_v \mid v.\mathbf{some}_{u, \text{fv}(C)}; \bar{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x)) \\ &\xrightarrow{0} (\nu v)((\nu \tilde{y}, \tilde{z})(v(x); \bar{x}.\mathbf{some}; x(x^\ell); x(x^1); x()); \llbracket M''[\tilde{x} \leftarrow x] \rrbracket_v \\ &\quad \mid Q'') \mid \bar{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x)) &= Q_1 \\ &\xrightarrow{0} (\nu x)((\nu v)((\nu \tilde{y}, \tilde{z})(\bar{x}.\mathbf{some}; x(x^\ell); x(x^1); x()); \llbracket M''[\tilde{x} \leftarrow x] \rrbracket_v \\ &\quad \mid Q'') \mid [v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x) &= Q_2 \\ &\xrightarrow{0} (\nu x)((\nu \tilde{y}, \tilde{z})(\bar{x}.\mathbf{some}; x(x^\ell); x(x^1); x()); \llbracket M''[\tilde{x} \leftarrow x] \rrbracket_u \mid Q'') \\ &\quad \mid \llbracket C \star U \rrbracket_x) &= Q_3 \end{aligned}$$

We also have that

$$\begin{aligned} N &= (\lambda x.(M''[\tilde{x} \leftarrow x])) \langle C_1/\tilde{y}_1 \rangle \cdots \langle C_p/\tilde{y}_p \rangle \llbracket U_1/z_1^! \rrbracket \cdots \llbracket U_q/z_q^! \rrbracket (C \star U) \\ &\equiv_\lambda (\lambda x.(M''[\tilde{x} \leftarrow x]) (C \star U)) \langle C_1/\tilde{y}_1 \rangle \cdots \langle C_p/\tilde{y}_p \rangle \llbracket U_1/z_1^! \rrbracket \cdots \llbracket U_q/z_q^! \rrbracket \\ &\longrightarrow M''[\tilde{x} \leftarrow x] \langle (C \star U)/x \rangle \langle C_1/\tilde{y}_1 \rangle \cdots \langle C_p/\tilde{y}_p \rangle \llbracket U_1/z_1^! \rrbracket \cdots \llbracket U_q/z_q^! \rrbracket = M \end{aligned}$$

Furthermore, we have:

$$\begin{aligned} \llbracket M \rrbracket_u &= \llbracket M''[\tilde{x} \leftarrow x] \langle (C \star U)/x \rangle \langle C_1/\tilde{y}_1 \rangle \cdots \langle C_p/\tilde{y}_p \rangle \llbracket U_1/z_1^! \rrbracket \cdots \llbracket U_q/z_q^! \rrbracket \rrbracket_u \\ &= (\nu x)((\nu \tilde{y}, \tilde{z})(\bar{x}.\mathbf{some}; x(x^\ell); x(x^1); x()); \llbracket M''[\tilde{x} \leftarrow x] \rrbracket_u \mid Q'') \mid \llbracket C \star U \rrbracket_x \end{aligned}$$

We consider different possibilities for $n \geq 1$; in all the cases, the result follows.

When $n = 1$:

We have $I = \{a\}$, $Q = Q_1$, $\llbracket N \rrbracket_u \xrightarrow{1} Q_1$. We also have that

$$- Q_1 \xrightarrow{2} Q_3 = Q_a,$$

- $N \rightarrow^1 M''[\tilde{x} \leftarrow x] \langle\langle (C \star U)/x \rangle\rangle \langle C_1/\tilde{y}_1 \rangle \cdots \langle C_p/\tilde{y}_p \rangle \llbracket U_1/z_1^! \rrbracket \cdots \llbracket U_q/z_q^! \rrbracket = N'$
- and $\llbracket M''[\tilde{x} \leftarrow x] \langle\langle (C \star U)/x \rangle\rangle \langle C_1/\tilde{y}_1 \rangle \cdots \langle C_p/\tilde{y}_p \rangle \llbracket U_1/z_1^! \rrbracket \cdots \llbracket U_q/z_q^! \rrbracket \rrbracket_u = Q_3$.

When $n = 2, 3$: the analysis is similar.

When $n \geq 4$:

We have $\llbracket N \rrbracket_u \rightarrow^3 Q_3 \rightarrow^* Q$. We also know that $N \rightarrow M$, $Q_3 = \llbracket M \rrbracket_u$. By the IH, there exist $\{Q_i\}_{i \in I}$, N' such that $Q \rightarrow^* \{Q_i\}_{i \in I}$, $M \rightarrow^* N'$ and $\llbracket N' \rrbracket_u \succeq_{\perp} Q_i \quad \forall i \in I$. Finally, $\llbracket N \rrbracket_u \rightarrow^3 Q_3 \rightarrow^l Q \rightarrow_I Q_i$ and $N \rightarrow M \rightarrow^* N'$.

ii. $M' = \text{fail}^{\tilde{z}}$.

Then, $\llbracket M' \rrbracket_v = \llbracket \text{fail}^{\tilde{z}} \rrbracket_v = \bar{v}.\text{none} \mid \bar{\tilde{z}}.\text{none}$. With this shape for M , we have:

$$\begin{aligned} \llbracket N \rrbracket_u &= \llbracket (M' (C \star U)) \rrbracket_u = (\nu v)(\llbracket M' \rrbracket_v \mid v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x)) \\ &= (\nu v)(\bar{v}.\text{none} \mid \bar{\tilde{z}}.\text{none} \mid v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x)) \\ &\rightarrow \bar{u}.\text{none} \mid \bar{\tilde{z}}.\text{none} \mid \overline{\text{fv}(C)}.\text{none} \end{aligned}$$

We also have that $N = \text{fail}^{\tilde{z}} C \star U \rightarrow \text{fail}^{\tilde{z} \cup \text{fv}(C)} = N'$. Furthermore,

$$\llbracket N' \rrbracket_u = \llbracket \text{fail}^{\tilde{z} \cup \text{fv}(C)} \rrbracket_u = \llbracket \text{fail}^{\tilde{z} \cup \text{fv}(C)} \rrbracket_u = \bar{u}.\text{none} \mid \bar{\tilde{z}}.\text{none} \mid \overline{\text{fv}(C)}.\text{none}.$$

(b.2) Case $m \geq 1$ and $n \geq 0$:

We distinguish two cases:

i. When $n = 0$:

Then, $(\nu v)(R \mid v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x)) = Q$ and $\llbracket M' \rrbracket_u \rightarrow^m R$ where $m \geq 1$. Then by the IH there exist $\{R'_i\}_{i \in I}$ and M'' such that $R \rightarrow^* \{R'_i\}_{i \in I}$, $M' \rightarrow^* M''$, and $\llbracket M'' \rrbracket_u \succeq_{\perp} R_i \quad \forall i \in I$. Hence we have that

$$\begin{aligned} \llbracket N \rrbracket_u &= (\nu v)(\llbracket M' \rrbracket_v \mid v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x)) \\ &\rightarrow^m (\nu v)(R \mid v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x)) = Q \end{aligned}$$

We also know that

$$Q \rightarrow^* \{(\nu v)(R'_i \mid v.\text{some}_{u, \text{fv}(C)}; \bar{v}[x].([v \leftrightarrow u] \mid \llbracket C \star U \rrbracket_x))\}_{i \in I} = \{Q_i\}_{i \in I}$$

and so the λ_C term can reduce as follows: $N = (M' (C \star U)) \rightarrow^* M'' (C \star U) = N'$ and $\llbracket N' \rrbracket_u \succeq_{\perp} Q_i \quad \forall i \in I$ via the \succeq_{\perp} rules.

ii. When $n \geq 1$:

Then R has an occurrence of an unguarded $\bar{v}.\text{some}$ or $\bar{v}.\text{none}$, hence it is of the form $\llbracket (\lambda x.(M''[\tilde{x} \leftarrow x]) \langle\langle N_1/y_1 \rangle\rangle \cdots \langle\langle N_p/y_p \rangle\rangle \llbracket U_1/z_1^! \rrbracket \cdots \llbracket U_q/z_q^! \rrbracket) \rrbracket_v$ or $\llbracket \text{fail}^{\tilde{z}} \rrbracket_v$.

This concludes the analysis for the case $N = (M' (C \star U))$.

3. $N = M'[\tilde{x} \leftarrow x]$. The sharing variable x is not free and the result follows by vacuity.
4. $N = M'[\tilde{x} \leftarrow x] \langle\langle C \star U/x \rangle\rangle$. Then we have

$$\llbracket N \rrbracket_u = \llbracket M'[\tilde{x} \leftarrow x] \langle\langle C \star U/x \rangle\rangle \rrbracket_u = (\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u \mid \llbracket C \star U \rrbracket_x$$

Let us consider three cases.

(a) When $\text{size}(\tilde{x}) = \text{size}(C)$. Then let us consider the shape of the bag C .

i. When $C = 1$.

We have the following

$$\begin{aligned}
\llbracket N \rrbracket_u &= (\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[\leftarrow x] \rrbracket_u \mid \llbracket \mathbf{1} \star U \rrbracket_x \\
&= (\nu x)(\bar{x}.\text{some}; x(x^\ell); x(x^1); x()); \llbracket M'[\leftarrow x] \rrbracket_u \mid x.\text{some}_{\text{fv}(C)}; \bar{x}[x^\ell]. \\
&\quad (\llbracket \mathbf{1} \rrbracket_{x^\ell} \mid \bar{x}[x^1].(!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid \bar{x}[])) \\
&\longrightarrow (\nu x)(x(x^\ell); x(x^1); x()); \llbracket M'[\leftarrow x] \rrbracket_u \mid \bar{x}[x^\ell].(\llbracket \mathbf{1} \rrbracket_{x^\ell} \mid \bar{x}[x^1].(!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid \bar{x}[])) = Q_1 \\
&\longrightarrow (\nu x, x^\ell)(x(x^1); x()); \llbracket M'[\leftarrow x] \rrbracket_u \mid \llbracket \mathbf{1} \rrbracket_{x^\ell} \mid \bar{x}[x^1].(!x^1(x_i); \llbracket U \rrbracket_{x_i} \mid \bar{x}[]) = Q_2 \\
&\longrightarrow (\nu x, x^\ell, x^1)(x()); \llbracket M'[\leftarrow x] \rrbracket_u \mid \llbracket \mathbf{1} \rrbracket_{x^\ell} \mid !x^1(x_i); \llbracket U \rrbracket_{x_i} \mid \bar{x}[] = Q_3 \\
&\longrightarrow (\nu x^\ell, x^1)(\llbracket M'[\leftarrow x] \rrbracket_u \mid \llbracket \mathbf{1} \rrbracket_{x^\ell} \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) = Q_4 \\
&= (\nu x^\ell, x^1)(\bar{x}^\ell.\text{some}; \bar{x}^\ell[y_i].(y_i.\text{some}_{u, \text{fv}(M')}); y_i(); \llbracket M' \rrbracket_u \mid \bar{x}^\ell.\text{none}) \mid \\
&\quad x^\ell.\text{some}_\emptyset; x^\ell(y_n); (\bar{y}_n.\text{some}; \bar{y}_n[] \mid x^\ell.\text{some}_\emptyset; \bar{x}^\ell.\text{none}) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i} \\
&\longrightarrow (\nu x^\ell, x^1)(\bar{x}^\ell[y_i].(y_i.\text{some}_{u, \text{fv}(M')}); y_i(); \llbracket M' \rrbracket_u \mid \bar{x}^\ell.\text{none}) \mid \\
&\quad x^\ell(y_n); (\bar{y}_n.\text{some}; \bar{y}_n[] \mid x^\ell.\text{some}_\emptyset; \bar{x}^\ell.\text{none}) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i} = Q_5 \\
&\longrightarrow (\nu x^\ell, x^1, y_i)(y_i.\text{some}_{u, \text{fv}(M')}); y_i(); \llbracket M' \rrbracket_u \mid \bar{x}^\ell.\text{none} \mid \bar{y}_i.\text{some}; \bar{y}_i[] \\
&\quad \mid x^\ell.\text{some}_\emptyset; \bar{x}^\ell.\text{none} \mid !x^1(x_i); \llbracket U \rrbracket_{x_i} = Q_6 \\
&\longrightarrow (\nu x^\ell, x^1, y_i)(y_i(); \llbracket M' \rrbracket_u \mid \bar{x}^\ell.\text{none} \mid \bar{y}_i[] \mid x^\ell.\text{some}_\emptyset; \bar{x}^\ell.\text{none} \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) = Q_7 \\
&\longrightarrow (\nu x^\ell, x^1)(\llbracket M' \rrbracket_u \mid \bar{x}^\ell.\text{none} \mid x^\ell.\text{some}_\emptyset; \bar{x}^\ell.\text{none} \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) = Q_8 \\
&\longrightarrow (\nu x^1)(\llbracket M' \rrbracket_u \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) = \llbracket M' \llbracket U/x^1 \rrbracket \rrbracket_u = Q_9
\end{aligned}$$

Notice how Q_8 has a choice however the x^ℓ name can be closed at any time so for simplicity we only perform communication across this name once all other names have completed their reductions.

Now we proceed by induction on the number of reductions $\llbracket N \rrbracket_u \longrightarrow^k Q$.

A. When $k = 0$, the result follows trivially. Just take $I = \{a\}$, $\mathbb{N} = \mathbb{N}'$ and $\llbracket N \rrbracket_u = Q = Q_a$.

B. When $k = 1$.

We have $Q = Q_1$, $\llbracket N \rrbracket_u \longrightarrow^1 Q_1$. Let us take $I = a$, we also have that $Q_1 \longrightarrow^8 Q_9 = Q_a$, $N \longrightarrow M' \llbracket U/x^1 \rrbracket = M$ and $\llbracket M' \rrbracket_u \succeq_{\dagger} Q_9$

C. When $2 \leq k \leq 8$.

Proceeds similarly to the previous case

D. When $k \geq 9$.

We have $\llbracket N \rrbracket_u \longrightarrow^9 Q_9 \longrightarrow^l Q$, for $l \geq 1$. Since $Q_9 = \llbracket M' \rrbracket_u$ we apply the induction hypothesis we have that there $\exists \{Q_i\}_{i \in I}$, N' s.t. $Q \longrightarrow^* \{Q_i\}_{i \in I}$, $M' \longrightarrow^* N'$ and $\llbracket N' \rrbracket_u \succeq_{\dagger} Q_i \quad \forall i \in I$. Then, $\llbracket N \rrbracket_u \longrightarrow^9 Q_9 \longrightarrow^l Q \longrightarrow^* \{Q_i\}_{i \in I}$ and by the contextual reduction rule it follows that $N = (M'[\leftarrow x]) \langle \langle 1/x \rangle \rangle \longrightarrow^* N'$ and the case holds.

ii. When $C = \lambda N_1 \} \dots \} \lambda N_l \}$, for $l \geq 1$. Then,

$$\begin{aligned}
\llbracket N \rrbracket_u &= \llbracket M'[\tilde{x} \leftarrow x] \langle C \star U/x \rangle \rrbracket_u \\
&= (\nu x)(\overline{x}.\text{some}; x(x^\ell); x(x^1); x(); \llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u \mid \llbracket C \star U \rrbracket_x) \\
&\rightarrow^4 (\nu x^1)((\nu x^\ell)(\llbracket M'[\tilde{x} \leftarrow x] \rrbracket_u \mid \llbracket C \rrbracket_{x^\ell} \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&= (\nu x^1)((\nu x^\ell)(\overline{x}^\ell.\text{some}; \overline{x}^\ell[y_1]; (y_1.\text{some}_\emptyset; y_1()); \mathbf{0} \\
&\quad \mid \overline{x}^\ell.\text{some}; x^\ell.\text{some}_{u, \text{fv}(M') \setminus \tilde{x}}; \prod_{x_{i_1} \in \tilde{x}} x(x_{i_1}); \dots \\
&\quad \overline{x}^\ell.\text{some}; \overline{x}^\ell[y_i]; (y_i.\text{some}_\emptyset; y_i()); \mathbf{0} \\
&\quad \mid \overline{x}^\ell.\text{some}; x^\ell.\text{some}_{u, \text{fv}(M') \setminus (\tilde{x} \setminus (x_{i_1}, \dots, x_{i_{l-1}}))}; \prod_{x_{i_l} \in \tilde{x}} x(x_{i_l}); \\
&\quad \overline{x}^\ell.\text{some}; \overline{x}^\ell[y_{l+1}]; (y_{l+1}.\text{some}_{u, \text{fv}(M')}; y_{l+1}()); \llbracket M' \rrbracket_u \mid \overline{x}^\ell.\text{none})) \dots) \\
&\quad \mid \\
&\quad x^\ell.\text{some}_{\text{fv}(C)}; x(y_1); x^\ell.\text{some}_{y_1, \text{fv}(C)}; \overline{x}^\ell.\text{some}; \overline{x}^\ell[z_1]; \\
&\quad (z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \overline{y_1}.\text{none} \mid \\
&\quad \dots x^\ell.\text{some}_{\text{fv}(M_l)}; x(y_l); x^\ell.\text{some}_{y_l, \text{fv}(M_l)}; \overline{x}^\ell.\text{some}; \overline{x}^\ell[z_l]; \\
&\quad (z_l.\text{some}_{\text{fv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \overline{y_l}.\text{none} \mid \\
&\quad x^\ell.\text{some}_\emptyset; x(y_{l+1}); (\overline{y_{l+1}}.\text{some}; \overline{y_{l+1}}[] \mid x^\ell.\text{some}_\emptyset; \overline{x}^\ell.\text{none})) \dots) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&\rightarrow^{6l} (\nu x^1)((\nu x^\ell)((\nu z_1)(\dots (\nu z_l)(\overline{x}^\ell.\text{some}; \overline{x}^\ell[y_{l+1}]; (y_{l+1}.\text{some}_{u, \text{fv}(M')}; \\
&\quad y_{l+1}()); \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \dots \{z_l/x_{i_l}\} \mid \overline{x}^\ell.\text{none} \\
&\quad \mid z_l.\text{some}_{\text{fv}(M_{j_l)}; \llbracket M_{j_l} \rrbracket_{z_l}) \dots \mid z_1.\text{some}_{\text{fv}(M_{j_1)}; \llbracket M_{j_1} \rrbracket_{z_1}) \mid \\
&\quad x^\ell.\text{some}_\emptyset; x^\ell(y_{l+1}).(\overline{y_{l+1}}.\text{some}; \overline{y_{l+1}}[] \mid x^\ell.\text{some}_\emptyset; \overline{x}^\ell.\text{none})) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&\rightarrow^5 (\nu x^1)((\nu z_1)(x_1.\text{some}_{\text{fv}(M_{j_1)}; \llbracket M_{j_1} \rrbracket_{x_1} \mid \dots \\
&\quad (\nu z_l)(x_l.\text{some}_{\text{fv}(M_{j_l)}; \llbracket M_{j_l} \rrbracket_{x_l} \mid \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \dots \{z_l/x_{i_l}\}) \dots) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i})
\end{aligned}$$

The proof follows by induction on the number of reductions $\llbracket N \rrbracket_u \rightarrow^k Q$.

- A. When $k = 0$, the result follows trivially. Just take $I = \{a\}$, $\mathbb{N} = \mathbb{N}'$ and $\llbracket N \rrbracket_u = Q = Q_a$.
B. When $1 \leq k \leq 6l + 9$.

Let $I = \{a\}$, Q_k be such that $\llbracket N \rrbracket_u \rightarrow^k Q_k$. We also have that $Q_k \rightarrow^{6l+9-k} Q_{6l+9} = Q_a$,
 $N \rightarrow^1 M' \langle C/x_1, \dots, x_l \rangle \llbracket U/x^1 \rrbracket = N'$ and

$$\begin{aligned}
\llbracket M' \langle C/x_1, \dots, x_l \rangle \llbracket U/x^1 \rrbracket \rrbracket_u &= (\nu x^1) \left((\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \right. \\
&\quad \dots (\nu z_l)(z_l.\text{some}_{\text{fv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \\
&\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_l\}} \dots \prod_{x_{i_l} \in \{x_1, \dots, x_l \setminus x_{i_1}, \dots, x_{i_{l-1}}\}} \\
&\quad \left. \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \dots \{z_l/x_{i_l}\}) \dots \mid !x^1(x_i); \llbracket U \rrbracket_{x_i} \right) \\
&\succeq_{\neq} (\nu x^1) \left((\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \right. \\
&\quad \dots (\nu z_l)(z_l.\text{some}_{\text{fv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \\
&\quad \mid \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \dots \{z_l/x_{i_l}\}) \dots \mid \\
&\quad \left. !x^1(x_i); \llbracket U \rrbracket_{x_i} \right) = Q_{6l+9}
\end{aligned}$$

- C. When $k > 6l + 9$.

Then, $\llbracket N \rrbracket_u \rightarrow^{6l+9} Q_{6l+9} \rightarrow^n Q$ for $n \geq 1$. Also,
 $N \rightarrow^1 M' \langle C/x_1, \dots, x_l \rangle \llbracket U/x^1 \rrbracket$ and

$$\llbracket M' \langle C/x_1, \dots, x_l \rangle \llbracket U/x^1 \rrbracket \rrbracket_u \succeq_{\neq} Q_{6l+9}.$$

$\exists \{P_j\}_{j \in J}, N'$, s.t. $M' \langle C/x_1, \dots, x_l \rangle \llbracket U/x^1 \rrbracket \rightarrow^* N'$, $\llbracket M' \langle C/x_1, \dots, x_l \rangle \llbracket U/x^1 \rrbracket \rrbracket_u \rightarrow^n P \rightarrow^* \{P_j\}_{j \in J}$
and $\llbracket N' \rrbracket_u \succeq_{\neq} P_j \quad \forall j \in J$. We also have by Prop.2 that as $\llbracket M' \langle C/x_1, \dots, x_l \rangle \llbracket U/x^1 \rrbracket \rrbracket_u \succeq_{\neq} Q_{6l+9}$ and
 $\llbracket M' \langle C/x_1, \dots, x_l \rangle \llbracket U/x^1 \rrbracket \rrbracket_u \rightarrow^* \{Q_j\}_{j \in J}$ implies $\exists \{Q_i\}_{i \in I}$ s.t. $Q_{6l+9} \rightarrow^* Q_{i \in I}$, $I \subset J$ and $P_i \succeq_{\neq} Q_i$, $\forall i \in I$

- (b) When $\text{size}(\tilde{x}) > \text{size}(C)$.

Then we have $N = M'[x_1, \dots, x_k \leftarrow x] \langle\langle C \star U/x \rangle\rangle$ with $C = \wr M_1 \wr \dots \wr M_l \wr$ $k > l$, $N \longrightarrow \mathbf{fail}^{\tilde{z}} = M$ and $\tilde{z} = (\mathbf{fv}(M') \setminus \{x_1, \dots, x_k\}) \cup \mathbf{fv}(C)$. On the one hand, we have: Hence $k = l + m$ for some $m \geq 1$

$$\begin{aligned}
\llbracket N \rrbracket_u &= \llbracket M'[x_1, \dots, x_k \leftarrow x] \langle\langle C \star U/x \rangle\rangle \rrbracket_u \\
&= (\nu x)(\overline{x}.\mathbf{some}; x(x^\ell); x(x^1); x()); \llbracket M'[x_1, \dots, x_k \leftarrow x] \rrbracket_u \mid \llbracket C \star U \rrbracket_x \\
&\longrightarrow^4 (\nu x^1)((\nu x^\ell)(\llbracket M'[x_1, \dots, x_k \leftarrow x] \rrbracket_u \mid \llbracket C \rrbracket_{x^\ell} \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&= (\nu x^1)((\nu x^\ell)(\overline{x}^\ell.\mathbf{some}; \overline{x}^\ell[y_1]; (y_1.\mathbf{some}_\emptyset; y_1()); \mathbf{0} \\
&\quad \mid \overline{x}^\ell.\mathbf{some}; x^\ell.\mathbf{some}_{u, \mathbf{fv}(M') \setminus \tilde{x}}; \prod_{x_{i_1} \in \tilde{x}} x(x_{i_1}); \dots \\
&\quad \overline{x}^\ell.\mathbf{some}; \overline{x}^\ell[y_k]; (y_k.\mathbf{some}_\emptyset; y_k()); \mathbf{0} \\
&\quad \mid \overline{x}^\ell.\mathbf{some}; x^\ell.\mathbf{some}_{u, \mathbf{fv}(M') \setminus \tilde{x}}; \prod_{x_{i_k} \in (\tilde{x} \setminus x_{i_1}, \dots, x_{i_{k-1}})} x(x_{i_k}); \\
&\quad \overline{x}^\ell.\mathbf{some}; \overline{x}^\ell[y_{k+1}]; (y_{k+1}.\mathbf{some}_{u, \mathbf{fv}(M')}; y_{k+1}()); \llbracket M' \rrbracket_u \mid \overline{x}^\ell.\mathbf{none})) \dots) \\
&\quad \mid x^\ell.\mathbf{some}_{\mathbf{fv}(C)}; x(y_1); x^\ell.\mathbf{some}_{y_1, \mathbf{fv}(C)}; \overline{x}^\ell.\mathbf{some}; \overline{x}^\ell[z_1]; \\
&\quad (z_1.\mathbf{some}_{\mathbf{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \overline{y_1}.\mathbf{none} \mid \\
&\quad \dots x^\ell.\mathbf{some}_{\mathbf{fv}(M_l)}; x(y_l); x^\ell.\mathbf{some}_{y_l, \mathbf{fv}(C)}; \overline{x}^\ell.\mathbf{some}; \overline{x}^\ell[z_l]; \dots \\
&\quad (z_l.\mathbf{some}_{\mathbf{fv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \overline{y_l}.\mathbf{none} \mid \\
&\quad x^\ell.\mathbf{some}_\emptyset; x(y_{l+1}); (\overline{y_{l+1}}.\mathbf{some}; \overline{y_{l+1}} \mid x^\ell.\mathbf{some}_\emptyset; \overline{x}^\ell.\mathbf{none})) \dots) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i} \\
&\longrightarrow^{6k} (\nu x^1)((\nu x^\ell)((\nu z_1)(z_1.\mathbf{some}_{\mathbf{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_k)(z_k.\mathbf{some}_{\mathbf{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \mid \\
&\quad \overline{x}^\ell.\mathbf{some}; \overline{x}^\ell[y_{k+1}]; (y_{k+1}.\mathbf{some}_{u, \mathbf{fv}(M')}; y_{k+1}()); \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\} \mid \overline{x}^\ell.\mathbf{none})) \dots) \\
&\quad \mid x^\ell.\mathbf{some}_{\mathbf{fv}(C) \setminus (M_1, \dots, M_k)}; x(y_{k+1}); x^\ell.\mathbf{some}_{y_{k+1}, \mathbf{fv}(C) \setminus (M_1, \dots, M_k)}; \overline{x}^\ell.\mathbf{some}; \overline{x}^\ell[z_{k+1}]; \\
&\quad (z_{k+1}.\mathbf{some}_{\mathbf{fv}(M_{k+1})}; \llbracket M_{k+1} \rrbracket_{z_{k+1}} \mid \overline{y_1}.\mathbf{none} \mid \\
&\quad \dots x^\ell.\mathbf{some}_{\mathbf{fv}(M_l)}; x(y_l); x^\ell.\mathbf{some}_{y_l, \mathbf{fv}(M_l)}; \overline{x}^\ell.\mathbf{some}; \overline{x}^\ell[z_l]; \dots \\
&\quad (z_l.\mathbf{some}_{\mathbf{fv}(M_l)}; \llbracket M_l \rrbracket_{z_l} \mid \overline{y_l}.\mathbf{none} \mid \\
&\quad x^\ell.\mathbf{some}_\emptyset; x(y_{l+1}); (\overline{y_{l+1}}.\mathbf{some}; \overline{y_{l+1}} \mid x^\ell.\mathbf{some}_\emptyset; \overline{x}^\ell.\mathbf{none})) \dots) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i} \\
&\longrightarrow^6 (\nu x^1)((\nu z_1)(z_1.\mathbf{some}_{\mathbf{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_k)(z_k.\mathbf{some}_{\mathbf{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \mid \\
&\quad \overline{u}.\mathbf{none} \mid \overline{(\mathbf{fv}(M') \setminus \tilde{x}).\mathbf{none}} \mid \overline{z_1}.\mathbf{none} \mid \dots \mid \overline{z_k}.\mathbf{none} \mid \overline{(\mathbf{fv}(C) \setminus (M_1, \dots, M_k)).\mathbf{none}}) \dots) \\
&\quad \mid !x^1(x_i); \llbracket U \rrbracket_{x_i} \\
&\longrightarrow^k \overline{u}.\mathbf{none} \mid \overline{(\mathbf{fv}(M') \setminus \{x_1, \dots, x_k\}).\mathbf{none}} \mid \overline{\mathbf{fv}(C).\mathbf{none}} \\
&= \llbracket \mathbf{fail}^{\tilde{z}} \rrbracket_u = Q_{7l+10}
\end{aligned}$$

The rest of the proof is by induction on the number of reductions $\llbracket N \rrbracket_u \longrightarrow^j Q$.

i. When $j = 0$, the result follows trivially. Just take $I = \{a\}$ $\mathbb{N} = \mathbb{N}'$ and $\llbracket N \rrbracket_u = Q = Q_a$.

ii. When $1 \leq j \leq 7k + 10$.

Let $I = \{a\}$ and Q_j be such that $\llbracket N \rrbracket_u \longrightarrow^j Q_j$. By the steps above one has

$$Q_j \longrightarrow^{7k+10-j} Q_{7k+10} = Q_a,$$

$$N \longrightarrow^1 \mathbf{fail}^{\tilde{z}} = N'; \text{ and } \llbracket \mathbf{fail}^{\tilde{z}} \rrbracket_u = Q_{7k+10}.$$

iii. When $j > 7k + 10$.

In this case, we have $\llbracket N \rrbracket_u \longrightarrow^{7k+10} Q_{7k+10} \longrightarrow^n Q$, for $n \geq 1$. We also know that $N \longrightarrow^1 \mathbf{fail}^{\tilde{z}}$. However no further reductions can be performed.

(c) When $\text{size}(\tilde{x}) < \text{size}(C)$, the proof proceeds similarly to the previous case.

5. $N = M' \langle C/x_1, \dots, x_k \rangle$.

In this case we let $C = \wr M_1 \wr \dots \wr M_k \wr$,

$$\begin{aligned}
\llbracket M' \langle C/x_1, \dots, x_k \rangle \rrbracket_u &= (\nu z_1)(z_1.\mathbf{some}_{\mathbf{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \dots (\nu z_k)(z_k.\mathbf{some}_{\mathbf{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \dots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \\
&\quad \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \dots \{z_k/x_{i_k}\} \dots)
\end{aligned}$$

Therefore,

$$\begin{aligned}
\llbracket N \rrbracket_u &= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\} \cdots) \\
&\longrightarrow^m (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} R\{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\} \cdots) \\
&\longrightarrow^n Q
\end{aligned}$$

for some process R . Where \longrightarrow^n is a reduction that initially synchronizes with $x_i.\mathbf{some}_{\text{fv}(M_{J_i})}$ for some $i \in \{1, \dots, k\}$, when $n \geq 1$, $n + m = k \geq 1$. Type preservation in sr^+ ensures reducing $\llbracket M' \rrbracket_u \longrightarrow^m$ does not consume possible synchronizations with $\overline{x_i}.\mathbf{some}$, if they occur. Let us consider the the possible sizes of both m and n .

(a) For $m = 0$ and $n \geq 1$.

We have that $R = \llbracket M' \rrbracket_u$ as $\llbracket M' \rrbracket_u \longrightarrow^0 \llbracket M' \rrbracket_u$.

Notice that there are two possibilities of having an unguarded $\overline{x_i}.\mathbf{some}$ or $\overline{x_i}.\mathbf{none}$ without internal reductions:

i. $M' = \mathbf{fail}^{x_i, \tilde{y}}$.

$$\begin{aligned}
\llbracket N \rrbracket_u &= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\} \cdots) \\
&= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \llbracket \mathbf{fail}^{x_i, \tilde{y}} \rrbracket_u \{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\} \cdots) \\
&= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \\
&\quad \quad \overline{u}.\mathbf{none} \mid \overline{x_i}.\mathbf{none} \mid \overline{\tilde{y}}.\mathbf{none} \{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\} \cdots)
\end{aligned}$$

by type preservation we have that $\tilde{y} = \{x_1, \dots, x_k\} \setminus x_i, \tilde{y}'$ for some \tilde{y}'

$$\begin{aligned}
&= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \overline{u}.\mathbf{none} \mid \overline{\tilde{y}'}.\mathbf{none} \\
&\quad \mid \overline{\{x_1, \dots, x_k\}}.\mathbf{none} \{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\} \cdots) \\
&= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \overline{u}.\mathbf{none} \mid \overline{\tilde{y}'}.\mathbf{none} \\
&\quad \mid \overline{\{z_1, \dots, z_k\}}.\mathbf{none} \cdots) \\
&\longrightarrow^k \overline{u}.\mathbf{none} \mid \overline{\tilde{y}'}.\mathbf{none} \mid \overline{\text{fv}(C)}.\mathbf{none}
\end{aligned}$$

Notice that no further reductions can be performed. Thus we take $I = \{a\}$ and,

$$\llbracket N \rrbracket_u \longrightarrow \overline{u}.\mathbf{none} \mid \overline{\tilde{y}}.\mathbf{none} \mid \overline{\text{fv}(C)}.\mathbf{none} = Q_a.$$

We also have that $N \longrightarrow \mathbf{fail}^{\tilde{y} \cup \text{fv}(C)} = N'$ and $\llbracket \mathbf{fail}^{\tilde{y} \cup \text{fv}(C)} \rrbracket_u = Q_a$.

ii. $\text{head}(M') = x_i$ with $i \in \{1, \dots, k\}$.

Then we have the following

$$\begin{aligned}
\llbracket N \rrbracket_u &= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\} \cdots) \\
&= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} (\nu \tilde{y})(\llbracket x_i \rrbracket_j \mid P)\{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\} \cdots) \\
&= (\nu z_1)(z_1.\mathbf{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\mathbf{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \prod_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \prod_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \\
&\quad \quad (\nu \tilde{y})(\overline{x_i}.\mathbf{some}; [x_i \leftrightarrow j] \mid P)\{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\} \cdots)
\end{aligned}$$

Let us consider a arbitrary sum where $x_{i_k} = x_i$, other cases follow similarly.

$$\begin{aligned}
&= (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\text{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \amalg_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \amalg_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \\
&\quad (\nu \tilde{y})(\bar{x}_i.\text{some}; [x_i \leftrightarrow j] \mid P)\{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\} \cdots) \\
&\rightarrow (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(\llbracket M_k \rrbracket_{z_k} \\
&\quad \mid (\nu \tilde{y})([x_k \leftrightarrow j] \mid P)\{z_1/x_{i_1}\} \cdots \{z_{k-1}/x_{i_{k-1}}\} \cdots) = Q_1 \\
&\rightarrow (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_{k-1})(z_{k-1}.\text{some}_{\text{fv}(M_{k-1})}; \llbracket M_{k-1} \rrbracket_{z_{k-1}} \\
&\quad \mid (\nu \tilde{y})(\llbracket M_k \rrbracket_j \mid P)\{z_1/x_{i_1}\} \cdots \{z_{k-1}/x_{i_{k-1}}\} \cdots) = Q_2
\end{aligned}$$

In addition, $N = M' \langle C/x_1, \dots, x_k \rangle \rightarrow M' \{C_i/x_k\} \langle (C \setminus C_i)/x_1, \dots, x_{k-1} \rangle = M$. Finally,

$$\begin{aligned}
\llbracket M \rrbracket_u &= \llbracket M' \{C_i/x_k\} \langle (C \setminus C_i)/x_1, \dots, x_{k-1} \rangle \rrbracket_u \\
&= (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_{k-1})(z_{k-1}.\text{some}_{\text{fv}(M_{k-1})}; \llbracket M_{k-1} \rrbracket_{z_{k-1}} \mid \\
&\quad \amalg_{x_{i_1} \in \{x_1, \dots, x_{k-1}\}} \cdots \amalg_{x_{i_{k-1}} \in \{x_1, \dots, x_{k-1} \setminus x_{i_1}, \dots, x_{i_{k-2}}\}} \\
&\quad (\nu \tilde{y})(\llbracket M_k \rrbracket_j \mid P)\{z_1/x_{i_1}\} \cdots \{z_{k-1}/x_{i_{k-1}}\} \cdots) \\
&\succeq_{\dagger} Q_2.
\end{aligned}$$

A. When $n = 1$:

Then, $I = \{a\}$, $Q = Q_1$ and $\llbracket N \rrbracket_u \rightarrow^1 Q_1$. Also, $Q_1 \rightarrow^1 Q_2 = Q_a$, $N \rightarrow^1 M' \{C_i/x_k\} \langle (C \setminus C_i)/x_1, \dots, x_{k-1} \rangle = N'$ and $\llbracket M' \{C_i/x_k\} \langle (C \setminus C_i)/x_1, \dots, x_{k-1} \rangle \rrbracket_u \succeq_{\dagger} Q_a$.

B. When $n = 2$:

Then, $I = \{a\}$, $Q = Q_2$ and $\llbracket N \rrbracket_u \rightarrow^2 Q_2$. Also, $Q_2 \rightarrow^0 Q_2 = Q_a$, $N \rightarrow^1 M' \{C_i/x_k\} \langle (C \setminus C_i)/x_1, \dots, x_{k-1} \rangle = N'$ and $\llbracket M' \{C_i/x_k\} \langle (C \setminus C_i)/x_1, \dots, x_{k-1} \rangle \rrbracket_u \succeq_{\dagger} Q_a$.

C. When $n > 2$:

Then $\llbracket N \rrbracket_u \rightarrow^2 Q_2 \rightarrow^l Q$, for $l \geq 1$. Also, $N \rightarrow^1 M$, $\llbracket M \rrbracket_u \succeq_{\dagger} Q_2$. $\exists \{P_j\}_{j \in J}, N', \text{ s.t. } M \rightarrow^* N'$, $\llbracket M \rrbracket \rightarrow^l P \rightarrow^* \{P_j\}_{j \in J}$ and $\llbracket N' \rrbracket_u \succeq_{\dagger} P_j \quad \forall j \in J$. We also have by Prop.2 that as $\llbracket M \rrbracket \succeq_{\dagger} Q_2$ and $\llbracket M \rrbracket \rightarrow^* \{Q_j\}_{j \in J}$ implies $\exists \{Q_i\}_{i \in I} \text{ s.t. } Q_2 \rightarrow^* \{Q_i\}_{i \in I}$, $I \text{ subseteq } J$ and $P_i \succeq_{\dagger} Q_i, \forall i \in I$

(b) For $m \geq 1$ and $n \geq 0$.

i. When $n = 0$.

Then $(\nu x)(R \mid x.\text{some}_{\text{fv}(N')}; \llbracket N' \rrbracket_x) = Q$ and $\llbracket M' \rrbracket_u \rightarrow^m R$ where $m \geq 1$. By the IH $\exists \{R_i\}_{i \in I}, M'', \text{ s.t. } M' \rightarrow^* M'', R \rightarrow^* \{R_i\}_{i \in I}$ and $\llbracket M'' \rrbracket_u \succeq_{\dagger} R_i \quad \forall i \in I$. Thus,

$$\begin{aligned}
\llbracket N \rrbracket_u &= (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\text{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \amalg_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \amalg_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} \llbracket M' \rrbracket_u \{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\} \cdots) \\
&\rightarrow^m (\nu z_1)(z_1.\text{some}_{\text{fv}(M_1)}; \llbracket M_1 \rrbracket_{z_1} \mid \cdots (\nu z_k)(z_k.\text{some}_{\text{fv}(M_k)}; \llbracket M_k \rrbracket_{z_k} \\
&\quad \mid \amalg_{x_{i_1} \in \{x_1, \dots, x_k\}} \cdots \amalg_{x_{i_k} \in \{x_1, \dots, x_k \setminus x_{i_1}, \dots, x_{i_{k-1}}\}} R \{z_1/x_{i_1}\} \cdots \{z_k/x_{i_k}\} \cdots) \\
&\rightarrow^n Q
\end{aligned}$$

Also, $Q \rightarrow^* \{(\nu x)(R_i \mid x.\text{some}_{\text{fv}(N')}; \llbracket N' \rrbracket_x)\}_{i \in I} = Q_i$, and the term can reduce as follows:

$N = M' \langle C/x_1, \dots, x_k \rangle \rightarrow^* M'' \langle C/x_1, \dots, x_k \rangle = N'$ and $\llbracket N' \rrbracket_u \succeq_{\dagger} Q_i \quad \forall i \in I$

ii. When $n \geq 1$. Then R has an occurrence of an unguarded $\bar{x}.\text{some}$ or $\bar{x}.\text{none}$, this case follows by IH and applying Proposition 2.

6. $N = M' \llbracket U/x^! \rrbracket$.

In this case, $\llbracket M' \llbracket U/x^! \rrbracket \rrbracket_u = (\nu x^!)(\llbracket M' \rrbracket_u \mid !x^!(x_i); \llbracket U \rrbracket_{x_i})$. Then,

$$\llbracket N \rrbracket_u = (\nu x^!)(\llbracket M' \rrbracket_u \mid !x^!(x_i); \llbracket U \rrbracket_{x_i}) \rightarrow^m (\nu x^!)(R \mid !x^!(x_i); \llbracket U \rrbracket_{x_i}) \rightarrow^n Q.$$

for some process R . Where \rightarrow^n is a reduction initially synchronises with $!x^!(x_i)$ when $n \geq 1$, $n + m = k \geq 1$. Type preservation in $\text{s}\pi^+$ ensures reducing $\llbracket M' \rrbracket_v \rightarrow^m$ doesn't consume possible synchronisations with $!x^!(x_i)$ if they occur. Let us consider the the possible sizes of both m and n .

(a) For $m = 0$ and $n \geq 1$.

In this case, $R = \llbracket M' \rrbracket_u$ as $\llbracket M' \rrbracket_u \longrightarrow^0 \llbracket M' \rrbracket_u$.

Notice that the only possibility of having an unguarded $?x^1[x_i]$ without internal reductions is when $\text{head}(M') = x[\text{ind}]$. By the diamond property we will be reducing each non-deterministic choice of a process simultaneously. Then we have the following:

$$\begin{aligned}
\llbracket N \rrbracket_u &= (\nu x^1)((\nu \tilde{y})(\llbracket x[\text{ind}] \rrbracket_j \mid P) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&= (\nu x^1)((\nu \tilde{y})(?x^1[x_i]; \overline{x_i}. \text{ind}; [x_i \leftrightarrow j] \mid P) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&\longrightarrow (\nu x^1)((\nu \tilde{y})(\nu x_i)(\overline{x_i}. \text{ind}; [x_i \leftrightarrow j] \mid \llbracket U \rrbracket_{x_i}) \mid P) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i} = Q_1 \\
&= (\nu x^1)((\nu \tilde{y})(\nu x_i)(\overline{x_i}. \text{ind}; [x_i \leftrightarrow j] \mid x_i. \text{case}(\text{ind}. \llbracket U_{\text{ind}} \rrbracket_{x_i})) \mid P) \\
&\quad \mid !x^1(x_i); \llbracket U \rrbracket_{x_i} \\
&\longrightarrow (\nu x^1)((\nu \tilde{y})(\nu x_i)([x_i \leftrightarrow j] \mid \llbracket U_{\text{ind}} \rrbracket_{x_i}) \mid P) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i} = Q_2 \\
&\longrightarrow (\nu x^1)((\nu \tilde{y})(\llbracket U_{\text{ind}} \rrbracket_j \mid P) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) = Q_3
\end{aligned}$$

We consider the two cases of the form of U_{ind} and show that the choice of U_{ind} is inconsequential

– When $U_{\text{ind}} = \lambda N^1$:

In this case, $N = M' \llbracket U/x^1 \rrbracket \longrightarrow M' \{N/x^1\} \llbracket U/x^1 \rrbracket = M$. and

$$\llbracket M \rrbracket_u = \llbracket M' \{N/x^1\} \llbracket U/x^1 \rrbracket \rrbracket_u = (\nu x^1)((\nu \tilde{y})(\llbracket \lambda N \rrbracket_j \mid P) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) = Q_3$$

– When $U_i = 1^1$:

In this case, $N = M' \llbracket U/x^1 \rrbracket \longrightarrow M' \{\text{fail}^0/x^1\} \llbracket U/x^1 \rrbracket = M$.

Notice that $\llbracket 1^1 \rrbracket_j = \overline{j}. \text{none}$ and that $\llbracket \text{fail}^0 \rrbracket_j = \overline{j}. \text{none}$. In addition,

$$\begin{aligned}
\llbracket M \rrbracket_u &= \llbracket M' \{\text{fail}^0/x^1\} \llbracket U/x^1 \rrbracket \rrbracket_u \\
&= (\nu x^1)((\nu \tilde{y})(\llbracket \text{fail}^0 \rrbracket_j \mid P) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \\
&= (\nu x^1)((\nu \tilde{y})(\llbracket 1^1 \rrbracket_j \mid P) \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) = Q_3
\end{aligned}$$

Both choices give an M that are equivalent to Q_3 .

i. When $n \leq 2$.

In this case, $Q = Q_n$ and $\llbracket N \rrbracket_u \longrightarrow^n Q_n$.

Also, $Q_n \longrightarrow^{3-n} Q_3 = Q'$, $N \longrightarrow^1 M = N'$ and $\llbracket M \rrbracket_u = Q_2$.

ii. When $n \geq 3$.

We have $\llbracket N \rrbracket_u \longrightarrow^3 Q_3 \longrightarrow^l Q$ for $l \geq 0$. We also know that $N \rightarrow M$, $Q_3 = \llbracket M \rrbracket_u$. By the IH, there exist $\{Q_i\}_{i \in I}$, N' such that $Q \longrightarrow^* \{Q_i\}_{i \in I}$, $M \longrightarrow^* N'$ and $\llbracket N' \rrbracket_u \succeq_{\perp} Q_i \quad \forall i \in I$. Finally, $\llbracket N \rrbracket_u \longrightarrow^3 Q_3 \longrightarrow^l Q \longrightarrow \{Q_i\}_{i \in I}$ and $N \rightarrow M \longrightarrow^* N'$.

(b) For $m \geq 1$ and $n \geq 0$.

(b.1) Case $n = 0$.

Then $(\nu x^1)(R \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) = Q$ and $\llbracket M' \rrbracket_u \longrightarrow^m R$ where $m \geq 1$. Then by the IH there exist $\{R'_i\}_{i \in I}$ and M'' such that $R \longrightarrow^* \{R'_i\}_{i \in I}$, $M' \longrightarrow^* M''$, and $\llbracket M'' \rrbracket_u \succeq_{\perp} R_i \quad \forall i \in I$. Hence we have that

$$\llbracket N \rrbracket_u = (\nu x^1)(\llbracket M' \rrbracket_u \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) \longrightarrow^m (\nu x^1)(R \mid !x^1(x_i); \llbracket U \rrbracket_{x_i}) = Q$$

We also know that

$$Q \longrightarrow^* \{(\nu x^1)(R'_i \mid !x^1(x_i); \llbracket U \rrbracket_{x_i})\}_{i \in I} = \{Q_i\}_{i \in I}$$

and so the λ_c term can reduce as follows: $N = M' \llbracket U/x^1 \rrbracket \longrightarrow^* M'' \llbracket U/x^1 \rrbracket = N'$ and $\llbracket N' \rrbracket_u \succeq_{\perp} Q_i \quad \forall i \in I$ via the \succeq_{\perp} rules

(b.2) Case $n \geq 1$.

Then R has an occurrence of an unguarded $?x^1[x_i]$, and the case follows by IH. \square

Theorem 10 (Loose Weak Soundness (Under \longrightarrow)). *If $\llbracket N \rrbracket_u \longrightarrow^* Q$ for a well-formed closed λ_c -term N , then there exist N' and Q' such that (i) $N \longrightarrow^* N'$ and (ii) $Q \longrightarrow^* Q'$ with $\llbracket N' \rrbracket_u \succeq_{\perp} Q'$.*

Proof. Immediate from Lemma 1. \square

H.3 Success Sensitivity

Theorem 23 (Success Sensitivity (Under \longrightarrow)). $M \Downarrow \checkmark_\lambda$ iff $\llbracket M \rrbracket_u \Downarrow \checkmark_\pi$ for well-formed closed terms M .

Proof. We proceed with the proof in two parts.

1. Suppose that $M \Downarrow \checkmark$. We will prove that $\llbracket M \rrbracket_u \Downarrow \checkmark$.

By Def. 17, there exists M' such that $M \longrightarrow^* M'$ and $\text{head}(M') = \checkmark$. By completeness, if $M \longrightarrow M'$ then there exist Q, Q' such that $\llbracket M \rrbracket_u \equiv Q \longrightarrow^* Q'$ and $\llbracket M' \rrbracket_u \succeq_{\#} Q'$.

We wish to show that there exists Q'' such that $Q' \longrightarrow^* Q''$ and Q'' has an unguarded occurrence of \checkmark .

By Proposition 4 (1) we have that $\text{head}(M') = \checkmark \implies \llbracket M' \rrbracket_u = (\nu \tilde{x})(P \mid \checkmark)$. Finally $\llbracket M' \rrbracket_u = (\nu \tilde{x})(P \mid \checkmark) \succeq_{\#} Q'$ hence Q must be of the form $(\nu \tilde{x})(P' \mid \checkmark)$ where $P \succeq_{\#} P'$. Hence Q reduces to a process that has an unguarded occurrence of \checkmark .

2. Suppose that $\llbracket M \rrbracket_u \Downarrow \checkmark$. We will prove that $M \Downarrow \checkmark$.

By operational soundness we have that if $\llbracket M \rrbracket_u \longrightarrow^* Q$ then there exist M' and $\{Q_i\}_{i \in I}$ such that: (i) $M \longrightarrow^* M'$ and (ii) $Q \longrightarrow^* \{Q_i\}_{i \in I}$ with $\llbracket M' \rrbracket_u \succeq_{\#} Q_j$, for all $j \in I$.

Since $\llbracket M \rrbracket_u \longrightarrow^* P_1 \# \dots \# P_k$ ($:= Q$), and $P_j = P'_j \mid \checkmark$ and $Q \longrightarrow^* \{Q_i\}_{i \in I}$ we must have that each Q_i is of the form $P'_{i_1} \# \dots \# P'_{i_k}$ with $P'_{i_j} = P''_{i_j} \mid \checkmark$. As $\llbracket M' \rrbracket_u \succeq_{\#} Q_j$, for all $j \in I$ we have that $\llbracket M' \rrbracket_u = (\prod_{i \in I} Q_i) \# R$ for some R . Finally applying Proposition 4 (2) we have that $\llbracket M' \rrbracket_u$ is itself a term with unguarded \checkmark , then M is itself headed with \checkmark . \square

I Proof of Separation of Lazy and Eager Semantics

Definition 19 (Dual Prefix). Given prefixes α and β (Definition 2), we say α and β are duals, denoted $\alpha \boxtimes \beta$, if and only if $\text{subj}\{\alpha\} \cap \text{subj}\{\beta\} \neq \emptyset$.

Lemma 18. Given $P \vdash \emptyset$, if $P \Downarrow_\alpha$, then there exist P', β such that $P \longrightarrow^* P' \Downarrow_\beta$ and $\alpha \boxtimes \beta$.

Proof. By well-typedness, there appears β in P with $\alpha \boxtimes \beta$. However, we may have $P \not\Downarrow_\beta$, because β is blocked by other prefixes. Hence, we need to find reductions from P such that we unblock β . However, the prefixes blocking β are connected to dual prefixes, that may be blocked themselves. The crux of this proof is thus to show that we can reduce P , such that we eventually unblock β .

The proof is by induction on the number of names that may block β (IH₁). Initially, this number corresponds to the total number of names appearing in P . Suppose β is blocked by n prefixes γ_i , where γ_n blocks β , and γ_1 is not blocked. We apply another layer of induction on n (IH₂).

In the inductive case, $n \geq 1$. The goal is to perform a reduction that synchronizes γ_1 with its dual, say $\overline{\gamma_1}$. The prefix $\overline{\gamma_1}$ may be blocked by a number of prefixes itself. However, the type system of $\text{s}\pi^+$ is based on CUT, so $\overline{\gamma_1}$ appears in parallel with the duals of $\gamma_2, \dots, \gamma_n$ and α . We then may apply IH₁ to find $P \longrightarrow^* P_0 \Downarrow_{\overline{\gamma_1}}$. We can then reduce P_0 by synchronizing between γ_1 and $\overline{\gamma_1}$: $P \longrightarrow^* P_0 \longrightarrow P_1$. In P_1 , β is blocked by one less prefix. Hence, by IH₂, $P \longrightarrow^* P_0 \longrightarrow P_1 \longrightarrow^* P' \Downarrow_\beta$, proving the thesis.

In the base case, β is not blocked: $P \Downarrow_\beta$. Let $P' := P$; trivially, $P \longrightarrow^* P' \Downarrow_\beta$, proving the thesis. \square

Theorem 11. Take $R \equiv \mathbb{N}[\alpha_1; (P \# Q)] \vdash \emptyset$ and $S \equiv \mathbb{N}[\alpha_2; P \# \alpha_3; Q] \vdash \emptyset$, where $\alpha_1 \boxtimes \alpha_2 \boxtimes \alpha_3$ and $\alpha_1, \alpha_2, \alpha_3$ require a continuation. Suppose that $P \not\sim_{\text{L}} Q$ and $P \not\sim_{\text{E}} Q$. Then (i) $R \sim_{\text{L}} S$ but (ii) $R \not\sim_{\text{E}} S$.

Proof. For (i), we construct a relation \mathbb{B} as follows:

$$\begin{aligned} \text{Id}^{\equiv} &:= \{(T, U) \mid T \equiv U\} \\ \mathbb{B}' &:= \{(T, U) \mid T \equiv \mathbb{M}[\beta_1; (V \# W)] \vdash \emptyset \text{ and} \\ &\quad U \equiv \mathbb{M}[\beta_2; V \# \beta_3; W] \vdash \emptyset \text{ and} \\ &\quad V \not\sim_{\text{L}} W \text{ and} \\ &\quad \beta_1 \boxtimes \beta_2 \boxtimes \beta_3 \text{ and} \\ &\quad \beta_1, \beta_2, \beta_3 \text{ require a continuation}\} \\ \mathbb{B} &:= \text{Id}^{\equiv} \cup \mathbb{B}' \end{aligned}$$

We prove that \mathbb{B} is a strong ready-prefix bisimulation w.r.t. the lazy semantics by proving the three conditions of Definition 9 for each $(T, U) \in \mathbb{B}$. We distinguish cases depending on whether $(T, U) \in \text{Id}^{\equiv}$ or $(T, U) \in \mathbb{B}'$.

– $(T, U) \in \text{Id}^{\equiv}$. The three conditions hold trivially.

- $(T, U) \in \mathbb{B}'$. Then $T \equiv \mathbf{M}[\beta_1; (V \parallel W)]$, $U \equiv \mathbf{M}[\beta_2; V \parallel \beta_3; W]$, $V \not\sim_{\mathbf{L}} W$, and $\beta_1 \bowtie \beta_2 \bowtie \beta_3$. We prove each condition separately.
 1. Suppose $T \rightsquigarrow T'$. Note that the hole in \mathbf{M} may appear inside a non-deterministic choice. We distinguish three cases:
 - (a) the reduction is inside \mathbf{M} and maintains the branch with the hole, (b) the reduction is inside \mathbf{M} and discards the branch with the hole, or (c) the reduction synchronizes on β_1 .
 - (a) The reduction is inside \mathbf{M} and maintains the branch with the hole. Then $T' \equiv \mathbf{M}'[\beta_1; (V \parallel W)]$ and $U \rightsquigarrow U' \equiv \mathbf{M}'[\beta_2; V \parallel \beta_3; W]$. Clearly, $(T', U') \in \mathbb{B}'$, so $(T', U') \in \mathbb{B}$.
 - (b) The reduction is inside \mathbf{M} and discards the branch with the hole. Then there exists U' such that $U \rightsquigarrow U' \equiv T'$, so $(T', U') \in \text{Id}^{\equiv}$, and thus $(T', U') \in \mathbb{B}$.
 - (c) The reduction synchronizes on β_1 . Then $T' \equiv \mathbf{M}'[V \parallel W]$ and, since $\beta_1 \bowtie \beta_2 \bowtie \beta_3$, $U \rightsquigarrow U' \equiv \mathbf{M}'[V \parallel W]$. Then $T' \equiv U'$, so $(T', U') \in \text{Id}^{\equiv}$ and thus $(T', U') \in \mathbb{B}$.
 2. Suppose $U \rightsquigarrow U'$. By reasoning analogous to above, $T \rightsquigarrow T'$ and $(T', U') \in \mathbb{B}$.
 3. Suppose $T \downarrow_{\gamma}$. If the prefix γ appears in \mathbf{M} , then clearly also $U \downarrow_{\gamma}$. Otherwise, $\gamma = \beta_1$. We have, e.g., $\gamma \bowtie \beta_2$ and clearly $U \downarrow_{\beta_2}$. The other direction is analogous.

It remains to show that $(R, S) \in \mathbb{B}$ which trivially holds.

For (ii), toward a contradiction, assume there exists a strong ready-prefix bisimulation w.r.t. $\longrightarrow \mathbb{B}$ where $(R, S) \in \mathbb{B}$.

By Lemma 18, there exist R', β_1 such that $R \longrightarrow^* R' \downarrow_{\beta_1}$, and $\alpha_1 \bar{\bowtie} \beta_1$. By the well-typedness of R and S , β_1 must appear in \mathbf{N} , and the reduction $R \longrightarrow^* R'$ takes place in \mathbf{N} . Take $x \in \text{subj}\{\alpha_1\} \cap \text{subj}\{\beta_1\}$ (which is non-empty by Lemma 18). Then $R' \equiv \mathbf{N}'_1[(\nu x)(\mathbf{N}'_2[\beta_1; R'_2] \mid \mathbf{N}'_3[\alpha_1; (P \parallel Q)])]$. Moreover, clearly $S \longrightarrow^* S'$ following the same reductions, resulting in $S' \equiv \mathbf{N}'_1[(\nu x)(\mathbf{N}'_2[\beta_1; R'_2] \mid \mathbf{N}'_3[\alpha_2; P \parallel \alpha_3; Q])]$; note that, since $\alpha_1 \bowtie \alpha_2 \bowtie \alpha_3$, also $\alpha_2 \bar{\bowtie} \beta_1$ and $\alpha_3 \bar{\bowtie} \beta_1$. At this point, we must have $(R', S') \in \mathbb{B}$.

The synchronization between β_1 and α_1 gives $R' \longrightarrow R'' \equiv \mathbf{N}''[P \parallel Q]$. Then by the bisimulation, there exists S'' such that $S' \longrightarrow S''$ with $(R'', S'') \in \mathbb{B}$. By clause 3 of the bisimulation, R'' and S'' must have the same ready-prefixes, so clearly the reduction $S' \longrightarrow S''$ results from a synchronization between β_1 and either of α_2 and α_3 . W.l.o.g., let us assume this was α_3 . Then $S'' \equiv \mathbf{N}''[Q]$. By assumption, $P \not\sim_{\mathbf{E}} Q$ and thus $P \parallel Q \not\sim_{\mathbf{E}} Q$, so clearly $R'' \not\sim_{\mathbf{E}} S''$. Hence, \mathbb{B} cannot be a strong ready-prefix bisimulation w.r.t. \longrightarrow . In other words, $R \not\sim_{\mathbf{E}} S$. \square