

Flavor- and CP-safe explanation of $g_\mu - 2$ anomaly

Jason L. Evans^a, Tsutomu T. Yanagida^{a,b}, and Norimi Yokozaki^c

^a*Tsung-Dao Lee Institute, Shanghai Jiao Tong University, Shanghai 200240, China*

^b*Kavli IPMU (WPI), UTIAS, University of Tokyo, Kashiwa, Chiba 277-8583, Japan*

^c*Zhejiang Institute of Modern Physics and Department of Physics, Zhejiang University, Hangzhou, Zhejiang 310027, China*

ABSTRACT

Supersymmetry is still a viable explanation for the muon $g - 2$ anomaly, if the sleptons and electroweak gauginos are $\mathcal{O}(100)$ GeV. However, for supersymmetry breaking masses this light, the SUSY flavor and CP-problem are exacerbated. To address this issue, we consider a flavor-safe gauge mediated explanation of the muon $g - 2$ with additional Higgs soft supersymmetry breaking mass parameters. The setup provides a generic parameter space within minimal gauge mediation. Furthermore, we show that the problematic CP violating phase can be dynamically suppressed. We find that gauge mediation models have large portions of parameter space where the muon $g - 2$ can be explained at 1σ level. The interplay between the slepton and the CP-odd Higgs masses also makes the majority of this model's parameter space testable at the LHC through searches for sleptons or additional Higgs bosons.

1 Introduction

The disagreement between the standard model prediction of $g_\mu - 2$ and the measured value has a long history, with the discrepancy first showing up more than 20 years ago [1]. The recent experimental measurement at Fermilab [2], when compared with recent theoretical predictions [3–7], has further increased the significance of this disagreement which is now at 4.2 standard deviations. Although not at the level of observation,¹ this deviation warrants our attention.

Although explaining $g_\mu - 2$ is not so difficult phenomenologically, it is non-trivial to find a model which is motivated. One of the best explanations for the deviation of $g_\mu - 2$, which is motivated beyond just its prediction of $g_\mu - 2$, is supersymmetry. In fact, $g_\mu - 2$ merely constrains the mass parameters of supersymmetry, since it already contains the necessary particles and order one couplings with the muon to explain $g_\mu - 2$ [9–11]. Furthermore, these constraints on the spectra prefer light soft masses and so tend to support the other nice features of supersymmetry not contradict them. For example, in [12–15] it was shown that supersymmetric dark matter could be realized in models which explain $g_\mu - 2$. And, of course, naturalness is unaffected by these constraints since it prefers light soft masses. Clearly, supersymmetry is a prime candidate for explaining $g_\mu - 2$. The main challenge for explaining $g_\mu - 2$ in supersymmetry is avoiding exclusion limits or theoretical problems. For example, the LHC slepton searches [16, 17], the chargino/neutralino searches [18–20] and the vacuum stability of the stau-Higgs potential [21–23] place strong constraints on the light needed slepton spectra. These constraints on the spectra are relatively easy to avoid if the correction to $g_\mu - 2$ arises from a chargino loop.² An important theoretical consideration, which we do not consider here, is can $g_\mu - 2$ be explained in models consistent with grand unification. It was shown that, at least in some unification models, it can be rectified [14, 28–31].

Although recent studies of the supersymmetric explanation for $g_\mu - 2$ are interesting, they tend to make assumptions about the sleptons and often neglect explaining key features of the needed spectrum like the lack of flavor and CP violation. These features of the spectrum are unavoidable when attempting to explaining $g_\mu - 2$, since the slepton masses must be less than about 1 TeV. If there are large amounts of flavor mixing or CP violation, the models are completely excluded. Since, generically, we expect order one flavor and CP violation in supersymmetry an explanation of their absence is needed. The flavor changing neutral current (FCNC) problem can be easily solved if supersymmetry breaking is mediated by gauge interactions.³ Gauge mediation provides a reliable solution to the FCNC problem based on four dimensional quantum field theory within perturbation theory. In gauge mediated models,

¹Some lattice calculations have weakened the significance of this deviation [8] but are ongoing.

²If the flavor universality of sleptons is violated and the staus are significantly heavier than the other sleptons at low energy, the LHC and vacuum stability constraints can be avoided even when $g_\mu - 2$ anomaly is explained by a bino-loop [23–25]. A low scale stau mass, which is heavy enough and does not lead to flavor problems, can be realized through loop effects if the down-type Higgs boson soft mass is tachyonic and large at the ultraviolet scale [26, 27].

³Gaugino mediation also offers a solution to the FCNC problem. However, it is in tension with the seesaw mechanism [32–34] if the sleptons are light enough to explain $g_\mu - 2$ [35].

since sfermion masses are generated by gauge interactions, they have no generation mixing and viable models can be found [36, 37] which avoid FCNC. CP violation, on the other hand, is not so simple of a problem to solve, even within gauge mediation since the gauginos and Higgs bilinear mass can still have order one CP violating phase.⁴ In fact, the constraints on the CP violating phase associated with these parameter are so severe that their phases need to be less than about 10^{-5} , as will be shown later, or the electric dipole moment of the electron will exceed experimental limits. Mindlessly setting the CP violation to zero in the SUSY breaking sector seems unnatural since we know that CP is almost maximally broken in the SM sector and generation of the CP violating phase in the Yukawa couplings from CP conserving models is non-trivial [39–43].

An important question that remains for the supersymmetric explanation of $g_\mu - 2$ is can an intelligent model produce the needed features in the slepton mass spectrum. This is the question we will examine here placing particular emphasis of the effects of CP violation. As we will take advantage of here, gauge mediation offers a dynamical means for suppressing the CP violating phases of the gaugino masses and Higgs B -term [44].⁵ This solves the last major hurdle for explaining $g_\mu - 2$, since gauge mediation already solves the FCNC problem and can be made consistent with all other relevant constraints. The aim of this work is to extend this original CP-safe gauge mediation model so that it can consistently explain the observed $g_\mu - 2$ anomaly. In particular, we will show that the CP-safe mechanism proposed in Ref. [44] becomes simpler and more generic by including anomaly mediated effects, which are usually neglected in gauge mediated models.

2 $g - 2$ anomaly and CP-violation

In this section, we review the gauge mediation model in [37] and its explanation of the $g_\mu - 2$ anomaly. At the end of this section, we will show that this model, like most models, has problematically large CP violating phases unless the phases are taken to be unnaturally small.

The gauge mediation sector of the model in [37] has an effective superpotential:

$$W = (\kappa_L Z + M_L) \Psi_L \Psi_{\bar{L}} + (\kappa_D Z + M_D) \Psi_D \Psi_{\bar{D}}, \quad (1)$$

where Z is a SUSY breaking field whose F -component, F_Z , has a non-zero vacuum expectation value, and Ψ_L and $\Psi_{\bar{D}}$ have the same SM charges as the left-handed leptons and down-type quarks, respectively, and $\Psi_{\bar{L}}$ and Ψ_D are their conjugate fields. We have formulated the messenger sector so that $(\Psi_{\bar{D}}, \Psi_L)$ transform as 5^* of $SU(5)_{\text{GUT}}$. However, we do not assume the existence of a grand unified theory (GUT). In fact, violation of the GUT relations, $\kappa_L/M_L \neq \kappa_D/M_D$,⁶ is necessary for explaining $g_\mu - 2$ and a 125 GeV Higgs boson

⁴In [28], it was shown that both the flavor and CP-problem are solved within gaugino mediation. The CP-problem is solved by assuming a shift symmetry where the imaginary part of the SUSY breaking field receives a constant shift [38]. See also [31] for another construction of a CP-safe gaugino mediation model.

⁵In Ref. [45], another mechanism to suppress the CP violating phases is proposed within gauge mediation.

⁶This combination is invariant under renormalization group evolution. Therefore, if we impose $\kappa_L = \kappa_D$ and $M_L = M_D$ at the ultraviolet scale (e.g., the GUT scale), $\kappa_L/M_L = \kappa_D/M_D$ holds at all scales.

mass becomes impossible due to competing requirements on the masses of sleptons and electroweak gauginos. This incongruence arises because the slepton masses are proportional to $\kappa_L \langle F_Z \rangle / M_L$ while the squark and gluino masses are proportional to $\kappa_D \langle F_Z \rangle / M_D$. The large stop mass needed to enhance the Higgs mass is inconsistent with light sleptons and the electroweak gaugino masses required for $g_\mu - 2$ unless $|\kappa_D/M_D| \gg |\kappa_L/M_L|$. Therefore, the GUT relations on these couplings and masses must be broken such that $|\kappa_D/M_D| \gg |\kappa_L/M_L|$. (We have neglected the $U(1)_Y$ gauge coupling in this discussion since it is a subleading effect, see Eqs. (5) and (7) below for the complete formulas.) Note, although one may consider messenger fields in other representations of $SU(5)_{\text{GUT}}$ such as $\mathbf{10}(+\mathbf{10})$ and a $\mathbf{24}$, gauge mediated models with these messenger fields are not advantageous for explaining $g_\mu - 2$ [36].

Here, we show the necessity of additional contributions to the Higgs soft masses beyond those provided by gauge mediation. These additional soft masses are vital for decoupling the explanation of a 125 GeV Higgs boson mass from the value of the μ parameter. To explain a Higgs boson mass of 125 GeV in gauge mediation, which has a relatively small stop trilinear mass term, the stop mass needs to be at least 5 TeV. Stop masses this large generate large radiative corrections to the Higgs soft masses. The electroweak symmetry breaking conditions then force $\mu \gtrsim 2$ TeV. With μ this large, the chargino-loop contribution to the $g_\mu - 2$ is quite suppressed. The bino loop contribution, on the other hand, is enhanced by $\mu \tan \beta$ and so might seem like a viable alternative for explaining $g_\mu - 2$. However, if the bino loop contribution to $g_\mu - 2$ is to explain the $g_\mu - 2$ anomaly, the sleptons need to be particularly light and $\mu \tan \beta$ needs to be rather large. Here, $\tan \beta$ is a ratio of the Higgs vacuum expectation values (VEVs). This spectrum is, unfortunately, excluded by the wino mass limits ($\gtrsim 660$ GeV) of the LHC [46] and/or the vacuum stability constraints from the stau-Higgs potential [21–23]. The strong limits on the wino mass arise because $|\kappa_D/M_D| \gg |\kappa_L/M_L|$ is needed to get the stop mass large enough to explain the Higgs boson mass. In this limit, the NLSP is an (almost) pure wino and so is long-lived. Long lived charge particles are strongly constrained by disappearing charged track searches at the LHC [46, 47]. Furthermore, if the trilinear coupling between the staus and Higgs boson, which is proportional to $\mu \tan \beta$, is too large, a charge breaking minimum which is deeper than the electroweak symmetry breaking minimum emerges. For staus as light as a few hundred GeV, $\mu \tan \beta$ is strongly constrained by the requirement that the life-time of the electroweak symmetry breaking minimum is longer than the age of the universe. By taking μ of $\mathcal{O}(100)$ GeV, both these problems are avoided. Then, the $g_\mu - 2$ can be explained by the chargino loop contribution. To achieve a small μ -term, our model generates an additional contribution to the soft SUSY breaking mass squared of the up-type Higgs, $m_{H_u}^2$, beyond the standard gauge mediated piece [36, 37, 48]. (See also Appendix A for an example of how to generate $m_{H_u}^2$ and $m_{H_d}^2$.)

As it turns out, the down type Higgs soft mass, $m_{H_d}^2$, also needs additional contributions to realize a viable model. In simple gauge mediation models, m_{H_d} is equal to the left-handed slepton mass at the messenger scale, which needs to be of $\mathcal{O}(100)$ GeV to explain $g_\mu - 2$. This is not simultaneously consistent with electroweak symmetry breaking conditions or the LHC constraints.

The problem with a small $m_{H_d}^2$ can be seen by examining one of the electroweak symmetry

breaking conditions, which we write as

$$m_{H/A/H^\pm}^2 \simeq |B_\mu(Q)|(\tan\beta + \cot\beta) \simeq m_{H_d}^2(Q) + |\mu(Q)|^2, \quad (2)$$

where H , A and H^\pm are the CP-even heavy Higgs, CP-odd Higgs and charged Higgs, respectively. The renormalization scale Q is taken to be around the stop mass scale, m_{stop} ; B_μ is defined by $V \ni B_\mu H_u H_d + h.c.$ with the superpotential $W = \mu H_u H_d$. Here, we neglect radiative corrections to the Higgs potential and terms suppressed by $\mathcal{O}(\tan^{-2}\beta)$. We have removed the argument of $B_\mu(Q)$ by the field redefinition of the Higgs fields, $|B_\mu(Q)| \exp(i\theta_{B_\mu}) \rightarrow |B_\mu(Q)|$, which rotates the phase of the μ term, $\mu \rightarrow \mu \exp(-i\theta_{B_\mu})$. In simple gauge mediation cases, $B_\mu(Q) = B(Q)\mu(Q)$ and $B(Q = m_{\text{stop}})$ is radiatively generated through the gaugino masses. Since we need $\tan\beta \gtrsim \mathcal{O}(10)$ to explain $g_\mu - 2$, $m_{H/A}$ should be at least 1 TeV to satisfy the LHC constraints [49]. The larger $\tan\beta$ is the heavier H and A need to be. For example, $m_{H/A}$ should be $\gtrsim 2$ TeV for $\tan\beta = 60$. In this case, $m_{H_d}^2(Q = m_{\text{stop}})$ needs to be $\mathcal{O}(10^6)$ GeV². This is not achievable in gauge mediation models unless there is an additional source of $m_{H_d}^2$. Lastly, we also consider an additional source for the B -term, since a $B_\mu(Q = m_{\text{stop}})$ generated by only gaugino loops in low scale gauge mediation tends to be inconsistent with the electroweak symmetry breaking and/or the LHC constraints on $m_{H/A}$. Generally, the process that generates a μ term will generate a B_μ term unless additional symmetries are imposed which forbid operators like $K = |Z|^2 H_u H_d + h.c.$.

In summary, the generic parameters of the gauge mediation model for explaining $g_\mu - 2$ are as follows:⁷

$$M_{\text{mess}}, \Lambda_D \equiv \kappa_D F_Z / M_D, r_L \equiv (\kappa_L / M_L) / (\kappa_D / M_D), \Delta m_{H_u}^2, \Delta m_{H_d}^2, B_\mu, N_5, \quad (3)$$

where $|M_D| = |M_L| = M_{\text{mess}}$ for simplicity; N_5 is the number of the messenger pairs ($\Psi_{\bar{D}}, \Psi_L$), which we included solely for generality. The Higgs soft mass parameters and B_μ can be related to μ , m_A and $\tan\beta$ through the electroweak symmetry breaking conditions. This gives us an effective set of parameters which we will use below of

$$M_{\text{mess}}, \Lambda_D \equiv \kappa_D F_Z / M_D, r_L \equiv (\kappa_L / M_L) / (\kappa_D / M_D), \mu, m_A, \tan\beta, N_5. \quad (4)$$

Furthermore, since we apply the electroweak symmetry breaking conditions at the weak scale, the boundary conditions on the μ , m_A , and $\tan\beta$ are set at the weak scale, and thus m_A corresponds to the physical mass of the pseudo-scalar Higgs boson.

In Eq. (3), Λ_D , r_L and B_μ are in general complex and their phases plus the phase of the μ parameter cannot be simultaneously removed by any combination of Higgs and messenger field redefinitions and a $U(1)_R$ transformation. Because of this, the predicted electron electric dipole moment (EDM) will easily exceeds the experimental bound [50] unless the phase is

⁷The previous models shown in Ref. [37] and [45], can only reproduce a portion of the parameter space presented here. In Ref. [37], only the case $m_{H_u}^2 = m_{H_d}^2$ (with $B \neq 0$ at the messenger scale) is considered while in Ref. [45], the authors only consider the case where the B -term is essentially zero at the messenger scale. The differences may seem slight at first glance, but the phenomenological and cosmological implications for gravitino dark matter and the masses of the heavy Higgs bosons can be quite extreme.

drastically suppressed. For the moment, we assume Λ_D , r_L , B_μ and μ can be made real. We will revisit this issue at the end of this section.

We now give expressions for the soft masses at the messenger scale, which are

$$M_{\tilde{b}} \simeq N_5 \frac{g_1^2}{16\pi^2} \Lambda_D \left(\frac{3}{5} r_L + \frac{2}{5} g(x) \right), \quad M_{\tilde{w}} \simeq N_5 \frac{g_2^2}{16\pi^2} r_L \Lambda_D, \quad M_{\tilde{g}} \simeq N_5 \frac{g_3^2}{16\pi^2} \Lambda_D g(x), \quad (5)$$

where g_1 , g_2 and g_3 are the SM gauge couplings of $U(1)_Y$, $SU(2)$ and $SU(3)$, $x = |\kappa_D F_Z / M_D^2|$ and $g(x)$ is a loop function given in Ref. [51],

$$g(x) = \frac{1}{x^2} [(1+x) \ln(1+x)] + (x \rightarrow -x), \quad (6)$$

and the factor of r_L in the gaugino masses comes from expanding $g(x)$ in the limit $r_L \ll 1$. The soft SUSY breaking masses for sleptons (\tilde{L}, \tilde{E}), squarks ($\tilde{Q}, \tilde{U}, \tilde{D}$) and Higgs fields (H_u, H_d) are

$$\begin{aligned} m_{\tilde{L}}^2 &\simeq N_5 \frac{|\Lambda_D|^2}{(16\pi^2)^2} \left(\frac{3}{2} |r_L|^2 g_2^4 + \frac{1}{4} g_1^4 \Lambda_{1,\text{eff}}^2 \right), \\ m_{\tilde{E}}^2 &\simeq N_5 \frac{|\Lambda_D|^2}{(16\pi^2)^2} (g_1^4 \Lambda_{1,\text{eff}}^2), \\ m_{\tilde{Q}}^2 &\simeq N_5 \frac{|\Lambda_D|^2}{(16\pi^2)^2} \left(\frac{8}{3} g_3^4 f(x) + \frac{3}{2} |r_L|^2 g_2^4 + \frac{1}{36} g_1^4 \Lambda_{1,\text{eff}}^2 \right), \\ m_{\tilde{U}}^2 &\simeq N_5 \frac{|\Lambda_D|^2}{(16\pi^2)^2} \left(\frac{8}{3} g_3^4 f(x) + \frac{4}{9} g_1^4 \Lambda_{1,\text{eff}}^2 \right), \\ m_{\tilde{D}}^2 &\simeq N_5 \frac{|\Lambda_D|^2}{(16\pi^2)^2} \left(\frac{8}{3} g_3^4 f(x) + \frac{1}{9} g_1^4 \Lambda_{1,\text{eff}}^2 \right), \\ m_{H_u}^2 &\simeq m_{\tilde{L}}^2 + \Delta m_{H_u}^2, \\ m_{H_d}^2 &\simeq m_{\tilde{L}}^2 + \Delta m_{H_d}^2, \end{aligned} \quad (7)$$

where

$$\begin{aligned} \Lambda_{1,\text{eff}}^2 &= \frac{6}{5} \left[\frac{3}{5} |r_L|^2 + \frac{2}{5} f(x) \right], \\ f(x) &= \frac{(1+x)}{x^2} \left[\ln(1+x) - 2\text{Li}_2 \left(\frac{x}{1+x} \right) + \frac{1}{2} \text{Li}_2 \left(\frac{2x}{1+x} \right) \right] + (x \rightarrow -x). \end{aligned} \quad (8)$$

The $|r_L|^2$ in the sfermion masses again comes from expanding $f(x)$ in the limit that $r_L \ll 1$

From these expressions, it is clear that light sleptons and electroweak gauginos are obtained for $|r_L| = |\kappa_L / \kappa_D| \ll 1$ while the squarks and gluino remain heavy. It is sometimes important to include the full function $g(x)$ and $f(x)$ for the contributions coming from Ψ_D and $\Psi_{\tilde{D}}$ but it is never important for the Ψ_L and $\Psi_{\tilde{L}}$ contribution since we will always take $|r_L| \ll 1$. We also see from these expressions that as N_5 increases the masses of the electroweak gauginos become larger for fixed slepton masses. This is particularly true when

M_{mess} is small.⁸ If we consider the LHC constraints on slepton masses, it becomes clear that it is marginally easier to explain $g_\mu - 2$ for $N_5 = 1$ than $N_5 > 1$. For $N_5 > 1$, we need an even larger $\tan\beta$, which in turn makes the staus lighter as we discuss later. Therefore, we will focus on the $N_5 = 1$ case in what follows.




Figure 1: The $M_{\tilde{w}} - \mu$ plane for $\Lambda_D = 900$ TeV. The black lines are contours of $m_{\tilde{\nu}_\mu}$ and the orange (yellow) shaded regions are consistent with $g_\mu - 2$ at the $1\text{-}\sigma$ ($2\text{-}\sigma$) level. The gray shaded region is excluded due to a tachyonic (or NLSP) stau. In the regions above the red solid lines, $m_{\chi_{2^\pm}} > m_{\tilde{\tau}_1}$. The blue-dotted regions are excluded due to the LHC constraint in Ref. [20]. The green lines correspond to $B_\mu(Q = M_{\text{mess}}) = 0$.

In Fig. 1, we show regions of parameter space which are consistent with $g_\mu - 2$ in the $M_{\tilde{w}}(Q = m_{\text{stop}}) - \mu$ plane.⁹ We take $\mu > 0$ and $r_L > 0$ (and $\Lambda_D > 0$). Otherwise, the relative signs of $M_{\tilde{w}}$ and $M_{\tilde{b}}$ is opposite leading to some cancellation between the chargino and bino-loop contributions. This gives a suppression of about $\sim 10\%$ to the prediction for $g_\mu - 2$.

⁸When M_{mess} is large, this effect is offset by the RG running of the soft masses

⁹In the figure, when we calculate $M_{\tilde{w}}$, the threshold correction to $g_2^2(Q = m_{\text{stop}})$ from the higgsino loop is evaluated with $\mu = 200$ GeV. In the range of μ taken in the figure, the difference between the corrections for $\mu = 200$ GeV and $\mu = 700$ GeV is less than 0.5%.




Figure 2: The lightest stau mass as a function of the messenger scale for different m_A , $m_A = (2300, 2500, 2700)$ GeV. In the left (right) panel, $\tan\beta = 40$ (50).




Figure 3: The maximum value of m_H (left) and corresponding $\tan\beta$ (right) as a function of $m_{\tilde{\nu}_\mu}$ when we require $19.2 \leq (a_\mu)_{\text{SUSY}} \times 10^{10}$ and the NLSP is a neutralino.

For these figures, we take $\Lambda_D = 900$ TeV, and r_L and μ are varied. The other parameters are taken to be $(M_{\text{mess}}/\text{GeV}, \tan \beta, m_A/\text{GeV}) = (10^6, 50, 2500), (10^8, 40, 2100), (10^{10}, 40, 2350)$ and $(10^{12}, 40, 2450)$ for the four figures.¹⁰ In the orange (yellow) regions, the $g_\mu - 2$ is explained at the 1σ (2σ) level.¹¹ For $m_{\tilde{\nu}_{e,\mu}} \gtrsim 660$ GeV, the LHC constraints on the first and second generation sleptons can be avoided [16, 17, 45]. In the gray region of the top-left panel, the stau is either the NLSP or tachyonic. This region is essentially excluded since the parameter sets where the stau is the NLSP and tachyonic are almost identical. For larger messenger scales, we have regions consistent with $B_\mu(Q = M_{\text{mess}}) = 0$, which are seen as the green lines. The blue dotted regions are excluded due to the constraints from LHC chargino/neutralino searches with $W/Z/h$ final states [20]. The constraints are applied when the left-handed stau is heavier than the wino. In the case where the wino-like chargino/neutralino decays to the stau or the tau sneutrino, the effective cross section with the $W/Z/h$ final states decreases to about 50% and the constraints in Ref. [20] are not directly applicable anymore. The predicted mass of the standard model-like Higgs (h) is ≈ 124.5 -125.5 GeV in the region where a_μ is in the 1σ range.¹² In the analysis, we use `SOFTSUSY 4.1.12` [54] to evaluate the SUSY mass spectra, `FeynHiggs 2.18.0`[55–61] to calculate the lightest Higgs boson mass, and `GM2Calc 2.1.0` [62, 63] to evaluate the SUSY contribution to the $g_\mu - 2$.

Now, we discuss the upper bound on $m_{H/A}$. A meaningful upper bound is obtained when the sleptons are heavier, since explaining the $g_\mu - 2$ then requires $\tan \beta$ to be pushed to larger values. The upper bound on $m_{H/A}$ then comes from the requirement that the lightest stau is not the next-to-lightest SUSY particle (NSLP) or tachyonic.¹³ Although the vacuum stability constraint from the stau-Higgs potential can be avoided easily in the small μ case, the soft SUSY breaking masses for the staus still give a meaningful constraint since the stau becomes the NSLP or even tachyonic when $\tan \beta$ and/or $m_{H_d}^2$ are large. This can be understood by examining the renormalization group equations:

$$\begin{aligned}
\frac{m_{H_d}^2}{d \ln Q} &\ni \frac{1}{16\pi^2} [6(m_{Q_3}^2 + m_{D_3}^2 + m_{H_d}^2)Y_b^2 + 2m_{H_d}^2 Y_\tau^2] \\
\frac{m_{L_3}^2}{d \ln Q} &\ni \frac{1}{16\pi^2} (2m_{H_d}^2 Y_\tau^2) \\
\frac{m_{E_3}^2}{d \ln Q} &\ni \frac{1}{16\pi^2} (4m_{H_d}^2 Y_\tau^2),
\end{aligned} \tag{9}$$

where Y_b and Y_τ are the bottom and tau Yukawa couplings, respectively; m_{Q_3} and m_{D_3} (m_{L_3} and m_{E_3}) are left and right handed sbottom (stau) masses. From the first equation

¹⁰We take these values of m_A so that the $B_\mu = 0$ line appears on the figure. The other aspects of these figures are fairly insensitive to these small changes.

¹¹There are regions where the higgsino and wino are as light as 150-200 GeV. In these regions, the predicted W -boson mass can be larger than the SM prediction by 3-6 MeV [52, 53]. The large mass shift recently reported by CDFII is very hard to explain in the present model. This will be checked in future experiments.

¹²For the small μ case, the negative contribution to m_h^2 from the sbottom loop is proportional to $(\mu \tan \beta)^4$ and so suppressed. Therefore, the stop mass required to explain a 125 GeV Higgs boson mass becomes somewhat smaller.

¹³These are effectively the same upper bound.

we see that $m_{H_d}^2(Q = M_{\text{mess}})$ must be much larger than $m_{H_d}^2(Q = m_{\text{stop}})$ as it receives large corrections from sbottom loops.¹⁴ This is particularly true when $\tan\beta$ is large since Y_b and Y_τ grow with $\tan\beta$. Since the stau masses receive corrections proportional to $|Y_\tau|^2 m_{H_d}^2$, they tend to be suppressed as the messenger scale is increased. In Fig. 2, the lightest stau mass as a function of the messenger scale is shown. The stau mass monotonically decreases as the messenger scale increases. For large $m_{H/A} \sim m_{H_d}(Q = m_{\text{stop}})$ and large $\tan\beta$, the staus mass will be too small or tachyonic. As the slepton masses are increased, the required $\tan\beta$ for explaining $g_\mu - 2$ becomes larger leading to an upper bound on $m_{H/A}$. Since the mass suppression discussed above is definitely smaller for the lower messenger scales, we will take a low messenger scale as a reference point. (The complications are much more severe for larger SUSY breaking mediation scales like the Planck or GUT scales. In this sense, gauge mediation is advantageous for explaining $g_\mu - 2$ via the chargino loop contribution.)

In the left panel of Fig. 3, we show the upper bound on $m_H(\simeq m_A)$ as a function of the muon sneutrino mass, $m_{\tilde{\nu}_\mu}$. The right panel shows the value of $\tan\beta$ for the maximal value of m_H . In these figures, we scan the following parameter ranges:

$$0.1 \leq r_L \leq 0.3, \quad 20 \leq \tan\beta \leq 70, \quad 1000 \leq m_A/\text{GeV} \leq 6000. \quad (10)$$

We require that $(a_\mu)_{\text{SUSY}}$ explain the experimental value of the $g_\mu - 2$ at 1σ level and the NLSP is a neutralino. The messenger scale and μ are fixed to be $M_{\text{mess}} = 10^6 \text{ GeV}$ and $\mu = 200 \text{ GeV}$ so that the chargino contribution to the $g_\mu - 2$ is (almost) maximized and the negative contribution to the square of the stau masses from renormalization group running is minimized. We also take $\Lambda_D = 900 \text{ TeV}$. This shows that for larger $m_{\tilde{\nu}_\mu}$, the maximum value of $m_{H/A}$ becomes smaller while $\tan\beta$ becomes larger. In particular, for $m_{\tilde{\nu}_\mu} \gtrsim 900 \text{ GeV}$, we get $m_{H/A} \leq 3 \text{ TeV}$ and $\tan\beta \gtrsim 56$. This shows the possibility to detect the heavy Higgs bosons, H and A , at the LHC, using the channel $H/A \rightarrow \tau\tau$ is quite good since Y_b and Y_τ are quite large.

In table 1, we show the mass spectra, $(a_\mu)_{\text{SUSY}}$, the life-time of the lightest chargino, and production cross section (bbH) of H/A at $\sqrt{s} = 14 \text{ TeV}$ and branching ratios of H/A for four different mass spectra. We also show the required $\Delta m_{H_u}^2$, $\Delta m_{H_d}^2$ and B_μ at the messenger scale. The production cross sections are calculated using `SusHi 1.7.0` package [64, 65] with `NNPDF4.0` [66] used for the parton distribution functions. At the point P4, the production cross section of H/A is quite large. In fact, P4 is marginally consistent with the current bound from the heavy Higgs search [49] and it is expected that the point will be checked soon [67].

Concerning the constraints from the chargino/neutralino searches at the LHC, for P2 and P4, with the left-handed stau lighter than the wino, about 26-27% of wino-like charginos and neutralinos (χ_2^\pm) decay to the stau or tau sneutrino. Therefore, the exclusion limits using the $W/Z/h$ final states become weaker as the effective cross section decreases to about 50%. Note, we cannot directly apply the exclusion limit in Ref. [20]. In order to check the LHC constraints, including those in Ref. [20], we used `SModelS` package [68–75] with the wino production cross sections at NLO+NLL estimated using `resumino` [76–82]. We found P2 and P4 were not excluded.

¹⁴Because the weak gauginos are light, they have a subdominant effect.

Lastly, we discuss the problem of CP violation. Since the smuons and electroweak gauginos need to be as light as $\mathcal{O}(100)$ GeV to explain the $g_\mu - 2$ and we consider gauge mediation, the selectrons must be light as well. The electron (EDM) constraints then require B_μ, r_L and Λ_D to be almost completely real.

To analyze the electron EDM, we rotate the fields so that $B_\mu(Q)$ is real and $\langle H_u^0 \rangle$ and $\langle H_d^0 \rangle$ are in turn real and positive. We then take $\mu = |\mu| \exp(-i\theta_B)$. In this basis, the potentially problematic phases are:¹⁵

$$\begin{aligned}\theta_{\tilde{w}} &= \text{Arg}(M_{\tilde{w}}|\mu| \exp(-i\theta_B)) = \text{Arg}(M_{\tilde{w}}(B_\mu/\mu)^*), \\ \theta_{\tilde{b}} &= \text{Arg}(M_{\tilde{b}}(B_\mu/\mu)^*),\end{aligned}\tag{11}$$

where all relevant parameters are defined at the low energy scale. The phases of these mass combinations are important because the same diagrams with the same combination of masses contribute to $g_\mu - 2$ and the electron (EDM) calculation.¹⁶ As is the case for the $g_\mu - 2$, the chargino diagram gives the largest contribution to the electron EDM and $\theta_{\tilde{w}}$ is most strongly constrained.

Furthermore, since a_e and the electric dipole moment of the electron, d_e , come from the real and imaginary parts of the same amplitude, $\tan(\text{Arg}(\text{Amplitude}))a_e$ is equal to $2m_e d_e/e$. Combining these relations, we find

$$\begin{aligned}(d_e/e)_{\text{SUSY}} &\simeq \tan \theta_{\tilde{w}} \frac{m_e}{2m_\mu^2} (a_\mu)_{\text{SUSY}} \\ &\approx \theta_{\tilde{w}} \left(\frac{(a_\mu)_{\text{SUSY}}}{2.2 \times 10^{-9}} \right) \times 10^{-24} \text{ cm},\end{aligned}\tag{12}$$

where we used the fact that the chargino contribution dominates in our model.¹⁷

In order to satisfy the ACME experimental constraint [50], $|d_e/e| < 1.1 \times 10^{-29}$ cm, $\theta_{\tilde{w}}$ should be smaller than about 10^{-5} . The constraint on $\theta_{\tilde{b}}$ is somewhat weaker, $\theta_{\tilde{b}} \leq \mathcal{O}(10^{-4})$, since the bino loop contribution only makes up $\mathcal{O}(10\%)$ of the anomalous magnetic moment calculation. It is extremely difficult to achieve such a small CP violating phase unless the complex arguments of $\Lambda_D, r_L \Lambda_D$ and B_μ/μ are identical to a very high level of precision. The requirement on Λ_D is due to the fact that the complex arguments of $M_{\tilde{b}}, M_{\tilde{w}}, M_{\tilde{g}}$ are mixed through two-loop RG running. Thus, the phases of all gaugino masses feed into the phase of B_μ/μ from RG running. This leaves only one possible conclusion; the complex arguments of $M_{\tilde{b}}, M_{\tilde{w}}, M_{\tilde{g}}$ and B_μ/μ must be aligned at some higher energy scale, i.e. the messenger scale.

¹⁵Although the complex argument of the scalar trilinear coupling, the A -term, is also relevant, we focus on the above mentioned arguments as the contribution to the EDM from picking up the A -term is not enhanced by $\tan \beta$.

¹⁶The one-loop SUSY contribution to a_e , the anomalous magnetic moment of the electron, is $(m_e/m_u)^2$ times a_μ as the selectron and smuon masses are nearly degenerate.

¹⁷The phase $\theta_{\tilde{b}}$ is also constrained by the electron EDM measurement, but to a lesser degree. However, the model we give below will suppress all problematic phases.

To estimate this effect, we can approximate the Higgs B -term at the low scale as [83]

$$\begin{aligned}
B_\mu(Q)/\mu(Q) &\approx (B_\mu/\mu) - \frac{1}{16\pi^2} \left(6g_2^2 M_{\tilde{w}} + \frac{6}{5}g_1^2 M_{\tilde{b}} \right) \ln \frac{M_{\text{mess}}}{Q} \\
&+ \frac{1}{8\pi^4} g_3^2 (Y_t^2 + Y_b^2) M_{\tilde{g}} \left[\ln \frac{M_{\text{mess}}}{Q} \right]^2,
\end{aligned} \tag{13}$$

as long as the messenger scale is not too high. Also, we have taken $Q = m_{\text{stop}}$. Although the gluino mass contribution is loop suppressed compared to the wino and bino mass contribution, the loop suppression of the gluino contribution is offset by a larger mass and larger couplings and so can be comparable or even larger than the bino and wino contributions in the parameter space of interest. From Eq. (13) it becomes clear, $B_\mu(Q)/\mu(Q)$ picks up a portion of all the complex arguments of the gaugino masses. This, combined with the RG mixing of the gaugino masses, tells us we need a model that predicts the complex arguments of Λ_D , $r_L \Lambda_D$ and B_μ/μ are aligned. How to achieve the needed alignment/suppression of the phases of Λ_D , $r_L \Lambda_D$, B_μ and μ will be explained in the next section.

3 A CP-safe gauge mediation model for the $g-2$ anomaly

In this section, we give an explicit model, which explains $\theta_{\tilde{w}} = \theta_{\tilde{b}} = 0$, after the gravitino mass is rotated to be real and positive, $m_{3/2} \in \mathbb{R}_{>0}$. Our model is based on a discrete symmetry, $Z(3)_1 \times Z(3)_2 \times Z(3)_3 \times Z_{4R}$ with the following superpotential interactions:

$$W = \frac{\lambda_i}{3} S_i^3 + k_i S_i \Psi_i \bar{\Psi}_i (i = 1, 2, 3). \tag{14}$$

We also introduce tachyonic soft SUSY breaking masses for S_i of

$$V_{\text{soft}} = -m_{S_i}^2 |S_i|^2. \tag{15}$$

An example model for generating $m_{S_i}^2$ is given in Appendix A. Here, $\Psi_1 = H_u$ ($\bar{\Psi}_1 = H_d$), $\Psi_2 = \Psi_L$, $\Psi_3 = \Psi_{\bar{D}}$, where $\Psi_{\bar{D}}$ and Ψ_L are the anti-quark and lepton-like messengers with charge assignments of a 5^* representation in an $SU(5)_{\text{GUT}}$ theory. However, we do not assume grand unification in this paper. The charges of S_i , Ψ_i and $\bar{\Psi}_i$ under $Z(3)_1 \times Z(3)_2 \times Z(3)_3 \times Z_{4R}$ are given in Table 2 in Appendix A. The discrete R symmetry, Z_{4R} , is anomaly free with one pair of $\Psi_{2,3}$ and $\bar{\Psi}_{2,3}$ [84]. Therefore it could be a gauge symmetry and left unbroken even with gravitational effects [85–90] included. The λ_i can be made real and positive by a field redefinition of the S_i without any loss of generality. Using a $U(1)_R$ rotation, we take the gravitino mass, $m_{3/2} = \langle W \rangle^* / M_{\text{Pl}}^2$, to be real and positive. Due to the tachyonic mass, both the scalar and F component of the superfield S_i obtain a non-zero VEV, giving the SUSY preserving and SUSY breaking masses for the messenger and Higgs fields. Note that, as shown in the previous section, there are some regions of parameter space which can explain $g_\mu - 2$ and the Higgs B -term vanishes at the messenger scale. In this case, we only introduce a higgsino mass term, $\mu H_u H_d$, instead of S_1 , and the Z_{4R} charge of $H_u H_d$ should be 2. Since this modification is straightforward, we do not include any details here.

Eqs. (14) and (15) give the following tree-level potential for S_i :

$$V = \lambda_i^2 |S_i|^4 - |m_{S_i}^2| |S_i|^2, \quad (16)$$

with a minimum at $\langle |S_i| \rangle = \sqrt{|m_{S_i}^2| / \sqrt{2\lambda_i^2}}$.

The phase direction of the $\langle S_i \rangle$ are not determined by the tree-level potential and are only determined once loop corrections are taken into account. At the tree level, the scalar potential for S_i in the framework of supergravity is given by

$$V = F_{S_i}^\dagger F_{S_i} - 3|W|^2/M_{\text{PL}}^2, \quad (17)$$

where

$$F_{S_i} = - [\lambda_i S_i^{*2} + S_i W^* / M_{\text{PL}}^2], \quad (18)$$

and we have assumed that the SUSY breaking field has a negligibly small scalar vacuum expectation value. We will return to this point later. As is clear from these expressions, the scalar potential in Eq. (17) does not determine $\text{Arg}(S_i)$.

The phases of the vevs are determined by the A -term contributions to the potential induced by anomaly mediated SUSY breaking (AMSB),

$$V = (A_i)_{\text{amsb}} \frac{\lambda_i}{3} S_i^3 + h.c. = \frac{m_{3/2}}{16\pi^2} (16\pi^2 \beta_{\lambda_i} / \lambda_i) \frac{\lambda_i}{3} S_i^3 + h.c., \quad (19)$$

where β_{λ_i} is the beta-function for λ_i , given by

$$(16\pi^2) \beta_{\lambda_i} / \lambda_i = 3(N_c |k_i|^2 + 2\lambda_i^2), \quad (20)$$

and $N_c = 2$ for $i = 1, 2$ and $N_c = 3$ for $i = 3$. In the previous paper [44], the effects from anomaly mediation were not considered.¹⁸ For S_1 , the tree-level contribution, $(A_1)_{\text{tree}}$, shown in Appendix C may also be considered, which dominates over the contribution from anomaly mediation $(A_1)_{\text{amsb}}$ and has the opposite sign. By writing $\langle S_i \rangle = |\langle S_i \rangle| e^{i\theta_{S_i}}$ ($\theta_{S_i} = \text{Arg}(\langle S_i \rangle)$), in Eq. (19) (and Eq. (56)), θ_{S_i} has a potential

$$V \ni A_i \frac{\lambda_i}{3} S_i^3 + h.c. \rightarrow \frac{2}{3} A_i |\langle S_i \rangle|^3 \lambda_i \cos(3\theta_{S_i}), \quad (21)$$

where A_i corresponds to $(A_{2,3})_{\text{amsb}}$ or $(A_1)_{\text{amsb}} + (A_1)_{\text{tree}}$, and is a real number. The above potential fixes $3\theta_{S_i} = \pi$ ($3\theta_{S_i} = 0$) for $A_i > 0$ ($A_i < 0$). Since the SUSY masses for Ψ_i are

¹⁸In the previous paper, the authors introduced higher dimensional operators, $K \ni |S_i|^4 / \Lambda_4^2$, to fix the phase direction of the S_i . If the cut-off Λ_4 is larger than $\sim 10 \langle S_i \rangle$, the anomaly mediation effects are dominant.

$k_i \langle S_i \rangle$ and the B -terms are $k_1 \lambda_1 \langle S_1^{*2} \rangle$ for $i = 1$ and $-k_i \lambda_i \langle S_i^{*2} \rangle$ for $i = 2$ and 3 ,¹⁹ we have

$$\begin{aligned} B_\mu/\mu &= \pm \lambda_1 |\langle S_1 \rangle|, \\ \Lambda_L (= r_L \Lambda_D) &= \lambda_2 |\langle S_2 \rangle|, \\ \Lambda_D &= \lambda_3 |\langle S_3 \rangle|, \end{aligned} \tag{22}$$

where B_μ/μ is positive (negative) when the tree level contribution (AMSB contribution) to S_1^3 term dominates. All of the above parameters are real. Therefore, there is no CP violating phase.

However, the above mechanism is not viable generically. This is because, as shown in the previous section, the phases of the A_i 's should be aligned at the order of 10^{-5} . The required conditions are summarized as follows:

- For SUSY breaking fields which are charged under some symmetry,²⁰ $|\langle Z_i^* F_{Z_i} \rangle / M_{\text{PL}}^2| \ll 10^{-7} |m_{3/2}|$.
- For moduli type SUSY breaking fields (e.g. axion) where the shift symmetry breaking is small, mixings with other SUSY breaking fields could be problematic. This effect is safe if the previous condition is satisfied.
- There are no true singlets, i.e. no charge under any symmetry, SUSY breaking fields since they generically will have an F -term of $O(m_{3/2} M_{\text{PL}})$ and will couple to S_i , as well as other fields, via a dimension 5 operator with a complex coefficient.

To derive these conditions, we have assumed that higher dimensional operators containing the SUSY breaking fields, matter fields and/or moduli fields are suppressed by the Planck scale. If these operators are suppressed by a lower mass scales than the Planck scale, the conditions of above become even more restrictive. For a class of motivated models, we have checked that our CP-safe mechanism still holds (see Appendix B).

To realize the above conditions on our mechanism, we assume that the SUSY breaking field, Z , which has the largest vev, i.e. $|\langle F_Z \rangle| \sim |m_{3/2} M_{\text{PL}}|$, is charged under some symmetry so that it cannot couple to the matter fields (e.g. S_i , Ψ_i , $\bar{\Psi}_i$ and the quark and lepton chiral superfields) through operators like $K = c_{IJ} Z Q_I^\dagger Q_J / M_{\text{PL}} + h.c.$ and $W = c'_{IJK} Z Q_I Q_J Q_K / M_{\text{PL}}$, and gauge fields through operators like $\int d^2\theta c_k Z W^k W^k / M_{\text{PL}} + h.c.$ ($k = \{U(1)_Y, SU(2)_L, SU(3)_c\}$). These operators are dangerous since they will generate additional contributions to the A_i with unaligned phases. If Z couples to S_i through these operators, it leads to a tree-level $A_i \sim m_{3/2}$ and an unconstrained phase, which clearly spoils our mechanism for solving the CP problem. If Z couples to Ψ_i or $\bar{\Psi}_i$, it induces A_i at the

¹⁹Following the standard convention, the definitions of the B -terms are $W = (\mu - B_\mu \theta^2) H_u H_d$ for the Higgs fields and $W = ((M_{\text{mess}})_{2,3} + B_{2,3} \theta^2) \Psi_{2,3} \bar{\Psi}_{2,3}$ for the messenger fields. Note that for $\lambda_1 \sim k_1 \sim 1$, S_1 significantly mixes with the Higgs doublets after electroweak symmetry breaking. Therefore, it affects the electroweak symmetry breaking as well as the collider and cosmological/astrophysical phenomena, which should be carefully investigated (see e.g., Refs. [91, 92]). On the other hand, for $\lambda_1, k_1 \ll 1$, S_1 decouples from the standard model, and the effects are limited.

²⁰SUSY breaking field means any field whose F component has a non-zero vacuum expectation value.

one-loop level. This contribution is comparable to that from anomaly mediation in Eq. (19). It, therefore, generically will spoil our mechanism. For the case where Z couples to the gauge fields, A_i is induced at the two-loop level which is still dangerous since it can give an $\mathcal{O}(10^{-2})$ correction to the phase of Eq. (19). Lastly, when Z couples to the quarks and leptons, it generates a three-loop level correction to A_i , which can still be dangerous. Thus, the supersymmetry breaking field must not be a true singlet.

Thus far, we have argued that operators like $K = Z|Q_I|^2 + h.c.$ must be forbidden. However, even a minimal Kähler potential $K = |Z|^2 + |Q_I|^2$ can be problematic. In this case, we still have effective couplings between $|Z|^2$ and $|Q_I|^2$, through the supergravity corrections to potential, $f = -3M_{\text{PL}}^2 \exp[-K/(3M_{\text{PL}}^2)]$. This induces Planck suppressed operators which generate trilinear couplings of order $A_i = -\langle Z \rangle^* \langle F_Z \rangle / M_{\text{PL}}^2$. For $|\langle Z \rangle| \sim M_{\text{PL}}$, $A_i = \mathcal{O}(m_{3/2})$. At first glance, this might seem fine since this generates the same A -term for each S_i , phase and all. However, the anomaly mediated contribution, which always exists, is dependent on λ_i^2 and $|k_i|^2$ and so is not universal for the A_i . The sum of these contributions then gives a unique phase for each A_i , all of which cannot be removed by field redefinitions. Thus, we restrict ourselves to models with $|\langle Z \rangle| \ll 10^{-7} M_{\text{PL}}$.²¹

For a SUSY breaking field, Z' , with $\langle F'_Z \rangle \ll \langle F_Z \rangle$, the above arguments still apply, but the effects are suppressed by $\langle F'_Z \rangle / \langle F_Z \rangle$. This SUSY breaking field, Z' , is required to satisfy $|\langle Z'^* F'_Z \rangle| / M_{\text{PL}}^2 \ll 10^{-7} |m_{3/2}|$. If Z' is a singlet of the symmetries, $|\langle F'_Z \rangle / M_{\text{PL}}| \ll 10^{-7} |m_{3/2}|$ is further required. However this condition is very unlikely to be satisfied as this singlet field has $K = c' M_{\text{PL}} Z' + h.c.$ ($c' = \mathcal{O}(1)$), which in turn leads to $W = c' m_{3/2} M_{\text{PL}} Z'$ after a Kähler transformation.

If Z' is a moduli-type field and the shift symmetry is not broken, it will couple to matter and SUSY breaking fields with real coefficients such as $K = c_I^n (Z' + Z'^*)^n |Q'_I|^2 / M_{\text{PL}}^n$, where Q'_I represents the matter and SUSY breaking fields that are charged under some symmetry. Including this term in the Kähler potential gives

$$\begin{aligned} -e^{K/2} \langle F'_Z \rangle &= (K^{-1})_{xx} \left\langle \frac{\partial K}{\partial x} W^* / M_{\text{PL}}^2 \right\rangle \\ &+ (K^{-1})_{xi} \langle W_i^* + K_i^* W^* / M_{\text{PL}}^2 \rangle, \end{aligned} \quad (23)$$

where $x = Z' + Z'^*$. The first term is proportional to $m_{3/2} M_{\text{PL}}$ with a real coefficient, and the second term is $\sim \langle Z^* \cdot F_Z \rangle / M_{\text{PL}} \ll 10^{-7} m_{3/2} M_{\text{PL}}$. Since the couplings to the matter fields, including S_i , are given by real constants this does not induce different phases for different S_i .

Lastly, let us discuss the cosmological domain wall problem. In our model, domain walls

²¹If $\langle Z \rangle$ is non-zero, the F -term of the conformal compensator Φ becomes,

$$\langle F \rangle = \left(m_{3/2} + \frac{1}{3M_{\text{PL}}^2} \left\langle \frac{\partial K}{\partial Z} F_Z \right\rangle \right),$$

where $\Phi = \phi(1 + F\theta^2)$ and we choose the gauge $\phi = \exp(\langle K \rangle / 6M_{\text{PL}}^2)$ to recover Einstein gravity. In this case the anomaly mediated effects are changed slightly. However, this does not affect our discussion since we require the second term to be smaller than $\mathcal{O}(10^{-7})m_{3/2}$ for any supersymmetry breaking field.

are formed once the Z_{3_i} symmetries are spontaneously broken. The domains eventually follow a scaling solution and about one domain wall will exist within a Hubble horizon [93–96]. In this case, their energy density can be approximated by $\sigma H^{-2}/H^{-3} = \sigma H$, where σ is the domain wall tension and H is the Hubble parameter. By comparing this to the total energy density, $3H^2 M_{\text{PL}}^2$, it turns out the domain walls dominate the energy density of the universe for

$$H_{\text{dom}} \sim \frac{\sigma}{M_{\text{PL}}^2} \sim \frac{\langle S_i \rangle^3}{M_{\text{PL}}^2}. \quad (24)$$

If the domain walls are stable enough to last until today, the energy density of the domain walls will dominate. In order to avoid this situation, we need a small Z_{3_i} breaking term, which removes the degeneracy of the three vacua by an amount we label ΔV . The domain walls collapse when their energy density becomes similar in size to ΔV . The conditions

$$H_{\text{col}} \sim \frac{\Delta V}{\sigma} \sim \frac{\Delta V}{\langle S_i \rangle^3} > H_{\text{dom}}, \quad (25)$$

should then be satisfied. Also, it is safer to require the domain walls collapse before big-bang nucleosynthesis (BBN) starts, which gives

$$H_{\text{col}} > \frac{\mathcal{O}(\text{MeV}^2)}{M_{\text{PL}}}. \quad (26)$$

ΔV should then satisfy the following condition:

$$\Delta V > \max \left(\frac{\langle S_i \rangle^6}{M_{\text{PL}}^2}, \frac{\mathcal{O}(\text{MeV}^2)}{M_{\text{PL}}} \langle S_i \rangle^3 \right). \quad (27)$$

Let us consider the following Z_{3_i} breaking term, which conserves Z_{4R} symmetry,

$$W \ni \mu_i^2 S_i \rightarrow V \ni (\mu_i^2 \lambda_i S_i^{2*} - m_{3/2} \mu_i^2 S_i) + h.c. \quad (28)$$

The first term gives the dominant contribution to ΔV . The conditions on ΔV lead to

$$|\mu_i^2| > \mathcal{O}(10^{-13}) \text{GeV}^2 \left(\frac{|\langle S_i \rangle|}{10^6 \text{GeV}} \right)^4, \quad (29)$$

and

$$|\mu_i^2| > \mathcal{O}(10^{-22}) \text{GeV}^2 \left(\frac{|\langle S_i \rangle|}{300 \text{GeV}} \right). \quad (30)$$

The latter condition from BBN is always weaker than the former one for $\langle S_2 \rangle$ and $\langle S_3 \rangle$. For $\langle S_1 \rangle$, the BBN constraint can be stronger. On the other hand, in order not to induce dangerous CP violating phases, $|\mu_i^2|$ is bounded from above as

$$|\mu_i^2| \ll \mathcal{O}(10^{-5}) (A_i)_{\text{AMSB}} |\langle S_i \rangle| \approx \mathcal{O}(10^{-3}) \text{GeV}^2 \left(\frac{m_{3/2}}{10 \text{MeV}} \right) \left(\frac{|\langle S_i \rangle|}{10^6 \text{GeV}} \right). \quad (31)$$

These conditions Eqs. (29)-(31) are satisfied as long as $M_{\text{mess}} < \mathcal{O}(10^{10}\text{-}10^{11} \text{ GeV})$. Here, $m_{3/2} = 10 \text{ MeV}$ is a reference value for $S_{2,3} \simeq 10^6 \text{ GeV}$ and $m_{3/2}$ is roughly proportional to $\langle S_{2,3} \rangle = M_{\text{mess}}/k_{2,3}$.

We need to be a bit careful in our treatment of S_1 as it mixes with the Higgs doublets, which could potentially affect Higgs physics. Focusing on

$$W = \frac{\lambda_1}{3} S_1^3 + k_1 S_1 H_u H_d, \quad (32)$$

the superpotential and soft SUSY breaking terms are exactly the same as those of Z_3 invariant Next-to-Minimal Supersymmetric Standard Model (NMSSM) (see Refs. [97, 98] for a review). The CP-even Higgs mass matrix contains

$$\begin{aligned} V &\ni \frac{1}{2} m_h^2 h^2 + v k_1 [2\mu - \sin 2\beta (B_{\text{eff}} + \lambda_1 \langle S_1 \rangle)] h s + \frac{1}{2} (4\lambda_1^2 \langle S_1 \rangle^2 + \frac{k_1 A'_1}{2} \frac{v^2}{\langle S_1 \rangle} \sin 2\beta) s^2, \\ &\approx \frac{1}{2} m_h^2 h^2 + (2k_1 v \mu) h s + \frac{1}{2} (2B_{\text{eff}} - 2A'_1)^2 s^2, \end{aligned} \quad (33)$$

where $h = -\sin \beta h_u^0 + \cos \beta h_d^0$ and $m_h \approx 125 \text{ GeV}$; $\mu = k_1 \langle S_1 \rangle$, $B_{\text{eff}} = B_\mu/\mu \simeq A'_1 + \lambda_1 \langle S_1 \rangle$, $V \ni A'_1 k_1 S H_u H_d + h.c.$; $A'_1 \sim 100 \text{ GeV}$ through gaugino loops at the SUSY mass scale; Here, we neglect A_1 . The Higgs B -term, B_{eff} , is estimated to be

$$B_{\text{eff}} \simeq \frac{m_A^2}{\mu \tan \beta} \approx 296 \text{ GeV} \times \left(\frac{m_A}{2000 \text{ GeV}} \right)^2 \left(\frac{300 \text{ GeV}}{\mu} \right) \left(\frac{45}{\tan \beta} \right). \quad (34)$$

To avoid the large mixing between h and s , $k_1 < \mathcal{O}(0.1)$, is required, and $\lambda_1 \sim k_1$ due to $B_{\text{eff}} \sim \mu$. The mass of the singlet-like CP-odd Higgs is

$$m_{a_1}^2 \simeq \frac{k_1 \sin 2\beta}{2} (4B_{\text{eff}} - 3A'_1) \frac{v^2}{\langle S_1 \rangle} - 3A_1 \lambda_1 \langle S_1 \rangle \quad (35)$$

$$\approx -3A_1 \lambda_1 \langle S_1 \rangle \approx 3(B_{\text{eff}} - A'_1) |A_1|, \quad (36)$$

which can be larger than $\mathcal{O}(1) \text{ GeV}$. This a_1 as well as the imaginary parts of $S_{2,3}$ (i.e. axion-like particles) can safely decay before the BBN.

4 Discussion and conclusions

We presented a CP-safe version of the gauge mediated model in [37] by extending the mechanism proposed in Ref. [44] to include anomaly mediation effects. First, we showed that large portions of the parameter space are still consistent with the $g_\mu - 2$ and LHC constraints. For heavy sleptons, the CP-odd Higgs and the Heavy CP-even Higgs tend to be light and, therefore, discoverable at the LHC. This analysis is for a generic minimal gauge mediation model with messengers in the fundamental representation of $SU(5)$.

We also showed that the dangerous CP violating phases can be drastically suppressed by applying the mechanism first proposed in [44]. This mechanism allows us to dynamically

suppress the phases of all relevant supersymmetry breaking parameters. This mechanism aligns the soft mass phases with that of $m_{3/2}$ through anomaly mediated A -terms in the messengers and Higgs bosons potential. Then, when the gravitino phase is rotated away all soft masses become real to a high level of precision leaving no dangerous CP violating phase. Furthermore, in Appendix A, we give a consistent ultraviolet model for this mechanism.

Acknowledgments

We thank Andre Lessa for useful discussions about ATLAS-SUSY-2018-41 constraints and the usage of `SmodelS`. N. Y. is supported by a start-up grant from Zhejiang University. J. L. E. is supported by a start-up grant from Shanghai Jiao-Tong University. T. T. Y. is supported in part by the China Grant for Tal ent Scientific Start-Up Project and by Natural Science Foundation of China (NSFC) under grant No. 12175134 as well as by World Premier International Research Center Initiative (WPI Initiative), MEXT, Japan.

A Generation of soft SUSY breaking masses beside the gauge mediation

Let us first discuss the tachyonic mass generation for S_i ($i = 1 \dots 3$), which determines the messenger mass scale, the messenger B -terms (therefore $\Lambda_{L,D}$), the μ -term and the Higgs B -term. For this purpose, we introduce new Yukawa couplings for the S_i ,

$$W = \frac{k'_i}{2} S_i E_i^2, \quad (37)$$

where E_i are gauge singlet fields. The tachyonic masses for S_i are generated by one-loop diagrams involving E_i , if the soft SUSY breaking mass squared of E_i is positive. This is analogous to how the stop loops generate a tacyonic Higgs soft SUSY breaking mass which generates electroweak symmetry breaking radiatively.

A positive supersymmetry breaking mass is generated for E_i from the following superpotential:

$$W = \frac{\kappa}{2} Z \Psi^2 + M_0 \Psi \bar{\Psi} + M_i E'_i \bar{E}'_i + g_i E_i E'_i \Psi + W_{\text{YIT}}, \quad (38)$$

where $M_0 \gg M_i$ is assumed, and Ψ , $\bar{\Psi}$, E'_i and \bar{E}'_i are all gauge singlet fields. A consistent set of charge assignment for these fields under $Z(3)_1 \times Z(3)_2 \times Z(3)_3 \times Z_{4R}$ is shown in table 2. We assume the non-anomalous Z_{4R} is a discrete gauge symmetry while the others are global symmetries. The charges of the matter multiplets in the standard model sector under $Z(3)_1 \times Z(3)_2 \times Z(3)_3 \times Z_{4R}$ are $\mathbf{5}^*(-2, 0, 0, 1)$, $\mathbf{10}(-1, 0, 0, 1)$ and $N(0, 0, 0, 1)$. Here, we again use $SU(5)$ notation for simplicity and N is for the right-handed neutrinos. These charge assignments are consistent with the standard model Yukawa interactions.

The last term in Eq.(38), W_{IYIT} , represents the effective superpotential for dynamical SUSY breaking based on an IYIT model [99, 100] with an $SP(1) = SU(2)$ gauge theory. After the quark superfields condense, the superpotential takes the following form:

$$W_{\text{IYIT}} = \frac{1}{2}(Q_1 Q_2 + Q_3 Q_4)Z = \Lambda^2 Z. \quad (39)$$

Here, Q_a ($a = 1 \dots 4$) are doublet quarks, and Λ is the dynamical scale. This superpotential breaks supersymmetry, i.e. $F_Z = \Lambda^2$. The IYIT model with the $SP(1)$ gauge symmetry has a discrete $Z(4)$ symmetry, where the doublet quarks Q_a carry +1 charges and Z has +2. Accordingly, the gauge singlet fields, Ψ , $\bar{\Psi}$, E'_i , \bar{E}'_i carry charges -1 , $+1$, $+1$ and -1 , respectively. All other fields carry a charge of zero under this $Z(4)$ symmetry. Importantly, this discrete $Z(4)$ symmetry prohibits a linear term involving $\bar{\Psi}$ and mass mixing between E'_i and E_i . The E_i and \bar{E}'_i can still decay before BBN through Planck-suppressed operators which break $S(3)_i$ and $Z(4)$ through operators such as $K = E_i^\dagger H_u H_d / M_{\text{PL}}$, $\bar{E}'_i^\dagger H_u H_d / M_{\text{PL}}$.

The positive squared masses for E'_i , which are generated after integrating out $\Psi, \bar{\Psi}$, are estimated using the one-loop formula for the effective Kähler potential [101]. The field dependent mass matrix for $\Psi, \bar{\Psi}$ and E'_i is

$$\mathcal{M} = \begin{pmatrix} \kappa Z & M_0 & g_i E_i \\ M_0 & 0 & 0 \\ g_i E_i & 0 & 0 \end{pmatrix}. \quad (40)$$

The matrix, $\mathcal{M}^\dagger \mathcal{M}$, has two non-zero eigenvalues, m_+^2 and m_-^2 , which are

$$m_\pm^2 = \frac{1}{2} \left(2|M_0|^2 + 2|g_i E_i|^2 + |\kappa Z|^2 \pm \sqrt{|\kappa Z|^2} \sqrt{|\kappa Z|^2 + 4(|M_0|^2 + |g_i E_i|^2)} \right). \quad (41)$$

Using this mass matrix, the effective Kähler potential can be written as

$$\begin{aligned} K_{\text{eff}} &= -\frac{1}{32\pi^2} \text{Tr} \left(\mathcal{M}^\dagger \mathcal{M} \ln \frac{\mathcal{M}^\dagger \mathcal{M}}{e^{c_r} Q^2} \right) \\ &= -\frac{1}{32\pi^2} \left(m_+^2 \ln \frac{m_+^2}{e^{c_r} Q^2} + m_-^2 \ln \frac{m_-^2}{e^{c_r} Q^2} \right). \end{aligned} \quad (42)$$

where Q is the renormalization scale and c_r is a scheme dependent constant. We then obtain the following soft SUSY breaking masses for $m_{E_i}^2$:

$$m_{E_i}^2 \simeq - \frac{\partial^2 K_{\text{eff}}}{\partial(|E_i|^2) \partial(|Z|^2)} \Big|_{|E_i|^2=|Z|^2=0} |F_Z|^2 = \frac{|g_i \kappa|^2}{32\pi^2} \left| \frac{F_Z}{M_0} \right|^2. \quad (43)$$

Thanks to the fact that this mass squared is positive, a tachyonic mass for the S_i is generated by E_i loops:²²

$$m_{S_i}^2 \simeq - \frac{|k'_i|^2}{16\pi^2} m_{E_i}^2 \ln \frac{|M_0|^2}{|m_{E_i}^2|}. \quad (44)$$

²²The operator in the Kähler potential giving the tachyonic mass, $K \sim 1/(16\pi^2)^2 |Z|^2 |S_i|^2 / M_0^2$, also induce A-terms for S_i , with $\langle Z \rangle = \mathcal{O}(m_{3/2})$ induced by effects from supergravity. However, $\langle Z F_Z^* \rangle$ is real, and it does not change the phase direction of $\langle S_i \rangle$.

To generate the soft masses for H_u and H_d , we introduce additional pairs of massive Higgs multiplets H'_u and \bar{H}'_u , and H'_d and \bar{H}'_d . The $Z(4)$ charges of H'_u , \bar{H}'_u , H'_d and \bar{H}'_d are $-1, +1, -1$ and $+1$, respectively. The superpotential for these massive Higgs multiplets is given by

$$W = \lambda_u H_u \bar{H}'_u \Psi + \lambda_d H_d \bar{H}'_d \Psi + M_u \bar{H}'_u H'_u + M_d \bar{H}'_d H'_d. \quad (45)$$

The superpotential which generates the Higgs soft masses is then the same as Eq. (38) with $E_i \rightarrow H_{u,d}$, $E'_i \rightarrow \bar{H}'_{u,d}$, and $\bar{E}'_i \rightarrow H'_{u,d}$. In the limit $M_u \sim M_d \ll M_0$, we then get

$$\Delta m_{H_{u,d}}^2 \simeq \frac{|\lambda_{u,d}\kappa|^2}{32\pi^2} \left| \frac{F_Z}{M_0} \right|^2. \quad (46)$$

B Some motivated extensions

Let us consider the following superpotential:

$$W = \lambda_{X_0} X_0 (X \bar{X} - v^2) + \lambda_X X \psi \bar{\psi}, \quad (47)$$

where X_0 and $\Psi \bar{\Psi}$ has a $U(1)_R$ (or a discrete \mathbb{R}) charge of 2. We consider the case where X and \bar{X} break some symmetries like a global $U(1)_{\text{PQ}}$ or gauged $U(1)_{B-L}$. As we will show,

$$B_{\psi\bar{\psi}}/M_{\psi\bar{\psi}} = \text{real constant} \times m_{3/2} = \text{real}, \quad (48)$$

where the B -term and SUSY preserving mass terms are given by

$$\mathcal{L} = B_{\psi\bar{\psi}} \psi \bar{\psi} + h.c., \quad M_{\psi\bar{\psi}} = \lambda_X \langle X \rangle. \quad (49)$$

Because the ratio of these terms is real, this extension has no effect on our mechanism for suppressing CP phases. If this were not the case, these extensions could generate dangerously large CP phases. For instance, in the case of a $U(1)_{\text{PQ}}$, a gluino mass is induced at the one-loop level and in turn generate S_3^3 potential term at the three-loop level. For $U(1)_{B-L}$ breaking, an A -term is induced for $LH_u \bar{N}$ at the one-loop level and S_1^3 is induced at the three-loop level as well.

The B -term is given by

$$\mathcal{L} = - [(\lambda_{X_0} X_0 \bar{X})^* + a' m_{3/2} X] \lambda_X \psi \bar{\psi} + h.c., \quad (50)$$

where a' is a real coefficient given by anomaly mediation. First, we assume $\langle X \rangle \sim \langle \bar{X} \rangle \sim v$, which is satisfied when the soft SUSY breaking masses for X and \bar{X} are the same order, $\mathcal{O}(m_{3/2})$, and from the minimization conditions we get $\text{Arg} \langle X \bar{X} \rangle = \text{Arg}(v^2)$. The potential for X_0 is

$$V = m_{X_0}^2 |X_0|^2 + (-2m_{3/2} \lambda_{X_0} v^2 X_0 + a'' m_{3/2} \lambda_{X_0} X_0 X \bar{X}) + h.c., \quad (51)$$

where a'' is a real coefficient induced by anomaly mediation and $m_{X_0}^2 = 2|\lambda_{X_0}|^2(|X|^2 + |\bar{X}|^2)$ with $|X| = |\bar{X}| = v$. The VEV of X_0 is fixed to be

$$\langle |X_0| \rangle \simeq |\lambda_{X_0} m_{3/2}| \quad \text{and} \quad \text{Arg}(\langle X_0 \rangle) = -\text{Arg}(v^2) - \text{Arg}(\lambda_{X_0}), \quad (52)$$

so that

$$\text{Arg}((\lambda_{X_0} X_0 \bar{X})^*) = \text{Arg}(\langle X \rangle), \quad (53)$$

and $B_{\psi\bar{\psi}}/M_{\psi\bar{\psi}}$ is real.

C The tree-level contribution to A_1

Here, we show a model which generates a tree-level contribution to A_1 with the opposite sign of $(A_1)_{\text{AMSB}}$. The superpotential is given by

$$W = k_Y S_1^2 Y / 2 + M_Y Y \bar{Y} + \lambda_1 S_1^3 / 3. \quad (54)$$

The scalar potential is

$$\begin{aligned} V &= |M_Y|^2 |Y|^2 + |S_1|^2 |\lambda_1 S_1 + k_Y Y|^2 + \frac{1}{4} |k_Y S_1^2 + 2M_Y \bar{Y}|^2 \\ &\quad - (m_{3/2} M_Y \bar{Y} + h.c.) + \mathcal{O}(M_{\text{PL}}^{-2}). \end{aligned} \quad (55)$$

By integrating out Y and \bar{Y} using the equations of motion, $\frac{\partial V}{\partial Y} = 0$ and $\frac{\partial V}{\partial \bar{Y}} = 0$, the effective scalar potential is

$$\begin{aligned} V_{\text{eff}} &= \lambda_1^2 |S_1|^4 \frac{|M_Y|^2}{|M_Y|^2 + |k_Y|^2 |S_1|^2} - \left[\frac{3}{2} m_{3/2} \lambda_1 S_1^3 / 3 \frac{|k_Y|^2 |S_1|^2}{|M_Y|^2 + |k_Y|^2 |S_1|^2} + h.c. \right] \\ &= \lambda_1^2 |S_1|^4 \frac{|M_Y|^2}{|M_Y|^2 + |k_Y|^2 |S_1|^2} - [(A_1)_{\text{tree}} \lambda_1 S_1^3 / 3 + h.c.], \end{aligned} \quad (56)$$

where we have neglected $\mathcal{O}(m_{3/2}^2)$ terms. It can be seen that the sign of S_1^3 is opposite to AMSB. The A -term in Eq. (56) can be larger than $(A_1)_{\text{amsb}}$. For instance, by taking $|M_Y|^2 = 10|k_Y|^2|S_1|^2$, the A -term is $\approx -0.14m_{3/2}$, which is larger than that of AMSB.

References

- [1] **Muon g-2** Collaboration, H. N. Brown *et al.*, ‘‘Precise measurement of the positive muon anomalous magnetic moment,’’ *Phys. Rev. Lett.* **86** (2001) 2227–2231, [arXiv:hep-ex/0102017](#).
- [2] **Muon g-2** Collaboration, B. Abi *et al.*, ‘‘Measurement of the Positive Muon Anomalous Magnetic Moment to 0.46 ppm,’’ *Phys. Rev. Lett.* **126** no. 14, (2021) 141801, [arXiv:2104.03281 \[hep-ex\]](#).
- [3] T. Aoyama *et al.*, ‘‘The anomalous magnetic moment of the muon in the Standard Model,’’ *Phys. Rept.* **887** (2020) 1–166, [arXiv:2006.04822 \[hep-ph\]](#).
- [4] G. Colangelo, M. Hoferichter, and P. Stoffer, ‘‘Two-pion contribution to hadronic vacuum polarization,’’ *JHEP* **02** (2019) 006, [arXiv:1810.00007 \[hep-ph\]](#).

- [5] M. Hoferichter, B.-L. Hoid, and B. Kubis, “Three-pion contribution to hadronic vacuum polarization,” *JHEP* **08** (2019) 137, [arXiv:1907.01556 \[hep-ph\]](#).
- [6] M. Davier, A. Hoecker, B. Malaescu, and Z. Zhang, “A new evaluation of the hadronic vacuum polarisation contributions to the muon anomalous magnetic moment and to $\alpha(\mathbf{m}_Z^2)$,” *Eur. Phys. J. C* **80** no. 3, (2020) 241, [arXiv:1908.00921 \[hep-ph\]](#). [Erratum: *Eur.Phys.J.C* 80, 410 (2020)].
- [7] A. Keshavarzi, D. Nomura, and T. Teubner, “ $g - 2$ of charged leptons, $\alpha(M_Z^2)$, and the hyperfine splitting of muonium,” *Phys. Rev. D* **101** no. 1, (2020) 014029, [arXiv:1911.00367 \[hep-ph\]](#).
- [8] S. Borsanyi *et al.*, “Leading hadronic contribution to the muon magnetic moment from lattice QCD,” *Nature* **593** no. 7857, (2021) 51–55, [arXiv:2002.12347 \[hep-lat\]](#).
- [9] J. L. Lopez, D. V. Nanopoulos, and X. Wang, “Large $(g-2)$ -mu in SU(5) x U(1) supergravity models,” *Phys. Rev. D* **49** (1994) 366–372, [arXiv:hep-ph/9308336](#).
- [10] U. Chattopadhyay and P. Nath, “Probing supergravity grand unification in the Brookhaven $g-2$ experiment,” *Phys. Rev. D* **53** (1996) 1648–1657, [arXiv:hep-ph/9507386](#).
- [11] T. Moroi, “The Muon anomalous magnetic dipole moment in the minimal supersymmetric standard model,” *Phys. Rev. D* **53** (1996) 6565–6575, [arXiv:hep-ph/9512396](#). [Erratum: *Phys.Rev.D* 56, 4424 (1997)].
- [12] M. Endo, K. Hamaguchi, S. Iwamoto, and K. Yanagi, “Probing minimal SUSY scenarios in the light of muon $g - 2$ and dark matter,” *JHEP* **06** (2017) 031, [arXiv:1704.05287 \[hep-ph\]](#).
- [13] P. Cox, C. Han, and T. T. Yanagida, “Muon $g - 2$ and dark matter in the minimal supersymmetric standard model,” *Phys. Rev. D* **98** no. 5, (2018) 055015, [arXiv:1805.02802 \[hep-ph\]](#).
- [14] P. Cox, C. Han, T. T. Yanagida, and N. Yokozaki, “Gaugino mediation scenarios for muon $g - 2$ and dark matter,” *JHEP* **08** (2019) 097, [arXiv:1811.12699 \[hep-ph\]](#).
- [15] P. Cox, C. Han, and T. T. Yanagida, “Muon $g-2$ and coannihilating dark matter in the minimal supersymmetric standard model,” *Phys. Rev. D* **104** no. 7, (2021) 075035, [arXiv:2104.03290 \[hep-ph\]](#).
- [16] **ATLAS** Collaboration, G. Aad *et al.*, “Searches for electroweak production of supersymmetric particles with compressed mass spectra in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector,” *Phys. Rev. D* **101** no. 5, (2020) 052005, [arXiv:1911.12606 \[hep-ex\]](#).

- [17] CMS Collaboration, A. M. Sirunyan *et al.*, “Search for supersymmetry in final states with two oppositely charged same-flavor leptons and missing transverse momentum in proton-proton collisions at $\sqrt{s} = 13$ TeV,” *JHEP* **04** (2021) 123, [arXiv:2012.08600 \[hep-ex\]](#).
- [18] ATLAS Collaboration, G. Aad *et al.*, “Search for chargino–neutralino pair production in final states with three leptons and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector,” *Eur. Phys. J. C* **81** no. 12, (2021) 1118, [arXiv:2106.01676 \[hep-ex\]](#).
- [19] CMS Collaboration, A. Tumasyan *et al.*, “Search for chargino-neutralino production in events with Higgs and W bosons using 137 fb^{-1} of proton-proton collisions at $\sqrt{s} = 13$ TeV,” *JHEP* **10** (2021) 045, [arXiv:2107.12553 \[hep-ex\]](#).
- [20] ATLAS Collaboration, G. Aad *et al.*, “Search for charginos and neutralinos in final states with two boosted hadronically decaying bosons and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector,” *Phys. Rev. D* **104** no. 11, (2021) 112010, [arXiv:2108.07586 \[hep-ex\]](#).
- [21] J. Hisano and S. Sugiyama, “Charge-breaking constraints on left-right mixing of stau’s,” *Phys. Lett. B* **696** (2011) 92–96, [arXiv:1011.0260 \[hep-ph\]](#). [Erratum: *Phys.Lett.B* 719, 472–473 (2013)].
- [22] T. Kitahara and T. Yoshinaga, “Stau with Large Mass Difference and Enhancement of the Higgs to Diphoton Decay Rate in the MSSM,” *JHEP* **05** (2013) 035, [arXiv:1303.0461 \[hep-ph\]](#).
- [23] M. Endo, K. Hamaguchi, T. Kitahara, and T. Yoshinaga, “Probing Bino contribution to muon $g - 2$,” *JHEP* **11** (2013) 013, [arXiv:1309.3065 \[hep-ph\]](#).
- [24] M. Ibe, T. T. Yanagida, and N. Yokozaki, “Muon $g-2$ and 125 GeV Higgs in Split-Family Supersymmetry,” *JHEP* **08** (2013) 067, [arXiv:1303.6995 \[hep-ph\]](#).
- [25] M. Ibe, M. Suzuki, T. T. Yanagida, and N. Yokozaki, “Muon $g - 2$ in Split-Family SUSY in light of LHC Run II,” *Eur. Phys. J. C* **79** no. 8, (2019) 688, [arXiv:1903.12433 \[hep-ph\]](#).
- [26] M. Yamaguchi and W. Yin, “A novel approach to finely tuned supersymmetric standard models: The case of the non-universal Higgs mass model,” *PTEP* **2018** no. 2, (2018) 023B06, [arXiv:1606.04953 \[hep-ph\]](#).
- [27] W. Yin and N. Yokozaki, “Splitting mass spectra and muon $g - 2$ in Higgs-anomaly mediation,” *Phys. Lett. B* **762** (2016) 72–79, [arXiv:1607.05705 \[hep-ph\]](#).
- [28] K. Harigaya, T. T. Yanagida, and N. Yokozaki, “Higgs boson mass of 125 GeV and $g - 2$ of the muon in a gaugino mediation model,” *Phys. Rev. D* **91** no. 7, (2015) 075010, [arXiv:1501.07447 \[hep-ph\]](#).

- [29] J. Ellis, J. L. Evans, N. Nagata, D. V. Nanopoulos, and K. A. Olive, “Flipped $\mathfrak{g}_\mu - 2$,” *Eur. Phys. J. C* **81** no. 12, (2021) 1079, [arXiv:2107.03025 \[hep-ph\]](#).
- [30] J. Ellis, J. L. Evans, N. Nagata, D. V. Nanopoulos, and K. A. Olive, “Flipped SU(5) GUT phenomenology: proton decay and $\mathfrak{g}_\mu - 2$,” *Eur. Phys. J. C* **81** no. 12, (2021) 1109, [arXiv:2110.06833 \[hep-ph\]](#).
- [31] K. Agashe, M. Ekhterachian, Z. Liu, and R. Sundrum, “Sleptonic SUSY: From UV Framework to IR Phenomenology,” [arXiv:2203.01796 \[hep-ph\]](#).
- [32] T. Yanagida, in “Proceedings of the Workshop on Unified Theory and Baryon Number of the Universe,” eds. O. Sawada and A. Sugamoto (KEK, 1979), *Conf. Proc. C* **7902131** (1979) 95-99.
- [33] T. Yanagida, “Horizontal Symmetry and Mass of the Top Quark,” *Phys. Rev. D* **20** (1979) 2986.
- [34] M. Gell-Mann, P. Ramond, and R. Slansky, “Complex Spinors and Unified Theories,” *Conf. Proc. C* **790927** (1979) 315–321, [arXiv:1306.4669 \[hep-th\]](#).
- [35] R. Nagai and N. Yokozaki, “Lepton flavor violations in SUSY models for muon $g - 2$ with right-handed neutrinos,” *JHEP* **01** (2021) 099, [arXiv:2007.00943 \[hep-ph\]](#).
- [36] T. T. Yanagida and N. Yokozaki, “Muon $g - 2$ in MSSM gauge mediation revisited,” *Phys. Lett. B* **772** (2017) 409–414, [arXiv:1704.00711 \[hep-ph\]](#).
- [37] G. Bhattacharyya, T. T. Yanagida, and N. Yokozaki, “An extended gauge mediation for muon ($g - 2$) explanation,” *Phys. Lett. B* **784** (2018) 118–121, [arXiv:1805.01607 \[hep-ph\]](#).
- [38] S. Iwamoto, T. T. Yanagida, and N. Yokozaki, “CP-safe gravity mediation and muon $g - 2$,” *PTEP* **2015** (2015) 073B01, [arXiv:1407.4226 \[hep-ph\]](#).
- [39] A. E. Nelson, “Naturally Weak CP Violation,” *Phys. Lett. B* **136** (1984) 387–391.
- [40] S. M. Barr, “Solving the Strong CP Problem Without the Peccei-Quinn Symmetry,” *Phys. Rev. Lett.* **53** (1984) 329.
- [41] M. Dine, R. G. Leigh, and A. Kagan, “Supersymmetry and the Nelson-Barr mechanism,” *Phys. Rev. D* **48** (1993) 2214–2223, [arXiv:hep-ph/9303296](#).
- [42] G. Hiller and M. Schmaltz, “Strong Weak CP Hierarchy from Nonrenormalization Theorems,” *Phys. Rev. D* **65** (2002) 096009, [arXiv:hep-ph/0201251](#).
- [43] J. Evans, C. Han, T. T. Yanagida, and N. Yokozaki, “Complete solution to the strong CP problem: Supersymmetric extension of the Nelson-Barr model,” *Phys. Rev. D* **103** no. 11, (2021) L111701, [arXiv:2002.04204 \[hep-ph\]](#).

- [44] J. L. Evans, M. Sudano, and T. T. Yanagida, “A CP-safe solution of the $\mu/B\mu$ problem of gauge mediation,” *Phys. Lett. B* **696** (2011) 348–351, [arXiv:1008.3165 \[hep-ph\]](#).
- [45] M. Ibe, S. Kobayashi, Y. Nakayama, and S. Shirai, “Muon $g - 2$ in gauge mediation without SUSY CP problem,” *JHEP* **07** (2021) 098, [arXiv:2104.03289 \[hep-ph\]](#).
- [46] **ATLAS** Collaboration, G. Aad *et al.*, “Search for long-lived charginos based on a disappearing-track signature using 136 fb⁻¹ of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector,” [arXiv:2201.02472 \[hep-ex\]](#).
- [47] **CMS** Collaboration, A. M. Sirunyan *et al.*, “Search for disappearing tracks in proton-proton collisions at $\sqrt{s} = 13$ TeV,” *Phys. Lett. B* **806** (2020) 135502, [arXiv:2004.05153 \[hep-ex\]](#).
- [48] M. Ibe, S. Matsumoto, T. T. Yanagida, and N. Yokozaki, “Heavy Squarks and Light Sleptons in Gauge Mediation ~From the viewpoint of 125 GeV Higgs Boson and Muon $g-2$,” *JHEP* **03** (2013) 078, [arXiv:1210.3122 \[hep-ph\]](#).
- [49] **ATLAS** Collaboration, G. Aad *et al.*, “Search for heavy Higgs bosons decaying into two tau leptons with the ATLAS detector using pp collisions at $\sqrt{s} = 13$ TeV,” *Phys. Rev. Lett.* **125** no. 5, (2020) 051801, [arXiv:2002.12223 \[hep-ex\]](#).
- [50] **ACME** Collaboration, V. Andreev *et al.*, “Improved limit on the electric dipole moment of the electron,” *Nature* **562** no. 7727, (2018) 355–360.
- [51] S. P. Martin, “Generalized messengers of supersymmetry breaking and the sparticle mass spectrum,” *Phys. Rev. D* **55** (1997) 3177–3187, [arXiv:hep-ph/9608224](#).
- [52] G. Marandella, C. Schappacher, and A. Strumia, “Supersymmetry and precision data after LEP2,” *Nucl. Phys. B* **715** (2005) 173–189, [arXiv:hep-ph/0502095](#).
- [53] A. Strumia, “Interpreting electroweak precision data including the W -mass CDF anomaly,” [arXiv:2204.04191 \[hep-ph\]](#).
- [54] B. C. Allanach, “SOFTSUSY: a program for calculating supersymmetric spectra,” *Comput. Phys. Commun.* **143** (2002) 305–331, [arXiv:hep-ph/0104145](#).
- [55] S. Heinemeyer, W. Hollik, and G. Weiglein, “FeynHiggs: A Program for the calculation of the masses of the neutral CP even Higgs bosons in the MSSM,” *Comput. Phys. Commun.* **124** (2000) 76–89, [arXiv:hep-ph/9812320](#).
- [56] S. Heinemeyer, W. Hollik, and G. Weiglein, “The Masses of the neutral CP - even Higgs bosons in the MSSM: Accurate analysis at the two loop level,” *Eur. Phys. J. C* **9** (1999) 343–366, [arXiv:hep-ph/9812472](#).

- [57] G. Degrandi, S. Heinemeyer, W. Hollik, P. Slavich, and G. Weiglein, “Towards high precision predictions for the MSSM Higgs sector,” *Eur. Phys. J. C* **28** (2003) 133–143, [arXiv:hep-ph/0212020](#).
- [58] M. Frank, T. Hahn, S. Heinemeyer, W. Hollik, H. Rzehak, and G. Weiglein, “The Higgs Boson Masses and Mixings of the Complex MSSM in the Feynman-Diagrammatic Approach,” *JHEP* **02** (2007) 047, [arXiv:hep-ph/0611326](#).
- [59] T. Hahn, S. Heinemeyer, W. Hollik, H. Rzehak, and G. Weiglein, “High-Precision Predictions for the Light CP -Even Higgs Boson Mass of the Minimal Supersymmetric Standard Model,” *Phys. Rev. Lett.* **112** no. 14, (2014) 141801, [arXiv:1312.4937 \[hep-ph\]](#).
- [60] H. Bahl and W. Hollik, “Precise prediction for the light MSSM Higgs boson mass combining effective field theory and fixed-order calculations,” *Eur. Phys. J. C* **76** no. 9, (2016) 499, [arXiv:1608.01880 \[hep-ph\]](#).
- [61] H. Bahl, T. Hahn, S. Heinemeyer, W. Hollik, S. Paßehr, H. Rzehak, and G. Weiglein, “Precision calculations in the MSSM Higgs-boson sector with FeynHiggs 2.14,” *Comput. Phys. Commun.* **249** (2020) 107099, [arXiv:1811.09073 \[hep-ph\]](#).
- [62] P. Athron, M. Bach, H. G. Fargnoli, C. Gnendiger, R. Greifenhagen, J.-h. Park, S. Paßehr, D. Stöckinger, H. Stöckinger-Kim, and A. Voigt, “GM2Calc: Precise MSSM prediction for $(g - 2)$ of the muon,” *Eur. Phys. J. C* **76** no. 2, (2016) 62, [arXiv:1510.08071 \[hep-ph\]](#).
- [63] P. Athron, C. Balazs, A. Cherchiglia, D. H. J. Jacob, D. Stöckinger, H. Stöckinger-Kim, and A. Voigt, “Two-loop prediction of the anomalous magnetic moment of the muon in the Two-Higgs Doublet Model with GM2Calc 2,” *Eur. Phys. J. C* **82** no. 3, (2022) 229, [arXiv:2110.13238 \[hep-ph\]](#).
- [64] R. V. Harlander, S. Liebler, and H. Mantler, “SusHi: A program for the calculation of Higgs production in gluon fusion and bottom-quark annihilation in the Standard Model and the MSSM,” *Comput. Phys. Commun.* **184** (2013) 1605–1617, [arXiv:1212.3249 \[hep-ph\]](#).
- [65] R. V. Harlander, S. Liebler, and H. Mantler, “SusHi Bento: Beyond NNLO and the heavy-top limit,” *Comput. Phys. Commun.* **212** (2017) 239–257, [arXiv:1605.03190 \[hep-ph\]](#).
- [66] **NNPDF** Collaboration, R. D. Ball *et al.*, “An open-source machine learning framework for global analyses of parton distributions,” *Eur. Phys. J. C* **81** no. 10, (2021) 958, [arXiv:2109.02671 \[hep-ph\]](#).
- [67] **ATLAS Collaboration** Collaboration, “Prospects for the search for additional Higgs bosons in the ditau final state with the ATLAS detector at HL-LHC,” tech.

- rep., CERN, Geneva, Dec, 2018. <https://cds.cern.ch/record/2652284>. All figures including auxiliary figures are available at <https://atlas.web.cern.ch/Atlas/GROUPS/PHYSICS/PUBNOTES/ATL-PHYS-PUB-2018-050>.
- [68] S. Kraml, S. Kulkarni, U. Laa, A. Lessa, W. Magerl, D. Proschofsky-Spindler, and W. Waltenberger, “SModelS: a tool for interpreting simplified-model results from the LHC and its application to supersymmetry,” *Eur. Phys. J. C* **74** (2014) 2868, [arXiv:1312.4175](https://arxiv.org/abs/1312.4175) [hep-ph].
 - [69] F. Ambrogio, S. Kraml, S. Kulkarni, U. Laa, A. Lessa, V. Magerl, J. Sonneveld, M. Traub, and W. Waltenberger, “SModelS v1.1 user manual: Improving simplified model constraints with efficiency maps,” *Comput. Phys. Commun.* **227** (2018) 72–98, [arXiv:1701.06586](https://arxiv.org/abs/1701.06586) [hep-ph].
 - [70] J. Dutta, S. Kraml, A. Lessa, and W. Waltenberger, “SModelS extension with the CMS supersymmetry search results from Run 2,” *LHEP* **1** no. 1, (2018) 5–12, [arXiv:1803.02204](https://arxiv.org/abs/1803.02204) [hep-ph].
 - [71] J. Heisig, S. Kraml, and A. Lessa, “Constraining new physics with searches for long-lived particles: Implementation into SModelS,” *Phys. Lett. B* **788** (2019) 87–95, [arXiv:1808.05229](https://arxiv.org/abs/1808.05229) [hep-ph].
 - [72] F. Ambrogio *et al.*, “SModelS v1.2: long-lived particles, combination of signal regions, and other novelties,” *Comput. Phys. Commun.* **251** (2020) 106848, [arXiv:1811.10624](https://arxiv.org/abs/1811.10624) [hep-ph].
 - [73] C. K. Khosa, S. Kraml, A. Lessa, P. Neuhuber, and W. Waltenberger, “SModelS Database Update v1.2.3” *LHEP* **2020** (2020) 158, [arXiv:2005.00555](https://arxiv.org/abs/2005.00555) [hep-ph].
 - [74] G. Alguero, S. Kraml, and W. Waltenberger, “A SModelS interface for pyhf likelihoods,” *Comput. Phys. Commun.* **264** (2021) 107909, [arXiv:2009.01809](https://arxiv.org/abs/2009.01809) [hep-ph].
 - [75] G. Alguero, J. Heisig, C. K. Khosa, S. Kraml, S. Kulkarni, A. Lessa, H. Reyes-González, W. Waltenberger, and A. Wongel, “Constraining new physics with SModelS version 2,” *JHEP* **08** (2022) 068, [arXiv:2112.00769](https://arxiv.org/abs/2112.00769) [hep-ph].
 - [76] B. Fuks, M. Klasen, D. R. Lamprea, and M. Rothering, “Precision predictions for electroweak superpartner production at hadron colliders with Resummino,” *Eur. Phys. J. C* **73** (2013) 2480, [arXiv:1304.0790](https://arxiv.org/abs/1304.0790) [hep-ph].
 - [77] J. Debove, B. Fuks, and M. Klasen, “Transverse-momentum resummation for gaugino-pair production at hadron colliders,” *Phys. Lett. B* **688** (2010) 208–211, [arXiv:0907.1105](https://arxiv.org/abs/0907.1105) [hep-ph].

- [78] J. Debove, B. Fuks, and M. Klasen, “Threshold resummation for gaugino pair production at hadron colliders,” *Nucl. Phys. B* **842** (2011) 51–85, [arXiv:1005.2909 \[hep-ph\]](#).
- [79] J. Debove, B. Fuks, and M. Klasen, “Joint Resummation for Gaugino Pair Production at Hadron Colliders,” *Nucl. Phys. B* **849** (2011) 64–79, [arXiv:1102.4422 \[hep-ph\]](#).
- [80] B. Fuks, M. Klasen, D. R. Lamprea, and M. Rothering, “Gaugino production in proton-proton collisions at a center-of-mass energy of 8 TeV,” *JHEP* **10** (2012) 081, [arXiv:1207.2159 \[hep-ph\]](#).
- [81] J. Fiaschi and M. Klasen, “Neutralino-chargino pair production at NLO+NLL with resummation-improved parton density functions for LHC Run II,” *Phys. Rev. D* **98** no. 5, (2018) 055014, [arXiv:1805.11322 \[hep-ph\]](#).
- [82] J. Fiaschi and M. Klasen, “Higgsino and gaugino pair production at the LHC with aNNLO+NNLL precision,” *Phys. Rev. D* **102** no. 9, (2020) 095021, [arXiv:2006.02294 \[hep-ph\]](#).
- [83] S. P. Martin and M. T. Vaughn, “Two loop renormalization group equations for soft supersymmetry breaking couplings,” *Phys. Rev. D* **50** (1994) 2282, [arXiv:hep-ph/9311340](#). [Erratum: *Phys.Rev.D* 78, 039903 (2008)].
- [84] K. Kurosawa, N. Maru, and T. Yanagida, “Nonanomalous R symmetry in supersymmetric unified theories of quarks and leptons,” *Phys. Lett. B* **512** (2001) 203–210, [arXiv:hep-ph/0105136](#).
- [85] L. M. Krauss and F. Wilczek, “Discrete Gauge Symmetry in Continuum Theories,” *Phys. Rev. Lett.* **62** (1989) 1221.
- [86] J. Preskill and L. M. Krauss, “Local Discrete Symmetry and Quantum Mechanical Hair,” *Nucl. Phys. B* **341** (1990) 50–100.
- [87] T. Banks and M. Dine, “Note on discrete gauge anomalies,” *Phys. Rev. D* **45** (1992) 1424–1427, [arXiv:hep-th/9109045](#).
- [88] L. E. Ibanez and G. G. Ross, “Discrete gauge symmetry anomalies,” *Phys. Lett. B* **260** (1991) 291–295.
- [89] L. E. Ibanez and G. G. Ross, “Discrete gauge symmetries and the origin of baryon and lepton number conservation in supersymmetric versions of the standard model,” *Nucl. Phys. B* **368** (1992) 3–37.
- [90] L. E. Ibanez, “More about discrete gauge anomalies,” *Nucl. Phys. B* **398** (1993) 301–318, [arXiv:hep-ph/9210211](#).

- [91] D. Sabatta, A. S. Cornell, A. Goyal, M. Kumar, B. Mellado, and X. Ruan, “Connecting muon anomalous magnetic moment and multi-lepton anomalies at LHC,” *Chin. Phys. C* **44** no. 6, (2020) 063103, [arXiv:1909.03969 \[hep-ph\]](#).
- [92] G. Beck, R. Temo, E. Malwa, M. Kumar, and B. Mellado, “Connecting multi-lepton anomalies at the LHC and in Astrophysics with MeerKAT/SKA,” 2, 2021. [arXiv:2102.10596 \[astro-ph.HE\]](#).
- [93] W. H. Press, B. S. Ryden, and D. N. Spergel, “Dynamical Evolution of Domain Walls in an Expanding Universe,” *Astrophys. J.* **347** (1989) 590–604.
- [94] T. Garagounis and M. Hindmarsh, “Scaling in numerical simulations of domain walls,” *Phys. Rev. D* **68** (2003) 103506, [arXiv:hep-ph/0212359](#).
- [95] J. C. R. E. Oliveira, C. J. A. P. Martins, and P. P. Avelino, “The Cosmological evolution of domain wall networks,” *Phys. Rev. D* **71** (2005) 083509, [arXiv:hep-ph/0410356](#).
- [96] A. M. M. Leite and C. J. A. P. Martins, “Scaling Properties of Domain Wall Networks,” *Phys. Rev. D* **84** (2011) 103523, [arXiv:1110.3486 \[hep-ph\]](#).
- [97] M. Maniatis, “The Next-to-Minimal Supersymmetric extension of the Standard Model reviewed,” *Int. J. Mod. Phys. A* **25** (2010) 3505–3602, [arXiv:0906.0777 \[hep-ph\]](#).
- [98] U. Ellwanger, C. Hugonie, and A. M. Teixeira, “The Next-to-Minimal Supersymmetric Standard Model,” *Phys. Rept.* **496** (2010) 1–77, [arXiv:0910.1785 \[hep-ph\]](#).
- [99] K.-I. Izawa and T. Yanagida, “Dynamical supersymmetry breaking in vector - like gauge theories,” *Prog. Theor. Phys.* **95** (1996) 829–830, [arXiv:hep-th/9602180](#).
- [100] K. A. Intriligator and S. D. Thomas, “Dynamical supersymmetry breaking on quantum moduli spaces,” *Nucl. Phys. B* **473** (1996) 121–142, [arXiv:hep-th/9603158](#).
- [101] M. T. Grisaru, M. Rocek, and R. von Unge, “Effective Kahler potentials,” *Phys. Lett. B* **383** (1996) 415–421, [arXiv:hep-th/9605149](#).

Table 1: The mass spectra and $(a_\mu)_{\text{SUSY}}$. The (lightest) squark and gluino masses as well as Λ_D are shown in units of TeV while the others are shown in units of GeV. We also show $\sqrt{\Delta m_{H_u}^2}$, $\sqrt{\Delta m_{H_d}^2}$ and $\sqrt{B_\mu}$ at the messenger scale.

	P1	P2	P3	P4
Λ_D	900	980	950	980
r_L	0.17	0.25	0.14	0.26
M_{mess}	10^6	10^6	10^8	1.3×10^6
μ	210	260	441	250
$m_A(\text{pole})$	2500	2500	2300	2000
$\tan \beta$	40	60	43	64.5
\tilde{g}	7.0	8.2	6.2	7.1
\tilde{q}	8.3	9.2	8.0	8.9
$\tilde{\tau}_1$	581	300	459	280
$\tilde{\tau}_2$	830	584	760	583
$\tilde{\mu}_R$	985	1091	1118	1177
$\tilde{\nu}_\mu$	687	935	736	987
$(\chi_{1,2}^0, \chi_{1,2}^\pm)$	(200.2, 206.4)	(257.6, 262.5)	(322.3, 323.3)	(247.2, 252.3)
χ_2^\pm	427	667	480	694
χ_2^0	223	271	456	260
χ_3^0	426	666	471	694
χ_4^0	735	923	636	823
m_h	125.2	125.2	124.8	125.0
$(a_\mu)_{\text{SUSY}}/10^{-10}$	22.7	19.3	19.8	19.7
$\tau_{\chi_1^\pm}(\text{ns})$	5.7×10^{-8}	2.1×10^{-7}	1.8×10^{-4}	1.7×10^{-7}
$\sigma_{H/A}/10^{-4}(\text{pb})$	8.1	18.0	17.4	110.3
$\text{Br}(H \rightarrow \tau\bar{\tau})$	0.10	0.14	0.12	0.15
$\text{Br}(H \rightarrow b\bar{b})$	0.53	0.66	0.56	0.70
$\sqrt{\Delta m_{H_u}^2}$	3887	4139	4878	4137
$\sqrt{\Delta m_{H_d}^2}$	3480	4946	3949	5084
$\sqrt{B_\mu}$	385	274	≈ 0	171

Table 2: Charge assignment for all the fields.

Operators	Z_{3_1}	Z_{3_2}	Z_{3_3}	Z_{4R}	Z_4
Z	0	0	0	2	2
Q_a	0	0	0	0	1
Ψ	0	0	0	0	-1
$\bar{\Psi}$	0	0	0	2	1
E'_1	2	0	0	2	1
\bar{E}'_1	1	0	0	0	-1
E'_2	0	2	0	2	1
\bar{E}'_2	0	1	0	0	-1
E'_3	0	0	2	2	1
\bar{E}'_3	0	0	1	0	-1
E_1	1	0	0	0	0
E_2	0	1	0	0	0
E_3	0	0	1	0	0
H'_u	2	0	0	0	0
\bar{H}'_u	1	0	0	2	0
H'_d	0	0	0	0	0
\bar{H}'_d	0	0	0	2	0
S_1	1	0	0	2	0
S_2	0	1	0	2	0
S_3	0	0	1	2	0
$\Psi_{\bar{D}}\Psi_D$	0	0	2	0	0
$\Psi_L\Psi_{\bar{L}}$	0	2	0	0	0
H_u	2	0	0	0	0
H_d	0	0	0	0	0