

Emergence of microphysical viscosity in binary neutron star post-merger dynamics

Elias R. Most,^{1,2,3,*} Alexander Haber,^{4,†} Steven P. Harris,^{5,‡} Ziyuan Zhang,^{4,6,§} Mark G. Alford,^{4,¶} and Jorge Noronha^{7,**}

¹*Princeton Center for Theoretical Science, Princeton University, Princeton, NJ 08544, USA*

²*Princeton Gravity Initiative, Princeton University, Princeton, NJ 08544, USA*

³*School of Natural Sciences, Institute for Advanced Study, Princeton, NJ 08540, USA*

⁴*Physics Department, Washington University in Saint Louis, 63130 Saint Louis, MO, USA*

⁵*Institute for Nuclear Theory, University of Washington, Seattle, WA 98195, USA*

⁶*McDonnell Center for the Space Sciences, Washington University in St. Louis, St. Louis, MO 63130, USA*

⁷*Illinois Center for Advanced Studies of the Universe & Department of Physics,
University of Illinois at Urbana-Champaign, Urbana, IL, 61801, USA*

(Dated: 1 July 2022)

In nuclear matter in neutron stars the flavor content (e.g., proton fraction) is subject to weak interactions, establishing flavor (β -)equilibrium. During the merger of two neutron stars there can be deviations from this equilibrium. By incorporating Urca processes into general-relativistic hydrodynamics simulations, we study the resulting out-of-equilibrium dynamics during the collision. We provide the first direct evidence that microphysical transport effects at late times reach a hydrodynamic regime with a nonzero bulk viscosity, making neutron star collisions intrinsically viscous. Finally, we identify signatures of this process in the post-merger gravitational wave emission.

I. INTRODUCTION

Binary neutron star mergers [1, 2] offer exciting prospects for constraining the dense matter equation of state (EoS) [3–5] (e.g., [6–11]). This can be done either by using the tidal deformability encoded in the inspiral gravitational wave signal [12–19], the potential presence of very massive neutron stars in gravitational wave events [20–26], or by using the combined information from electromagnetic counterparts [27–31]. Critically, some of these constraints heavily rely on properties of the mass ejected after merger (e.g., [32–38]), which crucially depends on models for the post-merger evolution, including the lifetime of the remnant neutron star (see, e.g., [39–41] for recent reviews). Future detection of post-merger gravitational wave signals [42–48] have the potential for putting even more stringent constraints on the EoS [49–58].

Beyond uncertainties in the cold EoS, finite-temperature effects [59–61], the appearance of exotic degrees of freedom or phase transitions [26, 62–69], or effects from weak interactions [70, 71] could alter the post-merger dynamics and systematically bias our ability to infer dense matter properties [72]. The latter remains largely not understood as it requires additional physics in the form of weak interactions [73–76] to be specified. These are important as the weak interaction (β -)equilibrium that is present in the inspiraling neutron stars can be dynamically violated in mergers [77, 78]. Weak interactions can then restore β -equilibrium on the dynamical timescale of post-merger gravitational wave emission [70], and it is expected that the corresponding back-reaction on the fluid can be described in terms of an effective bulk viscosity

[73]. Previous work has largely been devoted to understanding the properties [73, 74] and the potential appearance of bulk viscosity [70, 72, 77] in the remnant, or to studying the idealized limit of perfect β -equilibrium [79, 80]. While effects of microphysical viscosity in neutron-star merger simulations have largely been ignored, they have been well studied in the related context of dense matter formed heavy-ion collisions [81–87], see also [88] for a review.

Going beyond all of the above, in this paper we make essential steps towards fully self-consistent numerical simulations of the out-of-equilibrium dynamics of the flavor content of the matter in the post-merger phase of a neutron star collision. We begin by briefly reviewing the relevant processes (Urca processes) active in the core of neutron stars [89]. We then provide a detailed discussion of the dynamics of relaxation to β -equilibrium after the merger. For the first time, we directly demonstrate the emergence of microphysical bulk viscosity in post-merger simulations and provide a first quantification of its effects. We conclude by discussing how this alters the post-merger gravitational wave emission.

II. METHODS

We model the global dynamics of the system using general-relativistic (magneto-)hydrodynamics [90] in a dynamically evolved spacetime in the Z4c formulation [91]. More specifically, we evolve the baryon current $J_B^\mu = n_B u^\mu$ and the perfect-fluid energy-momentum tensor $T^{\mu\nu}$ [92] according to

$$\nabla_\alpha J_B^\alpha = 0, \quad \nabla_\alpha T^{\alpha\beta} = Q_{\nu_e}^\beta, \quad (1)$$

where we have included an effective energy sink term $Q_{\nu_e}^\beta$ associated with neutrino energy-losses [93, 94]. Here, n_B and u^μ respectively are the baryon number density and fluid four-velocity. In order to track the relative number of protons, we

* emost@princeton.edu

† ahaber@physics.wustl.edu

‡ harrissp@uw.edu

§ ziyuan.z@wustl.edu

¶ alford@physics.wustl.edu

** jn0508@illinois.edu





FIG. 1. Electron fraction, Y_e , present in the system after merger at time $t = 5$ ms as a function of density. Shown are results for the simulations using the SFHo (top) and TMA (bottom) equations of state, with and without Urca rates.


evolve the relative fraction of electrons (e)

$$u^\mu \nabla_\mu Y_e = \mathcal{R}_{\nu_e} + \Gamma_{mUrca} + \Gamma_{dUrca}. \quad (2)$$

Assuming npe matter, the electron fraction, Y_e , can only change by weak decays of neutrons (n) and protons (p). We model these in two ways. First, we include a standard set of weak interactions, \mathcal{R}_{ν_e} , commonly used in leakage scheme approaches to neutron star mergers [95, 96]. These are predominantly important for densities far below saturation [93, 94, 96], and we have confirmed that they do not impact our results. Secondly, we consistently include—for the first time—direct, Γ_{dUrca} , and modified, Γ_{mUrca} , Urca (net) rates [75, 97] that operate in the dense nuclear matter core of the merger. Specifically, we consider the direct Urca (dUrca),



and modified Urca (mUrca) processes,



where X is a spectator nucleon [89, 97]. We treat the Urca processes (3) and (4) in the Fermi surface approximation where the net rate is proportional to $\delta\mu(n_B, T, Y_e) = \mu_n - \mu_p - \mu_e$, with n_B and T being the baryon number density and temperature while μ_X is the chemical potential for particle X . The system therefore relaxes towards $\delta\mu = \delta\mu^{eq} = 0$. In npe -matter the direct Urca process dominates over modified Urca at proton fraction $Y_e \gtrsim 0.11$ [97, 98] (the “direct Urca threshold”). At lower Y_e direct Urca is suppressed and modified Urca becomes the leading contributor to β equilibration. To probe the influence of the dUrca threshold we consider two EoSs,




FIG. 2. Electron fraction, Y_e , present in the merger remnant at time $t = 5$ ms, when systematically including (top) or excluding (bottom) Urca processes for a GW170817-like neutron star merger using the TMA equation of state. Density contours (green) are denoted in units of nuclear saturation density, n_{sat} .

TMA [99] and SFHo [75, 100]. Beta-equilibrated SFHo matter always has $Y_e < 0.11$ and, therefore, it predominantly equilibrates via mUrca. Beta-equilibrated TMA matter reaches the dUrca threshold at a density just below $2n_{\text{sat}}$ (where $n_{\text{sat}} \approx 0.16 \text{ cm}^{-3}$ is the nuclear saturation density). More information can be found in the Appendix.

We solve the coupled system using the Frankfurt/IllinoisGRMHD code [101, 102], which utilizes the EinsteinToolkit infrastructure [103], where the Urca processes are included via a fully implicit operator-split approach. More details on the numerical setup and the choice of initial data can be found in [78].

III. RESULTS

In this first study we consider a system with fixed binary parameters loosely modeled after the GW170817 event [1]. We fix the total mass $M = 2.74 M_\odot$ with a mass ratio of $q = 0.85$. This results in a hot, (meta-)stable neutron star remnant after the merger, which does not immediately collapse to a black hole [104–110]. A detailed description of the (post-)merger dynamics and gravitational wave emission of this system for different EoSs can be found in [57, 78]. In this work, we exclusively focus on the impact of violating and dynamically restoring β -equilibrium [77, 78].

While weak interactions have sufficient time to establish equilibrium for all but the late stages of inspiral [111], the violent dynamics of the merger can drive the electron fraction away from its equilibrium value [77, 112]. This is displayed in Fig. 1, which shows the distribution of Y_e as a function of n_B 5 ms after merger. Without including Urca reactions, there can be considerable scatter in Y_e (as we see for SFHo), and the




FIG. 3. Effective out-of-beta-equilibrium pressure contribution Π to the equilibrium pressure P^{eq} during the merger and post-merger phase of two neutron stars when using the TMA equation of state with Urca rates. Shown is the orbital plane. The green contours denote the rest mass density of the merger remnant in units of nuclear saturation. All times, t , are stated relative to the time of merger t_{mer} .

system is far from chemical equilibrium at densities beyond saturation. Urca processes restore equilibrium on a time scale that depends on the density and temperature. Fig. 1 shows a clear correlation between n_B and Y_e in β -equilibrium for $n_B > n_{\text{sat}}$. In the case of the TMA EoS, this affects the structure of the star since in the presence of Urca processes the matter reaches higher densities, $n_B > 2.5 n_{\text{sat}}$. In the case of SFHo, the merger produces some regions with an excess of neutrons and some with an excess of protons; for TMA there is mostly an excess of neutrons (see Fig. 1). If weak interactions are correctly included, this leads to a rapid onset of neutron \leftrightarrow proton conversion in the star, as shown in Fig. 2. In the case of SFHo, this will be largely driven by mUrca, as most fluid elements have Y_e values below the dUrca threshold (Fig. 1), whereas for TMA it is a combination of dUrca and mUrca. We can also anticipate that this change in structure and composition might lead to a change in the (early) mass ejection (e.g., [113–121]), but leave the details to future work.

We now move on to the main part of our work, which is to quantify the influence on the global dynamics of the Urca-driven protonization/neutronization of the dense matter inside the neutron star. We begin by characterizing the impact of weak interactions on the fluid as dissipative corrections to a β -equilibrated flow [72, 122]. In practice, this amounts to splitting the total pressure, P , into

$$P(n_B, T, Y_e) = P^{\text{eq}}(n_B, T) + \Pi(n_B, T, Y_e), \quad (5)$$

where $P^{\text{eq}} = P(n_B, T, Y_e = Y_e^{\text{eq}})$ is the pressure in β -equilibrium, and Π is the bulk scalar pressure correction, which quantifies how the system deviates from β equilibrium. It can be shown that the equations of motion (1) to linear order in $\delta\mu$ can be approximated as a bulk-viscous hydrodynamic system of equations [122–124]. In the Navier-Stokes limit [92, 122–124] the dissipative pressure correction becomes $\Pi = \Pi_{\text{NS}} + \mathcal{O}(u^\nu \nabla_\nu \delta\mu)$, where

$$\Pi_{\text{NS}} = -\zeta \nabla_\mu u^\mu, \quad (6)$$

and $\zeta = \zeta(n_B, T)$ is the weak-interaction driven bulk viscosity [73, 74]. Applications of semi-analytic expressions for the bulk viscosity [73] to the background of a non-viscous neutron star merger calculation have projected that large viscous corrections could arise after the merger [72]. Here, we go beyond

such estimates, as well as first-order bulk-viscous approximations [122], and instead compute the full dynamics of the Urca process (in the Fermi Surface approximation) and its influence on dense matter beyond leading order in $\delta\mu$ by directly extracting Π from the total pressure (5) in our simulations. Using our self-consistent approach, we can provide a first quantification of the out-of-equilibrium dynamics in terms of the relative bulk pressure contribution Π/P^{eq} as shown in Fig. 3.

In the very late inspiral, right before the collision, tidal forces drive the material inside the stars out of equilibrium. While we only show the last orbit (Fig. 3, left panel), this could potentially set in even earlier [111]. Overall, we find that $\Pi/P^{\text{eq}} \gtrsim 5\%$ throughout the stars during initial stages of the collision (Fig. 3, left panel). The subsequent evolution now strongly depends on whether weak-interaction effects are included. If Urca processes are not included, the matter does not re-equilibrate on the dynamical time scale of the merger and the out-of-equilibrium pressure contribution, Π , remains constant on average for both EoSs (Fig. 4). If instead Urca processes are included, the system begins to re-equilibrate at a rate that depends on density and temperature. One can see that especially in higher density regions $n_B > 2 n_{\text{sat}}$, equilibration due to dUrca proceeds quickly with the core of the object having reduced Π/P^{eq} by more than one order of magnitude, with the hottest regions re-equilibrating first.

A short time after the merger, at $t = 4.5$ ms, the TMA system has largely re-equilibrated. However, post-merger oscillations [125] continue to locally drive the system out of β -equilibrium. It is especially this regime where we expect the feedback on the dense matter to behave like a bulk viscosity [72].

We will now provide a more thorough quantification of this dynamics by considering density averages over the system in the equatorial plane. For this, we utilize the concept of a dynamic inverse Reynolds number [72, 126], $\chi = \Pi/(e^{\text{eq}} + P^{\text{eq}})$, with e^{eq} being the total energy density of the fluid in β -equilibrium. We will in particular rely on a density-weighted average value $\langle \chi \rangle$. For reference, gravitational wave emission alone causes an effective viscous damping of density oscillations corresponding to $\chi \simeq 0.01$ [72]. With this in mind, we can now correlate chemical equilibration and its hydrodynamic feedback in Fig. 4. For TMA,




FIG. 4. Effective inverse Reynolds number $\chi \equiv \Pi / (e^{\text{eq}} + P^{\text{eq}})$ and density-averaged chemical potential difference $\delta\mu$ normalized by the temperature T at times t relative to the time of merger t_{mer} .

which is expected to equilibrate faster via dUrca as well as mUrca, we see that $\langle \chi \rangle$ briefly jumps up after the merger, following the no-Urca simulation, but then follows an approximately exponential decay with a lifetime of about 0.9 ms as Urca processes establish β -equilibrium. For SFHo, which is expected to equilibrate more slowly via mUrca processes, the behavior is more complicated: (a) $\langle \chi \rangle$ almost immediately reaches a plateau at a value about 10 times smaller than in the no-Urca simulations; (b) the plateau persists for about 4 ms before giving way to exponential decay with lifetime ~ 1.7 ms. This is partly because SFHo equilibrates slowly via mUrca processes, but by examining the simulation we can uncover interesting additional phenomena.

Some regions of SFHo matter reach temperatures $T > 90$ MeV, so that they equilibrate almost instantaneously. However, this creates feedback on the stellar structure which drives the outer layers of the star further out of chemical equilibrium, which is an additional reason why, for SFHo matter, $\langle \chi \rangle$ and $\langle \delta\mu/T \rangle$ in Fig. 3 do not decay immediately but feature an extend period of out-of-equilibrium dynamics, which only ends at $t - t_{\text{mer}} \simeq 5$ ms. The initial SFHo pressure correction Π is also considerably smaller than for the TMA case, possibly associated with stronger tidal effects in the TMA case. This underlines the fact that not all EoSs lead to strong violation of β -equilibrium during merger [78].

In both cases, β -equilibrium is restored when Urca processes are included, and remains constantly violated if they are absent. This overall trend in $\langle \chi \rangle$ is matched when considering density-averaged values of $\langle \delta\mu/T \rangle$ in Fig. 4. There is an initial rise at merger matching the appearance of out-of-equilibrium pressure corrections. In the absence of Urca rates the violation of chemical equilibrium remains constant throughout the simulation. For both EoSs the decay times are shorter than the timescale of post-merger gravitational wave emission. This implies that for long-term evolutions of the merger remnant (e.g., [127, 128]), the matter can be treated as being in chemical equilibrium. As the system approaches chemical equilibrium, $\delta\mu/T \ll 1$, at times $t > 5$ ms, post-merger oscillations of the stellar remnant only mildly drive the system out of equilibrium, so $\langle \chi \rangle \simeq 10^{-5}$ and $\delta\mu/T \simeq 10^{-2}$.

One of the goals of our work is to clarify how much of the merger region is in a regime where β -equilibration

can be treated hydrodynamically, i.e., when Π is largely determined by a bulk viscosity, see the discussion around (6). We call this regime the “bulk viscous regime”, and define it as the regime where deviations from chemical equilibrium can be treated via linear response theory, i.e., $\Pi \propto \delta\mu$. We note, however, that our total net Urca rate Γ (see Appendix) is applicable at any value of $\delta\mu/T$. The bulk viscous regime is where $\delta\mu/T$ is small enough so that the linear approximation $\Gamma \approx \Gamma_{\text{bulk}} = (d\Gamma/d\delta\mu)|_{\delta\mu=0} \delta\mu$ is valid. [122–124]. In Fig. 5, we determine in which regime we are by directly comparing the weak-interaction rates, plotting $\Gamma/\Gamma_{\text{bulk}}$. One can see that shortly after merger, when the out-of-equilibrium corrections for the SFHo system are large, there are small regions that are outside the bulk viscous regime, with Urca rates $10\times$ faster than in linear response. Those are present at low and high densities, and even in the tidal tail of the secondary neutron star that got partially disrupted in this unequal mass merger. In line with the evolution of $\langle \delta\mu/T \rangle$, these corrections decay rapidly over a timescale $\tau_{\delta\mu} \simeq 0.5$ ms, leaving the whole central region in the bulk viscous regime. In the case of SFHo, we extract effective viscosities of $\zeta_{\text{eff}} \lesssim 10^{28}$ g/(cm s) directly from the simulation by computing $\zeta_{\text{eff}} \approx \Pi/D_i u^i$ under the assumption of the Navier-Stokes limit (6). This appearance of dissipation also leads to the production of entropy, increasing it on average by $\Delta s \sim 0.1 k_B/\text{baryon}$ compared to a merger without Urca processes (see also [129]). This opens up the interesting prospect of studying these late-time effects in a complementary way using general-relativistic viscous hydrodynamics [130–138].

Ideally, one would like to identify a signature for the appearance of dissipative re-equilibration in the gravitational wave signal emitted after the collision. To this end, in Fig. 6 we compare the gravitational emission for all four simulations. Looking at the TMA case, we find good agreement between the gravitational wave strains in the late inspiral and the early merger phase. This is consistent with the Urca processes not enforcing β -equilibrium immediately, as shown in Fig. 4. However, after the characteristic re-equilibration time $\simeq 2$ ms, a phase difference begins to build up and continues to grow until the end of the simulation $t \simeq 10$ ms. The SFHo case overall is very similar. Due to the delayed onset




FIG. 5. Ratio of the total to linear Urca reaction rates $\Gamma/\Gamma_{\text{bulk}}$ (see Appendix) at time $t = 2.3$ ms after merger for the SFHo simulation. Blue regions are in the linear response “bulk viscous” regime, red regions require higher-order corrections. Density contours are shown in units of nuclear saturation.

of re-equilibration, see the discussion around Fig. 4, the dephasing does not set in until $t \gtrsim 2$ ms. Overall, we find frequency shifts $\Delta f \simeq 110(40)$ Hz for the TMA(SFHo) EoSs. These results are consistent with a study recently reported by [80] where an infinitely fast rate, $\Gamma \rightarrow \infty$, was considered. While this is appropriate for our late-time behavior, it crucially misses out on the dynamical re-equilibration and the emergence of bulk viscosity (see Fig. 5).

IV. CONCLUSIONS.

In this work, we have presented the first study of weak-interaction-driven out-of-equilibrium dynamics in a binary neutron star merger. We have shown how Urca processes acting on a millisecond timescale restore the departures from β -equilibrium that arise during the initial stages of the merger. We show that large parts of the merger remnant end up in the bulk-viscous regime, in which this re-equilibration leads to an effective bulk viscosity. We further show that this causes the gravitational wave frequencies to shift by up to $\Delta f \lesssim 110$ Hz, which is potentially discernible by next-generation facilities [80]. Further work will be required to fully map out the impact of out-of-equilibrium dynamics on the post-merger evolution. Firstly, not all equations of state lead to strong violation of β -equilibrium at merger [77, 78]. Those that do will be subject to uncertainties in the finite-temperature part [61], critically affecting what fraction of the merger will be in the neutrino trapped regime [75, 110, 112], in which re-equilibration happens almost instantaneously [74, 76]. Similarly, trapped neutrino pressure contributions might be of similar strength, but will affect hot instead of cold regions [112]. Furthermore, it remains to be




FIG. 6. Normalized gravitational wave strain, h_+^{22} , for the $l = m = 2$ component. Differences due to weak interactions in the post-merger are clearly visible. All times, t , are stated relative to the merger time and extraction radius r . The vertical lines indicate times of black hole formation.

seen what impact the change in remnant structure has on the mass ejection and lifetime of the post-merger system [139] (see also black hole formation times in Fig. 6). Effects that are not captured by the Fermi Surface approximation, such as finite-temperature blurring of the direct Urca threshold and the resultant modification of the beta equilibrium condition [75, 140], should be explored. Different phases of matter, such as hyperonic and quark matter, have different channels of chemical equilibration giving rise to bulk viscosities with different dependencies on temperature and density [141, 142]. The presence of a bulk-viscous phase in the merger opens a new window towards connecting dense matter in neutron star mergers to that in heavy-ion collisions [143], where out-of-equilibrium viscous corrections are relevant [88].

ACKNOWLEDGMENTS

ERM thanks F. Foucart, J. Noronha-Hostler, A. Pandya, F. Pretorius and C. Raithel for insightful discussions related to this work. ERM gratefully acknowledges support from a joint fellowship at the Princeton Center for Theoretical Science, the Princeton Gravity Initiative and the Institute for Advanced Study. ERM acknowledges support for compute time allocations on the NSF Frontera supercomputer under grants AST21006. This work used the Extreme Science and Engineering Discovery Environment (XSEDE) [144] through Expanse at SDSC and Bridges-2 at PSC through allocations PHY210053 and PHY210074. The simulations were also in part performed on computational resources managed and supported by Princeton Research Computing, a consortium of groups including the Princeton Institute for Computational Science and Engineering (PICSciE) and the Office of Informa-

tion Technology's High Performance Computing Center and Visualization Laboratory at Princeton University. ERM also acknowledges the use of high-performance computing at the Institute for Advanced Study. MGA, AH, and ZZ are partly supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under Award No. #DE-FG02-

05ER41375. SPH is supported by the U. S. Department of Energy grant DE-FG02-00ER41132 as well as the National Science Foundation grant No. PHY-1430152 (JINA Center for the Evolution of the Elements). JN is partially supported by the U.S. Department of Energy, Office of Science, Office for Nuclear Physics under Award No. DE-SC0021301.

Appendix A: Direct Urca rates

In this initial work, we use the Fermi surface approximation to calculate the Urca rates. In this assumption, the nuclear matter is treated as strongly degenerate [75, 140]. The net rate of direct Urca is given by (see also [124, 138])

$$\Gamma_{\text{dUrca}} \equiv \Gamma_{dU,nd} - \Gamma_{dU,ec} = \frac{G^2(1+3g_A^2)}{240\pi^5} E_{Fn}^* E_{Fp}^* p_{Fe} \theta_{dU} \delta\mu (17\pi^4 T^4 + 10\pi^2 \delta\mu^2 T^2 + \delta\mu^4), \quad (\text{A1})$$

where $G \equiv G_F \cos \theta_c$, $G_F = 1.166 \times 10^{-11} \text{MeV}^{-2}$ is the Fermi constant, $\theta_c = 13.04^\circ$ is the Cabbibo angle, $g_A = 1.26$, and $E_{FN}^* = \sqrt{k_{FN}^2 + m_N^{*2}}$ are the nucleon Fermi energies, and $\delta\mu \equiv \mu_n - \mu_p - \mu_e$ quantifies the departure of the system from chemical equilibrium. For simplicity, we fix the effective mass $m_N^* \approx 0.7 m_n$ relative to the neutron mass m_n , although in reality the effective mass drops with increasing density [75]. In the Fermi surface approximation, the direct Urca process operates only above a threshold density, where

$$\theta_{dU} = \begin{cases} 1 & p_{Fn} < p_{Fp} + p_{Fe} \\ 0 & \text{otherwise.} \end{cases} \quad (\text{A2})$$

We neglect finite temperature corrections to the Fermi surface approximation which would blur this threshold and lead to a nonzero $\delta\mu^{\text{eq}}$ [75, 140]. Above temperatures of $T \approx 10 \text{MeV}$ we expect neutrinos to be trapped [140, 145]. Since the associated Urca timescales are significantly shorter than the time steps of our simulations [76], we capture this regime approximately by enforcing cold β -equilibrium. Due to the implicit time-stepping, all rates acting on time scales shorter than the numerical time step, will lead to an effective instantaneous re-equilibration. Since this happens automatically for large T we use the same rates also in hot matter.

Appendix B: Modified Urca rates

The modified Urca rate is also calculated using the Fermi surface approximation. The individual rates are presented in Sec. 5 of [75]. The net rates of modified Urca, with a neutron or a proton spectator, are

$$\Gamma_{\text{mUrca},n} \equiv \Gamma_{mU,nd(n)} - \Gamma_{mU,ec(n)} = \frac{1}{5760\pi^9} G^2 g_A^2 f^4 \frac{(E_{Fn}^*)^3 E_{Fp}^*}{m_\pi^4} \frac{k_{Fn}^4 k_{Fp}}{(k_{Fn}^2 + m_\pi^2)^2} \theta_n \times \delta\mu (1835\pi^6 T^6 + 945\pi^4 \delta\mu^2 T^4 + 105\pi^2 \delta\mu^4 T^2 + 3\delta\mu^6). \quad (\text{B1})$$

$$\Gamma_{\text{mUrca},p} \equiv \Gamma_{mU,nd(p)} - \Gamma_{mU,ec(p)} = \frac{1}{40320\pi^9} G^2 g_A^2 f^4 \frac{E_{Fn}^* (E_{Fp}^*)^3}{m_\pi^4} \frac{(k_{Fn} - k_{Fp})^4 k_{Fn}}{((k_{Fn} - k_{Fp})^2 + m_\pi^2)^2} \theta_p \times \delta\mu (1835\pi^6 T^6 + 945\pi^4 \delta\mu^2 T^4 + 105\pi^2 \delta\mu^4 T^2 + 3\delta\mu^6). \quad (\text{B2})$$

Here we have introduced

$$\theta_n = \begin{cases} 1 & k_{Fn} > k_{Fp} + k_{Fe} \\ 1 - \frac{3(k_{Fp} + k_{Fe} - k_{Fn})^2}{8 k_{Fp} k_{Fe}} & k_{Fn} < k_{Fp} + k_{Fe}. \end{cases} \quad (\text{B3})$$

and

$$\theta_p = \begin{cases} 0 & k_{Fn} > 3k_{Fp} + k_{Fe} \\ \frac{(3k_{Fp} + k_{Fe} - k_{Fn})^2}{4(3k_{Fp} - k_{Fn})} \frac{k_{Fn}k_{Fe}}{k_{Fn}k_{Fe}} & 3k_{Fp} + k_{Fe} > k_{Fn} > 3k_{Fp} - k_{Fe} \\ 2 + \frac{k_{Fn}}{3(2k_{Fp} - k_{Fn})} - \frac{3(k_{Fp} - k_{Fe})^2}{k_{Fn}k_{Fe}} & 3k_{Fp} - k_{Fe} > k_{Fn} > k_{Fp} + k_{Fe} \\ & k_{Fn} < k_{Fp} + k_{Fe}. \end{cases} \quad (\text{B4})$$

The total modified Urca rate is given by the sum

$$\Gamma_{\text{mUrca}} \equiv \Gamma_{\text{mUrca},n} + \Gamma_{\text{mUrca},p}. \quad (\text{B5})$$

Appendix C: Total Urca rate

The total net Urca rate in the Fermi Surface approximation is

$$\Gamma = \Gamma_{\text{dUrca}} + \Gamma_{\text{mUrca}}, \quad (\text{C1})$$

Since bulk viscosity is typically defined in the linear (also sometimes called ‘‘subthermal’’ [146, 147]) regime where $\delta\mu \ll T$, it is natural to write this as

$$\Gamma = \Gamma_{\text{bulk}} + \text{higher order in } \delta\mu/T, \quad (\text{C2})$$

$$\text{where } \Gamma_{\text{bulk}} \equiv \left. \frac{d\Gamma}{d\delta\mu} \right|_{\delta\mu=0} \delta\mu. \quad (\text{C3})$$

-
- [1] B. P. Abbott *et al.* (LIGO Scientific, Virgo), GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral, *Phys. Rev. Lett.* **119**, 161101 (2017), [arXiv:1710.05832 \[gr-qc\]](#).
- [2] B. P. Abbott *et al.* (LIGO Scientific, Virgo), GW190425: Observation of a Compact Binary Coalescence with Total Mass $\sim 3.4M_{\odot}$, *Astrophys. J. Lett.* **892**, L3 (2020), [arXiv:2001.01761 \[astro-ph.HE\]](#).
- [3] J. M. Lattimer and M. Prakash, The Equation of State of Hot, Dense Matter and Neutron Stars, *Phys. Rept.* **621**, 127 (2016), [arXiv:1512.07820 \[astro-ph.SR\]](#).
- [4] M. Oertel, M. Hempel, T. Klöhn, and S. Typel, Equations of state for supernovae and compact stars, *Rev. Mod. Phys.* **89**, 015007 (2017), [arXiv:1610.03361 \[astro-ph.HE\]](#).
- [5] F. Özel and P. Freire, Masses, Radii, and the Equation of State of Neutron Stars, *Ann. Rev. Astron. Astrophys.* **54**, 401 (2016), [arXiv:1603.02698 \[astro-ph.HE\]](#).
- [6] E. E. Flanagan and T. Hinderer, Constraining neutron star tidal Love numbers with gravitational wave detectors, *Phys. Rev. D* **77**, 021502 (2008), [arXiv:0709.1915 \[astro-ph\]](#).
- [7] J. S. Read, C. Markakis, M. Shibata, K. Uryu, J. D. E. Creighton, and J. L. Friedman, Measuring the neutron star equation of state with gravitational wave observations, *Phys. Rev. D* **79**, 124033 (2009), [arXiv:0901.3258 \[gr-qc\]](#).
- [8] C. A. Raithel, Constraints on the Neutron Star Equation of State from GW170817, *Eur. Phys. J. A* **55**, 80 (2019), [arXiv:1904.10002 \[astro-ph.HE\]](#).
- [9] F. Hernandez Vivanco, R. Smith, E. Thrane, P. D. Lasky, C. Talbot, and V. Raymond, Measuring the neutron star equation of state with gravitational waves: The first forty binary neutron star merger observations, *Phys. Rev. D* **100**, 103009 (2019), [arXiv:1909.02698 \[gr-qc\]](#).
- [10] P. Landry, R. Essick, and K. Chatziioannou, Nonparametric constraints on neutron star matter with existing and upcoming gravitational wave and pulsar observations, *Phys. Rev. D* **101**, 123007 (2020), [arXiv:2003.04880 \[astro-ph.HE\]](#).
- [11] K. Chatziioannou, Uncertainty limits on neutron star radius measurements with gravitational waves, *Phys. Rev. D* **105**, 084021 (2022), [arXiv:2108.12368 \[gr-qc\]](#).
- [12] A. Bauswein, O. Just, H.-T. Janka, and N. Stergioulas, Neutron-star radius constraints from GW170817 and future detections, *Astrophys. J. Lett.* **850**, L34 (2017), [arXiv:1710.06843 \[astro-ph.HE\]](#).
- [13] E. Annala, T. Gorda, A. Kurkela, and A. Vuorinen, Gravitational-wave constraints on the neutron-star-matter Equation of State, *Phys. Rev. Lett.* **120**, 172703 (2018), [arXiv:1711.02644 \[astro-ph.HE\]](#).
- [14] E. R. Most, L. R. Weih, L. Rezzolla, and J. Schaffner-Bielich, New constraints on radii and tidal deformabilities of neutron stars from GW170817, *Phys. Rev. Lett.* **120**, 261103 (2018), [arXiv:1803.00549 \[gr-qc\]](#).
- [15] B. P. Abbott *et al.* (LIGO Scientific, Virgo), GW170817: Measurements of neutron star radii and equation of state, *Phys. Rev. Lett.* **121**, 161101 (2018), [arXiv:1805.11581 \[gr-qc\]](#).
- [16] C. Raithel, F. Özel, and D. Psaltis, Tidal deformability from GW170817 as a direct probe of the neutron star radius, *As-*

- trophys. J. Lett. **857**, L23 (2018), [arXiv:1803.07687 \[astro-ph.HE\]](#).
- [17] S. De, D. Finstad, J. M. Lattimer, D. A. Brown, E. Berger, and C. M. Biwer, Tidal Deformabilities and Radii of Neutron Stars from the Observation of GW170817, *Phys. Rev. Lett.* **121**, 091102 (2018), [Erratum: *Phys.Rev.Lett.* 121, 259902 (2018)], [arXiv:1804.08583 \[astro-ph.HE\]](#).
- [18] K. Chatziioannou, C.-J. Haster, and A. Zimmerman, Measuring the neutron star tidal deformability with equation-of-state-independent relations and gravitational waves, *Phys. Rev. D* **97**, 104036 (2018), [arXiv:1804.03221 \[gr-qc\]](#).
- [19] Z. Carson, A. W. Steiner, and K. Yagi, Constraining nuclear matter parameters with GW170817, *Phys. Rev. D* **99**, 043010 (2019), [arXiv:1812.08910 \[gr-qc\]](#).
- [20] F. J. Fattoyev, C. J. Horowitz, J. Piekarewicz, and B. Reed, GW190814: Impact of a 2.6 solar mass neutron star on the nucleonic equations of state, *Phys. Rev. C* **102**, 065805 (2020), [arXiv:2007.03799 \[nucl-th\]](#).
- [21] E. R. Most, L. J. Papenfort, L. R. Weih, and L. Rezzolla, A lower bound on the maximum mass if the secondary in GW190814 was once a rapidly spinning neutron star, *Mon. Not. Roy. Astron. Soc.* **499**, L82 (2020), [arXiv:2006.14601 \[astro-ph.HE\]](#).
- [22] H. Tan, J. Noronha-Hostler, and N. Yunes, Neutron Star Equation of State in light of GW190814, *Phys. Rev. Lett.* **125**, 261104 (2020), [arXiv:2006.16296 \[astro-ph.HE\]](#).
- [23] V. Dexheimer, R. O. Gomes, T. Klähn, S. Han, and M. Salinas, GW190814 as a massive rapidly rotating neutron star with exotic degrees of freedom, *Phys. Rev. C* **103**, 025808 (2021), [arXiv:2007.08493 \[astro-ph.HE\]](#).
- [24] I. Tews, P. T. H. Pang, T. Dietrich, M. W. Coughlin, S. Antier, M. Bulla, J. Heinzel, and L. Issa, On the Nature of GW190814 and Its Impact on the Understanding of Supranuclear Matter, *Astrophys. J. Lett.* **908**, L1 (2021), [arXiv:2007.06057 \[astro-ph.HE\]](#).
- [25] A. Nathanail, E. R. Most, and L. Rezzolla, GW170817 and GW190814: tension on the maximum mass, *Astrophys. J. Lett.* **908**, L28 (2021), [arXiv:2101.01735 \[astro-ph.HE\]](#).
- [26] H. Tan, T. Dore, V. Dexheimer, J. Noronha-Hostler, and N. Yunes, Extreme matter meets extreme gravity: Ultraheavy neutron stars with phase transitions, *Phys. Rev. D* **105**, 023018 (2022), [arXiv:2106.03890 \[astro-ph.HE\]](#).
- [27] D. Radice, A. Perego, F. Zappa, and S. Bernuzzi, GW170817: Joint Constraint on the Neutron Star Equation of State from Multimessenger Observations, *Astrophys. J. Lett.* **852**, L29 (2018), [arXiv:1711.03647 \[astro-ph.HE\]](#).
- [28] B. Margalit and B. D. Metzger, Constraining the Maximum Mass of Neutron Stars from Multi-Messenger Observations of GW170817, *Astrophys. J. Lett.* **850**, L19 (2017), [arXiv:1710.05938 \[astro-ph.HE\]](#).
- [29] L. Rezzolla, E. R. Most, and L. R. Weih, Using gravitational-wave observations and quasi-universal relations to constrain the maximum mass of neutron stars, *Astrophys. J. Lett.* **852**, L25 (2018), [arXiv:1711.00314 \[astro-ph.HE\]](#).
- [30] M. Ruiz, S. L. Shapiro, and A. Tsokaros, GW170817, General Relativistic Magnetohydrodynamic Simulations, and the Neutron Star Maximum Mass, *Phys. Rev. D* **97**, 021501 (2018), [arXiv:1711.00473 \[astro-ph.HE\]](#).
- [31] M. Shibata, E. Zhou, K. Kiuchi, and S. Fujibayashi, Constraint on the maximum mass of neutron stars using GW170817 event, *Phys. Rev. D* **100**, 023015 (2019), [arXiv:1905.03656 \[astro-ph.HE\]](#).
- [32] P. S. Cowperthwaite *et al.*, The Electromagnetic Counterpart of the Binary Neutron Star Merger LIGO/Virgo GW170817. II. UV, Optical, and Near-infrared Light Curves and Comparison to Kilonova Models, *Astrophys. J. Lett.* **848**, L17 (2017), [arXiv:1710.05840 \[astro-ph.HE\]](#).
- [33] R. Chornock *et al.*, The Electromagnetic Counterpart of the Binary Neutron Star Merger LIGO/VIRGO GW170817. IV. Detection of Near-infrared Signatures of r-process Nucleosynthesis with Gemini-South, *Astrophys. J. Lett.* **848**, L19 (2017), [arXiv:1710.05454 \[astro-ph.HE\]](#).
- [34] V. A. Villar *et al.*, The Combined Ultraviolet, Optical, and Near-Infrared Light Curves of the Kilonova Associated with the Binary Neutron Star Merger GW170817: Unified Data Set, Analytic Models, and Physical Implications, *Astrophys. J. Lett.* **851**, L21 (2017), [arXiv:1710.11576 \[astro-ph.HE\]](#).
- [35] M. Nicholl *et al.*, The Electromagnetic Counterpart of the Binary Neutron Star Merger LIGO/VIRGO GW170817. III. Optical and UV Spectra of a Blue Kilonova From Fast Polar Ejecta, *Astrophys. J. Lett.* **848**, L18 (2017), [arXiv:1710.05456 \[astro-ph.HE\]](#).
- [36] E. Troja, L. Piro, G. Ryan, H. van Eerten, R. Ricci, M. Wieringa, S. Lotti, T. Sakamoto, and S. B. Cenko, The outflow structure of GW170817 from late-time broad-band observations, *Mon. Not. Roy. Astron. Soc.* **478**, L18 (2018), [arXiv:1801.06516 \[astro-ph.HE\]](#).
- [37] N. R. Tanvir *et al.*, The Emergence of a Lanthanide-Rich Kilonova Following the Merger of Two Neutron Stars, *Astrophys. J. Lett.* **848**, L27 (2017), [arXiv:1710.05455 \[astro-ph.HE\]](#).
- [38] M. R. Drout *et al.*, Light Curves of the Neutron Star Merger GW170817/SSS17a: Implications for R-Process Nucleosynthesis, *Science* **358**, 1570 (2017), [arXiv:1710.05443 \[astro-ph.HE\]](#).
- [39] L. Baiotti and L. Rezzolla, Binary neutron star mergers: a review of Einstein's richest laboratory, *Rept. Prog. Phys.* **80**, 096901 (2017), [arXiv:1607.03540 \[gr-qc\]](#).
- [40] B. D. Metzger, Kilonovae, *Living Rev. Rel.* **23**, 1 (2020), [arXiv:1910.01617 \[astro-ph.HE\]](#).
- [41] D. Radice, S. Bernuzzi, and A. Perego, The Dynamics of Binary Neutron Star Mergers and GW170817, *Ann. Rev. Nucl. Part. Sci.* **70**, 95 (2020), [arXiv:2002.03863 \[astro-ph.HE\]](#).
- [42] M. Shibata, Constraining nuclear equations of state using gravitational waves from hypermassive neutron stars, *Phys. Rev. Lett.* **94**, 201101 (2005), [arXiv:gr-qc/0504082](#).
- [43] M. Punturo *et al.*, The Einstein Telescope: A third-generation gravitational wave observatory, *Class. Quant. Grav.* **27**, 194002 (2010).
- [44] D. Reitze *et al.*, Cosmic Explorer: The U.S. Contribution to Gravitational-Wave Astronomy beyond LIGO, *Bull. Am. Astron. Soc.* **51**, 035 (2019), [arXiv:1907.04833 \[astro-ph.IM\]](#).
- [45] K. Ackley *et al.*, Neutron Star Extreme Matter Observatory: A kilohertz-band gravitational-wave detector in the global network, *Publ. Astron. Soc. Austral.* **37**, e047 (2020), [arXiv:2007.03128 \[astro-ph.HE\]](#).
- [46] M. Wijngaarden, K. Chatziioannou, A. Bauswein, J. A. Clark, and N. J. Cornish, Probing neutron stars with the full premerger and postmerger gravitational wave signal from binary coalescences, *Phys. Rev. D* **105**, 104019 (2022), [arXiv:2202.09382 \[gr-qc\]](#).
- [47] M. Breschi, R. Gamba, S. Borhanian, G. Carullo, and S. Bernuzzi, Kilohertz Gravitational Waves from Binary Neutron Star Mergers: Inference of Postmerger Signals with the Einstein Telescope, (2022), [arXiv:2205.09979 \[gr-qc\]](#).
- [48] H. Yu, D. Martynov, R. X. Adhikari, and Y. Chen, Exposing Gravitational Waves below the Quantum Shot Noise, (2022), [arXiv:2205.14197 \[gr-qc\]](#).
- [49] A. Bauswein, H. T. Janka, K. Hebeler, and A. Schwenk,

- Equation-of-state dependence of the gravitational-wave signal from the ring-down phase of neutron-star mergers, *Phys. Rev. D* **86**, 063001 (2012), [arXiv:1204.1888 \[astro-ph.SR\]](#).
- [50] A. Bauswein, N. Stergioulas, and H. T. Janka, Revealing the high-density equation of state through binary neutron star mergers, *Phys. Rev. D* **90**, 023002 (2014), [arXiv:1403.5301 \[astro-ph.SR\]](#).
- [51] S. Bernuzzi, T. Dietrich, and A. Nagar, Modeling the complete gravitational wave spectrum of neutron star mergers, *Phys. Rev. Lett.* **115**, 091101 (2015), [arXiv:1504.01764 \[gr-qc\]](#).
- [52] K. Takami, L. Rezzolla, and L. Baiotti, Constraining the Equation of State of Neutron Stars from Binary Mergers, *Phys. Rev. Lett.* **113**, 091104 (2014), [arXiv:1403.5672 \[gr-qc\]](#).
- [53] K. Takami, L. Rezzolla, and L. Baiotti, Spectral properties of the post-merger gravitational-wave signal from binary neutron stars, *Phys. Rev. D* **91**, 064001 (2015), [arXiv:1412.3240 \[gr-qc\]](#).
- [54] L. Rezzolla and K. Takami, Gravitational-wave signal from binary neutron stars: a systematic analysis of the spectral properties, *Phys. Rev. D* **93**, 124051 (2016), [arXiv:1604.00246 \[gr-qc\]](#).
- [55] S. Bose, K. Chakravarti, L. Rezzolla, B. S. Sathyaprakash, and K. Takami, Neutron-star Radius from a Population of Binary Neutron Star Mergers, *Phys. Rev. Lett.* **120**, 031102 (2018), [arXiv:1705.10850 \[gr-qc\]](#).
- [56] S. Vretinaris, N. Stergioulas, and A. Bauswein, Empirical relations for gravitational-wave asteroseismology of binary neutron star mergers, *Phys. Rev. D* **101**, 084039 (2020), [arXiv:1910.10856 \[gr-qc\]](#).
- [57] C. A. Raithel and E. R. Most, Characterizing the breakdown of quasi-universality in the post-merger gravitational waves from binary neutron star mergers, (2022), [arXiv:2201.03594 \[astro-ph.HE\]](#).
- [58] M. Breschi, S. Bernuzzi, K. Chakravarti, A. Camilletti, A. Prakash, and A. Perego, Kilohertz Gravitational Waves From Binary Neutron Star Mergers: Numerical-relativity Informed Postmerger Model, (2022), [arXiv:2205.09112 \[gr-qc\]](#).
- [59] A. Bauswein, H. T. Janka, and R. Oechslin, Testing Approximations of Thermal Effects in Neutron Star Merger Simulations, *Phys. Rev. D* **82**, 084043 (2010), [arXiv:1006.3315 \[astro-ph.SR\]](#).
- [60] A. Figura, J. J. Lu, G. F. Burgio, Z. H. Li, and H. J. Schulze, Hybrid equation of state approach in binary neutron-star merger simulations, *Phys. Rev. D* **102**, 043006 (2020), [arXiv:2005.08691 \[gr-qc\]](#).
- [61] C. Raithel, V. Paschalidis, and F. Özel, Realistic finite-temperature effects in neutron star merger simulations, *Phys. Rev. D* **104**, 063016 (2021), [arXiv:2104.07226 \[astro-ph.HE\]](#).
- [62] R. Oechslin, K. Uryu, G. S. Poghosyan, and F. K. Thielemann, The Influence of quark matter at high densities on binary neutron star mergers, *Mon. Not. Roy. Astron. Soc.* **349**, 1469 (2004), [arXiv:astro-ph/0401083](#).
- [63] E. R. Most, L. J. Papenfort, V. Dexheimer, M. Hanauske, S. Schramm, H. Stöcker, and L. Rezzolla, Signatures of quark-hadron phase transitions in general-relativistic neutron-star mergers, *Phys. Rev. Lett.* **122**, 061101 (2019), [arXiv:1807.03684 \[astro-ph.HE\]](#).
- [64] A. Bauswein, N.-U. F. Bastian, D. B. Blaschke, K. Chatziioannou, J. A. Clark, T. Fischer, and M. Oertel, Identifying a first-order phase transition in neutron star mergers through gravitational waves, *Phys. Rev. Lett.* **122**, 061102 (2019), [arXiv:1809.01116 \[astro-ph.HE\]](#).
- [65] E. R. Most, L. Jens Papenfort, V. Dexheimer, M. Hanauske, H. Stoecker, and L. Rezzolla, On the deconfinement phase transition in neutron-star mergers, *Eur. Phys. J. A* **56**, 59 (2020), [arXiv:1910.13893 \[astro-ph.HE\]](#).
- [66] L. R. Weih, M. Hanauske, and L. Rezzolla, Postmerger Gravitational-Wave Signatures of Phase Transitions in Binary Mergers, *Phys. Rev. Lett.* **124**, 171103 (2020), [arXiv:1912.09340 \[gr-qc\]](#).
- [67] K. Chatziioannou and S. Han, Studying strong phase transitions in neutron stars with gravitational waves, *Phys. Rev. D* **101**, 044019 (2020), [arXiv:1911.07091 \[gr-qc\]](#).
- [68] A. Prakash, D. Radice, D. Logoteta, A. Perego, V. Nedora, I. Bombaci, R. Kashyap, S. Bernuzzi, and A. Endrizzi, Signatures of deconfined quark phases in binary neutron star mergers, *Phys. Rev. D* **104**, 083029 (2021), [arXiv:2106.07885 \[astro-ph.HE\]](#).
- [69] S. Tootle, C. Ecker, K. Topolski, T. Demircik, M. Järvinen, and L. Rezzolla, Quark formation and phenomenology in binary neutron-star mergers using V-QCD, (2022), [arXiv:2205.05691 \[astro-ph.HE\]](#).
- [70] M. G. Alford, L. Bovard, M. Hanauske, L. Rezzolla, and K. Schwenzer, Viscous Dissipation and Heat Conduction in Binary Neutron-Star Mergers, *Phys. Rev. Lett.* **120**, 041101 (2018), [arXiv:1707.09475 \[gr-qc\]](#).
- [71] D. Radice, S. Bernuzzi, A. Perego, and R. Haas, A new moment-based general-relativistic neutrino-radiation transport code: Methods and first applications to neutron star mergers, *Mon. Not. Roy. Astron. Soc.* **512**, 1499 (2022), [arXiv:2111.14858 \[astro-ph.HE\]](#).
- [72] E. R. Most, S. P. Harris, C. Plumberg, M. G. Alford, J. Noronha, J. Noronha-Hostler, F. Pretorius, H. Witek, and N. Yunes, Projecting the likely importance of weak-interaction-driven bulk viscosity in neutron star mergers, *Mon. Not. Roy. Astron. Soc.* **509**, 1096 (2021), [arXiv:2107.05094 \[astro-ph.HE\]](#).
- [73] M. G. Alford and S. P. Harris, Damping of density oscillations in neutrino-transparent nuclear matter, *Phys. Rev. C* **100**, 035803 (2019), [arXiv:1907.03795 \[nucl-th\]](#).
- [74] M. Alford, A. Harutyunyan, and A. Sedrakian, Bulk viscosity of baryonic matter with trapped neutrinos, *Phys. Rev. D* **100**, 103021 (2019), [arXiv:1907.04192 \[astro-ph.HE\]](#).
- [75] M. G. Alford, A. Haber, S. P. Harris, and Z. Zhang, Beta Equilibrium Under Neutron Star Merger Conditions, *Universe* **7**, 399 (2021), [arXiv:2108.03324 \[nucl-th\]](#).
- [76] M. Alford, A. Harutyunyan, and A. Sedrakian, Bulk viscosity from Urca processes: $npe\mu$ matter in the neutrino-trapped regime, *Phys. Rev. D* **104**, 103027 (2021), [arXiv:2108.07523 \[astro-ph.HE\]](#).
- [77] P. Hammond, I. Hawke, and N. Andersson, Thermal aspects of neutron star mergers, *Phys. Rev. D* **104**, 103006 (2021), [arXiv:2108.08649 \[astro-ph.HE\]](#).
- [78] E. R. Most and C. A. Raithel, Impact of the nuclear symmetry energy on the post-merger phase of a binary neutron star coalescence, *Phys. Rev. D* **104**, 124012 (2021), [arXiv:2107.06804 \[astro-ph.HE\]](#).
- [79] R. Ardevol-Pulpillo, H. T. Janka, O. Just, and A. Bauswein, Improved Leakage-Equilibration-Absorption Scheme (ILEAS) for Neutrino Physics in Compact Object Mergers, *Mon. Not. Roy. Astron. Soc.* **485**, 4754 (2019), [arXiv:1808.00006 \[astro-ph.HE\]](#).
- [80] P. Hammond, I. Hawke, and N. Andersson, Detecting the impact of nuclear reactions on neutron star mergers through gravitational waves, (2022), [arXiv:2205.11377 \[astro-ph.HE\]](#).
- [81] A. Monnai and T. Hirano, Effects of Bulk Viscosity at Freeze-out, *Phys. Rev. C* **80**, 054906 (2009), [arXiv:0903.4436 \[nucl-th\]](#).

- [82] H. Song and U. W. Heinz, Interplay of shear and bulk viscosity in generating flow in heavy-ion collisions, *Phys. Rev. C* **81**, 024905 (2010), arXiv:0909.1549 [nucl-th].
- [83] P. Bozek, Bulk and shear viscosities of matter created in relativistic heavy-ion collisions, *Phys. Rev. C* **81**, 034909 (2010), arXiv:0911.2397 [nucl-th].
- [84] K. Dusling and T. Schäfer, Bulk viscosity, particle spectra and flow in heavy-ion collisions, *Phys. Rev. C* **85**, 044909 (2012), arXiv:1109.5181 [hep-ph].
- [85] J. Noronha-Hostler, G. S. Denicol, J. Noronha, R. P. G. Andrade, and F. Grassi, Bulk Viscosity Effects in Event-by-Event Relativistic Hydrodynamics, *Phys. Rev. C* **88**, 044916 (2013), arXiv:1305.1981 [nucl-th].
- [86] S. Ryu, J. F. Paquet, C. Shen, G. S. Denicol, B. Schenke, S. Jeon, and C. Gale, Importance of the Bulk Viscosity of QCD in Ultrarelativistic Heavy-Ion Collisions, *Phys. Rev. Lett.* **115**, 132301 (2015), arXiv:1502.01675 [nucl-th].
- [87] S. Ryu, J.-F. Paquet, C. Shen, G. Denicol, B. Schenke, S. Jeon, and C. Gale, Effects of bulk viscosity and hadronic rescattering in heavy ion collisions at energies available at the BNL Relativistic Heavy Ion Collider and at the CERN Large Hadron Collider, *Phys. Rev. C* **97**, 034910 (2018), arXiv:1704.04216 [nucl-th].
- [88] P. Romatschke and U. Romatschke, *Relativistic Fluid Dynamics In and Out of Equilibrium*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, 2019) arXiv:1712.05815 [nucl-th].
- [89] D. G. Yakovlev and K. P. Levenfish, Modified URCA process in neutron star cores., *Astron. & Astrophys.* **297**, 717 (1995).
- [90] M. D. Duez, Y. T. Liu, S. L. Shapiro, and B. C. Stephens, Relativistic magnetohydrodynamics in dynamical spacetimes: Numerical methods and tests, *Phys. Rev. D* **72**, 024028 (2005), arXiv:astro-ph/0503420.
- [91] D. Hilditch, S. Bernuzzi, M. Thierfelder, Z. Cao, W. Tichy, and B. Bruegmann, Compact binary evolutions with the Z4c formulation, *Phys. Rev. D* **88**, 084057 (2013), arXiv:1212.2901 [gr-qc].
- [92] G. S. Denicol and D. H. Rischke, *Microscopic Foundations of Relativistic Fluid Dynamics* (Springer, 2021).
- [93] Y. Sekiguchi, K. Kiuchi, K. Kyutoku, and M. Shibata, Current Status of Numerical-Relativity Simulations in Kyoto, *PTEP* **2012**, 01A304 (2012), arXiv:1206.5927 [astro-ph.HE].
- [94] F. Galeazzi, W. Kastaun, L. Rezzolla, and J. A. Font, Implementation of a simplified approach to radiative transfer in general relativity, *Phys. Rev. D* **88**, 064009 (2013), arXiv:1306.4953 [gr-qc].
- [95] M. H. Ruffert, H. T. Janka, and G. Schaefer, Coalescing neutron stars: A Step towards physical models. 1: Hydrodynamic evolution and gravitational wave emission, *Astron. Astrophys.* **311**, 532 (1996), arXiv:astro-ph/9509006.
- [96] S. Rosswog and M. Liebendörfer, High-resolution calculations of merging neutron stars - II. Neutrino emission, *Mon. Not. Roy. Astron. Soc.* **342**, 673 (2003), arXiv:astro-ph/0302301 [astro-ph].
- [97] D. G. Yakovlev, A. D. Kaminker, O. Y. Gnedin, and P. Haensel, Neutrino emission from neutron stars, *Phys. Rept.* **354**, 1 (2001), arXiv:astro-ph/0012122.
- [98] J. M. Lattimer, M. Prakash, C. J. Pethick, and P. Haensel, Direct URCA process in neutron stars, *Phys. Rev. Lett.* **66**, 2701 (1991).
- [99] M. Hempel and J. Schaffner-Bielich, Statistical Model for a Complete Supernova Equation of State, *Nucl. Phys. A* **837**, 210 (2010), arXiv:0911.4073 [nucl-th].
- [100] A. W. Steiner, M. Hempel, and T. Fischer, Core-collapse supernova equations of state based on neutron star observations, *Astrophys. J.* **774**, 17 (2013), arXiv:1207.2184 [astro-ph.SR].
- [101] E. R. Most, L. J. Papenfort, and L. Rezzolla, Beyond second-order convergence in simulations of magnetized binary neutron stars with realistic microphysics, *Mon. Not. Roy. Astron. Soc.* **490**, 3588 (2019), arXiv:1907.10328 [astro-ph.HE].
- [102] Z. B. Etienne, V. Paschalidis, R. Haas, P. Mösta, and S. L. Shapiro, IllinoisGRMHD: An Open-Source, User-Friendly GRMHD Code for Dynamical Spacetimes, *Class. Quant. Grav.* **32**, 175009 (2015), arXiv:1501.07276 [astro-ph.HE].
- [103] F. Löffler *et al.*, The Einstein Toolkit: A Community Computational Infrastructure for Relativistic Astrophysics, *Class. Quant. Grav.* **29**, 115001 (2012), arXiv:1111.3344 [gr-qc].
- [104] A. Bauswein, T. W. Baumgarte, and H. T. Janka, Prompt merger collapse and the maximum mass of neutron stars, *Phys. Rev. Lett.* **111**, 131101 (2013), arXiv:1307.5191 [astro-ph.SR].
- [105] S. Köppel, L. Bovard, and L. Rezzolla, A General-relativistic Determination of the Threshold Mass to Prompt Collapse in Binary Neutron Star Mergers, *Astrophys. J. Lett.* **872**, L16 (2019), arXiv:1901.09977 [gr-qc].
- [106] A. Bauswein, S. Blacker, V. Vijayan, N. Stergioulas, K. Chatziioannou, J. A. Clark, N.-U. F. Bastian, D. B. Blaschke, M. Cierniak, and T. Fischer, Equation of state constraints from the threshold binary mass for prompt collapse of neutron star mergers, *Phys. Rev. Lett.* **125**, 141103 (2020), arXiv:2004.00846 [astro-ph.HE].
- [107] S. D. Tootle, L. J. Papenfort, E. R. Most, and L. Rezzolla, Quasi-universal Behavior of the Threshold Mass in Unequal-mass, Spinning Binary Neutron Star Mergers, *Astrophys. J. Lett.* **922**, L19 (2021), arXiv:2109.00940 [gr-qc].
- [108] R. Kashyap *et al.*, Numerical relativity simulations of prompt collapse mergers: Threshold mass and phenomenological constraints on neutron star properties after GW170817, *Phys. Rev. D* **105**, 103022 (2022), arXiv:2111.05183 [astro-ph.HE].
- [109] M. Kölsch, T. Dietrich, M. Ujevic, and B. Bruegmann, Investigating the mass-ratio dependence of the prompt-collapse threshold with numerical-relativity simulations, (2021), arXiv:2112.11851 [gr-qc].
- [110] A. Perego, D. Logoteta, D. Radice, S. Bernuzzi, R. Kashyap, A. Das, S. Padamata, and A. Prakash, Probing the incompressibility of nuclear matter at ultra-high density through the prompt collapse of asymmetric neutron star binaries, (2021), arXiv:2112.05864 [astro-ph.HE].
- [111] P. Arras and N. N. Weinberg, Urca reactions during neutron star inspiral, *Mon. Not. Roy. Astron. Soc.* **486**, 1424 (2019), arXiv:1806.04163 [astro-ph.HE].
- [112] A. Perego, S. Bernuzzi, and D. Radice, Thermodynamics conditions of matter in neutron star mergers, *Eur. Phys. J. A* **55**, 124 (2019), arXiv:1903.07898 [gr-qc].
- [113] S. Rosswog, M. Liebendoerfer, F. K. Thielemann, M. B. Davies, W. Benz, and T. Piran, Mass ejection in neutron star mergers, *Astron. Astrophys.* **341**, 499 (1999), arXiv:astro-ph/9811367.
- [114] A. Bauswein, S. Goriely, and H. T. Janka, Systematics of dynamical mass ejection, nucleosynthesis, and radioactively powered electromagnetic signals from neutron-star mergers, *Astrophys. J.* **773**, 78 (2013), arXiv:1302.6530 [astro-ph.SR].
- [115] Y. Sekiguchi, K. Kiuchi, K. Kyutoku, and M. Shibata, Dynamical mass ejection from binary neutron star mergers: Radiation-hydrodynamics study in general relativity, *Phys. Rev. D* **91**, 064059 (2015), arXiv:1502.06660 [astro-ph.HE].
- [116] Y. Sekiguchi, K. Kiuchi, K. Kyutoku, M. Shibata, and K. Taniguchi, Dynamical mass ejection from the merger of

- asymmetric binary neutron stars: Radiation-hydrodynamics study in general relativity, *Phys. Rev. D* **93**, 124046 (2016), [arXiv:1603.01918 \[astro-ph.HE\]](#).
- [117] D. Radice, F. Galeazzi, J. Lippuner, L. F. Roberts, C. D. Ott, and L. Rezzolla, Dynamical Mass Ejection from Binary Neutron Star Mergers, *Mon. Not. Roy. Astron. Soc.* **460**, 3255 (2016), [arXiv:1601.02426 \[astro-ph.HE\]](#).
- [118] L. Lehner, S. L. Liebling, C. Palenzuela, O. L. Caballero, E. O'Connor, M. Anderson, and D. Neilsen, Unequal mass binary neutron star mergers and multimessenger signals, *Class. Quant. Grav.* **33**, 184002 (2016), [arXiv:1603.00501 \[gr-qc\]](#).
- [119] L. Bovard, D. Martin, F. Guercilena, A. Arcones, L. Rezzolla, and O. Korobkin, r -process nucleosynthesis from matter ejected in binary neutron star mergers, *Phys. Rev. D* **96**, 124005 (2017), [arXiv:1709.09630 \[gr-qc\]](#).
- [120] D. Radice, A. Perego, K. Hotokezaka, S. A. Fromm, S. Bernuzzi, and L. F. Roberts, Binary Neutron Star Mergers: Mass Ejection, Electromagnetic Counterparts and Nucleosynthesis, *Astrophys. J.* **869**, 130 (2018), [arXiv:1809.11161 \[astro-ph.HE\]](#).
- [121] V. Nedora, S. Bernuzzi, D. Radice, A. Perego, A. Endrizzi, and N. Ortiz, Spiral-wave wind for the blue kilonova, *Astrophys. J. Lett.* **886**, L30 (2019), [arXiv:1907.04872 \[astro-ph.HE\]](#).
- [122] T. Celora, I. Hawke, P. C. Hammond, N. Andersson, and G. L. Comer, Formulating bulk viscosity for neutron star simulations, *Phys. Rev. D* **105**, 103016 (2022), [arXiv:2202.01576 \[astro-ph.HE\]](#).
- [123] L. Gavassino, M. Antonelli, and B. Haskell, Bulk viscosity in relativistic fluids: from thermodynamics to hydrodynamics, *Class. Quant. Grav.* **38**, 075001 (2021), [arXiv:2003.04609 \[gr-qc\]](#).
- [124] G. Camelio, L. Gavassino, M. Antonelli, S. Bernuzzi, and B. Haskell, Simulating bulk viscosity in neutron stars I: formalism, (2022), [arXiv:2204.11809 \[gr-qc\]](#).
- [125] N. Stergioulas, A. Bauswein, K. Zagkouris, and H.-T. Janka, Gravitational waves and nonaxisymmetric oscillation modes in mergers of compact object binaries, *Mon. Not. Roy. Astron. Soc.* **418**, 427 (2011), [arXiv:1105.0368 \[gr-qc\]](#).
- [126] G. S. Denicol, H. Niemi, E. Molnar, and D. H. Rischke, Derivation of transient relativistic fluid dynamics from the Boltzmann equation, *Phys. Rev. D* **85**, 114047 (2012), [Erratum: *Phys.Rev.D* 91, 039902 (2015)], [arXiv:1202.4551 \[nucl-th\]](#).
- [127] S. Fujibayashi, Y. Sekiguchi, K. Kiuchi, and M. Shibata, Properties of Neutrino-driven Ejecta from the Remnant of a Binary Neutron Star Merger: Pure Radiation Hydrodynamics Case, *Astrophys. J.* **846**, 114 (2017), [arXiv:1703.10191 \[astro-ph.HE\]](#).
- [128] S. Fujibayashi, K. Kiuchi, N. Nishimura, Y. Sekiguchi, and M. Shibata, Mass Ejection from the Remnant of a Binary Neutron Star Merger: Viscous-Radiation Hydrodynamics Study, *Astrophys. J.* **860**, 64 (2018), [arXiv:1711.02093 \[astro-ph.HE\]](#).
- [129] E. R. Most, A. Motornenko, J. Steinheimer, V. Dexheimer, M. Hanauske, L. Rezzolla, and H. Stoecker, Probing neutron-star matter in the lab: connecting binary mergers to heavy-ion collisions, (2022), [arXiv:2201.13150 \[nucl-th\]](#).
- [130] F. S. Bemfica, M. M. Disconzi, and J. Noronha, Causality and existence of solutions of relativistic viscous fluid dynamics with gravity, *Phys. Rev. D* **98**, 104064 (2018), [arXiv:1708.06255 \[gr-qc\]](#).
- [131] F. S. Bemfica, M. M. Disconzi, and J. Noronha, Causality of the Einstein-Israel-Stewart Theory with Bulk Viscosity, *Phys. Rev. Lett.* **122**, 221602 (2019), [arXiv:1901.06701 \[gr-qc\]](#).
- [132] F. S. Bemfica, M. M. Disconzi, and J. Noronha, Nonlinear Causality of General First-Order Relativistic Viscous Hydrodynamics, *Phys. Rev. D* **100**, 104020 (2019), [Erratum: *Phys.Rev.D* 105, 069902 (2022)], [arXiv:1907.12695 \[gr-qc\]](#).
- [133] F. S. Bemfica, M. M. Disconzi, and J. Noronha, First-Order General-Relativistic Viscous Fluid Dynamics, *Phys. Rev. X* **12**, 021044 (2022), [arXiv:2009.11388 \[gr-qc\]](#).
- [134] A. Pandya and F. Pretorius, Numerical exploration of first-order relativistic hydrodynamics, *Phys. Rev. D* **104**, 023015 (2021), [arXiv:2104.00804 \[gr-qc\]](#).
- [135] M. Chabanov, L. Rezzolla, and D. H. Rischke, General-relativistic hydrodynamics of non-perfect fluids: 3+1 conservative formulation and application to viscous black hole accretion, *Mon. Not. Roy. Astron. Soc.* **505**, 5910 (2021), [arXiv:2102.10419 \[gr-qc\]](#).
- [136] E. R. Most and J. Noronha, Dissipative magnetohydrodynamics for nonresistive relativistic plasmas: An implicit second-order flux-conservative formulation with stiff relaxation, *Phys. Rev. D* **104**, 103028 (2021), [arXiv:2109.02796 \[astro-ph.HE\]](#).
- [137] A. Pandya, E. R. Most, and F. Pretorius, Conservative finite volume scheme for first-order viscous relativistic hydrodynamics, *Phys. Rev. D* **105**, 123001 (2022), [arXiv:2201.12317 \[gr-qc\]](#).
- [138] G. Camelio, L. Gavassino, M. Antonelli, S. Bernuzzi, and B. Haskell, Simulating bulk viscosity in neutron stars II: evolution in spherical symmetry, (2022), [arXiv:2204.11810 \[gr-qc\]](#).
- [139] R. Gill, A. Nathanail, and L. Rezzolla, When Did the Remnant of GW170817 Collapse to a Black Hole?, *Astrophys. J.* **876**, 139 (2019), [arXiv:1901.04138 \[astro-ph.HE\]](#).
- [140] M. G. Alford and S. P. Harris, Beta equilibrium in neutron star mergers, *Phys. Rev. C* **98**, 065806 (2018), [arXiv:1803.00662 \[nucl-th\]](#).
- [141] M. G. Alford and A. Haber, Strangeness-changing Rates and Hyperonic Bulk Viscosity in Neutron Star Mergers, *Phys. Rev. C* **103**, 045810 (2021), [arXiv:2009.05181 \[nucl-th\]](#).
- [142] A. Schmitt and P. Shternin, Reaction rates and transport in neutron stars, *Astrophys. Space Sci. Libr.* **457**, 455 (2018), [arXiv:1711.06520 \[astro-ph.HE\]](#).
- [143] E. R. Most and A. A. Philippov, Electromagnetic precursor flares from the late inspiral of neutron star binaries, (2022), [arXiv:2205.09643 \[astro-ph.HE\]](#).
- [144] J. Towns *et al.*, XSEDE: Accelerating Scientific Discovery, *Comput. Sci. Eng.* **16**, 62 (2014).
- [145] L. F. Roberts and S. Reddy, Charged current neutrino interactions in hot and dense matter, *Phys. Rev. C* **95**, 045807 (2017), [arXiv:1612.02764 \[astro-ph.HE\]](#).
- [146] P. Haensel, K. P. Levenfish, and D. G. Yakovlev, Adiabatic index of dense matter and damping of neutron star pulsations, *Astron. Astrophys.* **394**, 213 (2002), [arXiv:astro-ph/0208078](#).
- [147] M. G. Alford, S. Mahmoodifar, and K. Schwenzer, Large amplitude behavior of the bulk viscosity of dense matter, *J. Phys. G* **37**, 125202 (2010), [arXiv:1005.3769 \[nucl-th\]](#).