Braiding with Majorana lattices: Groundstate degeneracy and supersymmetry

Pasquale Marra[®],^{1,2,*} Daisuke Inotani[®],² and Muneto Nitta[®]²

¹Graduate School of Mathematical Sciences, The University of Tokyo, 3-8-1 Komaba, Meguro, Tokyo, 153-8914, Japan

²Department of Physics, and Research and Education Center for Natural Sciences,

Keio University, 4-1-1 Hiyoshi, Yokohama, Kanagawa, 223-8521, Japan

(Dated: September 19, 2022)

Majorana-based topological qubits are expected to exploit the nonabelian braiding statistics of Majorana modes in topological superconductors to realize fault-tolerant topological quantum computation. Scalable qubit designs require several Majorana modes localized on quantum wires networks, with braiding operations relying on the presence of the groundstate degeneracy of the topologically nontrivial superconducting phase. However, this degeneracy is lifted due to the hybridization between Majorana modes localized at a finite distance. Here, we describe a braiding protocol in a trijunction where each branch consists of a lattice of Majorana modes overlapping at a finite distance. We find that the energy splitting between the groundstate and the lowest-energy state decreases exponentially with the number of Majorana modes if the system is in its topologically nontrivial regime. This result does not rely on the specific braiding geometry and on the details of the braiding scheme but is a consequence of the supersymmetry and nontrivial topology of the effective low-energy Hamiltonian describing the Majorana lattice.

I. INTRODUCTION

Topologically-protected Majorana zero modes localized at the boundaries of the nontrivial phase of a one-dimensional (1D) topological superconductor [1–3] are expected to exhibit nonabelian exchange statistics, which can be exploited to realize the quantum gates of a topological quantum computer [4–6]. Motivated by their potential impact in the field of quantum computing, there is an ongoing experimental effort to realize Majorana modes in realistic and scalable devices, including semiconductor planar heterostructures, and in arrays of magnetic adatoms on a conventional superconductor substrate (see Refs. [7, 8]). In principle, the nonabelian braiding statistics can be probed by adiabatically exchanging Majorana modes in physical space in a trijunction [9–12]. The exchange operations of a number 2*N* of Majorana modes are unitary transformations acting on the corresponding 2^{*N*}-fold degenerate groundstate. However, this groundstate degeneracy is exact only for Majorana modes localized at an infinite distance one from the other. In the real world, however, Majorana modes gain a finite but exponentially small energy $\propto e^{-l/\xi_M}$, where *l* is the distance between the two Majorana modes and ξ_M their localization length. This energy splitting is a source of quantum decoherence [13], which is a fundamental hindrance to the realization of fault-tolerant topological quantum gates. Moreover, scalable qubit designs require several Majorana modes localized on 2D or even 3D networks containing multiple trijunctions as building blocks [9, 10, 14, 15]. Hence, it is crucial to determine the stability and robustness of the groundstate degeneracy in braiding designs containing several Majorana modes overlapping at a finite distance.

Recently, we proposed the realization of arrays of several partially-overlapping, zero-dimensional Majorana modes in proximitized nanowires via periodically-modulated magnetic fields superimposed by a rotating field in the same plane [16, 17]. This lattice of Majorana modes can realize a "Majorana pump" with the adiabatic translation of one Majorana mode for each half rotation of the externally applied field. Remarkably, the groundstate degeneracy becomes exact in loop geometries or for an infinite number of Majorana modes [16, 17]. This exact degeneracy is a direct consequence of the quantum mechanical supersymmetry (SUSY) of the groundstate [16–19], which describes the symmetry between many-body states with different fermion parity, and which is generally present in topological superconductors [18–23] and their mechanical analogues [24, 25]. Here, by exploiting these ideas, we describe the realization of a braiding protocol in a trijunction, where each branch realizes a sliding lattice of Majorana modes overlapping at a finite distance, described by an effective \mathbb{Z}_2 topological invariant. We find that, in the topologically nontrivial phase, the presence of several Majorana modes does not increase but exponentially suppresses the energy splitting, restoring the groundstate degeneracy in the limit of infinitely many Majorana modes.

II. TRIJUNCTION WITH MAJORANA LATTICES

We consider a T-shaped or Y-shaped trijunction with a lattice of 2N Majorana modes on each branch. In order to obtain a lattice of regularly-spaced Majorana modes, one can employ periodical magnetic textures induced by nanomagnets [16, 17]. In this setup, the boundary between trivial and nontrivial segments can be smoothly controlled by rotating an applied field, inducing the adiabatic translation of the Majorana modes, i.e., and adiabatic "Majorana pump" [16, 17]. Alternatively, regularly-spaced Majorana modes can be produced in the presence of arrays of tunable spin-valves [26] or magnetic tunnel junctions [27, 28], or periodic arrays of electric gates [9, 29, 30]. In these configurations, by switching on and off the spin-valves, magnetic tunnel



FIG. 1. (a) Piano keyboard setup, where Majorana modes localized at the boundaries between trivial and nontrivial segments can be moved by manipulating magnetic textures or electric gates. (b) Majorana modes on a trijunction described by the effective Hamiltonian in Eq. (1). (c) Translational-invariant bipartite lattice with alternate couplings w and v.

junctions, or electric gates, one can drive different segments of the wire in and out of the topologically nontrivial phase to obtain a regular lattice of Majorana modes localized at the boundaries between trivial and nontrivial segments. In this "piano keyboard" setup [9, 29, 30], the Majorana modes can slide by slowly rearranging the magnetic texture or the electric gates configurations, as shown in Fig. 1(a). Alternatively, one can consider a Y-junction in planar Josephson junctions formed by depositing superconducting islands on top of a topological insulator, with Majorana modes localized at the superconducting vortex cores, which can be moved by applied currents, voltages, or phase differences [31].

If the lengths of the trivial and nontrivial segments are comparable to the Majorana localization length, there is a small overlap between the Majorana modes wavefunctions. In this case the low-energy Hamiltonian of the system can be obtained by projecting onto the subspace of Majorana operators, which in a trijunction configuration gives

$$\mathcal{H}_{\text{eff}} = i\Gamma H\Gamma^{\mathsf{T}} = i\left(u_{1,2}\gamma_{1,1}\gamma_{2,1} + u_{2,3}\gamma_{2,1}\gamma_{3,1} + u_{3,1}\gamma_{3,1}\gamma_{1,1}\right) + i\sum_{m=1}^{3}\sum_{n=1}^{2N-1} w_{m,n}\gamma_{m,n}\gamma_{m,n+1},\tag{1}$$

where $\gamma_{m,n}$ are the Majorana operators on each branch m = 1, 2, 3 and with $n = 1, 2, \ldots, 2N$ counting outward from the center, as shown in Fig. 1(b), $\Gamma = (\gamma_{1,1}, \gamma_{2,1}, \gamma_{3,1}, \ldots, \gamma_{1,2N}, \gamma_{2,2N}, \gamma_{3,2N})$ the vector of the Majorana operators, $u_{m,m'} \in \mathbb{R}$ the coupling between Majorana modes $\gamma_{m,1}$ and $\gamma_{m',1}$ at the center of the junction, and $w_{m,n} \in \mathbb{R}$ the couplings between contiguous Majorana modes $\gamma_{m,n}$ and $\gamma_{m,n+1}$ on each branch. Let us also assume that all the couplings can be written as $w_{m,n} = E_0 e^{-l_{m,n}/\xi_M}$ in terms of the Majorana localization length ξ_M where $l_{m,n}$ is the distance between contiguous Majorana modes $\gamma_{m,n}$ and E_0 a characteristic energy scale of the junction. We will later consider a simpler case where each branch forms a translational-invariant bipartite lattice with $w_{m,n} = w$ or $w_{m,n} = v$ for n odd and even, respectively, and with $u_{1,2} = u_{2,3} = u_{3,1} = u$, as in Fig. 1(c).

III. BRAIDING PROTOCOL

The Majorana modes can move along the three branches of the trijunction by manipulating magnetic textures or electric gates, as already mentioned [see Fig. 1(a)]. A simple braiding protocol which exchanges two Majorana modes at the center of the junction can be engineered by sliding the lattices of Majorana modes separately on the three different branches, as illustrated in Fig. 2. This braiding protocol is similar to the braiding protocol of Majorana modes in planar Josephson junctions introduced in Ref. 31. As a first step, we slide the Majorana lattices on the upper left and lower branches in the downward direction, such that one Majorana mode from the upper left branch crosses the center of the junction, reaching the lower branch. We then slide the Majorana lattices on the upper right and left branches in the leftward direction, such that one Majorana mode from the right branch. Finally, we slide the Majorana lattices on the upper right and lower branch. Finally, we slide the Majorana modes which were sitting close to the center of the junction into the right branch. At the end of the day, the Majorana modes which were sitting close to the center of the junction on the upper left and right branches are exchanged. Analogously, one can exchange the Majorana modes from the upper left and lower branches, and from the upper right and lower branches. If performed adiabatically, these processes can be described as the unitary braiding operators $U_{m,m'} = \frac{1}{\sqrt{2}} (1 + \gamma_{m,1} \gamma_{m',1}) = \exp(\frac{\pi}{4} \gamma_{m1} \gamma_{m',1})$ with $m, m' \in \{1, 2, 3\}$, which correspond to the adiabatic exchange of the two Majorana modes $\gamma_{m,1}$ and $\gamma_{m',1}$ at the center of the junction.



FIG. 2. Braiding of two Majorana modes in a trijunction obtained by sliding the Majorana modes at the boundaries between trivial and nontrivial segments. We move the Majorana modes on the upper left and lower branches downwards, then move the modes on the upper right and branches leftwards, and finally, move the modes on the upper right and lower branches upwards.

IV. GROUNDSTATE DEGENERACY AND SUPERSYMMETRY

Intuitively, one may expect that the presence of many overlapping Majorana modes would spoil the topological protection by lifting the degeneracy of the groundstate. However, this is not the case. To estimate the lowest energy level $E_1 = \Delta E$ of the Hamiltonian above we notice that, since the minimum of a list of non-negative numbers is less than or equal to the geometric average of all the 3N positive energy levels (corresponding to 6N Majorana modes) of the Hamiltonian above, which yields $\Delta E \leq (\prod_{n=1}^{3N} E_i)^{1/3N}$. Let us then recall that the product of the all energy levels is equal to the determinant of the Hamiltonian matrix, which is an antisymmetric matrix, giving $|\det H| = |\operatorname{pf} H|^2$, and that, due to the particle-hole symmetry of the superconductor, every energy level has a particle-hole symmetric level with opposite energy, which gives $|\det H| = \prod_{n=1}^{3N} E_i^2$. Hence, by explicitly calculating the the pfaffian of the Hamiltonian, one obtains

$$\Delta E \le \left(\prod_{i=1}^{3N} E_i\right)^{\frac{1}{3N}} = |\operatorname{pf} H|^{\frac{1}{3N}} = \left(\prod_{m=1}^{3} \prod_{n=1}^{N} w_{m,2n-1}\right)^{\frac{1}{3N}} = E_0 \mathrm{e}^{-\frac{1}{3N\xi_{\mathrm{M}}} \sum_{m=1}^{3} \sum_{n=1}^{N} l_{m,2n-1}}.$$
(2)

This equation gives an upper bound to the energy splitting between the groundstate and the lowest-energy level. Surprisingly, the geometric average does not depend on the couplings between the three central Majorana modes $\gamma_{1,1}$, $\gamma_{2,1}$, and $\gamma_{3,1}$, and depends only on the couplings $w_{m,2n-1}$ between the contiguous Majorana modes $\gamma_{m,2n-1}$ and $\gamma_{m,2n}$, but not on the couplings $w_{m,2n}$ of the contiguous modes $\gamma_{m,2n-1}$ and $\gamma_{m,2n}$ and $\gamma_{m,2n+1}$. This mandates that, if the coupling between a single pair of contiguous Majorana modes $\gamma_{m,2n-1}$ and $\gamma_{m,2n}$ on one of the branches is zero, there exists at least one energy level which is exactly zero. Hence, one can recover an exactly degenerate groundstate by tuning a single parameter. The presence of a zero-energy level in a system of several Majorana modes coupled together is a manifestation of quantum mechanical SUSY [16, 17, 19].

Let us consider the simpler case where each branch forms a translational-invariant bipartite lattice with $w_{m,n} = w = E_0 e^{-l_w/\xi_M} > 0$ and $w_{m,n} = v = E_0 e^{-l_v/\xi_M} > 0$ where l_w and l_v are the distances between contiguous Majorana modes $\gamma_{m,n}$ and $\gamma_{m,n+1}$ with n odd and even, respectively, and with $u_{1,2} = u_{2,3} = u_{3,1} = u$, as in Fig. 1(c). In this case the Hamiltonian in Eq. (1) can be written as

$$\mathcal{H}_{\text{eff}} = i\Gamma H\Gamma^{\mathsf{T}} = iu\left(\gamma_{1,1}\gamma_{2,1} + \gamma_{2,1}\gamma_{3,1} + \gamma_{3,1}\gamma_{1,1}\right) + i\sum_{m=1}^{3} \left(\sum_{n=1}^{N} w\gamma_{m,2n-1}\gamma_{m,2n} + \sum_{n=1}^{N-1} v\gamma_{m,2n}\gamma_{m,2n+1}\right), \quad (3)$$

The geometric average of all positive energy levels in this case becomes $|\operatorname{pf} H|^{\frac{1}{3N}} = w$, which gives an upper bound to the energy splitting $\Delta E \leq w$. This seems to suggest that, in a trijunction with branches of fixed length L = Nl with $l = l_w + l_v$ and $\Delta l = l_w - l_v$, the energy splitting would increase to a finite limit as $\Delta E \propto w = E_0 e^{-\Delta l/2\xi_M} e^{-L/2N\xi_M}$ with the number of Majorana modes 2N. As anticipated, this is not always the case, as we will show below.

The trijunction exhibits two topologically distinct phases realized respectively for |w| > |v| and |v| > |w|, which correspond to the trivial and nontrivial phases of the three Majorana lattices of the three branches of the trijunction. If the branches are decoupled u = 0, indeed, each branch can be described as a bipartite lattice of 0D Majorana modes with an effective \mathbb{Z}_2 topological invariant $\mathcal{P}_{\text{eff}} = \text{sgn}(|w| - |v|)$ (see Refs. [16, 17]). In the topologically trivial phase |w| > |v|, the energy splitting ΔE increases to a finite limit as $\Delta E \propto w \propto e^{-L/2N\xi_M}$ as the number of Majorana modes increases $N \to \infty$ (as expected from the above argument). At the topological phase transition |v| = |w|, the Majorana lattices exhibit two degenerate groundstates in the limit of infinite Majorana modes $N \to \infty$ (or, equivalently, in a finite lattice with periodic boundary conditions), which are SUSY multiplets with opposite fermion parity (the SUSY is spontaneously broken) [16–18]. The two degenerate groundstates



FIG. 3. Particle density ρ of the lowest-energy state of a trijunction with branches of length $L = 20\xi_{\rm M}$ with 2N = 40 Majorana modes, plotted on a single branch and on the three branches, and corresponding energy splitting ΔE as a function of the number of Majorana modes 2N. The data point corresponding to odd $\gamma_{m,2n-1}$ and even $\gamma_{m,2n}$ Majorana modes are grouped together. (a) Trivial phase |w| > |v| (w/v = 1.2). The Majorana modes on odd and even sites hybridize along the entire width of the trijunction branches. The energy splitting between the lowest-energy state and the groundstates increases and becomes proportional to the overlap w between contiguous Majorana modes $\gamma_{m,2n-1}$ and $\gamma_{m,2n}$, increasing to a finite limit as $\Delta E \propto w \propto e^{-L/2N\xi_{\rm M}}$. The continuous line at a constant value is a guide for the eye. (b) Topological phase transition |v| = |w|. The Majorana modes on odd and even sites become partially decoupled forming two Majorana modes delocalized on the whole lattice. In the limit $N \to \infty$, SUSY mandates that the $\Delta E \to 0$, with each branch of the trijunction exhibiting two degenerate groundstates which are SUSY multiplets with opposite fermion parity. As verified numerically, the energy splitting decreases polynomially as $\Delta E \propto w/N$. The continuous line is the best fit for $\Delta E/w \propto 1/N$. (c) Nontrivial phase |v| > |w| (w/v = 0.8). The Majorana modes on odd and even sites decouple and hybridize into left and right Majorana end modes exponentially localized respectively at the center and at the outer ends of the three branches, with localization length $\xi_{\rm eff}$. The energy splitting decreases exponentially as $\Delta E \propto e^{-2N/\xi_{\rm eff}}$. The continuous line is the best fit for $\log(\Delta E/w) \propto -2N/\xi_{\rm eff}$.

correspond to the vacuum and to counterpropagating and dispersive 1D Majorana modes delocalized on the whole length of the lattice [16, 17]. Hence, one expects that the energy splitting decreases with the number of Majorana modes, approaching zero for $N \to \infty$. We verify numerically that the energy splitting decreases polynomially with the inverse of the number of Majorana modes as $\Delta E \propto w/N$ for |v| = |w|, even for finite |u| > 0. In the topologically nontrivial phase |v| > |w| instead, each branch exhibits two Majorana end modes given by the hybridization of the 0D Majorana modes, given by the superposition of 0D Majorana modes on odd and even sites, are localized at the opposite ends, i.e., respectively at the center and at the outer ends of the trijunction, with effective localization length $\xi_{\text{eff}} = l/|\log |w/v|| = (l/|\Delta l|)\xi_M \ge \xi_M$. If the coupling between the three branches is finite |u| > 0, the three Majorana modes at the center hybridize into an unpaired Majorana mode at low energy [9]. The hybridization of the outer and inner Majorana modes yields a finite energy splitting $\Delta E \propto e^{-2N/\xi_{\text{eff}}}$

To numerically verify our findings, we diagonalize the Hamiltonian in Eq. (3) considering a trijunction with fixed length L and overlaps $w = E_0 e^{-\Delta l/2\xi_M} e^{-L/2N\xi_M}$ and $v = E_0 e^{+\Delta l/2\xi_M} e^{-L/2N\xi_M}$ with fixed Δl . Figure 3 shows the particle density ρ of the lowest-energy state and energy splitting ΔE as a function of the number of Majorana modes 2N. In the trivial phase |w| > |v|(i.e., $\Delta l > 0$), Majorana modes on odd and even sites hybridize along the trijunction, and the energy splitting increases to a finite limit as $\Delta E \propto w \propto e^{-L/2N\xi_M}$ [see Fig. 3(a)]. At the topological phase transition |v| = |w| (i.e., $\Delta l = 0$), the Majorana modes on odd and even sites hybridize into two Majorana modes, and the energy splitting decreases polynomially as $\Delta E \propto w/N$ [see Fig. 3(b)]. In the nontrivial phase |v| > |w| (i.e., $\Delta l < 0$), the Majorana modes on odd and even sites decouple and hybridize into left and right Majorana end modes, exponentially localized at the center and at the outer ends of the trijunction, with the energy splitting decreasing exponentially as $\Delta E \propto e^{-2N/\xi_{\text{eff}}}$ [see Fig. 3(c)]. In all these regimes, we numerically verified that the asymptotic behavior of the energy splitting is independent on the choice of the coupling u between the Majorana modes at the center of the junction.

The scaling of the energy splitting fully characterizes the different regimes of the trijunction: Indeed, the energy splitting increases to a finite limit, decreases exponentially, or decreases polynomially, respectively, in the trivial, nontrivial, and at the topological phase transition. One can also expect that this asymptotic behavior is preserved when the translational invariance is broken, i.e., in the presence of random disorder in the coupling between contiguous Majorana modes.

V. CONCLUSIONS

In this work, we analyzed the groundstate properties and described a braiding protocol for a trijunction where each branch contains an array of Majorana modes overlapping at a finite distance. We found that, in this setup, the energy splitting between the groundstate and the lowest-energy many-body state decreases exponentially with the number of Majorana modes if the effective low-energy Hamiltonian of the Majorana lattices is topologically nontrivial. This result does not depend on the geometry of the junction and on the details of the braiding scheme, and suggests that, in the nontrivial regime, the more Majorana modes there are in a topological qubit, the smaller the energy splitting.

ACKNOWLEDGMENTS

P. M. is supported by the Japan Science and Technology Agency (JST) of the Ministry of Education, Culture, Sports, Science and Technology (MEXT), JST CREST Grant No. JPMJCR19T2 and Japan Society for the Promotion of Science (JSPS) Grantin-Aid for Early-Career Scientists Grant No. 20K14375. M. N. is partially supported by the JSPS Grant-in-Aid for Scientific Research Grants No. JP18H01217 and No. JP22H01221.

- [5] A. Kitaev, Fault-tolerant quantum computation by anyons, Ann. Phys. 303, 2 (2003).
- [6] C. Nayak, S. Simon, A. Stern, M. Freedman, and S. Das Sarma, Non-Abelian anyons and topological quantum computation, Rev. Mod. Phys. 80, 1083 (2008).
- [7] S. Frolov, M. Manfra, and J. Sau, Topological superconductivity in hybrid devices, Nat. Phys. 16, 718 (2020).
- [8] K. Flensberg, F. von Oppen, and A. Stern, Engineered platforms for topological superconductivity and Majorana zero modes, Nat. Rev. Mater. 6, 944 (2021).
- [9] J. Alicea, Y. Oreg, G. Refael, F. Von Oppen, and M. Fisher, Non-Abelian statistics and topological quantum information processing in 1D wire networks, Nat. Phys. 7, 412 (2011).
- [10] D. Clarke, J. Sau, and S. Tewari, Majorana fermion exchange in quasi-one-dimensional networks, Phys. Rev. B 84, 035120 (2011).
- [11] A. Romito, J. Alicea, G. Refael, and F. Von Oppen, Manipulating Majorana fermions using supercurrents, Phys. Rev. B 85, 020502 (2012).

^{*} pmarra@ms.u-tokyo.ac.jp

^[1] A. Y. Kitaev, Unpaired Majorana fermions in quantum wires, Phys. Usp. 44, 131 (2001).

^[2] R. Lutchyn, J. Sau, and S. Das Sarma, Majorana fermions and a topological phase transition in semiconductor-superconductor heterostructures, Phys. Rev. Lett. 105, 077001 (2010).

^[3] Y. Oreg, G. Refael, and F. Von Oppen, Helical liquids and Majorana bound states in quantum wires, Phys. Rev. Lett. 105, 177002 (2010).

^[4] D. Ivanov, Non-Abelian statistics of half-quantum vortices in p-wave superconductors, Phys. Rev. Lett. 86, 268 (2001).

- [12] B. Halperin, Y. Oreg, A. Stern, G. Refael, J. Alicea, and F. Von Oppen, Adiabatic manipulations of Majorana fermions in a threedimensional network of quantum wires, Phys. Rev. B 85, 144501 (2012).
- [13] C. Knapp, T. Karzig, R. Lutchyn, and C. Nayak, Dephasing of Majorana-based qubits, Phys. Rev. B 97, 125404 (2018).
- [14] J. Sau, S. Tewari, and S. Das Sarma, Universal quantum computation in a semiconductor quantum wire network, Phys. Rev. A 82, 052322 (2010).
- [15] J. Sau, D. Clarke, and S. Tewari, Controlling non-Abelian statistics of Majorana fermions in semiconductor nanowires, Phys. Rev. B 84, 094505 (2011).
- [16] P. Marra, D. Inotani, and M. Nitta, 1D Majorana Goldstinos and partial supersymmetry breaking in quantum wires, Commun. Phys. 5, 149 (2022).
- [17] P. Marra, D. Inotani, and M. Nitta, Dispersive one-dimensional Majorana modes with emergent supersymmetry in one-dimensional proximitized superconductors via spatially modulated potentials and magnetic fields, Phys. Rev. B 105, 214525 (2022).
- [18] T. H. Hsieh, G. B. Halász, and T. Grover, All Majorana models with translation symmetry are supersymmetric, Phys. Rev. Lett. 117, 166802 (2016).
- [19] Z. Huang, S. Shimasaki, and M. Nitta, Supersymmetry in closed chains of coupled Majorana modes, Phys. Rev. B 96, 220504 (2017).
- [20] T. Grover, D. N. Sheng, and A. Vishwanath, Emergent space-time supersymmetry at the boundary of a topological phase, Science 344, 280 (2014).
- [21] A. Rahmani, X. Zhu, M. Franz, and I. Affleck, Emergent supersymmetry from strongly interacting Majorana zero modes, Phys. Rev. Lett. 115, 166401 (2015).
- [22] A. Rahmani and M. Franz, Interacting Majorana fermions, Rep. Prog. Phys. 82, 084501 (2019).
- [23] A. Rahmani, X. Zhu, M. Franz, and I. Affleck, Phase diagram of the interacting Majorana chain model, Phys. Rev. B 92, 235123 (2015).
- [24] K. Roychowdhury, J. Attig, S. Trebst, and M. J. Lawler, Supersymmetry on the lattice: Geometry, topology, and spin liquids, arXiv:2207.09475 [cond-mat.str-e] (2022).
- [25] P.-W. Lo, C.-M. Jian, and M. J. Lawler, Topology shared between classical metamaterials and interacting superconductors, arXiv:2207.10045 [cond-mat.str-e] (2022).
- [26] T. Zhou, N. Mohanta, J. E. Han, A. Matos-Abiague, and I. Žutić, Tunable magnetic textures in spin valves: From spintronics to Majorana bound states, Phys. Rev. B 99, 134505 (2019).
- [27] G. L. Fatin, A. Matos-Abiague, B. Scharf, and I. Žutić, Wireless Majorana bound states: From magnetic tunability to braiding, Phys. Rev. Lett. 117, 077002 (2016).
- [28] A. Matos-Abiague, J. Shabani, A. D. Kent, G. L. Fatin, B. Scharf, and I. Žutić, Tunable magnetic textures: From Majorana bound states to braiding, Solid State Commun. 262, 1 (2017).
- [29] M. Scheurer and A. Shnirman, Nonadiabatic processes in Majorana qubit systems, Phys. Rev. B 88, 064515 (2013).
- [30] B. Bauer, T. Karzig, R. V. Mishmash, A. E. Antipov, and J. Alicea, Dynamics of Majorana-based qubits operated with an array of tunable gates, SciPost Phys. 5, 4 (2018).
- [31] S. S. Hegde, G. Yue, Y. Wang, E. Huemiller, D. J. Van Harlingen, and S. Vishveshwara, A topological Josephson junction platform for creating, manipulating, and braiding Majorana bound states, Ann. Phys. 423, 168326 (2020).
- [32] For a list of non-negative numbers $0 \le x_1 \le x_2 \le \ldots \le x_n$, the geometric average is $\sqrt[n]{x_1 x_2 \cdots x_n} = e^{(\ln x_1 + \ln x_2 + \cdots + \ln x_n)/n} \ge e^{(\ln x_1 + \ln x_1 + \cdots + \ln x_1)/n} = x_1$.