The LSST era of supermassive black holes accretion-disk reverberation mapping

ANDJELKA B. KOVAČEVIĆ D,¹ VIKTOR RADOVIĆ D,^{1,2} DRAGANA ILIĆ D,^{1,3} LUKA Č. POPOVIĆ D,^{1,4} ROBERTO J. ASSEF D,⁵ PAULA SÁNCHEZ-SÁEZ D,^{6,7} ROBERT NIKUTTA D,⁸ CLAUDIA M. RAITERI D,⁹ ILSANG YOON D,¹⁰ YASAMAN HOMAYOUNI D,¹¹ YAN-RONG LI D,¹² NEVEN CAPLAR D,¹³ BOZENA CZERNY D,¹⁴ SWAYAMTRUPTA PANDA D,^{14, 15, 16} CLAUDIO RICCI,^{17, 18} ISIDORA JANKOV,¹ HERMINE LANDT,¹⁹ CHRISTIAN WOLF D,^{20, 21} JELENA KOVAČEVIĆ-DOJČINOVIĆ,⁴ MAŠA LAKIĆEVIĆ D,⁴ ĐORĐE V. SAVIĆ D,^{22, 4} OLIVER VINCE,⁴ SAŠA SIMIĆ D,²³ IVA ČVOROVIĆ-HAJDINJAK,¹ AND SLADJANA MARČETA-MANDIĆ⁴

¹University of Belgrade-Faculty of Mathematics, Department of astronomy, Studentski trg 16 Belgrade, Serbia

²PIFI Research Fellow, Key Laboratory for Particle Astrophysics, Institute of High Energy Physics, Chinese Academy of Sciences, 19B

Yuquan Road, 100049 Beijing, China

³Hamburger Sternwarte, Universitat Hamburg, Gojenbergsweg 112, 21029 Hamburg, Germany

⁴Astronomical Observatory, Volgina 7, 11000 Belgrade, Serbia

 5N úcleo de Astronomía de la Facultad de Ingeniería y Ciencias

Universidad Diego Portales, Av. Ejército Libertador 441, Santiago, Chile

⁶Directorate for Science, European Southern Observatory, Karl-Schwarzschild-Strasse 2, Garching bei München, Germany

⁷ Millennium Institute of Astrophysics (MAS), Nuncio Monseñor Sótero Sanz 100, Providencia, Santiago, Chile

⁸NSF's NOIRLab, 950 N. Cherry Ave., Tucson, AZ 85719, USA

⁹INAF-Osservatorio Astrofisico di Torino, Via Osservatorio 20, I-10025 Pino Torinese, Italy

¹⁰ The National Radio Astronomy Observatory, 520 Edgemont Road Charlottesville, VA 22903, USA

¹¹ The Pennsylvania State University, 201 Old Main, University Park, PA 16802, USA

¹²Key Laboratory for Particle Astrophysics, Institute of High Energy Physics, Chinese Academy of Sciences, 19B Yuquan Road, Beijing

100049, China

¹³Astrophysical Department, Princeton University, 4 Ivy Lane, Princeton, NJ, USA

¹⁴Center for Theoretical Physics, Polish Academy of Sciences, Al.Lotnikow 32/46, 02-668 Warsaw, Poland

¹⁵Nicolaus Copernicus Astronomical Center, Polish Academy of Sciences, ul. Bartycka 18, 00-716 Warsaw, Poland

¹⁶Laboratório Nacional de Astrofísica - MCTIC, R. dos Estados Unidos, 154 - Nações, Itajubá - MG, 37504-364, Brazil

¹⁷Núcleo de Astronomía de la Facultad de Ingeniería, Universidad Diego Portales, Av. Ejército Libertador 441, Santiago, Chile

¹⁸Kavli Institute for Astronomy and Astrophysics, Peking University, Beijing 100871, China

¹⁹Centre for Extragalactic Astronomy, Department of Physics, Durham University, South Road, Durham DH1 3LE, UK

²⁰Research School of Astronomy and Astrophysics, Australian National University, Weston Creek ACT 2611, Australia

²¹Centre for Gravitational Astrophysics, Australian National University, Canberra ACT 2600, Australia

²²Institut d'Astrophysique et de Géophysique, Université de Liège, Allée du 6 Août 19c, 4000 Liège, Belgium

²³ Faculty of Science, University of Kragujevac, Radoja Domanovića 12, 34000 Kragujevac, Serbia

ABSTRACT

The Vera C. Rubin Observatory's Legacy Survey of Space and Time (LSST) will detect an unprecedentedly large sample of actively accreting supermassive black holes with typical accretion disk (AD) sizes of a few light days. This brings us to face challenges in the reverberation mapping (RM) measurement of AD sizes in active galactic nuclei (AGNs) using interband continuum delays. We examine the effect of LSST cadence strategies on AD RM using our metric AGN_TimeLagMetric. It accounts for redshift, cadence, the magnitude limit, and magnitude corrections for dust extinction. Running our metric on different LSST cadence strategies, we produce an atlas of the performance estimations for LSST photometric RM measurements. We provide an upper limit on the estimated number of quasars for which the AD time lag can be computed within 0 < z < 7 using the features of our metric. We forecast that the total counts of such objects will increase as the mean sampling rate of the survey decreases. The AD time lag measurements are expected for > 1000 sources in each Deep Drilling field (DDF, (10 deg²)) in any filter, with the redshift distribution of these sources peaking at $z \approx 1$. We find the LSST observation strategies with a good cadence (≤ 5 days) and a long cumulative season (~ 9 yr), as proposed for LSST DDF, are favored for the AD size measurement. We create synthetic LSST light curves for the most suitable DDF cadences and determine RM time lags to demonstrate the impact of the best cadences based on the proposed metric.

Keywords: Quasars(1319) — Accretion(14) —Time domain astronomy(2109)— Reverberation mapping(2019)

1. INTRODUCTION

The first Event Horizon Telescope (EHT, Event Horizon Telescope Collaboration et al. 2019) observations of M87 revealed a prominent ring of ~ 40μ as in diameter, compatible with a lensed photon ring encircling the shadow of a supermassive black hole (SMBH). This iconic image, in fact, portrays the accretion disk (AD) surrounding the SMBH. As a result, we are now closer than ever to constructing a definitive model of accretion disks. Inflowing gas quickly accretes from a few parsecs down to a few Schwarzschild radii of the SMBH forming a ring-like structure (the AD) that characterizes the phenomenon of an active galactic nucleus (AGN). There is only a short (~2 Myr) active phase¹ for these objects, followed by prolonged periods of inactivity (see Anglés-Alcázar et al. 2021) ². Besides theoretical investigations, a variety of activity cycles have been observed in radio and optical active galactic nuclei (AGNs). The longest timescales of $10^7 - 10^8$ years have been traced using radio wavelengths, and short timescales of $10^4 - 10^5$ years (and down to years) have been estimated using radio wavelengths (~ 10^6 yr, Biava et al. 2021), radio and optical wavelengths (central engine already quiescent, but the remnant of the past activity is on scales of ~ 10^4 yr, Ichikawa et al. 2019; Morganti 2017, review of multiple cycles), optical wavelengths (e.g., changing look AGN where the continuum changes over a time-scale of years, Kollatschny & Fricke 1985; Ilić et al. 2020; Oknyansky et al. 2019; Panda 2022).

The most luminous AGNs, known as quasars (Matthews & Sandage 1963; Schmidt 1963), are thought to be SMBH accreting matter at high rates through accretion disks that are both geometrically thin and optically thick (Sun & Malkan 1989). In contrast, most AGNs in the local Universe³, such as the one in M87, correspond to SMBHs fed by hot plasma at significantly lower accretion rates than quasars (Ichimaru 1977; Blandford & Begelman 1999; Yuan & Narayan 2014).

We already have a typical several-component model of AGN that includes the AD, broad line region (BLR), corona, molecular torus, narrow line region (NLR), jets, and captures the observed features of spectral energy distributions for AGN. Nevertheless, the precise geometry and behavior of these components are mostly unknown (for a review, see Padovani et al. 2017). Even though the entirety of the large scale structure has been identified, the features of the central engine are unclear due to its small dimensions, even for the nearest AGNs. As a result, additional information must be gathered indirectly, through the comparison of theoretical model predictions and observations. Recent developments in instrumentation have allowed the BLR to be spatially resolved in the case of nearby active galaxies (Gravity Collaboration et al. 2018; Event Horizon Telescope Collaboration et al. 2019). However, at least in the near future, it will be difficult to resolve a significant sample of more distant AGNs. Fortunately, the SMBH AD may be examined using either microlensing studies of gravitationally lensed quasars to determine the accretion disk's half light radii at different wavelengths (Morgan et al. 2012; MacLeod et al. 2015), or reverberation mapping (RM) investigations of the SMBH's immediate surroundings (Blandford & McKee 1982). RM offers information on the structure and kinematics of the BLR and accretion disk (see early works of Cherepashchuk & Lyutyi 1973; Gaskell & Sparke 1986) in the time domain. According to the RM premises, the main driver of fluctuations in the BLR lines is accretion disk emission variability. Observing the time delays between disk and broad line emission can be used to calculate the BLR dimension which in turn allows to estimate the SMBH mass. More than 100 robust dynamical black hole mass estimates have been published in the literature, slowly building up to a statistically significant sample (Du et al. 2015; Kaspi et al. 2000; Thater et al. 2019; Panda et al. 2019; Peterson et al. 2004).

The Shakura & Sunyaev (1973) model was the first to provide a more comprehensive picture of an optically thick, geometrically thin accretion disk, albeit in the context of stellar-mass black holes. Soon, Galeev et al. (1979) proposed magnetic reconnection in a corona above a Shakura & Sunyaev (1973) disk to explain the hard X-ray emission observed

¹ For luminous quasars, the activity period can be of order 10 Myrs (e.g., Hopkins et al. 2005).

² Continuous accretion is proposed to cope with the demands of growing black holes in the early universe (Zubovas & King 2021).

³ We denote as local Universe the regions within 300 Mpc (z < 0.1) from Earth (Tekhanovich & Baryshev 2016).

in AGN. The central corona can directly illuminate and heat the outer AGN disk (e.g., Frank et al. 2002), resulting in the "lamppost/reprocessing" paradigm⁴. According to this hypothesis, the fluctuating X-ray energy from the corona can be reprocessed and detected in the UV/optical light from the disk.

Also, Sahu et al. (2019a) and Sahu et al. (2019b) proposed a relation between SMBH mass and central velocity dispersion that is dependent on morphological type, capable of more precisely estimating SMBH masses in other galaxies, and has implications for calibrating virial f-factors. Similarly to the case of the BLR, the light travel time across the AD can be inferred through RM time delays (see e.g., Collier et al. 1998). In particular, in classic AD theory, effective temperature is coupled to black hole mass and accretion rate (Cackett et al. 2007; Liu et al. 2008; Morgan et al. 2010; Fausnaugh et al. 2016; Panda et al. 2018). When compared to the outer and cooler AD regions, the reprocessing model demonstrates that radiation from the innermost part of the accretion disk, closer to the SMBH, has a peak of emission at shorter wavelengths, and its variability is detected (Cackett et al. 2007; Chelouche 2013; Edelson et al. 2015) with a mean delay scaling $\tau \sim \lambda^{4/3}$ (Collier et al. 1998; Starkey et al. 2016). An alternate AGN variability model proposed by Dexter & Agol (2011) assumes large local temperature fluctuations in the AD as the origin of flux variability. While it can explain the amplitude and time scales of observed variability (MacLeod et al. 2010), it cannot explain the wavelength dependence of the time lags (Starkey et al. 2016). Yu et al. (2022) model AGN variability as a Damped Harmonic Oscillator and offer more insight; the modeled short-term variability behaves differently in the vicinity of 2500A, which could be an imprint of a different accretion flow geometry rather than a homogeneous thin disk.

While thin ADs are the most robust explanation for AGN luminosity and variability, multiple studies have found that the AD sizes depart from the immediate prediction of the model. Estimated accretion disk sizes from RM campaigns are three times larger than the expected based on thin-disk theory (Sergeev et al. 2005; Shappee et al. 2014; Edelson et al. 2015; Fausnaugh et al. 2016; Jiang et al. 2017; Cackett et al. 2018; Mudd et al. 2018). Noticeably, microlensing studies of luminous lensed quasars have independently obtained comparable conclusions (Pooley et al. 2006, 2007; Morgan et al. 2010; Mosquera et al. 2013; Chartas et al. 2016).

Using data from the Neil Gehrels Swift Observatory (see Edelson et al. 2019), Shappee et al. (2014) and McHardy et al. (2014) found that the UV leads the optical emission in NGC 2617 and NGC 5548, respectively. Following these campaigns, Edelson et al. (2017) and Edelson et al. (2019) observed four AGNs (NGC 5548, NGC 4151, NGC 4593, and Mrk 509) and found significantly correlated UV/optical fluctuations, and the same overall pattern of interband delays growing from UV to optical wavelengths. While these lags are two times larger than expected from classical theory, the uncertainties on the predicted lags are large, so it is unclear whether this is inconsistent with the usual thin disk picture. Using Pan-STARRS light curves, Jiang et al. (2017) measured the continuum lags of ~ 200 quasars (the mean lags between the g band and the r, i, and z bands for the whole sample are 1.1, 2.1, and 3.0 days), whereas Mudd et al. (2018) used the Dark Energy Survey's (DES) photometric data to measure the disk sizes of 15 quasars (e.g. lags \leq 3.0 days in r,i band, and \leq 5.0 days in z band relative to g-band). Homayouni et al. (2019) recently measured the continuum lags of 95 quasars using photometric data from the Sloan Digital Sky Survey (SDSS) RM project. They found that average size of measured disks is consistent with the Shakura & Sunyaev (1973) analytic thin-disk model. However, it was shown that there is a wide variety of disk sizes, both small and large, that are in excess of the measurement uncertainties.

Yu et al. (2020) assessed the time lags and accretion disk sizes of 22 quasars with SMBH masses ranging from 10^7 to $10^9 M_{\odot}$ with the data from from the Dark Energy Survey (DES) standard-star fields and the supernova C fields. They concluded that when the disk variability is taken into account, most of the reported accretion disk diameters are compatible with the predictions of the classic thin-disk model. Guo et al. (2022) compiled a sample of 92 AGNs at z < 0.75 with gri photometric light curves from the archival data of the Zwicky Transient Facility and found that AD sizes are systematically larger than expected, although the ratio of the measured to theoretical disk sizes depend on using the emissivity – or responsivity – weighted disk radius.

According to a recent study conducted by Kara et al. (2021), short CIV and Ly α delays in the spectra of Mrk 817 show that the accretion disk extends beyond the UV broad-line region. Also, Netzer (2022) suggested that the large bulk of observed continuum delays are due to varying diffuse BLR emission rather than from disk emission (see also Korista & Goad 2001; Lawther et al. 2018).

⁴ Electrons heated by a magnetic field transfer their energy to optical/UV photons and upscatter them into the X-band

We are about to enter an era in which an abundance of new data provides the opportunity to settle the disk size dispute and unravel diverse effects. The Vera C. Rubin Observatory opens a window of discovery in our dynamic universe through its Legacy Survey of Space and Time (LSST). The LSST project is intended to investigate dark energy and dark matter, the solar system inventory, the transient optical sky, and to map the Milky Way (see Ivezić et al. 2019). The distribution of revisit times (i.e., the cadence of observations) is among the main constraints on data and system properties placed by the four main science themes (see Ivezić et al. 2019). The LSST community has conducted a thorough investigation of prospective cadences (Connolly et al. 2014). Metrics developed by the LSST community provide a scientific testbed for evaluating a suite of cadence simulations (see details in Bianco et al. 2022) employing the Metrics Analysis Framework (MAF Jones et al. 2014, 2020) and The Operations Simulator (OpSim Reuter et al. 2016). Both codes are open source. The OpSim generates databases containing 10-year pointing histories, complete with weather, seeing, and sky brightness information. It is more often now known as the 'Feature Based Scheduler' or FBS, and its fifth version is used for this analysis. The MAF offers an application programming interface (API) for inferring image properties (e.g., seeing, proper motion) over various spatial scales, as well as tools for predicting the properties of photometric measurements for objects and light curves inserted in survey simulations (see details in Bianco et al. 2022). LSST is expected to tightly constrain the radial dependence of the effective temperature in accretion disks through its u,q,r,i,z,y light curves with daily cadences of a large sample of AGNs (see details in AGN SC Roadmap Shemmer et al. 2018). Assuming a 5σ depth of 24.6 mag in the r-band, De Cicco et al. (2021) estimated a sky density of 6.2×10^6 AGNs for the LSST main survey of $\sim 18000 \text{deg}^2$. Specific metrics on the number of quasars that will be detected in the 10-year co-added *i*-band observations as a function of OpSim run have been developed by Assef et al. (2021a).

The LSST AGN and Transient and Variable Stars (TVS) Scientific Collaborations (SCs) devised a variety of metrics⁵ to quantify the impact of LSST observing strategies on different variability and photometric redshift measurements as well as quasar counts. The Blazar Variability metric by Raiteri et al. (2022) evaluates the performance of LSST observing strategies for extracting information from data linked to the shortest time and spatial scales of blazars (less than 1 hr to years). Related to largest scales, Assef et al. (2021b,a) have developed two metrics to evaluate the 10-year co-added depths expected for each band in the context of photometric redshifts for type-1 quasars. Yu & Richards (2021) have created a model-independent metric that quantifies the degree to which we can deduce the structure function of variable sources (e.g., AGNs) observed by LSST). The metric described in the following sections complements this suite by quantifying the impact of LSST observing strategies on the AD and BLR photometric reverberation mapping, covering time scales from a few up to 400 light days. In addition, as LSST will be able to measure AGN dust sublimation radii which recovery critically depends on the cadence as well as the continuity of observations (see Hönig 2014), our metric could be applied on the corresponding time scales of several tens of days. The caveat, however, is that if most accretion discs dominate the AGN emission in the NIR (see e.g., Landt et al. 2011), the dust sublimation radius reverberation signal will be difficult to isolate.

Kovačević et al. (2021b) developed a metric to assess the impact of different LSST cadences on the detection of periodicity and time lags for AGN. While modeling specific object classes is useful for assessing the science implications, they may not cover the entire discovery space. When paired with survey design, metrics that provide quantized estimates rather than continuous probability distributions can be useful for survey design optimization (Bellm et al. 2022).

Based on the AGN SC roadmap premises as well as a need for metrics, the goal of this work is to evaluate different LSST cadences for the science case of measuring AGN interband time delays, specifically in the Deep Drilling Fields (DDFs), and to identify suboptimal cadences. The LSST Deep Drilling Fields will have the deepest co-add depths of the survey (see Brandt et al. 2018), and are expected to exceed those achieved by 8m class observatories in the same areas (see Aihara et al. 2022). Roughly 10-20% of the LSST observing time will be used in mini-surveys and in the six DDFs. Through a combination of depth and cadence, the DDFs are expected to provide the most complete AGN catalogs and their host galaxies (Brandt et al. 2018; Zou et al. 2022).

We used our metric (Kovačević et al. 2021b) simulation in the LSST MAF to test various cadence strategies. For the best strategies revealed by our metric MAF simulations, we simulated AGN light curves using a damped random walk (DRW) model, as the simplest Gaussian random process model that can describe the optical light curves of quasars

⁵ The metrics have been presented at the LSST Project and Community Workshop 2021 in Survey Strategy III: Domain-Specific Cadence Optimization Discussions project.lsst.org/meetings/rubin2021/content/survey-strategy-iii-domain-specific-cadence-optimization-discussions. Joint session of AGN, TVS microlensing, and Strong Lensing scientific collaborations titles Together we stand! was held during The second SCOC science collaboration workshop https://project.lsst.org/meetings/scoc-sc-workshop2/agenda.

more-or-less reasonably well (DRW, Kelly et al. 2009; Kozłowski et al. 2010; Zu et al. 2013). However, the following limitation of the DRW model should be kept in mind: as Kozłowski (2017) identified, it is important that the time baseline of the light curve is considerably longer than the damping timescale. Also, Stone et al. (2022) imply that even a 20-year baseline is unlikely to be long enough to constrain the damping timescale in some of the quasars in their sample. This could be due to long-term trends in the light curve, or that these processes are more complex than a basic DRW with only one characteristic timescale (see Stone et al. 2022). We also used the AGN lampost⁶ AD model (Cackett et al. 2007) to account for wavelength dependence and replicate delayed LSST multiband light curves. Following that, we used OpSim 1.7-2.0 to produce data points for the delayed multiband light curves based on observational sequences of different cadences. We used a shifting method inspired by PyCS (Tewes et al. 2013) but with Partial Curve Mapping (PCM, Jekel et al. 2019) to determine the similarity of shifted curves in order to calculate the time delay from the simulated observations.

The following is a general outline of the structure of the article. Section 2 answers the question of what lags are expected for different black hole masses. Section 3 describes the assumed accretion disk model, and how the synthetic light curves, which are used to probe the optimal cadences indicated by our metric, are constructed. Section 4 describes two tiers of estimators of cadence strategies. The first tier shows our metric run, which quantifies the optimal LSST cadences for AD time lag measurements using the LSST MAF. In the second, LSST multiband light curves are created under the assumption of generic quasar parameters and optimal LSST cadences, and their time delays are assessed with respect to the u-band. Section 5 presents the results, which are interpreted and discussed in Section 6 in more detail. Section 7 concludes the article and presents further implications of this work.

2. DISTRIBUTION IN TIME LAG OF LSST OBSERVABLE AGNS

For reverberation mapping, it is not only interesting to know the efficiency of the survey observing strategies, but also what lags are expected for different black hole masses. For this purpose, we use the quasar luminosity function (QLF, Hopkins et al. 2007) to derive a probability density of AGNs per measured time lag. The probability for measuring the time lags $p_{\tau} = p(M, L, z)$ is estimated from a uniform sample of SMBH masses $M \in (10^6, 10^{10}) M_{\odot}$, whose luminosities are inferred according to procedure described in Kovačević et al. (2021b) and use the standard thin-disk model to calculate the expected mean time lags corresponding to LSST filters. The probability p_{τ} in our approach equals to our metric (containing expected mean time lags with respect to u band) divided by total number of measurements. The distribution in time lag (τ) of LSST observable AGNs is given by :

$$\frac{dN}{d\tau} = 4\pi \frac{d^2 V}{d\tau dz} \int_{\log L(z)}^{+\infty} \frac{d^2 \mathcal{N}}{d\log L dV} p_\tau d\log L dz \tag{1}$$

where $\frac{d^2 N}{d \log L dV}$ is a double-power-law QLF with redshift-dependent slopes from Shen et al. (2020, see their Table 4, second column) and $\frac{dV}{dz}$ is the co-moving volume per redshift (calculated with CosmoloPy ⁷).

Based on this, we summarize the expected distribution in time lag of LSST-observable AGNs in Figure 1 (left). Besides this general distribution, as a forecast for the distribution of the observed time lags in LSST DDFs, we built a sample of 208 broad-line AGNs located within the COSMOS field, with estimated optical spectral parameters from the SDSS DR7 catalogs of Liu et al. (2019) and Shen et al. (2011). If an object is identified in both catalogues, we used the data from Liu et al. (2019) as it provides the most up-to-date measurements of the AGN spectral parameters. The resulting sample covers a large range of redshifts (z = 0.02 - 3.82) and black hole masses ($\log(M/M_{\odot}) = 6.6 - 10.3$). As in the case of an ideal uniform set of SMBH masses, we use the standard thin-disk model to calculate the expected time lags in LSST *ugriz* bands. The right panel of Figure 1 shows the probability density function (PDF) of the predicted time lag. Both distributions show that a majority of AGNs will have lags less than 5 days. We can expect that higher observational cadences, such as multi_short, may increase the rate of detected time lags. As the LSST will provide higher-resolution images and a multi-visit-multi band data sets, our estimate is most likely a lower threshold.

3. SYNTHETIC LIGHT CURVES FOR PROBING LSST OPTIMAL CADENCES

Here, we outline the calculation of synthetic time-delayed AD light curves using a combination of a theoretical AD transfer function (Section 3.1) and a damped random walk (DRW, Section 3.2).

⁶ The lamppost is typically considered a point-like source hovering over the black hole spin axis and associated with the X-ray corona.

⁷ http://roban.github.com/CosmoloPy/



Figure 1. Left: Distribution of quasars per expected time lag with respect to *u*-band filter based on ideal set of uniform distributed masses in the range $10^6 - 10^{10} M_{\odot}$. Right: PDF of expected time lag with respect to *u*-band for the 208 quasars sample from Liu et al. (2019) and Shen et al. (2011) catalogues located in COSMOS.



Figure 2. Flat accretion disk time delay distributions $(\tau(r, \theta, i) = r(1 + \cos\theta \sin i)/c)$, as seen by distant observer, for various inclinations *i* given on top of each plot. The theoretical observer position is to the bottom at infinity. The colorbar (in units of days) indicates time lag values. Distributions are deformed from circular to elliptical and becomes parabolic for the extreme inclination $i = \pi/2$. The observer will witness the re-emitted light arriving with delays ranging from 0 to $2r(1 + \cos\theta \sin i)/c$, with a bulk average delay of $r(1 + \cos\theta \sin i)/c$ for radius *r* and azimuth θ of the reprocessing site, with $r \in [0, 10]$ light days. Our disk model reiterates the concept of isodelay surfaces as described in Peterson (2006); Starkey et al. (2016).

3.1. Accretion disk model and expected time lag

We assume a non-relativistic thin-disk model (Shakura & Sunyaev 1973) that emits black-body radiation and has a compact variable source (a "lamppost"), whose photons hit the disk and cause a local temperature increase; full details can be found in the literature (Cackett et al. 2007; Pozo Nuñez et al. 2015; Starkey et al. 2016, 2017). Here, we summarize the key points. We assume an optically thick and geometrically thin accretion disk. The total observed energy flux Ξ from the accretion disk is a sum of the energy flux radiated due to a viscous heating process Ξ_V from a surface unit and irradiated flux Ξ_I of the disk photo sphere, $\Xi = \Xi_V + \Xi_I$ (Pozo Nuñez et al. 2019).

If R is the radial distance from the disk's innermost radius, $R_0 = \frac{6GM}{c^2}$ for a non-spinning SMBH, G is the gravitational constant, M is the SMBH's mass, and \dot{M} is the mass accretion rate, then

$$\Xi_V = \frac{3GM\dot{M}}{8\pi R^3} \Big(1 - \sqrt{\frac{R_0}{R}}\Big).$$

Because the geometry of the emitting source is unknown in the lamppost model, a crude approximation is established by setting the source of luminosity $L^{* 8}$ at a height h^{*} along the SMBH's rotational axis. Then the irradiative flux might be calculated using the disk's albedo A as follows (Pozo Nuñez et al. 2019):

$$\Xi_I = \frac{L^\star (1-A)}{4\pi R^3} h^\star \cos\theta$$

⁸ Vasudevan et al. (2009) estimated that about 5% of total flux comes from X-ray source.

so that θ is the incident angle between the disk surface normal and radiation of the emitting source (Netzer 2013). For viscous heating alone, when $R >> R_0$ the disk temperature profile should follow

$$T(R) = \left(\frac{3GM\dot{M}}{8\pi R^3\sigma_{\rm SB}}\right)^{1/4}$$

where σ_{SB} is the Stefan–Boltzmann constant. Similarly, in the case of irradiation of the disk from a height h^* above the disk, the total temperature profile is a superposition of the temperature due to irradiation and the temperature due to viscous heating

$$T^{4}(R) = \left(\frac{3GM\dot{M}}{8\pi R^{3}\sigma_{\rm SB}}\right) + \frac{L^{\star}(1-A)}{4\pi R^{3}\sigma_{\rm SB}}h^{\star}\cos\theta$$

That is,

$$T(R) \propto \left(\frac{L^{\star}h^{\star}}{R^3}\right)^{1/4}$$

for $R >> h^*$ and the viscous heating flux is much smaller than the irradiated flux (Pozo Nuñez et al. 2019). Thus, in both cases, we expect that the temperature profile follows the relation $T(R) \propto R^{-3/4}$ (Cackett et al. 2007). We assume that reprocessing has a minor effect on deviations from the viscous temperature and the $T(R) \propto R^{-3/4}$ temperature structure (see Collier et al. 1998; Starkey et al. 2016):

$$T(R) = \left(\frac{3GM\dot{M}}{8\pi\sigma_{\rm SB}R_{\lambda}^{3}}\right)^{1/4} \left(\frac{R_{\lambda}}{R}\right)^{3/4} \sim \left(\frac{3GM^{2}L}{8\pi\eta c^{2}L_{\rm Edd}\sigma_{\rm SB}R_{\lambda}^{3}}\right)^{\frac{1}{4}} \left(\frac{R_{\lambda}}{R}\right)^{3/4}.$$
(2)

Here, L_{Edd} is the Eddington luminosity $(L_{Edd} \approx 1.3 \cdot 10^{38} \frac{M}{M_{\odot}} \text{ergs}^{-1})$, η is the accretion efficiency. If we are observing at rest wavelength λ^{rest} , the radius R_{λ} is usually approximated as the radius where the disk temperature matches the photon wavelength

$$kT = \frac{hc}{\lambda^{rest}};$$

here, h and k are the Planck and Boltzmann constants, respectively. Then the radius corresponding to the photon wavelength of LSST filters is

$$R_{\lambda} \sim \left(\frac{k\lambda}{hc}\right)^{\frac{4}{3}} \left(\frac{3GM\dot{M}}{8\pi\sigma_{\rm SB}}\right)^{1/3} \sim \left(\frac{k\lambda}{hc}\right)^{\frac{4}{3}} \left(\frac{3GM^2L}{8\pi\eta c^2 L_{\rm Edd}\sigma_{\rm SB}}\right)^{\frac{1}{3}}.$$
(3)

The disk transfer function $\psi(\tau, \lambda)$ describes the response of the disk to changes at a given wavelength and time delay (Cackett et al. 2007):

$$\psi(\tau,\lambda) = \int_{R_{in}}^{R_{out}} \int_0^{2\pi} \frac{\partial B}{\partial T} \frac{\partial T}{\partial L} \frac{R dR d\theta \cos i}{D^2} \delta(\tau - \frac{R}{c} (1 + \sin i \cos \theta))$$
(4)

where L is the driving luminosity, *i* is the inclination angle of the disk, θ is the azimuthal angle of emission patch, D is the distance of the object, and (R_{in}, R_{out}) are inner and outer boundary of integration (see Cackett et al. 2007, and references therein). Also, we assume the accretion disk flux arises from the blackbody radiation described by Planck's law

$$B(T) = \frac{2hc}{\lambda^5} \frac{1}{e^{\frac{hc}{\lambda kT} - 1}}$$

with the temperature profile given by Eq. 2.

If we assume the temperature variations to be small, then the derivative is

$$\left|\frac{\partial B}{\partial T}\frac{\partial T}{\partial L}\right|\sim \frac{q\cdot x^{-1/4}e^{qx^{3/4}}}{4(e^{qx^{3/4}}-1)^2},$$

where $x = \frac{R}{R_{\lambda}}$, $q = hc/\lambda k$.

For our blackbody disk model, the time-delay centroid $\langle \tau \rangle$ of the transfer function is⁹

$$\langle \tau \rangle = \frac{\int \psi(\tau, \lambda) \tau d\tau}{\int \psi(\tau, \lambda) d\tau} \sim \frac{\int \psi(\tau, \lambda) \frac{R^2}{c} dR}{\int \psi(\tau, \lambda) R dR}.$$
(5)

Figure 2 illustrates the differentiation of the time lag distributions within the disk with regard to different inclinations. Each region of distribution is labeled with the time delay it represents (so that we would observe relative to the continuum source). This is a well-known notion of isodelay surfaces, as described in Peterson (2006) and Starkey et al. (2016). The line of sight to the observer is oriented towards the bottom of each disk plot, except in the left panel, where it is exactly vertical to the plot plane.

Also, we plot example response functions showing the LSST wavelength dependence in Figure 3 for the following set of parameters: SMBH mass $M = 2 \cdot 10^8 M_{\odot}$, $L/L_{Edd} \sim 0.1$, $\eta = 0.1$, disk inclination of 45° and azimuth angle of 0°. As shown in Figure 3, the mean delay, indicated by the vertical lines, and the width of the response functions increase with wavelength. These effects occur due to the inverse scaling between temperature and radius (Starkey et al. 2016).

In addition, a larger inclination of the disk (see Figure 2) makes the response function more skewed (Starkey et al. 2016). The peaks will move toward shorter lags and develop a tail towards large lags. This occurs because larger inclinations increase the lag on the far side and shorten the lag of the near side of the disk (Starkey et al. 2016) relative to smaller inclination (see Figure 2). A simple sinusoidal with a period of 2.5 days was constructed to demonstrate how its peak positions and amplitudes change while convolving with the time-delay transfer function of a disk with an inclination of $\pi/4$ and for different LSST filters (see Figure 3).

Figure 4 demonstrates potential effects of a non-relativistic AD setup by comparing the observed luminosity normalized to the accretion rate for classical and Kerr-metric black-body disk emission as given by the Campitiello et al. (2018, see their Figure 2) as a function of SMBH spin a and viewing angle θ . It is easy to notice that for a < 0.8, the relativistic angular pattern of radiation is very similar to the classical behavior. The classical model, when observed at $\theta = 0$ is well approximated by the relativistic model with $a \sim 0.8$, but at larger viewing angles, the relativistic model is brighter because of the strong relativistic effects. When spin $a \rightarrow 1$, the radiation pattern is very steep, even for the largest viewing angles. This is due to the combination of different relativistic effects (Doppler beaming, gravitational redshift, and light-bending) along with the black hole spin: the trajectories of the energetic photons coming from the innermost region of the disk (close to the event horizon) are bent in all directions and the intensity of radiation is almost the same at all viewing angles.

To compare time lags in curved and flat spacetime, we calculate their ratio. Time lags in curved space are obtained by combining analytical approximation of time delay for Kerr black hole in flat spacetime (e.g., Campana & Stella 1995; Mastroserio et al. 2018) and approximation for light bending (Poutanen & Beloborodov 2006):

$$c\Delta t = ry \cdot \left(1 + \frac{uy}{8} + \frac{uy^2}{8}(\frac{1}{3} - \frac{u}{14})\right)$$
(6)

where r and θ are radius and azimuth of reprocessing site, $u = \frac{2GM}{c^2r}$, and $y = 1 + \cos(\theta)\sin(i)$ for a given inclination i. Note that $c\Delta t = ry$ corresponds to the time delay one would expect in flat space time (see Figure 4).

Figure 5 depicts a schematic view of the 2D distribution of the ratio between the relativistic time delay (Equation 6) and its classical equivalent. The ratio values are sharply increasing with decrease of distance to the center of mass. This is due to light bending which is largest there. For a non-rotating black hole $(a \rightarrow 0)$, the marginally stable radius is $r_{\rm ms} = 6 \frac{GM}{c^2} = 6r_{\rm g}$, whilst for a maximally rotating black hole $(a \rightarrow 1)$ it takes a value of $r_{\rm ms} = 1 \frac{GM}{c^2} = r_{\rm g}$. As theoretical predictions state that the maximal spin for a rotating black hole can not exceed 0.998, then the marginally stable radius of fast rotating black hole is $r_{ms} = 1.24r_{\rm g}$ (Thorne 1974). Figure 5 corresponds to fast rotating $a \rightarrow 1$ SMBH, so that the small white inner circle with radius of the marginally stable orbit $r_{\rm ms} = 1.24r_{\rm g}$ assumes that no source of light can orbit around SMBH. As for non-rotating $a \rightarrow 0$ SMBH, at $6r_{\rm g}$ and larger distances there will be no effects (dark green and dark blue color). The upper limit of the navy blue region is at $6r_{\rm g}$. Therefore, for a fast-rotating SMBH relativistic corrections should be taken into account, because $r_{\rm ms}$ is closer to the event horizon.

$$\langle \tau \rangle \sim \frac{R_{\lambda}}{c} \left(\frac{\lambda k}{hc}\right)^{4/3} \frac{I(8/3)}{I(4/3)}$$

where I(8/3) = 5.139, I(4/3) = 2.895.

⁹ Taking into account that the integral $I(a) \sim \int \frac{s^a s e^s ds}{(e^s - 1)^2}$ can be evaluated numerically (Cackett et al. 2007), and that the centroid time delay can be approximated as follows:



Figure 3. Top: Numerical accretion disk transfer function for LSST ugrizy filters calculated for the following set of parameters: SMBH mass $M = 2 \cdot 10^8 M_{\odot}$, $L/L_{\rm Edd} \sim 0.1$, $\eta = 0.1$, disk inclination of 45° and azimuth angle of 0°. Dashed vertical lines mark the theoretical centroid delay (Equation 5) for comparison with the peaks for each transfer function (values given in upper right corner). Middle: convolution (in colors) of transfer function and a sinusoid of 2.5 days period (shown in dashed black). Bottom: Synthetic LSST light curves obtained by convolution of driving light curve which flux is scaled (black line) and inclined AD transfer function shown in top panel. Different bands are plotted with the same color coding in all plots.

Even though it looks like an extreme possibility, there are well-known cases like MCG -6-30-15 and NGC 4051, whose



Figure 4. Schematic comparison of emission patterns across viewing angles for different AD models as given by the phenomenological approximation of Campitiello et al. (2018, see their Figure 2). The classical nonrelativistic model (dashed blue line) is compared with relativistic models with a = -1 (dark blue curve), a = 0 (dark purple curve), $a \sim 0.8$ (red curve) and a = 0.998(orange curve). The radial axis is the observed efficiency $\eta_o = \frac{L^o}{Mc^2}$, i.e. given by the observed luminosity normalized on the accretion rate (which is the same for each model). The plot plane is the accretion disk plane. The main axis of all models is in direction of 0° , assuming azimuthal symmetry with respect to this direction. A distant observer is at the left of the figure.

X-ray emission could be explained with light-bending effects (e.g., Dai et al. 2010; Chartas et al. 2012; Kara et al. 2014). Also, the published EHT observations of M87 and Sgr A^{*} are consistent with the Kerr paradigm, indicating that fast spinning black holes might be ubiquitously case (Event Horizon Telescope Collaboration et al. 2019, 2022).

3.2. Simulating driving and multiband light curves

We provide here a summary of our light curve modeling. A full description of synthetic light curve modeling, and of tests using these synthetic light curves, are presented by Kovačević et al. (2021b).

The time variability of AGN and quasar continuum emission can be approximated using a damped random walk (DRW, Kelly et al. 2009; Kozłowski et al. 2010; Zu et al. 2013). A DRW is defined by a stochastic differential equation with a broken power law power spectrum and a slope that steepens from 0 to 2 above a break time-scale. The break time-scale in AGNs measured with the SDSS is up to 100 days. It correlates favorably with black hole mass and wavelength, and is a weak function of brightness (MacLeod et al. 2010; Burke et al. 2021). However, light curves of AGNs observed with Kepler and ground-based surveys show steeper power spectral density (PSD) than the DRW model at timescales up to ~ 1 month (Mushotzky et al. 2011; Smith et al. 2018; Caplar et al. 2017; Sánchez-Sáez et al. Kozłowski (2017) demonstrated that light curves with insufficient time length cannot consistently constrain 2018). the damped variability timescale, and advised that light curves time length should be at least tenfold that of the true damped variability time scale. Based on 1384 variable AGNs from the QUEST-La Silla AGN variability survey, Sánchez-Sáez et al. (2018) showed that the DRW variability amplitude is influenced by the length of the light curve, and that the DRW model can represent the variability of 74% of AGNs in their specified sample. On the other hand, the work of Lu et al. (2019) suggests that only 35% of light curves in their RM AGN sample of 113 objects can be derived with accurate DRW model parameters. This proportion is much lower than that found by Sánchez-Sáez et al. (2018), which could be related to the RM AGN shorter light-curve length (one year) in Lu et al. (2019) than for the sample of Sánchez-Sáez et al. (2018). In this work, the probed damped variability time scales are short, up to 200 days with respect to LSST's 10 year time baseline, so that we can assume that DRW is an approximate working description



Figure 5. Distribution of ratio of the time delay calculated from a relativistic and a classical spacetime approximation, when $a \to 1$. The inclination of the observer is set to $i = \frac{\pi}{3}$, the region extent to $10r_{\rm g}$, and the SMBH mass is $10^8 M_{\odot}$. The dark green color depicts a ratio value of ~ 1.075. The small white circle is a 'plunging region' inside the marginally stable orbit of $r_{\rm ms} = 1.24r_{\rm g}$. The color bar shows the ratio between relativistic and classical time lags. The ratio is highest where the light bending effects are largest, i.e., near the SMBH. As the distance increases, the ratio approaches 1, because spacetime is almost flat.

of quasar variability for optical light curves (Suberlak et al. 2021). Although the damping mechanism is unknown yet, Kelly et al. (2009) show that the damping timescale might be connected to the thermal timescale of the accretion disk, assuming that its thermal fluctuations are responsible for the AGN variability. Because of this, we concentrate our analysis on the DRW model, based on its implementation in Zu et al. (2011, 2013) and Suberlak et al. (2021).

Neustadt & Kochanek (2022) investigated the effect of AD temperature fluctuations on AGN light curves, revealing that the light curves closely reflect the expectations for a lampost model, albeit with slower variability time-scales of the in-going waves. This also means that longer time-scale variability signals will increasingly diverge from conventional models, since the smoothing of slower-moving waves reduces significantly as their period or spatial wavelength grows. LSST will operate much longer than most RM campaigns, providing the first well-characterized multi-wavelength light curves to investigate what happens on long time scales.

We first generate a uniform time grid $\Delta t_i = 1$ day on which to evaluate the driving light curves F_{DRW} . This is convenient for evaluating the convolution integral. As the AD transfer functions asymmetry increases with an increase in the wavelengths of the filters, light curves features are still evident in the *u*-band for the shortest wavelength, but they are blurred out for the *y*-band (see Figure 3 and Section 6). Since the *y*-band light curve distortion is the greatest, determining the time lag requires additional care (due to large possible errors). Thus, for the sake of simplicity, we continue by generating quintets of multi-band light curves based on the LSST *ugriz* bands using a transfer function as described in Section 3.1.

The multiband continuum light curves $(\mathcal{F}(t, \lambda))$ are then obtained from the driver (F_{DRW}) by convolution with the asymmetric response function $\psi_{\lambda}(\tau)$ (see an example in Figure 3):

$$\mathcal{F}(t,\lambda) = \sum_{i=0}^{m} \psi(\tau_i,\lambda) F_{\text{DRW}}(t-\tau_i) \Delta \tau.$$
(7)

We assume the following set of parameters for the transfer function: SMBH mass $M = 2 \cdot 10^8 M_{\odot}$, $L/L_{\rm Edd} \sim 0.1$, $\eta = 0.1$, disk inclination of 45° and azimuth angle of 0°. For the driving source emission we assume a characteristic restframe timescale $\tau_{\rm DRW} \sim 200$ days and a structure function at infinity $\sigma_{\rm DRW} = 2.5$ (flux unit), which describes the

long-term variability of the light curve. These DRW parameters are typical of the quasars in the DES sample (Yu et al. 2020). In general, MacLeod et al. (2010) found that both parameters follow a power law relationship with AGN luminosity, measured wavelength, and black hole mass (see King et al. 2015, for a compact representation of these relations).

A realization of the driving source emission for given parameters is displayed in the bottom panel of Figure 3 in black. Using the transfer functions for LSST bands, we then generated the multiband light curves¹⁰ as shown in the bottom panel of Figure 3. In practice, light curves where the lags are not clearly visible are likely to have delays that are shorter than the cadence of the light curve.

4. ESTIMATOR OF CADENCE STRATEGIES

In this section, we characterize distinct LSST observing strategies. Section 4.1 describes our metric implemented in MAF that quantifies the impact of cadences for LSST RM time lag measurements. Additional calculations of time lag measurement of simulated LSST continuum light curves using the best cadences identified by our metric in two DDFs are outlined in Section 4.2.

4.1. Metric simulation in MAF

The Metrics Analysis Framework (MAF)¹¹ is a software package designed to enable writing code to evaluate the performance of different survey strategies. Our metric is integrated into MAF through the module AGN_TimeLagMetric¹².

The metric includes the quasar selection based on the limiting magnitude of the observation¹³ in the r filter. The magnitudes were corrected for the Galactic dust-extinction. The sky regions with dust extinction E(B-V) > 0.2 are excluded because this amount of dust extinction is problematic for extragalactic science for two reasons: it reduces the effective coadded 5σ depth¹⁴, and the total amount and wavelength dependence of dust extinction are not always well characterized, making it difficult to calibrate the effect on the background galaxies (see Jones et al. 2020).

The depth of LSST single visits is assumed to be similar to that of the COSMOS field observations by the VLT Survey Telescope (De Cicco et al. 2021). De Cicco et al. (2021) found that 91% of AGNs have r > 21 mag, putting them beyond the average depth of most existing variability studies. Thus we impose in our MAF simulations a cut for 5σ depth in the r band > 21.0 mag.¹⁵ This enables us to investigate how well AGN time lag measurements work for fainter objects. Our later analysis reveals that the most suitable DDF cadence strategies in the r-band have mean 5σ depths of 21.5 (see discussion in Section 6, and Figures 11-13). Application of our metric with concurrent limitations in all filters could eliminate the vast majority of visits for time lags up to 5 days due to the Nyquist criterion. Otherwise, for larger time delays (order of tens of days) simultaneous 5σ cuts can be used in more than one band. Thus, we relaxed the limitation in bands other than r in order to have a broader pool of feasible cadences in the other bands. When we compared the mean 5σ depths of r and other bands within the DDFs, we found that they are within the range of 18 and 21.5 mag in DDFs (Figures 11-13). This means that objects as bright as 21 mag will still be seen in those bands for these overlapping cadence strategies.

When analyzing observations of variable objects, two major issues arise. First, because of weather or other technical constraints, sampling times can be perturbed. Second, objects can have a wide range of variability characteristics and behaviors. As a result, it is important to establish a threshold to search for variability on the time scales of interest (Eyer & Bartholdi 1999). By assuming that there is no correlation between flux variation F_{var} (as defined in e.g., O'Brien et al. 1998), the measured time delay τ_{obs} and its uncertainty $\sigma_{\tau_{\text{obs}}}$, Kovačević et al. (2021b) developed the metric ϕ_{τ} :

$$\log \phi_{\tau} = \log \frac{\sigma_{\tau_{\rm obs}}}{\tau_{\rm obs}} \propto A + C_1 \frac{F_{\rm var}}{\sigma} + C_2 \frac{\tau_{\rm obs}}{(1+z)\Delta t^c},\tag{8}$$

where the time sampling (or cadence) of the light curve is Δt^c and the mean photometric error of light curve points is σ . If the observed variation is intrinsic to the quasars at cosmological distances, one would expect a time dilation effect. The time delay is reduced to the rest frame, taking into account the cosmological redshift z in Equation 8 by a factor $(1 + z)^{-1}$.

 $^{^{10}}$ In a work by Jankov et al. (2022), we determined which LSST filters contain continuum at different redshifts (see e.g. their Figure 1)

 $^{^{11} \}rm https://rubin-sim.lsst.io/rs_maf/index.html$

 $^{^{12}\,\}rm https://rubin-sim.lsst.io/rs_maf/metricList.html$

¹³ The 5 σ depth in the OpSim is the limiting magnitude of the visit – meaning, objects fainter than this limit will not be visible. However, objects brighter than that limit are definitely still visible, and at higher SNR than 5.

 $^{^{14} \ {\}rm Unit\ is\ magnitude,\ see\ https://confluence.lsstcorp.org/display/SIM/Summary+Table+Column+Descriptions.}$

¹⁵ Otherwise, adding a cut for 5σ depth in the r band < 21.0 mag means that the (vast majority) of visits with depth which is > 21.0 mag would not be counted, yet could still detect objects which are as bright as 21.0 mag.

As the expected size of the AD in the UV/optical is on the order of ≤ 10 light days where the Nyquist criterion becomes important, we modify our metric¹⁶ as follows:

$$\log \phi_{\tau} \propto \frac{\tau_{\rm obs}}{(1+z)\Delta t^c}.\tag{9}$$

Note that the fraction term on the right hand side of Equation 9 represents a sampling rate. There are different opinions in the literature when it comes to the sampling rate (Δt^c) for irregularly sampled data. Some authors (e.g., Horne & Baliunas 1986; Press et al. 1992) associate the Nyquist frequency with $1/2\langle t_{sam}\rangle$, where $\langle t_{sam}\rangle$ is the averaged sample rate, or with 1/(2s), where s is the smallest time sampling in the sample (Roberts et al. 1987; Scargle 1982).

For irregular sampling, Eyer & Bartholdi (1999) defined the Nyquist frequency as $f_{Ny} = 1/(2p) > 1/(2s)$, where p is a greatest common divisor (gcd) for all $t_i - t_1$ of time of observation t_i . Since gcd calculation is very time consuming, we will use the definition based on the smallest cadence in the sample (Roberts et al. 1987; Scargle 1982).

It is critical that the light curves are sufficiently well sampled for RM. Recalling Nyquist's Theorem, the sampling rate of the time lag measurement must be

$$\frac{\tau_{\rm obs}}{(1+z)\Delta t^c} > 2.2$$

in order for the time lag to be evaluated (Haas et al. 2011). As a result, our metric evaluates whether the sampling rate for a required time lag measurement is greater than the Nyquist threshold¹⁷.

4.2. Recovering time lags

For simulated quintets of AGN light curves in the LSST ugriz bands (Section 3.2), we estimate the time delay relative to *u*-band and compare it to the input value. We chose the *u* band as the reference band since its has the shortest wavelength. Moreover, as the cadence of *u*-band observations is a critical issue in the LSST observing strategy optimization (Brandt et al. 2018), we aim to probe if the *u*-band can be used to evaluate AD time lags in the DDFs.

In addition to the standard cross correlation methodologies for determining time lags (Gaskell & Sparke 1986; Gaskell & Peterson 1987; Alexander 1997), other methods have been developed as those built on the assumption of a DRW process (e.g., Zu et al. 2016), the DRW process combined with expansion of the transfer function into a series of Gaussian functions (Li et al. 2016), and shifting techniques (Tewes et al. 2013), to deal with complex light curves (Chan et al. 2020). For example, Kovačević et al. (2021a,b) adapted the functionality of the widely used ztransformed discrete correlation function technique (Alexander 1997) to take into account general Gaussian processes in the modeling of the light curves. We employed our shifting method to estimate time delays from the mock data, which is inspired by the PyCS toolbox developed by Tewes et al. (2013). One of the PyCS implementations is optimized to find the best set of time delays that minimizes the variability of the difference between Gaussian-process regressions on the light curves. This is motivated by the standard problem that there is a different number of light curve data points. An alternative objective function is needed to measure the similarity between the curves. As multiband light curves are mainly deformed due to the asymmetric transfer function, our basic idea of the optimizer is to shift the data iteratively in time and magnitude with respect to a reference band, and apply the Partial Curve Mapping ¹⁸ (PCM, Jekel et al. 2019) method to determine the similarity between curves and estimate a time delay through minimization of the PCM.

For measuring the time lag τ between two noisy light curves $f_1(t), f_2(t)$ we fit the relation $f_1(t) = a \cdot f_2(t-\tau)$ where a is a scaling coefficient. Denoting the LSST light curves as $X = \{x_j(t_i) | i = 1, ..., N, j \in (u, g, r, i, z)\}$, the scaling coefficients and the lags as a_k and $\tau_k, k \in g, r, i, z$, respectively, and finally the variances of the noise as v_k , we can fit the quartet of multiband light curves $\mathbf{x} = x_g, x_r, x_i, x_z$ with one predictor time series in the shortest wavelength x_u .

A prior estimate of the time delay range of light curves can be obtained through preprocessing with less precise algorithms (see e.g., Tewes et al. 2013), available prior knowledge (see an example of R-L relation use in Penton et al. 2022), or expected ranges obtained from assumed AD model $L = \text{math.floor}(\langle \tau \rangle_k - \langle \tau_u \rangle), k = g, r, i, z$ from AD model: [L, L + 1]. Considering the time lags that will be probed, we adopt the latter strategy, such that the uniform priors (in days) are: $\tau_{ug} = Uniform(0, 1), \tau_{ur} = Uniform(1, 2), \tau_{ui} = Uniform(2, 3), \tau_{uz} = Uniform(3, 4)$.

$$\frac{\tau_{\rm obs}}{(1+z)\Delta t^c} > 2.5.$$

 $^{^{16}}$ The other terms could be kept less than an arbitrary constant, e.g., assuming limits of $F_{\rm var} \sim 0.4$ and $\sigma \sim 0.005 {\rm mag}.$

¹⁷ Due to data loss to weather or technical difficulties in reality, practical sampling might require a stiffer criterion, known as the engineer's Nyquist (see e.g., Srinivasan et al. 1998):

¹⁸ https://github.com/cjekel/similarity_measures

They reflect the range of expected values of difference between mean time delays of the transfer function in different bands. These should only be considered as suggestions, and can be easily changed. Priors on expected time lags are non-informative, and their ranges do not overlap. Our emphasis is on the fact that multiple models for the AD model and its transfer function could be used in a real-world setting, and that the prior on the time lag should reflect this. To determine the similarity of curves after scaling and shifting, we apply PCM which combines arclength and area, because the choice of parameters can alter the overall length of the curve. We illustrate the principles of PCM, while details are given in (Jekel et al. 2019). First, the arc length of the one curve is superimposed onto a section of the other curve. After that, trapezoids are created between the curves, and the areas of the trapezoids are added together. The best fit PCM value is the one that minimizes the combined area of the trapezoids.

The PCM permits the use of non-resampled light curves; nevertheless, we followed the PyCS guideline and resampled them with homogeneous cadence of $\sim 0.5 - 1$ day. The time lag and its uncertainty is calculated by fitting the resampled, shifted light curves to the *u*-band light curve using a random value drawn from the prior. The PCM space for each light curve is quite irregular, with multiple local minima. To discern the different minima objectively and without expectation bias, we repeat the procedure 300 times for each of the resampled light curves, using a sample of 100 random-uniform initial time lags. The most common time lag (corresponding to 'dominant PCM minimum') is regarded as the best-fitting value for the specific light curve pair. Finally, from the distribution of fitted lag values, we compute PDFs, a mean time lag, and uncertainties as 16 and 84 quantile confidence intervals.

5. RESULTS - CADENCE STRATEGIES FOR AD TIME LAG MEASUREMENT

We use the $OpSim_v2.0$ and $OpSim_v1.7^{19}$ cadence simulations. The annotation adopted here follows the LSST nomenclature for the simulations. For example, the label baseline_v2.0_10yrs_5 days_r indicates that the metric is run on the observing strategy baseline, from the OpSim version 2.0, with a survey duration of 10 years, and that the time lag to be probed by the metric is 5 days, in the LSST r band.

The v2.0 simulations were developed in response to the Survey Cadence Optimization Committee's Phase 1 recommendations (Ivezić 2021). One of the most significant differences between previous survey simulations is that the baseline survey footprint has been slightly modified: the Wide Fast Deep area (WFD), which typically obtains on the order of 825 visits per pointing split across 6 filters, now explicitly includes a low-dust-extinction area as well as a Galactic Plane extension²⁰.

First, we provide the overall analysis in Section 5.1, which makes use of our metric AGN_TimeLagMetric implemented in MAF. The DDFs (DDFs, Brandt et al. 2018; Scolnic et al. 2018) are the most prominent locations for LSST RM, according to our initial analysis (Kovačević et al. 2021a). The existing baseline survey strategy contains five DDFs (Jones et al. 2020). Four of the DDF sites had been established for some time; the fifth field position is at present anticipated to overlap with the Euclid Deep Field South, which needs two LSST pointings to be covered entirely. The cadence of visits, as well as coadded depths of these DDFs, still needs to be finalized (Jones et al. 2020). The identifiers and equatorial coordinates of the DDFs are (see Jones et al. 2020): ELAISS1 ($9.45^{\circ}, -44.0^{\circ}$), XMM_LSS ($35.71^{\circ}, -4.75^{\circ}$), ECDFS ($53.13^{\circ}, -28.1^{\circ}$), COSMOS ($150.1^{\circ}, 2.18^{\circ}$), EDFS_a ($58.9^{\circ}, -49.32^{\circ}$), EDFS_b ($63.6^{\circ}, -47.6$)[°]. Note that the Euclid Deep Field South (EDFS) is divided into two sections (see details in Capak et al. 2019; Cuillandre et al. 2021) but for the sake of brevity, we only refer to the five unrelated DDFs when a broad discussion is required. Then, armed with the best strategies, we aim to simulate photometric time lag recovery as precisely as possible in Section 5.2. In our simulation, we chose a quasar redshift of z = 0.5 as for higher redshifts a careful choice of bands covering the continuum is needed (see details in Jankov et al. 2022).

5.1. Assessment of cadence strategies based on MAF simulations

Here, we use our metric to investigate the impact of different cadence strategies on the study of AD time lag measurement. We performed a simulation using the AGN_TimeLagMetric in MAF as described in Section 4.1. We probed fiducial time lags of 1, 2, 3, and 5 days because they are close to the expected mean time delay for our transfer function model. Aitoff projection of our metric realization in MAF simulation for LSST cadences are shown in Appendix (see Figures A1-A9).

 $^{^{19}\,\}rm https://github.com/lsst/sims_featureScheduler$

²⁰ https://github.com/lsst-pst/survey_strategy/blob/main/fbs_2.0/SummaryInfo_v2.0.ipynb

Table 1. Overview of the LSST cadence strategies. A 'visit' is currently defined to consists of two back-to-back 15 sec exposures, for a total of 30 sec of on-sky exposure time. These back-to-back exposures are always in the same filter, separated only by the 2 second readout time (Jones et al. 2020). From first to last column: observing strategy, OpSim version, the amount of area of WFD that receives at least 825 visits per pointing, the amount of area of WFD with low extinction, total number of visits per point on the sky, the median number of visits that the most frequently visited 18,000 square degrees receives, the median visits distributed per u and g band.

	OpSim	Area with	Area with	Nvisit	Median	Median Nvisit	
		> 825 visits	low extinction	total	Nvisit	u band	g band
		$[\deg^2]$	$[\deg^2]$				
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
baseline	2.0	12434.14	16842.94	2081749.0	838.0	54.0	68.0
carina	2.0	12519.74	16385.53	2087222.0	838.0	54.0	70.0
multi_short	2.0	151.07	16639.0	3584186.0	696.0	100.0	125.0
rolling_all_sky_ns2_rw0.9	2.0	12900.78	16599.55	2088105.0	839.0	54.0	69.0
short_exp	2.0	10664.07	16581.09	2180235.0	831.0	56.0	72.0
<pre>six_rolling_ns6_rw0.5</pre>	2.0	12342.65	16665.01	2084148.0	837.0	54.0	70.0
long_gaps_nightsof_delayed-1	2.0	10288.90	15954.13	2067452.0	830.0	50.0	70.0
euclid_dither1	1.7	17982.71	15174.43	2045493.0	888.0	55.0	79.0
euclid_dither3	1.7	17967.60	15161.84	2045144.0	888.0	55.0	78.0
footprint	1.7	15390.97	16492.96	2046219.0	853.0	52.0	72.0
u_long_ms_30	1.7	18091.81	15186.18	2080376.0	904.0	58.0	80.0

NOTE—More details are given in a collection of information relevant for survey strategy https://github.com/lsst-pst/ survey strategy. The cadence of visits, as well as coadded depths of DDFs, still needs to be finalized.

In our analysis, we searched for the shortest possible time lag that could be measured reliably by RM in all bands. In this and the following subsections, we show that the best cadences for simultaneous²¹ time lag measurements in *ugriz* bands are 5-day lags for the DDFs. Shorter time lags can be observed in *i* and *r* filters in DDFs, as will be seen in zoomed-in plots of metric values later on in this section. This should not come as a surprise given that these bands may contain cadences that are somewhat more dense (Kovačević et al. 2021a). Longer time lags, ≥ 7 days, are detectable in each of the filters, as we shall show in the following that mean cadence samplings are favorable across bands. As it could be an issue that the optimal solution matches the upper limit of the probed time-lags, we also tested these cadences by simulating RM measurements with shorter time delays in all bands.

Table 1 outlines some of the most important aspects of the LSST observing strategies used in this work. The baseline strategy (baseline_v2.0) features a modified²² survey footprint with an expanded dust-free area and WFD-level visits in the Galactic Bulge and Magellanic Clouds. Coverage of the Northern Ecliptic Spur, South Celestial Pole, and remainder of the Galactic Plane is maintained, at lower levels. The existing baseline used 2×15 sec exposures per visit, and most visits were in the same filter (although this was not enforced, Jones et al. 2020)²³. There are specific programs that, if given specialized observing time via "microsurveys," can produce additional science outside or in support of the basic Rubin LSST science drivers. One such program is that included in the strategy carina_2.0

²¹ The term simultaneity refers to the fact that our method determines all delays for the light curves quintet at the same time.

²² Filter balance is modified in different areas of the sky.

²³ As clarified by Jones et al. (2020): "A 'visit' here is an LSST default visit, which consists of two back-to-back 15 sec exposures, for a total of 30 sec of on-sky exposure time. These back-to-back exposures are always in the same filter, separated only by the 2 second readout time."

and is intended to observe the Carina nebula ²⁴. Both strategies show three prominent DDF fields-XMM_LS close to equator, ECDFS below previous and ELAISS1 close to polar axis (see Figure A1). For measuring time-lags smaller than 5 days carina cadences in r and z bands show a very small ares with Nyquist sampling or better at the sky location $\delta = 30^{\circ}$, $\alpha = 18 - 20h$ of the LSST equatorial sky plane.

OpSims are using a custom dither pattern for the Euclid deep drilling field to better match the Euclid field of view (EDFS_a, EDFS_b). Also, euclid_dither1 and euclid_dither3 exhibit three notable DDFs with moderate metric value in a tiny field at the position $\delta = 0^{\circ}, \alpha = 22h$ (Figure A2). Panel (d) in Figure A2 has an additional field of moderate metric value on the right side of main vertical axis of sky symmetry at the position $\delta > 0^{\circ}, \alpha = 8h$. The footprint cadence strategy allows various survey footprints based on increasing the low-extinction area of the WFD footprint. In all filters except the *u*-band filter, the cadence in footprint strategy exhibits a limited region which is shifting the location for time lags between 1 and 3 days (Figure A3). However, for the measurement of 5-day time lags in the *i* and *r* bands, the three DDFs (as in prior cadences) occur in panels (e) and (f) of Figure A4.

As the baseline survey strategies involves pairs of visits every few evenings, separated by around 33 min, long_gaps simulations include a third visit after a time interval of 2 to 7 hours. In the case of long_gaps_nightsoff, long gaps appear every seven nights starting after year five. DDFs are dominant in both the i and r bands for a measurement of 5-day time lags, which is comparable to prior occurrences in other metric realizations (see Figure A5).

Microsurveys multi_short_v2.0 cadences, emphasizes multiple (four) short (5 sec) exposures in each filter. For them metric simulations show a distinct topology so that suitable fields are clustered in a wavy band that is almost parallel to the sky equator but with varying densities (Figures A6-A7). Among the metric realizations, the *u*-band has the highest density of sky locations appropriate for RM, see Figure A6(d) and Figure A6(i), Figure A7(b) and Figure A7(g). The same three DDFs that were appearing in the prior cases are depicted in Figure A6(i) and Figure A7(h). Notably, four DDFs are found in the Figure A7(e) and Figure A7(f).

The rolling_all_sky_ns2 are two-band rolling all sky cadences with weights 0.5 and 0.9, where the metric realization is comparable to long_gaps_nightsoff, footprint, carina (see Figure A8) and euclid_dither (see Figure A9).

The microsurveys strategy short_exp takes up to three short (5 sec) visits in year one. As illustrated in Figures A10-A11, the topology of the metric realization is comparable to that of the metric realization on rolling_all_sky_ns2, with the exception that the *u*-band is more dominant.

As the updated baseline contains a two-band rolling cadence with strength $\sim 90\%$ in the dust-free WFD, six_rolling_ns6_rw0.5 simulations modify this rolling cadence in the dust-free WFD to use six regions instead of two, with variable weight ("strength") to the level of rolling. The u_long_ms_30 cadences take *u*-band observations as single snaps, and test increased *u*-band exposure times, in this case 30 secs. Interestingly, metric realizations on six_rolling_ns6_rw0.5 and u_long_ms_30_v1.7 cadences (Figure A12) are as prevalent in DDFs as they are in the case of metric realizations on euclid_dither cadences.

The LSST Science Advisory Council (SAC) report raised a series of questions and identified suggested simulation experiments to run. Among them are also listed Experiments with DDFs cadences (Jones et al. 2020). The observational cadence, which is currently being debated for the LSST DDFs (Jones et al. 2020), is indicated as one of the most critical elements that determine disk size measurements. The detailed investigation of the distribution of metric realizations (values above Nyquist criterion) on DDFs is shown in Figure 6. The highest quality time lag measurement is predicted to be obtained using data from the COSMOS, XMM_LSS and ECDFS whilst the lowest quality time lag measurement is expected to be obtained using data from the EDFS_a and EDFS_b fields. Notably, the metric value for the multi_short cadence in the *u* band is 20-30% lower than the threshold. As we will see in the following sections on examples of multi_short cadence, cadences with metric values nearing the threshold could be adequate to calculate time delays. The co-added depth varies across different filters and observing strategies in each DDF as illustrated in Figs.A13-A14. Currently, the deepest fields are ECDFS, XMM_LSS and ELAISS1. Interestingly, multishort strategy consistently produces the highest quality time delays and it also has the largest depth for all DDFs.

To demonstrate in greater detail the comparison shown in Figure 6, we perform MAF simulation in DDFs with significantly higher resolution (see Figures 7 and 8) than the original images in the Appendix. The HEALpygnomview is used to create the high resolution plots. The center of each plot is comprised of the point on the DDF itself (as mentioned above) plus a radius of 5 degrees (this defines the slice utilized to run the metric). The image itself displays



Figure 6. The comparison of metric values above the Nyquist threshold (y-axis) for the measurement of 5-day time lags in six DDFs, for various OpSim strategies indicated on the x-axis and six of the LSST filters stated in the legend. The solid lines serve as guides for the eye. The horizontal black line indicates the metric threshold. DDFs are indicated in the titles of panels. The passbands u, g, r, i, z, y are denoted with blue, orange, green, red, violet and brown color, respectively.

the resolution per pixel. The x-axis represents right ascension, and the y-axis represents declination. The metric realizations are given in two 3×3 grids of panels. The metric outperforms in all DDFs with the exception of EDFS_a and EDFS_b.

Based on the similarity of the metric value distributions, there are three types of DDFs: homogeneous DDFs with the highest metric values (first panel in the top row and middle panel in the middle row in the Figure 7), inhomogeneous DDFs with dispersed high metric values (second panel in the top row in panel in Figure 7; second and third panels in the top and middle row in Figure 8), and elliptically distributed DDFs (the last panel in the middle row in Figure 7; bottom rows in the Figures 7 and 8). This example, on the other hand, clearly indicates that observations in elliptical regions can be coupled with equivalent partitions in homogeneous DDFs in a straightforward manner.

In summary, we found that when the overall cumulative observational season length is long (~ 9 yr), the lag measurement likelihood increases because the multiband continuum light curves can track more light curve features. However, survey cadence strategy is also necessary for lag recovery because sampling the accretion disk light curves at the Nyquist frequency is required to resolve the lag accurately. As a result, objects having an estimated full width half-



Figure 7. Zoom-in of metric values (indicated in color bars) in DDFs for different cadence strategies (given in panel titles) for r, g, i bands. Upper row: COSMOS; Middle row: first and second panel ECDFS, third panel EDFS_a; Bottom row: EDFS_a.

maximum (FWHM) of the AD transfer function less than the minimum sampling are less likely to have a meaningful lag estimate. Larger gaps can be tolerated as long as the cadence is small and a long period can be covered, as proven by time lag measurements utilizing the COSMOS and ECDFS fields cadence strategies indicated by our metric.

5.2. Effective time lag measurement



Figure 8. The same as Figure 7, but for XMM in the top row; ELAISS1 in the middle row; EDFS_b in the bottom row.

Here, we further quantify the potential of measuring time lags using simulated LSST light curves for a subset of the best observational cadences as suggested by our metric realizations: the two DDFs (COSMOS and ECDFS) observed in multi_short_v2.0_10yrs cadence in the *ugrizy* bands.

We use typical parameters for quasars having median characteristics (SMBH mass $M = 2 \cdot 10^8 M_{\odot}$, $L/L_{\rm Edd} \sim 0.1$, $\eta = 0.1$, disk inclination of 45°, azimuth angle of 0°), characteristic DRW timescale $\tau_{\rm DRW} \sim 200$ days and a structure function at infinity $\sigma_{\rm DRW} = 2.5$, because we are interested in quantifying the performance of time lag measurement on a multishort cadence (having multiple (4) short (5s) exposures in each filter).

To create simulated light curves, we first produce the observation schedule and depth of each epoch using the LSST **OpSim** (Delgado & Reuter 2016). In each DDF we created a suite of quintets of multiband light curves by a convolution of the AD transfer function with the DRW driving light curve (see Section 3.2).

Examples of quintets for each DDF are given in Figure 9. The warping of light curves in different bands, due to transfer function skewness, is evident. For example, warping can be seen in **ELAISS1** DDF around MJD 61000 during one observational season. The photometric noise in all light curves is set to 0.005 mag (i.e. at the level of 0.02% as the best scenario found in our white paper²⁵ thorough simulations (Kovačević et al. 2021b). The first row in Table 2 lists the true values of AD time lags $\tau_{uj} = \langle \tau \rangle_j - \langle \tau \rangle_u$, $j \in (g, r, i, z)$ as calculated from Equation 5. The remainder show estimates using our procedure to determine time lags of the light curves quintets (see Section 4.2 for each details).



Figure 9. Examples of simulated quintets of LSST light curves for quasars in the LSST DDFs. We simulate the light curves over a period of 10 years with multi_short_v2.0_10yrs cadences. The error bars are set at 0.005 mag *rms*.

The estimated 1D probability distribution for each time lag and the joint probability distribution for each pair of time lags, are shown in Figure 10. We see that the inferred values of the time lags are both accurate (close to true values) and precise (narrow posterior distribution) for ECDS light curves in g, r and i bands. However, the PDF of τ_{ug} is deformed for COSMOS. In general, we see from the 1D histogram of Figure 10 that PDF intervals are much narrower than



Figure 10. Corner plot showing the PDFs of the time lags of the synthetic light curves with DDFs COSMOS (blue) and ECDFS (orange) sampling (left) and homogeneous 1-day (blue) and 5-day (orange) sampling (right). The distribution for each time lag independently is shown in the histograms along the diagonal, and the 2D distributions as contour plots in the other panels. The dashed black lines are the true time lag values inferred from the AD model. The contour levels of the 2D joint probability densities are at 1σ and 2σ regions.

the prior ranges (see Subsection 4.2). Table 2 lists the mean of the posterior and uncertainties corresponding to 16th and 84th quantile of the distribution of measured time lags. Because light curve distortion can make detecting delays in individual segments of light curves difficult, the AD time lag measurement is demonstrated on decadal observational time baseline (see e.g. Chan et al. 2020).

The light curves in OpSim observation schedules have breaks between a series of continuous observations, with the gaps ranging from several days to ≥ 80 days. To see how these gaps affect lag measurements, we created additional ideally sampled simulated light curves with homogeneous sampling of 1 and 5 days.

By examining the corner plot for recovered time lags of 1 and 5 day homogeneous light curves (right panel in Figure 10), we conclude that the lag distributions are affected by homogeneous sampling. For the 1 day sampling, the PDFs are narrower and more symmetric with respect to true values. Table 2 presents summary statistic of distribution of time lag measurement on these ideally sampled quintets of light curves.

Barth et al. (2011) have shown that for interpolation and cross correlation methods applied to real data, as long as the driving light-curve is well sampled (in their case almost nightly over the course of 200 days), poorer sampling of the output light-curves can be tolerated. We recall that, to first order, the centroid of the transfer function is simply proportional to the luminosity-weighted radius (Koratkar & Gaskell 1991). However, in the case of LSST multiband continuum transfer functions, their centroids are shifted closer to their FWHM (see Figure 3). Also, we see that interband continuum delays which are driven by DRW, have values below or close to transfer function centroids (e.g., Figure 10). So, it is important to have a sampling that is of the order of the FWHM of the AD transfer function. This condition is most relevant for measuring the time lag between u-band and g-band light curves as their interband time delay is the smallest.

6. DISCUSSION

Our MAF metric (AGN_TimeLagMetric) indicates that the best cadences (i.e., those with the largest values above the Nyquist threshold) in the DDFs are those in the multi_short family, taking into account fiducial time lags of 5 days

Table 2. Summary statistics of recovered time lags from the distributions for COSMOS, ECDFS and homogeneously sampled light curves given in Figure 10. The columns are: status of time lags, light curve sampling, the time lags of *g*-band vs. reference *u*-band light curves; *r*-band vs *u*-band light curves; *i*-band vs. *u*-band light curves; *z*-band vs. *u*-band light curves.

Status	LC sampling	$ au_{ug}$	$ au_{ur}$	$ au_{ui}$	$ au_{uz}$	
(1)				(5)	(6)	
(1)	(2)	(3)	(4)	(5)	(6)	
True	-	0.653	1.514	2.388	3.177	
Inferred	COSMOS	$0.643\substack{+0.06\\-0.04}$	$1.507^{+0.027}_{-0.02}$	$2.388\substack{+0.029\\-0.033}$	$3.166\substack{+0.017\\-0.017}$	
Inferred	ECDFS	$0.650\substack{+0.008\\-0.008}$	$1.540^{+0.012}_{-0.08}$	$2.377^{+0.017}_{-0.022}$	$3.168^{+0.012}_{-0.009}$	
Inferred	1-day	$0.649^{+0.01}_{-0.01}$	$1.508\substack{+0.01\\-0.01}$	$2.388\substack{+0.04\\-0.04}$	$3.168\substack{+0.01\\-0.01}$	
Inferred	5-day	$0.649\substack{+0.01\\-0.01}$	$1.516\substack{+0.03\\-0.04}$	$2.369^{+0.02}_{-0.03}$	$3.167\substack{+0.01\\-0.01}$	

NOTE—True values of mean relative time delays $\langle \tau \rangle_{uj}, j \in (g, r, i, z)$ calculated from AD model (see Equation 5). The inferred values represent the 50th quantile with errors corresponding to 16th and 84th quantiles of the respective probability distributions of calculated time lags.

in all LSST filters. This family of strategies belong to the micro-surveys. Micro-surveys aim to support and extend the LSST's primary science drivers, and they were submitted in response to the 2018 Call for White Papers (see Bianco et al. 2022). Many of the micro-surveys requiring between 0.3 - 3% of the overall survey time were simulated as part of OpSim v2.0 simulations. The multi_short takes four short (5 sec) exposures in each filter, which is about 20 sec in total per filter. Despite the short exposure time, the high observational cadence allows for photometric RM analysis as indicated by our simulations as well as other studies (see Yu et al. 2020). However, when considering micro-surveys, it is worthwhile to note that sometimes visits can be extremely short, which can affect RM observables. The DDFs currently receive about 1 percent of the total survey time per DDF location. There are five DDFs, so this accounts for 5 % of the total survey time. Each DDF is a single pointing, with the exception of the Euclid South DDF which consists of two shallower pointings.

However, there are simulations, where the amount of time allocated for DDFs is varied from 3 percent (ddf_fract_ddf_per0.6_v2.0_10yrs) to 8 percent (ddf_fract_ddf_per1.6_v2.0_10yrs). Our MAF simulations show that there is no significant difference between these two variants of DDFs when it comes to time lag measurements.

From our simulations (see also Kovačević et al. 2021b), we find that the most important parameters that affect the quality of time delay measurements in AGN ADs are the cumulative season length and the mean sampling (also referred to as the sampling frequency). A categorization of 14 observing strategies in all DDFs and LSST filters, based on mean 5σ depth, mean sampling, and mean cumulative season length are shown in Figures 11-13).

The mean cumulative season length, mean sampling and mean 5σ depth from a simulation of a given observing strategy are calculated by taking the mean of these parameters for the DDF under consideration (Figures 11-13). The cumulative season length is the sum of the duration of all observing seasons. A season gap for an LSST field is defined as no observations in any filter taking place for 85 days (Huber et al. 2019). Across all DDFs, the cumulative season length for the majority of strategies is ≥ 9.5 yr. We note that multi_short strategy has the fastest (smallest) sampling but the mean 5σ depth is shallower. The mean sampling exhibits a similar pattern of behavior per filter throughout all DDF fields, with subtle variations. Figure 11 shows COSMOS field as an example. The mean sampling for some of the observing strategies in the *u*-band in the COSMOS differs from other fields (e.g., compare mean sampling for families between carina and footprint in COSMOS and ECDFS, see Figure 11), which is related to time spent on other surveys such as the northern hemisphere, the southern Celestial Pole, and the Galactic Center. As seen in Figures 11-13, *y*-band has the best mean sampling; yet, *y*-band light curve distortion is the greatest due to AD transfer



Figure 11. Comparison of mean 5σ depth (left column), mean sampling (middle column) and mean cumulative season length (right column) for different observing strategies (x-axis) in DDFs and *ugrizy* filters. The solid lines guide the eye. The mean 5σ depths of the non-*r* and *r* bands are close. This means that objects as bright as 21 will still be seen in those bands for these overlapping cadences in DDF. Also, fainter objects in non-*r* band will be observable in *r*-band. The passbands *u*, *g*, *r*, *i*, *z*, *y* are denoted with blue, orange, green, red, violet and brown color respectively.

function asymmetry, making determining the time lag problematic (not impossible but with the largest error). Thus, for simplicity we do not consider the y-band lags.

We underline that our metric pointed to those strategies that would be appropriate for AD time lag measurements of the order of 5 light days, which is a severe limitation. We investigated alternative cuts for 5 σ magnitude depth for brighter (r < 21, g < 22) and fainter (r > 21, g > 22) sources. Adding a cut for 5 σ depth in the r band < 21.0 implies that the (vast majority) of visits with r > 21.0 will not be counted, but objects as bright as 21 in the \mathbf{r} band will still be detectable; on the other hand, using a cut of r < 21 implies that only visits with a very limited depth will be available (Jones, private communication). However, no significant adjustments have been made in the overall analysis, as strategies with mean 5 σ depth $r \sim 20 - 21, g \sim 20 - 21$ have been chosen.



Figure 12. The same as Figure 11 but for different DDFs.

Next, we obtain the differential number distribution of quasars with possible time lag measurements with respect to the u band by integrating the time lag measurement distribution over the source population:

$$\frac{dN}{dz} = \int_{L}^{\infty} dL \frac{d^2 \mathcal{N}}{dL dV} \Omega \frac{dV}{dz} p_{\tau}$$

where $\frac{d^2 \mathcal{N}}{dLdV}$ is a QLF, Ω is the fiducial survey area, and p_{τ} is the probability of a time lag measurement as defined in Section 2. We generated time lags using our AD model for 200 uniformly distributed SMBH masses $(10^6, 10^{10})M_{\odot}$ over 200 equal bins $(dz \sim 0.035)$ of redshifts $(z \sim 0 - 7)$. The total number of quasars for which time lags could be measured is

$$N = \int \frac{dN}{dz} dz \; .$$

In Figure 14, we plot the expected number of quasars (in 10deg^2 survey area) for which it will be possible to measure time lags in the main LSST bands simultaneously assuming a fiducial cadence ~ 1day as a function of redshift. The steepness of these number counts is shallower for u-band than for other bands, implying that for time lag measurements using u-band, the survey area is far more relevant than the redshift. Total counts will decrease in the same proportion



Figure 13. The same as Figure 11 but for different DDFs.

as cadence increases. As a result, the estimated number of quasars for a 5-day cadence will be 5 times lower than for a 1-day cadence. We emphasize that this is an upper bound (i.e. optimistic) estimate of expected number of quasars with possible time lag measurements.

Also, we approximate the total number of quasars whose time lag could be detected in a DDF (in any filter) with confidence CI assuming an accuracy of the time lag estimate of at least $\Delta = 10\%$ for blind search (no data about mass and redshift) as:

$$N \sim CIN_{rep} \Delta \frac{\Omega_{LSST}}{\Omega_{rep}} \frac{t_{LSST}^{cum}}{t_{rep}^{cum}} ,$$

where $N_{rep} \sim 344$ is the number of quasars in VST COSMOS estimated by De Cicco et al. (2021), $\Omega_{LSST} \sim 10 \text{deg}^2$ is a fiducial element of LSST area, $\Omega_{rep} \sim 1 \text{deg}^2$ is an area associated with N_{rep} , and $t_{rep}^{cum} \sim 3\text{yr}$ and $t_{LSST}^{cum} \sim 10\text{yr}$ are mean cumulative season lengths for VST COSMOS and LSST, respectively. Then the number of quasars for which time lags will be determined at least at $CI \sim 0.95$ and with accuracy rate $\Delta \sim 0.1$ is $N \sim 1089$ per 10 deg^2 in any filter.

Time lags measured with the actual DDF cadences are less accurate and precise than those estimated using homogeneously sampled light curves (see Figure 10). For instance, we see that the COSMOS and ECDFS PDFs are more skewed



Figure 14. The expected number of quasars for which time lag could be measured with respect to u-band, in main LSST filters as a function of the redshift, for fiducial 10deg^2 survey area and fiducial cadence of 1 day.

than for homogeneously sampled light curves. Furthermore, the PDF of measured time lags can be bimodal, as found for r-band light curves in ECDFS. This is not surprising given that our metric shows that r band sampling in ECDFS is of lower quality than in COSMOS, whereas it is the same for other bands (see Figure 6). Whilst the secondary peak on the left is much closer to the true value, it is also much less probable. We showed that using a shifting algorithm with a light curve similarity measure based on area and arc-length might yield temporal lag uncertainties in DDFs of the order of 10% or less (see Table 2). RM measurement errors are anticipated to be more tolerable in the DDFs, where sampling is expected to be an order of magnitude denser than in the main survey. By modelling many different observation tactics for the LSST DDFs, Yu et al. (2020) discovered that changing the cadence from 3 days to 2-1 days should greatly improve the yield of accretion disk size measurements. This is in accordance with other studies on reverberation mapping. For example, typical time lag errors of $\sim 30\%$ are typically reported by spectroscopic reverberation experiments (Haas et al. 2011; Pozo Nuñez et al. 2015). Note as well that estimated uncertainties of 30% for undersampled data may be optimistic (Haas et al. 2011). Enhanced sampling for a spectroscopic reverberation campaign on NGC 4051 revealed that its BLR size is actually a factor of three lower than previously assumed from undersampled data (Denney et al. 2009). Thus, with well-sampled LSST reverberation data, the time lag estimations can be improved and tensions with theoretical estimates may be resolved.

It has been reported that curve-shifting algorithms sensitive to sharp features and operating on the same time scales as the transfer function underestimate multi-band time delays by up to 20% (Chan et al. 2020). However, using machine learning methods for a similarity measure of light curves that do not require the same length and number of points, such as PCM, could be advantageous for LSST RM. Our preliminary tests show that using curve similarity measures has the potential to improve time lag determination so that a typical errors might be within 1% - 10% (see Table 2).

Because the disk variability for type I AGNs is almost unobscured, measuring intercontinuum delays for these objects will be easier than for type II (due to longer variability time scales and smaller variability amplitudes)²⁶. The LSST will also increase the number of type I AGNs, and low-luminosity AGNs with smaller black hole masses, so that shorter continuum delays will become more common. As the magnitude of optical variability on order of ~days could be anticorrelated with SMBH mass and AGN luminosity (Kelly et al. 2013), perhaps AGNs with smaller black hole masses will be better candidates for lag measurements (see also Yu et al. 2020).

An alternative to intensive high-cadence monitoring of a small number of objects is to follow up a larger sample of objects at a cost of a lower cadence. While individual objects are unlikely to provide high-fidelity lag-wavelength relationships, a large population of objects can be analyzed as a whole (see e.g., Jiang et al. 2017; Mudd et al. 2018; Yu et al. 2020; Homayouni et al. 2019). Both Homayouni et al. (2019) and Yu et al. (2020) report disk sizes that

²⁶ If type I Seyferts are host dominated, this issue may be applied to them as well.

are consistent with the standard disk model, whereas Homayouni et al. (2019) demonstrate that if only the best lag measurements are used, disk sizes are skewed to be higher than when the entire sample is taken into account. However, the multiband AD light curves are warped due to transfer function skewness. We demonstrate that when light curves are denser, as in the case of DDF, where sampling approaches the width of their transfer functions, the results are better. The *u*-band light curve has the shortest wavelength and the smallest width of the transfer function, implying that its higher cadence is very desirable. However, in order to examine X-ray to UV delays, the sample density of light curves in all bands should be greater than the width of their transfer function (see Kammoun et al. 2021).

Accretion disk investigations have been challenged by the fact that observations of their sizes are consistently larger than expectations from the thin-disk model by factors ranging from two to three (Chan et al. 2020). For an accurate measurement of disk size, it is necessary to take into consideration the appropriate transfer functions (Chan et al. 2020; Starkey et al. 2016). DDFs (Brandt et al. 2018; Scolnic et al. 2018) where substantial multiwavelength observations have been collected from earlier surveys (De Cicco et al. 2021) will be of greatest interest for detecting time lags by using these previous data to constrain the transfer functions. There are ongoing efforts of creating prime sample of quasars to further test the applicability of the DRW model and the stationarity of the stochastic variability process. Stone et al. (2022) investigated the optical continuum variability of 190 quasars within the SDSS Stripe 82 over 20 years long baseline. This sample contains hundreds of epochs from SDSS, PanSTARRS-1, the Dark Energy Survey, and dedicated follow-up photometric monitoring with DECam on the CTIO-4m Blanco telescope, making it one of the best-quality light curve data sets for studying quasar variability. Monitoring of this sample is a continuous endeavor to photometrically monitor deep extragalactic fields and collect a large amount of multi-wavelength and time-domain data (Stone et al. 2022). Extension of the baseline by another 5 years will allow these light curves to be smoothly combined with LSST data, so that this quasar sample will be premier for studying optical continuum variability.

We also briefly discuss the relation of our analysis to an important methodological study by Hu & Tak (2020) on multiband GP, the motivation of which is consistent with this work. Assuming correlations of the characteristic timescale (τ) and an asymptotic rms variability on long time-scales (SF_{∞}) among bands, they proposed a multivariate DRW model for modelling AGN light curves in five bands and with irregular cadences. Similar to our study, Hu & Tak (2020) firstly generated simulated light curves with measurement errors using the DRW model. Then Hu & Tak (2020) applied the univariate and multivariate DRW model on simulated light curves and a SDSS spectroscopically-confirmed quasar, and found that the multivariate model can help reveal the possible correlations of true time-scales across five bands and time delays between doubly lensed multiband light curves. Although our analysis does not include the more rigorous multiband GP time lag method, the time lag method in our study is sufficient for demonstrating RM within various OpSim runs in terms of optimizing observing cadence and strategy.

7. FINAL REMARKS

In this paper, we introduced a metric incorporated in MAF that quantifies the accuracy of LSST observational strategies for AGN continuum time lag measurements. Our metric relies on the Nyquist sampling criterion, which is advantageous when discerning information on small scales structures via electromagnetic signals. This enables us to identify specific LSST sky regions (compiled in an atlas) where observing strategies of high quality (satisfying the Nyquist criterion) could be effective for probing continuum time lags on scales of the AD and the transfer function.

A prominent example that will necessitate such spatial scales and will be explored in the future is the possibility that most observed continuum lags are driven by variable diffuse BLR emission rather than the irradiation disk (see Netzer 2022). We anticipate that light curves obtained from fields indicated by our metric will be useful for resolving what happens on long time-scales and how it diverges from the lamppost model (Neustadt & Kochanek 2022).

Unlike the spectroscopic approach, photometric RM does not typically allow the BLR emission line and continuum light curves to be separated. Under certain conditions, separation of these processes is possible so the lag can be measured (see details in Chelouche & Daniel 2012; Chelouche et al. 2012). LSST sky regions selected by our metric will be ideal for testing the photometric RM methodology on LSST data (Jankov et al. 2022).

It is also interesting to consider what deep learning algorithms (see e.g., Tachibana et al. 2020; Cvorović Hajdinjak et al. 2022) might reveal when trained on multicolor higher cadence data in optimal LSST sky regions (as defined by our metric). Another feature of our metric is that it can be simply converted to probability and hence combined with other metrics such as quasar counts (Assef et al. 2021a).

The following is a brief summary of our work:

- We compiled a comprehensive atlas of estimates of the performance of continuum continuum time lag measurements as a result of our metric simulation conducted on MAF. The metric points out as the best strategies those found in DDFs. The multi_short strategy has the best sampling (≤ 5 days) but the mean 5σ depth is shallower.
- To illustrate our finding, we simulate light curves using an AGN model based on a combination of thin-disk, lamppost and damped random walk for the driving function. We estimate the time lags using shifting method inspired by PyCS but with Partial Curve Mapping (PCM) method to determine the similarity between curves and identify a time delay measurement through minimization of PCM. Even with inferior sampling quality, like in the *r*-band of ECDFS, the time lags could be determined with accuracy of ~ 10%.
- Using our metric together with the artificially generated sample of quasar masses and their QLF, we predicted an upper bound of total quasar counts within 0 < z < 7 for which their time lags might be measured. We estimate that time lag measurements leading to AD size measurements for more than 1000 sources in each DDF per 10deg^2 in any filter will be possible. We found that the **baseline** family is suboptimal in all DDFs, as our metric is below the threshold in g and z bands. Similarly, the **rolling** family underperforms in the r and g bands. In contrast, the **good_seeing** family of cadences is mainly moderate for determining time lag in all DDFs.
- Based on preliminary numerical results, we propose that, in addition to methods that take into account the transfer function form in detail, LSST continuum reverberation mapping can apply deep learning techniques of similarity measures to potentially improve the time-lag measurement process.

We sincerely thank Rachel Street, Federica Bianco, Gordon T. Richards, and William N. Brandt for their essential 1 efforts in coordinating work on cadence evaluation within the Rubin-LSST Science Collaborations. This work was 2 conducted as joint action of the Rubin-LSST Active Galactic Nuclei (AGN) and Transients and Variable Stars (TVS) Science Collaborations. The authors express their gratitude to the Vera C. Rubin LSST AGN and TVS Science 4 Collaborations for fostering cooperation and the interchange of ideas and knowledge during their numerous meetings. 5 The authors would like to thank the Vera C. Rubin Observatory MAF team, specially Lynne Jones, for their significant support in designing and integrating our metric into MAF. The authors warmly acknowledge the LSST Corporation's 7 efforts to train and educate metric teams through many cadence tuning workshops and hackathons. 8 ABK, DI, IJ, ICH, LCP and VR acknowledge funding provided by University of Belgrade-Faculty of Mathematics (the contract 451-03-68/2022-14/200104), through the grants by the Ministry of Education, Science, and Technological 10 Development of the Republic of Serbia. ABK and LČP thank the support by Chinese Academy of Sciences President's 11 International Fellowship Initiative (PIFI) for visiting scientist. BC and SP acknowledge the financial support from the 12 Polish Funding Agency National Science Centre, project 2017/26/A/ST9/-00756 (MAESTRO 9) and MNiSW grant 13 DIR/WK/2018/12. CR acknowledges support from the Fondecyt Iniciacion grant 11190831 and ANID BASAL project 14 FB210003. DI acknowledges the support of the Alexander von Humboldt Foundation. HL acknowledges a Daphne 15 Jackson Fellowship sponsored by the Science and Technology Facilities Council (STFC), UK. JKD, LCP, ML, OV, 16 SMM acknowledge funding provided by Astronomical Observatory (the contract 451-03-68/2022-14/200002), through 17 the grants by the Ministry of Education, Science, and Technological Development of the Republic of Serbia. RJA 18 was supported by FONDECYT grant number 1191124 and by ANID BASAL project FB210003. PSS acknowledges 19 funds by ANID grant FONDECYT Postdoctorado N^o 3200250. SP acknowledges financial support from the Conselho 20 Nacional de Desenvolvimento Científico e Tecnológico (CNPq) Fellowship (164753/2020-6). SS acknowledges funding 21 provided by the University of Kragujevac - Faculty of Sciences (the contract 451-03-68/2022-14/200122) through 22 the grants by the Ministry of Education, Science, and Technological Development of the Republic of Serbia. YRL 23 acknowledges financial support from the National Science Foundation of China through grant No. 11922304 and from 24 the Youth Innovation Promotion Association CAS. 25

REFERENCES

Cuillandre, J. C., Rhodes, J., Stern, D., et al. 2021, LSST Cadence Note – Enhancing LSST Science with Euclid synergy and a mini-survey of the northern sky to Dec <+30 Aihara, H., AlSayyad, Y., Ando, M., et al. 2022,
Publications of the Astronomical Society of Japan, 74, 247, doi: 10.1093/pasj/psab122

28

- Alexander, T. 1997, in Astrophysics and Space Science Library, Vol. 218, Astronomical Time Series, ed.
 D. Maoz, A. Sternberg, & E. M. Leibowitz, 163, doi: 10.1007/978-94-015-8941-3 14
- Anglés-Alcázar, D., Quataert, E., Hopkins, P. F., et al. 2021, ApJ, 917, 53, doi: 10.3847/1538-4357/ac09e8
- Assef, R. J., Temple, M., Richards, G., Yu, W., & Bauer, F.
 2021a, LSST AGN SC Cadence Note: Quasar Counts
 —. 2021b, LSST AGN SC Cadence Note: Type-1 Quasar
- Colors in the Context of Photometric Redshifts
- Barth, A. J., Nguyen, M. L., Malkan, M. A., et al. 2011, ApJ, 732, 121, doi: 10.1088/0004-637X/732/2/121
- Bellm, E. C., Burke, C. J., Coughlin, M. W., et al. 2022, ApJS, 258, 13, doi: 10.3847/1538-4365/ac4602
- Bianco, F. B., Ivezić, Ž., Jones, R. L., et al. 2022, ApJS, 258, 1, doi: 10.3847/1538-4365/ac3e72
- Biava, N., Brienza, M., Bonafede, A., et al. 2021, A&A, 650, A170, doi: 10.1051/0004-6361/202040063
- Blandford, R. D., & Begelman, M. C. 1999, MNRAS, 303, L1, doi: 10.1046/j.1365-8711.1999.02358.x
- Blandford, R. D., & McKee, C. F. 1982, ApJ, 255, 419, doi: 10.1086/159843
- Brandt, W. N., Ni, Q., Yang, G., et al. 2018, arXiv e-prints, arXiv:1811.06542. https://arxiv.org/abs/1811.06542
- Burke, C. J., Shen, Y., Blaes, O., et al. 2021, Science, 373, 789, doi: 10.1126/science.abg9933
- Cackett, E. M., Chiang, C.-Y., McHardy, I., et al. 2018, ApJ, 857, 53, doi: 10.3847/1538-4357/aab4f7
- Cackett, E. M., Horne, K., & Winkler, H. 2007, MNRAS, 380, 669, doi: 10.1111/j.1365-2966.2007.12098.x
- Campana, S., & Stella, L. 1995, MNRAS, 272, 585, doi: 10.1093/mnras/272.3.585
- Campitiello, S., Ghisellini, G., Sbarrato, T., & Calderone, G. 2018, A&A, 612, A59,
- doi: 10.1051/0004-6361/201731897

Capak, P., Cuillandre, J.-C., Bernardeau, F., et al. 2019, arXiv e-prints, arXiv:1904.10439. https://arxiv.org/abs/1904.10439

- Caplar, N., Lilly, S. J., & Trakhtenbrot, B. 2017, ApJ, 834, 111, doi: 10.3847/1538-4357/834/2/111
- Chan, J. H. H., Millon, M., Bonvin, V., & Courbin, F. 2020, A&A, 636, A52, doi: 10.1051/0004-6361/201935423
- Chartas, G., Kochanek, C. S., Dai, X., et al. 2012, ApJ, 757, 137, doi: 10.1088/0004-637X/757/2/137
- Chartas, G., Rhea, C., Kochanek, C., et al. 2016, Astronomische Nachrichten, 337, 356, doi: 10.1002/asna.201612313

Chelouche, D. 2013, ApJ, 772, 9, doi: 10.1088/0004-637X/772/1/9

- Chelouche, D., & Daniel, E. 2012, ApJ, 747, 62, doi: 10.1088/0004-637X/747/1/62
- Chelouche, D., Daniel, E., & Kaspi, S. 2012, ApJL, 750, L43, doi: 10.1088/2041-8205/750/2/L43
- Cherepashchuk, A. M., & Lyutyi, V. M. 1973, Astrophys. Lett., 13, 165
- Collier, S. J., Horne, K., Kaspi, S., et al. 1998, ApJ, 500, 162, doi: 10.1086/305720
- Connolly, A. J., Angeli, G. Z., Chandrasekharan, S., et al. 2014, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 9150, Modeling, Systems Engineering, and Project Management for Astronomy VI, ed. G. Z. Angeli & P. Dierickx, 915014, doi: 10.1117/12.2054953
- Dai, X., Kochanek, C. S., Chartas, G., et al. 2010, ApJ, 709, 278, doi: 10.1088/0004-637X/709/1/278
- De Cicco, D., Bauer, F. E., Paolillo, M., et al. 2021, A&A, 645, A103, doi: 10.1051/0004-6361/202039193
- Delgado, F., & Reuter, M. A. 2016, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 9910, Observatory Operations: Strategies, Processes, and Systems VI, ed. A. B. Peck, R. L. Seaman, & C. R. Benn, 991013, doi: 10.1117/12.2233630
- Denney, K. D., Watson, L. C., Peterson, B. M., et al. 2009, ApJ, 702, 1353, doi: 10.1088/0004-637X/702/2/1353
- Dexter, J., & Agol, E. 2011, ApJL, 727, L24, doi: 10.1088/2041-8205/727/1/L24
- Du, P., Hu, C., Lu, K.-X., et al. 2015, ApJ, 806, 22, doi: 10.1088/0004-637X/806/1/22
- Edelson, R., Gelbord, J. M., Horne, K., et al. 2015, ApJ, 806, 129, doi: 10.1088/0004-637X/806/1/129
- Edelson, R., Gelbord, J., Cackett, E., et al. 2017, ApJ, 840, 41, doi: 10.3847/1538-4357/aa6890
- ---. 2019, ApJ, 870, 123, doi: 10.3847/1538-4357/aaf3b4
- Event Horizon Telescope Collaboration, Akiyama, K., Alberdi, A., Alef, W., & Algaba, J. C. 2022, ApJL, 930, L16, doi: 10.3847/2041-8213/ac6672
- Event Horizon Telescope Collaboration, Akiyama, K., Alberdi, A., et al. 2019, ApJL, 875, L4, doi: 10.3847/2041-8213/ab0e85
- Eyer, L., & Bartholdi, P. 1999, A&AS, 135, 1, doi: 10.1051/aas:1999102
- Fausnaugh, M. M., Denney, K. D., Barth, A. J., et al. 2016, ApJ, 821, 56, doi: 10.3847/0004-637X/821/1/56
- Frank, J., King, A., & Raine, D. J. 2002, Accretion Power in Astrophysics: Third Edition
- Galeev, A. A., Rosner, R., & Vaiana, G. S. 1979, ApJ, 229, 318, doi: 10.1086/156957

- Gaskell, C. M., & Peterson, B. M. 1987, ApJS, 65, 1, doi: 10.1086/191216
- Gaskell, C. M., & Sparke, L. S. 1986, ApJ, 305, 175, doi: 10.1086/164238
- Gravity Collaboration, Sturm, E., Dexter, J., et al. 2018, Nature, 563, 657, doi: 10.1038/s41586-018-0731-9
- Guo, W.-J., Li, Y.-R., Zhang, Z.-X., Ho, L. C., & Wang, J.-M. 2022, arXiv e-prints, arXiv:2201.08533. https://arxiv.org/abs/2201.08533
- Haas, M., Chini, R., Ramolla, M., et al. 2011, A&A, 535, A73, doi: 10.1051/0004-6361/201117325
- Homayouni, Y., Trump, J. R., Grier, C. J., et al. 2019, ApJ, 880, 126, doi: 10.3847/1538-4357/ab2638
- Hönig, S. F. 2014, ApJL, 784, L4, doi: 10.1088/2041-8205/784/1/L4
- Hopkins, P. F., Hernquist, L., Cox, T. J., et al. 2005, ApJ, 630, 705, doi: 10.1086/432438
- Hopkins, P. F., Richards, G. T., & Hernquist, L. 2007, ApJ, 654, 731, doi: 10.1086/509629
- Horne, J. H., & Baliunas, S. L. 1986, ApJ, 302, 757, doi: 10.1086/164037
- Hu, Z., & Tak, H. 2020, AJ, 160, 265, doi: 10.3847/1538-3881/abc1e2
- Huber, S., Suyu, S. H., Noebauer, U. M., et al. 2019, A&A, 631, A161, doi: 10.1051/0004-6361/201935370
- Ichikawa, K., Ueda, J., Bae, H.-J., et al. 2019, ApJ, 870, 65, doi: 10.3847/1538-4357/aaf233
- Ichimaru, S. 1977, ApJ, 214, 840, doi: 10.1086/155314
- Ilić, D., Oknyansky, V., Popović, L. Č., et al. 2020, A&A, 638, A13, doi: 10.1051/0004-6361/202037532
- Ivezić, Ž. 2021, Survey Cadence Optimization Committee's Phase 1 Recommendation
- Ivezić, Ž., Kahn, S. M., Tyson, J. A., et al. 2019, ApJ, 873, 111, doi: 10.3847/1538-4357/ab042c
- Jankov, I., Kovačević, A. B., Ilić, D., et al. 2022, Astronomische Nachrichten, 343, e210090, doi: https://doi.org/10.1002/asna.20210090
- Jekel, C. F., Venter, G., Venter, M. P., Stander, N., & Haftka, R. T. 2019, International Journal of Material Forming, doi: 10.1007/s12289-018-1421-8
- Jiang, Y.-F., Green, P. J., Greene, J. E., et al. 2017, ApJ, 836, 186, doi: 10.3847/1538-4357/aa5b91
- Jones, R. L., Yoachim, P., Ivezić, Ż., Neilsen, E. H., & Ribeiro, T. 2020, Survey Strategy and Cadence Choices for the Vera C. Rubin Observatory Legacy Survey of Space and Time (LSST) (v1.2), doi: 10.5281/zenodo.4048838

- Jones, R. L., Yoachim, P., Chandrasekharan, S., et al. 2014, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 9149, Observatory Operations: Strategies, Processes, and Systems V, ed.
 A. B. Peck, C. R. Benn, & R. L. Seaman, 91490B, doi: 10.1117/12.2056835
- Kammoun, E. S., Papadakis, I. E., & Dovčiak, M. 2021, MNRAS, 503, 4163, doi: 10.1093/mnras/stab725
- Kara, E., Cackett, E. M., Fabian, A. C., Reynolds, C., & Uttley, P. 2014, MNRAS, 439, L26, doi: 10.1093/mnrasl/slt173
- Kara, E., Mehdipour, M., Kriss, G. A., et al. 2021, ApJ, 922, 151, doi: 10.3847/1538-4357/ac2159
- Kaspi, S., Smith, P. S., Netzer, H., et al. 2000, ApJ, 533, 631, doi: 10.1086/308704
- Kelly, B. C., Bechtold, J., & Siemiginowska, A. 2009, ApJ, 698, 895, doi: 10.1088/0004-637X/698/1/895
- Kelly, B. C., Treu, T., Malkan, M., Pancoast, A., & Woo,
 J.-H. 2013, ApJ, 779, 187,
 doi: 10.1088/0004-637X/779/2/187
- King, A. L., Martini, P., Davis, T. M., et al. 2015, MNRAS, 453, 1701, doi: 10.1093/mnras/stv1718
- Kollatschny, W., & Fricke, K. J. 1985, A&A, 146, L11
- Koratkar, A. P., & Gaskell, C. M. 1991, ApJS, 75, 719, doi: 10.1086/191547
- Korista, K. T., & Goad, M. R. 2001, ApJ, 553, 695, doi: 10.1086/320964
- Kovačević, A. B., Ilić, D., Jankov, I., et al. 2021a, arXiv e-prints, arXiv:2105.12420.
 - https://arxiv.org/abs/2105.12420
- Kovačević, A. B., Ilić, D., Popović, L. Č., et al. 2021b, MNRAS, 505, 5012, doi: 10.1093/mnras/stab1595
- Kozłowski, S. 2017, A&A, 597, A128, doi: 10.1051/0004-6361/201629890
- Kozłowski, S., Kochanek, C. S., Udalski, A., et al. 2010, ApJ, 708, 927, doi: 10.1088/0004-637X/708/2/927
- Landt, H., Elvis, M., Ward, M. J., et al. 2011, MNRAS, 414, 218, doi: 10.1111/j.1365-2966.2011.18383.x
- Lawther, D., Goad, M. R., Korista, K. T., Ulrich, O., & Vestergaard, M. 2018, MNRAS, 481, 533, doi: 10.1093/mnras/sty2242
- Li, Y.-R., Wang, J.-M., & Bai, J.-M. 2016, ApJ, 831, 206, doi: 10.3847/0004-637X/831/2/206
- Liu, H. T., Bai, J. M., Zhao, X. H., & Ma, L. 2008, ApJ, 677, 884, doi: 10.1086/529361
- Liu, H.-Y., Liu, W.-J., Dong, X.-B., et al. 2019, ApJS, 243, 21, doi: 10.3847/1538-4365/ab298b
- Lu, K.-X., Huang, Y.-K., Zhang, Z.-X., et al. 2019, ApJ, 877, 23, doi: 10.3847/1538-4357/ab16e8

MacLeod, C. L., Ivezić, Ž., Kochanek, C. S., et al. 2010, ApJ, 721, 1014, doi: 10.1088/0004-637X/721/2/1014

- MacLeod, C. L., Morgan, C. W., Mosquera, A., et al. 2015, ApJ, 806, 258, doi: 10.1088/0004-637X/806/2/258
- Mastroserio, G., Ingram, A., & van der Klis, M. 2018, MNRAS, 475, 4027, doi: 10.1093/mnras/sty075
- Matthews, T. A., & Sandage, A. R. 1963, ApJ, 138, 30, doi: 10.1086/147615
- McHardy, I. M., Cameron, D. T., Dwelly, T., et al. 2014, MNRAS, 444, 1469, doi: 10.1093/mnras/stu1636
- Moravec, E., Svoboda, J., Borkar, A., et al. 2022, A&A, 662, A28, doi: 10.1051/0004-6361/202142870
- Morgan, C. W., Kochanek, C. S., Morgan, N. D., & Falco,
 E. E. 2010, ApJ, 712, 1129,
 doi: 10.1088/0004-637X/712/2/1129
- Morgan, C. W., Hainline, L. J., Chen, B., et al. 2012, ApJ, 756, 52, doi: 10.1088/0004-637X/756/1/52
- Morganti, R. 2017, Nature Astronomy, 1, 596, doi: 10.1038/s41550-017-0223-0
- Mosquera, A. M., Kochanek, C. S., Chen, B., et al. 2013, ApJ, 769, 53, doi: 10.1088/0004-637X/769/1/53
- Mudd, D., Martini, P., Zu, Y., et al. 2018, ApJ, 862, 123, doi: 10.3847/1538-4357/aac9bb
- Mushotzky, R. F., Edelson, R., Baumgartner, W., & Gandhi, P. 2011, ApJL, 743, L12, doi: 10.1088/2041-8205/743/1/L12

Netzer, H. 2013, The Physics and Evolution of Active Galactic Nuclei

- Neustadt, J. M. M., & Kochanek, C. S. 2022, arXiv e-prints, arXiv:2201.10565. https://arxiv.org/abs/2201.10565
- O'Brien, P. T., Dietrich, M., Leighly, K., et al. 1998, ApJ, 509, 163, doi: 10.1086/306464
- Oknyansky, V. L., Winkler, H., Tsygankov, S. S., et al. 2019, MNRAS, 483, 558, doi: 10.1093/mnras/sty3133
- Padovani, P., Alexander, D. M., Assef, R. J., et al. 2017, A&A Rv, 25, 2. doi: 10.1007/s00159-017-0102-9

Panda, S. 2022, arXiv e-prints, arXiv:2206.10056. https://arxiv.org/abs/2206.10056

- Panda, S., Czerny, B., Adhikari, T. P., et al. 2018, ApJ, 866, 115, doi: 10.3847/1538-4357/aae209
- Panda, S., Martínez-Aldama, M. L., & Zajaček, M. 2019, Frontiers in Astronomy and Space Sciences, 6, 75, doi: 10.3389/fspas.2019.00075
- Penton, A., Malik, U., Davis, T. M., et al. 2022, MNRAS, 509, 4008, doi: 10.1093/mnras/stab3027
- Peterson, B. M. 2006, The Broad-Line Region in Active Galactic Nuclei, ed. D. Alloin, Vol. 693, 77, doi: 10.1007/3-540-34621-X 3

- Peterson, B. M., Ferrarese, L., Gilbert, K. M., et al. 2004, ApJ, 613, 682, doi: 10.1086/423269
- Pooley, D., Blackburne, J. A., Rappaport, S., & Schechter,
 P. L. 2007, ApJ, 661, 19, doi: 10.1086/512115
- Pooley, D., Blackburne, J. A., Rappaport, S., Schechter,
 P. L., & Fong, W.-f. 2006, ApJ, 648, 67,
 doi: 10.1086/505860
- Poutanen, J., & Beloborodov, A. M. 2006, MNRAS, 373, 836, doi: 10.1111/j.1365-2966.2006.11088.x
- Pozo Nuñez, F., Ramolla, M., Westhues, C., et al. 2015, A&A, 576, A73, doi: 10.1051/0004-6361/201525910
- Pozo Nuñez, F., Gianniotis, N., Blex, J., et al. 2019, MNRAS, 490, 3936, doi: 10.1093/mnras/stz2830
- Press, W. H., Teukolsky, S. A., Vetterling, W. T., & Flannery, B. P. 1992, Numerical recipes in C. The art of scientific computing
- Raiteri, C. M., Carnerero, M. I., Balmaverde, B., et al. 2022, ApJS, 258, 3, doi: 10.3847/1538-4365/ac3bb0
- Reuter, M. A., Cook, K. H., Delgado, F., Petry, C. E., & Ridgway, S. T. 2016, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 9911, Modeling, Systems Engineering, and Project Management for Astronomy VI, ed. G. Z. Angeli & P. Dierickx, 991125, doi: 10.1117/12.2232680
- Roberts, D. H., Lehar, J., & Dreher, J. W. 1987, AJ, 93, 968, doi: 10.1086/114383
- Sahu, N., Graham, A. W., & Davis, B. L. 2019a, ApJ, 887, 10, doi: 10.3847/1538-4357/ab50b7
- ---. 2019b, ApJ, 876, 155, doi: 10.3847/1538-4357/ab0f32
- Sánchez-Sáez, P., Lira, P., Mejía-Restrepo, J., et al. 2018, ApJ, 864, 87, doi: 10.3847/1538-4357/aad7f9
- Scargle, J. D. 1982, ApJ, 263, 835, doi: 10.1086/160554
- Schmidt, M. 1963, Nature, 197, 1040,
- doi: 10.1038/1971040a0
- Scolnic, D. M., Lochner, M., Gris, P., et al. 2018, arXiv e-prints, arXiv:1812.00516.

https://arxiv.org/abs/1812.00516

Sergeev, S. G., Doroshenko, V. T., Golubinskiy, Y. V., Merkulova, N. I., & Sergeeva, E. A. 2005, ApJ, 622, 129, doi: 10.1086/427820

Shakura, N. I., & Sunyaev, R. A. 1973, A&A, 500, 33

- Shappee, B. J., Prieto, J. L., Grupe, D., et al. 2014, ApJ, 788, 48, doi: 10.1088/0004-637X/788/1/48
- Shemmer, O., Richards, G., Brandt, N., et al. 2018, LSST AGN Science Collaboration Roadmap
- Shen, X., Hopkins, P. F., Faucher-Giguère, C.-A., et al. 2020, MNRAS, 495, 3252, doi: 10.1093/mnras/staa1381
- Shen, Y., Richards, G. T., Strauss, M. A., et al. 2011, ApJS, 194, 45, doi: 10.1088/0067-0049/194/2/45

- Smith, K. L., Mushotzky, R. F., Boyd, P. T., et al. 2018, ApJ, 857, 141, doi: 10.3847/1538-4357/aab88d
- Srinivasan, R., Tucker, D. M., & Murias, M. 1998, Behavior Research Methods, Instruments, & Computers, 30, 11, doi: 10.3758/BF03209412
- Starkey, D., Horne, K., Fausnaugh, M. M., et al. 2017, ApJ, 835, 65, doi: 10.3847/1538-4357/835/1/65
- Starkey, D. A., Horne, K., & Villforth, C. 2016, MNRAS, 456, 1960, doi: 10.1093/mnras/stv2744
- Stone, Z., Shen, Y., Burke, C. J., et al. 2022, arXiv e-prints, arXiv:2201.02762. https://arxiv.org/abs/2201.02762
- Suberlak, K. L., Ivezić, Ż., & MacLeod, C. 2021, ApJ, 907, 96, doi: 10.3847/1538-4357/abc698
- Sun, W.-H., & Malkan, M. A. 1989, ApJ, 346, 68, doi: 10.1086/167986
- Tachibana, Y., Graham, M. J., Kawai, N., et al. 2020, ApJ, 903, 54, doi: 10.3847/1538-4357/abb9a9
- Tekhanovich, D. I., & Baryshev, Y. V. 2016, Astrophysical Bulletin, 71, 155, doi: 10.1134/S1990341316020048
- Tewes, M., Courbin, F., & Meylan, G. 2013, A&A, 553, A120, doi: 10.1051/0004-6361/201220123
- Thater, S., Krajnović, D., Cappellari, M., et al. 2019, A&A, 625, A62, doi: 10.1051/0004-6361/201834808
- Thorne, K. S. 1974, ApJ, 191, 507, doi: 10.1086/152991

- Vasudevan, R. V., Mushotzky, R. F., Winter, L. M., & Fabian, A. C. 2009, MNRAS, 399, 1553, doi: 10.1111/j.1365-2966.2009.15371.x
- Yu, W., & Richards, G. 2021, LSST AGN SC Cadence Note: Non-Parametric Structure Function Metric
- Yu, W., Richards, G. T., Vogeley, M. S., Moreno, J., & Graham, M. J. 2022, arXiv e-prints, arXiv:2201.08943. https://arxiv.org/abs/2201.08943
- Yu, Z., Martini, P., Davis, T. M., et al. 2020, ApJS, 246, 16, doi: 10.3847/1538-4365/ab5e7a
- Yuan, F., & Narayan, R. 2014, ARA&A, 52, 529, doi: 10.1146/annurev-astro-082812-141003
- Zou, F., Brandt, W. N., Chen, C.-T., et al. 2022, arXiv e-prints, arXiv:2206.06432. https://arxiv.org/abs/2206.06432
- Zu, Y., Kochanek, C. S., Kozłowski, S., & Peterson, B. M. 2016, ApJ, 819, 122, doi: 10.3847/0004-637X/819/2/122
- Zu, Y., Kochanek, C. S., Kozłowski, S., & Udalski, A. 2013, ApJ, 765, 106, doi: 10.1088/0004-637X/765/2/106
- Zu, Y., Kochanek, C. S., & Peterson, B. M. 2011, ApJ, 735, 80, doi: 10.1088/0004-637X/735/2/80
- Zubovas, K., & King, A. 2021, MNRAS, 501, 4289, doi: 10.1093/mnras/stab004
- Čvorović Hajdinjak, I., Kovačević, A. B., Ilić, D., et al. 2022, Astronomische Nachrichten, 343, e210103, doi: https://doi.org/10.1002/asna.20210103

APPENDIX

A. ATLAS OF METRIC AGN_TIMELAGMETRIC SIMULATION



Figure A1. Metric realization for cadences from baseline and carina observing strategies for observed time lags of 1,2,3, and 5 days. The colorbar displays metric values above the Nyquist threshold, which have been scaled for improved visual presentation in the Aitoff projection of celestial coordinates.



Figure A2. The same as Figure A1 but for euclid_dither1 and euclid_dither3 observing strategies. In all metric simulations, three DDFs stand out. For better color visibility in the right column, the metric threshold is multiplied by XXX.



Figure A3. Metric realization from footprint observing strategies. There is only one prominent field in each panel (except panel (g)) slightly varying in position on the sky.



Figure A4. Metric realization for cadences from footprint observing strategies, continued. In contrast to Figure A3, DDFs appear in panels (e) and (f).



Figure A5. Metric realizations on cadences from $long_gaps_nightsoff$ observing strategy. The first row of panels shows the fields (in the third and fourth area from the left of the main meridian) suitable for calculating 1- 3 day delays. DDFs appear in the bottom row with better metric realization in *i*-band then in *r*-band.



Figure A6. Metric realization on multishort cadences. DDFs appear in panel (l).



Figure A7. Metric realization on multishort cadences continued. DDFs appear in all panels except (b), (c) and (g).



41



Figure A8. Metric realization on rolling_all_sky_ns2 cadences. Panels (a)-(c) show the fields (in the fourth area from the left of the main meridian) suitable for calculating 1- 3 day delays. DDFs appear in panels (d) and (e).



Figure A9. Metric realization on rolling_all_sky_ns2 cadences continued. DDFs appear in all panels.



Figure A10. Metric realization on short_exp cadences. Metric run emphasizes two fields in u-band (panels (c), (g) and (k)).



Figure A11. Metric realization on short_exp cadences continued. Three DDFs are prominent in panels (b) and (c).



Figure A12. Metric realization on six_rolling and u_long_ms_30 cadences. DDFs are prominent in i band (panels (a) and (e)) and r band (panels (b) and (g)).



Figure A13. Co-added depth in deep drilling fields for different observing strategies in u, g, r, i filters. The best strategy for AD reverberation mapping is also the deepest in all bands.



Figure A14. The same as Fig A13, but for z, y filters.