

Juggler’s friezes

Roi Docampo and Greg Muller

ABSTRACT. This note generalizes $\mathrm{SL}(k)$ -friezes to configurations of numbers in which one of the boundary rows has been replaced by a ragged edge (described by a *juggling function*). We provide several equivalent definitions/characterizations of these **juggler’s friezes**, in terms of determinants, linear recurrences, and a dual juggler’s frieze. We generalize classic results, such as periodicity, duality, and a parametrization by part of a Grassmannian. We also provide a method of constructing such friezes from certain $k \times n$ matrices using the *twist* of a matrix defined in [MS16].

$\mathrm{SL}(k)$ -friezes are configurations of numbers in which certain diamonds have determinant 1 or 0 (when regarded as matrices). The theory began in [Cox71], when Coxeter introduced what would now be called $\mathrm{SL}(2)$ -friezes. This was followed by the (relatively) classical theory of $\mathrm{SL}(k)$ -friezes, including periodicity and many enumerative results [Cox71, CR72, CC73, Sha84].

The study of friezes recently resurged after a connection to cluster algebras was found in [CC06]. This fueled a number of generalizations and theorems; most notably, that $\mathrm{SL}(k)$ -friezes of height h are parametrized by a certain subvariety of the Grassmannian $\mathrm{Gr}(k, k + h)$ [MGOST14], and each cluster on this Grassmannian determines a unique positive integral frieze [BFG⁺21]. Many results on clusters can then be translated into friezes, such as enumerative results and mutation rules; see [MG15, BFG⁺18]. However, the Grassmannian has a much richer structure to exploit: the Grassmannian has a stratification into *positroid varieties*, each of which has a cluster structure. The subvariety and cluster structure corresponding to $\mathrm{SL}(k)$ -friezes is the ‘big cell’ in this stratification.

In this paper, we extend this connection to other positroid varieties by introducing a generalization of $\mathrm{SL}(k)$ -friezes, dubbed **juggler’s friezes** for the *juggling functions* that give them their shape. We prove that juggler’s friezes enjoy analogs of many of the vital properties of $\mathrm{SL}(k)$ -friezes.

- Juggler’s friezes are periodic (Theorem C).
- Juggler’s friezes are parametrized by the unimodular positroid variety (Theorem G).
- Juggler’s friezes are linear recurrences with superperiodic solutions (Theorem B).
- Juggler’s friezes are characterized by a dual transform (Theorem A).

Sections 1 and 2 review the theory of $\mathrm{SL}(k)$ -friezes, highlighting the results and techniques we later generalize. One original result here is a reformulation of the parametrization of $\mathrm{SL}(k)$ -friezes by the Grassmannian in terms of the *twist* of a matrix (Theorem 2.6).

Sections 3 and 4 introduce juggler’s friezes and review their aforementioned properties. The exposition intentionally mirrors the preceding sections, to highlight the connection between $\mathrm{SL}(k)$ -friezes and the broader theory of juggler’s friezes. Proofs are deferred until Section 5. While we describe the connection between juggler’s friezes and cluster algebras in Section 4.4, proofs of these results exceed the scope of this paper, and will appear in a subsequent paper [DM].

2020 *Mathematics Subject Classification.* Primary 05E99; Secondary 39A24, 14M15, 13F60.

Key words and phrases. Friezes, juggling functions, Grassmannians, positroid cells.

The research of the first author was partially supported by a grant from the Simons Foundation (638459, RD).

Contents

1. $\mathrm{SL}(k)$ -friezes	3
1.1. A breezy review of $\mathrm{SL}(k)$ -friezes	3
1.2. (Pre)friezes as matrices	4
1.3. Friezes and duality	5
1.4. Friezes as linear recurrences	6
2. Constructing $\mathrm{SL}(k)$ -friezes	7
2.1. Constructing friezes	7
2.2. Relation to the Grassmannian	10
3. Juggler's friezes	11
3.1. Juggling functions	11
3.2. π -prefriezes	12
3.3. Juggler's friezes	13
3.4. Visualizing the frieze condition	14
3.5. Visualizing the tameness condition	15
3.6. Recovering $\mathrm{SL}(k)$ -friezes	15
3.7. Juggler's friezes and duality	17
3.8. Juggler's friezes as linear recurrences	17
4. Constructing juggler's friezes	18
4.1. π -unimodular matrices	18
4.2. Constructing juggler's friezes	19
4.3. Relation to the Grassmannian	21
4.4. Positivity, enumeration, and clusters	21
Intermission: Computing a dual juggler's frieze via the twist	22
5. Proofs of Theorems	24
5.1. Duality for juggler's friezes (Theorem A)	24
5.2. Superperiodicity of solutions (Theorems B, C, and D)	26
5.3. Equivalence of constructions (Theorem E)	31
5.4. $F(\mathbf{A})$ is a π^\dagger -frieze (Theorem F)	32
5.5. Inverting F (Theorem G)	34
5.6. Duality and the twist (Theorem H)	35
Appendix A. Collected notational conventions	38
Acknowledgements	38
References	38

1. $SL(k)$ -friezes

1.1. A breezy review of $SL(k)$ -friezes. For a positive integers h and k , a $SL(k)$ -frieze of height h consists of $(h + 1)$ -many rows of numbers,¹ offset in a diamond grid, such that

- the top and bottom rows consist entirely of 1s,
- each solid $k \times k$ -minor is 1 (the **frieze** condition), and
- each solid $(k + 1) \times (k + 1)$ -minor is 0 (the **tameness** condition).

By a *solid minor*, we mean the determinant of a diamond of entries, regarded as a matrix by rotating 45° clockwise.²

Remark 1.1. Note that, by our definition, **every $SL(k)$ -frieze is tame**. Most authors define ‘friezes’ without the tameness condition, and use the tameness condition to define *tame friezes*. \triangle



Warning 1.2. The **height** h counts the gaps between rows, not the number of rows themselves!³ \triangle

Example 1.3. An $SL(2)$ -frieze of height 6.

	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
...	3	2	2	1	4	3	1	2	3	2	2	1	4	3	1	2	3	...	
	5	5	3	1	3	11	2	1	5	5	3	1	3	11	2	1	5	5	
...	8	7	1	2	8	7	1	2	8	7	1	2	8	7	1	2	8	...	
	3	11	2	1	5	5	3	1	3	11	2	1	5	5	3	1	3	11	
...	4	3	1	2	3	2	2	1	4	3	1	2	3	2	2	1	4	...	
	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	

Note that the **left submatrix** has determinant 1, and the **right submatrix** has determinant 0. \triangle

Example 1.4. An $SL(3)$ -frieze of height 5.

	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
...	3	3	3	1	18	1	5	2	3	3	3	1	18	1	5	2	3	...	
	3	6	5	2	7	16	1	8	3	6	5	2	7	16	1	8	3	6	
...	4	7	3	4	6	9	1	10	4	7	3	4	6	9	1	10	4	...	
	11	2	4	2	3	3	4	1	11	2	4	2	3	3	4	1	11	2	
...	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	

Coxeter [Cox71] introduced $SL(2)$ -friezes (called *frieze patterns* at the time) to understand and generalize the *pentagramma mirificum*, a recurrence observed by Gauss among the side lengths of a right-angled spherical pentagram, and showed they were always $(h + 2)$ -periodic (observe that Example 1.3 is 8-periodic). More generally, an $SL(k)$ -frieze of height h are $(h + k)$ -periodic [BR10, MGOST14, KV15] (observe that Example 1.4 is 8-periodic).

The first enumerative result was Coxeter-Conway [CC73], who showed that the positive integral $SL(2)$ -friezes of height h are in bijection with triangulations of an $(h + 2)$ -gon, and are therefore counted by a Catalan number. Despite this, there are infinitely many positive integral $SL(k)$ -friezes of height h except when $\min(k, h) \leq 2$, or $\min(k, h) = 3$ and $\max(k, h) \leq 6$ (see e.g. [BFG⁺21]).

¹By ‘numbers’, we mean elements in a fixed field \mathbb{k} , although all our examples will have values in the integers.
²FINE PRINT: The empty entries above and below the frieze are treated as 0s, and we only impose the minor conditions on diamonds whose center is strictly between the top and bottom rows. A more precise definition is in Definition 1.7.
³This varies among references. [CC73] call $h + 1$ the *order* of a frieze, and [MG15] calls $h - 1$ the *width* of a frieze.

Definition 1.7. An $SL(k)$ -frieze is a prefrieze C of height h such that

- (1) $\det(C_{[a,a+k-1],[b,b+k-1]}) = 1$ whenever $b \leq a \leq b + h$, and
- (2) $\det(C_{[a,a+k],[b,b+k]}) = 0$ whenever $b < a < b + h$.

1.3. Friezes and duality. We begin our alternate characterizations of friezes by observing that, instead of considering solid minors of size k and $k + 1$, one may instead consider solid minors centered on the second row with sizes between k and $k + h$.

Lemma 1.8. A prefrieze of height h is a $SL(k)$ -frieze iff

- (1) $\det(C_{[a+1,a+k],[a,a+k-1]}) = 1$ for all $a \in \mathbb{Z}$, and
- (2) $\det(C_{[a+1,a+\ell],[a,a+\ell-1]}) = 0$ for all $a \in \mathbb{Z}$ and all ℓ with $k < \ell \leq k + h$.

Proof sketch. A more general proof is given by Lemma 5.3, but we sketch a simpler argument here.

One direction follows from the observation that, if every solid $k \times k$ -minor of an $\ell \times \ell$ matrix is 1, then the determinant of that matrix is equal to the determinant of the matrix of its solid $(k + 1) \times (k + 1)$ -minors. This is a corollary of the Dodgson condensation algorithm, which ‘resets’ at the $(k + 1)$ th step if all the $k \times k$ -minors are 1. This observation implies that any $\ell \times \ell$ -minor of a $SL(k)$ frieze is 0, provided its $k \times k$ -minors stay in the region where they are forced to be 1.

The other direction follows by considering the adjugate matrix $\text{Adj}(C)$ of the matrix corresponding to C .⁵ The frieze and tameness conditions translate to $\text{Adj}(C)$ via the identities

$$\det(C_{[a,a+k-1],[b,b+k-1]}) = \pm \det(\text{Adj}(C)_{[b+k,a+k-1],[b,a-1]})$$

$$\det(C_{[a,a+k],[b,b+k]}) = \pm \det(\text{Adj}(C)_{[b+k+1,a+k],[b,a-1]})$$

Conditions (1) and (2) in the statement of the lemma imply the determinants on the right-hand-side are respectively upper unitriangular and strictly upper triangular (up to sign). After checking the signs work out, this shows that C satisfies the frieze and tameness conditions. \square

This lemma may seem arbitrary, but we can rephrase it in a more pleasing form by collecting all sufficiently small solid minors of C centered on the second row into a single object. Given a prefrieze C and a positive integer n , define the n -truncated dual C^\dagger to be the diamond grid of numbers (or equivalently, the $\mathbb{Z} \times \mathbb{Z}$ -matrix) defined by

$$C_{a,b}^\dagger := \begin{cases} \det(C_{[b+1,a],[b,a-1]}) & \text{if } b \leq a < b + n, \\ 0 & \text{otherwise.} \end{cases}$$

When $a = b$, the corresponding determinant is empty and is defined to be 1.

Example 1.9. The 8-truncated dual C^\dagger of Example 1.4 is below.

$$\begin{array}{cccccccccccccccccccc} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ \cdots & 3 & 3 & 3 & 1 & 18 & 1 & 5 & 2 & 3 & 3 & 3 & 1 & 18 & 1 & 5 & 2 & 3 & \cdots \\ & 3 & 3 & 4 & 1 & 11 & 2 & 4 & 2 & 3 & 3 & 4 & 1 & 11 & 2 & 4 & 2 & 3 & 3 \\ \cdots & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \cdots \end{array}$$

⁵While an infinite matrix may not have an adjugate, a unitriangular matrix does; see Lemma 5.5.

The entries depicted above are only the non-zero ones; there are infinitely many implicit rows of 0 above and below the strip above. Four of these rows correspond to solid minors of \mathbf{C} of sizes 4 to 7 that vanish; the rest are zero by the definition of \mathbf{C}^\dagger . \triangle

The minors in Lemma 1.8 are entries of \mathbf{C}^\dagger , and the conditions on these minors translate to the requirement that the k th row of \mathbf{C}^\dagger consists of 1s, and all lower rows consist of 0s; that is, that \mathbf{C}^\dagger is a prefrieze of height k .

Lemma 1.10. *A prefrieze \mathbf{C} of height h is an $\mathrm{SL}(k)$ -frieze iff the $(h+k)$ -truncated dual \mathbf{C}^\dagger is a prefrieze of height k .*

To reap the benefits of this result, note that $(\mathbf{C}^\dagger)^\dagger = \mathbf{C}$ whenever \mathbf{C} is a prefrieze of height less than n .⁶ Therefore, if \mathbf{C} is an $\mathrm{SL}(k)$ -frieze of height h , then the $(h+k)$ -truncated dual of \mathbf{C}^\dagger is a prefrieze of height k , and so \mathbf{C}^\dagger is an $\mathrm{SL}(h)$ -frieze of height k . We can restate this as follows.

Theorem 1.11. *Taking $(h+k)$ -truncated duals defines mutually inverse bijections between the sets of $\mathrm{SL}(k)$ -friezes of height h and $\mathrm{SL}(h)$ -friezes of height k .*

Remark 1.12. The frieze \mathbf{C}^\dagger differs from the *combinatorial Gale dual* of \mathbf{C} defined in [MGOST14] by a vertical reflection (called *projective duality* in *loc. cit.*) and a horizontal shift. \triangle

1.4. Friezes as linear recurrences. Given a prefrieze \mathbf{C} of height h , we may consider the matrix equation $\mathbf{C}\mathbf{x} = 0$, where $\mathbf{x} = (\dots, x_{-1}, x_0, x_1, x_2, \dots)$ denotes a vector of variables of length \mathbb{Z} . This is equivalent to a system of linear equations of the form

$$x_{a-h} + C_{a,a-h+1}x_{a-h+1} + C_{a,a-h+2}x_{a-h+2} + \cdots + C_{a,a-1}x_{a-1} + x_a = 0$$

running over all $a \in \mathbb{Z}$. Each such equation can be rewritten as a formula for x_a as a linear combination of earlier variables:

$$x_a = -C_{a,a-1}x_{a-1} - C_{a,a-2}x_{a-2} - \cdots - C_{a,a-h+1}x_{a-h+1} - x_{a-h}$$

For this reason, we call the matrix equation $\mathbf{C}\mathbf{x} = 0$ a **linear recurrence**.⁷

Theorem 1.13. [MGOST14] *A prefrieze \mathbf{C} of height h is a $\mathrm{SL}(k)$ -frieze iff every solution to the linear recurrence $\mathbf{C}\mathbf{x} = 0$ satisfies $x_{a+h+k} = (-1)^{k-1}x_a$ for all $a \in \mathbb{Z}$.*

Remark 1.14. [MGOST14] uses an alternating sign convention when associating a linear recurrence to a frieze, which we do not. As a result, the sign $(-1)^{k-1}$ appears different than in *loc. cit.* \triangle

The solutions to the linear recurrence $\mathbf{C}\mathbf{x} = 0$ are closely related to the dual \mathbf{C}^\dagger , as follows.

Theorem 1.15. *Let \mathbf{C} be a $\mathrm{SL}(k)$ -frieze of height h . For each b , the sequence \mathbf{x} defined by*

- $x_a := (-1)^{a+b}C_{a,b}^\dagger$ when $b \leq a < b+k+h$, and
- $x_{a+h+k} = (-1)^{k-1}x_a$ for all a

is a solution to $\mathbf{C}\mathbf{x} = 0$. Running over all b , these solutions span the space of solutions to $\mathbf{C}\mathbf{x} = 0$.

⁶This follows from the corresponding property of the adjugate of a matrix, or as a special case of Lemma 5.7.

⁷Also called a *linear recurrence relation* or a *system of linear difference equations*.

Conceptually, each northwest-southeast diagonal of C^\dagger determines a solution to $Cx = 0$ by first flipping half the signs and then extending superperiodically. This theorem is essentially [MGOST14, Prop. 5.1.1], and a special case of Theorem D.

Example 1.16. If C is the frieze in Example 1.4, then two solutions to $Cx = 0$ are

$$\begin{aligned} & \cdots, 0, 1, -3, 3, -1, 0, 0, 0, 0, 1, -3, 3, -1, 0 \cdots \\ & \cdots, 0, 0, 1, -4, 3, -1, 0, 0, 0, 0, 1, -4, 3, -1, \cdots \end{aligned}$$

Observe that they each satisfy the condition that $x_{a+8} = x_a$ (since $k - 1 = 2$). △

2. Constructing $SL(k)$ -friezes

While we now know several methods to recognize $SL(k)$ -friezes, our story lacks a method to construct them, which we remedy now. We give two equivalent constructions, and observe that this parametrizes $SL(k)$ -friezes of height h by a subvariety of the Grassmannian $Gr(h, k + h)$.

Notation 2.1. Given a $k \times n$ matrix A and $a \in \mathbb{Z}$, let A_a denote the \bar{a} th column of A , where $\bar{a} \equiv a \pmod n$. If $I \subset \mathbb{Z}$, let A_I denote the submatrix of A consisting of columns indexed by $I \pmod n$.⁸ △

2.1. Constructing friezes. A $k \times n$ -matrix A is **consecutively unimodular** if $\det(A_{[a, a+k-1]}) = 1$ for all $a \in \mathbb{Z}$; i.e. every submatrix on k -many cyclically consecutive⁹ columns has determinant 1.

Example 2.2. The following 3×8 matrix is consecutively unimodular, which can be verified by checking the six solid 3×3 -minors and the minors on columns 178 and 128.

$$A := \begin{bmatrix} 1 & 11 & 4 & 6 & 3 & 1 & 0 & 0 \\ 0 & 1 & 2 & 7 & 5 & 3 & 1 & 0 \\ 0 & 0 & 1 & 4 & 3 & 2 & 1 & 1 \end{bmatrix} \quad \triangle$$

Theorem 2.3 ([BFG⁺21]). *Each consecutively unimodular $k \times n$ -matrix A determines an $SL(k)$ -frieze $F(A)$ of height $h = n - k$, defined by*

$$F(A)_{a,b} = \begin{cases} \det(A_{[a+1, a+k-1] \cup \{b\}}) & \text{if } b \leq a \leq b + n - k, \\ 0 & \text{otherwise.} \end{cases}$$

The construction in Theorem 2.3 *a priori* requires computing $k(n - k)$ -many $k \times k$ -determinants, which is a daunting task even for small examples like Example 2.2. We can simplify this construction considerably by means of the *twist* of a matrix. The **(right) twist** of a consecutively unimodular $k \times n$ -matrix A is the $k \times n$ -matrix $\tau(A)$ whose columns are defined by the dot product equations

$$A_a \cdot \tau(A)_b = \begin{cases} 1 & \text{if } a = b, \\ 0 & \text{if } b < a < b + k. \end{cases}$$

Since the matrix is consecutively unimodular, the columns of A indexed by $[b, b + k - 1]$ are linearly independent, and so the above system uniquely determines each column of $\tau(A)$. Many interesting

⁸The columns of A_I remain in the same order as in A ; e.g. if the width $n = 5$, then $A_{\{4,5,6\}}$ would consist of the 1st, 4th, and 5th columns of A , in that order.

⁹A *cyclically consecutive* set is the image of a consecutive set under the quotient $\mathbb{Z} \rightarrow \mathbb{Z}/n\mathbb{Z}$.

properties of the twist can be found in [MS17, Section 6]; one relevant result is that the twist of a consecutively unimodular matrix is again consecutively unimodular [MS17, Prop. 6.6].



Warning 2.4. The twist may be defined more generally for $k \times n$ -matrices of rank k , but the definition must be generalized to ensure a linearly independent set of columns; see Section 4.2. \triangle

Remark 2.5. The twist of a matrix in this context was introduced in [MS16] and a normalized version was given in [MS17]. Since the two definitions differ by a factor of the determinant of cyclically consecutive columns; they coincide for a consecutively unimodular matrix. \triangle

Using the twist, we can describe an algorithm (Construction 2.8, on the following page) for constructing an infinite strip of numbers from a consecutively unimodular matrix. As the following theorem asserts, this construction produces the same $SL(k)$ -frieze as Theorem 2.3.

Theorem 2.6. *If A is a consecutively unimodular $k \times n$ -matrix, Construction 2.8 produces $F(A)$, the same $SL(k)$ -frieze as Theorem 2.3.*

Proof. Let C be the output of the construction, let $a \leq b < a + n$, and let \bar{a}, \bar{b} denote the residues of $a, b \bmod n$. Then steps (3) and (4) of the construction are equivalent to the following conditional.

$$\begin{aligned} C_{a,b} &= \begin{cases} (\tau(A)^\top A)_{\bar{a},\bar{b}} & \text{if } \bar{a} \geq \bar{b} \\ (-1)^{k-1} (\tau(A)^\top A)_{\bar{a},\bar{b}} & \text{if } \bar{a} < \bar{b} \end{cases} \\ &= \begin{cases} \tau(A)_{\bar{a}} \cdot A_{\bar{b}} & \text{if } \bar{a} \geq \bar{b} \\ (-1)^{k-1} \tau(A)_{\bar{a}} \cdot A_{\bar{b}} & \text{if } \bar{a} < \bar{b} \end{cases} \end{aligned}$$

The \bar{a} th column of $\tau(A)$ is defined to be the unique solution to $A_{[\bar{a},\bar{a}+k-1]}^\top x = e_{\bar{a}}$. Therefore, $\tau(A)_{\bar{a}} = (A_{[\bar{a},\bar{a}+k-1]}^\top)^{-1} e_{\bar{a}}$, and so

$$\tau(A)_{\bar{a}} \cdot A_{\bar{b}} = ((A_{[\bar{a},\bar{a}+k-1]}^\top)^{-1} e_{\bar{a}}) \cdot A_{\bar{b}} = e_{\bar{a}} \cdot (A_{[\bar{a},\bar{a}+k-1]})^{-1} A_{\bar{b}}$$

Equivalently, $\tau(A)_{\bar{a}} \cdot A_{\bar{b}}$ is the \bar{a} th entry of the solution to $A_{[\bar{a},\bar{a}+k-1]} x = A_{\bar{b}}$. By Cramer's Rule, this is the determinant of $A_{[\bar{a},\bar{a}+k-1]}$ after replacing the \bar{a} th column of A by the \bar{b} th column.¹⁰ This is *almost* $\det(A_{[\bar{a}+1,\bar{a}+k-1] \cup \{\bar{b}\}})$; however, our convention for ordering the columns in $\det(A_{[\bar{a}+1,\bar{a}+k-1] \cup \{\bar{b}\}})$ means there is an additional factor of $(-1)^{k-1}$ if $\bar{a} < \bar{b}$. That is,

$$\tau(A)_{\bar{a}} \cdot A_{\bar{b}} = \begin{cases} \det(A_{[\bar{a}+1,\bar{a}+k-1] \cup \{\bar{b}\}}) & \text{if } \bar{a} \geq \bar{b}, \\ (-1)^{k-1} \det(A_{[\bar{a}+1,\bar{a}+k-1] \cup \{\bar{b}\}}) & \text{if } \bar{a} < \bar{b}. \end{cases}$$

Therefore, $C_{a,b} = \det(A_{[\bar{a}+1,\bar{a}+k-1] \cup \{\bar{b}\}})$. Since this is 0 whenever $\bar{b} \in [\bar{a} + 1, \bar{a} + k - 1]$, we obtain the formula in Theorem 2.3. \square

Remark 2.7. Theorem 2.6 provides an alternate proof of Theorem 2.3. By Cauchy-Binet,

$$\det(F(A)_{[a,a+k-1],[b,b+k-1]}) = \det((\tau(A)^\top A)_{[a,a+k-1],[b,b+k-1]}) = \det(\tau(A)_{[a,a+k-1]}) \det(A_{[b,b+k-1]})$$

Both of the final determinants are 1, as A and $\tau(A)$ are consecutively unimodular. By a similar argument, any $(k+1) \times (k+1)$ minor of C vanishes, since the product $\tau(A)^\top A$ has rank k . \triangle

¹⁰There is also a denominator of $\det(A_{[\bar{a},\bar{a}+k-1]})$ in Cramer's Rule, but this determinant is 1 by assumption.

Construction 2.8. Let A be a consecutively unimodular $k \times n$ -matrix. As a running example:

$$A := \begin{bmatrix} 1 & 11 & 4 & 6 & 3 & 1 & 0 & 0 \\ 0 & 1 & 2 & 7 & 5 & 3 & 1 & 0 \\ 0 & 0 & 1 & 4 & 3 & 2 & 1 & 1 \end{bmatrix}$$

(1) Compute the twist $\tau(A)$ of A .


$$\tau(A) = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 0 & 0 \\ -11 & -10 & -6 & -3 & -1 & 0 & 1 & 0 \\ 18 & 16 & 9 & 4 & 1 & 0 & 0 & 1 \end{bmatrix}$$

(2) Compute the matrix product $\tau(A)^\top A$.


$$\tau(A)^\top A = \begin{bmatrix} 1 & 0 & 0 & 1 & 2 & 4 & 7 & 18 \\ 1 & 1 & 0 & 0 & 1 & 3 & 6 & 16 \\ 1 & 5 & 1 & 0 & 0 & 1 & 3 & 9 \\ 1 & 8 & 2 & 1 & 0 & 0 & 1 & 4 \\ 1 & 10 & 3 & 3 & 1 & 0 & 0 & 1 \\ 1 & 11 & 4 & 6 & 3 & 1 & 0 & 0 \\ 0 & 1 & 2 & 7 & 5 & 3 & 1 & 0 \\ 0 & 0 & 1 & 4 & 3 & 2 & 1 & 1 \end{bmatrix}$$

Remark 2.9. The entries of $\tau(A)^\top A$ are dot products between columns of $\tau(A)$ and columns of A , so $\tau(A)^\top A$ has a strip of 0s above the diagonal bounded by two diagonals of ± 1 s. \triangle

(3) Take the entries above the diagonal of $\tau(A)^\top A$, multiply them by $(-1)^{k-1}$, and slide them to the left of the rest of the matrix to form a parallelogram.



(4) Rotate the resulting parallelogram 45° counter-clockwise, delete any 0s below the bottom row of 1s, and duplicate the rest n -periodically in an infinite horizontal strip.



2.2. Relation to the Grassmannian. For all $G \in \mathrm{SL}(k)$, if A is consecutively unimodular, then so is GA , and $F(GA) = F(A)$. Therefore, F descends to a well-defined map

$$\mathrm{SL}(k) \backslash \{\text{consecutively unimodular } k \times n\text{-matrices}\} \xrightarrow{F} \{\mathrm{SL}(k)\text{-friezes of height } (n - k)\}$$

Theorem 2.10. *The map F is a bijection.*

Proof sketch. An inverse of F is given by sending C to the submatrix $C_{[n-k+1, n], [1, n]}$. □

This can be related to the Grassmannian $\mathrm{Gr}(k, n)$ of k -planes in n -space, via the bijection

$$\mathrm{GL}(k) \backslash \{k \times n\text{-matrices of rank } k\} \xrightarrow{\text{rowspan}} \mathrm{Gr}(k, n)$$

Restricting this to $\mathrm{SL}(k)$ -orbits of consecutively unimodular matrices gives a closed inclusion

$$\mathrm{SL}(k) \backslash \{\text{consecutively unimodular } k \times n\text{-matrices}\} \xrightarrow{\text{rowspan}} \mathrm{Gr}(k, n)$$

If we let $\mathrm{Gr}_{\mathrm{uni}}(k, n)$ denote the image of this map, then Theorem 2.10 descends to a bijection

$$F : \mathrm{Gr}_{\mathrm{uni}}(k, n) \xrightarrow{\sim} \{\mathrm{SL}(k)\text{-friezes of height } (n - k)\}$$

Remark 2.11. This parametrization of $\mathrm{SL}(k)$ -friezes by a subvariety of a Grassmannian has appeared in [MGOST14, Prop. 3.2.1] and [BFG⁺21, Theorem 3.1]. △

Now that we have introduced the twist of a consecutively unimodular matrix, we can relate it to the $(n$ -truncated) dual of a frieze. Given a $k \times n$ matrix A of rank k , define a **positive complement** of A to be any $(n - k) \times n$ matrix A^\ddagger such that, for every k -element set $I \subset [n]$,

$$\det(A_I) = \det(A^\ddagger_{[n] \setminus I})$$

Such a matrix may be constructed by first finding a matrix B whose rows span the orthogonal complement to the rowspan of A , negating odd numbered columns of B , and then rescaling until the above identities are satisfied. The following facts are straightforward from the definition.

- If A^\ddagger is a positive complement of A , then A is a positive complement of A^\ddagger .
- Positive complements exist and are unique up to left multiplication by $\mathrm{SL}(n - k)$.
- The positive complement of a consecutively unimodular matrix is consecutively unimodular.

Therefore, sending a matrix to its positive complement descends to well-defined isomorphisms

$$\ddagger : \mathrm{Gr}(k, n) \xrightarrow{\sim} \mathrm{Gr}(n - k, n) \text{ and } \ddagger : \mathrm{Gr}_{\mathrm{uni}}(k, n) \xrightarrow{\sim} \mathrm{Gr}_{\mathrm{uni}}(n - k, n)$$

whose inverse is \ddagger . This is related to \ddagger and τ by the following commutative diagram of bijections.

$$\begin{array}{ccccc} & & \mathrm{Gr}_{\mathrm{uni}}(k, n) & \xleftarrow{\ddagger} & \mathrm{Gr}_{\mathrm{uni}}(n - k, n) \\ & \nearrow \tau & & & \nwarrow \tau \\ \mathrm{Gr}_{\mathrm{uni}}(k, n) & & & \xleftarrow{\ddagger} & \mathrm{Gr}_{\mathrm{uni}}(n - k, n) \\ & \searrow \ddagger & & & \swarrow \tau \\ & & \mathrm{Gr}_{\mathrm{uni}}(n - k, n) & \xleftarrow{\ddagger} & \mathrm{Gr}_{\mathrm{uni}}(k, n) \\ & & & & \\ & & \mathrm{F} \downarrow & & \mathrm{F} \downarrow \\ & & \{\mathrm{SL}(k)\text{-friezes of height } (n - k)\} & \xleftarrow{\ddagger} & \{\mathrm{SL}(n - k)\text{-friezes of height } k\} \end{array}$$

The commutativity will be proven as a special case of Theorem H and Proposition 5.21.

3. Juggler’s friezes

Theorems in Sections 3 and 4 are proven in Section 5.

We now describe a generalization of $SL(k)$ -friezes, which we call *juggler’s friezes*, for which analogs of each of the preceding definitions and constructions hold. The simplest motivation for this generalization is to observe that Construction 2.8 can be applied to a more general class of matrices (*π -unimodular matrices*), yielding grids of numbers with certain minors necessarily equal to 0, 1, or -1 . However, instead of starting with this construction, we mirror the order of the preceding exposition: a definition in terms of certain minors, followed by several equivalent characterizations, and then conclude with the analog of Construction 2.8.

3.1. Juggling functions. First, we need a combinatorial object to describe the shape of our generalized friezes. A **juggling function of period**¹¹ n is a bijection $\pi : \mathbb{Z} \rightarrow \mathbb{Z}$ such that, for all i , $i \leq \pi(i) \leq i + n$ and $\pi(i + n) = \pi(i) + n$. Juggling functions are a set of combinatorial objects in bijection with *positroids*; matroids with positive representations (see [KLS13]).

Remark 3.1. A juggling function π can be thought of as a transcription of how a juggler catches and throws balls over time. At each moment a , the juggler catches the ball they threw at moment $\pi^{-1}(a)$ and immediately throws it again to be caught at moment $\pi(a)$...unless $\pi(a) = a$, in which case they neither catch nor throw a ball. △

The **number of balls** in a juggling function of period n is

$$\frac{1}{n} \sum_{i=1}^n (\pi(i) - i)$$

which is always an integer. The **siteswap notation** for a juggling function is the list of integers

$$(\pi(1) - 1, \pi(2) - 2, \dots, \pi(n) - n)$$

A **loop** (resp. a **coloop**) of a juggling function is an $a \in \mathbb{Z}$ such that $\pi(a) = a$ (resp. $\pi(a) = a + n$).¹²

Remark 3.2. Our definitions and theorems will frequently have extra cases for loops or coloops, but these rarely arise in interesting examples and can be safely ignored in a first reading. △

Example 3.3. The juggling function with siteswap notation ‘53635514’ is the function with values

i	1	2	3	4	5	6	7	8
$\pi(i)$	6	5	9	7	10	11	8	12

extended to the rest of \mathbb{Z} via the rule that $\pi(i + 8) = \pi(i) + 8$. △

Given a juggling function π of period n , the **dual juggling function** π^\dagger

$$\pi^\dagger(a) := \pi^{-1}(a) + n$$

is a juggling function with period n . If π has h -many balls, then π^\dagger has $(n - h)$ -many balls.

¹¹We emphasize that the choice of a distinguished period n is part of the data of a juggling function.

¹²This terminology is inherited from the matroid associated to π .

Example 3.4. The dual juggling function of the preceding example is 8-periodic with values

i	1	2	3	4	5	6	7	8
$\pi^\dagger(i)$	3	5	6	8	10	9	12	15

which has siteswap notation ‘23345357’. △

3.2. π -prefriezes. Before defining juggler's friezes, we generalize ‘prefriezes’ to configurations of numbers with a ragged lower edge. Unlike prefriezes in the previous sense, which always had 1s on the boundary, this new object may have 1s or -1 s on the boundary, depending on the parity of the following set. Given a juggling function π and integers $a, b \in \mathbb{Z}$, define

$$S_\pi(a, b) := \{i \in \mathbb{Z} \mid a < i \text{ and } \pi(i) < b\} = (a, b) \cap \pi^{-1}(a, b)$$

Remark 3.5. In the juggling perspective, $S_\pi(a, b)$ is the set of times a ball is thrown after moment a that will be caught before moment b . △

Definition 3.6. Given a juggling function π , a π -**prefrieze** consists of rows of numbers offset in a diamond grid \mathbb{C} with entries indexed as in Section 1.2¹³, such that

$$\mathbb{C}_{a,b} = \left\{ \begin{array}{ll} 1 & \text{if } a = b \\ (-1)^{|S_\pi(b,a)|} & \text{if } a = \pi(b) \\ 0 & \text{if } a \notin [b, \pi(b)] \text{ or } b \notin [\pi^{-1}(a), a] \end{array} \right\}$$

Note that this definition allows $\mathbb{C}_{a,b}$ to be arbitrary if $\pi^{-1}(a) < b < a < \pi(b)$.




Warning 3.7. The period n of the juggling function π is not used in the definition of a π -prefrieze; however, this data will be essential for the *frieze* and *tameness* conditions in the next section. △

Construction 3.8. We outline a more visually-oriented construction of a π -prefrieze. First, construct the **diagram of π** as follows.

- Draw a row of green circles (i.e. at position $\mathbb{C}_{a,a}$ for each $a \in \mathbb{Z}$).
- Draw a blue circle at position $\mathbb{C}_{\pi(a),a}$ for each $a \in \mathbb{Z}$.¹⁴
- Draw a line between each pair of circles which share a diagonal (i.e. in the same row or column of the corresponding matrix).

As a running example, the juggling function π with siteswap notation ‘53635514’ has diagram



The doubled green circle above denotes the position $\mathbb{C}_{1,1}$.


¹³Or, equivalently, a $\mathbb{Z} \times \mathbb{Z}$ -matrix rotated 45° counterclockwise.

¹⁴If $\pi(a) = a$, the circle $\mathbb{C}_{a,a}$ will be both blue and green, with no adjacent lines.

A π -prefrieze can be constructed from the diagram of π by putting:

- a 1 in each green circle,
- a $(-1)^s$ in each blue circle, where s is the number of blue circles in the cone above it, and
- any number at each intersection between lines.

All other entries are implicitly zero. An example of a π -prefrieze on the diagram above is below.



Of the two doubled blue circles, $C_{6,1} = (-1)^1$, as there is a single blue circle in the cone above (at $C_{5,2}$), and $C_{9,3} = (-1)^2$, as there are two blue circles in the cone above (at $C_{7,4}$ and $C_{8,7}$). \triangle

3.3. Juggler’s friezes. We can now state the primary definition of this note: a generalization of friezes to shapes defined by juggling functions.

Definition 3.9. Given a juggler’s function π of period n , a π -**frieze** is a π -prefrieze satisfying the following conditions.

- (the **frieze** condition) For all $a, b \in \mathbb{Z}$ with $\pi^\dagger(a) \leq b < a + n \leq \pi(b)$,

$$\det(C_{[a,b] \setminus I, [a,b] \setminus \pi^\dagger(I)}) = 1$$

where $I := \{i \in \mathbb{Z} \mid a \leq i \text{ and } \pi^\dagger(i) \leq b\}$; note this coincides with $S_{\pi^\dagger(a-1, b+1)}$.

- (the **tameness** condition) For all $a, b \in \mathbb{Z}$ with $\pi^\dagger(a) < b < a + n$ or $b < a + n < \pi(b)$,

$$\det(C_{(a,b] \setminus J, [a,b] \setminus \pi^\dagger(J)}) = 0$$

where $J := \{i \in \mathbb{Z} \mid a < i \text{ and } \pi^\dagger(i) < b\}$; note this coincides with $S_{\pi^\dagger(a, b)}$.

While the determinants in the definition are not solid minors, a diagrammatic construction of these determinants is given in the next two sections. Absent an explicit choice of π , we refer to this genre of friezes as **juggler’s friezes**.

Example 3.10. The running example of a π -prefrieze in Construction 3.8 is, in fact, a π -frieze. \triangle




Warning 3.11. Note that we include tameness in the definition, where others might call these *tame juggler’s friezes*. One reason for this is that *wild juggler’s friezes* (π -prefriezes only satisfying the frieze condition) can be poorly behaved, even compared with wild $SL(k)$ -friezes. E.g. tame π -friezes may not form an irreducible component of the space of wild π -friezes. \triangle


Remark 3.12. If C is a π -prefrieze, then $\det(C_{[a,b] \setminus I, [a,b] \setminus \pi^\dagger(I)}) = 1$ for any a, b with $a \leq b < a + n$; that is, the conditions $\pi^\dagger(a) \leq b$ and $a + n \leq \pi(b)$ may be dropped from the frieze condition. However, this larger class of determinants immediately reduces to the given ones. When $\pi^\dagger(a) \not\leq b$

or $a + n \not\leq \pi(b)$, the matrix $C_{[a,b] \setminus I, [a,b] \setminus \pi^\dagger(I)}$ can be block decomposed into a lower unitriangular matrix and a matrix of the form $C_{[a',b'] \setminus I', [a',b'] \setminus \pi^\dagger(I')}$, where $\pi^\dagger(a') \leq b'$ and $a' + n \leq \pi(b')$.¹⁵ \triangle

3.4. Visualizing the frieze condition. To visualize the frieze condition, first take the diagram of π^\dagger , reflect it vertically, make the blue circles red, and attach it to the π -prefrieze C along the green circles. In the running example:



For each a, b with $b \leq a < b - n$, consider $C_{[a,b], [a,b]}$ (the diamond centered on the green circles indexed by the interval $[a, b]$) and delete any diagonal containing a red circle. The resulting matrix is $C_{[a,b] \setminus I, [a,b] \setminus \pi^\dagger(I)}$ and the frieze condition is that all matrices of this form have determinant 1.




Depicted above are the submatrices corresponding to the intervals $[-2, 1]$, $[2, 6]$, and $[7, 12]$; one may check that all three have determinant 1.

Remark 3.13. This construction makes it easy to check whether $\pi^\dagger(a) \leq b$ and $a + n \leq \pi(b)$. The first inequality (respectively, second inequality) is equivalent to the presence of a red circle along the upper left diagonal (resp. upper right diagonal) of the diamond. Therefore, these inequalities are equivalent to requiring that the upper left and upper right diagonals are deleted.¹⁶ However, as observed in Remark 3.12, ignoring these inequalities does not change the frieze condition. \triangle

¹⁵However, the inequalities in the tameness condition cannot be dropped or even weakened to ‘and’; see Section 3.5.

¹⁶Read as a matrix, this is equivalent to requiring that the top row and the right column of $C_{[a,b], [a,b]}$ are deleted.

3.5. Visualizing the tameness condition. As before, attach the reflected diagram of π^\dagger to the top of the π -prefrieze. For each a, b with $b \leq a < b - n$, consider $C_{[a,b],[a,b]}$ (the diamond centered on the green circles indexed by the interval $[a, b]$) as before. Next, delete the upper left diagonal and the upper right diagonal (regardless of whether they contain a red circle). Then, delete any diagonal containing a red circle in the remaining matrix. The resulting matrix is $C_{(a,b) \setminus J, [a,b] \setminus \pi^\dagger(J)}$, and the tameness condition is that this determinant 0 whenever $\pi^\dagger(a) < b$ or $a + n < \pi(b)$.



Depicted above are the submatrices corresponding to the intervals $[-3, 1]$, $[2, 7]$, and $[8, 12]$; one may check that they have determinants 0, 0, and 1, respectively. The latter determinant does not contradict the tameness condition because it fails both conditions: $\pi^\dagger(8) \not\leq 12$ and $8 + 8 \not\leq \pi(12)$.

Remark 3.14. A pair (a, b) satisfies $\pi^\dagger(a) < b$ or $a + n < \pi(b)$ if the corresponding diamond contains a red circle which is in the upper left diagonal or the upper right diagonal, but not both. Note that the diamond associated to $(8, 12)$ does not have a red circle in either top diagonal. \triangle

3.6. Recovering $SL(k)$ -friezes. While the definition of a juggler’s frieze is less elegant than that of a $SL(k)$ -frieze, the latter is a special case of the former. A juggling function is **uniform** if there is some h for which $\pi(a) = a + h$ for all a ; note that this h is the number of balls of π .



Warning 3.15. With uniform juggling functions, it is particularly important to remember the choice of period is part of the data of a juggling function, since $\pi(a + m) = \pi(a) + m$ for all $m \in \mathbb{Z}$! \triangle

Proposition 3.16. *Let π be a uniform juggling function with h -many balls and period n .*

- (1) *A π -prefrieze is the same as a prefrieze of height h .*
- (2) *A π -frieze is the same as an $SL(n - h)$ -frieze of height h .*

Proof. Since $S_\pi(a, \pi(a)) = S_\pi(a, a + k) = \emptyset$, a π -prefrieze consists of $(h + 1)$ -many rows in a diamond grid, in which the top and bottom rows consist of 1s; that is, a prefrieze of height h .

For any a , $\pi^\dagger(a) = \pi^{-1}(a) + n = a + n - k$; that is, π^\dagger is the uniform juggling function with $(n - h)$ -many balls and period n . Then there is a π -frieze condition for each pair a, b with

$$a + n - k \leq b < a + n \leq b + k$$

which can be rewritten as $a < b - n + k + 1 < a + k$. The corresponding identity in the π -frieze condition is then

$$\det(C_{[b-n+k+1,b],[a,a+n-k-1]}) = 1$$


The π -frieze condition coincides with Condition (1) in Definition 1.7 for $k = n - h$.

As for tameness, observe that $\pi^\dagger(a) < b < a + n$ iff $b < a_n < \pi(b)$. Assuming either holds, the corresponding identity in the tameness condition is


$$\det(C_{[b-n+k,b],[a,a+n-k]}) = 0$$

The tameness condition for π -friezes coincides with Condition (2) in Definition 1.7 for $k = n - h$. \square


Example 3.17. Let π be the uniform 8-periodic juggler function with 5 balls. The $SL(3)$ -frieze of height 5 from Example 1.4 is seen to be a π -prefrieze by drawing the diagram of π .



For this choice of π , the construction in Section 3.4 produces 3×3 solid submatrices.



Similarly, the construction in Section 3.5 produces 4×4 solid submatrices.



We see that this $SL(5)$ -frieze is also a π -frieze. Note that the dashed diamonds above do not satisfy $\pi^\dagger(a) < b$ or $a + n < \pi(b)$, so they do not contribute a minor to the tameness condition. \triangle

Using the dual juggler's frieze and superperiodicity, we can describe a spanning family of solutions to the linear recurrence.

Theorem D. *Let π be an n -periodic juggling function with h -many balls, and let C be a π -frieze. For any b with $b < \pi(b)$, the sequence x defined by*

- $x_a := (-1)^{a+b} C_{a,b}^\dagger$ when $b \leq a < b + k + h$, and
- $x_{a+n} = (-1)^{n-h-1} x_a$ for all a

is a solution to $Cx = 0$. These solutions collectively span the space of solutions to $Cx = 0$.

When $b = \pi(b)$, the sequence defined above is not a solution to $Cx = 0$.

Remark 3.20. This spanning set can be refined to a basis; see Remark 5.13. △

4. Constructing juggler's friezes

As in Section 2, we may construct all juggler's friezes by twisting certain matrices.

4.1. π -unimodular matrices. First, we need a class of matrices that parametrize juggler's friezes. Given a juggling function π , the **landing schedule at $a \in \mathbb{Z}$** is the set

$$L_a := \{b \in \mathbb{Z} \mid \pi^{-1}(b) < a \leq b\}$$

Intuitively, L_a consists of the times at which the balls in the air right before time a will be caught. Matching earlier notation, we let $\overline{L}_a \subset \{1, 2, \dots, n\}$ denote the residues of $L_a \bmod n$.

We list some basic properties of landing schedules (see [KLS13]).

- (1) For all $a \in \mathbb{Z}$, $|L_a|$ equals the number of balls in π .
- (2) If $a < \pi(a)$, then $a \in L_a$ and $L_{a+1} = L_a \setminus \{a\} \cup \{\pi(a)\}$.
- (3) If $a = \pi(a)$, then $a \notin L_a$ and $L_{a+1} = L_a$.

Remark 4.1. The list $(\overline{L}_1, \overline{L}_2, \dots, \overline{L}_n)$ is the *Grassmann necklace*, a type of combinatorial object in bijection with positroids and juggling functions (see [KLS13]). △

Definition 4.2. Let π be n -periodic with k -many balls. A $k \times n$ -matrix A is **π -unimodular** if¹⁷

- $\det(A_{L_a}) = 1$ for all $a \in \mathbb{Z}$, and
- for all intervals $[a, b] \subset \mathbb{Z}$ with $b < a + n$, $\text{rank}(A_{[a,b]}) \leq |L_a \cap [a, b]|$.

Remark 4.3. The second condition is one of many equivalent ways to state that the rowspan of A is in the positroid variety corresponding to π ; see [KLS13, Section 5]. △

Example 4.4. If π is the juggling function with siteswap notation 23345357, then

a	1	2	3	4	5	6	7	8
\overline{L}_a	1247	2347	3457	4567	5678	2678	1278	1248

¹⁷Recall that, for any $I \subset \mathbb{Z}$, A_I denotes the submatrix of A with columns congruent to $I \bmod n$, in the order in which they appear in A .

The following matrix is π -unimodular; note the 8 minors indexed by the above $\overline{L_a}$ are 1.

$$\begin{bmatrix} 1 & 0 & -1 & 0 & 1 & 2 & 0 & -3 \\ 0 & 1 & 2 & 0 & -1 & -1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 2 & 1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \end{bmatrix} \quad \triangle$$

4.2. Constructing juggler's friezes. Given a π -unimodular $k \times n$ -matrix \mathbf{A} , define $F(\mathbf{A})$ to be the diamond grid of numbers (or $\mathbb{Z} \times \mathbb{Z}$ matrix) with the following entries.

$$F(\mathbf{A})_{a,b} := \begin{cases} (-1)^{|S_{\pi^\dagger}(b,a)|} \det(\mathbf{A}_{L_a \setminus \{a\} \cup \{b\}}) & \text{if } b \leq a < b+n \text{ and } a < \pi(a), \\ 1 & \text{if } \pi(a) = a = b, \\ (-1)^k & \text{if } \pi(a) = a = b+n, \\ 0 & \text{otherwise.} \end{cases} \quad (4.1)$$

In this definition, if $b \in L_a \setminus \{a\}$, we set $\det(\mathbf{A}_{L_a \setminus \{a\} \cup \{b\}}) = 0$. If there are no loops in π (that is, no a with $\pi(a) = a$), then this definition simplifies to

$$F(\mathbf{A})_{a,b} = \begin{cases} (-1)^{|S_{\pi^\dagger}(b,a)|} \det(\mathbf{A}_{L_a \setminus \{a\} \cup \{b\}}) & \text{if } b \leq a < b+n, \\ 0 & \text{otherwise.} \end{cases}$$

Theorem F. *If \mathbf{A} is a π -unimodular matrix, then $F(\mathbf{A})$ is a π^\dagger -frieze.*

Example 4.5. If \mathbf{A} is the π -unimodular matrix in the preceding example, then a few entries of $F(\mathbf{A})$ may be computed as follows.

$$\begin{aligned} F(\mathbf{A})_{2,1} &= (-1)^0 \det(\mathbf{A}_{L_2 \setminus \{2\} \cup \{1\}}) = \det(\mathbf{A}_{\{1347\}}) = 2 \\ F(\mathbf{A})_{2,-2} &= (-1)^1 \det(\mathbf{A}_{L_2 \setminus \{2\} \cup \{-2\}}) = (-1) \det(\mathbf{A}_{\{3467\}}) = -3 \\ F(\mathbf{A})_{6,1} &= (-1)^1 \det(\mathbf{A}_{L_6 \setminus \{6\} \cup \{1\}}) = (-1) \det(\mathbf{A}_{L_7}) = -1 \end{aligned} \quad \triangle$$

The juggler's frieze $F(\mathbf{A})$ defined in Theorem F can be constructed from (a more general version of) the twist. The **(right) twist** of a π -unimodular $k \times n$ -matrix \mathbf{A} is the $k \times n$ -matrix $\tau(\mathbf{A})$ whose columns are defined by the dot product equations

$$\tau(\mathbf{A})_a \cdot \mathbf{A}_b = \begin{cases} 1 & \text{if } b \in L_a \cap \{a\} \\ 0 & \text{if } b \in L_a \setminus \{a\} \end{cases}$$

Since $\det(\mathbf{A}_{L_a}) = 1$ for each a , the above system uniquely determines each column of $\tau(\mathbf{A})$. Note that, if a is not a loop of π , then $a \in L_a$ and the first condition above reduces to $\tau(\mathbf{A})_a \cdot \mathbf{A}_a = 1$. If a is a loop of π , then $\tau(\mathbf{A})_a$ is orthogonal to a basis and so $\tau(\mathbf{A})_a = 0$.

Remark 4.6. If \mathbf{A} is π -unimodular, then $\tau(\mathbf{A})$ is π -unimodular. \triangle

The twist can be used to give an alternate construction of $F(\mathbf{A})$ (Construction 4.7, next page), which is both faster in practice and the original motivation for the definition of juggler's friezes.

Theorem E. *If π is loop-free and \mathbf{A} is a π -unimodular matrix, then Construction 4.7 produces $F(\mathbf{A})$, the π^\dagger -frieze defined in Theorem F.*

See Remark 5.16 for a modification of this construction which covers loops.

4.3. Relation to the Grassmannian. Since the formula for $F(A)$ in Theorem F only depends on determinants of $k \times k$ -submatrices of A , the juggler’s frieze $F(A)$ only depends on the left $SL(k)$ -orbit of A within the set of π -unimodular matrices. This can be sharpened to a bijection.

Theorem G. *For each juggling function π with k -many balls, the map F descends to a bijection*

$$SL(k) \setminus \{\pi\text{-unimodular matrices}\} \xrightarrow{F} \{\pi^\dagger\text{-friezes}\}$$

As a consequence, F factors through a well-known description of the Grassmannian.

$$SL(k) \setminus \{\pi\text{-unimodular matrices}\} \subset GL(k) \setminus \{k \times n\text{-matrices of rank } k\} \xrightarrow{\text{rowspan}} Gr(k, n)$$

Since the π -unimodular matrices may be defined in terms of $k \times k$ -minors, the image of this inclusion is a closed subvariety of $Gr(k, n)$, which we denote by $Gr_{uni}(\pi)$. That is,

$$Gr_{uni}(\pi) := \{\text{rowspans of } \pi\text{-unimodular matrices}\} \subset Gr(k, n)$$

Corollary 4.8. *The map F induces a bijection $Gr_{uni}(\pi) \xrightarrow{\sim} \{\pi^\dagger\text{-friezes}\}$.*

Remark 4.9. The variety $Gr_{uni}(\pi)$ is related to the *positroid variety* of π [KLS13], which we denote $Gr(\pi)$, consisting of rowspans of matrices satisfying only the second condition in Definition 4.2. The variety $Gr_{uni}(\pi)$ is then the subset of $Gr(\pi)$ on which the Plücker coordinates of the form Δ_{L_a} are all equal to 1; it was dubbed the *critical variety* in [Gal21]. The bijection F can be extended to $Gr(\pi)$, by replacing π^\dagger -friezes with *quasiperiodic reduced recurrence matrices with juggling function π^\dagger* . Many of the results in this paper extend, as we hope to show in [DM]. \triangle

Sending a matrix to its positive complement descends to isomorphisms

$$\ddagger : Gr(\pi) \xrightarrow{\sim} Gr(\pi^\dagger) \text{ and } \ddagger : Gr_{uni}(\pi) \xrightarrow{\sim} Gr_{uni}(\pi^\dagger)$$

with inverses given by applying \ddagger again. Duality for friezes is then related to the twist as follows.

Theorem H. *Let A be a π -unimodular matrix. Then*

$$F(A)^\dagger = F(\tau(A)^\ddagger) = F(\tau^{-1}(A^\ddagger))$$

The preceding theorem implies the commutativity of the following diagram of bijections.

$$\begin{array}{ccccc}
 & & Gr_{uni}(\pi) & & \\
 & \tau \nearrow & & \ddagger \nwarrow & \\
 Gr_{uni}(\pi) & & & & Gr_{uni}(\pi^\dagger) \\
 & \ddagger \searrow & & \tau \swarrow & \\
 & & Gr_{uni}(\pi^\dagger) & & \\
 F \downarrow & & & & F \downarrow \\
 \{\pi^\dagger\text{-friezes}\} & \xleftarrow{\ddagger} & & \xrightarrow{\ddagger} & \{\pi\text{-friezes}\}
 \end{array}$$

Example 4.10. A computation of A^\ddagger and $F(\tau^{-1}(A^\ddagger))$ for the matrix A in Example 4.4 is given in the next section. The latter may be seen to coincide with $F(A)^\dagger$, appearing in Example 3.18. \triangle

4.4. Positivity, enumeration, and clusters. As mentioned in Section 1.1, many notable results count positive integral $SL(k)$ -friezes, often via bijections with other interesting sets of objects.

To pose an analogous problem for juggler’s friezes, we need to amend the notion of positivity.

Definition 4.11. A π -prefrieze over \mathbb{R} is **positive** if $(-1)^{|S_\pi(a,b)|} C_{a,b} > 0$ when not forced to be 0.

Remark 4.12. When π is uniform, $S_\pi(a,b)$ is empty whenever $C_{a,b}$ is not zero, and so the positivity condition becomes $C_{a,b} > 0$ when $C_{a,b}$ is not forced to be 0. \triangle

Many of the constructions in this paper have been defined so as to preserve positivity. For example, the dual of a positive juggler's frieze is positive, and F defines a bijection between the positive part of $\text{Gr}_{\text{uni}}(\pi)$ and the set of positive π^\dagger -friezes.

Restricting to the positive integral friezes gives a potentially interesting enumerative problem.

Problem 4.13. For which juggling functions π are there finitely many positive integral π -friezes? Given such π , how many are there? Are they in bijection with other objects of interest?

We claim that one robust source of juggler's friezes is a *cluster structure* on $\text{Gr}_{\text{uni}}(\pi)$, which we hope to prove in a subsequent work [DM].

Claim 4.14. *The variety $\text{Gr}_{\text{uni}}(\pi)$ admits a cluster structure, each cluster on $\text{Gr}_{\text{uni}}(\pi)$ determines a positive integral π^\dagger -frieze, and the map from clusters to positive integral π^\dagger -friezes is an inclusion.*

Remark 4.15. We can say a bit about this cluster structure. The positroid variety $\text{Gr}(\pi)$ has a cluster structure [Sco06, Pos06, Lec16, SSBW20, GL19]. Since $\text{Gr}_{\text{uni}}(\pi) \subset \text{Gr}(\pi)$ is the subvariety on which the frozen cluster variables of $\text{Gr}(\pi)$ are 1, the cluster structure descends to $\text{Gr}_{\text{uni}}(\pi)$ and has the same mutation type. In particular, the above claim implies that there are infinitely many positive integral π^\dagger -friezes whenever $\text{Gr}(\pi)$ has infinite mutation type. \triangle

Intermission: Computing a dual juggler's frieze via the twist

As an excuse to postpone the (rather technical) proofs of the main theorems, we provide a worked example of Theorem H. Consider the matrix A from Example 4.4, copied below.

$$A := \begin{bmatrix} 1 & 0 & -1 & 0 & 1 & 2 & 0 & -3 \\ 0 & 1 & 2 & 0 & -1 & -1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 2 & 1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \end{bmatrix}$$

Since A is in reduced row echelon form, a basis of the kernel of A may be computed via back substitution. We put this basis together into the rows of the following matrix.

$$B = \begin{bmatrix} 1 & -2 & 1 & 0 & 0 & 0 & 0 & 0 \\ -1 & 1 & 0 & -2 & 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & -1 & 0 & 1 & 0 & 0 \\ 3 & -1 & 0 & 1 & 0 & 0 & -1 & 1 \end{bmatrix}$$

A positive complement of A may be given by negating columns 1, 3, 5, and 7 of B . However, it will be convenient to also negate rows 1 and 2 (as this will make the entries of the inverse twist

5. Proofs of Theorems

In this section, we prove the aforementioned theorems. In the interest of precision, we will exclusively use ‘prefriezes’, ‘ π -prefriezes’, and related terms to refer to $\mathbb{Z} \times \mathbb{Z}$ -matrices, rather than grids of numbers in a diamond pattern. This primarily affects the meaning of the terms ‘row’, ‘column’, and ‘diagonal’; e.g. in this section we would say a prefrieze has two diagonals of 1s.



Warning 5.1. The reader is cautioned that a majority of what follows consists of verifying that various signs are correct. These arguments can be skipped without losing any key ideas. \triangle

5.1. Duality for juggler's friezes (Theorem A). A $\mathbb{Z} \times \mathbb{Z}$ -matrix C is **lower unitriangular** if $C_{a,b} = 0$ for all $a < b$ and $C_{a,b} = 1$ for all $a = b$. Rotated counterclockwise and deleting 0s above the diagonal, such a matrix becomes a diamond grid of numbers whose top row consists of 1s, but may continue infinitely downwards (as well as left and right, of course). This definition covers half of the definition of a π -prefrieze. The other half can be split off of the definition as follows.

Lemma 5.2. *A lower unitriangular $\mathbb{Z} \times \mathbb{Z}$ -matrix C is a π -prefrieze iff, for all a, b with $b \leq \pi^{-1}(a)$ or $\pi(b) \leq a$,*

$$C_{a,b} = \begin{cases} (-1)^{|S_\pi(b,a)|} & \text{if } a = \pi(b), \\ 0 & \text{if } b < \pi^{-1}(a) \text{ or } \pi(b) < a. \end{cases} \quad (5.1)$$

This condition on entries can be replaced by a condition on larger minors, via the following lemma. Recall that $S_\pi(a, b) := \{i \in \mathbb{Z} \mid a < i \text{ and } \pi(i) < b\}$.

Lemma 5.3. *A lower unitriangular $\mathbb{Z} \times \mathbb{Z}$ -matrix C is a π -prefrieze iff the following hold.*

(1) For all $a < b$,

$$\det(C_{\pi(S_\pi(a,b)), S_\pi(a,b)}) = 1 \quad (5.2)$$

(2) For all $a < b$ such that $a < \pi^{-1}(b)$ or $\pi(a) < b$,

$$\det(C_{\pi(S_\pi(a,b)) \cup \{b\}, \{a\} \cup S_\pi(a,b)}) = 0 \quad (5.3)$$

We note that Condition (1) is redundant and can be optimized; see Remark 5.4.

Proof. Note that if $|S_\pi(a, b)| = 0$, Equation (5.2) is vacuous and Equations (5.1) and (5.3) coincide.

(\Rightarrow) Assume that C is a π -prefrieze, and that C satisfies Equations (5.2) and (5.3) for any a, b with $|S_\pi(a, b)| < k$. Choose $a < b$ with $|S_\pi(a, b)| = k$, and index the elements in increasing order:

$$S_\pi(a, b) = \{i_1 < i_2 < \cdots < i_k\}$$

Note that, for all j , $S_\pi(i_j, b) = \{i_{j+1}, i_{j+2}, \dots, i_k\}$ and $S_\pi(i_1, \pi(i_j)) \subset S_\pi(i_1, b) \subsetneq S_\pi(a, b)$.

Therefore, $|S_\pi(i_1, \pi(i_j))| < |S_\pi(a, b)| = k$, and so $C_{\pi(i_1), i_1} = (-1)^{|S_\pi(i_1, \pi(i_1))|}$ and $C_{\pi(i_j), i_1} = 0$ for $j > 1$ by Equation (5.1). Taking a cofactor expansion of $\det(C_{\pi(S_\pi(a,b)), S_\pi(a,b)})$ along the i_1 th column and using that $C_{\pi(i_1), i_1}$ is in the $(|S_\pi(i_1, \pi(i_1))| + 1)$ th row of $C_{\pi(S_\pi(a,b)), S_\pi(a,b)}$,

$$\det(C_{\pi(S_\pi(a,b)), S_\pi(a,b)}) = (-1)^{|S_\pi(i_1, \pi(i_1))|} C_{\pi(i_1), i_1} \det(C_{\pi(S_\pi(i_1,b)), S_\pi(i_1,b)}) = \det(C_{\pi(S_\pi(i_1,b)), S_\pi(i_1,b)})$$

Since $S_\pi(i_1, b) \subsetneq S_\pi(a, b)$, the right-hand side is 1 by the assumed case of Equation (5.3). Therefore, Equation (5.2) holds for this a, b .

Now, assume that $a < \pi^{-1}(b)$ or $\pi(a) < b$, so that $C_{b,a} = 0$ by Equation (5.1). Since $|S_\pi(i_j, b)| < |S_\pi(a, b)| = k$ and $\pi(i_j) < b$, $C_{b,i_j} = 0$ for all j by Equation (5.1). Taking a cofactor expansion of $\det(C_{\pi(S_\pi(a,b)) \cup \{b\}, \{a\} \cup S_\pi(a,b)})$ along the b th row,

$$\det(C_{\pi(S_\pi(a,b)) \cup \{b\}, \{a\} \cup S_\pi(a,b)}) = (-1)^{|S_\pi(a,b)|} C_{b,a} \det(C_{\pi(S_\pi(a,b)), S_\pi(a,b)}) = 0$$

Therefore, Equation (5.3) holds for this a, b . Taking the limit as k goes to infinity implies that Equations (5.2) and (5.3) hold in general.

(\Leftarrow) Assume that C satisfies Equations (5.2) and (5.3) in general, and that C satisfies Equation (5.1) for any a, b with $|S_\pi(a, b)| < k$. Choose some $a < b$ such that $|S_\pi(a, b)| = k$. Again taking a cofactor expansion of $\det(C_{\pi(S_\pi(a,b)) \cup \{b\}, \{a\} \cup S_\pi(a,b)})$ along the b th row,

$$\det(C_{\pi(S_\pi(a,b)) \cup \{b\}, \{a\} \cup S_\pi(a,b)}) = (-1)^{|S_\pi(a,b)|} C_{b,a} \det(C_{\pi(S_\pi(a,b)), S_\pi(a,b)}) = (-1)^{|S_\pi(a,b)|} C_{b,a}$$

If $a < \pi^{-1}(b)$ or $\pi(a) < b$, then Equation (5.3) implies that the left-hand-side is 0, and so $C_{b,a} = 0$. If $\pi(a) = b$, then $S_\pi(a, b) \cup \{a\} = S_\pi(a-1, b+1)$ and Equation (5.2) implies that the left-hand-side is 1; and so $C_{b,a} = (-1)^{|S_\pi(a,b)|}$. Therefore, Equation (5.1) holds for this a, b . Taking the inductive limit over all k implies that Equation (5.1) holds in general, and so C is a π -prefrieze. \square

Remark 5.4. Condition (1) in Lemma 5.3 is redundant as stated, because many pairs (a, b) may yield the same set $S_\pi(a, b)$ and therefore the same Equation (5.2). One may restrict to a irredundant family by imposing Equation (5.2) for a, b such that $a + 1 \leq \pi^{-1}(b - 1)$ and $\pi(a + 1) \leq b - 1$. Reparametrizing $a + 1 \mapsto a$ and $b - 1 \mapsto b$, this condition becomes: $\det(C_{\pi(S), S}) = 1$ for all a, b with $a \leq \pi^{-1}(b)$ and $\pi(a) \leq b$, where $S = \{i \in \mathbb{Z} \mid a \leq i \text{ and } \pi(i) \leq b\} = S_\pi(a - 1, b + 1)$. \triangle

We would like a characterization of when C^\dagger is a π^\dagger -prefrieze. To do this, we need a relation between minors of C and C^\dagger .

Lemma 5.5. *Given subsets I, J of an interval $[a, b]$ with $b - a < n$,*

$$\det(C_{I, J}^\dagger) = \det(C_{[a, b] \setminus J, [a, b] \setminus I})$$

Proof. A lower unipotent matrix C has a well-defined adjugate matrix $\text{Adj}(C)$, defined by

$$\text{Adj}(C)_{a,b} := (-1)^{a+b} \det(C_{[a+1, b], [a, b-1]})$$

This adjugate is lower unitriangular and enjoys many nice properties, including

$$\det(\text{Adj}(C)_{I, J}) = (-1)^{\sum I + \sum J} \det(C_{[a, b] \setminus J, [a, b] \setminus I})$$

Here, $\sum I$ and $\sum J$ denote the sum of the elements of I and J , respectively.

For $b \leq a < b + n$, the dual coincides with the adjugate up to sign.

$$C_{a,b}^\dagger = \det(C_{[a+1, b], [a, b-1]}) = (-1)^{a+b} \text{Adj}(C)_{a,b}$$

Therefore,

$$\det(C_{I, J}^\dagger) = (-1)^{\sum I + \sum J} \det(\text{Adj}(C)_{I, J}) = \det(C_{[a, b] \setminus J, [a, b] \setminus I}) \quad \square$$

Lemma 5.6. *A π -prefrieze C is a π -frieze iff C^\dagger is a π^\dagger -prefrieze.*

Proof. Let C be a π -prefrieze. By Lemma 5.3 and Remark 5.4, C^\dagger is π^\dagger -prefrieze iff

- For all a, b with $a \leq \pi^{\dagger-1}(b)$ and $\pi^{\dagger}(a) \leq b$, and $b < a + n$,

$$\det(\mathbf{C}_{\pi^{\dagger}(I), I}^{\dagger}) = 1$$

where $I := \{i \in \mathbb{Z} \mid a \leq i \text{ and } \pi^{\dagger}(i) \leq b\} = S_{\pi^{\dagger}}(a - 1, b + 1)$.

- For all a, b with $a < \pi^{\dagger-1}(b)$ or $\pi^{\dagger}(a) < b$, and $b < a + n$,

$$\det(\mathbf{C}_{\pi^{\dagger}(J) \cup \{b\}, \{a\} \cup J}^{\dagger}) = 0$$

where $J := \{i \in \mathbb{Z} \mid a < i \text{ and } \pi^{\dagger}(i) < b\} = S_{\pi^{\dagger}}(a, b)$.

Using Lemma 5.5 to translate this into \mathbf{C} , \mathbf{C}^{\dagger} is a π^{\dagger} -prefrieze iff

- For all a, b with $a \leq \pi^{\dagger-1}(b)$ and $\pi^{\dagger}(a) \leq b$, and $b < a + n$,

$$\det(\mathbf{C}_{[a,b] \setminus I, [a,b] \setminus \pi^{\dagger}(I)}) = 1$$

where $I := \{i \in \mathbb{Z} \mid a \leq i \text{ and } \pi^{\dagger}(i) \leq b\} = S_{\pi^{\dagger}}(a - 1, b + 1)$.

- For all a, b with $a < \pi^{\dagger-1}(b)$ or $\pi^{\dagger}(a) < b$, and $b < a + n$,

$$\det(\mathbf{C}_{(a,b] \setminus J, [a,b] \setminus \pi^{\dagger}(J)}) = 0$$

where $J := \{i \in \mathbb{Z} \mid a < i \text{ and } \pi^{\dagger}(i) < b\} = S_{\pi^{\dagger}}(a, b)$.

These are precisely the frieze and tameness conditions in Definition 3.9. \square

As a consequence of Lemma 5.6, we can take the dual of the dual of a π -frieze.

Lemma 5.7. *If \mathbf{C} is a π -frieze, then $(\mathbf{C}^{\dagger})^{\dagger} = \mathbf{C}$.*

Proof. If $a \geq b > a - n$, then $(\mathbf{C}^{\dagger})_{a,b}^{\dagger} := \det(\mathbf{C}_{[a+1,b], [a,b-1]}^{\dagger})$, which equals $\mathbf{C}_{a,b}$ by Lemma 5.5.

Consider a with $\pi(a) = a + n$ (a loop of π), so that $\pi^{\dagger}(a) = a$ (a coloop of π^{\dagger}). Then

$$S_{\pi^{\dagger}}(a, a + n) = \{i \in \mathbb{Z} \mid a < i \text{ and } \pi^{\dagger}(i) < a + n\} = \{i \in \mathbb{Z} \mid a < i \text{ and } \pi^{-1}(i) < a\} = L_a \setminus \{a\}$$

In particular, the cardinality is one less than the number of balls in π , and so

$$(\mathbf{C}^{\dagger})_{a+n,a}^{\dagger} := (-1)^{(\# \text{ of balls in } \pi) - 1} = (-1)^{|S_{\pi^{\dagger}}(a, a+n)|} =: \mathbf{C}_{a+n,a}$$

All other entries in $(\mathbf{C}^{\dagger})^{\dagger}$ and \mathbf{C} are zero by definition, and so $(\mathbf{C}^{\dagger})^{\dagger} = \mathbf{C}$. \square

Theorem A. *If \mathbf{C} is a π -frieze, then the dual \mathbf{C}^{\dagger} is a π^{\dagger} -frieze, and $(\mathbf{C}^{\dagger})^{\dagger} = \mathbf{C}$.*

Proof. If \mathbf{C} is a π -frieze, then \mathbf{C}^{\dagger} is a π^{\dagger} -prefrieze by Lemma 5.6. Taking the dual again, $(\mathbf{C}^{\dagger})^{\dagger} = \mathbf{C}$ by Lemma 5.7. Then $(\mathbf{C}^{\dagger})^{\dagger}$ is a $(\pi^{\dagger})^{\dagger} = \pi$ -prefrieze, so Lemma 5.6 implies that \mathbf{C}^{\dagger} is a π^{\dagger} -frieze. \square

5.2. Superperiodicity of solutions (Theorems B, C, and D). This theorem follows from a number of results about linear recurrences proven in [Mul19]. The following lemma translates these results into the context and language of this paper.

Lemma 5.8. *Let \mathbf{C} be a π -prefrieze.*

- (1) *If \mathbf{C}' is another π -prefrieze with $\ker(\mathbf{C}) = \ker(\mathbf{C}')$, then $\mathbf{C} = \mathbf{C}'$.*
- (2) *There is a unique matrix $\mathbb{Z} \times \mathbb{Z}$ -matrix $\text{Sol}(\mathbf{C})$, called the solution matrix of \mathbf{C} , such that*
 - (a) $\mathbf{C}\text{Sol}(\mathbf{C}) = 0$,

- (b) $\text{Sol}(\mathbf{C})_{a,a} = 1$ whenever $\pi(a) \neq a$, and
- (c) $\text{Sol}(\mathbf{C})_{a,b} = 0$ whenever $\pi(a) = b = a$ or $a < b < \pi(a)$.
- (3) $\text{Sol}(\mathbf{C})$ is also the unique $\mathbb{Z} \times \mathbb{Z}$ -matrix such that
 - (a) $\text{Sol}(\mathbf{C})\mathbf{C} = 0$,
 - (b) $\text{Sol}(\mathbf{C})_{a,a} = 1$ whenever $\pi(a) \neq a$, and
 - (c) $\text{Sol}(\mathbf{C})_{a,b} = 0$ whenever $\pi(b) = a = b$ or $\pi^{-1}(b) < a < b$.
- (4) The columns of $\text{Sol}(\mathbf{C})$ span the space of solutions to $\mathbf{C}x = 0$, whose dimension equals the number of balls in π .
- (5) For all $b < a$, $\text{Sol}(\mathbf{C})_{a,b} = (-1)^{a+b} \det(\mathbf{C}_{[a+1,b],[a,b-1]})$.

Combining (4) with the definition of \mathbf{C}^\dagger , we see that $\text{Sol}(\mathbf{C})_{a,b} = (-1)^{a+b} \mathbf{C}_{a,b}^\dagger$ when $b \leq a < \pi(b)$.

Proof. A π -prefrieze is a reduced recurrence matrix of shape π^{-1} in the language of [Mul19].

- (1) is [Mul19, Theorem 6.5]; specifically, the uniqueness of the reduced recurrence matrix in a given equivalence class.
- (2) is [Mul19, Theorem 8.3].
- (3) is [Mul19, Theorem 10.5], since π^{-1} is bijective.
- (4) is [Mul19, Theorems 8.12 and 7.14]. Note that the solution space was shown to be the closure of the span of the columns of $\text{Sol}(\mathbf{C})$ in the product topology. However, since π has finitely many balls, the solution space is finite dimensional and therefore closed.
- (5) is [Mul19, Proposition 8.6]. □

Before the proof of Theorem B, the following formula will ensure the signs work out.

Lemma 5.9. *Let π be a juggling function of period n . Then, for all a ,*

$$\pi(a) - a + |S_{\pi^\dagger}(\pi(a), a + n)| - |S_\pi(a, \pi(a))| = (\# \text{ of balls in } \pi)$$

Proof. First, we show that the value of the expression

$$\pi(a) - a + |S_{\pi^\dagger}(\pi(a), a + n)| - |S_\pi(a, \pi(a))| \tag{5.4}$$

is independent of a .

Consider the case that $\pi(a) < \pi(a + 1)$. Then $S_\pi(a, \pi(a)) \subseteq S_\pi(a + 1, \pi(a + 1))$ and

$$S_\pi(a + 1, \pi(a + 1)) \setminus S_\pi(a, \pi(a)) = \{i \in \mathbb{Z} \mid a + 1 < i \text{ and } \pi(a) < \pi(i) < \pi(a + 1)\}$$

Similarly, $S_{\pi^\dagger}(\pi(a), a + n) \supseteq S_{\pi^\dagger}(\pi(a + 1), a + n + 1)$ and

$$S_{\pi^\dagger}(\pi(a), a + n) \setminus S_{\pi^\dagger}(\pi(a + 1), a + n + 1) = \{i \in \mathbb{Z} \mid \pi(a) < i < \pi(a + 1) \text{ and } \pi^\dagger(i) < a + n\}$$

Acting on this set by π^{-1} gives

$$\begin{aligned} & \pi^{-1}(S_{\pi^\dagger}(\pi(a), a + n) \setminus S_{\pi^\dagger}(\pi(a + 1), a + n + 1)) \\ &= \{i \in \mathbb{Z} \mid \pi(a) < \pi(i) < \pi(a + 1) \text{ and } i + n < a + n\} \end{aligned}$$

Comparing sets, we see that the interval $(\pi(a), \pi(a + 1))$ decomposes as a disjoint union

$$(S_\pi(a + 1, \pi(a + 1)) \setminus S_\pi(a, \pi(a))) \sqcup (\pi^{-1}(S_{\pi^\dagger}(\pi(a), a + n) \setminus S_{\pi^\dagger}(\pi(a + 1), a + n + 1)))$$

Counting elements on either side, $\pi(a+1) - \pi(a) - 1$ equals

$$|S_\pi(a+1, \pi(a+1))| - |S_\pi(a, \pi(a))| + |S_{\pi^\dagger}(\pi(a), a+n)| - |S_{\pi^\dagger}(\pi(a+1), a+n+1)|$$

Therefore, the value of Expression (5.4) is the same for a and $a+1$. An analogous argument applies when $\pi(a) > \pi(a+1)$, and so (5.4) is constant for all a .

Let h denote the value of (5.4). Averaging (5.4) over a from 1 to n ,

$$\begin{aligned} h &= \frac{1}{n} \sum_{a=1}^n \pi(a) - a + |S_{\pi^\dagger}(\pi(a), a+n)| - |S_\pi(a, \pi(a))| \\ &= (\# \text{ of balls in } \pi) + \frac{1}{n} \left(\sum_{a=1}^n |S_{\pi^\dagger}(\pi(a), a+n)| - \sum_{a=1}^n |S_\pi(a, \pi(a))| \right) \end{aligned}$$

Observe that $\sum_{a=1}^n |S_\pi(a, \pi(a))|$ counts orbits of *inversions* in π ; that is, pairs (a, b) with $a < b$ and $\pi(a) > \pi(b)$, up to the relation $(a, b) \sim (a+n, b+n)$. Similarly, $\sum_{a=1}^n |S_{\pi^\dagger}(\pi(a), a+n)|$ counts orbits of inversions in π^\dagger . To see the two sums cancel, observe that (a, b) is an inversion in π iff $(\pi(b), \pi(a))$ is an inversion in π^\dagger . It follows that h equals the number of balls in π . \square

Theorem B. *Let π be an n -periodic juggling function with h -many balls. A π -prefrieze C is a π -frieze iff every solution to the linear recurrence $Cx = 0$ satisfies $x_{a+n} = (-1)^{n-h-1}x_a$, $\forall a \in \mathbb{Z}$.*

Proof. For simplicity, let $k := (\# \text{ of balls in } \pi^\dagger) = n - (\# \text{ of balls in } \pi)$. Let Σ denote the permutation matrix with $\Sigma_{a, a+1} = 1$ and all other entries 0. Then an infinite sequence x (regarded as a vector of height \mathbb{Z}) satisfies

$$x_{a+n} = (-1)^{k-1}x_a \tag{5.5}$$

for all a iff $\Sigma^n x = (-1)^{k-1}x$.

Let C be a π -prefrieze and let $\text{Sol}(C)$ be the solution matrix of C . By Lemma 5.8.4, the columns of $\text{Sol}(C)$ span of the solutions to $Cx = 0$, and so every solution to $Cx = 0$ satisfies Equation (5.5) iff every column of $\text{Sol}(C)$ does. Therefore, by Theorem A, to prove Theorem B it will suffice to show that C^\dagger is a π^\dagger -prefrieze iff $\Sigma^n \text{Sol}(C) = (-1)^{k-1} \text{Sol}(C)$, which we will show via a lemma.

Lemma 5.10. *$\Sigma^n \text{Sol}(C) = (-1)^{k-1} \text{Sol}(C)$ iff the following conditions hold.*

- (1) $\text{Sol}(C)_{a+n, b} = 0$ whenever $a < b < \pi(a)$,
- (2) $\text{Sol}(C)_{a+n, b} = 0$ whenever $\pi^{-1}(b) < a < b$, and
- (3) $\text{Sol}(C)_{a+n, a} = (-1)^{k-1}$ whenever $\pi(a) \neq a$.

Proof. Observe that, by Lemma 5.8.3,¹⁸

$$\left((-1)^{k-1} \Sigma^n \text{Sol}(C) \right) C = (-1)^{k-1} \Sigma^n (\text{Sol}(C)C) = 0$$

Therefore, $(-1)^{k-1} \Sigma^n \text{Sol}(C)$ satisfies Condition (a) in Lemma 5.8.3. If $\pi(b) = b$, then the b th column of $\text{Sol}(C)$ consists entirely of zeros, by [Mul19, Lemma 8.9] (this also follows from $\text{Sol}(C)C = 0$), and so $\text{Sol}(C)_{a+n, b} = 0$ whenever $\pi(b) = a = b$.

¹⁸Associating products of infinite matrices is not always possible. It is allowed here because $(-1)^{k-1} \Sigma^n$ is a generalized permutation matrix.

Using that $((-1)^{k-1}\Sigma^n\text{Sol}(\mathbf{C}))_{a,b} = (-1)^{k-1}\text{Sol}(\mathbf{C})_{a+n,b}$, the remaining conditions of Lemma 5.8.3 imply that $((-1)^{k-1}\Sigma^n\text{Sol}(\mathbf{C})) = \text{Sol}(\mathbf{C})$ if and only if Conditions (2) and (3) above hold. Once we know $((-1)^{k-1}\Sigma^n\text{Sol}(\mathbf{C})) = \text{Sol}(\mathbf{C})$, then Condition (1) follows from Lemma 5.8.2. \square

Consider $b < a + n < b + n$. Then

$$\text{Sol}(\mathbf{C})_{a+n,b} = (-1)^{a+b+n} \det(\mathbf{C}_{[a+n+1,b],[a+n,b-1]}) = (-1)^{a+b+n} \mathbf{C}_{a+n,b}^\dagger$$

Then Conditions (1) and (2) in Lemma 5.10 may be rewritten as

- (1) $\mathbf{C}_{a,b}^\dagger = 0$ whenever $a < b + n < \pi^\dagger(a)$, and
- (2) $\mathbf{C}_{a,b}^\dagger = 0$ whenever $\pi^\dagger(b) < a < b + n$.

These equalities coincide with the vanishing part of the π^\dagger -prefrieze condition for \mathbf{C}^\dagger (the second case of Equation (5.1)).

For a with $\pi(a) \neq a$, consider the $(a + n, a)$ th entry in the equation $\text{Sol}(\mathbf{C})\mathbf{C} = 0$, which is

$$\sum_{b \in \mathbb{Z}} \text{Sol}(\mathbf{C})_{a+n,b} \mathbf{C}_{b,a} = 0$$

Since \mathbf{C} is a π -prefrieze, $\mathbf{C}_{b,a} = 0$ whenever $b \notin [a, \pi(a)]$. Assuming Condition (1) in Lemma 5.10, the above sum reduces to

$$\text{Sol}(\mathbf{C})_{a+n,a} + (-1)^{a+\pi(a)+n} \mathbf{C}_{a+n,\pi(a)}^\dagger \mathbf{C}_{\pi(a),a} = 0$$

By the π -prefrieze condition, $\mathbf{C}_{\pi(a),a} = (-1)^{|S_\pi(a,\pi(a))|}$ and so

$$\text{Sol}(\mathbf{C})_{a+n,a} = (-1)^{a+\pi(a)+n-1+|S_\pi(a,\pi(a))|} \mathbf{C}_{\pi^\dagger(\pi(a)),\pi(a)}^\dagger$$

Using Lemma 5.9,

$$\text{Sol}(\mathbf{C})_{a+n,a} = (-1)^{|S_{\pi^\dagger}(\pi(a),a+n)|+k-1} \mathbf{C}_{\pi^\dagger(\pi(a)),\pi(a)}^\dagger$$

Therefore, $\text{Sol}(\mathbf{C})_{a+n,a} = (-1)^{k-1}$ iff $\mathbf{C}_{\pi^\dagger(\pi(a)),\pi(a)}^\dagger = (-1)^{|S_{\pi^\dagger}(\pi(a),a+n)|}$, and so Condition (3) in Lemma 5.10 is equivalent to the non-vanishing part of the π^\dagger -prefrieze condition for \mathbf{C}^\dagger (the first case of Equation (5.1)). Therefore, \mathbf{C}^\dagger is a π^\dagger -prefrieze iff the conditions in Lemma 5.10 hold; equivalently, if every solution to $\mathbf{C}\mathbf{x} = 0$ satisfies Equation (5.5). \square

Theorem C. *If π is an n -periodic juggling function, then the entries of a π -frieze \mathbf{C} are n -periodic; that is, $\mathbf{C}_{a,b} = \mathbf{C}_{a+n,b+n}$.*

Proof. Let \mathbf{C} be a π -frieze, and let \mathbf{C}' denote the π -frieze defined by translating the entries by n :

$$(\mathbf{C}')_{a,b} := \mathbf{C}_{a+n,b+n}$$

Let $\mathbf{v} \in \ker(\mathbf{C})$. By Theorem B, $v_{a+n} = (-1)^{n-h-1}x_a$ for all a .

$$(\mathbf{C}'\mathbf{v})_a = \sum_{b \in \mathbb{Z}} (\mathbf{C}')_{a,b} v_b = \sum_{b \in \mathbb{Z}} \mathbf{C}_{a+n,b+n} \left((-1)^{n-h-1} v_{b+n} \right) = (-1)^{n-h-1} (\mathbf{C}\mathbf{v})_{a+n}$$

Since $\mathbf{v} \in \ker(\mathbf{C})$, this is always 0, and so $\mathbf{v} \in \ker(\mathbf{C}')$. A symmetric argument shows that $\ker(\mathbf{C}) = \ker(\mathbf{C}')$. By Lemma 5.8.1, $\mathbf{C} = \mathbf{C}'$. \square

As a consequence, the solution matrix of a π -frieze can be constructed by ‘tiling’ the dual \mathbf{C}^\dagger , as follows. Given an n -periodic juggling function π with h -many balls, define the **tiling** of a π -frieze

\mathbf{C} to be the $\mathbb{Z} \times \mathbb{Z}$ -matrix $\text{Tiling}(\mathbf{C})$ defined by

$$\text{Tiling}(\mathbf{C})_{a,b} = \sum_{i \in \mathbb{Z}} (-1)^{a+b+i(n-h-1)} \mathbf{C}_{a+in,b}$$

This first modifies the sign of each $\mathbf{C}_{a,b}$ by $(-1)^{a+b}$, and then extends the columns superperiodically.

Remark 5.11. [MGOST14] and [MG15] define tilings of $\text{SL}(k)$ -friezes without modifying the signs by $(-1)^{a+b}$ first. This is because the linear recurrence they associate to a frieze is not $\mathbf{C}\mathbf{x} = 0$, but instead use an alternating sign convention. \triangle

Proposition 5.12. *If \mathbf{C} is a juggler's frieze, then $\text{Sol}(\mathbf{C}) = \text{Tiling}(\mathbf{C}^\dagger)$.*

Proof. First, assume $b \leq a < b + n$. If $b \neq \pi(b)$, then by Lemma 5.8.5,

$$\text{Sol}(\mathbf{C})_{a,b} = (-1)^{a+b} \mathbf{C}_{a,b}^\dagger = \text{Tiling}(\mathbf{C}^\dagger)_{a,b}$$

If $b = \pi(b)$, then $\text{Sol}(\mathbf{C})_{a,b} = 0$. If $a > b$, then

$$\text{Tiling}(\mathbf{C}^\dagger)_{a,b} = \mathbf{C}_{a,b}^\dagger = 0$$

If $a = b = \pi(b)$, then

$$\text{Tiling}(\mathbf{C}^\dagger)_{a,b} = \mathbf{C}_{a,b}^\dagger - \mathbf{C}_{a+n,b}^\dagger = 1 - 1 = 0$$

In both cases, $\text{Sol}(\mathbf{C})_{a,b} = \text{Tiling}(\mathbf{C}^\dagger)_{a,b}$.

Next, we check that the columns of $\text{Tiling}(\mathbf{C}^\dagger)_{a,b}$ satisfy Equation (5.5):¹⁹

$$\begin{aligned} \text{Tiling}(\mathbf{C}^\dagger)_{a+n,b} &= \sum_{i \in \mathbb{Z}} (-1)^{a+n+b+i(n-k-1)} \mathbf{C}_{a+n+in,b}^\dagger \\ &= \sum_{i \in \mathbb{Z}} (-1)^{a+b+i(n-k-1)-(k+1)} \mathbf{C}_{a+in,b}^\dagger = (-1)^{k-1} \text{Tiling}(\mathbf{C}^\dagger)_{a,b} \end{aligned}$$

Since the columns of $\text{Sol}(\mathbf{C})$ also satisfy this equation, the two matrices coincide everywhere. \square

Theorem D. *Let π be an n -periodic juggling function with h -many balls, and let \mathbf{C} be a π -frieze. For any b with $b < \pi(b)$, the sequence \mathbf{x} defined by*

- $x_a := (-1)^{a+b} \mathbf{C}_{a,b}^\dagger$ when $b \leq a < b + k + h$, and
- $x_{a+n} = (-1)^{n-h-1} x_a$ for all a

is a solution to $\mathbf{C}\mathbf{x} = 0$. These solutions collectively span the space of solutions to $\mathbf{C}\mathbf{x} = 0$.

Proof. By Lemma 5.8.4, the columns of $\text{Sol}(\mathbf{C})$ span the space of solutions to $\mathbf{C}\mathbf{x} = 0$. When $b = \pi(b)$, the b th column of $\text{Sol}(\mathbf{C})$ is identically zero, so it suffices to consider the columns for which $b < \pi(b)$. By Proposition 5.12, the b th column of $\text{Sol}(\mathbf{C})$ coincides with the b th column of $\text{Tiling}(\mathbf{C}^\dagger)$, which is the sequence defined in the statement of Theorem D. \square

Remark 5.13. By [Mul19, Prop. 7.12], a basis of solutions to $\mathbf{C}\mathbf{x} = 0$ is given by the columns of $\text{Sol}(\mathbf{C})$ indexed by any *schedule* of π , such as a landing schedule L_a . \triangle

¹⁹Note that the number of balls in π^\dagger is $k := n - h$, not h .

5.3. Equivalence of constructions (Theorem E). The relation between the determinantal construction of $F(\mathbf{A})$ and Construction 4.7 follows primarily from the following lemma.

Lemma 5.14. *Let \mathbf{A} be a π -unimodular $k \times n$ -matrix. For any $1 \leq a, b \leq n$,*

$$\tau(\mathbf{A})_a \cdot \mathbf{A}_b = \begin{cases} (-1)^{|S_{\pi^\dagger}(b,a)|} \det(\mathbf{A}_{L_a \setminus \{a\} \cup \{b\}}) & \text{if } a \geq b \text{ and } a < \pi(a), \\ (-1)^{k-1+|S_{\pi^\dagger}(b,a+n)|} \det(\mathbf{A}_{L_a \setminus \{a\} \cup \{b\}}) & \text{if } a < b \text{ and } a < \pi(a), \\ 0 & \text{if } a = \pi(a). \end{cases}$$

Proof. If $a = \pi(a)$ (i.e. a is a loop of π), then $a \notin L_a$ and so the a th column of $\tau(\mathbf{A})$ is defined to be the unique solution to $(\mathbf{A}_{L_a})^\top \mathbf{x} = \mathbf{0}$. Therefore, $\tau(\mathbf{A})_a = \mathbf{0}$, and so $\tau(\mathbf{A})_a \cdot \mathbf{A}_b = \mathbf{0}$ for all b .

If $a < \pi(a)$, then $a \in L_a$ and so the a th column of $\tau(\mathbf{A})$ is defined to be the unique solution to $(\mathbf{A}_{L_a})^\top \mathbf{x} = \mathbf{e}_a$. Therefore, $\tau(\mathbf{A})_a = (\mathbf{A}_{L_a})^{\top-1} \mathbf{e}_a$, and so

$$\tau(\mathbf{A})_a \cdot \mathbf{A}_b = ((\mathbf{A}_{L_a})^{\top-1} \mathbf{e}_a) \cdot \mathbf{A}_b = \mathbf{e}_a \cdot (\mathbf{A}_{L_a})^{-1} \mathbf{A}_b$$

Equivalently, $\tau(\mathbf{A})_a \cdot \mathbf{A}_b$ is the a th entry of the solution to $\mathbf{A}_{L_a} \mathbf{x} = \mathbf{A}_b$. By Cramer's Rule, $\tau(\mathbf{A})_a \cdot \mathbf{A}_b$ equals the determinant of \mathbf{A}_{L_a} after replacing the a th column of \mathbf{A} by the b th column of \mathbf{A} .²⁰ By our convention for ordering the columns²¹ in $\mathbf{A}_{L_a \setminus \{a\} \cup \{b\}}$,

$$\tau(\mathbf{A})_a \cdot \mathbf{A}_b = (-1)^s \det(\mathbf{A}_{L_a \setminus \{a\} \cup \{b\}})$$

where s is the number of columns of \mathbf{A}_{L_a} between the a th column and the b th column.

If $b \in \overline{L_a}$ (i.e. b is congruent to an element of L_a), then $\det(\mathbf{A}_{L_a \setminus \{a\} \cup \{b\}}) = 0$ and so s is irrelevant.

If $b \leq a$ and $b \notin \overline{L_a}$, then s is the number of columns of \mathbf{A}_{L_a} right of the b th column and left of the a th column. Lifting to \mathbb{Z} , this is the cardinality of the set $L_{a-n} \cap [b+1, a-1]$.

$$\begin{aligned} s &= |L_{a-n} \cap [b+1, a-1]| = |\{i \in [b+1, a-1] \mid \pi^{-1}(i) < a-n\}| \\ &= |\{i \in [b+1, a-1] \mid \pi^\dagger(i) < a\}| = |[b+1, a-1] \cap \pi^\dagger([b+1, a-1])| =: |S_{\pi^\dagger}(b, a)| \end{aligned}$$

If $b > a$ and $b \notin \overline{L_a}$, then s is the number of columns of \mathbf{A}_{L_a} right of the a th column and left of the b th column. Lifting to \mathbb{Z} , this is the cardinality of the set $L_a \cap [a+1, b-1]$. Since $\{a\} \in L_a$ and $\{b\} \notin L_a$, L_a decomposes as a disjoint union

$$L_a = \{a\} \sqcup (L_a \cap [a+1, b-1]) \sqcup (L_a \cap [b+1, a+n-1])$$

The cardinality of L_a is the number of balls of π , which is k . By the same argument as the previous case, the cardinality of $L_a \cap [b+1, a+n-1]$ is $|S_{\pi^\dagger}(b, a+n)|$. Therefore, in this case,

$$s = k - 1 - |S_{\pi^\dagger}(b, a+n)|$$

This completes the proof. □

This allows us to restate the definition of $F(\mathbf{A})$ as follows.

²⁰There is also a denominator of $\det(\mathbf{A}_{L_a})$ in Cramer's Rule, but this determinant is 1 by π -unimodularity.

²¹Specifically, the columns in \mathbf{A}_{L_a} retain the same ordering as in \mathbf{A} .

Corollary 5.15. *Let $F(\mathbf{A})$ be defined as in Theorem F. Then*

$$F(\mathbf{A})_{a,b} := \begin{cases} \tau(\mathbf{A})_{\bar{a}} \cdot \mathbf{A}_{\bar{b}} & \text{if } b \leq a < b+n, \bar{b} \leq \bar{a}, \text{ and } a < \pi(a), \\ (-1)^{k-1} \tau(\mathbf{A})_{\bar{a}} \cdot \mathbf{A}_{\bar{b}} & \text{if } b \leq a < b+n, \bar{b} > \bar{a}, \text{ and } a < \pi(a), \\ 1 & \text{if } \pi(a) = a = b, \\ (-1)^k & \text{if } \pi(a) = a = b+n, \\ 0 & \text{otherwise.} \end{cases}$$

where $1 \leq \bar{a}, \bar{b} \leq n$ denote the respective residues of a, b mod n .

Theorem E. *If π is loop-free and \mathbf{A} is a π -unimodular matrix, then Construction 4.7 produces $F(\mathbf{A})$, the π^\dagger -frieze defined in Theorem F.*

Proof of Theorem E. If π is loop-free, then the corollary specializes to

$$F(\mathbf{A})_{a,b} := \begin{cases} \tau(\mathbf{A})_{\bar{a}} \cdot \mathbf{A}_{\bar{b}} & \text{if } b \leq a < b+n \text{ and } \bar{b} \leq \bar{a}, \\ (-1)^{k-1} \tau(\mathbf{A})_{\bar{a}} \cdot \mathbf{A}_{\bar{b}} & \text{if } b \leq a < b+n \text{ and } \bar{b} > \bar{a}, \\ 0 & \text{otherwise.} \end{cases}$$

Since $(\tau(\mathbf{A})^\top \mathbf{A})_{\bar{a}, \bar{b}} = \tau(\mathbf{A})_{\bar{a}} \cdot \mathbf{A}_{\bar{b}}$, Construction 4.7 produces the above entries. \square

Remark 5.16. Corollary 5.15 indicates how to adapt Construction 4.7 to the case when π has loops. Specifically, if $\pi(a) = a$, then $(\tau(\mathbf{A})^\top \mathbf{A})_{a,a} = 0$. In Step (3) of Construction 4.7, this 0 should be split into $1 + (-1)$. The 1 stays in place and the (-1) moves with the above-diagonal entries to become a $(-1)^k$ in position $F(\mathbf{A})_{a+n,a}$. \triangle

5.4. $F(\mathbf{A})$ is a π^\dagger -frieze (Theorem F). We will prove that $F(\mathbf{A})$ is a π^\dagger -frieze by first showing it is a π^\dagger -prefrieze, and then showing its kernel consists of superperiodic sequences.

Lemma 5.17. *If \mathbf{A} is π -unimodular, then $F(\mathbf{A})$ is a π^\dagger -prefrieze.*

Proof. Since $F(\mathbf{A})$ is lower unitriangular, it suffices to show that $F(\mathbf{A})$ satisfies Lemma 5.2 for π^\dagger . Given $a \in \mathbb{Z}$, we split into two cases.

- If $a = \pi(a)$ (i.e. a is a loop of π), then $\pi^\dagger(a) = a + n$ and so $F(\mathbf{A})_{\pi^\dagger(a), a} := (-1)^k$. Then

$$|S_{\pi^\dagger(a), a}| = n - (\# \text{ of balls in } \pi^\dagger) = (\# \text{ of balls in } \pi) = k$$

Therefore, $F(\mathbf{A})_{\pi^\dagger(a), a} = (-1)^{|S_{\pi^\dagger(a), a}|}$.

- If $a < \pi(a)$, then $\pi^\dagger(a) < a + n$. Then last inequality is equivalent to $\pi^{-1}(a) < a$, and so $\pi^{-1}(a) \in L_{\pi^{-1}(a)}$. Since $a = \pi(\pi^{-1}(a))$,

$$\begin{aligned} L_{\pi^{-1}(a)+1} &= L_{\pi^{-1}(a)} \setminus \{\pi^{-1}(a)\} \cup \{a\} \\ F(\mathbf{A})_{\pi^\dagger(a), a} &:= (-1)^{|S_{\pi^\dagger(a), \pi^\dagger(a)}|} \det(\mathbf{A}_{L_{\pi^\dagger(a)} \setminus \{\pi^\dagger(a)\} \cup \{a\}}) \\ &= (-1)^{|S_{\pi^\dagger(a), \pi^\dagger(a)}|} \det(\mathbf{A}_{L_{\pi^{-1}(a)} \setminus \{\pi^{-1}(a)\} \cup \{a\}}) \\ &= (-1)^{|S_{\pi^\dagger(a), \pi^\dagger(a)}|} \det(\mathbf{A}_{L_{\pi^{-1}(a)+1}}) = (-1)^{|S_{\pi^\dagger(a), \pi^\dagger(a)}|} \end{aligned}$$

We have used that \mathbf{A}_I only depends on I modulo n , and that \mathbf{A} is π -unimodular.

In all cases, we see that $F(\mathbf{A})_{\pi^\dagger(a),a} = (-1)^{|S_{\pi^\dagger(a,\pi^\dagger(a))}|}$, satisfying the first case of Equation (5.1).

Next, consider a, b such that $a - n < b < \pi^\dagger(a)$ or $\pi^\dagger(b) < a < b + n$. By Corollary 5.15,

$$F(\mathbf{A})_{a,b} = \pm(\tau(\mathbf{A})_a \cdot \mathbf{A}_b)$$

The dot product on the right-hand-side always vanishes, e.g. by [MS17, Lemma 6.4].²² Therefore, $F(\mathbf{A})$ satisfies the conditions of Lemma 5.2 for π^\dagger . \square

Given a vector $\mathbf{v} \in \mathbb{k}^n$ and integers (k, n) , the **superperiodic extension** of \mathbf{v} to be the infinite sequence $q_{(k,n)}(\mathbf{v}) \in \mathbb{k}^{\mathbb{Z}}$ defined by

$$q_{(k,n)}(\mathbf{v})_a := (-1)^{(k-1)\frac{a-\bar{a}}{n}} \mathbf{v}_{\bar{a}}$$

Note that \mathbf{v} can be recovered from $q_{(k,n)}(\mathbf{v})$ by restricting to the terms indexed by $[n] \subset \mathbb{Z}$.

Lemma 5.18. *Let \mathbf{A} be a π -unimodular $k \times n$ -matrix. Then $\mathbf{A}\mathbf{v} = 0$ iff $F(\mathbf{A})q_{(k,n)}(\mathbf{v}) = 0$.*

I.e. a vector \mathbf{v} is in the kernel of \mathbf{A} iff its superperiodic extension $q_{(k,n)}(\mathbf{v})$ is in the kernel of $F(\mathbf{A})$.

Proof. Let $\mathbf{v} \in \ker(\mathbf{A})$, and consider the a th entry in the product $F(\mathbf{A})q_{(k,n)}(\mathbf{v})$, which is

$$\sum_{b \in \mathbb{Z}} F(\mathbf{A})_{a,b} q_{(k,n)}(\mathbf{v})_b = \sum_{b \in \mathbb{Z}} F(\mathbf{A})_{a,b} (-1)^{(k-1)\frac{b-\bar{b}}{n}} \mathbf{v}_{\bar{b}}$$

If $a = \pi(a)$, then the only non-zero terms occur when $b \in \{a, a - n\}$, in which case

$$\sum_{b \in \mathbb{Z}} F(\mathbf{A})_{a,b} q_{(k,n)}(\mathbf{v})_b = (-1)^{(k-1)\frac{a-\bar{a}}{n}} \mathbf{v}_{\bar{a}} + (-1)^{(k-1)\frac{a-n-\bar{a}}{n}} \mathbf{v}_{\bar{a}} = (-1)^{(k-1)\frac{a-\bar{a}}{n}} (\mathbf{v}_{\bar{a}} - \mathbf{v}_{\bar{a}}) = 0$$

If $a < \pi(a)$, then the only non-zero terms occur when $b \in [a - n + 1, a]$. Let i be the unique integer with $a - n < in \leq a$. Then

$$\begin{aligned} \sum_{b \in \mathbb{Z}} F(\mathbf{A})_{a,b} q_{(k,n)}(\mathbf{v})_b &= \sum_{b=a-n+1}^a F(\mathbf{A})_{a,b} (-1)^{(k-1)\frac{b-\bar{b}}{n}} \mathbf{v}_{\bar{b}} \\ &= \sum_{b=a-n+1}^{in} (\tau(\mathbf{A})_{\bar{a}} \cdot \mathbf{A}_{\bar{b}}) (-1)^{(k-1)i} \mathbf{v}_{\bar{b}} + \sum_{b=in}^a (-1)^{k-1} (\tau(\mathbf{A})_{\bar{a}} \cdot \mathbf{A}_{\bar{b}}) (-1)^{(k-1)(i-1)} \mathbf{v}_{\bar{b}} \\ &= (-1)^{(k-1)i} \sum_{b=a-n+1}^{an} (\tau(\mathbf{A})_{\bar{a}} \cdot \mathbf{A}_{\bar{b}}) \mathbf{v}_{\bar{b}} = (-1)^{(k-1)i} \sum_{\bar{b}=1}^n (\tau(\mathbf{A})_{\bar{a}} \cdot \mathbf{A}_{\bar{b}}) \mathbf{v}_{\bar{b}} \\ &= (-1)^{(k-1)i} (\tau(\mathbf{A})^\top \mathbf{A} \mathbf{v})_{\bar{a}} \end{aligned}$$

Since $\mathbf{A}\mathbf{v} = 0$, this expression is 0. Therefore, $F(\mathbf{A})q_{(k,n)}(\mathbf{v})$ is the infinite sequence of zeroes.

This establishes that the superperiodic extension $q_{(k,n)}(\ker(\mathbf{A}))$ of $\ker(\mathbf{A})$ is contained in $\ker(F(\mathbf{A}))$; we show equality next. Recall that $F(\mathbf{A})$ is a π^\dagger -prefrieze (by Lemma 5.17), and so the dimension of the kernel of $F(\mathbf{A})$ equals the number of balls in π^\dagger (by Lemma 5.8.4), which is $n - k$. Since \mathbf{A} is $k \times n$ with rank k , the dimension of $\ker(\mathbf{A})$ is $n - k$. Since $q_{(k,n)} : \mathbb{k}^n \rightarrow \mathbb{k}^{\mathbb{Z}}$ is an inclusion, $q_{(k,n)}(\ker(\mathbf{A}))$ has the same dimension as $\ker(F(\mathbf{A}))$, and so the two spaces are equal. \square

²²It is also not too hard to show directly that $\det(\mathbf{A}_{L_a \setminus \{a\} \cup \{b\}}) = 0$ whenever $a - n < b < \pi^\dagger(a)$ or $\pi^\dagger(b) < a < b + n$.

Theorem F. *If A is a π -unimodular matrix, then $F(A)$ is a π^\dagger -frieze.*

Proof of Theorem F. By Lemma 5.17, $F(A)$ is a π^\dagger -prefrieze, and by Lemma 5.18, it has superperiodic kernel. By Theorem B, $F(A)$ is a π^\dagger -frieze. \square

5.5. Inverting F (Theorem G). In this section, we show that every juggler's frieze can be constructed using F . First, a useful formula relating minors of A to projections of $\ker(A)$.

Lemma 5.19. *Let A be a $k \times n$ matrix of rank k . Then for any subset $I \subset [n]$,*

$$\text{rank}(A_I) = \dim(\ker(A)_{[n] \setminus I}) + |I| - (n - k)$$

where $\ker(A)_{[n] \setminus I}$ is the image of $\ker(A)$ under the restriction map $\mathbb{k}^n \rightarrow \mathbb{k}^{[n] \setminus I}$.

Proof. Since the restriction map $\mathbb{k}^n \rightarrow \mathbb{k}^{[n] \setminus I}$ is the quotient map by the subspace $\mathbb{k}^I \subset \mathbb{k}^n$,

$$\ker(A)_{[n] \setminus I} \simeq \frac{\ker(A)}{\ker(A) \cap \mathbb{k}^I}$$

Since $\ker(A) \cap \mathbb{k}^I \simeq \ker(A_I)$, applying the Rank-Nullity Theorem gives the following

$$\dim(\ker(A)_{[n] \setminus I}) = \dim(\ker(A)) - \overbrace{\dim(\ker(A_I))}^{=k} = (n - \text{rank}(A)) - (|I| - \text{rank}(A_I)) \quad \square$$

Lemma 5.20. *Let π be an n -periodic juggling function with k -many balls and let C be a π^\dagger -frieze. If A is a $k \times n$ matrix such that, for all vectors $v \in \mathbb{k}^n$,*

$$Av = 0 \Leftrightarrow Cq_{(k,n)}(v) = 0$$

then there is a π -unimodular matrix A' with the same kernel as A and $F(A') = C$.

Proof. Let $[a, b] \subset \mathbb{Z}$ be any interval with $b - a < n$, and let $\overline{[a, b]} \subset [n]$ denote the residues mod n . By Lemma 5.19,

$$\text{rank}(A_{\overline{[a, b]}}) = \dim(\ker(A)_{[n] \setminus \overline{[a, b]}}) + (b - a + 1) - (n - k)$$

Since $[n] \setminus \overline{[a, b]} = \overline{[b + 1, a + n - 1]}$,

$$\dim(\ker(A)_{[n] \setminus \overline{[a, b]}}) = \dim(\ker(A)_{\overline{[b + 1, a + n - 1]}}) = \dim(\ker(C)_{[b + 1, a + n - 1]})$$

By [Mul19, Prop. 7.12], this is equal to the number of π^\dagger -balls in the interval $[b + 1, a + n - 1]$; i.e.

$$\begin{aligned} \dim(\ker(C)_{[b + 1, a + n - 1]}) &= |\{i \in \mathbb{Z} \mid b < i < a + n \text{ and } a + n \leq \pi^\dagger(i)\}| \\ &= |\{i \in \mathbb{Z} \mid b < i < a + n \text{ and } a \leq \pi^{-1}(i)\}| \\ &= (a + n - b - 1) - |\{i \in \mathbb{Z} \mid b < i < a + n \text{ and } a > \pi^{-1}(i)\}| \\ &= (a + n - b - 1) - |L_a \cap [b + 1, a + n - 1]| \\ &= (a + n - b - 1) - (|L_a| - |L_a \cap [a, b]|) \end{aligned}$$

Since $|L_a| = k$, the previous formulas yield that $\text{rank}(A_{\overline{[a, b]}}) = |L_a \cap [a, b]|$; that is, A satisfies the second condition of being π -unimodular.

Define a $\mathbb{Z} \times \mathbb{Z}$ -matrix $F(\mathbf{A})$ via the formula (identical to 4.1 but without assuming π -unimodular)

$$F(\mathbf{A})_{a,b} := \begin{cases} \tau(\mathbf{A})_{\bar{a}} \cdot \mathbf{A}_{\bar{b}} & \text{if } b \leq a < b+n, \bar{b} \leq \bar{a}, \text{ and } a < \pi(a), \\ (-1)^{k-1} \tau(\mathbf{A})_{\bar{a}} \cdot \mathbf{A}_{\bar{b}} & \text{if } b \leq a < b+n, \bar{b} > \bar{a}, \text{ and } a < \pi(a), \\ 1 & \text{if } \pi(a) = a = b, \\ (-1)^k & \text{if } \pi(a) = a = b+n, \\ 0 & \text{otherwise.} \end{cases}$$

If $a > \pi^\dagger(b)$, then either $a \geq b+n$ or $a < b+n$. In the first case, $F(\mathbf{A})_{a,b}$ is defined to be 0. In the second case, $\pi^{-1}(b) < a-n < b$, and so $\bar{b} \in \overline{L_a} \setminus \{\bar{a}\}$. Therefore, $F(\mathbf{A})_{a,b} = \pm(\tau(\mathbf{A})_{\bar{a}} \cdot \mathbf{A}_{\bar{b}}) = 0$. If $b < (\pi^\dagger)^{-1}(a)$, then either $a \geq b+n$ or $a < b+n$. In the first case, $F(\mathbf{A})_{a,b}$ is defined to be 0. In the second case, $a < b+n < \pi(a)$, and so $\tau(\mathbf{A})_{\bar{a}} \cdot \mathbf{A}_{\bar{b}} = 0$ by [MS17, Lemma 6.4], and $F(\mathbf{A})_{a,b} = 0$.

Therefore, $F(\mathbf{A})$ is a *reduced recurrence matrix* in the sense of [Mul19]. Since \mathbf{C} is another reduced recurrence matrix with the same kernel, $F(\mathbf{A}) = \mathbf{C}$ by [Mul19, Theorem 6.5]. Then for any $b \in \mathbb{Z}$,

$$(-1)^{|S_\dagger(b, \pi^\dagger(b))|} = \mathbf{C}_{\pi^\dagger(b), b} = F(\mathbf{A})_{\pi^\dagger(b), b}$$

Assuming $\pi^\dagger(b) < b+n$, then $\pi(\pi^\dagger(b)) > \pi^\dagger(b)$ and $\overline{\pi^\dagger(b)} = \pi^{-1}(\bar{b})$ so

$$(-1)^{|S_\dagger(b, \pi^\dagger(b))|} = \begin{cases} \tau(\mathbf{A})_{\pi^\dagger(\bar{b})} \cdot \mathbf{A}_{\bar{b}} & \text{if } \bar{b} \leq \bar{a}, \\ (-1)^{k-1} \tau(\mathbf{A})_{\pi^\dagger(\bar{b})} \cdot \mathbf{A}_{\bar{b}} & \text{if } \bar{b} > \bar{a}. \end{cases}$$

By an analogous computation to the proof of Lemma 5.14,

$$\tau(\mathbf{A})_{\pi^\dagger(\bar{b})} \cdot \mathbf{A}_{\bar{b}} = (-1)^{|S_\dagger(b, \pi^\dagger(b))|} \begin{cases} \frac{\det(\mathbf{A}_{L_{b-1}})}{\det(\mathbf{A}_{L_b})} & \text{if } \bar{b} \leq \bar{a}, \\ (-1)^{k-1} \frac{\det(\mathbf{A}_{L_{b-1}})}{\det(\mathbf{A}_{L_b})} & \text{if } \bar{b} > \bar{a}. \end{cases}$$

Therefore, $\det(\mathbf{A}_{L_{b-1}}) = \det(\mathbf{A}_{L_b})$. In the missing case when $\pi^\dagger(b) = b+n$, $L_{b-1} = L_b$ and so $\det(\mathbf{A}_{L_{b-1}}) = \det(\mathbf{A}_{L_b})$. It follows that $\det(\mathbf{A}_{L_a}) = \det(\mathbf{A}_{L_b})$ for all $a, b \in \mathbb{Z}$. Choose any $k \times k$ matrix \mathbf{G} with $\det(\mathbf{G}) = \det(\mathbf{A}_{L_a})^{-1}$ and set $\mathbf{A}' := \mathbf{G}\mathbf{A}$. Then \mathbf{A}' is π -unimodular and $\ker(\mathbf{A}') = \ker(\mathbf{A})$. \square

Theorem G. *For each juggling function π with k -many balls, the map F descends to a bijection*

$$\mathrm{SL}(k) \setminus \{\pi\text{-unimodular matrices}\} \xrightarrow{F} \{\pi^\dagger\text{-friezes}\}$$

Proof. Lemma 5.20 gives an explicit preimage of each π^\dagger -frieze, so F is surjective.

Next, we show that F is injective. Let \mathbf{A} and \mathbf{A}' be π -unimodular matrices with $F(\mathbf{A}) = F(\mathbf{A}')$. Then $\ker(F(\mathbf{A})) = \ker(F(\mathbf{A}'))$, and so Lemma 5.18 implies $\ker(\mathbf{A}) = \ker(\mathbf{A}')$. Then there is an invertible $k \times k$ -matrix \mathbf{G} such that $\mathbf{G}\mathbf{A} = \mathbf{A}'$. Choosing any a ,

$$\det(\mathbf{A}'_{L_a}) = \det((\mathbf{G}\mathbf{A})_{L_a}) = \det(\mathbf{G}) \det(\mathbf{A}_{L_a})$$

By π -unimodularity, $\det(\mathbf{G}) = 1$, and so \mathbf{A} and \mathbf{A}' are in the same $\mathrm{SL}(k)$ -orbit. \square

5.6. Duality and the twist (Theorem H). In this section, we demonstrate the relation between the twist, the positive complement, and the dual of a frieze.

Proposition 5.21. *Let \mathbf{A} be π -unimodular, and let \mathbf{A}^\dagger be a positive complement of \mathbf{A} . Then $\tau(\mathbf{A})$ is a positive complement to $\tau^{-1}(\mathbf{A}^\dagger)$.*

Equivalently, we may write $\tau(\mathbf{A})^\dagger \equiv \tau^{-1}(\mathbf{A}^\dagger)$ (modulo left multiplication by $\mathrm{SL}(k)$).

Proof. This proof uses notation and terminology from [MS17] that has not been introduced and won't be used elsewhere. Consider a reduced plabic graph Γ with juggling function π , and assume there is an edge weighting $w \in (\mathbb{k}^\times)^E$ of Γ whose image $\mathbb{D}(w)$ under the boundary measurement map is the rowspan of \mathbf{A} . By [MS17, Theorem 7.1],

$$\overset{\bullet \rightarrow}{\mathbb{F}}(\tau(\mathbf{A})) = \overset{\bullet \rightarrow}{\mathbb{F}} \tau \mathbb{D}(w) = \overrightarrow{\mathbb{M}}(w)$$

where $\overset{\bullet \rightarrow}{\mathbb{F}}$ denote the source-labeled Plücker coordinates of Γ and $\overrightarrow{\mathbb{M}}(w)$ is the face weighting of Γ determined by the downstream wedges of the edges in w .

Let Γ^\dagger denote the reduced plabic graph obtained by swapping the colors of the vertices of Γ . The edge weighting w may be identified with an edge weighting of Γ^\dagger , and the image $\mathbb{D}^\dagger(w)$ under the boundary measurement map of Γ^\dagger is the rowspan of \mathbf{A}^\dagger . By [MS17, Theorem 7.1],

$$\overset{\bullet \leftarrow \dagger}{\mathbb{F}}(\tau^{-1}(\mathbf{A}^\dagger)) = \overset{\bullet \leftarrow \dagger}{\mathbb{F}} \tau^{-1} \mathbb{D}^\dagger(w) = \overleftarrow{\mathbb{M}}^\dagger(w)$$

where $\overset{\bullet \leftarrow \dagger}{\mathbb{F}}$ denote the target-labeled Plücker coordinates of Γ^\dagger and $\overleftarrow{\mathbb{M}}^\dagger(w)$ is the face weighting of Γ^\dagger determined by the upstream wedges of the edges in w . Since the upstream wedges in Γ^\dagger are the downstream wedges in Γ , $\overleftarrow{\mathbb{M}}^\dagger(w) = \overrightarrow{\mathbb{M}}(w)$ and so

$$\overset{\bullet \rightarrow}{\mathbb{F}}(\tau(\mathbf{A})) = \overset{\bullet \leftarrow \dagger}{\mathbb{F}}(\tau^{-1}(\mathbf{A}^\dagger))$$

Since the target labellings of Γ^\dagger are the complements of the source labellings of Γ ,

$$\overset{\bullet \leftarrow \dagger}{\mathbb{F}}(\tau^{-1}(\mathbf{A}^\dagger)) = \overset{\bullet \rightarrow}{\mathbb{F}}(\tau^{-1}(\mathbf{A}^\dagger)^\dagger)$$

where $\tau^{-1}(\mathbf{A}^\dagger)^\dagger$ is a positive complement to $\tau^{-1}(\mathbf{A}^\dagger)$. Since $\overset{\bullet \rightarrow}{\mathbb{F}}$ is bijective on the image of $\tau \circ \mathbb{D}$ [MS17, Theorem 7.1], $\tau(\mathbf{A})$ is a positive complement of $\tau^{-1}(\mathbf{A}^\dagger)$.

This verifies the proposition when the rowspan of \mathbf{A} is in the image of the boundary measurement map of Γ . By [MS17, Prop. 7.6], this set is dense in the entire positroid variety, and so the proposition holds in general by continuity. \square

Lemma 5.22. *If \mathbf{A} is a π -unimodular $k \times n$ -matrix, then for all $a, b \in \mathbb{Z}$,*

$$\mathrm{Tiling}(\mathbf{F}(\mathbf{A}))_{a,b} = (-1)^{a+b+(k-1)\frac{a+b-\bar{a}-\bar{b}}{n}} \tau(\mathbf{A})_{\bar{a}} \cdot \mathbf{A}_{\bar{b}}$$

Proof. Since $\mathbf{F}(\mathbf{A})$ is a π^\dagger -frieze with $(n-k)$ -many balls, the (a,b) th entry of the tiling is defined as

$$\mathrm{Tiling}(\mathbf{F}(\mathbf{A}))_{a,b} = \sum_{i \in \mathbb{Z}} (-1)^{a+b+i(k-1)} \mathbf{F}(\mathbf{A})_{a+in,b} \quad (5.6)$$

The latter sum can be evaluated using the cases in Corollary 5.15.

If $\pi^\dagger(a) < a+n$ (that is, a is not a loop of π), then every term in the sum (5.6) must vanish except for the $i \in \mathbb{Z}$ such that $b \leq a+in < b+n$. Equivalently, $1 \leq a-b+1+in \leq n$, and so

$i = \frac{(a-b+1)-\overline{a-b+1}}{n}$. Corollary 5.15 gives the value of the remaining term as

$$\text{Tiling}(\mathbf{F}(\mathbf{A}))_{a,b} = (-1)^{a+b+i(k-1)} \begin{cases} \tau(\mathbf{A})_{\overline{a}} \cdot \mathbf{A}_{\overline{b}} & \text{if } \overline{b} \leq \overline{a}, \\ (-1)^{k-1} \tau(\mathbf{A})_{\overline{a}} \cdot \mathbf{A}_{\overline{b}} & \text{if } \overline{b} > \overline{a}. \end{cases}$$

If $\overline{b} \leq \overline{a}$, then $\overline{a-b+1} = \overline{a} - \overline{b} + 1$, and so $i = \frac{(a-b)-(\overline{a}-\overline{b})}{n}$. If $\overline{b} > \overline{a}$, $\overline{a-b+1} = (\overline{a} - \overline{b} + 1) + n$, and so $i = \frac{(a-b)-(\overline{a}-\overline{b})}{n} - 1$. Therefore, the lemma holds whenever $\pi^\dagger(a) < a + n$.

If $\pi^\dagger(a) = a + n$, then $\pi(a) = a$ and so $\mathbf{A}_{\overline{a}} = 0$ and $\tau(\mathbf{A})_{\overline{a}} = 0$. If $b \not\equiv a \pmod{n}$, then every term in the sum (5.6) vanishes. If $b = a + in$ for some i , then the sum (5.6) has two terms, and

$$\begin{aligned} \text{Tiling}(\mathbf{F}(\mathbf{A}))_{a,b} &= (-1)^{a+b+i(k-1)} \mathbf{F}(\mathbf{A})_{a+in,b} + (-1)^{a+b+(i+1)(k-1)} \mathbf{F}(\mathbf{A})_{a+(i+1)n,b} \\ &= (-1)^{a+b+i(k-1)} (1) + (-1)^{a+b+(i+1)(k-1)} (-1)^k = 0 \end{aligned}$$

Therefore, the lemma holds whenever $\pi^\dagger(a) = a + n$. \square

Theorem H. *Let \mathbf{A} be a π -unimodular matrix. Then*

$$\mathbf{F}(\mathbf{A})^\dagger = \mathbf{F}(\tau(\mathbf{A})^\dagger) = \mathbf{F}(\tau^{-1}(\mathbf{A}^\dagger))$$

Proof. Let \mathbf{A} be a π -unimodular $k \times n$ -matrix, and let \mathbf{B} be the negation of the odd numbered columns of \mathbf{A} , so that the rows of \mathbf{B} span the kernel of \mathbf{A}^\dagger . By Prop. 5.21, $\tau(\mathbf{A})$ is a positive complement of $\tau^{-1}(\mathbf{A}^\dagger)$ and the rows of $\tau(\mathbf{B})$ span the kernel of $\tau^{-1}(\mathbf{A}^\dagger)$.

Given a $k \times n$ -matrix \mathbf{M} , let $q_{(n-k,n)}(\mathbf{M})$ denote the $k \times \mathbb{Z}$ -matrix whose rows are the $(n-k, n)$ superperiodic extensions of the rows of \mathbf{M} ; that is,

$$q_{(n-k,n)}(\mathbf{M})_{a,b} = (-1)^{(n-k-1)\frac{b-\overline{b}}{n}} \mathbf{M}_{a,\overline{b}}$$

Since $\tau^{-1}(\mathbf{A}^\dagger)$ is $(n-k) \times n$, by Lemma 5.18, the rows of $q_{(n-k,n)}(\tau(\mathbf{B}))$ span the kernel of $\mathbf{F}(\tau^{-1}(\mathbf{A}^\dagger))$. By [Mul19, Theorem 9.6], the solution matrix of this frieze factors as

$$\text{Sol}(\mathbf{F}(\tau(\mathbf{A}^\dagger))) = q_{(n-k,n)}(\tau(\mathbf{B}))^\top q_{(n-k,n)}(\mathbf{B})$$

The (a, b) th entry of $\text{Sol}(\mathbf{F}(\tau(\mathbf{A}^\dagger)))$ is therefore

$$\begin{aligned} \left(q_{(n-k,n)}(\tau(\mathbf{B}))^\top q_{(n-k,n)}(\mathbf{B}) \right)_{a,b} &= q_{(n-k,n)}(\tau(\mathbf{B}))_a \cdot q_{(n-k,n)}(\mathbf{B})_b \\ &= \left((-1)^{(n-k-1)\frac{a-\overline{a}}{n}} \tau(\mathbf{B})_{\overline{a}} \right) \cdot \left((-1)^{(n-k-1)\frac{b-\overline{b}}{n}} \tau(\mathbf{B})_{\overline{b}} \right) \\ &= \left((-1)^{(n-k-1)\frac{a-\overline{a}}{n} + \overline{a}} \tau(\mathbf{A})_{\overline{a}} \right) \cdot \left((-1)^{(n-k-1)\frac{b-\overline{b}}{n} + \overline{b}} \mathbf{A}_{\overline{b}} \right) \\ &= (-1)^{(n-k-1)\frac{a+b-\overline{a}-\overline{b}}{n} + \overline{a} + \overline{b}} (\tau(\mathbf{A})_{\overline{a}} \cdot \mathbf{A}_{\overline{b}}) \\ &= (-1)^{(k-1)\frac{a+b-\overline{a}-\overline{b}}{n} + a+b} (\tau(\mathbf{A})_{\overline{a}} \cdot \mathbf{A}_{\overline{b}}) \end{aligned}$$

By Lemma 5.22, this coincides with the (a, b) th entry of $\text{Tiling}(\mathbf{F}(\mathbf{A}))$. With Proposition 5.12,

$$\text{Sol}(\mathbf{F}(\tau^{-1}(\mathbf{A}^\dagger))) = \text{Tiling}(\mathbf{F}(\mathbf{A})) = \text{Sol}(\mathbf{F}(\mathbf{A})^\dagger)$$

Since the friezes $\mathbf{F}(\tau^{-1}(\mathbf{A}^\dagger))$ and $\mathbf{F}(\mathbf{A})^\dagger$ have the same solution matrix, they have the same kernel by Lemma 5.8.4, and so they are equal by Lemma 5.8.1. \square

Appendix A. Collected notational conventions

We collect a number of notational conventions here for the reader's convenience.

- \mathbb{k} denotes an arbitrary choice of field.
- For an integer $n > 0$, $[n]$ denotes the interval of consecutive integers from 1 to n ; that is,

$$[n] := \{1, 2, 3, \dots, n\}$$

- For $a, b \in \mathbb{Z}$, $[a, b]$ denotes the interval of consecutive integers from a to b ; that is,

$$[a, b] := \{a, a + 1, a + 2, \dots, b\}$$

Similarly, $[a, b)$, $(a, b]$, and (a, b) denote consecutive integers save one or both of a and b .

- When a choice of n is clear from context, we use the following notation.
 - For each $a \in \mathbb{Z}$, \bar{a} denotes the unique integer in $[n]$ congruent to $a \pmod n$.
 - For each $I \subset \mathbb{Z}$, \bar{I} denotes the set of integers in $[n]$ congruent to elements of $I \pmod n$.
- When A is a $k \times n$ -matrix, we use the following notation.
 - Given $a \in [n]$, A_a denotes the a th column of A .
 - Given $a \in \mathbb{Z}$, $A_{\bar{a}}$ denotes the \bar{a} th column of A .
 - Given $I \subset [n]$, A_I denotes the $k \times |I|$ -submatrix of A whose columns are indexed by I .
 - Given $I \subset \mathbb{Z}$, $A_{\bar{I}}$ denotes the $k \times |\bar{I}|$ -submatrix of A whose columns are indexed by \bar{I} .

Warning: The columns of $A_{\bar{I}}$ are ordered according to the indices order in \bar{I} , not in I .

Acknowledgements. This paper owes its existence to conversations with two people.

- David Speyer, with whom the second author discovered several curious properties of the twist; most notably the $\mathrm{GL}(k)$ -invariance of the function $A \mapsto \tau(A)^\top A$.
- Karin Baur, whose talks on their work in [BFG⁺18, BFG⁺21] connected these curious properties to friezes and inspired this project.

We are also grateful to Emily Gunawan and Khrystyna Serhiyenko for helpful conversations.

References

- [BFG⁺18] Karin Baur, Eleonore Faber, Sira Gratz, Khrystyna Serhiyenko, and Gordana Todorov, *Mutation of friezes*, Bull. Sci. Math. **142** (2018), 1–48. (cit. on pp. 1 and 38)
- [BFG⁺21] ———, *Friezes satisfying higher SL_k -determinants*, Algebra Number Theory **15** (2021), no. 1, 29–68. (cit. on pp. 1, 3, 7, 10, and 38)
- [BR10] François Bergeron and Christophe Reutenauer, *SL_k -tilings of the plane*, Illinois J. Math. **54** (2010), no. 1, 263–300. (cit. on p. 3)
- [CC73] John H. Conway and Harold S. M. Coxeter, *Triangulated polygons and frieze patterns*, Math. Gaz. **57** (1973), no. 400, 87–94. (cit. on pp. 1 and 3)
- [CC06] Philippe Caldero and Frédéric Chapoton, *Cluster algebras as Hall algebras of quiver representations*, Comment. Math. Helv. **81** (2006), no. 3, 595–616. (cit. on p. 1)
- [Cox71] Harold S. M. Coxeter, *Frieze patterns*, Acta Arith. **18** (1971), 297–310. (cit. on pp. 1 and 3)
- [CR72] Craig M. Cordes and D. P. Roselle, *Generalized frieze patterns*, Duke Math. J. **39** (1972), 637–648. (cit. on p. 1)
- [DM] Roi Docampo and Greg Muller, *Spaces of quasiperiodic sequences*. In preparation. (cit. on pp. 1, 21, and 22)
- [GL19] Pavel Galashin and Thomas Lam, *Positroid varieties and cluster algebras*, posted on 2019, available at [arXiv:1906.03501](https://arxiv.org/abs/1906.03501). (cit. on p. 22)

- [Gal21] Pavel Galashin, *Critical varieties in the Grassmannian*, posted on 2021, available at [arXiv:2102.13339](https://arxiv.org/abs/2102.13339). (cit. on p. 21)
- [KV15] Rinat Kedem and Panupong Vichitkunakorn, *T-systems and the pentagram map*, J. Geom. Phys. **87** (2015), 233–247. (cit. on p. 3)
- [KLS13] Allen Knutson, Thomas Lam, and David E. Speyer, *Positroid varieties: juggling and geometry*, Compos. Math. **149** (2013), no. 10, 1710–1752. (cit. on pp. 11, 18, and 21)
- [Lec16] Bernard Leclerc, *Cluster structures on strata of flag varieties*, Adv. Math. **300** (2016), 190–228. (cit. on p. 22)
- [MG15] Sophie Morier-Genoud, *Coxeter’s frieze patterns at the crossroads of algebra, geometry and combinatorics*, Bull. Lond. Math. Soc. **47** (2015), no. 6, 895–938. (cit. on pp. 1, 3, and 30)
- [MGOST14] Sophie Morier-Genoud, Valentin Ovsienko, Richard Evan Schwartz, and Serge Tabachnikov, *Linear difference equations, frieze patterns, and the combinatorial Gale transform*, Forum Math. Sigma **2** (2014), Paper No. e22, 45. (cit. on pp. 1, 3, 6, 7, 10, and 30)
- [MS16] Bethany R. Marsh and Jeanne S. Scott, *Twists of Plücker coordinates as dimer partition functions*, Comm. Math. Phys. **341** (2016), no. 3, 821–884. (cit. on pp. 1 and 8)
- [MS17] Greg Muller and David E. Speyer, *The twist for positroid varieties*, Proc. Lond. Math. Soc. (3) **115** (2017), no. 5, 1014–1071. (cit. on pp. 8, 33, 35, and 36)
- [Mul19] Greg Muller, *Linear recurrences indexed by \mathbb{Z}* , posted on 2019, available at [arXiv:1906.04311](https://arxiv.org/abs/1906.04311). (cit. on pp. 26, 27, 28, 30, 34, 35, and 37)
- [Pos06] Alexander Postnikov, *Total positivity, Grassmannians, and networks*, posted on 2006, available at [arXiv:math/0609764](https://arxiv.org/abs/math/0609764). (cit. on p. 22)
- [Sco06] J. S. Scott, *Grassmannians and cluster algebras*, Proc. London Math. Soc. (3) **92** (2006), no. 2, 345–380. (cit. on p. 22)
- [Sha84] Louis W. Shapiro, *Positive definite matrices and Catalan numbers, revisited*, Proc. Amer. Math. Soc. **90** (1984), no. 3, 488–496. (cit. on p. 1)
- [SSBW20] Khrystyna Serhiyenko, Melissa Sherman-Bennett, and Lauren Williams, *Combinatorics of cluster structures in Schubert varieties*, Sém. Lothar. Combin. **82B** (2020), Art. 8, 12. (cit. on p. 22)

Roi Docampo

Department of Mathematics
University of Oklahoma
601 Elm Avenue, Room 423
Norman, OK 73019 (USA)
roi@ou.edu

Greg Muller

Department of Mathematics
University of Oklahoma
601 Elm Avenue, Room 423
Norman, OK 73019 (USA)
gmuller@ou.edu