

Isotope Shifts in Cadmium as a Sensitive Probe for Physics Beyond the Standard Model

B. Ohayon*

Institute for Particle Physics and Astrophysics, ETH Zürich, CH-8093 Zürich, Switzerland

S. Hofsäss, J.E. Padilla-Castillo, S. C. Wright, G. Meijer, and S. Truppe†

Fritz-Haber-Institut der Max-Planck-Gesellschaft, Faradayweg 4-6, 14195 Berlin, Germany

K. Gibble‡

Department of Physics, Pennsylvania State University University Park, PA, 16802 USA

B. K. Sahoo§

Atomic and Molecular Physics Division, Physical Research Laboratory, Navrangpura, Ahmedabad 380009, India

Isotope shifts (ISs) in atomic energy levels are sensitive probes of nuclear structure and new physics beyond the Standard Model. We present an analysis of the ISs of the cadmium atom (Cd I) and singly charged cadmium ion (Cd II). ISs of the 229 nm, 326 nm, 361 nm and 480 nm lines of Cd I are measured with a variety of techniques; buffer-gas-cooled beam spectroscopy, capturing atoms in a magneto-optic-trap, and optical pumping. IS constants for the D₁ and D₂ lines of Cd II are calculated with high accuracy by employing analytical response relativistic coupled-cluster theory in the singles, doubles and triples approximations. Combining the calculations for Cd II with experiments, we infer IS constants for all low-lying transitions in Cd I. We benchmark these constants as calculated via different many-body methods. Our calculations for Cd II enable nuclear charge radii of Cd isotopes to be extracted with unprecedented accuracy. The combination of our precise calculations and measurements shows that King Plots for Cd I can improve the state-of-the-art sensitivity to a new heavy boson by up to two orders of magnitude.

I. INTRODUCTION

The most accurately determined quantities are transition frequencies of optical clocks [1], which are measured to better than 10^{-18} . This superb spectroscopic precision enables stringent searches for new physics [2–5]. One approach compares the isotope shifts (IS) of two or more transitions as in a King Plot (KP) [6], in which a deviation from a linear behaviour [7–10] can be a signature for beyond Standard Model (BSM) physics, such as a new boson mediating a force between electrons and neutrons. King Plot searches are applicable to systems possessing narrow optical transitions for which ISs can be measured with high precision, even with extraordinary sub-Hz accuracy using common-mode noise rejection [11, 12]. Searches for deviations from linearity with KPs require a minimum of four stable (or long-lived) even-even isotopes, which severely restricts the number of candidate elements.

BSM searches through KP non-linearity were performed for two elements in different regimes. On the low atomic number (Z) side are measurements in calcium ($Z = 20$); the lightest element with more than three

stable even-even isotopes. A KP comprised of two transitions with different relativistic effects for Ca II, with a characteristic precision of 20 Hz, shows no non-linearity, translating directly to limits on several BSM scenarios [13]. As experimental precision improves, the potential for new physics searches using KPs in light elements such as Ca is expected to be limited by the difficulty of calculating the standard model (SM) contribution to the KP non-linearity resulting from high-order recoil effects in a light many-electron system [14, 15].

In the high Z regime, the effects of a new massive boson are more pronounced [7, 15], while recoil effects are heavily suppressed. However, other sources of SM non-linearity become dominant making the interpretation of an observed non-linearity more involved. For Yb ($Z = 70$), ISs in several ionic and atomic lines were measured with high-precision [16–19]. A recent analysis encompassing all of the experimental data indicates a strong non-linearity attributed to nuclear deformation [19], which currently cannot be calculated *ab initio* with sufficient precision. This behaviour is particularly pronounced and follows from Yb isotopes being amongst the most deformed in the stable region of the nuclear chart [20]. A non-linearity from an additional source was identified, whose origin is currently being studied [17, 19]. Considering the cases above, once a non-linearity is observed, it is not straightforward to interpret it as a signature of new physics. Pronounced non-linearities in several systems will add clarity. Moreover, there is a trade-off between lower values of Z , where both the non-linearities

* bohayon@ethz.ch

† truppe@fhi-berlin.mpg.de. Current address: Centre for Cold Matter, Blackett Laboratory, Imperial College London, Prince Consort Road, London SW7 2AZ, United Kingdom

‡ kgibble@psu.edu

§ bijaya@prl.res.in




FIG. 1: a: Low-lying energy levels of Cd I. The isotope shifts of the indicated transitions are discussed in this work. Broken lines represent transitions whose ISs are deduced from combinations of the ISs of measured ISs. b: Magnitudes of the difference of field shift constants. The top boxes corresponds to the 69% confidence interval. The transitions are ordered for clarity, with the narrow transitions in bold, and the 214 nm transition is for Cd II.

arising from the Standard Model (SM) and those that come from new physics effects are less pronounced; and higher values of Z , where several sources of non-linearity may be difficult to disentangle due theoretical intractability.

Cadmium ($Z = 48$) sits between these two limits. Having a relatively simple atomic structure and narrow transitions [21, 22], it is a prime candidate for new physics searches via KP non-linearities. Being close to the $Z = 50$ proton shell closure makes Cd nuclei much less deformed than e.g. Yb, suppressing a major source of SM non-linearity. Cd possesses six stable even-even isotopes, which could allow up to three different sources of non-linearity to be identified. In this work, we present measurements of ISs in the neutral cadmium atom (Cd I). We combine these measurements with literature values and highly accurate calculations of IS constants for transitions in the singly charged cadmium ion (Cd II). This enables us to map out all of the IS constants of the low-lying transitions in Cd I shown in Fig. 1, and identify the promising combinations for BSM searches. We compare the obtained IS constants with recent calculations and discuss the current state of the art in calculating these constants for two-valence systems. Such calculations are needed to assess the sensitivity of KPs to specific new physics models. We also combine calculated IS constants with measurements in a long chain of short-lived isotopes to yield highly accurate charge radii differences, which are needed to determine the nuclear contributions to KP non-linearities [17, 23]. Finally, we discuss the prospects for searches for BSM physics via KPs in Cd in light of

our analysis.

II. THEORY

The IS $\delta\nu_i^{A,A'}$ between isotopes with mass numbers A, A' , at the precision of the measurements analyzed here, can be written as [24]

$$\delta\nu_i^{A,A'} = F_i \delta\lambda^{A,A'} + K_i \mu^{A,A'}. \quad (1)$$

Here $\mu^{A,A'} = 1/M_A - 1/M_{A'}$ the inverse nuclear mass difference, i denotes a particular atomic transition, and $\delta\lambda^{A,A'} = \sum_k S_{2k} (\delta r_c^{2k})^{A,A'}$ is the nuclear parameter [25], which is expanded in a series of even charge moments

$$r_c^k = \frac{\int r^k \rho(r) r^2 dr}{\int \rho(r) r^2 dr}. \quad (2)$$

For brevity, we denote the root mean square (RMS) charge radius $\sqrt{r_c^2}$ here as r_c . The constants F_i and K_i are the field shift and mass shift, which depend only on the transition in question at this level. To calculate these constants, the nuclear charge distribution can be approximated using a Fermi distribution

$$\rho(r) = \frac{\rho_0}{1 + e^{(r-b)/a}}, \quad (3)$$

where ρ_0 is a normalization constant, and b and a are determined from electron scattering experiments [26, 27].

The operator for the field shift is given by

$$F = - \sum_e \frac{\delta V_{nuc}(r_c, r_e)}{\delta \lambda} \approx - \sum_e \frac{\delta V_{nuc}(r_c, r_e)}{\delta \langle r_c^2 \rangle} (1 + f_\lambda) \quad (4)$$

where the electrostatic potential V_{nuc} is a function of r_c , the radial distance of an electron is r_e , and the sum is over all electrons of the system. The contribution from higher nuclear moments f_λ is estimated as

$$f_\lambda \approx -S_4 \frac{r_c^4}{r_e^2} - S_6 \frac{r_c^6}{r_e^2} = 2.8(3)\%, \quad (5)$$

where the Seltzer coefficients S_4 and S_6 are estimated for the 5S levels of Cd II [25], and the ratio of moments is taken from two parameterizations of electron scattering experiments [26, 27]. The uncertainty of f_λ is estimated from the model-dependence of the charge distribution of Eq. 3, and its variation between isotopes.

The mass shift constant can further be split into a normal mass shift (NMS) and specific mass shift (SMS). The operators to determine the NMS and SMS constants are defined in a relativistic framework as [28, 29]

$$K^{\text{NMS}} = \frac{1}{2} \sum_i \left(\vec{p}_i^2 - \frac{\alpha Z}{r_i} \vec{\alpha}_i^D \cdot \vec{p}_i - \frac{\alpha Z}{r_i} (\vec{\alpha}_i^D \cdot \vec{C}_i^1) \vec{C}_i^1 \cdot \vec{p}_i \right) \quad (6)$$

and

$$K^{\text{SMS}} = \frac{1}{2} \sum_{i \neq j} \left(\vec{p}_i \cdot \vec{p}_j - \frac{\alpha Z}{r_i} \vec{\alpha}_i^D \cdot \vec{p}_j - \frac{\alpha Z}{r_i} (\vec{\alpha}_i^D \cdot \vec{C}_i^1) (\vec{p}_j \cdot \vec{C}_j^1) \right), \quad (7)$$

where the p_i 's are the components of the momentum operator, α^D is the Dirac matrix, and C^1 is the Racah operator. We note that there are slight modifications in the definitions of the F , K^{NMS} and K^{SMS} operators if we include quantum electrodynamics (QED) interactions. These contributions are expected to be smaller than our reported uncertainties.

A linear relation results after applying Eq. 1 to the measured ISs of two transitions (i, j):

$$\delta \bar{\nu}_i^{A, A'} = F_{ij} \delta \bar{\nu}_j^{A, A'} + K_{ij} \quad (8)$$

with the modified ISs $\delta \bar{\nu}_i^{A, A'} = \delta \nu_i^{A, A'} / \mu^{A, A'}$, the slope $F_{ij} = F_i / F_j$, and the offset $K_{ij} = K_i - F_{ij} K_j$. In this work we find that the linear relation of Eq. 8 holds at the \sim MHz-level for all Cd transition pairs. It may therefore be used to project IS constants from one transition to another. We calculate IS constants for transitions in Cd II using the analytical response (AR) relativistic coupled-cluster (RCC) method, and project them using Eq. 8 to atomic transitions.

III. THE AR-RCC METHOD

We begin with the Dirac-Coulomb (DC) Hamiltonian:

$$H_{DC} = \sum_i \left[c \vec{\alpha}_i^D \cdot \vec{p}_i + (\beta_i - 1) c^2 + V_{nuc}(r_i) \right] + \sum_{i, j > i} \frac{1}{r_{ij}}, \quad (9)$$

where c is the speed of light, $\vec{\alpha}^D$ and β are the Dirac matrices, \vec{p} is the single particle momentum operator. $\sum_{i, j} \frac{1}{r_{ij}}$ represents the Coulomb potential between the electrons located at the i^{th} and j^{th} positions. Corrections from the Breit and QED interactions are estimated by adding the corresponding potential terms as in Ref. [30], yielding the atomic Hamiltonian H_0 .

We consider the nuclear charge distribution of Eq. (3) to define the potential V_{nuc} [31]

$$V_{nuc}(r) = - \frac{Z}{\mathcal{N}r} \times \begin{cases} \frac{1}{b} \left(\frac{3}{2} + \frac{a^2 \pi^2}{2b^2} - \frac{r^2}{2b^2} + \frac{3a^2}{b^2} P_2^+ \frac{6a^3}{b^3} (S_3 - P_3^+) \right) & \text{for } r_i \leq b \\ \frac{1}{r_i} \left(1 + \frac{a^2 \pi^2}{b^2} - \frac{3a^2}{b^3} P_2^- + \frac{6a^3}{b^3} (S_3 - P_3^-) \right) & \text{for } r_i > b, \end{cases} \quad (10)$$

with

$$\begin{aligned} \mathcal{N} &= 1 + \frac{a^2 \pi^2}{b^2} + \frac{6a^3}{b^3} S_3, \\ S_k &= \sum_{l=1}^{\infty} \frac{(-1)^{l-1}}{l^k} e^{-lb/a}, \\ P_k^\pm &= \sum_{l=1}^{\infty} \frac{(-1)^{l-1}}{l^k} e^{\pm l(r-b)/a}. \end{aligned} \quad (11)$$

We determine these constants for the ground state, $[4d^{10}]5s$, and the first two excited states, $[4d^{10}]5p_{1/2}$ and $[4d^{10}]5p_{3/2}$ of Cd II to study the ISs of its D₁ and D₂ lines.

In the RCC theory *ansatz*, the wave function of the above atomic states are constructed as [32–34]

$$|\Psi_v\rangle = e^T \{1 + S_v\} |\Phi_v\rangle, \quad (12)$$

where $|\Phi_v\rangle$ is a mean-field wave function from a Dirac-Hartree-Fock (DHF) treatment, and T and S_v are the excitation operators that account for electron correlations from the core orbitals and valence orbital, respectively. The subscript v denotes the valence orbital associated with the respective state. It is introduced to uniquely identify the states having the common closed-shell configuration $[4d^{10}]$. Considering the IS operators, F , K^{NMS} and K^{SMS} , denoted as O in the total Hamiltonian as $H = H_0 + \eta O$, we express the above wave function in the AR-RCC formulation as

$$|\Psi_v\rangle = |\Psi_v^{(0)}\rangle + \eta |\Psi_v^{(1)}\rangle + \mathcal{O}(\eta^2), \quad (13)$$

with an energy

$$E_v = E_v^{(0)} + \eta E_v^{(1)} + \mathcal{O}(\eta^2). \quad (14)$$

Here η is nominally equal to one and is introduced for the perturbation expansion in orders of O , denoted by the superscripts. $E_v^{(0)}$ corresponds to the contribution from H_0 and $E_v^{(1)}$ includes first-order contributions from O and electron correlations. The above can be implemented in the RCC theory by expanding the RCC operators as

$$T = T^{(0)} + \eta T^{(1)} + \mathcal{O}(\eta^2) \quad (15)$$

and

$$S_v = S_v^{(0)} + \eta S_v^{(1)} + \mathcal{O}(\eta^2). \quad (16)$$

The $\mathcal{O}(\eta^2)$ contributions are usually small and neglected in IS calculations. However, these non-linear contributions can be significant when probing BSM physics. The zeroth-order RCC operator amplitudes follow from

$$\langle \Phi_0^* | \bar{H}_0 | \Phi_0 \rangle = 0 \quad (17)$$

and

$$\langle \Phi_v^* | \{ (\bar{H}_0 - E_v^{(0)}) S_v^{(0)} \} + \bar{H}_0 | \Phi_v \rangle = 0, \quad (18)$$

where $\langle \Phi_{0,v}^* |$ denotes for all possible excited Slater determinants and $\bar{H}_0 = \left(H_0 e^{T^{(0)}} \right)_{conn}$, where the subscript *conn* denotes for only connected terms in the expansion. With these amplitudes of the RCC operators, we calculate the zeroth-order energies as

$$E_0^{(0)} = \langle \Phi_0 | \bar{H}_0 | \Phi_0 \rangle \quad (19)$$

and

$$E_v^{(0)} = \langle \Phi_v | \bar{H}_0 \{ 1 + S_v^{(0)} \} | \Phi_v \rangle, \quad (20)$$

where $E_0^{(0)}$ is the energy of the common closed-core of the considered atomic states, $[4d^{10}]$ for Cd II. In the actual calculations, we consider a normal-ordered Hamiltonian defined with respect to $[4d^{10}]$. Therefore, $E_v^{(0)}$ corresponds to the electron affinity, $E_v^{(0)} - E_0^{(0)}$, of the valence orbital relative to the $[4d^{10}]$ configuration. We note that Eqs. (18) and (20) are coupled.

In the AR-RCC theory, we calculate the desired FS, NMS and SMS constants as the first-order energy corrections $E_v^{(1)} \equiv \langle \Psi_v^{(0)} | O | \Psi_v^{(0)} \rangle$ using the following expression [30, 35]

$$E_v^{(1)} = \langle \Phi_v | \bar{H}_0 S_v^{(1)} + (\bar{H}_0 T^{(1)} + \bar{O}) \{ 1 + S_v^{(0)} \} | \Phi_v \rangle \quad (21)$$

where $\bar{O} = (O e^{T^{(0)}})_{conn}$. Here, the normal-ordered form of operators again yield calculated values relative to the contributions from the $[4d^{10}]$ configuration. The amplitudes of the first-order perturbed RCC operators are given by

$$\langle \Phi_0^* | \bar{H}_0 T^{(1)} + \bar{O} | \Phi_0 \rangle = 0. \quad (22)$$

and

$$\langle \Phi_v^* | (\bar{H}_0 - E_v^{(0)}) S_v^{(1)} + (\bar{H}_0 T^{(1)} + \bar{O}) \{ 1 + S_v^{(0)} \} | \Phi_v \rangle = 0. \quad (23)$$

We use Gaussian type orbitals (GTOs) [36] to construct the single particle DHF wave functions. The large and small radial components of the DHF orbitals, $P(r)$ and $Q(r)$, are expressed using these GTOs as

$$P(r) = \sum_{k=1}^{N_k} c_k^{\mathcal{L}} \zeta_{\mathcal{L}} r^l e^{-\eta_0 \gamma^k r^2} \quad (24)$$

and

$$Q(r) = \sum_{k=1}^{N_k} c_k^{\mathcal{S}} \zeta_{\mathcal{L}} \zeta_{\mathcal{S}} \left(\frac{d}{dr} + \frac{\kappa}{r} \right) r^l e^{-\eta_0 \gamma^k r^2}, \quad (25)$$

where l is the orbital quantum number, κ is the relativistic angular momentum quantum number, $c_k^{\mathcal{L}(\mathcal{S})}$ are the expansion coefficients, $\zeta_{\mathcal{L}(\mathcal{S})}$ are the normalization factors of GTOs, η_0 and γ are optimized GTO parameters for a given orbital, and N_k represents the number of GTOs. To construct the GTOs, we use $\eta_0 = 0.00715, 0.0057, 0.0072, 0.0052$ and 0.0072 for s, p, d, f and g orbitals, respectively, with corresponding γ values 1.92, 2.04, 1.97, 2.07 and 2.54. Since our orbitals are not bounded, we integrate to an upper radial limit of $r = 500$ a.u. on a grid using a 10-point Newton-Cotes Gaussian quadrature method. Our numerical calculations use exponential grids with a step-size of 0.016 a.u. and 1200 grid points, and the coefficients are determined by the Roothan equation in the relativistic framework. To reduce the required computational resources, we have limited the virtual space by considering all possible single and double excitations in the AR-RCC theory (AR-RCCSD method) for the 1–20s, 2–20p, 3–20d, 4–17f, and 5–14g orbitals. The AR-RCCSDT method adds triple excitations to the above single and double excitations only for the 16s, 16p, 14d, 10f, and 5g orbitals.

IV. CALCULATED IS CONSTANTS FOR CD II

An important test of *ab initio* methods to obtain accurate atomic wave functions is to compare the calculated energies with the measured ones. In Table Ia we present the calculated electron affinities of the ground and first two excited states of Cd II. From these values, we determined the excitation energies of the D₁ and D₂ transitions given in Table Ia. The RCCSDT results show that triple excitations contribute more to the electron affinities than to the excitation energies, and that the Breit and QED corrections are significant. The uncertainties of the calculated energies are estimated from the convergence. We reproduce the experimental transition energies to 0.2%, an order of magnitude of improvement over CI-MBPT calculations for these transitions [39]. The high accuracy with which experimental energies are reproduced using the RCC method validates the accuracy of our wave functions for the following IS constants calculation. However, accurate wavefunctions are a necessary but insufficient condition for obtaining accurate IS constants. Various many-body methods for calculating the operators produce significantly different results for the IS constants; the AR method performs well against stringent benchmarks [30, 48].

We present IS constants using the DHF, AR-RCCSD and AR-RCCSDT methods in Table I. The AR-RCCSDT uncertainty follows from the numerical convergence as well as a perturbative estimation of partial quadrupole

TABLE I: Cd II Energies and IS constants for the ground and first two excited states, and their transitions for the D₁ (226.5 nm) and D₂ (214.4 nm) lines.

(a) Electron affinities and transition energies (cm ⁻¹)					
Method	E_{5S}	$E_{5P1/2}$	$E_{5P3/2}$	E_{D1}	E_{D2}
DHF	124568	84903	82871	39666	41698
RCCSD	136013	91858	89357	44155	46657
RCCSDT	136717	92401	98974	44316	46843
+ Δ Breit	-83	-83	-50	0.	-33
+ Δ QED	-100	-4	-6	-97	-94
SUM	136533(200)	92314(150)	89818(150)	44218(97)	46715(104)
Exp [37, 38]	136374.7(1)	92238.7(1)	89756.2(1)	44136.1(1)	46618.6(1)

(b) Normal mass shift (GHz u)					
Method	$K_{NMS,5S}$	$K_{NMS,5P1/2}$	$K_{NMS,5P3/2}$	$K_{NMS,D1}$	$K_{NMS,D2}$
DHF	5952	3374	3163	2578	2789
AR-RCCSD	2113	1381	1339	732	774
AR-RCCSDT	2204(20)	1470(15)	1426(15)	734(7)	778(8)
+ Δ Breit	-4	-4	-2	0.	-2
+ Δ QED	-7(2)	0.	0.	-7(2)	-7(2)
SUM	2194(20)	1466(15)	1424(15)	727(8)	769(8)
Scaling	2243	1517	1476	726	767

(c) Specific Mass Shift (GHz u)					
Method	$K_{SMS,5S}$	$K_{SMS,5P1/2}$	$K_{SMS,5P3/2}$	$K_{SMS,D1}$	$K_{SMS,D2}$
DHF	-2775	-1753	-1488	-1022	-1287
AR-RCCSD	1288	123	256	1165	1032
AR-RCCSDT	1343(15)	129(5)	260(7)	1214(16)	1083(17)
+ Δ Breit	12	5	4	7	8
+ Δ QED	4(1)	-2(0.)	-2(0.)	6(1)	5(1)
SUM	1359(15)	132(5)	263(7)	1226(16)	1096(17)

(d) Total Mass Shift (GHz u)					
Method	K_{5S}	$K_{5P1/2}$	$K_{5P3/2}$	K_{D1}	K_{D2}
$K_{SMS}+K_{NMS}$	3552(25)	1598(16)	1686(17)	1954(18)	1866(19)
CI-MBPT [39]				1770(300)	1667(300)
CKP [39]					2199(507)

(e) Field Shift (MHz fm ⁻²)					
Method	F_{5S}	$F_{5P1/2}$	$F_{5P3/2}$	F_{D1}	F_{D2}
DHF	-4778	-59	-0.	-4719	-4778
AR-RCCSD	-6140	-177	-100	-5963	-6040
AR-RCCSDT	-6227(20)	-232(5)	-152(5)	-5995(21)	-6075(21)
+ Δ Breit	22	1	1	21	21
+ Δ QED	147(37)	4(1)	3(1)	143(36)	144(36)
- f_λ	-172(17)	0	0	-172(17)	-172(17)
SUM	-6230(45)	-277(5)	-149(5)	-6003(45)	-6082(45)
CI-MBPT [39]				-6067(300)	-6144(300)
CKP [39]					-6621(530)

excitations. An uncertainty of 25% is assigned to the QED corrections and 10% to the higher moment corrections. Focusing first on the NMS constants, they can be obtained in the nonrelativistic limit by invoking the Virial theorem [49]. This results in the scaling law

$K_{NMS} \approx E \cdot m_e$, where E is the experimental energy and m_e the electron mass in atomic units. However, for a medium mass system such as Cd II, it is not *a priori* clear how applicable this method is. Table Ib gives our results for K_{NMS} calculated using the corresponding rel-

TABLE II: Isotope shifts of Cd I in MHz, relative to ^{114}Cd . Derived values are in italics; for the 332 nm clock line, from this work and previous measurements of 468 nm ISs [40], and similarly for the narrow 314 nm transition using 509 nm ISs [41].

A	229 nm	326 nm	480 nm	361 nm	332 nm	314 nm
106	1818.1(3.5) ^a 1748(11) [42]	1911.2(3.3) ^a 1913.0(1.0) ^b 1921(26) [43, 44]	-798.5(1.0) ^c	-607.6(3.0) ^b	<i>1911.4(4.7)</i>	<i>1915.5(3.9)</i>
108	1336.5(3.4) ^a 1258(9) [42]	1399.4(3.3) ^a 1402.4(1.0) ^b 1410(40) [44]	-586.7(1.0) ^c	-447.4(3.0) ^b	<i>1404.5(3.7)</i>	<i>1403.9(3.0)</i>
110	865.0(3.3) ^a 826(6) [42] 906(35) [45]	909.3(3.3) ^a 914.7(1.0) ^b 909(13) [43, 46, 47]	-383.4(1.0) ^c	-293.9(1.0) ^b	<i>914.1(2.1)</i>	<i>913.7(2.1)</i>
112	407.5(3.3) ^a 392(5) [42] 396(30) [45]	426.3(3.3) ^a 429.9(1.0) ^b 403(11) [43, 46, 47]	-183.1(1.0) ^c	-142.2(1.0) ^b	<i>428.9(1.7)</i>	<i>429.7(1.6)</i>
116	-316.1(3.3) ^a -299(4) [42]	-326.9(3.3) ^a -321.5(1.0) ^b -279(12) [43, 47]	152.7(1.0) ^c	122.0(1.0) ^b	<i>-320.6(1.7)</i>	<i>-320.2(1.7)</i>

^a Beam measurement, ^b Blue MOT edge, ^c MOT optical pumping rate

ativistic operator in Eq. 6. We see that for the D_1 and D_2 transitions, the scaling law agrees with our calculation within the 1% numerical accuracy. This behaviour is attributed to strong cancellations of relativistic contributions to the NMS in transitions with the same principal quantum number [50]. For the ground state the scaling law is accurate to 2 – 3%. Triple excitations contribute significantly to the NMS constants for the electron affinities, even though these quantities are evaluated with a one-body operator. We thus expect that for similar transitions in lighter systems, for which the contribution of electron correlations to the NMS is more pronounced, using the scaling-law could be preferable to a fully relativistic calculation.

The specific mass shift is associated with the two-body operator of Eq. 7 and thus strongly affected by high-order electron correlations, which are challenging to estimate [51]. Even in modern calculations, an uncertainty of 10 – 20% in K_{SMS} is usually given (see e.g. [35, 39, 52–54]). Our results in Table Ic are quoted with an accuracy of 1 – 2%, attributed to higher-order electron correlations, which we estimate perturbatively. This is in contrast to K_{NMS} and the energies, whose uncertainties follow from numerical convergence. We see that the sign of K_{SMS} changes between the DHF and AR-RCCSD calculations. This points to the importance of strong electron correlations in K_{SMS} , with a 10% difference coming from triple excitations. Owing to the two-body nature of the SMS operator, triple excitations in the AR-RCC method take several months to calculate at a typical high-performance computing facility. The total mass shift constant $K = K_{\text{NMS}} + K_{\text{SMS}}$ in Table Id agrees well with, and is an order of magnitude more precise than, a recent CI-MBPT calculation for both ionic transitions [39], as well as a calibrated King plot (CKP) estimation combin-

ing muonic x-ray and electron scattering measurements with precise isotope shifts for the D_2 transition [39]. The contributions of the Breit and QED interaction to K are found to be negligible at the current level of precision.

The calculation of the field shift operator is considered robust for a number of systems. Various methods typically agree at the few percent level [16, 30, 48, 55–60]. In Cd II, F convergences quickly, with triple excitations contributing only 0.6%. This yields an 0.35% uncertainty of F at the AR-RCCSDT level, which could be reduced further with more computational resources. However, we find that QED effects, which are often not taken into account in such calculations, are not negligible. In fact, our final uncertainty for F is dominated by the systematic uncertainty associated with QED correction only being present at the Hamiltonian level. This is expected to be even more pronounced in heavier systems such as Yb, where calculations of F are given with sub-percent numerical precision [61]. Our results in Table Ie agree with the CI-MBPT and a CKP results of [39]. To our knowledge, this is the first calculation of F with sub-percent accuracy (computational and systematic) for a system with more than 19 electrons.

V. IS MEASUREMENTS IN THE CD ATOM

In this section we describe our IS measurements of four transitions of Cd I for all stable bosonic isotopes. Measurements are done with two experimental systems employing different frequency calibration procedures. A more detailed description of the experimental systems, as well as results for the fermionic isotopes and absolute wavenumbers, will be given in upcoming publications [62, 63]. Our results are given in Table II, along with




FIG. 2: Isotope shifts of the Cd I bosons at 326 nm, measured with a cryogenic helium-cooled atomic beam and a frequency-doubled dye laser. The fermionic hyperfine transitions are beyond this spectral range.

available previous measurements, which disagree significantly in several cases. The two independent experimental results at 326 nm agree to within 0.5 – 1.6 combined standard errors, whereas the disagreements with these previous measurements are as large as four combined errors.

A. Atomic Beam Measurements of 229 nm and 326 nm ISs

We carried out laser induced fluorescence spectroscopy with a buffer-gas-cooled beam. A pulsed beam is produced by laser ablation of a Cd target, which is mounted in a buffer gas cell. Cryogenic helium gas (3 K) flows continuously through the cell, cools the hot atoms and extracts them into an atomic beam with a mean forward velocity of roughly 120 m/s. The laser light intersects the atomic beam perpendicularly, and fluorescence is collected on a UV-sensitive photomultiplier tube. The transverse velocity of the atoms is reduced to below about 0.5 m/s by a slit directly before they enter the fluorescence detector. The resulting Doppler broadenings are 1.5 MHz (326 nm) and 2.1 MHz (229 nm).

Due to the large natural linewidth of the 229 nm line, $\Gamma/(2\pi) = 100(3)\text{MHz}$ [62], a high-precision IS measurement is hampered by the overlap of the different isotopes in the spectrum. To improve the accuracy, we use enriched targets to precisely determine the lineshape. A Lorentzian function fits the data well, suggesting that Doppler broadening is indeed negligible. We then measure the IS of the (110, 112) pair, for which the overlap in the spectrum is most severe. This is done by taking multiple spectra while alternating two separate enriched ablation targets for ^{110}Cd and ^{112}Cd . This IS is then fixed

in a fit to the spectrum recorded using an ablation target with natural abundance. To resolve the frequencies of the fermionic isotopes in the spectrum, we vary their amplitude relative to the bosonic isotopes by changing the polarisation angle of the excitation laser with respect to the detector axis. The weighted means for each isotope are averaged and the final result is given in Table II and plotted in Fig. 3a. Our results are more accurate and differ considerably from those of a recent measurement [42]. This discrepancy manifests itself as a horizontal offset in the fits to the experimental data portrayed in Fig. 3a.

The 326 nm intercombination line of Cd has a 69 kHz natural linewidth that enables both cooling to low temperatures and precise spectroscopy. To measure the ISs of this line in an atomic beam, we use a frequency-doubled dye laser (Sirah Matisse 2DX with a Spectra Physics Wavetrain doubling module), whose linewidth is stabilised to better than 100 kHz short term and 1 MHz shot-to-shot stability. A typical spectrum is shown in Fig. 2, where the larger Doppler broadened linewidth of 5.6(2) MHz (Gaussian FWHM) arises from inclusion of atoms with higher velocity in the spectrum. The isotope shifts of five independent measurements have an average standard deviation of 0.8 MHz, consistent with the wavemeter (HighFinesse WS8-10) resolution of 0.4 MHz at the fundamental wavelength. This statistical uncertainty is small compared with the systematic uncertainty in the wavemeter measurement of the laser frequency. To place an upper bound on this systematic uncertainty, we measure nearby optical transitions of atomic copper in the same beam machine. The D_1 (327.5 nm) and D_2 (324.8 nm) lines lie roughly 1.3 nm to either side of the 326 nm line in Cd. We reproduce the precisely measured ground state hyperfine intervals




FIG. 3: King Plots of selected Cd transitions. The vertical axis is the modified isotope shift of the 214 nm line from [39], and the AR-RCCSDT theoretical prediction from Table I. The data for the horizontal axes includes the experimental measurements from this work in Table II, the theory prediction for 229 nm is from CI-MBPT [39], and for 326 and 332 nm from MCDHF [64]. Fits to previous experimental data are given for the 229 nm [42] and 326 nm lines [43, 44, 46, 47, 65]. The bands correspond to 68% confidence intervals.

of ^{65}Cu (12568.81(1) MHz) and ^{63}Cu (11733.83(1) MHz) [66] with a mean deviation of 3.1 ± 2.0 MHz for ^{63}Cu and 1.8 ± 3.7 MHz for ^{65}Cu . By measuring the wavemeter linearity with a temperature and pressure stabilized Zerodur cavity we find a systematic uncertainty of 3.3 MHz. We assign this as our systematic frequency uncertainty for the 326 nm and 229 nm spectra of Cd I.

B. Measurements of 326, 480 and 361 nm ISs with magneto-optic traps

We determine the ISs of the 326 line by measuring the sharp blue-edge of the fluorescence [67] from a 326 nm magneto-optic trap (MOT) [63]. The blue edge, where the cooling and trapping breaks down, is of order 20 kHz wide (90% to 10%) and serves as a precise reference for ISs. We directly load a 326 nm MOT of the abundant isotopes, ^{110}Cd to ^{116}Cd , from an effusive 1.2 cm diameter source of natural Cd that is 2.2 cm from the MOT. To capture the atoms, we use 60 to 150 mW of 326 nm light that is frequency modulated with a peak-to-peak amplitude of 8.6 MHz [22, 63]. The three retro-reflected MOT beams have a diameter of 8.5 mm. A lower-intensity cooling stage follows the MOT loading with no applied magnetic field gradient or frequency modulation, during which fluorescence is measured.

We produce 326 nm light with doubly-resonant sum frequency generation (SFG) of 542 nm and 820 nm light in a Beta Barium Borate crystal [21]. The 542 nm light is produced by frequency doubling a fiber-amplified 1083 nm extended cavity diode laser (ECDL). An 820 nm ECDL and a tapered amplifier supply the 820 nm light.

Isotope shifts are measured relative to various modes of a temperature-tuned, 4-mirror, rectangular reference cavity. Its free-spectral range is 905.03 MHz, its tangential and sagittal transverse mode splittings are 201.04 MHz and 140.84 MHz, and the cavity linewidth is 350 kHz. The 1083 nm ECDL is locked to the 00, 01 and 10 modes with an adjustable frequency offset provided by a double-passed acousto-optic modulator (AOM). The frequency of the 820 nm ECDL is measured relative to resonances of the same reference cavity [21], by slowly frequency modulating a double-passed AOM with a zero-to-peak amplitude of 1.2 MHz or less. Within 1 to 5 minutes after measuring the 326 nm frequency offset for each isotope, the absolute frequency of a 1083 nm cavity resonance is checked relative to a molecular iodine line via saturated absorption at 542 nm. We correct for the cavity drift between the measurements for each isotope, which is typically of order 1 MHz, and had a maximum drift of 4.2 MHz. The measurement uncertainty is given by the less than 1 MHz centering precision of the 820 nm light on a cavity resonance.

For ^{106}Cd and ^{108}Cd , with 1.25% and 0.89% natural abundances, we enhance the 326 nm MOT loading rate by capturing atoms on the 23 MHz wide 361 nm 3P_2 - 3D_3 transition, after optical pumping from 3P_1 and 3P_0 to 3P_2 via the 3S_1 state (see Fig. 1a). The 361 nm light is generated via doubly-resonant SFG of 1083 nm and 542 nm light, producing 70 to 200 mW [21, 63]. To load the 326 nm MOT, the 361 nm MOT is inhibited by turning off the 480 nm 3P_1 - 3S_1 optical pumping light and allowing the atoms to equilibrate in the 326 nm MOT for 57 ms before the start of the 326 nm cooling and detection phase. Our results for the ISs for the 326 nm line

are given in Table II and Fig. 3b. They are more accurate than our measurement of the same line with a beam, and comprise an order of magnitude of improvement compared with the weighted average of previous measurements. A disagreement of 4 combined errors is found for the (114,116) pair, previously measured with interferometry of fluorescence [43, 47]. This disagreement manifests as different slopes of the King Plot portrayed in Fig. 3b.

To measure ISs for the 480 nm transition, we use all four laser sources to enhance the MOT loading and then inhibit the 480 nm light for 28 ms to transfer atoms to the 326 nm MOT and cool them. We then turn on a 4.4 ms low intensity pulse of 480 nm light and observe the resulting 361 nm fluorescence. We generate the 468 nm and 480 nm optical pumping light using single-pass SFG of 1083 nm light and 823 nm and 862 nm light in fiber-coupled PPLN waveguides. The 468 nm and 480 nm light are combined on a beamsplitter and the two output beams illuminate the atoms on three nearly orthogonal axes before being retro-reflected. The 862 nm laser, and thereby the frequency of the 480 nm light, is slowly stabilized with a wavemeter and is monitored with the same reference cavity, using a number of transverse modes and a zero-to-peak frequency modulation of 1.2 MHz via a double-passed AOM. We determine the center frequency of the 17.5 MHz wide transition with a resolution of 0.8 MHz using a slow square wave frequency modulation of the 480 nm light by ± 4 MHz with a double-passed AOM. Our 480 nm ISs in Table II are the first measurements for this transition. By combining them with our 326 nm intercombination line ISs along with previous measurements for 468 nm [40], we determine the ISs for the 332 nm clock transition (see Fig. 3c), and similarly for the narrow 314 nm transition using measured 509 nm ISs [41].

Finally, we measure the ISs of the the 361 nm 3P_2 - 3D_3 transition. We again use the sharp blue-edge of the cooling as the reference for the ISs [67]. Here, we use the reference cavity stability to measure frequency offsets relative to 542 nm iodine saturated absorption resonances [68]. The blue edge is determined to better than 0.4 MHz and we estimate that the blue edges, especially for the slower loading ^{106}Cd and ^{108}Cd MOTs, may have systematic errors of order 3 MHz, much less than the 23 MHz 361 nm transition linewidth.

VI. BENCHMARKING IS CONSTANTS CALCULATIONS

The constants of eq. 8 for the D_1 and D_2 ionic transitions can be extracted from our calculation (Table I) with high accuracy because the uncertainties for each transition are highly correlated. We find $F_{D_2,D_1} = 1.0131(2)$ and $K_{D_2,D_1} = -114(2)$ GHz u. A Monte-Carlo fit to the available experimental data [41, 47, 70–72] yields $F_{D_2,D_1} = 1.02(6)$ and $K_{D_2,D_1} = -43(229)$ GHz u, which

TABLE III: IS constants of atomic transitions deduced from our calculations (Table I) projected using King Plots with experimental data from Table II and the literature [39, 41–44, 47].

i (nm)	F_i (MHz fm $^{-2}$)	K_i (GHz u)	Method
229	-3764(134)	1199(104)	
	-4024(200)	1428(220)	CI-MBPT [39]
332	-4391(86)	1712(43)	
	-4660(160)	1629(19)	MCDHF [64]
326	-4354(62)	1673(43)	
	-4680(160)	1616(17)	MCDHF [64]
	-4559(230)	1865(400)	CI-MBPT [39]
	-4420(340)	1717(330)	CKP [69]
314	-4421(86)	1739(61)	
468	1150(49)	-1(47)	
480	1122(40)	30(41)	
509	1180(19)	-32(24)	
	1228(60)	-63(400)	CI-MBPT [39]
361	-654(42)	-237(59)	

agrees with our calculation, within the large experimental uncertainty. To better test our calculation, recent ion-trap IS measurements for the D_2 transition [39] could be extended to the D_1 transition.

To test calculation in Cd I, we transform our constants for the Cd II D_2 transition to transitions in Cd I using eq. 8, fitting experimental ISs from this work and the literature [39, 41–44, 47]. Here, we do not consider fermionic isotopes, for which eq. 8 may not be a good approximation at the few MHz level [64]. The resulting IS constants are given in Table III. Their uncertainty is dominated by that of the measured isotope shifts of the atomic lines. The IS constants of tables I and III agree with, and are up to an order of magnitude more precise than, the CI-MBPT calculations [39] for both F and K for all tested ionic and atomic transitions. We thus validate both the central values and the uncertainty estimation of CI-MBPT, which is applicable for calculations of both linear and non-linear IS constants in a variety of atomic systems.

For the intercombination line, our measurements produce a significantly larger KP slope than prior results, increasing from $F_{214,326} = 0.98(8)$ to $F_{214,326} = 1.397(17)$, corresponding to the green solid and red dashed lines in figure 8b. Combining the slope with $F_{214} = -6082(45)$ MHz fm $^{-2}$ from Table Ie shifts F_{326} from $-6180(478)$ MHz fm $^{-2}$ to $-4354(62)$ MHz fm $^{-2}$, closer to the value obtained for the 229 nm transition which is expected as they share an s-state. Our F_{326} and K_{326} constants agree with and are more accurate than CI-MBPT calculations [39] and an empirical CKP determination [69]. However, Fig. 3b shows a slight disagreement with a recent calculation via MCDHF [64]. As summarized in Table Ie, the deviation for F_{326} is two combined standard errors and for K_{326} by one. A similar deviation is apparent for the clock transition at 332 nm (Fig. 3c). The changes in F_{326} and K_{326} and the updated intercombina-

TABLE IV: Updated differences of charge radii of even-even Cd isotopes in fm². This work and [39] use measurements for the D2 line from [39, 41], while the radii extracted using atomic factors from the MCDHF method [64] utilize measurements in the intercombination line summarized in [69]. The charge radius is obtained by adding $(r_c)^{114} = 4.614(3)$ fm in quadrature.

A	$(\delta r^2)^{A,114}$	CKP [39]	MCDHF [64]	$(\delta r^2)^{A+2,A}$	Ref [69]
100	-1.466(12)	-1.409(19)		0.2876(23)	
102	-1.178(9)	-1.135(14)		0.2480(20)	0.28(12)
104	-0.930(8)	-0.897(10)		0.2149(18)	0.235(95)
106	-0.714(6)	-0.690(8)	-0.662(24)	0.1899(15)	0.182(16)
108	-0.524(4)	-0.506(6)	-0.478(19)	0.1817(15)	0.183(15)
110	-0.343(3)	-0.331(4)	-0.310(12)	0.1791(14)	0.174(13)
112	-0.164(1)	-0.158(2)	-0.145(6)	0.1635(13)	0.157(10)
114	0	0	0	0.1360(11)	0.129(10)
116	0.136(1)	0.134(2)	0.115(5)	0.1108(27)	0.083(22)
118	0.248(2)	0.244(6)		0.1003(25)	0.084(50)
120	0.349(3)	0.344(8)		0.0892(19)	
122	0.439(4)	0.434(11)		0.0825(31)	
124	0.520(5)	0.514(14)		0.0760(9)	
126	0.596(5)	0.590(18)		0.0755(9)	
128	0.671(6)	0.666(21)		0.0443(8)	
130	0.716(7)	0.713(24)			

tion line ISs account for the 20 – 50% differences between the ISs calculated in [64] and those summarized in Ref. [69]. Whereas the range of values of F_i is a few percent, we note that the many-body QED corrections are estimated to contribute 3%. Further improvements in field shift calculations should include these QED effects, especially for heavier elements.

VII. IMPROVED CHARGE RADII

High-precision optical isotope and isomer shifts for the Cadmium chain were measured at ISOLDE using collinear laser spectroscopy [41, 73]. To extract charge radii differences from spectroscopic data, eq. 1 is often calibrated with the radii of stable isotopes extracted from a combination of muonic x-ray energies and electron scattering data [69]. This commonly used CKP method has several disadvantages: it is currently applicable only to elements possessing at least three stable or long-lived isotopes; the determined IS constants are limited by the knowledge of nuclear polarization corrections to muonic levels, as well as their unknown correlations; the calibration coefficients from electron scattering experiments are in many cases unknown (e.g. for ^{106,108,111,113}Cd) or nuclear-model-dependent (e.g. for ^{110,112,114,116}Cd); and the resulting IS constants K and F are highly correlated, leading to diverging uncertainties of extracted radii outside of the calibrated region.

The advent of high-accuracy many-body calculations of IS constants allows differences of charge radii to be extracted directly from optical measurements with no input, beyond small higher-moment corrections, from muonic x-ray or electron scattering experiments. In Table IV and Fig. 4 we compare $(\delta r^2)^{A,114} = r_A^2 - r_{114}^2$

extracted from experimental data [39, 41] and our calculated F_{214} and K_{214} to those determined recently using a CKP method [39]. The radii of odd isotopes and isomers [73] could also be extracted after calculating off-diagonal hyperfine elements. For ^{118,120,122}Cd, measured only using an atomic transition at 509 nm, we use F_{509} and K_{509} given in Table III. Our reported uncertainty is 0.7 – 0.9%, dominated by the calculated F_{214} uncertainty, in turn stemming from a systematic uncertainty in the QED correction. For all $(\delta r^2)^{A,114} = r_A^2 - r_{114}^2$, it is smaller than the uncertainty from the CKP method, especially for neutron-rich isotopes, for which the uncertainty can be as much as a factor of 4 smaller. A slight disagreement is found for mass numbers 112 and lower as seen in Fig. 4a, which we ascribe to the limitations the CKP method listed above. Our improved charge radii increase the disagreement with state-of-the-art nuclear theory calculations [39, 41] for the most neutron rich isotopes (Fig. 4b), while improving the agreement for those that are proton-rich (Fig. 4a).

The benefit of extracting radii directly from calculated atomic constants is even more pronounced for the ladder differences $(\delta r^2)^{A+2,A} = r_{A+2}^2 - r_A^2$, which are highly sensitive to the uncertainties of muonic x-ray measurements. Such differences are useful, e.g. for identifying quantum phase transitions in nuclear shapes [74, 75]. Our results in Table IV are up to 60 times more accurate than those obtained with a CKP method [69].

VIII. TOWARDS BSM PHYSICS

We can use the above results to assess the requirements for King Plots in Cd I ISs to be sensitive to new physics. An interaction between electrons and neutrons




FIG. 4: Squared RMS charge radii of even-even Cd isotopes relative to ^{114}Cd . a: proton rich, b: neutron rich. The data points correspond to columns 2 – 4 of Table IV. The nuclear theory lines are from the Fayans functional (Fy), extended by information in the radii of Ca [41], and Skyrme-Hartree-Fock-Bogoliubov mass formula (HFB-24) [39].

that is mediated by a heavy boson has a short range, less than a Bohr radius. To leading order, it is a contact interaction, proportional to the wave function overlap with the nucleus, and therefore proportional to the FS. Such a contribution is absorbed into charge radii differences, so it does not create deviations of linearity in KPs. In the next order, and in a hydrogenic approximation, the deviation from linearity turns out to be proportional to the FS [8]. Therefore, a KP for transitions i and j where $|F_i - F_j|$ is large, can have a high sensitivity to such new physics. In Fig. 1b we plot $|F_i - F_j|$ for important transition pairs, whose atomic constants are given in Table III. As expected, large differences are between pairs that do not share the same S state. However, narrow transitions couple the ground S state with the $5p^3P_j$ states, enabling exceptional precisions for their measurements. Focusing on the clock and intercombination line pair, a KP returns $F_{332,326} = 1.008(13)$, with an uncertainty largely from the 468 nm measurement [40] and our 480 nm measurement. Multiplying by $F_{326} = -4354(62) \text{ MHz fm}^{-2}$ from Table III, we find $|F_{332} - F_{326}| \leq 100 \text{ MHz fm}^{-2}$. For the $S - D$ transitions used in the KP of Ca II, $|F_{732} - F_{729}| = 0.55(2) \text{ MHz fm}^{-2}$ [13], which is up to two orders of magnitude smaller. We therefore surmise that when searching for new interactions mediated by a heavy boson, the sensitivity of a KP between the ISs of the clock and intercombination lines in Cd could be as much as two orders more sensitive than the current limit set by IS measurements in Ca II. As the accuracy of measurements in the latter are of order 20 Hz, accuracies of at least few kHz would be required for Cd I. Considering that the natural linewidth of the intercombination line is 69 kHz, such an accuracy should be straightforward with trapped samples.

It is difficult to attribute an observed KP non-linearity

to new physics without estimating non-linear SM contributions, such as the quadratic mass shift (QMS). Due to the similarity between nuclear masses and mass numbers, the deviation pattern of the QMS is nearly indistinguishable from the pattern induced by a new boson that couples electrons and neutrons, and so it sets a stringent limit on the new physics reach. The order of magnitude of the QMS can be estimated with a hydrogenic approximation to be $K_{ij}^{\text{QMS}} = K_{ij}(m/M_A + m/M_{A'})$ [76], with $|K_{332,326}| \leq 60 \text{ GHz u}$ from our KP. A linear fit of $K_{332,326}^{\text{QMS}}$ versus $\delta\bar{V}_{326}^{A,A+2}$ returns an offset of 0.7 MHz u, a slope of 6×10^{-10} , and, most importantly, isotope-dependent non-linearities. The largest deviation from the linear fit is 4 Hz, for the $^{106,108}\text{Cd}$ pair and the maximum value of $K_{332,326}$. It is comparable to the estimated 3 Hz QMS-induced non-linearity in Ca II [77].

For the medium mass Cd system, with its small nuclear deformations, the largest expected SM non-linearity is the quadratic field shift (QFS) [15]. In the notation of Eq. 8, it takes the form $F_{ij}^{\text{QFS}} = G_{ij}(\delta(r^2)^{AA'})^2/\mu_{AA'}$, where $G_{ij} = G_i - F_{ij}G_j$. G_{ij} sets the scale of the non-linearity and is a transition-dependent and nucleus-independent constant [16, 17]. This scale is difficult to estimate and different calculations yield a wide range of sensitivities [16]. Assuming that the QFS is approximately quadratic in $|F_i - F_j|$, the absence of observed KP non-linearities in this work already indicates that the non-linearity induced by the QFS of the narrow transition pairs is less than 1 kHz. Moreover, the QFS may be identified from the deviation pattern from linearity [17]. The precisions with which $\delta(r^2)$ are known set the limits for this identification, thus emphasizing the need for the improved radii in Table IV.

Other sources of non-linearity, such as nuclear polarization and nuclear deformation, could be non-negligible

at the anticipated level of precision [15, 78, 79]. Calculating their magnitude and pattern is difficult due to our limited understanding of nuclear structure. However, like the QFS, their magnitude increases with the wave function overlap with the nucleus. By combining the ISs of three or more transitions with different FSs, these non-linearities can be isolated [10, 80]. Based on the above considerations, the ISs of the clock and intercombination lines can be measured to the order of ~ 100 Hz, limited by the natural linewidth of the intercombination line. If no non-linearity is observed, the ultra-narrow 314 nm line may provide higher precision with reasonably controlled tensor light shift. If a non-linearity is present at this level, the IS of a transition that does not include the ground state can lead to a sensitive multi-dimensional KP. As suggested by Fig. 1b, the $|F_i - F_j|$ for these transitions are at least two orders of magnitude larger than for ground state transitions. Narrow transitions between excited states, such as from 3P_0 to long-lived low-lying Rydberg states, can thus increase the sensitivity of searches for BSM physics.

IX. SUMMARY

The measurements reported here significantly improve the isotope shifts of several low-lying transitions of Cd I, including the first measurements of two transitions. For the wide 229 nm and the narrow 326 nm transitions, we utilize a cryogenic beam with enriched samples, combined with a detection method that distinguishes between the emission patterns of bosonic and fermionic isotopes. We also measured ISs of the 326 and 361 nm transitions using the sharp blue edge of laser-cooling in a magneto-optical trap, and of the 480 nm transition via optical pumping of the trapped atoms. Combining our 480 and 326 nm ISs with previous ISs for the 486 nm [40] and 509 nm [41] transitions, we predict the ISs of the two ultra-narrow transitions in Cd, which have not yet been measured. Our results are significantly more accurate, often in disagreement with previous measurements, and highlight the benefits of isotope-selective and cold sources for IS measurements.

We also present high-precision calculations of IS constants of Cd II. By projecting our calculations from Cd II to Cd I, using King Plots with measured ISs, we find the IS constants for all low-lying transitions in Cd I, including those of the narrow intercombination and ultra-narrow lines. Our resulting IS constants largely agree with, and are more accurate than those obtained from the CI-MBPT [39] and MCDHF methods [64], setting stringent benchmarks for improving the theory. Some disagreements are observed with recent calculations [64], potentially from underestimating the uncertainty of high-order electron correlations as well as the importance of the QED contributions to the field shift constants.

By combining our calculated IS constants with measurements of a long chain of short-lived ions, we extract accurate charge radii differences, without the limitations from muonic atom measurements. To our knowledge, this is the first extraction of charge radii differences for a chain of isotopes with an accuracy better than 1%, which opens opportunities to improve our knowledge of nuclear sizes far from stability. Beyond benchmarking nuclear models, these are important to identify the patterns of non-linearities in future high-precision measurements [23]. This work sets the stage for new physics searches using generalized KPs in the Cd system. Our results suggest that precise future Cd IS measurements can improve, by as much as two orders of magnitude, the current best bounds on new electron-neutron interactions, obtained from a KP for Ca II transitions.

ACKNOWLEDGEMENT

We thank Ronnie Kosloff for stimulating conversations, Eberhard Tiemann for helpful assistance, and Yotam Soreq for valuable suggestions. We gratefully acknowledge the use of the Vikram-100 HPC cluster of Physical Research Laboratory, Ahmedabad (B.K.S.) and financial support from the US National Science Foundation (K.G.) and the European Research Council (ERC) under the European Union’s Horizon 2020 Research and Innovation Programme (CoMoFun, S. T.).

-
- [1] BACON collaboration, Frequency ratio measurements at 18-digit accuracy using an optical clock network, *Nature* **591**, 564 (2021).
 - [2] A. Derevianko and M. Pospelov, Hunting for topological dark matter with atomic clocks, *Nature Physics* **10**, 933 (2014).
 - [3] M. S. Safronova, D. Budker, D. DeMille, D. F. J. Kimball, A. Derevianko, and C. W. Clark, Search for new physics with atoms and molecules, *Rev. Mod. Phys.* **90**, 025008 (2018).
 - [4] M. Takamoto, I. Ushijima, N. Ohmae, T. Yahagi, K. Kokado, H. Shinkai, and H. Katori, Test of general relativity by a pair of transportable optical lattice clocks, *Nature Photonics* **14**, 411 (2020).
 - [5] R. Lange, N. Huntemann, J. M. Rahm, C. Sanner, H. Shao, B. Lipphardt, C. Tamm, S. Weyers, and E. Peik, Improved limits for violations of local position invariance from atomic clock comparisons, *Phys. Rev. Lett.* **126**, 011102 (2021).
 - [6] W. H. King, Comments on the article “peculiarities of the isotope shift in the samarium spectrum”, *J. Opt. Soc. Am.* **53**, 638 (1963).
 - [7] C. Delaunay, R. Ozeri, G. Perez, and Y. Soreq, Probing atomic Higgs-like forces at the precision frontier, *Phys.*

- Rev. D* **96**, 093001 (2017).
- [8] J. C. Berengut, D. Budker, C. Delaunay, V. V. Flambaum, C. Fruguele, E. Fuchs, C. Grojean, R. Harnik, R. Ozeri, G. Perez, and Y. Soreq, Probing new long-range interactions by isotope shift spectroscopy, *Phys. Rev. Lett.* **120**, 091801 (2018).
- [9] Y. V. Stadnik, Probing long-range neutrino-mediated forces with atomic and nuclear spectroscopy, *Phys. Rev. Lett.* **120**, 223202 (2018).
- [10] J. C. Berengut, C. Delaunay, A. Geddes, and Y. Soreq, Generalized King linearity and new physics searches with isotope shifts, *Phys. Rev. Research* **2**, 043444 (2020).
- [11] T. Manovitz, R. Shaniv, Y. Shapira, R. Ozeri, and N. Akerman, Precision measurement of atomic isotope shifts using a two-isotope entangled state, *Phys. Rev. Lett.* **123**, 203001 (2019).
- [12] X. Zheng, J. Dolde, V. Lochab, B. N. Merriman, H. Li, and S. Kolkowitz, Differential clock comparisons with a multiplexed optical lattice clock, *Nature* **602**, 425 (2022).
- [13] C. Solaro, S. Meyer, K. Fisher, J. C. Berengut, E. Fuchs, and M. Drewsen, Improved isotope-shift-based bounds on bosons beyond the standard model through measurements of the ${}^2D_{3/2} - {}^2D_{5/2}$ interval in Ca^+ , *Phys. Rev. Lett.* **125**, 123003 (2020).
- [14] G. W. F. Drake, H. S. Dhindsa, and V. J. Marton, King and second-King plots with optimized sensitivity for lithium ions, *Phys. Rev. A* **104**, L060801 (2021).
- [15] P. Munro-Laylim, V. A. Dzuba, and V. V. Flambaum, Nuclear polarisation and relativistic effects contributions to King plot non-linearity (2021), [arXiv:2112.14313 \[physics.atom-ph\]](https://arxiv.org/abs/2112.14313).
- [16] I. Counts, J. Hur, D. P. L. Aude Craik, H. Jeon, C. Leung, J. C. Berengut, A. Geddes, A. Kawasaki, W. Jhe, and V. Vuletić, Evidence for nonlinear isotope shift in Yb^+ search for new boson, *Phys. Rev. Lett.* **125**, 123002 (2020).
- [17] K. Ono, Y. Saito, T. Ishiyama, T. Higomoto, T. Takano, Y. Takasu, Y. Yamamoto, M. Tanaka, and Y. Takahashi, Observation of nonlinearity of generalized King plot in the search for new boson, *Phys. Rev. X* **12**, 021033 (2022).
- [18] N. L. Figueroa, J. C. Berengut, V. A. Dzuba, V. V. Flambaum, D. Budker, and D. Antypas, Precision determination of isotope shifts in ytterbium and implications for new physics, *Phys. Rev. Lett.* **128**, 073001 (2022).
- [19] J. Hur, D. P. L. Aude Craik, I. Counts, E. Knyazev, L. Caldwell, C. Leung, S. Pandey, J. C. Berengut, A. Geddes, W. Nazarewicz, P.-G. Reinhard, A. Kawasaki, H. Jeon, W. Jhe, and V. Vuletić, Evidence of two-source King plot nonlinearity in spectroscopic search for new boson, *Phys. Rev. Lett.* **128**, 163201 (2022).
- [20] S. Ebata and T. Nakatsukasa, Octupole deformation in the nuclear chart based on the 3D Skyrme Hartree-Fock plus BCS model, *Physica Scripta* **92**, 064005 (2017).
- [21] D. Schussheim and K. Gibble, Laser system to laser-cool and trap cadmium: towards a cadmium optical lattice clock, in *Frontiers in Optics / Laser Science* (Optical Society of America, 2018) p. LTh1F.2.
- [22] A. Yamaguchi, M. S. Safronova, K. Gibble, and H. Katori, Narrow-line cooling and determination of the magic wavelength of Cd, *Phys. Rev. Lett.* **123**, 113201 (2019).
- [23] R. A. Müller, V. A. Yerokhin, A. N. Artemyev, and A. Surzhykov, Nonlinearities of King's plot and their dependence on nuclear radii, *Phys. Rev. A* **104**, L020802 (2021).
- [24] W. H. King, *Isotope shifts in atomic spectra* (Springer Science & Business Media, 2013).
- [25] S. A. Blundell, P. E. G. Baird, C. W. P. Palmer, D. N. Stacey, and G. K. Woodgate, A reformulation of the theory of field isotope shift in atoms, *Journal of Physics B: Atomic and Molecular Physics* **20**, 3663 (1987).
- [26] W. A. Gillespie, S. W. Brain, A. Johnston, E. W. Lees, R. P. Singhal, A. G. Slight, M. W. S. M. Brimicombe, D. N. Stacey, V. Stacey, and H. Huhnermann, Measurements of the nuclear charge distribution in the cadmium isotopes, *Journal of Physics G: Nuclear Physics* **1**, L6 (1975).
- [27] J. W. Lightbody, S. Penner, S. P. Fivozinsky, P. L. Hollowell, and H. Crannell, Electron scattering from vibrational nuclei, *Phys. Rev. C* **14**, 952 (1976).
- [28] C. W. P. Palmer, Reformulation of the theory of the mass shift, *Journal of Physics B: Atomic and Molecular Physics* **20**, 5987 (1987).
- [29] V. M. Shabaev, Qed theory of the nuclear recoil effect in atoms, *Phys. Rev. A* **57**, 59 (1998).
- [30] B. K. Sahoo and B. Ohayon, Benchmarking many-body approaches for the determination of isotope-shift constants: Application to the Li , Be^+ , and Ar^{15+} isoelectronic systems, *Phys. Rev. A* **103**, 052802 (2021).
- [31] G. Estévez and L. B. Bhuiyan, Electrostatic potential due to a Fermi-type charge distribution, *American Journal of Physics* **53**, 450 (1985).
- [32] I. Lindgren and J. Morrison, *Atomic many-body theory*, Vol. 13 (1982).
- [33] D. Mukhopadhyay and D. Mukherjee, Size-extensive effective hamiltonian formalisms using quasi-Hilbert and quasi-Fock space strategies with incomplete model spaces, *Chemical Physics Letters* **163**, 171 (1989).
- [34] B. K. Sahoo, S. Majumder, R. K. Chaudhuri, B. P. Das, and D. Mukherjee, Ab initio determination of the lifetime of the $6^2P_{3/2}$ state for ${}^{207}\text{Pb}^+$ by relativistic many-body theory, *Journal of Physics B: Atomic, Molecular and Optical Physics* **37**, 3409 (2004).
- [35] B. K. Sahoo, A. R. Vernon, R. F. G. Ruiz, C. L. Binnersley, J. Billowes, M. L. Bissell, T. E. Cocolios, G. J. Farooq-Smith, K. T. Flanagan, W. Gins, R. P. de Groote, Á. Koszorús, G. Neyens, K. M. Lynch, F. Parnefjord-Gustafsson, C. M. Ricketts, K. D. A. Wendt, S. G. Wilkins, and X. F. Yang, Analytic response relativistic coupled-cluster theory: the first application to indium isotope shifts, *New Journal of Physics* **22**, 012001 (2020).
- [36] S. F. Boys and J. E. Lennard-Jones, Electronic wave functions III. some theorems on integrals of antisymmetric functions of equivalent orbital form, *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences* **206**, 489 (1951).
- [37] A. G. Shenstone and J. T. Pittenger, Cadmium spectra, *J. Opt. Soc. Am.* **39**, 219 (1949).
- [38] K. Burns and K. B. Adams, Energy levels and wavelengths of natural cadmium and of cadmium-114, *J. Opt. Soc. Am.* **46**, 94 (1956).
- [39] J. Z. Han, C. Pan, K. Y. Zhang, X. F. Yang, S. Q. Zhang, J. C. Berengut, S. Goriely, H. Wang, Y. M. Yu, J. Meng, J. W. Zhang, and L. J. Wang, Isotope shift factors for the $\text{Cd}^+ 5s^2S_{1/2} \rightarrow 5p^2P_{3/2}$ transition and determination of Cd nuclear charge radii, *Phys. Rev. Research* **4**, 033049 (2022).

- [40] R. Wenz, A. Timmermann, and E. Matthias, Subshell effect in mean square charge radii of stable even cadmium isotopes, *Zeitschrift für Physik A Atoms and Nuclei* **303**, 87 (1981).
- [41] M. Hammen, W. Nörtershäuser, D. L. Balabanski, M. L. Bissell, K. Blaum, I. Budinčević, B. Cheal, K. T. Flanagan, N. Frömmgen, G. Georgiev, C. Geppert, M. Kowalska, K. Kreim, A. Krieger, W. Nazarewicz, R. Neugart, G. Neyens, J. Papuga, P.-G. Reinhard, M. M. Rajabali, S. Schmidt, and D. T. Yordanov, From calcium to cadmium: Testing the pairing functional through charge radii measurements of $^{100-130}\text{Cd}$, *Phys. Rev. Lett.* **121**, 102501 (2018).
- [42] J. N. Tinsley, S. Bandarupally, J.-P. Penttinen, S. Manzoor, S. Ranta, L. Salvi, M. Guina, and N. Poli, Watt-level blue light for precision spectroscopy, laser cooling and trapping of strontium and cadmium atoms, *Opt. Express* **29**, 25462 (2021).
- [43] F. M. Kelly and E. Tomchuk, Isotope shift in the CdI intercombination resonance line λ 3261 Å, *Proceedings of the Physical Society* **74**, 689 (1959).
- [44] R. Kloch, P. Baird, M. Boshier, M. Macpherson, C. Palmer, D. Stacey, and V. Stacey, Isotope shifts in λ 326.1 nm of CdI, *Zeitschrift für Physik D Atoms, Molecules and Clusters* **6**, 315 (1987).
- [45] F. M. Kelly and E. Tomchuk, Isotope shift in the CdI resonance line 2288 nm, *Proceedings of the Physical Society* **78**, 1304 (1961).
- [46] P. Brix and A. Steudel, Die Isotopieverschiebung in der Cd I-Resonanzlinie λ 3261Å, *Zeitschrift für Physik* **128**, 260 (1950).
- [47] F. Leś, *Acta Physica Polonica* **24**, 951 (1964).
- [48] B. Ohayon, R. F. G. Ruiz, Z. H. Sun, G. Hagen, T. Papenbrock, and B. K. Sahoo, Nuclear charge radii of Na isotopes: Interplay of atomic and nuclear theory, *Phys. Rev. C* **105**, L031305 (2022).
- [49] B. Finkelstein, Über den virialsatz in der wellenmechanik, *Zeitschrift für Physik* **50**, 293 (1928).
- [50] J. Li, C. Nazé, M. Godefroid, G. Gaigalas, and P. Jönsson, On the breakdown of the Dirac kinetic energy operator for estimating normal mass shifts, *The European Physical Journal D* **66**, 1 (2012).
- [51] E. Lindroth and A.-M. Mårtensson-Pendrill, Calculation of the isotope shift in Na, *Zeitschrift für Physik A Atoms and Nuclei* **309**, 277 (1983).
- [52] H. Heylen, C. Babcock, R. Beerwerth, J. Billowes, M. L. Bissell, K. Blaum, J. Bonnard, P. Campbell, B. Cheal, T. Day Goodacre, D. Fedorov, S. Fritzsche, R. F. Garcia Ruiz, W. Geithner, C. Geppert, W. Gins, L. K. Grob, M. Kowalska, K. Kreim, S. M. Lenzi, I. D. Moore, B. Maass, S. Malbrunot-Ettenauer, B. Marsh, R. Neugart, G. Neyens, W. Nörtershäuser, T. Otsuka, J. Papuga, R. Rossel, S. Rothe, R. Sánchez, Y. Tsunoda, C. Wraith, L. Xie, X. F. Yang, and D. T. Yordanov, Changes in nuclear structure along the Mn isotopic chain studied via charge radii, *Phys. Rev. C* **94**, 054321 (2016).
- [53] H. Heylen, C. S. Devlin, W. Gins, M. L. Bissell, K. Blaum, B. Cheal, L. Filippin, R. F. G. Ruiz, M. Godefroid, C. Gorges, J. D. Holt, A. Kanellakopoulos, S. Kaufmann, A. Koszorús, K. König, S. Malbrunot-Ettenauer, T. Miyagi, R. Neugart, G. Neyens, W. Nörtershäuser, R. Sánchez, F. Sommer, L. V. Rodríguez, L. Xie, Z. Y. Xu, X. F. Yang, and D. T. Yordanov, High-resolution laser spectroscopy of $^{27-32}\text{Al}$, *Phys. Rev. C* **103**, 014318 (2021).
- [54] Á. Koszorús, X. Yang, W. Jiang, S. Novario, S. Bai, J. Billowes, C. Binnersley, M. Bissell, T. E. Cocolios, B. Cooper, et al., Charge radii of exotic potassium isotopes challenge nuclear theory and the magic character of $N=32$, *Nature Physics* **17**, 439 (2021).
- [55] V. A. Dzuba, W. R. Johnson, and M. S. Safronova, Calculation of isotope shifts for cesium and francium, *Phys. Rev. A* **72**, 022503 (2005).
- [56] C. Shi, F. Gebert, C. Gorges, S. Kaufmann, W. Nörtershäuser, B. Sahoo, A. Surzhykov, V. Yerokhin, J. Berengut, F. Wolf, et al., Unexpectedly large difference of the electron density at the nucleus in the $4p$ $^2P_{1/2,3/2}$ fine-structure doublet of Ca^+ , in *Exploring the World with the Laser* (Springer, 2018) pp. 1–19.
- [57] M. S. Safronova and W. R. Johnson, Third-order isotope-shift constants for alkali-metal atoms and ions, *Phys. Rev. A* **64**, 052501 (2001).
- [58] I. I. Tupitsyn, V. M. Shabaev, J. R. Crespo López-Urrutia, I. Draganić, R. Soria Orts, and J. Ullrich, Relativistic calculations of isotope shifts in highly charged ions, *Phys. Rev. A* **68**, 022511 (2003).
- [59] J. C. Berengut, V. A. Dzuba, and V. V. Flambaum, Isotope-shift calculations for atoms with one valence electron, *Phys. Rev. A* **68**, 022502 (2003).
- [60] A. Dorne, B. K. Sahoo, and A. Kastberg, Relativistic coupled-cluster calculations of isotope shifts for the low-lying states of Ca II in the finite-field approach, *Atoms* **9**, 10.3390/atoms9020026 (2021).
- [61] J. S. Schelfhout and J. J. McFerran, Isotope shifts for $^1S_0 - ^3P_{0,1}$ Yb lines from multiconfiguration Dirac-Hartree-Fock calculations, *Phys. Rev. A* **104**, 022806 (2021).
- [62] S. Hofsäss, E. Padilla, S. C. Wright, S. Kray, R. Thomas, B. Sartakov, G. Meijer, and S. Truppe, In preparation, (2022).
- [63] K. Gibble, In preparation (2022).
- [64] J. S. Schelfhout and J. J. McFerran, Multiconfiguration Dirac-Hartree-Fock calculations for Hg and Cd with estimates for unknown clock-transition frequencies, *Phys. Rev. A* **105**, 022805 (2022).
- [65] F. Buchinger, P. Dabkiewicz, H.-J. Kluge, A. Mueller, and E.-W. Otten, The n -dependence of Cd mean square charge radii ($54 \leq N \leq 72$) and the nuclear moments of ^{103}Cd , *Nuclear Physics A* **462**, 305 (1987).
- [66] Y. Ting and H. Lew, Hyperfine structure of Cu^{63} and Cu^{65} , *Phys. Rev.* **105**, 581 (1957).
- [67] M. Walhout, H. J. L. Megens, A. Witte, and S. L. Rolston, Magneto-optical trapping of metastable xenon: Isotope-shift measurements, *Phys. Rev. A* **48**, R879 (1993).
- [68] B. Bodermann, H. Knöckel, and E. Tiemann, Widely usable interpolation formulae for hyperfine splittings in the $^{127}\text{I}_2$ spectrum, *The European Physical Journal D-Atomic, Molecular, Optical and Plasma Physics* **19**, 31 (2002).
- [69] G. Fricke and K. Heilig, *Nuclear charge radii · 48-Cd cadmium: Datasheet from Landolt-Börnstein - group I elementary particles, nuclei and atoms · volume 20*, springer-Verlag Berlin Heidelberg.
- [70] R. Contreras and F. Kelly, Isotope shift in the second spectrum of cadmium, *Canadian Journal of Physics* **47**, 1979 (1969).

- [71] D. C. Bishop and W. H. King, Isotope shifts in the resonance lines of Cd II and Sn IV and the size of specific mass shifts in alkali like resonance lines, *Journal of Physics B: Atomic and Molecular Physics* **4**, 1798 (1971).
- [72] M. Brimicombe, D. Stacey, V. Stacey, H. Hühnermann, and N. Menzel, Optical isotope shifts and hyperfine structure in Cd, *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences* **352**, 141 (1976).
- [73] D. T. Yordanov, D. L. Balabanski, M. L. Bissell, K. Blaum, I. Budinčević, B. Cheal, K. Flanagan, N. Frömmgen, G. Georgiev, C. Geppert, M. Hammen, M. Kowalska, K. Kreim, A. Krieger, J. Meng, R. Neugart, G. Neyens, W. Nörtershäuser, M. M. Rajabali, J. Papuga, S. Schmidt, and P. W. Zhao, Simple nuclear structure in $^{111-129}\text{Cd}$ from atomic isomer shifts, *Phys. Rev. Lett.* **116**, 032501 (2016).
- [74] S. Zerguine, P. Van Isacker, and A. Bouldjedri, Consistent description of nuclear charge radii and electric monopole transitions, *Phys. Rev. C* **85**, 034331 (2012).
- [75] N. Gavrielov, A. Leviatan, and F. Iachello, Zr isotopes as a region of intertwined quantum phase transitions, *Phys. Rev. C* **105**, 014305 (2022).
- [76] V. A. Yerokhin, R. A. Müller, A. Surzhykov, P. Micke, and P. O. Schmidt, Nonlinear isotope-shift effects in b-like, b-like, and c-like argon, *Phys. Rev. A* **101**, 012502 (2020).
- [77] V. V. Flambaum, A. J. Geddes, and A. V. Viatkina, Isotope shift, nonlinearity of King plots, and the search for new particles, *Phys. Rev. A* **97**, 032510 (2018).
- [78] V. V. Flambaum, I. B. Samsonov, H. B. T. Tan, and A. V. Viatkina, Nuclear polarization effects in atoms and ions, *Phys. Rev. A* **103**, 032811 (2021).
- [79] S. O. Allehabi, V. A. Dzuba, V. V. Flambaum, and A. V. Afanasjev, Nuclear deformation as a source of the nonlinearity of the King plot in the yb^+ ion, *Phys. Rev. A* **103**, L030801 (2021).
- [80] C. Frugiuele, E. Fuchs, G. Perez, and M. Schlaffer, Constraining new physics models with isotope shift spectroscopy, *Phys. Rev. D* **96**, 015011 (2017).