On Nonparametric Estimation in Online Problems

Man Fung Leung

Department of Statistics, University of Illinois at Urbana-Champaign

and

Kin Wai Chan *

Department of Statistics, The Chinese University of Hong Kong

Abstract

Offline estimators are often inadequate for real-time applications. Nevertheless, many online estimators are found by sacrificing some statistical efficiency. This paper presents a general framework to understand and construct efficient nonparametric estimators for online problems. Statistically, we choose long-run variance as an exemplary estimand and derive the first set of sufficient conditions for O(1)-time or O(1)-space update, which allows methodological generation of estimators. Our asymptotic theory shows that the generated estimators dominate existing alternatives. Computationally, we introduce mini-batch estimation to accelerate online estimators for real-time applications. Implementation issues such as automatic optimal parameters selection are discussed. Practically, we demonstrate how to use our framework with recent development in change point detection, causal inference, and stochastic approximation. We also illustrate the strength of our estimators in some classical problems such as Markov chain Monte Carlo convergence diagnosis and confidence interval construction.

Keywords: Batch-means estimator; Kernel estimator; Long-run variance; Nonparametric estimation; Time series

^{*}Correspondence email: kinwaichan@sta.cuhk.edu.hk

1 Introduction

1.1 Motivation

Online problems arise naturally in many fields of statistics. On top of them, modern computing allows intractable offline problems to be approached with online techniques. To name a few, multidimensional integrals can be approximated by Markov chain Monte Carlo (MCMC) methods (Hastings, 1970); deep neural networks can be trained using stochastic approximation (Robbins and Monro, 1951); week learners can be strengthened through boosting (Schapire, 1990). Nevertheless, many online estimators are found by sacrificing some statistical efficiency and little is known about the reason behind. In light of it, we propose a general framework to understand and construct efficient nonparametric estimators for online problems. Long-run variance (LRV) is an exemplary estimand but our framework applies not only to LRV; see Section 5 for further discussion.

Statistically, we open the door to methodological studies of online estimators. Prior to this paper, online estimators were defined directly and their algorithms were provided to show their computational properties. In LRV estimation, existing works (Wu, 2009; Zheng et al., 2016; Chan and Yau, 2016, 2017a; Chen et al., 2020; Zhu et al., 2021) only focus on how to select subsamples so that a batch means estimator can be updated online. Nevertheless, their proposals are all dominated by an offline benchmark that they used and none of them know the reason. Using the proposed framework, we pinpoint the source of sub-efficiency in their proposals. While some refinements can be made already, we go beyond that and derive sufficient conditions for O(1)-time or O(1)-space update, which allows us to construct online estimators that even dominate the offline benchmark in the literature.

Computationally, we introduce mini-batch estimation to accelerate online estimators for real-time applications. In practice, users may not be interested in every estimate but estimates at predetermined points $n_1, n_2, ...$ instead. For instance, one may assess the convergence of a MCMC sample every m iterations for a fixed $m \in \mathbb{Z}^+$ (Brooks et al., 2011). On the other hand, high frequency data may arrive faster than executing an online algorithm. To solve these problems, we bring the concept of mini-batch training in machine learning (Goodfellow et al., 2016) to LRV estimation. We also present an automatic optimal parameters selector so that users need not to run any offline pilot procedure.

Practically, we demonstrate how to use our framework with recent development in change

point detection, causal inference, and stochastic approximation. A notable issue with online techniques is that a reasonable terminal sample size is not known *a priori*. In light of it, statisticians perform inference to terminate their procedures; see, e.g., Flegal and Jones (2010) and Vats et al. (2019). Lesser known to the statistics community, engineers perform inference to improve their learning algorithms; see, e.g., Lang et al. (2019). Apart from illustrating the strength of our framework in these applications, we hope to promote inference-based techniques developed by one community to another.

The rest of this paper is organized as follows. Section 1.2 reviews relevant literature and introduces some notation. Section 2 proposes a general framework for efficient nonparametric estimation, from which we construct a LRV estimator methodologically. Section 3 illustrates its statistical efficiency through asymptotic theory and Monte Carlo experiment. Section 4 demonstrates its computational efficiency in five interesting applications from different communities. Section 5 discusses our findings. The R-package **rlaser**, which implements our estimators, and the supplement, which contains most algorithms, proofs and ancillary results, are available online. All experiments are performed on Red Hat Enterprise Linux 7.9 with an Intel Xeon CPU E5-2680 v4 and R version 4.1.1.

1.2 Review

Suppose the sample mean $\bar{X}_n \coloneqq n^{-1} \sum_{i=1}^n X_i$ is of interest in an online problem. When the data $\{X_i\}$ are independent and identically distributed (IID), Welford (1962) proposed the first online estimator of $n \operatorname{Var}(\bar{X}_n)$:

$$\hat{\sigma}_{n,\mathsf{WFD}}^{2} = \frac{1}{n} \sum_{i=1}^{n} (X_{i} - \bar{X}_{n})^{2} = \frac{n-1}{n} \left\{ \hat{\sigma}_{n-1,\mathsf{WFD}}^{2} + \frac{1}{n} (X_{n} - \bar{X}_{n-1})^{2} \right\},$$
(1.1)

where $\hat{\sigma}_{0,\text{WFD}}^2 = 0$. The $\hat{\sigma}_{n,\text{WFD}}^2$ is an online estimator because only a finite number of arithmetic operations are involved (O(1)-time update) and a finite number of statistics need to be stored (O(1)-time update). Unfortunately, the IID assumption does not hold in many online problems. To be more realistic, consider $\{X_i\}_{i\in\mathbb{Z}}$ to be stationary and ergodic with mean $\mu \coloneqq \mathbb{E}(X_1)$ and autocovariance function (ACVF) $\gamma_k \coloneqq \mathbb{E}\{(X_0 - \mu)(X_k - \mu)\}, k \in \mathbb{N}_0 \coloneqq \{0, 1, \ldots\}$. Under some suitable conditions (see, e.g., Hannan (1979)), the LRV

$$\lim_{n \to \infty} n \operatorname{Var}(\bar{X}_n) = \sum_{k \in \mathbb{Z}} \gamma_k =: \sigma^2$$
(1.2)

accounts for the serial dependence. In contrast, $\hat{\sigma}_{n,\mathsf{WFD}}^2$ only estimates γ_0 .

LRV estimation has a long history because it is crucial to the inference of dependent data. Let ℓ_n be the batch size (also known as the bandwidth parameter) that is monotonically increasing in n. A classical estimator utilizing the overlapping batch means (obm) (Meketon and Schmeiser, 1984) is

$$\hat{\sigma}_{n,\text{obm}}^{2} \coloneqq \frac{\sum_{i=\ell_{n}}^{n} \left(\sum_{j=i-\ell_{n}+1}^{i} X_{j} - \ell_{n} \bar{X}_{n}\right)^{2}}{\sum_{i=\ell_{n}}^{n} \ell_{n}}.$$
(1.3)

However, it is an offline estimator because there is no relation like (1.1) between $\hat{\sigma}_{n+1,\text{obm}}^2$ and $\hat{\sigma}_{n,\text{obm}}^2$ (Wu, 2009).

Online LRV estimation starts with a series of literature on dynamic batch means (DBM), which is proposed in Yeh and Schmeiser (2000). DBM allows O(1)-time and O(1)-space update but may lead to inconsistency. The seminal work of Wu (2009) proposed the first consistent online estimator based on triangular selection rule (\triangle SR). Specifically, a subsample selection rule modifies $\hat{\sigma}_{n,\text{obm}}^2$ by replacing ℓ_n with a sequence of batch sizes $\{\ell_i\}_{i=1}^n$, which leads to

$$\hat{\sigma}_{n,\mathsf{sub}}^2 \coloneqq \frac{\sum_{i=1}^n \left(\sum_{j=i-\ell_i+1}^i X_j - \ell_i \bar{X}_n \right)^2}{\sum_{i=1}^n \ell_i}.$$
(1.4)

The \triangle SR is an algorithm for constructing $\{\ell_i\}_{i=1}^n$. However, the asymptotic mean squared error (AMSE) of \triangle SR is much higher than that of $\hat{\sigma}_{n,\text{obm}}^2$. Chan and Yau (2016, 2017a) proposed trapezoidal (TSR) and parallelogrammatic (PSR) selection rules to improve the AMSE from 1.78 times (\triangle SR) to 1.20 times (TSR) to 1.12 times (PSR) of $\hat{\sigma}_{n,\text{obm}}^2$.

Now, we introduce some notation. For $a, b \in \mathbb{R}$, define $\sum_{k=a}^{b} x_k = \sum_{a \le k \le b, k \in \mathbb{Z}} x_k$ and $a \lor b = \max(a, b)$. Denote the floor, ceiling and indicator functions by $\lfloor \cdot \rfloor$, $\lceil \cdot \rceil$ and $\mathbb{1}_{\{\cdot\}}$. For real sequences $\{a_n\}$ and $\{b_n\}$, write $a_n \sim b_n$ if $\lim_{n \to \infty} a_n/b_n = 1$. For $p \ge 1$, denote the \mathscr{L}^p -norm by $\lVert \cdot \rVert_p$. A random variable ζ is said to be in \mathscr{L}^p if $\lVert \zeta \rVert_p < \infty$.

Remark 1. Existing online LRV estimation literature also includes Zheng et al. (2016), Chen et al. (2020) and Zhu et al. (2021). To be specific, Zheng et al. (2016) applied prewhitening to \triangle SR in Wu (2009). Chen et al. (2020) proposed increasing batch sizes for inference related to stochastic gradient descent, which is same as the generalized PSR in Chan and Yau (2017a) without overlapping under stationarity. Zhu et al. (2021) adopted a fully online approach compared with Chen et al. (2020), which is same as the generalized \triangle SR in Chan and Yau (2017a) under stationarity.

2 General Framework

2.1 Window Decomposition

Existing literature often regards online update as an algorithmic property and studies online estimators on an algorithm-by-algorithm basis. This is not only true for LRV but also other estimands, e.g., in Xiao and Wu (2011) and Huang et al. (2014). Consequently, the search for efficient online estimators is not methodological as the relation between the statistical efficiency, computational efficiency and form of estimators remains an open problem. In this section, we study this problem in an exemplary setting of LRV estimation. Consider the following quadratic form:

$$\hat{\sigma}_n^2 = \hat{\sigma}_n^2(W) \coloneqq \frac{1}{n} \sum_{i=1}^n \sum_{j=1}^n W_n(i,j) (X_i - \bar{X}_n) (X_j - \bar{X}_n), \qquad (2.1)$$

where $W_n(i, j)$ is a window function (also known as a kernel). This choice is motivated by Song and Schmeiser (1993), who used (2.1) to show that the offline benchmark $\hat{\sigma}_{n,\text{obm}}^2$ and the Bartlett kernel estimator (Newey and West, 1987),

$$\hat{\sigma}_{n,\text{bart}}^2 \coloneqq \sum_{k=-\ell_n}^{\ell_n} \left(1 - \frac{|k|}{\ell_n} \right) \frac{1}{n} \sum_{i=|k|+1}^n (X_i - \bar{X}_n) (X_{i-|k|} - \bar{X}_n),$$
(2.2)

are asymptotically equivalent. Now, we propose a new decomposition that underlies our framework:

$$W_n(i,j) = T\left(d_n^T(i,j)\right) S\left(d_n^S(i,j)\right),\tag{2.3}$$

where $T(\cdot) : [0, \infty) \to \mathbb{R}$ is called a tapering function, $S(\cdot) : [0, \infty) \to \{0, 1\}$ is called a subsampling function, and $d_n^T(i, j) > 0$ and $d_n^S(i, j) > 0$ are distances between times *i* and *j* when the sample size is *n*. Since (2.3) disentangles the tapering weight and subsampling frequency, it allows us to understand and construct efficient online estimators methodologically. For instance, Example 2 shows that it can easily identify how to improve the state-of-the-art PSR. Moreover, (2.3) can be considered as a generalization of kernel and subsampling methods in both offline and online settings. It can be used with many innovations in the literature, which we are going to see a few later.

To get a grasp of the strength of (2.3), the distances $d_n^T(i,j)$ and $d_n^S(i,j)$ in (2.3) are naturally

$$d_n^T(i,j) = \frac{|i-j|}{t_n(i,j)}$$
 and $d_n^S(i,j) = \frac{|i-j|}{s_n(i,j)}$ (2.4)

in order to measure the standardized time lag |i - j| between observations X_i and X_j , where

 $t_n(i,j) > 0$ and $s_n(i,j) > 0$ are some smoothing parameters. Now, recall the fact that $\hat{\sigma}_{n,\text{obm}}^2$ and $\hat{\sigma}_{n,\text{bart}}^2$ are asymptotically equivalent (Song and Schmeiser, 1993). Under (2.3), this can be explained by both of them having the same window that satisfies (2.4). In contrast, the existing online LRV estimators are sub-efficient because they violate (2.4); see Example 2.

The tapering function $T(d_n^T(i, j))$ mainly controls the statistical efficiency as it determines how much weight to be placed on X_iX_j , which estimates the ACVF at lag |i - j|. To reduce variance, it is sensible to assign a lighter weight when the distance $d_n^T(i, j)$ is large. However, if $T(d_n^T(i, j))$ tapers X_iX_j too much, a larger bias is introduced. Hence, the taper governs the bias-variance tradeoff and affects the statistical efficiency.

The subsampling function $S(d_n^S(i,j))$ mainly controls the computational efficiency as it determines which observations to be included in each subsample. Since $\{X_j\}_{j>i}$ is not observed at time *i* in an online setting, consider

$$\hat{\sigma}_n^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n) \sum_{j=1}^i W'_n(i,j) (X_j - \bar{X}_n), \qquad (2.5)$$

where $W'_n(i,j) = (2 - \mathbb{1}_{i=j})W_n(i,j)$ assuming $W_n(i,j)$ is symmetric. Then, the *i*-th subsample can be interpreted as $\mathcal{S}_n(i) = \{X_j : S(d_n^S(i,j)) = 1, 1 \le j \le i\}$. Intuitively, the computational cost increases with the subsample size. Moreover, if $\mathcal{S}_n(1), \ldots, \mathcal{S}_n(n)$ are defined in an "irregular" way, it is hard to update the subsamples, and directly affects the time and space complexity.

The next two examples use (2.3) to identify the structural problem of existing online LRV estimators.

Example 1 (Offline estimators). Let T(x) = 1-x; $S(x) = \mathbb{1}_{x \leq 1}$; and $d_n^T(i,j) = d_n^S(i,j) = |i-j|/\ell_n$, where $\ell_n = \Lambda n^{1/3}$ for some $\Lambda \in \mathbb{R}^+$. Then, it gives celebrated Bartlett kernel:

$$W_{\mathsf{bart}} = W_{n,\mathsf{bart}}(i,j) = \left(1 - \frac{|i-j|}{\ell_n}\right) \mathbb{1}_{|i-j| \le \ell_n}.$$
(2.6)

Other kernel functions $\mathscr{K}(x) = T(x)\mathbb{1}_{x\leq 1}$ are formed similarly, e.g., $T(x) = \{1 + \cos(\pi x)\}/2$ leads to the Tukey–Hanning kernel. Hence, the classical kernel estimators (Andrews, 1991) are our special cases. For these estimators, there are two important features:

- The distances that determine the tapering and subsampling behaviors, i.e., $d_n^T(i,j)$ and $d_n^S(i,j)$, are identical, and *simultaneously* controlled by the same bandwidth ℓ_n .
- The distances $d_n^T(i,j)$ and $d_n^S(i,j)$ depend not only on i,j but also on n because both distances are standardized by the global bandwidth ℓ_n .

Example 2 (Online estimators). Let T(x) = 1 - x; $S(x) = \mathbb{1}_{x \le 1}$; $d_n^T(i, j) = (|i - j| + \overline{\ell}_n - \ell_{i \lor j})/\overline{\ell}_n$; and $d_n^S(i, j) = |i - j|/\ell_{i \lor j}$, where $\ell_n = \Lambda n^{1/3}$, and $\overline{\ell}_n = n^{-1} \sum_{i=1}^n \ell_i$ for some $\Lambda \in \mathbb{R}^+$. Then, the window of PSR can be written as

$$W_{\mathsf{PSR}} = W_{n,\mathsf{PSR}}(i,j) = \left(1 - \frac{|i-j| + \overline{\ell}_n - \ell_{i \lor j}}{\overline{\ell}_n}\right) \mathbb{1}_{|i-j| \le \ell_{i \lor j}}.$$
(2.7)

The window of other existing online estimators in the form of (1.4) can be similarly represented as in (2.7); see Supplement F.2. For these estimators, there are also two important features:

- The natural distance |i j| in (2.7) is *distorted* by an amount of $\overline{\ell}_n \ell_{i \vee j}$, which is counterintuitive and inflexible.
- The distances $d_n^T(i,j)$ and $d_n^S(i,j)$ are separately regularized by the global average bandwidth $\overline{\ell}_n$ and the local bandwidth $\ell_{i\vee j}$, respectively.

Example 2 suggests some possible ways to refine existing online LRV estimators. Indeed, a substitution of $\bar{\ell}_n$ with $\ell_{i\vee j}$ will already improve the AMSE; see Corollary 3(a). However, we can achieve more by finding sufficient conditions for O(1)-time or O(1)-space update. In this way, efficient LRV estimators can be constructed methodologically.

Remark 2. The decomposition in (2.3) is not unique as $S(d_n^S(i,j))$ can be absorbed into the definition of $T(d_n^T(i,j))$, which reduces to the classical way to define a window; see Example 1. Nevertheless, (2.3) is flexible for studying the statistical and computational properties. The decomposed functions $T(\cdot)$ and $S(\cdot)$ are named as tapering and subsampling to stay close with the literature; see, e.g., Wu (2011) and Chan and Yau (2017a).

Remark 3. While we only consider (2.4) in this paper, there might be cases where other distances are natural; see Rho and Vogelsang (2019) for an example in a missing data context.

2.2 Principle-driven Sufficient Conditions

In this subsection, we characterize efficient online $\hat{\sigma}_n^2$ in a principle-driven way and construct an estimator. In contrast to standard texts in online estimation such as Ljung and Söderström (1983) and Young (2011), which regard O(1)-time and O(1)-space update collectively as online update, we discuss both updates separately as constant memory may be nonessential and adversely affect the statistical efficiency.

LASER principles. Consider (2.3). The following step-by-step estimation principles, which

can be summarized as Local Alike Separated Exterior Ramping, unite the statistical and computational properties of $\hat{\sigma}_n^2$.

- (L) An online estimator should utilize local subsamples.
- (A) Under stationarity, if (X_i, X_j) and $(X_{i'}, X_{j'})$ are included in the subsamples and |i j| = |i' j'|, they should be weighted alike.
- (S) The tapering and subsampling parameters should be separately chosen.
- (E) An O(1)-time estimator should be able to exteriorize the tapering parameter from the weights of every subsample.
- (R) An O(1)-space estimator should ramp up the subsampling parameter until it is too large.

Philosophically, Principle L means that online estimates should be adapted to the present stage, i.e., the future (e.g., the future sample size n) should not affect the already computed statistics. In terms of LRV estimation, the subsamples $S_n(i)$ should be constructed based on their local time i and unaffected when the global time n increases.

Principle A states that when the distances are the same, the data pairs contain the same amount of information under stationarity and so should be treated equally. This is sensible in time series and helps to explain the sub-efficiency of existing online estimators. For example, the state-of-the-art PSR in (2.7) violates Principle A.

Principle S introduces more flexibility to meet different user requirements. In Examples 1 and 2, $W_n(i, j)$ uses the same smoothing parameter ℓ_n for both tapering and subsampling. Consequently, tuning ℓ_n affects the statistical and computational efficiency simultaneously, which is usually undesirable. By choosing separate parameters for tapering and subsampling, $\hat{\sigma}_n^2$ can be tailored for different tasks.

Principle E focuses on the relation between O(1)-time update and the taper. By Principle A, $t_n(i,j)$ should be a function of n such that the distances are standardized by the same factor. If the *n*-related part of $t_n(i,j)$ can be "exteriorized" out of the weights assigned to every subsample, the weighted sums will be local and O(1)-time update is possible. For illustration, consider T(x) = 1 - x in Example 1 with S(x) = 1 and omit $2 - \mathbb{1}_{i=j}$ in (2.5). Then,

$$\frac{1}{n}\sum_{i=1}^{n} (X_i - \bar{X}_n) \sum_{j=1}^{i} \left(1 - \frac{|i-j|}{\ell_n}\right) (X_j - \bar{X}_n)$$
$$= \frac{\ell_n}{n} \sum_{i=1}^{n} (X_i - \bar{X}_n) \sum_{j=1}^{i} (X_j - \bar{X}_n) - \frac{1}{n\ell_n} \sum_{i=1}^{n} (X_i - \bar{X}_n) \sum_{j=1}^{i} (i-j) (X_j - \bar{X}_n)$$

can be updated in O(1) time because the weighted sums $\sum_{j=1}^{i} (X_j - \bar{X}_n)$ and $\sum_{j=1}^{i} (i-j)(X_j - \bar{X}_n)$

are local. In contrast, we cannot exteriorize ℓ_n out of $T(|i-j|/\ell_n)$ when $T(x) = \{1 + \cos(\pi x)\}/2$.

Principle R suggests a relation between O(1)-space update and the subsample. If the subsamples change far observations too frequently, e.g., $(X_1, X_2), (X_2, X_3), (X_3, X_4), (X_4, X_5), \ldots$, the memory cost is high as individual past observations must be remembered to be removed from the summary statistics. In contrast, if we ramp up the subsamples' sizes moderately, e.g., $(X_1, X_2), (X_1, X_2, X_3), (X_1, X_2, X_3, X_4), (X_4, X_5), \ldots, O(1)$ -space update is possible. In this spirit, we define the effective (ramped) subsampling parameter as

$$s'_{n} := \begin{cases} s'_{n-1} + 1, & \text{if } s_{n-1} \le s'_{n-1} + 1 < \phi s_{n-1}; \\ s_{n}, & \text{if } s'_{n-1} + 1 \ge \phi s_{n-1} \end{cases}$$
(2.8)

where s_n is the intended subsampling parameter, and $\phi \in [1, \infty)$ is the memory parameter.

Before revealing the sufficient conditions, we shall state the constructed estimator for easy reference:

$$\hat{\sigma}_{n,\mathsf{LASER}(q,\phi)}^{2} = \hat{\sigma}_{n}^{2}(W_{\mathsf{LASER}(q,\phi)}) \qquad \text{with} \qquad W_{n,\mathsf{LASER}(q,\phi)}(i,j) = \left(1 - \frac{|i-j|^{q}}{t_{n}^{q}}\right) \mathbb{1}_{|i-j| \le s_{i \lor j}'}, \quad (2.9)$$

where $\Theta, \Psi \in \mathbb{R}^+$; $\theta, \psi \in (0,1)$; $q \in \mathbb{Z}^+$; $\phi \in [1,\infty)$; $t_n \sim \Theta n^{\theta}$; $s_n \sim \Psi n^{\psi}$; and s'_n can be found in (2.8). If there is no ambiguity, we will refer to (2.9) as $\mathsf{LASER}(q,\phi)$, whose name comes from the LASER principles. We remark that $\mathsf{LASER}(q,\phi)$ reduces to $\hat{\sigma}^2_{n,\mathsf{WFD}}$ when $s_i = 0$ for $i \in \mathbb{Z}^+$, and $\hat{\sigma}^2_{n,\mathsf{bart}}$ when $q = \phi = 1$ and $s_i = t_n = \ell_n$ for $i \in \mathbb{Z}^+$.

Proposition 1 (O(1)-time update of $\hat{\sigma}_n^2$). Consider (2.3) with standardized distances in (2.4) and a subsampling function $S(x) = \mathbb{1}_{x \leq 1}$. Let $q \in \mathbb{Z}^+$; $a_0, \ldots, a_q \in \mathbb{R}$; and $\{s_1, s_2, \ldots\}, \{t_1, t_2, \ldots\}, \{t'_1, t'_2, \ldots\}$ be some non-zero sequences. If

- (L) the subsampling parameter is local, i.e., $s_n(i,j) = s_{i \lor j}$;
- (A) the tapering parameter is separable in $i \lor j$ and n, i.e., $t_n(i, j) = t_n t'_{i\lor j}$, with $\{t'_i\}$ preferably a constant sequence;
- (S) the sequences $\{s_i\}$ and $\{t_i\}$ are preferably different;
- (E) the tapering function is of the form $T(x) = \sum_{r=0}^{q} a_r x^r$, preferably with $a_0 = 1$, $a_q = -1$, and $a_1 = \cdots = a_{q-1} = 0$; and

(R) the subsampling parameter in (L) is preferably replaced by $s'_{i \vee j}$ in (2.8) with $\phi \in [1, \infty)$, then $\hat{\sigma}_n^2$ can be updated in O(1) time.

Proposition 1 considers a polynomial T(x) for two reasons. First, it is theoretically interesting as other windows can be approximated by a polynomial basis; see, e.g., Sun et al. (2011).

Estimator	or			Statistical Properties			Computational Properties		
$\hat{\sigma}_n^2$	q	ϕ	Flexibility	Optimal AMSE	Automatic	Time	Space	Mini-batch	
bart (Newey and West, 1987)	1	/	medium	$B_n = 2.29\sigma^4 \kappa_1^{2/3} n^{-2/3}$	×	O(n)	O(n)	X	
parz (Parzen, 1957)	3	/	medium	$P_n = 3.39\sigma^4 \kappa_3^{2/7} n^{-6/7}$	×	O(n)	O(n)	×	
WFD (Welford, 1962)	1	/	low	inconsistent for σ^2	/	O(1)	O(1)	/	
DBM (Yeh and Schmeiser, 2000)	1	/	low	inconsistent for σ^2	×	O(1)	O(1)	×	
△SR (Wu, 2009)	1	/	medium	$1.78B_{n}$	×	O(1)	O(1)	×	
TSR (Chan and Yau, 2016)	1	/	medium	$1.20B_{n}$	×	O(1)	O(1)	×	
PSR (Chan and Yau, 2016)	1	/	medium	$1.12B_{n}$	×	O(1)	$O(n^{1/3})$	×	
PSR (Chan and Yau, 2017a)	3	/	medium	$1.06P_{n}$	1	O(1)	$O(n^{1/7})$	X	
LASER (proposal)	1	2	high	$1.01B_{n}$	1	O(1)	O(1)	X	
LASER (proposal)	1	1	high	$0.96B_n$	1	O(1)	$O(n^{1/3})$	1	
LASER (proposal)	3	1	high	$0.98P_{n}$	1	O(1)	$O(n^{1/7})$	1	

Table 1: Summary of properties of different LRV estimators. For the discussion of different properties, see Sections 2.1 (Flexibility), 3.2 (Optimal AMSE), 3.3 (Automatic), 4.1 (Time), 4.2 (Space), and 4.3 (Mini-batch).

Second, it is compatible with the existing selection rule framework as \triangle SR, TSR and PSR can be written as a linear combination of several parts that satisfy Proposition 1. However, we notice that a separate set of statistics is needed for online updates of each term in the polynomial. As the flatness at origin determines the convergence rate, it is more practical to consider $T(x) = 1 - x^q$, which was also suggested by Parzen (1957). Along with the other preferable conditions, we construct LASER (q, ϕ) .

Proposition 2 (O(1)-space update of $\hat{\sigma}_n^2$). Consider (2.3) with standardized distances in (2.4) and a subsampling function $S(x) = \mathbb{1}_{x \leq 1}$. Let $\{s_n \in \mathbb{N}_0\}_{n \in \mathbb{Z}^+}$ be a fixed sequence or a monotonically increasing random sequence. Suppose the estimator $\hat{\sigma}_n^2$ can be updated in O(1) time. If, in addition,

(R) the subsampling parameter is ramped with $\phi \ge 2$, i.e., $s_n(i,j) = s'_{i \lor j}$ in (2.8), then $\hat{\sigma}_n^2$ can also be updated in O(1) space.

Note that Proposition 2 applies to any O(1)-time $\hat{\sigma}_n^2$. It considers O(1)-space update as an additional property to O(1)-time update because the latter is more fundamental in online problems. According to Principle R, we need to prepare for resets of the effective subsampling parameter s'_n . This is possible if we know the intended s_n in advance, e.g., when $\{s_n\}$ is prespecified, or data-driven but monotonically increasing. By Propositions 1 and 2, LASER (q, ϕ) is always O(1)-time, and O(1)-space when $\phi \geq 2$.

Before proceeding, we summarize the properties of different LRV estimators in Table 1. While we have restricted our attention to online estimators, we emphasize that our construction can be used with many innovations in the offline estimation literature; see Example 5. Our framework can also be extended easily to estimands other than LRV; see Section 5. For the other properties, we highlight that automatic update frees users from smoothing parameters selection so that they only need to decide the characteristic exponent q and the memory parameter ϕ with respect to their problems. In general, we recommend LASER(1,1). If memory is scarce, we recommend LASER(1,2).

Remark 4. Without affecting any properties, $LASER(q, \phi)$ can be extended to estimate the long-run covariance matrix (LRCM). Supplement A.9 describes the multivariate algorithm implemented in our R-package.

Remark 5. The finite-sample estimates of $LASER(q, \phi)$ may not be positive definite. This is because we are motivated to search for efficient online estimators rather than efficient positive semi-definite estimators in Andrews (1991). To resolve this issue, we follow Jentsch and Politis (2015) to provide an adjustment that retains asymptotic properties in our R-package; see also Vats and Flegal (2021). Supplement F.1 restates their procedure and points out that it can preserve the computational properties of $LASER(q, \phi)$.

3 Statistical Properties

3.1 Consistency

Hereafter, we develop the asymptotic theory of $\hat{\sigma}_n^2$ based on the dependence measures of Wu (2005). Let $X_i = g(\mathscr{F}_i)$ for some measurable function g, where $\mathscr{F}_i \coloneqq (\dots, \epsilon_{i-1}, \epsilon_i)$ is the shift process of IID innovations $\{\epsilon_i\}_{i\in\mathbb{Z}}$. Let ϵ'_j be an IID copy of ϵ_j , $\mathscr{F}_{i,\{j\}} \coloneqq (\mathscr{F}_{j-1}, \epsilon'_j, \epsilon_{j+1}, \dots, \epsilon_i)$ and $X_{i,\{j\}} = g(\mathscr{F}_{i,\{j\}})$. Wu (2005) defined:

- physical dependence measure: $\delta_{i,p} \coloneqq \|X_i X_{i,\{0\}}\|_p$; and
- predictive dependence measure: $\omega_{i,p} \coloneqq \|\mathbb{E}(X_i \mid \mathscr{F}_0) \mathbb{E}(X_i \mid \mathscr{F}_{0,\{0\}})\|_p$.

This framework has a wide range of applications in practice, which include but are not limited to LRV estimation. Interested readers are referred to Wu (2011). We also use this framework in coherence with the online LRV estimation literature. For the consistency of $\hat{\sigma}_n^2$, the following regularity conditions are imposed.

Assumption 1 (α -stability). For some $\alpha > 2$, $\Delta_{\alpha} \coloneqq \sum_{i=0}^{\infty} \delta_{i,\alpha} < \infty$.

Assumption 2 (Summability of window). Let $\alpha' = \min(\alpha/2, 2)$. Define

$$G_{1,n} \coloneqq \max_{1 \le i \le n} \sum_{j=1}^{n} \left\{ \left| W_n(i,j) \right| + W_n^2(i,j) \right\},\$$

$$G_{2,n} \coloneqq \max_{1 \le i,j \le n} \left| W_n(i,j) \right| + \max_{1 \le i \le n} \left\{ \sum_{j=2}^{n} \left| W_n(i,j) - W_n(i,j-1) \right|^{\alpha'} \right\}^{\frac{1}{\alpha'}}$$

The window function in (2.1) satisfies $G_{1,n} = o(n^{2-2/\alpha'})$ and $G_{2,n} = o(n^c)$ for some constant $c \in (0, 1 - 1/\alpha')$.

Assumption 3 (General window for LRV estimation). For $n \in \mathbb{Z}^+$ and k = 0, ..., n-1, define $w_{n,k} \coloneqq n^{-1} \sum_{i=k+1}^{n} W_n(i, i-k)$. The window function in (2.1) satisfies

- (a) $W_n(i,j) = W_n(j,i)$ for all $n \in \mathbb{Z}^+$ and $i, j \in \{1,\ldots,n\}$;
- (b) $W_n(i,i) = 1$ for all $n \in \mathbb{Z}^+$ and $i \in \{1,\ldots,n\}$; and
- (c) there exists an increasing sequence $\{b_n\}_{n \in \mathbb{Z}^+}$ that diverges to ∞ as $n \to \infty$ such that $\max_{1 \le k \le b_n} |w_{n,k} - 1| = o(1)$ and $\max_{b_n < k < n} |w_{n,k}| = O(1)$.

Assumption 1 is known as the stability condition in Wu (2005), which ensures that $\sigma^2 < \infty$. Assumptions 2 and 3 regulate the behavior of a general window so that $\hat{\sigma}_n^2$ is guaranteed to be precise and accurate, respectively. These assumptions are easily verifiable once the window $W_n(i, j)$ and the smoothing parameters $s_n(i, j)$ and $t_n(i, j)$ are specified. We emphasize that they are mild and satisfied by many windows and smoothing parameters. For instance, they hold under Definition 1; see Corollary 1. The next theorem shows the $\mathscr{L}^{\alpha/2}$ -consistency of $\hat{\sigma}_n^2$.

Theorem 1 (Consistency of $\hat{\sigma}_n^2$). Let $\alpha > 2$. Suppose that $X_1 \in \mathscr{L}^{\alpha}$. If Assumptions 1–3 hold, then $\|\hat{\sigma}_n^2 - \sigma^2\|_{\alpha/2} = o(1)$.

We highlight that Theorem 1 applies to general $\hat{\sigma}_n^2$, which covers both offline and online estimators, e.g., all consistent estimators stated in Table 1. Future research which propose new principles based on our framework may utilize Theorem 1 to prove consistency. For the proposed LASER principles, we have the following recommendation.

Assumption 4 (\tilde{q} -th order serial dependence). For some $\tilde{q} \in \mathbb{Z}^+$, $u_{\tilde{q}} \coloneqq \sum_{k \in \mathbb{Z}} |k|^{\tilde{q}} |\gamma_k| < \infty$.

Definition 1 (Parameters of LASER (q, ϕ)). Let $q \in \mathbb{Z}^+$ and $\phi \in [1, \infty)$ be fixed; the effective subsampling parameter s'_n depends on s_n according to (2.8); and $\alpha' = \min(\alpha/2, 2)$. The window function takes the form in (2.9). The intended subsampling parameter and tapering parameter take the form $s_n = \min(\lfloor \Psi n^{\psi} \rfloor, n-1)$ and $t_n = \min(\lceil \Theta n^{\theta} \rceil, n)$. The coefficients satisfy $\Psi, \Theta \in \mathbb{R}^+$, and the exponents satisfy either (a) $0 < \psi \le \theta < \min\{2 - 2/\alpha', 1/(1+q)\};$ or

(b) $0 < \psi < \min\{2 - 2/\alpha', 1/(1+q)\}\ \text{and}\ \max\{\psi + (\psi - 2 + 2/\alpha')/(2q), (q - \tilde{q})\psi/q\} < \theta < \psi,\$

where \tilde{q} is the order of serial dependence in Assumption 4.

By replacing Assumptions 2 and 3 with Definition 1, the next Corollary shows the $\mathscr{L}^{\alpha/2}$ consistency of LASER (q, ϕ) .

Corollary 1 (Consistency of LASER (q, ϕ)). Let $\alpha > 2$. Suppose that $X_1 \in \mathscr{L}^{\alpha}$ and Assumption 1 holds. Under Definition 1(a), or Definition 1(b) with Assumption 4, $\|\hat{\sigma}_{n, \mathsf{LASER}(q, \phi)}^2 - \sigma^2\|_{\alpha/2} = o(1)$.

Since $\{X_i\}$ are dependent, some form of mixing is needed to establish consistency. Assumption 1 is one of such conditions and cannot be replaced in the corollary. For Definition 1, (a) is in line with the literature. When $\alpha \ge 4$, we have $\alpha' = 2$ so $\min\{2 - 2/\alpha', 1/(1+q)\} = 1/(1+q)$. This resembles the same upper bound of the bandwidth for $\hat{\sigma}_{n,\text{bart}}^2$ in Andrews (1991). When $2 < \alpha < 4$, we have $1 < \alpha' < 2$ so the moment condition may affect the feasible range of (ψ, θ) . When $\psi \le \theta$, i.e., case (a), the estimator is always consistent. However, if $\psi > \theta$, i.e., case (b), s_n/t_n diverges so the window is not absolutely bounded. In this case, the estimator is inconsistent unless some condition is imposed on the serial dependence. Definition 1(b) handles it under Assumption 4. Consequently, θ admits the lower bound $(q - \tilde{q})\psi/q$. We remark that consistency under $\psi \neq \theta$ is not established in the literature because of the novelty of our framework.

3.2 MSE-optimal Convergence Rate

In this subsection, we establish the exact convergence rates for the variance and bias of $LASER(q, \phi)$. Consider

Assumption 5 (q-th order weak α -stability). For some $\alpha \ge 1$ and $q \in \mathbb{Z}^+$, $\Omega_{\alpha}^{(q)} \coloneqq \sum_{j=0}^{\infty} j^q \omega_{j,\alpha} < \infty$.

Assumption 5 is satisfied by a broad class of linear and nonlinear time series, e.g., ARMA model, Bilinear model, Generalized Autoregressive Conditional Heteroskedastic (GARCH) model; see Wu (2011) and the references therein. This assumption also implies $u_q = \sum_{k \in \mathbb{Z}} |k|^q |\gamma_k| < \infty$ and $v_q \coloneqq \sum_{k \in \mathbb{Z}} |k|^q \gamma_k$ is well-defined; see Wu (2009). Then, the following theorem shows the exact \mathscr{L}^2 -convergence rate of LASER (q, ϕ) .

Theorem 2 (Exact convergence rate of $LASER(q, \phi)$). Let $q \in \mathbb{Z}^+$ be fixed. Suppose that $X_1 \in \mathscr{L}^{\alpha}$ and Definition 1 holds. (a) Let $\alpha \ge 4$. If Assumption 1 holds, then as $n \to \infty$,

$$\begin{split} n^{1-\psi-\max\{2q(\psi-\theta),0\}} \mathrm{Var}(\hat{\sigma}_{n}^{2}) &\to \mathscr{V}_{\psi,\Psi,\theta,\Theta,q,\phi}\sigma^{4}, \qquad where \qquad \mathscr{V}_{\psi,\Psi,\theta,\Theta,q,\phi} := \\ \left\{ \begin{array}{l} \frac{2\Psi(\phi+1)}{\psi+1} \mathbbm{1}_{\psi\leq\theta} - \frac{8\Psi^{q+1}\Theta^{-q}(\phi^{q+2}-1)}{(\phi-1)(q+1)(q+2)(\psi q+\psi+1)} \mathbbm{1}_{\psi=\theta} + \frac{2\Psi^{2q+1}\Theta^{-2q}(\phi^{2q+2}-1)}{(\phi-1)(q+1)(2q+1)(2\psi q+\psi+1)} \mathbbm{1}_{\psi\geq\theta}, \quad \phi > 1; \\ \frac{4\Psi}{\psi+1} \mathbbm{1}_{\psi\leq\theta} - \frac{8\Psi^{q+1}\Theta^{-q}}{(q+1)(\psi q+\psi+1)} \mathbbm{1}_{\psi=\theta} + \frac{4\Psi^{2q+1}\Theta^{-2q}}{(2q+1)(2\psi q+\psi+1)} \mathbbm{1}_{\psi\geq\theta}, \qquad \phi = 1. \end{split} \right.$$

(b) Let $\alpha \geq 2$. If Assumption 5 holds for the fixed q, then as $n \to \infty$,

$$\operatorname{Bias}(\hat{\sigma}_n^2) \sim \begin{cases} o(n^{-q\psi}), & \psi < \theta; \\ -\Theta^{-q} n^{-q\theta} v_q, & \psi \ge \theta. \end{cases}$$

Similar to the consistency under $\psi \neq \theta$, the variance under $\alpha = 4$ is not established in the literature. We weaken this moment condition based on the *m*-dependent process approximation in Liu and Wu (2010). Useful lemmas that extend their Propositions 1 and 2 are proved in Supplement E. It follows from Theorem 2 that $\psi = \theta = 1/(1+2q)$ optimize the order of $MSE(\hat{\sigma}_n^2)$ for each q; see Remark 6. The other MSE-optimal parameters are summarized below.

Corollary 2 (MSE-optimal parameters). Define $\kappa_q := |v_q|/\sigma^2$. Suppose that the conditions in Theorem 2(a) and (b) hold. If $\psi = \theta = 1/(1+2q)$, the AMSE-optimal Ψ is

$$\Psi_{\star} = \begin{cases} \left\{ \frac{(\phi+1)(2q+1)}{2q(q+1)} - \frac{4(\phi^{q+2}-1)(2q+1)}{(\phi-1)q(q+1)(q+2)(3q+2)} + \frac{(\phi^{2q+2}-1)}{2(\phi-1)q(q+1)(2q+1)} \right\}^{-\frac{1}{1+2q}} \kappa_q^{\frac{2}{1+2q}}, \quad \phi > 1; \\ \left\{ \frac{2q+1}{q(q+1)} - \frac{4(2q+1)}{q(q+1)(3q+2)} + \frac{1}{q(2q+1)} \right\}^{-\frac{1}{1+2q}} \kappa_q^{\frac{2}{1+2q}}, \qquad \phi = 1. \end{cases}$$

In addition, if $\Theta = \rho \Psi_{\star}$ is allowed for any $\rho \in \mathbb{R}^+$, then the AMSE-optimal Θ is

$$\Theta_{\star} = \begin{cases} \left\{ \frac{(q+2)(3q+2)(\phi^{2q+2}-1)}{4(2q+1)^{2}(\phi^{q+2}-1)} + \frac{\Psi_{\star}^{-2q-1}\kappa_{q}^{2}(\phi-1)(q+1)(q+2)(3q+2)}{4(2q+1)(\phi^{q+2}-1)} \right\}^{\frac{1}{q}} \Psi_{\star}, \quad \phi > 1; \\ \left\{ \frac{(q+1)(3q+2)}{2(2q+1)^{2}} + \frac{\Psi_{\star}^{-2q-1}\kappa_{q}^{2}(q+1)(3q+2)}{4(2q+1)} \right\}^{\frac{1}{q}} \Psi_{\star}, \qquad \phi = 1. \end{cases}$$

If O(1)-space update is required, the AMSE-optimal ϕ is $\phi_{\star} = 2$. Otherwise, the AMSE-optimal ϕ is $\phi_{\star} = 1$.

Corollary 3 (Benefits of our framework). Let $\alpha \ge 4$ and $B_n = 2.29\sigma^4 \kappa_1^{2/3} n^{-2/3}$, where κ_1 is defined in Corollary 2.

- (a) (Understanding the structural problem of existing estimators). Suppose that $X_1 \in \mathscr{L}^{\alpha}$, and Assumptions 1 and 5 hold for q = 1. If $\overline{\ell}_n$ is substituted with $\ell_{i \vee j}$ in (2.7), the AMSEoptimal Λ is $\Lambda_{\star} = (9/2)^{1/3} \kappa_1^{2/3}$. The AMSE improves from $1.12B_n$ (PSR) to $1.08B_n$.
- (b) (Constructing O(1)-time estimator methodologically). Under Corollary 2, the AMSEoptimal parameters for LASER(1,1) are $\psi_{\star} = \theta_{\star} = 1/3$, $\Psi_{\star} = (30/19)^{1/3} \kappa_1^{2/3}$ and $\Theta_{\star} =$

 $(13/12)\Psi_{\star}$. The AMSE improves $1.12B_n$ (PSR) to $0.97B_n$.

(c) (Constructing O(1)-time and O(1)-space estimator methodologically). Under Corollary 2, the AMSE-optimal parameters for LASER(1,2) are $\psi_{\star} = \theta_{\star} = 1/3$, $\Psi_{\star} = (10/7)^{1/3} \kappa_1^{2/3}$ and $\Theta_{\star} = (8/7)\Psi_{\star}$. The AMSE improves $1.20B_n$ (TSR) to $1.01B_n$.

Corollary 2 allows us to compare the optimal AMSEs of $LASER(q, \phi)$ for different q and ϕ with existing offline and online estimators in Table 1. Corollary 3 focuses on q = 1 and confirms the structural problem of existing online estimators highlighted in Example 2. Surprisingly, LASER(1,1) and LASER(1,2) are super and nearly optimal in the sense of Chan and Yau (2017a); see Remark 7. While it is one of the many advantages of $LASER(q, \phi)$, we emphasize that we are proposing a general framework and $LASER(q, \phi)$ is constructed from this framework. Therefore, similar improvements can be expected in other online estimation problems that utilize windows.

Example 3 (Online LRV estimation). To study the finite-sample efficiency, consider the following time series models:

- (I) ARMA(1,1): Let $X_i \mu = a(X_{i-1} \mu) + b\varepsilon_{i-1} + \varepsilon_i$, where $\varepsilon_i \stackrel{\text{iid}}{\sim} N(0,\nu^2)$. Take a = 0.5, b = 0.5, $\nu = 1$ and $\mu = 0$, which results in a mildly autocorrelated linear time series.
- (II) *Bilinear*: Same as (4.1), which yields a strongly autocorrelated nonlinear time series.
- (III) Fractional Gaussian Noise Process: Let X_i be a zero-mean Gaussian processes with ACVF $\mathbb{E}(X_0X_k) = a(k+b)^{-c}$. Take a = 100, b = 5 and c = 5, which leads to a nonlinear time series where $u_3 < \infty$ but $u_4 = \infty$.
- (IV) ARMA(1,1): Same as Model I but take a = 0.2, b = -0.6, $\nu = 1$ and $\mu = 0$, which results in a negatively autocorrelated time series.

Under Models I–IV, we compute the MSEs of different online LRV estimators, which includes TSR , PSR , $\mathsf{LASER}(1,1)$ and $\mathsf{LASER}(1,2)$, based on 1000 replications; see Table 1 for a summary of these estimators. Oracle parameters are used to separate the effect of smoothing parameters selection, which will be investigated in Example 4. Figure 1 confirms the improvements brought by our framework. While $\mathsf{LASER}(1,2)$ performs slightly worse under Model IV, the other O(1)-space estimator TSR is similar. This is because their subsample sizes are non-monotone and vary frequently. Since LRV estimators are like weighted sums of the ACVFs with bounds of summation determined by the subsample sizes, the finite-sample estimates vary more when the data is negatively autocorrelated.



Figure 1: LRV estimation using different online estimators: (a) TSR (dotted gray); (b) PSR (dashed red); (c) LASER(1,1) (longdash blue); (d) LASER(1,2) (dotdash purple). The plots show the value of $MSE(\cdot)/MSE(\hat{\sigma}_{n,bart}^2)$.

Remark 6. When $1/(1 + 2q) = \psi < \theta$, the order of $MSE(\hat{\sigma}_n^2)$ is also optimized. However, the window in Definition 1 becomes rectangular asymptotically. Consequently, there is no known expression for the leading-order term of $MSE(\hat{\sigma}_n^2)$ and the MSE-optimal parameters cannot be found (Chan and Yau, 2017b).

Remark 7. We are aware that some offline estimators have a lower AMSE than bart for q = 1. Nevertheless, the literature has discussed the optimality of online estimators by comparing with obm (which is asymptotically equivalent to bart) because they admit a similar form. We follow the same practice to consider bart and parz in Table 1.

3.3 Automatic Optimal Parameters Selection

By Corollary 2, the MSE-optimal parameters depend on $\kappa_q = |v_q|/\sigma^2$ for some $q \in \mathbb{Z}^+$. To choose q accounting for the empirical serial dependence, we recommend q = 1 (q = 3) for a relatively strongly (weakly) auto-correlated time series. Since the online estimate of σ^2 is available from

the last iteration, it remains to handle $v_q = \sum_{k \in \mathbb{Z}} |k|^q \gamma_k$ for a fixed q. Classical methods such as parametric plug-in (Andrews, 1991) and correlogram threshold (Paparoditis and Politis, 2012) are computationally inefficient. The computational properties of an online estimator can be preserved by applying them on a pilot sample (Wu, 2009), but it may be shortsighted in some online problems. To fully utilize the increasing sample size in online problems, a natural solution is to also estimate v_q online. Consider

$$\hat{v}_{q,n} = \hat{v}_{q,n}(W) \coloneqq \frac{1}{n} \sum_{i=1}^{n} \sum_{j=1}^{n} W_n(i,j) |i-j|^q (X_i - \bar{X}_n) (X_j - \bar{X}_n).$$
(3.1)

Since $\hat{v}_{q,n}$ has a similar form as $\hat{\sigma}_n^2$, our framework, particularly the LASER principles, suggests

$$\hat{v}_{q,n,\mathsf{LASER}(p,\phi)} = \hat{v}_{q,n}(W_{\mathsf{LASER}(p,\phi)}) \tag{3.2}$$

where the window function takes the form in (2.9), q, ϕ are inherited from $\hat{\sigma}_{n,\text{LASER}(q,\phi)}^2$, and p is recommended to be 1; see Remark 8. As $W_{\text{LASER}(p,\phi)}$ satisfies Principle E, it follows from Propositions 1 and 2 immediately that $\hat{v}_{q,n,\text{LASER}(p,\phi)}$ can be updated in O(1) time or space; see Supplement A.7. Updating $\hat{\sigma}_{n,\text{LASER}(q,\phi)}^2$ becomes automatic in the sense that users only need to supply the incoming observations, but the smoothing parameters are optimally selected and the computational properties are preserved. We detail this automatic optimal parameters selector in Supplement A.8 and compare it with some existing methods in the next example.

Example 4 (Smoothing parameter selection). Consider the following smoothing parameters selectors for LASER(1,1):

- (a) best existing: PSR with the automatic update in Chan and Yau (2017a) is included.
- (b) *pilot*: κ_1 is estimated with the first 500 observations. Two Bartlett kernel estimators with asymptotically rate-optimal bandwidths $\lceil n^{1/5} \rceil$ and $\lceil n^{1/3} \rceil$ are used for v_1 and σ^2 , respectively.
- (c) *auto*: κ_1 is handled with the automatic update in this paper.
- (d) oracle: the theoretical value of κ_1 , which is unknown in practice, is used.

Under (4.1), we obtain the trajectories of LASER(1,1) of a typical realization with 10^5 observations. We also compute the efficiency gained (in terms of $\ln MSE(\cdot)$) for m = 500 (see Section 4.3) based on 1000 replications. Figure 2 shows that the *auto* trajectory is very close to that with the *oracle*. Meanwhile, the *pilot* trajectory performs worse in the sense that it is obviously off from the true LRV in finite sample. This is because realizations of (4.1)



Figure 2: LRV estimation using different smoothing parameter selectors: (a) *best existing* (dotted gray); (b) *pilot* (dashed red); (c) *auto* (longdash blue); (d) *oracle* (dotdash purple). The horizontal line in plot (i) is the true LRV of (4.1).

are strongly autocorrelated so 500 observations and asymptotically rate-optimal bandwidths may be insufficient to obtain a good estimate of κ_1 . Although automatic update doubles the computation cost, it is worth noting that the relative cost is negligible compared with the state of the art, and a sufficient sample size for pilot study is unknown *a priori*.

Remark 8. By Theorem 2, a larger q improves the convergence rate of $\hat{\sigma}_{n,\text{LASER}(q,\phi)}^2$ subject to the empirical serial dependence. A larger p improves $\hat{v}_{q,n,\text{LASER}(p,\phi)}$ similarly but this improvement has little effect on the LRV estimate.

4 Computational Properties and Applications

4.1 *O*(1)-time Update and Change Point Detection

Among the properties that characterize efficient online estimators, O(1)-time update is fundamental because the time complexity to update a LRV estimator should not be higher than \bar{X}_n . Otherwise, time spent in updating the LRV estimator may be used to improve \bar{X}_n in online problems. For example, one may generate more observations in a simulation (Yeh and Schmeiser, 2000). While modern hardware allows classical LRV estimators to be computed more quickly, this does not change the fact that their update costs O(n) time (Meketon and Schmeiser, 1984). In contrast, LASER (q, ϕ) can be updated in O(1) time, which is demonstrated with the following condensed version of the algorithm for updating LASER(1, 1). The derivation and other algorithms can be found in Supplement A. Algorithm 1 (LASER(1,1)). Initialize $\mathscr{C}_1 = \{1, 0, 1, X_1^2, X_1, \{0, \dots, 0\}_{b=0,1}, \{X_1\}\}$. At time $n \in \mathbb{Z}^+$, store $\mathscr{C}_n = \{n, s_n, t_n, Q_n, \bar{X}_n, \{K_{n,b}, R_{n,b}, k_{n,b}, r_{n,b}, U_{n,b}, V_{n,b}\}_{b=0,1}, \{X_{n-b}\}_{b=0,\dots,s_n}\}$. At time n + 1, update \mathscr{C}_n to \mathscr{C}_{n+1} by:

$$\begin{array}{ll} \text{(a)} & s_{n+1} = \min\{[\Psi(n+1)^{\psi}], n\}; \\ \text{(b)} & t_{n+1} = \min\{[\Theta(n+1)^{\theta}], n+1\}; \\ \text{(c)} & Q_{n+1} = Q_n + X_{n+1}^2; \\ \text{(d)} & \bar{X}_{n+1} = (n\bar{X}_n + X_{n+1})/(n+1); \\ \text{(e)} & K_{n+1,0} = K_{n,0} + X_n - X_{n-s_n}\mathbb{1}_{s_{n+1}=s_n}; \\ \text{(f)} & K_{n+1,1} = K_{n,1} + K_{n+1,0} - s_n X_{n-s_n}\mathbb{1}_{s_{n+1}=s_n}; \\ \text{(b)} & V_{n+1,b} = V_{n,b} + K_{n+1,b}X_{n+1}; \\ \text{(c)} & V_{n+1,b} = V_{n,b} + K_{n+1,b}X_{n+1}; \\ \end{array}$$

where $D_{n+1}^* = D_{n+1,0} - D_{n+1,1}/t_{n+1}$ for $D \in \{R, r, U, V\}$, e.g., $r_{n+1}^* = r_{n+1,0} - r_{n+1,1}/t_{n+1}$.

The next example shows the usefulness of O(1)-time LRV estimators.

Example 5 (Online CP detection). Recently, Gösmann et al. (2021) developed a new approach for online CP detection in an open-end scenario. In their outlook, they mentioned that the standard approach to LRV estimation in the field was to employ only the initial data but updating the LRV estimate was logical particularly for stronger dependent model (Gösmann et al., 2021). We echo their view in this example. Consider a Bilinear model

$$X_i = (0.9 + 0.1\varepsilon_i)X_{i-1} + \varepsilon_i, \quad \text{where} \quad \varepsilon_i \stackrel{\text{nd}}{\sim} N(0, 1). \tag{4.1}$$

....

Let $\mu_i = \mathbb{E}(X_i)$; $k^* \in \mathbb{Z}^+$; and m be the initial sample size. We are interested in testing

$$H_0: \mu_1 = \dots = \mu_m = \mu_{m+1} = \dots$$
 against $H_1: \mu_1 = \dots = \mu_{m+k^*-1} \neq \mu_{m+k^*} = \mu_{m+k^*+1} = \dots$

In this case, the online CP monitoring scheme in Gösmann et al. (2021) can be written as

$$\hat{E}_m(k) = m^{-1/2} \max_{0 \le j \le k-1} (k-j) \left| \bar{X}_{1,m+j} - \bar{X}_{m+j+1,m+k} \right| /\hat{\sigma},$$

where $\bar{X}_{a,b} = (b-a+1)^{-1} \sum_{i=a}^{b} X_i$ and $\hat{\sigma}^2$ is a LRV estimator. Following examples in Gösmann et al. (2021), we use the same threshold function $w(t) = (1+t)^{-1}$, nominal size $\alpha = 0.05$ and stopping point $n^* = 4000$ to monitor $w(k/m)\hat{E}_m(k)$. We also simulate H_1 by $X_t^{(\delta)} = X_t + \delta \mathbb{1}_{t \ge m+k^*}$ with 100 burn-in and m = 400 initial observations. However, we consider different LRV estimation methods:



Figure 3: Online CP detection at 5% nominal size using different LRV estimation methods: (a) fix (dotted gray); (b) offline (dashed red); (c) online (longdash blue).

- (a) fix: the LRV is estimated with the initial data only, which is the standard approach. $\hat{\sigma}_{m,\text{bart}}^2$ implemented in the R-package sandwich (Zeileis et al., 2020) without prewhitening and adjustment is used.
- (b) offline: the LRV estimate is computed as in (a). However, the estimate is updated as data arrive. For robustness, the first order difference statistics in Chan (2022) instead of the raw data are used. In favor of offline update, the smoothing parameter is first selected based on the initial data and then scaled according to Chan (2022).
- (c) online: the LRV estimate is updated as in (b) but using LASER(1,1).

We conduct the simulation for $m + k^* = 601, 1001, 1401$ and $\delta = 0, 1, \ldots, 5$ each with 1000 replications. Figure 3 reports the results for $m + k^* = 1401$. When the LRV estimate is updated, the type I error is considerably closer to 5%. Furthermore, the positive predictive value improves substantially and is the best using *online* with negligible time cost. The results for $m + k^* = 601, 1001$ are similar and so deferred to the supplement.

Remark 9. We state that the smoothing parameter selection in Example 5 is favorable to *offline* because it reduces the time cost of *offline* much more than *online*. The scaling factor in Chan (2022) due to the use of difference statistics is also optimal to *offline* only. Investigating the scaling factor for *online* is of interest but beyond the scope here.

4.2 *O*(1)-space Update and Convergence Diagnosis

An online estimator is more attractive if it uses less memory. Indeed, data compression is one of the reasons to perform online estimation (Ljung and Söderström, 1983). Algorithm 1 shows that LASER(1,1) reduces the space complexity from O(n) to $O(n^{\psi})$ as only $\{X_{n-b}\}_{b=0,...,s_n}$ and a fixed number of statistics are needed at time n. This is satisfactory because Corollary 2 shows that $\psi = 1/3$ is MSE-optimal. However, one may want to further reduce $O(n^{\psi})$ to O(1) in some real-time applications where memory is limited. To this end, LASER(q, ϕ) provides a memory parameter ϕ so that $\{X_{n-b}\}_{b=0,...,s_n}$ can be replaced by another fixed yet smaller number of statistics with minimal loss of efficiency when $\phi = 2$. The next example illustrates the strength of our O(1)-space LRV estimators.

Example 6 (Online CI construction/Convergence diagnosis). In MCMC methods, a common goal is to estimate $\mathbb{E}_{\zeta \sim \pi} \{h(\zeta)\}$ using $\bar{h}_n \coloneqq n^{-1} \sum_{i=1}^n h(X_i)$ by generating a Markov chain $\{X_i\}$ that satisfies certain conditions for some distribution π and target function h (Flegal and Jones, 2010). However, the terminal sample size n^* for a reasonably precise \bar{h}_{n^*} is unknown a priori. In light of it, Jones et al. (2006) proposed the half-width test to terminate a simulation at

$$n^* = \inf\left\{n \in \mathbb{Z}^+ : z_{1-\alpha/2}\hat{\sigma}_n/\sqrt{n} + p(n) < \epsilon\right\},\tag{4.2}$$

where $\alpha \in (0, 1)$ is the significance level, $z_{1-\alpha/2}$ is the $100(1 - \alpha/2)\%$ lower quantile of N(0,1), p(n) is a penalty function for n that is too small, and $\epsilon > 0$ is the maximum tolerable error. Essentially, the half-width test is based on the central limit theorem (CLT) for \bar{h}_n and stops when the $100(1-\alpha)\%$ confidence interval (CI) is short enough; see also Glynn and Whitt (1992) and Vats et al. (2019). Since \bar{h}_n can be updated in O(1) space, it is sensible to construct the CI using an O(1)-space LRV estimator for data reduction, which we demonstrate with a classical example in Hastings (1970). To sample from N(0,1), Hastings (1970) used a Metropolis– Hastings algorithm with a random walk on $[-\delta, \delta]$ as the proposal. Given a MCMC sample $\{X_i\}_{i=1}^n$ generated with $\delta = 1$, we can estimate $\mathbb{E}_{\zeta \sim N(0,1)}(\zeta^2)$ by \bar{h}_n , where $h(x) = x^2$. Set $\alpha = 0.05$ and $p(n) = \epsilon \mathbb{1}_{n \leq 500}$, we conduct the half-width test for $\epsilon = 0.13, 0.12, \ldots, 0.04$ and 1000 replications using different LRV estimators ("R-packages"):

- (a) obm ("mcmcse"): the LRV is estimated by the mcse function (Flegal et al., 2021), which was rewritten in Rcpp (Eddelbuettel and François, 2011) in Google Summer of Code 2021.
 obm with lugsail parameter r=1 is used.
- (b) TSR ("rTACM"): the LRV is estimated by the rTACM function (Chan and Yau, 2017a). The best existing O(1)-space (and O(1)-time) estimator TSR is used.
- (c) LASER(1,2) ("rlaser"): the LRV is estimated by the lrv function with LASER(1,2).Arguments that are not mentioned, such as the smoothing parameter selector, are left as



Figure 4: Online 95%-half-width test using different LRV estimators: (a) obm (dotted gray); (b) TSR (dashed red); (c) LASER(1,2) (longdash blue).

default in all R-functions. Figure 4 shows that the coverage rates using LASER(1,2) are comparable to obm and always better than TSR, which is consistent with their optimal AMSE in Table 1. Moreover, LASER(1,2) is considerably faster than obm, and this speed improvement is not due to software implementation as the lrv function is written in R (R Core Team, 2021). Finally, we emphasize that LASER(1,2) requires a constant amount of memory only.

Remark 10. An alternative approach to online CI construction is self-normalization (SN) (Shao, 2010), which generally leads to a better coverage but wider width. However, this does not apply to Example 6 because SN is inconsistent for σ^2 . For a fixed ϵ in (4.2), using SN will result in a non-degenerate terminal sample size and so an incorrect coverage.

4.3 Mini-batch Estimation and Causal Inference

In practice, many problems are approached in a way similar to the MCMC convergence assessment described in Section 1.1 instead of in a completely online manner. For these problems, the LRV estimator can be updated every m iterations rather than every iteration. Nevertheless, none of the existing LRV estimators address this issue. In light of it, we propose mini-batch estimation, which consists of two algorithmic modifications:

(a) removing redundant operations: we take Algorithm 1 as an example. When LASER(1,1) is updated at time n_{j+1} from n_j , some intermediate statistics still need to be computed at all $n_j + 1, n_j + 2, ..., n_{j+1}$. For example, $K_{i,0}$ should be updated by

$$K_{i,0} = K_{i-1,0} + X_{i-1} - X_{i-s_i} \mathbb{1}_{s_i = s_{i-1}}$$
 for $i = n_j + 1, n_j + 2, \dots, n_{j+1}$

as they are needed to compute other statistics, e.g., $R_{n_{j+1},0}$. However, other statistics that are only related to the output can be computed at time n_{j+1} directly.

(b) applying vectorized operations: computation time can be further reduced through vectorization; see, e.g., Wickham (2019). While offline estimators can also be vectorized, online estimators have an edge that each update is computed from n_j to n_{j+1} only. In contrast, offline estimators may need to compute an update from 1 to n_{j+1} .

The next example demonstrates the practical value of mini-batch LRV estimators.

Example 7 (Online moment selection/Causal inference). To deal with multiple data sources in causal inference, Gupta et al. (2021) proposed online moment selection (OMS), a strategy that applies the generalized method of moments (GMM) (Hansen, 1982) to estimate the average treatment effect (ATE) and decide which data sources to query. Their best policy, OMS via explore-then-greedy (OMS-ETG), updates the data source selection ratio in batches based on the asymptotic variance estimate. However, they have only considered the sample variance estimator. Motivated by Hansen (1982), we compare different variance estimators in the presence of serial dependence here. Consider

$$X = aZ + \eta, \qquad Y = bX + \xi, \qquad \eta \downarrow \xi, \qquad \eta \perp Z, \qquad \xi \perp Z, \tag{4.3}$$

where Z is the instrument, X is the treatment, Y is the outcome, b is the target ATE, $\{\{Z, X\}, \{Z, Y\}\}\$ are data sources, and \perp denotes independence. Everything is same as the instrumental variable example in Gupta et al. (2021) except that $\{Z_i, X_i, Y_i\}\$ are not IID; see Supplement F.4 for detail. Then, we run an OMS-ETG algorithm that collects 20% of the sample per round for 5 rounds and $n = 1000, 1500, \ldots, 5000$ with different variance estimators:

- (a) sample: the sample variance estimator, whose univariate online version is WFD, is used.
- (b) PSR: PSR with automatic and adaptive update in Chan and Yau (2017a) is used.
- (c) LASER(1,1): mini-batch LASER(1,1) with automatic update and m = [0.02n] is used; see also Remark 4.
- (d) oracle: the data source selection ratio is pre-estimated based on 10^6 observations using GMM with bart.

Based on 1000 replications, we compute the ATE MSEs and time costs relative to oracle. Figure 5 shows that LASER(1,1) brings noticeable statistical improvement to OMS-ETG with negligible time cost. Meanwhile, PSR gives similar improvement with a much higher time cost. In line with Hansen (1982), sample leads to a consistent but not the most efficient ATE estimate



Figure 5: ATE estimation using OMS-ETG with different variance estimators: (a) sample (dotted gray); (b) PSR (dashed red); (c) LASER(1,1) (longdash blue).

under dependence.

4.4 Stochastic Approximation

Stochastic approximation (Robbins and Monro, 1951) is a popular class of optimization algorithms, particularly in machine learning (Goodfellow et al., 2016). Commonly, users consider the optimization problem $\min_{w \in \mathbb{R}^d} \mathbb{E}_{\zeta \sim \Pi} \{h(w, \zeta)\}$, where ζ denotes the data sampled from some (unknown) distribution Π , w represents the model parameters, and $h(w, \zeta)$ is a loss function (Lang et al., 2019). In the vanilla stochastic gradient descent (SGD), the model parameters are updated by $w_i = w_{i-1} - \eta_i g_{i-1}$, where w_0 is the initial point, η_i is the learning rate at the *i*-th iteration, and g_i denotes the gradient of $h(w, \zeta_i)$ with respect to w at $w = w_i$ (Zhu et al., 2021). Since SGD is an O(1)-time, O(1)-space and anytime algorithm, it is widely used in real-time applications.

Despite its popularity and advantages, SGD still suffers from the convergence problem of many online techniques: a reasonable terminal sample size is not known *a priori*. In Examples 8 and 9, we present two different views to this problem recently developed by the statistics and engineering communities, respectively. Coincidentally, they both utilize LRV estimators to perform some kind of real-time inference, where our framework fits in seamlessly.

Example 8 (Convergence of SGD: a statistical view). In light of the convergence problem, Zhu et al. (2021) proposed quantifying the uncertainty of the averaged SGD (ASGD), whose limit theorems had been established (Polyak and Juditsky, 1992), in a fully online fashion; see also Remark 1. Building on \triangle SR in Wu (2009), they provide an alternative non-overlapping version and derive both versions' rate-optimal batch size under the non-stationarity in SGD. To be specific, suppose the learning rate is chosen to be $\eta_i = \eta_0 i^{-\alpha}$ for some $\eta_0 > 0$ and $\alpha \in (1/2, 1)$. Then, the starting point of the k-th block in \triangle SR should be $a_k = \lfloor Ak^{2/(1-\alpha)} \rfloor$ for some A > 0. In several Monte Carlo experiments, Zhu et al. (2021) showed that the performance is relatively insensitive to the choice of A.

On the other hand, it is certain that TSR and PSR dominate \triangle SR under stationarity (Chan and Yau, 2017a). Using our framework, Example 2 further reveals the structural problem of these selection rules. Wondering whether similar results hold under non-stationarity, we consider a logistic regression example. Let $\{\zeta_i = (x_i, y_i)\}_{i \in \mathbb{Z}^+}$ and w^* denote an IID sequence of data and the true model parameter, where $x_i \sim N(0, \mathbf{I}_d)$, $y_i \mid x_i \sim \text{Bernoulli}(\{1 + \exp(-x_i^{\mathsf{T}}w^*)\}^{-1})$, w^* is a *d*-dimensional vector linearly spaced between -1 and 1, and d = 10. The loss function is $h(w, x_i, y_i) = (1 - y_i)x_i^{\mathsf{T}}w + \ln\{1 + \exp(-x_i^{\mathsf{T}}w^*)\}$. Following experiments in Zhu et al. (2021), the learning rate is chosen as $\eta_i = 0.5i^{-\alpha}$ with $\alpha = 0.505$, and the starting point of the k-th block is set to be $a_k = \lfloor Ak^{2/(1-\alpha)} \rfloor$ with A = 1 for:

- (a) \triangle SR ("rTACM"): TSR with $\xi = 0$ in Chan and Yau (2017a) is used, which is essentially the overlapping \triangle SR in Zhu et al. (2021) as a_k is set to be the same. Automatic update does not apply because of non-stationarity.
- (b) online LASER ("rlaser"): "LASER" with q = 1 and m = 1 is used. While (2.8) does not apply, blocking with a_k is compatible with the form in (2.3). In the k-th block, the subsampling parameter increases by 1 per iteration until it reaches the block maximum $[2A(1-\alpha)^{-1}k^{(1+\alpha)/(1-\alpha)}]$, where it remains unchanged until the next block, and the tapering parameter is chosen to be the block maximum scaled according to Corollary 2.
- (c) mini-batch LASER ("rlaser"): the LRV is estimated as in (b) except that $m = 10^4$.
- (d) TSR ("rTACM"): TSR with $\xi = 1$ is used, which is proven to be more efficient than \triangle SR under stationarity.

We compare the above in constructing 95% CIs for the summed coefficient $\mathbf{1}^{\mathsf{T}}w^*$ with a burn-in stage (Chen et al., 2020) of size 500 based on 1000 replications. Figure 6 shows that the performance of \triangle SR is consistent with the empirical findings in Zhu et al. (2021). Furthermore, there are notable improvements according to the theory under stationarity. Real-time inference is enabled by mini-batch estimation without loss of statistical efficiency. These indicate that our framework can be used as a novel starting point for research in the same area. Finally, we



Figure 6: CI construction for the summed ASGD estimates using different LRV estimators: (a) \triangle SR (dotted gray); (b) online LASER (dashed red); (c) mini-batch LASER (longdash blue); (d) TSR (dotdash purple).

remark that CIs constructed with LRV estimators can be used as convergence diagnostics; see Example 6 and Zhu et al. (2021).

Example 9 (Convergence of SGD: an engineering view). Lang et al. (2019) took a different approach to the convergence problem of SGD. As the vanilla SGD or its popular variants typically require problem-specific hyperparameter hand-tuning to achieve good performance (Wilson et al., 2017), Lang et al. (2019) proposed the statistical adaptive stochastic approximation (SASA) method to automatically find good learning rate schedules for different problems. Fundamentally, SASA tries to identify whether training progress has stalled by testing stationarity of $\{w_j\}$. It applies not only to the vanilla SGD but also a wider class of stochastic approximation method; see Zhang et al. (2020).

Here, we shall focus on the vanilla case as it is studied in Example 8. Let $d_j = w_{j-1}^{\mathsf{T}} g_{j-1} - 0.5\eta_j g_{j-1}^{\mathsf{T}} g_{j-1}$. If $\{w_j\}$ starts under non-stationarity and converges to a stationary state, we have $\lim_{j\to\infty} \mathbb{E}(d_j) = 0$ as long as g_j is generated with some time-homogeneous dynamics (Zhang et al., 2020), which is satisfied by SGD. Based on the Markov chain CLT in Jones et al. (2006), SASA tests this condition and decreases the learning rate geometrically if stationarity is not rejected. For more details, we refer readers to Zhang et al. (2020) as they extend SASA to SASA+ using the same idea. Now, we revisit the logistic regression in Example 8 with the same setting but different learning rate schedules:

- (a) Vanilla SGD: the learning rate $\eta_i = 0.5i^{-\alpha}$ with $\alpha = 0.505$ in Zhu et al. (2021) is used.
- (b) SASA+: the learning rate is chosen by SASA+ in Zhang et al. (2020); see Supplement F.5



Figure 7: Learning rate tuning in SGD: (a) Vanilla SGD in Zhu et al. (2021) (dotted gray); (b) SASA+ in Zhang et al. (2020) with mini-batch LASER(1,1) (longdash blue).

for the complete algorithm, where we change the offline obm used in the original SASA+ to mini-batch LASER(1,1).

Figure 7 shows that SASA+ can drastically improve the statistical efficiency. This is not surprising because of two reasons. First, SASA(+) is inspired by the theoretical and empirical success of geometrically decaying learning rate in some (strongly convex) scenarios; see, e.g., Ge et al. (2019). Second, SASA(+) is shown to match with the best hand-tuned learning rate schedules on several deep learning tasks (Lang et al., 2019; Zhang et al., 2020). While SASA(+) needs additional time to perform inference, the cost is small with mini-batch LASER(1,1).

5 Discussion

Ljung and Söderström (1983) mentioned that "many people regard the area (of recursive identification/estimation) to be a 'bag of tricks' rather than a theory". Indeed, existing online estimators are often found by sacrificing some statistical efficiency and studied on an algorithmby-algorithm basis. Contrary to this approach, we presented a novel framework to understand and construct efficient nonparametric estimators for online problems. Our framework consisted of a flexible window decomposition, a principle-driven theoretical approach, and a common practical acceleration. To the best of our knowledge, this is the first attempt to characterize the form of online (LRV) estimators through sufficient conditions. The study of necessary conditions would be a theoretically important future direction.

Using LRV as an exemplary estimand, we demonstrated how to construct an estimator that

could be super or nearly optimal in the sense of Chan and Yau (2017a). It is impressive to see that requiring O(1)-time update does not harm the statistical efficiency, while further requiring O(1)-space update has practically negligible cost. The constructed estimator LASER (q, ϕ) is also general as it was shown to be compatible with difference statistics (Chan, 2022), automatic update (Chan and Yau, 2017a), and blocking for SGD (Zhu et al., 2021). Other innovations that may be added, which are all potential directions, include but are not limited to prewhitening (Andrews and Monahan, 1992), weighted batch means (Liu and Flegal, 2018), and lugsail lag window (Vats and Flegal, 2021).

Apart from LRV estimation, our framework can be applied to a considerable number of tasks. One example is the estimation of v_q discussed in Section 3.3. Other examples include spectral density estimation (Xiao and Wu, 2011), nonparametric regression (Huang et al., 2014), and asymptotic variance estimation for other estimators such as sample quantile (Doss et al., 2014). This is because our window decomposition generalizes the existing kernel and subsampling methods in both offline and online settings. Any estimator built upon these methods can be reconstructed and likely improved under our framework as in Example 2. For example, a classical kernel density estimator takes the form

$$\hat{f}_{n,1}(x) = \frac{1}{n\ell_n} \sum_{i=1}^n W_n(x, X_i) = \frac{1}{n\ell_n} \sum_{i=1}^n T\left(\frac{|x - X_i|}{\ell_n}\right) \mathbb{1}_{|x - X_i| \le \ell_n}.$$
(5.1)

Huang et al. (2014) noticed that $\hat{f}_{n,1}(x)$ can only be updated in O(n) time when $\ell_n \neq \ell_{n+1}$ so they proposed

$$\hat{f}_{n,2}(x) = \frac{1}{\sum_{i=1}^{n} \ell_i} \sum_{i=1}^{n} T\left(\frac{|x - X_i|}{\ell_i}\right) \mathbb{1}_{|x - X_i| \le \ell_i},\tag{5.2}$$

which can be updated in O(1) time. Nevertheless, our Principle A suggests that

$$\hat{f}_{n,3}(x) = \frac{1}{nt_n} \sum_{i=1}^n T\left(\frac{|x - X_i|}{t_n}\right) \mathbb{1}_{|x - X_i| \le s_i}$$
(5.3)

is likely more efficient but still O(1)-time if T(x) is a polynomial as in Proposition 1. Note that users can also propose new principles on top of our framework. For instance, one may investigate variable tapers in Terrell and Scott (1992):

$$\hat{f}_{n,4}(x) = \frac{1}{n} \sum_{i=1}^{n} \frac{1}{t_n(x, X_i)} T\left(\frac{|x - X_i|}{t_n(x, X_i)}\right) \mathbb{1}_{|x - X_i| \le s_i},\tag{5.4}$$

where the tapering parameter may depend on the point of estimation x or observation X_i . This is something that the existing frameworks, e.g., subsample selection rule, cannot come up with. Last but not least, we selected several interesting applications from different communities. They included change point detection, MCMC convergence diagnosis, causal inference, and stochastic approximation. One useful message is that offline problems can be approached and sometimes better solved with online techniques. Another interesting observation is that different communities may develop different attitudes to inference. We hope this paper broadens the audience of online techniques and serves as a bridge between different communities regarding inference.

A Algorithms

A.1 Overview of Notations

In this subsection, we state and explain the notations that we are going to use throughout Section A. Most of them will be recalled later when they are used. Note that these notations only apply to Section A in general. We begin with some simple components:

$$\begin{split} K_{n,b} &\coloneqq \sum_{k=1}^{s_n} k^b X_{n-k}, & k_{n,b} \coloneqq \sum_{k=1}^{s_n} k^b, \\ R_{n,b} &\coloneqq \sum_{i=1}^n \sum_{k=1}^{s_i} k^b X_i X_{i-k}, & r_{n,b} \coloneqq \sum_{i=1}^n \sum_{k=1}^{s_i} k^b, \\ U_{n,b} &\coloneqq \sum_{i=1}^n \sum_{k=1}^{s_i} k^b X_i, & V_{n,b} \coloneqq \sum_{i=1}^n \sum_{k=1}^{s_i} k^b X_{i-k} \\ Q_n &\coloneqq \sum_{i=1}^n X_i^2, & \bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i, \end{split}$$

where $b \in \mathbb{N}_0 = \{0, 1, \ldots\}$. Note the first subscript *n* indicates the sample size and the second subscript *b* indicates the exponent in the summation, which is usually related to the characteristic exponent *q*. We capitalize the variables when they depends on the observations X_1, \ldots, X_n . To lighten notations in the algorithms, we define the shorthands

$$R_n^{(a,b,c)} \coloneqq R_{n,a} - \frac{R_{n,b}}{t_n^c}, \quad R_n^{(a,b)} = R_n^{(a,b,b)} = R_{n,a} - \frac{R_{n,b}}{t_n^b},$$

and similarly for $r_n^{(a,b,c)}, U_n^{(a,b,c)}, V_n^{(a,b,c)}$, e.g., $U_n^{(a,b,c)} = U_{n,a} - U_{n,b}/t_n^c$. After we introduce the characteristic exponent q, we need the backward finite difference operator $\nabla^{(b)}$, which is defined by

$$\nabla^{(1)}f(k) \coloneqq f(k) - f(k-1)$$
 and $\nabla^{(b)}f(k) \coloneqq \nabla^{(b-1)}f(k) - \nabla^{(b-1)}f(k-1),$

where $b \in \mathbb{Z}^+$ and f is a function that takes an integer input k. Then, we have

$$d_{k,q}^{(b)} = \nabla^{(b)} k^q$$
 and $D_{n,q}^{(b)} \coloneqq \sum_{k=1}^{s_n} d_{k,q}^{(b)} X_{n-k}$,

where b = 1, 2, ..., q. Note that the superscript (b) indicates the number of backward finite differences, which is used in $d_{k,q}^{(b)}$ and $D_{k,q}^{(b)}$ only. When the memory parameter $\phi > 1$, the intended subsampling parameter s_i is ramped up as s'_i defined in (2.8). The components become

$$\begin{split} K'_{n,b} &\coloneqq \sum_{k=1}^{s'_n} k^b X_{n-k}, & k'_{n,b} &\coloneqq \sum_{k=1}^{s'_n} k^b, \\ R'_{n,b} &\coloneqq \sum_{i=1}^n \sum_{k=1}^{s'_i} k^b X_i X_{i-k}, & r'_{n,b} &\coloneqq \sum_{i=1}^n \sum_{k=1}^{s'_i} k^b, \\ U'_{n,b} &\coloneqq \sum_{i=1}^n \sum_{k=1}^{s'_i} k^b X_i, & V'_{n,b} &\coloneqq \sum_{i=1}^n \sum_{k=1}^{s'_i} k^b X_{i-k}, \end{split}$$

where $b \in \mathbb{N}_0$. Since the prime symbol is used in the ramped subsampling parameter s'_i , we reserve it for the corresponding components. The shorthands become

$$R_n^{\prime(a,b,c)} \coloneqq R_{n,a}^{\prime} - \frac{R_{n,b}^{\prime}}{t_n^c}, \quad R_n^{\prime(a,b)} = R_n^{\prime(a,b,b)} = R_{n,a}^{\prime} - \frac{R_{n,b}^{\prime}}{t_n^b},$$

and similarly for $r'^{(a,b,c)}_{n}, U'^{(a,b,c)}_{n}, V'^{(a,b,c)}_{n}$, e.g., $U'^{(a,b,c)}_{n} = U'_{n,a} - U'_{n,b}/t^{c}_{n}$. To perform O(1)-space update when $\phi \ge 2$, we need some "pre-calculated" components (denoted by the double prime symbol):

$$a_{n} \coloneqq [\phi s_{n}],$$

$$K_{n,b}'' \coloneqq \begin{cases} K_{n-1,b}'' + (a_{n-1} - s_{n}')^{b} X_{n}, & s_{n}' \ge a_{n-1} - s_{n-1} \text{ and } s_{n}' > 0; \\ 0, & s_{n}' < a_{n-1} - s_{n-1} \text{ or } s_{n}' = 0, \end{cases}$$

$$k_{n,b}'' \coloneqq \begin{cases} k_{n-1,b}'' + (a_{n-1} - s_{n}')^{b}, & s_{n}' \ge a_{n-1} - s_{n-1} \text{ and } s_{n}' > 0; \\ 0, & s_{n}' < a_{n-1} - s_{n-1} \text{ or } s_{n}' = 0. \end{cases}$$

In a mini-batch setting, we are interested in updating the components at $n_0 = 1 < n_1 < n_2 < \cdots$ instead of at $n = 1, 2, \ldots$ Therefore, we use the notations

$$\begin{split} R_{n_{j},b} &= \sum_{i=1}^{n_{j}} \sum_{k=1}^{s_{i}} k^{b} X_{i} X_{i-k}, \quad r_{n_{j},b} = \sum_{i=1}^{n_{j}} \sum_{k=1}^{s_{i}} k^{b}, \\ U_{n_{j},b} &= \sum_{i=1}^{n_{j}} \sum_{k=1}^{s_{i}} k^{b} X_{i}, \qquad V_{n_{j},b} = \sum_{i=1}^{n_{j}} \sum_{k=1}^{s_{i}} k^{b} X_{i-k}, \\ Q_{n_{j}} &= \sum_{i=1}^{n_{j}} X_{i}^{2}, \qquad \bar{X}_{n_{j}} = \frac{1}{n_{j}} \sum_{i=1}^{n_{j}} X_{i}, \end{split}$$

where $b, j \in \mathbb{N}_0$. For $K_{n,b}$ and $k_{n,b}$, they need to be computed at $n_{j-1} + 1, n_{j-1} + 2, \dots, n_j$ and so we introduce the vector notations

$$\vec{K}_{n_j,b} \coloneqq (K_{n_{j-1}+1,b}, \dots, K_{n_j,b})^{\mathsf{T}} \text{ and } \vec{k}_{n_j,b} \coloneqq (k_{n_{j-1}+1,b}, \dots, k_{n_j,b})^{\mathsf{T}}.$$

A.2 LASER(1,1): Known Zero-mean

Since the constructed estimator $LASER(q, \phi)$ is based on a novel framework, we shall show the derivation of the algorithm of LASER(1, 1) when $\mu = 0$ is known for illustrative purpose first. The other algorithms can be derived using a similar procedure and will be discussed later. To begin with, note that the estimator can be written as

$$\bar{\sigma}_{n,\mathsf{LASER}(1,1)}^{2} = \frac{1}{n} \sum_{i=1}^{n} X_{i}^{2} + \frac{2}{n} \sum_{i=1}^{n} \sum_{k=1}^{s_{i}} \left(1 - \frac{k}{t_{n}}\right) X_{i} X_{i-k}$$
$$= \frac{1}{n} \left(Q_{n} + 2R_{n,0} - \frac{2}{t_{n}} R_{n,1}\right),$$

where

$$Q_n = \sum_{i=1}^n X_i^2, \quad R_{n,0} = \sum_{i=1}^n \sum_{k=1}^{s_i} X_i X_{i-k} \quad \text{and} \quad R_{n,1} = \sum_{i=1}^n \sum_{k=1}^{s_i} k X_i X_{i-k}.$$

The recursive formula for Q_n is trivial so we shall focus on $R_{n,0}$ and $R_{n,1}$. To lighten notations, write the subsampling parameter at the last iteration as $s = s_{n-1}$. For $R_{n,0}$, we have

$$R_{n,0} = \sum_{i=1}^{n-1} \sum_{k=1}^{s_i} X_i X_{i-k} + X_n \sum_{k=1}^{s_n} X_{n-k}$$
$$= R_{n-1,0} + X_n K_{n,0},$$

where $K_{n,0} = \sum_{k=1}^{s_n} X_{n-k}$. Note that $K_{n,0}$ can be updated easily since

$$K_{n,0} = \begin{cases} 0, & s_n = s = 0; \\ K_{n-1,0} + X_{n-1} - X_{n-s-1}, & s_n = s > 0; \\ K_{n-1,0} + X_{n-1}, & s_n = s + 1. \end{cases}$$

Similarly, for $R_{n,1}$, observe that

$$R_{n,1} = \sum_{i=1}^{n-1} \sum_{k=1}^{s_i} k X_i X_{i-k} + X_n \sum_{k=1}^{s_n} k X_{n-k}$$
$$= R_{n-1,1} + X_n K_{n,1},$$

where $K_{n,1} = \sum_{k=1}^{s_n} k X_{n-k}$. We can update $K_{n,1}$ recursively by

$$K_{n,1} = \begin{cases} 0, & s_n = s = 0; \\ K_{n-1,1} + K_{n,0} - sX_{n-s-1}, & s_n = s > 0; \\ K_{n-1,1} + K_{n,0}, & s_n = s + 1. \end{cases}$$

We summarize the recursive formulas for different components in the order of their updates:

Proposition A.1. Suppose the subsampling parameter $\{s_i \in \mathbb{N}_0\}_{i \in \mathbb{Z}^+}$ is a monotonically increasing sequence such that $\sup_{i \in \mathbb{Z}^+} |s_{i+1} - s_i| \leq 1$. Write its value at the last iteration as $s = s_{n-1}$. Then, the following components can be updated in O(1) time:

$$K_{n,0} = \sum_{k=1}^{s_n} X_{n-k} = \begin{cases} 0, & s_n = s = 0; \\ K_{n-1,0} + X_{n-1} - X_{n-s-1}, & s_n = s > 0; \\ K_{n-1,0} + X_{n-1}, & s_n = s + 1, \end{cases}$$

$$K_{n,1} = \sum_{k=1}^{s_n} k X_{n-k} = \begin{cases} 0, & s_n = s = 0; \\ K_{n-1,1} + K_{n,0} - s X_{n-s-1}, & s_n = s > 0; \\ K_{n-1,1} + K_{n,0}, & s_n = s + 1, \end{cases}$$

$$R_{n,b} = \sum_{i=1}^{n} \sum_{k=1}^{s_i} k^b X_i X_{i-k} = R_{n-1,b} + X_n K_{n,b} \quad for \quad b = 0, 1, \end{cases}$$

$$Q_n = \sum_{i=1}^{n} X_i^2 = Q_{n-1} + X_n^2.$$

When the data arrive sequentially, we need to maintain a vector \vec{x} in order to store the last s_n observations to update the moving sum $K_{n,0}$. This can be implemented efficiently using a queue data structure as it is first-in-first-out. Hence the recursive algorithm for $\bar{\sigma}_{n,\text{LASER}(1,1)}^2$ is given by Algorithm A.1.

A.3 LASER(1,1): Unknown General Mean

When $\mu \in \mathbb{R}$ is unknown, we can subtract the sample mean $\overline{X}_n = n^{-1} \sum_{i=1}^n X_i$ from each observation. Therefore, the estimator can be written as

$$\hat{\sigma}_{n,\mathsf{LASER}(1,1)}^{2} = \frac{1}{n} \left\{ \sum_{i=1}^{n} (X_{i} - \bar{X}_{n})^{2} + 2 \sum_{i=1}^{n} \sum_{k=1}^{s_{i}} \left(1 - \frac{k}{t_{n}} \right) (X_{i} - \bar{X}_{n}) (X_{i-k} - \bar{X}_{n}) \right\}$$
$$= \frac{1}{n} \left\{ \sum_{i=1}^{n} X_{i}^{2} - n\bar{X}_{n}^{2} + 2 \sum_{i=1}^{n} \sum_{k=1}^{s_{i}} \left(1 - \frac{k}{t_{n}} \right) (X_{i}X_{i-k} - X_{i}\bar{X}_{n} - X_{i-k}\bar{X}_{n} + \bar{X}_{n}^{2}) \right\}$$
$$= \frac{1}{n} \left[Q_{n} - n\bar{X}_{n}^{2} + 2R_{n,0} - \frac{2}{t_{n}}R_{n,1} - 2\bar{X}_{n} \left\{ \sum_{i=1}^{n} \sum_{k=1}^{s_{i}} \left(1 - \frac{k}{t_{n}} \right) (X_{i} + X_{i-k} - \bar{X}_{n}) \right\} \right]$$

Algorithm A.1: LASER(1, 1), known zero-mean

[1] initalization: [2] Set $n = 1, s = 0, Q_n = X_1^2, K_{n,0} = K_{n,1} = R_{n,0} = R_{n,1} = 0$ [3] Push X_1 into \vec{x} begin [4] Receive X_{n+1} [5] Set n = n + 1[6] Compute s_n, t_n [7] Retrieve X_{n-1}, X_{n-s-1} from \vec{x} /* last and first elements in $ec{x}$ */ [8] if $\underline{s_n = s}$ then [9] Update $K_{n,0}, K_{n,1}$ with Prop. A.1 [10] Pop X_{n-s-1} from \vec{x} [11] else [12] Update $K_{n,0}, K_{n,1}$ with Prop. A.1 [13] Push X_n into \vec{x} [14] Update $R_{n,0}, R_{n,1}, Q_n$ with Prop. A.1 [15] [16] Set $s = s_n$ Output $\bar{\sigma}_{n,\text{LASER}(1,1)}^2 = (Q_n + 2R_{n,0} - 2R_{n,1}/t_n)/n$ [17]

$$= \frac{1}{n} \left\{ Q_n + 2R_{n,0} - \frac{2}{t_n} R_{n,1} + \left(2r_{n,0} - \frac{2}{t_n} r_{n,1} - n \right) \bar{X}_n^2 - 2\bar{X}_n \left(U_{n,0} - \frac{1}{t_n} U_{n,1} + V_{n,0} - \frac{1}{t_n} V_{n,1} \right) \right\},$$

where the additional components are

$$r_{n,b} = \sum_{i=1}^{n} \sum_{k=1}^{s_i} k^b, \quad U_{n,b} = \sum_{i=1}^{n} \sum_{k=1}^{s_i} k^b X_i \quad \text{and} \quad V_{n,b} = \sum_{i=1}^{n} \sum_{k=1}^{s_i} k^b X_{i-k} \quad \text{for} \quad b = 0, 1.$$

To lighten notations, denote $R_n^{(a,b)} = R_{n,a} - R_{n,b}/t_n^b$ and similarly for $r_n^{(a,b)}, U_n^{(a,b)}, V_n^{(a,b)}$, e.g., $U_n^{(a,b)} = U_{n,a} - U_{n,b}/t_n^b$. Then, the estimator can be expressed as

$$\hat{\sigma}_{n,\mathsf{LASER}(1,1)}^2 = \frac{1}{n} \left\{ Q_n + 2R_n^{(0,1)} + (2r_n^{(0,1)} - n)\bar{X}_n^2 - 2\bar{X}_n (U_n^{(0,1)} + V_n^{(0,1)}) \right\}$$

Using the same procedure as in Section A.2, we can derive the recursive formulas for the additional components:

Proposition A.2. Suppose the subsampling parameter $\{s_i \in \mathbb{N}_0\}_{i \in \mathbb{Z}^+}$ is a monotonically increasing sequence such that $\sup_{i \in \mathbb{Z}^+} |s_{i+1} - s_i| \leq 1$. Write its value at the last iteration as $s = s_{n-1}$. For each b = 0, 1, the following components can be updated in O(1) time:

$$k_{n,b} = \sum_{k=1}^{s_n} k^b = \begin{cases} 0, & s_n = s = 0; \\ k_{n-1,b}, & s_n = s > 0; \\ k_{n-1,b} + s_n^b, & s_n = s + 1, \end{cases}$$
$$r_{n,b} = \sum_{i=1}^n \sum_{k=1}^{s_i} k^b = r_{n-1,b} + k_{n,b},$$

Algorithm A.2: LASER(1,1), unknown general mean

[1] initalization: [2] Set $n = 1, s = 0, Q_n = X_1^2, \bar{X}_n = X_1$ [3] Set $K_{n,b} = R_{n,b} = k_{n,b} = r_{n,b} = U_{n,b} = V_{n,b} = 0$ for b = 0, 1[4] Push X_1 into \vec{x} begin [5] Receive X_{n+1} [6] Set n = n + 1[7] Compute s_n, t_n [8] /* last and first elements in $ec{x}$ */ Retrieve X_{n-1}, X_{n-s-1} from \vec{x} [9] if $\underline{s_n = s}$ then [10] Update $K_{n,b}$ with Prop. A.1 for b = 0, 1[11] Pop X_{n-s-1} from \vec{x} [12] [13] else Update $K_{n,b}$ with Prop. A.1 and $k_{n,b}$ with Prop. A.2 for b = 0, 1[14] Push X_n into \vec{x} [15] Update $R_{n,b}, Q_n$ with Prop. A.1 and $r_{n,b}, U_{n,b}, V_{n,b}, \bar{X}_n$ with Prop. A.2 for b = 0, 1[16] Set $s = s_n$ [17] Output $\hat{\sigma}_{n,\text{LASER}(1,1)}^2 = \{Q_n + 2R_n^{(0,1)} + (2r_n^{(0,1)} - n)\bar{X}_n^2 - 2\bar{X}_n(U_n^{(0,1)} + V_n^{(0,1)})\}/n$ [18]

$$\begin{split} U_{n,b} &= \sum_{i=1}^{n} \sum_{k=1}^{s_i} k^b X_i = U_{n-1,b} + k_{n,b} X_n, \\ V_{n,b} &= \sum_{i=1}^{n} \sum_{k=1}^{s_i} k^b X_{i-k} = V_{n-1,b} + K_{n,b}, \\ \bar{X}_n &= \frac{1}{n} \sum_{i=1}^{n} X_i = \frac{(n-1)\bar{X}_{n-1} + X_n}{n}. \end{split}$$

We notice that it may not be necessary to maintain the components separately, e.g., $U_{n,0}$ and $U_{n,1}$. Instead, we can maintain the components like $\mathscr{U}_n = \sum_{i=1}^n \sum_{k=1}^{s_i} (1 - k/t_n) X_i$ directly. However, it may violate Principle E. Consequently, the recursive formulas for these components will depend on the tapering parameter t_n and cannot be generalized to the settings in Sections A.4 or A.6 without restricting the changes in t_n . Therefore, it is better to maintain the components separately with an exterior t_n . The recursive algorithm for $\hat{\sigma}_{n,\text{LASER}(1,1)}^2$ is given by Algorithm A.2.

A.4 LASER(q, 1): Characteristic Exponent

In this subsection, we illustrate how to derive recursive formulas for $\hat{\sigma}_{n,\mathsf{LASER}(q,1)}^2$. We are going to focus on the q-th order term as similar formulas apply to lower order terms except that a separate set of components is needed for each term.

To begin with, we follow the same procedure as in Sections A.2 and A.3 to write the estimator

as

$$\hat{\sigma}_{n,\mathsf{LASER}(q,1)}^{2} = \frac{1}{n} \sum_{i=1}^{n} (X_{i} - \bar{X}_{n})^{2} + \frac{2}{n} \sum_{i=1}^{n} \sum_{k=1}^{s_{i}} \left(1 - \frac{k^{q}}{t_{n}^{q}} \right) (X_{i} - \bar{X}_{n}) (X_{i-k} - \bar{X}_{n})$$
$$= \frac{1}{n} \left\{ Q_{n} + 2R_{n}^{(0,q)} + (2r_{n}^{(0,q)} - n)\bar{X}_{n}^{2} - 2\bar{X}_{n} (U_{n}^{(0,q)} + V_{n}^{(0,q)}) \right\},$$

where the shorthands, e.g., $R_n^{(0,q)} = R_{n,0} - R_{n,q}/t_n^q$, are introduced in Section A.3; see Section A.1 for an overview of notations. Now, we need to work on a recursive formula for $K_{n,q} = \sum_{k=1}^{s_n} k^q X_{n-k}$ in order to update $R_{n,q}$. To this end, we can use the backward finite difference to be discussed in Section D.1. Recall the backward finite difference operator $\nabla^{(b)}$ is defined by

$$\nabla^{(1)}f(k) = f(k) - f(k-1)$$
 and $\nabla^{(b)}f(k) = \nabla^{(b-1)}f(k) - \nabla^{(b-1)}f(k-1)$,

where $b \in \mathbb{Z}^+$ and f is a function that takes an integer input k. When $f(k) = k^q$, we have the recursive relation

$$\nabla^{(1)}k^q = k^q - (k-1)^q$$
 and $\nabla^{(b)}k^q = \nabla^{(b-1)}k^q - \nabla^{(b-1)}(k-1)^q$.

To lighten notations, denote $d_{k,q}^{(b)} = \nabla^{(b)} k^q$. From Section D.1, we know that $d_{k,q}^{(q)}$ is a non-zero constant and well-defined when k > q. Therefore, we can expand the recursive relation and put k = q + 1 to obtain

$$d_{q+1,q}^{(q)} = (q+1)^q - {\binom{q}{1}}q^q + {\binom{q}{2}}(q-1)^q - \dots + (-1)^q 1^q$$
$$= \sum_{r=0}^q (-1)^r {\binom{q}{r}}(q-r+1)^q.$$

Similarly, the initial values $d_{b+1,q}^{(b)}$ for b = 1, 2, ..., q-1 are given by

$$d_{b+1,q}^{(b)} = \sum_{r=0}^{b} (-1)^r {\binom{b}{r}} (b-r+1)^q.$$

It remains to find $d_{k,q}^{(b)}$ when k > b+1 for b = 1, 2, ..., q-1. By rearranging $\nabla^{(b+1)}k^q = \nabla^{(b)}k^q - \nabla^{(b)}(k-1)^q$, we have

$$d_{k,q}^{(b)} = \nabla^{(b)} k^q = \nabla^{(b)} (k-1)^q + \nabla^{(b+1)} k^q$$
$$= d_{k-1,q}^{(b)} + d_{k,q}^{(b+1)}.$$

When k increases, we can update $d_{k,q}^{(b)}$ recursively from b = q - 1 to b = 1 since $d_{k-1,q}^{(b)}$ comes from the last iteration and $d_{k,q}^{(b+1)}$ is a constant or has been updated. We summarize the recursive formulas for $d_{k,q}^{(b)}$ below:

$$d_{k,q}^{(b)} = \begin{cases} 0, & k < b+1 \text{ and } b = 1, 2, \dots, q; \\ \sum_{r=0}^{b} (-1)^r {b \choose r} (b-r+1)^q, & k = b+1 \text{ and } b = 1, 2, \dots, q; \\ d_{q+1,q}^{(q)}, & k > b+1 \text{ and } b = q; \\ d_{k-1,q}^{(b)} + d_{k,q}^{(b+1)}, & k > b+1 \text{ and } b = 1, 2, \dots, q-1. \end{cases}$$

We notice that it is also possible to compute $d_{k,q}^{(b)}$ directly using $d_{k,q}^{(b)} = \sum_{r=0}^{b} (-1)^r {b \choose r} (k-r)^q$; see, e.g., Zia (1991). Nonetheless, this approach requires more arithmetic operations when k changes. Therefore, it is more desirable to use the recursive formulas for $d_{k,q}^{(b)}$. Analogous to Proposition A.1, the recursive formulas for the zero-mean components are summarized below.

Proposition A.3. Suppose the subsampling parameter $\{s_i \in \mathbb{N}_0\}_{i \in \mathbb{Z}^+}$ is a monotonically increasing sequence such that $\sup_{i \in \mathbb{Z}^+} |s_{i+1} - s_i| \leq 1$. Write its value at the last iteration as $s = s_{n-1}$. Then, the following components can be updated in O(1) time:

$$\begin{split} c_q^{(b)} &= \sum_{r=0}^{b} (-1)^r \binom{b}{r} (b-r+1)^q \quad for \quad b=1,2,\ldots,q, \\ d_{s_n,q}^{(b)} &= \begin{cases} 0, & s_n < b+1 \; and \; b=1,2,\ldots,q; \\ c_q^{(b)}, & s_n = b+1 \; and \; b=1,2,\ldots,q; \\ d_{s,q}^{(b)}, & s_n = s > b+1 \; and \; b=1,2,\ldots,q; \\ d_{s,q}^{(b)}, & s_n = s+1 > b+1 \; and \; b=q; \\ d_{s,q}^{(b)}, & s_n = s+1 > b+1 \; and \; b=1,2,\ldots,q-1, \end{cases} \\ D_{n,q}^{(q)} &= \sum_{k=1}^{s_n} d_{k,q}^{(q)} X_{n-k} \\ &= \begin{cases} 0, & s_n < q+1; \\ D_{n-1,q}^{(q)} + c_q^{(q)} X_{n-q-1} - d_{s,q}^{(q)} X_{n-s-1}, \quad s_n = s \ge q+1; \\ D_{n-1,q}^{(q)} + c_q^{(q)} X_{n-q-1}, & s_n = s+1 \ge q+1, \end{cases} \\ D_{n,q}^{(b)} &= \sum_{k=1}^{s_n} d_{k,q}^{(b)} X_{n-k} \quad for \quad b=q-1,q-2,\ldots,1, \\ &= \begin{cases} 0, & s_n < b+1; \\ D_{n-1,q}^{(b)} + c_q^{(b)} X_{n-b-1} + D_{n,q}^{(b+1)} - d_{s,q}^{(b)} X_{n-s-1}, & s_n = s+1 \ge b+1, \end{cases} \end{split}$$
$$K_{n,q} = \sum_{k=1}^{s_n} k^q X_{n-k} = \begin{cases} 0, & s_n = s = 0; \\ K_{n-1,q} + X_{n-1} + D_{n,q}^{(1)} - s^q X_{n-s-1}, & s_n = s > 0; \\ K_{n-1,q} + X_{n-1} + D_{n,q}^{(1)}, & s_n = s + 1, \end{cases}$$

$$K_{n,0} = \sum_{k=1}^{s_n} X_{n-k} = \begin{cases} 0, & s_n = s = 0; \\ K_{n-1,0} + X_{n-1} - X_{n-s-1}, & s_n = s > 0; \\ K_{n-1,0} + X_{n-1}, & s_n = s + 1, \end{cases}$$

$$R_{n,b} = \sum_{i=1}^{n} \sum_{k=1}^{s_i} k^b X_i X_{i-k} = R_{n-1,b} + X_n K_{n,b} \quad for \quad b = 0, q,$$

$$Q_n = \sum_{i=1}^{n} X_i^2 = Q_{n-1} + X_n^2.$$

Analogous to Proposition A.2, the recursive formulas for the additional components are as follows:

Proposition A.4. Suppose the subsampling parameter $\{s_i \in \mathbb{N}_0\}_{i \in \mathbb{Z}^+}$ is a monotonically increasing sequence such that $\sup_{i \in \mathbb{Z}^+} |s_{i+1} - s_i| \leq 1$. Write its value at the last iteration as $s = s_{n-1}$. For each b = 0, q, the following components can be updated in O(1) time:

$$\begin{split} k_{n,b} &= \sum_{k=1}^{s_n} k^b = \begin{cases} 0, & s_n = s = 0; \\ k_{n-1,b}, & s_n = s > 0; \\ k_{n-1,b} + s_n^b, & s_n = s + 1, \end{cases} \\ r_{n,b} &= \sum_{i=1}^n \sum_{k=1}^{s_i} k^b = r_{n-1,b} + k_{n,b}, \\ U_{n,b} &= \sum_{i=1}^n \sum_{k=1}^{s_i} k^b X_i = U_{n-1,b} + k_{n,b} X_n, \\ V_{n,b} &= \sum_{i=1}^n \sum_{k=1}^{s_i} k^b X_{i-k} = V_{n-1,b} + K_{n,b}, \\ \bar{X}_n &= \frac{1}{n} \sum_{i=1}^n X_i = \frac{(n-1)\bar{X}_{n-1} + X_n}{n}. \end{split}$$

The recursive algorithm for $\hat{\sigma}_{n,\mathsf{LASER}(q,1)}^2$ is given by Algorithm A.3.

A.5 LASER (q, ϕ) : Memory Parameter

In this subsection, we illustrate how to derive recursive formulas for $\hat{\sigma}_{n,\mathsf{LASER}(q,\phi)}^2$ with q = 1and $\phi \ge 2$. We take these values to focus on how to perform O(1)-space update. The general algorithm for $q \in \mathbb{Z}^+$ and $\phi \ge 2$, which is implemented in our R package, can be obtained by combining the ideas in Section A.4 and here. Algorithm A.3: LASER(q, 1)

[1] initalization: Set $n = 1, s = 0, Q_n = X_1^2, \bar{X}_n = X_1$ [2] [3] Set $K_{n,b} = R_{n,b} = k_{n,b} = r_{n,b} = U_{n,b} = V_{n,b} = 0$ for b = 0, qCompute $c_q^{(b)}$, and set $D_{n,q}^{(b)} = d_{s_n,q}^{(b)} = 0$ for b = 1, 2, ..., q[4] Push X_1 into \vec{x} [5] begin [6] Receive X_{n+1} [7] Set n = n + 1[8] Compute s_n, t_n [9] Retrieve $X_{n-1}, \ldots, X_{n-1-\max\{\min(s_n-1,q),0\}}$ and X_{n-s-1} from \vec{x} [10] if $\underline{s_n} = \underline{s}$ then [11] Update $D_{n,q}^{(b)}$ with Prop. A.3 for $b = q, q - 1, \dots, 1$ [12] Update $K_{n,b}$ with Prop. A.3 for b = 0, q[13] Pop X_{n-s} from \vec{x} [14] else [15] Update $d_{s_n,q}^{(b)}, D_{n,q}^{(b)}$ with Prop. A.3 for $b = q, q - 1, \dots, 1$ [16] Update $K_{n,b}$ with Prop. A.3 and $k_{n,b}$ with Prop. A.4 for b = 0, q[17] Push X_n into \vec{x} [18] Update $R_{n,b}, Q_n$ with Prop. A.3 and $r_{n,b}, U_{n,b}, V_{n,b}, \overline{X}_n$ with Prop. A.4 for b = 0, q[19] Set $s = s_n$ [20] Output $\hat{\sigma}_{n,\mathsf{LASER}(q,1)}^2 = \{Q_n + 2R_n^{(0,q)} + (2r_n^{(0,q)} - n)\bar{X}_n^2 - 2\bar{X}_n(U_n^{(0,q)} + V_n^{(0,q)})\}/n$ [21]

To begin with, we follow the same procedure as in Sections A.2 and A.3 to write the estimator as

$$\hat{\sigma}_{n,\mathsf{LASER}(1,\phi)}^{2} = \frac{1}{n} \sum_{i=1}^{n} (X_{i} - \bar{X}_{n})^{2} + \frac{2}{n} \sum_{i=1}^{n} \sum_{k=1}^{s'_{i}} \left(1 - \frac{k}{t_{n}}\right) (X_{i} - \bar{X}_{n}) (X_{i-k} - \bar{X}_{n})$$
$$= \frac{1}{n} \left\{ Q_{n} + 2R_{n}^{\prime(0,1)} + (2r_{n}^{\prime(0,1)} - n)\bar{X}_{n}^{2} - 2\bar{X}_{n} (U_{n}^{\prime(0,1)} + V_{n}^{\prime(0,1)}) \right\},$$

where the shorthands, e.g., $R'_{n}^{(0,1)} = R'_{n,0} - R'_{n,1}/t_n$, are similarly introduced in Section A.3; see Section A.1 for an overview of notations. Since q = 1, the components are analogous to those in Propositions A.1 and A.2 with the only difference in the use of the ramped subsampling parameter s'_i . When s'_i is increasing, there is no difference in the recursive formulas. However, when s'_i is too large that it resets to the value of s_i , we need another way to update the components recursively. For $\phi \ge 2$, this can be done by "pre-calculating" some components. We summarize the recursive formulas for the zero-mean components below.

Proposition A.5. Suppose the intended subsampling parameter $\{s_i \in \mathbb{N}_0\}_{i \in \mathbb{Z}^+}$ is a monotonically increasing sequence such that $\sup_{i \in \mathbb{Z}^+} |s_{i+1} - s_i| \leq 1$. Furthermore, suppose the ramped subsampling parameter defined in (2.8), i.e., $\{s'_i\}_{i \in \mathbb{Z}^+}$, is ramped with $\phi \geq 2$. Write its value at the last iteration as $s' = s'_{n-1}$. Denote the ramping upper bound by $a_n = [\phi s_n]$. To keep the branching simple, assume s_n and a_n only increase when s'_n resets. Then, the following components can be

updated in O(1) time and O(1) space:

Note that $K''_{n-1,0}, K''_{n-1,1}, X'_{n-1}, X''_{n-1}$ are the "pre-calculated" versions of $K'_{n-1,0}, K'_{n-1,1}, X_{n-1}, X_{n-s_n}$, respectively, when s'_n resets to s_n . Using these components, we do not need to store the past observations explicitly, e.g., by maintaining the vector \vec{x} in Algorithm A.1. As a result, only a constant amount of memory is involved. For the restriction that s_n and a_n only increase when s'_n resets, we remark that the statistical efficiency is not affected asymptotically as it just slightly delay the increment of s_n . For the additional components, their recursive formulas are as follows:

Proposition A.6. Suppose the intended subsampling parameter $\{s_i \in \mathbb{N}_0\}_{i \in \mathbb{Z}^+}$ is a monotonically increasing sequence such that $\sup_{i \in \mathbb{Z}^+} |s_{i+1} - s_i| \leq 1$. Furthermore, suppose the ramped subsampling parameter defined in (2.8), i.e., $\{s'_i\}_{i \in \mathbb{Z}^+}$, is ramped with $\phi \geq 2$. Write its value at the last iteration as $s' = s'_{n-1}$. Denote the ramping upper bound by $a_n = \lfloor \phi s_n \rfloor$. To keep the branching simple, assume s_n and a_n only increase when s'_n resets. For each b = 0, 1, the following components can be updated in O(1) time and O(1) space:

$$\begin{split} k_{n,b}' &= \sum_{k=1}^{s'_n} k^b = \begin{cases} 0, & 0 = s'_n = s'; \\ k_{n-1,b}' + (s'_n)^b, & 0 < s'_n = s' + 1; \\ k_{n-1,b}', & 0 < s'_n = s_{n-1} < s'; \\ k_{n-1,b}' + s_n^b, & 0 < s'_n = s_{n-1} + 1 < s', \end{cases} \\ k_{n,b}'' &= \begin{cases} k_{n-1,b}'' + (a_{n-1} - s'_n)^b, & s'_n \ge a_{n-1} - s_{n-1} \ and \ s'_n > 0; \\ 0, & s'_n < a_{n-1} - s_{n-1} \ or \ s'_n = 0, \end{cases} \\ r_{n,b}' &= \sum_{i=1}^n \sum_{k=1}^{s'_i} k^b = r_{n-1,b}' + k_{n,b}', \end{cases} \\ U_{n,b}' &= \sum_{i=1}^n \sum_{k=1}^{s'_i} k^b X_i = U_{n-1,b}' + k_{n,b}' X_n, \\ V_{n,b}' &= \sum_{i=1}^n \sum_{k=1}^{s'_i} k^b X_{i-k} = V_{n-1,b}' + K_{n,b}', \\ \bar{X}_n &= \frac{1}{n} \sum_{i=1}^n X_i = \frac{(n-1)\bar{X}_{n-1} + X_n}{n}. \end{split}$$

The recursive algorithm for $\hat{\sigma}_{n,\mathsf{LASER}(1,\phi)}^2$ is given by Algorithm A.4.

A.6 Mini-batch Estimation

In mini-batch estimation, the traditional recursive formulas are extended to updating at $n_0 = 1 < n_1 < n_2 < \cdots$ instead of at $n = 1, 2, \ldots$. In this subsection, we illustrate how to derive recursive formulas for $\hat{\sigma}_{n_j, \text{LASER}(q, \phi)}^2$ with q = 1 and $\phi = 1$. The general algorithm for $q \in \mathbb{Z}^+$, which is implemented in our R package, can be obtained by combining the ideas in Section A.4 and here. Note that we do not develop algorithm for $\phi > 1$ because fewer operations can be vectorized under ramping. In that case, the improvement in computational efficiency brought by mini-batch estimation is less significant.

To begin with, we follow the same procedure as in Sections A.2 and A.3 to write the estimator as

$$\hat{\sigma}_{n_j,\mathsf{LASER}(1,1)}^2 = \frac{1}{n_j} \sum_{i=1}^{n_j} (X_i - \bar{X}_{n_j})^2 + \frac{2}{n_j} \sum_{i=1}^{n_j} \sum_{k=1}^{s_i} \left(1 - \frac{k}{t_{n_j}} \right) (X_i - \bar{X}_{n_j}) (X_{i-k} - \bar{X}_{n_j})$$
$$= \frac{1}{n_j} \left\{ Q_{n_j} + 2R_{n_j}^{(0,1)} + (2r_{n_j}^{(0,1)} - n_j) \bar{X}_{n_j}^2 - 2\bar{X}_{n_j} (U_{n_j}^{(0,1)} + V_{n_j}^{(0,1)}) \right\},$$

where the shorthands, e.g., $R_{n_j}^{(0,1)} = R_{n_j,0} - R_{n_j,1}/t_{n_j}$, are similarly introduced in Section A.3;

Algorithm A.4: LASER $(1, \phi)$

[1] initalization: [2] Set $n = 1, s = 0, Q_n = X_1^2, \bar{X}_n = X_1$ [3] Set $K'_{n,b} = K''_{n,b} = R'_{n,b} = k'_{n,b} = k''_{n,b} = r'_{n,b} = U'_{n,b} = V'_{n,b} = 0$ for b = 0, 1[4] Set $s' = a = 0, X'_n = X''_n = X_1$ begin [5] Receive X_{n+1} [6] Set n = n + 1[7] Compute s_n, t_n, a_n [8] if s' + 1 < a then [9] Set $s'_n = s' + 1$, $s_n = s$, $a_n = a$ /* restrict s_n when $s'_n > s'$ */ [10] Update $K'_{n,b}$ with Prop. A.5 and $k'_{n,b}$ with Prop. A.6 for b = 0, 1[11] if $\underline{s'_n \ge a-s}$ then [12] Update $K''_{n,b}$ with Prop. A.5 and $k''_{n,b}$ with Prop. A.6 for b = 0, 1[13] [14] else Set $s'_n = s_n, a = a_n$ ۲15<u>]</u> if $\underline{s'_n} == \underline{s}$ then [16] Update $K'_{n,b}$ with Prop. A.5 and $k'_{n,b}$ with Prop. A.6 for b = 0, 1[17] else [18] Update $K'_{n,b}$ with Prop. A.5 and $k'_{n,b}$ with Prop. A.6 for b = 0, 1[19] Set $K_{n,b}'' = k_{n,b}'' = 0$ for b = 0, 1[20] Update $R'_{n,b}, Q_n$ with Prop. A.5 and $r'_{n,b}, U'_{n,b}, V'_{n,b}, \overline{X}_n$ with Prop. A.6 for b = 0, 1[21] if $\underline{s'_n = a_n - s_n}_{\text{Set } X''_n = X'_n}$ then [22] [23] Set $s' = s'_n, s = s_n, X'_n = X_n$ [24] Output $\hat{\sigma}_{n,\text{LASER}(1,\phi)}^2 = \{Q_n + 2R_n^{\prime(0,1)} + (2r_n^{\prime(0,1)} - n)\bar{X}_n^2 - 2\bar{X}_n(U_n^{\prime(0,1)} + V_n^{\prime(0,1)})\}/n$ [25]

see Section A.1 for an overview of notations. Note that the components are identical to those in Propositions A.1 and A.2. However, we need to work on computationally efficient formulas for $\vec{K}_{n_j,b} = (K_{n_{j-1}+1,b}, \ldots, K_{n_j,b})^{\mathsf{T}}$ and $\vec{k}_{n_j,b} = (k_{n_{j-1}+1,b}, \ldots, k_{n_j,b})^{\mathsf{T}}$ where b = 0, 1. To this end, we can utilize vectorized operations such as cumulative sums. The corresponding recursive formulas are stated below.

Proposition A.7. Suppose the subsampling parameter $\{s_i \in \mathbb{N}_0\}_{i \in \mathbb{Z}^+}$ is a monotonically increasing sequence such that $\sup_{i \in \mathbb{Z}^+} |s_{i+1} - s_i| \leq 1$. Write its value at the last iteration as $s = s_{n_{j-1}}$, where $n_0 = 1 < n_1 < n_2 < \cdots$ are the sample points that are of interest. Let $f(i) = \inf \{k : \sum_{h=2}^k \mathbb{1}_{s_h=s_{h-1}} \geq i\}$. To keep the branching simple, assume $s_{n_{j-1}+1} = s_{n_{j-1}}$ for all $j \in \mathbb{Z}^+$. For $n_{j-1} < n \leq n_j$, the following components can be updated in $O(n_j - n_{j-1})$ time:

$$K_{n,0} = K_{n_{j-1},0} + \sum_{i=n_{j-1}}^{n-1} X_i - \sum_{i=n_{j-1}-s}^{n-s_n-1} X_i,$$

$$K_{n,1} = K_{n_{j-1},1} + \sum_{i=n_{j-1}+1}^n K_{i,0} - \sum_{i=n_{j-1}-s}^{n-s_n-1} s_{f(i)} X_i,$$

$$k_{n,b} = k_{n_{j-1},b} + \sum_{i=s+1}^{s_n} i^b \quad for \quad b = 0, 1.$$

Note that we fix $s_{n_{j-1}+1} = s_{n_{j-1}}$ for all $j \in \mathbb{Z}^+$ to ensure the subsampling parameter at the beginning of each mini-batch update agrees with that at the end of the previous update. This does not affect the statistical efficiency asymptotically as it just delay the increment of $s_{n_{j-1}+1}$ (if any) by one observation. For the definition of f(i), it is used to handle changes in the subsampling parameter when we compute cumulative sums. Once $\vec{K}_{n_{j},b}$ and $\vec{k}_{n_{j},b}$ are available, only the values at the end point n_{j} are required for the remaining components. We summarize their recursive formulas as follows:

Proposition A.8. Suppose the subsampling parameter $\{s_i \in \mathbb{N}_0\}_{i \in \mathbb{Z}^+}$ is a monotonically increasing sequence such that $\sup_{i \in \mathbb{Z}^+} |s_{i+1} - s_i| \leq 1$. Further suppose $n_0 = 1 < n_1 < n_2 < \cdots$ are the sample points that are of interest. To keep the branching simple, assume $s_{n_{j-1}+1} = s_{n_{j-1}}$ for all $j \in \mathbb{Z}^+$. For each b = 0, 1, the following components can be updated in $O(n_j - n_{j-1})$ time:

$$\begin{split} R_{n_{j},b} &= R_{n_{j-1},b} + \sum_{i=n_{j-1}+1}^{n_{j}} X_{i}K_{i,b}, \\ Q_{n_{j}} &= Q_{n_{j-1}} + \sum_{i=n_{j-1}+1}^{n_{j}} X_{i}^{2}, \\ r_{n_{j},b} &= r_{n_{j-1},b} + \sum_{i=n_{j-1}+1}^{n_{j}} k_{i,b}, \\ U_{n_{j},b} &= U_{n_{j-1},b} + \sum_{i=n_{j-1}+1}^{n_{j}} k_{i,b}X_{i}, \\ V_{n_{j},b} &= V_{n_{j-1},b} + \sum_{i=n_{j-1}+1}^{n_{j}} K_{i,b}, \\ \bar{X}_{n_{j}} &= \frac{1}{n_{j}} \left(n_{j-1}\bar{X}_{n_{j-1}} + \sum_{i=n_{j-1}+1}^{n_{j}} X_{i} \right) \end{split}$$

The recursive algorithm for $\hat{\sigma}_{n_i, \text{LASER}(1,1)}^2$ is given by Algorithm A.5.

A.7 Nuisance Parameter Estimation

In this subsection, we illustrate how to derive recursive formulas for $\hat{v}_{q,n,\mathsf{LASER}(p,\phi)}$ with $p,q \in \mathbb{Z}^+$ and $\phi = 1$. The general algorithms under ramping or mini-batch estimation, which are implemented in our R package, can be obtained by combining the ideas in Sections A.5, A.6 and here.

To begin with, we follow the same procedure as in Sections A.2 and A.3 to write the estimator as

$$\hat{v}_{q,n,\mathsf{LASER}(p,1)} = \frac{2}{n} \sum_{i=1}^{n} \sum_{k=1}^{s_i} \left(1 - \frac{k^p}{t_n^p} \right) k^q (X_i - \bar{X}_n) (X_{i-k} - \bar{X}_n)$$

Algorithm A.5: Mini-batch LASER(1,1)

[1] initalization: [2] Set $n = 1, s = j = 0, Q_{n_j} = X_1^2, \bar{X}_{n_j} = X_1$ [3] Set $K_{n_j,b} = R_{n_j,b} = k_{n_j,b} = r_{n_j,b} = U_{n_j,b} = V_{n_j,b} = 0$ for b = 0, 1[4] Set $\vec{x} = (X_1)^{\mathsf{T}}$ [5] begin Receive $X_{n+1}, X_{n+2}, ..., X_{n_{i+1}}$ [6] Set j = j + 1[7] Compute $n_j, \vec{s}_{n_j} = (s_{n_{j-1}+1}, \dots, s_{n_j})^{\mathsf{T}}, t_{n_j}$ Set $s_{n_{j-1}+1} = s$ /* ensure the subsampling parameters agree */ [8] [9] Compute $\vec{K}_{n_i,b}$, $\vec{k}_{n_i,b}$ with Prop. A.7 for b = 0, 1 using cumulative sums [10] Update $R_{n_j,b}, Q_{n_j}, r_{n_j,b}, U_{n_j,b}, V_{n_j,b}, \overline{X}_{n_j}$ with Prop. A.8 for b = 0, 1[11] Retrieve $K_{n_i,b}, k_{n_i,b}$ from $\bar{K}_{n_i,b}, \bar{k}_{n_i,b}$ for b = 0, 1[12] Set $n = n_j, s = s_{n_j}$ [13] Set $\vec{x} = (X_{n-s}, \dots, X_n)^{\mathsf{T}}$ [14] $\text{Output} \ \hat{\sigma}_{n_{j},\text{LASER}(1,1)}^{2} = \{ Q_{n_{j}} + 2R_{n_{j}}^{(0,1)} + (2r_{n_{j}}^{(0,1)} - n_{j})\bar{X}_{n_{j}}^{2} - 2\bar{X}_{n_{j}}(U_{n_{j}}^{(0,1)} + V_{n_{j}}^{(0,1)}) \} / n_{j}$ [15]

$$= \frac{2}{n} \left\{ R_n^{(q,p+q,p)} + r_n^{(q,p+q,p)} \bar{X}_n^2 - \bar{X}_n (U_n^{(q,p+q,p)} + V_n^{(q,p+q,p)}) \right\}$$

where the shorthands, e.g., $R_n^{(q,p+q,p)} = R_{n,q} - R_{n,p+q}/t_n^p$, are the full versions of those introduced in Section A.3; see Section A.1 for an overview of notations. The components are analogous to those in Propositions A.3 and A.4 with the only difference in the new subscript p + q. We summarize the recursive formulas for the zero-mean components below.

Proposition A.9. Suppose the subsampling parameter $\{s_i \in \mathbb{N}_0\}_{i \in \mathbb{Z}^+}$ is a monotonically increasing sequence such that $\sup_{i \in \mathbb{Z}^+} |s_{i+1} - s_i| \leq 1$. Write its value at the last iteration as $s = s_{n-1}$. For each $\rho = q, p + q$, the following components can be updated in O(1) time:

$$\begin{split} c_{\rho}^{(b)} &= \sum_{r=0}^{b} (-1)^{r} {\binom{b}{r}} (b-r+1)^{\rho} \quad for \quad b=1,2,\ldots,\rho, \\ \\ d_{s_{n},\rho}^{(b)} &= \begin{cases} 0, & s_{n} < b+1 \ and \ b=1,2,\ldots,\rho; \\ c_{\rho}^{(b)}, & s_{n} = b+1 \ and \ b=1,2,\ldots,\rho; \\ d_{s,\rho}^{(b)}, & s_{n} = s > b+1 \ and \ b=1,2,\ldots,\rho; \\ d_{s,\rho}^{(b)}, & s_{n} = s+1 > b+1 \ and \ b=\rho; \\ d_{s,\rho}^{(b)} + d_{s+1,\rho}^{(b+1)}, & s_{n} = s+1 > b+1 \ and \ b=1,2,\ldots,\rho-1, \end{cases} \\ \\ D_{n,\rho}^{(\rho)} &= \sum_{k=1}^{s_{n}} d_{k,\rho}^{(\rho)} X_{n-k} \\ &= \begin{cases} 0, & s_{n} < \rho+1; \\ D_{n-1,\rho}^{(\rho)} + c_{\rho}^{(\rho)} X_{n-\rho-1} - d_{s,\rho}^{(\rho)} X_{n-s-1}, & s_{n} = s \geq \rho+1; \\ D_{n-1,\rho}^{(\rho)} + c_{\rho}^{(\rho)} X_{n-\rho-1}, & s_{n} = s+1 \geq \rho+1, \end{cases} \end{split}$$

$$\begin{split} D_{n,\rho}^{(b)} &= \sum_{k=1}^{s_n} d_{k,\rho}^{(b)} X_{n-k} \quad for \quad b = \rho - 1, \rho - 2, \dots, 1, \\ &= \begin{cases} 0, & s_n < b + 1; \\ D_{n-1,\rho}^{(b)} + c_{\rho}^{(b)} X_{n-b-1} + D_{n,\rho}^{(b+1)} - d_{s,\rho}^{(b)} X_{n-s-1}, & s_n = s \ge b + 1; \\ D_{n-1,\rho}^{(b)} + c_{\rho}^{(b)} X_{n-b-1} + D_{n,\rho}^{(b+1)}, & s_n = s + 1 \ge b + 1, \end{cases} \\ K_{n,\rho} &= \sum_{k=1}^{s_n} k^{\rho} X_{n-k} = \begin{cases} 0, & s_n = s = 0; \\ K_{n-1,\rho} + X_{n-1} + D_{n,\rho}^{(1)} - s^{\rho} X_{n-s-1}, & s_n = s > 0; \\ K_{n-1,\rho} + X_{n-1} + D_{n,\rho}^{(1)}, & s_n = s + 1, \end{cases} \\ R_{n,\rho} &= \sum_{i=1}^{n} \sum_{k=1}^{s_i} k^{\rho} X_i X_{i-k} = R_{n-1,\rho} + X_n K_{n,\rho}. \end{split}$$

For the additional components, their recursive formulas are as follows:

Proposition A.10. Suppose the subsampling parameter $\{s_i \in \mathbb{N}_0\}_{i \in \mathbb{Z}^+}$ is a monotonically increasing sequence such that $\sup_{i \in \mathbb{Z}^+} |s_{i+1} - s_i| \leq 1$. Write its value at the last iteration as $s = s_{n-1}$. For each $\rho = q, p + q$, the following components can be updated in O(1) time:

$$\begin{aligned} k_{n,\rho} &= \sum_{k=1}^{s_n} k^{\rho} = \begin{cases} 0, & s_n = s = 0; \\ k_{n-1,\rho}, & s_n = s > 0; \\ k_{n-1,\rho} + s_n^{\rho}, & s_n = s + 1, \end{cases} \\ r_{n,\rho} &= \sum_{i=1}^n \sum_{k=1}^{s_i} k^{\rho} = r_{n-1,\rho} + k_{n,\rho}, \\ U_{n,\rho} &= \sum_{i=1}^n \sum_{k=1}^{s_i} k^{\rho} X_i = U_{n-1,\rho} + k_{n,\rho} X_n, \\ V_{n,\rho} &= \sum_{i=1}^n \sum_{k=1}^{s_i} k^{\rho} X_{i-k} = V_{n-1,\rho} + K_{n,\rho}, \\ \bar{X}_n &= \frac{1}{n} \sum_{i=1}^n X_i = \frac{(n-1)\bar{X}_{n-1} + X_n}{n}. \end{aligned}$$

The recursive algorithm for $\hat{v}_{q,n,\mathsf{LASER}(p,1)}$ is similar to Algorithm A.3 once we have Propositions A.9 and A.10. Therefore, we omit its pseudocode here.

A.8 Automatic Optimal Parameters Selection

In this subsection, we discuss how to select the smoothing parameters s_n and t_n automatically when we update $\hat{\sigma}_{n,\mathsf{LASER}(q,\phi)}^2$ with $q \in \mathbb{Z}^+$ and $\phi = 1$. The general algorithms under ramping or mini-batch estimation are similar and thus omitted.

To begin with, denote the set of components used to update $\hat{\sigma}_n^2$ and $\hat{v}_{q,n}$ by $\mathscr{C}(\hat{\sigma}_n^2)$ and

Algorithm A.6: Automatic optimal parameters selector for LASER(q, 1)

[1] input for the selector: (i) n, s_n, t_n – the sample size, subsampling and tapering parameters from the last iteration [2] (ii) $\Psi_{\diamond}, \psi_{\star}$ – the optimal coefficient (excluding κ) and exponent for computing s_n [3] (iii) $\Theta_{\diamond}, \theta_{\star}$ – the optimal coefficient (excluding κ) and exponent for computing t_n [4] (iv) s_0, t_0 – the minimum values of subsampling and tapering parameters [5] input for the estimators: (v) X_{n+1} – the new observation [7] (vi) $\mathscr{C}(\hat{\sigma}_{n,\mathsf{LASER}(q,1)}^2), \mathscr{C}(\hat{v}_{q,n,\mathsf{LASER}(p,1)})$ – the components for recursive estimation [8] begin [9] Set $\hat{\kappa} = \left| \hat{v}_{q,n,\mathsf{LASER}(p,1)} \right| / \hat{\sigma}_{n,\mathsf{LASER}(q,1)}^2$ [10] Set $s_{n+1} = \lfloor \Psi_{\diamond} \hat{\kappa}^{2/(1+2q)} (n+1)^{\psi_{\star}} \rfloor$ and $t_{n+1} = \lfloor \Theta_{\diamond} \hat{\kappa}^{2/(1+2q)} (n+1)^{\theta_{\star}} \rfloor$ [11] if $s_n < \max(s_{n+1}, s_0)$ then [12] Set $s_{n+1} = s_n + 1$ [13] else [14] Set $s_{n+1} = s_n$ ۲15<u>]</u> $\mathbf{if} \underbrace{t_n < \max(t_{n+1}, t_0)}_{\text{Set } t_{n+1} = t_n + 1} \mathbf{then}$ [16] [17] else [18] Set $t_{n+1} = t_n$ [19] Update $\hat{\sigma}_{n, \mathsf{LASER}(q, 1)}^2$ with $X_{n+1}, \mathscr{C}(\hat{\sigma}_{n, \mathsf{LASER}(q, 1)}^2), s_{n+1}, t_{n+1}$ [20] Update $\hat{v}_{q,n,\mathsf{LASER}(p,1)}$ with $X_{n+1}, \mathscr{C}(\hat{v}_{q,n,\mathsf{LASER}(p,1)})$ [21]

 $\mathscr{C}(\hat{v}_{q,n})$. For instance,

$$\mathscr{C}(\hat{\sigma}_{n,\mathsf{LASER}(1,1)}^2) = \left\{ n, s, Q_n, \bar{X}_n, \left\{ K_{n,b}, R_{n,b}, k_{n,b}, r_{n,b}, U_{n,b}, V_{n,b} \right\}_{b=0,1} \right\};$$

see Algorithm A.2. We also define $\Psi_{\diamond} = \Psi_{\star}/\kappa^{2/(1+2q)}$ and $\Theta_{\diamond} = \Theta_{\star}/\kappa^{2/(1+2q)}$, which denote the optimal coefficients for computing s_n and t_n excluding the unknown κ ; see Corollary 2. Then, the automatic optimal parameters selector is given by Algorithm A.6.

There are several remarks about Algorithm A.6. First, the nuisance parameter estimate $\hat{\kappa}$ seems to use fewer observations than the updated LRV estimate $\hat{\sigma}_{n+1}^2$. This is natural because the selector uses the previous LRV estimate $\hat{\sigma}_n^2$ as the denominator for $\hat{\kappa}$. If the numerator $\hat{v}_{q,n}$ was updated first, the nuisance parameter estimate could be unstable in a mini-batch setting where $\hat{\kappa} = |\hat{v}_{q,n_{j+1}}| / \hat{\sigma}_{n_j}^2$. Alternatively, we can update $\hat{\kappa}$ as follows:

- (1) Set $\hat{\kappa}_n = |\hat{v}_{q,n}| / \hat{\sigma}_n^2$.
- (2) Update $\hat{\sigma}_n^2$ with $X_{n+1}, \hat{\kappa}_n$.
- (3) Update $\hat{v}_{q,n}$.
- (4) Set $\hat{\kappa}_{n+1} = |\hat{v}_{q,n+1}| / \hat{\sigma}_{n+1}^2$.
- (5) Reupdate $\hat{\sigma}_n^2$ with $X_{n+1}, \hat{\kappa}_{n+1}$.

Using a similar procedure, it is further possible to update $\hat{\kappa}$ iteratively. However, we do

not recommend doing so because the computation time will increase significantly with little improvement in statistical efficiency.

Second, updating $\hat{v}_{q,n}$ also requires smoothing parameters selection. Denote these ancillary smoothing parameters by s_n^{\diamond} and t_n^{\diamond} . We notice that it is not possible to select the coefficients of s_n^{\diamond} and t_n^{\diamond} based on asymptotic theory as another nuisance parameter κ_{p+q} will arise. This circular problem is common and well-known in nonparametric estimation. If we just consider the exponents, choosing $s_n^{\diamond} = \lfloor n^{1/\{1+2(p+q)\}} \rfloor$ may be too small and lead to poor empirical performance. Therefore, we suggest

$$s_n^{\diamond} = \begin{cases} \lfloor n^{1/2} \rfloor, & n \le n_0; \\ \\ \lfloor \max\left[n_0^{1/2}, (p+q)n^{1/\{1+2(p+q)\}} \right] \end{bmatrix}, & n > n_0, \end{cases}$$

where $n_0 = 1000$ is a threshold for small sample size and the empirical adjustment $\lfloor n^{1/2} \rfloor$ is recommended by Jones et al. (2006). The ancillary tapering parameter t_n^{\diamond} may admit the same form except that the floor functions are replaced by the ceiling functions as t_n^{\diamond} cannot be zero. Although it is possible to select the coefficients according to some benchmark models such as AR(q) instead of using p + q, this kind of benchmarking is still optimal to a particular stochastic process only. Hence we recommend the simple choice p + q to solve the problem of small exponents when p + q is large.

Third, the main smoothing parameters s_n and t_n can be lower bounded by s_0 and t_0 in finite sample. This is because if users wrongly choose a large characteristic exponent q without accounting for the empirical serial dependence, the computed s_n and t_n may be too small in finite sample. In light of a similar problem, Jones et al. (2006) suggested using $\lfloor n^{1/2} \rfloor$. However, this is not adaptive and Theorem 2 states that this is not MSE-optimal. To balance theoretical and practical considerations, we provide the possibility to lower bound s_n and t_n by s_0 and t_0 in our R package, where the default is $s_0 = t_0 = 5$. The empirical adjustment $\lfloor n^{1/2} \rfloor$ is implemented for the aforementioned ancillary smoothing parameters s_n^{\diamond} and t_n^{\diamond} in small sample only. Example 4 shows that recursive estimation with Algorithm A.6 performs very close to that with the oracle and much better than that with a pilot study.

A.9 Multivariate Extension

In the main text, we focus on LRV estimation to introduce new ideas. Nevertheless, our framework is also applicable to long-run covariance matrix (LRCM) estimation. Consider a

d-dimensional time series $\{ \boldsymbol{X}_i = (X_i^{(1)}, \dots, X_i^{(d)})^{\mathsf{T}} \}_{i \in \mathbb{Z}}$, where $d \in \mathbb{Z}^+$. The LRCM is defined as $\Sigma := \lim_{n \to \infty} n \operatorname{Var}(\bar{\boldsymbol{X}}_n)$, where $\Gamma_k = \operatorname{Cov}(X_0, X_k)$. We can estimate Σ by

$$\hat{\Sigma}_n = \hat{\Sigma}_n(W) \coloneqq \frac{1}{n} \sum_{i=1}^n \sum_{j=1}^n W_n(i,j) (\boldsymbol{X}_i - \bar{\boldsymbol{X}}_n) (\boldsymbol{X}_j - \bar{\boldsymbol{X}}_n)^{\mathsf{T}}.$$
(A.1)

Using the symmetry of (A.1), the (h, k)-th entry of $\hat{\Sigma}_n$ can be expressed as

$$\hat{\Sigma}_{n}^{(h,k)} = \frac{1}{2n} \sum_{i=1}^{n} \sum_{j=1}^{n} W_{n}(i,j) \left(\hat{X}_{i}^{(h)} \hat{X}_{j}^{(k)} + \hat{X}_{i}^{(k)} \hat{X}_{j}^{(h)} \right)
= \frac{1}{2n} \sum_{i=1}^{n} \sum_{j=1}^{n} W_{n}(i,j) \left\{ \left(\hat{X}_{i}^{(h)} + \hat{X}_{i}^{(k)} \right) \left(\hat{X}_{j}^{(h)} + \hat{X}_{j}^{(k)} \right) - \hat{X}_{i}^{(h)} \hat{X}_{j}^{(h)} - \hat{X}_{i}^{(k)} \hat{X}_{j}^{(k)} \right\}
= \frac{1}{2} \left\{ \hat{\sigma}_{n}^{2} (W; X^{(h)} + X^{(k)}) - \hat{\sigma}_{n}^{2} (W; X^{(h)}) - \hat{\sigma}_{n}^{2} (W; X^{(k)}) \right\},$$
(A.2)

where $\hat{X}_{i}^{(k)} = X_{i}^{(k)} - \bar{X}_{n}^{(k)}$ is the demeaned version of $X_{i}^{(k)}$ for i = 1, ..., n and k = 1, ..., d; and $\hat{\sigma}_{n}^{2}(W; Y)$ denotes the LRV estimator $\hat{\sigma}_{n}^{2}(W)$ if (univariate) data $Y_{1:n}$ are used. In view of (A.2), $\hat{\Sigma}_{n}$ can be expressed as a function of d(d+1)/2 LRV estimators $\hat{\sigma}_{n}^{2}$ with different input samples. Hence, all computational and statistical properties discussed before are also enjoyed by $\hat{\Sigma}_{n}$ without the need of deriving new formulas or theories.

In practice, users are usually interested in a scalar statistic $\mathbf{c}^{\mathsf{T}} \bar{\mathbf{X}}_n$, instead of the whole vector $\bar{\mathbf{X}}_n$, for some $\mathbf{c} \in \mathbb{R}^d$. Then, the smoothing parameters in $\hat{\Sigma}_n$ can be selected so that the LRV estimator of the reference statistic, i.e., $\hat{\sigma}_n^2(W; \mathbf{c}^{\mathsf{T}} \mathbf{X})$, is optimized. The vector \mathbf{c} can be interpreted as the relative importance of the d entries in \mathbf{X} . The selected smoothing parameters can be used for all entries, which enable efficient matrix operations and enhance computational efficiency. An additional advantage of this approach is that users only need to specify the relative importance of the d entries in \mathbf{X} . In contrast, Chan and Yau (2017a) required users to specify the relative importance of $d \times d$ entries in Σ . It is not only difficult to interpret but also hard to determine the relative importance.

B Proof of Theorems

B.1 Proof of Theorem 1

The proof is divided into 4 steps, which are stated in Sections B.1.1 to B.1.4. Each step requires some technical lemmas, whose proofs are deferred to Section E.

B.1.1 Replacement of Sample Mean

By Assumption 3(b) and the symmetry of γ_k , we can focus on the following rewritten form:

$$\hat{\sigma}_n^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n)^2 + \frac{2}{n} \sum_{i=2}^n \sum_{j=1}^{i-1} W_n(i,j) (X_i - \bar{X}_n) (X_j - \bar{X}_n).$$
(B.1)

Define the known-mean version of $\hat{\sigma}_n^2$ as

$$\bar{\sigma}_n^2 \coloneqq \frac{1}{n} \sum_{i=1}^n (X_i - \mu)^2 + \frac{2}{n} \sum_{i=2}^n \sum_{j=1}^{i-1} W_n(i,j) (X_i - \mu) (X_j - \mu).$$
(B.2)

By Minkowski inequality, we have

$$\left\|\hat{\sigma}_n^2 - \sigma^2\right\|_{\frac{\alpha}{2}} \leq \left\|\hat{\sigma}_n^2 - \bar{\sigma}_n^2\right\|_{\frac{\alpha}{2}} + \left\|\bar{\sigma}_n^2 - \sigma^2\right\|_{\frac{\alpha}{2}}.$$

Lemma E.1 states that $\|\hat{\sigma}_n^2 - \bar{\sigma}_n^2\|_{\alpha/2} = o(1)$. Hence it suffices to show that the known-mean version satisfies $\|\bar{\sigma}_n^2 - \sigma^2\|_{\alpha/2} = o(1)$.

Without loss of generality, assume $\mu = 0$ for the remaining of this proof. Otherwise, consider the demeaned series $\{X_i - \mu\}$.

B.1.2 Approximation by *m*-dependent Process

Define the *m*-dependent process and *m*-dependent process approximated version of $\bar{\sigma}_n^2$ as

$$\tilde{X}_i \coloneqq \mathbb{E}(X_i \mid \epsilon_{i-m}, \dots, \epsilon_i), \tag{B.3}$$

$$\tilde{\sigma}_n^2 \coloneqq \frac{1}{n} \sum_{i=1}^n \tilde{X}_i^2 + \frac{2}{n} \sum_{i=2}^n \sum_{j=1}^{i-1} W_n(i,j) \tilde{X}_i \tilde{X}_j, \tag{B.4}$$

respectively. By Minkowski inequality, we have

$$\left\|\bar{\sigma}_n^2 - \sigma^2\right\|_{\frac{\alpha}{2}} \le \left\|\bar{\sigma}_n^2 - \mathbb{E}(\bar{\sigma}_n^2) - \tilde{\sigma}_n^2 + \mathbb{E}(\tilde{\sigma}_n^2)\right\|_{\frac{\alpha}{2}} + \left\|\tilde{\sigma}_n^2 - \mathbb{E}(\tilde{\sigma}_n^2) + \mathbb{E}(\bar{\sigma}_n^2) - \sigma^2\right\|_{\frac{\alpha}{2}}.$$

Lemma E.2 states that $\|\bar{\sigma}_n^2 - \mathbb{E}(\bar{\sigma}_n^2) - \tilde{\sigma}_n^2 + \mathbb{E}(\tilde{\sigma}_n^2)\|_{\alpha/2} = o(1)$. Thus, it remains to show that $\|\tilde{\sigma}_n^2 - \mathbb{E}(\tilde{\sigma}_n^2) + \mathbb{E}(\bar{\sigma}_n^2) - \sigma^2\|_{\alpha/2} = o(1)$.

B.1.3 Approximation by Martingale Difference

For $i = 1, 2, \ldots, n$, define

$$M_i \coloneqq \sum_{h=0}^{\infty} \mathbb{E}(\tilde{X}_{i+h} \mid \mathscr{F}_i), \tag{B.5}$$

$$D_i \coloneqq M_i - \mathbb{E}(M_i \mid \mathscr{F}_{i-1}). \tag{B.6}$$

Note that $\{D_i\}$ is a martingale difference sequence with respect to the filtration \mathscr{F}_i . Therefore, the martingale difference approximated version of $\tilde{\sigma}_n^2$ is

$$\breve{\sigma}_n^2 \coloneqq \frac{1}{n} \sum_{i=1}^n D_i^2 + \frac{2}{n} \sum_{i=2}^n \sum_{j=1}^{i-1} W_n(i,j) D_i D_j.$$
(B.7)

Applying Minkowski inequality again,

$$\begin{split} \left\| \tilde{\sigma}_n^2 - \mathbb{E}(\tilde{\sigma}_n^2) + \mathbb{E}(\bar{\sigma}_n^2) - \sigma^2 \right\|_{\frac{\alpha}{2}} &\leq \left\| \tilde{\sigma}_n^2 - \mathbb{E}(\tilde{\sigma}_n^2) - \breve{\sigma}_n^2 + \mathbb{E}(\breve{\sigma}_n^2) \right\|_{\frac{\alpha}{2}} \\ &+ \left\| \breve{\sigma}_n^2 - \mathbb{E}(\breve{\sigma}_n^2) + \mathbb{E}(\bar{\sigma}_n^2) - \sigma^2 \right\|_{\frac{\alpha}{2}} \end{split}$$

By Lemma E.3, $\|\tilde{\sigma}_n^2 - \mathbb{E}(\tilde{\sigma}_n^2) - \check{\sigma}_n^2 + \mathbb{E}(\check{\sigma}_n^2)\|_{\alpha/2} = o(1)$. Therefore, it suffices to prove that $\|\check{\sigma}_n^2 - \mathbb{E}(\check{\sigma}_n^2) + \mathbb{E}(\bar{\sigma}_n^2) - \sigma^2\|_{\alpha/2} = o(1)$.

B.1.4 Application of Ergodic Theorem

Since D_i 's are martingale differences, $\mathbb{E}(D_i D_j) = 0$ for $i \neq j$. We can also choose m such that $m \to \infty$ as $n \to \infty$ by Assumption 2; see Section E.3. Therefore, by Lemma E.4, $\mathbb{E}(\check{\sigma}_n^2) \to \sigma^2$ as $n \to \infty$. Using the Ergodic Theorem (see, e.g., Durrett (2019)), we obtain $\|\check{\sigma}_n^2 - \sigma^2\|_{\alpha/2} = o(1)$. On the other hand,

$$\mathbb{E}(\bar{\sigma}_n^2) = \gamma_0 + \frac{2}{n} \sum_{i=2}^n \sum_{j=1}^{i-1} W_n(i,j) \gamma_{i-j} = \gamma_0 + \frac{2}{n} \sum_{i=2}^n \sum_{k=1}^{i-1} W_n(i,i-k) \gamma_k$$
$$= \gamma_0 + 2 \sum_{k=1}^{n-1} \frac{1}{n} \sum_{i=k+1}^n W_n(i,i-k) \gamma_k$$
$$= \gamma_0 + 2 \sum_{1 \le k \le b_n} w_{n,k} \gamma_k + 2 \sum_{b_n < k < n} w_{n,k} \gamma_k,$$

where $w_{n,k} = n^{-1} \sum_{i=k+1}^{n} W_n(i, i-k)$ is defined in Assumption 3. By the definition of LRV $\sigma^2 = \gamma_0 + 2 \sum_{k \in \mathbb{Z}^+} \gamma_k$ and Minkowski inequality, we have

$$\left| \mathbb{E}(\bar{\sigma}_{n}^{2}) - \sigma^{2} \right| = \left| 2 \sum_{1 \le k \le b_{n}} (w_{n,k} - 1) \gamma_{k} + 2 \sum_{b_{n} < k < n} (w_{n,k} - 1) \gamma_{k} - 2 \sum_{k=n}^{\infty} \gamma_{k} \right|$$

$$\leq 2 \sum_{1 \le k \le b_{n}} |w_{n,k} - 1| |\gamma_{k}| + 2 \sum_{b_{n} < k < n} (|w_{n,k}| + 1) |\gamma_{k}| + 2 \sum_{k=n}^{\infty} |\gamma_{k}|.$$

Note that under Assumption 1, $\sum_{k \in \mathbb{Z}} |\gamma_k| < \infty$; see Wu (2009). Therefore,

$$\sum_{1 \le k \le b_n} |w_{n,k} - 1| |\gamma_k| \le \max_{1 \le k \le b_n} |w_{n,k} - 1| \sum_{1 \le k \le b_n} |\gamma_k| = o(1),$$

$$\sum_{b_n < k < n} (|w_{n,k}| + 1) |\gamma_k| = \left(\max_{b_n < k < n} |w_{n,k}| + 1\right) \sum_{b_n < k < n} |\gamma_k| = o(1),$$

$$\sum_{k=n}^{\infty} |\gamma_k| = o(1).$$

It follows that $\left|\mathbb{E}(\bar{\sigma}_n^2) - \sigma^2\right| = o(1)$ and

$$\left|\mathbb{E}(\bar{\sigma}_n^2) - \mathbb{E}(\breve{\sigma}_n^2)\right| \le \left|\mathbb{E}(\bar{\sigma}_n^2) - \sigma^2\right| + \left|\mathbb{E}(\breve{\sigma}_n^2) - \sigma^2\right| = o(1).$$

Combining the results with Minkowski inequality,

$$\left\|\breve{\sigma}_{n}^{2} - \mathbb{E}(\breve{\sigma}_{n}^{2}) + \mathbb{E}(\bar{\sigma}_{n}^{2}) - \sigma^{2}\right\|_{\frac{\alpha}{2}} \leq \left\|\breve{\sigma}_{n}^{2} - \sigma^{2}\right\|_{\frac{\alpha}{2}} + \left|\mathbb{E}(\bar{\sigma}_{n}^{2}) - \mathbb{E}(\breve{\sigma}_{n}^{2})\right| = o(1),$$

which completes the proof.

B.2 Proof of Theorem 2

The proof is divided into 6 steps, which are stated in Sections B.2.1 to B.2.6. Each step requires some technical lemmas, whose proofs are deferred to Section E. Before beginning the proof, we explain some blocking variables that will be used later:

$$\nu_k \coloneqq \left[\frac{1}{(\phi - 1)\psi} \left(\frac{1}{\Psi} \right)^{\frac{1}{\psi}} k^{\frac{1}{\psi} - 2} \right], \tag{B.8}$$

$$\varpi_k \coloneqq \lceil (\phi - 1)k \rceil, \tag{B.9}$$

$$\eta_k \coloneqq \begin{cases} 1 + \sum_{h=1}^{k-1} \varpi_h \cdot \nu_h, & \phi \in (1, \infty); \\ \left\lceil \left(\frac{k}{\Psi}\right)^{\frac{1}{\psi}} \right\rceil, & \phi = 1. \end{cases}$$
(B.10)

Note that we group both $\{i \in \mathbb{Z}^+ : s_i = 0\}$ and $\{i \in \mathbb{Z}^+ : s_i = 1\}$ in the first block. For $\phi > 1$, we can partition $\{1, \ldots, n\}$ into s_n blocks of data with ramps as the subsample size will be ramped:

$$\{1,\ldots,n\} = \underbrace{\left\{\bigcup_{h=1}^{s_n-1}\bigcup_{i=1}^{\nu_h}\bigcup_{j=\eta_h+(i-1)\varpi_h}^{\eta_h+i\varpi_h-1}\{j\}}_{s_n-1 \text{ complete block}} \bigcup_{\text{last incomplete block}} \left\{\bigcup_{j=\eta_{s_n}}^{n}\{j\}\right\},$$

where $\bigcup_{i=i_0}^{i_1} E_i = \emptyset$ if $i_0 > i_1$, for any sets E_i 's. We visualize an example in Figure 8. Note that the physical meaning of η_k is the minimum sample size such that the intended subsample size is k. For $\phi = 1$, this can be seen from

$$\Psi \eta_k^{\psi} \sim k \iff \eta_k \sim \left(\frac{k}{\Psi}\right)^{\frac{1}{\psi}}$$

Blocking Partitions



Figure 8: A blocking example with n = 100, $\Psi = 1$, $\psi = 1/3$ and $\phi = 2$. The 4th block (solid blue) is incomplete so we do not need to find ν_4 .

For $\phi > 1$, this interpretation also holds but we define η_k based on ν_k and ϖ_k instead, where ν_k is the number of ramps and ϖ_k is the width of a ramp in the k-th block:

$$(\phi-1)k\cdot\nu_k \sim \eta_{k+1}-\eta_k \iff \nu_k \sim \frac{1}{(\phi-1)\psi}\left(\frac{1}{\Psi}\right)^{\frac{1}{\psi}}k^{\frac{1}{\psi}-2}.$$

B.2.1 Replacement of Sample Mean

We find the exact convergence rate of variance first. Recall the definitions of unknown-mean version $\hat{\sigma}_n^2$ and known-mean version $\bar{\sigma}_n^2$ in (B.1) and (B.2), respectively. For simplicity, write $\|\cdot\|_2 = \|\cdot\|$. We are going to approximate $\|\hat{\sigma}_n^2 - \mathbb{E}(\hat{\sigma}_n^2)\|$ by $\|\bar{\sigma}_n^2 - \mathbb{E}(\bar{\sigma}_n^2)\|$. To this end, we can use the following procedure:

First, by Minkowski inequality,

$$\begin{aligned} \left\| \hat{\sigma}_{n}^{2} - \mathbb{E}(\hat{\sigma}_{n}^{2}) \right\| &\leq \left\| \bar{\sigma}_{n}^{2} - \mathbb{E}(\bar{\sigma}_{n}^{2}) \right\| + \left\| \hat{\sigma}_{n}^{2} - \bar{\sigma}_{n}^{2} \right\| + \left| \mathbb{E}(\hat{\sigma}_{n}^{2}) - \mathbb{E}(\bar{\sigma}_{n}^{2}) \right| \\ &=: \left\| \bar{\sigma}_{n}^{2} - \mathbb{E}(\bar{\sigma}_{n}^{2}) \right\| + H_{1} + H_{2}. \end{aligned}$$
(B.11)

By similar steps as in (C.3),

$$G_{3,n} = O(n^{2\psi + \max\{2q(\psi-\theta), 0\}+1})$$
 and $G_{4,n} = O(n^{\psi + \max\{q(\psi-\theta), 0\}+1}),$

where $G_{3,n}$ and $G_{4,n}$ are defined in Lemma E.1. Therefore, by Lemma E.1 and Jensen's inequality,

$$H_{1} = O(n^{-1}) + O\left(n^{-\frac{3}{2}}G_{3,n}^{\frac{1}{2}}\right) + O\left(n^{-2}G_{4,n}\right) = O\left(n^{\psi+\max\{q(\psi-\theta),0\}-1}\right),$$

$$H_{2} \le \mathbb{E}\left|\hat{\sigma}_{n}^{2} - \bar{\sigma}_{n}^{2}\right| \le H_{1} = O\left(n^{\psi+\max\{q(\psi-\theta),0\}-1}\right).$$

(B.11) now becomes

$$\left\|\hat{\sigma}_{n}^{2} - \mathbb{E}(\hat{\sigma}_{n}^{2})\right\| \leq \left\|\bar{\sigma}_{n}^{2} - \mathbb{E}(\bar{\sigma}_{n}^{2})\right\| + O\left(n^{\psi + \max\{q(\psi-\theta), 0\}-1}\right).$$
(B.12)

On the other hand, applying Minkowski inequality to $\|\bar{\sigma}_n^2 - \mathbb{E}(\bar{\sigma}_n^2)\|$ yields

$$\begin{split} \left\| \bar{\sigma}_{n}^{2} - \mathbb{E}(\bar{\sigma}_{n}^{2}) \right\| &\leq \left\| \hat{\sigma}_{n}^{2} - \mathbb{E}(\hat{\sigma}_{n}^{2}) \right\| + H_{1} + H_{2} \\ &= \left\| \hat{\sigma}_{n}^{2} - \mathbb{E}(\hat{\sigma}_{n}^{2}) \right\| + O\left(n^{\psi + \max\{q(\psi - \theta), 0\} - 1} \right), \end{split}$$

which implies

$$\left\|\hat{\sigma}_{n}^{2} - \mathbb{E}(\hat{\sigma}_{n}^{2})\right\| \ge \left\|\bar{\sigma}_{n}^{2} - \mathbb{E}(\bar{\sigma}_{n}^{2})\right\| + O\left(n^{\psi + \max\{q(\psi-\theta), 0\} - 1}\right).$$
(B.13)

Check that $2\psi + \max\{2q(\psi - \theta), 0\} - 2 < \psi + \max\{2q(\psi - \theta), 0\} - 1$ always holds under Definition 1 because $\psi < 1/(1+q) < 1$. Hence, it suffices to show that

$$\left\|\bar{\sigma}_{n}^{2} - \mathbb{E}(\bar{\sigma}_{n}^{2})\right\|^{2} \sim \mathscr{V}_{\psi,\Psi,\theta,\Theta,q,\phi} \sigma^{4} n^{\psi + \max\{2q(\psi-\theta),0\}-1}.$$
(B.14)

Without loss of generality, assume $\mu = 0$ for the remaining of this proof. Otherwise, consider the demeaned series $\{X_i - \mu\}$.

B.2.2 Approximation by Blocking

When $\phi > 1$, we can partition the ramped subsamples using (B.8) to (B.10) into s_n blocks:

$$\bar{\sigma}_n^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 + \frac{2}{n} \sum_{i=2}^n \sum_{j=1}^{i-1} W_n(i,j) X_i X_j$$
$$= \frac{1}{n} \sum_{i=1}^n X_i^2 + \frac{2}{n} \sum_{i=2}^n \sum_{k=1}^{s'_i} \left(1 - \frac{k^q}{t_n^q}\right) X_i X_{i-k}$$
$$= \frac{2}{n} \sum_{h=1}^{s_n-1} \sum_{i=1}^{\nu_h} \sum_{j=\eta_h+(i-1)\varpi_h}^{\eta_h+i\varpi_h-1} \sum_{k=1}^{j-\eta_h-(i-1)\varpi_h+h} \left(1 - \frac{k^q}{t_n^q}\right) X_j X_{j-k}$$

$$+ \frac{1}{n} \left\{ \sum_{i=1}^{n} X_{i}^{2} + 2 \sum_{j=\eta_{s_{n}}}^{n} \sum_{k=1}^{s'_{j}} \left(1 - \frac{k^{q}}{t_{n}^{q}} \right) X_{j} X_{j-k} \right\}$$

=: $\bar{\sigma}_{1,n}^{2} + \bar{\sigma}_{2,n}^{2}$, (B.15)

where $\bar{\sigma}_{2,n}^2$ is constructed from the sum of squared terms and the last incomplete block. Then, we can use the same procedure from (B.11) to (B.14) to argue that $\bar{\sigma}_{2,n}^2$ is asymptotically negligible in finding the variance. First, by Minkowski inequality,

$$\|\bar{\sigma}_{n}^{2} - \mathbb{E}(\bar{\sigma}_{n}^{2})\| \leq \|\bar{\sigma}_{1,n}^{2} - \mathbb{E}(\bar{\sigma}_{1,n}^{2})\| + \|\bar{\sigma}_{2,n}^{2} - \mathbb{E}(\bar{\sigma}_{2,n}^{2})\|.$$

In view of Lemma E.6,

$$\|\bar{\sigma}_{2,n}^2 - \mathbb{E}(\bar{\sigma}_{2,n}^2)\|^2 = o(n^{\psi + \max\{2q(\psi-\theta), 0\}-1}).$$

Repeating the same arguments in (B.12) and (B.13), it remains to prove

$$\left\|\bar{\sigma}_{1,n}^2 - \mathbb{E}(\bar{\sigma}_{1,n}^2)\right\|^2 \sim \mathscr{V}_{\psi,\Psi,\theta,\Theta,q,\phi} \sigma^4 n^{\psi+\max\{2q(\psi-\theta),0\}-1},$$

for the case $\phi > 1$. If $\phi = 1$, we do not need partition the subsamples as in (B.15) because the subsampling parameter is monotonically increasing instead of being ramped. In that case, we consider

$$\bar{\sigma}_{n}^{2} = \frac{1}{n} \sum_{i=1}^{n} X_{i}^{2} + \frac{2}{n} \sum_{i=2}^{n} \sum_{k=1}^{s_{i}'} \left(1 - \frac{k^{q}}{t_{n}^{q}}\right) X_{i} X_{i-k}$$

$$= \frac{2}{n} \sum_{h=1}^{s_{n}} \sum_{i=\eta_{h}}^{n} \left(1 - \frac{h^{q}}{t_{n}^{q}}\right) X_{i} X_{i-h} + \frac{1}{n} \sum_{i=1}^{n} X_{i}^{2}$$

$$=: \bar{\sigma}_{3,n}^{2} + \bar{\sigma}_{4,n}^{2}, \qquad (B.16)$$

where $\bar{\sigma}_{4,n}^2$ is constructed from the sum of squared terms only. Since $\|\bar{\sigma}_{4,n}^2 - \mathbb{E}(\bar{\sigma}_{4,n}^2)\|^2 = O(n^{-1})$ by Lemma E.6, we can similarly argue that $\bar{\sigma}_{4,n}^2$ is asymptotically negligible in finding the variance.

With slight abuse of notation, we only write $\bar{\sigma}_n^2$ in Sections B.2.3 and B.2.4 for tidiness. This is possible because the results on $\bar{\sigma}_n^2$ also apply to $\bar{\sigma}_{1,n}^2$ and $\bar{\sigma}_{3,n}^2$, which is equivalent to say that the last incomplete block or sum of squared terms does not exist. We will separate the two cases when we try to find $\mathcal{V}_{\psi,\Psi,\theta,\Theta,q,\phi}$.

B.2.3 Approximation by *m*-dependent Process

Recall the definition of *m*-dependent process approximated version $\tilde{\sigma}_n^2$ in (B.4). We are going to approximate $\|\bar{\sigma}_n^2 - \mathbb{E}(\bar{\sigma}_n^2)\|$ by $\|\tilde{\sigma}_n^2 - \mathbb{E}(\tilde{\sigma}_n^2)\|$. To this end, we can use the same procedure from (B.11) to (B.14) again. By Minkowski inequality,

$$\left\|\bar{\sigma}_n^2 - \mathbb{E}(\bar{\sigma}_n^2)\right\| \le \left\|\tilde{\sigma}_n^2 - \mathbb{E}(\tilde{\sigma}_n^2)\right\| + \left\|\bar{\sigma}_n^2 - \mathbb{E}(\bar{\sigma}_n^2) - \tilde{\sigma}_n^2 + \mathbb{E}(\tilde{\sigma}_n^2)\right\|.$$

Since $\psi > 0$, we can always choose $m = o(n^{\psi/6})$ such that $m \to \infty$ as $n \to \infty$. Then, $d_{m,\alpha} = o(1)$ as $n \to \infty$ by Lemma E.5, and $G_{5,n} = O(n^{\psi+\max\{2q(\psi-\theta),0\}})$ by (C.3), where $d_{m,\alpha}$ and $G_{5,n}$ are defined in (E.4) and Lemma E.2, respectively. For $\alpha \ge 4$, Lemma E.2 states that

$$\left\|\bar{\sigma}_{n}^{2} - \mathbb{E}(\bar{\sigma}_{n}^{2}) - \tilde{\sigma}_{n}^{2} + \mathbb{E}(\tilde{\sigma}_{n}^{2})\right\| = O\left(n^{-\frac{1}{2}}d_{m,\alpha}G_{5,n}^{\frac{1}{2}}\right) = o\left(n^{\frac{\psi}{2} + \max\{q(\psi-\theta), 0\} - \frac{1}{2}}\right).$$

Repeating the same arguments as in (B.12) and (B.13), it remains to show that

$$\left\|\tilde{\sigma}_{n}^{2} - \mathbb{E}(\tilde{\sigma}_{n}^{2})\right\|^{2} \sim \mathcal{V}_{\psi,\Psi,\theta,\Theta,q,\phi} \sigma^{4} n^{\psi + \max\{2q(\psi-\theta),0\}-1}$$

B.2.4 Approximation by Martingale Difference

Recall the definition of martingale difference approximated version $\check{\sigma}_n^2$ in (B.7). We can use the same procedure from (B.11) to (B.14) again to approximate $\| \tilde{\sigma}_n^2 - \mathbb{E}(\tilde{\sigma}_n^2) \|$ by $\| \check{\sigma}_n^2 - \mathbb{E}(\check{\sigma}_n^2) \|$. By Minkowski inequality,

$$\left\|\tilde{\sigma}_n^2 - \mathbb{E}(\tilde{\sigma}_n^2)\right\| \le \left\|\check{\sigma}_n^2 - \mathbb{E}(\check{\sigma}_n^2)\right\| + \left\|\tilde{\sigma}_n^2 - \mathbb{E}(\tilde{\sigma}_n^2) - \check{\sigma}_n^2 + \mathbb{E}(\check{\sigma}_n^2)\right\|.$$

Note that $G_{6,n} = O(n^{\max\{q(\psi-\theta),0\}})$ by (C.5), where $G_{6,n}$ is defined in Lemma E.3. In Section B.2.3, we have chosen $m = o(n^{\psi/6})$. It follows from Lemma E.3 that

$$\|\tilde{\sigma}_{n}^{2} - \mathbb{E}(\tilde{\sigma}_{n}^{2}) - \breve{\sigma}_{n}^{2} + \mathbb{E}(\breve{\sigma}_{n}^{2})\| = O\left(m^{\frac{5}{2}}n^{-\frac{1}{2}}G_{6,n}\right) = o\left(n^{\frac{\psi}{2} + \max\{q(\psi-\theta), 0\} - \frac{1}{2}}\right),$$

as $\alpha \ge 4$ implies $\alpha' = 2$. Repeating the same arguments as in (B.12) and (B.13), it suffices to prove

$$\left\|\breve{\sigma}_n^2 - \mathbb{E}(\breve{\sigma}_n^2)\right\|^2 \sim \mathscr{V}_{\psi,\Psi,\theta,\Theta,q,\phi} \sigma^4 n^{\psi + \max\{2q(\psi-\theta),0\}-1}.$$

B.2.5 Exact Convergence Rate of Variance

Consider the case $\phi > 1$. We similarly define $\breve{\sigma}_{1,n}^2$ as in (B.15) with X_j replaced by martingale difference D_j . By Section B.2.2, we can work on $\|\breve{\sigma}_{1,n}^2 - \mathbb{E}(\breve{\sigma}_{1,n}^2)\|$ instead of $\|\breve{\sigma}_n^2 - \mathbb{E}(\breve{\sigma}_n^2)\|$. Then,

by $\mathbb{E}(D_i D_j) = 0$ when $i \neq j$ (see Section E.4) and Minkowski inequality,

$$\begin{split} \|\breve{\sigma}_{1,n}^{2} - \mathbb{E}(\breve{\sigma}_{1,n}^{2})\| \\ &= \frac{2}{n} \left\| \sum_{h=1}^{s_{n-1}} \sum_{i=1}^{\nu_{h}} \sum_{j=\eta_{h}+(i-1)\varpi_{h}}^{\eta_{h}+i\varpi_{h}-1} \sum_{k=1}^{j-\eta_{h}-(i-1)\varpi_{h}+h} \left(1 - \frac{k^{q}}{t_{n}^{q}}\right) D_{j} D_{j-k} \right\| \\ &\leq \frac{2}{n} \left\| \sum_{h=\lceil(m+1)/\phi\rceil}^{s_{n}-1} \sum_{j=\eta_{h}+(i-1)\varpi_{h}+m+1}^{\nu_{h}} \sum_{k=m+1}^{j-\eta_{h}-(i-1)\varpi_{h}+h} \left(1 - \frac{k^{q}}{t_{n}^{q}}\right) D_{j} D_{j-k} \right\| \\ &+ \frac{2}{n} \left\| \sum_{h=1}^{s_{n}-1} \sum_{i=1}^{\nu_{h}} \sum_{j=\eta_{h}+(i-1)\varpi_{h}}^{\eta_{h}+i\varpi_{h}-1} \min\{j-\eta_{h}-(i-1)\varpi_{h}+h,m\}} \sum_{k=1}^{(1-1)m} \left(1 - \frac{k^{q}}{t_{n}^{q}}\right) D_{j} D_{j-k} \right\| \\ &=: H_{3} + H_{4}, \end{split}$$
(B.17)

where *m* is the ancillary variable used in *m*-dependent process approximation. In Section B.2.3, we have chosen $m = o(n^{\psi/6})$, which means $m = o(s_n)$ under Definition 1. As the physical meaning of $j - \eta_h - (i-1)\varpi_h + h$ in the above is the ramped subsample size at the *j*-th observation, $m = o(s_n)$ implies that $\min\{j - \eta_h - (i-1)\varpi_h + h - 1, m\} = m$ most of the time. Consequently, $H_4 = o(H_3)$. Repeating the same arguments as in (B.12) and (B.13), we only need to prove

$$H_3^2 \sim \mathscr{V}_{\psi,\Psi,\theta,\Theta,q,\phi} \sigma^4 n^{\psi+\max\{2q(\psi-\theta),0\}-1}.$$
(B.18)

For $\phi = 1$, we can similarly define $\breve{\sigma}_{3,n}^2$ as in (B.16) so that

$$\begin{aligned} \left\| \breve{\sigma}_{3,n}^{2} - \mathbb{E}(\breve{\sigma}_{3,n}^{2}) \right\| \\ &= \frac{2}{n} \left\| \sum_{h=1}^{s_{n}} \sum_{i=\eta_{h}}^{n} \left(1 - \frac{h^{q}}{t_{n}^{q}} \right) D_{i} D_{i-h} \right\| \\ &\leq \frac{2}{n} \left\| \sum_{h=m+1}^{s_{n}} \sum_{i=\eta_{h}}^{n} \left(1 - \frac{h^{q}}{t_{n}^{q}} \right) D_{i} D_{i-h} \right\| + \frac{2}{n} \left\| \sum_{h=1}^{m} \sum_{i=\eta_{h}}^{n} \left(1 - \frac{h^{q}}{t_{n}^{q}} \right) D_{i} D_{i-h} \right\| \\ &=: H_{5} + H_{6}. \end{aligned}$$
(B.19)

Repeating the arguments from (B.17) to (B.18), we only need to prove

$$H_5^2 \sim \mathscr{V}_{\psi,\Psi,\theta,\Theta,q,\phi} \sigma^4 n^{\psi+\max\{2q(\psi-\theta),0\}-1}.$$
(B.20)

Lemma E.7 confirms (B.18) and (B.20).

B.2.6 Exact Convergence Rate of Bias

Now, we find the exact convergence rate of bias. To approximate $\text{Bias}(\hat{\sigma}_n^2)$ by $\text{Bias}(\bar{\sigma}_n^2)$, we use the same procedure from (B.11) to (B.14). Note that

$$\operatorname{Bias}(\hat{\sigma}_n^2) = \operatorname{Bias}(\bar{\sigma}_n^2) + \mathbb{E}(\hat{\sigma}_n^2) - \mathbb{E}(\bar{\sigma}_n^2).$$

By (B.11),

$$|\mathbb{E}(\hat{\sigma}_n^2) - \mathbb{E}(\bar{\sigma}_n^2)| = H_2 = O\left(n^{\psi + \max\{q(\psi - \theta), 0\} - 1}\right).$$

Check that $\psi - 1 < -\psi$ and $\psi + q(\psi - \theta) - 1 < -q\theta$ always hold under Definition 1 because $\psi < 1/(1+q) \le 1/2$. Repeating the same arguments as in (B.12) and (B.13), we can assume $\mu = 0$, and it suffices to prove

$$\operatorname{Bias}(\bar{\sigma}_n^2) \sim \begin{cases} o(n^{-\psi}), & \psi < \theta; \\ -\Theta^{-q} n^{-q\theta} v_q, & \psi \ge \theta. \end{cases}$$

To better illustrate the idea, we consider the case $\phi = 1$ first. Recall that we can rearrange the order of summations in (B.16). Taking expectation yields

$$\mathbb{E}(\bar{\sigma}_{n}^{2}) = \frac{1}{n} \sum_{i=1}^{n} \gamma_{0} + \frac{2}{n} \sum_{h=1}^{s_{n}} \sum_{i=\eta_{h}}^{n} \left(1 - \frac{h^{q}}{t_{n}^{q}}\right) \gamma_{h}$$
$$= \gamma_{0} + 2 \sum_{h=1}^{s_{n}} \frac{n - \eta_{h} + 1}{n} \left(1 - \frac{h^{q}}{t_{n}^{q}}\right) \gamma_{h}$$
$$\sim \gamma_{0} + 2 \sum_{h=1}^{s_{n}} \left\{1 - \frac{h^{q}}{t_{n}^{q}} - \left(\frac{h}{s_{n}}\right)^{\frac{1}{\psi}} \left(1 - \frac{h^{q}}{t_{n}^{q}}\right)\right\} \gamma_{h}, \tag{B.21}$$

where we have used the order of η_h in (B.10) to arrive at (B.21). As $n \to \infty$, $\gamma_0 + 2\sum_{h=1}^{s_n} \gamma_h \to \sigma^2$. On the other hand, by Kronecker's lemma and Assumption 5 that $u_q = \sum_{j \in \mathbb{Z}} |j|^q |\gamma_j| < \infty$,

$$\begin{split} \sum_{h=1}^{s_n} \left(\frac{h}{s_n}\right)^{\frac{1}{\psi}} \gamma_h &= \frac{1}{s_n^q} \sum_{h=1}^{s_n} \frac{h^{1/\psi}}{s_n^{1/\psi-q}} \gamma_h \\ &= \frac{1}{s_n^q} \sum_{h=1}^{\lceil s_n^{1-q\psi} \rceil} \frac{h^{1/\psi}}{\left(s_n^{1-q\psi}\right)^{1/\psi}} \gamma_h + \frac{1}{s_n^q} \sum_{h=\lceil s_n^{1-q\psi} \rceil+1}^{s_n} \frac{\left(h^{1-q\psi}\right)^{1/\psi}}{\left(s_n^{1-q\psi}\right)^{1/\psi}} h^q \gamma_h \\ &= \frac{1}{s_n^q} o(1) + \frac{1}{s_n^q} o(1) = o(s_n^{-q}). \end{split}$$

Check that $1 - q\psi > 0$ always holds under Definition 1. Similarly,

$$\sum_{h=1}^{s_n} \left(\frac{h}{s_n}\right)^{\frac{1}{\psi}} \frac{h^q}{t_n^q} \gamma_h = \frac{1}{t_n^q} \sum_{h=1}^{s_n} \left(\frac{h}{s_n}\right)^{\frac{1}{\psi}} h^q \gamma_h$$
$$= \frac{1}{t_n^q} o(1) = o(t_n^{-q}).$$
(B.22)

Consequently, (B.21) becomes

$$\mathbb{E}(\bar{\sigma}_n^2) \sim \begin{cases} \sigma^2 - 2\sum_{h=1}^{s_n} (h/s_n)^{\frac{1}{\psi}} \gamma_h + O(t_n^{-q}), & \psi < \theta; \\ \sigma^2 - 2t_n^{-q} \sum_{h=1}^{s_n} h^q \gamma_h + o(t_n^{-q}), & \psi \ge \theta \end{cases}$$
$$= \begin{cases} \sigma^2 + o(n^{-q\psi}), & \psi < \theta; \\ \sigma^2 - \Theta^{-q} n^{-q\theta} v_q + o(n^{-q\theta}), & \psi \ge \theta. \end{cases}$$

For $\phi > 1$, note that the partitions in (B.15) cannot give a form like (B.21) after taking expectation. By rearranging the order of summations, approximating the sums by Riemann integrals and using the order of ν_i in (B.8),

$$\begin{split} \mathbb{E}(\bar{\sigma}_{n}^{2}) \\ &= \gamma_{0} + \frac{2}{n} \sum_{h=1}^{s_{n}-1} \sum_{i=1}^{\nu_{h}} \sum_{j=\eta_{h}+(i-1)\varpi_{h}}^{\eta_{h}+i\varpi_{h}-1} \sum_{k=1}^{j-\eta_{h}-(i-1)\varpi_{h}+h} \left(1 - \frac{k^{q}}{t_{n}^{q}}\right) \gamma_{k} + \frac{2}{n} \sum_{j=\eta_{sn}}^{n} \sum_{k=1}^{s'_{j}} \left(1 - \frac{k^{q}}{t_{n}^{q}}\right) \gamma_{k} \\ &\sim \gamma_{0} + \frac{2}{n} \sum_{k=1}^{\left\lceil \phi s_{n} \right\rceil} \gamma_{k} \int_{k/\phi}^{s_{n}} \int_{0}^{\nu_{h}} \int_{\max(k-h,0)}^{(\phi-1)h} \left(1 - \frac{k^{q}}{t_{n}^{q}}\right) dj di dh \\ &= \gamma_{0} + \frac{2}{n} \sum_{h=1}^{\left\lceil \phi s_{n} \right\rceil} \gamma_{h} \int_{h/\phi}^{s_{n}} \int_{0}^{\nu_{i}} \int_{\max(h-i,0)}^{(\phi-1)i} \left(1 - \frac{h^{q}}{t_{n}^{q}}\right) dk dj di \\ &= \gamma_{0} + \frac{2}{n} \sum_{h=1}^{\left\lceil \phi s_{n} \right\rceil} \left(1 - \frac{h^{q}}{t_{n}^{q}}\right) \gamma_{h} \int_{h/\phi}^{s_{n}} \nu_{i} \{(\phi - 1)i - \max(h - i, 0)\} di \\ &\sim \gamma_{0} + \frac{2}{(\phi - 1)\psi} \left(\frac{1}{s_{n}}\right)^{\frac{1}{\psi}} \sum_{h=1}^{s_{n}} \left(1 - \frac{h^{q}}{t_{n}^{q}}\right) \gamma_{h} \left\{\int_{h/\phi}^{h} (\phi i - h)i^{\frac{1}{\psi}-2} di + \int_{h}^{s_{n}} (\phi - 1)i^{\frac{1}{\psi}-1} di\right\} \\ &+ \frac{2}{(\phi - 1)\psi} \left(\frac{1}{s_{n}}\right)^{\frac{1}{\psi}} \sum_{h=s_{n}+1}^{\left\lceil \phi s_{n} \right\rceil} \left(1 - \frac{h^{q}}{t_{n}^{q}}\right) \gamma_{h} \int_{h/\phi}^{s_{n}} (\phi i - h)i^{\frac{1}{\psi}-2} di. \end{split}$$

Evaluating the integrals, we have

$$\mathbb{E}(\bar{\sigma}_{n}^{2}) \sim \gamma_{0} + 2\sum_{h=1}^{s_{n}} \left(1 - \frac{h^{q}}{t_{n}^{q}}\right) \left\{ 1 - \frac{\psi\left(\phi^{1-1/\psi} - 1\right)}{(\phi - 1)\left(\psi - 1\right)} \left(\frac{h}{s_{n}}\right)^{\frac{1}{\psi}} \right\} \gamma_{h} + 2\sum_{h=s_{n}+1}^{\left\lceil \phi s_{n} \right\rceil} \left(1 - \frac{h^{q}}{t_{n}^{q}}\right) \left\{ \frac{\phi}{\phi - 1} + \frac{1}{(\phi - 1)(\psi - 1)} \cdot \frac{h}{s_{n}} - \frac{\phi^{1-1/\psi}\psi}{(\phi - 1)\left(\psi - 1\right)} \left(\frac{h}{s_{n}}\right)^{\frac{1}{\psi}} \right\} \gamma_{h}.$$
(B.23)

Repeating the arguments from (B.21) to (B.22), we have

$$\mathbb{E}(\bar{\sigma}_n^2) \sim \begin{cases} \sigma^2 + o(n^{-q\psi}), & \psi < \theta; \\ \sigma^2 - \Theta^{-q} n^{-q\theta} v_q + o(n^{-q\theta}), & \psi \ge \theta, \end{cases}$$

which completes the proof.

C Proof of Corollaries

C.1 Proof of Corollary 1

By Theorem 1, it suffices to verify Assumptions 2 and 3 under Definition 1. We find the order of $G_{1,n}$ and $G_{2,n}$ first. For $G_{1,n}$, we have

$$G_{1,n} = \max_{1 \le i \le n} \sum_{j=1}^{n} \left\{ \left| 1 - \frac{|i-j|^q}{t_n^q} \right| \mathbb{1}_{|i-j| \le s'_{i\vee j}} + \left(1 - \frac{|i-j|^q}{t_n^q} \right)^2 \mathbb{1}_{|i-j| \le s'_{i\vee j}} \right\}$$

$$\leq \max_{1 \le i \le n} \sum_{j=1}^{n} \left| 1 - \frac{|i-j|^q}{t_n^q} \right| \mathbb{1}_{|i-j| \le \phi s_n} + \max_{1 \le i \le n} \sum_{j=1}^{n} \left(1 - \frac{|i-j|^q}{t_n^q} \right)^2 \mathbb{1}_{|i-j| \le \phi s_n}$$
(C.1)
$$\leq 2 \sum_{k=0}^{\phi s_n} \left| 1 - \frac{k^q}{t_n^q} \right| + 2 \sum_{k=0}^{\phi s_n} \left(1 - \frac{k^q}{t_n^q} \right)^2$$

$$\leq O(1) \cdot \int_0^{\phi s_n} \left| 1 - \frac{k^q}{t_n^q} \right| \,\mathrm{d}k + O(1) \cdot \int_0^{\phi s_n} \left(1 - \frac{k^q}{t_n^q} \right)^2 \,\mathrm{d}k \tag{C.2}$$

$$\leq O(n^{\psi} + n^{q(\psi-\theta)+\psi}) + O(n^{\psi} + n^{2q(\psi-\theta)+\psi}) = O(n^{\psi+\max\{2q(\psi-\theta),0\}}),$$
(C.3)

where (C.1) follows from the fact that $s'_{i\vee j} \leq \phi s_n$ for $1 \leq i, j \leq n$, and (C.2) follows from approximating the sums by Riemann integrals. When $\psi \leq \theta$, check that $\psi + \max\{2q(\psi - \theta), 0\} < 2 - 2/\alpha'$. When $\psi > \theta$, we have

$$\psi + \max\{2q(\psi - \theta), 0\} < \psi - 2q \cdot \frac{\psi - 2 + 2/\alpha'}{2q} = 2 - \frac{2}{\alpha'}.$$

For $G_{2,n}$, observe that

$$G_{2,n} = \max_{1 \le i,j \le n} |W_n(i,j)| + \max_{1 \le i \le n} \left\{ \sum_{j=2}^n |W_n(i,j) - W_n(i,j-1)|^{\alpha'} \right\}^{\frac{1}{\alpha'}} \\ \le \max\left(\left| 1 - \frac{\phi^q s_n^q}{t_n^q} \right|, 1 \right) + 2 \sum_{k=0}^{\phi s_n} \left| \frac{(k+1)^q - k^q}{t_n^q} \right| \\ \le O(n^{\max\{q(\psi-\theta), 0\}}) + O(1) \cdot \int_0^{\phi s_n} \frac{k^{q-1}}{t_n^q} \, \mathrm{d}k$$

$$(C.4)$$

$$= O(n^{\max\{q(\psi-\theta),0\}} + n^{q(\psi-\theta)}) = O(n^{\max\{q(\psi-\theta),0\}}),$$
(C.5)

where (C.4) follows from approximating the sum by a Riemann integral. When $\psi \leq \theta$, check that $c \in (0, 1 - 1/\alpha')$ satisfies $\max\{q(\psi - \theta), 0\} = 0 < c$. When $\psi > \theta$, there exists $c \in (1 - 1/\alpha' - \psi/2, 1 - 1/\alpha')$ such that

$$\max\{q(\psi-\theta),0\} < -q \cdot \frac{\psi-2+2/\alpha'}{2q} < c.$$

Now we verify Assumption 3. It is clear that $W_n(i,j) = W_n(j,i)$ and $W_n(i,i) = 1$ for all $n \in \mathbb{Z}^+$ and $i, j \in \{1, ..., n\}$ under Definition 1. In addition, note that

$$\begin{split} \max_{1 \le k \le b_n} |w_{n,k} - 1| &= \max_{1 \le k \le b_n} \left| \frac{1}{n} \sum_{i=k+1}^n \left(1 - \frac{k^q}{t_n^q} \right) \mathbb{1}_{k \le s'_i} - 1 \right| \\ &= \max_{1 \le k \le b_n} \left| -\frac{1}{n} \sum_{i=k+1}^n \frac{k^q}{t_n^q} \mathbb{1}_{k \le s'_i} - \frac{1}{n} \sum_{i=k+1}^n \mathbb{1}_{k > s'_i} - \frac{k}{n} \right| \\ &\leq \max_{1 \le k \le b_n} \left| \frac{1}{n} \sum_{i=k+1}^n \frac{k^q}{t_n^q} \mathbb{1}_{k \le s'_i} \right| + \frac{1}{n} \sum_{i=1}^n \mathbb{1}_{b_n > s'_i} + \frac{b_n}{n}, \\ &\max_{b_n < k < n} |w_{n,k}| = \max_{b_n < k < n} \left| \frac{1}{n} \sum_{i=k+1}^n \frac{k^q}{t_n^q} \mathbb{1}_{k \le s'_i} \right| + \frac{1}{n} \sum_{i=1}^n \mathbb{1}_{b_n \le s'_i}. \end{split}$$

When $\psi \leq \theta$, note that $s'_i \geq Cn^{2\psi/3}$ for some $C \in \mathbb{R}^+$ and $i > n^{2/3}$. We can choose $b_n = O(n^{\psi/2})$ such that

$$\frac{1}{n}\sum_{i=1}^{n}\mathbb{1}_{b_n>s'_i} \leq \frac{n^{2/3}}{n} + \frac{1}{n}\sum_{n^{2/3}< i\leq n}\mathbb{1}_{b_n>Cn^{2\psi/3}} = o(1).$$

Therefore,

$$\max_{1 \le k \le b_n} \left| \frac{1}{n} \sum_{i=k+1}^n \frac{k^q}{t_n^q} \mathbb{1}_{k \le s_i'} \right| + \frac{1}{n} \sum_{i=1}^n \mathbb{1}_{b_n > s_i'} + \frac{b_n}{n} \le \frac{1}{n} \sum_{i=1}^n \frac{b_n^q}{t_n^q} + o(1) + o(1) = o(1)$$
$$\max_{b_n < k < n} \left| \frac{1}{n} \sum_{i=k+1}^n \frac{k^q}{t_n^q} \mathbb{1}_{k \le s_i'} \right| + \frac{1}{n} \sum_{i=1}^n \mathbb{1}_{b_n \le s_i'} \le \frac{\phi^q}{n} \sum_{i=1}^n \frac{s_n^q}{t_n^q} + O(1) = O(1).$$

When $\psi > \theta$, Assumption 3(c) does not hold. However, Assumption 3(c) is only used to show that $|\mathbb{E}(\bar{\sigma}_n^2) - \sigma^2| = o(1)$ in Section B.1.4. We prove the same thing under Definition 1(b) and Assumption 4. Consider the case $\phi = 1$, by (B.21) we have

$$\mathbb{E}(\bar{\sigma}_n^2) \sim \sigma^2 - 2\sum_{h=1}^{s_n} \frac{h^q}{t_n^q} \gamma_h - 2\sum_{h=1}^{s_n} \left(\frac{h}{s_n}\right)^{\frac{1}{\psi}} \left(1 - \frac{h^q}{t_n^q}\right) \gamma_h.$$

Since $u_{\tilde{q}} < \infty$ under Assumption 4, Kronecker's lemma gives

$$\begin{split} \sum_{h=1}^{s_n} \frac{h^q}{t_n^q} \gamma_h &= \frac{s_n^{q-\tilde{q}}}{t_n^q} \sum_{h=1}^{s_n} \frac{h^{q-\tilde{q}}}{s_n^{q-\tilde{q}}} h^{\tilde{q}} \gamma_h = O(n^{q(\psi-\theta)-\tilde{q}\psi}) o(1) = o(n^{q(\psi-\theta)-\tilde{q}\psi}), \\ \sum_{h=1}^{s_n} \left(\frac{h}{s_n}\right)^{\frac{1}{\psi}} \gamma_h &= \frac{1}{s_n^{\tilde{q}}} \sum_{h=1}^{s_n} \frac{h^{1/\psi}}{s_n^{1/\psi-\tilde{q}}} \gamma_h \\ &= \frac{1}{s_n^{\tilde{q}}} \sum_{h=1}^{s_n^{n-\tilde{q}\psi}} \frac{h^{1/\psi}}{(s_n^{1-\tilde{q}\psi})^{1/\psi}} \gamma_h + \frac{1}{s_n^{\tilde{q}}} \sum_{h=|s_n^{1-\tilde{q}\psi}|+1}^{s_n} \frac{(h^{1-\tilde{q}\psi})^{1/\psi}}{(s_n^{1-\tilde{q}\psi})^{1/\psi}} h^{\tilde{q}} \gamma_h \\ &= O(n^{-\tilde{q}\psi}) o(1) + O(n^{-\tilde{q}\psi}) o(1) = o(n^{-\tilde{q}\psi}), \\ \\ \sum_{h=1}^{s_n} \left(\frac{h}{s_n}\right)^{\frac{1}{\psi}} \frac{h^q}{t_n^q} \gamma_h = \frac{s_n^{q-\tilde{q}}}{t_n^q} \sum_{h=1}^{s_n} \frac{h^{q-\tilde{q}+1/\psi}}{s_n^{q-\tilde{q}+1/\psi}} h^{\tilde{q}} \gamma_h = o(n^{q(\psi-\theta)-\tilde{q}\psi}). \end{split}$$

Check that

$$q(\psi - \theta) - \tilde{q}\psi < 0 \iff \frac{q - \tilde{q}}{q}\psi < \theta,$$

which is always true under Definition 1(b). The case $\phi > 1$ can be similarly proved using (B.23). Therefore, we have $|\mathbb{E}(\bar{\sigma}_n^2) - \sigma^2| = o(1)$, which completes the proof.

C.2 Proof of Corollary 2

We try to find Ψ_{\star} first assuming we do not separate the smoothing parameters, i.e., $\Psi = \Theta$. If $\psi = \theta = 1/(1+2q)$, then as $n \to \infty$, the standardized AMSE is given by

$$\begin{split} &n^{2q/(1+2q)}\mathrm{MSE}(\hat{\sigma}_n^2)\\ &\rightarrow \begin{cases} \frac{\kappa_q^2}{\Psi^{2q}} + \frac{\Psi(\phi+1)(2q+1)}{q+1} - \frac{8\Psi(\phi^{q+2}-1)(2q+1)}{(\phi-1)(q+1)(q+2)(3q+2)} + \frac{\Psi(\phi^{2q+2}-1)}{(\phi-1)(q+1)(2q+1)}, &\phi>1;\\ \frac{\kappa_q^2}{\Psi^{2q}} + \frac{2\Psi(2q+1)}{q+1} - \frac{8\Psi(2q+1)}{(q+1)(3q+2)} + \frac{2\Psi}{2q+1}, &\phi=1. \end{cases}$$

Differentiating the standardized AMSE with respect to Ψ , we can derive the minimizer Ψ_{\star} easily:

$$\Psi_{\star} = \begin{cases} \left\{ \frac{(\phi+1)(2q+1)}{2q(q+1)} - \frac{4(\phi^{q+2}-1)(2q+1)}{(\phi-1)q(q+1)(q+2)(3q+2)} + \frac{(\phi^{2q+2}-1)}{2(\phi-1)q(q+1)(2q+1)} \right\}^{-\frac{1}{1+2q}} \kappa_q^{\frac{2}{1+2q}}, \quad \phi > 1; \\ \left\{ \frac{2q+1}{q(q+1)} - \frac{4(2q+1)}{q(q+1)(3q+2)} + \frac{1}{q(2q+1)} \right\}^{-\frac{1}{1+2q}} \kappa_q^{\frac{2}{1+2q}}, \qquad \phi = 1. \end{cases}$$

When the smoothing parameters are separated, we can allow $\Theta = \rho \Psi_{\star}$, where $\rho \in \mathbb{R}^+$, to view Θ as an improvement to the optimal value without separation of parameters. Consequently, we find ρ that minimize

$$\begin{split} &n^{2q/(1+2q)} \mathrm{MSE}(\hat{\sigma}_n^2) \\ & \rightarrow \begin{cases} \frac{\kappa_q^2}{\rho^{2q} \Psi_*^{2q}} + \frac{\Psi_\star(\phi+1)(2q+1)}{q+1} - \frac{8\rho^{-q} \Psi_\star(\phi^{q+2}-1)(2q+1)}{(\phi-1)(q+1)(q+2)(3q+2)} + \frac{\rho^{-2q} \Psi_\star(\phi^{2q+2}-1)}{(\phi-1)(q+1)(2q+1)}, & \phi > 1; \\ \frac{\kappa_q^2}{\rho^{2q} \Psi_\star^{2q}} + \frac{2\Psi_\star(2q+1)}{q+1} - \frac{8\rho^{-q} \Psi_\star(2q+1)}{(q+1)(3q+2)} + \frac{2\rho^{-2q} \Psi_\star}{2q+1}, & \phi = 1. \end{cases}$$

Differentiating the above with respect to ρ , the minimizer is

$$\rho_{\star} = \begin{cases} \left\{ \frac{(q+2)(3q+2)(\phi^{2q+2}-1)}{4(2q+1)^2(\phi^{q+2}-1)} + \frac{\Psi_{\star}^{-2q-1}\kappa_q^2(\phi-1)(q+1)(q+2)(3q+2)}{4(2q+1)(\phi^{q+2}-1)} \right\}^{\frac{1}{q}}, \quad \phi > 1; \\ \left\{ \frac{(q+1)(3q+2)}{2(2q+1)^2} + \frac{\Psi_{\star}^{-2q-1}\kappa_q^2(q+1)(3q+2)}{4(2q+1)} \right\}^{\frac{1}{q}}, \quad \phi = 1. \end{cases}$$

Note that the κ_q^2 from Ψ_{\star}^{-2q-1} will cancel with the κ_q^2 in ρ_{\star} . Therefore, the AMSE-optimal Θ is

$$\Theta_{\star} = \begin{cases} \left\{ \frac{(q+2)(3q+2)(\phi^{2q+2}-1)}{4(2q+1)^{2}(\phi^{q+2}-1)} + \frac{\Psi_{\star}^{-2q-1}\kappa_{q}^{2}(\phi-1)(q+1)(q+2)(3q+2)}{4(2q+1)(\phi^{q+2}-1)} \right\}^{\frac{1}{q}} \Psi_{\star}, \quad \phi > 1; \\ \left\{ \frac{(q+1)(3q+2)}{2(2q+1)^{2}} + \frac{\Psi_{\star}^{-2q-1}\kappa_{q}^{2}(q+1)(3q+2)}{4(2q+1)} \right\}^{\frac{1}{q}} \Psi_{\star}, \qquad \phi = 1. \end{cases}$$

Using Ψ_{\star} and Θ_{\star} , we can check that the standardized AMSE decreases when ϕ decreases, ceteris paribus. Hence the AMSE-optimal ϕ is $\phi_{\star} = 1$. If O(1)-space update is required, the AMSE-optimal ϕ is $\phi_{\star} = 2$ by Proposition 2.

C.3 Proof of Corollary 3

If $\overline{\ell}_n$ is substituted with $\ell_{i \vee j}$ in (2.7) so that the window becomes

$$W_{\mathsf{PSR}'} = W_{n,\mathsf{PSR}'}(i,j) = \left(1 - \frac{|i-j|}{\ell_{i \lor j}}\right) \mathbb{1}_{|i-j| \le \ell_{i \lor j}},$$

Theorem 2 does not apply directly. Nevertheless, we can use the same procedure as in Section B.2 to prove a similar theorem particularly for $W_{\mathsf{PSR}'}$. After going through the steps, we have

$$n^{1-\lambda} \operatorname{Var}(\hat{\sigma}_{n,\mathsf{PSR}'}^2) \to \frac{4\Lambda}{3(\lambda+1)} \quad \text{and} \quad \operatorname{Bias}(\hat{\sigma}_{n,\mathsf{PSR}'}^2) \sim -\frac{\Lambda^{-1}}{(1-\lambda)} n^{-\lambda} v_1$$

Clearly, $\lambda_{\star} = 1/3$ optimize the order of $\text{MSE}(\hat{\sigma}_{n,\text{PSR}'}^2)$. We can also find that $\Lambda_{\star} = (9/2)^{1/3} \kappa_1^{2/3}$ by differentiating $\text{MSE}(\hat{\sigma}_{n,\text{PSR}'}^2)$ and the improvement in AMSE follows. For the other items, Theorem 2 and Corollary 2 apply directly.

D Proof of Propositions

D.1 Proof of Proposition 1

From (2.3) and (2.5), we have

$$\hat{\sigma}_n^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n) \left\{ \sum_{j=1}^i (2 - \mathbb{1}_{i=j}) T\left(\frac{|i-j|}{t_n(i,j)}\right) S\left(\frac{|i-j|}{s_n(i,j)}\right) (X_j - \bar{X}_n) \right\}.$$

Under the conditions in Proposition 1, $\hat{\sigma}_n^2$ can be further written as

$$\hat{\sigma}_{n}^{2} = \frac{1}{n} \sum_{i=1}^{n} (X_{i} - \bar{X}_{n}) \left\{ \sum_{k=0}^{s_{i}} (2 - \mathbb{1}_{k=0}) \sum_{r=0}^{q} a_{r} \left(\frac{k}{t_{n} t_{i}'} \right)^{r} (X_{i-k} - \bar{X}_{n}) \right\}$$
$$= \frac{1}{n} \sum_{r=0}^{q} \frac{a_{r}}{t_{n}^{r}} \sum_{i=1}^{n} \frac{X_{i}}{(t_{i}')^{r}} (K_{i,r} - k_{i,r} \bar{X}_{n}) - \frac{1}{n} \sum_{r=0}^{q} \frac{a_{r}}{t_{n}^{r}} \sum_{i=1}^{n} \frac{\bar{X}_{n}}{(t_{i}')^{r}} (K_{i,r} - k_{i,r} \bar{X}_{n}), \quad (D.1)$$

where $K_{i,r} = \sum_{k=0}^{s_i} (2-\mathbb{1}_{k=0}) k^r X_{i-k}$ and $k_{i,r} = \sum_{k=0}^{s_i} (2-\mathbb{1}_{k=0}) k^r$. To prove Proposition 1, it suffices to find a way such that $K_{i,r}$ in (D.1) can be computed in O(1) time as $k_{i,r}$ is a special case. To do so, note that

$$K_{i,r} = \sum_{k=0}^{s_i} (2 - \mathbb{1}_{k=0}) k^r X_{i-k} = 0^r X_i + 2 \sum_{j=i-s_i}^{i-1} (i-j)^r X_j$$

Algorithm D.1: Pratical O(1)-time update

[1] input:

(i) n, s – the sample size and subsampling parameter from the last iteration [2] (ii) $\mathscr{C}_{n,r}, r = 0, 1, \dots, q$ – the components for the r-th order term [3] (iii) \vec{x} – the vector storing the last s observations [4] begin [5] Receive X_{n+1} [6] Set n = n + 1[7] Compute s_n, t_n [8] Retrieve $X_{n-1}, \ldots, X_{n-1-\max\{\min(s_n-1,q),0\}}$ and X_{n-s-1} from \vec{x} [9] if $s_n = s$ then [10] Update $\mathscr{C}_{n,r}$ with Prop. A.3 and A.4 for $r = 0, 1, \ldots, q$ [11] Pop X_{n-s} from \vec{x} [12] [13] else Update $\mathscr{C}_{n,r}$ with Prop. A.3 and A.4 for $r = 0, 1, \ldots, q$ [14] Push X_n into \vec{x} [15] Set $s = s_n$ [16] Output the online estimate computed with $\mathscr{C}_{n,r}$, where $r = 0, 1, \ldots, q$ [17]

$$= 0^{r} X_{i} + 2 \sum_{u=0}^{r} {r \choose u} i^{r-u} (-1)^{u} \sum_{j=i-s_{i}}^{i-1} j^{u} X_{j}$$
$$= 0^{r} X_{i} + 2 \sum_{u=0}^{r} {r \choose u} i^{r-u} (-1)^{u} (P_{i-1,u} - P_{i-s_{i}-1,u})$$

where $P_{i,u} = \sum_{j=1}^{i} j^{u} X_{j}$. If we store $P_{i,u}$ for i = 1, ..., n-1 and u = 0, ..., q, we can update $P_{n,u}$ by $P_{n-1,u} + n^{u} X_{n}$ for u = 0, ..., q, and consequently compute $A_{n,r}$ in O(1) time, which completes the proof.

In the remaining of this subsection, we discuss the implications of the conditions in Proposition 1. To begin with, note that (D.1) requires O(n) space as it has to store all $P_{i,u}$'s. Furthermore, to support access of $P_{i,u}$ in O(1) time, an array or hash table of unlimited size is necessary. Otherwise, we may not be able to perform direct access, which is needed for computing $P_{n-1,u} - P_{n-s_n-1,u}$ with unrestricted s_n , or we may need to resize the data structure, which typically costs O(n) time under direct access; see Remark D.1. To perform practical O(1) time update, we can assume $t'_i = 1$ for all $i \in \mathbb{Z}^+$, and replace the subsampling parameter condition by "the subsampling parameter $\{s_i \in \mathbb{N}_0\}_{i \in \mathbb{Z}^+}$ is a monotonically increasing sequence such that $\sup_{i \in \mathbb{Z}^+} |s_{i+1} - s_i| \leq 1$ ". Denote the set of components used to update the *r*-th order term in the polynomial by $\mathscr{C}_{n,r}$. For instance,

$$\mathscr{C}_{n,r} = \left\{ K_{n,r}, R_{n,r}, k_{n,r}, r_{n,r}, U_{n,r}, V_{n,r}, \left\{ d_{s_n,r}^{(b)}, D_{n,r}^{(b)} \right\}_{b=1,\dots,r} \right\};$$

see Algorithm A.3. Then, we have the Algorithm D.1.

Now, we interpret the conditions in Proposition 1 one by one. Recall the backward finite

difference operator $\nabla^{(b)}$ is defined by

$$\nabla^{(1)}f(k) = f(k) - f(k-1)$$
 and $\nabla^{(b)}f(k) = \nabla^{(b-1)}f(k) - \nabla^{(b-1)}f(k-1)$,

where $b \in \mathbb{Z}^+$ and f is a function that takes an integer input k. The implications of the polynomial T(x) condition are twofold. First, the tapering parameter can be exteriorzed, which has been briefly discussed in Section 2.2. This is important because the tapering parameter should depend on n according to Principle A. However, such tapered sum, e.g., $\sum_{k=1}^{s_i} \cos(\pi k/t_n) X_{i-k}$, may not be updated in O(1) time when t_n changes. Polynomial T(x) is free of this problem as we can work on $\sum_{k=1}^{s_i} T(k) X_{i-k}$ instead of $\sum_{k=1}^{s_i} T(k/t_n) X_{i-k}$.

Second, the backward finite difference method can be used to derive recursive formulas for polynomial T(x). This is because if $\nabla^q T(k)$ is a constant, we can recursively update $\sum_{k=1}^{s_i} T(k) X_{i-k}$ by rearranging the finite differences:

$$\nabla^{q-1}T(k) = \nabla^{q}T(k) + \nabla^{q-1}T(k-1),$$

$$\nabla^{q-2}T(k) = \nabla^{q-1}T(k) + \nabla^{q-2}T(k-1)$$

$$\vdots$$

$$T(k) = \nabla T(k) + T(k-1).$$

Since the terms on right hand side are either constants, updated by their above formulas, or coming from the last iteration (for those with T(k-1)), we can update $\sum_{k=1}^{s_i} T(k)X_{i-k}$ in O(1)time. A working example has been illustrated in Section A.4. In contrast, if $\nabla^q T(k)$ is not a constant for any $q \in \mathbb{Z}^+$, the backward finite difference method will fail. Consequently, it is uncertain whether $\sum_{k=1}^{s_i} \nabla^q T(k)X_{i-k}$ can be updated in O(1) time. A typical example is $T(k) = \exp(k)$ as its derivative, which is the continuous analogue of finite difference, is always $\exp(k)$. Restricting T(k) to be a q-th order polynomial avoids this problem as the constancy of $\nabla^q T(k)$ is well-known; see, e.g., Beji (2018).

Next, we investigate the separable tapering parameter $t_n(i, j) = t_n t'_{i \lor j}$ condition. Note that $i \lor j$ is taken in the smoothing parameter to make the window symmetric only. If we do not assume $t'_i = 1$ for all $i \in \mathbb{Z}^+$, O(1)-time update is still possible in Algorithm D.1 by a similar decomposition as in (D.1). However, we do not consider it in Algorithm D.1 because it may violate Principle A. In contrast, if the tapering parameter is not separable, e.g., $t_n(i, j) =$

 $n + (i \lor j)$, the decomposition in (D.1) becomes

$$\hat{\sigma}_n^2 = \frac{1}{n} \sum_{r=0}^q a_r \sum_{i=1}^n \frac{X_i}{(n+i)^r} (K_{i,r} - k_{i,r} \bar{X}_n) - \frac{1}{n} \sum_{r=0}^q a_r \sum_{i=1}^n \frac{\bar{X}_n}{(n+i)^r} (K_{i,r} - k_{i,r} \bar{X}_n),$$

Consequently, we need to re-compute $X_i K_{i,r}/(n+i)^r$ for i = 1, ..., n and r = 0, ..., q, which likely cannot be done in O(1) time.

Then, we discuss the subsampling function $S(x) = \mathbb{1}_{x \leq 1}$ condition. It is possible to use other subsampling functions such as $S(x) = \mathbb{1}_{x \leq C}$ for some $C \in \mathbb{R}^+$. Nevertheless, the form $S(x) = \mathbb{1}_{x \leq 1}$ is commonly studied in the literature. It also offers a direct relationship between $s_n(i, j)$ and the subsample size since

$$S_n(i) = \left\{ X_j : S\left(\frac{|i-j|}{s_n(i,j)}\right) = 1, 1 \le j \le i \right\} = \left\{ X_j : i - s_n(i,j) \le j \le i \right\},\$$

which means that the subsample size is approximately $s_n(i, j)$. Note that we use the term "approximately" because $s_n(i, j)$ may not be an integer. If we use other S(x) such as $\mathbb{1}_{x < C}$, we will lose this nice interpretation of $s_n(i, j)$.

Finally, for the subsampling parameter condition, the importance of local subsample has been briefly elaborated in Section 2.2. In short, it makes online estimation possible by adapting the estimates to their corresponding points in time. By changing the condition to "the subsampling parameter $\{s_i \in \mathbb{N}_0\}_{i\in\mathbb{Z}^+}$ is a monotonically increasing sequence such that $\sup_{i\in\mathbb{Z}^+} |s_{i+1}-s_i| \leq$ 1", Algorithm D.1 only requires $O(s_n)$ space in contrast to the O(n) space requirement of (D.1). In addition, Algorithm D.1 can be implemented using a queue data structure, which supports insertions and the necessary access in O(1) time. We remark that the condition $\sup_{i\in\mathbb{Z}^+} |s_{i+1}-s_i| \leq 1$ is sufficient throughout Section A because the subsample size grows sublinearly and cannot exceed the sample size in practice; see Definition 1.

Remark D.1. When an online algorithm exhausts the initial memory, its theoretically O(1)time operations may need more than O(1) time to deal with memory reallocation. Therefore, the choice of data structure matters in implementation, and O(1)-space update is desirable under memory constraint.

Remark D.2. We believe that the conditions in Proposition 1 are close to be necessary given their implications. In particular, it is clear that online estimation is closely related to the backward finite difference method. However, it will be difficult to prove that an online algorithm does not exist without certain conditions in general.

Algorithm D.2: Practical O(1)-space update

[1] input:

(i) n, s – the sample size and subsampling parameter from the last iteration [2] (ii) $\mathscr{C}'_n, \mathscr{C}''_n$ – the original components and their pre-calculated versions [3] (iii) s', a – the ramped subsampling parameter and its upper bound from the last iteration [4] begin [5] Receive X_{n+1} [6] Set n = n + 1[7] Compute s_n, t_n, a_n [8] if s' + 1 < a then [9] /* restrict s_n when $s_n' > s'$ */ Set $s'_n = s' + 1$, $s_n = s$, $a_n = a$ Update \mathscr{C}'_n with their recursive formulas [10] [11] if $\underline{s'_n \ge a-s}$ then [12] Update \mathscr{C}''_n with their recursive formulas [13] [14] else Re-initialize \mathscr{C}''_n with their recursive formulas [15] else [16] Set $s'_n = s_n, a = a_n$ [17] Replace the applicable components in \mathscr{C}'_n with \mathscr{C}''_n [18] Update \mathscr{C}'_n with their recursive formulas [19] Re-initialize \mathscr{C}''_n with their recursive formulas [20] Set $s' = s'_n, s = s_n$ [21] Output the recursive estimate computed with \mathscr{C}'_n [22]

D.2 Proof of Proposition 2

To prove that the conditions are sufficient for having O(1)-space update, we consider the case that the original subsampling parameter s_i is monotonically increasing. Denote the set of original components and the set of pre-calculated components under ramping by \mathscr{C}'_n and \mathscr{C}''_n , respectively. For instance,

$$\mathscr{C}'_{n} = \left\{ Q_{n}, \bar{X}_{n}, \left\{ K'_{n,b}, R'_{n,b}, k'_{n,b}, r'_{n,b}, U'_{n,b}, V'_{n,b} \right\}_{b=0,1} \right\};$$

see Algorithm A.4. Since the estimator can be updated in O(1) time, recursive formulas for updating \mathscr{C}'_n and \mathscr{C}''_n exists. Then, the proof is completed by Algorithm D.2. For the case that s_i is fixed or prespecified, the algorithm is similar as the ramping upper bound a_i is also prespecified.

In the remaining of this subsection, we discuss the implications of the conditions in Proposition 2. The O(1)-time update condition ensures the existence of recursive formulas for updating \mathscr{C}'_n and \mathscr{C}''_n . Consequently, we can use the pre-calculation technique to perform O(1)-time and O(1)-space update. A working example has been illustrated in Section A.5.

Next, we discuss the subsampling function $S(x) = \mathbb{1}_{x \leq 1}$ condition. As in Proposition 1, this condition offers a direct relationship between the ramped subsampling parameter $s'_{i \lor j}$ and the

subsample size since

$$S_n(i) = \left\{ X_j : S\left(\frac{|i-j|}{s'_i}\right) = 1, 1 \le j \le i \right\} = \left\{ X_j : i - s'_i \le j \le i \right\},\$$

which means that the subsample size is approximately $s'_{i \lor j}$; see Section D.1.

Finally, for the subsampling parameter condition, it ensures the correctness of the precalculation technique in Algorithm D.2. This is because we need to re-initialize \mathscr{C}''_n to prepare for the next reset of s'_n . If re-initialization cannot be done, we need to drop far observations in \mathscr{C}''_n , which ruins the O(1)-space update. To re-initialize properly, we need $s'_n \leq a - s$ for some $s'_n \in [s, a)$, which is guaranteed by the $\phi \geq 2$ condition. If the original subsampling parameter s_i is not monotonically increasing but prespecified, we can still pre-calculate the components by computing s_n at the next point of reset in advance.

E Proof of Lemmas

E.1 Lemma E.1

The following lemma shows that the unknown-mean version $\hat{\sigma}_n^2$ and the known-mean version $\bar{\sigma}_n^2$ are asymptotically equivalent in the $\mathscr{L}^{\alpha/2}$ -sense.

Lemma E.1. Let $\alpha > 2$. Suppose that $X_1 \in \mathscr{L}^{\alpha}$. If Assumption 1 holds, then

$$\left\|\hat{\sigma}_{n}^{2}-\bar{\sigma}_{n}^{2}\right\|_{\frac{\alpha}{2}}=O(n^{-1})+O\left(n^{-\frac{3}{2}}G_{3,n}^{\frac{1}{2}}\right)+O\left(n^{-2}G_{4,n}\right)$$

where

$$G_{3,n} \coloneqq \sum_{i=2}^{n} \left| \sum_{j=1}^{i-1} W_n(i,j) \right|^2 + \sum_{j=1}^{n-1} \left| \sum_{i=j+1}^{n} W_n(i,j) \right|^2 \quad and \quad G_{4,n} \coloneqq \left| \sum_{i=2}^{n} \sum_{j=1}^{i-1} W_n(i,j) \right|.$$

In addition, if Assumptions 2 and 3(a) also hold, then

$$\left\|\hat{\sigma}_n^2 - \bar{\sigma}_n^2\right\|_{\frac{\alpha}{2}} = o(1).$$

Proof. By Minkowski inequality and Hölder's inequality,

$$\begin{split} \left\| \hat{\sigma}_{n}^{2} - \bar{\sigma}_{n}^{2} \right\|_{\frac{\alpha}{2}} &\leq \\ \left\| \frac{1}{n} \sum_{i=1}^{n} (2X_{i} - \bar{X}_{n} - \mu)(\mu - \bar{X}_{n}) \right\|_{\frac{\alpha}{2}} \\ &+ \\ \left\| \frac{2}{n} \sum_{i=2}^{n} \sum_{j=1}^{i-1} W_{n}(i,j) \left\{ \bar{X}_{n}^{2} - (\bar{X}_{n} - \mu)(X_{i} + X_{j}) - \mu^{2} \right\} \right\|_{\frac{\alpha}{2}} \\ &\leq \\ \left\| (\bar{X}_{n} - \mu)(\mu - \bar{X}_{n}) \right\|_{\frac{\alpha}{2}} \\ &+ \\ \left\| \frac{2}{n} \sum_{i=2}^{n} \sum_{j=1}^{i-1} W_{n}(i,j) \left\{ (\bar{X}_{n} - \mu)^{2} - (\bar{X}_{n} - \mu)(X_{i} + X_{j} - 2\mu) \right\} \right\|_{\frac{\alpha}{2}} \\ &\leq \\ \left\| \bar{X}_{n} - \mu \right\|_{\alpha}^{2} + \frac{2}{n} \left\| \sum_{i=2}^{n} \sum_{j=1}^{i-1} W_{n}(i,j) \right\| \left\| \bar{X}_{n} - \mu \right\|_{2\alpha}^{2} \\ &+ \\ &+ \frac{2}{n} \left\| \sum_{i=2}^{n} \sum_{j=1}^{i-1} W_{n}(i,j)(X_{i} + X_{j} - 2\mu) \right\|_{\alpha} \left\| \bar{X}_{n} - \mu \right\|_{\alpha}. \end{split}$$
(E.1)

By the moment inequality in Theorem 3 of Wu (2011) and Assumption 1, $\|\bar{X}_n - \mu\|_{\alpha} = O(n^{-1/2})$. For the last term in (E.1), we can use Minkowski inequality to decompose the sum:

$$\left\|\sum_{i=2}^{n}\sum_{j=1}^{i-1}W_{n}(i,j)(X_{i}+X_{j}-2\mu)\right\|_{\alpha} \leq \left\|\sum_{i=2}^{n}\sum_{j=1}^{i-1}W_{n}(i,j)(X_{i}-\mu)\right\|_{\alpha} + \left\|\sum_{j=1}^{n-1}\sum_{i=j+1}^{n}W_{n}(i,j)(X_{j}-\mu)\right\|_{\alpha}.$$

By Lemma 1 of Liu and Wu (2010),

$$\left\|\sum_{i=2}^{n}\sum_{j=1}^{i-1}W_{n}(i,j)(X_{i}-\mu)\right\|_{\alpha} \leq C_{\alpha}\Delta_{\alpha}\sqrt{\sum_{i=2}^{n}\left|\sum_{j=1}^{i-1}W_{n}(i,j)\right|^{2}} \leq C_{\alpha}\Delta_{\alpha}G_{3,n}^{\frac{1}{2}},$$
$$\left\|\sum_{j=1}^{n-1}\sum_{i=j+1}^{n}W_{n}(i,j)(X_{j}-\mu)\right\|_{\alpha} \leq C_{\alpha}\Delta_{\alpha}\sqrt{\sum_{j=1}^{n-1}\left|\sum_{i=j+1}^{n}W_{n}(i,j)\right|^{2}} \leq C_{\alpha}\Delta_{\alpha}G_{3,n}^{\frac{1}{2}},$$

where $C_{\alpha} \in \mathbb{R}^+$ is some constant that only depends on α and may change from line to line in the proofs. Hence under Assumption 1,

$$\left\|\sum_{i=2}^{n}\sum_{j=1}^{i-1}W_n(i,j)(X_i+X_j-2\mu)\right\|_{\alpha}=O(G_{3,n}^{\frac{1}{2}}).$$

Combining the results, we have

$$\begin{split} \left\|\hat{\sigma}_{n}^{2} - \bar{\sigma}_{n}^{2}\right\|_{\frac{\alpha}{2}} &= O(n^{-1}) + n^{-1}G_{4,n}O(n^{-1}) + n^{-1}O\left(G_{3,n}^{\frac{1}{2}}\right)O\left(n^{-\frac{1}{2}}\right) \\ &= O(n^{-1}) + O\left(n^{-\frac{3}{2}}G_{3,n}^{\frac{1}{2}}\right) + O\left(n^{-2}G_{4,n}\right). \end{split}$$

In addition, if Assumptions 2 and 3(a) also hold, Minkowski inequality gives

$$\begin{aligned} G_{3,n} &\leq 2\sum_{i=1}^{n} \left(\sum_{j=1}^{n} \left| W_n(i,j) \right| \right)^2 \leq 2\sum_{i=1}^{n} G_{1,n}^2 = O(n)o\left(n^{4-\frac{4}{\alpha'}}\right) = o(n^3), \\ G_{4,n} &\leq \sum_{i=1}^{n} \sum_{j=1}^{n} \left| W_n(i,j) \right| \leq \sum_{i=1}^{n} G_{1,n} = O(n)o\left(n^{2-\frac{2}{\alpha'}}\right) = o(n^2), \end{aligned}$$

since $\alpha' \in (1,2]$. Therefore, $\|\hat{\sigma}_n^2 - \bar{\sigma}_n^2\|_{\alpha/2} = o(1)$.

E.2 Lemma E.2

Recall that we can assume $\mu = 0$ after applying Lemma E.1. Define

$$\Delta_{m,\alpha} \coloneqq \sum_{i=m}^{\infty} \delta_{i,\alpha}, \tag{E.2}$$

$$\Upsilon_{m,\alpha} \coloneqq \sqrt{\sum_{i=m}^{\infty} \delta_{i,\alpha}^2},\tag{E.3}$$

$$d_{m,\alpha} \coloneqq \sum_{i=0}^{\infty} \min(\delta_{i,\alpha}, \Upsilon_{m+1,\alpha}), \tag{E.4}$$

$$\mathscr{P}_{i} := \mathbb{E}(\cdot \mid \mathscr{F}_{i}) - \mathbb{E}(\cdot \mid \mathscr{F}_{i-1}).$$
(E.5)

For simplicity, write $\Delta_{0,\alpha} = \Delta_{\alpha}$. The following lemma shows that the known-mean version $\bar{\sigma}_n^2$ and the *m*-dependent process approximated version $\tilde{\sigma}_n^2$ are asymptotically equivalent in the $\mathscr{L}^{\alpha/2}$ -sense.

Lemma E.2. Let $\alpha > 2$ and $\alpha' = \min(\alpha/2, 2)$. Suppose that $X_1 \in \mathscr{L}^{\alpha}$. If Assumption 1 holds, then

$$\left\|\bar{\sigma}_n^2 - \mathbb{E}(\bar{\sigma}_n^2) - \tilde{\sigma}_n^2 + \mathbb{E}(\tilde{\sigma}_n^2)\right\|_{\frac{\alpha}{2}} = O\left(n^{\frac{1}{\alpha'}-1}d_{m,\alpha}G_{5,n}^{\frac{1}{2}}\right),$$

where

$$G_{5,n} \coloneqq \max_{1 \le i \le n} \sum_{j=1}^{n} \left\{ W_n^2(i,j) + W_n^2(j,i) \right\}.$$

In addition, if Assumptions 2 and 3(a) also hold, then

$$\left\|\bar{\sigma}_n^2 - \mathbb{E}(\bar{\sigma}_n^2) - \tilde{\sigma}_n^2 + \mathbb{E}(\tilde{\sigma}_n^2)\right\|_{\frac{\alpha}{2}} = o(1).$$

Remark E.1. The proof of Lemma E.2 follows a similar procedure as the proof of Proposition 1 in Liu and Wu (2010). However, their Proposition 1 cannot be applied directly since our window functions may depend on the running indices i, j in addition to the difference i - j. In addition, their Proposition 1 does not provide explicit approximation of $\sum_{i=1}^{n} X_i^2$, which is included in our estimators. Finally, their Proposition 1 does not consider the case of $2 < \alpha < 4$,

which we will show later for proving consistency of our estimators.

Proof. Recall that the *m*-dependent process \tilde{X}_j is defined in (B.3). Let $Z_i = 2 \sum_{j=1}^{i-1} W_n(i,j) X_j$, and $\tilde{Z}_i = 2 \sum_{j=1}^{i-1} W_n(i,j) \tilde{X}_j$. Define the intermediate approximated version

$$\tilde{\sigma}_n^* \coloneqq \frac{1}{n} \sum_{i=1}^n X_i \tilde{X}_i + \frac{2}{n} \sum_{i=2}^n X_i \sum_{j=1}^{i-1} W_n(i,j) \tilde{X}_j = \frac{1}{n} \sum_{i=1}^n X_i \tilde{X}_i + \frac{1}{n} \sum_{i=2}^n X_i \tilde{Z}_i$$

In addition, recall that $X_{i,\{h\}} = g(\mathscr{F}_{i,\{h\}})$ represents a coupled version of $X_i = g(\mathscr{F}_i)$ with ϵ_h replaced by an IID copy ϵ'_h in \mathscr{F}_i . We similarly define $Z_{i,\{h\}}$ and $\tilde{Z}_{i,\{h\}}$ as the coupled versions of Z_i and \tilde{Z}_i , respectively. Note that for any $X_i \in \sigma(\mathscr{F}_i)$,

$$\left\|\mathscr{P}_{h}X_{i}\right\|_{\alpha} \leq \left\|X_{i} - X_{i,\{h\}}\right\|_{\alpha};$$

see Wu (2005). Therefore, by Minkowski inequality,

$$n \left\| \mathscr{P}_{h}(\bar{\sigma}_{n}^{2} - \tilde{\sigma}_{n}^{*}) \right\|_{\frac{\alpha}{2}} \leq \left\| \sum_{i=1}^{n} \left\{ X_{i}(X_{i} - \tilde{X}_{i}) - X_{i,\{h\}}(X_{i,\{h\}} - \tilde{X}_{i,\{h\}}) \right\} \right\|_{\frac{\alpha}{2}} \\ + \sum_{i=2}^{n} \left\{ X_{i}(Z_{i} - \tilde{Z}_{i}) - X_{i,\{h\}}(Z_{i,\{h\}} - \tilde{Z}_{i,\{h\}}) \right\} \right\|_{\frac{\alpha}{2}} \\ \leq \sum_{i=1}^{n} \left\| X_{i,\{h\}} \left\{ (X_{i} - \tilde{X}_{i}) - (X_{i,\{h\}} - \tilde{X}_{i,\{h\}}) \right\} \right\|_{\frac{\alpha}{2}} \\ + \sum_{i=1}^{n} \left\| (X_{i} - X_{i,\{h\}})(X_{i} - \tilde{X}_{i}) \right\|_{\frac{\alpha}{2}} \\ + \left\| \sum_{i=2}^{n} X_{i,\{h\}} \left\{ (Z_{i} - \tilde{Z}_{i}) - (Z_{i,\{h\}} - \tilde{Z}_{i,\{h\}}) \right\} \right\|_{\frac{\alpha}{2}} \\ + \sum_{i=2}^{n} \left\| (X_{i} - X_{i,\{h\}})(Z_{i} - \tilde{Z}_{i}) \right\|_{\frac{\alpha}{2}} \\ =: I_{1} + I_{2} + I_{3} + I_{4}.$$
(E.6)

By Jensen's inequality,

$$\begin{split} \left\| \tilde{X}_{i} - \tilde{X}_{i,\{0\}} \right\|_{\alpha} &= \left\| \mathbb{E} \left(X_{i} - X_{i,\{0\}} \mid \epsilon_{i-m}, \dots, \epsilon_{i}, \epsilon_{0}' \right) \right\|_{\alpha} \\ &\leq \left\| X_{i} - X_{i,\{0\}} \right\|_{\alpha} = \delta_{i,\alpha}. \end{split}$$

For $\alpha > 2$, Burkholder inequality and Theorem 1 of Wu (2005) give

$$\begin{aligned} \left\|X_{i} - \tilde{X}_{i}\right\|_{\alpha}^{2} &= \left\|\sum_{l=0}^{\infty} \mathscr{P}_{i-l} X_{i} - \sum_{l=0}^{\infty} \mathscr{P}_{i-l} \tilde{X}_{i}\right\|_{\alpha}^{2} \\ &= \left\|\sum_{l=m+1}^{\infty} \mathscr{P}_{i-l} X_{i}\right\|_{\alpha}^{2} \end{aligned}$$

$$\leq C_{\alpha} \sum_{l=m+1}^{\infty} \|\mathscr{P}_{i-l}X_i\|_{\alpha}^2$$
$$\leq C_{\alpha} \sum_{l=m+1}^{\infty} \delta_{l,\alpha}^2 = C_{\alpha}\Upsilon_{m+1,\alpha}^2$$

Combining the results yields

$$\left\|X_{i} - \tilde{X}_{i} - X_{i,\{h\}} + \tilde{X}_{i,\{h\}}\right\|_{\alpha} \le C_{\alpha} \min(\delta_{i-h,\alpha}, \Upsilon_{m+1,\alpha}).$$
(E.7)

Now we begin to derive upper bounds for I_1 to I_4 . By Hölder's inequality, (E.7) and stationarity,

$$I_{1} \leq \sum_{i=1}^{n} \|X_{i,\{h\}}\|_{\alpha} \|X_{i} - \tilde{X}_{i} - X_{i,\{h\}} + \tilde{X}_{i,\{h\}}\|_{\alpha}$$
$$\leq C_{\alpha} \max_{1 \leq i \leq n} \|X_{i,\{h\}}\|_{\alpha} \sum_{i=1}^{n} \min(\delta_{i-h,\alpha}, \Upsilon_{m+1,\alpha})$$
$$\leq C_{\alpha} \|X_{1}\|_{\alpha} d_{m,\alpha}.$$

Observe that $d_{m,\alpha} \leq \sum_{i=0}^{\infty} \delta_{i,\alpha} = \Delta_{\alpha}$. Hence under Assumption 1 and $X_1 \in \mathscr{L}^{\alpha}$, $I_1 = O(1)$. For the second term in (E.6), by Hölder's inequality we have

$$I_{2} \leq \sum_{i=1}^{n} \left\| X_{i} - X_{i,\{h\}} \right\|_{\alpha} \left\| X_{i} - \tilde{X}_{i} \right\|_{\alpha}$$
$$\leq \max_{1 \leq i \leq n} \left\| X_{i} - \tilde{X}_{i} \right\|_{\alpha} \sum_{i=1}^{n} \delta_{i-h,\alpha}$$
$$\leq C_{\alpha} \Upsilon_{m+1,\alpha} \Delta_{\alpha}.$$

Note that

$$\Upsilon_{m+1,\alpha} = \sqrt{\sum_{i=m+1}^{\infty} \delta_{i,\alpha}^2} \le \sqrt{\sum_{i=1}^{\infty} \delta_{i,\alpha}^2} \le \sqrt{\left|\sum_{i=1}^{\infty} \delta_{i,\alpha}\right|^2} = \Delta_{\alpha}.$$

So under Assumption 1, $I_2 = O(1)$. For the third term in (E.6), we first rearrange the order of summation:

$$I_{3} = \left\| 2 \sum_{i=2}^{n} X_{i,\{h\}} \sum_{j=1}^{i-1} W_{n}(i,j) (X_{j} - \tilde{X}_{j} - X_{j,\{h\}} + \tilde{X}_{j,\{h\}}) \right\|_{\frac{\alpha}{2}}$$
$$= \left\| 2 \sum_{j=1}^{n-1} (X_{j} - \tilde{X}_{j} - X_{j,\{h\}} + \tilde{X}_{j,\{h\}}) \sum_{i=j+1}^{n} W_{n}(i,j) X_{i,\{h\}} \right\|_{\frac{\alpha}{2}}.$$

By Minkowski inequality, Hölder's inequality, Lemma 1 of Liu and Wu (2010) and (E.7),

$$I_{3} \leq 2 \sum_{j=1}^{n-1} \left\| X_{j} - \tilde{X}_{j} - X_{j,\{h\}} + \tilde{X}_{j,\{h\}} \right\|_{\alpha} \left\| \sum_{i=j+1}^{n} W_{n}(i,j) X_{i,\{h\}} \right\|_{\alpha}$$
$$\leq C_{\alpha} \Delta_{\alpha} \sum_{j=1}^{n} \min(\delta_{j-h,\alpha}, \Upsilon_{m+1,\alpha}) \sqrt{\sum_{i=j+1}^{n} W_{n}^{2}(i,j)}$$

$$\leq C_{\alpha} \Delta_{\alpha} d_{m,\alpha} \max_{1 \leq j \leq n} \sqrt{\sum_{i=j+1}^{n} W_n^2(i,j)}$$
$$\leq C_{\alpha} \Delta_{\alpha} d_{m,\alpha} G_{5,n}^{\frac{1}{2}}.$$

Under Assumption 1, $I_3 = O(d_{m,\alpha}G_{5,n}^{1/2})$. For the last term in (E.6), note that by Lemma 1 of Liu and Wu (2010),

$$\begin{aligned} \left\| Z_i - \tilde{Z}_i \right\|_{\alpha} &= \left\| 2 \sum_{j=1}^{i-1} W_n(i,j) (X_j - \tilde{X}_j) \right\|_{\alpha} \\ &\leq C_{\alpha} \Delta_{m+1,\alpha} \sqrt{\sum_{j=1}^{i-1} W_n^2(i,j)} \\ &\leq C_{\alpha} \Delta_{m+1,\alpha} G_{5,n}^{\frac{1}{2}}. \end{aligned}$$

Thus, by Hölder's inequality and Assumption 1,

$$I_{4} \leq \sum_{i=1}^{n} \left\| X_{i} - X_{i,\{h\}} \right\|_{\alpha} \left\| Z_{i} - \tilde{Z}_{i} \right\|_{\alpha}$$
$$\leq C_{\alpha} \Delta_{m+1,\alpha} G_{5,n}^{\frac{1}{2}} \sum_{i=1}^{n} \delta_{i-h,\alpha}$$
$$\leq C_{\alpha} \Delta_{\alpha} \Delta_{m+1,\alpha} G_{5,n}^{\frac{1}{2}}.$$

Note that

$$\sum_{i=0}^{\infty} \Upsilon_{m+1,\alpha} - \Delta_{m+1,\alpha} = \sum_{i=0}^{\infty} \sqrt{\sum_{i=m+1}^{\infty} \delta_{i,\alpha}^2} - \sum_{i=m+1}^{\infty} \sqrt{\delta_{i,\alpha}^2} \ge 0,$$

which implies $\Delta_{m+1,\alpha} \leq d_{m,\alpha}$. Hence we have $I_4 = O(d_{m,\alpha}G_{5,n}^{1/2})$. Combining the results, we can bound (E.6) by

$$\begin{aligned} \left\| \mathscr{P}_{h}(\bar{\sigma}_{n}^{2} - \tilde{\sigma}_{n}^{*}) \right\|_{\frac{\alpha}{2}} &= \frac{1}{n} \left\{ O(1) + O\left(d_{m,\alpha} G_{5,n}^{\frac{1}{2}}\right) \right\} \\ &= O\left(n^{-1} d_{m,\alpha} G_{5,n}^{\frac{1}{2}}\right). \end{aligned}$$

Finally, by the decomposition $X_n - \mathbb{E}(X_n) = \sum_{h=-\infty}^n \mathscr{P}_h X_n$ for stationary $\{X_i\}$ and Burkholder inequality,

$$\begin{split} \left\|\bar{\sigma}_{n}^{2} - \mathbb{E}(\bar{\sigma}_{n}^{2}) - \tilde{\sigma}_{n}^{*} + \mathbb{E}(\tilde{\sigma}_{n}^{*})\right\|_{\frac{\alpha}{2}}^{\alpha'} &\leq C_{\alpha} \sum_{h=-\infty}^{n} \left\|\mathscr{P}_{h}(\bar{\sigma}_{n}^{2} - \tilde{\sigma}_{n}^{*})\right\|_{\frac{\alpha}{2}}^{\alpha'} \\ &= O\left(n^{1-\alpha'}d_{m,\alpha}^{\alpha'}G_{5,n}^{\frac{\alpha'}{2}}\right), \\ \left\|\bar{\sigma}_{n}^{2} - \mathbb{E}(\bar{\sigma}_{n}^{2}) - \tilde{\sigma}_{n}^{*} + \mathbb{E}(\tilde{\sigma}_{n}^{*})\right\|_{\frac{\alpha}{2}} &= O\left(n^{\frac{1}{\alpha'}-1}d_{m,\alpha}G_{5,n}^{\frac{1}{2}}\right). \end{split}$$

A similar inequality for $\|\tilde{\sigma}_n^* - \mathbb{E}(\tilde{\sigma}_n^*) - \tilde{\sigma}_n^2 + \mathbb{E}(\tilde{\sigma}_n^2)\|_{\alpha/2}$ completes the proof of the general case. In addition, note that $d_{m,\alpha} = o(1)$ as $m \to \infty$ in view of Lemma E.5. To fulfill this condition, we can choose m such that $m \to \infty$ as $n \to \infty$ by Assumption 2; see Section E.3. It also follows from Assumptions 2 and 3(a) that

$$G_{5,n} = 2 \max_{1 \le i \le n} \sum_{j=1}^{n} W_n^2(i,j) \le 2G_{1,n} = o(n^{2-\frac{2}{\alpha'}}).$$

Therefore, $\|\bar{\sigma}_n^2 - \mathbb{E}(\bar{\sigma}_n^2) - \tilde{\sigma}_n^* + \mathbb{E}(\tilde{\sigma}_n^*)\|_{\alpha/2} = o(1).$ A similar inequality for $\|\tilde{\sigma}_n^* - \mathbb{E}(\tilde{\sigma}_n^*) - \tilde{\sigma}_n^2 + \mathbb{E}(\tilde{\sigma}_n^2)\|_{\alpha/2}$ completes the proof.

E.3 Lemma E.3

The following lemma shows that the *m*-dependent process approximated version $\tilde{\sigma}_n^2$ and the martingale difference approximated version $\check{\sigma}_n^2$ are asymptotically equivalent in the $\mathscr{L}^{\alpha/2}$ -sense.

Lemma E.3. Let $\alpha > 2$ and $\alpha' = \min(\alpha/2, 2)$. Suppose that $\mathbb{E}|X_1|^{\alpha} < \infty$. If Assumption 1 holds, then

$$\left\|\tilde{\sigma}_n^2 - \mathbb{E}(\tilde{\sigma}_n^2) - \breve{\sigma}_n^2 + \mathbb{E}(\breve{\sigma}_n^2)\right\|_{\frac{\alpha}{2}} = O\left(m^{3-\frac{1}{\alpha'}}n^{\frac{1}{\alpha'}-1}G_{6,n}\right),$$

where

$$G_{6,n} \coloneqq \max_{1 \le i,j \le n} |W_n(i,j)| + \max_{1 \le i \le n} \left[\sum_{j=2}^n \left\{ \left| \nabla_1 W_n(j,i) \right|^{\alpha'} + \left| \nabla_2 W_n(i,j) \right|^{\alpha'} \right\} \right]^{\frac{1}{\alpha'}},$$

$$\nabla_1 W_n(j,i) = W_n(j,i) - W_n(j-1,i) \quad and \quad \nabla_2 W_n(i,j) = W_n(i,j) - W_n(i,j-1).$$

In addition, if Assumptions 2 and 3(a) also hold, then

$$\left\|\tilde{\sigma}_n^2 - \mathbb{E}(\tilde{\sigma}_n^2) - \breve{\sigma}_n^2 + \mathbb{E}(\breve{\sigma}_n^2)\right\|_{\frac{\alpha}{2}} = o(1).$$

Remark E.2. The proof of Lemma E.3 follows a similar procedure as the proof of Proposition 2 in Liu and Wu (2010). However, their Proposition 2 cannot be applied directly for similar reasons stated in Remark E.1. It is also worth noting that $W_n(i,j)$ is implicitly assumed to be real throughout our paper. Therefore, we do not consider the case of complex coefficients in contrast to Proposition 2 of Liu and Wu (2010).

Proof. We first show some simple results that will be used later in this proof. They were similarly established in the proof of Proposition 2 in Liu and Wu (2010) but we try to give more details here. Recall that \tilde{X}_i , M_i and D_i are defined in (B.3), (B.5) and (B.6), respectively. By
the construction of \tilde{X}_i , we can rewrite M_i by

$$M_i = \sum_{h=0}^m \mathbb{E}(\tilde{X}_{i+h} \mid \mathscr{F}_i),$$

since $\mathbb{E}(\tilde{X}_{i+h} | \mathcal{F}_i) = 0$ when h > m. In addition, D_i 's are *m*-dependent and form martingale differences with respect to \mathcal{F}_i . By Minkowski inequality, Jensen's inequality and stationarity of X_i , we have

$$\|M_i\|_{\alpha} = \left\| \sum_{h=0}^{m} \mathbb{E}(\tilde{X}_{i+h} \mid \mathscr{F}_i) \right\|_{\alpha} \le 2m \|X_1\|_{\alpha}.$$
(E.8)

We also have

$$M_{i} - \mathbb{E}(M_{i+1} \mid \mathscr{F}_{i}) = \sum_{h=0}^{m} \mathbb{E}(\tilde{X}_{i+h} \mid \mathscr{F}_{i}) - \sum_{h=0}^{m} \mathbb{E}(\tilde{X}_{i+h+1} \mid \mathscr{F}_{i})$$
$$= \mathbb{E}(\tilde{X}_{i} \mid \mathscr{F}_{i}) = \tilde{X}_{i}.$$

As a result,

$$\tilde{X}_{i} - D_{i} = M_{i} - \mathbb{E}(M_{i+1} \mid \mathscr{F}_{i}) - M_{i} + \mathbb{E}(M_{i} \mid \mathscr{F}_{i-1})$$
$$= \mathbb{E}(M_{i} \mid \mathscr{F}_{i-1}) - \mathbb{E}(M_{i+1} \mid \mathscr{F}_{i}).$$
(E.9)

Now we begin the main part of this proof. Let

$$Y_{1,i} = \tilde{X}_i \sum_{j=1}^{i-8m} W_n(i,j) (\tilde{X}_j - D_j), \quad Y_{2,i} = \tilde{X}_i \sum_{j=i-8m+1}^{i-1} W_n(i,j) (\tilde{X}_j - D_j),$$

$$Y_i = 2\tilde{X}_i \sum_{j=1}^{i-1} W_n(i,j) (\tilde{X}_j - D_j) = 2(Y_{1,i} + Y_{2,i}),$$

$$Y'_i = 2(\tilde{X}_i - D_i) \sum_{j=1}^{i-1} W_n(i,j) D_j.$$

By Minkowski inequality,

$$n \|\tilde{\sigma}_{n}^{2} - \mathbb{E}(\tilde{\sigma}_{n}^{2}) - \check{\sigma}_{n}^{2} + \mathbb{E}(\check{\sigma}_{n}^{2})\|_{\frac{\alpha}{2}} \leq \left\|\sum_{i=1}^{n} \left\{\tilde{X}_{i}^{2} - \mathbb{E}(\tilde{X}_{i}^{2})\right\}\right\|_{\frac{\alpha}{2}} + \left\|\sum_{i=1}^{n} \left\{D_{i}^{2} - \mathbb{E}(D_{i}^{2})\right\}\right\|_{\frac{\alpha}{2}} + \left\|\sum_{i=1}^{n} \left\{Y_{i} - \mathbb{E}(Y_{i})\right\}\right\|_{\frac{\alpha}{2}} + \left\|\sum_{i=1}^{n} \left\{Y_{i}' - \mathbb{E}(Y_{i}')\right\}\right\|_{\frac{\alpha}{2}} = :I_{5} + I_{6} + I_{7} + I_{8}.$$
(E.10)

To upper bound the first term in (E.10), note that

$$\begin{aligned} \left\| \tilde{X}_{i}^{2} - \tilde{X}_{i,\{0\}}^{2} \right\|_{\frac{\alpha}{2}} &\leq \left\| \mathbb{E} (X_{i} - X_{i,\{0\}} | \epsilon_{i-m}, \dots, \epsilon_{i}, \epsilon'_{0}) \right\|_{\alpha} \left\| \mathbb{E} (X_{i} + X_{i,\{0\}} | \epsilon_{i-m}, \dots, \epsilon_{i}, \epsilon'_{0}) \right\|_{\alpha} \\ &\leq \delta_{i,\alpha} \left\| X_{i} + X_{i,\{0\}} \right\|_{\alpha}. \end{aligned}$$

Hence under Assumption 1 and $X_i \in \mathscr{L}^{\alpha}$,

$$\sum_{i=0}^{\infty} \left\| \tilde{X}_i^2 - \tilde{X}_{i,\{0\}}^2 \right\|_{\frac{\alpha}{2}} \le \sup_{i \in \mathbb{N}_0} \left\| X_i + X_{i,\{0\}} \right\|_{\alpha} \sum_{i=0}^{\infty} \delta_{i,\alpha}$$
$$= \Delta_{\alpha} \sup_{i \in \mathbb{N}_0} \left\| X_i + X_{i,\{0\}} \right\|_{\alpha} < \infty.$$

This means that $\{\tilde{X}_i^2\}$ is $\alpha/2$ -stable (Wu, 2011). Thus, it follows from Lemma 1 of Liu and Wu (2010) that

$$I_5 = O\left\{\left(\sum_{i=1}^n 1^{\alpha'}\right)^{\frac{1}{\alpha'}}\right\} = O(n^{\frac{1}{\alpha'}}).$$

To upper bound the second term in (E.10), we apply Minkowski inequality to obtain

$$I_{6} \leq \sum_{i=1}^{4m-1} \left\| \sum_{h=0}^{\lfloor (n-i)/(4m) \rfloor} \left\{ D_{i+4mh}^{2} - \mathbb{E}(D_{i+4mh}^{2}) \right\} \right\|_{\frac{\alpha}{2}}.$$
 (E.11)

Since D_i^2 's are also *m*-dependent, (E.11) becomes

$$I_6 = m \cdot O\left\{ \left(\frac{n}{m}\right)^{\frac{1}{\alpha'}} \right\} = O(m^{1 - \frac{1}{\alpha'}} n^{\frac{1}{\alpha'}}).$$

For the third term in (E.10), we try to find an upper bound for $Y_{1,i}$ first. By (E.9), Minkowski inequality, Jensen's inequality and (E.8),

$$\begin{aligned} \left\| \sum_{j=1}^{i-8m} W_{n}(i,j)(\tilde{X}_{j} - D_{j}) \right\|_{\frac{\alpha}{2}} &= \left\| \sum_{j=1}^{i-8m} W_{n}(i,j) \left\{ \mathbb{E}(M_{j} \mid \mathscr{F}_{j-1}) - \mathbb{E}(M_{j+1} \mid \mathscr{F}_{j}) \right\} \right\|_{\frac{\alpha}{2}} \\ &\leq C_{\alpha} m \left\| X_{1} \right\|_{\frac{\alpha}{2}} \max_{1 \leq i,j \leq n} \left| W_{n}(i,j) \right| \\ &+ \left\| \sum_{j=2}^{i-8m} \left\{ W_{n}(i,j) - W_{n}(i,j-1) \right\} \mathbb{E}(M_{j} \mid \mathscr{F}_{j-1}) \right\|_{\frac{\alpha}{2}} \\ &=: C_{\alpha} m \left\| X_{1} \right\|_{\frac{\alpha}{2}} \max_{1 \leq i,j \leq n} \left| W_{n}(i,j) \right| + I_{9}. \end{aligned}$$
(E.12)

To upper bound I_9 , note that $\mathbb{E}(M_j \mid \mathscr{F}_{j-1}) = \sum_{h=1}^m \mathscr{P}_{j-h} \mathbb{E}(M_j \mid \mathscr{F}_{j-1})$ and $\mathscr{P}_{j-h} \mathbb{E}(M_j \mid \mathscr{F}_{j-1})$ are martingale differences with respect to \mathscr{F}_{j-h} . They can be verified since

$$\sum_{h=1}^{m} \mathscr{P}_{j-h} \mathbb{E}(M_j \mid \mathscr{F}_{j-1}) = \sum_{h=1}^{m} \left\{ \mathbb{E}(M_j \mid \mathscr{F}_{j-h}) - \mathbb{E}(M_j \mid \mathscr{F}_{j-h-1}) \right\}$$
$$= \mathbb{E}(M_j \mid \mathscr{F}_{j-1}) - \mathbb{E}(M_j \mid \mathscr{F}_{j-m-1}) = \mathbb{E}(M_j \mid \mathscr{F}_{j-1}),$$

and

$$\mathbb{E}\left\{\mathscr{P}_{j-h}\mathbb{E}(M_{j} \mid \mathscr{F}_{j-1}) \mid \mathscr{F}_{j-h-1}\right\} = \mathbb{E}(M_{j} \mid \mathscr{F}_{j-h-1}) - \mathbb{E}(M_{j} \mid \mathscr{F}_{j-h-1}) = 0.$$

Hence, by Burkholder inequality, Minkowski inequality, Jensen's inequality and (E.8),

$$\begin{split} I_{9}^{\alpha'} &\leq C_{\alpha} \sum_{j=2}^{i-8m} \left\| \left\{ W_{n}(i,j) - W_{n}(i,j-1) \right\} \sum_{h=1}^{m} \mathscr{P}_{j-h} \mathbb{E}(M_{j} \mid \mathscr{F}_{j-1}) \right\|_{\frac{\alpha}{2}}^{\alpha'} \\ &\leq C_{\alpha} \sum_{j=2}^{i-8m} \left| W_{n}(i,j) - W_{n}(i,j-1) \right|^{\alpha'} \left\{ \sum_{h=1}^{m} \left\| \mathscr{P}_{j-h} \mathbb{E}(M_{j} \mid \mathscr{F}_{j-1}) \right\|_{\frac{\alpha}{2}} \right\}^{\alpha'} \\ &\leq C_{\alpha} m^{2\alpha'} \left\| X_{1} \right\|_{\frac{\alpha}{2}}^{\alpha'} \sum_{j=2}^{i-8m} \left| W_{n}(i,j) - W_{n}(i,j-1) \right|^{\alpha'}, \end{split}$$

which implies that

$$I_{9} \leq C_{\alpha} m^{2} \|X_{1}\|_{\frac{\alpha}{2}} \left\{ \sum_{j=2}^{i-8m} \left| W_{n}(i,j) - W_{n}(i,j-1) \right|^{\alpha'} \right\}^{\frac{1}{\alpha'}}.$$

(E.12) becomes

$$\left\|\sum_{j=1}^{i-8m} W_n(i,j)(\tilde{X}_j - D_j)\right\|_{\frac{\alpha}{2}} \le C_{\alpha} m^2 G_{6,n} \|X_1\|_{\frac{\alpha}{2}}.$$
(E.13)

Similarly, we have

$$\left\|\sum_{j=i-8m+1}^{i-1} W_n(i,j)(\tilde{X}_j - D_j)\right\|_{\frac{\alpha}{2}} \le C_{\alpha} m^2 G_{6,n} \|X_1\|_{\frac{\alpha}{2}}.$$
(E.14)

Now to apply the idea of independence as in (E.11), we need to check that $Y_{1,i}, Y_{1,i+4m}, Y_{1,i+8m}, \ldots$ are martingale differences. This can be verified since

$$\mathbb{E}(Y_{1,i+4m} \mid \mathscr{F}_i) = \mathbb{E}\left\{\tilde{X}_{i+4m} \sum_{j=1}^{i-4m} W_n(i,j)(\tilde{X}_j - D_j) \mid \mathscr{F}_i\right\}$$
$$= \sum_{j=1}^{i-4m} W_n(i,j)(\tilde{X}_j - D_j)\mathbb{E}(\tilde{X}_{i+4m}) = 0.$$

Using Hölder's inequality and (E.13),

$$\|Y_{1,i}\|_{\frac{\alpha}{2}} \leq \|\tilde{X}_i\|_{\alpha} \left\| \sum_{j=1}^{i-8m} W_n(i,j) (\tilde{X}_j - D_j) \right\|_{\alpha}$$
$$\leq C_{\alpha} m^2 G_{6,n} \|X_1\|_{\alpha}^2.$$

Thus, by Burkholder inequality,

$$\left\| \sum_{h=0}^{\lfloor (n-i)/(4m) \rfloor} Y_{1,i+4mh} \right\|_{\frac{\alpha}{2}}^{\alpha'} \leq C_{\alpha} \sum_{h=0}^{\lfloor (n-i)/(4m) \rfloor} \left\| Y_{1,i+4mh} \right\|_{\frac{\alpha}{2}}^{\alpha'}$$
$$\leq C_{\alpha} m^{2\alpha'-1} n G_{6,n}^{\alpha'} \left\| X_1 \right\|_{\alpha}^{2\alpha'},$$

which implies

$$\left\| \sum_{h=0}^{\lfloor (n-i)/(4m) \rfloor} Y_{1,i+4mh} \right\|_{\frac{\alpha}{2}} \le C_{\alpha} m^{2-\frac{1}{\alpha'}} n^{\frac{1}{\alpha'}} G_{6,n} \left\| X_1 \right\|_{\alpha}^2$$

By Minkowski inequality,

$$\left\|\sum_{i=1}^{n} Y_{1,i}\right\|_{\frac{\alpha}{2}} \leq \sum_{i=1}^{4m-1} \left\|\sum_{h=0}^{\lfloor (n-i)/(4m) \rfloor} Y_{1,i+4mh}\right\|_{\frac{\alpha}{2}}$$
$$\leq C_{\alpha} m^{3-\frac{1}{\alpha'}} n^{\frac{1}{\alpha'}} G_{6,n} \left\|X_{1}\right\|_{\alpha}^{2}.$$
(E.15)

For $Y_{2,i}$, observe that

$$\mathbb{E}\left\{Y_{2,i+12m} - \mathbb{E}(Y_{2,i+12m}) \mid \mathscr{F}_i\right\} = \mathbb{E}(Y_{2,i+12m}) - \mathbb{E}(Y_{2,i+12m}) = 0.$$

Therefore, by (E.14) and similar arguments as in (E.15), we have

$$\left\|\sum_{i=1}^{n} \left\{ Y_{2,i} - \mathbb{E}(Y_{2,i}) \right\} \right\|_{\frac{\alpha}{2}} \le C_{\alpha} m^{3-\frac{1}{\alpha'}} n^{\frac{1}{\alpha'}} G_{6,n} \left\| X_1 \right\|_{\alpha}^2.$$

By Minkowski inequality and $\mathbb{E}|X_1|^{\alpha} < \infty$, we now have

$$I_{7} \leq \left\|\sum_{i=1}^{n} Y_{1,i}\right\|_{\frac{\alpha}{2}} + \left\|\sum_{i=1}^{n} \left\{Y_{2,i} - \mathbb{E}(Y_{2,i})\right\}\right\|_{\frac{\alpha}{2}} = O\left(m^{3-\frac{1}{\alpha'}} n^{\frac{1}{\alpha'}} G_{6,n}\right).$$
(E.16)

To upper bound the last term in (E.10), note that we can rearrange the order of summation by

$$\sum_{i=1}^{n} Y'_{i} = 2 \sum_{i=2}^{n} \left(\tilde{X}_{i} - D_{i} \right) \sum_{j=1}^{i-1} W_{n}(i,j) D_{j}$$
$$= 2 \sum_{j=1}^{n-1} D_{j} \sum_{i=j+1}^{n} W_{n}(i,j) \left(\tilde{X}_{i} - D_{i} \right).$$

Therefore, by similar arguments as in (E.13), (E.15) and (E.16),

$$I_8 = O\left(m^{3-\frac{1}{\alpha'}} n^{\frac{1}{\alpha'}} G_{6,n}\right).$$

Combining the results in (E.10) gives us

$$\begin{split} \left\| \tilde{\sigma}_{n}^{2} - \mathbb{E}(\tilde{\sigma}_{n}^{2}) - \breve{\sigma}_{n}^{2} + \mathbb{E}(\breve{\sigma}_{n}^{2}) \right\|_{\frac{\alpha}{2}} &= n^{-1} \left\{ O(n^{\frac{1}{\alpha'}}) + O(m^{1 - \frac{1}{\alpha'}} n^{\frac{1}{\alpha'}}) + O\left(m^{3 - \frac{1}{\alpha'}} n^{\frac{1}{\alpha'}} G_{6,n}\right) \right\} \\ &= O\left(m^{3 - \frac{1}{\alpha'}} n^{\frac{1}{\alpha'} - 1} G_{6,n}\right), \end{split}$$

since $\alpha' \in (1,2]$. For the additional case, note that m needs to be chosen such that $m \to \infty$ when $n \to \infty$ for m-dependent process approximation to work well. Under Assumptions 2 and 3(a),

$$G_{6,n} \le 2G_{2,n} = o(n^c),$$

for some constant $c \in (0, 1 - 1/\alpha')$. Therefore, We can choose $m = O(n^{(1-1/\alpha'-c)/3})$ such that $m \to \infty$ as $n \to \infty$, and $\|\tilde{\sigma}_n^2 - \mathbb{E}(\tilde{\sigma}_n^2) - \check{\sigma}_n^2 + \mathbb{E}(\check{\sigma}_n^2)\|_{\alpha/2} = o(1)$.

E.4 Lemma E.4

To establish the consistency or variance of our estimators under martingale difference approximation, we have the following lemma.

Lemma E.4. Let $\alpha > 2$. If Assumption 1 holds, then as $m \to \infty$,

$$||D_i|| \to \sigma \quad and \quad \operatorname{Var}(D_i D_j) \to \sigma^4,$$

for positive integers i and j where |i - j| > m.

Proof. Recall that the LRV can be expressed in a probabilistic representation $\sigma = \|\sum_{h=i}^{\infty} \mathscr{P}_i X_h\|$; see, e.g., Wu (2009). By Minkowski inequality, Theorem 1 of Wu (2005) and (E.7),

$$\begin{aligned} \left\| D_i - \sum_{h=i}^{\infty} \mathscr{P}_i X_h \right\| &= \left\| \sum_{h=0}^{\infty} \left\{ \mathbb{E}(\tilde{X}_{i+h} \mid \mathscr{F}_i) - \mathbb{E}(\tilde{X}_{i+h} \mid \mathscr{F}_{i-1}) \right\} - \sum_{h=i}^{\infty} \mathscr{P}_i X_h \right| \\ &= \left\| \sum_{h=i}^{\infty} \mathscr{P}_i (\tilde{X}_h - X_h) \right\| \\ &\leq \sum_{h=i}^{\infty} \left\| \tilde{X}_h - X_h - \tilde{X}_{h,\{i\}} + X_{h,\{i\}} \right\| \le C_2 d_{m,2}, \end{aligned}$$

where $d_{m,2}$ is defined in (E.4). As $m \to \infty$, $d_{m,2} = o(1)$ in view of Lemma E.5. Thus, we have

$$\lim_{n \to \infty} \left\| D_i - \sum_{h=i}^{\infty} \mathscr{P}_i X_h \right\| = 0.$$

γ

By Minkowski inequality,

$$\limsup_{m \to \infty} \|D_i\| \le \limsup_{m \to \infty} \left\| D_i - \sum_{h=i}^{\infty} \mathscr{P}_i X_h \right\| + \limsup_{m \to \infty} \left\| \sum_{h=i}^{\infty} \mathscr{P}_i X_h \right\| = \sigma.$$

At the same time,

$$\liminf_{m \to \infty} \left\| \sum_{h=i}^{\infty} \mathscr{P}_i X_h \right\| \le \liminf_{m \to \infty} \|D_i\| + \liminf_{m \to \infty} \left\| \sum_{h=i}^{\infty} \mathscr{P}_i X_h - D_i \right\|$$

which implies $\liminf_{m\to\infty} \|D_i\| \ge \sigma$. By Sandwich Theorem, we have $\lim_{m\to\infty} \|D_i\| = \sigma$. For $\operatorname{Var}(D_i D_j)$, recall that D_i 's form martingale differences with respect to \mathscr{F}_i and are *m*-dependent. Without loss of generality, assume i > j. Then $\mathbb{E}(D_i D_j) = \mathbb{E}\{D_j \mathbb{E}(D_i \mid \mathscr{F}_j)\} = 0$ and

$$\operatorname{Var}(D_i D_j) = \mathbb{E}(D_i^2 D_j^2) - \{\mathbb{E}(D_i D_j)\}^2$$
$$= \mathbb{E}(D_i^2)\mathbb{E}(D_j^2)$$

$$= \|D_i\|^2 \|D_j\|^2 \to \sigma^4,$$

where |i - j| > m as $m \to \infty$.

E.5 Lemma E.5

The following lemma provides an asymptotic upper bound of $d_{m,\alpha}$.

Lemma E.5. Let $\alpha > 2$. If Assumption 1 holds, then as $m \to \infty$, $d_{m,\alpha} = o(1)$.

Proof. Let M = f(m) such that $\Upsilon_{m+1,\alpha} > \sup_{i \ge M} \delta_{i,\alpha}$ and $M \to \infty$ as $m \to \infty$. Then

$$d_{m,\alpha} = \sum_{i=0}^{\infty} \min\left(\delta_{i,\alpha}, \Upsilon_{m+1,\alpha}\right)$$
$$\leq \sum_{i=0}^{M-1} \Upsilon_{m+1,\alpha} + \sum_{i=M}^{\infty} \delta_{i,\alpha} = M\Upsilon_{m+1,\alpha} + \sum_{i=M}^{\infty} \delta_{i,\alpha}.$$

As $m \to \infty$, $\sum_{i=M}^{\infty} \delta_{i,\alpha} \to 0$ and $\Upsilon_{m+1,\alpha} \to 0$ by Assumption 1. It suffices to show that $M\Upsilon_{m+1,\alpha} \to 0$, which we may apply the L'Hôpital's rule:

$$\lim_{m \to \infty} M \Upsilon_{m+1,\alpha} = \lim_{m \to \infty} \frac{f(m)}{1/\Upsilon_{m+1,\alpha}} = \lim_{m \to \infty} -\frac{f'(m)\Upsilon_{m+1,\alpha}^2}{\Upsilon_{m+1,\alpha}'},$$

which is 0 if

$$f'(m) = -\frac{\Upsilon'_{m+1,\alpha}}{\Upsilon_{m+1,\alpha}} \iff f(m) = -\ln(\Upsilon_{m+1,\alpha}) + C,$$

where C is an arbitrary constant. We check that $-\ln(\Upsilon_{m+1,\alpha}) \to \infty$ since $\Upsilon_{m+1,\alpha} \to 0$. To ensure f(m) is always positive, we can take $C = \ln(\Upsilon_{0,\alpha}) + 1$.

E.6 Lemma E.6

The following lemma provides asymptotic upper bounds of the sum of squared terms and the last incomplete block $\bar{\sigma}_{2,n}^2$ when $\phi > 1$, and the sum of squared terms $\bar{\sigma}_{4,n}^2$ when $\phi = 1$.

Lemma E.6. Let $\alpha \geq 4$. Suppose that $X_1 \in \mathscr{L}^{\alpha}$. If Assumption 1 holds, then

$$\left\|\bar{\sigma}_{2,n}^{2} - \mathbb{E}(\bar{\sigma}_{2,n}^{2})\right\|^{2} = o\left(n^{\psi + \max\{2q(\psi-\theta), 0\}-1}\right) \quad and \quad \left\|\bar{\sigma}_{4,n}^{2} - \mathbb{E}(\bar{\sigma}_{4,n}^{2})\right\|^{2} = O(n^{-1})$$

Proof. We first establish the stability of $\{X_i X_{i-k}\}$ for a fixed $k \in [0, i) \cap \mathbb{Z}$. By Minkowski inequality and Hölder's inequality,

$$\begin{split} \|X_i X_{i-k} - X_{i,\{0\}} X_{i-k,\{0\}} \|_{\frac{\alpha}{2}} &= \|X_i (X_{i-k} - X_{i-k,\{0\}}) + X_{i-k,\{0\}} (X_i - X_{i,\{0\}}) \|_{\frac{\alpha}{2}} \\ &\leq \delta_{i-k,\alpha} \|X_i\|_{\alpha} + \delta_{i,\alpha} \|X_{i-k,\{0\}}\|_{\alpha} \,. \end{split}$$

Hence under Assumption 1 and $X_i \in \mathcal{L}^{\alpha}$,

$$\sum_{i=0}^{\infty} \|X_i X_{i-k} - X_{i,\{0\}} X_{i-k,\{0\}}\|_{\frac{\alpha}{2}} \le \Delta_{\alpha} \left(\sup_{i \in \mathbb{N}_0} \|X_i\|_{\alpha} + \sup_{i \in \mathbb{N}_0} \|X_{i-k,\{0\}}\|_{\alpha} \right) < \infty,$$

which means $\{X_i X_{i-k}\}$ is $\alpha/2$ -stable for a fixed $k \in [0, i) \cap \mathbb{Z}$ (Wu, 2011). For $\alpha \ge 4$, it follows from Lemma 1 of Liu and Wu (2010) that

$$\left\|\bar{\sigma}_{4,n}^2 - \mathbb{E}(\bar{\sigma}_{4,n}^2)\right\| = \frac{1}{n} \left\|\sum_{i=1}^n \left\{X_i^2 - \mathbb{E}(X_i^2)\right\}\right\| = \frac{1}{n} \cdot O\left\{\left(\sum_{i=1}^n 1^2\right)^{\frac{1}{2}}\right\} = O(n^{-\frac{1}{2}}).$$

In addition, by Minkowski inequality and Lemma 1 of Liu and Wu (2010),

$$\begin{split} \left\|\bar{\sigma}_{2,n}^{2} - \mathbb{E}(\bar{\sigma}_{2,n}^{2})\right\| &\leq \frac{2}{n} \left\|\sum_{j=\eta_{s_{n}}}^{n} \sum_{k=1}^{s_{j}'} \left(1 - \frac{k^{q}}{t_{n}^{q}}\right) \{X_{j}X_{j-k} - \mathbb{E}(X_{j}X_{j-k})\}\right\| + O(n^{-\frac{1}{2}}) \\ &\leq \frac{2}{n} \sum_{k=1}^{\left\lceil \phi_{s_{n}} \right\rceil} \left\|\sum_{j=\eta_{s_{n}}}^{n} \left(1 - \frac{k^{q}}{t_{n}^{q}}\right) \mathbb{1}_{k \leq s_{j}'} \{X_{j}X_{j-k} - \mathbb{E}(X_{j}X_{j-k})\}\right\| + O(n^{-\frac{1}{2}}) \\ &\leq O(n^{-1}) \sum_{k=1}^{\left\lceil \phi_{s_{n}} \right\rceil} \sqrt{\sum_{j=\eta_{s_{n}}}^{n} \left|1 - \frac{k^{q}}{t_{n}^{q}}\right|^{2}} + O(n^{-\frac{1}{2}}) \\ &= O(n^{\psi + \max\{q(\psi - \theta, 0\} - 1\})} + O(n^{-\frac{1}{2}}). \end{split}$$

Check that $2\psi + \max\{2q(\psi-\theta), 0\} - 2 < \psi + \max\{2q(\psi-\theta), 0\} - 1 \text{ and } -1 < \psi + \max\{2q(\psi-\theta), 0\} - 1$ always hold under Definition 1.

E.7 Lemma E.7

Define

$$H_{7,\phi} \coloneqq \begin{cases} \frac{2}{n} \sum_{h=\lceil (m+1)/\phi \rceil}^{s_n-1} \sum_{i=1}^{\nu_h} \sum_{j=\eta_h+(i-1)\varpi_h+m+1}^{\eta_h+i\varpi_h-1} \sum_{k=m+1}^{j-\eta_h-(i-1)\varpi_h+h} \left(1-\frac{k^q}{t_n^q}\right) D_j D_{j-k}, & \phi > 1; \\ \frac{2}{n} \sum_{h=m+1}^{s_n} \sum_{i=\eta_h}^n \left(1-\frac{h^q}{t_n^q}\right) D_i D_{i-h}, & \phi = 1. \end{cases}$$
(E.17)

Note that $\operatorname{Var}(H_{7,\phi}) = H_3^2$ for $\phi > 1$, and $\operatorname{Var}(H_{7,\phi}) = H_5^2$ for $\phi = 1$, where H_3 and H_5 are defined in (B.17) and (B.19), respectively. The following lemma provides the exact convergence rate of $\operatorname{Var}(H_{7,\phi})$.

Lemma E.7. Let $\alpha \geq 4$. Suppose that $X_1 \in \mathscr{L}^{\alpha}$. If Assumption 1 holds, then as $n \to \infty$,

$$\operatorname{Var}(H_{7,\phi}) \sim \mathscr{V}_{\psi,\Psi,\theta,\Theta,q,\phi} \sigma^4 n^{\psi+\max\{2q(\psi-\theta),0\}-1}$$

where $\mathscr{V}_{\psi,\Psi,\theta,\Theta,q,\phi}$

$$= \begin{cases} 4 \int_{0}^{\Psi} \int_{0}^{\nu_{w}} \int_{0}^{(\phi-1)w} \int_{0}^{w+y} \left(\mathbb{1}_{\psi \leq \theta} - 2z^{q} \Theta^{-q} \mathbb{1}_{\psi=\theta} + z^{2q} \Theta^{-2q} \mathbb{1}_{\psi \geq \theta} \right) dz dy dx dw, \quad \phi > 1; \\ 4 \int_{0}^{\Psi} \int_{(w/\Psi)^{1/\psi}}^{1} \left(\mathbb{1}_{\psi \leq \theta} - 2w^{q} \Theta^{-q} \mathbb{1}_{\psi=\theta} + w^{2q} \Theta^{-2q} \mathbb{1}_{\psi \geq \theta} \right) dx dw, \qquad \phi = 1 \end{cases}$$
$$= \begin{cases} \frac{2\Psi(\phi+1)}{\psi+1} \mathbb{1}_{\psi \leq \theta} - \frac{8\Psi^{q+1}\Theta^{-q}(\phi^{q+2}-1)}{(\phi-1)(q+1)(q+2)(\psi q+\psi+1)} \mathbb{1}_{\psi=\theta} + \frac{2\Psi^{2q+1}\Theta^{-2q}(\phi^{2q+2}-1)}{(\phi-1)(q+1)(2q+1)(2\psi q+\psi+1)} \mathbb{1}_{\psi \geq \theta}, \quad \phi > 1; \\ \frac{4\Psi}{\psi+1} \mathbb{1}_{\psi \leq \theta} - \frac{8\Psi^{q+1}\Theta^{-q}}{(q+1)(\psi q+\psi+1)} \mathbb{1}_{\psi=\theta} + \frac{4\Psi^{2q+1}\Theta^{-2q}}{(2q+1)(2\psi q+\psi+1)} \mathbb{1}_{\psi \geq \theta}, \qquad \phi = 1. \end{cases}$$

Proof. Consider $\phi > 1$ first. Recall that D_i 's form martingale differences with respect to \mathscr{F}_i and are *m*-dependent. Therefore, $\mathbb{E}(D_i D_{i-k}) = 0$ if k > m, and $\mathbb{E}(D_i D_{i-k} D_j D_{j-k}) = 0$ if k > m and $i \neq j$. By Lemma E.4, we have $\operatorname{Var}(D_i D_{i-k}) \to \sigma^4$ for k > m as $m \to \infty$. Thus, as $n \to \infty$,

$$\operatorname{Var}(H_{7,\phi}) = \frac{4}{n^2} \operatorname{Var}\left\{ \sum_{h=\lceil (m+1)/\phi \rceil}^{s_n-1} \sum_{i=1}^{\nu_h} \sum_{j=\eta_h+(i-1)\varpi_h+m+1}^{\eta_h+i\varpi_h-1} \sum_{k=m+1}^{j-\eta_h-(i-1)\varpi_h+h} \left(1 - \frac{k^q}{t_n^q}\right) D_j D_{j-k} \right\}$$
$$\sim \frac{4\sigma^4}{n^2} \sum_{h=\lceil (m+1)/\phi \rceil}^{s_n-1} \sum_{i=1}^{\nu_h} \sum_{j=\eta_h+(i-1)\varpi_h+m+1}^{\eta_h+i\varpi_h-1} \sum_{k=m+1}^{j-\eta_h-(i-1)\varpi_h+h} \left(1 - \frac{k^q}{t_n^q}\right)^2.$$
(E.18)

Since we can choose $m = o(n^{\psi/6})$ as in Section B.2.3, we can approximate (E.18) by Riemann integrals and use change of variables to obtain

$$\begin{aligned} \operatorname{Var}(H_{7,\phi}) &\sim \frac{4\sigma^4}{n^2} \int_0^{s_n} \int_0^{\nu_h} \int_{\eta_h + (i-1)\varpi_h}^{\eta_h + i\varpi_h} \int_0^{j-\eta_h - (i-1)\varpi_h + h} \left(1 - \frac{k^q}{t_n^q}\right)^2 \, \mathrm{d}k \, \mathrm{d}j \, \mathrm{d}i \, \mathrm{d}h \\ &= \frac{4\sigma^4}{n^2} \int_0^{s_n} \int_0^{\nu_h} \int_0^{\varpi_h} \int_0^{h+j} \left(1 - \frac{k^q}{t_n^q}\right)^2 \, \mathrm{d}k \, \mathrm{d}j \, \mathrm{d}i \, \mathrm{d}h \\ &= \begin{cases} 4\sigma^4 n^{\psi-1} \int_0^{\Psi} \int_0^{\nu_w} \int_0^{(\phi-1)w} \int_0^{w+y} 1 \, \mathrm{d}z \, \mathrm{d}y \, \mathrm{d}x \, \mathrm{d}w, & \psi < \theta; \\ 4\sigma^4 n^{\psi-1} \int_0^{\Psi} \int_0^{\nu_w} \int_0^{(\phi-1)w} \int_0^{w+y} (1 - z^q \Theta^{-q})^2 \, \mathrm{d}z \, \mathrm{d}y \, \mathrm{d}x \, \mathrm{d}w, & \psi = \theta; \\ 4\sigma^4 n^{\psi+2q(\psi-\theta)-1} \int_0^{\Psi} \int_0^{\nu_w} \int_0^{(\phi-1)w} \int_0^{w+y} z^{2q} \Theta^{-2q} \, \mathrm{d}z \, \mathrm{d}y \, \mathrm{d}x \, \mathrm{d}w, & \psi > \theta \\ &= \mathscr{V}_{\psi,\Psi,\theta,\Theta,q,\phi} \sigma^4 n^{\psi+\max\{2q(\psi-\theta),0\}-1}. \end{aligned}$$

Evaluating the integral gives

$$\mathscr{V}_{\psi,\Psi,\theta,\Theta,q,\phi} = \frac{2\Psi(\phi+1)}{\psi+1} \mathbb{1}_{\psi \le \theta} - \frac{8\Psi^{q+1}\Theta^{-q}(\phi^{q+2}-1)}{(\phi-1)(q+1)(q+2)(\psi q+\psi+1)} \mathbb{1}_{\psi=\theta} + \frac{2\Psi^{2q+1}\Theta^{-2q}(\phi^{2q+2}-1)}{(\phi-1)(q+1)(2q+1)(2\psi q+\psi+1)} \mathbb{1}_{\psi \ge \theta}.$$

Now consider $\phi = 1$. By similar arguments as in (E.18),

$$\operatorname{Var}(H_{7,\phi}) \sim \frac{4\sigma^4}{n} \sum_{h=m+1}^{s_n} \sum_{i=\eta_h}^n \left(1 - \frac{h^q}{t_n^q}\right)^2.$$

Using Riemann integral approximation and change of variables,

$$\operatorname{Var}(H_{7,\phi}) \sim \frac{4\sigma^4}{n^2} \int_0^{s_n} \int_{\eta_h}^n \left(1 - \frac{h^q}{t_n^q}\right)^2 \mathrm{d}i \, \mathrm{d}h$$

$$= \begin{cases} 4\sigma^4 n^{\psi-1} \int_0^{\Psi} \int_{(w/\Psi)^{1/\psi}}^1 1 \, \mathrm{d}x \, \mathrm{d}w, & \psi < \theta; \\ 4\sigma^4 n^{\psi-1} \int_0^{\Psi} \int_{(w/\Psi)^{1/\psi}}^1 (1 - w^q \Theta^{-q})^2 \, \mathrm{d}x \, \mathrm{d}w, & \psi = \theta; \\ 4\sigma^4 n^{\psi+2q(\psi-\theta)-1} \int_0^{\Psi} \int_{(w/\Psi)^{1/\psi}}^1 w^{2q} \Theta^{-2q} \, \mathrm{d}x \, \mathrm{d}w, & \psi > \theta \end{cases}$$
$$= \mathscr{V}_{\psi,\Psi,\theta,\Theta,q,\phi} \sigma^4 n^{\psi+\max\{2q(\psi-\theta),0\}-1}.$$

Evaluating the integral gives

$$\mathcal{V}_{\psi,\Psi,\theta,\Theta,q,\phi} = \frac{4\Psi}{\psi+1} \mathbb{1}_{\psi \le \theta} - \frac{8\Psi^{q+1}\Theta^{-q}}{(q+1)(\psi q + \psi + 1)} \mathbb{1}_{\psi=\theta} + \frac{4\Psi^{2q+1}\Theta^{-2q}}{(2q+1)(2\psi q + \psi + 1)} \mathbb{1}_{\psi \ge \theta}.$$

F Ancillary Results

F.1 Positive Definiteness Adjustment

A practical concern in LRCM estimation is that the finite-sample estimates may not be positive definite. Indeed, classical estimators such as those utilizing the overlapping batch means and Bartlett kernel are only guaranteed to be positive semidefinite. To address this issue, we follow Jentsch and Politis (2015) to provide an adjustment that retains asymptotic properties of LRCM estimators:

- (1) Obtain the diagonal matrix $\hat{V} = \text{diag}(\hat{\Sigma}_n)$.
- (2) Compute the correlation matrix $\hat{\boldsymbol{R}}_n = \hat{\boldsymbol{V}}^{-1/2} \hat{\boldsymbol{\Sigma}}_n \hat{\boldsymbol{V}}^{-1/2}$.
- (3) Eigendecompose the correlation matrix into $\hat{\boldsymbol{R}}_n = \hat{\boldsymbol{Q}} \hat{\Lambda}_n \hat{\boldsymbol{Q}}^{\mathsf{T}}$, where $\hat{\boldsymbol{Q}}$ is an orthogonal matrix whose columns are the eigenvectors and $\hat{\Lambda}_n = \text{diag}(\hat{\lambda}_1, \dots, \hat{\lambda}_d)$ is the diagonal matrix whose diagonal elements are the eigenvalues.
- (4) Set $\hat{\Lambda}_n^+ = \operatorname{diag}(\hat{\lambda}_1^+, \dots, \hat{\lambda}_d^+)$, where $\hat{\lambda}_i^+ = \max(\hat{\lambda}_i, an^{-b})$ for some user-specified a > 0 and b > 1/2. We use $a = \{\ln(n)/d\}^{1/2}$ and b = 9/10 as suggested in Vats and Flegal (2021).
- (5) Return the positive definite estimate $\hat{\Sigma}_n^+ = \hat{V}^{1/2} \hat{Q}^\top \hat{\Lambda}_n^+ \hat{Q} \hat{V}^{1/2}$.

Since this procedure costs O(d) time and space, it is able to preserve the computational properties of online estimators.

F.2 Example 2: Window of Selection Rules

In this subsection, we try to find the window associated with the subsampling estimator defined in (1.4). Recall that

$$\hat{\sigma}_{n,\mathsf{sub}}^2 \coloneqq \frac{\sum_{i=1}^n \left\{ \sum_{j=i-\ell_i+1}^i (X_j - \bar{X}_n) \right\}^2}{\sum_{i=1}^n \ell_i}.$$

Let $\bar{\ell}_n = n^{-1} \sum_{i=1}^n \ell_i$ be the global average batch size. Assume that $t \mapsto t - \ell_t$ is a monotonically increasing sequence, which is satisfied by $\triangle SR$, TSR and PSR. Then, we can write

$$\hat{\sigma}_{n,\mathsf{sub}}^{2} = \frac{1}{n\bar{\ell}_{n}} \sum_{i=1}^{n} \sum_{j=i-\ell_{i}+1}^{i} \sum_{j'=i-\ell_{i}+1}^{i} (X_{j} - \bar{X}_{n}) (X_{j'} - \bar{X}_{n})$$
$$= \frac{1}{n\bar{\ell}_{n}} \sum_{j=1}^{n} \sum_{j'=1}^{n} \sum_{i\in I_{n}(j,j')}^{n} (X_{j} - \bar{X}_{n}) (X_{j'} - \bar{X}_{n})$$
$$= \frac{1}{n} \sum_{i=1}^{n} \sum_{j=1}^{n} \frac{|I_{n}(i,j)|}{\bar{\ell}_{n}} (X_{i} - \bar{X}_{n}) (X_{j} - \bar{X}_{n}),$$

where $I_n(i,j) = \{t \in \{1,\ldots,n\} : i, j \in \{t - \ell_t + 1,\ldots,t\}\}$. Denote $f_n(i) = \max\{t \in \{1,\ldots,n\} : i \in \{t - \ell_t + 1,\ldots,t\}\}$ and $b_n(i) = f_n(i) - i + 1$. Physically, the meaning of $f_n(i)$ is the largest index such that the *i*-th observation is still included in the subsample, while $b_n(i)$ is the subsample size of the $f_n(i)$ -th subsample. We can express the cardinality of $I_n(i,j)$ as

$$|I_n(i,j)| = \left\{b_n(i \wedge j) - |i-j|\right\} \mathbb{1}_{|i-j| \le b_n(i \wedge j)}.$$

Consequently, $\hat{\sigma}_{n,\mathsf{sub}}^2$ is exactly equivalent to $\hat{\sigma}_n^2(W_{\mathsf{sub}})$ with

$$W_{\mathsf{sub}} = W_{n,\mathsf{sub}}(i,j) = \left(1 - \frac{|i-j| + \bar{\ell}_n - b_n(i \wedge j)}{\bar{\ell}_n}\right) \mathbb{1}_{|i-j| \le b_n(i \wedge j)}.$$

Note that the subsampling parameter admits the form $b_n(i \wedge j)$ because the way to subsample differs under the subsampling form in (1.4) and the quadratic form in (2.1). Nevertheless, we can approximate W_{sub} under PSR asymptotically by

$$W_{\mathsf{PSR}} = W_{n,\mathsf{PSR}}(i,j) = \left(1 - \frac{|i-j| + \bar{\ell}_n - \ell_{i \lor j}}{\bar{\ell}_n}\right) \mathbb{1}_{|i-j| \le \ell_{i \lor j}}$$

We formalize this result with the following proposition.

Proposition F.1. Let $\alpha \ge 4$. Suppose that $X_1 \in \mathscr{L}^{\alpha}$. If Assumptions 1 and 5 hold for q = 1, and $\ell_i = \min(\lfloor \Lambda i^{\lambda} \rfloor, i)$ for some $\Lambda \in \mathbb{R}^+$ and $\lambda \in (0, 1/2)$, i.e., PSR is used, then

$$\frac{\left\|\hat{\sigma}_{n}^{2}(W_{\mathsf{sub}}) - \hat{\sigma}_{n}^{2}(W_{\mathsf{PSR}})\right\|^{2}}{\left\|\hat{\sigma}_{n}^{2}(W_{\mathsf{sub}}) - \sigma^{2}\right\|^{2}} = o(1).$$

Proof. When Assumptions 1 and 5 hold for q = 1, we can apply Theorem 3 of Chan and Yau

(2017a) to obtain $\|\hat{\sigma}_n^2(W_{sub}) - \sigma^2\|^2 = O(n^{\max(-2\lambda,\lambda-1)})$. It suffices to show that

$$\left\|\hat{\sigma}_n^2(W_{\mathsf{sub}}) - \hat{\sigma}_n^2(W_{\mathsf{PSR}})\right\| = o(n^{\max(-\lambda,\frac{\lambda-1}{2})}).$$

Now, recall the definition of the known-mean version $\bar{\sigma}_n^2$ in (B.2). We can follow Section B.2.1 to replace the sample mean similarly and assume $\mu = 0$ without loss of generality. This is because our Lemma E.1 and the Lemma E.1 in Chan and Yau (2017a) give

$$\left\|\hat{\sigma}_{n}^{2}(W_{\mathsf{PSR}}) - \bar{\sigma}_{n}^{2}(W_{\mathsf{PSR}})\right\| = O(n^{\lambda - 1}) \quad \text{and} \quad \left\|\hat{\sigma}_{n}^{2}(W_{\mathsf{sub}}) - \bar{\sigma}_{n}^{2}(W_{\mathsf{sub}})\right\| = O(n^{\lambda - 1})$$

Check that $\lambda - 1 < \max(-\lambda, \lambda - 1/2)$ always hold because $\lambda \in (0, 1/2)$. Therefore, we only need to prove

$$\left\|\bar{\sigma}_n^2(W_{\mathsf{sub}}) - \bar{\sigma}_n^2(W_{\mathsf{PSR}})\right\| = o(n^{\max(-\lambda,\frac{\lambda-1}{2})})$$

Note that we can express $\bar{\sigma}_n^2(W_{\sf sub})$ and $\bar{\sigma}_n^2(W_{\sf PSR})$ by

$$\bar{\sigma}_{n}^{2}(W_{\mathsf{sub}}) = \frac{1}{n\bar{\ell}_{n}} \sum_{i=1}^{n} \sum_{j=1}^{n} \left\{ b_{n}(i \wedge j) - |i-j| \right\} \mathbb{1}_{|i-j| \le b_{n}(i \wedge j)} X_{i} X_{j},$$
$$\bar{\sigma}_{n}^{2}(W_{\mathsf{PSR}}) = \frac{1}{n\bar{\ell}_{n}} \sum_{i=1}^{n} \sum_{j=1}^{n} \left(\ell_{i \vee j} - |i-j| \right) \mathbb{1}_{|i-j| \le \ell_{i \vee j}} X_{i} X_{j}.$$

Denote $F_n(i) = \max\{t \in \mathbb{Z}^+ : i \in \{t - \ell_t + 1, \dots, t\}\}, \ \Xi_{1,n} = \{i \in \{1, \dots, n\} : b_n(i) - \ell_i > 1\}, \ \Xi_{2,n} = \{i \in \{1, \dots, n\} : f_n(i) < F_n(i)\}, \ \text{and} \ \Xi_{3,n} = \{1, \dots, n\} \setminus \{\Xi_{1,n} \cup \Xi_{2,n}\}.$ The meaning of $\Xi_{1,n}$ is the set of indices in the beginning where ℓ_i is strictly increasing, while the meaning of $\Xi_{2,n}$ is the set of indices that belongs to the last incomplete block where $F_n(i)$ is truncated by n. Then, we partition $\overline{\sigma}_n^2(W_{sub})$ into

$$\begin{split} \bar{\sigma}_{1,n}^{2}(W_{\mathsf{sub}}) &= \frac{1}{n\bar{\ell}_{n}} \sum_{i \in \Xi_{1,n}} \sum_{j=1}^{n} \left\{ b_{n}(i \wedge j) - |i - j| \right\} \mathbb{1}_{|i - j| \leq b_{n}(i \wedge j)} X_{i} X_{j} \\ &+ \frac{1}{n\bar{\ell}_{n}} \sum_{i \in \Xi_{3,n}} \sum_{j \in \Xi_{1,n}} \left\{ b_{n}(i \wedge j) - |i - j| \right\} \mathbb{1}_{|i - j| \leq b_{n}(i \wedge j)} X_{i} X_{j}, \\ \bar{\sigma}_{2,n}^{2}(W_{\mathsf{sub}}) &= \frac{1}{n\bar{\ell}_{n}} \sum_{i \in \Xi_{2,n}} \sum_{j=1}^{n} \left\{ b_{n}(i \wedge j) - |i - j| \right\} \mathbb{1}_{|i - j| \leq b_{n}(i \wedge j)} X_{i} X_{j} \\ &+ \frac{1}{n\bar{\ell}_{n}} \sum_{i \in \Xi_{3,n}} \sum_{j \in \Xi_{2,n}} \left\{ b_{n}(i \wedge j) - |i - j| \right\} \mathbb{1}_{|i - j| \leq b_{n}(i \wedge j)} X_{i} X_{j}, \\ \bar{\sigma}_{3,n}^{2}(W_{\mathsf{sub}}) &= \frac{1}{n\bar{\ell}_{n}} \sum_{i \in \Xi_{3,n}} \sum_{j \in \Xi_{3,n}} \left\{ b_{n}(i \wedge j) - |i - j| \right\} \mathbb{1}_{|i - j| \leq b_{n}(i \wedge j)} X_{i} X_{j}, \end{split}$$

and similarly for $\bar{\sigma}_n^2(W_{\mathsf{PSR}})$ with $\{b_n(i \wedge j) - |i-j|\}\mathbb{1}_{|i-j| \leq b_n(i \wedge j)}$ replaced by $(\ell_{i \vee j} - |i-j|)\mathbb{1}_{|i-j| \leq \ell_{i \vee j}}$.

By Minkowski inequality,

$$\begin{split} \left\| \bar{\sigma}_{n}^{2}(W_{\mathsf{sub}}) - \bar{\sigma}_{n}^{2}(W_{\mathsf{PSR}}) \right\| &\leq \left\| \bar{\sigma}_{1,n}^{2}(W_{\mathsf{sub}}) - \bar{\sigma}_{1,n}^{2}(W_{\mathsf{PSR}}) \right\| + \left\| \bar{\sigma}_{2,n}^{2}(W_{\mathsf{sub}}) - \bar{\sigma}_{2,n}^{2}(W_{\mathsf{PSR}}) \right\| \\ &+ \left\| \bar{\sigma}_{3,n}^{2}(W_{\mathsf{sub}}) - \bar{\sigma}_{3,n}^{2}(W_{\mathsf{PSR}}) \right\| \\ &=: H_{8} + H_{9} + H_{10}. \end{split}$$

For H_8 , since $\Xi_{1,n}$ is the set of indices in the beginning where ℓ_i is strictly increasing, $\max_{i \in \Xi_{1,n}} i \leq C$ for some constant $C \in \mathbb{R}^+$ that may change from line to line in the remaining of this proof. Without loss of generality, consider $i \geq j$ as in (2.5). By Minkowski inequality and the meaning of $\Xi_{1,n}$,

$$H_{8} \leq \frac{2}{n\bar{\ell}_{n}} \left\| \sum_{i=1}^{C+\ell_{n}} \sum_{j=1}^{C} \left\{ b_{n}(i \wedge j) - |i-j| \right\} \mathbb{1}_{|i-j| \leq b_{n}(i \wedge j)} X_{i} X_{j} \right\| \\ + \frac{2}{n\bar{\ell}_{n}} \left\| \sum_{i=1}^{C+\ell_{n}} \sum_{j=1}^{C} \left(\ell_{i \vee j} - |i-j| \right) \mathbb{1}_{|i-j| \leq \ell_{i \vee j}} X_{i} X_{j} \right\| \\ =: H_{11} + H_{12}.$$

Consider H_{11} , Minkowski inequality gives

$$H_{11} \leq \frac{2}{n\bar{\ell}_n} \left\| \sum_{i=1}^{C+\ell_n} \sum_{j=1}^{C} \left\{ b_n(i \wedge j) - |i-j| \right\} \mathbb{1}_{|i-j| \leq b_n(i \wedge j)} \left\{ X_i X_j - \mathbb{E}(X_i X_j) \right\} \right\| \\ + \frac{2}{n\bar{\ell}_n} \left\| \sum_{i=1}^{C+\ell_n} \sum_{j=1}^{C} \left\{ b_n(i \wedge j) - |i-j| \right\} \mathbb{1}_{|i-j| \leq b_n(i \wedge j)} \mathbb{E}(X_i X_j) \right\| \\ =: H_{13} + H_{14}.$$
(F.1)

Since the stability of $\{X_iX_j\}$ for $j \in \{1, ..., i\}$ has been established in Section E.6, it follows from Assumption 1 and Lemma 1 of Liu and Wu (2010) that

$$H_{13} \le O(n^{-1-\lambda}) \sqrt{\sum_{i=1}^{C+\ell_n} \left| \sum_{j=1}^{C} C \right|^2} = O(n^{-1-\lambda}) O(n^{\frac{\lambda}{2}}) = o(n^{\max(-\lambda, \frac{\lambda-1}{2})}).$$
(F.2)

On the other hand, $\sum_{k \in \mathbb{Z}} |\gamma_k| < \infty$ under Assumption 1; see Wu (2009). By Minkowski inequality,

$$H_{14} \le O(n^{-1-\lambda}) \sum_{i=1}^{C+\ell_n} \sum_{j=1}^{C} C|\gamma_{|i-j|}| = O(n^{-1-\lambda})O(1) = o(n^{\max(-\lambda, \frac{\lambda-1}{2})}).$$
(F.3)

Using the same procedure from (F.1) to (F.3), we have

$$H_{12} \le O(n^{-1-\lambda}) \sqrt{\sum_{i=1}^{C+\ell_n} \left|\sum_{j=1}^{C} \ell_n\right|^2} + O(n^{-1-\lambda}) \sum_{i=1}^{C+\ell_n} \sum_{j=1}^{C} \ell_n |\gamma_{|i-j|}|$$

$$= O(n^{-1-\lambda})O(n^{\frac{3\lambda}{2}}) + O(n^{-1-\lambda})O(n^{\lambda}) = o(n^{\max(-\lambda,\frac{\lambda-1}{2})}).$$

For H_9 , since the meaning of $\Xi_{2,n}$ is the set of indices that belongs to the last incomplete block where $F_n(i)$ is truncated by n, $\min_{i \in \Xi_{2,n}} i \ge (\ell_n / \Lambda)^{1/\lambda} - C = n - C$; see (B.10). Repeating the same arguments, we have

$$H_9 = o(n^{\max(-\lambda, \frac{\lambda-1}{2})}).$$

Finally, note that for $i, j \in \Xi_{3,n}$, we have $\{b_n(i \wedge j) - \ell_{i \vee j}\}\mathbb{1}_{|i-j| \leq b_n(i \wedge j)} = 1$ or $\{b_n(i \wedge j) - \ell_{i \vee j}\}\mathbb{1}_{|i-j| \leq b_n(i \wedge j)} = 0$. By Minkowski inequality,

$$H_{10} = \frac{1}{n\bar{\ell}_n} \left\| \sum_{i \in \Xi_{3,n}} \sum_{j \in \Xi_{3,n}} \left\{ b_n(i \wedge j) - \ell_{i \vee j} \right\} \mathbb{1}_{|i-j| \leq b_n(i \wedge j)} X_i X_j \right\|$$

$$\leq \frac{1}{n\bar{\ell}_n} \left\| \sum_{i \in \Xi_{3,n}} \sum_{j \in \Xi_{3,n}} \left\{ b_n(i \wedge j) - \ell_{i \vee j} \right\} \mathbb{1}_{|i-j| \leq b_n(i \wedge j)} \left\{ X_i X_j - \mathbb{E}(X_i X_j) \right\} \right\|$$

$$+ \frac{1}{n\bar{\ell}_n} \left| \sum_{i \in \Xi_{3,n}} \sum_{j \in \Xi_{3,n}} \left\{ b_n(i \wedge j) - \ell_{i \vee j} \right\} \mathbb{1}_{|i-j| \leq b_n(i \wedge j)} \mathbb{E}(X_i X_j) \right|$$

$$=: H_{15} + H_{16}.$$

Without loss of generality, consider $i \ge j$ again. By similar reasons as in (F.2) and (F.3),

$$H_{15} \leq O(n^{-1-\lambda}) \sqrt{\sum_{i \in \Xi_{3,n}} \left| \sum_{j=\min_{k \in \Xi_{3,n}} k}^{i} \left\{ b_{n}(i \wedge j) - \ell_{i \vee j} \right\} \mathbb{1}_{|i-j| \leq b_{n}(i \wedge j)} \right|^{2} }$$

= $O(n^{-\frac{1}{2}}) = o(n^{\max(-\lambda, \frac{\lambda-1}{2})}),$
 $H_{16} \leq O(n^{-1-\lambda}) \sum_{i \in \Xi_{3,n}} \sum_{j \in \Xi_{3,n}} \mathbb{1}_{|i-j| \leq \ell_{n}} |\gamma_{|i-j|}|$
= $O(n^{-1-\lambda})O(n^{\lambda}) = o(n^{\max(-\lambda, \frac{\lambda-1}{2})}).$

Combining the results completes the proof.

F.3 Example 5: Other Cases

Figure 9 reports the results for $m + k^* = 601$ (top panel) and $m + k^* = 1001$ (bottom panel), which are similar to the results for $m + k^* = 1401$ in Example 5.



Figure 9: Online CP detection at 5% nominal size using different LRV estimation methods: (a) fix (dotted gray); (b) offline (dashed red); (c) online (longdash blue). The locations of the CP are $m + k^* = 601$ (top panel) and $m + k^* = 1001$ (bottom panel).

F.4 Example 7: Details of Model

Consider an ARMA(1,1) model

$$X_i - \mu = \alpha (X_{i-1} - \mu) + \beta \varepsilon_{i-1} + \varepsilon_i \quad \text{where} \quad \varepsilon_i \stackrel{\text{ind}}{\sim} \mathcal{N}(0, \nu^2).$$

Data in Example 7 is simulated as follows:

$$\begin{split} W &\sim \mathrm{N}(0,1), \\ Z &\sim \mathrm{ARMA}(1,1) & \text{with} & \alpha = 0.5, \ \beta = 0.5, \ \nu = 1, \ \mu = 0, \\ e_x &\sim \mathrm{ARMA}(1,1) & \text{with} & \alpha = 0.6, \ \beta = 0.6, \ \nu = 1, \ \mu = 0, \\ e_y &\sim \mathrm{ARMA}(1,1) & \text{with} & \alpha = -0.7, \ \beta = -0.7, \ \nu = 1, \ \mu = 0 \\ X &= aZ + W + e_x, \end{split}$$

Algorithm F.1: Modified SASA+

[1] input: (i) η_0, τ – the initial learning rate and its drop factor [2] (ii) $T_{\rm burn}$, $N_{\rm min}$ – the burn-in size and minimum sample size for testing [3] (iii) K_{test} – the period to perform hypothesis test [4] (iv) α – the significance level of hypothesis testing [5] begin [6] Set $k = 0, t = 0, \eta = \eta_0$ [7] while not the end of training do [8] Randomly sample ζ_k [9] Compute the gradient g_k [10] Set $w_{k+1} = w_k - \eta g_k$ [11] Set $d_k = w_k \cdot g_k - 0.5\eta \|g_k\|_2^2$, where \cdot denotes the dot product [12] Set $N=t-T_{\rm burn}+1$ [13] if $t > T_{\text{burn}}$ and $k \mod K_{\text{test}} == 0$ then [14] Update $\bar{d}_N, \hat{\sigma}_N^2$ with $d_{k+1-\min(K_{\text{test}}, t-T_{\text{burn}})}, \ldots, d_k$ [15] else if $t == T_{\text{burn}}$ then [16] L Initialize $\overline{d_N}, \hat{\sigma}_N^2$ with d_k [17] Set t = t + 1[18] if $N > N_{\min}$ and $k \mod K_{\text{test}} == 0$ then [19] if $\underbrace{0 \in \bar{d}_N \pm z_{1-\alpha/2} \hat{\sigma}_N / \sqrt{N}}_{\text{Set } t = 0, \eta = \tau \eta}$ then [20] [21] Set k = k + 1[22]

$$Y = bX + W + e_y.$$

Same as Gupta et al. (2021), a = 1 and b = 1 are used.

F.5 Example 9: Details of SASA+

In Example 9, we change the offline LRV estimator used in the original SASA+ Zhang et al. (2020) to our mini-batch LASER(1,1). Now, we state the modified SASA+ in Algorithm F.1.

In SASA, d_N was a moving average so as to "diminish the effect of 'initialization bias' due to starting outside of the stationary distribution" (Lang et al., 2019). As seen in Example 6, another reason could be the computational burden of offline LRV estimators. To deal with the these issues, Zhang et al. (2020) further reduced the sample size of the moving average and conducted the hypothesis test every K_{test} iterations in their SASA+. In our modified SASA+, we use burn-in and mini-batch LRV estimators instead. The parameter K_{test} perfectly matches the idea of our mini-batch size. We summarize the parameters used in Example 9 below:

• Initial learning rate: the average of the first 5000 learning rates used in Vanilla SGD, i.e., $\eta_0 = 5000^{-1} \sum_{k=1}^{5000} 0.5k^{-0.505}$. This allows a relatively fair start when $N_{\min} = 5000$. Note that in addition to SASA+, Zhang et al. (2020) proposed an automatic method to select η_0 , which was not included in Example 9.

- Learning rate drop factor: $\tau = 0.8$.
- Burn-in size: $T_{\text{burn}} = 2500$.
- Minimum sample size for testing: $N_{\min} = 5000$.
- Period to perform test: $K_{\text{test}} = 500$.
- Significance level: $\alpha = 0.05$.

References

- Andrews, D. W. K. (1991) Heteroskedasticity and autocorrelation consistent covariance matrix estimation. *Econometrica*, 59, 817–858.
- Andrews, D. W. K. and Monahan, J. C. (1992) An improved heteroskedasticity and autocorrelation consistent covariance matrix estimator. *Econometrica*, **60**, 953–966.
- Beji, S. (2018) A fundamental relationship of polynomials and its proof. Advances in Pure Mathematics, 8, 559–563.
- Brooks, S., Gelman, A., Jones, G. and Meng, X.-L. (2011) Handbook of Markov Chain Monte Carlo. London: Chapman & Hall/CRC.
- Chan, K. W. (2022) Optimal difference-based variance estimators in time series: A general framework. The Annals of Statistics, 1376–1400.
- Chan, K. W. and Yau, C. Y. (2016) New recursive estimators of the time-average variance constant. Statistics and Computing, 26, 609–627.
- (2017a) Automatic optimal batch size selection for recursive estimators of time-average covariance matrix. Journal of the American Statistical Association, **112**, 1076–1089.
- (2017b) High-order corrected estimator of asymptotic variance with optimal bandwidth. Scandinavian Journal of Statistics, 44, 866–898.
- Chen, X., Lee, J. D., Tong, X. T. and Zhang, Y. (2020) Statistical inference for model parameters in stochastic gradient descent. *The Annals of Statistics*, **48**, 251–273.
- Doss, C. R., Flegal, J. M., Jones, G. L. and Neath, R. C. (2014) Markov chain Monte Carlo estimation of quantiles. *Electronic Journal of Statistics*, 8, 2448 – 2478.
- Durrett, R. (2019) Probability: Theory and Examples. Cambridge: Cambridge University Press.
- Eddelbuettel, D. and François, R. (2011) Rcpp: Seamless R and C++ integration. *Journal of Statistical Software*, **40**, 1–18.

- Flegal, J. M., Hughes, J., Vats, D., Dai, N., Gupta, K. and Maji, U. (2021) mcmcse: Monte Carlo Standard Errors for MCMC. Riverside, CA, and Kanpur, India. R package version 1.5-0.
- Flegal, J. M. and Jones, G. L. (2010) Batch means and spectral variance estimators in Markov chain Monte Carlo. The Annals of Statistics, 38, 1034–1070.
- Ge, R., Kakade, S. M., Kidambi, R. and Netrapalli, P. (2019) The step decay schedule: A near optimal, geometrically decaying learning rate procedure for least squares. In Advances in Neural Information Processing Systems (eds. H. Wallach, H. Larochelle, A. Beygelzimer, F. d'Alché-Buc, E. Fox and R. Garnett), vol. 32, 14977–14988. Curran Associates, Inc.
- Glynn, P. W. and Whitt, W. (1992) The asymptotic validity of sequential stopping rules for stochastic simulations. The Annals of Applied Probability, 2, 180–198.
- Goodfellow, I., Bengio, Y. and Courville, A. (2016) Deep learning. Cambridge, MA: MIT Press.
- Gupta, S., Lipton, Z. and Childers, D. (2021) Efficient online estimation of causal effects by deciding what to observe. In Advances in Neural Information Processing Systems (eds. M. Ranzato, A. Beygelzimer, Y. Dauphin, P. Liang and J. W. Vaughan), vol. 34, 20995–21007. Curran Associates, Inc.
- Gösmann, J., Kley, T. and Dette, H. (2021) A new approach for open-end sequential change point monitoring. *Journal of Time Series Analysis*, 42, 63–84.
- Hannan, E. J. (1979) The central limit theorem for time series regression. Stochastic Processes and their Applications, 9, 281–289.
- Hansen, L. P. (1982) Large sample properties of generalized method of moments estimators. *Econometrica*, **50**, 1029–1054.
- Hastings, W. K. (1970) Monte Carlo sampling methods using Markov chains and their applications. *Biometrika*, 57, 97–109.
- Huang, Y., Chen, X. and Wu, W. B. (2014) Recursive nonparametric estimation for time series. *IEEE Transactions on Information Theory*, **60**, 1301–1312.
- Jentsch, C. and Politis, D. N. (2015) Covariance matrix estimation and linear process bootstrap for multivariate time series of possibly increasing dimension. The Annals of Statistics, 43, 1117–1140.
- Jones, G. L., Haran, M., Caffo, B. S. and Neath, R. (2006) Fixed-width output analysis for Markov chain Monte Carlo. Journal of the American Statistical Association, 101, 1537–1547.
- Lang, H., Xiao, L. and Zhang, P. (2019) Using statistics to automate stochastic optimization. In Advances in Neural Information Processing Systems (eds. H. Wallach, H. Larochelle, A. Beygelzimer, F. d'Alché-Buc, E. Fox and R. Garnett), vol. 32, 9540–9550. Curran Associates, Inc.
- Liu, W. and Wu, W. B. (2010) Asymptotics of spectral density estimates. Econometric Theory, 26,

1218 - 1245.

- Liu, Y. and Flegal, J. M. (2018) Weighted batch means estimators in Markov chain Monte Carlo. *Electronic Journal of Statistics*, **12**, 3397–3442.
- Ljung, L. and Söderström, T. (1983) *Theory and practice of recursive identification*. MIT Press series in signal processing, optimization, and control. 4. Cambridge, MA: MIT Press.
- Meketon, M. S. and Schmeiser, B. (1984) Overlapping batch means: something for nothing? In Proceedings of the Winter Simulation Conference, 226–230. IEEE.
- Newey, W. K. and West, K. D. (1987) A simple, positive semi-definite, heteroskedasticity and autocorrelation consistent covariance matrix. *Econometrica*, **55**, 703–708.
- Paparoditis, E. and Politis, D. N. (2012) Nonlinear spectral density estimation: thresholding the correlogram. Journal of Time Series Analysis, 33, 386–397.
- Parzen, E. (1957) On consistent estimates of the spectrum of a stationary time series. Annals of Mathematical Statistics, 28, 329–348.
- Polyak, B. T. and Juditsky, A. B. (1992) Acceleration of stochastic approximation by averaging. SIAM Journal on Control and Optimization, 30, 838–855.
- R Core Team (2021) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
- Rho, S.-H. and Vogelsang, T. J. (2019) Heteroskedasticity autocorrelation robust inference in time series regressions with missing data. *Econometric Theory*, **35**, 601–629.
- Robbins, H. and Monro, S. (1951) A stochastic approximation method. The Annals of Mathematical Statistics, 22, 400–407.
- Schapire, R. (1990) The strength of weak learnability. Machine Learning, 5, 197–227.
- Shao, X. (2010) A self-normalized approach to confidence interval construction in time series. Journal of the Royal Statistical Society: Series B (Statistical Methodology), 72, 343–366.
- Song, W. T. and Schmeiser, B. W. (1993) Variance of the sample mean: Properties and graphs of quadratic-form estimators. Operations Research, 41, 501–517.
- Sun, Y., Phillips, P. C. and Jin, S. (2011) Power maximization and size control in heteroskedasticity and autocorrelation robust tests with exponentiated kernels. *Econometric Theory*, 27, 1320–1368.
- Terrell, G. R. and Scott, D. W. (1992) Variable kernel density estimation. The Annals of Statistics, 20, 1236–1265.
- Vats, D. and Flegal, J. M. (2021) Lugsail lag windows for estimating time-average covariance matrices.

- Vats, D., Flegal, J. M. and Jones, G. L. (2019) Multivariate output analysis for Markov chain Monte Carlo. *Biometrika*, **106**, 321–337.
- Welford, B. P. (1962) Note on a method for calculating corrected sums of squares and products. Technometrics, 4, 419–420.
- Wickham, H. (2019) Advanced R. Boca Raton, FL: Taylor & Francis, CRC Press.
- Wilson, A. C., Roelofs, R., Stern, M., Srebro, N. and Recht, B. (2017) The marginal value of adaptive gradient methods in machine learning. In *Advances in Neural Information Processing Systems* (eds. I. Guyon, U. V. Luxburg, S. Bengio, H. Wallach, R. Fergus, S. Vishwanathan and R. Garnett), vol. 30, 4148–4158. Curran Associates, Inc.
- Wu, W. B. (2005) Nonlinear system theory: Another look at dependence. Proceedings of the National Academy of Sciences, 102, 14150–14154.
- (2009) Recursive estimation of time-average variance constants. The Annals of Applied Probability, 19, 1529–1552.
- (2011) Asymptotic theory for stationary processes. Statistics and Its Interface, 4, 207–226.
- Xiao, H. and Wu, W. B. (2011) A single-pass algorithm for spectrum estimation with fast convergence. IEEE transactions on information theory, 57, 4720–4731.
- Yeh, Y. and Schmeiser, B. (2000) Simulation output analysis via dynamic batch means. In Proceedings of the Winter Simulation Conference, vol. 1, 637–645. IEEE.
- Young, P. C. (2011) Recursive Estimation and Time-Series Analysis: An Introduction for the Student and Practitioner. Berlin, Heidelberg: Springer.
- Zeileis, A., Köll, S. and Graham, N. (2020) Various versatile variances: An object-oriented implementation of clustered covariances in R. Journal of Statistical Software, 95, 1–36.
- Zhang, P., Lang, H., Liu, Q. and Xiao, L. (2020) Statistical adaptive stochastic gradient methods. Tech. Rep. MSR-TR-2020-3, Microsoft.
- Zheng, W., Jin, Y. and Zhang, G. (2016) Recursive estimation of time-average variance constants through prewhitening. Statistics & Probability Letters, 114, 30–37.
- Zhu, W., Chen, X. and Wu, W. B. (2021) Online covariance matrix estimation in stochastic gradient descent. To appear in Journal of the American Statistical Association.
- Zia, L. (1991) Using the finite difference calculus to sum powers of integers. The College Mathematics Journal, 22, 294–300.