Descent for sheaves on compact Hausdorff spaces

Peter J. Haine

October 4, 2022

Abstract

These notes explain some descent results for ∞ -categories of sheaves on compact Hausdorff spaces and derive some consequences. Specifically, given a compactly assembled ∞ -category \mathcal{E} , we show that the functor sending a locally compact Hausdorff space X to the ∞ -category Sh^{post}($X; \mathcal{E}$) of *Postnikov complete* \mathcal{E} -valued sheaves on X satisfies descent for proper surjections. This implies proper descent for *left complete* derived ∞ -categories and that the functor Sh^{post}($-; \mathcal{E}$) is a sheaf on the category of compact Hausdorff spaces equipped with the topology of finite jointly surjective families. Using this, we explain how to embed Postnikov complete sheaves on a locally compact Hausdorff space into condensed objects. This implies that the condensed and sheaf cohomologies of a locally compact Hausdorff space agree.

Contents

0	Introduction		2
	0.1	Postnikov completion	2
	0.2	Descent for Postnikov complete sheaves	3
	0.3	The comparison between sheaf and condensed cohomology	3
	0.4	Linear overview	4
1	Background		
	1.1	Hypercompleteness	5
	1.2	Postnikov completeness	7
	1.3	Condensed/pyknotic mathematics	9
2	Proper descent		10
	2.1	Proper descent for Postnikov sheaves	11
	2.2	Consequences in shape theory	12
	2.3	The necessity of Postnikov completion	13
3	The comparison functor		14
	3.1	Descent for open covers	14
	3.2	The comparison functor	17
	3.3	Naturality of the comparison functor	18
4	Full faithfulness of the comparison functor		19
	4.1	Full faithfulness for extremally disconected profinite sets	19
	4.2	Full faithfulness and descent	20
	4.3	The necessity of Postnikov completion	22
Re	References		

0 Introduction

The first goal of these notes is to explain some descent results for ∞ -categories of sheaves on locally compact Hausdorff spaces. Our motivation comes from condensed/pyknotic mathematics developed by Clausen–Scholze [20; 21; 22; 23], in our joint work with Barwick [2, Chapter 13; 3], and in Lurie's work on ultracategories [17; 18, Chapter 4]. Write **Comp** for the category of compact Hausdorff spaces. The category **Comp** has a Grothendieck topology where the covering families are finite families of jointly surjective maps. Because of the simplicity of the Grothendieck topology, the sheaf condition is very explicit: a presheaf on **Comp** is a sheaf if and only if it carries finite disjoint unions of compact Hausdorff spaces to finite products and satisfies descent for surjections.

Our first goal is to answer the following question:

0.1 Question. Is the functor Sh : $\operatorname{Comp}^{\operatorname{op}} \to \operatorname{Cat}_{\infty}$ that assigns a compact Hausdorff space *K* the ∞ -category Sh(*K*) of sheaves of spaces on *K* a sheaf with respect to this topology?

Perhaps surprisingly, the answer to Question 0.1 is *negative* (see Corollary 2.18). Moreover, if one replaces sheaves by hypersheaves, the answer to Question 0.1 is still negative. The reason for this failure of descent is that every compact Hausdorff space admits a surjection from a profinite set, and the ∞ -category of sheaves on a profinite set satisfies a strong completeness property which the ∞ -category of (hyper)sheaves on a general compact Hausdorff space does not satisfy. So it is not reasonable to ask for the ∞ -category of sheaves on a general compact Hausdorff space to be expressible as a limit of ∞ -categories satisfying this completeness property.

0.1 Postnikov completion

Since this completeness property is central to these notes, before stating the main results, let us briefly introduce it. See $\S1.2$ for more details.

0.2 Definition. Let *X* be a topological space. The *Postnikov completion* of the ∞ -category of sheaves of spaces on on *X* is the inverse limit

$$\operatorname{Sh}^{\operatorname{post}}(X) \coloneqq \lim \left(\cdots \xrightarrow{\tau_{\leq n+1}} \operatorname{Sh}(X)_{\leq n+1} \xrightarrow{\tau_{\leq n}} \operatorname{Sh}(X)_{\leq n} \xrightarrow{\tau_{\leq n-1}} \cdots \right)$$

of the ∞ -categories of sheaves of *n*-truncated spaces along the truncation functors.

Objects of $Sh^{post}(X)$ are towers

$$\cdots \to F_{n+1} \to F_n \to \cdots \to F_0$$

where F_n is an sheaf of *n*-truncated spaces on *X* such that $\tau_{\leq n} F_{n+1} \cong F_n$. There is a natural left adjoint $Sh(X) \to Sh^{\text{post}}(X)$ sending a sheaf *F* to its Postnikov tower $\{\tau_{\leq n} F\}_{\geq 0}$. We say that Sh(X) is *Postnikov complete* if this functor $Sh(X) \to Sh^{\text{post}}(X)$ is an equivalence.

0.3 Example. The ∞ -topos of sheaves on a profinite set is Postnikov complete. On the other hand, the ∞ -topos of sheaves on the Hilbert cube $\prod_{i>1}[0,1]$ is not Postnikov complete.

For a presentable ∞ -category \mathcal{E} , we write $\operatorname{Sh}^{\operatorname{post}}(X; \mathcal{E})$ for the tensor product $\operatorname{Sh}^{\operatorname{post}}(X) \otimes \mathcal{E}$. With stable coefficients this recovers the *left-complete* derived ∞ -category of sheaves: **0.4 Example.** Let *X* be a topological space and let *R* be a ring. Write D(X; R) for the derived ∞ -category of the abelian category of sheaves of *R*-modules on *X*. Then Sh^{post}(*X*; D(R)) is the *left completion* of D(X; R) with respect to the standard t-structure.¹ That is, Sh^{post}(*X*; D(R)) is the limit of the diagram of ∞ -categories

$$\cdots \xrightarrow{\tau_{\leq n+1}} \mathbf{D}(X; R)_{< n+1} \xrightarrow{\tau_{\leq n}} \mathbf{D}(X; R)_{< n} \xrightarrow{\tau_{\leq n-1}} \cdots$$

along the truncation functors with respect to the standard t-structure.

0.2 Descent for Postnikov complete sheaves

The following is the main descent result of these notes. Note that all compactly generated ∞ -categories are compactly assembled (see Recollection 1.10).

0.5 Theorem (Corollary 2.8). Let \mathcal{E} be a compactly assembled ∞ -category. Then for every proper surjection of locally compact Hausdorff spaces $p : X \rightarrow Y$, natural functor

$$\operatorname{Sh}^{\operatorname{post}}(Y;\mathcal{E}) \longrightarrow \lim \left(\operatorname{Sh}^{\operatorname{post}}(X;\mathcal{E}) \stackrel{\operatorname{pr}_1^*}{\underset{\operatorname{pr}_2^*}{\rightleftharpoons}} \operatorname{Sh}^{\operatorname{post}}(X \times_Y X;\mathcal{E}) \stackrel{\longrightarrow}{\underset{\underset{\underset{\underset{\underset{\underset{}}{\longrightarrow}}}{\longleftrightarrow}}{\longleftrightarrow}} \cdots \right)$$

is an equivalence in Cat_{∞} . Consequently, the functor $Sh^{post}(-; \mathcal{E})$: $Comp^{op} \rightarrow Cat_{\infty}$ is a hypersheaf of ∞ -categories on the site of compact Hausdorff spaces.

0.6 Example. Let *R* be a ring. Then the functor $\widehat{D}(-; R)$: **Comp**^{op} \rightarrow **Cat**_{∞} carrying a compact Hausdorff space to its left complete derived ∞ -category is a hypersheaf. Hence the functor

$$D(-;R)_{<\infty}$$
: **Comp**^{op} \rightarrow **Cat** _{∞}

that sends a compact Hausdorff space *K* to its bounded-above derived ∞ -category² is also a hypersheaf of ∞ -categories.

Passing to global sections shows that sheaf cohomology also satisfies proper descent.

0.7 Corollary. Let R be a connective E_1 -ring spectrum and M a bounded-above left R-module spectrum. The functor

$$R\Gamma_{\text{sheaf}}(-;M)$$
: **Comp**^{op} \rightarrow LMod(*R*)

is a hypersheaf.

Note that if *R* is an ordinary ring, then the ∞ -category LMod(*R*) is the derived ∞ -category D(*R*).

0.3 The comparison between sheaf and condensed cohomology

Part of our motivation for proving Theorem 0.5 is that it has a number of consequences. One application is a generalization of work of Dyckhoff and Clausen–Scholze that compares sheaf cohomology with condensed cohomology. Let *X* be a locally compact Hausdorff space. We can also regard *X* as an object of the ∞ -category Sh(**Comp**) via the restricted Yoneda embedding.

¹We use *homological* indexing for our t-structures.

²What we write as $D(K; R)_{<\infty}$ is often written as $D^+(K; R)$.

Dyckhoff [7, Theorem 3.11; 8] and Clausen–Scholze [23, Theorem 3.2] showed that if *A* is an abelian group, and *X* is *compact* then there is an isomorphism

$$\mathrm{H}^*_{\mathrm{sheaf}}(X;A) \xrightarrow{\sim} \mathrm{H}^*_{\mathrm{cond}}(X;A)$$

from the sheaf cohomology of X to the cohomology of X regarded as an object Sh(**Comp**).

We extend this result in two directions: to locally compact Hausdorff spaces and to very general coefficients. The comparison map between sheaf and condensed cohomology is induced by a natural geometric morphism

$$c_{X,*}$$
: Sh(**Comp**)_{/X} \rightarrow Sh(X)

given by sending a sheaf G: **Comp**^{op} \rightarrow **Spc** to the sheaf on *X* defined by

 $c_{X,*}(G)(U) \coloneqq \operatorname{Map}_{\operatorname{Sh}(\operatorname{\mathbf{Comp}})_{/X}}(U,G)$.

(See §3.2 for details.)

Since cohomology is computed by derived global sections, to show that the sheaf and condensed cohomologies of X agree, it suffices to show that c_X^* is fully faithful. Again, this is generally only true *after* Postnikov completion (see Warnings 4.16 and 4.19).

0.8 Proposition (Corollary 4.11). Let X be a locally compact Hausdorff space and let \mathcal{E} be a compactly assembled ∞ -category. Then the pullback functor

$$c^{*,\text{post}}$$
: $\text{Sh}^{\text{post}}(X;\mathcal{E}) \to \text{Sh}^{\text{post}}(\text{Comp}_{/X};\mathcal{E})$

is fully faithful.

Proposition 0.8 implies, for example:

0.9 Corollary (Corollary 4.12). Let X be locally compact Hausdorff space. Let R be a connective E_1 -ring spectrum and let M be a bounded-above left R-module spectrum. Then the natural map

$$R\Gamma_{\text{sheaf}}(X;M) \to R\Gamma_{\text{cond}}(X;M)$$

is an equivalence in the ∞ -category LMod(R) of left R-module spectra.

As a consequence, the condensed, singular, and sheaf cohomologies of a topological space admitting a locally finite CW structure all agree (see Remark 4.13 and (4.14)).

0.4 Linear overview

We imagine that the reader might be interested in condensed/pyknotic mathematics but not necessarily familiar with all of the intricacies about the theory of ∞ -topoi. With this in mind, in §1, we review the basics of hypercomplete and Postnikov complete ∞ -topoi; the familiar reader can safely skip this section. Section 2 proves Theorem 0.5 and derives some consequences in shape theory. In §3, we construct the comparison geometric morphism

$$c_*: \operatorname{Sh}(\operatorname{\mathbf{Comp}}_{/L}) \to \operatorname{Sh}(L)$$

and record its basic properties. Section 4 is dedicated to proving Proposition 0.8.

Acknowledgments. We thank Ko Aoki, Clark Barwick, Marc Hoyois, Jacob Lurie, Mark Macerato, Zhouhang Mao, Denis Nardin, Piotr Pstrągowski, Marco Volpe, Sebastian Wolf, and Tong Zhou for helpful comments and conversations around the contents of these notes. Special thanks are due to Marc Hoyois and Jacob Lurie for explaining Example 2.15 to us. These notes are clearly highly influenced by Dustin Clausen and Peter Scholze's ideas; we would like to thank them too.

We gratefully acknowledge support from the UC President's Postdoctoral Fellowship and NSF Mathematical Sciences Postdoctoral Research Fellowship under Grant #DMS-2102957.

1 Background

Recall the following fundamental results about the ∞ -category of spaces.

- (1) Whitehead's Theorem: A map f: X → Y of spaces is an equivalence if and only if f induces a bijection on connected components and isomorphisms on homotopy groups at each basepoint. Said differently, f is an equivalence if and only if for each n ≥ 0, the induced map on n-truncations τ_{<n}(f): τ_{<n}(X) → τ_{<n}(Y) is an equivalence.
- (2) Convergence of Postnikov towers: Every space X is the limit of its Postnikov tower. That is, the the natural map $X \to \lim_{n>0} \tau_{< n}(X)$ is an equivalence.

The statement of Whitehead's Theorem and the convergence of Postnikov towers make can be formulated in an arbitrary ∞ -topos. However, even for the ∞ -topos of sheaves on a compact Hausdorff space, neither result need hold (see Example 2.15). The purpose of this section is to review two completion procedures (*hypercompletion* and *Postnikov completion*) that force Whitehead's Theorem to hold and Postnikov towers to converge, respectively. In the higher-categorical world, these give rise to three natural 'sheaf theories' (sheaves, hypersheaves, and Postnikov complete sheaves) extending the classical theory of sheaves on a topological space. They all have the same truncated objects, so the subtle differences between these theories only appears when considering 'unbounded' objects.

Subsection 1.1 reviews the basics of hypercompleteness; in the process, we set some notation. In § 1.2 we review Postnikov completeness. In § 1.3, we recall the basic setup of condensed/pyknotic mathematics.

1.1 Hypercompleteness

In this subsection, we set up some notation and review the basics of hypercompletions of ∞ -topoi. We refer the reader unfamiliar with hypercomplete objects and hypercompletion to [HTT, §§6.5.2–6.5.4], [2, §3.11], or [11, §1.2] for further reading on the subject.

1.1 Notation. Write **Spc** for the ∞ -category of spaces and **Cat**_{∞} for the ∞ -category of ∞ -categories.

1.2 Notation. Let \mathcal{C} be an ∞ -site and \mathcal{E} a presentable ∞ -category. We write

$$PSh(\mathcal{C};\mathcal{E}) \coloneqq Fun(\mathcal{C}^{op},\mathcal{E})$$

for the ∞ -category of \mathcal{E} -valued presheaves on \mathcal{C} . We write $Sh(\mathcal{C}; \mathcal{E}) \subset PSh(\mathcal{C}; \mathcal{E})$ for the full subcategory spanned by \mathcal{E} -valued sheaves. When $\mathcal{E} = Spc$, we simply write

$$PSh(\mathcal{C}) \coloneqq PSh(\mathcal{C}; \mathbf{Spc})$$
 and $Sh(\mathcal{C}) \coloneqq Sh(\mathcal{C}; \mathbf{Spc})$.

1.3. The ∞ -categories PSh($\mathcal{C}; \mathcal{E}$) and Sh($\mathcal{C}; \mathcal{E}$) are naturally identified with the tensor products of presentable ∞ -categories PSh(\mathcal{C}) $\otimes \mathcal{E}$ and Sh(\mathcal{C}) $\otimes \mathcal{E}$ [SAG, Remark 1.3.1.6 & Proposition 1.3.1.7].

1.4 Notation. Let *X* be a topological space. We write Open(X) the poset of open subsets of *X*, ordered by inclusion. We regard Open(X) as a site with the covering families given by open covers. We write

 $PSh(X; \mathcal{E}) := PSh(Open(X); \mathcal{E})$ and $Sh(X; \mathcal{E}) := Sh(Open(X); \mathcal{E})$.

1.5 Notation. Let \mathcal{E} be a presentable ∞ -category and $f_* : \mathcal{X} \to \mathcal{Y}$ a geometric morphism of ∞ -topoi. For simplicity, we also denote the tensor product $f_* \otimes \mathcal{E} : \mathcal{X} \otimes \mathcal{E} \to \mathcal{Y} \otimes \mathcal{E}$ by f_* .

1.6 Recollection (hypercompleteness). Let \mathcal{X} be an ∞ -topos. The ∞ -category of *hypercomplete* objects of \mathcal{X} is the full subcategory $\mathcal{X}^{hyp} \subset \mathcal{X}$ spanned by those objects that are local with respect to the ∞ -connected morphisms. The inclusion $\mathcal{X}^{hyp} \subset \mathcal{X}$ admits a left exact left adjoint; hence \mathcal{X}^{hyp} is also an ∞ -topos.

As the name suggests, \mathcal{X}^{hyp} can also be identified as the full subcategory of \mathcal{X} spanned by those objects that satisfy descent for hypercovers; see [HTT, Corollary 6.5.3.13; 6, §1; 24, Corollary 3.4.7].

1.7 Notation. Let \mathcal{X} be an ∞ -topos and $n \ge 0$ an integer. Write $\mathcal{X}_{\le n} \subset \mathcal{X}$ for the full subcategory spanned by the *n*-truncated objects, and write $\tau_{\le n} : \mathcal{X} \to \mathcal{X}_{\le n}$ for the left adjoint to the inclusion.

1.8 Remark. For each integer $n \ge 0$, the inclusion $\mathcal{X}^{hyp} \hookrightarrow \mathcal{X}$ restricts to an equivalence $(\mathcal{X}^{hyp})_{\le n} \cong \mathcal{X}_{\le n}$ on subcategories of *n*-truncated objects [HTT, Lemma 6.5.2.9]. As a consequence, given an integer $n \ge 0$ and presentable *n*-category \mathcal{E} , we have $\mathcal{X}^{hyp} \otimes \mathcal{E} \simeq \mathcal{X} \otimes \mathcal{E}$ [HA, Example 4.8.1.22]. In particular, every sheaf of sets is hypercomplete.

1.9 Notation. Let \mathcal{C} be a site and \mathcal{E} a presentable ∞ -category. We write $\mathrm{Sh}^{\mathrm{hyp}}(\mathcal{C}) \coloneqq \mathrm{Sh}(\mathcal{C})^{\mathrm{hyp}}$ and

$$\operatorname{Sh}^{\operatorname{hyp}}(\mathcal{C};\mathcal{E}) \coloneqq \operatorname{Sh}^{\operatorname{hyp}}(\mathcal{C}) \otimes \mathcal{E}$$

We refer to objects of $Sh^{hyp}(\mathcal{C}; \mathcal{E})$ as \mathcal{E} -valued *hypersheaves*. We use analogous notation for hypersheaves on a topological space.

For reasonable coefficients, hypersheaves on a topological space can be identified very explicitly.

1.10 Recollection (compactly assembled ∞ -categories). A presentable ∞ -category \mathcal{E} is *compactly assembled* if \mathcal{E} is a retract of a compactly generated ∞ -category regarded as an object of the ∞ -category \mathbf{Pr}^{L} of presentable ∞ -categories and left adjoints [SAG, Definition 21.1.2.1 & Theorem 21.1.2.18]. A *stable* presentable ∞ -category \mathcal{E} is compactly assembled if and only if \mathcal{E} is dualizable in the symmetric monoidal ∞ -category \mathbf{Pr}_{st}^{L} of stable presentable ∞ -categories and left adjoints [SAG, Proposition D.7.3.1].

1.11 Remark. Let *X* be a topological space and \mathcal{E} a compactly assembled ∞ -category. Then the subcategory

$$\operatorname{Sh}^{\operatorname{nyp}}(X;\mathcal{E}) \subset \operatorname{Sh}(X;\mathcal{E})$$

is the localization obtained by inverting all morphisms that induce equivalences on stalks [HA, Lemma A.3.9; 10, Lemma 2.11].

1.12 Notation. Write $LTop_{\infty}$ for the ∞ -category with objects ∞ -topoi and morphisms left exact left adjoints.

1.13. We repeatedly use the fact that the forgetful functor $LTop_{\infty} \rightarrow Cat_{\infty}$ preserves limits [HTT, Proposition 6.3.2.3].

1.14. Hypercompletion defines a functor $(-)^{hyp}$: $LTop_{\infty} \rightarrow LTop_{\infty}$ left adjoint to the inclusion of hypercomplete ∞ -topoi into $LTop_{\infty}$.

1.2 Postnikov completeness

In this subsection, we review the basics of *Postnikov completions* of ∞ -topoi. We refer the unfamiliar reader to [HTT, §5.5.6; SAG, §§A.7.2 & A.7.2; 2, §3.2] for more background. The ∞ -category of spaces actually satisfies a stronger property than the requirement that every object be the limit of its Postnikov tower: the entire ∞ -category can be recovered as the limit of the subcategories of *n*-truncated spaces along the truncation functors. This is the property that we want to generalize to arbitrary ∞ -topoi.

1.15 Definition. Let \mathcal{X} be an ∞ -topos. The *Postnikov completion* of \mathcal{X} is the limit

$$\mathcal{X}^{\text{post}} \coloneqq \lim \left(\cdots \longrightarrow \mathcal{X}_{\leq n+1} \xrightarrow{\tau_{\leq n}} \mathcal{X}_{\leq n} \longrightarrow \cdots \xrightarrow{\tau_{\leq 0}} \mathcal{X}_{\leq 0} \right)$$

formed in Cat_{∞} . Thus objects of \mathcal{X}^{post} are given by towers

$$\cdots \to V_{n+1} \to V_n \to \cdots \to V_0$$

in \mathcal{X} , where V_n is *n*-truncated and the map $V_{n+1} \to V_n$ exhibits V_n as the *n*-truncation of V_{n+1} .

1.16. The Posnikov completion $\mathcal{X}^{\text{post}}$ is also an ∞ -topos. Moreover, there is a natural left exact left adjoint t^* : $\mathcal{X} \to \mathcal{X}^{\text{post}}$ defined by sending an object to its Postnikov tower:

$$t^*(U) \coloneqq \{\tau_{\leq n} U\}_{n \geq 0}$$

See [SAG, Theorem A.7.2.4]. The right adjoint $t_* : \mathcal{X}^{\text{post}} \to \mathcal{X}$ sends a tower $\{V_n\}_{n \ge 0}$ to the limit $\lim_{n \ge 0} V_n$ formed in \mathcal{X} [SAG, Remark A.7.3.6].

1.17 Observation. The functor t^* : $\mathcal{X} \to \mathcal{X}^{\text{post}}$ is fully faithful if and only if for each object $U \in \mathcal{X}$, the natural map $U \to \lim_{n \ge 0} \tau_{\le n} U$ is an equivalence. That is, t^* if fully faithful if and only if every object of \mathcal{X} is the limit of its Postnikov tower.

1.18 Remark. For each integer $n \ge 0$, the functor $t^* : \mathcal{X} \to \mathcal{X}^{\text{post}}$ restricts to an equivalence $\mathcal{X}_{\le n} \cong (\mathcal{X}^{\text{post}})_{\le n}$ [SAG, Corollary A.7.3.8]. Moreover, given an integer $n \ge 0$ and presentable *n*-category \mathcal{E} , we have

$$\mathcal{X}^{\text{post}} \otimes \mathcal{E} \simeq \mathcal{X}^{\text{hyp}} \otimes \mathcal{E} \simeq \mathcal{X} \otimes \mathcal{E} .$$

1.19 Warning. The right adjoint t_* : $\mathcal{X}^{\text{post}} \to \mathcal{X}$ is fully faithful if and only if $\mathcal{X}^{\text{hyp}} = \mathcal{X}^{\text{post}}$.

1.20 Definition. We say that an ∞ -topos \mathcal{X} is *Postnikov complete* if the functor $t^* : \mathcal{X} \to \mathcal{X}^{\text{post}}$ is an equivalence of ∞ -categories.

1.21 Remark. The natural geometric morphism $\mathcal{X}^{hyp} \hookrightarrow \mathcal{X}$ induces an equivalence

 $(\mathcal{X}^{\text{hyp}})^{\text{post}} \xrightarrow{\sim} \mathcal{X}^{\text{post}}$.

Moreover, if \mathcal{X} is Postnikov complete, then \mathcal{X} is hypercomplete. However, the converse is false (see Example 2.15).

1.22 Observation. Let \mathcal{X} be an ∞ -topos. Then \mathcal{X} is Postnikov complete if and only if the pushforward $t_* : \mathcal{X}^{\text{post}} \to \mathcal{X}$ is conservative and the pullback t^* is fully faithful. From the explicit descriptions of t^* and t_* , we see that \mathcal{X} is Postnikov complete if and only if the following conditions are satisfied:

(1.22.1) For each $U \in \mathcal{X}$, the natural map $U \to \lim_{n \ge 0} \tau_{\le n} U$ is an equivalence.

(1.22.2) For each integer $n \ge 0$, the functor $t_* : \mathcal{X}^{\text{post}} \to \mathcal{X}$ commutes with *n*-truncation.

See also [HTT, Proposition 5.5.6.26].

1.23 Warning. As far as we are aware, it is not known if there exists an ∞ -topos \mathcal{X} such that every object of \mathcal{X} is the limit of its Postnikov tower, but \mathcal{X} is not Postnikov complete.

1.24 Notation. Write $LTop_{\infty}^{post} \subset LTop_{\infty}$ for the full subcategory spanned by the Postnikov complete ∞ -topoi.

1.25. Postnikov completion defines a functor

$$(-)^{\text{post}}$$
: LTop _{∞} \rightarrow LTop _{∞} ^{post}

which is left adjoint to the inclusion [SAG, Corollary A.7.2.6]. The functor $(-)^{\text{post}}$ is also a right adjoint [SAG, Corollary A.7.2.7]. Hence the full subcategory $\text{LTop}_{\infty}^{\text{post}} \subset \text{LTop}_{\infty}$ is closed under limits. As a consequence of (1.13), the forgetful functor $\text{LTop}_{\infty}^{\text{post}} \rightarrow \text{Cat}_{\infty}$ preserves limits.

To prove Theorem 0.5, use the following reformulation of what it means for a diagram of Postnikov complete ∞ -topoi to be a limit diagram:

1.26 Lemma. Let \mathcal{I} be an ∞ -category and \mathcal{X}_{\bullet} : $\mathcal{I}^{\triangleleft} \to \mathbf{LTop}_{\infty}$ a diagram of ∞ -topoi. The following are equivalent:

(1.26.1) For each integer $n \ge 0$, the diagram $(\mathcal{X}_{\bullet})_{\le n} : \mathcal{I}^{\triangleleft} \to \mathbf{Cat}_{\infty}$ is a limit diagram.

(1.26.2) The diagram of Postnikov complete ∞ -topoi $\mathcal{X}^{\text{post}}_{\bullet}$: $\mathcal{I}^{\triangleleft} \to \mathbf{Cat}_{\infty}$ is a limit diagram.

Proof. By the definition of Postnikov completion and the fact that limits commute, we see that $(1.26.1) \Rightarrow (1.26.2)$. To prove that $(1.26.2) \Rightarrow (1.26.1)$, tensor the limit diagram $\mathcal{X}_{\bullet}^{\text{post}}$: $\mathcal{I}^{\triangleleft} \rightarrow \mathbf{Cat}_{\infty}$ with $\mathbf{Spc}_{\leq n}$ and apply [10, Lemma 2.15].

We finish this subsection with some examples of Postnikov complete ∞ -topoi from topology.

1.27 Notation. Let \mathcal{C} be an ∞ -site and \mathcal{E} a presentable ∞ -category. Write $Sh^{post}(\mathcal{C}) \coloneqq Sh(\mathcal{C})^{post}$ and

$$\operatorname{Sh}^{\operatorname{post}}(\mathcal{C};\mathcal{E}) \coloneqq \operatorname{Sh}^{\operatorname{post}}(\mathcal{C}) \otimes \mathcal{E}$$

We refer to objects of $\text{Sh}^{\text{post}}(\mathcal{C}; \mathcal{E})$ as *Postnikov complete sheaves* on \mathcal{C}^3 . We use analogous notation for Postnikov complete sheaves on a topological space.

³Since Sh^{post}($\mathcal{C}; \mathcal{E}$) is not generally a subcategory of Sh($\mathcal{C}; \mathcal{E}$), this is slightly abusive.

1.28 Example. Let *S* be a profinite set. Since *S* is 0-dimensional, by [HTT, Corollary 7.2.1.10, Theorem 7.2.3.6, & Remark 7.2.4.18] the ∞ -topos Sh(*S*) is Postnikov complete.

1.29 Example [13]. Let *X* be a topological space. If *X* admits a CW structure, then the ∞ -topos Sh(*X*) is Postnikov complete.

For hypersheaves and Postnikov complete sheaves, pullbacks along surjections are conservative:

1.30 Observation. Let $p: X \twoheadrightarrow Y$ be a surjection of topological spaces. Since $\text{Sh}^{\text{hyp}}(Y)$ has enough points and the points of $\text{Sh}^{\text{hyp}}(Y)$ are in natural bijection with the underlying set of *Y*, the pullback functor

$$p^{*,\text{hyp}}$$
: $\text{Sh}^{\text{hyp}}(Y) \to \text{Sh}^{\text{hyp}}(X)$

is conservative. In particular, the functor p^* : $Sh(Y) \to Sh(X)$ is conservative when restricted to the subcategory of truncated objects. Hence $p^{*,post}$: $Sh^{post}(Y) \to Sh^{post}(X)$ is also conservative.

1.3 Condensed/pyknotic mathematics

In this subsection, we briefly recall the formalism of condensed/pyknotic mathematics. We refer the reader to [2, §13.3; 3; 5; 20; 21; 22, Lecture I; 23] for more details and motivation.

1.31 Notation. Write **Top** for the category of topological spaces and **Comp** \subset **Top** for the full subcategory spanned by the compact Hausdorff spaces. We regard **Comp** as a site where the covering families are finite families of jointly surjective maps.

1.32 Remark (set theory). Since **Comp** is a large category, one has to be careful about talking about sheaves on **Comp**. To do this, we adopt the set-theoretic conventions of [2, §13.3; 3]; this uses universes to deal with the set theory. Clausen and Scholze [23] use alternative set-theoretic foundations that avoid using universes. This minor difference is only in the set theory and does not affect any arguments in an essential way, so we will not mention it again.

In this setting, the sheaf condition is particularly easy to formulate:

1.33 Observation. Let \mathcal{D} be an ∞ -category. A presheaf F: **Comp**^{op} $\rightarrow \mathcal{D}$ is a sheaf if and only if the following conditions are satisfied:

- (1.33.1) The functor F preserves finite products. That is, F carries finite coproducts of compact Hausdorff spaces to finite products in D.
- (1.33.2) For every surjection of compact Hausdorff spaces $p: X \twoheadrightarrow Y$, the augmented cosimplicial diagram

$$F(Y) \xrightarrow{p^*} F(X) \overleftrightarrow{\longrightarrow} F(X \times_Y X) \overleftrightarrow{\longleftrightarrow} \cdots$$

obtained by applying *F* to the Čech nerve of *p* exhibits F(Y) as the limit of its restriction to $\Delta \subset \Delta_+$.

See [SAG, Proposition A.3.3.1].

1.34. Every representable presheaf on **Comp** is a sheaf. Moreover, the topology on **Comp** is designed exactly so that the Yoneda embedding **Comp** \hookrightarrow Sh(**Comp**) preserves finite coproducts and carries surjections to effective epimorphisms.

1.35 Notation. Write Extr \subset Comp for the full subcategory spanned by the *extremally disconnected* profinite sets. The extremally disconnected profine sets are exactly the projective objects of the category Comp [9; 16, Chapter III, §3.7].

1.36. Let *K* be a compact Hausdorff space. Write K^{disc} for the underlying set of *K* equipped with the discrete topology. There is a natural surjection $\beta(K^{\text{disc}}) \twoheadrightarrow K$ from the Stone–Čech compactification of K^{disc} to *K*. Since the profinite set $\beta(K^{\text{disc}})$ is extremally disconnected, the subcategory **Extr** \subset **Comp** is a basis for the Grothendieck topology on **Comp**. Therefore, restriction of presheaves defines an equivalence of ∞ -categories

$$\mathrm{Sh}^{\mathrm{hyp}}(\mathbf{Comp}) \cong \mathrm{Sh}^{\mathrm{hyp}}(\mathbf{Extr})$$

with inverse given by right Kan extension [1, Corollary A.8; 2, Corollary 3.12.14].

1.37. Since every surjection of extremally disconnected profinite sets admits a section, a presheaf F on **Extr** is a sheaf if and only if F preserves finite products. Moreover, by [3, Lemma 2.4.10], the ∞ -topos Sh(**Extr**) is Postnikov complete. Hence Sh^{hyp}(**Comp**) is also Postnikov complete.

Most reasonable topological spaces embed into sheaves on **Comp**:

1.38 Notation. Write \downarrow : **Top** \rightarrow Sh(**Comp**) for the restricted Yoneda functor defined by

$$\sharp(X)(K) := \operatorname{Map}_{\operatorname{Top}}(K, X)$$

When it does not cause confusion, we also simply denote $\downarrow(X) \in Sh(Comp)$ by *X*.

1.39. The functor \downarrow is not fully faithful. However, \downarrow is fully faithful when restricted to the the full subcategory of *compactly generated* topological spaces.

1.40. Note that the functor \sharp : **Top** \rightarrow Sh(**Comp**) preserves limits. The functor \sharp does not preserve arbitrary colimits. However, in §3.1 we show that \sharp behaves well with open covers, coproducts, and proper surjections.

2 Proper descent

Let \mathcal{E} be a compactly assembled ∞ -category (see Recollection 1.10). In this section, we show that the functor sending a locally compact Hausdorff space *X* to the ∞ -category Sh^{post}(*X*; \mathcal{E}) of Postnikov complete sheaves on *X* satisfies descent for proper surjections in the following sense.

2.1 Notation. Write LCH \subset Top for the full subcategory spanned by the locally compact Hausdorff spaces.

2.2 Definition. Let \mathcal{D} be an ∞ -category. We say that a functor $F : \mathbf{LCH}^{\mathrm{op}} \to \mathcal{D}$ satisfies proper *descent* if for every proper surjection of locally compact Hausdorff spaces $p : X \twoheadrightarrow Y$, the augmented cosimplicial diagram

$$F(Y) \xrightarrow{p^*} F(X) \overleftrightarrow{\longrightarrow} F(X \times_Y X) \overleftrightarrow{\bigoplus} \cdots$$

obtained by applying *F* to the Čech nerve of *p* exhibits F(Y) as the limit of its restriction to $\Delta \subset \Delta_+$.

Subsection 2.1 proves Theorem 0.5 (see Corollary 2.8). In §2.2 we record some shape-theoretic consequences. In §2.3, we explain why Theorem 0.5 does not hold before Postnikov completion.

2.1 Proper descent for Postnikov sheaves

To prove that the functor $X \mapsto Sh^{post}(X)$ satisfies proper descent we apply the following criterion:

2.3 Proposition [HA, Corollary 4.7.5.3]. Let $S^{\bullet} : \Delta_{+} \to \mathbf{Cat}_{\infty}$ be an augmented cosimplicial ∞ -category. Let $G : S^{-1} \to S^{0}$ denote the agumentation. Assume that:

- (2.3.1) The ∞ -category S^{-1} admits totalizations of *G*-split cosimplicial objects, and those totalizations are preserved by *G*.
- (2.3.2) For every morphism α : $[m] \rightarrow [n]$ in Δ_+ , the square

$$\begin{array}{ccc} S^m & \stackrel{d^0}{\longrightarrow} & S^{m+1} \\ \alpha^* \downarrow & & \downarrow ([0] \star \alpha)^* \\ S^n & \stackrel{d^0}{\longrightarrow} & S^{n+1} \end{array}$$

is horizontally right adjointable. (Note that, in particular, this requires that the coface functors d⁰ be left adjoints.)

Then the natural functor θ : $S^{-1} \to \lim_{[n] \in \Delta} S^n$ admits a fully faithful right adjoint. Moreover, if *G* is conservative, then θ is an equivalence.

We are interested in applying Proposition 2.3 in the case where S^* is obtained by applying sheaves to the Čech nerve of a proper surjection.

2.4 Example. Let \mathcal{E} be a presentable ∞ -category which is compactly generated or stable, and let $p: X \twoheadrightarrow Y$ be a proper surjection of locally compact Hausdorff spaces. The Proper Basechange Theorem [HTT, Corollary 7.3.1.18; 10, Subexample 3.15] implies that the augmented cosimplicial diagram

(2.5)
$$\operatorname{Sh}(Y;\mathcal{E}) \xrightarrow{p^*} \operatorname{Sh}(X;\mathcal{E}) \rightleftharpoons \operatorname{Sh}(X \times_Y X;\mathcal{E}) \rightleftharpoons \cdots$$

satisfies hypothesis (2.3.2).

2.6 Observation. In the setting of Example 2.4, the functor p^* : $Sh(Y; \mathcal{E}) \rightarrow Sh(X; \mathcal{E})$ is left exact. If there is an integer $n \ge 0$ such that \mathcal{E} is an *n*-category, then the totalizations in (2.3.1) can be computed by finite limits [12, Proposition A.1]. Hence, in this case, the diagram (2.5) also satisfies (2.3.1).

2.7 Lemma. Let $n \ge 0$ be an integer. Then:

(2.7.1) The functor $Sh(-)_{\leq n}$: LCH^{op} \rightarrow Cat_{∞} satisfies proper descent.

(2.7.2) The functor $Sh(-)_{\leq n}$: **Comp**^{op} \rightarrow **Cat**_{∞} is a hypersheaf.

Proof. For (2.7.1), note that since $\mathbf{Spc}_{\leq n}$ is compactly generated, by Example 2.4 and Observation 2.6 it suffices to check that for each proper surjection of locally compact Hausdorff spaces $p: X \twoheadrightarrow Y$, the pullback functor $p^*: \mathrm{Sh}(Y)_{\leq n} \to \mathrm{Sh}(X)_{\leq n}$ is conservative. This follows from the assumption that p is a surjection (Observation 1.30).

Since $\text{Sh}(-)_{\leq n}$ factors through the full subcategory $\operatorname{Cat}_{n+1} \subset \operatorname{Cat}_{\infty}$ spanned by the (n + 1)-categories, and Cat_{n+1} is an (n + 2)-category, by Remark 1.8, item (2.7.2) is equivalent to the claim that $\operatorname{Sh}(-)_{\leq n}$ is a sheaf. Thus (2.7.2) follows from (2.7.1) and the fact that the functor $\operatorname{Sh}(-)_{\leq n}$ carries coproducts of topological spaces to products of ∞ -categories.

2.8 Corollary. Let \mathcal{E} be a compactly assembled ∞ -category. Then:

(2.8.1) The functor $Sh^{post}(-; \mathcal{E})$: LCH^{op} \rightarrow Cat_{∞} satisfies proper descent.

(2.8.2) The functor $\operatorname{Sh}^{\operatorname{post}}(-; \mathcal{E})$: Comp^{op} \rightarrow Cat_{∞} is a hypersheaf.

Proof. Since \mathcal{E} is compactly assembled, the functor $\mathcal{E} \otimes (-)$: $\mathbf{Pr}^{L} \to \mathbf{Pr}^{L}$ commutes with limits of diagrams where the transition functors are left exact [10, Lemma 2.15]. Hence it suffices to prove the claims for $\mathcal{E} = \mathbf{Spc}$. In this case, the claims follow from Lemmas 1.26 and 2.7.

2.2 Consequences in shape theory

We now record some consequences of Corollary 2.8 regarding the shape of compact Hausdorff spaces. The reader unfamiliar with the shape is encouraged to consult [HTT, §7.1.6; HA, §A.1; SAG, §E.2; 2, Chapter 4]. We follow the notations of [2, Chapter 4].

Since the shape Π_{∞} : **RTop**_{∞} \rightarrow Pro(**Spc**) is a left adjoint, Corollary 2.8 implies:

2.9 Corollary. Let $p: X \twoheadrightarrow Y$ be a proper surjection of locally compact Hausdorff spaces. Then the natural map of prospaces

$$\operatorname{colim}_{[n]\in\Delta^{\operatorname{op}}}\Pi_{\infty}\operatorname{Sh}^{\operatorname{post}}(X^{\times_{Y}n})\to\Pi_{\infty}\operatorname{Sh}^{\operatorname{post}}(Y)$$

is an equivalence.

Recall that the natural geometric morphism $\text{Sh}^{\text{post}}(X) \to \text{Sh}(X)$ induces an equivalences on protruncated shapes. Combining Corollary 2.9 with work of Hoyois [14], we deduce that the protruncated shape preserves many (co)limits of compact Hausdorff spaces.

2.10 Corollary. The protruncated shape $\Pi_{<\infty}$ Sh : Comp \rightarrow Pro(Spc_{< ∞}) preserves:

(2.10.1) Finite coproducts.

(2.10.2) Cofiltered limits and arbitrary products.

(2.10.3) Geometric realizations of Čech nerves of surjections.

Proof. For (2.10.1), note that the functor Sh : **Comp** \rightarrow **RTop**_{∞} preserves finite coproducts and the protruncated shape

$$\Pi_{<\infty}: \mathbf{RTop}_{\infty} \to \operatorname{Pro}(\mathbf{Spc}_{<\infty})$$

is a left adjoint. Item (2.10.2) is [14, Example 2.12], and Corollary 2.9 implies (2.10.3).

2.11 Warning. The protruncated shape does *not* preserve pullbacks of compact Hausdorff spaces. To see this, let S¹ denote the topological circle, and choose a point $t \in S^1$. The shapes of Sh({*t*}) and Sh(S¹) are the point and the homotopy type B**Z**, respectively (see [HTT, Corollary 7.2.1.12, Theorem 7.2.3.6 & Remark 7.2.4.18; 11, Corollary 3.5]). The pullback of topological spaces {*t*} ×_{S¹} {*t*} is a point, hence

$$\Pi_{\infty} \operatorname{Sh}(\{t\} \times_{\mathrm{S}^1} \{t\}) \simeq * .$$

On the other hand,

$$\Pi_{\infty} \operatorname{Sh}(\{t\}) \underset{\Pi_{\infty} \operatorname{Sh}(\mathrm{S}^{1})}{\times} \underset{\mathrm{B}\mathbf{Z}}{\Pi_{\infty} \operatorname{Sh}(\{t\})} \simeq * \underset{\mathrm{B}\mathbf{Z}}{\times} * \simeq \mathbf{Z} \,.$$

2.3 The necessity of Postnikov completion

We conclude this section by explaining why Postnikov completion is necessary in the statement of Corollary 2.8. To do this, we give an example of a compact Hausdorff space K for which $Sh^{hyp}(K)$ is not Postnikov complete. We begin with some preliminary observations.

2.12. Let \mathcal{X} be an ∞ -topos. For each $m \ge 1$, write $K_{\mathcal{X}}(\mathbf{Z}, m)$ for the constant object of \mathcal{X} at the Eilenberg–MacLane space $K(\mathbf{Z}, m)$. Note that since $K(\mathbf{Z}, m)$ is *m*-truncated, $K_{\mathcal{X}}(\mathbf{Z}, m)$ is a hypercomplete object of \mathcal{X} . Moreover, since the hypercomplete objects of \mathcal{X} are closed under limits, the object $\prod_{m>1} K_{\mathcal{X}}(\mathbf{Z}, m)$ is also a hypercomplete object of \mathcal{X} .

2.13 Observation. If the ∞ -topos \mathcal{X} is Postnikov complete, then

$$\tau_{\leq n}\left(\prod_{m\geq 1} \mathcal{K}_{\mathcal{X}}(\mathbf{Z},m)\right) \simeq \prod_{m=1}^{n} \mathcal{K}_{\mathcal{X}}(\mathbf{Z},m).$$

In particular, $\tau_{\leq 0} \left(\prod_{m \geq 1} K_{\mathcal{X}}(\mathbf{Z}, m) \right)$ is a terminal object of \mathcal{X} , so that $\prod_{m \geq 1} K_{\mathcal{X}}(\mathbf{Z}, m)$ is connected.

2.14 Observation. Let *X* be a topological space and write $F := \prod_{m \ge 1} K_{Sh(X)}(\mathbf{Z}, m)$. Then

$$\pi_0 \Gamma(X; F) \simeq \prod_{m \ge 1} \mathcal{H}^m_{\text{sheaf}}(X; \mathbf{Z})$$

Note that if the sheaf *F* is connected, then any element $(e_m)_{m\geq 1} \in \prod_{m\geq 1} \operatorname{H}^m_{\operatorname{sheaf}}(X; \mathbb{Z})$ vanishes *locally* in the following sense: any point $x \in X$ has an open neighborhood $U \subset X$ such that for each $m \geq 1$, the class $e_m|_U \in \operatorname{H}^m_{\operatorname{sheaf}}(U; \mathbb{Z})$ is zero.

2.15 Example. Let $X \coloneqq \prod_{m \ge 1} S^m$ be the product of positive-dimensional spheres. We claim that the ∞ -topos Sh^{hyp}(X) is not Postnikov complete. In light of Observation 2.13, to see this it suffices to show that the hypersheaf $\prod_{m \ge 1} K_{Sh(X)}(\mathbf{Z}, m)$ is not connected. Observation 2.14 shows that it suffices to construct cohomology classes $e_i \in H^i_{sheaf}(X; \mathbf{Z})$ that do not simultaneously vanish on any nonempty open of X. For this, let e_i be the pullback of the generator of

$$\mathrm{H}^{i}_{\mathrm{sheaf}}(\mathrm{S}^{i};\mathbf{Z})\cong\mathbf{Z}$$

under the projection $pr_i : \prod_{m \ge 1} S^m \to S^i$.

2.16 Example. Not only is the ∞ -topos of hypersheaves on $X = \prod_{m \ge 1} S^m$ not Postnikov complete, but there are also objects which are not limits of their Postnikov towers. To see this, consider the filtered colimit

$$G \coloneqq \operatorname{colim}\left(\mathrm{K}_{\operatorname{Sh}(X)}(\mathbf{Z},1) \longrightarrow \cdots \longrightarrow \prod_{m=1}^{n} \mathrm{K}_{\operatorname{Sh}(X)}(\mathbf{Z},m) \longrightarrow \prod_{m=1}^{n+1} \mathrm{K}_{\operatorname{Sh}(X)}(\mathbf{Z},m) \longrightarrow \cdots\right)$$

formed in $Sh^{hyp}(X)$. Here, the transition map

$$\prod_{m=1}^{n} \mathrm{K}_{\mathrm{Sh}(X)}(\mathbf{Z},m) \to \mathrm{K}_{\mathrm{Sh}(X)}(\mathbf{Z},n+1) \times \prod_{m=1}^{n} \mathrm{K}_{\mathrm{Sh}(X)}(\mathbf{Z},m)$$

is the constant map at the basepoint of $K_{\operatorname{Sh}(X)}(\mathbf{Z}, n + 1)$ on the first factor and the identity on the second factor. Since connected objects are closed under filtered colimits, *G* is connected. The functor $\tau_{\leq n}$: Sh^{hyp}(X) \rightarrow Sh^{hyp}(X) preserves filtered colimits and finite products, hence

$$\tau_{\leq n} G \simeq \prod_{m=1}^{n} \mathrm{K}_{\mathrm{Sh}(X)}(\mathbf{Z}, m) \, .$$

As a consequence, we have

$$\lim_{n\geq 0}\tau_{\leq n}\,G\simeq\prod_{m\geq 1}\mathrm{K}_{\mathrm{Sh}(X)}(\mathbf{Z},m)\,.$$

Example 2.15 shows that the limit $\lim_{n\geq 0} \tau_{\leq n} G$ is not connected. In particular, the natural map $G \to \lim_{n\geq 0} \tau_{\leq n} G$ is not an equivalence.

Example 2.15 lets us see that the presheaves Sh, Sh^{hyp} : $Comp^{op} \rightarrow Cat_{\infty}$ are not sheaves.

2.17 Lemma. Let $S: \operatorname{Comp}^{\operatorname{op}} \to \operatorname{LTop}_{\infty}$ be a sheaf and assume that the restriction of S to profinite sets factors through $\operatorname{LTop}_{\infty}^{\operatorname{post}}$. Then for each compact Hausdorff space K, the ∞ -topos S(K) is Postnikov complete.

Proof. Since every compact Hausdorff space admits a surjection from a profinite set, this follows from the fact that the subcategory $LTop_{\infty}^{post} \subset LTop_{\infty}$ is closed under limits (1.25).

Since the functors Sh and Sh^{post} agree on profinite sets (Example 1.28), Example 2.15 shows:

2.18 Corollary. The presheaves Sh, Sh^{hyp}: $Comp^{op} \rightarrow LTop_{\infty}$ are not sheaves.

3 The comparison functor

Let X be a topological space. In this section, we construct a comparison geometric morphism

$$c_X^*$$
: Sh(X) \rightarrow Sh(Comp)_{/X}.

The idea is to define c_X^* by left Kan extending the functor $d_X : \operatorname{Open}(X) \to \operatorname{Sh}(\operatorname{Comp})_{/X}$ to presheaves on *X* along the Yoneda embedding. A priori, the right adjoint to this left Kan extension functor only lands in presheaves on *X*. To see that it factors through sheaves on *X* amounts to showing that the ∞ -topos Sh(Comp) has descent for open covers of topological spaces. We prove this in §3.1. Subsection 3.2 constructs the comparison functor and gives an alternative description when *X* is a profinite set. In §3.3, we explain two naturality properties of the comparison functor.

3.1 Descent for open covers

3.1 Lemma. Let X be a topological space and $\{U_i\}_{i \in I}$ an open cover of X. Then the natural map $\prod_{i \in I} \& (U_i) \to \& (X)$ is an effective epimorphism in Sh(**Comp**).

Proof. First we prove the claim under the assumption that X is compact Hausdorff. In this case, since X is compact, there exists a finite subset $I_0 \subset I$ such that $\{U_i\}_{i \in I_0}$ covers X. Since X is compact Hausdorff, there exists a cover of X by closed subsets $\{Z_i\}_{i \in I_0}$ such that $Z_i \subset U_i$.

Since \downarrow : **Comp** \hookrightarrow Sh(**Comp**) preserves finite coproducts and carries surjections to effective epimorphisms, the composite map

is an effective epimorphism. Hence the second map $\prod_{i \in I} \sharp(U_i) \to \sharp(X)$ is also an effective epimorphism.

Now we prove the claim in general. We need to show that for each compact Hausdorff space K and map $\tilde{f} : \sharp(K) \to \sharp(X)$ in Sh(**Comp**), the induced map

$$\sharp(K) \underset{\sharp(X)}{\times} \coprod_{i \in I} \sharp(U_i) \to \sharp(K)$$

is an effective epimorphism. Since *K* is compact Hausdorff, by the Yoneda lemma the map $\tilde{f} : {}_{\&}(K) \to {}_{\&}(X)$ is induced by a map of topological spaces $f : K \to X$. Since coproducts in Sh(**Comp**) are universal and ${}_{\&}: \mathbf{Top} \to \mathrm{Sh}(\mathbf{Comp})$ preserves limits, we see that

$$\begin{split} \sharp(K) \underset{\bigstar(X)}{\times} \coprod_{i \in I} \sharp(U_i) &\simeq \coprod_{i \in I} \sharp(K) \underset{\bigstar(X)}{\times} \sharp(U_i) \\ &\simeq \coprod_{i \in I} \sharp(f^{-1}(U_i)) \,. \end{split}$$

Since $\{f^{-1}(U_i)\}_{i \in I}$ is an open cover of *K*, the claim now follows from the compact case.

3.2 Corollary. Let X be a topological space and let $\{U_i\}_{i \in I}$ be an open cover of X. Then the induced augmented simplicial object

$$\cdots \stackrel{\longrightarrow}{\Longrightarrow} \coprod_{i_1,i_2 \in I} \measuredangle(U_{i_1} \cap U_{i_2}) \stackrel{\longrightarrow}{\Longrightarrow} \coprod_{i \in I} \measuredangle(U_i) \longrightarrow \measuredangle(X)$$

exhibits $\sharp(X)$ as the geometric realization of its restriction to $\Delta^{\text{op}} \subset \Delta^{\text{op}}_+$.

Proof. This follows from the fact that $\coprod_{i \in I} \sharp(U_i) \to \sharp(X)$ is an effective epimorphism and the computation

$$\begin{split} \left(\coprod_{i \in I} \measuredangle(U_i) \right) \underset{\bigstar(X)}{\times} \cdots \underset{\bigstar(X)}{\times} \left(\coprod_{i \in I} \measuredangle(U_i) \right) &\simeq \coprod_{i_1, \dots, i_n \in I} \measuredangle(U_{i_1}) \underset{\bigstar(X)}{\times} \cdots \underset{\bigstar(X)}{\times} \measuredangle(U_{i_n}) \\ &\simeq \coprod_{i_1, \dots, i_n \in I} \measuredangle(U_{i_1} \cap \dots \cap U_{i_n}) \,. \end{split}$$

The usual cofinality argument shows:

3.3 Corollary. Let X be a topological space and let \mathcal{U} be a covering sieve of X. Then the natural map

$$\operatorname{colim}_{U \in \mathcal{U}} \sharp(U) \to \sharp(X)$$

is an equivalence in Sh(Comp).

Since the methods of proof are similar, we conclude this subsection showing that the functor &: **Top** \rightarrow Sh(**Comp**) carries proper surjections to effective epimorphisms and preserves coproducts. These results are not used in the rest of the paper.

3.4 Lemma. Let $p: X \rightarrow Y$ be a proper surjection of locally compact Hausdorff spaces. Then $\sharp(p): \sharp(X) \rightarrow \sharp(Y)$ is an effective epimorphism in Sh(**Comp**).

Proof. Since \downarrow preserves pullbacks and is fully faithful on locally compact Hausdorff spaces, we need to show that for every compact Hausdorff space *K* and map $K \rightarrow Y$, the induced map

is an effective epimorphism. Since p is a proper surjection, the pullback $\bar{p} : K \times_Y X \to K$ is also a proper surjection. Since K is compact and \bar{p} is proper, $K \times_Y X$ is also compact. The fact that &: **Comp** \hookrightarrow Sh(**Comp**) carries surjections to effective epimorphisms completes the proof. \Box

3.5 Lemma. The functor \downarrow : **Top** \rightarrow Sh(**Comp**) preserves coproducts.

Proof. Let $\{X_i\}_{i \in I}$ be a collection of topological spaces. We need to show that for any compact Hausdorff space *K* and map $f : K \to \prod_{i \in I} X_i$, the induced map

(3.6)
$$\qquad \qquad \boldsymbol{\natural}(K) \underset{\boldsymbol{\natural}(\coprod_{j \in I} X_j)}{\times} \coprod_{i \in I} \boldsymbol{\natural}(X_i) \to \boldsymbol{\natural}(K)$$

is an equivalence. Since \downarrow preserves pullbacks and coproducts in Sh(**Comp**) are universal, we see that

$$\begin{split} & \downarrow(K) \underset{\& (\coprod_{j \in I} X_j)}{\times} \coprod_{i \in I} \& (X_i) \simeq \coprod_{i \in I} \& (K) \underset{\& (\coprod_{j \in I} X_j)}{\times} \& (X_i) \\ & \simeq \coprod_{i \in I} \& \left(K \underset{\coprod_{j \in I} X_j}{\times} X_i \right) \\ & = \coprod_{i \in I} \& (f^{-1}(X_i)) \,. \end{split}$$

Since *K* is compact, there is a finite subset $I_0 \subset I$ such that $f^{-1}(X_i) \neq \emptyset$ if and only if $i \in I_0$. Hence

$$\sharp(K) \underset{\sharp(\coprod_{j \in I} X_j)}{\times} \simeq \coprod_{i \in I_0} \sharp(f^{-1}(X_i))$$

and the map $\prod_{i \in I_0} \sharp(f^{-1}(X_i)) \to \sharp(K)$ is induced by the inclusions $f^{-1}(X_i) \hookrightarrow K$. Note that

$$K \cong \coprod_{i \in I_0} f^{-1}(X_i)$$

Moreover, each $f^{-1}(X_i)$ is clopen in *K*, hence compact Hausdorff. Thus the fact that the Yoneda embedding \downarrow : **Comp** \hookrightarrow Sh(**Comp**) preserves finite coproducts shows that the map (3.6) is an equivalence.

3.2 The comparison functor

In this subsection, we define the comparison functor and give an alternative description of this functor in the case of a profinite set.

3.7 Definition. Let *X* be a topological space. Write

$$c_X^*$$
: PSh(X) \rightarrow Sh(**Comp**)_{/X}

for the left Kan extension of the functor

$$\sharp$$
: Open(X) \rightarrow Sh(**Comp**)_{/X}

along the Yoneda embedding $Open(X) \hookrightarrow PSh(X)$.

3.8. Since the functor \sharp : Open(X) \rightarrow Sh(Comp)_{/X} is left exact, by [HTT, Proposition 6.1.5.2], the functor c_x^* is also left exact.

3.9 Observation. The functor c_x^* has a right adjoint

$$c_{X,*}$$
: Sh(**Comp**)_{/X} \rightarrow PSh(X)

given by the formula

$$c_{X,*}(G)(U) \coloneqq \operatorname{Map}_{\operatorname{Sh}(\operatorname{\mathbf{Comp}})_{/Y}}({}^{\sharp}(U),G).$$

By Corollary 3.3, we see that for each $G \in \text{Sh}(\text{Comp})_{/X}$, the presheaf $c_{X,*}(G)$ is a sheaf. Hence the adjunction $c_X^* \dashv c_{X,*}$ restricts to a geometric morphism

$$\operatorname{Sh}(X) \xrightarrow[c_{X,*}]{c_{X,*}} \operatorname{Sh}(\operatorname{\mathbf{Comp}})_{/X}.$$

3.10 Remark. If X is not compactly generated, then the functor

$$\&$$
 : Open(*X*) → Sh(**Comp**; **Set**)_{/*X*}

need not be fully faithful. Hence, the functor c_X^* : Sh(X; Set) \rightarrow Sh(Comp; Set)_{/X} need not be fully faithful.

In the remainder of this subsection, we give an alternative description of the comparison geometric morphism when X is a profinite set. To do so, we use the following description of $Sh(Comp)_{/X}$ as sheaves on a site.

3.11 Recollection. Let $F \in \text{Sh}(\text{Comp})$ be a sheaf of sets. Write $\text{Comp}_{/F} \subset \text{Sh}(\text{Comp})_{/F}$ for the full subcategory spanned by morphisms from representable sheaves. Give $\text{Comp}_{/F}$ the finest Grothendieck toplogy that contains the sieves that become covering sieves after applying the forgetful functor $\text{Comp}_{/F} \to \text{Comp}$. With respect to this topology, the natural functor

$$\operatorname{Sh}(\operatorname{\mathbf{Comp}}_{/F}) \to \operatorname{Sh}(\operatorname{\mathbf{Comp}})_{/F}$$

is an equivalence of ∞ -categories [SGA 4₁, Exposé III, §5, Proposition 5.4].

3.12. Let *X* be a topological space. To simplify notation for sheaves with coefficients, we implicitly make use of the identification

$$\operatorname{Sh}(\operatorname{\mathbf{Comp}}_{/X}) \simeq \operatorname{Sh}(\operatorname{\mathbf{Comp}})_{/X}$$

provided by Recollection 3.11 repeatedly throughout the rest of these notes.

3.13 Recollection. Let *S* be a profinite set, and write $\operatorname{Clop}(S) \subset \operatorname{Open}(S)$ for the poset of clopen subsets of *S*. Then $\operatorname{Clop}(S)$ is a basis closed under finite intersection. Hence for any presentable ∞ -category \mathcal{E} , restriction along the inclusion $\operatorname{Clop}(S)^{\operatorname{op}} \subset \operatorname{Open}(S)^{\operatorname{op}}$ defines an equivalence of ∞ -categories

$$\operatorname{Sh}(S; \mathcal{E}) \xrightarrow{\sim} \operatorname{Sh}(\operatorname{Clop}(S); \mathcal{E})$$

The inverse equivalence is given by right Kan extension [1, Corollary A.8; 2, Corollary 3.12.14].

3.14. For a profinite set *S*, the inclusion $Clop(S) \subset Comp_{/S}$ is a morphism of sites.

3.15 Observation. Let *S* be a profinite set and \mathcal{E} a presentable ∞ -category. The composite

$$\operatorname{Sh}(\operatorname{\mathbf{Comp}}_{/S}) \simeq \operatorname{Sh}(\operatorname{\mathbf{Comp}})_{/S} \xrightarrow{c_{S,*}} \operatorname{Sh}(S; \mathcal{E}) \xrightarrow{\sim} \operatorname{Sh}(\operatorname{Clop}(S); \mathcal{E})$$

carries a sheaf G: $(\mathbf{Comp}_{/S})^{\mathrm{op}} \to \mathbf{Spc}$ to its restriction to $\mathrm{Clop}(S)^{\mathrm{op}} \subset (\mathbf{Comp}_{/S})^{\mathrm{op}}$. As a consequence, the geometric morphism

$$\operatorname{Sh}(S) \xrightarrow[c_{S,*}]{c_{S,*}} \operatorname{Sh}(\operatorname{\mathbf{Comp}}_{/S}).$$

is induced by the inclusion of sites $Clop(S) \subset Comp_{/S}$.

3.3 Naturality of the comparison functor

3.16 Lemma. Let $f : X \to Y$ be a map of topological spaces. Then the square

(3.17)
$$\begin{array}{ccc} \operatorname{Sh}(Y) & \xrightarrow{c_Y^*} & \operatorname{Sh}(\operatorname{Comp})_{/Y} \\ f^* & & & \downarrow_{X \times_Y(-)} \\ & \operatorname{Sh}(X) & \xrightarrow{c_X^*} & \operatorname{Sh}(\operatorname{Comp})_{/X} \end{array}$$

canonically commutes.

Proof. Since both composites are left adjoints and Sh(Y) is generated under colimits by Open(Y), it suffices to show that these composites agree on opens. For this, note for $V \subset Y$ open, we have

$$c_X^* f^*(V) = \pounds(f^{-1}(V)) \qquad \text{(by definition)}$$

$$\simeq \pounds(V) \underset{\&(Y)}{\times} \pounds(X) \qquad (\& \text{ preserves pullbacks})$$

$$= c_Y^*(V) \times_Y X \qquad (by definition) \qquad . \qquad \Box$$

3.18. In light of Lemma 3.16, the comparison functors assemble into a natural transformation $Sh(-) \rightarrow Sh(Comp)_{/(-)}$ of presheaves of ∞ -categories on **Top**.

For open embeddings, the square (3.17) is also vertically left adjointable:

3.19 Lemma. Let $j: U \hookrightarrow X$ be an open embedding of topological spaces. Then the square

$$\begin{array}{ccc} \operatorname{Sh}(U) & \stackrel{c_U^*}{\longrightarrow} & \operatorname{Sh}(\operatorname{\mathbf{Comp}})_{/U} \\ & & & & \\ j_! \int & & & & \\ \operatorname{Sh}(X) & \stackrel{c_X^*}{\longrightarrow} & \operatorname{Sh}(\operatorname{\mathbf{Comp}})_{/X} \end{array}$$

canonically commutes.

Proof. Since both composites are left adjoints and Sh(U) is generated under colimits by the opens of U, it suffices to show that these composites agree on opens. For this, note that for $V \subset U$ open, by definition we have

$$c_X^* j_!(V) = c_X^*(V) = \downarrow(V) = c_U^*(V)$$

as objects of $Sh(Comp)_{/X}$.

4 Full faithfulness of the comparison functor

Let \mathcal{E} be a compactly assembled ∞ -category (Recollection 1.10). In this section, we show that if *X* is a locally compact Hausdorff space, then the comparison functor

$$c^{*,\text{post}}$$
: $\text{Sh}^{\text{post}}(X;\mathcal{E}) \to \text{Sh}^{\text{post}}(\text{Comp}_{/X};\mathcal{E})$

is fully faithful (Corollary 4.11). Since cohomology is computed by global sections, this implies that the sheaf cohomology and condensed cohomology of *X* agree (see Corollary 4.12 and Remark 4.13).

In §4.1, we begin by proving the claim when X is an extremally disconnected profinite set. In §4.2, we use proper descent (Corollary 2.8) and the functoriality of the comparison morphism to prove the result in general. Subsection 4.3 explains why the functor c_X^* : $Sh(X) \rightarrow Sh(Comp)_{/X}$ is not generally fully faithful before Postnikov completion.

4.1 Full faithfulness for extremally disconected profinite sets

Let *S* be an extremally disconnected profinite set. To prove that the functor c_S^* is fully faithful, we verify that the inclusion $\operatorname{Clop}(S) \subset \operatorname{Comp}_{/S}$ satisfies the *covering lifting property*.⁴

4.1 Recollection. If *S* is an extremally disconnected topological space, and $U \subset S$ is open, then *U* is also extremally disconnected. However, closed subsets of extremally disconnected topological spaces need not be extremally disconnected [4, Remark 2.4.11].

4.2 Lemma. Let S be an extremally disconnected profinite set. Then the inclusion of sites

 $Clop(S) \hookrightarrow Comp_{/S}$

satisfies the covering lifting property.

⁴The reader unfamiliar with the covering lifting property should consult [19, Definition A.12].

Proof. Let $U \subset S$ be a clopen subset and let $\{f_i : K_i \to U\}_{i \in I}$ be a finite jointly surjective family in **Comp**_{/S}. We claim that there exits a cover $\{V_i\}_{i \in i}$ of U by clopens such that each inclusion $V_i \hookrightarrow U$ factors through $f_i : K_i \to U$. To see this, first note that since $U \subset S$ is clopen, U is also an extremally disconnected profinite set. Thus the surjection

$$\coprod_{i\in I}f_i:\coprod_{i\in I}K_i\twoheadrightarrow U$$

admits a section $s: U \to \coprod_{i \in I} K_i$. For each $i \in I$, define $V_i := s^{-1}(K_i)$. Since K_i is a clopen subset of $\coprod_{i \in I} K_i$, the subset $V_i \subset U$ is also clopen. Moreover, $\{V_i\}_{i \in I}$ covers U. Lastly, by construction, for each $i \in I$, the composite

$$V_i \xrightarrow{s|_{V_i}} K_i \xrightarrow{f_i} U$$

coincides with the inclusion $V_i \subset U$.

The following is an application of Observation 3.15, Lemma 4.2, and [19, Proposition A.13].

4.3 Notation. Let *S* be a profinite set and \mathcal{E} a presentable ∞ -category. Write

$$\mathcal{C}_{S}^{\sharp}$$
: PSh(Clop(S); \mathcal{E}) \hookrightarrow PSh(**Comp**_{/S}; \mathcal{E})

for the functor given by right Kan extension the inclusion $\operatorname{Clop}(S)^{\operatorname{op}} \hookrightarrow (\operatorname{Comp}_{/S})^{\operatorname{op}}$.

4.4 Corollary. Let *S* be an extremally disconnected profinite set and \mathcal{E} a presentable ∞ -category. *Then:*

(4.4.1) The functor c_s^{\sharp} preserves sheaves. In particular, c_s^{\sharp} restricts to a fully faithful right adjoint to

$$c_{S,*}$$
: Sh(**Comp**_{/S}; \mathcal{E}) \rightarrow Sh(S; \mathcal{E}).

(4.4.2) The left adjoint c_s^* : Sh $(S; \mathcal{E}) \hookrightarrow$ Sh $(\mathbf{Comp}_{/S}; \mathcal{E})$ is fully faithful.

4.2 Full faithfulness and descent

In this subsection, we extend (4.4.2) to Postnikov complete sheaves on a locally compact Hausdorff space. We begin by extending to compact Hausdorff spaces by proper descent. To do so, we make use of the following well-known lemma:

4.5 Lemma. Let \mathcal{I} be an ∞ -category, let $\mathcal{X}_{\bullet}, \mathcal{Y}_{\bullet} : \mathcal{I} \to \mathbf{Cat}_{\infty}$ be diagrams of ∞ -categories, and let $f_{\bullet} : \mathcal{X}_{\bullet} \to \mathcal{Y}_{\bullet}$ be a natural transformation. If for each $i \in \mathcal{I}$, the functor $f_i : \mathcal{X}_i \to \mathcal{Y}_i$ is fully faithful, then the induced functor on limits

$$\lim_{i \in I} f_i : \lim_{i \in I} \mathcal{X}_i \to \lim_{i \in I} \mathcal{Y}_i$$

is fully faithful.

4.6 Corollary. Let f^* : $\mathcal{Y} \to \mathcal{X}$ be a left exact left adjoint functor between ∞ -topoi. The following are equivalent:

(4.6.1) The induced functor on Postnikov completions $f^{*,\text{post}}$: $\mathcal{Y}^{\text{post}} \rightarrow \mathcal{X}^{\text{post}}$ is fully faithful.

(4.6.2) For each integer $n \ge 0$, the restriction f^* : $\mathcal{Y}_{\le n} \to \mathcal{X}_{\le n}$ is fully faithful.

Proof. Remark 1.18 shows that $(4.6.1) \Rightarrow (4.6.2)$. Lemma 4.5 shows that $(4.6.2) \Rightarrow (4.6.1)$.

4.7 Corollary. Let S be an extremally disconnected profinite set and \mathcal{E} a compactly assembled ∞ -category. Then the comparison functor

$$c_S^{*,\text{post}}$$
: $\text{Sh}(S;\mathcal{E}) \to \text{Sh}^{\text{post}}(\text{Comp}_{/S};\mathcal{E})$

is fully faithful.

Proof. Since $c^{*,\text{post}}$: Sh(*S*) \rightarrow Sh^{post}(**Comp**_{*/S*}) is left exact and \mathcal{E} is compactly assembled, by tensoring with \mathcal{E} it suffices to prove the claim for Postnikov sheaves valued in the ∞ -category of spaces [10, Lemma 2.14]. The claim now follows from Corollaries 4.4 and 4.6.

4.8 Recollection (van Kampen colimits). A colimit in an ∞ -category \mathcal{X} with pullbacks is *van Kampen* if the functor $\mathcal{X}^{op} \rightarrow \mathbf{Cat}_{\infty}$ given by $U \mapsto \mathcal{X}_{/U}$ transforms it into a limit in \mathbf{Cat}_{∞} . A presentable ∞ -category \mathcal{X} is an ∞ -topos if and only if all colimits in \mathcal{X} are van Kampen; see [HTT, Proposition 5.5.3.13, Theorem 6.1.3.9(3), & Proposition 6.3.2.3; 15].

4.9 Corollary. Let K be a compact Hausdorff space and \mathcal{E} a compactly assembled ∞ -category. Then the comparison functor

$$c_K^{*,\text{post}}$$
: $\text{Sh}^{\text{post}}(K;\mathcal{E}) \to \text{Sh}^{\text{post}}(\text{Comp}_{/K};\mathcal{E})$

is fully faithful.

Proof. Since the functors

$$K \mapsto \operatorname{Sh}^{\operatorname{post}}(K; \mathcal{E})$$
 and $K \mapsto \operatorname{Sh}^{\operatorname{post}}(\operatorname{Comp}_{/K}; \mathcal{E})$

are hypersheaves on **Comp** (Corollary 2.8 and Recollection 4.8, respectively), by choosing a hypercover of *K* by extremally disconnected profinite sets and applying Lemma 4.5, we are reduced to the case where *K* is extremally disconnected. We conclude by Corollary 4.7. \Box

The full faithfulness of $c_{\chi}^{*,\text{post}}$ is preserved by passing to open subspaces:

4.10 Lemma. Let $j : U \hookrightarrow X$ be an open embedding of topological spaces and let \mathcal{E} be a compactly assembled ∞ -category. If $c_X^{*,\text{post}} : \operatorname{Sh}^{\operatorname{post}}(X) \to \operatorname{Sh}^{\operatorname{post}}(\operatorname{Comp}_{/X})$ is fully faithful, then the functor

$$c_U^{*,\text{post}}$$
: $\text{Sh}^{\text{post}}(U;\mathcal{E}) \to \text{Sh}^{\text{post}}(\text{Comp}_{/U};\mathcal{E})$

is also fully faithful.

Proof. Since $c_U^{*,\text{post}}$: Sh(U) \rightarrow Sh^{post}(**Comp**_{/U}) is left exact and \mathcal{E} is compactly assembled, by tensoring with \mathcal{E} it suffices to prove the claim for Postnikov sheaves valued in the ∞ -category of spaces [10, Lemma 2.14]. In this case, Lemma 3.19 shows that for each integer $n \ge 0$, the diagram

$$\begin{array}{ccc} \operatorname{Sh}(U)_{\leq n} & \stackrel{c_U}{\longrightarrow} & (\operatorname{Sh}(\operatorname{\mathbf{Comp}})_{/U})_{\leq n} \\ & & & & \\ j_! \int & & & & \\ \operatorname{Sh}(X)_{\leq n} & \stackrel{c_X^*}{\longrightarrow} & (\operatorname{Sh}(\operatorname{\mathbf{Comp}})_{/X})_{\leq n} \end{array}$$

commutes. Since $c_X^{*,\text{post}}$ is fully faithful, the bottom horizontal functor is fully faithful. Hence the top horizontal functor is also fully faithful. We conclude by Corollary 4.6.

Since every locally compact Hausdorff space embeds as an open in a compact Hausdorff space, Corollary 4.9 and Lemma 4.10 show:

4.11 Corollary. Let X be a locally compact Hausdorff space and \mathcal{E} a compactly assembled ∞ -category. Then the functor

$$c_X^{*,\text{post}}$$
: $\text{Sh}^{\text{post}}(X;\mathcal{E}) \to \text{Sh}^{\text{post}}(\mathbf{Comp}_{/X};\mathcal{E})$

is fully faithful.

We conclude this subsection with some cohomological consequences of Corollary 4.11. The following generalizes [7, Theorem 3.11; 23, Theorem 3.2].

4.12 Corollary. Let X be a locally compact Hausdorff space, let R be a connective E_1 -ring spectrum, and let M be a bounded-above left R-module spectrum. Then the natural comparison map

$$R\Gamma_{\text{sheaf}}(X;M) \to R\Gamma_{\text{cond}}(X;M)$$

is an equivalence in the ∞ -category LMod(R) of left R-module spectra.

4.13 Remark (condensed, singular, and sheaf cohomology). Let *R* be a ring and *X* a topological space that is locally compact Hausdorff and locally weakly contractible. Write $C_*(X; R) \in D(R)$ for the complex of *singular chains* on *X*. Given an object $M \in D(R)$, write

$$C^{-*}(X; M) \coloneqq \operatorname{RHom}_R(C_*(X; R), M)$$

If *M* is an ordinary *R*-module, then $C^{-*}(X; M)$ is what is usually referred to as the complex of *singular cochains* on *X* with values in *M*.

If *M* is t-bounded-above, then Corollary 4.12 and [11, Corollary 3.31] provide natural equivalences

$$\mathrm{R}\Gamma_{\mathrm{cond}}(X;M) \leftarrow \mathrm{R}\Gamma_{\mathrm{sheaf}}(X;M) \xrightarrow{\sim} \mathrm{C}^{-*}(X;M)$$
.

Hence the condensed, singular, and sheaf cohomologies of *X* all agree.

4.14. Let *X* be a topological space that admits a locally finite CW structure. Since Sh(X) is Postnikov complete, Corollary 4.12 and [11, Corollary 3.31] actually imply that for *any* object $M \in D(R)$, there are natural equivalences

$$R\Gamma_{cond}(X;M) \leftarrow R\Gamma_{sheaf}(X;M) \rightarrow C^{-*}(X;M)$$

4.3 The necessity of Postnikov completion

We conclude by explaining why Corollary 4.11 is generally false before passing to Postnikov completions. First we explain why Corollary 4.11 is false if Sh^{post} is replaced by Sh.

4.15 Lemma. Let K be a compact Hausdorff space. Then the following are equivalent:

(4.15.1) The functor c_K^* : Sh(K) \rightarrow Sh(**Comp**_{/K}) is conservative.

(4.15.2) The ∞ -topos Sh(K) is hypercomplete.

Proof. Choose an extremally disconnected profinite set *S* equipped with a surjection $p: S \rightarrow K$. Consider the commutative square

$$\begin{array}{ccc} \operatorname{Sh}(K) & \stackrel{c_K^*}{\longrightarrow} & \operatorname{Sh}(\operatorname{\mathbf{Comp}})_{/K} \\ p^* & & & \downarrow_{S \times_K(-)} \\ \operatorname{Sh}(S) & \stackrel{c_S^*}{\longrightarrow} & \operatorname{Sh}(\operatorname{\mathbf{Comp}})_{/S} . \end{array}$$

Since *S* is an extremally disconnected profinite set, c_S^* is fully faithful (Corollary 4.4). Since *p* is surjective and the functor Sh(Comp)_{/(-)} satisfies descent for surjections of compact Hausdorff spaces, the right-hand vertical functor is conservative. Hence c_K^* is conservative if and only if p^* is conservative. To conclude, note that since *p* is surjective and Sh(*S*) is hypercomplete, the pullback functor p^* is conservative if and only if Sh(*K*) is hypercomplete.

4.16 Warning. Since there exist compact Hausdorff spaces *K* for which Sh(K) is not hypercomplete [HTT, Counterexample 6.5.4.8], Lemma 4.15 implies that the functor c_K^* is *not* generally fully faithful.

Now we explain why Corollary 4.11 is false if Sh^{post} is replaced by Sh^{hyp}.

4.17 Lemma. Let X be a topological space. If $c_X^{*,hyp}$: $\operatorname{Sh}^{hyp}(X) \to \operatorname{Sh}^{hyp}(\operatorname{Comp})_{/X}$ fully faithful, then every object of $\operatorname{Sh}^{hyp}(X)$ is the limit of its Postnikov tower.

4.18 Lemma. Let $f^* : \mathcal{Y} \hookrightarrow \mathcal{X}$ be a fully faithful left exact left adjoint between ∞ -topoi. If every object of \mathcal{X} is the limit of its Postnikov tower, then every object of \mathcal{Y} is the limit of its Postnikov tower.

Proof. By Observation 1.17, we need to show that $t_y^* : \mathcal{Y} \to \mathcal{Y}^{\text{post}}$ is fully faithful. Consider the commutative square

$$\begin{array}{ccc} \mathcal{Y} & \xrightarrow{t_{\mathcal{Y}}^*} & \mathcal{Y}^{\text{post}} \\ f^* & & & \downarrow f^{*,\text{post}} \\ \mathcal{X} & \xrightarrow{t_{\mathcal{X}}^*} & \mathcal{X}^{\text{post}} \end{array}.$$

By Corollary 4.6, the functor $f^{*,\text{post}}$ is fully faithful, and by assumption $t_{\mathcal{X}}^*$ is fully faithful. Hence $t_{\mathcal{Y}}^*$ is also fully faithful.

Proof of Lemma 4.17. Combine Lemma 4.18 with the fact that $Sh^{hyp}(Comp)_{/X}$ is Postnikov complete (1.37).

4.19 Warning. Example 2.16 and Lemma 4.17 show that for $X = \prod_{m \ge 1} S^m$, the functor $c_X^{*,hyp}$ is *not* fully faithful.

References

- HTT J. Lurie, *Higher topos theory*, Annals of Mathematics Studies. Princeton, NJ: Princeton University Press, 2009, vol. 170, pp. xviii+925, ISBN: 978-0-691-14049-0; 0-691-14049-9.
- HA _____, *Higher algebra*, Preprint available at math.ias.edu/~lurie/papers/HA.pdf, Sep. 2017.

- SAG _____, Spectral algebraic geometry, Preprint available at math.ias.edu/~lurie/papers/SAGrootfile.pdf, Feb. 2018.
- SGA 4, Théorie des topos et cohomologie étale des schémas. Tome 1: Théorie des topos, Séminaire de Géométrie Algébrique du Bois Marie 1963–64 (SGA 4). Dirigé par M. Artin, A. Grothendieck, J.-L. Verdier. Avec la collaboration de N. Bourbaki, P. Deligne, B. Saint–Donat. Lecture Notes in Mathematics, Vol. 269. Berlin: Springer-Verlag, 1963–64, pp. xix+525.
 - 1. K. Aoki, *Tensor triangular geometry of filtered objects and sheaves*, Preprint available at arXiv: 2001.00319, Jan. 2020.
 - 2. C. Barwick, S. Glasman, and P. J. Haine, *Exodromy*, Preprint available at arXiv:1807.03281v7, Aug. 2020.
 - 3. C. Barwick and P. J. Haine, *Pyknotic objects, I. Basic notions*, Preprint available at arXiv:1904. 09966, Apr. 2019.
 - 4. B. Bhatt and P. Scholze, *The pro-étale topology for schemes*, Astérisque, no. 369, pp. 99–201, 2015.
 - 5. D. Clausen and P. Scholze, *Condensed mathematics and complex geometry*, Lecture notes available at people.mpim-bonn.mpg.de/scholze/Complex.pdf, Apr. 2022.
 - 6. D. Dugger, S. Hollander, and D. C. Isaksen, *Hypercovers and simplicial presheaves*, Math. Proc. Cambridge Philos. Soc., vol. 136, no. 1, pp. 9–51, 2004. DOI: 10.1017/S0305004103007175.
 - 7. R. Dyckhoff, Categorical methods in dimension theory, in Categorical topology (Proc. Conf., Mannheim, 1975), 1976, 220–242. Lecture Notes in Math., Vol. 540.
 - 8. _____, *Projective resolutions of topological spaces*, J. Pure Appl. Algebra, vol. 7, no. 1, pp. 115–119, 1976. DOI: 10.1016/0022-4049(76)90069-4.
 - 9. A. M. Gleason, Projective topological spaces, Illinois J. Math., vol. 2, pp. 482–489, 1958.
 - 10. P. J. Haine, *From nonabelian basechange to basechange with coefficients*, Preprint available at arXiv:2108.03545, Sep. 2022.
 - 11. P. J. Haine, M. Porta, and J.-B. Teyssier, *The homotopy-invariance of constructible sheaves*, Preprint available at arXiv:2010.06473, Sep. 2022.
 - 12. L. Hesselholt and P. Pstrągowski, *Dirac geometry I: Commutative algebra*, Preprint available at arXiv: 2207.09256, Jul. 2022.
 - 13. M. Hoyois, Answer to Math Overflow Question 168526: Is the ∞-topos Sh(X) hypercomplete whenever X is a CW complex? M0:168526, Oct. 2016.
 - 14. *Higher Galois theory*, J. Pure Appl. Algebra, vol. 222, no. 7, pp. 1859–1877, 2018. DOI: 10.1016/j.jpaa.2017.08.010.
 - 15. _____, Topoi of parametrized objects, Theory Appl. Categ., vol. 34, Paper No. 9, 243–248, 2019.
 - P. T. Johnstone, *Stone spaces*, Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1986, vol. 3, pp. xxii+370, Reprint of the 1982 edition, ISBN: 0-521-33779-8.
 - 17. J. Lurie, Course notes for *Categorical Logic (278x)*, Available at math.ias.edu/~lurie/278x. html, Spring 2018.
 - , Ultracategories, Preprint available at math.ias.edu/~lurie/papers/Conceptual. pdf, 2018.
 - 19. P. Pstrągowski, *Synthetic spectra and the cellular motivic category*, Preprint available at arXiv: 1803.01804, Feb. 2022.
 - P. Scholze, Condensed mathematics, Talk at MSRI, msri.org/workshops/873/schedules/ 26349, Mar. 2019.

- 21. _____, *Condensed mathematics*, Notes for an MSRI talk taken by Ian Coley, available at msri. org/workshops/873/schedules/26349/documents/50220/assets/85849, Mar. 2019.
- 22. ____, *Lectures on analytic geometry*, Lecture notes available at math.uni-bonn.de/people/ scholze/Analytic.pdf, Oct. 2019.
- 23. _____, Lectures on condensed mathematics, Lecture notes available at math.uni-bonn.de/ people/scholze/Condensed.pdf, Apr. 2019.
- 24. B. Toën and G. Vezzosi, *Homotopical algebraic geometry. I. Topos theory*, Adv. Math., vol. 193, no. 2, pp. 257–372, 2005.