# On the distance spectrum and distance-based topological indices of central vertex-edge join of three graphs

Haritha T<sup>1</sup><sup>\*</sup>, Chithra A. V<sup>1†</sup>

1 Department of Mathematics, National Institute of Technology Calicut, Calicut-673 601, Kerala, India

#### Abstract

Topological indices are molecular descriptors that describe the properties of chemical compounds. These topological indices correlate specific physico-chemical properties like boiling point, enthalpy of vaporization, strain energy, and stability of chemical compounds. This article introduces a new graph operation based on central graph called central vertex-edge join and provides its results related to graph invariants like eccentricconnectivity index, connective eccentricity index, total-eccentricity index, average eccentricity index, Zagreb eccentricity indices, eccentric geometric-arithmetic index, eccentric atom-bond connectivity index, and Wiener index. Also, we discuss the distance spectrum of the central vertex-edge join of three regular graphs. Furthermore, we obtain new families of D-equienergetic graphs, which are non D-cospectral.

Keywords: Distance matrix; Distance eigenvalues; Distance equienergetic graphs; Topological indices.

2020 Mathematics Subject Classification: 05C50, 05C76, 05C12.

#### 1 Introduction

All graphs considered in this paper are simple connected and un-directed. Let  $G_n$  =  $(V(G_n), E(G_n))$  be a graph of order n and size m, with vertices  $v_i, 1 \leq i \leq n$  and edges  $e_j$ ,  $1 \leq j \leq m$ . The *adjacency matrix* of a graph  $G_n$  is a square symmetric matrix  $A(G_n) = (a_{ij})_{n \times n}$ , where  $a_{ij} = 1$  when  $v_i$  is adjacent to  $v_j$  and  $a_{ij} = 0$  otherwise. Denote the eigenvalues of  $A(G_n)$  by  $\theta_1 \ge \theta_2 \ge \cdots \ge \theta_n$ , then the collection of all the eigenvalues of  $A(G_n)$  is said to be the adjacency spectrum of  $G_n$ . The *incidence matrix* of a graph  $G_n$  is the  $n \times m$  matrix  $Q(G_n) = (q_{ij})_{n \times m}$ , where  $q_{ij} = 1$  when  $v_i$  is incident to  $e_j$  and  $q_{ij} = 0$  otherwise. The line graph  $L(G_n)$  of  $G_n$  is a graph with  $V(L(G_n)) = E(G_n)$ , in which two vertices in  $L(G_n)$  are adjacent only if the corresponding edges in  $G_n$  have an end vertex in common. For a partitioned matrix we denote its *equitable quotient matrix* by  $M$  [\[3\]](#page-9-0) and all one entry matrix by  $J_{n\times m}$ . We represent by  $C_n$  the cycle,  $K_{p,q}$   $(p+q=n)$  the complete bipartite graph, and  $K_n$  the complete graph each of order n.

<sup>∗</sup> harithathottungal96@gmail.com

<sup>†</sup> chithra@nitc.ac.in

The *distance*  $d_G(v_i, v_j)$  between two vertices is the length of the shortest path connecting them in  $G_n$  [\[4\]](#page-9-1). In particular, for any  $v_i \in V(G_n)$ ,  $d_{G_n}(v_i, v_i) = 0$ . The eccentricity of a vertex  $v_i \in V(G_n)$  is defined as  $ec_{G_n}(v_i) = max{d(v_i, v_j) : v_j \in V(G_n)}$ . The square symmetric matrix  $D(G_n) = (d_{ij})_{n \times n} = d_{G_n}(v_i, v_j)$  is called the *distance matrix* of  $G_n$  and its eigenvalues are said to be the distance eigenvalues (D-eigenvalues). Let  $\mu_1 \geq \mu_2 \geq \cdots \geq \mu_t$ be the distinct D-eigenvalues of  $G_n$  with multiplicities  $s_1, s_2, \ldots, s_t$ . Then the *distance* spectrum(D-spectrum) of  $D(G_n)$  is given by  $Spec_D(G_n) = \begin{pmatrix} \mu_1 & \mu_2 & \cdots & \mu_t \\ \vdots & \vdots & \ddots & \vdots \\ \vdots$  $s_1$   $s_2$  ...  $s_t$  . The nonisomorphic graphs of the same order having identical D-spectrum are called D-cospectral *graphs.* The transmission  $Tr(v_i)$  of a vertex  $v_i$  is the sum of all distances from  $v_i$  to all other vertices in G. See [\[2\]](#page-9-2) for a survey of the distance spectra of graphs.

In graph theory, operations on graphs play a significant role. For significant works on distance spectra of graph operations see [\[5,](#page-9-3)[11,](#page-10-0)[12\]](#page-10-1). The *central graph*  $C(G_n)$  of  $G_n$  [\[1\]](#page-9-4), is the graph obtained by adding new vertex to each edge of  $G_n$  and joining all the non adjacent vertices in  $G_n$ . In [\[10\]](#page-10-2), the authors defined the central vertex(edge) join of two graphs and their adjacency spectra are investigated when  $G_{n_1}$  and  $G_{n_2}$  are both regular graphs. Inspired by these works, we introduce a new graph operation based on central graph and join.

**Definition 1.1.** Let  $G_{n_1}$ ,  $G_{n_2}$  and  $G_{n_3}$  be any three graphs of orders  $n_1$ ,  $n_2$ ,  $n_3$  and sizes  $m_1, m_2, m_3$  respectively. The *central vertex-edge join* (CVE-join) of  $G_{n_1}$  with  $G_{n_2}$  and  $G_{n_3}$ , is the graph  $G_{n_1}^C \triangleright (G_{n_2}^V \cup G_{n_3}^E)$ , is obtained from  $C(G_{n_1})$ ,  $G_{n_2}$  and  $G_{n_3}$  by joining each vertex of  $G_{n_1}$  with every vertex of  $G_{n_2}$  and each vertex corresponding to the edges of  $G_{n_1}$  with every vertex of  $G_{n_3}$ . The set of vertices in  $C(G_{n_1})$  corresponding to the edges of  $G_{n_1}$  is denoted by  $I(G_{n_1}).$ 

The order and size of  $G_{n_1}^C \triangleright (G_{n_2}^V \cup G_{n_3}^E)$  are  $n_1 + m_1 + n_2 + n_3$  and  $m_1 + m_2 + m_3 +$  $n_1 n_2 + m_1 n_3 + \frac{n_1(n_1-1)}{2}$  respectively.



Figure 1.1.  $C_4^C \triangleright (K_2^V \cup K_2^E)$ .

In general, finding the D-spectrum of complicated networks is a difficult problem. In Section 2, we obtain the D-spectrum of CVE-join of three regular graphs. Using the results mentioned above, we can find the D-spectra of more complicated networks.

The distance energy (D-energy) [\[7\]](#page-10-3) was introduced by Indulal et al., and it is defined as  $E_D(G_n) = \sum_{i=1}^n |\mu_i|$ . Two graphs of the same order having equal D-energy are called D-equienergetic graphs. In the literature  $[8, 9, 12]$  $[8, 9, 12]$  $[8, 9, 12]$ , we can see the constructions of different D-equienergetic graphs. In Section 2, we provide a new class of D-equienergetic graphs of diameter 3 using the above operation.

The study of graphs through their topological descriptors is beneficial for deriving their underlying topologies. This process has many applications in modeling physico-chemical properties of molecules by QSAR/QSPR analysis (quantitative structure- activity relationships/quantitative structure-property relationships). In the studies, QSARs and QSPRs, the topological descriptors of graphs are used to approximate the biological activities and properties of chemical compounds. Various topological indices of chemical structures have emerged as a result of successful applications of QSAR/QSPR studies. This article gives some distance-based topological indices of the new graph operation central vertex-edge join of three graphs.

The *eccentric connectivity index* of  $G_n$  is introduced in [\[13\]](#page-10-6) as

$$
\xi^{c}(G_n) = \sum_{v_i \in V(G_n)} deg_{G_n}(v_i) e_{G_n}(v_i). \tag{1}
$$

The *connective eccentricity index* [\[15\]](#page-10-7) of  $G_n$  is

$$
\xi^{ce}(G_n) = \sum_{v_i \in V(G_n)} \frac{deg_{G_n}(v_i)}{ec_{G_n}(v_i)}.
$$
\n
$$
(2)
$$

By using the eccentric connectivity index, the total-eccentricity index [\[16\]](#page-10-8) is given by

$$
\tau(G_n) = \sum_{v_i \in V(G_n)} e c_{G_n}(v_i). \tag{3}
$$

The mean value of eccentricity of vertices in  $V(G_n)$  is called the *average eccentricity*  $aveg(G_n)$  [\[17\]](#page-10-9),

$$
aveg(G_n) = \frac{1}{n} \sum_{v_i \in V(G_n)} ec_{G_n}(v_i) = \frac{\tau(G_n)}{n}.
$$
\n(4)

The Zagreb indices [\[18\]](#page-10-10) of  $G_n$  in terms of eccentricities is

$$
M_1(G_n) = \sum_{v_i \in V(G_n)} e c_{G_n}^2(v_i),
$$
  
\n
$$
M_2(G_n) = \sum_{v_i v_j \in E(G_n)} e c_{G_n}(v_i) e c_{G_n}(v_j).
$$
\n(5)

The geometric-arithmetic index [\[19\]](#page-10-11) in terms of eccentricity is given by

$$
GA_4(G_n) = \sum_{v_i v_j \in E(G_n)} \frac{2\sqrt{ec_{G_n}(v_i)ec_{G_n}(v_j)}}{ec_{G_n}(v_i) + ec_{G_n}(v_j)}.
$$
(6)

The atom-bond connectivity index [\[20\]](#page-10-12) in terms of eccentricity is given by

$$
ABC_5(G_n) = \sum_{v_i v_j \in E(G_n)} \sqrt{\frac{ec_{G_n}(v_i) + ec_{G_n}(v_j) - 2}{ec_{G_n}(v_i)ec_{G_n}(v_j)}}.
$$
(7)

The *Wiener index*  $W(G)$  [\[14\]](#page-10-13) of  $G_n$  is defined as

$$
W(G_n) = \frac{1}{2} \sum_{v_i, v_j \in V(G_n)} d_G(v_i, v_j) = \frac{1}{2} \sum_{v_i \in V(G_n)} Tr(v_i).
$$
 (8)

Section 3 gives the eccentric connectivity index, connective eccentricity index, totaleccentricity index, average eccentricity index, Zagreb indices, geometric-arithmetic index, atom-bond connectivity index and Wiener index of central vertex-edge join of three graphs.

## 2 Distance spectrum of CVE-join of three regular graphs

In this section, we discuss the D-spectrum of  $G_{n_1}^C \triangleright (G_{n_2}^V \cup G_{n_3}^E)$ , where  $G_{n_1}$  is a triangle-free regular graph and  $G_{n_2}$ ,  $G_{n_3}$  are two regular graphs.

**Lemma 2.1.** [\[6\]](#page-9-5) Let  $G_n$  be a k-regular graph of order n and size m with  $Spec(G_n)$  =  $\{\theta_1, \theta_2, \ldots, \theta_n\}$ . Then

$$
Spec(L(G_n)) = \begin{pmatrix} 2k-2 & \theta_2 + k - 2 & \dots & \theta_n + k - 2 & -2 \\ 1 & 1 & \dots & 1 & m - n \end{pmatrix}.
$$

Moreover,  $-2$  is an eigenvalue of  $L(G_n)$  with eigenvector V if and only if  $Q(G_n)V=0$ .

**Theorem 2.2.** For  $i = 1, 2, 3$ , let  $G_{n_i}$  be a  $k_i$ -regular graph of order  $n_i$  and size  $m_i$ , where  $G_{n_1}$  is triangle-free. Then the D-eigenvalues of  $G_{n_1}^C \triangleright (G_{n_2}^V \cup G_{n_3}^E)$  are the following

(i) 
$$
\frac{-3+ \theta_{1j} \pm \sqrt{(\theta_{1j}+1)^2+4(\theta_{1j}+k_1)}}{2}, \, j=2,3,\ldots,n_1.
$$

(ii)  $-2$  with multiplicity  $m_1 - n_1$ .

(iii)  $-\theta_{ij} - 2, i = 2, 3 \text{ and } j = 2, 3, ..., n_i.$ 

(iv) eigenvalues of the matrix

$$
\begin{bmatrix} n_1 - 1 + k_1 & 2m_1 - k_1 & n_2 & 2n_3 \ 2n_1 - 2 & 2m_1 - 2 & 2n_2 & n_3 \ n_1 & 2m_1 & 2n_2 - k_2 - 2 & 3n_3 \ 2n_1 & m_1 & 3n_2 & 2n_3 - k_3 - 2 \end{bmatrix}
$$

.

*Proof.* By a proper ordering of vertices in  $G_{n_1}^C \triangleright (G_{n_2}^V \cup G_{n_3}^E)$ , the distance matrix  $D(G_{n_1}^C \triangleright$  $(G_{n_2}^V \cup G_{n_3}^E))$  can be written as

$$
D(G_{n_1}^C \triangleright (G_{n_2}^V \cup G_{n_3}^E))
$$
\n
$$
= \begin{bmatrix}\nJ_{n_1} - I_{n_1} + A(G_{n_1}) & 2J_{n_1 \times m_1} - Q(G_{n_1}) & J_{n_1 \times n_2} & 2J_{n_1 \times n_3} \\
2J_{m_1 \times n_1} - Q(G_{n_1})^T & 2(J_{m_1} - I_{m_1}) & 2J_{m_1 \times n_2} & J_{m_1 \times n_3} \\
J_{n_2 \times n_1} & 2J_{n_2 \times m_1} & 2(J_{n_2} - I_{n_2}) - A(G_{n_2}) & 3J_{n_2 \times n_3} \\
2J_{n_3 \times n_1} & J_{n_3 \times n_1} & 3J_{n_3 \times n_2} & 2(J_{n_3} - I_{n_3}) - A(G_{n_3})\n\end{bmatrix}.
$$

Let  $k_i = \theta_{i1} \ge \theta_{i2} \ge \ldots \ge \theta_{in_i}$  be the A-eigenvalues of  $G_{n_i}$ , where  $i = 1, 2, 3$ . Let U be a vector such that  $A(G_{n_1})U = \theta_{1j}U$  (for  $j = 2, 3, ..., n_1$ ), then by Perron-Frobenius theory  $J_{1\times n_1}U=0.$ 

Now let,  $\phi_1 =$  $\sqrt{ }$  $\Big\}$ tU  $Q(G_{n_1})^T U$ 0 0 1  $\parallel$ be an eigenvector of  $D(G_{n_1}^C \triangleright (G_{n_2}^V \cup G_{n_3}^E))$  for the eigenvalue  $\mu$ ,

where t is a real number. Then  $D(G_{n_1}^C \triangleright (G_{n_2}^V \cup G_{n_3}^E))\phi_1 = \mu \phi_1$ . That is,

$$
\begin{bmatrix}\nJ_{n_1} - I_{n_1} + A(G_{n_1}) & 2J_{n_1 \times m_1} - Q(G_{n_1}) & J_{n_1 \times n_2} & 2J_{n_1 \times n_3} \\
2J_{m_1 \times n_1} - Q(G_{n_1})^T & 2(J_{m_1} - I_{m_1}) & 2J_{m_1 \times n_2} & J_{m_1 \times n_3} \\
J_{n_2 \times n_1} & 2J_{n_2 \times m_1} & 2(J_{n_2} - I_{n_2}) - A(G_{n_2}) & 3J_{n_2 \times n_3} \\
2J_{n_3 \times n_1} & J_{n_3 \times m_1} & 3J_{n_3 \times n_2} & 2(J_{n_3} - I_{n_3}) - A(G_{n_3})\n\end{bmatrix}\n\begin{bmatrix}\ntU \\
Q(G_{n_1})^T U \\
0 \\
0\n\end{bmatrix}
$$
\n
$$
= \mu \begin{bmatrix}\ntU \\
Q(G_{n_1})^T U \\
0 \\
0\n\end{bmatrix}
$$
\n
$$
\begin{bmatrix}\n(-t + \theta_{1j}t - \theta_{1j} - k_1)U \\
0 \\
0 \\
0\n\end{bmatrix} = \begin{bmatrix}\n\mu U \\
\mu Q(G_{n_1})^T U \\
0 \\
0\n\end{bmatrix}.
$$
\nTherefore,

$$
-t + \theta_{1j}t - \theta_{1j} - k_1 = \mu t
$$

$$
-t - 2 = \mu
$$

from this, we get

$$
t^2 + (\theta_{1j} + 1)t - (\theta_{1j} + k_1) = 0
$$

so that  $t$  has two values

$$
t_1 = \frac{-(\theta_{1j} + 1) + \sqrt{(\theta_{1j} + 1)^2 + 4(\theta_{1j} + k_1)}}{2},
$$
  

$$
t_2 = \frac{-(\theta_{1j} + 1) - \sqrt{(\theta_{1j} + 1)^2 + 4(\theta_{1j} + k_1)}}{2}.
$$

That is,  $-t_1 - 2$  and  $-t_2 - 2$  are eigenvalues of  $D(G_{n_1}^C \triangleright (G_{n_2}^V \cup G_{n_3}^E))$  corresponding to the eigenvalues  $\theta_{1j} \neq k_1$  of  $A(G_{n_1})$ . Thus, we get  $2(n_1 - 1)$  eigenvalues of  $D(G_{n_1}^C \triangleright (G_{n_2}^V \cup G_{n_3}^E))$ . By Lemma 2.1, we get  $Q(G_{n_1})V = 0$ , where V is an eigenvector of  $L(G_{n_1})$  corresponding to the eigenvalue  $-2$  (repeated  $m_1 - n_1$  times).

If 
$$
\phi_2 = \begin{bmatrix} 0 \\ V \\ 0 \\ 0 \end{bmatrix}
$$
, then we have

$$
D(G_{n_1}^C \triangleright (G_{n_2}^V \cup G_{n_3}^E))\phi_2 = -2\phi_2.
$$

Therefore,  $-2$  is an eigenvalue of  $D(G_{n_1}^C \triangleright (G_{n_2}^V \cup G_{n_3}^E))$  repeated  $m_1 - n_1$  times. Consider the eigenvalue  $\theta_{2j}$  (for  $j = 2, 3, ..., n_2$ ) of  $G_{n_2}$  with an eigenvector W such that  $J_{1\times n_2}W=0.$ 

Put 
$$
\phi_3 = \begin{bmatrix} 0 \\ 0 \\ W \\ 0 \end{bmatrix}
$$
, then we have

$$
D(G_{n_1}^C \triangleright (G_{n_2}^V \cup G_{n_3}^E))\phi_3 = -(\theta_{2j} + 2)\phi_3.
$$

This shows that  $-(\theta_{2j}+2)$  are eigenvalues of  $D(G_{n_1}^C \triangleright (G_{n_2}^V \cup G_{n_3}^E))$  corresponding to the eigenvalues  $\theta_{2j} \neq k_2$  of  $A(G_{n_2})$ .

Similarly, for each eigenvalue  $\theta_{3j}$  (for  $j = 2, 3, ..., n_3$ ) of  $G_{n_3}$ , we get an eigenvalue  $-(\theta_{3j} + 2)$ of  $D(G_{n_1}^C \triangleright (G_{n_2}^V \cup G_{n_3}^E))$  with eigenvector  $\sqrt{ }$   $\theta$  $\theta$  $\theta$ X 1 . Thus, we get  $2n_1-2+m_1-n_1+n_2-1+n_3-1$  =  $n_1 + m_1 + n_2 + n_3 - 4$  eigenvalues of  $D(G_{n_1}^C \triangleright (G_{n_2}^V \cup G_{n_3}^E))$ . All the other eigenvalues of  $D(G_{n_1}^C \triangleright (G_{n_2}^V \cup G_{n_3}^E))$  are the eigenvalues of the equitable quotient matrix

$$
M = \begin{bmatrix} n_1 - 1 + k_1 & 2m_1 - k_1 & n_2 & 2n_3 \\ 2n_1 - 2 & 2m_1 - 2 & 2n_2 & n_3 \\ n_1 & 2m_1 & 2n_2 - k_2 - 2 & 3n_3 \\ 2n_1 & m_1 & 3n_2 & 2n_3 - k_3 - 2 \end{bmatrix}.
$$

Using the above theorem we present some new families of  $D$ -equienergetic graphs.

For a fixed  $a \in \mathbb{N}$ , let  $\mathcal{P}_a$  be the family of all integer partitions of 'a', denoted by  $p_1, p_2, \ldots, p_s$ , each of size at least 3. For  $P = \{p_1, p_2, \ldots, p_s\} \in \mathcal{P}_a$ , let  $C_P$  be the union of cycles with vertices  $p_1, p_2, \ldots, p_s$ .

**Theorem 2.3.** Let  $H_{n_1}$  be a  $k_1$ -regular triangle free graph of order  $n_1$  and size  $m_1$  and  $H_{n_2}$ be a  $k_2$ - regular graph of order  $n_2$  and size  $m_2$  with the least eigenvalue being at least  $-2$ . Then for every  $a \in \mathbb{N}$ ,  $H_{n_1}^C \triangleright (C_P^V \cup H_{n_2}^E)$  forms a family of D-equienergetic graphs.

*Proof.* Let  $k_1 = \theta_1 \ge \theta_2 \ge \cdots \ge \theta_{n_1}$ ,  $2 = \lambda_1 \ge \lambda_2 \ge \cdots \ge \lambda_a$  and  $k_2 = \gamma_1 \ge \gamma_2 \ge \cdots \ge \gamma_{n_2}$ be the A-eigenvalues of  $H_{n_1}$ ,  $C_P$  and  $H_{n_2}$  respectively. From Theorem 2.2, the D-eigenvalues of  $H_{n_1}^C \triangleright (C_P^V \cup H_{n_2}^E)$  are  $\frac{-3+\theta_j \pm \sqrt{(\theta_j+1)^2+4(\theta_j+k_1)}}{2}$  $\frac{1}{2^{n+1}}\frac{\Gamma(1)}{\Gamma(2)}\frac{\Gamma(1)}{\Gamma(2)}$ ,  $j = 2, 3, ..., n_1; -2$  with multiplicity  $m_1 - n_1;$  $-\lambda_{j'}-2, j'=2,3,\ldots,a; -\gamma_{j''}-2, j''=2,3,\ldots,n_2$  and along with the eigenvalues of the equitable quotient matrix M of  $H_{n_1}^C \triangleright (C_P^V \cup H_{n_2}^E)$ ,

$$
\begin{bmatrix} n_1 + k_1 - 1 & 2m_1 - k_1 & a & 2n_2 \ 2n_1 - 2 & 2m_1 - 2 & 2a & n_2 \ n_1 & 2m_1 & 2a - 4 & 3n_2 \ 2n_1 & m_1 & 3a & 2n_2 - k_2 - 2 \end{bmatrix}
$$

for every partition  $P$  of  $'a'$ .

Since  $\lambda_{j'} \geq -2$ , for  $j' = 2, 3, ..., a$ , we have  $-(\lambda_{j'} + 2) \leq 0$ . Similarly, we get  $-(\gamma_{j''} + 2) \leq$  $0, j'' = 2, 3, \ldots, n_2.$ 

Therefore,

$$
\sum_{j'=2}^{a} \left| -\lambda_{j'} - 2 \right| + \sum_{j''=2}^{n_2} \left| -\gamma_{j''} - 2 \right| = \sum_{j'=2}^{a} (\lambda_{j'} + 2) + \sum_{j''=2}^{n_2} (\gamma_{j''} + 2)
$$
  
= -2 + 2(a - 1) - k<sub>2</sub> + 2(n<sub>2</sub> - 1)  
= 2a - k<sub>2</sub> + 2n<sub>2</sub> - 6.

Hence for every partition of  $'a'$ , the energy remains the same.

 $\Box$ 

 $\Box$ 

### 3 Distance-based topological indices of CVE-join of three graphs

For three graphs  $G_{n_1}, G_{n_2}$ , and  $G_{n_3}$ , their CVE-join  $\mathcal{G} = G_{n_1}^C \vee (G_{n_2}^V \cup G_{n_2}^E)$  has the following properties,

$$
deg_G(v) = \begin{cases} n_1 + n_2 - 1, & \text{if } v \in V(G_{n_1}) \\ n_3 + 2, & \text{if } v \in I(G_{n_1}) \\ deg_{G_{n_2}}(v) + n_1, & \text{if } v \in V(G_{n_2}) \\ deg_{G_{n_3}}(v) + m_1, & \text{if } v \in V(G_{n_3}) \end{cases}
$$
(9)

Now we give the eccentricity of vertices in  $\mathcal G$  by two different cases; Case 1. If  $G_{n_1}$  is a triangle free graph,

$$
ecg(v) = \begin{cases} 2, & \text{if } v \in V(G_{n_1}) \text{or } v \in I(G_{n_1}) \\ 3, & \text{if } v \in V(G_{n_2}) \text{ or } v \in V(G_{n_3}) \end{cases} (10)
$$

Case 2. If  $G_{n_1}$  is a graph which is not triangle free, then

$$
ec_{\mathcal{G}}(v) = 3, \text{ for every vertex } v \in V(\mathcal{G}) \tag{11}
$$

This section gives some graph invariants of  $G = G_{n_1}^C \vee (G_{n_2}^V \cup G_{n_2}^E)$  like Wiener index, Zagreb indices, etc.

**Theorem 3.1.** For  $i = 1, 2, 3$ , let  $G_{n_i}$  be a  $k_i$ - regular graph of order  $n_i$  and size  $m_i$ , where  $G_{n_1}$  is triangle-free. Then the Wiener index of  $\mathcal{G} = G_{n_1}^C \vee (G_{n_2}^V \cup G_{n_2}^E)$  is given by,  $W(G) = \frac{1}{2}(n_1^2 - n_1 + 2(n_2^2 + n_3^2 + n_1n_2 - n_2 - n_3 + m_1n_3 + m_1^2) + 4(n_1n_3 + m_1n_1 + m_1n_2$  $m_1) - (n_2k_2 + n_3k_3) + 6n_2n_3).$ 

*Proof.* Let the vertex set of  $\mathcal{G} = G_{n_1}^C \vee (G_{n_2}^V \cup G_{n_2}^E)$  be  $V(G) = \{v_1, v_2, \ldots, v_{n_1}, v_{e_1}, v_{e_2}, \ldots, v_{e_{m_1}}, u_1, u_2, \ldots, u_{n_2}, w_1, w_2, \ldots, w_{n_3}\},$  where  $v_1, v_2, \ldots, v_{n_1}$ are the vertices in  $G_{n_1}, v_{e_1}, v_{e_2}, \ldots, v_{e_{m_1}}$  are the vertices corresponding to the edges of  $G_{n_1}$  in  $C(G_{n_1}), u_1, u_2, \ldots, u_{n_2}$  are the vertices in  $G_{n_2}$ , and  $w_1, w_2, \ldots, w_{n_3}$  are the vertices in  $G_{n_3}$ . Then

$$
Tr(v_i) = n_1 + n_2 + 2n_3 + 2m_1 - 1, \ i = 1, ..., n_1.
$$
  
\n
$$
Tr(v_{e_j}) = 2n_1 + 2n_2 + n_3 + 2m_1 - 4, \ j = 1, ..., m_1.
$$
  
\n
$$
Tr(u_k) = n_1 + 2n_2 + 3n_3 + 2m_1 - k_2 - 2, \ k = 1, ..., n_2.
$$
  
\n
$$
Tr(w_l) = 2n_1 + 3n_2 + 2n_3 + m_1 - k_3 - 2, \ l = 1, ..., n_3.
$$

From (8),

$$
W(G) = \frac{1}{2} \sum_{v_i \in V(G)} Tr(v_i)
$$
  
=  $\frac{1}{2} (n_1^2 - n_1 + 2(n_2^2 + n_3^2 + n_1 n_2 - n_2 - n_3 + m_1 n_3 + m_1^2)$   
+  $4(n_1 n_3 + m_1 n_1 + m_1 n_2 - m_1) - (n_2 k_2 + n_3 k_3) + 6 n_2 n_3).$ 

**Theorem 3.2.** For  $i = 1, 2, 3$ , let  $G_{n_i}$  be a graph of order  $n_i$  and size  $m_i$ , then

(i) 
$$
\xi^{c}(\mathcal{G}) = \begin{cases} 2n_{1}^{2} + 5n_{1}n_{2} + 5m_{1}n_{3} - 2n_{1} + 4m_{1} + 6m_{2} + 6m_{3}, & \text{if } G_{n_{1}} \text{ is triangle-free} \\ 3n_{1}^{2} + 6n_{1}n_{2} + 6m_{1}n_{3} - 3n_{1} + 6(m_{1} + m_{2} + m_{3}), & \text{otherwise} \end{cases}
$$
  
(ii) 
$$
\xi^{ce}(\mathcal{G}) = \begin{cases} \frac{1}{6} \left( 3n_{1}^{2} + 5(n_{1}n_{2} + m_{1}n_{3}) + 6m_{1} + 4(m_{2} + m_{3}) - 3n_{1} \right), & \text{if } G_{n_{1}} \text{ is triangle-free} \\ \frac{1}{3} \left( n_{1}^{2} + 2(n_{1}n_{2} + m_{1}n_{3}) - n_{1} + 2(m_{1} + m_{2} + m_{3}) \right), & \text{otherwise} \end{cases}
$$

.

 $\Box$ 

*Proof.* (i) Let  $\mathcal{G} = G_{n_1}^C \vee (G_{n_2}^V \cup G_{n_2}^E)$ , where  $G_{n_1}$  is triangle-free, then from (1) by using  $(9)$  and  $(10)$ , we have

$$
\xi^{c}(\mathcal{G}) = \sum_{v \in V(G_{n_1})} 2(n_1 + n_2 - 1) + \sum_{v \in I(G_{n_1})} 2(n_3 + 2)
$$
  
+ 
$$
\sum_{v \in V(G_{n_2})} 3(deg_{G_{n_2}}(v) + n_1) + \sum_{v \in V(G_{n_3})} 3(deg_{G_{n_3}}(v) + m_1)
$$
  
= 
$$
2n_1^2 + 5n_1n_2 + 5m_1n_3 - 2n_1 + 4m_1 + 6m_2 + 6m_3.
$$

Similarly in the case when  $G_{n_1}$  is not a triangle-free graph by using (9) and (10) we get,

$$
\xi^{c}(\mathcal{G})=3n_1^2+6n_1n_2+6m_1n_3-3n_1+6(m_1+m_2+m_3).
$$

(ii) Let  $G_{n_1}$  be a triangle-free graph. Then from  $(2)$ , by using  $(9)$  and  $(10)$ , we get

$$
\xi^{ce}(\mathcal{G}) = \sum_{v \in V(G_{n_1})} \frac{n_1 + n_2 - 1}{2} + \sum_{v \in I(G_{n_1})} \frac{n_3 + 2}{2} + \sum_{v \in V(G_{n_2})} \frac{deg_{G_{n_2}} + n_1}{3} + \sum_{v \in V(G_{n_3})} \frac{deg_{G_{n_3}} + m_1}{3} = \frac{1}{6} \left( 3n_1^2 + 5(n_1n_2 + m_1n_3) + 6m_1 + 4(m_2 + m_3) - 3n_1 \right).
$$

Similarly in the case when  $G_{n_1}$  is not a triangle-free graph by using (9) and (10) we get,

$$
\xi^{ce}(\mathcal{G}) = \frac{1}{3} \left( n_1^2 + 2(n_1 n_2 + m_1 n_3) - n_1 + 2(m_1 + m_2 + m_3) \right).
$$

**Theorem 3.3.** For  $i = 1, 2, 3$ , let  $G_{n_i}$  be a graph of order  $n_i$  and size  $m_i$ , then

(i) 
$$
\tau(G) = \begin{cases} 2n_1 + 2m_1 + 3n_2 + 3n_3, & \text{if } G_{n_1} \text{ is triangle-free} \\ 3(n_1 + n_2 + n_3 + m_1), & \text{otherwise} \end{cases}.
$$
  
(ii) 
$$
aveg(G) = \begin{cases} \frac{2n_1 + 2m_1 + 3n_2 + 3n_3}{n_1 + n_2 + n_3 + m_1}, & \text{if } G_{n_1} \text{ is triangle-free} \\ 3, & \text{otherwise} \end{cases}.
$$

*Proof.* (i) Consider  $G = G_{n_1}^C \vee (G_{n_2}^V \cup G_{n_2}^E)$ , when  $G_{n_1}$  is triangle-free, then from (3) by using  $(10)$ , we get

$$
\tau(\mathcal{G}) = 2n_1 + 2m_1 + 3n_2 + 3n_3.
$$

Similarly when  $G_{n_1}$  is not a triangle-free graph we get

$$
\tau(\mathcal{G}) = 3(n_1 + n_2 + n_3 + m_1).
$$

(ii) From  $(4)$ , we get

$$
aveg(\mathcal{G}) = \frac{\tau(\mathcal{G})}{n_1 + n_2 + n_3 + m_1}
$$
  
= 
$$
\begin{cases} \frac{2n_1 + 2m_1 + 3n_2 + 3n_3}{n_1 + n_2 + n_3 + m_1}, & \text{if } G_{n_1} \text{ is triangle-free} \\ 3, & \text{otherwise} \end{cases}.
$$

**Theorem 3.4.** For  $i = 1, 2, 3$ , let  $G_{n_i}$  be a graph of order  $n_i$  and size  $m_i$ , then

(i) 
$$
M_1(\mathcal{G}) = \begin{cases} 4n_1 + 4m_1 + 9n_2 + 9n_3, & \text{if } G_{n_1} \text{ is triangle-free} \\ 9(n_1 + n_2 + n_3 + m_1), & \text{otherwise} \end{cases}
$$
  
(ii)  $M_2(\mathcal{G}) = \begin{cases} 4m_1 + 9(m_2 + m_3) + 2(n_1^2 - n_1) + 6(n_1n_2 + m_1n_3), & \text{if } G_{n_1} \text{ is triangle-free} \\ 9(n_1 + m_2 + m_3 + n_1n_2 + m_1n_3 + \frac{n_1^2 - n_1}{2}), & \text{otherwise} \end{cases}$ 

.

 $\Box$ 

 $\Box$ 

*Proof.* (i) Let  $G = G_{n_1}^C \vee (G_{n_2}^V \cup G_{n_2}^E)$ , where  $G_{n_1}$  is triangle-free, then from (5) by using  $(10)$  we get

$$
M_1(\mathcal{G}) = \sum_{v \in V(G_{n_1})} 2^2 + \sum_{v \in I(G_{n_1})} 2^2 + \sum_{v \in V(G_{n_2})} 3^2 + \sum_{v \in V(G_{n_3})} 3^2
$$
  
=  $4n_1 + 4m_1 + 9n_2 + 9n_3$ .

Similarly when  $G_{n_1}$  is not a triangle-free graph we get,

 $M_1(\mathcal{G}) = 9(n_1 + n_2 + n_3 + m_1).$ 

(ii) Let  $G_{n_1}$  be a triangle-free graph. Then from (5) by using (10) we get

$$
M_2(\mathcal{G}) = \sum_{uv \in E(G_{n_2})} 3 \times 3 + \sum_{uv \in E(G_{n_3})} 3 \times 3 + \sum_{u \in V(G_{n_1}), \atop v \in V(G_{n_2})} 2 \times 3 + \sum_{\substack{u \in I(G_{n_1}), \atop v \in V(G_{n_3})}} 2 \times 3 + \sum_{\substack{uv \in E(\mathcal{G}), \atop v \in V(G_{n_1}), v \in I(G_{n_1})}} 2 \times 2 + \sum_{\substack{uv \in E(\mathcal{G}), \atop u, v \in V(G_{n_1})}} 2 \times 2
$$
  
= 4m<sub>1</sub> + 9m<sub>2</sub> + 9m<sub>3</sub> + 2n<sub>1</sub><sup>2</sup> - 2n<sub>1</sub> + 6n<sub>1</sub>n<sub>2</sub> + 6m<sub>1</sub>n<sub>3</sub>.

Similarly when  $G_{n_1}$  is not a triangle-free graph we get,

$$
M_2(\mathcal{G}) = 9\left(m_1 + m_2 + m_3 + n_1n_2 + m_1n_3 + \frac{n_1^2 - n_1}{2}\right).
$$

 $\Box$ 

**Theorem 3.5.** For  $i = 1, 2, 3$ , let  $G_{n_i}$  be a graph of order  $n_i$  and size  $m_i$ , then

(i) 
$$
GA_4(\mathcal{G}) = \begin{cases} m_1 + m_2 + m_3 + \frac{2\sqrt{6}}{5}(n_1n_2 + m_1n_3) + \frac{n_1^2 - n_1}{2}, & \text{if } G_{n_1} \text{ is triangle-free} \\ m_1 + m_2 + m_3 + n_1n_2 + m_1n_3 + \frac{n_1^2 - n_1}{2}, & \text{otherwise} \end{cases}
$$

(ii) 
$$
ABC_5(\mathcal{G}) = \begin{cases} \frac{2}{3}(m_2 + m_3) + \frac{1}{\sqrt{2}} \left( n_1 n_2 + m_1 n_3 + m_1 + \frac{n_1(n_1 - 1)}{2} \right), & \text{if } G_{n_1} \text{ is triangle-free} \\ \frac{2}{3}(m_1 + m_2 + m_3 + n_1 n_2 + m_1 n_3) + \frac{n_1(n_1 - 1)}{3}, & \text{otherwise} \end{cases}
$$

*Proof.* (i) Let  $\mathcal{G} = G_{n_1}^C \vee (G_{n_2}^V \cup G_{n_2}^E)$ , where  $G_{n_1}$  is triangle-free, then from (6) by using (10) we have

$$
GA_4(\mathcal{G}) = \sum_{uv \in E(G_{n_2})} 2\frac{\sqrt{3 \times 3}}{3+3} + \sum_{uv \in E(G_{n_3})} 2\frac{\sqrt{3 \times 3}}{3+3} + \sum_{u \in V(G_{n_1}), \atop v \in V(G_{n_2})} 2\frac{\sqrt{2 \times 3}}{2+3} + \sum_{u \in V(G_{n_1}) \atop v \in V(G_{n_2})} 2\frac{\sqrt{2 \times 3}}{2+3} + \sum_{u \in V(G_{n_1}) \atop v \in U(G_{n_1})} 2\frac{\sqrt{2 \times 2}}{2+2} + \sum_{uv \in E(\mathcal{G}), \atop uv \in E(\mathcal{G})} 2\frac{\sqrt{2 \times 2}}{2+2} + \sum_{uv \in E(G, n_1) \atop uv \in E(\mathcal{G})} 2\frac{\sqrt{2 \times 2}}{2+2} + \sum_{uv \in E(G, n_1) \atop uv \in E(\mathcal{G})} 2\frac{\sqrt{2 \times 2}}{2+2} + \sum_{uv \in E(G, n_1) \atop uv \in E(\mathcal{G})} 2\frac{\sqrt{2 \times 3}}{2+2} + \sum_{uv \in E(G, n_1) \atop uv \in E(\mathcal{G})} 2\frac{\sqrt{2 \times 3}}{2+2} + \sum_{uv \in E(G, n_1) \atop v \in V(G_{n_1})} 2\frac{\sqrt{2 \times 3}}{2+3} + \sum_{uv \in V(G, n_2) \atop v \in V(G, n_2)} 2\frac{\sqrt{2 \times 3}}{2+3} + \sum_{uv \in V(G, n_1) \atop v \in V(G, n_1)} 2\frac{\sqrt{2 \times 3}}{2+3} + \sum_{uv \in E(G, n_1) \atop uv \in E(\mathcal{G})} 2\frac{\sqrt{2 \times 3}}{2+3} + \sum_{uv \in E(G, n_1) \atop uv \in E(\mathcal{G})} 2\frac{\sqrt{2 \times 3}}{2+3} + \sum_{uv \in E(G, n_1) \atop uv \in E(\mathcal{G})} 2\frac{\sqrt{2 \times 3}}{2+3} + \sum_{uv \in E(G, n_1) \atop uv \in E(\mathcal{G})} 2\frac{\sqrt{2 \times 3}}{
$$

Similarly when  $G_{n_1}$  is not a triangle-free graph we get,

$$
GA_4(\mathcal{G}) = m_1 + m_2 + m_3 + n_1n_2 + m_1n_3 + \frac{n_1^2 - n_1}{2}.
$$

(ii) Let  $G_{n_1}$  be a triangle-free graph. Then from (7) by using (10) we get

$$
ABC_{5}(\mathcal{G}) = \sum_{uv \in E(G_{n_{2}})} \sqrt{\frac{3+3-2}{3\times3}} + \sum_{uv \in E(G_{n_{3}})} \sqrt{\frac{3+3-2}{3\times3}} + \sum_{u \in V(G_{n_{1}})} \sqrt{\frac{2+3-2}{2\times3}} + \sum_{v \in V(G_{n_{2}})} \sqrt{\frac{2+3-2}{2\times3}} + \sum_{\substack{uv \in E(\mathcal{G}) \\ v \in V(G_{n_{3}})}} \sqrt{\frac{2+3-2}{2\times3}} + \sum_{\substack{uv \in E(\mathcal{G}) \\ u \in V(G_{n_{1}}), \ v \in I(G_{n_{1}})}} \sqrt{\frac{2+2-2}{2\times2}} + \sum_{\substack{uv \in E(\mathcal{G}) \\ u, v \in V(G_{n_{1}})}} \sqrt{\frac{2+2-2}{2\times2}} + \sum_{\substack{uv \in V(G_{n_{1}}) \\ u, v \in V(G_{n_{1}})}} \sqrt{\frac{2+2-2}{2\times2}} + \sum_{\substack{uv \in V(G_{n_{1}}) \\ u, v \in V(G_{n_{1}})}} \sqrt{\frac{2+2-2}{2\times2}} + \sum_{\substack{uv \in V(G_{n_{1}}) \\ u, v \in V(G_{n_{1}})}} \sqrt{\frac{2+2-2}{2\times2}} + \sum_{\substack{uv \in E(\mathcal{G}) \\ v \in I(G_{n_{1}})}} \sqrt{\frac{2+2-2}{2\times2}} + \sum_{\substack{uv \in V(G_{n_{1}}) \\ v \in V(G_{n_{1}})}} \sqrt{\frac{2+3-2}{2\times2}} + \sum_{\substack{uv \in V(G_{n_{1}}) \\ v \in I(G_{n_{1}})}} \sqrt{\frac{2+3-2}{2\times2}} + \sum_{\substack{uv \in V(G_{n_{1}}) \\ v \in V(G_{n_{1}})}} \sqrt{\frac{2+2-2}{2\times2}} + \sum_{\substack{uv \in V(G_{n_{1}}) \\ v \in I(G_{n_{1}})}} \sqrt{\frac{2+2-2}{2\times2}} + \sum_{\substack{uv \in V(G_{n_{1}}) \\ v \in V(G_{n_{1}})}} \sqrt{\frac{2+2-2}{2\times2}} + \sum_{\substack{uv \in V(G_{n_{1}}) \\ v \in I(G_{n_{1}})}} \sqrt{\frac{2+2-2}{2\times
$$

Similarly when  $G_{n_1}$  is not a triangle-free graph we get,

$$
ABC_5(\mathcal{G}) = \frac{2}{3}(m_1 + m_2 + m_3 + n_1n_2 + m_1n_3) + \frac{n_1(n_1 - 1)}{3}.
$$

 $\Box$ 

# 4 Conclusion

Topological indices play an important role in mathematical chemistry. In QSAR/QSPR study, topological indices can be used to estimate the bioactivity of chemical compounds. In modeling the chemical and other properties of molecules, we can use topological indices as a tool. This article gives the results related to topological indices like the eccentric connectivity index, connective eccentricity index, total- eccentricity index, average eccentricity index, Zagreb indices, geometric-arithmetic index, atom-bond connectivity index, and Wiener index of the newly defined graph operation called central vertex-edge join (CVE-join). Also, the Dspectrum of CVE-join of three regular graphs  $G_{n_1}$ ,  $G_{n_2}$  and  $G_{n_3}$  when  $G_{n_1}$  is a triangle-free are obtained. These results enable us to construct new D-equienergetic graph families.

### Acknowledgements

The authors would like to thank the DST, Government of India, for providing support to carry out this work under the scheme 'FIST'(No.SR /FST/MS-I/2019/40).

#### <span id="page-9-4"></span>References

- <span id="page-9-2"></span>[1] J. V. Vivin, M. M. Akbar Ali and K. Thilagavathi, On harmonious coloring of central graphs, Advances and applications in discrete mathematics, 2(1), 17—33, 2008.
- [2] M. Aouchiche and P. Hansen, Distance spectra of graphs: A survey, Linear algebra and its applications, 458, 301—386, 2014.
- <span id="page-9-1"></span><span id="page-9-0"></span>[3] A. E. Brouwer and W. H. Haemers, Spectra of graphs, Springer: New york, USA, 2012.
- [4] F. Buckley and F. Harary, Distance in graphs, volume 2, Addison-Wesley Redwood City, CA, 1990.
- <span id="page-9-3"></span>[5] G. Indulal, C. S. Deena and X. Liuc, The distance spectrum of the subdivision vertex join and subdivision edge join of two regular graphs, Discrete Math. Lett, 1, 36-–41, 2019.
- <span id="page-9-5"></span>[6] D. M. Cvetković, M. Doob and H. Sachs, Spectra of graphs- Theory and applications,  $3^{rd}$  edition, Johann Ambrosius Barth Verlag, Heidelberg, Leipzig, 1995.
- <span id="page-10-4"></span><span id="page-10-3"></span>[7] G. Indulal, I. Gutman and A. Vijayakumar, On distance energy of graphs, MATCH Commun. Math. Comput. Chem., 60(2), 461—472, 2008.
- <span id="page-10-5"></span>[8] H. S. Ramane, I. Gutman and D. S. Revankar, Distance equienergetic graphs, MATCH Commun. Math. Comput. Chem. , 60, 473-–484, 2008.
- <span id="page-10-2"></span>[9] A. Ili´c, Distance spectra and distance energy of integral circulant graphs, Linear algebra and its applications, 433(5), 1005—1014, 2010.
- <span id="page-10-0"></span>[10] T. K. Jahfar and A. V. Chithra, Central vertex join and central edge join of two graphs, AIMS Mathematics, 5(6), 7214—7233, 2020.
- [11] G. Aalipour, A. Abiad, Z. Berikkyzy, J. Cummings, D. S. Jessica, W. Gao, K. Heysse, L. Hogben, F. H. J. Kenter, M. Tait, Jephian C. H. Lin, On the distance spectra of graphs, Linear Algebra and its Applications, 497, 66–87, 2016.
- <span id="page-10-6"></span><span id="page-10-1"></span>[12] Haritha. T, and Chithra. A. V, On the distance spectra of central vertex join and central edge join of two regular graphs, Ricerche di Matematica, 1-19, 2022.
- [13] V. Sharma, R. Goswami, and A. K. Madan, Eccentric Connectivity Index: A Novel Highly Discriminating Topological Descriptor for Structure-Property and Structure-Activity Studies, J. Chem. Inf. Comput. Sci., 37, 2, 273–282, 1997.
- <span id="page-10-13"></span>[14] H. Wiener, Structural determination of paraffin boiling points, Journal of the American chemical society, 69(1):17-20, 1947.
- <span id="page-10-7"></span>[15] S. Gupta, M. Singh, and A. K. Madan, Connective eccentricity index: A novel topological descriptor for predicting biological activity, J. Mol. Graph. Model., 18, 1, 18–25, 2000.
- <span id="page-10-8"></span>[16] A. R. Ashrafi, M. Ghorbani, and M. A. Hossein-Zadeh, The eccentric connectivity polynomial of some graph operations, Serdica J. Comput., 5, 2, 101–116, 2011.
- <span id="page-10-10"></span><span id="page-10-9"></span>[17] V. A. Skorobogatov and A. A. Dobrynin, Metric analysis of graphs, MATCH Commun. Math. Comput. Chem., 23, 1, 105–151, 1988.
- <span id="page-10-11"></span>[18] M. Ghorbani and M. A. Hosseinzadeh, A new version of Zagreb indices, Filomat, 26, 93–100, 2012.
- <span id="page-10-12"></span>[19] M. Ghorbani and A. Khaki, A note on the fourth version of geometric-arithmetic index, Optoelectron. Adv. Mat., 4, 12, 2212–2215, 2010.
- [20] M. R. Farahani, Eccentricity version of atom-bond connectivity index of benzenoid family  $ABC_5(H_k)$ , World Appl. Sci. J. Chem., 21, 9, 1260–1265, 2013.