SEMI-COARSE SPACES, HOMOTOPY AND HOMOLOGY

ANTONIO RIESER AND JONATHAN TREVIÑO-MARROQUÍN

ABSTRACT. We begin the study the algebraic topology of semi-coarse spaces, which are generalizations of coarse spaces that enable one to endow non-trivial 'coarse-like' structures to compact metric spaces, something which is impossible in coarse geometry. We first study homotopy in this context, and we then construct homology groups which are invariant under semi-coarse homotopy equivalence. We further show that any undirected graph $G = (V, E)$ induces a semi-coarse structure on its set of vertices V_G , and that the respective semicoarse homology is isomorphic to the Vietoris-Rips homology. This, in turn, leads to a homotopy invariance theorem for the Vietoris-Rips homology of undirected graphs.

CONTENTS

This material is based in part upon work supported by the US National Science Foundation under Grant No. DMS-1928930 while the authors participated in a program supported by the Mathematical Sciences Research Institute. The program was held in the summer of 2022 in partnership with the the Universidad Nacional Autónoma de México. This work was also supported by the CONACYT Investigadoras y Investigadores por México Project #1076, the CONACYT Ciencia de Fronteras grant CF-2019-217392, and by the grant N62909-19-1-2134 from the US Office of Naval Research Global and the Southern Office of Aerospace Research and Development of the US Air Force Office of Scientific Research. The second author was also supported by the CONACYT postgraduate studies scholarship number 839062.

1. INTRODUCTION

Coarse geometry [\[14\]](#page-55-2) is often referred to as 'geometry in the large', or 'large scale geometry', for the reason that, on a metric space endowed with the natural coarse structure, all bounded phenomena are trivial from the coarse point of view. The study of coarse geometry has led to a number of important advances, in particular in group theory, where it has enabled many questions about groups to be addressed geometrically via the analysis of their Cayley graphs (see, for instance, [\[6\]](#page-55-3) for a recent book-length discussion of this approach). Nonetheless, the large-scale nature of coarse geometry forces all finite-diameter spaces to be coarsely-equivalent to a point. It is not difficult, however, to imagine scenarios where one might like to 'coarsen' a space up to a certain scale, but where one does not want to erase all bounded phenomena at every scale. Many problems of topological data analysis, for instance, in which one would like to deduce the topological invariants of a space from algebraic invariants built from a finite subset of the space, present themselves naturally as problems of 'medium-scale' coarsening. Several possibilities for a formal context for 'coarsening' a space up to a scale were developed by the first author in $[12]$ and $[13]$, where he studied the problem from the point of view of Cech closure spaces and semi-uniform spaces, respectively. The connection to coarse geometry in these earlier works, however, remained unclear, and this question forms the primary motivation for the work presented here.

Whereas the closure structures and semi-uniform structures studied in [\[12\]](#page-55-4) and [\[13\]](#page-55-5) are generalizations of topological structures and uniform structures, respectively, in this article, we focus instead on a generalization of coarse structures, modifying the axioms of coarse geometry [\[14\]](#page-55-2) to allow 'coarsenings' only up to a preferred size. Doing so links the homotopy of spaces coarsened up to a fixed scale to the study of coarse spaces, and, in particular, the development of homotopy on semi-coarse spaces that we have begun here has been used by the second author in [\[17\]](#page-55-6) to construct new homotopical invariants on coarse spaces themselves, something which is unclear how to accomplish through either closure or semi-uniform spaces directly. Technically, this 'coarsening up to a scale' is achieved by eliminating the product axiom in the definition of the coarse structure, which, indeed, has a similar flavor to the elimination of the idempotence axiom in the passage from Kurotowski (topological) closure structures to more general Cech closure structures, or to the elimination of the product axiom in the passage from uniform spaces to semi-uniform spaces. While removing the product axiom effectively destroys the notion of coarse equivalence, we observe in the following that coarse equivalence can be profitably generalized to a useful notion of homotopy equivalence in this setting. The resulting structures and spaces are called semi-coarse structures and semi-coarse spaces, respectively, and were first introduced in [\[18\]](#page-55-7). Many examples of semi-coarse spaces exist. The examples of most interest to topological data analysis may be built from pseudo-metric spaces and a preferred scale $r > 0$, but many others exist as well, in particular those constructed from semi-uniform spaces. Furthermore, when specialized to graphs on lattices, the semi-coarse homotopy introduced here is also similar to the digital homotopy studied in [\[2–](#page-55-8)[4,](#page-55-9) [7–](#page-55-10)[11,](#page-55-11) [15\]](#page-55-12).

The outline of this article is as follows. In Section [2,](#page-2-0) we give the basic definitions, point-set properties, and principal examples of semi-coarse spaces. We will also show in that section how to obtain a coarse space from a semi-coarse space through a certain limiting process. In Section [3,](#page-13-0) we begin the study of homotopy invariants in the semi-coarse category. Unfortunately, the interval does not appear to have a natural non-trivial semi-coarse structure, significantly complicating the construction. We are able circumvent this shortcoming by adapting the homotopy construction from [\[1\]](#page-55-13), using finite-length subsequences of $\mathbb Z$ in place of the interval to construct cylinders. While technically delicate, this nonetheless allows us to define homotopy groups, prove a long-exact sequence of pairs, and compute the fundamental group of a 'semi-coarse circle' with four points. Finally, in Section [4,](#page-35-0) we construct homology groups for semi-coarse spaces, inspired by the Vietoris-Rips construction now commonly used in topological data analysis. We give a construction for any semi-coarse space, demonstrate the invariance of the homology groups with respect to the homotopy introduced in Section [3,](#page-13-0) and show that, for a countable semi-coarse space, this homology is exactly the Vietoris-Rips homology of an associated graph.

Many of the results in this paper were first presented in the Master's thesis of the second author [\[16\]](#page-55-14), written under the supervision of the first author, where semi-coarse spaces and structures were called pseudo-coarse.

2. Semi-Coarse Spaces

In this section, we define semi-coarse spaces, bornologous functions, and we give examples of semi-coarse structures constructed from a metric space and a scale parameter $r > 0$. We then define semi-coarse quotients, disjoint unions, and products, and we show how to build an induced coarse structure from a semi-coarse structure.

2.1. Fundamental Concepts and Examples. We begin by setting some notation which we will use throughout the article.

Definition 2.1.1. Let X be a set. We denote by $\mathcal{P}(X)$ the collection of all subsets of X , and

$$
n
$$
-times

- (1) X^n will denote $\overline{X \times X} \times \cdots \times \overline{X}$.
- (2) $\Delta_X := \{(x, x) \in X \times X \mid x \in X\}$ will be called the diagonal of X.
- (3) For $V \in \mathcal{P}(X \times X)$, we define

$$
V^{-1} := \{ (y, x) \in X \times X : (x, y) \in V \},\
$$

which we call the inverse of V .

(4) For $V, W \in \mathcal{P}(X \times X)$, we define

$$
V \circ W := \{(x, y) \in X \times X \mid \exists z \in X, (x, z) \in V \text{ and } (z, y) \in W\},\
$$

which will be called the set product of V and W.

n−times

Given an integer $n \geq 2$, we write $V^{\circ n}$ for $\widetilde{V} \circ \ldots \circ \widetilde{V}$.

(5) Let X and Y be sets, $f: X \to Y$ a set function, and $V \in \mathcal{P}(X \times X)$. Then

$$
(f \times f)(V) := \{ (f(x), f(x')) \in Y \times Y \mid (x, x') \in V \},
$$

and we call $(f \times f)(V)$ the image of V under $f \times f$.

We now define semi-coarse spaces, our principal object of interest.

Definition 2.1.2 (Semi-coarse space). Let X be a set, and let $\mathcal{V} \subset \mathcal{P}(X \times X)$ be a collection of subsets of $X \times X$ which satisfies

$$
(\mathrm{sc}1) \ \Delta_X \in \mathcal{V},
$$

(sc2) If $B \in \mathcal{V}$ and $A \subset B$, then $A \in \mathcal{V}$,

(sc3) If $A, B \in \mathcal{V}$, then $A \cup B \in \mathcal{V}$,

(sc4) If $A \in \mathcal{V}$, then $A^{-1} \in \mathcal{V}$.

We call V a semi-coarse structure on X, and we say that the pair (X, V) is a semi-coarse space.

If, in addition, $\mathcal V$ satisfies

(sc5) If $A, B \in \mathcal{V}$, then $A \circ B \in \mathcal{V}$.

then V will be called a *coarse structure*, and (X, V) will be called a *coarse space*, as in [\[14\]](#page-55-2).

The elements of V will be called *controlled sets*. Moreover, if there exist a, b such that $\{(a, b)\}\in \mathcal{V}$ we will say that a and b are adjacent. If V and V' are semi-coarse structures on X such that $V \subset V'$, then we say that V' is finer than V and V is coarser than V' . Finally, when the structure V is unambiguous, we will sometimes refer to the semi-coarse space only by X .

The functions of interest between semi-coarse spaces will be those which preserve the semi-coarse structure.

Definition 2.1.3. We will say that $f : X \to Y$ is a (V, W) -bornologous function, or simply bornologous, if $f \times f$ maps each controlled set $V \in \mathcal{V}$ to a controlled set $(f \times f)(V) \in \mathcal{W}$.

Since the composition of set maps is associative, the composition of bornologous maps is bornologous, and the identity is bornologous for every semi-coarse space (X, V) , we have

Theorem 2.1.4. Semi-coarse spaces and bornologous functions form a category.

We denote the category of semi-coarse spaces and bornologous functions by SCoarse.

For our first collection of examples, we show how undirected graphs induce semicoarse structures. We begin with a pair of lemmas.

Lemma 2.1.5. Let X be a set, and let $W \subset X \times X$ such that

(1) $\Delta_X \subset W$, (2) $W = W^{-1}$.

Then $(X, \mathcal{P}(W))$ is a semi-coarse space.

Proof. We verify directly that the axioms $(\text{sc}1)$ - $(\text{sc}4)$ from Definition [2.1.2](#page-3-2) are satisfied.

- (sc1) $\Delta_X \subset W$, so $\Delta_X \in \mathcal{P}(W)$.
- (sc2) $\mathcal{P}(W)$ is closed under taking subsets by definition.
- (sc3) Since any two sets $A, B \in \mathcal{P}(W)$ are subsets of W, the union $A \cup B \subset W$, and therefore $A \cup B \in \mathcal{P}(W)$.
- (sc4) Suppose $A \in \mathcal{P}(W)$. Then $A \subset W$. However, by hypothesis on W, if $(a, b) \in W$ then $(b, a) \in W$. Therefore $A^{-1} \subset W$, so $A^{-1} \in \mathcal{P}(W)$ as well.

It follows from the above that $(X, \mathcal{P}(W))$ is a semi-coarse space, as desired. \Box

It will sometimes be convenient to use this lemma in the following alternative form.

Lemma 2.1.6. Let X be a set and suppose that $\mathcal{U} \subset \mathcal{P}(X \times X)$ is a collection of subsets of $X \times X$ such that

(1)
$$
U \in U \implies U^{-1} \in U
$$
, and
(2) $\Delta_X \in U$.

Let $W := \bigcup_{U \in \mathcal{U}} U$. Then $(X, \mathcal{P}(W))$ is a semi-coarse space.

Proof. By construction, W satisfies the hypotheses of Lemma [2.1.5.](#page-3-3) The conclusion follows. \Box

We now use these lemmas to give several important examples of semi-coarse spaces.

Example 2.1.6.1. Let $G = (V, E)$ be an undirected graph (i.e. $(u, v) \in E \iff$ $(v, u) \in E$). Define $V \subset \mathcal{P}(V \times V)$ to be

$$
\mathcal{V}_G := \mathcal{P}(E \cup \Delta_V),
$$

where Δ_V is the diagonal of $V \times V$. By Lemma [2.1.6](#page-4-0) above, the pair (V, V_G) is a semi-coarse space.

Definition 2.1.7. Given a graph $G = (V, E)$, we say that the semi-coarse space (V, V_G) constructed in Example [2.1.6.1](#page-4-1) is the semi-coarse space generated by the graph G.

Another important class of examples may be constructed from metric spaces combined with a positive scale parameter $r > 0$.

Example 2.1.7.1. (a) Let (X, d) be a metric space. Let $r > 0$ be a positive real number and define

$$
U_r := \{ (x, x') \in X \times X \mid d(x, x') \le r \},\
$$

and let $\mathcal{V}_r := \mathcal{P}(U_r)$. Then (X, \mathcal{V}_r) is a semi-coarse space by Lemma [2.1.5.](#page-3-3) (b) Similarly, defining U_r^{\lt} by

$$
U_r^<\coloneqq\{(x,x')\in X\times X\mid d(x,x')
$$

and let $\mathcal{V}_r^{\leq} := \mathcal{P}(U_r^{\leq})$. Lemma [2.1.5](#page-3-3) gives that $(X, \mathcal{V}_r^{\leq})$ is a semi-coarse space.

For the next example, which generalizes the ones in Example [2.1.7.1](#page-4-2) above, we introduce semi-pseudometric spaces.

Definition 2.1.8 (Semi-Pseudometric; [\[5\]](#page-55-15), 18 A.1.). Let X be a set and $d: X \times$ $X \to \mathbb{R}$ be a function, we will say d is a semi-pseudometric on X if they satisfies the next conditions

(m1) For each $x \in X$, $d(x, x) = 0$.

(m2) For every $x, y \in X$, $d(x, y) = d(y, x) \geq 0$.

A semi-pseudometric on X is a *pseudometric on* X if also

(m3) For each $x, y, z \in X$, $d(x, z) \leq d(x, y) + d(y, z)$, i.e. d satisfies the triangle inequality.

A semi-pseudometric will be called a semi-metric, if it also satisfies

 $(m4)$ $d(x, y) = 0$ implies $x = y$.

Finally, a semi-pseudometric d will be called a *metric* iff it satisfies $(m1)$ - $(m4)$, that is, iff d is both a semi-metric and a pseudometric.

A semi-pseudometric space is an ordered pair (X, d) where X is a set and d is a semi-pseudometric on X. Similarly, (X, d) is a semi-metric, pseudometric, or metric space when d is a semi-metric, pseudometric, or metric, respectively.

We will also need the following semi-pseudometrics constructed from a metric and a pre-determined scale $r > 0$.

Definition 2.1.9. Let (X, d) be a metric space and let $r > 0$ be a non-negative real number. We will define the functions $d_r, d_r \leq d_{\overline{r}} \colon X \times X \to [0, \infty)$ by

$$
d_r^{\leq}(x, y) = \begin{cases} 0 & \text{if } d(x, y) \leq r, \\ 1 & \text{if } d(x, y) > r, \end{cases}
$$

$$
d_r^{\leq}(x, y) = \begin{cases} 0 & \text{if } d(x, y) < r, \\ 1 & \text{if } d(x, y) \geq r. \end{cases}
$$

$$
d_r(x, y) = \max\{0, d(x, y) - r\}
$$

Remark 2.1.10. For $r \geq 0$, the function d_r satisfies $d_r(x, x) = 0$ and $d_r(x, y) = 0$ $d_r(y, x)$, by the symmetry of d. Therefore, d_r is a semi-pseudometric on X for any $r \geq 0$. Similarly, the functions d_r^{\le} and d_r^{\le} are semi-pseudometrics on X for $r \leq 0$ and $r > 0$, respectively.

Example 2.1.10.1. Let $d: X \times X \rightarrow [0, \infty)$ be a semi-pseudometric, and define $U := \{(x, x') \in X \times X \mid d(x, x') = 0\}.$ Then $(X, \mathcal{P}(U))$ is a semi-coarse space by Lemma [2.1.5.](#page-3-3) The semi-pseudometrics $d_{\overline{r}} \leq$ and $d_{\overline{r}} \leq$ from Definition [2.1.9](#page-5-1) give the semi-coarse structures in Examples [2.1.7.1](#page-4-2)[\(a\)](#page-4-4) and 2.1.7.1[\(b\),](#page-4-5) respectively.

The following examples are independent of the Lemmas [2.1.5](#page-3-3) and [2.1.6,](#page-4-0) and give examples of semi-coarse spaces which are not coarse.

Example 2.1.10.2. (a) Let (X, d) be a metric space which is at least countably infinite. Define $\mathcal{V} \subset \mathcal{P}(X \times X)$ by

 $V \in \mathcal{V} \iff \{ |V| < \infty \mid \forall (x, x') \in V, d(x, x') < 1 \}.$

Then (X, V) is a semi-coarse space. If $X = \mathbb{Q}$ with the Euclidean metric, then $(\mathbb{Q}, \mathcal{V})$ is a semi-coarse space which is not coarse.

(b) Let (X, c) be an uncountable metric space. Define $\mathcal{V} \subset \mathcal{P}(X \times X)$ by

$$
V \in \mathcal{V} \iff \{ |V| \le \aleph_0 \mid \forall (x, x') \in V, \ d(x, x') < 1 \}.
$$

Then (X, V) is a semi-coarse space. If $X = \mathbb{R}$ with the Euclidean metric, then $(\mathbb{R}, \mathcal{V})$ is a semi-coarse space which is not coarse.

2.2. Subspaces. We will now proceed to build new semi-coarse spaces from existing ones, in particular, constructing subspaces, products, and quotient spaces in the semi-coarse category. Our first step, in this section, will be to construct semi-coarse subspaces of a semi-coarse space (X, V) .

We first require the following lemma.

Lemma 2.2.1. Let X be a set, and let $\{A_{\lambda}\}_{{\lambda \in \Lambda}}$ a collection of elements of $\mathcal{P}(X \times$ X), where Λ is an index set. Then

 (i) U $\bigcup_{\lambda \in \Lambda} (A_{\lambda})^{-1} = \left(\bigcup_{\lambda \in \Lambda}$ $\bigcup_{\lambda \in \Lambda} A_{\lambda} \bigg)^{-1},$ $(ii) \cap$ $\bigcap_{\lambda \in \Lambda} (A_{\lambda})^{-1} = \left(\bigcap_{\lambda \in \Lambda}$ $\bigcap_{\lambda \in \Lambda} A_{\lambda} \bigg)^{-1}.$

Proof. Let X be a set, let Λ be a index set, and let $\{A_{\lambda}\}_{\lambda \in \Lambda}$ be a collection of elements of $\mathcal{P}(X \times X)$.

(i) We see that

$$
(x, y) \in \bigcup_{\lambda \in \Lambda} (A_{\lambda})^{-1} \iff \exists \lambda_0 \in \Lambda \text{ such that } (x, y) \in (A_{\lambda_0})^{-1} \iff (y, x) \in A_{\lambda_0}
$$

$$
\iff (y, x) \in \bigcup_{\lambda \in \Lambda} A_{\lambda} \iff (x, y) \in \left(\bigcup_{\lambda \in \Lambda} A_{\lambda}\right)^{-1}.
$$

Therefore \bigcup $\bigcup_{\lambda \in \Lambda} (A_{\lambda})^{-1} = \left(\bigcup_{\lambda \in \Lambda}$ $\bigcup_{\lambda \in \Lambda} A_\lambda \bigg)^{-1}$ (ii) Now note that

$$
(x, y) \in \bigcap_{\lambda \in \Lambda} (A_{\lambda})^{-1} \iff \forall \lambda \in \Lambda, (x, y) \in (A_{\lambda})^{-1} \iff (y, x) \in A_{\lambda} \forall \lambda \in \Lambda
$$

$$
\iff (y, x) \in \bigcap_{\lambda \in \Lambda} A_{\lambda} \iff (x, y) \in \left(\bigcap_{\lambda \in \Lambda} A_{\lambda}\right)^{-1}.
$$

Therefore, $\bigcap (A_{\lambda})^{-1} = \left(\bigcap A_{\lambda}\right)^{-1}$ as desired.

The λ∈Λ (A_λ) $^{-1}$ = ∩ $\bigcap_{\lambda \in \Lambda} A_{\lambda}$

Proposition 2.2.2. Let (X, V) be a semi-coarse space and suppose that $Y \subset X$. Define the collection $\mathcal{V}_Y \in \mathcal{P}(Y \times Y)$ by

$$
\mathcal{V}_Y \coloneqq \{ V \cap (Y \times Y) \mid V \in \mathcal{V} \}
$$

Then the pair (Y, V_Y) is a semi-coarse space.

Proof. Let (X, V) , $Y \subset X$, and V_Y be as in the statement of the proposition. We check that the axioms for a semi-coarse space are satisfied by (Y, V_Y) .

(sc1) We observe that $\Delta_Y = \Delta_X \cap (Y \times Y)$. Therefore $\Delta_Y \in V_Y$.

(sc2) Let $A \in V_Y$ and suppose that $A' \subset A$. By definition, there is a $B \in V$ such that $A = B \cap (Y \times Y)$. We define $B' := A' \cap B$ and we observe that $B' \in V$ since $B' \subset B \in \mathcal{V}$. However, $B' \cap (Y \times Y) = A' \cap B \cap (Y \times Y) = A' \cap A = A'$, so $A' \in \mathcal{V}_Y$.

(sc3) If $A, B \in V_Y$, then there are $A', B' \in V$ such that $A = A' \cap (Y \times Y)$ and $B = B' \cap (Y \times Y)$. Therefore

$$
A \cup B = (A' \cap (Y \times Y)) \cup (B' \cap Y \times Y)
$$

= $(A' \cup B') \cap (Y \times Y)$

Since $A' \cup B' \in V$, it follows that $A \cup B \in V_Y$.

(sc4) If $A \in \mathcal{V}_Y$, then there is $A' \in \mathcal{V}_Y$ such that $A = A' \cap (Y \times Y)$, then by [Lemma 2.2.1](#page-6-1) we get $A^{-1} = (A \cap (Y \times Y)^{-1} = A'^{-1} \cap (Y \times Y)^{-1} = A'^{-1} \cap (Y \times Y)$. Therefore $A^{-1} \in V_Y$.

It follows that (Y, V_Y) is a semi-coarse space, as desired. \Box

Definition 2.2.3 (Semi-Coarse Subspace). Let (X, V) be a semi-coarse space and let $Y \subset X$. The ordered pair (Y, V_Y) from [Proposition 2.2.2](#page-6-2) will be called a semicoarse subspace of X. When the structure \mathcal{V}_Y is clear from the context, we will simply refer to the subspace (Y, V_Y) as Y.

The following proposition gives a useful criterion for checking whether a function is bornologous on a semi-coarse space (X, V) .

Proposition 2.2.4. Let (X, V) and (Y, W) be semi-coarse spaces, and suppose that $(X_i, \mathcal{V}_i) \subset (X, \mathcal{V}), i \in \{1, \ldots, n\},$ are subspaces of (X, \mathcal{V}) such that $\bigcup_{i=1}^n X_i = X$ and every set $V \in V$ may be written in the form

$$
V = \bigcup_{i=n}^{n} V_i
$$

where each $V_i \in \mathcal{V}_i$. Now suppose that $f: X \to Y$ is a map such that the restrictions $f|_{X_i}:(X_i,\mathcal{V}_i)\to(Y_i\mathcal{W})$ are bornologous for all $i\in\{1,\ldots,n\}$. Then $f:(X,\mathcal{V})\to Y_i$ (Y, W) is bornologous.

Proof. Let $V \in \mathcal{V}$. Then, by hypothesis, $V \in \mathcal{V}_i$ for some $i \in \{1, \ldots, n\}$, and since $f|_{X_i}$ is bornologous, we have that $(f \times f)(V) = (f|_{X_i} \times f|_{X_i})(V) \in W$. Since $V \in V$ is arbitrary, it follows that f is bornologous. \Box

2.3. Product Semi-Coarse Spaces. In this section, we construct product semicoarse structure on the product of sets. We begin with the following definition and several preliminary results.

Definition 2.3.1. Let X and Y be sets, and suppose that $V \in \mathcal{P}(X \times X)$ and $W \in \mathcal{P}(Y \times Y)$. We define

 $V \boxtimes W := \{((a, b), (c, d)) \in (X \times Y) \times (X \times Y) : (a, c) \in V, (b, d) \in W\}$

which we call the Cartesian cross product of V and W.

Lemma 2.3.2. The Cartesian cross product is associative.

Proof. Let X, Y, and Z be sets, and suppose that $U \in \mathcal{P}(X \times X)$, $V \in \mathcal{P}(Y \times Y)$, and $W \in \mathcal{P}(Z \times Z)$. We wish to show that $(V \boxtimes W) \boxtimes U \cong V \boxtimes (W \boxtimes U)$. By definition, we have that

$$
(U \boxtimes V) \boxtimes W = \{(((a, c), e), ((b, d), f)) \in ((X \times Y) \times Z) \times ((X \times Y) \times Z)
$$

$$
\mid (a, b) \in (X \times X), (c, d) \in (Y \times Y), \text{ and } (e, f) \in (Z \times Z) \}, \text{ and}
$$

$$
U \boxtimes (V \boxtimes W) = \{(((a, (c, e)), (b, (d, f))) \in (X \times (Y \times Z)) \times (X \times (Y \times Z))
$$

$$
\mid (a, b) \in (X \times X), (c, d) \in (Y \times Y), \text{ and } (e, f) \in (Z \times Z) \}.
$$

Since the sets $(X \times Y) \times Z \cong X \times (Y \times Z)$ are isomorphic, this proves the result. \Box

Lemma 2.3.3. Let X and Y be sets, $A \subset X \times X$, $B \subset Y \times Y$, $\{A_{\lambda}\}_{\lambda \in \Lambda}$ a collection of subsets of $X \times X$ indexed by Λ and $\{B_{\gamma}\}_{{\gamma} \in \Gamma}$ a collection of subsets of $Y \times Y$ indexed by Γ . Then

(i)
$$
(A \boxtimes B)^{-1} = A^{-1} \boxtimes B^{-1}
$$
.
\n(ii) $\bigcup_{\lambda \in \Lambda} (A_{\lambda} \boxtimes B) = \left(\bigcup_{\lambda \in \Lambda} A_{\lambda}\right) \boxtimes B$.
\n(iii) $\bigcup_{\gamma \in \Gamma} (A \boxtimes B_{\gamma}) = A \boxtimes \left(\bigcup_{\gamma \in \Gamma} B_{\gamma}\right)$.

Proof. Let X, Y, $A \subset X \times X$, $B \subset Y \times Y$, $\{A_{\lambda}\}_{\lambda \in \Lambda}$, and $\{B_{\gamma}\}_{\gamma \in \Gamma}$ be as in the statement of the proposition.

(i) We observe that

$$
(A \boxtimes B)^{-1} = \{ ((x', y'), (x, y)) : (x, x') \in A, (y, y') \in B \}
$$

=
$$
\{ ((x', y'), (x, y)) : (x', x) \in A^{-1}, (y', y) \in B^{-1} \}
$$

= $A^{-1} \boxtimes B^{-1}$.

(ii) If $((a, c), (b, d)) \in \bigcup$ $\bigcup_{\lambda \in \Lambda} (A_{\lambda} \boxtimes B)$, then there exists a $\lambda_0 \in \Lambda$ such that $((a, b), (c, d)) \in A_{\lambda_0} \boxtimes B$, so $(a, b) \in A_{\lambda_0}$ and $(c, d) \in B$. This implies that $(a, b) \in A_{\lambda_0}$ U $\bigcup_{\lambda \in \Lambda} A_{\lambda}$ and $(b, d) \in B$, from which we conclude that $((a, c), (b, d)) \in \left(\bigcup_{\lambda \in A} A_{\lambda}\right)$ $\bigcup_{\lambda \in \Lambda} A_{\lambda} \bigg) \boxtimes B.$

Conversely, if $((a, c), (b, d)) \in \left(\bigcup \right)$ $\bigcup_{\lambda \in \Lambda} A_{\lambda} \bigg) \boxtimes B$, then $(a, b) \in \bigcup_{\lambda \in A}$ $\bigcup_{\lambda \in \Lambda} A_{\lambda}$ and $(c, d) \in$ B. This implies that there is $\lambda_0 \in \Lambda$ such that $(a, b) \in A_{\lambda_0}$, which gives that $((a, c), (b, d)) \in A_{\lambda_0} \boxtimes B$. This implies that $((a, c), (b, d)) \in \bigcup$ $\bigcup_{\lambda \in \Lambda} (\mathring{A}_{\lambda} \boxtimes B)$, as desired.

(iii) The proof of this point is analogous to the proof of part [\(ii\)](#page-8-0) above. \Box

In the next proposition, we construct the product of two semi-coarse spaces.

Proposition 2.3.4. Let (X, Y) and (Y, W) be semi-coarse spaces, and let $V \times W$ be the collection of all subsets of finite unions of sets of the form $V \boxtimes W$, where $V \in \mathcal{V}$ and $W \in \mathcal{W}$, i.e.

$$
\mathcal{V} \times \mathcal{W} := \{ U \in \mathcal{P}((X \times Y) \times (X \times Y))
$$

\n
$$
| \exists n \in \mathbb{N} \text{ such that } U \subset \bigcup_{i=1}^{n} V_i \boxtimes W_i,
$$

\nwhere $V_i \in \mathcal{V}, W_i \in \mathcal{W} \forall i \in \{1, ..., n\} \}.$

Then $(X \times Y, V \times W)$ is a semi-coarse space.

Proof. Let (X, V) and (Y, W) be semi-coarse spaces and let the collection $V \times W$ be as in the statement of the proposition. We check that $(X \times Y, V \times W)$ satisfies the axioms for a semi-coarse space.

(sc1) We observe that $\Delta_X \boxtimes \Delta_Y = \{((x, y), (x, y)) \in (X \times Y) \times (X \times Y) \mid x \in$ $X, y \in Y$ } = $\Delta_{X \times Y}$.

(sc2) Let $A \in \mathcal{V} \times \mathcal{W}$ and $B \subset A$. Then there are $n \in \mathbb{N}$, $\{V_1, ..., V_n\} \subset \mathcal{V}$ and $\{W_1, ..., W_n\} \subset \mathcal{W}$ such that $B \subset A \subset \bigcup$ $\bigcup_{1\leq k\leq n} (V_k \boxtimes W_k)$, and therefore $B \in \mathcal{V} \times \mathcal{W}$.

(sc3) If $A, B \in V \times W$. Then there are $m, n \in \mathbb{N}, \{V_1, \ldots, V_m\}, \{V'_1, \ldots, V'_n\} \subset V$ and $\{W_1, \ldots, W_m\}$, $\{W'_1, \ldots, W'_n\} \subset \mathcal{W}$ such that

$$
A \subset \bigcup_{1 \leq k \leq m} (V_k \boxtimes W_k),
$$

$$
B \subset \bigcup_{1 \leq k \leq n} (V_k' \boxtimes W_k').
$$

It follows that $A \cup B \subset$ $\sqrt{2}$ U $\bigcup_{1\leq j\leq m}(V_j\boxtimes W_j)\right)$ U $\sqrt{2}$ U $1 \leq k \leq n$ $(V'_k \boxtimes W'_k$ \setminus , which implies that $A \cup B \in \mathcal{V} \times \mathcal{W}$.

(sc4) If $A \in \mathcal{V} \times \mathcal{W}$, then there is an $n \in \mathbb{N}$, $\{V_1, ..., V_n\} \subset \mathcal{V}$, and $\{W_1, ..., W_n\} \subset$ W such that $A \subset \bigcup$ $\bigcup_{1 \leq k \leq n} (V_k \boxtimes W_k)$, so $A^{-1} \subset \bigcup_{1 \leq k \leq n}$ $1 \leq k \leq n$ $(V_k^{-1} \boxtimes W_k^{-1})$ by [Lemma 2.2.1](#page-6-1) and [Lemma 2.3.3,](#page-8-1) and therefore $A^{-1} \in \mathcal{V} \times \mathcal{W}$

We conclude that $\mathcal{V} \times \mathcal{W}$ is a semi-coarse structure on the set $V \times W$.

Definition 2.3.5 (Product Semi-Coarse Space). Let (X, V) and (Y, W) be semicoarse spaces, and let $V \times W$ be the semi-coarse structure on $X \times Y$ constructed in [Proposition 2.3.4](#page-8-2) We call the ordered pair $(X \times Y, \mathcal{V} \times \mathcal{W})$ the **product semi-coarse** space of (X, V) and (Y, W) .

It is important to observe that this product is associative, which is established in the next proposition.

Proposition 2.3.6. Let (X, V) , (Y, W) and (Z, Z) be semi-coarse spaces. Then $(\mathcal{V} \times \mathcal{W}) \times \mathcal{Z} = \mathcal{V} \times (\mathcal{W} \times \mathcal{Z}).$

Proof. Let (X, V) , (Y, W) and (Z, Z) be semi-coarse spaces. Consider $A \in (V \times V)$ $W \times \mathcal{Z}$. Then there is a natural number *n* and sets $\{D_k\}_{k=1}^n \subset \mathcal{V} \times \mathcal{W}$ and ${E_k}_{k=1}^n \subset \mathcal{Z}$ such that $A \subset \bigcup^n$ $\bigcup_{k=1}^{n} (D_k \boxtimes E_k)$. It follows that, for each $k \in \{1, \ldots, n\}$, there are natural numbers $\{m_k\}_{k=1}^n \subset \mathbb{N}$ and sets $\{B_i\}_{i\in I}^{m_k} \subset \mathcal{V}$ and $\{C_i\}_{i\in I}^{m_k} \subset \mathcal{W}$ such that $D_k \subset \bigcup^{m_k}$ $\bigcup_{i=1}^{m_k} (B_i \boxtimes C_i)$. This gives that $A \subset \bigcup_{k=1}^{n}$ $k=1$ \bigcup^m $\bigcup_{i=1}^{n} (B_i \boxtimes C_i \boxtimes E_k)$, by [Lemma 2.3.2](#page-7-1) and [Lemma 2.3.3.](#page-8-1) Since this is a finite union, and, moreover, since $(C_i \boxtimes E_k) \in W \times \mathcal{Z}$ and $B_i \in V$ for each $i \in \{1, \dots, m_k\}, k \in \{1, \dots, n\}$, we have that $A \in \mathcal{V} \times (\mathcal{W} \times \mathcal{Z})$.

The proof that $(\mathcal{V} \times \mathcal{W}) \times \mathcal{Z} \supset \mathcal{V} \times (\mathcal{W} \times \mathcal{Z})$ is analogous. Thus, the two structures are identical. \Box

The following alternative characterization of the product structure will be useful for characterizing bornologous functions.

Proposition 2.3.7. Let (X, V) and (Y, W) be semi-coarse spaces. The product structure $V \times W$ is the collection of subsets of sets of the form $V \boxtimes W$, where $V \in V$ and $W \in \mathcal{W}$.

Proof. Let (X, V) and (Y, W) be semi-coarse spaces, and let $V \boxtimes W$ denote the collection of subsets of sets of the form $V \boxtimes W$, where $V \in V$, $W \in W$.

By definition, it is clear that $V \boxtimes W \subset V \times W$. On the other hand, if $A \in$ $V \times W$, then there is a natural number $n \in \mathbb{N}$ and sets $\{V_1, \dots, V_n\} \in V$ and

$$
\{W_1, \dots, W_n\} \in \mathcal{W} \text{ such that } A = \bigcup_{k=1}^n (V_k \boxtimes W_k). \text{ However, we also have that}
$$

$$
\bigcup_{k=1}^n V_k \in \mathcal{V}, \bigcup_{k=1}^n W_k \in \mathcal{W}, \text{ and}
$$

$$
A \subset \bigcup_{k=1}^n (V_k \boxtimes W_k) \subset \left(\bigcup_{k=1}^n V_k\right) \boxtimes \left(\bigcup_{k=1}^n W_k\right).
$$

Therefore $A \in V \boxtimes W$, and the result follows. \Box

The following corollaries are now an immediate consequence of the above proposition.

Corollary 2.3.8. Let (X, V) , (Y, W) , and (Z, A) be semi-coarse spaces. Then $f: (X \times Y, V \times W) \rightarrow (Z, \mathcal{A})$ is a bornologous function iff $f(B \boxtimes C) \in \mathcal{A}$ for each $B \in \mathcal{V}$ and $C \in \mathcal{W}$.

Corollary 2.3.9. Let $(X, Y), (Y, W)$, and (Z, A) be semi-coarse spaces, and suppose that (Y, W) is roofed [\(Definition A.1.11\)](#page-44-0). Then $f : (X \times Y, V \times W) \rightarrow (Z, \mathcal{A})$ is a bornologous function iff $(f \times f)(V \times \mathfrak{R}(\mathcal{W})) \in \mathcal{A}$ for all $V \in \mathcal{V}$.

Proof. If $(f \times f)(V \times \mathfrak{R}(\mathcal{W})) \in \mathcal{A}$ for all $V \in \mathcal{V}$, then for any $W \in \mathcal{W}$, $W \subset \mathfrak{R}(\mathcal{V})$, so $(f \times f)(V \times W) \subset (f \times f)(W \times \mathfrak{R}(V) \in \mathcal{A}$. Therefore, $(f \times f)(V \times W) \in \mathcal{A}$ and f is bornologous by [Proposition 2.3.7.](#page-9-0)

Conversely, suppose that f is bornologous. Then $(f \times f)(V \times \mathfrak{F}(W)) \in \mathcal{A}$. \Box

2.4. Quotient Spaces. In this section, we study quotient spaces of semi-coarse spaces, i.e. the semi-coarse structures constructed on a set Y given a semi-coarse space (X, V) and a surjective map $g: X \to Y$. We begin with the following definition.

Definition 2.4.1. Let (X, Y) be a semi-coarse space, let Y be a set, let $q: X \to Y$ be a surjective function, and define

$$
\mathcal{V}_g := \{ (g \times g)(V) : V \in \mathcal{V} \}.
$$

We call \mathcal{V}_g the semi-coarse quotient structure on Y induced by g, and we call (Y, \mathcal{V}_g) the semi-coarse quotient (space) of X induced by g .

Remark 2.4.2. The set V_q constructed above will occasionally be referred to as the semi-coarse structure inductively generated by the function $g: X \to Y$, and (Y, V_q) will likewise also be called the *semi-coarse space inductively generated by* the function $g: X \to Y$.

The following proposition justifies this choice of terminology.

Proposition 2.4.3. Let (X, V) , Y , $g: X \rightarrow Y$, and V_g be as in [Definition 2.4.1.](#page-10-1) Then (Y, V_g) is a semi-coarse space. Moreover, V_g is the coarsest semi-coarse structure which makes the function g bornologous.

Proof. Let (X, V) , Y , $g: X \to Y$, and V_g be as in the statement of the proposition. We verify that V_q satisfies the axioms of a semi-coarse structure.

(sc1) Since g is surjective, $g(X) = Y$, and therefore $(g \times g)(\Delta_X) = \Delta_Y$, which gives that $\Delta_Y \in \mathcal{V}_q$.

(sc2) Let $B \in V_q$ and suppose that $A \subset B$. Then there is a set $W \in V$ such that $(f \times f)(W) = B$, and for each $(y, y') \in A$, there is an $(x, x') \in W$ such that $g(x) = y$ and $g(x') = y'$. Defining

$$
A_g := \{ (x, x') \in W : (g \times g)(x, x') \in A \},
$$

we have that $A_g \subset W$ and $(g \times g)(A_g) = A$. Since $A_g \in V$, it follows that $A \in V_g$.

(sc3) Let $A, B \in V_g$. Then there are $A', B' \in V$ such that $(g \times g)(A') = A$ and $(g \times g)(B') = B$. Now note that $A \cup B = (g \times g)(A') \cup (g \times g)(B') = (g \times g)(A' \cup B')$, so $A \cup B \in \mathcal{V}_q$.

(sc4) Let $A \in \mathcal{V}_g$, then there is $W \in \mathcal{V}$ such that $(g \times g)(A) = W$. We observe that

$$
(g \times g)(A^{-1}) = \{(x, y) \mid \exists (x', y') \in A^{-1}, (g \times g)(x', y') = (x, y)\}
$$

=\{(x, y) \mid \exists (y', x') \in A, (g \times g)(x', y') = (x, y)\}
=\{(y, x) \mid \exists (x', y') \in A, (g \times g)(x', y') = (x, y)\}
=((g \times g)(A))^{-1}.

We conclude that V_g is a semi-coarse structure on Y.

We now show that V_g is the coarsest semi-coarse structure making g bornologous. By definition, g is a bornologous function iff for each $A \in \mathcal{V}$ we have that $(g \times g)(A) \in$ \mathcal{V}_g . However, by definition, $\{(g \times g)(A) : A \in \mathcal{V}\} = \mathcal{V}_g$, so it follows that any semicoarse structure W making g bornologous must contain V_g , and therefore V_g is the coarsest such structure.

□

The following property follows directly from the definition of the quotient structure.

Proposition 2.4.4. Let (Y, Y_q) be a semi-coarse space inductively generated by the function $g: X \to Y$ and let (Z, \mathcal{Z}) be a semi-coarse space. A function $f: (Y, \mathcal{V}_q) \to Y$ (Z, \mathcal{Z}) is bornologous iff $f \circ g : X \to Z$ is bornologous.

Proof. Suppose that $f : (Y, V_g) \to (Z, \mathcal{Z})$ is a bornologous function and let $A \subset X$, $A \in \mathcal{V}$. By definition, $(g \times g)(A) \in \mathcal{V}_g$, so $(f \times f) \circ (g \times g)(A) = (f \circ g \times f \circ g)(A) \in \mathcal{Z}$.

Now suppose that $f \circ g : (X, V) \to (Z, Z)$ is a bornologous function, and let $A \in \mathcal{V}_g$. Then there is a set $W \in \mathcal{V}$ such that $(g \times g)(W) = A$. It follows that $(f \times f)(A) = (f \circ g \times f \circ g)(W) \in \mathcal{Z}$, and we conclude that f is bornologous. \square

Definition 2.4.5. Let $p:(X,\mathcal{V}_X)\to(Y,\mathcal{V}_Y)$ be a surjective bornologous map. We say that p is a quotient map if, in addition, $V_Y = V_p$, where V_p is the semi-coarse structure defined in [Definition 2.4.1.](#page-10-1)

Remark 2.4.6. It follows immediately from the definition of quotient map that saying that $g: (X, V_X) \to (Y, V_Y)$ is a quotient map is equivalent to the statement that $B \in V_Y$ if and only if there exists $A \in V_X$ such that $B = (g \times g)(A)$.

The following theorem expresses the universal property of the quotient space (Y, V_q) .

Theorem 2.4.7. Let (X, \mathcal{V}_X) , (Y, \mathcal{V}_g) , $g : X \rightarrow Y$ be as in [Definition 2.4.1.](#page-10-1) Suppose that (Z, V_Z) is a semi-coarse space and that there exists a (not necessarily bornologous) map $h: X \to Z$ which is constant on each set $g^{-1}(\{y\})$. Then h

induces a map $f: Y \to Z$ such that $f \circ g = h$. The induced map f is bornologous from $(Y, V_q) \rightarrow (Z, V_Z)$ iff h is bornologous, and f is a quotient map iff h is a quotient map.

Proof. We first observe that the part of the theorem saying that f is bornologous if and only if h is bornologous is a direct implication of Proposition $2.4.4$.

Now we prove the statement about quotient maps. Suppose that f is a quotient map, then by [Remark 2.4.6,](#page-11-1) for every $C \in V_Z$ there exists $B \in V_q$ such that $C = (f \times f)(B)$. At the same time, g is a quotient map by hypothesis, so there exists $A \in \mathcal{V}_X$ such that $B = (g \times g)(A)$. Thus, there exists $A \in \mathcal{V}_X$ such that $C = (f \times f) \circ (g \times g)(A) = (h \times h)(A)$, obtaining that h is a quotient map. Observe that this fragment of the proof also says that the composition of quotient maps is a quotient map.

On the other hand, let h be a quotient map. Then for every $C \in V_Z$ there exists $A \in V_X$ such that $C = (h \times h)(A)$. Since g is a bornologous map, then $(g \times g)(A) \in \mathcal{V}_g$. Since $C = (h \times h)(A) = (f \times f) \circ (g \times g)(A)$, we may conclude that for every $C \in V_Z$ there exists $B = (g \times g)(A) \in V_g$ such that $C = (f \times f)(B)$. Therefore, f is a quotient map, as desired. \Box

An important example of an inductively generated semi-coarse space is the quotient space generated by an equivalence relation.

Example 2.4.7.1 (Quotient Semi-Coarse Space). Let (X, V) be a semi-coarse space and let \sim be an equivalence relation on X. Let $p : X \to X/\sim$ be the map $x \mapsto [x]$ sending each point $x \in X$ to its equivalence class $[x] \in X/\sim$. Then the semi-coarse space $(X/\sim, \mathcal{V}_p)$ is called the quotient space of X induced by the equivalence relation ∼.

It will be useful in the following to have an alternate formulation of the quotient semi-coarse structure induced by an equivalence relation, which we provide in the next proposition.

Proposition 2.4.8. Let (X, V) be a semi-coarse space and let ∼ be an equivalence relation on X. Furthermore, extend the relation \sim to $X \times X$ by defining $(x, y) \sim$ (x', y') iff $x \sim x'$ and $y \sim y'$. We denote by $[x, y]$ the equivalence class of (x, y) , and for a subset $A \subset X \times X$, we define

$$
[A] := \{ [x, y] \mid (x, y) \in A \},
$$

and we let \mathcal{V}/\sim denote the collection

$$
\mathcal{V}/\!\!\sim\; := \{[B] \mid B \in \mathcal{V}\}.
$$

Finally, let $p : X \to X/\sim$ be the map $x \mapsto [x]$ sending a point $x \in X$ to its equivalence class $[x] \in X/\sim$.

Then $\mathcal{V}_p = \mathcal{V}/\!\!\sim$.

Proof. First, let $A \in \mathcal{V}/\sim$. Then there is a set $A' \in \mathcal{V}$ such that $A = [A']$. Furthermore, $A = [A'] = (p \times p)(A')$, and we conclude that $A \in \mathcal{V}_p$, so $\mathcal{V}/\sim \mathcal{L} \mathcal{V}_p$.

Now suppose that $A \in \mathcal{V}_p$. Then there is a set $A' \in \mathcal{V}$ such that $(p \times p)(A') = A$. However, by the definition of p, $(p \times p)(A') = [A'] = A$, and therefore $A \in \mathcal{V}/\sim$, so $\mathcal{V}_p \subset \mathcal{V}/\sim$ as well. \Box

3. HOMOTOPY

In this section, we develop the basics of homotopy theory in semi-coarse spaces. One of the key difficulties in doing so is that there is no natural semi-coarse structure on the topological interval $[0, 1]$. Naively, one could try to endow the set $[0, 1]$ with any of the semi-coarse structures \mathcal{V}_r , \mathcal{V}_r^{\lt} , $0 < r < 1$ defined in [2.1.7.1,](#page-4-2) and then use the product $(X, V) \times ([0, 1], V_r)$ as the cylinder for the space (X, V) . However, in the following proposition, we see that the concatenation of two semi-coarse intervals $([0,1], \mathcal{V}_{r'})$ and $([0,1], \mathcal{V}_{r''})$ does not result in an interval of the form $([0,1], \mathcal{V}_r)$. A consequence of this is that proceeding in this manner will not allow us to define a product between two paths by simply concatenating them, complicating the development of homotopy theory for semi-coarse spaces.

Proposition 3.0.1. Let $r, r', r'' \in (0, 1)$. Any bijective map

 $\phi: ([0,1], \mathcal{V}_r) \to ([0,1], \mathcal{V}_{r'}) \sqcup_{1 \sim 1} ([1,2], \mathcal{V}_{r''}),$

where the latter space is the pushout of the two intervals, glued at one endpoint, is not bornologous at the point $1 \in [0,2]$ where the two intervals are glued. Then ϕ cannot be a bornologous map, and the spaces $([0,1], \mathcal{V}_r)$ and $([0,1], \mathcal{V}_{r'}) \sqcup_{1 \sim 1}$ $([1,2], \mathcal{V}_{r''})$ are not homeomorphic.

Proof. Denote $([0,1], \mathcal{V}_{r'}) \sqcup_{1 \sim 1} ([1,2], \mathcal{V}_{r''})$ by (Z, \mathcal{V}_Z) . Suppose that $\phi : ([0,1], \mathcal{V}_r) \to$ (Z, V_Z) is bijective and bornologous. By the Archimedean property, there exists an integer n such that $(n - 1)(r/2) < 1 \leq n(r/2)$; in particular, we can compute n as the least integer that is greater or equal to $2/r$.

Consider the set $\{x_0, x_1, \ldots, x_{n-1}, x_n\}$ with $x_i = i(r/2)$ if $i < n$ and $x_n = 1$. Observe that $|x_i - x_{i+1}| < r$ for every i, then

$$
(\phi \times \phi)([x_i, x_{i+1}] \times [x_i, x_{i+1}]) \in \mathcal{V}_r
$$

There is no pair $z_i, z'_i \in [x_i, x_{i+1}]$ such that $\phi(z_i) < 1$ and $\phi(z'_i) > 1$ because $\{(\phi(z_i), \phi(z'_i))\} \notin V_Z$ and ϕ would not be a bornologus map. Thus, we obtain that either $\phi([x_i, x_{i+1}]) \subset [0,1]$ or $\phi([x_i, x_{i+1}]) \subset [1,2]$. Without loss of generality, suppose that $\phi([0, x_1]) \subset [1, 2]$; this implies that every $\phi([x_i, x_{i+1}]) \subset [1, 2]$ and then $\phi([0,1]) \subset [1,2]$, from which we conclude that ϕ is not bijective, a contradiction.

Therefore ϕ is either not bornologous or not bijective. \Box

In order to resolve the problem highlighted by the [Proposition 3.0.1,](#page-13-2) we adapt the construction in [\[1\]](#page-55-13) to the semi-coarse category, using finite intervals in $\mathbb Z$ endowed with their 'nearest neighbor' semi-coarse structure in place of the interval for homotopical constructions. In contrast to [\[1\]](#page-55-13), however, the product which we use throughout is the categorical product for semi-coarse spaces, which gives the resulting theory a more simplicial, rather than cubical, character.

3.1. Homotopy and Homotopy Equivalence for Semi-Coarse Spaces. Before we define homotopy, we define the semi-coarse structure on $\mathbb Z$ which will be used throughout.

Definition 3.1.1. We call the semi-coarse structure on \mathbb{Z} generated by the graph $G = (\mathbb{Z}, E)$, where $E \coloneqq \bigcup_{z \in \mathbb{Z}} \{ (z, z - 1), (z, z), (z, z + 1) \},$ i.e.

the the canonical semi-coarse structure of \mathbb{Z} , and we denote this structure by \mathcal{Z} .

Remark 3.1.2. Note that the canonical semi-coarse structure on \mathbb{Z} is the roofed semi-coarse structure [\(Definition A.1.11\)](#page-44-0) on $\mathbb Z$ with roof E.

With the semi-coarse structure on $\mathbb Z$ in place, we now define homotopy between bornologous maps.

Definition 3.1.3. Let (X, V) and (Y, W) be semi-coarse spaces and let f, g : $(X, V) \rightarrow (Y, W)$ be bornologous functions. We will say that f is homotopic to g, written $f \simeq_{sc} g$, iff there is a bornologous function $H : (X \times \mathbb{Z}, V \times \mathcal{Z}) \to (Y, W)$ and integers $N, M \in \mathbb{Z}$ with $M \langle N, \text{where } H(x,n) = f(x)$ if $n \leq M$ and $H(x, n) = g(x)$ if $n \geq N$.

We will say that two semi-coarse spaces (X, V) and (Y, W) are *homotopy equiv*alent when there are bornologous functions $f: X \to Y$ and $g: Y \to X$ such that $g \circ f \simeq_{sc} id_X$ and $f \circ g \simeq_{sc} id_Y$.

Proposition 3.1.4. Homotopy equivalence \simeq_{sc} is an equivalence relation on bornologous functions.

Proof. We check that \simeq_{sc} satisfies the axioms of an equivalence relation.

Reflexivity: Let (X, V) and (Y, W) be semi-coarse spaces, let $f : (X, V) \to (Y, W)$ be a bornologous function, and suppose that $H : X \times \mathbb{Z} \to Y$ is a map such that $H(x, z) = f(x)$ for each $z \in \mathbb{Z}$. Since $H(x, n) = x$ for all $n \in \mathbb{Z}$, to see that $f \simeq_{sc} f$, it is enough to prove that H is bornologous. From the definition of H , we have

$$
(H \times H)(V \boxtimes \mathfrak{R}(\mathcal{Z})) = (f \times f)(V) \in \mathcal{W},
$$

so that H is bornologous by [Corollary 2.3.9.](#page-10-2) It follows that $f \simeq_{sc} f$.

Symmetry: Suppose that $f \simeq_{sc} g$. Then there exists a bornologous function $H: X \times \mathbb{Z} \to Y$ and integers $M \lt N$ such that $H(x,n) = f(x)$ for all $n \leq M$ and $H(x, x) = g(x)$ for all $n \geq N$. Defining $H'(x, n) = H(x, -n)$, we have that $H'(x, n) = g(x)$ for each $x \in X$ if $n \leq -N$, and $H'(x, n) = f(x)$ for each $x \in X$ is $n' \geq -M$. Moreover, the function $h : (X \times \mathbb{Z}, V \times \mathbb{Z}) \to (X \times \mathbb{Z}, V \times \mathbb{Z})$ given by $h(x, n) := (x, -n)$ is bornologous by [Corollary 2.3.9,](#page-10-2) since $(h \times h)(V \times \mathfrak{R}) =$ $V \times \mathfrak{R} \in V \times \mathcal{Z}$. Therefore, $H' = H \circ h$ is a bornologous function, from which it follows that $g \simeq_{sc} f$.

Transitivity: Suppose that $f \simeq_{sc} g$ and $g \simeq_{sc} h$. Then there are bornologous functions $H_{fg}: (X \times \mathbb{Z}, V \times \mathbb{Z} \to Y \text{ and } H_{gh}: X \times \mathbb{Z} \to Y \text{ and pairs of integers})$ $M_f < N_q$ and $M_q < N_h$ such that

$$
H_{fg}(x, n) = \begin{cases} f(x) & n \le M_f \\ g(x) & n \ge N_g, \end{cases}
$$
 and

$$
H_{gh}(x, n) = \begin{cases} g(x) & n \le M_g \\ h(x) & n \le N_h. \end{cases}
$$

Without loss of generality, we take $M_f < -1 = N_g$ and $M_g = 1 < N_h$. Define $H_{fh}: X \times \mathbb{Z} \to Y$ by

$$
H_{fh}(x,n) = \begin{cases} H_{fg}(x,n) & n \le 0\\ H_{gh}(x,n) & n > 0, \end{cases}
$$

and observe that $H_{fh}(x, n) = f(x)$ for all $n \leq M_f$, $H_{fh}(x, n) = h(x)$ for all $n \geq N_h$, and $H_{fh}(x, n) = g(x) = H_{fg}(x, n) = H_{gh}(x, n)$ for $-1 = N_g \le n \le M_g = 1$. H_{fh} is bornologous by [Proposition 2.2.4,](#page-7-2) where $X_1 = X \times ((-\infty, 0] \times \mathbb{Z})$ and $X_2 = X \times ([0, \infty) \cap \mathbb{Z})$, and X_1 and X_2 are endowed with the respective subspace semi-coarse structures induced from $\mathcal{V} \times \mathcal{Z}$.

Let $\mathbb{Z}_{-}^{2} := \{(a,b) \in \mathbb{Z}^{2} \mid a \le 0\}$, and observe that, for any $V \in \mathcal{V}$, if $(x, n, x', n') \in \mathcal{V}$ $V \boxtimes (\mathfrak{R}(\mathcal{Z}) \cap \mathbb{Z}_{-}^2)$ then $n' = n + k$, where $k \in \{-1, 0, 1\}$. In this case, we also have

$$
H_{fh}(V \boxtimes (\mathfrak{R}(\mathcal{Z}) \cap \mathbb{Z}_{-}^{2})) = H_{fg}(V \boxtimes (\mathfrak{R}(\mathcal{Z}) \cap \mathbb{Z}_{-}^{2}))
$$

and therefore $H_{fh}(V \boxtimes (\mathfrak{R}(\mathcal{Z}) \cap \mathbb{Z}_-^2)) \in \mathcal{W}$.

Similarly, let $\mathbb{Z}_+^2 \coloneqq \{(a, b) \in \mathbb{Z}^2 \mid a \geq 0\}$, and observe that, for any $V \in \mathcal{V}$), if $(x, z, x', z') \in V \boxtimes (\mathfrak{R}(\mathcal{Z}) \cap \mathbb{Z}_+^2)$, then $z' = z + k$, where $k \in \{-1, 0, 1\}$. In this case, we also have

$$
H_{fh}(V \boxtimes (\mathfrak{R}(\mathcal{Z}) \cap \mathbb{Z}_+^2)) = H_{gh}(V \boxtimes ((\mathcal{Z}) \cap \mathbb{Z}_+^2)),
$$

and therefore $H_{fh}(V \boxtimes (\mathfrak{R}(\mathcal{Z}) \cap \mathbb{Z}_+^2)) \in \mathcal{W}$. Since

$$
H_{fh}(V \boxtimes \mathfrak{R}(\mathbb{Z})) = H_{fh}(V \boxtimes (\mathfrak{R}(\mathcal{Z}) \cap \mathbb{Z}_-^2)) \cup H_{fh}(V \boxtimes (\mathfrak{R}(\mathcal{Z}) \cap \mathbb{Z}_+^2))
$$

for each $V \in \mathcal{V}$, it follows from [Corollary 2.3.9](#page-10-2) that H_{fh} is a bornologous function. We conclude that $f \simeq_{sc} h$. □

The following proposition gives a useful condition for determining whether a potential homotopy is bornologous.

Proposition 3.1.5. Let (X, V) and (Y, W) be semi-coarse spaces, let $f, g: (X, V) \rightarrow$ (Y, W) be bornologous maps, and suppose that $H : X \times \mathbb{Z} \to Y$ is a (possibly nonbornologous) map such that there exist integers $M_f \langle N_g \text{ with } H(x,n) = f(x)$ for all $n \leq M_f$ and $H(x, n) = g(x)$ for all $n \geq N_g$.

If $(H \times H)(V \boxtimes (n, n + 1)), (H \times H)(V \boxtimes (n, n)),$ and $(H \times H)(V \boxtimes (n, n - 1))$ are controlled sets of (Y, W) for each $V \in V$ and $n \in \mathbb{Z}$, then H is bornologous.

Proof. Let $\mathbb{Z}_{fg}^2 := \Re(\mathcal{Z}) \cap \{(a,b) \in \mathbb{Z}^2 \mid M_f \le a \le N_g\}$. Since $H(x,n) = f(x)$ for $n \leq M_f$ and $H(x, n) = g(x)$ for $n \geq N_g$, it follows that, for any $V \in \mathcal{V}$,

$$
(H \times H)(V \boxtimes \mathfrak{R}(\mathcal{Z})) = (H \times H)(V \boxtimes \mathbb{Z}_{fg}^2)
$$

=
$$
(H \times H) \left(\bigcup_{(a,b) \in \mathbb{Z}_{fg}^2} (V \boxtimes (a,b)) \right).
$$

Since the finite union of sets in W are in W, H is bornologous iff, for all $(a, b) \in \mathbb{Z}_{fg}^2$, $(H\times H)(V\boxtimes (a, b))\in W$, which is true iff $(H\times H)(V\boxtimes (n, n+1)), (H\times H)(V\boxtimes (n, n)),$ and $(H \times H)(V \boxtimes (n, n-1))$ are controlled sets of (Y, W) for every $n \in \mathbb{Z}$.

In the following, we define the homotopy of maps of pairs and triples for semicoarse spaces and show that they are also an equivalence relation.

Definition 3.1.6 (Homotopy of Maps of Pairs and Triples). Let (X, V) and (Y, W) be semi-coarse spaces, let $B \subset A \subset X$ and $D \subset C \subset Y$ be endowed with the subspace semi-coarse structures, and let $f, g : (X, A, B; V) \rightarrow (Y, C, D; W)$ be bornologous functions of a triple, i.e. such that $f|_A \subset C$ and $f|_B \subset D$. We say that *relatively homotopic to g* and write $f \simeq_{sc} g$ iff there is a homotopy $H: (X \times \mathbb{Z}, \mathcal{V} \times \mathcal{Z}) \to (Y, \mathcal{W})$ such that $H|_{A \times \mathbb{Z}} \subset C$ and $H|_{B \times \mathbb{Z}} \subset D$.

We define a homotopy between maps of pairs $f, g: (X, A; \mathcal{V}) \to (Y, C; \mathcal{W})$ to be the homotopy between maps of a triple as above with $B = A$ and $C = D$.

Lemma 3.1.7. Let (X, V) and (Y, W) be semi-coarse spaces, and $B \subset A \subset X$, $D \subset C \subset Y$ be endowed with the subspace semi-coarse structures. Then the relative homotopy of triples \cong_{sc} is an equivalence relation.

Proof. The proof is nearly identical to the proof of [Definition 3.1.3,](#page-14-0) with the addition that we need to observe that the $H_{fh}|_A \subset C$ (and $H_{fh}|_D \subset B$). However, this follows the fact that H_{fg} and H_{gh} are homotopies of triples and from the definition of H_{fh} .

We end this section with the definition of homotopy equivalence.

Definition 3.1.8. Two semi-coarse spaces (X, V) and (Y, W) are said to be homotopy equivalent, written $(X, V) \simeq (Y, W)$, iff there exist bornologous maps $f:(X,\mathcal{V})\to (Y,\mathcal{W})$ and $g:(Y,\mathcal{W})\to (X,\mathcal{V})$ such that $f\circ g\simeq_{sc}1_Y$ and $g \circ f \simeq_{sc} 1_X$.

3.2. Homotopy Groups. As mentioned in Section [1,](#page-1-0) to define the homotopy groups for semi-coarse spaces, we will largely follow the ideas in the construction of the A-homotopy groups from [\[1\]](#page-55-13), with the difference that the product we use is the categorical product rather than a semi-coarse version of the graph product. We begin this section with the construction of the relative homotopy classes of maps of cubes in \mathbb{Z}^n to a semi-coarse space (X, V) . We set some notation in the next definition.

Definition 3.2.1. Let n and m be natural numbers, let I_m be the set $\{0, 1, \dots, m\}$, and let \mathcal{Z}_m be the subspace semi-coarse structure induced by the inclusion of I_m in (\mathbb{Z}, \mathbb{Z}) . We denote by (I_m, \mathbb{Z}_m) the corresponding semi-coarse space, and we denote by (I_m^n, \mathcal{Z}_m^n) the Cartesian product of n copies of (I_m, \mathcal{Z}_m) . We write a_i for the *i*-th coordinate of a point $a \in \mathbb{Z}^n$, where $i \in \{1, \dots, n\}$, unless otherwise stated. Finally, we denote by

$$
\partial I_m^n \coloneqq \{(x_1, \ldots, x_n) \in I_m^n \mid \exists i \in \{1, \ldots, n\} \text{ such that } x_i \in \{0, m\}\},\
$$

the boundary of the discrete *n*-cube with sides of length m , and we define $J_m^{n-1} \subset$ ∂I_m^n to be the "discrete (upside-down) open *n*-dimensional box with sides of length m", i.e. the set

 $J_m^{n-1} = \{(x_1, \ldots, x_n) \in \partial I_m^n \mid \text{If } x_n = 0, \text{ then } \exists i \neq n \text{ such that } x_i \in \{0, m\}\}.$

To define the relative homotopy groups of a pointed semi-coarse space pair $(X, A, *; \mathcal{V})$ (i.e. $* \in A \subset X$), we will consider maps of triples

$$
f:(I_m^n, \partial_m^n, J_m^{n-1}; \mathcal{Z}_m^n) \to (X, A, *; \mathcal{V}),
$$

and we define $\pi_n^{sc}(X, A, *; \mathcal{V})$ to be a certain direct limit which we construct below. First, however, we require the following definition and lemmas.

Definition 3.2.2. Let $m < m'$ be two positive integers. We define the retraction $\phi_m^{m'} : I^n_{m'} \to I^n_m$ by

$$
(\phi_m^{m'}(x))_k \coloneqq \begin{cases} x_k & x_k \le m, \\ m & m < x_k \le m', \end{cases}
$$

where $(\phi_m^{m'}(x))_k$ is the k-th coordinate of $\phi_m^{m'}(x)$. We will typically abuse notation and write $\phi_m^{m'}$ for the retraction $I_{m'}^n \to I_m^n$ for any dimension n, as the dimension will usually be clear from context.

Lemma 3.2.3. The function $\phi_m^{m'} : (I^n_{m'}, \mathcal{Z}^n_{m'}) \to (I^n_m, \mathcal{Z}^n_m)$ is bornologous.

Proof. The result follows from [Proposition 2.2.4,](#page-7-2) [Proposition A.1.17,](#page-46-0) and the definition of the semi-coarse structures $\mathcal{Z}_{m'}^n$ and \mathcal{Z}_m^n \Box m.

Lemma 3.2.4. Let $f:(I_m^n, \partial I_m^n, J_m^{n-1}; \mathcal{Z}^n) \to (X, A, *, \mathcal{V})$ be a bornologous function of triples. Then for any $m' \in \mathbb{Z}$ with $m' > m$, f has a bornologous extension of triples

$$
f': (I^n_{m'}, \partial I^n_{m'}, J^{n-1}_{m'}; \mathcal{Z}^n) \to (X, A, *; \mathcal{V})
$$

given by $f'(z) = f \circ \phi_m^{m'}(z)$.

Furthermore, if $f \simeq_{sc} g$ as maps of triples $(I_m^n, \partial I_m^n, J_m^{n-1}; \mathcal{Z}_m^n) \to (X, A, *, \mathcal{V}),$ then for the extensions f' and g' to $I_{m'}^n$ of f and g, respectively, we have $f' \simeq_{sc} g'$ as well.

Proof. The extension f' of f is bornologous since the composition of bornologous functions is bornologous.

If $f \simeq_{sc} g$ then there exists a homotopy of triples $H: (I_m^n \times \mathbb{Z}, \mathbb{Z}_m^n \times \mathbb{Z}) \to (X, V)$ between f and g. We define $H': (I^n_{m'} \times \mathbb{Z}, \mathcal{Z}^n_{m} \times \mathcal{Z}) \to (\widetilde{X}, \mathcal{V})$ by

$$
H'(x,n) \coloneqq H(\phi_m^{m'}(x),n).
$$

Since $\phi_m^{m'} \times Id$ and H are bornologous, it follows that H' is bornologous, and therefore H' is a homotopy from f' to g' by definition.

[Lemma 3.2.4](#page-17-0) implies that extending a map from I_m^n to $I_{m'}^n$ induces a map on the relative homotopy classes

$$
i_m^{m^\prime}: [(I^n_m,\partial I^n_m,J^{n-1}_m;\mathcal{Z}^n_m),(X,A,*;\mathcal{V})] \rightarrow [(I^n_{m^\prime},\partial I^n_{m^\prime},J^{n-1}_m;\mathcal{Z}^n_m),(X,A,*;\mathcal{V})]
$$

such that $i_m^{m'} = i_{m'}^{m'} \circ i_m^{m'}$. The maps $i_m^{m'}$ therefore make the homotopy classes into a directed system of sets $\left([(I_m^n, \partial I_m^n, \mathcal{Z}^n), (X, \ast, \mathcal{V})], i_m^{m'}, \mathbb{N} \right)$. We define the classes $\pi_n^{sc}(X, A^*; \mathcal{V})$ in the following manner.

Definition 3.2.5. Given the set $\{*,1\}$ with the diagonal semi-coarse structure $\mathcal{D}_{*,1}$, i.e. $\mathcal{D}_{*,1} = \mathcal{P}(\{(*, *), (1, 1)\})$. We define the set $\pi_0^{sc}(X, A, *, \mathcal{V})$ by

$$
\pi_0^{sc}(X, A, *; \mathcal{V}) \coloneqq [(\{*, 1\}, *, *; \mathcal{D}_{0,1}), (X, A, *; \mathcal{V})].
$$

For $n \geq 1$, we define $\pi_n^{sc}(X, A, *, \mathcal{V})$ by

$$
\pi_n^{sc}(X, A, *; \mathcal{V}) \coloneqq \varinjlim \left([(I^n_m, \partial I^n_m, J^{n-1}_m; \mathcal{Z}^n_m), (X, A, *; \mathcal{V})], i^{m'}_m, \mathbb{N} \right),
$$

For a homotopy class $[f] \in [(I_m^n, \partial I_m^n, J_m^{n-1}; \mathcal{Z}_m^n), (X, A, *; \mathcal{V})]$, we write $\langle f \rangle$ for its image in $\pi_n^{sc}(X, \ast, \mathcal{V})$.

When $A = *$, we write $\pi_n^{sc}(X, *, *; \mathcal{V})$ as $\pi_n^{sc}(X, *, \mathcal{V})$ or $\pi_n^{sc}(X; \mathcal{V})$.

Remark 3.2.6. One may also construct the homotopy classes $\pi_n^{sc}(X, \ast, \mathcal{V})$ as a direct limit of homotopy classes of maps of pairs $f:(\mathbb{Z}^n, J_m^n, \mathcal{Z}^n) \to (X, \ast, \mathcal{V})$, where $J_m^n := \mathbb{Z} - I_{m-1}^n$. One sees that these formulations are equivalent by extending the maps $f:(I_m^n, \partial I_m^n, \mathcal{Z}_m^n) \to (X, \ast, \mathcal{V})$ to \mathbb{Z}^n , where the extension f' sends all of J_m^n

to $* \in X$. The same argument as in [Lemma 3.2.4](#page-17-0) shows that f' is bornologous, and it follows from the definitions that

$$
f \simeq_{sc} g \iff f' \simeq_{sc} g'.
$$

Therefore the homotopy classes $\pi_n^{sc}(X, *; \mathcal{V})$ may be defined as

$$
\pi^{sc}_n(X,\ast;\mathcal{V})\coloneqq\varinjlim\left([(\mathbb{Z}^n,E^n_m,\mathcal{Z}^n),(X,\ast,\mathcal{V})],i^{m'}_m,\mathbb{N}\right),
$$

where, as before, the $i_m^{m'}$ are the maps on the respective homotopy classes induced by the interpreting a map $f: (\mathbb{Z}^n, E_m^n, \mathcal{Z}^n) \to (X, \ast, \mathcal{V})$ as a map $f: (\mathbb{Z}^n, E_{m'}^n, \mathcal{Z}^n) \to$ $(X, \ast, \mathcal{V}).$

Remark 3.2.7. Let $\langle f \rangle$, $\langle g \rangle \in \pi_n^{sc}(X, A, *, \mathcal{V})$. Observe that $\langle f \rangle = \langle g \rangle$ iff there exist $m \in \mathbb{N}$ and bornologous functions $\tilde{f}, \tilde{g} : (I_m^n, \mathcal{Z}_m^n) \to (X, \mathcal{V})$ such that $[\tilde{f}] \in \langle f \rangle$, $[\tilde{q}] \in \langle q \rangle$ and $\tilde{f} \simeq_{sc} \tilde{q}$.

We will now define an operation \star on the homotopy classes $\pi_n^{sc}(X, *; \mathcal{V})$ and $\pi(X, A, *; \mathcal{V})$ and show this operation makes these classes into a group $n \geq 1$ and $n \geq 2$, respectively.

Definition 3.2.8 (Operation \star). Let X be a set, let $n \in \mathbb{N}$, m and m' be nonnegative integers, and suppose that $f: I_m^n \to (X, V)$ and $g: I_{m'}^n \to X$ are bornologous functions. Let $\alpha = (\alpha_1, \alpha_2, \ldots, \alpha_n) \in I^n_{m+m'}$, and let $e_1 = (1, 0, \ldots, 0) \in \mathbb{Z}^n$. We define the \star -product $f \star g : I^n_{m+m'} \to X$ such that

$$
f \star g(\alpha) = \begin{cases} f(\phi_m^{m+m'}(\alpha)) & \text{if } 0 \le \alpha_1 \le m \\ g(\phi_{m'}^{m+m'}(\alpha - me_1)) & \text{if } m < \alpha_1 \le m+m'. \end{cases}
$$

The following proposition tells us that the result of applying the \star operation to a pair of bornologous functions which agree on the boundary is bornologous.

Proposition 3.2.9. Let $n \in \mathbb{N}$, let m and m' be non-negative integers, and suppose that $f:(I_m^n,\mathcal{Z}_m^n)\to (X,\mathcal{V})$ and $g:(I_{m'}^n,\mathcal{Z}_m^n)\to (X,\mathcal{V})$ are bornologous functions such that

 $f((m, \alpha_2, \ldots, \alpha_n)) = g((0, \alpha_2, \ldots, \alpha_n))$

is satisfied for all $\alpha_2, \ldots, \alpha_n$ where both $f((m, \alpha_2, \ldots, \alpha_n))$ and $g((0, \alpha_2, \ldots, \alpha_n))$ are defined. Then $f \star g$ is bornologous.

Proof. Let $n \in \mathbb{N}$, m and m' be non-negative integers, and suppose that f: $(I_m^n, \mathcal{Z}_m^n) \to (X, V)$ and $g: (I_{m'}^n, \mathcal{Z}_{m'}^n) \to (X, V)$ satisfy the hypothesis of the proposition. Let K_1 and K_2 be the sets

$$
K_1 = \{ \alpha \in I_{m+m'}^n \mid 0 \le \alpha_1 \le m
$$

$$
K_2 = \{ \alpha \in I_{m+m'}^n \mid m \le \alpha_1 \le m' + m.
$$

Then the restrictions $f \star g|_{K_1}$ and $f \star g|_{K_2}$ are bornologous, and K_1 and K_2 satisfy the hypothesis of [Proposition 2.2.4.](#page-7-2) Therefore $f \star g$ is bornologous by Proposition 2.2.4. □

Corollary 3.2.10. Suppose that $f: (I_m^n, \partial I_m^n, J_m^{n-1}; \mathcal{Z}_m^n) \to (X, A, *, \mathcal{V})$ and g: $(I_{m'}^n, \partial I_{m'}^n, J_{m'}^{n-1}; \mathcal{Z}_{m'}^n) \rightarrow (X, A, *; \mathcal{V})$ are bornologous functions of triples. Then $f \star g: (I^n_{m+m'}, \partial I^n_{m+m'}, J^{n-1}_{m+m'}; \mathcal{Z}^n_{m+m'}) \to (X, V)$ is bornologous.

Proof. Since f and q are maps of triples, it follows by definition that

$$
f((m, \alpha_2, \ldots, \alpha_n)) = * = g(-m', \alpha_2, \ldots, \alpha_n))
$$

whenever both sides are defined, so the hypotheses of [Proposition 3.2.9](#page-18-0) are satisfied. The conclusion follows. \Box

Corollary 3.2.11. Let $f:(I_m^n, \partial I_m^n; \mathcal{Z}_m^n) \to (X, \ast; \mathcal{V})$ and $g:(I_{m'}^n, \partial I_{m'}^n; \mathcal{Z}_{m'}^n) \to$ $(X, *; \mathcal{V})$ be bornologous functions of triples. Then $f \star g : (I^n_{m+m'}, \partial I^n_{m+m'}; \mathcal{Z}^n_{m+m'}) \to$ (X, V) is bornologous.

Proof. The result follows from [Corollary 3.2.10](#page-18-1) with $A = *$. □

The next lemma enables us to to extend the definition of \star from functions to homotopy classes.

Lemma 3.2.12. Let $n \geq 2$ and suppose that $\langle f \rangle, \langle g \rangle \in \pi_n^{sc}(X, A, *)$, then the product $\langle f \rangle \star \langle g \rangle := \langle f \star g \rangle$ is well-defined. Similarly, if $n \geq 1$ and $\langle f \rangle, \langle g \rangle \in$ $\pi_n^{sc}(X, *),$ then the product $\langle f \rangle \star \langle g \rangle := \langle f \star g \rangle$ is well-defined.

Proof. Let $n \geq 2$, and suppose that $\langle f \rangle$, $\langle f' \rangle$, $\langle g \rangle$, $\langle g' \rangle \in \pi_n^{sc}(X, A, *, \mathcal{V})$ such that $\langle f \rangle = \langle f' \rangle$ and $\langle g \rangle = \langle g' \rangle$. Then, by [Remark 3.2.7,](#page-18-2) there exist non-negative integers m and m' and functions $\tilde{f}, \tilde{f}' : (I_m^n, \partial I_m^n, J_m^{n-1}; \mathcal{Z}_m^n) \to (X, A, *, \mathcal{V})$ and \tilde{g}, \tilde{g}' : $(I_{m'}^n, \partial I_{m'}^n, J_{m'}^{n-1}; \mathcal{Z}_{m'}^n) \to (X, A, *, \mathcal{V})$ such that $\tilde{f} \simeq_{sc} \tilde{f}'$ and $\tilde{g} \simeq_{sc} \tilde{g}'$. We wish to show that $\tilde{f} \star \tilde{g} \simeq_{sc} \tilde{f}' \star \tilde{g}'.$

Let H_1 be a homotopy between \tilde{f} and \tilde{f}' and H_2 be a homotopy between \tilde{g} and \tilde{g}' . Define a function

$$
H: (I^n_{m+m'} \times \mathbb{Z}, \partial I^n_{m+m'} \times \mathbb{Z}, J^{n-1}_{m+m'} \times \mathbb{Z}; \mathcal{Z}^n_{m+m'} \times \mathcal{Z}) \to (X, A, *, \mathcal{V})
$$

by

$$
H(\alpha, n) = \begin{cases} H_1(\alpha + m'e_1, n) & -m' - m \le \alpha_1 \le -m' + m \\ H_2(\alpha + me_1, n) & -m' + m < \alpha_1 \le m' + m. \end{cases}
$$

Then H is a map of triples by definition, and H is bornologous by [Proposition 2.2.4](#page-7-2) and H is a homotopy between $\tilde{f} \star \tilde{g}$ and $\tilde{f}' \star \tilde{g}'$. By [Remark 3.2.7,](#page-18-2) it follows that $\langle f \rangle \star \langle g \rangle = \langle f \star g \rangle$ is well-defined.

For $\pi_n^{sc}(X, *; \mathcal{V})$, the above proof also applies for any $n \geq 1$, with the modification that all of the maps are bornologous maps of pairs instead of maps of triples. \Box

Given [Lemma 3.2.12,](#page-19-0) we make the following definition.

Definition 3.2.13. For $n \geq 2$ and $\langle f \rangle$, $\langle g \rangle \in \pi_n^{sc}(X, A, *, \mathcal{V})$, we define $\langle f \rangle \star \langle g \rangle \coloneqq$ $\langle f \star g \rangle$.

Similarly, for $n \geq 1$ and $\langle f \rangle, \langle g \rangle \in \pi_n^{sc}(X, *; \mathcal{V})$, we define $\langle f \rangle \star \langle g \rangle := \langle f \star g \rangle$.

We will now show that the homotopy classes π_n^{sc} with the \star operation form a group given assumptions on n identical to those in the topological case. This theorem is stated below as [Theorem 3.2.18,](#page-23-0) and the proof is an adaptation of the proof for topological spaces. The semi-coarse setting introduces a number of technical subtleties, however, which must be dealt with in order to give a complete proof of this theorem. While not difficult, they are nonetheless intricate, and are handled in [Lemma 3.2.15](#page-20-0)[-Lemma 3.2.17](#page-22-0) below. The next definition establishes notation for these lemmas.

Definition 3.2.14. In the following definitions, we denote by a the point

$$
\mathbf{a}=(a_1^\leftarrow,a_1^\rightarrow,a_2^\leftarrow,a_2^\rightarrow,\cdots,a_n^\leftarrow,a_n^\rightarrow)\in I_m^{2n},
$$

where $a_i^{\leftarrow} \leq a_i^{\rightarrow}$ for all $i \in \{1, \dots, n\}$.

(i) We call the set $\mathcal{B}_{\mathbf{a}}$

$$
\mathcal{B}_{\mathbf{a}} := \{ a \in I_m^n : a_i^{\leftarrow} \le a_i \le a_i^{\rightarrow}, i \in \{1, \dots, n\} \}.
$$

the block delimited by **a**. We write $\mathcal{B}_{\mathbf{a}}^{\hat{n},k}$ for the set $\{a \in \mathcal{B}_{\mathbf{a}} : a_{\hat{n}} = k\}$. When $a_{\hat{n}}^{\leftarrow} = a_{\hat{n}}^{\rightarrow}$ for some $\hat{n} \in \{1, \ldots, n\}$, we denote $\mathcal{B}_{\mathbf{a}}$ by $\mathcal{P}_{\mathbf{a}}^{\hat{n}}$, which we call a plate. (ii) Let \mathcal{B}_a be the block delimited by a. We denote by $\partial \mathcal{B}_a$ the set

$$
\{a \in \mathcal{B}_\mathbf{a} : \exists k \in \{1, \dots, n\} \text{ with } a_k \in \{a_k^{\leftarrow}, a_k^{\rightarrow}\}\}\
$$

which we call the *boundary* of \mathcal{B}_{a} .

(iii) Let \mathcal{B}_a be the block delimited by a, let $\hat{n} \in \{1, \dots, n\}$, and suppose that $\epsilon \in \{\leftarrow, \rightarrow\}.$ We will call $J_{\mathbf{a}}^{\hat{n}, \epsilon} \subset \partial \mathcal{B}_{\mathbf{a}}$ the (\hat{n}, ϵ) -open box of $\mathcal{B}_{\mathbf{a}}$, which we define to be the set

$$
J_{\mathbf{a}}^{\hat{n},\epsilon} := \{ a \in \partial \mathcal{B}_{\mathbf{a}} \mid \text{If } a_{\hat{n}} = a_{\hat{n}}^{\epsilon}, \text{ then } \exists k \neq \hat{n} \text{ such that } a_k \in \{ a_k^{\leftarrow}, a_k^{\rightarrow} \} \}.
$$

In particular, J_m^{n-1} from [Definition 3.2.1](#page-16-1) is $J_{\mathbf{a}}^{n, \leftarrow}$, where **a** is given by $\mathbf{a} =$ $\{0, m, 0, m, \ldots, 0, m\}.$

- (iv) $e_k \in I_m^n$ is such that $(e_k) = \delta_{jk}$, the Kroenecker delta. For $a, b \in I_m^n$, $a + b$ is such that $(a+b)_k = a_k + b_k$. For $A \subset I_m^n$ and $a \in I_m^n$, $A + a := \{b + a : b \in A\}$.
- (v) Let $\rho_{\epsilon}(\cdot): \{\rightarrow, \leftarrow\} \rightarrow \{-1, 1\}$ be the set function defined by

$$
\rho(\epsilon) = \begin{cases} -1, & \epsilon = \leftarrow \\ 1, & \epsilon = \rightarrow \end{cases}
$$

We will often refer to $\rho(\epsilon)$ as ρ_{ϵ} .

(vi) For every element $\beta \in I_m^n$, let $Nbh(\beta) := \{(\gamma, \beta) \mid (\gamma, \beta) \in \mathcal{I}_m^n\}.$

When we move a plate one step, this plate will leave a copy of the plate in the original position, as well as shifting a second copy one step. The next lemma shows that this process is bornologous.

Lemma 3.2.15. Let X be a semi-coarse space, let $f: I_m^n \to (X, V)$ be a bornologous function, and let \mathcal{P}_a be a plate on n^* delimited by a . Suppose that $\epsilon \in \{\leftarrow, \rightarrow\}.$ Suppose that, for every $\alpha \in \mathcal{P}_a$, $f(\alpha)$ is adjacent to every point in the set $\{f(\beta) \mid$ β is adjacent to $\alpha + \rho_{\epsilon} e_{n^*}$, and define

$$
f_{\mathcal{P}_{\mathbf{a}}}^{\epsilon}(\beta) \coloneqq \begin{cases} f(\beta - \rho_{\epsilon}e_{n^*}) & \text{if } \beta \in \mathcal{P}_{\mathbf{a}} + \rho_{\epsilon}e_{n^*} \\ f(\beta) & \text{if } \beta \notin \mathcal{P}_{\mathbf{a}} + \rho_{\epsilon}e_{n^*}. \end{cases}
$$

Then $f_{\mathcal{P}_a}^{\epsilon}: I_m^n \to X$ bornologous and $f \simeq_{sc} f_{\mathcal{P}_a}^{\epsilon}$.

Proof. We show that $f_{\mathcal{P}_a}^{\epsilon}$ is bornologous. We check that $f_{P_a}^{\epsilon}$ sends adjacent points in I_m^n to adjacent points in X.

First, let $\beta \notin \mathcal{P}_\mathbf{a} \cup (\mathcal{P}_\mathbf{a} + \rho_\epsilon e_{n^*}) \cup (\mathcal{P}_\mathbf{a} + 2\rho_\epsilon e_{n^*})$. Then $f_{\mathcal{P}_\mathbf{a}}^{\epsilon}|_{\text{Nbh}(\beta)} = f|_{\text{Nbh}(\beta)}$.

Now suppose that $\beta \in \mathcal{P}_a$.

$$
f_{\mathcal{P}_{\mathbf{a}}}^{\epsilon}(\mathrm{Nbh}(\beta)) = f_{\mathcal{P}_{\mathbf{a}}}^{\epsilon}(\{(\gamma,\beta) \in \mathrm{Nbh}(\beta) \mid \gamma \in \mathcal{P}_{\mathbf{a}} + \rho_{\epsilon}e_{n^*}\})
$$

$$
\cup f_{\mathcal{P}_{\mathbf{a}}}^{\epsilon}(\{(\gamma,\beta) \in \mathrm{Nbh}(\beta) \mid \gamma \notin \mathcal{P}_{\mathbf{a}} + \rho_{\epsilon}e_{n^*}\})
$$

$$
= f(\{(\gamma - \rho_{\epsilon}e_{n^*}), \beta) \mid (\gamma,\beta) \in \mathrm{Nbh}(\beta), \gamma \in \mathcal{P}_{\mathbf{a}} + \rho_{\epsilon}e_{n^*}\})
$$

$$
\cup f(\{(\gamma,\beta) \in \mathrm{Nbh}(\beta) \mid \gamma \notin \mathcal{P}_{\mathbf{a}} + \rho_{\epsilon}e_{n^*}\})
$$

$$
\subset f(\mathrm{Nbh}(\beta)) \in \mathcal{V}.
$$

Next, let $\beta \in \mathcal{P}_a + \rho_{\epsilon} e_{n^*}$. Then

$$
\begin{aligned} f^{\epsilon}_{\mathcal{P}_{\mathbf{a}}}(\mathrm{Nbh}(\beta)) &= f^{\epsilon}_{\mathcal{P}_{\mathbf{a}}}(\{(\gamma, \beta) \in \mathrm{Nbh}(\beta) \mid \gamma \in \mathcal{P}_{\mathbf{a}} \cup \mathcal{P}_{\mathbf{a}} + 2\rho_{\epsilon}e_{n^*}\}) \\ & \cup f^{\epsilon}_{\mathcal{P}_{\mathbf{a}}}(\{(\gamma, \beta) \in \mathrm{Nbh}(\beta) \mid \gamma \in \mathcal{P}_{\mathbf{a}} + \rho_{\epsilon}e_{n^*}\}) \\ &= \{(f(\gamma), f(\beta - \rho_{\epsilon}e_{n^*})) \mid (\gamma, \beta) \in \mathrm{Nbh}(\beta), \gamma \in \mathcal{P}_{\mathbf{a}} \cup \mathcal{P}_{\mathbf{a}} + 2\rho_{\epsilon}e_{n^*}\} \\ & \cup \{(f(\gamma - \rho_{\epsilon}e_{n^*}), f(\beta - \rho_{\epsilon}e_{n^*})) \mid (\gamma, \beta) \in \mathrm{Nbh}(\beta), \gamma \in \mathcal{P}_{\mathbf{a}}\} \\ & \subset f(\mathrm{Nbh}(\beta - \rho_{\epsilon}e_{n^*})), \end{aligned}
$$

where the last set is in V by hypothesis. Then

Finally, let $\beta \in \mathcal{P}_a + 2\rho_{\epsilon}e_{n^*}$.

$$
\begin{aligned} f^{\epsilon}_{\mathcal{P}_{\mathbf{a}}}(\mathrm{Nbh}(\beta)) &= f^{\epsilon}_{\mathcal{P}_{\mathbf{a}}}(\{(\gamma, \beta) \in \mathrm{Nbh}(\beta) \mid \gamma \in \mathcal{P}_{\mathbf{a}} + \rho_{\epsilon} e_{n^*}\}) \\ &\quad \cup f^{\epsilon}_{\mathcal{P}_{\mathbf{a}}}(\{(\gamma, \beta) \in \mathrm{Nbh}(\beta) \mid \gamma \notin \mathcal{P}_{\mathbf{a}} + \rho_{\epsilon} e_{n^*}\}) \\ &= \{ (f(\gamma - \rho_{\epsilon} e_{n^*}), f(\beta)) \mid (\gamma, \beta) \in \mathrm{Nbh}(\beta), \gamma \in \mathcal{P}_{\mathbf{a}} + \rho_{\epsilon} e_{n^*} \} \\ &\quad \cup f(\{(\gamma, \beta) \in \mathrm{Nbh}(\beta) \mid \gamma \notin \mathcal{P}_{\mathbf{a}} + \rho_{\epsilon} e_{n^*}\}) \\ &\subset f(\mathrm{Nbh}(\beta)), \end{aligned}
$$

where the last set is in V by hypothesis. Since I_m^n is a roofed semi-coarse space and has a finite number of elements, the above verifications imply that $f_{P_{\bf a}}^{\epsilon}$ is bornologous by [Proposition 2.2.4.](#page-7-2)

It remains to show that $f \simeq_{sc} f_{\mathcal{P}_a}^{\epsilon}$. Let $H: I_m^n \times \mathbb{Z} \to X$ such that $H(\cdot, k) = f(\cdot)$ for every $k \leq 0$ and $H(\cdot, k) = f_{\mathcal{P}_a}^{\epsilon}(\cdot)$ for every $k \geq 1$. We wish to show that H is bornologous. By [Proposition 2.2.4,](#page-7-2) it is enough to show that the restrictions of H to $I_m^n \times \{z \leq 0\}$, $I_m^n \times \{0,1\}$, and $I_m^n \times \{z \geq 1\}$ are bornologous. By definition, $H(\mathcal{I}_{m}^{n}\boxtimes \{z \leq 0\}^{2}) = f(\mathcal{I}_{m}^{n})$ and $H(\mathcal{I}_{m}^{n}\boxtimes \{z \geq 1\}^{2}) = f_{\mathcal{P}_{a}}^{\epsilon}(\mathcal{I}_{m}^{n})$, so H is bornologous on these regions. It remains to check the behavior of H on $I_m^n \times \{0,1\}$. Since $I_m^n\times \{0,1\}$ has a finite number of elements, we proceed as above by checking that ${\cal H}$ sends adjacent elements.

Note that $H(\mathcal{I}_m^n \boxtimes (0,0)) = f(\mathcal{I}_m^n)$ and $H(\mathcal{I}_m^n \boxtimes (1,1)) = f_{\mathcal{P}_a}^{\epsilon}(\mathcal{I}_m^n)$, so H is also bornologous on these regions. Therefore, it only remains to check H on $\mathcal{I}_m^n \boxtimes (0,1)$. Let $\beta \in I^n_m$. We consider the following two cases separately: either $\beta \in \mathcal{P}_a + \rho_{\epsilon} e_{n^*}$ or $\beta \notin \mathcal{P}_a + \rho_{\epsilon} e_{n^*}$.

- (1) Case 1: $\beta \in \mathcal{P}_a + \rho_{\epsilon} e_{n*}$. Then $H(\text{Nbh}(\beta) \boxtimes (0,1)) = \{ (f(\gamma), f_{\mathcal{P}_a}^{\epsilon}(\beta)) \mid$ $(\gamma, \beta) \in \text{Nbh}(\beta)$ = $\{(f(\gamma), f(\beta - \rho_{\epsilon}e_{n^*})) \mid (\gamma, \beta) \in \text{Nbh}(\beta)\}\in \mathcal{V}$, which was shown in the first part of the proof.
- (2) Case 2: $\beta \notin \mathcal{P}_a + \rho_{\epsilon} e_{n^*}$. Then $H(\mathrm{Nbh}(\beta) \boxtimes (0,1)) = \{ (f(\gamma), f_{\mathcal{P}_a}^{\epsilon}(\beta)) \mid$ $(\gamma, \beta) \in \text{Nbh}(\beta)$ } = { $(f(\gamma), f(\beta)) | (\gamma, \beta) \in \text{Nbh}(\beta)$ } $\in \mathcal{V}$.

From this, we conclude that the image of H of the roof of $\mathcal{I}_m^n \boxtimes \mathcal{Z}$ is controlled, so H is bornologous. \Box **Lemma 3.2.16** (Displacement with wake of a block.). Let (X, V) be a semi-coarse space, $f: I_m^n \to X$ a bornologous function, let B be a block delimited by a, and let $n^* \in \{1, \cdots, n\}$. Define

$$
f_{\mathcal{B}}^{\epsilon}(\beta) = \begin{cases} f(\beta - \rho_{\epsilon}e_{n^*}) & \text{if } \beta \in \mathcal{B} + \rho_{\epsilon}e_{n^*} \\ f(\beta) & \text{if } \beta \notin \mathcal{B} + \rho_{\epsilon}e_{n^*} \end{cases}
$$

Then $f^*: I_m^n \to X$ is bornologous and $f^* \simeq_{sc} f$ if for each $\alpha \in J_{\mathbf{a}}^{n^*, \epsilon}$, $f(\alpha)$ is adjacent to the image under f for each adjacent element to $\alpha + \rho_{\epsilon}e_{n^*}$ which are not in B.

Proof. Let (X, V) , $f: I_m^n \to X$, \mathcal{B} , and $n^* \in \{1, ..., n\}$ be as in the statement of the lemma. For $j \in \{a_{n^*}^{\leftarrow}, a_{n^*}^{\leftarrow} + 1, \ldots, a_{n^*}^{\rightarrow}\},\$ define $f_j: I_m^n \rightarrow X$ by

$$
f_j(\beta) = \begin{cases} f_{j+1}(\beta - \rho_{\epsilon}e_{n^*}) & \text{if } \beta \in \mathcal{B}_{n^*}^j + \rho_{\epsilon}e_{n^*} \\ f_{j+1}(\beta) & \text{if } \beta \notin \mathcal{B}_{n^*}^j + \rho_{\epsilon}e_{n^*} \end{cases}
$$

where $f_{a_{n^*} \to 1} := f$.

We proceed by induction.

Base case: By hypothesis, for each $\alpha \in \mathcal{B}_{n^*}^{\alpha_{n^*}^*}$, $f(\alpha)$ is adjacent to the image under f of adjacent elements of $\alpha + \rho_{\epsilon} e_{n^*}$. By [Lemma 3.2.15,](#page-20-0) $f_{a_{n^*}^{\epsilon}}$ is bornologous and homotopy equivalent to f .

Inductive step: Suppose that f_{J+1} is bornologous and homotopy equivalent to f. We show that this implies that the same is true for f_J . Note that for $\alpha \in \mathcal{B}_{n^*}^{J+2}$ we have that $f_{J+1}(\alpha) = f_{J+1}(\alpha - \rho_{\epsilon}e_{n^*}) = f_J(\alpha - \rho_{\epsilon}e_{n^*})$, and therefore f_J restricted to $\mathcal{B}_{n^*}^J$ satisfies [Lemma 3.2.15.](#page-20-0) It follows that f_J is bornologous and $f_J \simeq_{sc} f_{J+1} \simeq_{sc} f.$ □

We now use the above to move blocks a finite number of steps.

Lemma 3.2.17 (Displacement with wake k-steps of a block.). Let (X, V) be a semi-coarse space, $f: I_m^n \to X$ a bornologous function, $\mathcal B$ be a block delimited by $\mathcal{A}, k \in \mathbb{N}$ and $n^* \in \{1, \cdots, n\}.$

Define

$$
f_{\mathcal{B}}^{\epsilon}(\beta) = \begin{cases} f(\beta - k\rho_{\epsilon}e_{n^*}) & \text{if } \beta \in \mathcal{B} + k\rho_{\epsilon}e_{n^*} \\ f(\beta - v\rho_{\epsilon}e_{n^*}) & \text{if } \beta \in \mathcal{B}_{n^*}^{\alpha_{n^*}} + v\rho_{\epsilon}e_{n^*}, \ v \in \{1, \dots, k-1\} \\ f(\beta) & \text{otherwise} \end{cases}
$$

Then $f_{\mathcal{B}}^{\epsilon}: I_{m}^{n} \to X$ is bornologous and $f_{\mathcal{B}}^{\epsilon} \simeq_{sc} f$ if for each $\alpha \in J_{\mathbf{a}}^{n^{*}, \epsilon}$ we have that $f(\alpha)$ is adjacent to the image under f of the set

$$
\bigcup_{v \in \{1, \ldots, k-1\}} Nbh(\alpha + v\rho_{\epsilon}e_{n^*}) \qquad \qquad \text{if } \alpha \in \mathcal{B}_{\mathbf{a}}^{n^*, a_{n^*}^{\epsilon}} \cup \bigcup_{v \in \{1, \ldots, k-1\}} Nbh(\alpha + v\rho_{\epsilon}e_{n^*}) - \bigcup_{v \in \{1, \ldots, k-1\}} (\mathcal{B} + v\rho_{\epsilon}e_{n^*}) \qquad \text{if } \alpha \notin \mathcal{B}_{\mathbf{a}}^{n^*, a_{n^*}^{\epsilon}}.
$$

Proof. Let (X, V) , $f: I_m^n \to X$, \mathcal{B} , $k \in \mathcal{N}$, and $n^* \in \{1, ..., n\}$ be as in the hypothesis of the lemma. Assume that for each $\alpha \in J_{\mathbf{a}}^{n^*,\epsilon}$ we have that $f(\alpha)$ is adjacent to the image under f of the adjacent elements of $\alpha + v e_{n^*}, v \in \{1, \dots, w\}$.

For $j \in \{0, \ldots, k-1\}$, we define

$$
\mathcal{B}^{\langle j \rangle} = \mathcal{B} + j \rho_{\epsilon} e_{n^*}
$$

$$
f_{j+1} := \begin{cases} f_j(\beta - \rho_{\epsilon} e_{n^*}) & \text{if } \beta \in \mathcal{B}^{\langle j \rangle} + \rho_{\epsilon} e_{n^*} \\ f_j(\beta) & \text{if } \beta \notin \mathcal{B}^{\langle j \rangle} + \rho_{\epsilon} e_{n^*} \end{cases}
$$

Let $f_0 := f$ and observe that $\mathcal{B}^{\langle 0 \rangle} = \mathcal{B}$. We proceed by induction.

Base case: Observe that, for $v = 1$, the functions f, f_1 and the block B satisfy the hypotheses of [Lemma 3.2.16,](#page-22-1) from which it follows that f_1 is bornologous and $f_1 \simeq_{sc} f$.

Inductive step: Assume that f_{J-1} is bornologous and homotopy equivalent to f. Observe that, if $v = J$, then f_{J-1} , f_J and $\mathcal{B}^{\langle J-1\rangle}$ satisfy the hypotheses of [Lemma 3.2.16,](#page-22-1) from which it follows that f_J is bornologous and $f_J \simeq_{sc} f_{J-1}$. Since $f_{\mathcal{B}}^{\epsilon} = f_w$, we conclude that $f_{\mathcal{B}}^{\epsilon}$ is bornologus and $f_{\mathcal{B}}^{\epsilon} \simeq_{sc} f$.

The above lemmas now allow us to prove the following theorem. The proof is analogous to that in the topological case, replacing the *n*-disk $Dⁿ$ with the blocks I^n_m .

Theorem 3.2.18 (Semi-coarse Homotopy Groups). Let (X, Y) be a semi-coarse space, $A \subset X$, and $n \in \mathbb{N}$.

- If $n \geq 1$, then $(\pi_n^{sc}(X, *), \star)$ is a group.
- If $n \geq 2$, then $(\pi_n^{sc}(X, *), *)$ is an abelian group.
- If $n \geq 2$, then $(\pi_n^{sc}(X, A, *), *)$ is a group.
- If $n \geq 3$, then $(\pi_n^{sc}(X, A, *), \star)$ is an abelian group.

Proof. Let (X, V) be a semi-coarse space, let $n \geq 2$, and suppose that there are m, m', m'' such that $f : (I_m^n, \partial I_m^n, J_m^{n-1}) \to (X, A, *), g : (I_{m'}^n, \partial I_{m'}^n, J_{m'}^{n-1}) \to$ $(X, A, *)$ and $h: (I^n_{m''}, \partial I^n_{m''}, J^{n-1}_{m''}) \rightarrow (X, A, *)$. Recall from [Definition 3.2.8,](#page-18-3) that $(f \star g) \star h = f \star (g \star h)$. Thus, by [Lemma 3.2.12,](#page-19-0)

$$
\langle f \star g \rangle \star \langle h \rangle = \langle (f \star g) \star h \rangle = \langle f \star (g \star h) \rangle = \langle f \rangle \star \langle g \star h \rangle,
$$

so the product \star is associative.

Define $e: I_0^n \to X$ to be the constant map $e(x) = *$, and consider the function $f:(I_m^n, \partial I_m^n, J_m^{n-1}) \to (X, A, *),$ a bornologous map of triples. By [Definition 3.2.8](#page-18-3) of \star , $f \star e = f = e \star f$. Thus, $\langle f \star e \rangle = \langle f \rangle = \langle e \star f \rangle$.

We now show that every bornologous map $f: (I_m^n, \partial I_m^n, J_m^{n-1}) \rightarrow (X, A, *)$ has a homotopy inverse. Define $g: (I_m^n, \partial I_m^n, J_m^{n-1}) \to (X, A, *)$ to be the map $g(\alpha) = f(\beta)$, where $\alpha, \beta \in I_m^n$ and the coordinates of α and β satisfy

$$
\alpha_1 = m - \beta_1,
$$

\n
$$
\alpha_i = \beta_i \text{ for } i \neq 1.
$$

We show that $\langle f \star g \rangle = \langle e \rangle$ by induction, and we remark that we can proceed in the same way to show that $\langle g * f \rangle = \langle e \rangle$. Define $f_m := f * g$, and define $f_j: (I_{2m}^n, \partial I_{2m}^n, J_{2m-1}^n) \to (X, A, *)$ for $j \in \{1, \ldots, m-1\}$ recursively by

$$
f_{j-1}(\alpha) := \begin{cases} f_j(\alpha), & \text{if } 0 \le \alpha_1 \le j-1 \\ f_j(-\alpha + 2(j-1)e_1), & \text{if } j \le \alpha_1 \le 2(j-1) \\ \ast, & \text{elsewhere.} \end{cases}
$$

Let $\mathcal{B}^{\langle j \rangle}$ the block delimited by

$$
\mathbf{a}^{\langle j \rangle} := (j+1, 2j, 0, 2m, \dots, 0, 2m) \in I_{2m}^n
$$

Base case: f_{m-1} is a bornologous function and homotopy equivalent to $f' \star (f')^{-1}$ by [Lemma 3.2.17,](#page-22-0) taking $\mathcal{B} = \mathcal{B}^{\langle m \rangle}$, $k = 2$ and $f = f' \star (f')^{-1}$. To be able to use that lemma, observe that for all α such that $\alpha_1 = m + 1$, $f(\alpha)$ is adjacent to the image of elements adjacent to $\alpha - e_1$ (which is *) and to the image of elements adjacent to $\alpha - 2e_1$.

Induction step: Suppose that f_i is a bornologous function and that it is homotopy equivalent to f_{j+1} . By [Lemma 3.2.17,](#page-22-0) f_{j-1} is bornologus and homotopy equivalent to f_j , taking $\mathcal{B} = \mathcal{B}^{\langle j \rangle}$, $k = 2$, and $f = f_j$ in the lemma. Also observe that, for all α such that $\alpha_1 = j + 1$, $f_j(\alpha)$ is adjacent to the image of elements adjacent to $\alpha - e_1$ and $\alpha - 2e_1$.

The above implies that $f \star g \simeq_{sc} \ast$, and therefore $(\pi_n^{sc}(X, A, \ast), \star)$ is a group.

Now suppose, in addition, that $n \geq 3$, and let $\langle f \rangle$, $\langle g \rangle \in \pi_n^{sc}(X, A, *)$ such that m and m' are integers with $f: (I_m^n, \partial I_m^n, J_m^{n-1}) \to (X, A, *)$ and $g: (I_{m'}^n, \partial I_{m'}^n, J_{m'}^{n-1}) \to$ $(X, A, *)$.

For $i \in \{1, 2, 3, 4, 5\}$, let $\mathcal{B}^{\langle i \rangle}$ be the blocks delimited by $\mathbf{a}^{\langle i \rangle}$, where

$$
\mathbf{a}^{\langle 1 \rangle} := (m+1, m+m'-1, 0, m'-1, 0, m+m', \dots, 0, m+m')
$$

\n
$$
\mathbf{a}^{\langle 2 \rangle} := (m+1, m+m', m+1, m'-1, 0, m+m', \dots, 0, m+m')
$$

\n
$$
\mathbf{a}^{\langle 3 \rangle} := (1, m'-1, m+1, m+m', 0, m+m', \dots, 0, m+m')
$$

\n
$$
\mathbf{a}^{\langle 4 \rangle} := (0, m-1, 1, m-1, 0, m+m', \dots, 0, m+m')
$$

\n
$$
\mathbf{a}^{\langle 5 \rangle} := (m'+1, m+m'-1, 1, m-1, 0, m+m', \dots, 0, m+m').
$$

Define

$$
f_1(\beta) := \begin{cases} f \star g(\beta - me_2) & \text{if } \beta \in \mathcal{B}^{(1)} + me_2 \\ f \star g(\beta - ve_2) & \text{if } \beta \in (\mathcal{B}^{(1)})_2^0 + ve_2, \ v \in \{1, \dots, m-1\} \\ f \star g(\beta) & \text{anywhere else} \end{cases}
$$

Observe that $f \star g((\mathcal{B}^{\langle 1 \rangle} \cup \mathcal{B}^{\langle 4 \rangle})^c) = \{*\},\$ then by [Lemma 3.2.17](#page-22-0) we have that f_1 is bornologous and $f_1 \simeq_{sc} f \star g$. Let's define

$$
f_2(\beta) := \begin{cases} f_1(\beta + me_1) & \text{if } \beta \in \mathcal{B}^{(2)} - me_1 \\ f_1(\beta + ve_1) & \text{if } \beta \in (\mathcal{B}^{(2)})_1^{m+m'} - ve_1, \ v \in \{1, ..., m-1\} \\ f_1(\beta) & \text{anywhere else} \end{cases}
$$

Observe that $f_1((\mathcal{B}^{\langle 2 \rangle} \cup \mathcal{B}^{\langle 4 \rangle})^c) = \{*\}$, then by [Lemma 3.2.17](#page-22-0) we have that f_2 is bornologous and $f_2 \simeq_{sc} f_1$. Let's define

$$
f_3(\beta) := \begin{cases} f_2(\beta - m'e_1) & \text{if } \beta \in \mathcal{B}^{\langle 4 \rangle} + m'e_1 \\ f_2(\beta - ve_1) & \text{if } \beta \in (\mathcal{B}^{\langle 4 \rangle})_1^0 + ve_1, \ v \in \{1, ..., m' - 1\} \\ f_2(\beta) & \text{anywhere else} \end{cases}
$$

Observe that $f_2((\mathcal{B}^{\langle 3 \rangle} \cup \mathcal{B}^{\langle 4 \rangle})^c) = \{*\}$, then by [Lemma 3.2.17](#page-22-0) we have that f_3 is bornologous and $f_3 \simeq_{sc} f_2$. Let's define

$$
f_4(\beta) := \begin{cases} f_3(\beta + me_2) & \text{if } \beta \in \mathcal{B}^{(3)} - me_2 \\ f_3(\beta + ve_2) & \text{if } \beta \in (\mathcal{B}^{(3)})_2^{m+m'} - ve_2, v \in \{1, ..., m-1\} \\ f_3(\beta) & \text{anywhere else} \end{cases}
$$

Observe that $f_3((\mathcal{B}^{(3)} \cup \mathcal{B}^{(5)})^c) = \{*\}$, then by [Lemma 3.2.17](#page-22-0) we have that f_4 is bornologous and $f_4 \simeq_{sc} f_3$. Finally, note that $f_4 = g * f$, and we conclude that $\pi_n^{sc}(X, A, *)$ is commutative.

If $A = \{*\}$, then we have shown that $\pi_n^{sc}(X, *)$ is a group for $n \geq 2$ and abelian for $n \geq 3$. Consider the case where $n = 1$, and let $\langle f \rangle, \langle g \rangle, \langle h \rangle \in \pi_1^{sc}(X, *)$. Then there exist non-negative integers m, m', m'' such that $f : (I_m, \partial I_m) \to (X, *), g$: $(I_{m'}, \partial I_{m'}) \rightarrow (X, *)$ and $h : (I_{m''}, \partial I_{m''}) \rightarrow (X, *)$. By definition, $(f * g) * h =$ $f \star (g \star h)$, and therefore, by [Lemma 3.2.12,](#page-19-0)

$$
[\langle f' \star g' \rangle] \star [\langle h' \rangle] = [\langle (f \star g) \star h \rangle] = [\langle f \star (g \star h) \rangle] = [\langle f' \rangle] \star [\langle g' \star h' \rangle],
$$

so \star is associative.

Define $e: I_0 \to X$ to be the map which sends I_0 to *. Therefore, $\langle e \rangle \in \pi_1^{sc}(X,*)$. Let $\langle f \rangle \in \pi_1^{sc}(X, *),$ so there exists a non-negative integer m with $f: (I_m^n, \partial I_m^n) \to$ $(X, *)$. By definition of \star , $f \star e = f = e \star f$. Thus, $\langle f \star e \rangle = \langle f \rangle = \langle e \star f \rangle$ and $\langle e \rangle$ is the identity in $\pi_1^{sc}(X,*)$.

Let $\langle f \rangle \in \pi_1^{sc}(X,*)$. Then there exists a non-negative integer m with f: $(I_m, \partial I_m) \to (X, *)$. Define $(f')^{-1}$ such that $(f')^{-1}(\alpha) = f'(\beta)$ with $\alpha = m - \beta$.

We show that $\langle f' \star (f')^{-1} \rangle = \langle e \rangle$ by mathematical induction and remark that we can proceed in the same way for $\langle (f')^{-1} \star f' \rangle = \langle e \rangle$. Define $f_j : I_{2m} \to X$ for $j \in \{1, \ldots, m\}$ by

$$
f_{j-1}(\alpha) := \begin{cases} f'_j(\alpha) & \text{if } 0 \le \alpha \le j-1 \\ f'_j(-\alpha + 2(j-1)e_1) & \text{if } j \le \alpha \le 2(j-1) \\ * & \text{anywhere else} \end{cases}
$$

with $f_m = f' \star (f')^{-1}$, and observe that $f_{j-1} : (I_{2m}, \partial I_{2m}) \to (X, *)$ is bornologous. Let $\mathcal{B}^{\langle j \rangle}$ be the block delimited by

$$
\mathbf{a}^{\langle j \rangle} := (j+1, 2j).
$$

Base case: f_{m-1} is a bornologous function and homotopy equivalent to f' \star $(f')^{-1}$ by [Lemma 3.2.17](#page-22-0) with $\mathcal{B} = \mathcal{B}^{\langle m \rangle}$, $w = 2$ and $f = f' \star (f')^{-1}$. Observe, in addition, that for $\alpha = m + 1$, $f(\alpha)$ is adjacent to the image of elements adjacent to $\alpha - e_1$ (which is *) and to the image of elements adjacent to $\alpha - 2e_1$.

Induction step: Suppose that f_j is a bornologous function and homotopy equivalent to f_{j+1} . By [Lemma 3.2.17,](#page-22-0) f_{j-1} is bornologous and homotopy equivalent to f_j , taking $\mathcal{B} = \mathcal{B}^{(j)}$, $w = 2$, and $f = f_j$ in the lemma. Observe that, for $\alpha = j+1$, $f_j(\alpha)$ is adjacent to the image of elements adjacent to $\alpha - e_1$ and $\alpha - 2e_1$.

We conclude that $(\pi_1^{sc}(X, *), \star)$ is a group.

Finally, if $n = 2$, then we may show that $\pi_2^{sc}(X, *)$ is abelian by repeating the above proof of this fact for $\pi_n^{sc}(X, A, *)$ for $n \geq 3$. Observe that the proof is not valid for $\pi_2^{sc}(X, A, *)$ because $f \star g(\alpha) \in A$ when $\alpha_2 = 0$. However, if $A = \{*\}$, we have that $f \star g(\alpha) = \ast$ when $\alpha_2 = 0$.

In the case of coarse spaces, the next theorem shows that the semi-coarse homotopy groups are trivial.

Theorem 3.2.19. Let (X, V) be a coarse space with $A \subset X$. Then for any $n \in \mathbb{N}$, we have $\pi_n^{sc}(X, A, *) \cong \pi_n^{sc}(X, *) \cong \{1\}.$

Proof. Let (X, V) be a coarse space, $n \in \mathbb{N}$, and suppose that $h: I_m^n \to X$ is the constant map $h(\alpha) = *$ for all $\alpha \in I_m^n$.

Let $\langle f \rangle \in \pi_n^{sc}(X, A, *)$. By definition, $f(J^{n-1}) = \{ * \}$, in particular $f(\alpha) = *$ when the first coordinate $\alpha_1 = 0$. Note that, for all $\alpha \in I_m^n$ satisfies that $f(\alpha - ke_1)$ is adjacent to $f(\alpha - (k+1)e_1)$ for any $k \in \{0, \ldots, \alpha_1 - 1\}$, that is

$$
\{(f(\alpha), f(\alpha - e_1)), (f(\alpha - e_1), f(\alpha - 2e_1)), \dots, (f(\alpha - (\alpha_1 - 1)e_1, f(\alpha - \alpha_1e_1)))\}\in \mathcal{V}
$$

because f is bornologous. Thereby, since (X, V) is a coarse space and $f(\alpha - \alpha_1 e_1) =$ ^{*}, we conclude that ${(f(α), *)} ∈ V$ for every $α ∈ Iⁿ_m$, and therefore

$$
f \simeq_{sc} *
$$

Therefore, $\pi_n^{sc}(X, A, *) \cong \{1\}$. The proof for $\pi_n^{sc}(X, *)$ is analogous. \Box

While the above theorem says that semi-coarse homotopy groups are trivial for coarse spaces, we expect there to be other semi-coarse invariants, perhaps defined relative to infinity, which are non-trivial for both coarse and non-coarse semi-coarse spaces. However, we leave this question open for future work.

3.3. Connectedness.

Definition 3.3.1 (Semi-Coarse *n*-Connected Space). Let (X, V) be a semi-coarse space. We will say that (X, V) is an *n*-connected (semi-coarse) space iff $\pi_n^{sc}(X, *) \cong$ {0}. In particular, a 0-connected space will be called a connected (semi-coarse) space.

In the next proposition we will observe that, when the semi-coarse structure is coarse, then semi-coarse connectedness is equivalent to coarse connectedness. We first recall the notion of coarse connectedness. See also [\[14\]](#page-55-2) for more details on coarse connectedness.

Definition 3.3.2 (Coarse connected space). Let (X, V) be a coarse space. We say that (X, V) is coarsely connected iff every point in $X \times X$ belongs to some controlled set $V \in \mathcal{V}$.

Proposition 3.3.3. Let (X, V) be a coarse space. Then (X, V) is coarsely connected iff $\pi_0^{sc}(X, *; \mathcal{V}) \cong \{0\}.$

Proof. Let (X, V) be a coarse space.

(←) If (X, V) is coarsely connected, then for each $(x, y) \in X \times X$ there exists $E_{x,y} \in \mathcal{V}$ such that $(x, y) \in E_{x,y}$, that is, $\{(x, y)\}\in \mathcal{V}$ for any $x, y \in X$. Let f : $(\{*,1\},*,\mathcal{D}_{\{*,1\}}) \to (X,*,\mathcal{V})$ and $g: (\{*,1\},*,\mathcal{D}_{\{*,1\}}) \to (X,*,\mathcal{V})$ be bornologous maps. Note that, since $\mathcal{D}_{\{\ast,1\}}$ is the diagonal semi-coarse structure, $f(1)$ and $g(1)$ may be arbitrary points of X. Define $H : \{*, 1\} \times \mathbb{Z} \to X$ such that $H(\alpha, n) = f(\alpha)$ when $n \leq 0$ and $H(\alpha, n) = g(\alpha)$ when $n \geq 1$. Since X is coarsely connected, we have that $\{(f(1), g(1))\}\in V$, and therefore H is bornologous. It follows that $f \simeq_{sc} g$. However, f and g were arbitrary, and we conclude that $\pi_0^{sc}(X,*) \cong \{0\}.$

(\Rightarrow) If (X, V) is a coarse space with $\pi_0^{sc}(X, *; V) = \{0\}$, then for each f: $({\{*,1\}}, *) \rightarrow (X, *)$ and $g : ({\{*,1\}}, *) \rightarrow (X, *)$ there exist $N, M \in \mathbb{Z}$ with $N < M$ and a bornologous function $H: \{*,1\} \times \mathbb{Z} \to X$ such that $H(e, z) = f(z)$ when $z \leq N$ and $H(e, z) = g(z)$ when $z \geq M$. Given that H is bornologous, then $H(1, z)$ is adjacent to $H(1, z + 1)$ for each $z \in \mathbb{Z}$. Therefore

$$
\{(f(1), H(1, N+1))\}, \{(H(1, N+1), H(1, N+2))\}, \dots \{(H(1, M-1), g(1))\} \in \mathcal{V}
$$

Then, $\{(f(1), g(1))\}\in \mathcal{V}$. How f and g are arbitrary, $f(1)$ and $g(1)$ are any element of x. Thus, for each $(x, y) \in X \times X$, $\{(x, y)\}\in \mathcal{V}$.

The highlight in this section is being able to note that the base point does not matter when we have a connected space. We are going to prove a lemma which makes that easy.

Lemma 3.3.4. Let (X, V) be a semi-coarse space. If $x, y \in X$ are adjacent elements, then $\pi_n^{sc}(X, x) \cong \pi_n^{sc}(X, y)$ for $n \geq 1$.

Proof. Let (X, V) be a semi-coarse space, $n \geq 1$ and $x, y \in X$ be adjacent elements. Let $\vert \langle f \rangle \vert : (I^n, \partial I^n) \to (X, x)$, then there exists a non-negative integer m such that $f': (I_m^n, \partial I_m^n) \to (X, x)$ and $f' \in \langle f \rangle$. By definition, $(\phi_m^{m+2})^n f' \in \langle f \rangle$.

Let's define β the block delimited by $\mathcal A$ such that $a_i^{\leftarrow} = 1$ and $a_i^{\rightarrow} = m - 1$ and

$$
f_1(\beta) := \begin{cases} (\phi_m^{m+2})^n f'(\beta - \sum_{i=1}^n e_i) & \text{if } \beta \in \mathcal{B} + \sum_{i=1}^n e_i \\ x & \text{anywhere else} \end{cases}
$$

We get that $(\phi_m^{m+2})^n f'$ is bornologous and $(\phi_m^{m+2})^n f' \simeq_{sc} f_1$. Let's define

$$
f_2(\beta) := \begin{cases} y & \text{if } \beta \in \partial I_{m+2}^n \\ f_1(\beta) & \text{anywhere else} \end{cases}
$$

which is bornologous because x is adjacent to y .

Observe that we get a function $\Psi : \pi_n^{sc}(X, x) \to \pi_n^{sc}(X, y)$ which is well-defined by $(\phi_m^{m+2})^n f' \simeq_{sc} f_1$. We need to prove that this is a group homomorphism.

Let $[\langle f \rangle], [\langle g \rangle] \in \pi_n^{sc}(X, x)$, then there exist non-negative integers m, m' such that $f': (I_m^n, \partial I_m^n) \to (X, x), g': (I_{m'}^n, \partial I_{m'}^n) \to (X, x), f' \in \langle f \rangle$ and $g' \in \langle g \rangle$. We just need to prove that $\langle \Psi(f' \star g') \rangle \simeq_{sc} \langle \Psi(f') \star \Psi(g') \rangle$. Define $\mathcal{B}^{\langle 1 \rangle}$ the block delimited by $\mathcal{A}^{\langle 1 \rangle}$ such that

$$
a_1^{\leftarrow} = m + m' + 2, \ a_1^{\rightarrow} = m + m' + 3
$$

$$
a_i^{\leftarrow} = 1, \ a_i^{\rightarrow} = m' + 1 \ i \neq 1
$$

and $\mathcal{B}^{\langle 2 \rangle}$ the block delimited by $\mathcal{A}^{\langle 2 \rangle}$ such that

$$
\begin{aligned} a_1^\leftarrow &= m+1, \ a_1^\rightarrow &= m+m'+1 \\ a_i^\leftarrow &= 1, \ a_i^\rightarrow &= m'+1 \ i \neq 1 \end{aligned}
$$

Let's define $h_1: I_{m+m'+4} \to X$ such that

$$
h_1(\beta) := \begin{cases} x & \text{if } \beta \in \mathcal{B}^{\langle 1 \rangle} \\ (\phi_{m+m'+2}^{m+m'+4})^n \Psi(f' \star g')(\beta) & \text{anywhere else} \end{cases}
$$

Note that $h_1 \simeq_{sc} (\phi_{m+m'+2}^{m+m'+4})^n \Psi(f' \star g')$. Let's define

$$
h_2(\beta) := \begin{cases} h_1(\beta - 2e_1) & \text{if } \beta \in \mathcal{B}^{\langle 2 \rangle} + 2e_1 \\ h_1(\beta - e_1) & \text{if } \beta \in (\mathcal{B}^{\langle 2 \rangle})_1^{m+1} + e_1 \\ h_1(\beta) & \text{anywhere else} \end{cases}
$$

which is a bornologous function and $h_1 \simeq_{sc} h_2$. Let's define $\mathcal{B}^{\langle 3 \rangle}$ and $\mathcal{B}^{\langle 4 \rangle}$ delimited by $\mathcal{A}^{\langle 3 \rangle}$ and $\mathcal{A}^{\langle 4 \rangle}$ such that

$$
\mathcal{A}^{\langle 3 \rangle} := \{ \alpha \in I_{m+m'+4}^n : 1 \le \alpha_i \le m+1 \}
$$

$$
\mathcal{A}^{\langle 4 \rangle} := \{ \alpha \in I_{m+m'+4}^n : m+3 \le \alpha_1 \le m+m'+3, 1 \le \alpha_i \le m'+1, i \ne 1 \}
$$

Let's define

$$
h_3(\beta) := \begin{cases} h_2(\beta) & \text{if } \beta \in \mathcal{B}^{\langle 3 \rangle} \cup \mathcal{B}^{\langle 4 \rangle} \\ y & \text{anywhere else} \end{cases}
$$

which is bornologous and $h_3 \simeq_{sc} h_4$ because x is adjacent to y. Finally, observe that $h_3 = \Psi(f' \star g')$, getting what we want. Thus, Ψ is a group homomorphism.

Under the same argument, we are able to define Θ : $\pi_n^{sc}(X, y) \to \pi_n^{sc}(X, x)$. So, we just need to prove that $\Psi^{-1} = \Theta$ and we will get our isomorphism.

Let $[\langle f \rangle] \in \pi_n^{sc}(X, x)$, then there exists a non-negative integer m such that f' : $(I_m, \partial I_m) \to (X, x)$ and $f' \in \langle f \rangle$. Since x and y are adjacent, we are able to replace every y for x in $\Theta \Psi(f')$ calling that function h, then h is a bornologous function, $h \simeq_{sc} \Theta \Psi(f')$ and $h \in \langle f' \rangle = \langle f \rangle$. Thus, $\Theta \Psi[\langle f \rangle] = [\langle f \rangle]$, that is, $\Theta \Psi = 1_{\pi_n^{sc}(X,x)}$. Under the same argument, $\Psi \Theta = 1_{\pi_n^{sc}(X,y)}$, getting that $\Psi^{-1} = \Theta$.

Theorem 3.3.5. Let (X, V) be a connected semi-coarse space. Then, for every $x, y \in X$ we get $\pi_n^{sc}(X, x) \cong \pi_n^{sc}(X, y)$.

Proof. Let (X, V) be a connected semi-coarse space and $x, y \in X$. Then, there exist $N, M \in \mathbb{Z}$ such that $N < M$ and a bornologous function $H: \{*, 1\} \times \mathbb{Z} \to X$ such that $H(*, z) = *$ when $z \in \mathbb{Z}$, $H(1, z) = x$ when $z \leq N$ and $H(1, z) = y$ when $z \geq M$. Since H is bornologous, then $\{(x, H(1, N+1), \{(H(1, N+1), H(1, N+1), \}(H(1, N+1)), H(1, N+1)\})$ 2)) $\}, \ldots \{(H(1, M-1), y)\} \in \mathcal{V}$. Thus,

$$
\pi_n^{sc}(X, x) \cong \pi_n^{sc}(X, H(1, N+1)) \cong \dots \cong \pi_n^{sc}(X, H(1, M-1)) \cong \pi_n^{sc}(X, y).
$$

Getting that
$$
\pi_n^{sc}(X, x) \cong \pi_n^{sc}(X, y).
$$

3.4. The Semi-Coarse Fundamental Group of Cyclic Graphs. In this section, we will compute the semi-coarse fundamental group of cyclic graphs with different structures, and, in particular, this will provide a class of examples where semi-coarse homotopy is non-trivial in dimensions greater than zero. (For the 0 homotopy class, we get that $\#(\pi_0^{sc}(X,*)) = \#X$, the number of semi-coarse connected components of X .)

Definition 3.4.1. Let $C_n := \{0, 1, 2, \ldots, n-1\}$ and m a positive integer. Let's define \mathcal{C}_n^m as the graph with vertices C_n and edges $(k, k-i)$ mod $n, (k, k+i)$ mod n for every $k \in C_n$ and $i \in \{1, 2, ..., m\}$. When $m = 1$, we call C_n^1 an *n*-cycle and write \mathcal{C}_n .

Example 3.4.1.1. For $n = 4, m = 1$ the resulting graph is

For the remainder of this section, we denote by $C_{n,m}$ the semi-coarse space (C_n, \mathcal{C}_n^m) , where C_n is given the semi-coarse structure induced by the graph \mathcal{C}_n^m . We denote by c to the map from I_{n+1} onto C_n such that $c(i) = i \mod(n)$, and by c^{-1} the map such that $c^{-1}(i) = c(n - i)$.

We will now define a class of functions which we can see as being endowed with an orientation. They will be particularly useful in the following.

Definition 3.4.2. We say that $f: I_k \to C_{n,m}$ is unidirectional if for every $i \in \{1, \ldots, k-2\}$ we have that $f(i)$ is not a neighbor of $f(i+2)$ (i.e. $\{(f(i), f(i+2))\}$ $2))\}\notin \mathcal{C}_n^m$).

Lemma 3.4.3. For every bornologous function $f: I_k \to C_n$ there exists a bornologous function $f': I_{k'} \to C_n$ such that $\langle f \rangle \simeq \langle f' \rangle$ and f' is unidirectional.

Proof. Assume f is not a unidirectional function, then there exists $i \in \{1, \ldots, k-2\}$ such that $f(i)$ is neighbor of $f(i+2)$. We are able to define $f_1 : I_k \to C_n$ such that $f_1(j) = f(j)$ with $j \neq i + 1$, $f_1(i + 1) = f(i + 2)$. Clearly $f_1 \simeq_{sc} f$.

We now define $\hat{f}_1: I_{k-1} \to C_n$ such that $\hat{f}_1(j) = f_1(j)$ if $j \leq i$, $\hat{f}_1(j) = f_1(j+1)$ if $j > i$. It is not difficult to show that $\langle f \rangle \simeq_{sc} \langle \hat{f}_1 \rangle$.

We repeat this process on f_1 , and so on, until one is left with a unidirectional function. Observe this is a finite process since each iteration reduces the length of th domain of f_1 by 1. \Box

Remark 3.4.4. If $\lceil \frac{n}{m} \rceil \leq 2$, then no function $f: I_k \to C_n$ is unidirectional for $k > 0$, because \mathcal{C}_n^m is a complete graph.

The following result shows that we are able to reduce the map c to a standard form.

Proposition 3.4.5. Let $\hat{c}: I_{\lceil \frac{n}{m} \rceil} \to C_n$ be the map $\hat{c}(k) = (k-1)m$ if $1 \leq i \leq \lceil \frac{n}{m} \rceil$, and $\hat{c}(\lceil \frac{n}{m} \rceil) = 0$. Then $\hat{c} \simeq_{sc} c$.

Proof. Define $\dot{c}: I_n \to C_n$ as follows

$$
\dot{c}(i) := \begin{cases}\n0 & 0 \leq i \leq m-1 \text{ or } i = n \\
m & m \leq i \leq 2m-1 \\
2m & 2m \leq i \leq 3m-1 \\
\vdots & \vdots \\
(\lceil \frac{n}{m} \rceil - 1)m & (\lceil \frac{n}{m} \rceil - 1)m \leq i \leq n-1\n\end{cases}
$$

The \mathcal{C}_n^m structure gives immediately that $c \simeq_{sc} \dot{c}$. It now follows that $\langle c \rangle \simeq_{sc} \langle \hat{c} \rangle$ \square

Our final proposition shows us that every unidirectional function has a standard form, up to homotopy.

Proposition 3.4.6. Every unidirectional function $f: I_k \to C_n$ is homotopy equivalent to a function of one of the following forms: 1, $c \star c \star ... \star c$ or $c^{-1} \star ... \star c^{-1}$.

Proof. We will assume that $\frac{n}{m} > 2$, otherwise we have complete graph which is homotopy equivalent a point, and the conclusion of the proposition is satisfied. We also assume that $1 \neq f$.

Let $f: I_k \to C_n$ be unidirectional. Then $f(0) = 0$, and either $f(1) \in \{1, ..., m\}$ or $f(1) \in \{n-1,\ldots,n-m\}$. We will work with the first case. The other case is analogous.

Let $\hat{f}_1: I_{k+f(1)} \to C_n$ be defined by

$$
\hat{f}_1(i) := \begin{cases} i & 1 \le i \le f(1) \\ f(i - f(1)) & f(1) < i \le f(1) + k \end{cases}
$$

We observe that $\langle f \rangle \simeq_{sc} \langle \hat{f}_1 \rangle$

We call the transformation of f into \hat{f}_1 a flattening of f from $i = 0$ to $i = 1$. We now flatten \hat{f}_1 from $f(1)$ to $f(1) + 1$, and we repeat this process until we arrive at a function \hat{f} such that $\hat{f}(i) - \hat{f}(i+1) = 1$. Note that this takes a finite number of steps, and that $\langle f \rangle \simeq_{sc} \langle \hat{f} \rangle$, which proves the result. \Box

Combining the above two propositions, we get that the first homotopy group of (C_n, \mathcal{C}_n^m) is isomorphic to a the fundamental group of a different cyclic graph with a simpler structure, i.e.

Corollary 3.4.7. $\pi_1^{sc}(C_n, \mathcal{C}_n^m) \cong \pi_1^{sc}(C_{\lceil \frac{n}{m} \rceil}, \mathcal{C}_{\lceil \frac{n}{m} \rceil}).$

Remark 3.4.8. Let (C_n, \mathcal{C}_n) be the *n*-cycle and $n \in \{1, 2, 3\}$, then it follows that \mathcal{C}_n is a connected coarse structure, precisely $\mathcal{C}_n = \mathcal{P}(C_n \times C_n)$. So, their homotopy groups are trivial.

In the rest of this subsection, we compute the fundamental group of (C_n, \mathcal{C}_n) .

Theorem 3.4.9. Let (C_n, C_n) be the semi-coarse space induced by the n-cycle graph. Then, $\pi_1^{sc}(C_n, \mathcal{C}_n) \cong \mathbb{Z}$.

Before the proof, we are going to need proving the following three lemmas, for which we define the function $p : (\mathbb{Z}, \mathcal{Z}) \to (C_n, C_n)$ by $p(k) := k \mod n$ This function will be called projection of the integers onto n-cycle or simply the projection. We note that p is a bornologous function.

Lemma 3.4.10. Let $f: I_k \to C_n$, $n \geq 4$, be a bornologous function with $f(0) = 0$. Then, for each $m \in p^{-1}(0)$ there exists a unique bornologous function $\hat{f}: I_k \to \mathbb{Z}$ such that $\hat{f}(0) = m$ and $f = p \circ \hat{f}$.

Proof. Let $f: I_k \to C_n$ be a bornologous function with $f(0) = 0$, and let $m \in$ $p^{-1}(x_0)$.

We begin by proving that if there are two bornologous functions $\hat{f}, \hat{g} : I_k \to$ $(\mathbb{Z}, \mathcal{Z})$ such that $\hat{f}(0) = \hat{g}(0) = m$ and $f = p \circ \hat{f} = p \circ \hat{g}$, then $\hat{f} = \hat{g}$. Given that $p \circ \hat{f} = p \circ \hat{g}$, then $\hat{f}(i) = \hat{g}(i) + n \cdot i$ for each $i \in \{0, \dots, k-1\}$. Also, by hypothesis, $\hat{f}(0) = \hat{g}(0)$. Since both functions are bornologous, we have that $\hat{f}(1) = \hat{f}(0) + i_0$ and $\hat{g}(1) = \hat{g}(0) + j_0$ with $i_0, j_0 \in \{-1, 0, 1\}$, and $i_0 \mod n = j_0$ mod n, which implies that $i_0 = j_0$. We conclude that $\hat{f}(1) = \hat{g}(1)$. Inductively, if $\hat{f}(i-1) = \hat{g}(i-1)$, then $\hat{f}(k) = \hat{g}(k)$ by the same argument made for $i = 1$. Therefore, $\hat{f}(i) = \hat{g}(i)$ for each $i \in \{0, \dots, m-1\}$, and $\hat{f} = \hat{g}$ as desired.

We now construct a function $\hat{f}: I_k \to \mathbb{Z}$ which satisfies $\hat{f}(0) = m$ and $f = p \circ \hat{f}$. First, define $\hat{f}(0) = m$, and $i \in \{1, 2, \dots, m-1\}$ we define inductively $\hat{f}(i) =$ $\hat{f}(i - 1) + i_{i-1}$ where

$$
i_{i-1} = \begin{cases} 1 & \text{si } f(k) \equiv (f(k-1)+1) \mod n \\ 0 & \text{si } f(k) \equiv (f(k-1)) \mod n \\ -1 & \text{si } f(k) \equiv (f(k-1)-1) \mod n \end{cases}
$$

So it follows by construction that \hat{f} is a bornologous function, that $\hat{f}(0) = m$ and that $p \circ \hat{f} = f$. $\hat{f} = f.$

Lemma 3.4.11. Let $f, g: I_k \to C_n$ be bornologous functions such that $f(0) =$ $g(0) = 0$ and $f(k) = g(k) = x_k$. Assume we have a homotopy $H: I_k \times \mathbb{Z} \to C_n$, between f and g such that $H(0, z) = x_0$ and $H(k, z) = x_k$ for each $z \in \mathbb{Z}$. Then, for each $\hat{x}_0 \in p^{-1}(0)$, there exists a unique lift \hat{H} : $I_k \times \mathbb{Z} \to \mathbb{Z}$ of H such that $p \circ \hat{H} = H$, $\hat{H}(0, z) = \hat{x}_0$, and there exists an $N \in \mathbb{Z}$ with $\hat{H}(k, N) = \hat{H}(k, z)$ for each $z \in \mathbb{Z}$.

Proof. Let $f, g: I_k \to C_n$ be bornologous functions such that $f(0) = g(0) = x_0$ and $f(k) = g(k) = x_k$. Assume we have $H: I_k \times \mathbb{Z} \to C_k$, a homotopy between f and g. By definition, there exist integers $N, M \in \mathbb{Z}, N \lt M$, such that $H(m, i) = f(m)$ for $i \leq N$ and $H(m, i) = g(m)$ for $i \geq M$. Therefore $H(0, z) = x_0$ and $H(k, z) = x_k$ for each $z \in \mathbb{Z}$. Consider a point $\hat{x}_0 \in p^{-1}(x_0)$.

For each $z \in \mathbb{Z}$, we define $f_z: I_k \to C_n$ by $f(x) := H(x, z)$ with $x \in I_k$. By [Lemma 3.4.10,](#page-30-0) then there are unique bornologous functions $\hat{f}_z : I_m \to \mathbb{Z}$ for each $z \in \mathbb{Z}$ such that $p \circ \hat{f}_z = f_z$ and $\hat{f}_z(0) = \hat{x}_0$, so we can define $\hat{H}(x, z) \coloneqq \hat{f}_z(x)$ for each $z \in \mathbb{Z}$ and $x \in I_k$. Under this construction, it follows that $p \circ \hat{H} = H$, $H(0, z) = \hat{x}_0$ for each $z \in \mathbb{Z}$, so it remains to show that H is bornologous.

Let's observe first that by construction $\hat{H}(x, z) = \hat{f}(x)$ if $z \leq N$, $\hat{H}(x, z) = \hat{g}(x)$ if $z \geq M$ and $\{(\hat{H}(x, z), \hat{H}(x + 1, z))\}\in \mathcal{Z}$ for each $z \in \mathbb{Z}$ and $x \in I_k$. So, to prove H is bornologous is enough to show that

$$
\{(\hat{H}(x,z),\hat{H}(x,z+1))\}\in\mathcal{Z},\{(\hat{H}(x+1,z),\hat{H}(x+1,z+1))\}\in\mathcal{Z},\{\hat{H}(x,z),\hat{H}(x+1,z+1))\}\in\mathcal{Z},\{(\hat{H}(x,z+1),\hat{H}(x+1,z))\}\in\mathcal{Z}.
$$

for each $x \in I_k$ and $z \in \{N, N+1, \dots, M-2, M-1\}.$

Let $x \in I_k$ and $z \in \{N, N+1, \cdots, M-2, M-1\}$. For each of the pairs $(x, z), (x', z')$ considered, above, since H is bornologous, we have $p \circ \hat{H}(x, z) = p \circ \hat{H}(x, z)$ $\hat{H}(x',z') + i_{x,z,x',z'}$, where $i_{x,z,x',z'} \in \{-1,0,1\}$. Then $H(x,z) = H(x',z') + \hat{i}_{x,z,x',z'}$, where $\hat{i}_{x,z,x',z'} \in \{-1,0,1\} \mod n$. For $x = 0$, note that $\hat{f}_z(0) = \hat{f}_{z+1}(0) = \hat{x}_0$, and therefore $\hat{i}_{0,z,0,z'} = 0$ for any $z \in \mathbb{Z}$ and $z' \in \{z, z + 1\}$. Since $\hat{f}_z(x) = \hat{f}_z(x + 1)$ 1) + $i_{x,z,x+1,z}$ by construction, we also have that $i_{x,z,x+1,z} = i_{x,z,x'z} \in \{-1,0,1\}$ for all $x \in I_k$. It follows that, for $x, x' \in \{0, 1\}$, any z, and $z' \in \{z, z + 1\}$, we have $\hat{i}_{x,z,x',z'} \in \{-1,0,1\}$. We use the same argument and induction to show that $\hat{i}_{x,z,x',z'} = i_{x,z,x'z'} \in \{-1,0,1\}$ for any $x,x' \in I_k$, $z \in \mathbb{Z}$, and $z' \in \{z,z+1\}$. It follows that \hat{H} is bornologous. \Box

Remark 3.4.12. The homotopy is unique because f_z is unique for each $z \in \mathbb{Z}$. If we change any element, the relations $p \circ \hat{H} = H$, $\hat{f}_z(0) = \hat{x}_0$ will no longer be true.

Lemma 3.4.13. Let $f: I_p \to \mathbb{Z}$ and $g: I_q \to \mathbb{Z}$ be bornologous functions such that $f(0) = g(0)$ and $f(p) = g(q)$. Then $\langle f \rangle \simeq_{sc} \langle g \rangle$.

Proof. Let $f: I_p \to \mathbb{Z}$ and $g: I_q \to \mathbb{Z}$ be bornologous functions such that $f(0) =$ $g(0)$ and $f(p) = g(q)$.

Observe that there exist unidirectional functions $f': I_{p'} \to \mathbb{Z}, f' \in \langle f \rangle$, and $g': I_{q'} \to \mathbb{Z}, g' \in \langle g \rangle$. If $f(0) = f(p)$, then $m' = n' = 0$, and the result follows.

Now assume that $f(0) \neq f(p)$. Then f' must be the function $f' : I_{|f(p)-f(0)|} \to \mathbb{Z}$ such that $f'(i) = f(0) + sign(f(m) - f(0))i$ for every $i \in \{0, ..., |f(m) - f(0)|\}$, with $sign(\cdot)$ the sign function. Analogously for g'. However, by hypothesis, $f(p) - f(0) =$ $g(q) - g(0)$, thus $f' = g'$. Therefore $\langle f \rangle = \langle g \rangle$.

Remark 3.4.14. Let $f, g: I_k \to X$ be bornologous functions such that $f(k) = g(0)$. By [Proposition 3.2.9,](#page-18-0) $f \star g$ is bornologous.

[Theorem 3.4.9.](#page-30-1) Let (C_n, \mathcal{C}_n) be the semi-coarse space induced by the *n*-cycle graph. We define $\phi : \mathbb{Z} \to \pi_1^{sc}(C_n, \mathcal{C}_n)$ by $\phi(z) := [\langle c^{*z} \rangle]$ Note that, by construction, c^{*z} is a bornologous function for each $z \in \mathbb{Z}$. It is clear from the definition and the definition of the \star operation that ϕ is a group homomorphism.

To prove that ϕ is surjective, consider $[\langle f \rangle] \in \pi_1^{sc}(C_n, C_n)$, and, without loss of generality, assume that $f \in \langle f \rangle$ is a bornologous function $f: I_k \to C_n$ with $f(0) = f(k) = 0$, and note that $0 \in p^{-1}(\{0\}) = n\mathbb{Z}$. Thus, by [Lemma 3.4.10](#page-30-0) there is a unique bornologous function \hat{f} such that $\hat{f}(0) = 0$ and $f = p \circ \hat{f}$. Since

 $(p \circ \hat{f})(k) = f(k) = 0$, note that $\hat{f}(k) = nq$ for some $q \in \mathbb{Z}$. Thus, $\hat{f} \simeq_{sc} \hat{c}^{*q}$ by [Lemma 3.4.13,](#page-31-0) and therefore

$$
f = p \circ \hat{f} \cong_{sc} p \circ \widehat{c^{*n}} = c^{*n}
$$

We conclude that $[\langle f \rangle] = [\langle \widehat{c^{*q}} \rangle] = \phi(q)$, and therefore ϕ is surjective.

To prove that ϕ is injective, suppose that $\phi(p) = \phi(q)$ for some $p, q \in \mathbb{Z}$, so we have that $\langle c^{*q} \rangle \simeq_{sc} \langle c^{*p} \rangle$. Let $H: I_k \to C_n$ be a homotopy from c^{*q} to c^{*p} , where $k = |pqn|$, and, abusing notation, we understand c^{np} and c^{iq} here to be the extension of each map to I_k by defining $c^{*p}(i) = 0$ if $i > |pn|$, and similarly for c^{*q} . Note that $H(0, i) = 0 \in C_n$ for each $i \in \mathbb{Z}$ by definition. We also note that $0 \in p^{-1}(\{0\}) = n\mathbb{Z}.$

By [Lemma 3.4.11,](#page-30-2) there is a unique homotopy $\hat{H}: I_{pan} \to C_n$ such that $H = p \circ \hat{H}$ and $\hat{H}(0,n) = 0$ for each $n \in \mathbb{Z}$. By uniqueness of the homotopy, we have that $\widehat{c^{kp}}(i) = \widehat{H}(i,N)$ and $\widehat{c^{*q}}(i) = \widehat{H}(i,M)$. Furthermore, since $\widehat{c^{*q}}(|nq|) = \widehat{c^{*p}}(k) =$ $\hat{H}(k, N) = \hat{H}(k, M) = \hat{c}^{*q}(k) = \hat{c}^{*q}(nq)$, we obtain $p = q$, and we conclude that the function is injective.

3.5. Long Exact Sequence in Homotopy. With the same goal, we will do the following definitions which will help us to prove the fundamental result: the long exact sequence in relative homotopy.

Definition 3.5.1. Let (X, V) be a semi-coarse space and (A, V_A) be a semi-coarse subspace. Then,

- The bornologous function $r : X \to A$ is a retraction if satisfies that $r \circ i =$ id_A , where $i : A \to X$ such that $i(a) = a$ for each $a \in A$.
- A bornologous function $F : (X \times \mathbb{Z}, V \times \mathbb{Z}) \to (X, V)$ is a deformation retract of X onto A if there are $N < 0 < M$ such that, for each $x \in X$ and $a \in A$, $F(x, z) = x$ if $z \leq N$, $F(x, z) \in A$ and $F(a, z) = a$ if $z \geq M$.
- If F also satisfies that $F(a, z) = a$ for each $z \in \mathbb{Z}$, then F is called strong deformation retraction.

Lemma 3.5.2. Let $f: I_m^n \to X$ be a bornologous function, $k \in I_m^n$ and $w \in$ $\{1, \ldots, n\}$. Then $f \simeq g \simeq h$ such that

$$
g(y) := \begin{cases} g(y) & \text{if } y_w \le k_w \\ g(y - (y_w - k_w)e_w) & \text{if } y_w > k_w \end{cases}
$$

and

$$
h(y) := \begin{cases} h(y) & \text{if } y_w \ge k_w \\ h(y - (y_w - k_w)e_w) & \text{if } y_w < k_w \end{cases}
$$

Proof. Let $f: I_m^n \to X$ be a bornologous function, $k \in I_m^n$, $w \in \{1, ..., n\}$,

$$
g(y) := \begin{cases} g(y) & \text{if } y_w \le k_w \\ g(y - (y_w - k_w)e_w) & \text{if } y_w > k_w \end{cases}
$$

and

$$
h(y) := \begin{cases} h(y) & \text{if } y_w \ge k_w \\ h(y - (y_w - k_w)e_w) & \text{if } y_w < k_w \end{cases}
$$

We will just prove for g, the proof for h is analogous. We are going to proceed by math induction:

Base case: Let k such that $k_w = m - 1$, $f \simeq g$ such that

$$
g(y) = \begin{cases} g(y) & \text{if } y_w \le m - 1 \\ g(y - e_w) & \text{if } y_w = m \end{cases}
$$

Induction step: Assume that $f \simeq g_i$ for every $i \in \{1, ..., k_w - 1\}$ where

$$
g_i(y) := \begin{cases} g_i(y) & \text{if } y_w \leq i \\ g_i(y - (y_w - i)e_w) & \text{if } y_w > i \end{cases}
$$

once again $f \simeq q$ such that

$$
g(y) := \begin{cases} g(y) & \text{if } y_w \le k_w \\ g(y - (y_w - k_w)e_w) & \text{if } y_w > k_w \end{cases}
$$

.

Observe that the last lemma makes us being able to replace either the plates "above" or "below" $(I_m^n)_{w}^k$. Thereby, the following lemma is a corollary. With the intention of making the proof clear, allow us call g as $(f)_{w}^{\uparrow}$ and h as $(f)_{w}^{\downarrow}$.

Lemma 3.5.3. Let n be a natural number and m be a non-negative integer number, then there is a strong deformation retraction I_m^n onto $\{*\}$.

Proof. Let n be natural number and m be a non-negative integer number, let's consider $* \in I_m^n$, then $* := k = (k_1, \dots, k_n)$ where $k_i \in \{0, 1, \dots, m\}$ fixed for each $i \in \{1, \dots, n\}$. Denote $1_{I_m^n}$ the identity function, then $1_{I_m^n} \simeq (1_{I_m^n})_1 \to \infty$ $((1_{I_m^n})_1^{\rightarrow})_1^{\leftarrow}$ =: g_1 by [Lemma 3.5.2.](#page-32-1) Therefore, $g_i := ((g_{i-1})_i^{\rightarrow})_i^{\leftarrow})_i^{\leftarrow}$ g_{i-1} for each $i\{2,\ldots,n\}$ by [Lemma 3.5.2.](#page-32-1) Thus, $g_n \simeq f$. The strong deformation that we are looking for is precisely that homotopy. \Box

The following lemma is a tool to prove the Compression Criterion, which one is our goal to finally get the long exact sequence in our homotopy.

Lemma 3.5.4. Let $f, g: (I_m^n, \partial I_m^n, J_m^{n-1}) \to (X, A, *)$ such that $f \simeq_{sc} g rel(\partial I^n)$. Let H the homotopy with $N < M$ and $N' > 0$, then $\langle f \rangle \simeq_{sc} \langle h_{N'} \rangle$ rel(∂I^n) with $h_{N'} : (I_{m+N'}^n, \partial I_{m+N'}^n, J_{m+N'}^{n-1} \to (X, A, *)$ such that

$$
h_{N'}(x) := \begin{cases} H(x - N'e_n, N + N') & x \in I_m^n + N'e_n \\ H(x - ke_n, N + k) & x \in (I_m^n)_n^0 + ke_n \text{ and } k \in \{0, ..., N' - 1\} \\ * & otherwise \end{cases}
$$

where $(I_m^n)_n^0 \subset I_m^n \subset I_{m+N'}^n$.

Proof. Let $f, g: (I_m^n, \partial I_m^n, J_m^{n-1}) \to (X, A, *)$ such that $f \simeq_{sc} g rel(\partial I^n)$. Let H the homotopy with $N < M$. We are going to do the proof by mathematical induction:

Base case: We have that $\langle f \rangle \simeq_{sc} \langle g_1 \rangle rel(\partial I^n)$ where g_1 : $(I^n_{m+1}, \partial I^n_{m+1}, J^{n-1}_{m+1}) \rightarrow$ $(X, A, *)$

$$
g_1(x) = \begin{cases} f(x - e_n) & x \in I_m^n + e_n \\ f(x) & x \in (I_m)_n^0 \\ * & \text{otherwise} \end{cases}
$$

Since H is a bornologous function, we are able to replace $f(x - e_n)$ by $H(x - e_n)$ $e_n, N + 1$, calling that function h_1 , and we get $g \simeq_{sc} h_1 rel(\partial I_{m+1}^n)$. Note that $f(x) = H(x, N')$ when $x \in (I_m)_n^0$. So, $\langle f \rangle \simeq_{sc} \langle h_1 \rangle rel(\partial I^n)$.

□

Induction step: Suppose that for $w > 0$ we have $\langle f \rangle \simeq_{sc} \langle h_w \rangle rel(\partial I^n)$ with $h_{N'}: (I_{m+w}^n, \partial I_{m+w}^n, J_{m+w}^{n-1} \to (X, A, *)$ such that

$$
h_w(x) := \begin{cases} H(x - we_n, N + w) & x \in I_m^n + we_n \\ H(x - ke_n, N + k) & x \in (I_m^n)_n^0 + ke_n \text{ and } k \in \{0, ..., w - 1\} \\ * & \text{otherwise} \end{cases}
$$

We have that $\langle h_w \rangle \simeq_{sc} \langle g_{w+1} \rangle rel(\partial I^n)$ where g_{w+1} : $(I^n_{m+w+1}, \partial I^n_{m+w+1}, J^{n-1}_{m+w+1}) \rightarrow$ $(X, A, *)$ such that

$$
g_{w+1}(x) := \begin{cases} H(x - (w+1)e_n, N+w) & x \in I_m^n + (w+1)e_n \\ H(x - we_n, N+w) & x \in (I_m^n)_0^n + we_n \\ H(x - ke_n, N+k) & x \in (I_m^n)_n^0 + ke_n \text{ and } k \in \{0, \dots, w-1\} \\ * & \text{otherwise} \end{cases}
$$

Since H is a bornologous function, we are able to replace $H(x - (w + 1)e_n, N + W)$ by $H(x - (w + 1), N + w + 1)$, calling that function h_{w+1} , and we get $g_{w+1} \simeq_{sc}$ h_{w+1} rel(∂I_{m+w+1}^n). $\binom{n}{m+w+1}$.

Lemma 3.5.5 (Compression Criterion). Let (X, V) be a semi-coarse space and $*\in A\subset X$. Then a function $f:(I^n,\partial I^n,J^{n-1})\to (X,A,*)$ represents the zero in $\pi_n^{sc}((X, V), A, *)$ if, and only if, is relative homotopic in ∂I^n to a function with its image contained in A.

Proof. Let (X, V) be a semi-coarse space, $* \in A \subset X$ and $f : (I^n, \partial I^n, J^{n-1}) \to$ $(X, A, *)$.

- (\Rightarrow) If $[f] = 0$, then there exist $m \in \mathbb{N}$ with $f: (I_m^n, \partial I_m^n, J^{n-1}) \to (X, A, *)$ such that $f \simeq_{sc} *$. Let H be the homotopy with $N < M$, then by [Lemma 3.5.4,](#page-33-0) we get h_{M-N} as we write in such lemma which has its image contained in A and $\langle f \rangle \simeq_{sc} \langle h_{M-N} \rangle rel(\partial I^n)$.
- (←) If f is relative homotopic in ∂I^n to a function with its image contained in A, let's say g. As we do in [Lemma 3.5.3,](#page-33-1) with $w = (0, \ldots, 0, m)$, then $g \simeq_{sc} (g)^{\downarrow}_{n} = *$. Getting what we wanted.

$$
\qquad \qquad \Box
$$

Definition 3.5.6. Let (X, Y) and (Y, W) be semi-coarse spaces and $f: ((X, Y), A, *) \rightarrow$ $((Y.W), B, *)$ be a Bornologous function. Then, we define $f_* : \pi_*^{sc}((X, V), A, *) \rightarrow$ $\pi_*^{sc}((Y,W),B,*)$ as $f_*([\langle h \rangle]) = [\langle f \circ h \rangle]$ for each $[\langle h \rangle] \in \pi_*^{sc}((X,V),A,*)$.

Remark 3.5.7. Let (X, V) and (Y, W) be semi-coarse spaces and $f : ((X, V), A, *) \rightarrow$ $((Y.W), B, *)$ be a (based) Bornologous function. Then f_* is well-defined and is an homomorphism. It is well-defined because, if we take $\langle g \rangle \in [\langle h \rangle] \in \pi_*^{sc}((X, \mathcal{V}), A, \ast),$ then there is a homotopy H between them, so $f \circ H$ is a homotopy between $\langle f \circ h \rangle$ and $\langle f \circ g \rangle$. If $[\langle h \rangle], [\langle g \rangle] \in \pi_*^{sc}((X, V), A, *)$. It is a homomorphism because $f \circ (g \star h) = f \circ g(k_1, \cdots k_n)$ if $0 \leq k_i \leq m$ for each $i \in \{1, \cdots, n\},$ $f \circ (g \star h) = f \circ h(k_1, \cdots k_n)$ if $m < k_1 \leq 2m$ $0 \leq k_i \leq m$ for each $i \in \{2, \cdots, n\},$ and $f \circ (g \star h) = f(\star)$ anywhere else, which is the same as $(f \circ g) \star (f \circ h)$.

Theorem 3.5.8 (Long Exact Sequence in Homotopy). Let (X, V) be a semi-coarse space and $* \in A \subset X$. As well $i : (A, *) \rightarrow (X, *)$, $j : (X, *) \rightarrow (X, A)$ inclusions. Then there is a homomorphism $\partial_n : \pi_n^{sc}((X, V), A) \to \pi_n^{sc}(A, *)$ such that the long sequence

$$
\cdots \longrightarrow^{\quad j_*} \pi_{n+1}^{sc}(X, A) \longrightarrow^{\partial_{n+1}} \pi_n^{sc}(A, *) \longrightarrow^{\quad i_*} \pi_n^{sc}(X, *) \longrightarrow^{\quad j_*} \cdots
$$

$$
\cdots \longrightarrow^{\partial_2} \pi_1^{sc}(A, *) \longrightarrow^{\quad i_*} \pi_1^{sc}(X, *) \longrightarrow^{\quad j_*} \pi_1^{sc}(X, A)
$$

is exact.

Proof. Let (X, V) be a semi-coarse space and $* \in A \subset X$. As well $i : (A, *) \rightarrow$ $(X, *), j : (X, *) \to (X, A)$ inclusions. We are going to define $\partial_n : \pi_n^{sc}((X, V), A) \to$ $\pi_n^{sc}(A,*)$ as $f \mapsto f|_{I^{n-1}\times\{0\}},$ which is well-defined and is a homomorphism. Now we will prove that the long sequence is exact:

- $(Im(i_*) \subset Ker(j_*))$ If $n \geq 1$ and $f : (I^n, \partial I^n, J^{n-1}) \rightarrow (A, *, *) \in$ $\pi_n^{sc}(A, *),$ then $j \circ i \circ f \in \pi_n^{sc}(X, A)$, so by compression criterion we have that $[\langle f \rangle] = [\langle * \rangle]$.
- $(Im(i_*) \supset Ker(j_*)$ If $n \geq 1$ and $[\langle f \rangle] \in \pi_n^{sc}((X, V), *)$ such that $j_*[\langle f \rangle] =$ [⟨*)]. Then by compression criterion, $\langle f \rangle \simeq_{sc} \langle g \rangle rel(\partial I^n)$, where the image of g is contained in A, thus $[\langle g \rangle] = [\langle f \rangle] \in \pi_n^{sc}((X, V), *)$ is in the image of $i_{*}.$
- $(Im(j_*)\subset Ker(\partial))$ If $n \geq 2$ and $[\langle f \rangle] \in \pi_n^{sc}((X,\mathcal{V}),*)$, then $\partial j\langle f \rangle =$ $\langle f|_{I^{n-1}\times\{0\}}\rangle = \langle * \rangle$, so $\partial j_*[\langle f \rangle] = [\langle * \rangle]$.
- $(Im(j_*) \supseteq Ker(\partial))$ If $n \geq 2$ and $[\langle f \rangle] \in \pi_n^{sc}((X, V), A)$ such that $\partial[\langle f \rangle] =$ [⟨*)]. Then $f|_{I^{n-1}\times\{0\}}$ is homotopic to $\langle\ast\rangle$ by the compression theorem through a homotopy $H: I^{n-1}_m \times \mathbb{Z} \to A$ $rel(\partial I^{n-1}_m)$, $N < 0 < M$ such that $H(x, z) = *$ if $z \leq N$ and $H(x, z) = f|_{I^{n-1} \times \{0\}}(x, z)$ if $z \geq M$. Let's define $g: (I_{m+N-N}^n, \partial I_{m+N-N}^n, J_{m+N-N}^{n-1}) \to ((X, V), A, *)$ such that

$$
g(k_1,\cdots,k_n)=H(k_1,\cdots,k_{n-1},z=k_n)
$$

with $0 \leq k_i \leq m$ when $i \in \{1, \dots, n-1\}$ and $0 \leq k_n \leq M-N$,

$$
g(k_1, \dots, k_n) = f(k_1, \dots, k_{n-1}, k_n - M + N)
$$

with $0 \leq k_i \leq m$ when $i \in \{1, \dots, n-1\}$ and $M-N+1 \leq k_n \leq M-N+m$, and $g(k_1, \dots, k_n)$ anywhere else. So, we note that $g(\partial I_{m+M-N}^n) = *$, that is, $[\langle g \rangle] \in Im(j_*)$, and $[\langle f \rangle] = [\langle g \rangle]$.

- $(Im(\partial) \subset Ker(i_{*}))$ If $n \geq 2$ and $f : (I_{m}^{n}, \partial I_{m}^{n}, J_{m}^{n-1}) \rightarrow ((X, V), A, *)$. Then, defining $H: I^{n-1}_{m} \times Z \to (X, V)$ as $H(k_1, \dots, k_n) = f(k_1, \dots, k_n)$ with $0 \le k_n \le m$, $H(k_1, \dots, z) = f(k_1, \dots, k_{n-1}, 0)$ if $z \le 0$, and $H(k_1, \dots, z) =$ $f(k_1, \dots, k_{n-1}, m)$ if $z \geq m$, we get that $f|_{I^{n-1}_{m} \times \{0\}}$ is relative homotopic to $*$ through H. So, $i_*\partial[\langle f \rangle] = [\langle * \rangle]$ by [Lemma 3.5.5.](#page-34-0)
- $(Im(\partial) \supset Ker(i_{*}))$ If $n \geq 2$ and $f : (I_{m}^{n}, \partial I_{m}^{n}, J_{m}^{n-1}) \rightarrow (A, *, *)$ such that $i_*([f]) = [\langle * \rangle]$, then we have a homotpy between f and *, which gives us a function $F: (I_{M-N}^{n+1}, \partial I_{M-N}^{n+1}, J_{M-N}^n) \to ((X, V), A, *)$ such that $\partial[\langle F \rangle] = [\langle f \rangle].$

$$
\Box
$$

4. Homology

In the last section, we will construct a Vietoris-Rips homology for semi-coarse spaces and show that it is homotopy invariant. We finish with the observation

that the homology defined here for a finite semi-coarse space is isomorphic to the homology of the clique complex of a finite graph.

4.1. Simplicial Homology.

Definition 4.1.1. Let (X, V) be a semi-coarse and E be a controlled set, and let R be a relation on X such that $x \mathcal{R} y$ iff $(x, y) \in E$. We now define the following

- \bullet $\Sigma^{(0)}_E$ $E^{(0)}_E := \{\{x\} : x \in X\},\$
- \bullet $\Sigma^{(n)}_E$ $E_E^{(n)} := \{ \{x_0, \dots, x_n\} \subset X \mid \forall i, j \in \{0, \dots, n\} x_i \mathcal{R} x_j \text{ and } ((i \neq j) \implies$ $x_i \neq x_j$ }
- We call the collection $\cup_{i=0}^n \Sigma_E^{(i)}$ $E^{(i)}$ the *n-skeleton* of Σ .
- $\Sigma_E := \cup_n \Sigma_E^{(n)}$ will be called the simplicial complex associated to E.

Remark 4.1.2. We note that Σ_E satisfies the definition of simplicial complex for any element $E \in V$ in the semi-coarse structure X, [Definition 4.1.1,](#page-36-1) since $\Sigma_E^{(0)}$ contains all sets with one vertex and every subset of a simplex is a simplex having all their elements related.

Definition 4.1.3. We define $C_q(X, E)$ to be the free abelian groups generated by ordered simplicial chains, denoted $[v_0, v_1, \ldots, v_q]$, where $[v_0, v_1, \ldots, v_q] = 0$ if the vertices are not all pairwise different. We define the differential ∂_q by

$$
\partial_q[v_0, v_1, \dots, v_q] := \sum_{i=0}^q (-1)^i [v_0, \dots, \hat{v}_i, v_q]
$$

and we denote the chain complex by $C_*(X, E) := \{C_q(X, E), \partial_q\}$ and the resulting homology groups by $H_*(X, E)$.

The above definitions show how to construct homology groups from a single element of a semi-coarse structure $E \in \mathcal{V}$. The next lemma will allow us to construct a directed system from these homology groups.

Lemma 4.1.4. Let (X, V) be a semi-coarse space and E, E' controlled sets such that $E \subset E'$. Then there exists a homomorphism $i_* : H(X, E) \to H(X, E')$.

Proof. Let (X, V) be a semi-coarse space and let E, E' be controlled sets such that $E \subset E'$. If $i : E \to E'$ is the inclusion from E to E', then for a generator $\sigma = [v_0, \ldots, v_q]$ of $C_q(X, E)$, we define $(i_{\#})_q : C_q(X, E) \to C_q(X, E')$ by

$$
(i_{\#})_q[x_0, \cdots, x_q] := [i(x_0), \cdots, i(x_q)] = [x_0, \cdots, x_q] \in C_q(X, E').
$$

We then extend this by linearity. Note that the inclusion $E \subset E'$ ensures that $C_q(X, E) \subset C_q(X, E'),$ since $x \mathcal{R}_E y \implies x \mathcal{R}_{E'} y$ for any $x, y \in X$. The inclusion $i_{#}$ also satisfies $\partial_{E'} i_{#} = i_{#} \partial_{E}$, so $i_{#}$ is a chain map. The induced map i_{*} : $H_*(X, E) \to H_*(X, E')$ such that $i_*[\sigma] = [i_{\#}(\sigma)] = [\sigma] \in H_*(X, E')$ is the desired map in homology.

Definition 4.1.5. Let (X, Y) be a semi-coarse space, and consider V with the partial order given by inclusion of sets. We define

$$
H_*(X,\mathcal{V}):=\lim_{\rightarrow}\{H(X,E),\pi_E^{E'},\mathcal{V}\},\
$$

where $\pi_E^{E'} = i_* : H(X, E) \to H(X, E')$ from the last lemma. We call $H_*(X, V)$ the homology of the semi-coarse space (X, V) . We will sometimes refer to $H_*(X, V)$ as the Vietoris-Rips homology of (X, V) .

We can note that the set of symmetric sets $E \in \mathcal{V}$ is cofinal in \mathcal{V} , i.e. for every $E \subset \mathcal{V}$, we have $E \subset E \cup E^{-1} \cup \Delta_X \in \mathcal{V}$. Denote by $\mathcal{V}_{\mathcal{S}}$ the collection ${E \in \mathcal{V} \mid E = E^{-1}}$. By the cofinality We have that

$$
\lim_{\rightarrow} \{H(X, E), \pi_E^{E'}, \mathcal{V}\} \cong \lim_{\rightarrow} \{H(X, E), \pi_E^{E'}, \mathcal{V}_{\mathcal{S}}\},\
$$

so it is enough to consider symmetric elements of V when constructing the homology of (X, V) .

We will now show the semi-coarse homology is a covariant functor. We will start by showing that homology is functorial for a fixed controlled set $E \in \mathcal{V}$.

Lemma 4.1.6. Let (X, V) and (X, W) be semi-coarse spaces, E controlled by V and $f: (X, V) \to (Y, W)$ be a bornologous function. Then,

(i)
$$
f_{\#}: C(X, E) \to C(Y, (f \times f)(E))
$$
 where

$$
(f_{\#})_{n}[x_{0},\cdots,x_{n}]=[f(x_{0}),\cdots,f(x_{n})]
$$

is a chain map.

(ii) $f_*: H(X, E) \to H(Y, (f \times f)(E))$ is a group homomorphism, where $(f_*)_n[\sigma^n] =$ $[(f_{\#})_n \sigma^n].$

Proof. Let (X, V) and (Y, W) be semi-coarse spaces, E controlled by V and f: $(X, \mathcal{V}) \to (Y, \mathcal{W})$ be a bornologous function. Since f is bornologous, $\{(f(x), f(y))\} \in$ $(f \times f)(E) \in \mathbb{W}$ if $\{(x, y)\}\in E$, so f induces the simplicial map $f_{\#}$ defined by

$$
(f_{\#})_{n}[x_{0},\cdots,x_{n}]=[f(x_{0}),\cdots,f(x_{n})].
$$

Since $f_{\#}$ is a chain map, it induces the map $f :_{*}: H_{*}(X, E) \to H_{*}(Y, (f \times f)(E))$ on homology by

$$
(f_*)_n[\sigma^n] = [(f_{\#})_n \sigma^n]. \square
$$

Remark 4.1.7. The previous result is also true for $f_{\#}: C(X, E) \to C(Y, A)$ and $f_*: H(X, E) \to H(Y, A)$ such that $(f \times f)(E) \subset A$ is controlled by W. This follows from the previous lemma and [Lemma 4.1.4.](#page-36-2)

Theorem 4.1.8. Semi-coarse homology is a covariant functor H_* : SCoarse \rightarrow Ab.

Proof. Let (X, V) and (Y, W) be semi-coarse spaces and let $f : (X, V) \to (Y, W)$ be a bornologous function. Let $\{H(X, E), \pi_E^{E'}, \mathcal{V}\}\$ and $\{H(Y, E), \pi_E^{E'}, \mathcal{W}\}\$ be the directed systems of the homology groups, where the directed sets V and W are partially ordered by inclusion.

Note that, if $A \subset B \subset X \times X$, then $(f \times f)(A) \subset (f \times f)(B)$ so $(f \times f)$ is preserves the preorders on V and W. Therefore, for each $E, E' \in V$ such that $E \subset E'$, the diagram

$$
H(X, E) \longrightarrow^{\mathit{f}_{*}} H(Y, (f \times f)(E))
$$

\n
$$
\pi_{E}^{E'} \downarrow \pi_{(f \times f)(E')}^{\pi_{(f \times f)(E')}} \downarrow
$$

\n
$$
H(X, E') \longrightarrow^{\mathit{f}_{*}} H(Y, (f \times f)(E'))
$$

commutes. It follows by definition that there exists a homomorphism $F : H_*(X, \mathcal{V}) \to$ $H_*(X, \mathcal{W})$ between the directed limits, that is,

$$
F: \lim_{\longrightarrow} \{H(X, E), \pi_E^{E'}, \mathcal{V}\} \to \lim_{\longrightarrow} \{H(Y, E), \pi_E^{E'}, (f \times f)(\mathcal{V})\}.
$$

We now show that homotopic maps induce the same map on semi-coarse homology

Theorem 4.1.9. Let (X, V) and (Y, W) be semi-coarse spaces. If $f, g : (X, V) \rightarrow$ (Y, W) are homotopic functions, then the induced homomorphisms $f_*, g_* : H_*(X, V) \to$ $H_*(Y, W)$ are equal.

Proof. Let (X, V) and (Y, W) be semi-coarse spaces. If $f, g : (X, V) \to (Y, W)$ are homotopic functions, then there are a bornologous function

$$
H: (X \times \mathbb{Z}, \mathcal{V} \times \mathcal{Z}) \to (Y, \mathcal{W})
$$

and $N < 0 < M$ integer numbers such that $H(x, z) = f(x)$ if $z \leq N, x \in X$ and $H(x, z) = g(x)$ if $z \geq M, x \in X$. We will denote by $h_z(x) := H(x, z)$ for $x \in X$ and $z \in \{N+1, N+2, \cdots, M-2, M-1\}.$

We will observe what happens with the induced homomorphisms by f and h_{N+1} . Then, for each E controlled by V, we define $(\Psi_E)_q : C_q(X, E) \to C(Y, H(E, \mathbb{Z}))$ such that

$$
(\Psi_E)_q[x_0, \cdots, x_q] := \sum_{i=0}^q (-1)^i [f(x_0), \cdots, f(x_i), h_{N+1}(x_i), \cdots, h_{N+1}(x_q)].
$$

A straightforward calculation shows that Ψ_E is a chain homotopy between the induced chain maps $f_{\#}$ and $(h_{N+1})_{\#}$, so $f_{*} = (h_{N+1})_{*} : H(C(X, E)) \to H(C(Y, H(E, \mathbb{Z})))$. Thereby, by similar arguments to the previous theorem, we get that

$$
f_* = (h_{N+1})_* : H(C(X, V)) \to H(C(Y, W)).
$$

We will repeat the same for h_i and h_{i+1} with $i \in N+1, \cdots M-1$, finally arriving at

$$
f_* = g_* : H(C(X, V)) \to H(C(Y, W)),
$$

as desired. \Box

4.2. Graphs and Semi-Coarse Spaces. Now that we have introduced the basic concepts of semi-coarse homology, we recall every graph is a roofed semi-coarse space (where roofed semi-coarse space is defined in [Definition A.1.11\)](#page-44-0), and, given the similarities between the constructions of the semi-coarse homology and the Vietoris-Rips homology of a graph, is natural to ask whether they are isomorphic. We will answer this in the affirmative in this section, in addition to showing that the semi-coarse homology only depends on the roof of the semi-coarse structure. We begin with the following lemma.

Lemma 4.2.1. Let (X, V) be a (possibly non-roofed) semi-coarse space with roof $\mathfrak A$. Then every finite subset A of $\mathfrak A$ is an element of $\mathcal V$.

Proof. The lemma is immediate if (X, V) is roofed. Recall that, by definition, $\mathfrak{A} = \bigcup_{V \in \mathcal{V}} V$. If $A = \{a_0, \ldots, a_k\}$ is a finite subset of \mathfrak{A} , then each element a_i of A is contained in some set $V_i \in \mathcal{V}$. By the axioms of a semi-coarse structure, this implies that $\{a_i\} \in \mathcal{V}$ for each $a_i \in A$, and therefore that $A = \bigcup_{i=0}^k \{a_i\} \in \mathcal{V}$. \Box

We now use this to show that the semi-coarse homology only depends on the roof of the semi-coarse structure.

□

Theorem 4.2.2. Let (X, V) be a semi-coarse space. Then

$$
H(X, V) \cong H(X, \mathfrak{R}(X, V)).
$$

Proof. Let (X, V) be a roofed semi-coarse space with roof \mathfrak{A} . Then $\{H(X, \mathfrak{A})\}$ is cofinal in the directed system $\{H(X, E), \pi_E^{E'}, \mathcal{V}\}$, and the result follows.

Now assume that V is non-roofed with roof \mathfrak{A} , and let W be the the roofed semicoarse structure with roof \mathfrak{A} . Since $\mathcal{V} \subset \mathcal{W}$, then there exists a homomorphism

$$
\pi^{\mathcal{W}}_{\mathcal{V}}: \lim\limits_{\rightarrow}\{H(X,E), \pi^{E'}_E, \mathcal{V}\} \rightarrow \lim\limits_{\rightarrow}\{H(X,E), \pi^{E'}_E, \mathcal{W}\}
$$

such that $\pi_{\mathcal{V}}^{\mathcal{W}}\langle\sigma_E\rangle_{\mathcal{V}} = \langle\sigma_E\rangle_{\mathcal{W}}.$

Surjectivity: Let $n \in \mathbb{N}$ and $x \in \lim_{\to} \{H_n(X, E), \pi_E^{E'}, \mathcal{W}\}$, then there exists a set $E \in \mathcal{W}$ and $y \in H_n(X, E)$ such that $x = \langle y_E \rangle_{\mathcal{W}}$. If $E \in \mathcal{V}$, then $\pi_{\mathcal{V}}^{\mathcal{W}}[y_E]_{\mathcal{V}} = x$.

If $E \notin V$, we need to do something else. As $y \in H(X, E)$, then there exists a cycle $z \in \text{Ker}(\partial_n^E)$ such that $y = \langle z \rangle \in H_n(X, E)$ and

$$
z = \sum_{i=1}^{q} \alpha_i \sigma_i, \ \sigma_i = [z_0^i, \dots, z_n^i], \alpha_i \in \mathbb{Z},
$$

where, the σ_i are the elements of the chain complex corresponding to the ordered simplices $[z_0^i, \ldots, z_n^i]$. By [Lemma 4.2.1,](#page-38-1) each subset of vertices $A_i := \{z_0^i, \ldots, z_n^i\} \in$ V forming each ordered simplex σ_i is in V, and therefore $A := \cup_i A_i \in V$ as well.

It now follows that $z \in Ker(\partial_n^A)$, which the properties of the directed system It how follows that $z \in \mathbb{R}e_r(o_n)$, which the properties of the different system
imply that $\pi_A^{A \cup E}(\langle z \rangle_A) = \langle z \rangle_{A \cup E} = \pi_E^{A \cup E}(\langle z \rangle_E)$, and it follows that $\langle z \rangle_A$ and $\langle z \rangle_E$ represent the same element of $H_*(X,\mathcal{W})$, i.e. $\langle z \rangle_{A \cup E}|_{\mathcal{W}} = [\langle z \rangle_A]_{\mathcal{W}} = x$, i.e. $\langle z \rangle$. it now follows that $\pi_{\mathcal{V}}^{\mathcal{W}}[\langle z \rangle_A]_{\mathcal{V}} = x$. Thus, $\pi_{\mathcal{V}}^{\mathcal{W}}$ is surjective.

Injectivity: In this part of the proof we will abuse notation and denote by $\pi_{\mathcal{U}}^{\mathcal{W}}$ the homomorphism

$$
\pi^{\mathfrak{A}}_{\mathcal{V}}: \lim\limits_{\rightarrow}\{H(X,E), \pi^{E'}_E, \mathcal{V}\} \rightarrow H(X, \mathfrak{A})
$$

implicitly composing the original map with the isomorphism

$$
\lim_{\rightarrow} \{H(X, E), \pi_E^{E'}, \mathcal{W}\} \cong H(X, \mathfrak{A})
$$

Let $n \in \mathbb{N}$ and $x \in \lim_{\to} \{H_n(X, E), \pi_E^{E'}, \mathcal{V}\}\$ such that $\pi_{\mathcal{V}}^{\mathfrak{A}}(x) = 0$. Then there exists a set $E \in V$ and $z \in Ker(\partial_n^E)$ such that $x = [\langle z \rangle_E]_V$ and $\pi_E^{\mathfrak{A}} \langle z \rangle = 0$. Therefore, $z \in Im(\partial_{n+1}^{\mathfrak{A}})$, that is, there exists an element $z' \in C_{n+1}^{\mathfrak{A}}$ such that $\partial_{n+1}^{\mathfrak{A}}(z') = z$.

Since $\pi_V^{\mathfrak{A}}$ is surjective, there is a set $U \in \mathcal{V}$ such that $z' \in C_{n+1}^U$ and $\partial z' =$ $z \in C_n^U$. This implies that $\langle z \rangle_U = 0_U$ and $\pi_E^U \langle z \rangle_E = 0_U$. Thus, $[\langle z \rangle_E]_{\mathcal{V}} = 0 \in$ $\lim_{\longrightarrow} \{H(X, E), \pi_E^{E'}, \mathcal{V}\},$ and $\pi_{\mathcal{V}}^{\mathcal{W}}$ is injective.

It now follows that
$$
\lim_{\to} \{H(X, E), \pi_E^{E'}, \mathcal{V}\} \cong H(X, \mathfrak{A}).
$$

We now compare the semi-coarse homology of the vertices of a graph with the semi-coarse structure induced by the graph and the homology of the clique complex of a graph. We begin by recalling the definition of the clique complex.

Definition 4.2.3. A graph is called complete if each pair of vertices is adjacent. A k-clique in G is a complete subgraph of G with k vertices, and it is a maximal k-clique if it is not proper subgraph of another clique.

Definition 4.2.4. Given a graph $G = (V, E)$ the *clique complex* Σ_G of G is the simplicial complex such that $\Sigma_G^{(0)} = V$ and a finite set $\sigma := \{v_0, \ldots, v_k\} \subset V$ is a k-simplex in Σ_G iff the induced subgraph of G on the vertices in σ is a $(k +$ 1)-clique). The Vietoris-Rips homology of a graph G, denoted $H_*^{VR}(G)$, is the simplicial homology of the simplicial complex Σ_G .

Theorem 4.2.5. Let $G = (V, E)$ be a graph and let (V, V_G) be the semi-coarse space induced by G. Then $H_*(V, V_G) \cong H_*^{V \hat{R}}(G)$.

Proof. Let $G = (V, E)$ be a graph and (V, V_G) be semi-coarse space associated to G. Note that $\bigcup_{E\in\mathcal{V}}E=\mathfrak{R}(\mathcal{V})$, and therefore

$$
\lim_{\rightarrow} \{H(V,E), \pi_E^{E'}, \mathcal{V}\} = H\left(V, \bigcup_{E \in \mathcal{V}} E\right),\,
$$

by thereby we only have to examine the chain complex for the roof of V .

Let $C(G)$ denote the ordered chain complex generated by ordered simplices of the clique complex of G, and define $\iota : C(X, \cup_{E \in \mathcal{V}} E) \to C(G)$ such that $\iota_q([x_0,\dots,x_q])=[x_0,\dots,x_q]$ if all the elements are different and $\iota_q([x_0,\dots,x_q])=0$ it at least a pair of elements are equal. Note that ι is well-defined when all of elements are different because $x_i \mathcal{R} x_j$ for each $i, j \in \{0, 1, \dots, q\}$, thereby $\{x_i, x_j\} \in V$ when $i \neq j$, concluding $\{x_0, \dots, x_q\}$ is a clique with $q + 1$ elements in G.

On the other hand, let's define $\kappa : C(G) \to C(X, \cup_{E \in \mathcal{V}} E)$ such that

$$
\kappa_q([x_0,\cdots,x_q])=[x_0,\cdots,x_q].
$$

By the argument in the previous paragraph, it is clear that κ is well-defined. Moreover, $\iota \circ \kappa = 1_{C(X,\cup_{E \in \mathcal{V}} E)}$ and $\kappa \circ \iota = 1_{C(G)}$, so $C(X,\cup_{E \in \mathcal{V}} E) \cong C(G)$, thus $H(V(G), V) \cong H(G).$

5. Discussion and Future Work

In this paper, we have begun the development of algebraic topology for semicoarse spaces, a generalization of the category of coarse spaces which allows, in particular one to coarsen a metric space up to a predetermined scale. The axioms for a semi-coarse structure on a space are obtained by deleting the product axiom from the axioms for coarse spaces, and semi-coarse spaces receive their name in analogy with the loss of the triangle inequality when one passes from metric spaces to semi-metric spaces. Our main results are the development of a homotopy theory on semi-coarse spaces using the integers as an interval, and the construction of Vietoris-Rips homology for semi-coarse spaces. We further define roofed semicoarse spaces, which clarify the precise structure which determines the Vietoris-Rips homology of a semi-coarse space.

Since we are trying to adapt the algebraic topology of topological spaces to this category, it is tempting to try to construct homotopy invariants using the unit interval as a cylinder object. However, our first observation is that the closed interval [0, 1] is not the best candidate for the interval in a semi-coarse homotopy theory:

Proposition [3.0.1.](#page-13-2) Let $r, r' \in (0, 1)$. Any bijective map $\phi : ([0, 1], \mathcal{V}_{r'}) \rightarrow$ $([0,1], \mathcal{V}_{r'}) \sqcup_{1 \sim 1} ([1,2], \mathcal{V}_r)$ where the latter space is the pushout of the two intervals, glued at one endpoint, is not bornologous at the point 1 where the two intervals are glued.

After that, through a long construction, we obtain our homotopy classes and a binary operation for them, and show that they behave similarly to homotopy classes from topological spaces with their usual product.

Theorem [3.2.18.](#page-23-0) Let (X, V) be a semi-coarse space, $A \subset X$, and $n \in \mathbb{N}$.

- If $n \geq 1$, then $(\pi_n^{sc}(X, *), *)$ is a group.
- If $n \geq 2$, then $(\pi_n^{sc}(X, *), \star)$ is an abelian group.
- If $n \geq 2$, then $(\pi_n^{sc}(X, A, *), *)$ is a group.
- If $n \geq 3$, then $(\pi_n^{sc}(X, A, *), \star)$ is an abelian group.

At the same time, in [Theorem 3.2.19,](#page-25-0) we observe that this homotopy becomes trivial on coarse spaces.

Theorem [3.2.19.](#page-25-0) Let (X, V) be a coarse space with $A \subset X$. Then for any $n \in \mathbb{N}$, we have $\pi_n^{sc}(X, A, *) \cong \pi_n^{sc}(X, *) \cong \{1\}.$

We conclude the homotopy section with three important results: the existence of semi-coarse spaces whose homotopy is not trivial (the cyclic graphs, [3.4.9\)](#page-30-1), the compression criterion [\(3.5.5\)](#page-34-0) and the long exact sequence in homotopy [\(3.5.8\)](#page-34-1).

Theorem [3.4.9.](#page-30-1) Let (C_n, \mathcal{C}_n) be the semi-coarse space induced by the *n*-cycle graph. Then, $\pi_1^{sc}(C_n, \mathcal{C}_n) \cong \mathbb{Z}$.

Lemma [3.5.5.](#page-34-0) Let (X, V) be a semi-coarse space and $* \in A \subset X$. Then a function $f:(I^n,\partial I^n,J^{n-1})\to (X,A,*)$ represents the zero in $\pi_n^{sc}((X,\mathcal{V}),A,*)$ if, and only if, is relative homotopic in ∂I^n to a function with its image contained in A.

Theorem [3.5.8.](#page-34-1) Let (X, Y) be a semi-coarse space and $* \in A \subset X$. As well $i: (A, *) \rightarrow (X, *), j: (X, *) \rightarrow (X, A)$ inclusions. Then there is a homomorphism $\partial_n : \pi_n^{sc}((X, V), A) \to \pi_n^{sc}(A, *)$ such that the long sequence

$$
\cdots \longrightarrow^j \pi_{n+1}^{sc}(X, A) \longrightarrow^{\partial_{n+1}} \pi_n^{sc}(A, *) \longrightarrow^i \pi_n^{sc}(X, *) \longrightarrow^j \cdots
$$

$$
\cdots \longrightarrow^{\partial_2} \pi_1^{sc}(A, *) \longrightarrow^i \pi_1^{sc}(X, *) \longrightarrow^j \pi_1^{sc}(X, A)
$$

is exact.

After our discussion of semi-coarse homotopy groups, we develop a homology with the same flavor as the Vietoris-Rips homology of metric spaces at a given scale. We observe, in particular, that this homology is completely characterized by the roofed semi-coarse spaces [\(Theorem 4.2.2\)](#page-39-0), which are essentially graphs, and we conclude with a theorem showing that, for any graph, their Vietoris-Rips homology is isomorphic to the semi-coarse homology of the semi-coarse spaces induced by the graph [\(Theorem 4.2.5\)](#page-40-1). This latter theorem serves to show that the semi-coarse homology developed here is a strict generalization of the homology of the clique complex of a graph, and motivates the possibility of its use in topological data analysis. We further show that the semi-coarse homology is invariant with respect to semi-coarse homotopy equivalence which we defined earlier [\(Theorem 4.1.9\)](#page-38-2).

Theorem [4.2.2.](#page-39-0) Let (X, V) be a semi-coarse space. Then

$$
H(X, V) \cong H(X, \mathfrak{R}(X, V)).
$$

Theorem [4.2.5.](#page-40-1) Let $G = (V, E)$ be a graph and let (V, V_G) be the semi-coarse space induced by G. Then $H_*(V, V_G) \cong H_*^{\hat{V}R}(G)$.

Theorem [4.1.9.](#page-38-2) Let (X, V) and (Y, W) be semi-coarse spaces. If $f, g: (X, V) \rightarrow$ (Y, W) are homotopic functions, then the induced homomorphisms $f_*, g_* : H_*(X, V) \to$ $H_*(Y, \mathcal{W})$ are equal.

There are many directions in which these results may be· developed in future work, particularly as we expect much of algebraic topology to be transferable to this setting. Some specific possibilities are Hurewicz-type theorems, relations between the homologies in semi-uniform spaces and semi-coarse spaces, excision for semicoarse homology, the development of other semi-coarse homology theories, and the development of semi-coarse invariants which are non-trivial on coarse spaces. In the case of the latter, the second author has constructed a version of the fundamental groupoid for semi-coarse spaces in [\[17\]](#page-55-6) and further showed that it may be used to distinguish coarse structures in addition to semi-coarse structure.

Appendix A. Additional Results on Point-Set Topology of Semi-Coarse Spaces and Their Relations with Semi-Uniform and Coarse Spaces

A.1. Semi-uniform Spaces, Graphs and Roofed Semi-coarse Spaces. We introduce the notion of semi-uniform spaces

Finally, we show how to generate a semi-coarse structure from a semi-uniform space, generalizing Examples [2.1.6.1,](#page-4-1) [2.1.7.1,](#page-4-2) and [2.1.10.1](#page-5-2) above.

Semi-uniform spaces (described in detail in [\[5\]](#page-55-15), Chapter 23) are a generalization of uniform spaces which are no longer necessarily topological. After the following preliminary definition, we recall the definition of semi-uniform spaces and give several examples. We then show how to construct a semi-coarse space from a semiuniform space.

Definition A.1.1. A filter U on a set X is a non-empty collection of subsets of X such that

- (f1) $\emptyset \notin \mathcal{U}$,
- (f2) If $A, A' \in \mathcal{U}$, then $A \cap A' \in \mathcal{U}$,
- (f3) If $A \in \mathcal{U}$ and $A \subset A'$, then $A' \in \mathcal{U}$.

A subcollection \mathcal{U}_0 of \mathcal{U} is a *filter base* of \mathcal{U} iff each element of \mathcal{U} contains some element of \mathcal{U}_0 .

Remark A.1.2. A filter is sometimes defined in the literature without [\(f1\)](#page-42-2) above, in which case a filter which also satisfies [\(f1\)](#page-42-2) is called a proper filter. We will not make this distinction in the present article, and we assume that a filter always satisfies [\(f1\).](#page-42-2)

Definition A.1.3 (Semi-Uniform Space; [\[5\]](#page-55-15), 23 A.3.). Let X be a set and U be a filter on $X \times X$. We call U a semi-uniform structure on X and the pair (X,\mathcal{U}) a semi-uniform space iff

(su1) Each element $U \in \mathcal{U}$ contains the diagonal Δ of X,

(su2) If $A \in \mathcal{U}$, then A^{-1} contains an element of \mathcal{U} ,

Since U is a filter, as noted in [\[5\]](#page-55-15), Axiom [\(su2\)](#page-42-3) may be replaced by (su2') If $A \in \mathcal{U}$, then $A^{-1} \in \mathcal{U}$.

A semi-uniform space (X, \mathcal{U}) is called a *uniform space* iff, in addition to the above, U satisfies

(su3) For every $U \in \mathcal{U}$, there exists a $V \in \mathcal{U}$ such that $V \circ V \subset U$.

When context allows it, we will just represent the semi-uniform space by its set X . Let (X, \mathcal{U}) and (Y, \mathcal{T}) be semi-uniform spaces. We will say that $f : (X, \mathcal{U}) \to$ (Y, \mathcal{T}) is uniformly continuous iff for each $B \in \mathcal{T}$ there exists a $A \in \mathcal{U}$ such that $(f \times f)(A) \subset (B).$

Definition A.1.4. We denote by SUnif the category of semi-uniform spaces and uniformly continuous maps.

Following [\[5\]](#page-55-15), Example 23.A.7, we have the following construction of a semiuniform space from a semi-pseudometric space.

Example A.1.4.1. (1) Let (X, d) be a semi-pseudometric space, and let \mathcal{U}_d be the semi-uniform structure on $X \times X$ generated by the collection of sets ${B_q}_{q>0}$, where $B_q = \{(x, y) : d(x, y) < q\}.$

(2) Given a metric space (X, d) and a positive real number $r > 0$, the construction in [item \(1\)](#page-43-0) above applied to the semi-pseudometrics d_r, d_r^{\lt} , and d_r^{\le} [Defini](#page-5-1)[tion 2.1.9](#page-5-1) gives examples of semi-uniform spaces which are not uniform.

Definition A.1.5. Given a semi-pseudometric space (X, d) , We call \mathcal{U}_d the semiuniform structure induced by the semi-pseudometric d. If, in addition, $r \geq 0$ is a non-negative real number, then we write \mathcal{U}_r for the semi-uniform structure induced by the semi-pseudometric d_r from [Definition 2.1.9.](#page-5-1)

We now show how to build a semi-uniform structure from a semi-coarse structure and vice-versa. We first state the following simple lemma.

Lemma A.1.6. Let X be a set and suppose that $A, B \in \mathcal{P}(X \times X)$. Then

(i) $A = (A^{-1})^{-1}$, and (ii) If $A \subset B$, then $A^{-1} \subset B^{-1}$.

Proof. (i) By definition, we have that

$$
(A^{-1})^{-1} = \{(x, y) : (y, x) \in A^{-1}\}\
$$

$$
= \{(x, y) : (x, y) \in A\}
$$

$$
= A.
$$

(ii) Let $(x, y) \in A^{-1}$, then $(y, x) \in A$, so that $(y, x) \in B$. Therefore, $(x, y) \in B^{-1}$. □

Proposition A.1.7 (Semi-Coarse Space from a Semi-Uniform Space). Let (X, \mathcal{U}) be a semi-uniform space, and define the collection $\mathcal{U}^{\downarrow} \subset \mathcal{P}(X \times X)$ by

$$
\mathcal{U}^{\downarrow} := \left\{ B \subset X \times X : B \subset \bigcap_{A \in \mathcal{U}} A \right\}.
$$

Then $(X, \mathcal{U}^{\downarrow})$ is a semi-coarse space.

Proof. Let (X, \mathcal{U}) and \mathcal{U}^{\downarrow} be as in the statement of the Proposition. We check that the axioms for semi-coarse structures are satisfied by \mathcal{U}^{\downarrow} .

(sc1) By definition of a semi-uniform space, the diagonal $\Delta_X \subset U$ for each $U \in \mathcal{U}$. Therefore, $\Delta_X \in \mathcal{U}^{\downarrow}$.

(sc2) Let $B \in \mathcal{U}^{\downarrow}$ and $B' \subset B$. Then $B' \subset B \subset A$ for each $A \in \mathcal{U}$. So $B' \in \mathcal{U}^{\downarrow}$. (sc3) Let $B, B' \in \mathcal{U}^{\downarrow}$. By definition, we have that $B, B' \subset A$ for every $A \in \mathcal{U}$,

and therefore $B \cup B' \subset A$ for every $A \in \mathcal{U}$ as well. Therefore, $B \cup B' \in \mathcal{U}^{\downarrow}$.

(sc4) Let $B \in \mathcal{U}^{\downarrow}$, then $B \subset A$ for each $A \in \mathcal{U}$. By [Lemma A.1.6,](#page-43-1) we have that $B^{-1} \subset A^{-1}$ for each $A \in \mathcal{U}$, and since $A \in \mathcal{U}$ implies $A^{-1} \in \mathcal{U}$, [Lemma A.1.6](#page-43-1) further implies that $B^{-1} \subset A$ for each $A \in \mathcal{U}$. Therefore, $B^{-1} \in \mathcal{U}^{\downarrow}$. It now follows that $(X, \mathcal{U}^{\downarrow})$ is a semi-coarse space. \square

Definition A.1.8. The semi-coarse structure \mathcal{U}^{\downarrow} in [Proposition A.1.7](#page-43-2) is called the semi-coarse structure induced by the semi-uniform structure U.

We now construct a semi-uniform structure \mathcal{V}^{\uparrow} from a semi-coarse space (X, \mathcal{V}) .

Proposition A.1.9. Let (X, V) be a semi-coarse space, and define the collection $\mathcal{V}^{\uparrow} \subset \mathcal{P}(X \times X)$ by

$$
\mathcal{V}^{\uparrow} = \left\{ U \subset X \times X : \left(\bigcup_{V \in \mathcal{V}} V \right) \subset U \right\}.
$$

Then $(X, \mathcal{V}^{\uparrow})$ is a semi-uniform space.

Proof. Let (X, V) and V^{\uparrow} be as in the statement of the proposition. We first show that \mathcal{V}^{\uparrow} is a filter.

(f1) Since $\Delta_X \in \mathcal{V}, \Delta_X \subset (\cup_{V \in \mathcal{V}} V) \subset A \cap A'$, and, in particular, $\emptyset \neq A \cap A'$.

(f2) Suppose that $A, A' \in \mathcal{V}^{\uparrow}$ Then $(\cup_{V \in \mathcal{V}} V) \subset A \cap A'$ by construction, so $A \cap A' \in \mathcal{V}^{\uparrow}$.

(f3) Suppose that $A \in \mathcal{V}^{\uparrow}$ and $A' \subset X \times X$ such that $A \subset A'$. Then $(\cup_{V \in \mathcal{V}} V) \subset$ $A \subset A'$, and therefore $A' \in \mathcal{V}^{\uparrow}$ as well.

We now prove that \mathcal{V}^{\uparrow} satisfies the axioms of a semi-uniform structure.

(su1) Since $\Delta_X \in \mathcal{V}$, then $\Delta_X \subset (\cup_{V \in \mathcal{V}} V) \subset U$ for each $U \in \mathcal{V}^{\uparrow}$.

(su2') Since $V \in \mathcal{V} \iff V^{-1} \in \mathcal{V}$, it follows that $\bigcup_{V \in \mathcal{V}} V = \bigcup_{V \in \mathcal{V}} V^{-1}$. Now suppose that $A \in \mathcal{V}^{\uparrow}$. Then $\cup_{V \in \mathcal{V}} V = \cup_{V \in \mathcal{V}} V^{-1} \subset A^{-1}$, so $A^{-1} \in \mathcal{V}^{\uparrow}$.

It follows that $(X, \mathcal{V}^{\uparrow})$ is a semi-uniform space.

Definition A.1.10. We call the semi-uniform structure V^{\uparrow} in [Proposition A.1.9](#page-44-1) the Semi-uniform structure induced by the semi-coarse space (X, V) .

Propositions [A.1.7](#page-43-2) and [A.1.9](#page-44-1) motivate the following definitions.

Definition A.1.11. (Roof) Let (X, V) be a semi-coarse space. We call $\mathfrak{R}(V) :=$ $\bigcup_{V \in \mathcal{V}} V$ the roof of V, and we say that the semi-coarse space (X, V) is roofed iff $\mathfrak{R}(\mathcal{V}) \in \mathcal{V}$. Otherwise, we say that (X, \mathcal{V}) is non-roofed.

Example A.1.11.1. (1) The semi-coarse spaces in [Lemma 2.1.5](#page-3-3) are roofed with roof W.

(2) The semi-coarse spaces induced by graphs as defined in [Definition 2.1.7](#page-4-6) are roofed with roof E , the set of edges of the graph.

Proposition A.1.12. Let SCoarse, RSCoarse and Graphs the categories of semi-coarse spaces, roofed semi-coarse spaces and graphs, respectively. Let Φ : $SCoarse \rightarrow Graphs$ be a map of categories defined by

$$
\Phi(X, V) = (V, E),
$$

$$
\Phi(f) = f,
$$

where $V = X$ and $E = \{\{u, v\} \mid (u \neq v)$ and $(u, v) \in V\}.$ Given a graph $G = (V, E)$, define

$$
\mathfrak{R}_E := \Delta_V \cup \left(\bigcup_{\{u,v\} \in E} \{ (u,v), (v,u) \} \right),
$$

and let $\Psi: \mathbf{Graphs} \to \mathbf{SCoarse}$ be the map of categories defined by

$$
\Psi(V, E) = (V, \mathcal{P}(\mathfrak{R}_E))
$$

$$
\Psi(f) = f
$$

where $\mathcal{P}(\mathfrak{R}_E)$ is the power set of \mathfrak{R}_E .

Then Φ and Ψ are functors. Moreover, $\Phi|_{\text{RSCoarse}} \circ \Psi = 1_{\text{Graphs}}$ and $\Psi \circ$ $\Phi_{\text{RSCoarse}} = 1_{\text{RSCoarse}}$, so $\Psi(\text{Graphs}) = \text{RSCoarse}$ and $\Phi(\text{SCoarse}) = \text{Graphs}$.

Proof. We first need to show that Φ maps bornologous functions to graph morphisms, and that Ψ maps graph morphisms to bornologous functions.

Let (X, V) and (Y, W) be semi-coarse spaces, and let $f : X \to Y$ be a bornologous function. We recall that $f: G \to G'$ is a graph morphism iff, for all $(v, v') \in E_G$, either $f(v) = f(v')$ or $(f(v), f(v')) \in E_{G'}$. Let $x, y \in X$ be such that $f(x) \neq f(y)$. Then $\{(f(x), f(y))\}\in W$, so $\{f(x), f(y)\}\in E_Y$ by construction. Thus $\Phi(f)$ is a graph morphism. Since Φ preserves the identity and respects composition, it follows that Φ is a functor from **SCoarse** to **Graphs**.

Now let G and G' be graphs and $f : G \to G'$ a graph morphism. Suppose that $(v, w) \in G_E$. Then $(f(v), f(w)) \in \mathfrak{R}_{E_{G'}}$ by definition. Furthermore, by the definition of \mathfrak{R}_{E_G} and $\mathfrak{R}_{E_{G'}}$, it follows that $f(\mathfrak{R}_{E_G}) \subset \mathfrak{R}_{E_{G'}}$, and therefore f is bornologous. Since Ψ also preserves the identity and respects composition, it follows that Ψ is a functor from **Graphs** to **SCoarse**.

Note that, by construction, $\Psi(V, E)$ is the roofed semi-coarse structure on V with roof \mathfrak{R}_E . Finally, the last sentence of the theorem follows by construction. \Box

Definition A.1.13. Let (X, \mathcal{U}) be a semi-uniform space. We call $\mathfrak{F}(\mathcal{U}) := \bigcap_{U \in \mathcal{U}} U$ the foundation of the semi-uniform structure U . We say that (X, U) is cemented (from the Spanish *cimentado*) iff $\mathfrak{F}(\mathcal{U}) \in \mathcal{U}$. Otherwise, we say that (X, \mathcal{U}) is non-cemented.

Remark A.1.14. Note that the roof is the maximal in the semi-coarse structure with order relation $A < B$ iff $A \subset B$, while the foundation is the minimal in the semi-uniform structure with the same order.

Proposition A.1.15. (1) Let $U \subset X \times X$ such that $\Delta_X \subset U$ and $U = U^{-1}$, and denote by [U] the filter generated by U. Then $(X,[U])$ is a cemented semiuniform space with foundation U.

(2) Let U and T be semi-uniform structures on X. If $\mathfrak{F}(\mathcal{U}) = \mathfrak{F}(\mathcal{T})$ and (X, \mathcal{U}) is cemented, then $\mathcal{T} \subset \mathcal{U}$. It follows that, for each foundation, there exists a unique cemented semi-uniform space with that foundation.

Proof. (1) Let $U \subset X \times X$ such that $\Delta_X \subset U$ and $U = U^{-1}$, and denote by [U] the filter generated by U. Since $\Delta_X \subset U$, then $\Delta_X \subset A$ for all $A \in [U]$. Furthermore, if $A \in [U]$, then $U \subset A$ and $U = U^{-1} \subset A^{-1}$, from which it follows that $A^{-1} \in [U]$. Thus, $(X, [U])$ is a semi-uniform space.

More over, since $U \subset A$ for all $A \in [U]$, then

$$
U\subset \bigcap_{A\in [U]}A\subset U
$$

Thus $(X, [U])$ is a cemented semi-uniform space with foundation [U].

(2) Let U and T be semi-uniform structures on X such that $\mathfrak{F}(\mathcal{U}) = \mathfrak{F}(\mathcal{T})$, and suppose that (X, \mathcal{U}) is cemented. Therefore, for all $B \in \mathcal{T}$, $\mathfrak{F}(\mathcal{U}) \subset B$, and since $\mathfrak{F}(\mathcal{U})$, therefore $B \in \mathcal{U}$. Thus, $\mathcal{T} \subset \mathcal{U}$ and the result follows.

Proposition A.1.16. Let V and T be a semi-coarse structures on a set X. Suppose that V is roofed, and that $\Re(V) = \Re(\mathcal{T})$. Then $\mathcal{T} \subset V$, and for each roof, there exists a unique roofed semi-coarse structure with that roof.

Proof. For all $B \in \mathcal{T}$, $B \subset \mathfrak{R}(\mathcal{T}) = \mathfrak{R}(\mathcal{V}) \in \mathcal{V}$, and therefore $B \in \mathcal{V}$. The result follows. \Box

Proposition A.1.17. Let (X, V) and (Y, W) be semi-coarse spaces, and suppose that (X, V) is roofed. A map $f : X \to Y$ is bornologous iff $(f \times f)(\mathfrak{R}(V)) \in W$.

Proof. Any set $V \in \mathcal{V}$ is a subset of $\mathfrak{R}(\mathcal{V})$ by definition. Therefore, if $(f \times f)(V) \subset$ $(f \times f)(\mathfrak{R}(V)) \in \mathcal{W}$, then $(f \times f)(V) \in \mathcal{W}$, so f is bornologous. Conversely, suppose that f is bornologous. Then $(f \times f)(\mathfrak{R}(\mathcal{V})) \in \mathcal{W}$.

Similarly, for cemented semi-uniform spaces, we have

Proposition A.1.18. Let (X, \mathcal{U}) and (Y, \mathcal{T}) be semi-uniform spaces and suppose that (Y, \mathcal{T}) is cemented. A map $f : X \to Y$ is uniformly continuous iff $(f \times f)$ $f)(\mathfrak{F}(\mathcal{U}))\subset \mathfrak{F}(\mathcal{T}).$

Proof. Let (X, \mathcal{U}) and (Y, \mathcal{T}) be cemented semi-uniform spaces, and let $f : (X, \mathcal{U}) \rightarrow$ (Y, \mathcal{T}) be a uniformly continuous function. Then, for each $B \in \mathcal{T}$ there exists a set $A_B \in \mathcal{U}$ such that $(f \times f)(A_B) \subset B$. Therefore

$$
(f \times f)(\mathfrak{F}(\mathcal{U})) = (f \times f) \left(\bigcap_{A \in \mathcal{U}} A\right) \subset \bigcap_{A \in \mathcal{U}} (f \times f)(A)
$$

$$
\subset \bigcap_{B \in \mathcal{T}} (f \times f)(A_B) \subset \bigcap_{B \in \mathcal{T}} B
$$

$$
= \mathfrak{F}(\mathcal{T}).
$$

Conversely, assume that $(f \times f)(\mathfrak{F}(\mathcal{U})) \subset \mathfrak{F}(\mathcal{T})$. If $B \in \mathcal{T}$, then $\mathfrak{F}(\mathcal{T}) \subset B$, and therefore $(f \times f)(\mathfrak{F}(U)) \subset B$ as well. However, since (X, U) is cemented, we have that $\mathfrak{F}(\mathcal{U}) \in \mathcal{U}$, from which it follows that f is uniformly continuous.

Proposition A.1.19. Let X be a set, and suppose that V and U are semi-coarse and semi-uniform structures on X, respectively. Then

- (i) (X, U^{\downarrow}) is a roofed semi-coarse space.
- (ii) $(X, \mathcal{V}^{\uparrow})$ is a cemented semi-uniform space.
- (iii) If (X, \mathcal{U}) is a cemented semi-uniform space, then $(\mathcal{U}^{\downarrow})^{\uparrow} = \mathcal{U}$.
- (iv) If (X, V) is a roofed semi-coarse space, then $(V^{\uparrow})^{\downarrow} = V$.

Proof. Let X, V, and U be as in the hypothesis of the proposition.

(i) First note that, by definition, $\mathfrak{F}(\mathcal{U}) \in \mathcal{U}^{\downarrow}$. Now suppose that $A \in \mathcal{U}^{\downarrow}$. By construction, $A \subset B$ for every $B \in \mathcal{U}$, and therefore $A \subset \mathfrak{F}(\mathcal{U})$. Therefore, $\mathfrak{R}(\mathcal{U}^{\downarrow}) =$ $\mathfrak{F}(\mathcal{U}) \in \mathcal{U}^{\downarrow}$, and $(X, \mathcal{U}^{\downarrow})$ is a roofed semi-coarse space.

(ii) As before, note that $\mathfrak{R}(\mathcal{V}) \in \mathcal{V}^{\uparrow}$ by definition. Now let $B \in \mathcal{V}^{\uparrow}$. By construction, $A \subset B$ for every $A \in V$, and therefore $\mathfrak{R}(\mathcal{V} \subset B$. It follows that $\mathfrak{F}(\mathcal{V}^{\uparrow}) = \mathfrak{R}(\mathcal{V})$, so $(X, \mathcal{V}^{\uparrow})$ is a cemented semi-uniform space.

(iii) By points [\(i\)](#page-46-1) and [\(ii\)](#page-46-2) above, $\mathfrak{F}(\mathcal{U}) = \mathfrak{R}(\mathcal{U}^{\downarrow}) = \mathfrak{F}(\mathcal{U}^{\downarrow})^{\uparrow}$). Therefore, $\mathcal{U} =$ $(\mathcal{U}^{\downarrow})^{\uparrow}$ by [Proposition A.1.15.](#page-45-0)

(iv) By points [\(i\)](#page-46-1) and [\(ii\)](#page-46-2) above, $\mathfrak{R}(\mathcal{V}) = \mathfrak{F}(\mathcal{V}^{\uparrow}) = \mathfrak{R}(\mathcal{V}^{\uparrow})^{\downarrow}$. Therefore, $\mathcal{V} = (\mathcal{V}^{\uparrow})^{\downarrow}$ by [Proposition A.1.16.](#page-46-3)

□

We now show that the constructions above are functorial. Let **RSCoarse** and CSUnif denote the full subcategories of roofed semi-coarse spaces and cemented semi-uniform spaces, respectively.

Proposition A.1.20. Let Φ : **SCoarse** \to **SUnif** be the map $\Phi(X, V) = (X, V^{\uparrow})$ on objects and $\Phi(f : X \to Y) = f$ on morphisms, and let $\Psi : \mathbf{S}\mathbf{Unif} \to \mathbf{S}\mathbf{Coarse}$ be the map $\Psi(X, \mathcal{U}) = (X, \mathcal{U}^{\downarrow})$ on objects and $\Psi(f : X \to Y) = f$ on morphisms.

Then Φ and Ψ are functors, $\Phi(\text{SCoarse}) = \text{CSUnif}$, and $\Psi(\text{CSUnif}) =$ RSCoarse. Moreover, we have $\Phi|_{\text{RSCoarse}} \circ \Psi|_{\text{CSUnif}} = 1_{\text{CSUnif}}$, and $\Psi|_{\text{CSUnif}} \circ$ $\Phi|_{\rm RSCoarse} = 1_{\rm RSCoarse}.$

Proof. It suffices to prove that Φ maps bornologous function to uniformly continuous functions and that Ψ maps uniformly continuous functions to bornologous functions. The rest of the proposition is a direct consequence of [Proposition A.1.19.](#page-46-4)

Let $f: (X, V) \to (Y, W)$ be a bornologous function. Then $(f \times f)(A) \in W$ for every $A \in \mathcal{V}$. We therefore have that

$$
(f \times f) \left(\bigcup_{A \in \mathcal{V}} A\right) \subset \bigcup_{B \in \mathcal{W}} B.
$$

Thus, the foundation of \mathcal{W}^{\uparrow} contains the image of the foundation of \mathcal{V}^{\uparrow} under $f \times f$, i.e., $(f \times f)(\mathfrak{F}(\mathcal{V}^{\uparrow})) \subset U$ for every $U \in \mathcal{W}^{\uparrow}$. Therefore $f : (X, \mathcal{U}^{\uparrow}) \to (Y, \mathcal{W}^{\uparrow})$ is uniformly continuous by [Proposition A.1.18.](#page-46-5)

Now let $f: (X, \mathcal{U}) \to (Y, \mathcal{T})$ be a uniformly continuous function. By definition, for every $T \in \mathcal{T}$, there exists a $U_T \in \mathcal{U}$ such that $(f \times f)(U_T) \subset T$. We therefore have that

$$
(f \times f)(\mathfrak{F}(\mathcal{U})) = (f \times f) \left(\bigcap_{U \in \mathcal{U}} U\right) \subset \bigcap_{U \in \mathcal{U}} (f \times f)(U)
$$

$$
\subset \bigcap_{T \in \mathcal{T}} (f \times f)(U_T) \subset \bigcap_{T \in \mathcal{T}} T
$$

$$
= \mathfrak{F}(\mathcal{T}).
$$

Therefore, the roof of \mathcal{T}^{\downarrow} contains the image of the roof of U under $f \times f$. It follows that $(f \times f)(A) \subset \mathfrak{R}(Y, \mathcal{T}^{\downarrow}),$ or, equivalently, $(f \times f)(A) \in \mathcal{T}^{\downarrow}$ for every $A \in \mathcal{U}^{\downarrow}$. Therefore, $f : (X, \mathcal{U}^{\downarrow}) \to (Y, \mathcal{T}^{\downarrow})$ is a bornologous function.

In addition to the above, the next result shows that, when restricted to cemented semi-uniform spaces and roofed semi-coarse spaces, the functors in [Propo](#page-47-0)[sition A.1.20](#page-47-0) are adjoints of each other.

Proposition A.1.21. Φ and Ψ satisfy $\Phi \dashv \Psi$, where Φ and Ψ are the functors defined in [Proposition A.1.20.](#page-47-0)

Proof. Let (X, V) be a semi-coarse space, and let (Y, U) be a semi-uniform space. We first show that $\Phi \dashv \Psi$. Let $f \in Hom_{\text{SCoarse}}(X, \Psi(Y))$. Then, by definition, $(f \times f)(A) \in \mathcal{U}^{\downarrow}$ for all $A \in \mathcal{V}$. This equivalent to the statement that $(f \times f)(A) \subset$ $\mathfrak{F}(\mathcal{U})$ for every $A \in \mathcal{V}$. This, in turn, is equivalent to $(f \times f)(\mathfrak{R}(\mathcal{V})) \subset \mathfrak{F}(\mathcal{U})$, since $A \subset \bigcup_{A' \in \mathcal{V}} A'$ for every $A \in \mathcal{V}$. Finally, we have that $(f \times f)(A) \subset \mathfrak{F}(\mathcal{U})$ for every $A \in V$ implies $\cup_{A \in V} (f \times f)(A) = (f \times f)(\cup_{A \in V} A) \subset \mathfrak{F}(\mathcal{U})$. Therefore, $f \in Hom_{\text{SUnif}}(\Phi(X), Y)$ by [Proposition A.1.18.](#page-46-5)

Similarly, let $f \in Hom_{\text{SUnif}}(\Phi(X), Y)$, by definition for every $B \in \mathcal{U}$ there exists $A \in \mathcal{V}^{\uparrow}$ such that $(f \times f)(A) \subset B$. This equivalent to the fact that, for every $B \in \mathcal{U}$ we have $(f \times f)(\Re(\mathcal{V})) \subset B$, by definition of $\Re(\cdot)$ and the construction of \mathcal{U}^{\uparrow} . Since $\mathfrak{F}(\mathcal{U}) = \cap_{B \in \mathcal{U}} B$, this identical $(f \times f)(\mathfrak{R}(\mathcal{V})) \subset \mathfrak{F}(\mathcal{V})$. Therefore, $f \in Hom_{\text{SCoarse}}(X, \Psi(Y))$ by [Proposition A.1.17.](#page-46-0)

It now follows that $Hom_{\text{SCoarse}}(X, \Psi(Y)) \cong Hom_{\text{SUnif}}(\Phi(X), Y)$, so $\Phi \dashv \Psi$. □

A.2. Product Structure in Semi-coarse Spaces. We discuss infinite products of semi-coarse spaces. The next two propositions give semi-coarse analogues of the box product and (Tychonoff) product of topological spaces.

Proposition A.2.1. Let Λ be an index set and $\{(X_{\lambda}, V_{\lambda})\}$ be a collection of semicoarse spaces. Let $\prod_{\lambda \in \Lambda} \mathcal{V}_{\lambda}$ denote the collection of subsets of $(\prod_{\lambda \in \Lambda} X_{\lambda})^2$ that are contained in some set of the form

$$
\left\{ (x,y) \in \left(\prod_{\lambda \in \Lambda} X_{\lambda} \right)^2 : (\pi_{\lambda}(x), \pi_{\lambda}(y)) \in A_{\lambda} \text{ for each } \lambda \in \Lambda \right\}
$$

with $A_{\lambda} \in \mathcal{V}_{\lambda}$.

Then $\prod_{\lambda \in \Lambda} \mathcal{V}_{\lambda}$ is a semi-coarse structure on the set $\prod_{\lambda \in \Lambda} X_{\lambda}$.

Proof. Let Λ , $\{(X_{\lambda}, V_{\lambda})\}_{\lambda \in \Lambda}$, and $\prod_{\lambda \in \Lambda} V_{\lambda}$ be as in the statement of the proposition. We check that $\prod_{\lambda \in \Lambda} \mathcal{V}_{\lambda}$ satisfies the axioms of a semi-coarse structure. We have

(sc1) $\Delta_{(\prod_{\lambda \in \Lambda} X_{\lambda})} = \prod_{\lambda \in \Lambda} \Delta_{X_{\lambda}} \in \prod_{\lambda \in \Lambda} \mathcal{V}_{\lambda}$.

(sc2) If $V \in \prod_{\lambda \in \Lambda} V_{\lambda}$ and W is subset of V, then there are sets $\{A_{\lambda} \in V_{\lambda}\}_{\lambda \in \Lambda}$ such that $W \subset V \subset \prod_{\lambda \in \Lambda} A_{\lambda}$, and therefore $W \in \prod_{\lambda \in \Lambda} V_{\lambda}$.

 $(\text{sc}3)$ If $V, W \in \Pi$ Π $\lambda \in \Lambda$ \mathcal{V}_{λ} , then there are $\{A_{\lambda}, B_{\lambda} \in \mathcal{V}_{\lambda}\}_{\lambda \in \Lambda}$ such that $V \subset$ $\lambda \in \Lambda$ A and $W \subset \prod_{\lambda \in \Lambda} B_{\lambda}$. Therefore, $V \cup W \subset \prod_{\lambda \in \Lambda} (A_{\lambda} \cup B_{\lambda})$, and we conclude that $V \cup W \in \prod_{\lambda \in \Lambda} V_{\lambda}$.

(sc4) If $V \in \prod_{\lambda \in \Lambda} V_{\lambda}$, then there are $\{A_{\lambda} \in V_{\lambda}\}_{\lambda \in \Lambda}$ such that $V \subset \prod_{\lambda \in \Lambda} A_{\lambda}$, and therefore $V^{-1} \subset \prod_{\lambda \in \Lambda} (A_{\lambda})^{-1}$, which gives that $V^{-1} \in \prod_{\lambda \in \Lambda} \mathcal{V}_{\lambda}$.

Definition A.2.2. The semi-coarse structure $\prod_{\lambda \in \Lambda} \mathcal{V}_{\lambda}$ from [Proposition A.2.1](#page-48-1) will be called the *box product semi-coarse structure on* $(\prod_{\lambda \in \Lambda} X_{\lambda})$. We call the pair $\left(\prod_{\lambda\in\Lambda}X_{\lambda},\prod_{\lambda\in\Lambda}\mathcal{V}_{\lambda}\right)$ the box product of the family of semi-coarse spaces $\{(X_{\lambda},\mathcal{V}_{\lambda}\}.$

Proposition A.2.3. Let $\prod_{\lambda \in \Lambda}^{\tau} \mathcal{V}_{\lambda}$ be the collection of subsets of $(\prod_{\lambda \in \Lambda} X_{\lambda})^2$ such they are contained in a set of the form

$$
\left\{ (x,y) \in \left(\prod_{\lambda \in \Lambda} X_{\lambda} \right)^2 \mid (\pi_{\lambda}(x), \pi_{\lambda}(y)) \in A_{\lambda} \text{ if } \lambda \in F, \right\}
$$

$$
(\pi_{\lambda}(x), \pi_{\lambda}(y)) \in \Delta_{X_{\lambda}} \text{ if } \lambda \in \Lambda \backslash F \right\},\
$$

where $F \subset \Lambda$ is a finite set and $A_{\lambda} \in \mathcal{V}_{\lambda}$ for all $\lambda \in \Lambda$.

Then $\prod_{\lambda \in \Lambda}^{\tau} \mathcal{V}_{\lambda}$ is a semi-coarse structure on $\prod_{\lambda \in \Lambda} X_{\lambda}$.

Proof. Let $\prod_{\lambda \in \Lambda}^{\tau} V_{\lambda}$ be as in the statement of the proposition. We check that $\prod_{\lambda \in \Lambda}^{\tau} \mathcal{V}_{\lambda}$ the axioms for semi-coarse structure.

(sc1) $\Delta_{(\prod_{\lambda \in \Lambda} X_{\lambda})} = \prod_{\lambda \in \Lambda} \Delta_{X_{\lambda}} \in \prod_{\lambda \in \Lambda}^{\tau} \mathcal{V}_{\lambda}$.

(sc2) If $V \in \prod_{\lambda \in \Lambda}^{\tau} V_{\lambda}$ and W subset of V, then there are sets $\{V_{\lambda} \in V_{\lambda}\}_{\lambda \in \Lambda}$ such that a finite number of $V_{\lambda} \neq \Delta_{X_{\lambda}}$ and $W \subset V \subset \prod_{\lambda \in \Lambda} V_{\lambda}$, from which it follows that $W \in \prod_{\lambda \in \Lambda}^{\tau} \mathcal{V}_{\lambda}$.

(sc3) If $V, W \in \prod_{\lambda \in \Lambda}^{\infty} V_{\lambda}$, then there are sets $\{V_{\lambda} \in V_{\lambda}\}_{\lambda \in \Lambda}, \{W_{\lambda} \in V_{\lambda}\}_{\lambda \in \Lambda}$ such that a finite number $n \in \mathbb{N}$ of the V_{λ} satisfy $V_{\lambda} \neq \Delta_{X_{\lambda}}$ and a finite number m of the W_{λ} satisfy $W_{\lambda} \neq \Delta_{X_{\lambda}}$, and $V \subset \prod_{\lambda \in \Lambda} V_{\lambda}$, $W \subset \prod_{\lambda \in \Lambda} W_{\lambda}$. Therefore, $V \cup W \subset \prod_{\lambda \in \Lambda} (V_{\lambda} \cup W_{\lambda})$ and there are at most a finite number $m + n$ of unions $V_{\lambda} \cup W_{\lambda}$ which are not equal to the corresponding diagonal $\Delta_{X_{\lambda}}$. It follows that $V \cup W \in \prod_{\lambda \in \Lambda}^{\tau} \mathcal{V}_{\lambda}.$

(sc4) If $V \in \prod_{\lambda \in \Lambda} \mathcal{V}_{\lambda}$, then there are sets $\{V_{\lambda} \in \mathcal{V}_{\lambda}\}_{\lambda \in \Lambda}$ such that for a finite number of them, $V_{\lambda} \neq \Delta_{X_{\lambda}}$, and we have $V \subset \prod_{\lambda \in \Lambda} V_{\lambda}$. It follows that $V^{-1} \subset$ $\prod_{\lambda \in \Lambda} (V_{\lambda})^{-1}$, which implies that $V^{-1} \in \prod_{\lambda \in \Lambda}^{\tau} \mathcal{V}_{\lambda}$.

Definition A.2.4. We call the semi-coarse structure $\prod_{\lambda \in \Lambda}^{\tau} \mathcal{V}_{\lambda}$ from [Proposition A.2.3](#page-49-0) the Tychonoff product of the family $\{\mathcal{V}_\lambda\}_{\lambda \in \Lambda}$, and we call the pair $\left(\prod_{\lambda \in \Lambda} X_\lambda, \prod_{\lambda \in \Lambda}^{\tau} \mathcal{V}_\lambda\right)$ the Tychonoff product of the family of semi-coarse spaces $\{(X_\lambda, \mathcal{V}_\lambda)\}_{\lambda \in \Lambda}$.

The following proposition shows us that the box product is the preferred product for products of infinite families of semi-coarse spaces. Naturally, when $|\Lambda| < \infty$, the box product and the Tychonoff product coincide.

Proposition A.2.5. Let (Y, W) be a semi-coarse space, and let $\{(X_{\lambda}, V_{\lambda})\}_{\lambda \in \Lambda}$ be a collection of semi-coarse spaces indexed by the set Λ . A map $f : (Y, W) \to$ $(\prod_{\lambda \in \Lambda} X_{\lambda}, \prod_{\lambda \in \Lambda} V_{\lambda})$ is bornologous iff $\pi_{\lambda} \circ f : (Y, W) \to (X_{\lambda}, V_{\lambda})$ is bornologous for each $\lambda \in \Lambda$.

Proof. First, suppose that f is bornologous. Since $\pi_{\lambda} : (\prod_{\lambda \in \Lambda} X_{\lambda}, \prod_{\lambda \in \Lambda} V_{\lambda})$ is bornologous for any $\lambda \in \Lambda$, it follows that $\pi_{\lambda} \circ f$ is bornologous for each $\lambda \in \Lambda$, since the composition of bornologous functions is bornologous.

Now suppose that $\pi_{\lambda} \circ f$ is bornologous for each $\lambda \in \Lambda$, and let $V \in \mathcal{W}$. Then $\pi_{\lambda} \circ f(V) \in \mathcal{V}_{\lambda}$. So $\prod_{\lambda \in \Lambda} (\pi_{\lambda} \circ f(V)) \in \prod_{\lambda \in \Lambda} \mathcal{V}_{\lambda}$. Since $f(V) \subset \prod_{\lambda \in \Lambda} (\pi_{\lambda} \circ f(V)),$ f is a bornologous function. \Box

Our final discussion in this section will be about the relation between products and the functors sending semi-coarse spaces to semi-uniform spaces and vice-versa. We first recall the definition of the Cartesian product of semi-uniform spaces.

Definition A.2.6 (Product Semi-Uniform Space; [\[5\]](#page-55-15), 23.D.10.). The product of a family $\{(X_a, \mathcal{U}_a) : a \in A\}$ of semi-uniform spaces, denoted by $\prod_{a \in A} (X_a, \mathcal{U}_a)$ is defined to be the semi-uniform space (X, \mathcal{U}) where X is the cartesian product of the family $\{X_a\}$, and U, called the *product semi-uniformity*, is the collection of subsets of $X \times X$ containing a set of the form

$$
\{(x,y)\in X\times X : a\in F \Rightarrow (\pi_a(x),\pi_a(y))\in U_a\}
$$

where F is a finite subset of A and $U_a \in \mathcal{U}_a$. Sets of the above form are called the canonical elements of the product semi-uniformity.

Theorem A.2.7. Let $\{(X_a, \mathcal{U}_a)\}\$ be a collection of semi-uniform spaces indexed by a set A, V_a the semi-coarse structure induced by \mathcal{U}_a , U the product semi-uniformity and V the semi-coarse structure induced by U. Then, $V = \prod_{a \in A} V_a$, the box product semi-coarse structure.

Proof. Let $\{(X_a, \mathcal{U}_a)\}\$ be a collection of semi-uniform spaces indexed by a set A, V_a the semi-coarse structure induced by \mathcal{U}_a , U the product semi-uniformity and V the semi-coarse structure induced by \mathcal{U} .

Suppose that $V \in V$. Then V is contained in every canonical element of the product semi-uniformity, which implies that $(\pi_a \times \pi_a)(V) \in U_a$ for each $U_a \in \mathcal{U}_a$, and therefore $(\pi_a \times \pi_a)(V) \in \mathcal{V}_a$ for each $a \in A$. We conclude that $V \in \prod_{a \in A} \mathcal{V}_a$.

On the other hand, if $V \in \prod_{a \in A} \mathcal{V}_a$, then $(\pi_a \times \pi_a)(V) \subset U_a$ for each $U_a \in \mathcal{U}_a$. In particular, for every finite subset F of A we have $(\pi_a \times \pi_a)(V) \in U_a$ for each $U_a \in \mathcal{U}_a, a \in F$, and therefore U is contained in each canonical element of the product semi-uniformity. We conclude that $V \in \mathcal{V}$, and the result follows. \Box

Making the cartesian product of two semi-pseudometric spaces, we get the following and last result.

Lemma A.2.8. Let (X, d^X) and (Y, d^Y) be metric spaces, r and s non-negative real numbers. Suppose that d_r^X and d_s^Y are constructed as in [Definition 2.1.9,](#page-5-1) then

(i) Let \mathcal{U}_r^X , \mathcal{U}_s^Y and $\mathcal{U}_{r,s}$ denote the semi-uniform structures induced by d_r^X , d_s^Y (*i*) Let a_r , a_s and $a_{r,s}$ denote the semi-diagorm structures made by a_r , a_s
and $d_{r,s} := \max\{d_r, \delta_s\}$, respectively. Then $\mathcal{U}_r^X \times \mathcal{U}_s^Y = \mathcal{U}_{r,s}$.

(ii) Let \mathcal{V}_r^X , \mathcal{V}_s^Y and $\mathcal{V}_{r,s}$ be the semi-coarse structures induced by d_r^X , d_s^Y and $d_{r,s}$, respectively. Then $\mathcal{V}_r^X \times \mathcal{V}_s^Y = \mathcal{V}_{r,s}$.

Proof. Let (X, d^X) and (Y, d^Y) be metric spaces, let $r, s \geq 0$ be non-negative real numbers, and construct the semi-pseudometrics d_r^X and d_s^Y as in [Definition 2.1.9.](#page-5-1)

(i) Let \mathcal{U}_r^X , \mathcal{U}_s^Y and $\mathcal{U}_{r,s}$ be the semi-uniform structures induced by d_r^X , d_s^Y and $d_{r,s} := \max\{d_r^X, d_s^Y\}$, respectively. First note that, if $W \in \mathcal{U}_r^X \times \mathcal{U}_s^Y$, then, by [Definition A.2.6,](#page-50-0) there are $U \in \mathcal{U}_r^X$ and $U' \in \mathcal{U}_s^Y$ such that $U \boxtimes U' \subset W$. By the definition of the box product \mathbb{Z} , we have that the set $\{(x,x'): d_r^X(x,x')=$ $0\} \boxtimes \{(y,y') : d_s^Y(y,y') = 0\} \subset W$ is equal to $\{(x,y,x',y') : d_{r,s}((x,y),(x',y')) = 0\}$ 0} $\subset W$. It therefore follows that $W \in \mathcal{D}_{r,s}$.

Conversely, if $W \in \mathcal{V}_{r,s}$, then $\{(x, y, x', y') : d_{r,s}((x, y), (x', y')) = 0\} \subset W$, which is equivalent to the condition that $\{(x, x'): d_r^X(x, x') = 0\} \boxtimes \{(y, y') : d_s^Y(y, y') = 0\}$ $0\} \subset W$. Thus, $W \in \mathcal{U}_r^X \times \mathcal{U}_s^Y$.

(ii) Let \mathcal{V}_r^X , \mathcal{V}_s^Y and $\mathcal{V}_{r,s}$ be the semi-coarse structures induced by d_r^X , d_s^Y and $d_{r,s}$, respectively. Then item (ii) in the Lemma follows from [Theorem A.2.7](#page-50-1) and item (i) above. \Box A.3. Coarse Space Induced by a Semi-Coarse Space. In this section, we discuss a method of generating a coarse space from a semi-coarse space. In order to proceed with the construction, we will first discuss direct limits for semi-coarse spaces. We begin by constructing a semi-coarse structure on a disjoint union of semi-coarse spaces.

Proposition A.3.1. Let $\{(X_{\lambda}, \mathcal{V}_{\lambda})\}_{\lambda \in \Lambda}$ be a collection of semi-coarse spaces indexed by the set Λ , and let $\Box_{\lambda \in \Lambda}$ be the collection of sets of the form

$$
\sqcup_{\lambda\in\Lambda}A_\lambda,
$$

where each $A_{\lambda} \in V_{\lambda}$. (Note that any given A_{λ} may be the empty set.) Then $(\sqcup_{\lambda \in \Lambda} X_{\lambda}, \sqcup_{\lambda \in \Lambda} V_{\lambda})$ is a semi-coarse space.

Proof. We show that $\sqcup_{\lambda \in \Lambda} V_{\lambda}$ satisfies the axioms for a semi-coarse structure.

(sc1) $\Delta_{\Box_{\lambda\in\Lambda}X_{\lambda}}=\Box_{\lambda\in\Lambda}\Delta_{\lambda}\in\Box_{\lambda\in\Lambda}\mathcal{V}_{\lambda}.$

(sc2) If $A := \sqcup_{\lambda \in \Lambda} A_{\lambda} \in \sqcup_{\lambda \in \Lambda} V_{\lambda}$ and $W \subset A$, then $W = \sqcup_{\lambda \in \Lambda} W_{\lambda}$, where $W_{\lambda} \subset A_{\lambda}$ for every $\lambda \in \Lambda$. Therefore each $W_{\lambda} \in \mathcal{V}_{\lambda}$, and $W \in \sqcup_{\lambda \in \Lambda} \mathcal{V}_{\lambda}$.

(sc3) If $A, W \in \sqcup_{\lambda \in \Lambda} \mathcal{V}_{\lambda}$, then there are $\sqcup_{\lambda \in \Lambda} A_{\lambda}$ and $\sqcup_{\lambda \in \Lambda} W_{\lambda}$ with $A_{\lambda}, W_{\lambda} \in \mathcal{V}_{\lambda}$ such that $A = \sqcup_{\lambda \in \Lambda} A_{\lambda}$ and $W = \sqcup_{\lambda \in \Lambda} W_{\lambda}$. So $A \cup W \subset \sqcup_{\lambda \in \Lambda} (A_{\lambda} \cup W_{\lambda})$, from which we conclude that $A \cup W \in \sqcup_{\lambda \in \Lambda} \mathcal{V}_{\lambda}$.

(sc4) If $A \in \sqcup_{\lambda \in \Lambda} \mathcal{V}_{\lambda}$, then there is $\sqcup_{\lambda \in \Lambda} A_{\lambda}$ with $A_{\lambda} \in \mathcal{V}_{\lambda}$ such that $A =$ ⊔λ∈ΛAλ. So

$$
A^{-1} = (\sqcup_{\lambda \in \Lambda} A_{\lambda})^{-1} = \sqcup_{\lambda \in \Lambda} (A_{\lambda})^{-1},
$$

and therefore $A^{-1} \in \sqcup_{\lambda \in \Lambda} \mathcal{V}_{\lambda}$.

It follows that $(\sqcup_{\lambda \in \Lambda} X_{\lambda}, \sqcup_{\lambda \in \Lambda} V_{\lambda})$ is a semi-coarse space. □

Definition A.3.2. We call the semi-coarse structure $\sqcup_{\lambda \in \Lambda} \mathcal{V}_{\lambda}$ the *disjoint union* semi-coarse structure, and we call the semi-coarse space $(\sqcup_{\lambda \in \lambda} X_{\lambda}, \sqcup_{\lambda \in \Lambda} V_{\lambda})$ the disjoint union of the semi-coarse spaces $\{X_{\lambda}, \mathcal{V}_{\lambda}\}_{\lambda \in \Lambda}$.

We will now introduce the notion of a directed system in semi-coarse category, which we will use to construct a coarse space from a semi-coarse space.

Definition A.3.3. Let Λ be a directed set. We will call $\{(X_\alpha, \mathcal{V}_\alpha), f_\alpha^\beta, \Lambda\}$ a directed system of semi-coarse spaces if $\{X_\alpha, f_\alpha^\beta, \Lambda\}$ is a directed system of sets, $(X_\alpha, \mathcal{V}_\alpha)$ are semi-coarse spaces for each $\alpha \in \Lambda$, and each f_{α}^{β} is bornologous.

Proposition A.3.4. Let Λ be a directed set, let $\{(X_\alpha, \mathcal{V}_\alpha), f_\alpha^\beta, \Lambda\}$ be a directed system of semi-coarse spaces, and let \sim be the equivalence relation on $\sqcup_{\lambda \in \Lambda} X_{\lambda}$ such that for $x^{\alpha} \in X^{\alpha}$ and $x^{\beta} \in X^{\beta}$, $x^{\alpha} \sim x^{\beta}$ iff there is $a \gamma \in \Lambda$ satisfying $\alpha \leq \gamma$, $\beta \leq \gamma$ and where $f_{\alpha}^{\gamma} x^{\alpha} = f_{\beta}^{\gamma} x^{\beta}$. Then

$$
\lim_{\rightarrow} \{ (X_{\alpha}, \mathcal{V}_{\alpha}), f_{\alpha}^{\beta}, \Lambda \} = ((\sqcup_{\lambda \in \Lambda} X_{\lambda})/\sim, (\sqcup_{\lambda \in \Lambda} \mathcal{V}_{\lambda})/\sim),
$$

where the left hand side of the above equation is the direct limit of the directed system, and the right hand side is the quotient semi-coarse space from [Example 2.4.7.1](#page-12-0)

$$
\mathbb{L}^2
$$

Proof. It is enough to show that, for any (Y, W) and a collection of diagrams of solid arrows of the form

where $\alpha, \lambda \in \Lambda$ satisfy $\lambda \leq \alpha$ and the solid arrows are bornologous and commute, there exists a bornologous map

$$
g:((\sqcup_{\lambda\in\Lambda}X_{\lambda})/\sim,(\sqcup_{\lambda\in\Lambda}\mathcal{V}_{\lambda})/\sim)\to(Y,\mathcal{W})
$$

making the entire diagram commute for any choice of $\alpha, \lambda \in \Lambda$ with $\lambda \leq \alpha$. However, the existence of a set map g making the diagram commute is guaranteed by the fact that $\sqcup_{\lambda \in \Lambda} X_{\lambda}$ is the direct limit of the directed system viewed as sets, and g is bornologous by [Proposition 2.4.4.](#page-11-0) \Box

We now characterize the direct limit of some special direct systems which we will use to construct a coarse structure from a semi-coarse structure on a set.

Corollary A.3.5. Let X be a set, suppose that Λ is a totally ordered set, and let $\{\mathcal{V}_\lambda\}_{\lambda \in \Lambda}$ be a family of semi-coarse structures on X satisfying $\mathcal{V}_\lambda \subset \mathcal{V}_{\lambda'}$ if $\lambda \leq \lambda'$. Denote by $\{(X_\alpha, \mathcal{V}_\alpha), i_\alpha^\beta, \Lambda\}$ the directed system such that $X = X_\alpha$ for all $\alpha \in \Lambda$ and the maps i_{α}^{β} are all identity maps.

Then the direct limit of $\{(X_{\alpha},\mathcal{V}_{\alpha}),i_{\alpha}^{\beta},\Lambda\}$ is the space $(X, \bigcup$ $\bigcup_{\lambda \in \Lambda} \mathcal{V}_{\lambda}$.

Proof. We note that $X = (\sqcup_{\lambda \in \Lambda} X_{\lambda})/\sim$ and $\bigcup_{\lambda \in \Lambda} V_{\lambda} = (\sqcup_{\lambda \in \Lambda} V_{\lambda})/\sim$. The result now follows from [Proposition A.3.4.](#page-51-1) □

Corollary A.3.6. Let $\{(X_{\alpha}, V_{\alpha}), i_{\alpha}^{\beta}, \Lambda\}$ be the directed system such that $X = X_{\alpha}$ for all $\alpha \in \Lambda$, the i_{α}^{β} are the identity maps, and, in addition, suppose that the following condition holds.

(1) $\forall \lambda_1, \lambda_2 \in \Lambda \exists \lambda_3 \in \Lambda \text{ such that } V \circ W \in \mathcal{V}_{\lambda_3} \forall V \in \mathcal{V}_{\lambda_1}, W \in \mathcal{V}_{\lambda_2}.$

Then $\lim_{\longrightarrow} \mathcal{V}_{\lambda}$ is a coarse structure on X.

Proof. We must show that the semi-coarse structure $(\sqcup_{\lambda \in \Lambda} \mathcal{V}_{\lambda})/\sim$ satisfies axiom [\(sc5\)](#page-3-4) in [Definition 2.1.2,](#page-3-2) but this follows immediately from Condition [\(1\)](#page-52-0) in the statement of the corollary. \Box

Given a semi-coarse space (X, V) , we will proceed to construct a coarse space by repeatedly adding sets of the form $V \circ W$ to the semi-coarse structure $\mathcal V$. We first show that adding $\{V \circ W \mid V, W \in \mathcal{V}\}\)$ to a semi-coarse structure V gives another semi-coarse structure. To do so, we will require the following lemma about properties of the set product.

Lemma A.3.7. Let X and Y be sets and let $A, B, C, D \in \mathcal{P}(X \times X)$. Then

(i) $(A \circ B)^{-1} = B^{-1} \circ A^{-1}$. (ii) $(A \circ B) \cup (C \circ D) \subset (A \cup C) \circ (B \cup D).$ (iii) Let $f: X \to Y$ be a function of sets. Then

$$
(f \times f)(A \circ B) \subset (f \times f)(A) \circ (f \times f)(B).
$$

Proof. Let X and Y be sets and let $A, B, C, D \in \mathcal{P}(X \times X)$. Then

(i) Let $(x, y) \in (A \circ B)^{-1}$, that is, $(y, x) \in A \circ B$. This is equivalent to the existence of a point $z \in X$ such that $(y, z) \in A$ and $(z, x) \in B$. This, in turn, is true iff $(x, z) \in B^{-1}$ and $(z, y) \in A^{-1}$. By definition, we have $(x, y) \in B^{-1} \circ A^{-1}$.

(ii) Let $(x, y) \in (A \circ B) \cup (C \circ D)$. Then $(x, y) \in A \circ B$ or $(x, y) \in C \circ D$, and therefore there exists a point $z \in X$ such that $((x, z) \in A$ and $(z, y) \in B$ or $((x, z) \in C$ and $(z, y) \in D)$, which implies that $(x, z) \in A \cup C$ and $(z, y) \in B \cup D$. Thus, $(x, y) \in (A \cup C) \circ (B \cup D)$.

(iii) Suppose that $(a', b') \in (f \times f)(A \circ B)$. Then there exists an element pairs $(a, b) \in A \circ B$ such that $f(a) = a'$ and $f(b) = b'$, and therefore there exists $x \in X$ with $(a, x) \in A$ and $(x, b) \in B$. It follows that $(a', f(x)) \in (f \times f)(A)$ and $(f(x), b') \in (f \times f)(B)$, which implies that $(a', b') \in (f \times f)(A) \circ (f \times f)(B)$. Therefore, $(f \times f)(A \circ B) \subset (f \times f)(A) \circ (f \times f)(B)$, as desired. \square

Remark A.3.8. In [item \(ii\)](#page-53-0) of the previous lemma, the other inclusion is not necessarily true: Let A be a non-empty set, $D = A^{-1}$ and $B = C = \emptyset$, then $A \circ B = C \circ D = \emptyset$ and $(A \cup C) \circ (B \cup D) = A \circ A^{-1} \neq \emptyset$, so that $(A \circ B) \cup (C \circ D) \not\supseteq$ $(A \cup C) \circ (B \cup D).$

Proposition A.3.9. Let (X, V) be a semi-coarse space, and define

$$
\mathcal{V}^{PE} := \{ C \subset X \times X \mid \exists A, B \in \mathcal{V} \text{ with } C \subset A \circ B \}.
$$

Then (X, V^{PE}) is a semi-coarse space, and $V \subset V^{PE}$.

Proof. We first verify that V^{EP} satisfies the axioms of a semi-coarse structure.

(sc1) We observe that $\Delta_X = \Delta_X \circ \Delta_X$. Thus $\Delta_X \in \mathcal{V}^{PE}$.

(sc2) Let $B \in \mathcal{V}^{PE}$ and suppose that $A \subset B$. Then there are sets $A', B' \in \mathcal{V}$ such that $A \subset B \subset A' \circ B'$, so it follows that $A \in \mathcal{V}^{PE}$.

(sc3) Let $A, B \in \mathcal{V}^{PE}$. Then there are $A', B', A'', B'' \in \mathcal{V}$ such that $A \subset A' \circ A''$ and $B \subset B' \circ B''$. Since $A' \cup B', A'' \cup B'' \in V$, we have that $(A' \cup B') \circ (A'' \cup B'') \in$ \mathcal{V}^{EP} . By [Lemma A.3.7,](#page-52-1) it follows that $A \cup B \subset (A' \cup B') \circ (A'' \cup B'') \in \mathcal{V}^{PE}$. We conclude that $A \cup B \in \mathcal{V}^{PE}.$

(sc4) Let $C \in \mathcal{V}^{PE}$. Then there are $A, B \in \mathcal{V}$ such that $C \subset A \circ B$. Since $A^{-1}, B^{-1} \in V$, [Lemma A.3.7](#page-52-1) implies that $C^{-1} \subset B^{-1} \circ A^{-1}$, and we conclude that $C^{-1} \in \mathcal{V}^{PE}.$

Thus (X, V^{PE}) is a semi-coarse space.

Finally, for any $A \in \mathcal{V}$, $A \circ \Delta_X = A$, and therefore $A \in \mathcal{V}^{PE}$.

Definition A.3.10. Let (X, V) be a semi-coarse space. We call the structure V^{EP} in [Proposition A.3.9](#page-53-1) the set product extension of V, and the ordered pair (X, V^{EP}) will be called the set product extension of (X, V) . For any $k \in \mathbb{N}$, we recursively define \mathcal{V}^{kPE} to be the set product extension of $\mathcal{V}^{(k-1)PE}$.

Proposition A.3.11. If $f : (X, V) \to (Y, W)$ is bornologous, then $f : (X, V^{PE}) \to$ (Y, W^{PE}) is bornologous.

Proof. If $A \in \mathcal{V}^{PE}$, then there exist sets $A', A'' \in \mathcal{V}$ such that $A \subset A' \circ A''$, so $(f \times f)(A'), (f \times f)(A'') \in \mathcal{W}$. By [Lemma A.3.7](#page-52-1) [item \(iii\),](#page-53-2) we have

$$
(f \times f)(A) \subset (f \times f)(A' \circ A'') \subset (f \times f)(A') \circ (f \times f)(A'') \in \mathcal{V}^{PE}.
$$

Therefore, $(f \times f)(A) \in \mathcal{V}^{PE}$ and $f: X \to Y$ is a $(\mathcal{V}^{EP}, \mathcal{W}^{EP})$ -bornologous function. \Box

Remark A.3.12. Note that [Proposition A.3.11](#page-53-3) shows that the map

 $\Psi(f) = f$

$$
\Psi : \text{SCoarse} \to \text{SCoarse}
$$

$$
\Psi(X, \mathcal{V}) = (X, \mathcal{V}^{PE})
$$

is a functor.

As a final observation, we note that the set product of roofed semi-coarse spaces takes a special form.

Remark A.3.13. Let (X, V) be a roofed semi-coarse space. Then, for every $A, B \in$ $\mathcal{V}, A \circ B \subset \mathfrak{R}(X, \mathcal{V}) \circ \mathfrak{R}(X, \mathcal{V})$. Thus, (X, \mathcal{V}^{EP}) is a roofed semi-coarse space and $\mathfrak{R}(X,\mathcal{V}^{EP}) = \mathfrak{R}(X,\mathcal{V}) \circ \mathfrak{R}(X,\mathcal{V}).$

We are now ready to construct a coarse structure from a semi-coarse space (X, V) .

Theorem A.3.14. Let (X, V) be a semi-coarse space, and let $\{(X, V^{kPE}), f_i^j, \mathbb{N}\}_{k=0}^{\infty}$ be the directed system of semi-coarse structures such that all the $f_i^j : (X, V^{iPE}) \rightarrow$ (X, \mathcal{V}^{jPE}) are the identity map. Then $\underline{\lim} \{ (X, \mathcal{V}^{kPE}), f_i^j, \mathbb{N} \}$ is a coarse space.

Proof. This follows from [Corollary A.3.6,](#page-52-2) the definition of V^{kPE} , and the fact that $\mathcal{V}^{iPE} \subset \mathcal{V}^{jPE}$ for any $i < j$.

Definition A.3.15. We call the coarse structure constructed in [Theorem A.3.14](#page-54-0) the coarse structure induced by \mathcal{V} , and which we denote by \mathcal{V}^{∞} .

Let (X, d) be a metric space, and recall that the set $\mathcal{V}_r \subset X \times X$ is defined by

$$
\mathcal{V}_r = \{(x, y) \mid d(x, y) \le r\}.
$$

Now denote by $\mathcal{V}_{\infty} := \lim_{\to \infty} \mathcal{V}_r$, the direct limit of the directed system in [Corol](#page-52-3)[lary A.3.5.](#page-52-3)

We now have

Corollary A.3.16. (X, V_∞) is a coarse space.

Proof. Let $\Lambda = (\mathbb{R}, \leq),$ let $\{\mathcal{V}_r, f_r^{r'}, \Lambda\}$ be the directed system given by the collection of semi-coarse structures induced by a metric d. For any $r \leq r'$, define the map $f_r^{r'} = i : (X, \mathcal{V}_r) \to (X, \mathcal{V}_{r'})$. By definition, $\mathcal{V}_r \subset \mathcal{V}_{r'}$ for $r \leq r'$, and \mathcal{V}_{∞} is a semi-coarse structure by [Corollary A.3.5.](#page-52-3)

Now suppose that $V, W \in V_{\infty}$. Then there are $r, r' \in [0, \infty)$ such that $V \in V_r$ and $W \in \mathcal{V}_{r'}$, Without loss of generality, $r \leq r'$, and therefore $V, W \in \mathcal{V}_{r'}$.

So, if $(x, y) \in V \circ W$, then there is $z \in X$ such that $(x, z) \in V$ and $(z, y) \in W$, thereby $d(x, z) \leq r'$ and $d(z, y) \leq r'$. By the triangle inequality property, $d(x, y) \leq r'$ 2r ′ , and therefor,

$$
V \circ W \subset \{(x, y) \in X \times X : d(x, y) \le 2r'\}.
$$

We conclude that $V \circ W \in \mathcal{V}_{2r'}$, and by [Corollary A.3.6,](#page-52-2) $(X, \mathcal{V}_{\infty})$ is a coarse space. \Box

Acknowledgements

The authors would like to thank Omar Antolín Camarena, Noe Barcenas, Jesús Rodríguez, and Alejandra Trujillo for helpful comments on earlier versions of the manuscript. We are also grateful to have had the opportunity to present parts of this work at the annual meeting of the Sociedad Matemática Mexicana, the Topological Data Analysis Workshop at the Institute for Mathematical and Statistical Innovation, and the GEOTOP-A Seminar. Finally, we would like to thank the anonymous referee for a number of useful comments which helped to improve the manuscript.

REFERENCES

- [1] Eric Babson, Hélène Barcelo, Mark de Longueville, and Reinhard Laubenbacher, Homotopy *theory of graphs*, J. Algebraic Combin. **24** (2006), no. 1, 31–44. MR2245779
- [2] Laurence Boxer and P. Christopher Staecker, Fundamental groups and Euler characteristics of sphere-like digital images, Appl. Gen. Topol. 17 (2016), no. 2, 139–158. MR3562604
- $[3]$, Homotopy relations for digital images, Note Mat. 37 (2017), no. 1, 99-126. MR3733806
- $__$, Remarks on pointed digital homotopy, Topology Proc. 51 (2018), 19–37. MR3633236
- [5] Eduard Čech, *Topological spaces*, Revised edition by Zdeněk Frolíc and Miroslav Katětov. Scientific editor, Vlastimil Pták. Editor of the English translation, Charles O. Junge, Publishing House of the Czechoslovak Academy of Sciences, Prague; Interscience Publishers John Wiley & Sons, London-New York-Sydney, 1966. MR0211373
- [6] Cornelia Druty and Michael Kapovich, *Geometric group theory*, American Mathematical Society Colloquium Publications, vol. 63, American Mathematical Society, Providence, RI, 2018. With an appendix by Bogdan Nica. MR3753580
- [7] Greg Lupton, John Oprea, and Nicholas A. Scoville, The digital Hopf construction, Topology Appl. 326 (2023), Paper No. 108405, 19. MR4533985
- [8] Gregory Lupton, John Oprea, and Nicholas A. Scoville, A fundamental group for digital images, J. Appl. Comput. Topol. 5 (2021), no. 2, 249–311. MR4259435
- [9] , Homotopy theory in digital topology, Discrete Comput. Geom. 67 (2022), no. 1, 112– 165. MR4356246
- [10] $\qquad \qquad$, Subdivision of maps of digital images, Discrete Comput. Geom. 67 (2022), no. 3, 698–742. MR4393077
- [11] Gregory Lupton and Nicholas A. Scoville, Digital fundamental groups and edge groups of *clique complexes*, J. Appl. Comput. Topol. 6 (2022), no. 4, 529–558. MR4496690
- $[12]$ Antonio Rieser, Cech closure spaces: A unified framework for discrete and continuous homo $topy$, Topology and its Applications 296 (2021), 107613.
- [13] $\frac{1}{13}$ Vietoris-rips homology theory for semi-uniform spaces, 2021. submitted, arXiv:2008.05739.
- [14] John Roe, Lectures on coarse geometry, University Lecture Series, vol. 31, American Mathematical Society, Providence, RI, 2003. MR2007488
- [15] P. Christopher Staecker, Digital homotopy relations and digital homology theories, Appl. Gen. Topol. 22 (2021), no. 2, 223–250. MR4359765
- [16] Jonathan Treviño-Marroquín, Espacios pseuso-coarse, Master's Thesis, 2019.
- [17] , Semi-coarse spaces: Fundamental groupoid and the van kampen theorem, 2024.
- [18] Nicolò Zava, Generalisations of coarse spaces, Topology Appl. 263 (2019), 230-256. MR3968132