Thompson Sampling for High-Dimensional Sparse Linear Contextual Bandits

Sunrit Chakraborty * Saptarshi Roy * Ambuj Tewari

University of Michigan, Ann Arbor {sunritc, roysapta, tewaria}@umich.edu

January 31, 2023

Abstract

We consider the stochastic linear contextual bandit problem with high-dimensional features. We analyze the Thompson sampling algorithm using special classes of sparsity-inducing priors (e.g., spike-and-slab) to model the unknown parameter and provide a nearly optimal upper bound on the expected cumulative regret. To the best of our knowledge, this is the first work that provides theoretical guarantees of Thompson sampling in high-dimensional and sparse contextual bandits. For faster computation, we use variational inference instead of Markov Chain Monte Carlo (MCMC) to approximate the posterior distribution. Extensive simulations demonstrate the improved performance of our proposed algorithm over existing ones.

Keywords: linear contextual bandithigh-dimension sparsity Thompson sampling slab-and-spike prior variational Bayes.

1 Introduction

Sequential decision-making, including bandits problems and reinforcement learning, has been one of the most active areas of research in machine learning. It formalizes the idea of selecting actions based on current knowledge to optimize some long term reward over sequentially collected data. On the other hand, the abundance of personalized information allows the learner to make decisions while incorporating this contextual information, a setup that is mathematically formalized as contextual bandits. Moreover, in the big data era, the personal information used as contexts often has a much larger size, which can be modeled by viewing the contexts as high-dimensional vectors. Examples of such models cover internet marketing and treatment assignment in personalized medicine, among many others.

A particularly interesting special case of the contextual bandit problem is the linear contextual bandit problem, where the expected reward is a linear function of the features (Abe et al., 2003;

^{*}Alphabetical order: First two authors have equal contribution.

Auer, 2002). Under this setting, Dani et al. (2008), Chu et al. (2011) and Abbasi-Yadkori et al. (2011) showed polynomial dependence of the cumulative regret on ambient dimension d and time horizon T in low dimensional case. Specifically, Dani et al. (2008) and Abbasi-Yadkori et al. (2011) proved a regret upper bound scaling as $O(d\sqrt{T})$, while Chu et al. (2011) showed a regret upper bound of the order $O(\sqrt{dT})$. It is worthwhile to mention that all of the aforementioned algorithms fall under a certain class of algorithms known as upper confidence bound (UCB) type algorithms that rely on the construction of a specific confidence set for the unknown parameter. In contrast, Thompson Sampling (TS) maintains uncertainty about the unknown parameter in the form of a posterior distribution. The first TS algorithm under this setting was proposed by Agrawal and Goyal (2013) where they established a regret bound of the order $O(d^2\delta^{-1}\sqrt{T^{1+\delta}})$ for any $\delta \in (0, 1)$.

There is also a large body of work present in high-dimensional sparse linear contextual bandit setup, where the reward only depends on a small subset of features of the observed contexts. This area has recently attracted considerable attention due to its abundance in modern reinforcement learning applications (e.g, clinical trials, personalized recommendation systems, etc.) and has quite naturally spawned theoretical research in this direction. To mention a few, the LASSObandit algorithm in Bastani and Bayati (2020), doubly robust-LASSO (Kim and Paik, 2019) and MCP-bandit algorithm in Wang et al. (2018) are shown to have a regret bound with logarithmic dependence on d. Hao et al. (2020) proposed explore the sparsity and then commit (ESTC) algorithm which also enjoys the regret scaling as $O(T^{2/3} \log d)$ in "data poor" regime, i.e., when $T \ll d$. Very recently, Chen et al. (2022) proposed sparse-LinUCB algorithm and sparse-SupLinUCB algorithm for high dimensional contextual bandits with linear rewards. Both of the algorithms are based on the best subset selection approach and enjoy $O(\sqrt{T})$ scaling in the time horizon and polylogarithmic dependence in the ambient dimension d. However, there has been very limited work dedicated to analyzing TS algorithms in high-dimensional sparse bandit setups. Hao et al. (2021) proposed a sparse information-directed sampling (IDS) algorithm which under a special case reduces to a TS algorithm based on a spike-and-slab Gaussian-Laplace prior. However, the regret bound of IDS scales polynomially in d, which is sub-optimal in the high-dimensional regime. In related work, Gilton and Willett (2017) proposed a linear TS algorithm based on a relevance vector machine (RVM) which again suffers from sub-optimal dependence on d.

In this paper, we specifically focus on the high-dimensional sparse linear contextual bandit (SLCB) setup and propose a TS algorithm based on a sparsity-inducing prior that enjoys almost dimension-independent regret bound. While TS algorithms have been known to empirically perform better than optimism-based algorithms (Chapelle and Li, 2011; Kaufmann et al., 2012), theoretical understanding of these is challenging due to the complex dependence structure of the bandit environment. Moreover, posterior sampling in high-dimensional regression (which is the crucial step for TS in high-dimensional SLCB), using MCMC, generally suffers from computational bottleneck. Our work overcomes all these challenges and makes the following contributions:

- 1. We use the sparsity inducing prior proposed in Castillo et al. (2015) for posterior sampling and establish posterior contraction result for *non-i.i.d. observations* coming from bandit environment and for a wide class of noise distributions.
- 2. Using the posterior contraction result, we establish an almost *dimension free* regret bound for our proposed TS algorithm under different arm-separation regimes parameterized by ω . The algorithm enjoys minimax optimal performance for $\omega \in [0, 1)$. To the best of our knowledge,

this is the first work that proposes a novel TS algorithm with desirable regret guarantees in high-dimensional and sparse SLCB setup.

- 3. Our algorithm does *not* need the knowledge of model sparsity level, unlike other algorithms such as LASSO-bandit, MCP-bandit, ESTC, etc.
- 4. Finally, the prior allows us to design a computationally efficient TS algorithm based on Variational Bayes.

The rest of the paper is organized as follows. In Section 2, we introduce the problem formally and discuss the assumptions and prior distribution. In Section 3, we present the crucial posterior contraction result and the main regret bound for our proposed algorithm. In Section 4, we discuss the challenges of drawing samples from the posterior in such problems and present a faster alternative relying on variational inference. In Section 5, we present simulation studies under different setups comparing our proposed method with existing algorithms. Detailed proofs of results and technical lemmas are deferred to the appendix.

Notation: Let \mathbb{R} and \mathbb{R}^+ denote the set of real numbers and the set of non-negative real numbers respectively. Denote by \mathbb{R}^p the *p*-dimensional Euclidean space and $\mathbb{S}^{p-1} := \{x \in \mathbb{R}^p \mid ||x||_2 = 1\}$ the (p-1)-dimensional unit sphere. For a positive integer *K*, denote by [K] the set $\{1, 2, \ldots, K\}$. Regarding vectors and matrices, for a vector $v \in \mathbb{R}^p$, we denote by $||v||_0$, $||v||_1$, $||v||_2$, $||v||_\infty$ the $\ell_0, \ell_1, \ell_2, \ell_\infty$ norms of *v* respectively. We use \mathbb{I}_p to denote the *p*-dimensional identity matrix. For a $p \times q$ matrix *M* we define the $||M|| := \max_{j \in [q]} \sqrt{(M^\top M)_{j,j}}$.

Regarding distributions, $N(\mu, \sigma)$ denotes the Gaussian distribution with mean μ and standard deviation σ , Lap (λ) denotes the Laplace distribution with density $f(x) = (\lambda/2) \exp(-\lambda |x|)$, and Beta(a, b) denotes the beta distribution with parameters a, b.

Throughout the paper, let $O(\cdot)$ denote the standard big-O notation, i.e., we say $a_n = O(b_n)$ if there exists a universal constant C > 0, such that $a_n \leq Cb_n$ for all $n \in \mathbb{N}$. Sometimes for notational convenience, we write $a_n \leq b_n$ in place of $a_n = O(b_n)$. We write $a_n \asymp b_n$ or $a_n = \Theta(b_n)$ if $a_n = O(b_n)$ and $b_n = O(a_n)$.

2 Problem Formulation

We consider a linear stochastic contextual bandit with K arms. At time $t \in [T]$, context vectors $\{x_i(t)\}_{i \in [K]}$ are revealed for every arm i. We assume $x_i(t) \in \mathbb{R}^d$ for all $i \in [K], t \in [T]$ and for every $i, \{x_i(t)\}_{t \in [T]}$ are i.i.d. from some distribution \mathcal{P}_i . At every time step t, an action $a_t \in [K]$ is chosen by the learner and a reward r(t) is generated according to the following linear model:

$$r(t) = x_{a_t}(t)^\top \beta^* + \epsilon(t) \tag{1}$$

where $\beta^* \in \mathbb{R}^d$ is the unknown true signal and $\{\epsilon(t)\}_{t\in[T]}$ are independent sub-Gaussian random noise, also independent of all the other processes. We assume that the true parameter β^* is s^* -sparse, i.e., $\|\beta^*\|_0 = s^*$. We denote by S^* the true support of β^* , i.e., $S^* = \{j : \beta_j^* \neq 0\}$.

The goal is to design a sequential decision-making policy π that maximizes the expected cumulative reward over the time horizon. To formalize the notion, we define the history \mathcal{H}_t up to time t as follows:

$$\mathcal{H}_t := \left\{ (a_{\tau}, r(\tau), \{x_i(\tau)\}_{i \in [K]}) : \tau \in [t] \right\},\$$

and an admissible policy π generates a sequence of random variables a_1, a_2, \ldots taking values in [K] such that a_t is measurable with respect to the σ -algebra generated by the previous feature vectors from each arm, observed rewards of the chosen arms till the previous round and the current feature vectors, i.e., measurable with respect to the filtration $\mathcal{F}_t := \sigma (x_{a_\tau}(\tau), r(\tau), x_i(t); \tau \in [t-1], i \in [K])$.

Thus, an algorithm for contextual bandits is a policy π , which at every round t, chooses an action (arm) a_t based on history \mathcal{H}_{t-1} and current contexts. We note that although contexts of the previous round corresponding to arms that were not chosen are in \mathcal{F}_t , however, they do not provide useful information on the parameter, since we do not observe rewards corresponding to them under the bandit feedback, and hence are not included in the history \mathcal{H}_t . To measure the quality of performance, we compare it with the oracle policy π^* which uses the knowledge of the true β^* to choose the optimal action $a_t^* := \arg \max_{i \in [K]} x_i(t)^\top \beta^*$. Define $\Delta_i(t)$ to be the difference between the mean rewards of the optimal arm and *i*th arm at time t, i.e., $\Delta_i(t) = x_{a_t^*}(t)^\top \beta^* - x_i(t)^\top \beta^*$. Note that under the random-design assumption, a_t^* is also random. Then the regret at time t is defined as regret $(t) = \Delta_{a_t}(t)$ and the objective of the learner is to minimize the total regret till time T, defined as $R(T) = \sum_{t \in [T]} \text{regret}(t)$. We also define the matrix $X_t := (x_{a_1}(1), \ldots, x_{a_t}(t))^\top$. The time horizon T is finite but possibly unknown, but much smaller compared to the ambient dimension of the parameter, i.e. $d \gg T$. Hao et al. (2021) refers to this regime as "data-poor" regime; such a regime adds an extra layer of hardness on top of the difficulty incurred by the sparse structure of β . We also assume that K is fixed and much smaller compared to both d and T.

2.1 Assumptions

In this section, we discuss the assumptions of our model.

Definition 2.1 (Sparse Riesz Condition (SRC)). Let M be a $d \times d$ positive semi-definite matrix. The maximum and minimum sparse eigenvalues of M with parameter $s \in [d]$ are defined as follows:

$$\phi_{\min}(s; M) := \inf_{\delta: \delta \neq 0, \|\delta\|_0 \le s} \frac{\delta^{\top} M \delta}{\|\delta\|_2^2},$$
$$\phi_{\max}(s; M) := \sup_{\delta: \delta \neq 0, \|\delta\|_0 \le s} \frac{\delta^{\top} M \delta}{\|\delta\|_2^2}.$$

We say M has Sparse Riesz Condition if $0 < \phi_{\min}(s, M) \le \phi_{\max}(s; M) < \infty$.

Now we are ready to state the assumptions on the context distributions, which are as follows:

Assumption 2.1 (Assumptions on Context Distributions). *We assume that*

- (a) For some constant $x_{\max} \in \mathbb{R}^+$, we have that for all $i \in [K]$, $\mathcal{P}_i(||x||_{\infty} \leq x_{\max}) = 1$.
- (b) For all arms $i \in [K]$ the distribution \mathcal{P}_i is sub-Gaussian, i.e., $\max_{i \in [K]} \mathbb{E}_{x \sim \mathcal{P}_i} \{ \exp(tu^\top x) \} \le \exp(\vartheta^2 t^2/2), \text{ for all } t \in \mathbb{R} \text{ and } u \in \mathbb{S}^{d-1}.$
- (c) There exists a constant $\xi \in \mathbb{R}^+$ such that for each $u \in \mathbb{S}^{d-1} \cap \{v \in \mathbb{R}^d : \|v\|_0 \leq Cs^*\}$ and $h \in \mathbb{R}^+ \mathcal{P}_i(\langle x, u \rangle^2 \leq h) \leq \xi h$, for all $i \in [K]$, where $C \in (2, \infty)$.
- (d) The matrix $\Sigma_i := \mathbb{E}_{x \sim \mathcal{P}_i}[xx^\top]$ has bounded maximum sparse eigenvalue, i.e., $\phi_{\max}(Cs^*, \Sigma_i) \le \phi_u < \infty$, for all $i \in [K]$, where C is the same constant as in part (c).

Assumption 2.1(a) basically tells that the contexts are bounded; such assumptions are standard in the bandit literature to obtain results on regret bound that are independent of the scaling of the contexts (or parameter). Assumption 2.1(b) says that all the arm-contexts are generated from *mean zero* sub-Gaussian distribution with parameter ϑ , for all time point t. This is indeed a very mild assumption on the context distribution and a broad class of distributions enjoys such property. For example, truncated multivariate normal distribution with covariance matrix \mathbb{I}_d , where the truncation is over the set $\{u \in \mathbb{R}^d : ||u||_{\infty} \leq 1\}$ is a valid distribution for the contexts. Assumption 2.1(c) talks about anti-concentration condition that plays a critical role in controlling the estimation accuracy of β^* . Intuitively, this condition prohibits the context features to fall along a singular direction. In particular, this condition ensures that the directions of the arms are well spread across every direction and thus implicitly promotes exploration in the feature space. Moreover, this anticoncentration is very mild as it holds if the density of $x_i(t)^{\top}u$ is bounded above. Assumption 2.1(d) imposes an upper bound on the maximum sparse eigenvalue of Σ_i which is a common assumption in high-dimensional literature (Zhang and Huang, 2008; Zhang, 2010).

Next, we come to the assumptions on the true parameter β^*

Assumption 2.2 (Assumptions on the true parameter). We assume the followings:

- (a) Sparsity and Soft-sparsity: There exist positive constants $s^* \in \mathbb{N}$ and $b_{\max} \in \mathbb{R}^+$ such that $\|\beta^*\|_0 = s^*$ and $\|\beta^*\|_1 \leq b_{\max}$.
- (b) Margin condition: There exists positive constants Δ_* , A and $\omega \in [0, \infty]$, such that for $h \in \left[A\sqrt{\log(d)/T}, \Delta_*\right]$ and for all $t \in [T]$,

$$\mathbb{P}\left(x_{a_t^*}(t)^\top \beta^* \le \max_{i \ne a_t^*} x_i(t)^\top \beta^* + h\right) \le \left(\frac{h}{\Delta^*}\right)^{\omega}$$

The first part of the assumption requires boundedness of the true parameter β^* to make the final regret bound scale free. Such an assumption is also standard in bandit literature (Bastani and Bayati, 2020; Abbasi-Yadkori et al., 2011).

The second part of the assumption imposes a margin condition on the arm distributions. Essentially, this assumption controls the probability of the optimal arm falling into *h*-neighborhood of the sub-optimal arms. As ω increases, the margin condition becomes stronger as the sub-optimal arms are less likely to fall close to the optimal arms. As a result, it becomes easier for any bandit policy to distinguish the optimal arm. As an illustration, consider the two extreme cases $\omega = \infty$ and $\omega = 0$. The $\omega = \infty$ case tells that there is a deterministic gap between rewards corresponding to the optimal arm and sub-optimal arms. This is the same as the "gap assumption" in Abbasi-Yadkori et al. (2011). Thus, quite evidently it is easy for any bandit policy to recognize the optimal arm. This phenomenon is reflected in the regret bound of Theorem 5 in Abbasi-Yadkori et al. (2011), where the regret depends on the time horizon T only though poly-logarithmic terms. In contrast, $\omega = 0$ corresponds to the case when there is no apriori information about the separation between the arms, and as a consequence, we pay the price in regret bound by a \sqrt{T} term (Hao et al., 2021; Agrawal and Goyal, 2013; Chu et al., 2011).

The margin condition with $\omega = 1$ has been assumed in Goldenshluger and Zeevi (2013); Bastani and Bayati (2020); Wang et al. (2018) and will be satisfied when the density of $x_i(t)^{\top}\beta^*$ is uniformly bounded for all $i \in [K]$. Li et al. (2021) also discusses an example where the margin condition holds for different values of ω . The final assumption is on the noise variables:

Assumption 2.3 (Assumption on Noise). We assume that the random variables $\{\epsilon(t)\}_{t\in[T]}$ are independent and also independent of the other processes and each one is σ -Sub-Gaussian, i.e., $\mathbb{E}[e^{a\epsilon(t)}] \leq e^{\sigma^2 a^2/2}$ for all $t \in [T]$ and $a \in \mathbb{R}$.

Various families of distribution satisfy such a requirement, including normal distribution and bounded distributions, which are commonly chosen noise distributions. Note that such a requirement automatically implies that for every $t \in [T]$, $\mathbb{E}[\epsilon(t)] = 0$ and $\operatorname{Var}[\epsilon(t)] \leq \sigma^2$.

2.2 Thompson Sampling and Prior

We discuss the basics of Thompson sampling and introduce the specific structure of the prior that we use and analyze. Typically, we place a prior Π on the unknown parameter (β in our case) along with a *specified likelihood model* on the data, and do the following: while taking action, we draw a sample from the posterior distribution of the parameter given the data and use that as the proxy for the unknown parameter value, hence in our case at time t, we draw a sample $\hat{\beta}_t \sim \Pi(\beta | \mathcal{H}_{t-1})$ and choose $a_t = \arg \max_i x_i(t)^{\top} \hat{\beta}_t$ as the action. While simple enough to describe, Thompson sampling has been difficult to analyze theoretically, particularly because of the complex dependence between the observations due to the bandit structure. The choice of prior plays a crucial role, as we shall see, in providing the correct exploration-exploitation trade-off. In the high-dimensional sparse case that we are dealing with, this choice is specifically important since we do not wish to have a linear dependence on the dimension d in our regret bound - which would be incurred if we use the normal prior-likelihood setup of Agrawal and Goyal (2013), which analyzes Thompson sampling in contextual bandits.

While there is a rich literature on Bayesian priors for high dimensional regression, including horseshoe priors and slab-and-spike priors among others, we shall be using the complexity prior introduced in Castillo et al. (2015). Specifically, we consider a prior Π on β that first selects a dimension s from a prior π_d on the set [d], next a random subset $S \subset [d]$ of size |S| = s and finally, given S, a set of nonzero values $\beta_S := \{\beta_i : i \in S\}$ from a prior density g_S on \mathbb{R}^S . Formally, the prior on (S, β) can be written as

$$(S,\beta) \mapsto \pi_d(|S|) \frac{1}{\binom{d}{|S|}} g_S(\beta_S) \delta_0(\beta_{S^c}), \tag{2}$$

where the term $\delta_0(\beta_{S^c})$ refers to coordinates β_{S^c} being set to 0. Moreover, we choose g_S as a product of Laplace densities on \mathbb{R} with parameter λ/σ , i.e., $\beta_i \mapsto (2\sigma)^{-1}\lambda \exp(-\lambda|\beta_i|/\sigma)$ for all $i \in S$. Note that, here we assume that the noise level σ is known. In practice, one can add another level of hierarchy by setting a prior on σ but in this paper we do not pursue that direction.

The prior π_d plays the role of expressing the sparsity of the parameter. This is in contrast to other priors like product of independent Laplace densities over the coordinates (typically known as Bayesian Lasso), where the Laplace parameter plays the role of shrinking the coefficients towards 0. However, in our case, the scale parameter λ of the Laplace does not have this role and we assume that during the *t*th round we use $\lambda \in [(5/3)\overline{\lambda}_t, 2\overline{\lambda}_t]$, where $\overline{\lambda}_t \asymp \sqrt{t \log d}$, which is the usual order of the regularization parameter used in the LASSO.

The choice of the prior π_d is very critical; it should down weight big models but at the same time give enough mass to the true model. Following Castillo et al. (2015), we assume that there are constants $A_1, A_2, A_3, A_4 > 0$ such that $\forall s \in [d]$

$$A_1 d^{-A_3} \pi_d(s-1) \le \pi_d(s) \le A_2 d^{-A_4} \pi_d(s-1).$$
(3)

Complexity priors of the form $\pi_d(s) \propto c^{-s} d^{-as}$ for constants a, c satisfy the above requirement. Moreover, slab and spike priors of the form $(1-r)\delta_0 + r \text{Lap}(\lambda/\sigma)$ independently over the coordinates satisfy the requirement with hyperprior on r being $\text{Beta}(1, d^u)$.

Finally, we specify the data likelihood that is crucial for the TS algorithm. At each time point $t \in [T]$, given the observations coming from model (1), we model the $\{\epsilon(\tau)\}_{\tau \leq t}$ as i.i.d. $N(0, \sigma^2)$. We emphasize that this Gaussian assumption is *only required for likelihood modeling* and our main results hold under any true error distribution satisfying Assumption 2.3.

3 Main Results

3.1 Posterior contraction

Now, we present an informal version of the main posterior contraction result for the estimation of β . A more detailed version of the result with exact rates, along with the measure theoretic details, is in Appendix C.

Theorem 1 (Informal). Write $\mathbf{r}_t = (r(1), \ldots, r(t))^{\top}$, and let the Assumption 2.1–2.3 hold with $C = \Theta(\phi_u \vartheta^2 \xi K \log K)$, and $K \ge 2, d \ge T$. With $\lambda \asymp x_{\max}(t \log d)^{1/2}$ and $\varepsilon_{t,d,s} = s^* \{(\log d + \log t)/t\}\}^{1/2}$, the following holds as $t \to \infty$:

$$\mathbb{E}_{\mathbf{r}_{t}} \Pi \left(\left\| \beta - \beta^{*} \right\|_{1} \gtrsim \sigma \varepsilon_{t,d,s} \mid \mathbf{r}_{t}, X_{t} \right) \stackrel{a.s.}{\rightarrow} 0.$$

The above result is similar to Theorem 3 in Castillo et al. (2015) under classical linear regression setup with i.i.d. observations and Gaussian noise. However, we generalize their result under bandit setup and sub-Gaussian noise by carefully controlling the correlation between noise and observed contexts, which is crucial for our regret analysis.

3.2 Algorithm and regret bound

In this section we introduce the Thomson sampling algorithm for high-dimensional contextual bandit, a pseudo-code for which is provided below in Algorithm 1. Similar to the Thompson sampling algorithm in Agrawal and Goyal (2013), in the *t*th round Algorithm 1 sets the a specific prior on β and updates it sequentially based on the observed rewards and contexts. In particular, it chooses the prior described in (2) with an appropriate choice of round-specific prior scaling λ_t and updates the posterior using the observed rewards and contexts until (t-1)th round. Then a sample is generated from the posterior and an arm a_t is chosen greedily based on the generated sample.

Now, we show that the Thompson sampling algorithm achieves desirable regret upper bound.

Theorem 2. Let the Assumption 2.1–2.3 hold with $C = \Theta(\phi_u \vartheta^2 \xi K \log K)$, and $K \ge 2, d \ge T$. Define the quantity $\kappa(\xi, \vartheta, K) := \min\{(4c_3K\xi\vartheta^2)^{-1}, 1/2\}$ where c_3 is a universal positive constant. Also, set the prior scaling λ_t as follows:

$$(5/3)\overline{\lambda}_t \le \lambda_t \le 2\overline{\lambda}_t, \quad \overline{\lambda}_t = x_{\max}\sqrt{2t(\log d + \log t)}.$$

Then there exists a universal constant $C_0 > 0$ such that we have the following regret bound for Algorithm 1:

$$\mathbb{E}\{R(T)\} \lesssim I_b + I_\omega,$$

where,

$$I_{b} = \left\{ \frac{b_{\max} x_{\max} \phi_{u} \vartheta^{2} \xi(K \log K)}{\min \left\{ \kappa^{2}(\xi, \vartheta, K), \log K \right\}} \right\} s^{*} \log(Kd),$$

$$I_{\omega} = \begin{cases} \Phi^{1+\omega} \left(\frac{s^{*1+\omega} (\log d)^{\frac{1+\omega}{2}} T^{\frac{1-\omega}{2}}}{\Delta_{\omega}^{\omega}} \right), & \text{for } \omega \in [0, 1), \\ \Phi^{2} \left(\frac{s^{*2} [\log d + \log T] \log T}{\Delta_{\omega}} \right), & \text{for } \omega = 1, \\ \frac{\Phi^{2}}{(\omega-1)} \left(\frac{s^{*2} [\log d + \log T]}{\Delta_{\omega}} \right), & \text{for } \omega \in (1, \infty) \\ \Phi^{2} \left(\frac{s^{*2} [\log d + \log T]}{\Delta_{\omega}} \right), & \text{for } \omega = \infty, \end{cases}$$

$$P_{\max}^{2} \xi K \left(2 + 40A_{4}^{-1} + C_{0}K\xi x_{\max}^{2}A_{4}^{-1} \right).$$

and $\Phi = \sigma x_{\max}^2 \xi K \left(2 + 40A_4^{-1} + C_0 K \xi x_{\max}^2 A_4^{-1} \right)$.

Discussion on the above result: The regret bound provided by Theorem 2 shows that the regret of the algorithm grows poly-logarithmically in d, i.e., $\mathbb{E}\{R(T)\} = O((\log d)^{\frac{1+\omega}{2}})$, when $\omega \in [0, 1)$; logarithmically in d, i.e., $O(\log d)$ when $\omega \in [1, \infty]$. Meanwhile, the expected cumulative regret depends polynomially in T, i.e., $\mathbb{E}\{R(T)\} = O(T^{\frac{1-\omega}{2}})$ when $\omega \in [0, 1)$; ploy-logarithmically in T; i.e., $\mathbb{E}\{R(T)\} = O((\log T)^2)$, when $\omega = 1$. In $\omega \in (1, \infty]$ regime, the expected cumulative regret depends poly-logarithmically in both the time horizon T and ambient dimension d. As $T \ll d$, the expected regret ultimately scales as $O(\log d)$. Comparing our upper bound result with minimax regret lower bound established in Theorem 1 of Li et al. (2021), it follows that our algorithm enjoys optimal dependence on both ambient dimension d and time-horizon T when $\omega \in [0, 1)$. In $\omega = 1$ region, the regret upper bound in the above theorem is optimal up to a $O(\log T)$ term. To the best of our knowledge, there does not exist any result on minimax lower bound in the regime $\omega > 1$ in the high-dimensional linear contextual bandit literature. It is worth mentioning that this is an upper bound on the expected (frequentist) regret, as compared to Bayesian regret which is often considered for Thompson sampling based algorithms..

Intuitively, the initial term I_b in regret upper bound in Theorem 2 describes the regret caused by the "burn-in" period of exploring the space of contexts and it does not contribute to the asymptotic regret growth. Note that we consider running Thompson sampling from the very beginning, without an explicit random exploration phase, in contrast to most of the existing algorithms; the distinction between the burn-in phase and the subsequent phase is only a construct of our theoretical analysis. Furthermore, the constant Δ_* plays the role of gap parameter which commonly appears in a problem-dependent regret bound (Abbasi-Yadkori et al., 2011). Note that, for $\omega = 0$, we get a problem-independent regret bound of the order $O(s^*\sqrt{T} \log d)$. The appearance of the \sqrt{T} , term

Algorithm 1: Thompson Sampling Algorithm

Set $\mathcal{H}_0 = \emptyset$; for $t = 1, \dots, T$ do if $t \leq 1$ then | Choose action a_t uniformly over [K]; end else | Set $\overline{\lambda}_t = x_{\max}\sqrt{2t(\log d + \log t)}$ and choose $\lambda_t \in (5\overline{\lambda}_t/3, 2\overline{\lambda}_t)$; Generate sample $\tilde{\beta}_t \sim \Pi(\cdot \mid \mathcal{H}_{t-1})$ with prior Π in (2)-(3), $\lambda = \lambda_t$ and Gaussian likelihood; Play arm: $a_t = \arg \max_{i \in [K]} x_i(t)^{\top} \tilde{\beta}_t$; end Observe reward $r_{a_t}(t)$; Update $\mathcal{H}_t \leftarrow \mathcal{H}_{t-1} \cup \{(a_t, r_{a_t}(t), x_{a_t}(t))\}$. end

is not surprising, as the condition $\omega = 0$ poses no prior knowledge on the arm-separability, Thus, in the worst case, the context vectors may fall into each other, making the bandit environment harder to learn. In contrast, as ω increases the optimal arm becomes more distinguishable than the sub-optimal arms and the bandit environment becomes easier to learn. As a result, the effect of the time horizon becomes less and less severe as ω increases. In particular, when $\omega \in [1, \infty]$, the time horizon does not affect the asymptotic growth of the regret bound. Finally, as we mainly focus on the case when the number of arms is very small, the quantity Φ roughly has an inflating effect of O(1) on the regret bound.

Sketch of the proof of Theorem 2 While a self-contained and detailed proof of the above result is given in the Appendix, here we go through the main steps and ideas of the proof. The proof is broadly divided into 3 parts for clarity:

- (i) In Section B.1 we will first show that the estimated covariance matrix $\widehat{\Sigma}_t := X_t^{\top} X_t / t$ enjoys SRC condition with high probability for sufficiently large t. In our analysis, we carefully decouple this complex dependent structure and exploit the special temporal dependence structure of the bandit environment to establish SRC property of $\widehat{\Sigma}_t$.
- (ii) Next, in Section B.2 we will establish a compatibility condition for the matrix $\widehat{\Sigma}_t$. We use a Transfer Lemma (Lemma B.6) which essentially translates the uniform lower bound on $\phi_{\min}(Cs^*, \widehat{\Sigma}_t)$ to a certain compatibility number.
- (iii) Finally, in Section B.3, under the compatibility condition we use the posterior contraction result in Theorem 1 to give bound on the per round regret $\Delta_{a_t}(t)$.

3.3 Comparison with existing literature

The stochastic linear bandit problem was first introduced by Auer (2002), and later was subsequently studied by Dani et al. (2008); Chu et al. (2011) and many others. Later, Abbasi-Yadkori

et al. (2011) and Abbasi-Yadkori et al. (2012) proposed OFUL algorithm for both low-dimensional and high-dimensional settings. Although there are some similarities in our model parametrization with the setting considered in Abbasi-Yadkori et al. (2011) and Dani et al. (2008), there are some significant differences that need attention. To mention a few, the set of contexts considered in Abbasi-Yadkori et al. (2011); Dani et al. (2008) consists of infinitely many feature vectors, whereas in our setting the set of contexts consists of finitely many feature vectors coming from an underlying distribution. Moreover, in Dani et al. (2008), the set of contexts does not change over time, therefore the optimal arm remains the same in every round. In high dimensional bandit, a similar setting is also considered in Hao et al. (2020, 2021). In contrast, in our setting, due to the randomness of the observed contexts, the optimal arm does not necessarily remain the same in every round.

Now we shift focus to the regret bound analysis. In low-dimensional setting Rusmevichientong and Tsitsiklis (2010) proved a lower bound of $\Omega(d\sqrt{T})$ for both cumulative regret and Bayesian regret in linear bandit setting, where the set of contexts is a compact set of infinitely many feature vectors, e.g., the unit sphere $\mathbb{S}^{d-1} \subseteq \mathbb{R}^d$. Later, Chu et al. (2011) proposed LinUCB algorithm, which has near-optimal regret upper bound $O(\sqrt{Td\log(T\log(T)/\delta)})$ with probability $1-\delta$, which corresponds to $\omega = 0$ case. Abbasi-Yadkori et al. (2011) proved that the expected regret of OFUL algorithm scales as $O(d\sqrt{T})$ in both low-dimensional and high-dimensional setting. In all of these works, the regret bound analysis is based on the worst-case scenario, which leads to polynomial dependence on d. In high-dimension, such a polynomial dependence of d may lead to very suboptimal performance of the aforementioned algorithms. Later, Bastani and Bayati (2020) proposed LASSO-bandit algorithm, and Wang et al. (2018) proposed MCP bandit algorithm which enjoys improved regret upper bound scaling as $O(\log d)$ under the margin condition $\omega = 1$, but require forced sampling which could be costly in some practical settings such as medical and marketing applications. In comparison, our method does not need any forced sampling and does not require the knowledge of the gap parameter ω . On the other hand, our theoretical analysis also covers the regime $\omega \in [0, \infty]$, whereas the results of Bastani and Bayati (2020) and Wang et al. (2018) are only valid for $\omega = 1$ case, for which our algorithm enjoys the same $O(\log d)$ dependence in regret bound as LASSO-bandit and MCP-bandit.

Remark 1. It is worthwhile to point out that the setup in Bastani and Bayati (2020) and Wang et al. (2018) consider different $\beta_i^* \in \mathbb{R}^d$ for different arms $i \in [K]$ and a single context $x(t) \in \mathbb{R}^d$ every round, which is in sharp contrast to our setting. Their formulation can be mathematically reparametrized into our formulation. In particular, for the *i*th arm we construct the new context vectors $\tilde{x}_i(t) = (\mathbf{0}, \mathbf{0}, \dots, x(t)^\top, \mathbf{0}, \dots, \mathbf{0})^\top$ where the *i*th block is x(t), thus $x_i(t) \in \mathbb{R}^{Kd}$. The common parameter is $\beta^* = (\beta_1^{*\top}, \dots, \beta_K^{*\top})^\top$. However, in this case, the new contexts are highly degenerate and violate Assumption 2.1(c) and restrict us from directly applying our result in this case.

4 Computation

In this section, we discuss the computational challenges and how these are overcome by using Variational Bayes (VB). While priors as (2) have been shown to perform well, both empirically and in theory, the discrete model selection component of the prior makes it challenging to allow

computation and inference on the posterior. For $\beta \in \mathbb{R}^d$, inference using the slab and spike prior requires a combinatorial search over 2^d possible models, which in the case of high dimension is computationally infeasible. Fast algorithms are known only in the special diagonal design case and traditional Markov Chain Monte Carlo methods have very slow mixing in such high dimensional cases. Thus, following Ray and Szabó (2021) we use Variational Bayes to make computations faster. Specifically, in the sampling step of Algorithm 1, we consider the VB approximation of the posterior $\Pi(\cdot | \mathcal{H}_{t-1})$ arising from slab and spike prior with $Lap(\lambda/\sigma)$ slab in the mean-field family

$$\left\{\bigotimes_{j=1}^{d} [\gamma_j \mathsf{N}(\mu_j, \sigma_j^2) + (1 - \gamma_j)\delta_0] : (\mu_j, \sigma_j, \gamma_j) \in \mathcal{R}\right\},\$$

where $\mathcal{R} = \mathbb{R} \times \mathbb{R}^+ \times [0, 1]$. We use the sparsevb package (Clara et al., 2021) in R to use the Coordinate Ascent Variational Inference (CAVI) algorithm proposed in Ray and Szabó (2021) to obtain the VB posterior. This makes the Thompson sampling algorithm much faster as one can efficiently obtain samples from the VB posterior due to its structure. The details of the algorithm for the Variational Bayes Thompson Sampling (VBTS) are in the appendix (see Section E).

5 Numerical Experiment

In both simulations and real data experiments, we present results corresponding to $\lambda_t = 1$ for all $t \in [T]$. Recall that Theorem 2 suggests that in tth round $\lambda_t \approx \sqrt{t \log d}$ is a reasonable choice for the exact Thompson sampling algorithm. However, in practice, we noticed that such choices of λ_t lead to numerical instability. Some recent findings in Ray and Szabó (2021) suggest that λ_t in the order of $O(\sqrt{t \log d}/s^*)$ should be an appropriate choice, which is smaller than the predicted order of λ_t in our main theorem. Motivated by this, we also present the simulation results for synthetic data experiments with $\lambda_t = \lambda_* \sqrt{t}$ for $\lambda_* \in \{0.2, 0.3, 0.4, 0.5\}$ in the Appendix A.2. We found the performance of VBTS to be robust with respect to the choice of the tuning parameter λ_t .

5.1 Synthetic data

In this section, we illustrate the performance of the VBTS algorithm on a simulated data set. As a benchmark, we consider ESTC (Hao et al., 2020), LinUCB (Abbasi-Yadkori et al., 2011), DRLasso (Kim and Paik, 2019), Lasso-L1 confidence ball algorithm (Li et al., 2021), LinearTS (Agrawal and Goyal, 2013) and TS algorithm based on Bayesian Lasso (Park and Casella, 2008) (BLasso TS) to compare the performance of VBTS (Algorithm 2).

Equicorrelated (EC) structure

We set the number of arms K = 10 and we generate the context vectors $\{x_i(t)\}_{i=1}^K$ from multivariate *d*-dimensional Gaussian distribution $N_d(0, \Sigma)$, where $\Sigma_{ij} = \rho^{|i-j| \wedge 1}$ and $\rho = 0.3$. We consider d = 1000 and the sparsity $s^* = 5$. We choose the set of active indices S^* uniformly over all the subsets of [d] of size s^* . Next, for each choice of d, we consider two types generating scheme for β :




Figure 1. Cumulative regret of competing algorithms.

• Setup 1: $\{U_i\}_{i \in S^*} \stackrel{i.i.d.}{\sim} \text{Uniform}(0.3, 1) \text{ and set } \beta_j = U_j(\sum_{\ell \in S^*} U_\ell^2)^{-1/2} \mathbb{1}(j \in S^*).$

• Setup 2:
$$\{Z_i\}_{i\in S^*} \stackrel{i.i.d.}{\sim} N(0,1)$$
 and set $\beta_j = Z_j(\sum_{\ell\in S^*} Z_\ell^2)^{-1/2} \mathbb{1}(j\in S^*)$.

We run 40 independent simulations and plot the mean cumulative regret with 95% confidence band in Figure 1a-1b. In all the setups, we see that VBTS outperforms its competitors by a wide margin. VBTS also enjoys superior empirical performance under the autoregressive (AR) model (see Figure 1c-1d) with auto-correlation coefficient 0.3 and the details of the simulations can be found in Appendix A.1. Table 1 shows the mean execution time (across Setup 1 and 2) of all TS algorithms. Among the class of TS algorithms, VBTS outperforms its other competing algorithms.

5.2 Real data - gravier Breast Carcinoma Data

We consider breast cancer data gravier (microarray package in R) for 168 patients to predict metastasis of breast carcinoma based on 2905 gene expressions (bacterial artificial chromosome or BAC array). The goal of the learner is to identify the positive cases.

Similar to Kuzborskij et al. (2019); Chen et al. (2021), in our experimental setup, we convert the breast cancer classification problem into 2-armed contextual bandit problem. More details about the data and reward generation process are provided in Appendix A.3. We perform 10 independent Monte Carlo simulations and plot the expected regret of VBTS in Figure 2 along

Туре	Algorithm	Mean time of execution (seconds)		
		Equicorrelated	Auto-regressive	
TS	LinTS	1344.39	1346.46	
	BLasso TS	1511.68	1455.53	
	VBTS	29.33	27.65	

Table 1: Time comparison among the competing algorithms.

with its competitors. In this experiment, we omit LinUCB and LinTS algorithms as they were performing far worse compared to the existing ones in Figure 2. The figure shows that VBTS and Lasso-L1 confidence ball algorithms are by far the clear winners in terms of cumulative regret. However, upon closer look, we see that VBTS is slightly better than Lasso-L1 confidence in terms of cumulative regret. In terms of accuracy, Lasso-L1 and VBTS are in the same ball park as seen in Table 2.




Figure 2: Cumulative regret plot for breast cancer data set.

6 Conclusion

In this paper, we consider the stochastic linear contextual bandit problem with high-dimensional sparse features and a fixed number of arms. We propose a Thompson sampling algorithm for this problem by placing a suitable *sparsity-inducing* prior on the unknown parameter to induce sparsity. We also develop a crucial posterior contraction result for *non-i.i.d.* data that allows us to obtain an almost *dimension independent* regret bound for our proposed algorithm. We explicitly

Algorithm	DRLasso	Lasso-L1	ESTC	VBTS
Accuracy(%)	65.63	81.20	73.32	81.88

Table 2: Classification accuracy of competing algorithms.

point out the dependences on d and T for different arm-separation regimes parameterized by ω , which is also minimax optimal for $\omega \in [0, 1)$. Moreover, the choice of prior allows us to devise a Variational Bayes algorithm that enjoys computational expediency over traditional MCMC. We demonstrate the superior performance of our algorithm through extensive simulation studies. We finally perform an experiment on a gravier dataset by converting the classification problem into a 2-armed contextual bandit problem, for which our method performs better compared to other existing algorithms.

Now we point the readers toward some of the natural research directions that we plan to cover in our future works. In terms of regret analysis, similar to most of the recent works in high dimensional contextual bandits, relies on upper bounding the regret through estimation of the parameter, i.e., we rely on the estimation of β^* to be able to provide meaningful regret bound. However, this should not be required - as an example, consider the case where the first coordinate of $x_i(t)$ is 0 for all $i \in [K], t \in [T]$. Then the first coordinate of β^* is not estimable, however, this does not pose any problem to designing a sensible policy since this coordinate does not appear in the regret. Unfortunately, Assumption 2.1(c) is not satisfied for such degeneracy in the contexts and as a result, it would require a modified analysis of the regret bound. Secondly, we underscore the fact that in our setup we adopt the Variational Bayes framework only to sidestep the computational hurdles of MCMC arising from a myriad of challenges such as slow mixing times of the chains, lack of easy implementation, etc. However, in high-dimensional regression setup Yang et al. (2016) has proposed Metropolis-Hastings algorithms based on truncated sparsity priors that do not meet the above roadblocks. It could be very well possible that some other prior structure will allow us to design more efficient MCMC algorithms with faster mixing times in the high-dimensional SLCB setup along with theoretical guarantees. Finally, we also plan to analyze Thompson sampling for high-dimensional generalized contextual bandit problems.

Author Contribution: All authors conceived and carried out the research project jointly. S.C. and S.R. jointly wrote the paper and code for numerical experiments. A.T. helped edit the paper.

References

- Abbasi-Yadkori, Y., Pál, D., and Szepesvári, C. (2011). Improved algorithms for linear stochastic bandits. *Advances in neural information processing systems*, 24.
- Abbasi-Yadkori, Y., Pal, D., and Szepesvari, C. (2012). Online-to-confidence-set conversions and application to sparse stochastic bandits. In *Artificial Intelligence and Statistics*, pages 1–9. PMLR.

- Abe, N., Biermann, A. W., and Long, P. M. (2003). Reinforcement learning with immediate rewards and linear hypotheses. *Algorithmica*, 37(4):263–293.
- Agrawal, S. and Goyal, N. (2013). Thompson sampling for contextual bandits with linear payoffs. In *International Conference on Machine Learning*, pages 127–135. PMLR.
- Auer, P. (2002). Using confidence bounds for exploitation-exploration trade-offs. *Journal of Machine Learning Research*, 3(Nov):397–422.
- Bastani, H. and Bayati, M. (2020). Online decision making with high-dimensional covariates. *Operations Research*, 68(1):276–294.
- Bickel, P. J., Ritov, Y., and Tsybakov, A. B. (2009). Simultaneous analysis of lasso and dantzig selector. *The Annals of statistics*, 37(4):1705–1732.
- Castillo, I., Schmidt-Hieber, J., and Van der Vaart, A. (2015). Bayesian linear regression with sparse priors. *The Annals of Statistics*, 43(5):1986–2018.
- Chapelle, O. and Li, L. (2011). An empirical evaluation of thompson sampling. *Advances in neural information processing systems*, 24.
- Chen, C., Luo, L., Zhang, W., Yu, Y., and Lian, Y. (2021). Efficient and robust high-dimensional linear contextual bandits. In *Proceedings of the Twenty-Ninth International Conference on International Joint Conferences on Artificial Intelligence*, pages 4259–4265.
- Chen, Y., Wang, Y., Fang, E. X., Wang, Z., and Li, R. (2022). Nearly dimension-independent sparse linear bandit over small action spaces via best subset selection. *Journal of the American Statistical Association*, (just-accepted):1–31.
- Chu, W., Li, L., Reyzin, L., and Schapire, R. (2011). Contextual bandits with linear payoff functions. In *Proceedings of the Fourteenth International Conference on Artificial Intelligence and Statistics*, pages 208–214. JMLR Workshop and Conference Proceedings.
- Clara, G., Szabo, B., and Ray, K. (2021). sparsevb: Spike-and-slab variational bayes for linear and logistic regression: R package version 0.1. 0.
- Dani, V., Hayes, T. P., and Kakade, S. M. (2008). Stochastic linear optimization under bandit feedback.
- Gilton, D. and Willett, R. (2017). Sparse linear contextual bandits via relevance vector machines. In 2017 International Conference on Sampling Theory and Applications (SampTA), pages 518–522. IEEE.
- Goldenshluger, A. and Zeevi, A. (2013). A linear response bandit problem. *Stochastic Systems*, 3(1):230–261.
- Gravier, E., Pierron, G., Vincent-Salomon, A., Gruel, N., Raynal, V., Savignoni, A., De Rycke, Y., Pierga, J.-Y., Lucchesi, C., Reyal, F., et al. (2010). A prognostic dna signature for t1t2 node-negative breast cancer patients. *Genes, chromosomes and cancer*, 49(12):1125–1134.

- Hao, B., Lattimore, T., and Deng, W. (2021). Information directed sampling for sparse linear bandits. arXiv preprint arXiv:2105.14267.
- Hao, B., Lattimore, T., and Wang, M. (2020). High-dimensional sparse linear bandits. In Larochelle, H., Ranzato, M., Hadsell, R., Balcan, M. F., and Lin, H., editors, *Advances in Neural Information Processing Systems*, volume 33, pages 10753–10763. Curran Associates, Inc.
- Kaufmann, E., Korda, N., and Munos, R. (2012). Thompson sampling: An asymptotically optimal finite-time analysis. In *International conference on algorithmic learning theory*, pages 199–213. Springer.
- Kim, G.-S. and Paik, M. C. (2019). Doubly-robust lasso bandit. Advances in Neural Information Processing Systems, 32:5877–5887.
- Kuzborskij, I., Cella, L., and Cesa-Bianchi, N. (2019). Efficient linear bandits through matrix sketching. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 177–185. PMLR.
- Li, K., Yang, Y., and Narisetty, N. N. (2021). Regret lower bound and optimal algorithm for high-dimensional contextual linear bandit. *Electronic Journal of Statistics*, 15(2):5652–5695.
- Oliveira, R. I. (2013). The lower tail of random quadratic forms, with applications to ordinary least squares and restricted eigenvalue properties. *arXiv preprint arXiv:1312.2903*.
- Park, T. and Casella, G. (2008). The bayesian lasso. *Journal of the American Statistical Association*, 103(482):681–686.
- Ray, K. and Szabó, B. (2021). Variational bayes for high-dimensional linear regression with sparse priors. *Journal of the American Statistical Association*, pages 1–12.
- Rusmevichientong, P. and Tsitsiklis, J. N. (2010). Linearly parameterized bandits. *Mathematics of Operations Research*, 35(2):395–411.
- Vershynin, R. (2018). *High-dimensional probability: An introduction with applications in data science*, volume 47. Cambridge university press.
- Wainwright, M. J. (2019). *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge University Press.
- Wang, X., Wei, M., and Yao, T. (2018). Minimax concave penalized multi-armed bandit model with high-dimensional covariates. In *International Conference on Machine Learning*, pages 5200–5208. PMLR.
- Yang, Y., Wainwright, M. J., and Jordan, M. I. (2016). On the computational complexity of highdimensional bayesian variable selection. *The Annals of Statistics*, 44(6):2497–2532.
- Zhang, C.-H. (2010). Nearly unbiased variable selection under minimax concave penalty. *The Annals of statistics*, 38(2):894–942.
- Zhang, C.-H. and Huang, J. (2008). The sparsity and bias of the lasso selection in high-dimensional linear regression. *The Annals of Statistics*, 36(4):1567–1594.

Appendix A Details of Simulations

A.1 Simulation details for AR(1) structure

We set the number of arms K = 10 and we generate the context vectors $\{x_i(t)\}_{i=1}^K$ from multivariate *d*-dimensional Gaussian distribution $N_d(0, \Sigma)$, where $\Sigma_{ij} = \phi^{|i-j|}$ and $\phi = 0.3$. We consider d = 1000 and the sparsity $s^* = 5$. We choose the set of active indices S^* uniformly over all the subsets of [d] of size s^* . Next, for each choice of d, we consider two types generating scheme for β :

• Setup 1: $\{U_i\}_{i \in \mathcal{S}^*} \stackrel{i.i.d.}{\sim}$ Uniform(0.3, 1) and set β as the following:

$$\beta_j = \begin{cases} \frac{U_j}{\sqrt{\sum_{\ell \in \mathcal{S}^*} U_\ell^2}}, & \text{if } j \in \mathcal{S}^*, \\ 0, & \text{otherwise.} \end{cases}$$

• Setup 2: $\{Z_i\}_{i\in\mathcal{S}^*} \stackrel{i.i.d.}{\sim} \operatorname{Normal}(0,1)$ and set β as the following:

$$\beta_j = \begin{cases} \frac{Z_j}{\sqrt{\sum_{\ell \in \mathcal{S}^*} Z_\ell^2}}, & \text{if } j \in \mathcal{S}^*, \\ 0, & \text{otherwise.} \end{cases}$$

We run 40 independent simulations and plot the mean cumulative regret with 95% confidence band in Figure 1. In all the setups, we see that VBTS outperforms its competitors by a wide margin. Similar to the previous simulation example, in this case also Table 1 shows that VBTS is far better in terms of mean execution time than its competitors in the class of TS algorithms.

A.2 Siumlation for different choices of λ

As discussed in the first paragraph of Section 5, for each of these simulation settings, we tried a few choices for the tuning parameter λ_t . In addition to the default choice of $\lambda_t = 1$ (for all time points t), we also explored the performance of the algorithm under growing λ , as required by our theoretical results. In particular, we tried $\lambda_t = \lambda_* \sqrt{t}$ for $\lambda_* \in \{0.2, 0.3, 0.4, 0.5\}$. For comparison, we only kept the faster optimism based methods DRLasso, Lasso-L1 and ESTC. We found the results to be roughly robust to the choice of this tuning parameter. The results are summarized in Figure 3 and Figure 4 below. However, we found that larger values of λ_* lead to numerical issues, we conjecture that this is an artifact of the variational Bayes approximation, rather than the prior itself. For our simulation settings, the choice $\sqrt{t \log d}/s^* \approx 0.5\sqrt{t}$ and hence by the findings in Ray and Szabó (2021), values of λ_* higher than this may yield inaccurate Variational Bayes estimation.

A.3 Details of real data experiment

We consider breast cancer data gravier (microarray package in R) for 168 patients to predict metastasis of breast carcinoma based on 2905 gene expressions (bacterial artificial chromosome or BAC array). (Gravier et al., 2010) considered small, invasive ductal carcinomas without axillary




Figure 3: Regret bound for equi-correlated design for different tuning parameter choices




Figure 4: Regret bound for auto-regressive design for different tuning parameter choices

lymph node involvement (T1T2N0) to predict metastasis of small node-negative breast carcinoma. Using comparative genomic hybridization arrays, they examined 168 patients over a five-year period. The 111 patients with no event after diagnosis were labeled good (class 0), and the 57 patients with early metastasis were labeled poor (class 1). The 2905 gene expression levels were normalized with a \log_2 transformation.

Similar to Kuzborskij et al. (2019); Chen et al. (2021), in our experimental setup we convert the breast cancer classification problem into 2-armed contextual bandit problem as follows: Given the gravier data set with 2 classes, we first set Class 1 as the target class. In each round, the environment randomly draws one sample from each class and composes a set of contexts of 2 samples. The learner chooses one sample and observes the reward following a logit model. In particular, we model the reward as

$$r(t) := \log \left\{ \frac{\mathbb{P}(\text{Selected class} = 1)}{\mathbb{P}(\text{Selected class} = 0)} \right\} = x_{a_t}^\top \beta^* + \epsilon(t)$$

where $a_t \in \{1, 2\}$ is the selected arm at round t. Thus, small cumulative regret insinuates that the learner is able to differentiate the positive patients eventually. Such concepts can be used for constructing online classifiers to differentiate carcinoma metastasis from healthy patients based on gene expression data. However, in practice, we can not measure the regret defined in (2), unless we have the knowledge of β^* . To resolve this issue, we first fit a logit model on the whole gravier data set and consider the estimated $\hat{\beta}$ as the ground truth and report the expected regret with respect to the estimated β . As reported in Gravier et al. (2010), 24 (out of 2905) BACs showed statistically significant difference (comparing Cy3/Cy5 values) between the two groups, motivating the use of a sparse logit model in our case. The estimated β^* in our sparse logistic model on the dataset had a sparsity of 18 using the dataset. In addition to this, we also treat the estimated noise variance from the fitted logit model as the true noise variance of the error induced by the environment in each round.

Appendix B Proof of Theorem 2

In this section, we present the detailed proof of Theorem 2. First, for clarity of presentation, we introduce some notations. We use X_t to denote the matrix $(x_{a_1}(1), \ldots, x_{a_t}(t))^{\top} \in \mathbb{R}^{t \times d}$. Given this, we denote the covariance matrix $\hat{\Sigma}_t = X_t^{\top} X_t / t$. Next, we define the set

$$\mathbb{S}_0^{d-1}(s) \triangleq \mathbb{S}^{d-1} \cap \{ v : \|v\|_0 \le s \}.$$

We also define the following:

Definition B.1. For a index set $I \subseteq [d]$ and $\alpha \in \mathbb{R}^+$, we define the restricted cone as

$$\mathbb{C}_{\alpha}(I) := \{ v \in \mathbb{R}^d : \|v_{I^c}\|_1 \le \alpha \|v_I\|_1, v_I \neq 0 \}$$

In high-dimensional literature one typically assumes compatibility condition on the design matrix X, i.e.,

$$\phi_{\mathsf{comp}}(S^*; X) := \inf_{\delta \in \mathbb{C}_7(S^*)} \frac{\|X\delta\|_2 \, |S^*|^{1/2}}{t^{1/2} \, \|\delta\|_1} > 0, \tag{4}$$

where $S^* = \{j : \beta_j^* \neq 0\}$. This is mainly to guarantee the estimation accuracy of high-dimensional estimators like LASSO (Bickel et al., 2009) or to show the posterior consistency in Bayesian high dimensional literature (Castillo et al., 2015).

As discussed in the main paper, we prove the theorem in three parts, the subsequent sections deal with each part separately.

B.1 Proof of part (i)

In this section we will show that the matrix $\hat{\Sigma}_t$ enjoys the SRC condition with high probability. As a warm up, we recall the definition of Orlicz norms:

Definition B.2 (Orlicz norms). For random variable Z we have the followings:

(a) The sub-Gaussian norm of a random variable Z, denoted $||Z||_{\psi_2}$, is defined as

$$||Z||_{\psi_2} := \inf\{\lambda > 0 : \mathbb{E}\{\exp(Z^2/\lambda^2)\} \le 2\}.$$

(b) The sub-Exponential norm of a random variable Z, denoted $||Z||_{\psi_1}$, is defined as

$$||Z||_{\psi_1} := \inf\{\lambda > 0 : \mathbb{E}\{\exp(|Z|/\lambda)\} \le 2\}.$$

The details and related properties can be found in Section 2.5.2 and Section 2.7 in Vershynin (2018). The following is a relationship between sub-gaussian and sub-exponential random variables.

Lemma B.1 (Sub-Exponential and sub-Gaussian squared). A random variable Z is sub-Gaussian iff Z^2 is sub-Exponential. Moreover,

$$\left\| Z^2 \right\|_{\psi_1} = \left\| Z \right\|_{\psi_2}^2.$$

Proof. The proof can be found in Lemma 2.7.4 of Vershynin (2018)

Lemma B.2 (Bernstein's inequality). Let Z_1, \ldots, Z_N be independent mean-zero sub-exponential random variable. Then for every $\delta \ge 0$ we have

$$\mathbb{P}\left(\left|\frac{1}{N}\sum_{i=1}^{N}Z_{i}\right|\geq\delta\right)\leq2\exp\left\{-c_{2}\min\left(\frac{\delta^{2}}{K_{0}^{2}},\frac{\delta}{K_{0}}\right)N\right\},$$

where $K_0 = \max_{i \in [N]} \|Z_i\|_{\psi_1}$.

Proof. The proof can be found in Corollary 2.8.3 of Vershynin (2018).

Proposition 1 (Empirical SRC). Let $\varepsilon = \min\{1/4, 1/(\tilde{c}\phi_u \vartheta^2 \xi K \log K + 3)\}$ for some universal constant $\tilde{c} > 0$ and also define the quantity $\kappa(\xi, \vartheta, K) \triangleq \min\{(4c_3K\xi\vartheta^2)^{-1}, 1/2\}$ for some universal positive constants c_3 . Then the followings are true for any constant C > 0:

$$\mathbb{P}\left(\phi_{\max}(Cs^*; \widehat{\Sigma}_t) \ge \frac{c_9 \vartheta^2 \phi_u \log K}{1 - 3\varepsilon}\right) \le \exp\left\{-c_8 t \log K + Cs^* \log K + Cs^* \log(3d/\varepsilon)\right\},$$

$$\mathbb{P}\left(\phi_{\min}(Cs^*; \widehat{\Sigma}_t) \le \frac{1}{8K\xi}\right) \le 2\exp\{-c_2\kappa^2(\xi, \vartheta, K)t + Cs^*\log K + Cs^*\log(3d/\varepsilon)\} + \exp\{-c_8\log(K)t + Cs^*\log K + Cs^*\log(3d/\varepsilon)\},\$$

where all c_i 's in the above display are universal positive constants.

Proof. We will first show the SRC condition for a fixed vector $v \in \mathbb{S}_0^{d-1}(Cs^*)$. Then, the whole argument will be extended via a ε -net argument.

Analysis for a fixed vector v: Let $v \in \mathbb{S}_0^{d-1}(Cs^*)$ be a fixed vector. Now note that following fact:

$$v^{\top}\widehat{\Sigma}_{t}v = \frac{1}{t}\sum_{\tau=1}^{t} \{v^{\top}x_{a_{\tau}}(\tau)\}^{2} \ge \frac{1}{t}\sum_{\tau=1}^{t}\min_{i\in[K]} \{v^{\top}x_{i}(\tau)\}^{2}.$$

We define $Z_{\tau,v} \triangleq \min_{i \in [K]} \{v^{\top} x_i(\tau)\}^2$ and note that for a fixed $v \in \mathbb{S}_0^{d-1}(Cs^*)$, the random variables $\{Z_{\tau,v}\}_{\tau=1}^t$ are i.i.d. across the time points. Moreover, due to Assumption 2.1(b) and Lemma B.1, we have

$$||Z_{\tau,v}||_{\psi_1} = \left\|\min_{i\in[K]} |v^{\top}x_i(\tau)|\right\|_{\psi_2}^2 \le c_3\vartheta^2.$$

Thus, $\{Z_{\tau,v}\}_{\tau=1}^t$ are i.i.d sub-exponential random variables. First we will show that $\mathbb{E}(Z_{\tau,v})$ is uniformly lower bounded. Due to Assumption 2.1(c) we have

$$\mathbb{P}(Z_{\tau,v} \le h) \le \sum_{i=1}^{K} \mathbb{P}\{(v^{\top} x_i(\tau))^2 \le h\} \le K\xi h.$$

Thus we have the following:

$$\mathbb{E}(Z_{\tau,v}) = \int_0^\infty \mathbb{P}(Z_{\tau,v} \ge u) \, du$$

$$\ge \int_0^h \mathbb{P}(Z_{\tau,v} \ge u) \, du$$

$$\ge \int_0^h (1 - K\xi u) \, du$$

$$= h(1 - K\xi h/2).$$
 (5)

Setting $h = 1/(K\xi)$ in Equation (5) yields $\mathbb{E}(Z_{\tau,v}) \ge \frac{1}{2K\xi}$. Now, using Lemma B.2, we have the following for a $\mu \in (0, c_1/||Z_{1,v}||_{\psi_1})$ and $\delta > 0$:

$$\mathbb{P}\left(\frac{1}{t}\sum_{\tau=1}^{t} \{Z_{\tau,v} - \mathbb{E}(Z_{\tau,v})\} \ge \delta\right) \le 2\exp\left\{-c_2 \min\left(\frac{\delta^2}{\|Z_{1,v}\|_{\psi_1}^2}, \frac{\delta}{\|Z_{1,v}\|_{\psi_1}}\right)t\right\}$$
$$\le 2\exp\left\{-c_2 \min\left(\frac{\delta^2}{c_3^2\vartheta^4}, \frac{\delta}{c_3\vartheta^2}\right)t\right\}.$$

Now choose $\delta = \min\{(4K\xi)^{-1}, c_3\vartheta^2/2\}$ to finally get

$$\mathbb{P}\left(\left|\frac{1}{t}\sum_{\tau=1}^{t} \{Z_{\tau,v} - \mathbb{E}(Z_{\tau,v})\}\right| \ge \frac{1}{4K\xi}\right) \le 2\exp\left\{-c_2\kappa^2(\xi,\vartheta,K)t\right\},\tag{6}$$

where $\kappa(\xi, \vartheta, K) = \min\{(4c_3K\xi\vartheta^2)^{-1}, 1/2\}$. Now recall that $\mathbb{E}(Z_{\tau,v}) \ge 1/(2K\xi)$, which shows that

$$\mathbb{P}\left\{\frac{1}{t}\sum_{\tau=1}^{t} Z_{\tau,v} \ge \frac{1}{4K\xi}\right\} \le 2\exp\left\{-c_2\kappa^2(\xi,\vartheta,K)t\right\}, \quad \forall \ v \in \mathbb{S}_0^{d-1}(Cs^*).$$
(7)

 ε -net argument: We consider a ε -net of the space $\mathbb{S}_0^{d-1}(Cs^*)$ constructed in a specific way which will be described shortly. We denote it by $\mathcal{N}_{\varepsilon}$. Let $J \subseteq [d]$ such that $|J| = Cs^*$ and consider the set $E_J = \mathbb{S}^{d-1} \cap \operatorname{span}\{e_j : j \in J\}$. Here e_j denotes the *j*th canonical basis of \mathbb{R}^d . Thus we have

$$\mathbb{S}_0^{d-1}(Cs^*) = \bigcup_{J:|J|=Cs^*} E_J.$$

Now we describe the procedure of constructing a net for $\mathbb{S}_0^{d-1}(Cs^*)$ which is essential for controlling the parse eigenvalues.

Greedy construction of net:

- Construct a ε -net of E_J for each J of size Cs^* . We denote this net by $\mathcal{N}_{\varepsilon,J}$. Note that $|\mathcal{N}_{\varepsilon,J}| \leq (3/\varepsilon)^{Cs^*}$ (Vershynin, 2018, Corollary 4.2.13) for $\varepsilon \in (0, 1)$ as E_J can be viewed as an unit ball embedded in \mathbb{R}^{Cs^*} .
- Then the net $\mathcal{N}_{\varepsilon}$ is constructed by taking union over all the $\mathcal{N}_{\varepsilon,J}$, i.e.,

$$\mathcal{N}_{\varepsilon} = \bigcup_{J:|J|=Cs^*} \mathcal{N}_{\varepsilon,J}$$

Thus, from from the construction we have

$$|\mathcal{N}_{\varepsilon}| \le {\binom{d}{Cs^*}} {\binom{3}{\varepsilon}}^{Cs^*} \le \exp\{Cs^*\log(3d/\varepsilon)\},\tag{8}$$

whenever $\varepsilon \in (0, 1)$. Now, we state an useful lemma on evaluating minimum eigenvalue on ε -net. **Lemma B.3.** Let A be a $m \times m$ symmetric positive-definite matrix and $\varepsilon \in (0, 1)$. Then, for ε -net $\mathcal{N}_{\varepsilon}$ of $\mathbb{S}_{0}^{d-1}(s)$ constructed in greedy way, we have

$$\phi_{\min}(s; A) \ge \min_{u \in \mathcal{N}_{\varepsilon}} u^{\top} A u - 3\varepsilon \phi_{\max}(s; A).$$

The proof of the lemma is deferred to Appendix D.1. Note that from Equation (7) and an union bound argument we get

$$\mathbb{P}\left(\min_{u\in\mathcal{N}_{\varepsilon}}v^{\top}\widehat{\Sigma}_{t}v\geq\frac{1}{4K\xi}\right)\geq1-2\exp\{-c_{2}\kappa^{2}(\xi,\vartheta,K)t+Cs^{*}\log K+Cs^{*}\log(3d/\varepsilon)\}.$$
 (9)

If $\phi_{\max}(Cs^*, \widehat{\Sigma}_t)$ is bounded with high probability, then for small ε , then along with Lemma B.3 we will readily have an uniform lower bound on $\phi_{\min}(Cs^*, \widehat{\Sigma}_t)$.

Bounding $\phi_{\max}(Cs^*, \widehat{\Sigma}_t)$: Here we gain start with $v \in \mathcal{N}_{\varepsilon}$. Similar, to previous discussion we have

$$v^{\top} \widehat{\Sigma}_t v = \frac{1}{t} \sum_{\tau=1}^t \{ v^{\top} x_{a_{\tau}}(\tau) \}^2 \le \frac{1}{t} \sum_{\tau=1}^t \max_{i \in [K]} \{ v^{\top} x_i(\tau) \}^2.$$

We define $W_{\tau,v} \triangleq \max_{i \in [K]} \{v^{\top} x_i(\tau)\}^2$ and note that for a fixed $v \in \mathbb{S}_0^{d-1}(Cs^*)$, the random variables $\{W_{\tau,v}\}_{\tau=1}^t$ are i.i.d. across the time points. Moreover, due to Assumption 2.1(b) and Lemma B.1, we have

$$||W_{\tau,v}||_{\psi_1} = \left||\max_{i\in[K]} \{v^{\top} x_i(\tau)\}^2\right||_{\psi_1} \le c_4 K \vartheta^2.$$

Thus, $\{W_{\tau,v}\}_{\tau=1}^t$ are i.i.d sub-exponential random variables. Recall, that $\phi_{\max}(Cs^*, \Sigma_i) \leq \phi_u$ for all $i \in [K]$. The next lemma provides an upper bound on the moment generating function (MGF) of sub-Exponential random variables.

Lemma B.4. (Vershynin, 2018, Lemma 2.8.1) Let X be a mean-zero, sub-Exponential random variable. Then there exists positive constants c_5 , c_6 , such that for any λ with $|\lambda| \leq c_5 / ||X||_{\psi_1}$, the following is true:

$$\mathbb{E}\{\exp(\lambda X)\} \le \exp(c_6 \lambda^2 \|X\|_{\psi_1}^2).$$

Equipped with the above lemma we have the following;

$$\mathbb{P}\left(\frac{1}{t}\sum_{\tau=1}^{t}W_{\tau,v}-\phi_{u}\geq\delta\right) = \mathbb{P}\left(\sum_{\tau=1}^{t}\{W_{\tau,v}-\phi_{u}\}\geq\delta t\right) \\
= \mathbb{P}\left(\exp\left\{\mu\sum_{\tau=1}^{t}(W_{\tau,v}-\phi_{u})\right\}\geq e^{\mu\delta t}\right) \qquad (10) \\
\leq e^{-\mu\delta t}\prod_{\tau=1}^{t}\mathbb{E}\{e^{\mu(W_{\tau,v}-\phi_{u})}\}.$$

For a fixed $\tau \in [t]$ we note the following;

$$\mathbb{E}\{e^{\mu(W_{\tau,v}-\phi_u)}\} \le \sum_{i=1}^{K} \mathbb{E}\{e^{\mu[(v^{\top}x_i(\tau))^2 - \phi_u]}\}$$
$$\le \sum_{i=1}^{K} \mathbb{E}\{e^{\mu[(v^{\top}x_i(\tau))^2 - v^{\top}\Sigma_i v]}\}$$

For brevity let $\kappa_i \triangleq \|\{v^{\top} x_i(\tau)\}^2\|_{\psi_1}$. If we choose $\mu \leq c_5 / \max_{i \in [K]} \kappa_i$, then by Lemma B.4 we have

$$\mathbb{E}\left\{e^{\mu(W_{\tau,v}-\phi_u)}\right\} \le \exp\left(c_6\mu^2 \max_{i\in[K]}\kappa_i^2 + \log K\right)$$

Using the above inequality in Equation (10), it follows that

$$\mathbb{P}\left(\frac{1}{t}\sum_{\tau=1}^{t}W_{\tau,v}-\phi_{u}\geq\delta\right)\leq\exp\left(-\mu\delta t+c_{6}t\mu^{2}\max_{i\in[K]}\kappa_{i}^{2}+t\log K\right).$$
(11)

The right hand side of Equation (11) is minimized at $\mu = \delta/(2c_2 \max_{i \in [K]} \kappa_i^2)$ with the minimum value of

$$\exp\left(-\frac{\delta^2 t}{4c_6 \max_{i \in [K]} \kappa_i^2} + t \log K\right).$$

If $\delta/(2c_6 \max_{i \in [K]} \kappa_i^2) > c_5/\max_{i \in [K]} \kappa_i^2$, then the right hand side of Equation (11) is minimized at $\mu = c_5/\max_{i \in [K]} \kappa_i^2$ and we get

$$\mathbb{P}\left(\frac{1}{t}\sum_{\tau=1}^{t}W_{\tau,v}-\phi_{u}\geq\delta\right)\leq\exp\left(-\frac{c_{5}\delta t}{\max_{i}\kappa_{i}}+c_{6}c_{5}^{2}t+t\log K\right).$$

Using the fact that $\delta/(2c_6 \max_{i \in [K]} \kappa_i^2) > c_5/\max_{i \in [K]} \kappa_i^2$, the right hand side of the above display can be upper bounded by

$$\exp\left(-\frac{c_5\delta t}{2\max_i\kappa_i} + t\log K\right)$$

Thus we have for all $\delta > 0$

$$\mathbb{P}\left(\frac{1}{t}\sum_{\tau=1}^{t}W_{\tau,v} - \phi_u \ge \delta\right) \le \exp\left(-\min\left\{\frac{\delta^2 t}{4c_6 \max_{i \in [K]}\kappa_i^2}, \frac{c_5\delta t}{2\max_i\kappa_i}\right\} + t\log K\right).$$
(12)

Next we set $\delta = c_7 \vartheta^2 \phi_u \log K$ for sufficiently large $c_7 > 0$. Then Equation (12) yields

$$\mathbb{P}\left(\frac{1}{t}\sum_{\tau=1}^{t}W_{\tau,v}-\phi_{u}\geq\delta\right)\leq\exp\left(-c_{8}t\log K\right).$$

Finally taking union bond over all vectors in $\mathcal{N}_{\varepsilon}$ we get

$$\mathbb{P}\left(\forall v \in \mathcal{N}_{\varepsilon} : v^{\top} \widehat{\Sigma}_{t} v \ge c_{9} \vartheta^{2} \phi_{u} \log K\right) \le \exp\left\{-c_{8} t \log K + C s^{*} \log K + C s^{*} \log(3d/\varepsilon)\right\}.$$
(13)

Next, to prove the same for all $v \in \mathbb{S}_0^{d-1}(Cs^*)$ we need the following lemma.

Lemma B.5 (maximum sparse eigenvalue on net). Let A be a $m \times m$ symmetric positive-definite matrix and $\varepsilon \in (0, 1/3)$. Then, for ε -net $\mathcal{N}_{\varepsilon}$ of $\mathbb{S}_0^{d-1}(s)$ constructed in greedy way, we have

$$\max_{v \in \mathcal{N}_{\varepsilon}} v^{\top} A v \le \max_{v \in \mathbb{S}_{0}^{d-1}(Cs^{*})} v^{\top} A v \le \frac{1}{1 - 3\varepsilon} \max_{v \in \mathcal{N}_{\varepsilon}} v^{\top} A v.$$

The proof of the above lemma is deferred to Appendix D.2. Now we set some $\varepsilon \in (0, 1/3)$. In light of the above lemma we immediately have that

$$\mathbb{P}\left(\phi_{\max}(Cs^*; \widehat{\Sigma}_t) \ge \frac{c_9 \vartheta^2 \phi_u \log K}{1 - 3\varepsilon}\right) \le \exp\left\{-c_8 t \log K + Cs^* \log K + Cs^* \log(3d/\varepsilon)\right\}.$$
(14)

Finally using Equation (9), (14) and Lemma B.3 we have

$$\mathbb{P}\left(\phi_{\min}(Cs^*; \widehat{\Sigma}_t) \ge \frac{1}{4K\xi} - \frac{3\varepsilon c_9 \vartheta^2 \log K}{1 - 3\varepsilon} \phi_u\right) \\
\ge 1 - 2\exp\{-c_2\kappa^2(\xi, \vartheta, K)t + Cs^* \log K + Cs^* \log(3d/\varepsilon)\} \\
- \exp\{-c_8 t \log K + Cs^* \log K + Cs^* \log(3d/\varepsilon)\}.$$
(15)

Now the result follows from taking $\varepsilon = \min\{1/4, 1/(24\phi_u \vartheta^2 \xi K \log K + 3)\}.$

B.2 Proof of part (ii)

In this section we will show that the matrix $\widehat{\Sigma}_t$ enjoys the compatibility condition (4) with high probability. This is equivalent to showing that the quantity

$$\Psi(S^*; \widehat{\Sigma}_t) \triangleq \inf_{\delta \in \mathbb{C}_7(S^*)} \left(\frac{\delta^\top \widehat{\Sigma}_t \delta}{\|\delta\|_1^2} \right) s^* = \phi_{\mathsf{comp}}(S^*; X_t)^2.$$

is bounded away from 0 with high probability. First we present the Transfer lemma (Oliveira, 2013, Lemma 5) below.

Lemma B.6 (Transfer lemma). Suppose $\widehat{\Sigma}_t$ and Σ are matrix with non-negative diagonal entries, and assume $\eta \in (0, 1)$, $m \in [d]$ are such that

$$\forall v \in \mathbb{R}^d \text{ with } \|v\|_0 \le m, v^\top \widehat{\Sigma}_t v \ge (1 - \eta) v^\top \Sigma v.$$
(16)

Assume D is a diagonal matrix whose diagonal entries $D_{j,j}$ are non-negative and satisfy $D_{j,j} \ge (\widehat{\Sigma}_t)_{j,j} - (1 - \eta) \Sigma_{j,j}$. Then

$$\forall x \in \mathbb{R}^d, x^\top \widehat{\Sigma}_t x \ge (1 - \eta) x^\top \Sigma x - \frac{\left\| D^{1/2} x \right\|_1^2}{m - 1}.$$
(17)

Condition (16) basically demands that $\hat{\Sigma}_t$ enjoys SRC condition with the sparsity parameter m. Then under the proper choice of diagonal matrix D with sufficiently large diagonal elements $\{D_{j,j}\}_{j=1}^d$, Equation (17) will yield the desired compatibility condition for $\hat{\Sigma}_t$. We formally state the result in the following lemma:

Proposition 2 (Empirical compatibility condition). Assume the conditions of Proposition 1 hold. Also assume that Assumption 2.1(d) holds with $C = C_0 \phi_u \vartheta^2 \xi K \log K$ for some sufficiently large universal constant $C_0 > 0$. Then there exists a positive constant C_1 such that the following is true:

$$\mathbb{P}\left(\Psi(S^*; \widehat{\Sigma}_t) \ge \frac{1}{C_1 \xi K}\right)$$

= 1 - 2 exp{-c_2 \kappa^2(\xi, \vartheta, K)t + Cs^* log K + Cs^* log(3d/\varepsilon)}
- 2 exp{-c_8 t log K + Cs^* log K + Cs^* log(3d/\varepsilon)}

with $\varepsilon = \min\{1/4, 1/(\tilde{c}\phi_u \vartheta^2 \xi K \log K + 3)\}$ for the same universal constant $\tilde{c} > 0$ in Proposition 1 and $\kappa(\xi, \vartheta, K) = \min\{(4c_3K\xi\vartheta^2)^{-1}, 1/2\}.$

Proof. As suggested before we will make use of Lemma B.6. Towards this, we set $\Sigma = \frac{1}{4K\xi} \mathbb{I}_d$ and $D = \text{diag}(\widehat{\Sigma}_t)$. Next, we define the following two events:

$$\mathcal{G}_{t,1} := \left\{ \phi_{\max}(Cs^*; \widehat{\Sigma}_t) \le \frac{c_9 \vartheta^2 \phi_u \log K}{1 - 3\varepsilon} \right\},$$
$$\mathcal{G}_{t,2} := \left\{ \phi_{\min}(Cs^*; \widehat{\Sigma}_t) \ge \frac{1}{8K\xi} \right\},$$

where the constants ε and c_9 are same as in Proposition 1. Under $\mathcal{G}_{t,1}$ and $\mathcal{G}_{t,2}$, the inequality in Equation (16) holds with $\eta = 1/2$ and $m = Cs^*$. Also, due construction of D, we trivially have

$$D_{j,j} \ge (\widehat{\Sigma}_t)_{j,j} - (1-\eta)\Sigma_{j,j}.$$

Lastly, note that

$$\max_{j \in [d]} D_{j,j} = \max_{j \in [d]} (\widehat{\Sigma}_t)_{j,j} = \max_{j \in [d]} e_j^\top \widehat{\Sigma}_t e_j \le \frac{c_9 \vartheta^2 \phi_u \log K}{1 - 3\varepsilon}$$
(18)

under $\mathcal{G}_{t,1}$. Equipped with Lemma B.6, under $\mathcal{G}_{t,1}$ and $\mathcal{G}_{t,2}$, for all $x \in \mathbb{C}_7(S^*) \cap \mathbb{S}^{d-1}$ we have the following:

$$x^{\top} \widehat{\Sigma}_t x \ge \frac{1}{8K\xi} - \frac{\left\| D^{1/2} x \right\|_1^2}{Cs^* - 1}$$
$$\ge \frac{1}{8K\xi} - \frac{\left(\frac{c_9 \vartheta^2 \phi_u \log K}{1 - 3\varepsilon}\right) \|x\|_1^2}{Cs^* - 1}$$
$$\ge \frac{1}{8K\xi} - \frac{\left(\frac{c_9 \vartheta^2 \phi_u \log K}{1 - 3\varepsilon}\right) 64s^*}{Cs^* - 1}$$

The last inequality follows form the fact that

$$\|x\|_{1} = \|x_{S^{*}}\|_{1} + \|x_{(S^{*})^{c}}\|_{1} \le 8 \|x_{S^{*}}\|_{1} \le 8\sqrt{s^{*}} \|x_{S^{*}}\|_{2} \le 8\sqrt{s^{*}}.$$
(19)

Thus, if $C \gtrsim \frac{\phi_u \vartheta^2 \xi K \log K}{1-3\varepsilon} + \frac{1}{s^*}$ then $x^\top \widehat{\Sigma}_t x \ge 1/(16K\xi)$. Also, note that from the choice of ε in Proposition 1, we have $\varepsilon < 1/4$. This further tells that if $C = C_0 \phi_u \vartheta^2 \xi K \log K$ for large enough $C_0 > 0$, then

$$\inf_{\delta \in \mathbb{C}_7(S^*)} \frac{\delta^\top \Sigma_t \delta}{\|\delta\|_2^2} \ge \frac{1}{16K\xi}$$

Then, the result follows from Proposition 1. Using this and Equation (19) we also have

$$\Psi(S^*; \widehat{\Sigma}_t) \ge \frac{1}{64} \inf_{\delta \in \mathbb{C}_7(S^*)} \frac{\delta^\top \widehat{\Sigma}_t \delta}{\|\delta\|_2^2} \ge \frac{1}{C_1 \xi K}$$

where $C_1 = 1024$. Finally, the result follows from conditioning over the events $\mathcal{G}_{t,1}$ and $\mathcal{G}_{t,2}$ and using Proposition 1.

B.3 Proof of part (iii)

In this section we will establish the desired regret bound in Theorem 2. The main tools that has been used to prove the regret bound is the Bayesian contraction in high-dimensional linear regression problem. In particular, we will use Theorem 3 to control the ℓ_1 -distance between $\tilde{\beta}_t$ and β^* at each time point $t \in [T]$.

We recall that $X_t = (x_{a_1}(1), \dots, x_{a_t}(t))^{\top}$. Also, note that the sequence $\{x_{a_\tau}(\tau)\}_{\tau=1}^t$ forms an adapted sequence of observations, i.e., $x_{a_\tau}(\tau)$ may depend on the history $\{x_{a_u}(u), r(u)\}_{u=1}^{\tau-1}$. Also, recall that $\{\epsilon(\tau)\}_{\tau=1}^t$ are mean-zero σ -sub-Gaussian errors.

Lemma B.7 (Bernstein Concentration). Let $\{D_k, \mathcal{F}_k\}_{k=1}^{\infty}$ be a martingale difference sequence, and suppose that D_k is a σ -sub-Gaussian in adapted sense, i.e., for all $\alpha \in \mathbb{R}, \mathbb{E}[e^{\alpha D_k} | \mathcal{F}_{k-1}] \leq e^{\alpha^2 \sigma^2/2}$ almost surely. Then, for all $t \geq 0, \mathbb{P}(|\sum_{k=1}^t D_k| \geq \delta) \leq 2 \exp\{-\delta^2/(2t\sigma^2)\}$.

Proof. Proof of Lemma B.7 follows from Theorem 2.19 of Wainwright (2019) by setting $\alpha_k = 0$ and $\nu_k = \sigma$ for all k.

Lemma B.7 is the main tool that is used to control the correlation between $\epsilon_t := (\epsilon(1), \ldots, \epsilon(t))^{\top}$ and the chosen contexts X_t which is important to control the Bayesian contraction of the posterior distribution in each round. To elaborate, let $X_t^{(j)}$ be the *j*th column for $j \in [d]$ and define $D_{t,j} := \epsilon(t)x_{at,j}(t)$. Note that for a fixed $j \in [d]$, $\{D_{\tau,j}\}_{\tau=1}^t$ forms a martingale difference sequence with respect to the filtration $\{\mathcal{F}_{\tau}\}_{\tau=1}^{t-1}$ with $\mathcal{F}_{\tau} := \sigma(\mathcal{H}_{\tau})$ is the σ -algebra generated by \mathcal{H}_{τ} and $\mathcal{F}_1 = \emptyset$. Also note that

$$\mathbb{E}(e^{\alpha D_{t,j}} \mid \mathcal{F}_{t-1}) \leq \mathbb{E}\{e^{\alpha^2 \sigma^2 x_{a_t,j}^2(t)/2}\} \leq \mathbb{E}\{e^{\alpha^2 \sigma^2 x_{\max}^2/2}\}.$$

Thus, using Lemma B.7, we have the following proposition:

Proposition 3 (Lemma EC.2, Bastani and Bayati (2020)). Define the event

$$\mathcal{T}_t(\lambda_0(\gamma)) := \left\{ \max_{j \in [d]} \frac{\left| \epsilon_t^\top X_t^{(j)} \right|}{t} \le \lambda_0(\gamma) \right\},\,$$

where $\lambda_0(\gamma) = x_{\max}\sigma\sqrt{(\gamma^2 + 2\log d)/t}$. Then we have $\mathbb{P}\left\{\mathcal{T}_t(\lambda_0(\gamma))\right\} \ge 1 - 2\exp(-\gamma^2/2)$.

The proof of the above proposition mainly relies on the martingale difference structure and Lemma B.2. It is important to mention that the proof does not depend on any particular algorithm.

For notational brevity we define $||X_t|| := \max_{j \in [d]} \sqrt{(X_t^\top X_t)_{j,j}}$. Next, we will set $\gamma = \gamma_t := \sqrt{2 \log t}$. Hence by Proposition 3 we have $\mathbb{P} \{ \mathcal{T}_t(\lambda_0(\gamma_t)) \} \ge 1 - 2t^{-1}$. Also recall that, under $\mathcal{G}_{t,1}$ and $\mathcal{G}_{t,2}$, we have

$$\frac{1}{\sqrt{8K\xi}} \le \left\|X_t\right\| / \sqrt{t} \le \sqrt{4c_9 \phi_u \vartheta^2 \log K}.$$
(20)

Also, under $\mathcal{G}_{t,2} \cap \mathcal{T}_t(\lambda_0(\gamma_t))$ it follows that

$$\max_{j \in [d]} \frac{\left| \epsilon_t^\top X_t^{(j)} \right|}{\sigma} \le x_{\max} \sqrt{2t(\log d + \log t)} = \overline{\lambda}_t$$
(21)

Now, we are ready to present the proof of the main regret bound.

Main regret bound

Recall the definition of regret is $R(T) = \sum_{t=1}^{T} \Delta_{a_t}(t)$, where $\Delta_{a_t}(t) = x_{a_t^*}(t)^{\top} \beta^* - x_{a_t}(t)^{\top} \beta^*$. Next, we partition the whole time horizon [T] in to two parts, namely $\{t : 1 \leq t \leq T_0\}$ and $\{t : T_0 \leq t \leq T\}$, where T_0 will be chosen later. Thus, the regret can be written as

$$R(T) = \underbrace{\sum_{t=1}^{T_0} \Delta_{a_t}(t)}_{R(T_0)} + \underbrace{\sum_{t=T_0+1}^{T} \Delta_{a_t}(t)}_{\tilde{R}(T)}.$$

All notations for expectation operators and probability measures are given in Appendix C.

Now by Assumption 2.1(a) and 2.2(a) we have the following inequality:

$$\mathbb{E}\{R(T_0)\} \le 2x_{\max}b_{\max}T_0.$$
(22)

Next, we focus on the term $\tilde{R}(T)$. First, we define a few quantities below:

$$\overline{\phi}_{t}(s) := \inf_{\delta} \left\{ \frac{\|X_{t}\delta\|_{2} |S_{\delta}|^{1/2}}{t^{1/2} \|\delta\|_{1}} : 0 \neq |S_{\delta}| \leq s \right\},$$

$$\widetilde{\phi}_{t}(s) := \inf_{\delta} \left\{ \frac{\|X_{t}\delta\|_{2}}{t^{1/2} \|\delta\|_{2}} : 0 \neq |S_{\delta}| \leq s \right\}.$$
(23)

、

Now set

$$\begin{split} \overline{\psi}_t(S) &= \bar{\phi}_t \left(\left(2 + \frac{40}{A_4} + \frac{128A_4^{-1}x_{\max}^2}{\Psi(S,\widehat{\Sigma}_t)} \right) |S| \right), \\ \widetilde{\psi}_t(S) &= \widetilde{\phi}_t \left(\left(2 + \frac{40}{A_4} + \frac{128A_4^{-1}x_{\max}^2}{\Psi(S,\widehat{\Sigma}_t)} \right) |S| \right). \end{split}$$

Note that $\overline{\phi}_t(s) \ge \widetilde{\phi}_t(s)$, hence $\overline{\psi}_t(S) \ge \widetilde{\psi}_t(S)$.

Recall that

$$5\overline{\lambda}_t/3 \le \lambda_t \le 2\overline{\lambda}_t,\tag{24}$$

under $\mathcal{G}_{t,1}$ and $\mathcal{G}_{t,2}$. Also define the following events:

$$\mathcal{E}_t := \left\{ \left\| \tilde{\beta}_{t+1} - \beta^* \right\|_1 \le Q_4 \sigma x_{\max} K \xi(D_* + s^*) \sqrt{\frac{\log d + \log t}{t}} \right\}.$$

where $D_* = \{1 + (40/A_4) + 128A_4^{-1}x_{\max}^2/\Psi(S^*, \widehat{\Sigma}_t)\}s^*$ and Q_4 is large enough universal constant as specified in Theorem 3. Also we have

$$\begin{split} \bar{\psi}_t(S) &\leq \bar{\phi}_t \left(\left(2 + \frac{40}{A_4} + \frac{64A_4^{-1}x_{\max}^2}{\Psi(S,\widehat{\Sigma}_t)}\frac{\lambda}{\bar{\lambda}} \right) |S| \right), \text{and} \\ \tilde{\psi}_t(S) &\leq \tilde{\phi}_t \left(\left(2 + \frac{40}{A_4} + \frac{64A_4^{-1}x_{\max}^2}{\Psi(S,\widehat{\Sigma}_t)}\frac{\lambda}{\bar{\lambda}} \right) |S| \right). \end{split}$$

Next, by Proposition 2, the event

$$\mathcal{G}_{t,3} := \left\{ \Psi(S^*; \widehat{\Sigma}_t) \ge \frac{1}{C_1 \xi K} \right\}$$

holds with probability of at least $1 - 2\exp\{-c_2\kappa^2(\xi,\vartheta,K)t + Cs^*\log K + Cs^*\log(3d/\varepsilon)\}$ – $2\exp\{-c_8t\log K + Cs^*\log K + Cs^*\log(3d/\varepsilon)\}\}$. For, notational brevity, we define

$$\tilde{C} := C_1 \xi K.$$

Also, define the event

$$\mathcal{G}_{t,4} := \left\{ \widetilde{\psi}_t^2(S^*) \ge \frac{1}{8K\xi} \right\}.$$

Noting that $\widetilde{\psi}_t^2(S^*) = \phi_{\min}(\tilde{C}_1 s^*; \widehat{\Sigma}_t)$ with

$$\tilde{C}_1 = 2 + \frac{40}{A_4} + 128A_4^{-1}x_{\max}^2\tilde{C}_2$$

an argument similar to the proof of Proposition 1 yields

$$\mathbb{P}\left(\mathcal{G}_{t,4}\right) \geq 1 - 2\exp\left\{-c_{2}\kappa^{2}(\xi,\vartheta,K)t + \tilde{C}_{1}s^{*}\log K + \tilde{C}_{1}s^{*}\log(3d/\varepsilon)\right\} - \exp\left\{-c_{8}\log(K)t + \tilde{C}_{1}s^{*}\log K + \tilde{C}_{1}s^{*}\log(3d/\varepsilon)\right\}.$$
(25)

Finally, Using Proposition 3 with $\gamma = \gamma_d$ and the result of Theorem 3, we have the following:

$$\mathbb{P}(\mathcal{E}_{t}^{c}) = \mathbb{E}_{X_{t}} \mathbb{E}_{t}^{X} \left(\mathbb{1}_{\mathcal{E}_{t}^{c}} \right) \\
= \mathbb{E}_{X_{t}} \mathbb{E}_{t,\mathbf{r}_{t}}^{X} \left\{ \Pi_{t}^{X} \left(\mathcal{E}_{t}^{c} \mid \mathbf{r}_{t} \right) \right\} \\
= \mathbb{E}_{X_{t}} \mathbb{E}_{t,\mathbf{r}_{t}}^{X} \left\{ \Pi_{t}^{X} \left(\mathcal{E}_{t}^{c} \mid \mathbf{r}_{t} \right) \mathbb{1}_{\mathcal{T}_{t}(\lambda_{0}(\gamma_{t})) \cap \mathcal{G}_{t,2}} \right\} + \mathbb{E}_{X_{t}} \mathbb{E}_{t,\mathbf{r}_{t}}^{X} \left\{ \Pi_{t}^{X} \left(\mathcal{E}_{t}^{c} \mid \mathbf{r}_{t} \right) \mathbb{1}_{\mathcal{T}_{t}^{c}(\lambda_{0}(\gamma_{t})) \cup \cap \mathcal{G}_{t,2}^{c}} \right\} (26) \\
\leq \frac{M_{1}}{d^{s^{*}}} + \frac{2}{t} + 2 \exp\{-c_{2}\kappa^{2}(\xi, \vartheta, K)t + Cs^{*}\log K + Cs^{*}\log(3d/\varepsilon)\} \\
+ \exp\{-c_{8}t\log K + Cs^{*}\log K + Cs^{*}\log(3d/\varepsilon)\}$$

for some large universal constant $M_1 > 0$. Next, let $\mathcal{G}_t = \bigcap_{i=1}^4 \mathcal{G}_{t,i}$. Under the event $\mathcal{E}_t \cap \mathcal{G}_t$, we have

$$D_* + s^* \le \underbrace{\left(2 + \frac{40}{A_4} + \frac{128C_1 K \xi x_{\max}^2}{A_4}\right)}_{:=\rho} s^*,$$

and,

$$\left\|\tilde{\beta}_{t+1} - \beta^*\right\|_1 \le M_2 \rho \sigma x_{\max} \xi K \left\{\frac{s^{*2}(\log d + \log t)}{t}\right\}^{1/2},$$

where M_2 is an universal constant depending on A_4 . Now we set

$$\delta_t = M_2 \rho \sigma x_{\max}^2 \xi K \left\{ \frac{s^{*2} (\log d + \log t)}{t} \right\}^{1/2}.$$

It follows that under $\mathcal{E}_{t-1} \cap \mathcal{G}_{t-1}$, we have the following almost sure inequality:

$$\begin{aligned} \Delta_{a_{t}}(t) &= x_{a_{t}^{*}}^{\top}(t)\beta^{*} - x_{a_{t}}^{\top}(t)\beta^{*} \\ &= x_{a_{t}^{*}}^{\top}(t)\beta^{*} - x_{a_{t}^{*}}^{\top}(t)\tilde{\beta}_{t} + \underbrace{(x_{a_{t}^{*}}^{\top}(t)\tilde{\beta}_{t} - x_{a_{t}^{*}}^{\top}(t)\tilde{\beta}_{t})}_{\leq 0} + x_{a_{t}^{*}}^{\top}(t)\tilde{\beta}_{t} - x_{a_{t}^{*}}^{\top}(t)\beta^{*} \\ &\leq \left\| x_{a_{t}^{*}}(t) \right\|_{\infty} \|\tilde{\beta}_{t} - \beta^{*}\|_{1} + \| x_{a_{t}}(t) \|_{\infty} \|\tilde{\beta}_{t} - \beta^{*}\|_{1} \\ &\leq 2\delta_{t-1}. \end{aligned}$$

Finally define the event

$$\mathcal{M}_t := \left\{ x_{a_t^*}^\top \beta^* > \max_{i \neq a_t^*} x_{a_t}^\top \beta^* + h_{t-1} \right\}.$$

Under $\mathcal{M}_t \cap \mathcal{E}_{t-1} \cap \mathcal{G}_{t-1}$, we have the following for any $i \neq a_t^*$:

$$x_{a_t^*}^{\top}(t)\tilde{\beta}_t - x_i^{\top}(t)\tilde{\beta}_t = \left\langle x_{a_t^*}(t), \tilde{\beta}_t - \beta^* \right\rangle + \left\langle x_{a_t}(t) - x_i(t), \beta^* \right\rangle + \left\langle x_i(t), \beta^* - \tilde{\beta}_t \right\rangle$$
$$\geq -\delta_{t-1} + h_{t-1} - \delta_{t-1}.$$

Thus, if we set $h_{t-1} = 3\delta_{t-1}$ then $x_{a_t}^{\top}(t)\tilde{\beta}_t - \max_{i \neq a_t} x_i^{\top}(t)\tilde{\beta}_t \ge \delta_{t-1}$. As a result, in *t*th round the regret is 0 almost surely as the optimal arm will be chosen with probability 1. Thus, finally using Assumption 2.2(b), we have

$$\mathbb{E}(\Delta_{a_{t}}(t)) = \mathbb{E}\{\Delta_{a_{t}}(t)\mathbb{1}_{\mathcal{M}_{t}^{c}}\} \\
= \mathbb{E}\{\Delta_{a_{t}}(t)\mathbb{1}_{\mathcal{M}_{t}^{c}\cap\mathcal{E}_{t-1}\cap\mathcal{G}_{t-1}}\} + \mathbb{E}\{\Delta_{a_{t}}(t)\mathbb{1}_{\mathcal{M}_{t}^{c}\cap(\mathcal{E}_{t-1}\cap\mathcal{G}_{t-1})^{c}}\} \\
\leq 2\delta_{t-1}\mathbb{P}(\mathcal{M}_{t}^{c}) + 2x_{\max}b_{\max}\mathbb{P}(\mathcal{E}_{t}^{c}\cup\mathcal{G}_{t}^{c}) \\
\leq 2\delta_{t-1}\mathbb{P}(\mathcal{M}_{t}^{c}) + \frac{2M_{1}x_{\max}b_{\max}}{d^{s^{*}}} + \frac{2x_{\max}b_{\max}}{d} \\
+ M_{3}x_{\max}b_{\max}\exp\{-c_{2}\kappa^{2}(\xi,\vartheta,K)t + Ds^{*}\log K + Ds^{*}\log(3d/\varepsilon)\} \\
+ M_{4}x_{\max}b_{\max}\exp\{-c_{8}\log(K)t + Ds^{*}\log K + Ds^{*}\log(3d/\varepsilon)\},$$
(27)

where M_3 , M_4 are large enough universal positive constants and $D = \max\{C, \tilde{C}_1\} = \Theta(\phi_u \vartheta^2 \xi K \log K)$. Thus, if we set

$$T_0 = M_5 \max\left\{\frac{1}{\kappa^2(\xi, \vartheta, K)}, \frac{1}{\log K}\right\} (Ds^* \log K + Ds^* \log(3d/\varepsilon)),$$
(28)

for some large universal constant $M_5 > 0$. Thus, we have

$$\mathbb{E}\{\tilde{R}(T)\} \leq 2 \sum_{\substack{t=T_0+1\\I_{\omega}}}^{T} \delta_{t-1} \mathbb{P}(\mathcal{M}_t^c) + M_6 x_{\max} \exp\left\{-M_7 (Ds^* \log K + Ds^* \log(3d/\varepsilon))\right\} + O(\log T).$$

Recall that

$$\delta_t = M_2 \rho \sigma x_{\max}^2 \xi K \left\{ \frac{s^{*2} (\log d + \log t)}{t} \right\}^{1/2}.$$

For $\omega \in [0, 1]$ we have

$$I_{\omega} \leq 2 \sum_{t=T_{0}+1}^{T} \delta_{t-1} \left(\frac{3\delta_{t-1}}{\Delta_{*}}\right)^{\omega} \\ \approx \frac{\left\{3M_{2}\rho\sigma x_{\max}^{2}\xi K\right\}^{1+\omega} s^{*1+\omega}}{\Delta_{*}^{\omega}} \int_{T_{0}}^{T} (\log d + \log u)^{\frac{1+\omega}{2}} u^{-\frac{1+\omega}{2}} du \qquad (29) \\ \lesssim \begin{cases} \frac{\left[3M_{2}\rho\sigma x_{\max}^{2}\xi K\right]^{1+\omega} s^{*1+\omega} (\log d)^{\frac{1+\omega}{2}} T^{\frac{1-\omega}{2}}}{\Delta_{*}^{\omega}}, & \text{for } \omega \in [0,1), \\ \frac{\left[3M_{2}\rho\sigma x_{\max}^{2}\xi K\right]^{2} s^{*2} (\log d + \log T) \log T}{\Delta_{*}}, & \text{for } \omega = 1. \end{cases}$$

For $\omega \in (1,\infty)$ we have

$$I_{\omega} \le 2\sum_{t=T_0+1}^{T} \delta_{t-1} \min\left\{1, \left(\frac{3\delta_{t-1}}{\Delta_*}\right)^{\omega}\right\}.$$
(30)

Note that

$$\frac{3\delta_{t-1}}{\Delta_*} \le 1 \Rightarrow t \ge T_1 := \left[3M_2\rho\sigma x_{\max}^2\xi K\right]^2 \frac{s^{*2}\log d}{\Delta_*^2} + 1.$$

Thus. from Equation (30) we have

$$\begin{split} I_{\omega} &\leq 2 \sum_{t=T_{0}+1}^{T_{1}} \delta_{t-1} + 2 \sum_{t=T_{1}+1}^{T} \delta_{t-1} \left(\frac{3\delta_{t-1}}{\Delta_{*}}\right)^{\omega} \\ &\leq 2 \int_{T_{0}}^{T_{1}} M_{2} \rho \sigma x_{\max}^{2} \xi K \left\{\frac{s^{*2} (\log d + \log u)}{u}\right\}^{1/2} du + 2 \sum_{t=T_{1}+1}^{T} \delta_{t-1} \left(\frac{3\delta_{t-1}}{\Delta_{*}}\right)^{\omega} \\ &\lesssim 4 \left[M_{2} \rho \sigma x_{\max}^{2} \xi K\right]^{2} \left\{\frac{s^{*2} (\log d + \log T)}{\Delta_{*}}\right\} + 2 J_{\omega}, \end{split}$$

where $J_{\omega} := \sum_{t=T_1+1}^{T} \delta_{t-1} \left(\frac{3\delta_{t-1}}{\Delta_*}\right)^{\omega}$. Finally, we give bound on the term J_{ω} :

$$J_{\omega} = \sum_{t=T_{1}+1}^{T} \delta_{t-1} \left(\frac{3\delta_{t-1}}{\Delta_{*}}\right)^{\omega}$$

$$= \sum_{t=2}^{T} \delta_{t-1} \left(\frac{3\delta_{t-1}}{\Delta_{*}}\right)^{\omega} \mathbb{1}\{3\delta_{t-1}/\Delta_{*} \leq 1\}$$

$$\leq \left(\frac{3^{\frac{\omega}{1+\omega}}M_{2}\rho\sigma x_{\max}^{2}\xi K}{\Delta_{*}^{\frac{\omega}{1+\omega}}}\right)^{1+\omega} \int_{1}^{T} \{s^{*2}(\log d + \log u)\}^{\frac{1+\omega}{2}} u^{-\frac{1+\omega}{2}} \mathbb{1}\{u \geq T_{1}\} du$$

$$\leq \left(\frac{3^{\frac{\omega}{1+\omega}}M_{2}\rho\sigma x_{\max}^{2}\xi K}{\Delta_{*}^{\frac{\omega}{1+\omega}}}\right)^{1+\omega} \{s^{*2}(\log d + \log T)\}^{\frac{1+\omega}{2}} \int_{T_{1}}^{\infty} u^{-\frac{1+\omega}{2}} du$$

$$= 2\left(\frac{3^{\frac{\omega}{1+\omega}}M_{2}\rho\sigma x_{\max}^{2}\xi K}{\Delta_{*}^{\frac{\omega}{1+\omega}}}\right)^{1+\omega} \{s^{*2}(\log d + \log T)\}^{\frac{1+\omega}{2}} \frac{T_{1}^{-\frac{\omega-1}{2}}}{\omega-1}$$

$$\approx \left\{\frac{6\left[M_{2}\rho\sigma x_{\max}^{2}\xi K\right]^{2}}{(\omega-1)}\right\} \left(\frac{s^{*2}\log d}{\Delta_{*}}\right).$$
(31)

Finally, for $\omega = \infty$ it is easy to see that $J_{\omega} = 0$. Hence, the result follows from combing Equation (22), (29), (30) and (31).

Appendix C Posterior contraction result

We briefly describe the probability space under which we are working. Given β , the bandit environment (along with the specific policy π) gives rise to the chosen contexts X_t and rewards \mathbf{r}_t . Here we note that the chosen contexts depend not only on the arm-specific distributions, but also on the sequence of actions taken under π till time t. Let Q_t denote the joint distribution of $(\beta, X_t, \mathbf{r}_t)$ under $\beta \sim \Pi$ (prior) and $(X_t, \mathbf{r}_t) \mid \beta \sim \text{SLCB}_t(\beta, \pi, \mathcal{P}_\epsilon)$ where the latter indicates the joint distribution of the observed contexts and rewards (till time t) under the SLCB environment with policy π , true parameter β and \mathcal{P}_ϵ denotes the noise distribution. We work under a likelihood misspecified regime, which we now discuss.

We assume that the true parameter is β^* and the observations (X_t, \mathbf{r}_t) is generated from $\mathcal{Q}_t^* :=$ SLCB_t $(\beta^*, \pi, \mathcal{P}_{\epsilon}^*)$, where π is the policy given by the TS and \mathcal{P}_{ϵ}^* is an arbitrary sub-Gaussian distribution. We denote by $\mathcal{Q}_{t,\mathbf{r}_t}^{*X}$ the conditional distribution of \mathbf{r}_t given X_t arising from the joint \mathcal{Q}_t^* and $\mathbb{E}_{t,\mathbf{r}_t}^X$ to be the expectation under $\mathcal{Q}_{t,\mathbf{r}_t}^{*X}$. Furthermore, we denote by $\mathcal{Q}_{X_t}^*$ the marginal distribution of X_t under the joint \mathcal{Q}_t^* and \mathbb{E}_{X_t} to be the corresponding expectation.

For modelling purpose, we place prior Π on β and model the likelihood as $(X_t, \mathbf{r}_t) \mid \beta \sim$ SLCB_t $(\beta, \pi, \mathcal{P}_{\epsilon})$, where \mathcal{P}_{ϵ} is taken to be N $(0, \sigma^2)$. This gives rise to a joint distribution \mathcal{Q}_t , as discussed above. Now, let $\Pi_t^X(\cdot \mid \mathbf{r}_t)$ denote the posterior distribution of β given all others, i.e. it is the conditional measure of β given X_t , \mathbf{r}_t arising from the joint \mathcal{Q}_t .

Thus, given a measurable set B, $\Pi_t^X(B|\mathbf{r}_t)$ is a random measure, whose randomness is due to (X_t, r_t) . In the following result, we consider $\mathbb{E}_{t,\mathbf{r}_t}^X \Pi_t^X(B|\mathbf{r}_t)$, which is the expectation of the above under $\mathcal{Q}_{t,\mathbf{r}_t}^{*X}$. Thus, this quantity itself is a random variable, whose randomness is due to X_t . The following result shows that, for B taken as the complement of an appropriate ball around the true β^* , this random variable is small, almost surely $\mathcal{Q}_{X_t}^*$.

Theorem 3. Consider the bandit problem in (1) and let Assumption 2.1-2.3 hold. Also, assume that the prior on parameter β is modeled as (2) with

 $(5/3)\overline{\lambda}_t \le \lambda \le 2\overline{\lambda}_t, \quad \overline{\lambda}_t = x_{\max}\sqrt{2t(\log d + \log t)}$

Then the following is true:

$$\mathbb{E}_{t,\mathbf{r}_{t}}^{X}\left\{\Pi_{t}^{X}\left(\left\|\beta-\beta^{*}\right\|_{1}\geq Q_{4}\sigma x_{\max}K\xi(D_{*}+s^{*})\sqrt{\frac{\log d+\log t}{t}}\mid\mathbf{r}_{t}\right)\mathbb{1}_{\mathcal{G}_{t}\cap\mathcal{T}_{t}(\lambda_{0}(\gamma_{t}))}\right\}\lesssim d^{-s^{*}},$$

almost sure X_t , where Q_4 is a universal constant and $D_* = D_1 s^* + \frac{D_2 x_{\max}^2 s^*}{\phi_{\mathsf{comp}}^2 (S^*; X_t)}$ with $D_1 = 1 + (40/A_4)$ and $D_2 = 128A_4^{-1}$.

Proof. Without loss of generality we assume that $\sigma = 1$ as the bandit reward model can be viewed as

$$(\mathbf{r}_t/\sigma) = X_t(\beta^*/\sigma) + (\epsilon_t/\sigma).$$

In this case $\overline{\lambda}_t = x_{\max} \sqrt{2t(\log d + \log t)} =: \overline{\lambda}$.

Next, define the event

$$\mathcal{T}_0 := \left\{ \max_{j \in [d]} \left| \epsilon_t^\top X_t^{(j)} \right| \le \bar{\lambda} \right\}.$$

By Lemma 3 and condition (21), it follows that fro any measurable set $\mathcal{B} \subset \mathbb{R}^d$,

$$\mathbb{E}_{t,\mathbf{r}_{t}}\Pi_{t}^{X}(\mathcal{B} \mid \mathbf{r}_{t}) \leq \left[\mathbb{E}_{t,\mathbf{r}_{t}}\left\{\Pi_{t}^{X}(\mathcal{B} \mid \mathbf{r}_{t})\mathbb{1}_{\mathcal{T}_{0}}\right\}\right]^{1/2} + \frac{2}{t}.$$

Recall that the errors ϵ_t is modeled as isotropic standard Gaussian independent of the features. Thus, conditioned on the matrix X_t , model likelihood ration takes the following form:

$$\mathcal{L}_{t,\beta,\beta^*}(\mathbf{r}_t) := \exp\left\{-\frac{\|X_t\beta - X_t\beta^*\|_2^2}{2} + (\mathbf{r}_t - X_t\beta^*)^\top (X_t\beta - X_t\beta^*)\right\}.$$

Then by Lemma 2 of Castillo et al. (2015) it follows that

$$\int \mathcal{L}_{t,\beta,\beta^*}(\mathbf{r}_t) \ d\Pi(\beta) \ge \frac{\pi_d(s^*)}{p^{2s^*}} e^{-\lambda \|\beta^*\|_1} e^{-1},$$

where Π is given by (2). The only change that is needed in their proof to run the argument in our case is the following upper bound:

$$\|X\beta\|_2 \le \|\beta\|_1 \|X\| \le c_9 \vartheta^2 \phi_u \log K/(1-3\varepsilon).$$

The last inequality follows from the fact that we are on the event $\mathcal{G}_{t,1}$ by assumption. The rest of the proof follows from the fact that $\lambda \in (5\overline{\lambda}/3, 2\overline{\lambda})$.

Thus by Bayes's formula it follows that

$$\Pi_{t}^{X}(\mathcal{B} \mid \mathbf{r}_{t}) = \frac{\int_{\mathcal{B}} \mathcal{L}_{t,\beta,\beta^{*}}(\mathbf{r}_{t}) \, d\Pi(\beta)}{\int \mathcal{L}_{t,\beta,\beta^{*}}(\mathbf{r}_{t}) \, d\Pi(\beta)}$$

$$\leq \frac{ed^{2s^{*}}}{\pi_{d}(s^{*})} e^{\lambda \|\beta^{*}\|_{1}} \int_{\mathcal{B}} \exp\left\{-\frac{\|X_{t}\beta - X_{t}\beta^{*}\|_{2}^{2}}{2} + (\mathbf{r}_{t} - X_{t}\beta^{*})^{\top} (X_{t}\beta - X_{t}\beta^{*})\right\} \, d\Pi(\beta).$$
(32)

Using Holder's inequality, we see that on T_0 ,

$$(\mathbf{r}_{t} - X_{t}\beta^{*})^{\top}X_{t}(\beta - \beta^{*}) = \epsilon_{t}^{\top}X_{t}(\beta - \beta^{*})$$

$$\leq \|\epsilon_{t}^{\top}X_{t}\|_{\infty} \|\beta - \beta^{*}\|_{1}$$

$$\leq \bar{\lambda} \|\beta - \beta^{*}\|_{1}.$$
(33)

Therefore, on the event \mathcal{T}_0 , the expected value under \mathbb{E}_{β^*} of the integrand on the right hand side of (32) is bounded above by

$$e^{-(1/2)\|X_t(\beta-\beta^*)\|_2^2+\overline{\lambda}\|\beta-\beta^*\|_1}.$$

Thus, we have

$$\Pi_{t}^{X}(\mathcal{B} \mid \mathbf{r}_{t}) \mathbb{1}_{\tau_{0}} \leq \frac{ep^{2s^{*}}}{\pi_{d}(s^{*})} \int_{\mathcal{B}} e^{\lambda \|\beta\|_{1} - (1/2)\|X_{t}(\beta - \beta^{*})\|_{2}^{2} + \overline{\lambda}\|\beta - \beta^{*}\|_{1}} d\Pi(\beta).$$
(34)

Now, by triangle inequality,

$$\begin{split} \lambda \|\beta^*\|_1 + \overline{\lambda} \|\beta - \beta^*\|_1 &= \lambda \|\beta^*_{S^*}\|_1 + \overline{\lambda} \|\beta - \beta^*\|_1 \\ &\leq \lambda \|\beta^*_{S^*} - \beta_{S^*}\|_1 + \lambda \|\beta_{S^*}\|_1 + \overline{\lambda} \|\beta_{S^*} - \beta^*_{S^*}\|_1 + \overline{\lambda} \|\beta_{S^{*c}}\| \\ &= \lambda \|\beta^*_{S^*}\|_1 + \overline{\lambda} \|\beta^*_{S^{*c}}\|_1 + (\lambda + \overline{\lambda}) \|\beta^*_{S^*} - \beta^*_{S^*}\|_1 \\ &= \lambda \|\beta\|_1 + (\overline{\lambda} - \lambda) \|\beta^*_{S^{*c}}\|_1 + (\lambda + \overline{\lambda}) \|\beta^*_{S^*} - \beta^*_{S^*}\|_1. \end{split}$$
(35)

Case 1: Suppose $7 \|\beta_{S^*} - \beta^*_{S^*}\|_1 \le \|\beta_{S^{*c}}\|_1$. Then the following holds:

$$\begin{aligned} (\lambda + \overline{\lambda}) \|\beta_{S^*} - \beta_{S^*}^*\|_1 &= (\overline{\lambda} - 3\lambda/4) \|\beta_{S^*} - \beta_{S^*}^*\|_1 + (7\lambda/4) \|\beta_{S^*} - \beta_{S^*}^*\|_1 \\ &\leq (\overline{\lambda} - 3\lambda/4) \|\beta_{S^*} - \beta_{S^*}^*\|_1 + (\lambda/4) \|\beta_{S^{*c}}\|_1. \end{aligned}$$

Using the above inequality in (35) we get

$$\lambda \|\beta^{*}\|_{1} + \overline{\lambda} \|\beta - \beta^{*}\|_{1} \leq \lambda \|\beta\|_{1} + (\overline{\lambda} - 3\lambda/4) \|\beta_{S^{*c}}\|_{1} + (\overline{\lambda} - 3\lambda/4) \|\beta_{S^{*}} - \beta^{*}_{S^{*}}\|_{1}$$

$$= \lambda \|\beta\|_{1} + (\overline{\lambda} - 3\lambda/4) \|\beta - \beta^{*}\|_{1}$$
(36)

Case 2: Now assume $7 \|\beta_{S^*} - \beta^*_{S^*}\|_1 > \|\beta_{S^{*c}}\|_1$. We again focus on the inequality (35), i.e.,

$$\begin{split} \lambda \|\beta^{*}\|_{1} + \overline{\lambda} \|\beta - \beta^{*}\|_{1} &\leq \lambda \|\beta\|_{1} + (\overline{\lambda} - \lambda) \|\beta_{S^{*c}}\|_{1} + (\lambda + \overline{\lambda}) \|\beta_{S^{*}} - \beta^{*}_{S^{*}}\|_{1} \\ &= \lambda \|\beta\|_{1} + (\overline{\lambda} - \lambda) \|\beta_{S^{*c}}\|_{1} + (\overline{\lambda} - \lambda) \|\beta_{S^{*}} - \beta^{*}_{S^{*}}\|_{1} \\ &+ 2\lambda \|\beta_{S^{*}} - \beta^{*}_{S^{*}}\|_{1} \\ &= \lambda \|\beta\|_{1} + (\overline{\lambda} - \lambda) \|\beta - \beta^{*}\|_{1} + 2\lambda \|\beta_{S^{*}} - \beta^{*}_{S^{*}}\|_{1} \\ &\leq \lambda \|\beta\|_{1} + (\overline{\lambda} - 3\lambda/4) \|\beta - \beta^{*}\|_{1} + 2\lambda \|\beta_{S^{*}} - \beta^{*}_{S^{*}}\|_{1} . \end{split}$$
(37)

Finally, by compatibility condition and Young's inequality we get

$$2\lambda \|\beta_{S^*} - \beta_{S^*}^*\| \le 2\lambda \frac{\|X_t(\beta - \beta^*)\|_2 s^{*1/2}}{t^{1/2}\phi_{\mathsf{comp}}(S^*; X_t)} \le \frac{\|X_t(\beta - \beta^*)\|_2^2}{2} + \frac{2s^*\lambda^2}{t\phi_{\mathsf{comp}}^2(S^*; X_t)}$$

Using the above inequality in (37) we get

$$\lambda \|\beta^*\|_1 + \overline{\lambda} \|\beta - \beta^*\|_1 \le \lambda \|\beta\|_1 + (\overline{\lambda} - 3\lambda/4) \|\beta - \beta^*\|_1 + \frac{\|X_t(\beta - \beta^*)\|_2^2}{2} + \frac{2s^*\lambda^2}{t\phi_{\mathsf{comp}}^2(S^*; X_t)}.$$
(38)

Thus combining (36) and (38) we can conclude

$$\lambda \|\beta^*\|_1 + \overline{\lambda} \|\beta - \beta^*\|_1 \le \lambda \|\beta\|_1 + (\overline{\lambda} - 3\lambda/4) \|\beta - \beta^*\|_1 + \frac{\|X_t(\beta - \beta^*)\|_2^2}{2} + \frac{2s^*\lambda^2}{t\phi_{\mathsf{comp}}^2(S^*; X_t)}.$$
(39)

Using the above result and recalling that $5\overline{\lambda}/3 \le \lambda \le 2\overline{\lambda}$, we see that the right hand side of (34) is bounded by

$$\Pi_t^X(\mathcal{B} \mid \mathbf{r}_t) \mathbb{1}_{\mathcal{T}_0} \le \frac{ed^{2s^*}}{\pi_d(s^*)} e^{\frac{2s^*\lambda^2}{t\phi_{\mathsf{comp}}^2(S^*;X_t)}} \int_{\mathcal{B}} e^{\lambda \|\beta\|_1 - (\lambda/4)\|\beta - \beta^*\|_1} d\Pi(\beta)$$

Controlling sparsity: For the set $\mathcal{B} = \{\beta : |S_{\beta}| > L\}$ and $L \ge s^*$, the above integral can be bounded by

$$\sum_{S:s=|S|>L} \frac{\pi_d(s)}{\binom{d}{s}} \left(\frac{\lambda}{2}\right)^s \int e^{-(\lambda/4)} \left\|\beta_S - \beta_S^*\right\| d\beta_S$$
$$\leq \sum_{s=L+1}^{\infty} \pi_d(s) 4^s$$
$$\leq \pi_d(s^*) 4^{s^*} \left(\frac{4A_2}{d^{A_4}}\right)^{L+1-s^*} \sum_{j=0}^{\infty} \left(\frac{4A_2}{d^{A_4}}\right)^j$$

Thus, we have

$$\begin{split} & \mathbb{E}_{t,\mathbf{r}_{t}}^{X} \left\{ \Pi_{t}^{X}(\mathcal{B} \mid \mathbf{r}_{t}) \mathbb{1}_{\mathcal{T}_{0}} \right\} \\ & \lesssim \exp \left\{ 4s^{*} \log d + \frac{2s^{*}\lambda^{2}}{t\phi_{\mathsf{comp}}^{2}(S^{*};X_{t})} + s^{*} \log 4 - (A_{4}/4)(L+1-s^{*}) \log d \right\} \\ & \leq \exp \left\{ 5s^{*} \log d + \frac{4s^{*}\lambda\overline{\lambda}}{t\phi_{\mathsf{comp}}^{2}(S^{*};X_{t})} - (A_{4}/4)(L+1-s^{*}) \log d \right\} \end{split}$$

Now recall that $\overline{\lambda}^2 = 2tx_{\max}^2(\log d + \log t) \le 4tx_{\max}^2\log d$. Using this inequality in the above display we have

$$\mathbb{E}_{t,\mathbf{r}_t}^X \left\{ \Pi_t^X (\mathcal{B} \mid \mathbf{r}_t) \mathbb{1}_{\mathcal{T}_0} \right\} \lesssim \exp\left\{ 5s^* \log d + \frac{16s^* (\lambda/\overline{\lambda}) x_{\max}^2 \log d}{\phi_{\mathsf{comp}}^2 (S^*; X_t)} - (A_4/4)(L+1-s^*) \log d \right\}.$$

Thus, setting $L \ge 40s^*/A_4 + s^* + \frac{64A_4^{-1}s^*x_{\max}^2}{\phi_{\text{comp}}^2(S^*;X_t)}(\lambda/\overline{\lambda})$ then there exists a universal constant $Q_1 > 0$ such that

$$\mathbb{E}_{t,\mathbf{r}_t}^X \left\{ \Pi_t^X(\mathcal{B} \mid \mathbf{r}_t) \mathbb{1}_{\mathcal{T}_0} \right\} \le Q_1 d^{-s^*}.$$

Control on prediction: Recall that $\lambda/\overline{\lambda} \leq 2$. Using this and the result in the previous part, we can conclude that the posterior distribution is asymptotically supported on the even $\mathcal{B}_1 = \{\beta : |S_\beta| \leq D_*\}$, where $D_* = D_1 s^* + \frac{D_2 x_{\max}^2 s^*}{\phi_{\text{comp}}^2 (S^*; X_t)}$ where $D_1 = 1 + (40/A_4)$ and $D_2 = 128A_4^{-1}$. By combining (32), (33) and the inequality $\lambda \|\beta^*\|_1 \leq 2\overline{\lambda} \|\beta - \beta^*\|_1 + \lambda \|\beta\|_1$ we can conclude that any Borel set \mathcal{B} ,

$$\Pi_t^X(\mathcal{B} \mid \mathbf{r}_t) \mathbb{1}_{\tau_0} \le \frac{ed^{2s^*}}{\pi_d(s^*)} \int_{\mathcal{B}} \exp\left\{-\frac{\|X_t\beta - X_t\beta^*\|_2^2}{2} + 3\overline{\lambda} \|\beta - \beta^*\|_1 + \lambda \|\beta\|_1\right\} d\Pi(\beta).$$

By the definition in (23) we have,

$$(4-1)\overline{\lambda} \|\beta - \beta^*\|_1 \leq \frac{4\overline{\lambda} \|X_t(\beta - \beta^*)\|_2 |S_{\beta - \beta^*}|^{1/2}}{t^{1/2}\overline{\phi}_t(|S_{\beta - \beta^*}|)} - \overline{\lambda} \|\beta - \beta^*\|_1$$

$$\leq \frac{1}{4} \|X_t(\beta - \beta^*)\|_2^2 + \frac{16\overline{\lambda}^2 |S_{\beta - \beta^*}|}{t\overline{\phi}_t(|S_{\beta - \beta^*}|)^2} - \overline{\lambda} \|\beta - \beta^*\|_1.$$
(40)

Since $|S_{\beta-\beta^*}| \leq |S_{\beta}| + s^* \leq D_* + s^*$ on the event \mathcal{B}_1 , it follows that

$$\Pi_{t}^{X}(\mathcal{B} \mid \mathbf{r}_{t}) \mathbb{1}_{\mathcal{T}_{0}} \leq \frac{ed^{2s^{*}}}{\pi_{d}(s^{*})} e^{16\overline{\lambda}^{2}(D_{*}+s^{*})/(t\overline{\psi}_{t}(S^{*})^{2})} \times \int_{\mathcal{B}} e^{-(1/4)\|X_{t}(\beta-\beta^{*})\|_{2}^{2}-\overline{\lambda}\|\beta-\beta^{*}\|_{1}+\lambda\|\beta\|_{1}} d\Pi(\beta).$$

$$(41)$$

Now we set $\mathcal{B} = \mathcal{B}_2 := \{\beta \in \mathcal{B}_1 : \|X_t(\beta - \beta^*)\|_2 > L\}$, where *L* will be chosen shortly. Recall that $\pi_d(s^*) \ge (A_1 p^{-A_3})^{s^*} \pi_p(0)$. It follows that for set \mathcal{B} , the right hand side of (41) is upper bounded by

$$\frac{ed^{2s^*}}{\pi_d(s^*)}e^{16\overline{\lambda}^2(D_*+s^*)/(t\overline{\psi}_t(S^*)^2)}e^{-(1/4)L^2}\int e^{-\overline{\lambda}\|\beta-\beta^*\|_1+\lambda\|\beta\|_1} d\Pi(\beta)$$

$$\lesssim d^{(2+A_3)s^*}A_1^{-s^*}e^{16\overline{\lambda}^2(D_*+s^*)/(t\overline{\psi}_t(S^*)^2)}e^{-(1/4)L^2}\underbrace{\sum_{s=0}^d \pi_d(s^*)2^s}_{O(1)}.$$

Hence by a calculation similar to previous discussion yields that for

$$\frac{1}{4}L^2 = (3+A_3)s^*\log d + \frac{16\overline{\lambda}^2(D_*+s^*)}{t\overline{\psi}_t(S^*)^2} \le Q_2 x_{\max}^2(D_*+s^*)\frac{\log d + \log t}{\overline{\psi}_t(S^*)^2} =: L_*^2,$$

where $Q_2 > 0$ is sufficiently large universal constant, then we have

$$\mathbb{E}_{t,\mathbf{r}_t}^X \left\{ \Pi_t^X (\mathcal{B}_2 \mid \mathbf{r}_t) \mathbb{1}_{\mathcal{T}_0} \right\} \le \frac{1}{d^{s^*}}.$$

Control on estimation: Similar to (40) we have

$$\begin{split} \overline{\lambda} \|\beta - \beta^*\|_1 &\leq \frac{\overline{\lambda} \|X_t(\beta - \beta^*)\|_2 |S_{\beta - \beta^*}|^{1/2}}{t^{1/2} \overline{\phi}_t(|S_{\beta - \beta^*}|)} \\ &\leq \frac{1}{4} \|X_t(\beta - \beta^*)\|_2^2 + \frac{\overline{\lambda}^2 |S_{\beta - \beta^*}|}{t \overline{\phi}_t(|S_{\beta - \beta^*}|)^2}. \end{split}$$

On the event \mathcal{B}_2 , we thus have

$$\overline{\lambda} \left\| \beta - \beta^* \right\|_1 \le Q_3 x_{\max}^2 (D_* + s^*) \frac{\log d + \log t}{\overline{\psi}_t (S^*)^2}$$

Finally on the event $\mathcal{B}_2 \cap \mathcal{G}_t$ we have $\overline{\lambda} = x_{\max}\sqrt{2t(\log d + \log t)}$ and $\overline{\psi}_t(S^*)^2 \gtrsim (K\xi)^{-1}$ and it follows that

$$\|\beta - \beta^*\|_1 \le Q_4 K \xi x_{\max} (D_* + s^*) \sqrt{\frac{\log d + \log t}{t}}.$$

Appendix D Technical lemmas

D.1 Proof of Lemma B.3

As A is symmetric positive definite matrix, by Cholesky decomposition there exists as lower triangular matrix L such that $A = LL^{\top}$. Let $v \in \mathbb{S}_0^{d-1}(s)$. Then there exists a index set J of size s, such that $\sup(v) \subseteq J$. Hence we have $v \in E_J$. Now consider the net $\mathcal{N}_{\varepsilon,J}$ and let u be the nearest point to v in $\mathcal{N}_{\varepsilon,J}$. Thus we have $||v - u||_2 \leq \varepsilon < 1$ and $||v - u||_0 \leq s$. Then we have the following:

$$v^{\top}Av = (v-u)^{\top}A(v-u) + 2(v-u)^{\top}Au + u^{\top}Au$$

$$\geq u^{\top}Au - 3\varepsilon\phi_{\max}(s;A).$$
(42)

The second inequality follows from the following facts:

$$\begin{aligned} \left| (v-u)^{\top} A(v-u) \right| &\leq \|v-u\|_{2}^{2} \phi_{\max}(s;A) \leq \varepsilon \phi_{\max}(s;A), \\ \left| (v-u)^{\top} Au \right| &= \left| (v-u)^{\top} LL^{\top} u \right| \\ &\leq \sqrt{(v-u)^{\top} LL^{\top} (v-u)} \sqrt{u^{\top} LL^{\top} u} \\ &= \sqrt{(v-u)^{\top} A(v-u)} \sqrt{u^{\top} Au} \\ &\leq \varepsilon \phi_{\max}(s;A) \end{aligned}$$

Then the result follows from taking infimum over u and v in both sides.

D.2 Proof of Lemma B.5

The lower bound result is trivial. Hence, we focus on the upper bound part. As A is symmetric positive definite matrix, by Cholesky decomposition there exists as lower triangular matrix L such that $A = LL^{\top}$. Let $v \in \mathbb{S}_0^{d-1}(s)$. Then there exists a index set J of size s, such that $\text{supp}(v) \subseteq J$. Hence we have $v \in E_J$. Now consider the net $\mathcal{N}_{\varepsilon,J}$ and let u be the nearest point to v in $\mathcal{N}_{\varepsilon,J}$. Thus we have $||v - u||_2 \leq \varepsilon < 1/3$ and $||v - u||_0 \leq s$. By a similar argument as in the proof of Lemma B.3, we can conclude that

$$v^{\top}Av \leq 3\varepsilon\phi_{\max}(s;A) + \max_{u\in\mathcal{N}_{\varepsilon}}u^{\top}Au.$$

Thus by taking supremum over v on the left-hand side of the above display we get

$$(1 - 3\varepsilon)\phi_{\max}(s; A) \le \max_{u \in \mathcal{N}_{\varepsilon}} u^{\top} A u \Longleftrightarrow \phi_{\max}(s; A) \le \frac{1}{1 - 3\varepsilon} \max_{u \in \mathcal{N}_{\varepsilon}} u^{\top} A u.$$

Appendix E Pseudo code of VBTS and other tables

Algorithm 2: Variational Bayes Thompson Sampling

 $\begin{array}{l} \operatorname{Set} \mathcal{H}_{0} = \emptyset; \\ \operatorname{\textbf{for}} t = 1, \cdots, T \operatorname{\textbf{do}} \\ & \quad \operatorname{\textbf{if}} t \leq 1 \operatorname{\textbf{then}} \\ & \quad | \quad \operatorname{Choose \ action} a_{t} \ \mathrm{uniformly \ over} \ [K]; \\ & \quad \operatorname{\textbf{end}} \\ & \quad \operatorname{\textbf{else}} \\ & \quad | \quad \operatorname{Compute} \operatorname{VB \ posterior} \tilde{\Pi}_{t-1} \ \mathrm{from} \ \Pi(\cdot \mid \mathcal{H}_{t-1}) \ \mathrm{using} \ \operatorname{CAVI}; \\ & \quad \operatorname{Generate \ sample} \ \tilde{\beta}_{t} \sim \tilde{\Pi}_{t-1}; \\ & \quad \operatorname{Play \ arm:} \ a_{t} = \arg \max_{i \in [K]} x_{i}(t)^{\top} \tilde{\beta}_{t}; \\ & \quad \operatorname{\textbf{end}} \\ & \quad \operatorname{Observe \ reward} r_{a_{t}}(t); \\ & \quad \operatorname{Update} \ \mathcal{H}_{t} \leftarrow \mathcal{H}_{t-1} \cup \{(a_{t}, r_{a_{t}}(t), x_{a_{t}}(t))\}. \end{array}$

Table 3: Time comparison among the competing algorithms.

Туре	Algorithm	Mean time of execution (seconds)		
		Equicorrelated	Auto-regressive	
Non-TS	LinUCB	15.41	16.30	
	DR Lasso	3.12	3.13	
	Lasso-L1	3.57	3.59	
	ESTC	1.01	1.05	
TS	LinTS	1344.39	1346.46	
	BLasso TS	1511.68	1455.53	
	VBTS	29.33	27.65	