

Status of the W-boson mass averaging project

Simone Amoroso, on behalf of the Tevatron/LHC W-mass combination Working Group*

*Deutsches Elektronen-Synchrotron (DESY),
Notkestrasse 85, Hamburg, Germany*

E-mail: simone.amoroso@desy.de

We present the current status of the W-boson mass averaging project, an ongoing effort aimed at combining Tevatron and LHC measurements. Methods are presented to accurately evaluate the effect of PDFs and other modelling variations on existing measurements. Based on this approach, the measurements can be corrected to a common modelling reference and to the same PDFs, and subsequently combined accounting for PDF correlations in a quantitative way. We discuss the combination procedure, and the impact of improvements in the theoretical description of W-boson production and decay.

*41st International Conference on High Energy physics - ICHEP2022
6-13 July, 2022
Bologna, Italy*

*Speaker

1. Introduction

The W-boson mass, m_W , is a fundamental parameter of the Standard Model, and its value one of the most important inputs to global electroweak fits [1]. The W-boson mass has been precisely measured at hadron colliders by the CDF [2] and D0 [3] Collaborations at the Tevatron and by the ATLAS [4] and LHCb [5] Collaborations at the LHC. In this contribution, we report on the current status of the Tevatron/LHC W-boson mass combination effort, aiming to provide a world average of m_W determinations endorsed by the experimental collaborations. Such a combination is particularly motivated in light of the discrepancy between the most recent CDF measurement and the results from the other experiments.

At hadron colliders, measurements of m_W rely on template fits to kinematic peaks of distributions determined from the leptonic decays of the W boson, such as the lepton transverse momentum, p_T^l , or the W-boson transverse mass m_T^W . These final-state distributions carry information about the decaying particle mass, but are also affected by the description of W-boson production and decay, such as the rapidity and transverse momentum distributions and its polarization. Predictions of these observables are obtained using Monte Carlo (MC) event generators with input parton distribution functions (PDFs). With each measurement relying on a different generator and PDF choice, prior to combining the measurements a coherent theoretical treatment is required to estimate uncertainty correlations. These modelling differences further motivate small adjustments to the measured values or uncertainties, an effect which has been neglected in past combinations [6].

Beyond the interest of improving the overall measurement precision on m_W , through this combination effort we establish a methodology to combine present and upcoming measurements of m_W and enable possible updates of the physics modelling (e.g. PDFs, p_T^W) as our theoretical knowledge improves. In addition, as more electroweak measurements become dominated by PDF uncertainties (i.e. m_W and $\sin^2 \theta_{W,\ell}^{\text{eff.}}$), this methodology will permit to consistently correlate them in global electroweak fits.

2. Combination strategy

For a consistent combination of different measurements, their correlations need to be determined. The main source of correlations among the m_W measurements comes from the knowledge of the proton structure, embedded in the PDFs. Due to the different center-of-mass energies and initial states (pp vs $p\bar{p}$) at which the measurements are performed this correlation is however non-trivial. Uncertainties related to the p_T^W distribution, while significant in size, are evaluated independently in each experiment through a detailed analysis of Z-boson production, and are hence assumed to be uncorrelated. Other sources of theoretical uncertainties, such as electroweak corrections, are typically very small, and their detailed correlation knowledge do not affect the combination. Experimental uncertainties are expected to be uncorrelated between experiments.

The combination of m_W results is performed in a two-step procedure. At first, the results have to be translated to a common reference model. The full procedure to correct the published m_W values is decomposed into QCD and PDF effects, and allows to improve existing experimental results to include progress in theoretical prediction and PDF determinations. The published m_W

values $m_W^{\text{published}}$ of each experiment is subject to the correction:

$$m_W^{\text{updated}} = m_W^{\text{published}} - \delta m_W^{\text{QCD}} - \delta m_W^{\text{PDF}}, \quad (1)$$

where δm_W^{QCD} incorporates any eventual corrections to the QCD modelling beyond the originally quoted uncertainties, and δm_W^{PDF} brings all measurements to a common PDF set. The results such obtained are then combined properly across experiments including correlations. The model dependence of the result is evaluated by repeating this procedure for a relevant set of PDF determinations.

3. Detector Emulation

The original m_W measurements have been performed at detector level, and reproducing the analyses and detector simulations used by the experiments is a challenging task. Instead, we make use of a parametrized detector response, tuned on publicly available information (lepton energy/momentum scale and resolution, efficiencies; recoil response) and applied on large Monte Carlo event samples at particle-level, generated with varying QCD modelling assumptions and PDF sets. This approach has been found sufficient for a reliable evaluation of variations in the underlying generator-level distributions, such as PDF uncertainties and extrapolations, and effects in the lepton angular distributions. After event selection it can reproduce published distributions at the % level, with systematic uncertainty on δm_W estimated to be at the level of 1-2 MeV.

4. QCD modelling aspects

We concentrate here on the description of the angular distribution of the decay leptons. After integrating over azimuthal angle, we can express the W -boson cross-section to all orders in the strong coupling as:

$$d\sigma = \sigma_{\text{unpol}} \left[(1 + \cos^2 \theta_{CS}) + \frac{A_0}{2} (1 - 3 \cos^2 \theta_{CS}) + A_4 \cos \theta_{CS} \right], \quad (2)$$

with σ_{unpol} the unpolarised cross-section, $\cos \theta_{CS}$ the polar angle in the Collins-Soper frame [7], and A_0 and A_4 angular coefficients which will in general depend on the boson kinematics. Theoretical predictions at fixed-order for the angular coefficients are currently known up to $O(\alpha_S^3)$ [8, 9], and describe to a very good degree of accuracy the measured coefficients at the LHC [10].

Different theoretical predictions have been used for the various m_W measurements, reflecting the theoretical state of the art at the time of their preparation. For the descriptions of the W -boson production and decay, CDF used events generated with the Resbos-C [11] code at NLO+NLL while the Resbos-CP [12] code at NNLO+NNLL has been used by D0. ATLAS and LHCb relied on parton shower generated events from Pythia8 [13], but reweighted the angular distributions to $O(\alpha_S^2)$ fixed-order predictions from DYNNLO [14].

We generated event samples using the same MC generation chain as the original measurements. In addition, MC samples have been generated using Powheg Z_EW [15], MiNNLOPS [16, 17] and an updated version of Resbos (dubbed here Resbos2) with an improved treatment of spin correlations [18]. This allows to study the impact of computing δm_W^{QCD} and δm_W^{PDF} with predictions at different theoretical accuracies.

We compare in Fig. 1 different predictions for A_0 , A_4 a function of p_T^W . Large differences can be seen between the Resbos-CP version employed by D0 and the modern event generator or fixed-order codes. Similar differences are also observed with respect to the Resbos-C version used by CDF. This is explained as the legacy Resbos codes do not perform a consistent resummation of all the angular coefficients, but only A_4 and σ_{unpol} receive corrections from the resummation. This behavior has been corrected in ResBos2, which largely reproduces the fixed-order behavior.




Figure 1: The angular coefficients A_0 (left plot) and A_4 (right plot) in W-boson production shown as a function of p_T^W in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV. The predictions from the legacy D0 version of Resbos (black) are compared to predictions from an updated version of Resbos (red), fixed-order predictions at $O(\alpha_S)$ from DYNNLO (blue) and MC predictions at $O(\alpha_S^2)$ from MiNNLOPS (magenta).




Figure 2: Relative effect of the generator corrections δm_W^{QCD} on the lepton transverse momentum (left plot) and W-boson transverse mass distributions (right plot), after including detector effect and applying the p_T^W constraint. The correction is obtained by reweighting the angular coefficients to the Resbos 2 predictions (red) and shown with respect to the D0 legacy Resbos-CP code (black) and compared to the directly generated Resbos2 predictions (blue).

We evaluated a correction by reweighting the angular coefficients in the original D0 Resbos-CP sample to the $O(\alpha_S)$ prediction of the Resbos2 sample, and compare it with the shift obtained by directly fitting the Resbos2 distributions. The impact of this correction for the D0 measurement is shown in Tab. 1. When constraining the p_T^W distribution to stay unchanged under the corrections yields shifts of 7.3 ± 1.1 MeV and 8.4 ± 1.0 MeV for the p_T^l and m_T fits. If the p_T^W distribution is instead left unconstrained, shifts of 13.9 ± 1.1 MeV and 10.3 ± 1.0 MeV for the p_T^l and m_T fits

Correction	δm_W^{QCD} [MeV]					
	p_T^W -constrained			No constraint		
	p_T^l	m_T	p_T^y	p_T^l	m_T	p_T^y
Invariant mass	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1
Rapidity	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1
A_0	7.6	10.0	15.8	16.0	12.6	19.5
A_1	-2.4	-1.9	-1.8	-1.2	-1.6	-1.4
A_2	-3.0	-2.6	2.9	-4.2	-3.0	2.3
A_3	2.9	1.6	-0.5	3.5	1.8	-0.2
A_4	2.4	-0.1	-0.5	0.1	-0.7	-1.0
$A_0 - A_4$	7.6	7.0	16.0	14.1	9.1	18.9
Total	7.6	7.0	16.0	14.1	9.1	18.9
RESBos2	7.3±1.1	8.4±1.0	16.6±1.2	13.9±1.1	10.3±1.0	19.8±1.2
Non-closure	-0.3±1.1	1.4±1.0	0.6±1.2	-0.2±1.1	1.2±1.0	0.9±1.2

Table 1: Impact of the angular coefficients reweighting in the D0 RESBos-CP events to those of RESBos2, compared to a direct fit of RESBos1 to RESBos2. Results are shown for the p_T^l , p_T^y and m_T^W distributions with and without a constraint to leave the p_T^W unchanged.

are obtained. As the measurements constrain p_T^W only through the measured p_T^Z distribution, these numbers are to be considered as an upper bound of the effect.

5. Summary

We presented studies towards a first combination of Tevatron and LHC measurements of m_W . Measurement correlations are dominated by PDF uncertainties and are evaluated through a simplified emulation of the detector response. Extensive comparisons against state-of-the-art QCD predictions highlighted the need to correct the Tevatron m_W determinations for the treatment of lepton angular distributions in W-boson decays in the legacy Resbos codes. In the context of the D0 measurement, correcting the spin correlations affects the result at the level of -10 MeV, depending on the distribution used in the fit.

References

- [1] J. Erler and M. Schott, *Electroweak Precision Tests of the Standard Model after the Discovery of the Higgs Boson*, *Prog. Part. Nucl. Phys.* **106** (2019) 68 [1902.05142].
- [2] CDF collaboration, *High-precision measurement of the W boson mass with the CDF II detector*, *Science* **376** (2022) 170.
- [3] D0 collaboration, *Measurement of the W boson mass with the D0 detector*, *Phys. Rev. D* **89** (2014) 012005 [1310.8628].

- [4] ATLAS collaboration, *Measurement of the W-boson mass in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **78** (2018) 110 [1701.07240].
- [5] LHCb collaboration, *Measurement of the W boson mass*, *JHEP* **01** (2022) 036 [2109.01113].
- [6] LHC–TEVATRON W-BOSON MASS COMBINATION WORKING GROUP collaboration, *Towards a combination of LHC and TeVatron W-boson mass measurements*, CERN-LPCC–2022–06, FERMILAB-TM–2779–V.
- [7] J.C. Collins and D.E. Soper, *Angular Distribution of Dileptons in High-Energy Hadron Collisions*, *Phys. Rev. D* **16** (1977) 2219.
- [8] R. Gauld, A. Gehrmann-De Ridder, T. Gehrmann, E.W.N. Glover and A. Huss, *Precise predictions for the angular coefficients in Z-boson production at the LHC*, *JHEP* **11** (2017) 003 [1708.00008].
- [9] M. Pellen, R. Poncelet, A. Popescu and T. Vitos, *Angular coefficients in W + j production at the LHC with high precision*, *Eur. Phys. J. C* **82** (2022) 693 [2204.12394].
- [10] ATLAS collaboration, *Measurement of the angular coefficients in Z-boson events using electron and muon pairs from data taken at $\sqrt{s} = 8$ TeV with the ATLAS detector*, *JHEP* **08** (2016) 159 [1606.00689].
- [11] G.A. Ladinsky and C.P. Yuan, *The Nonperturbative regime in QCD resummation for gauge boson production at hadron colliders*, *Phys. Rev. D* **50** (1994) R4239 [hep-ph/9311341].
- [12] C. Balazs and C.P. Yuan, *Soft gluon effects on lepton pairs at hadron colliders*, *Phys. Rev. D* **56** (1997) 5558 [hep-ph/9704258].
- [13] C. Bierlich et al., *A comprehensive guide to the physics and usage of PYTHIA 8.3*, 2203.11601.
- [14] S. Catani, L. Cieri, G. Ferrera, D. de Florian and M. Grazzini, *Vector boson production at hadron colliders: a fully exclusive QCD calculation at NNLO*, *Phys. Rev. Lett.* **103** (2009) 082001 [0903.2120].
- [15] L. Barze, G. Montagna, P. Nason, O. Nicrosini, F. Piccinini and A. Vicini, *Neutral current Drell-Yan with combined QCD and electroweak corrections in the POWHEG BOX*, *Eur. Phys. J. C* **73** (2013) 2474 [1302.4606].
- [16] P.F. Monni, P. Nason, E. Re, M. Wiesemann and G. Zanderighi, *MiNNLO_{PS}: a new method to match NNLO QCD to parton showers*, *JHEP* **05** (2020) 143 [1908.06987].
- [17] P.F. Monni, E. Re and M. Wiesemann, *MiNNLO_{PS}: optimizing $2 \rightarrow 1$ hadronic processes*, *Eur. Phys. J. C* **80** (2020) 1075 [2006.04133].
- [18] J. Isaacson, Y. Fu and C.P. Yuan, *ResBos2 and the CDF W Mass Measurement*, 2205.02788.