EXTREME NATURE OF FOUR BLUE-EXCESS DUST-OBSCURED GALAXIES REVEALED BY OPTICAL SPECTROSCOPY

Akatoki Noboriguchi, 1,2,3 Tohru Nagao, 4 Yoshiki Toba, 5,6,7,4 Kohei Ichikawa, 2 Masaru Kajisawa, 3 Nanako Kato, 3 Toshihiro Kawaguchi, 8 Hideo Matsuhara, 9,10 Yoshiki Matsuoka, 4 Kyoko Onishi, 11 Masafusa Onoue, 12,13,14 Nozomu Tamada, 3 Koki Terao, 15,16 Yuichi Terashima, 3,4 Yoshihiro Ueda, 6 and Takuji Yamashita 5

ABSTRACT

We report optical spectroscopic observations of four blue-excess dust-obscured galaxies (BluDOGs) identified by Subaru Hyper Suprime-Cam. BluDOGs are a sub-class of dust-obscured galaxies (DOGs, defined with the extremely red color $(i - [22])_{AB} \ge 7.0$; Toba et al. 2015), showing a significant flux excess in the optical g- and r-bands over the power-law fits to the fluxes at the longer wavelengths. Noboriguchi et al. (2019) has suggested that BluDOGs may correspond to the blowing-out phase involved in a gas-rich major merger scenario. However the detailed properties of BluDOGs are not understood because of the lack of spectroscopic information. In this work, we carry out deep optical spectroscopic observations of four BluDOGs using Subaru/FOCAS and VLT/FORS2. The obtained spectra show broad emission lines with extremely large equivalent widths, and a blue wing in the C IV line profile. The redshifts are between 2.2 and 3.3. The averaged rest-frame equivalent widths of the C IV lines are 160 ± 33 Å, \sim 7 times higher than the average of a typical type-1 quasar. The FWHMs of their velocity profiles are between 1990 and 4470 km s⁻¹, and their asymmetric parameters are 0.05 and 0.25. Such strong C IV lines significantly affect the broad-band magnitudes, which is partly the origin of the blue excess seen in the spectral energy distribution of BluDOGs. Their estimated supermassive black hole masses are $1.1 \times 10^8 < M_{\rm BH}/M_{\odot} < 5.5 \times 10^8$. The inferred Eddington ratios of the BluDOGs

Corresponding author: Akatoki Noboriguchi akatoki@shinshu-u.ac.jp

¹School of General Education, Shinshu University, 3-1-1 Asahi, Matsumoto, Nagano 390-8621, Japan

² Frontier Research Institute for Interdisciplinary Sciences, Tohoku University, 6-3 Aramaki, Aoba-ku, Sendai, Miyagi 980-8578, Japan

³ Graduate School of Science and Engineering, Ehime University, 2-5 Bunkyo-cho, Matsuyama, Ehime 790-8577, Japan

⁴Research Center for Space and Cosmic Evolution, Ehime University, 2-5 Bunkyo-cho, Matsuyama, Ehime 790-8577, Japan

⁵ National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan

⁶ Department of Astronomy, Kyoto University, Kitashirakawa-Oiwake-cho, Kyoto 606-8502, Japan

⁷ Academia Sinica Institute of Astronomy and Astrophysics, 11F of Astronomy-Mathematics Building, AS/NTU, No.1, Section 4, Roosevelt Road, Taipei 10617, Taiwan

⁸ Department of Economics, Management and Information Science Onomichi City University, Hisayamada 1600-2, Onomichi, Hiroshima 722-8506, Japan

⁹ Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 3-1-1 Yoshinodai, Chuo-ku, Sagamihara, Kanagawa 252-5210, Japan

¹⁰Department of Space and Astronautical Science, The Graduate University for Advanced Studies, SOKENDAI, 3-1-1 Yoshinodai, Chuo-ku, Sagamihara, Kanagawa 252-5210, Japan

¹¹ Department of Space, Earth and Environment, Chalmers University of Technology, Onsala Observatory, SE-439 92 Onsala, Sweden

¹²Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU, WPI), The University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa, Chiba 277-8583, Japan

¹³Kavli Institute for Astronomy and Astrophysics, Peking University, Beijing 100871, China

¹⁴ Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117, Heidelberg, Germany

¹⁵Subaru Telescope, National Astronomical Observatory of Japan, 650 North A'ohoku Place, Hilo, HI 96720, USA

¹⁶ Astronomical Institute, Tohoku University, 6-3 Aramaki, Aoba-ku, Sendai, Miyagi 980-8578, Japan

are higher than 1 (1.1 < $\lambda_{\rm Edd}$ < 3.8), suggesting that the BluDOGs are in a rapidly evolving phase of supermassive black holes.

 $\it Keywords:$ galaxies: active — galaxies: evolution — infrared: galaxies — quasars: general — techniques: spectroscopic

1. INTRODUCTION

In the last two decades, observations of low-redshift galaxies have revealed tight correlations between the mass of supermassive black holes (SMBHs) and the host galaxy properties such as bulge mass (e.g., Magorrian et al. 1998; Ferrarese & Merritt 2000; Gebhardt et 2000; Tremaine et al. 2002; Marconi & Hunt 2003; Kormendy & Ho 2013; Ding et al. 2020). Such scaling relations suggest the so-called co-evolution between galaxies and SMBHs. It has been argued that a major merger of gas-rich galaxies triggers active starforming activity and subsequent mass accretion onto SMBHs (e.g., Sanders et al. 1988; Hopkins et al. 2008; Treister et al. 2012; Goulding et al. 2018). In this scenario, the merging two galaxies first evolve into a dusty star-forming (SF) galaxy. Then it evolves into a dusty active galactic nucleus (AGN) as gas accretion to the nuclear region triggers the activity of SMBHs. Finally, a dusty AGN evolves into an optically-thin quasar after the surrounding dust is blown out by the powerful AGN outflow. The most active period of such SF and AGN activity is generally heavily obscured by dust, which prevents us from investigating these phases observationally.

By combining optical, near-infrared (NIR), and mid-infrared (MIR) catalogs obtained from the Subaru Hyper Suprime-Cam (HSC; Miyazaki et al. 2018)-Subaru Strategic Program (SSP; Aihara et al. 2018), the VISTA Kilo-degree Infrared Galaxy survey (VIKING; Arnaboldi et al. 2007), and the Wide-field Infrared Survey Explorer (WISE; Wright et al. 2010) all-sky survey (ALLWISE; Cutri 2014), Toba et al. (2015, 2017b) and Noboriguchi et al. (2019) selected dusty SF galaxies and/or powerful AGNs as dust-obscured galaxies (DOGs: Dev et al. 2008; Fiore et al. 2008; Bussmann et al. 2009; Desai et al. 2009; Bussmann et al. 2011). DOGs are defined with a very red optical-MIR color $((i - [22])_{AB} \ge 7.0$; Toba et al. 2015). DOGs represent a transition phase from a gas-rich major merger to an optically-thin quasar in the gas-rich major merger scenario (Dev et al. 2008), suggesting that some DOGs are expected to have buried AGNs. Recently, eight blueexcess DOGs (BluDOGs; Noboriguchi et al. 2019) were discovered from the HSC-selected DOGs based on their optical spectral slopes (i.e., $\alpha_{\rm opt} < 0.4$, where $\alpha_{\rm opt}$ is the observed-frame optical spectral index for the HSC g-, r-, i-, z-, and y-bands in the power-law fit, $f_{\nu} \propto \lambda^{\alpha_{\rm opt}}$), and are a very rare population (eight BluDOGs out of 571 HSC-selected DOGs). Noboriguchi et al. (2019) suggested that the BluDOGs with such blue excess may be in the blowing-out phase involved in the gas-rich major merger scenario. However, the detailed properties of BluDOGs are not well understood because of the lack of spectroscopic information. Spectroscopic observations will give us accurate redshifts, and thus reliable AGN luminosities as a measure of the accretion rates, as well as the SMBH masses.

al. Another interesting population that may represent the transition phase between optically-thick AGNs and optically-thin quasars is extremely red quasars (ERQs; e.g., Ross et al. 2015; Hamann et al. 2017; Perrotta et al. 2019; Villar Martín et al. 2020). ERQs were identified by combining the optical photometric data of Sloan Digital Sky Survey (SDSS; York et al. 2000), the optical spectroscopic data from SDSS-III (Eisenstein et al. 2011) Baryon Oscillation Spectroscopic Survey (BOSS; Dawson et al. 2013), and the MIR photometric data of WISE catalog. are also defined with very red optical to MIR colors $(F_{\nu}(24\mu\text{m})/F_{\nu}(R) \geq 1000)$, and their spectra show broad emission lines with extremely large equivalent widths (Ross et al. 2015; Hamann et al. 2017). Hamann et al. (2017) refined the definition of ERQs as $(i - [12])_{AB} > 4.6^{1}$, and reported notable blue-wing features in their C IV profiles, which suggests the presence of powerful outflow. However, the ERQ sample is limited to optically bright objects since their selection requires SDSS spectra. Detailed studies of opticallyfaint populations in the transition phase between the optically-thick and optically-thin stages are required to understand the whole scenario of the merger-driven evolution of SMBHs. Therefore, it is important to execute the spectroscopic observations for BluDOGs and to research their spectroscopic properties.

In this work, we present the results of spectroscopic observations and subsequent analyses of four BluDOGs. This paper is organized as follows. We describe sample selection of our targets and observations in Section 2. In Section 3, we present properties of the detected emission lines, the estimated dust extinctions, bolometric luminosities of an AGN ($L_{\rm bol}^{\rm AGN}$), and SMBH masses ($M_{\rm BH}$). The discussion on the large equivalent widths of the C IV emission, their SMBH mass, and Eddington ratios is given in Section 4. Then we give a brief summary in Section 5. Throughout this paper, the adopted cosmology is a flat universe with $H_0=70~{\rm km~s^{-1}~Mpc^{-1}}$, $\Omega_M=0.3$, and $\Omega_\Lambda=0.7$. Unless otherwise noted, all magnitudes refer to the AB system.

2. SAMPLE AND THE DATA

 1 All of the BluDOGs also satisfy the criterion of the ERQ (see Table 1).

HSC r-band HSC i-band Name WISE W3-band WISE W4-band [AB mag] [AB mag] [AB mag] [AB mag] HSC J090705.64+020955.8 (HSC J0907) 22.56 ± 0.01 22.59 ± 0.01 16.06 ± 0.13 14.89 ± 0.34 HSC J120200.84-011846.4 (HSC J1202) 20.92 ± 0.00 20.87 ± 0.00 14.47 ± 0.04 $13.46 {\pm} 0.10$ HSC J120728.71-005808.4 (HSC J1207) 22.12 ± 0.01 22.31 ± 0.01 16.28 ± 0.16 15.01 ± 0.36 $17.24^{\rm a}$ HSC J141435.21+003547.4 23.32 ± 0.02 23.11 ± 0.02 $15.33 {\pm} 0.33$ $HSC\ J143727.40-011726.5$ 23.17 ± 0.02 23.10 ± 0.01 16.94 ± 0.23 15.37 ± 0.31 HSC J144333.84-000830.3 (HSC J1443) 22.34 ± 0.01 $22.24{\pm}0.01$ 16.14 ± 0.10 $15.04 {\pm} 0.23$ HSC J144813.65+002244.3 15.37 ± 0.34 23.55 ± 0.02 23.43 ± 0.02 16.83 ± 0.16

 23.95 ± 0.03

 23.74 ± 0.02

 17.15 ± 0.22

 $15.47 {\pm} 0.36$

Table 1. Photometric data of BluDOGs

HSC J144900.84+002350.2

https://wise2.ipac.caltech.edu/docs/release/allwise/expsup/sec2_1a.html

Name	Exp. time [s]	Date	Standard star	Instrument
HSC J0907	900×2	2019 October 8	G191-B2B	FOCAS (Subaru)
	600×1			
HSC J1202	900×6	2019 February 27	LTT 6248	FORS2 (VLT)
HSC J1207	900×12	2019 March 1, 2, 6	LTT 4816	FORS2 (VLT)
HSC J1443	900×12	2019 March 7 8	LTT 4816 EG 274	FORS2 (VLT)

Table 2. Observation log

2.1. Sample selection

In Noboriguchi et al. (2019), 571 DOGs were selected by combining ~105 deg² imaging data obtained from the survey of HSC-SSP² (g, r, i, z, and y), VIKING (Z, Y, J, H, and Ks), and ALLWISE (W1, W2, W3, and W4). The eight BluDOGs were defined among the DOG sample with the smallest observed-frame optical slope $(\alpha_{\text{opt}} < 0.4, \text{ where } \alpha_{\text{opt}} \text{ is the observed-frame optical spectral index of the power-law fits to the HSC } g, r, i, z, \text{ and } y\text{-band fluxes}, f_{\nu} \propto \lambda^{\alpha_{\text{opt}}}$). We selected the four brightest BluDOGs $(r_{\text{AB}} < 23: \text{ see Table 1})$ as the targets of our spectroscopic observations presented in this paper.

2.2. Spectroscopic observations and data reductions

We executed the observations by using Faint Object Camera and Spectrograph (FOCAS; Kashikawa et al. 2002) installed on the Subaru Telescope of National Astronomical Observatory of Japan, and FORS2 (Appenzeller et al. 1998) installed on Very Large Telescope (VLT-UT1) of European Southern Observatory (ESO). We present the observation log in Table 2.

2.2.1. Subaru FOCAS

By using FOCAS, we observed HSC J090705.64+020955.8 (hereafter J0907) on October 8th in 2019, with airmass ~ 1.76 and seeing ~ 0.5 arcsec. We used the 300B grism and the SY47 filter to cover $\lambda_{\rm obs} \sim 4700-9200$ Å, with the resultant spectral resolution of $R \sim 800$ for the used 0".8-width slit. To reduce the obtained data, we performed bias correction, flat fielding with dome flat, removal of cosmic-rays, spectral extraction, sky subtraction, wavelength calibration, and flux calibration

^aThe magnitude is a 95% confidence upper limit.

 $^{^2}$ We utilize the photometric data of S16A HSC-SSP, which was released internally within the HSC survey team and is based on data obtained from 2014 March to 2016 April.

with a standard star (G191-B2B) using the Python packages of Astropy and Numpy. For removing cosmic-rays, we utilized Astro-SCRAPPY (McCully & Tewes 2019). Astro-SCRAPPY is based on the algorithm of L.A.Cosmic, which removes cosmic-rays based on a variation of Laplacian edge detection (van Dokkum 2001). The final spectrum is an inverse-variance weighted mean of the individual shots, corrected for the Galactic extinction (Schlegel et al. 1998).

2.2.2. *VLT FORS2*

By using FORS2, we observed HSC J120200.84-011846.4, HSC J120728.71-005808.4, and HSC J144333.84-000830.3 (hereafter J1202, J1207, and J1443, respectively) between February 27th and March 8th, 2019. used the GRISM_600RI+19 and the GG435 filter to cover $\lambda_{\rm obs} \sim 5200-8000$ Å, which results in the spectral resolution of $R \sim 1500$ with 0".7-width slit. The typical airmasses of the observations for J1202, J1207, and J1443 were 1.17, 1.24, and 1.12, and the typical seeing sizes were ~ 1.0 , 0.5, and 0.5 arcsec, respectively. For the data reduction, we utilized the Recipe flexible execution workbench (Reflex; Freudling et al. 2013) software. Reflex performed bias correction, flat fielding with dome flat, sky subtraction, removing cosmic-rays, spectral extraction, wavelength calibration, and flux calibration with a standard star (LTT 6248, LTT 4816, and EG 274). The final spectrum of each target is the inverse-variance weighted mean of the individual shots, corrected for the Galactic extinction.

2.2.3. Spectrophotometric re-calibration

We re-calibrated the reduced spectra to match the HSC photometry, in order to correct for the effects of the slit loss of the flux, systematic errors in the photometric and spectroscopic calibrations, and any other possible systematic errors. In our observations, the spectra cover the wavelength range of the HSC r-band. We calculate the calibration factor, $f_{\rm photo_calib} = F_{\rm photo_r}/F_{\rm spec_r}$, where $F_{\rm photo_r}$ and $F_{\rm spec_r}$ are the photometric and spectroscopic fluxes in the HSC r-band. The derived calibration factors of J0907, J1202, J1207, and J1443 are 0.97, 1.50, 1.40, and 1.36, respectively. We multiply the spectra with the derived calibration factors.

3. RESULTS

3.1. Emission-line measurements

Figure 1 shows the reduced spectra of the four Blu-DOGs. In order to measure the emission-line properties, we divide emission lines into six groups as follows; (1) Ly α 1216, N v λ 1240, and Si II λ 1263, (2) Si IV λ 1397

and O IV] $\lambda 1402$, (3) He II $\lambda 1640$ and O III] $\lambda 1663$, and (4) Al III λ 1857, Si III] λ 1892, and C III] λ 1909, (5) C IV 1549, and (6) Mg II. We fit the emission lines in each group simultaneously, with a linear continuum model subtracted from the observed spectrum. We adopt a single-Gaussian profile for Ly α , N V, Si II, Si IV, O IV], O III], Al III, Si III and Mg II. The C III] of J1202 is fitted with a single-Gaussian profile, while those of J0907 and J1207 are fitted with a double-Gaussian profile. For the fit around the Si IV and O IV of J1202, we add an additional Gaussian profile to reproduce the observed broad component. We fit C IV and He II with double-Gaussian profiles, and denote the blue and red components with the suffixes of "_B" and "_R", respectively. Additionally, we fit the doublet absorption lines observed around the C IV emission lines of J1202 and J1443. The C IV absorption lines observed at $\lambda_{\rm obs} = 5916.8$ Å and 5926.6Å for J1202 and those at $\lambda_{\rm obs} = 6678.8$ Å and 6689.9 Å for J1443 are fitted using the Voigt profile, respectively. The doublet absorption line ratio is fixed as 2:1 (Feibelman 1983). The best values and standard deviations for emission and absorption lines parameters are estimated by using scipy.optimize.curve_fit³, while we calculate full width at half maximum (FWHM) of emission lines with double Gaussian by using a Monte Carlo method. For this Monte Carlo simulation, we created 10,000 mock spectra using the noise arrays of the observed spectra, and calculate the mean and standard deviation of the line properties. The results for emission lines are listed in Tables 3–6. For absorption lines, the observed-frame equivalent widths and redshifts of the doublet absorption lines on the J1202 C IV emission line are 47.7 ± 19.6 Å, 23.6 ± 9.7 Å, and 2.822, respectively, while the observed-frame equivalent widths and redshift of the doublet absorption lines on J1443 C IV emission line are 14.0 ± 6.3 Å, 6.94 ± 3.14 Å, and 3.314, respectively. Therefore, the co-moving distance between J1202 and its C IV absorber is 8.73 Mpc, while that between J1443 and its C IV absorber is 2.79 Mpc.

The flux ratios of N v/Ly α and N v/C IV for J1443 are 3.9 and 1.8, respectively, whereas the values for the typical quasar (Vanden Berk et al. 2001) are 0.02 and 0.10. One possible reason of these unusual flux ratios in J1443 is the presence of absorption lines, which absorb most of the Ly α and the C IV fluxes around the peak. The unusual flux ratios cannot be explained by the dust reddening, given too small wavelength separations among emission lines of Ly α , N V, and C IV.

³ https://docs.scipy.org/doc/scipy/reference/




Figure 1. The reduced spectra of the BluDOGs. The spectra are for J0907, J1202, J1207 and J1443 from the top to bottom. Detected lines are marked by arrows and labels.

Figure 2 showcases the best-fit models to the C IV emission lines in the four BluDOGs. We adopt the C IV redshift taking C IV_R + C IV_B into account as the systemic redshift of the targets. The determined systemic redshifts of J0907, J1202, J1207, and J1443 are 2.258 ± 0.002 , 2.830 ± 0.002 , 2.511 ± 0.001 , and 3.317 ± 0.006 , respectively.

3.2. Emission-line contributions to the HSC g- and r-band magnitudes

Figure 1 suggests the very large equivalent width (EW) of the emission lines. The average rest-frame EW (REW) of the C IV line of the four BluDOGs is 160 ± 33 Å, ~ 7 times higher than the average of SDSS type-1 quasars (23.8 \pm 0.1 Å; Vanden Berk et al. 2001). Here we investigate the effect of the large REWs on the HSC g- and r-band magnitudes.

First, we calculate the expected magnitudes at the gand r-bands from an extrapolation of the power-law fit
to the longer wavelength bands (i, z, y, Z, Y, J, H, Ks, W1, W2, W3, and W4). Figure 3 clearly shows
that the observed g- and r-band magnitudes exceed the
extrapolation of the power-law fit. The excesses of the gband magnitudes for J0907, J1202, J1207, and J1443 are

1.27, 1.13, 1.48, and 0.88 mag, and those for the r-band excesses are 0.47, 0.44, 0.65, and 0.40 mag, respectively.

Furthermore, we estimate the effect of the strong emission lines, based on their observed-frame EWs and the band widths (BW) of the HSC g and r-bands. The BWs of the HSC g and r-bands are 1468 and 1508 Å (Kawanomoto et al. 2018), respectively. By taking all of the emission lines (Tables 3-6) covered by the HSC g-band (4000–5500Å) and r-band (5500–7000Å) into account, the total observed-frame EWs for J0907, J1202, J1207, and J1443 in the g-band are 604, 258, 608, and 1380 Å respectively, while those in the r-band are 129, 836, 329, and 721 Å (Table 7). Note that the total observed-frame EWs in the q-band are lower limits, because our optical spectra do not cover the entire wavelength range of the band (Section 2.2) and thus some emission lines are not taken into account in the derived total observed-frame EWs. Especially Ly α , the strongest emission line in the rest-frame UV spectrum of typical AGNs, is not covered in our spectra of J0907, J1202, and J1207, thus the total observed-frame EWs




Figure 2. Spectral fits to the C IV emission lines of the BluDOGs. The top left, top right, bottom left, and bottom right show the C IV emission lines of J0907, J1202, J1207, and J1443, respectively. The green, magenta, red, blue, and black lines represent the observed spectrum, linear fit to the continuum emission, two Gaussians for the red and blue components, and best-fit model, respectively. The orange line on the J1202 and J1443 panels represents the C IV doublet absorption line. The horizontal black bars denote the wavelength range used to fit the continuum emission. In each panel, the lower part presents the residual of the best-fit, with the same flux scale as in the upper part.

for these 3 objects are largely underestimated⁴. Since the magnitude excess by the emission lines is given by Δ mag = 2.5 log (1 + EW/BW), the estimated effects in the g-band for J0907, J1202, J1207, and J1443 are 0.37, 0.18, 0.38, and 0.72 mag, respectively. Similarly, the estimated effects of emission lines to the r-band magnitudes are 0.09, 0.48, 0.21, and 0.42 mag, respectively (see Table 7 for a summary). We will discuss the implication from these estimates in Section 4.2.

3.3. Estimating the dust extinction

We need to estimate dust extinction, E(B-V) of AGN radiation, and $L_{\rm bol}^{\rm AGN}$ to calculate the SMBH mass and Eddington ratio. Since Balmer decrement or other spectral measures of E(B-V) is not available, we perform the SED fitting to the broad-band photometry to estimate the E(B-V) and $L_{\rm bol}^{\rm AGN}$. In this work, we utilize the new version of Code Investigating GAlaxy

Emission (CIGALE; Burgarella et al. 2005; Noll et al. 2009; Boquien et al. 2019) called X-CIGALE (Yang et al. 2020), to perform the SED fit in a self-consistent framework by considering an energy balance between the UV/optical absorption and IR emission. X-CIGALE generates the best-fit model including the stellar, AGN, and SF components that fits the photometric data in the rest-frame UV to far-infrared (FIR) bands. We utilize the Herschel Space Observatory (Pilbratt et al. 2010) Astrophysical Terahertz Large Area Survey (H-ATLAS; Eales et al. 2010; Valiante et al. 2016; Bourne et al. 2016) data observed with Photodetector Array Camera and Spectrometer (PACS; Poglitsch et al. 2010) at 100 and 160 μm and with Spectral and Photometric Imaging REceiver (SPIRE; Griffin et al. 2010) at 250, 350, and 500 μ m in the FIR, in addition to optical, NIR, and MIR data obtained by Subaru HSC, VISTA, and WISE. The 1σ limiting fluxes at 100, 160, 250, 350, and $500 \ \mu \text{m}$ are 44, 49, 7.4, 9.4, and 10.2 mJy, respectively (Valiante et al. 2016).

To search for the H-ATLAS counterpart of the four BluDOGs, we adopt a search radius of 10 arcsec by fol-

⁴ The Ly α line of J0907 is at the shorter edge of the HSC g-band coverage but the flux contribution to the g-band magnitude is likely to be significant owing to its broad nature.

Table 3. The detected lines of J0907

Line name	$\lambda_{\mathrm{rest}} \ [\mathrm{\mathring{A}}]$	$z_{ m line}$	$FWHM_{\text{rest}}$ [Å]	$F_{\rm line} [{\rm erg~s^{-1}~cm^{-2}}]$	EW_{rest} [Å]	$v_{\rm width} \; [{\rm km \; s^{-1}}]$
(1)	(2)	(3)	(4)	(5)	(6)	(7)
C IV_R	1549.5	2.258 ± 0.001	12.8±0.7	$(1.10\pm0.07)E-15$	118±10	2470±130
C IV_B	1549.5	$2.227{\pm}0.004$	13.8 ± 2.5	$(2.87\pm0.60)E-16$	$31.5 {\pm} 6.7$	2670 ± 490
C iv-R + C iv-B	1549.5	$2.258 {\pm} 0.002$	$15.2 {\pm} 0.8$	$(1.39\pm0.09)E-15$	$148 {\pm} 12$	$2940{\pm}150$
He II_R	1640.4	2.260 ± 0.002	$3.63{\pm}2.32$	$(3.50\pm2.36)E-17$	4.43 ± 3.01	663 ± 425
He II_B	1640.4	$2.235{\pm}0.005$	$38.1 {\pm} 5.9$	$(2.13\pm0.48)E-16$	$26.8 {\pm} 6.3$	6960 ± 1080
He II_R + He II_B	1640.4	$2.260{\pm}0.002$	$5.54{\pm}1.29$	$(2.48\pm0.53)E-16$	31.4 ± 7.0	1010 ± 240
O III]	1663.5	$2.264{\pm}0.001$	$3.29{\pm}1.69$	$(4.49\pm1.48)E-17$	5.84 ± 1.98	593 ± 305
Si III	1892.0	$2.258 {\pm} 0.002$	4.03 ± 4.16	$(2.97\pm2.55)E-17$	3.56 ± 3.06	639 ± 659
C III]_R	1908.7	$2.261 {\pm} 0.005$	30.3 ± 5.3	$(2.34\pm0.66)E-16$	$28.2 {\pm} 8.1$	4760 ± 830
C III]_R	1908.7	$2.258{\pm}0.002$	6.19 ± 3.04	$(6.50\pm3.21)E-17$	7.83 ± 3.89	971 ± 477
C III R + C III B	1908.7	$2.258 {\pm} 0.003$	$11.6 {\pm} 1.7$	$(2.99\pm0.73)E-16$	36.0 ± 9.0	1830 ± 260
Mg II	2799.1	2.259 ± 0.001	17.6 ± 1.6	$(2.44\pm0.29)E-16$	49.0 ± 7.3	1890 ± 170

NOTE—Column (1): Line name, (2): Rest-frame wavelength of the line, (3): Line redshift, (4): Rest-frame FWHM, (5): Line flux, (6): Rest-frame EW, (7): Velocity width after the correction for the instrumental broadening.

Table 4. The detected lines of J1202

Line name	$\lambda_{\mathrm{rest}} [\mathrm{\AA}]$	$z_{ m line}$	$FWHM_{\rm rest}$ [Å]	$F_{\rm line} [{\rm erg \ s^{-1} \ cm^{-2}}]$	EW_{rest} [Å]	$v_{\rm width} \ [{\rm km \ s^{-1}}]$
(1)	(2)	(3)	(4)	(5)	(6)	(7)
Si IV	1393.8	2.831 ± 0.001	$4.27{\pm}0.73$	$(1.56\pm0.33)E-16$	5.55 ± 1.17	919±157
Broad component ^a	_	_	_	$(1.23\pm0.13)E-15$	_	4970 ± 170
O IV]	1399.9	$2.842 {\pm} 0.001$	$9.90 {\pm} 0.87$	$(5.13\pm0.68)E-16$	18.1 ± 2.4	2120 ± 190
C IV_R	1549.5	2.831 ± 0.001	15.0 ± 0.4	$(5.46\pm0.26)E-15$	177±9	2900 ± 70
C IV_B	1549.5	2.793 ± 0.002	13.0 ± 1.0	$(7.98\pm1.03)E-16$	27.7 ± 3.6	2510 ± 190
$C \text{ iv_R} + C \text{ iv_B}$	1549.5	$2.830 {\pm} 0.002$	16.0 ± 0.5	$(6.26\pm0.28)E-15$	203 ± 10	3100 ± 90
He II_R	1640.4	2.838 ± 0.002	11.0 ± 1.7	$(1.68\pm0.47)E-16$	5.06 ± 1.43	2010 ± 310
Не п_В	1640.4	$2.806 {\pm} 0.005$	21.1 ± 3.4	$(3.16\pm0.55)E-16$	$9.31{\pm}1.64$	3870 ± 620
He II_R + He II_B	1640.4	$2.834 {\pm} 0.004$	$25.5 {\pm} 2.5$	$(4.84\pm0.73)E-16$	$14.5 {\pm} 2.2$	4650 ± 450
О ІІІ]	1663.5	$2.852 {\pm} 0.003$	$7.57{\pm}2.94$	$(2.06\pm1.07)E-17$	0.661 ± 0.343	1360 ± 530
Al III	1858.8	2.839 ± 0.002	21.5 ± 2.4	$(3.05\pm0.41)E-16$	$10.5 {\pm} 1.4$	3470 ± 390
Si III	1892.0	2.819 ± 0.004	$13.6 {\pm} 5.5$	$(9.40\pm4.81)E-17$	$3.19{\pm}1.64$	2160 ± 870
C 111]	1908.7	2.835 ± 0.002	30.6 ± 1.9	$(1.01\pm0.07)E-15$	33.5 ± 2.4	4810 ± 300

 $\mbox{\sc Note}\mbox{\sc See}$ Table 3 for the description of each column.

^aThe observed-frame wavelength, continuum flux, and FWHM of the broad component are 5349.5 ± 2.0 Å, (7.36 ± 0.12) E-18 erg s⁻¹ cm⁻² Å⁻¹, and 88.9 ± 2.9 Å, respectively. See Section 3.1 for details.

Table 5. The detected lines of J1207

Line name	$\lambda_{\mathrm{rest}} [\mathrm{\AA}]$	$z_{ m line}$	$FWHM_{\text{rest}}$ [Å]	$F_{\rm line} [{\rm erg \ s^{-1} \ cm^{-2}}]$	EW_{rest} [Å]	$v_{\rm width} \ [{\rm km \ s^{-1}}]$
(1)	(2)	(3)	(4)	(5)	(6)	(7)
C IV_R	1549.5	2.512 ± 0.001	7.87 ± 0.43	$(1.02\pm0.08)E-15$	74.2±6.0	1520±80
C IV_B	1549.5	2.500 ± 0.001	$20.2 {\pm} 0.5$	$(1.36\pm0.09)E-15$	$99.3 {\pm} 6.9$	3900 ± 90
C iv-R + C iv-B	1549.5	2.511 ± 0.001	10.3 ± 0.4	$(2.39\pm0.12)E-15$	173 ± 9	1990 ± 70
He II_R	1640.4	$2.511 {\pm} 0.001$	$6.53{\pm}1.12$	$(6.61\pm1.50)E-17$	$5.32{\pm}1.21$	1190 ± 210
Не 11_В	1640.4	2.499 ± 0.002	17.7 ± 1.3	$(1.47\pm0.24)E-16$	$11.7 {\pm} 1.9$	3240 ± 240
He II_R + He II_B	1640.4	2.509 ± 0.002	11.6 ± 1.2	$(2.13\pm0.28)E-16$	17.2 ± 2.3	2120 ± 220
O III]	1663.5	2.516 ± 0.002	12.2 ± 2.4	$(3.70\pm0.95)E-17$	$3.18 {\pm} 0.82$	2200 ± 430
Al III	1858.8	$2.508 {\pm} 0.002$	$19.7 {\pm} 1.7$	$(9.09\pm1.02)E-17$	$9.23{\pm}1.04$	3180 ± 280
Si III	1892.0	$2.511 {\pm} 0.003$	$10.7 {\pm} 4.9$	$(2.64\pm1.60)E-17$	$2.79{\pm}1.69$	1690 ± 770
C 111]_R	1908.7	$2.510 {\pm} 0.001$	$12.9 {\pm} 1.4$	$(1.69\pm0.25)E-16$	$18.2 {\pm} 2.7$	2030 ± 220
С III]_В	1908.7	2.503 ± 0.002	42.0 ± 2.9	$(4.02\pm0.65)E-16$	43.1 ± 7.0	6600 ± 450
C III] $R + C III]$ B	1908.7	2.509 ± 0.002	$19.6 {\pm} 1.2$	$(5.71\pm0.70)E-16$	61.4 ± 7.5	3080 ± 180

Note—See Table 3 for the description of each column.

Table 6. The detected lines of J1443

Line name	λ_{rest} [Å]	$z_{ m line}$	$FWHM_{\rm rest}$ [Å]	$F_{\rm line} [{\rm erg \ s^{-1} \ cm^{-2}}]$	EW_{rest} [Å]	$v_{\rm width} \ [{\rm km \ s^{-1}}]$
(1)	(2)	(3)	(4)	(5)	(6)	(7)
Ly α	1215.7	3.341 ± 0.003	10.7 ± 2.6	$(5.28\pm1.31)E-16$	64.3±17.3	2650 ± 640
NV	1240.8	3.312 ± 0.001	$16.8 {\pm} 0.6$	$(2.06\pm0.09)E-15$	242 ± 23	4070 ± 160
Si II	1262.6	$3.326{\pm}0.003$	16.0 ± 1.9	$(1.24\pm0.17)E-16$	13.6 ± 2.0	3790 ± 450
Si IV	1393.8	3.324 ± 0.010	16.4 ± 3.2	$(3.18\pm1.01)E-16$	$33.8 {\pm} 10.7$	3520 ± 700
O IV]	1399.9	3.339 ± 0.008	12.9 ± 2.5	$(1.74\pm1.11)E-16$	$18.6 {\pm} 11.8$	2760 ± 540
C IV_R	1549.5	3.328 ± 0.005	$14.2 {\pm} 1.6$	$(6.15\pm1.74)E-16$	60.0 ± 17.0	2740 ± 310
C IV_B	1549.5	$3.296 {\pm} 0.007$	15.7 ± 2.3	$(5.47\pm1.67)E-16$	$54.9 {\pm} 16.8$	3030 ± 440
$C \text{ IV_R} + C \text{ IV_B}$	1549.5	$3.317{\pm}0.006$	$23.1 {\pm} 1.8$	$(1.16\pm0.24)E-15$	114 ± 24	4470 ± 350
He II_R	1640.4	3.337 ± 0.001	$6.48{\pm}1.56$	$(3.23\pm1.01)E-17$	$3.24{\pm}1.02$	1180 ± 290
Не п₌В	1640.4	3.307 ± 0.004	20.3 ± 2.9	$(9.15\pm1.53)E-17$	$9.25{\pm}1.56$	3710 ± 530
He II $_{ m R}$ + He II $_{ m B}$	1640.4	$3.335{\pm}0.003$	$20.1 {\pm} 2.5$	$(1.24\pm0.18)E-16$	12.4 ± 1.9	3680 ± 460

Note—See Table 3 for the description of each column.

Table 7.	Emission-line	contribution	and ex	cess ma	gnitude to	the	power-law	fit in	the g -	and
r-band										

	g-band			r-band		
	Total EWs	$\Delta \mathrm{mag}$	Excess mag	Total EWs	$\Delta \mathrm{mag}$	Excess mag
	[Å]	[AB mag]	[AB mag]	[Å]	[AB mag]	[AB mag]
HSC J0907	>604	>0.37	1.27	129	0.09	0.47
${\rm HSC\ J1202}$	> 258	>0.18	1.13	836	0.48	0.44
${\rm HSC\ J1207}$	>608	>0.38	1.48	329	0.21	0.65
HSC J1443	1380	0.72	0.88	721	0.43	0.40

NOTE— Δ mag: 2.5 log (1 + EW/BW), Excess mag: The excesses of the g- and r- band magnitudes between the observed magnitudes and the expected magnitudes from an extrapolation of the power-law fit to the longer wavelength bands.

lowing Toba et al. (2019) (and Toba et al. 2022). Accordingly we found the counterparts of two BluDOGs (J1202 and J1207). The separation between the HSC position and the H-ATLAS counterpart position is 0.94 arcsec for J1202, and 9.7 arcsec for J1207. The relatively large separation in the latter case suggests the counterpart being a coincidental detection. There are two WISE sources around J1207 (Figure 4); one probably corresponds to J1207 itself (the angular separation between the HSC and WISE potions is 0.65 arcsec) and another is located at 19 arcsec away to the north-east direction. The H-ATLAS source is located between these two WISE sources, and thus the FIR fluxes given in the H-ATLAS catalog are possibly attributed to the two WISE sources. Therefore we regard the H-ATLAS fluxes of J1207 as the upper limit. For the remaining two Blu-DOGs (J0907 and J1443), we adopt the 5σ upper limit fluxes.

As for the optical–MIR photometric data, we utilize g, r, i, z, y (HSC-SSP), Z, Y, J, H, Ks (VIKING DR2), W1, W2, W3, and W4 (ALLWISE) bands (see Noboriguchi et al. 2019). Note that the SNRs in these bands are more than 5, except for the W4-band with SNR more than 3 because we adopted such SNR cut in the selection of DOGs (Noboriguchi et al. 2019). Since the g- and r-band photometry are significantly affected by the strong emission lines (see Figure 1 and Tables 3–6) which cannot be treated properly in X-CIGALE, we corrected for their contribution by referring to the estimates given in Table 7.

The models and parameters of X-CIGALE adopted in this work are summarized in Table 8. We assume a delayed star formation history (SFH; Ciesla et al. 2015) with the e-folding times of the main stellar population

 $(\tau_{\rm main})$ and late starburst population $(\tau_{\rm burst})$, mass fraction of the late burst population (f_{burst}) , and age of the main stellar population (Age_{main}) and the late burst (Age_{burst}). As the stellar population, we assume the initial mass function of Chabrier (2003), solar metallicity, and 10-Gyr separation between young and old stellar population (Age_{separation}). The nebular emission model (Inoue 2011) is characterized by the ionization parameter (U), fractions of Lyman continuum photons escaping the galaxy $(f_{\rm esc})$ and absorbed by dust (f_{dust}) , and line width. We utilize a modified dust attenuation model presented by Boquien et al. (2019). The dust attenuation model for the continuum is taken from Calzetti et al. (2000) with the extension taken from Leitherer et al. (2002) between the Lyman break and 1500 Å. The emission lines are attenuated with a Milky Way extinction with $R_V = 3.1$ (Cardelli et al. 1989). We assumed $E(B-V)_{\text{continuum}} = 0.44E(B-V)_{\text{line}}$, following Calzetti et al. (2000). The $E(B-V)_{line}$ is varied between 3 and 10. We utilize the SKIRTOR model as the AGN emission model, which takes geometric parameters of the AGN into account and also allows us to incorporate the effect of extinction by the polar dust. The parameters of the AGN model are the average edge-on optical depth at 9.7 μ m ($\tau_{9.7}$), the torus density parameters (p and q; Stalevski et al. 2016), the angle between the equatorial plane and the edge of the torus (oa), the ratio of the maximum to minimum radii of the dust torus $(R_{\rm ratio})$, the fraction of total dust mass inside clumps $(M_{\rm cl})$, the inclination (i), the AGN fraction $(f_{\rm AGN})$, the extinction law, color excess $(E(B-V)_{polar\ dust}^{AGN})$, dust temperature $(T_{\text{polar dust}}^{\text{AGN}})$, and emissivity index of the polar dust.

Table 8. Parameters adopted in the X-CIGALE fit

Parameter	Value
Delayed SFH (Cie	esla et al. 2015)
$ au_{ m main} \ [{ m Myr}]$	100, 250, 500
$ au_{ m burst} \; [m Myr]$	10, 50
$f_{ m burst}$	$0.0,\ 0.5,\ 0.99$
Age_{main} [Myr]	500, 800, 1000
Age_{burst} [Myr]	1, 5, 10
Single stellar population (1	Bruzual & Charlot 2003)
IMF	Chabrier (2003)
Metallicity	0.02
$Age_{separation}$ [Myr]	10
Nebular emission	n (Inoue 2011)
$\log U$	-2.0
$f_{ m esc}$	0.0
$f_{ m dust}$	0.0
Lines width $[\text{km s}^{-1}]$	300.0
Dust attenuation (C	alzetti et al. 2000)
$E(B-V)_{\text{line}}$	3, 4, 5, 6, 7, 8, 9, 10
$f_{E(B-V)}$	0.44
$\lambda_{ m UV,bump} \ [m nm]$	217.5
FWHM _{UV,bump} [nm]	35.0
$A_{ m UV,bump}$	0.0
δ	0.0
Extinction law of emission lines	the Milky Way
R_V	3.1
Dust emission (D	
AGN fraction	0.0
$lpha_{ m IR,AGN}$	0.0625, 0.2500, 2.0000
AGN model (Stale	
$ au_{9.7}$	3, 7
p	1.0
q	1.0
oa [deg]	10, 20, 30, 40, 50, 60, 70, 80
$R_{ m ratio}$	20
$M_{ m cl}$	0.97
$i [\deg]$	0, 10, 20, 30, 40,
, [O]	50, 60, 70, 80, 90
$f_{ m AGN}$	0.1, 0.3, 0.5, 0.7, 0.9
Extinction law of polar dust	Calzetti et al. 2000
$E(B-V)_{ m polar\ dust}^{ m AGN}$	0.1, 0.2, 0.3, 0.4, 0.5
$T_{ m polar~dust}^{ m AGN} \ [m K]$	600, 700, 800, 900,
-polar dust [11]	1000, 1100, 1200, 1300, 1400
Emissivity of polar dust	
Eminosivity of polar dust	1.6




Figure 3. The SED of J0907 (top), J1202 (middle upper), J1207 (middle lower), and J1443 (bottom). The red dots denote the g- and r-band magnitudes, while the blue dots denote the longer-wavelength optical and near-infrared magnitudes that are used for the power-law fit (black line). The green lines represent the observed spectra.

The best-fit SED models are shown in Figure 5. The reduced χ^2 of the fits are 1.38, 3.19, 0.93, and 1.74 for J0907, J1202, J1207, and J1443, respectively. The best-fit values and associated errors for $E(B-V)_{\rm polar\ dust}^{\rm AGN}$ are estimated with a Bayesian-like strategy presented in Noll et al. (2009), and are reported in Ta-




Figure 4. J1207 images in the WISE W1-band (left) and H-ATLAS 250 μ m-band (right). The orange stars, green crosses, and red cross denote source detections in the HSC-SSP, ALLWISE, and H-ATLAS catalogs, respectively. The size of each image is $60'' \times 60''$, centered at the HSC position of J1207.

ble 9. On the other hands, we cannot quantitatively constrain the parameters of the host galaxies because the E(B-V) values are too large and the optical parts in their SEDs are dominated by their AGN emission (see Figure 5).

3.4. Measurement of the SMBH mass

We have detected the C IV emission line for all the four BluDOGs and Mg II emission line for J0907, both of which are widely used to calculate the SMBH mass of type-1 AGNs. Note that the systematic uncertainty is larger in the C IV-based SMBH mass than in the Mg II-based SMBH mass, due to a powerful outflow sometimes seen in the C IV velocity profile (e.g., Baskin & Laor 2005; Netzer 2015; Coatman et al. 2017). We calculate the single-epoch mass of SMBHs with the C IV and Mg II emission lines, following the calibrations given in Vestergaard & Peterson (2006) and Vestergaard & Osmer (2009) respectively:

$$M_{\rm BH} = 10^{6.66} \left(\frac{\rm FWHM(C~IV)}{10^3~{\rm km~s^{-1}}}\right)^2 \left(\frac{\lambda L_{\lambda}(1350 \rm \mathring{A})}{10^{44}~{\rm erg~s^{-1}}}\right)^{0.53} M_{\odot}, \tag{1}$$

and

$$M_{\rm BH} = 10^{6.86} \left(\frac{\rm FWHM(Mg~II)}{10^3~\rm km~s^{-1}}\right)^2 \left(\frac{\lambda L_{\lambda}(3000\text{Å})}{10^{44}~\rm erg~s^{-1}}\right)^{0.5} M_{\odot},$$
(2)

where FWHM(C IV), FWHM(Mg II), $\lambda L_{\lambda}(1350\text{Å})$ and $\lambda L_{\lambda}(3000\text{Å})$ are the FWHM of the C IV and Mg II velocity profile, and the monochromatic luminosity at 1350 Å and 3000 Å, respectively. Note that we use the FWHM of C IV_R + C IV_B as the FWHM of the C IV. We cannot eliminate the possibility that the estimated

SMBH masses are overestimated because the C IV profiles are affected by nucleus outflows (Section 4.1). For estimating the reddening-corrected monochromatic luminosity, we use the optical spectra presented in Section 3.1. We converted the spectra to the rest-frame, de-reddened them with $E(B-V)_{\rm polar}^{\rm AGN}$ derived in the SED fit, and masked out emission and absorption lines as well as pixels with negative values. Then, we fit a power-law continuum model to the spectra and estimate the monochromatic luminosities from the best fits. The estimated $\lambda L_{\lambda}(1350)$ of J0907, J1202, J1207, and J1443 are $(1.54\pm0.05)\times10^{45}$, $(9.64\pm0.27)\times10^{45}$, $(3.06\pm0.06)\times10^{45}$, and $(2.93\pm0.03)\times10^{45}$ erg s⁻¹, respectively. The $\lambda L_{\lambda}(3000)$ of J0907 is estimated to be $(1.45\pm0.04)\times10^{45}$ erg s⁻¹.

The resultant SMBH masses are summarized in Table 9. It should be noted that the C IV-based $M_{\rm BH}$ and Mg II-based $M_{\rm BH}$ of J0907 is not consistent within the statistical error. This is probably attributed to a systematic error especially in the C IV-based $M_{\rm BH}$, known to be accompanied with a large systematic error (\sim 0.5 dex; see, e.g., Shen 2013). Hereafter we use only the C IV-based $M_{\rm BH}$, since it is measured in all the four BluDOGs.

4. DISCUSSION

4.1. Spectral features and nuclear outflows

We found that the redshifts of the four BluDOGs are in the range of 2.2 \lesssim $z_{\rm sp}$ \lesssim 3.3. They are systematically higher than the typical redshifts of DOGs $(z_{\rm sp} = 1.99 \pm 0.45; \text{ Dey et al. 2008}; \text{ Pope et al. 2008}).$ One possible reason for this systematically high redshift is a selection effect related to the blue-excess criterion. When we select BluDOGs from the parent DOG sample, the q- and r-band magnitudes show an excess of the expected magnitudes estimated by the power-law extrapolation from i-band to W4-band. Thus we may select DOGs in a preferred redshift range where strong emission lines such as Ly α and C IV shifts into the two bands (see Section 4.2 for more quantitative assessments). The reason for the underestimated photometric redshift (~ 1 ; Noboriguchi et al. 2019) is the unusual emission lines with the large REW.

The detected emission lines have large velocity widths, $\gtrsim 2000~\rm km~s^{-1}$ in most cases. This suggests that the broad-line region (BLR) of the BluDOGs is not completely obscured; in other words, the observed Blu-DOGs are classified as type-1 AGNs. This is an unexpected result, because their very red color between optical and mid-IR suggests the heavily obscured nature. One possible interpretation is that we are looking at a phase where the surrounding dust is just blown




Figure 5. The results of the SED fitting for the four BluDOGs. The upper left, upper right, lower left and lower right panels show the results of J0907, J1202, J1207, and J1443, respectively. The black, blue, green, red, and orange lines represent the best-fit model, stellar component (with dust attenuation), AGN component, SF component (FIR re-emission from the dust heated by SF), and nebular component, respectively. The magenta plots represent the photometric data. The arrows denote 5σ upper limit flux.

Table 9. Physical properties of the four BluDOGs

	HSC J0907	HSC J1202	HSC J1207	HSC J1443
Redshift	2.258 ± 0.002	2.830 ± 0.002	2.511 ± 0.001	3.317 ± 0.006
$E(B-V)_{\text{polar dust}}^{\text{AGN}}$	0.26 ± 0.05	0.33 ± 0.04	0.30 ± 0.02	0.20 ± 0.01
$L_{ m bol}^{ m AGN}/L_{\odot}$	$(6.11 \pm 0.95) \times 10^{12}$	$(6.11 \pm 1.18) \times 10^{13}$	$(7.95 \pm 1.47) \times 10^{12}$	$(2.52 \pm 0.13) \times 10^{13}$
$M_{ m BH}~({ m C~IV})/M_{\odot}$	$(1.69 \pm 0.17) \times 10^8$	$(4.95 \pm 0.30) \times 10^8$	$(1.11 \pm 0.08) \times 10^8$	$(5.48 \pm 0.86) \times 10^8$
$M_{ m BH}~({ m Mg~II})/M_{\odot}$	$(9.85 \pm 1.80) \times 10^7$	_	_	_
$\lambda_{ m Edd}$ (C IV)	1.10 ± 0.20	3.75 ± 0.76	2.19 ± 0.43	1.40 ± 0.23

away by the nuclear activity (outflow, radiation pressure, or both), as discussed more in Section 4.3. It should be noted that the type-1 nature is seen not only in the presented BluDOGs but also in some other DOGs (e.g., Toba & Nagao 2016; Toba et al. 2017a; Zou et al. 2020). Systematic spectroscopic observations for the whole populations of DOGs are required to study the nature of obscuration occurring in various populations of DOGs.

As shown in Figure 2, the velocity profile of the observed C IV lines show a notable excess feature in the blue wing. Such an excess in the C IV velocity profile has been observed in other type-1 AGNs, and interpreted as a result of powerful nuclear outflows (e.g., Baskin & Laor 2005; Netzer 2015; Coatman et al. 2017). To evaluate quantitatively how the nuclear outflow in BluDOGs is strong compared to ordinary AGNs, we examine the "asymmetry parameter (α_{β}) " defined by De Robertis (1985) as

$$\alpha_{\beta} = \frac{\lambda_c(3/4) - \lambda_c(1/4)}{\Delta\lambda(1/2)},\tag{3}$$

where $\lambda_c(h)$ and $\Delta\lambda(1/2)$ are the central wavelength at which the flux falls to a h time the peak flux and FWHM of the broad profile, respectively. The positive and negative values of α_{β} express the blue and red excesses, respectively. The derived values of α_{β} for J0907, J1202, J1207, and J1443 are 0.216, 0.102, 0.246, and 0.051, respectively. As a reference, the C IV velocity profile in the composite spectrum of SDSS type-1 quasars given by Vanden Berk et al. (2001) shows $\alpha_{\beta} = 0.110$. Thus J0907, and J1207 may possess a significant nuclear outflow that is more powerful than typical quasars.

In order to compare α_{β} of the BluDOG with that of another dusty AGN population, we fitted the C IV profile of 97 "core" ERQs (ERQs with REW(C IV) > 100 Å) in Hamann et al. (2017) and measured α_{β} by adopting a single or double Gaussian profile. The core ERQ sample consists of 80 objects without BAL and 17 objects with BAL, and we investigate the statistics of α_{β} for the two subsamples separately because the BAL feature can affect the C IV line profile. Here we exclude J1443 from the BluDOG sample when comparing the α_{β} index because its velocity profile is largely affected by narrow absorption lines (hereafter the limited-BluDOG sample to infer the 3 BluDOGs; i.e., J0907, J1202, and J1207). Figure 6 shows the cumulative fraction of α_{β} for the limited-BluDOGs, core ERQs without BAL, and core ERQs with BAL. The averaged values of the limited-BluDOGs, core ERQs without BAL, and core ERQs with BAL are 0.15 ± 0.08 , 0.02 ± 0.13 , and 0.01 ± 0.09 , respectively. We performed the Kolmogorov-Smirnov

test (KS-test) to examine the statistical significance of the difference in α_{β} among the samples. The p-values of the limited-BluDOGs-core ERQs without BAL, and limited-BluDOG-core ERQs with BAL are 0.0178 and 0.0175, respectively. Thus we conclude that the distributions of α_{β} of the limited-BluDOGs and core ERQs with/without BAL are marginally different with > 2 sigma significance. This suggests that the BluDOGs show nuclear outflow that is possibly more powerful than the nuclear outflow in core ERQs with/without BAL.




Figure 6. Cumulative distribution of the α_{β} indices for the limited-BluDOGs (see the main text; blue line), core ERQs without BAL (green line), and core ERQs with BAL (orange line). The red dashed line denotes the α_{β} index measured for the composite spectrum of SDSS type-1 quasars.




Figure 7. Cumulative distribution of the kt_{80} indices for the limited-BluDOGs (blue line), core ERQs without BAL (green line), and core ERQs with BAL (orange line). The red dashed line denotes the kt_{80} index for the single Gaussian profile.

We also focus on the kurtosis index (kt_{80}) defined as follows (see Hamann et al. 2017 for detailes): $kt_{80} = \Delta v(80\%) / \Delta v(20\%)$, where $\Delta v(x\%)$ is the velocity width at x% of the peak flux height. In addition to α_{β} , this kt_{80} index is useful to characterize the C IV wing (a more prominent blue wing results in smaller kt_{80}). By using the best-fit double Gaussian profile of the BluDOGs, kt_{80} of J0907, J1202, J1207, and J1443 are 0.276, 0.313, 0.252, and 0.440, respectively. Again we exclude J1443 from the BluDOG sample when comparing the kt_{80} index as the discussion of the α_{β} index. For comparison, kt_{80} of a single Gaussian is $\sqrt{1-\frac{2\ln(2)}{\ln(5)}}$ (~ 0.37), whereas most quasars have $kt_{80} \stackrel{\cdot}{\sim} 0.15 - 0.30$ (see Figure 7 in Hamann et al. 2017). The limited-BluDOGs, core ERQs without BAL, and core ERQs with BAL show $kt_{80} = 0.28 \pm 0.03$, 0.33 ± 0.06 , and 0.34 ± 0.05 , respectively (see also Figure 7). Note that the C IV profile of 41 core ERQs without BAL and 7 core ERQs with BAL is fitted by a single Gaussian, which is the reason why many objects have $kt_{80} \sim 0.37$ as shown in Figure 7. C IV velocity profiles of core ERQs with/without BAL are roughly consistent with the Gaussian without a blue wing. However, the kt_{80} index of the limited-BluDOGs is less than $\sqrt{1-\frac{2\ln(2)}{\ln(5)}}$, suggesting that their C IV line profile has a wing. We performed the KS-test to examine the statistical significance of the difference in kt_{80} among the samples. The p-values of the limited-BluDOGs-core ERQs without BAL, and limited-BluDOGs-core ERQs with BAL are 0.0637 and 0.0175, respectively. Therefore, we conclude that the distributions of kt_{80} between the samples of the limited-BluDOGs and core ERQs with/without BAL feature are marginally different with > 2 sigma significance.

It has been reported that AGNs with a high Eddington ratio tend to show a Lorentzian-line velocity profile in BLR lines (e.g., Moran et al. 1996; Véron-Cetty et al. 2001; Collin et al. 2006; Zamfir et al. 2010). Therefore, the small kt_{80} value of BluDOGs can be caused by the contribution of extended Lorentzian wings instead of the asymmetric blue wing. For a symmetric Lorentzian profile, $kt_{80} \sim 1/16$ (much smaller than a Gaussian profile, $kt_{80} \sim 0.37$) and $\alpha_{\beta} = 0$ are expected. However, the BluDOGs are inconsistent with this expectation (Figure 8). This Figure 8 also shows that the BluDOGs follow the trend made by core ERQs with/without BAL in the $kt_{80} - \alpha_{\beta}$ plane, while a systematic deviation of BluDOGs toward $(\alpha_{\beta}, kt_{80}) = (0, 0)$ is expected if a Lorentzian component significantly contributes to the C IV line of BluDOGs. Thus, we conclude that extended Lorentzian wings do not affect the C IV line profile of

BluDOGs, but the small kt_{80} of BluDOGs is caused by the asymmetric blue excess due to the stronger nuclear outflow than that of ERQs.




Figure 8. The kt_{80} vs. α_{β} plot for the limited-BluDOGs (blue dots), core ERQs without BAL (green dots), and core ERQs with BAL (orange dots). The black solid line and dashed line denote the kt_{80} values of single Gaussian and single Lorentzian, respectively. Note that 41 core ERQs without BAL and 7 core ERQs with BAL are fitted with a single Gaussian, and they are plotted at $(\alpha_{\beta}, kt_{80}) = (0, \sqrt{1 - \frac{2\ln(2)}{\ln(5)}})$.

4.2. Large equivalent widths of the CIV emission

As we summarized in Table 7, the blue excess in J1443 can be almost explained by the contribution of the strong emission lines. This is also the case for J1202 by taking into account of the additional contribution of unobserved Ly α to g-band. On the other hand, the blue excess of the remaining two BluDOGs cannot be explained only by the contribution of BLR emission lines. Figure 3 strongly suggests that a part of the excess flux comes from the continuum emission, which deviates at $\lesssim 7000$ Å from the extrapolation of the power-law fit. These results demonstrate the complexity and diversity of BluDOGs; systematic exploration of a larger sample is required to statistically understand the origin of the blue excess.

Not only REW(C IV), but the REW of other BLR emission lines are also systematically larger than observed in typical type-1 quasars (see Tables 3–6, Figure 9, and also Table 2 in Vanden Berk et al. 2001). Such a trend may be explained if the observed Blu-DOGs have lower UV luminosity than typical quasars owing to the Baldwin effect (Baldwin 1977; Kinney et al. 1990; Baskin & Laor 2004), i.e., the negative correlation between the REWs and the continuum luminosities of




Figure 9. Rest-frame EW ratio vs. rest-frame wavelength. REW ratios are defined as REWs of objects over REWs from the composite spectrum of SDSS type-1 quasar measured by Vanden Berk et al. (2001). The red, blue, green, and orange plots show the REW ratios of J0907, J1202, J1207, and J1443, respectively.




Figure 10. Rest-frame EW of the C IV vs. the monochromatic luminosity at 1350 Å. Blue and orange dots represent the four BluDOGs and WISSH quasars (Vietri et al. 2018). The grey 2D histogram represents the number density of the SDSS quasars (Shen et al. 2011). The green line represents the linear fit to the distribution of the SDSS quasars. The red plots show the mean and standard deviation in luminosity bins with 0.5 dex width. The numbers of the SDSS quasars in the individual bins are shown at the bottom of the panel.

quasars. Figure 10 shows the four BluDOGs on the C IV REW vs. $\lambda L_{\lambda}(1350\text{Å})$ diagram. Note that the REW of J1443 (114 ± 24 Å) is somewhat smaller than that of the remaining three BluDOGs (148 ± 12, 203 ± 10 and 173 ± 9 Å for J0907, J1202 and J1207, respectively; see Tables 3–6). This is partly because of an underestima-

tion of the C IV flux caused by the absorption features. The figure also shows SDSS type-1 quasars with reliable measurement of C IV REW (EWCIV/e_EWCIV > 5) and without broad absorption lines (BAL < 1) taken from Shen et al. (2011). Since the Baldwin effect does not significantly depend on redshift (e.g., Croom et al. 2002; Dietrich et al. 2002; Niida et al. 2016), we do not adopt any redshift criterion to select the SDSS quasars so that a wide luminosity range is covered. We also use another comparison sample taken from the WISE/SDSS selected hyper-luminous quasar sample (WISSH; Bischetti et al. 2017; Vietri et al. 2018)⁵, in order to add objects at the high-luminosity end.

Figure 10 clearly shows that the C IV REWs of Blu-DOGs are larger than the comparison samples at a given UV luminosity. The excess REW over the average relation of the Baldwin effect (shown with a green solid line in Figure 10) for J0907, J1202, J1207, and J1443 are 0.29, 0.66, 0.44, and 0.26 dex, respectively. This excess is larger than the scatter of the comparison samples (see red plots in Figure 10). Therefore the large REW seen in the BluDOGs are not due to the Baldwin effect.

The averages and standard deviations of REW(C IV) for core ERQs and ERQ-like objects are 178 ± 74 and 86 ± 45 Å, respectively (Hamann et al. 2017). The distributions of REW(C IV) and $(i-W3)_{AB}$ color for Blu-DOGs are consistent with these of core ERQs although the most of core ERQs and ERQ-like objects do not show a blue-wing profile in C IV (Section 4.1). Hamann et al. (2017) proposed a scenario that the large REW of ERQs are possibly due to the spatially extended geometry of BLRs caused by the powerful nuclear outflow. If the obscuration is heavier for the accretion disk than for the BLRs which have extended geometry, the continuum emission is more heavily extinct than the BLR emission lines and thus the observed-frame EW becomes larger. Such a scenario may also apply to BluDOGs. Unfortunately it is not observationally feasible to confirm this idea by resolving the spatial structure of BLRs in ERQs or BluDOGs due to the required angular resolution, even with the JWST or exisiting ground-based interferome-

 5 The C IV REW and C IV line luminosity of WISSH quasars are given by Vietri et al. (2018). To calculate $L_{\lambda}(1350\text{Å})$ of WISSH quasars, we assume that the continuum spectrum of WISSH quasars is a power-law and adopt the following formula:

$$L_{\lambda}(1350\text{Å}) = \frac{L^{\text{line}}(\text{C IV})}{REW(\text{C IV})} \times \left(\frac{1350}{1549}\right)^{\alpha_{\lambda}},\tag{4}$$

where $L_{\lambda}(1350 \text{Å})$, $L^{\text{line}}(\text{C IV})$, and α_{λ} are the monochromatic luminosity at 1350 Å, the line luminosity of C IV, and power-law index, respectively. Here we adopt $\alpha_{\lambda} = -1.7$ (Vanden Berk et al. 2001) as the power-law index.

ters. Without spatially resolving them, a possible approach is the velocity-resolved reverberation mapping of the geometry and kinematics of BLR clouds (e.g., Horne et al. 2004; Denney et al. 2009; Li et al. 2013; Kollatschny et al. 2014; Pancoast et al. 2014).

4.3. Possible extreme accretion and the nature of BluDOGs

To understand the nature of BluDOGs especially in the context of the major-merger scenario for the quasar evolution, we compare the SMBH accretion of the four BluDOGs with other AGN populations. Figure 11 is a diagram of $L_{\rm bol}$ vs. the SMBH mass. As in Section 4.2, SDSS quasars (Shen et al. 2011) and WISSH quasars (Vietri et al. 2018) are used as comparison samples. For the SDSS quasars, we select only non-BAL quasars (BAL < 1) with the uncertainty of $L_{\rm bol}$ and $M_{\rm BH}$ less than $0.5 \operatorname{dex} (e_{\log BHCV} < 0.5 \& e_{\log Lbol} < 0.5)$, and adopt the C IV-based SMBH mass for a fair comparison with those of the BluDOGs. We also plot samples of 28 ERQs (Perrotta et al. 2019), 5 Hot DOGs (Wu et al. 2018), 2 power-law DOGs (Melbourne et al. 2011), and 1 Compton-thick (CT) DOG (Toba et al. 2020). Hot DOGs are DOGs with a special color of WISE (very faint in the 3.4 μ m and 4.5 μ m bands, but bright in the longer bands; Eisenhardt et al. 2012; Wu et al. 2012), while power-law DOGs are DOGs with a featureless power-law SED from optical to mid-IR (e.g., Dev et al. 2008; Bussmann et al. 2012; Toba et al. 2015; Noboriguchi et al. 2019). The CT DOG was identified by Nuclear Spectroscopic Telescope Array (Harrison et al. 2013) from SDSS-WISE DOGs sample. All but the CT DOG have spectroscopic redshifts. The SMBH masses of the ERQs and the WISSH quasars are estimated from $H\beta$, while those of the Hot DOGs and the DOGs are estimated from H α . The SMBH mass of the CT DOG was estimated by Toba et al. (2020) from the stellar mass by using an empirical relation between the stellar mass and SMBH mass (Kormendy & Ho 2013). Since Perrotta et al. (2019) and Melbourne et al. (2011) did not correct the absorption of dust, the $M_{\rm BH}$ of ERQs and DOGs are lower limits.

Figure 11 shows that the four BluDOGs are more luminous than the other AGN populations at a given SMBH mass, or equivalently, they have lower-mass SMBHs than the other AGN populations at a given bolometric luminosity. This suggests that the SMBH growth in the BluDOGs is more rapid than AGNs in comparison samples. Indeed, the Eddington ratios ($\lambda_{\rm Edd}$) of J0907, J1202, J1207, and J1443 are 1.10 \pm 0.20, 3.89 \pm 0.78, 2.19 \pm 0.44, and 1.40 \pm 0.23, respectively (Table 9), with the average value of 2.26. In other words, the SMBHs




Figure 11. The SMBH mass vs. the bolometric luminosity diagram. The filled-blue stars and gray contour denote the BluDOGs and SDSS quasars, while the filled hexagons with green, orange, purple, cyan, and light green colors denote ERQs (Perrotta et al. 2019), WISSH quasars (Vietri et al. 2018), Hot DOGs (Wu et al. 2018), DOGs (Melbourne et al. 2011, 2012), and a CT DOG (Toba et al. 2020), respectively. The red dashed lines represent a constant Eddington ratio of $\lambda_{\rm Edd} = 0.01, \, 0.1, \, 1.0, \, {\rm and} \, \, 10.0.$

in the BluDOGs are now in the stage of the Eddingtonlimit or super-Eddington accretion. Even if the intrinsic SMBH masses are lower than those estimated (Section 3.4), the conclusion of this study remains qualitatively unchanged. The higher Eddington ratios compared to other populations suggest that the SMBHs in BluDOGs are in the most rapidly evolving phase during the whole evolutionary history of SMBHs. In the gas-rich major merger scenario of Hopkins et al. (2008), the peak of the AGN activity (i.e., the mass growth of SMBHs) corresponds to the transition phase from the optically thick to optically thin quasars, where the surrounding dust is blown out by the powerful AGN activity. Note that optically thick quasars in the major merger scenario should be recognized as type-2 quasars in optical (the BLR cannot be observed due to the heavy dust reddening). Since optically-thick quasars in the final stage of the evolution can be recognized as both type-1 and type-2 due to the orientation effect toward the dusty torus, the observed type-1 nature suggests the object is not in the early (optically-thick) stage in the major merger evolutionary scenario. Preferentially in AGNs with high $\lambda_{\rm Edd}$, a blue-wing feature tends to be observed (e.g., Aoki et al. 2005; Komossa et al. 2008). The observed characteristics of the BluDOGs such as the type-1 nature and the blue-wing feature of the C IV velocity profile are consistent with the picture that BluDOGs are in such a peak stage of the SMBH evolution.

To discuss the evolutionary relation among populations of dusty galaxies (BluDOGs, core ERQs, and hot DOGs), we focus on E(B-V) and kt_{80} . E(B-V) for BluDOGs, core ERQs (Hamann et al. 2017), and hot DOGs (Wu et al. 2018) are 0.273 ± 0.049 , 0.242 ± 0.127 , and 4.781 ± 1.986 , respectively. The E(B-V) of the hot DOGs is significantly larger than that of the Blu-DOG and core ERQ samples, suggesting the hot DOGs are thought to be in a heavily obscured phase. Since the kt_{80} of BluDOGs is smaller than that of core ERQs (Section 4.1), and the kt_{80} of Mid-IR detected quasars is close to that of BluDOGs (Figure 1 of Monadi & Bird 2022), the BluDOG phase is thought to be close to the optically-thin quasar phase. Therefore, it is suggested that the evolutionary path of various AGN populations is "Hot DOGs - core ERQs - BluDOGs - optically-thin quasars".

For AGNs in general, the mass accretion efficiency (η) is defined as following:

$$\eta = \frac{L_{\text{bol}}}{\dot{M}c^2},\tag{5}$$

where M is the mass accretion rate. By multiplying the \dot{M} and lifetime of BluDOGs ($t_{\rm life}$), we can roughly estimate the accreted mass (M_{Acc}) in the BluDOGs phase. Bian & Zhao (2003) estimated $\log \eta = -1.61$ of Seyfert 1 galaxies and Palomar-Green quasars by assuming that the geometrically-thin and optically-thick standard α prescription accretion disk model (Shakura & Sunyaev 1973). By assuming $\log \eta = -1.61$ and $t_{life} = 1$ Myr (Noboriguchi et al. 2019), the estimated M_{Acc} of J0907, J1202, J1207, and J1443 are about 1.68×10^7 , 1.68×10^8 , 2.19×10^7 , and $6.93 \times 10^7 M_{\odot}$. The SMBH masses reached when the SMBH masses of the BluDOGs are increased by the observed mass accretion rate during the typical lifetime of BluDOGs $(M_{\rm BH}^+ = M_{\rm BH} + M_{\rm Acc})$ of J0907, J1202, J1207, and J1443 are 1.86×10^8 , 6.63×10^8 , 1.32×10^8 , and $6.17 \times 10^8 M_{\odot}$, respectively. Therefore the SMBH mass of BluDOGs increases by $\sim 20\%$ during the short BluDOG phase, suggesting that BluDOGs are actually in a rapidly glowing phase.

Figure 12 shows the Eddington ratios of various populations of AGNs as a function of redshift. The excess of $\lambda_{\rm Edd}$ of the four BluDOGs is more significant than the scatter of the $\lambda_{\rm Edd}$ distribution, suggesting that BluDOGs are a special class of AGNs that harbor SMBHs in the most actively evolving phase. Then, why such a class of AGNs is found only in a limited redshift range, $2.2 < z_{\rm sp} < 3.3$? A possible reason comes from their selection criteria, as briefly mentioned in Sec 4.1. Since the BluDOGs are selected by the blue excesses which are largely caused by the contribution of strong BLR

emission lines, the resultant redshift distribution would be biased such that the blue bands contain strong emission lines. It is also not clear whether the whole population of DOGs have systematically larger $\lambda_{\rm Edd}$ than ordinary type-1 quasars, due to the paucity of the spectroscopic data. In order to reveal the total picture of the dust-enshrouded evolution of SMBHs, more systematic spectroscopic observations for various populations of BluDOGs and DOGs are needed.




Figure 12. The redshift vs. the Eddington ratio diagram. The filled-blue stars and hexagons are the same as in Figure 11. The gray 2D histogram represents the number density of SDSS quasars (Shen et al. 2011). The red plots show the mean and standard deviation of $\lambda_{\rm Edd}$ in redshift bins with the width of $\Delta z = 0.5$. The numbers shown at the upper part denote the numbers of SDSS quasars in the redshift bins.

5. CONCLUSION

We carried out spectroscopic observations of the four BluDOGs selected by Noboriguchi et al. (2019) using Subaru/FOCAS and VLT/FORS2. The analysis of the obtained spectroscopic data revealed the following spectroscopic properties of the BluDOGs:

- 1. The rest-frame UV spectra of the BluDOGs show broad ($\gtrsim 2000~{\rm km~s^{-1}}$) emission lines. This suggests that the BLRs of the BluDOGs are not completely obscured, albeit the very dusty nature inferred from their optical-IR SED.
- 2. The C IV lines of the BluDOGs show a significant blue wing, which is more prominent than in ordinary SDSS type-1 quasars. This suggests a

presence of powerful nuclear outflow at the spatial scale of the BLR in the BluDOGs.

- 3. The REWs of their BLR lines are very large, REW(C IV)~160 Å, ~7 times larger than the average of SDSS type-1 quasars. Such strong lines cause the flux excess of the two BluDOGs in the HSC g- and r-bands, while blue continuum emission also contributes the blue excess in the remaining two objects. The large REWs are not explained by the Baldwin effect. A possible origin is a powerful nuclear outflow in the BluDOGs causing a selective obscuration of the nuclear region, as suggested for ERQs.
- 4. The Eddington ratios of the BluDOGs are higher than 1.0 and are systematically larger than other AGN populations. The mass accretion onto the SMBH in BluDOGs is in the mode of the Eddington-limit or super-Eddington accretion.

All of the above results support the scenario that Blu-DOGs represent a population of AGNs in the transition phase from optically thick to optically thin quasars, i.e., in the blowing-out phase of the major-merger scenario for the SMBH evolution. The spectroscopic properties of the BluDOGs are similar to those of ERQs. For further understandings of the complete picture, more systematic spectroscopic observations are crucial, not only of BluDOGs but also of the whole population of DOGs.

The authors gratefully acknowledge the anonymous referee for a careful reading of the manuscript and very helpful comments. This study is based on data collected at Subaru Telescope, which is operated by the National Astronomical Observatory of Japan, and observations collected at the European Southern Observatory under a ESO programme 0102.B-0614(A). We are honored and grateful for the opportunity of observing the Universe from Maunakea, which has the cultural, historical and natural significance in Hawaii. The Hyper Suprime-Cam (HSC) collaboration includes the astronomical communities of Japan and Taiwan, and Princeton University. The HSC instrumentation and software were developed by the National Astronomical Observatory of Japan (NAOJ), the Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU), the University of Tokyo, the High Energy Accelerator Research Organization (KEK), the Academia Sinica Institute for Astronomy and Astrophysics in Taiwan (ASIAA), and Princeton University. Funding was contributed by the FIRST program from Japanese Cabinet Office, the Ministry of Education, Culture, Sports, Science and

Technology (MEXT), the Japan Society for the Promotion of Science (JSPS), Japan Science and Technology Agency (JST), the Toray Science Foundation, NAOJ, Kavli IPMU, KEK, ASIAA, and Princeton University. This paper makes use of software developed for the Large Synoptic Survey Telescope. We thank the LSST Project for making their code available as free software at http://dm.lsstcorp.org. The Pan-STARRS1 Surveys (PS1) have been made possible through contributions of the Institute for Astronomy, the University of Hawaii, the Pan-STARRS Project Office, the Max-Planck Society and its participating institutes, the Max Planck Institute for Astronomy, Heidelberg and the Max Planck Institute for Extraterrestrial Physics, Garching, The Johns Hopkins University, Durham University, the University of Edinburgh, Queen's University Belfast, the Harvard-Smithsonian Center for Astrophysics, the Las Cumbres Observatory Global Telescope Network Incorporated, the National Central University of Taiwan, the Space Telescope Science Institute, the National Aeronautics and Space Administration under Grant No. NNX08AR22G issued through the Planetary Science Division of the NASA Science Mission Directorate, the National Science Foundation under Grant No. AST-1238877, the University of Maryland, and Eotvos Lorand University (ELTE) This publication has made use of data from the VIKING survey from VISTA at the ESO Paranal Observatory, program ID 179.A-2004. Data processing has been contributed by the VISTA Data Flow System at CASU, Cambridge and WFAU, Edinburgh. This publication makes use of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, funded by the National Aeronautics and Space Administration. Herschel is an ESA space observatory with science instruments provided by Europeanled Principal Investigator consortia and with important participation from NASA. This research made use of Astropy, 6 a community-developed core Python package for Astronomy (Astropy Collaboration et al. 2013, 2018). We would like to thank Enago (www.enago.jp) for the English language review and Kohei Iwashita for helpful discussion. This study was financially supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI grant Nos. 19J10458, 21H01126 (A.N.), 16H03958, 17H01114, 19H00697, 20H01949 (T.N.), 18J01050, 19K14759, 22H01266 (Y. Toba), 20H01939 (K.I.), and 20K04014 (Y. Terashima).

⁶ http://www.astropy.org

Software: Astropy, Astro-SCRAPPY, X-CIGALE, Numpy, Scipy.optimaize.curve_fit, Recipe flexible execution workbench (Reflex)

Facilities: Subaru (FOCAS, HSC), VLT (FORS2), VISTA, WISE, and Herschel (PACS, SPIRE)

REFERENCES

- Aihara, H., Arimoto, N., Armstrong, R., et al. 2018, PASJ, 70, S4, doi: 10.1093/pasj/psx066
- Aoki, K., Kawaguchi, T., & Ohta, K. 2005, ApJ, 618, 601, doi: 10.1086/426075
- Appenzeller, I., Fricke, K., Fürtig, W., et al. 1998, The Messenger, 94, 1
- Arnaboldi, M., Neeser, M. J., Parker, L. C., et al. 2007, The Messenger, 127
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, A&A, 558, A33, doi: 10.1051/0004-6361/201322068
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, AJ, 156, 123,
 - doi: 10.3847/1538-3881/aabc4f
- Baldwin, J. A. 1977, ApJ, 214, 679, doi: 10.1086/155294
- Baskin, A., & Laor, A. 2004, MNRAS, 350, L31,
- doi: 10.1111/j.1365-2966.2004.07833.x
- —. 2005, MNRAS, 356, 1029,
 - doi: 10.1111/j.1365-2966.2004.08525.x
- Bian, W.-H., & Zhao, Y.-H. 2003, PASJ, 55, 599, doi: 10.1093/pasj/55.3.599
- Bischetti, M., Piconcelli, E., Vietri, G., et al. 2017, A&A, 598, A122, doi: 10.1051/0004-6361/201629301
- Boquien, M., Burgarella, D., Roehlly, Y., et al. 2019, A&A, 622, A103, doi: 10.1051/0004-6361/201834156
- Bourne, N., Dunne, L., Maddox, S. J., et al. 2016, MNRAS, 462, 1714, doi: 10.1093/mnras/stw1654
- Bruzual, G., & Charlot, S. 2003, MNRAS, 344, 1000, doi: 10.1046/j.1365-8711.2003.06897.x
- Burgarella, D., Buat, V., & Iglesias-Páramo, J. 2005, MNRAS, 360, 1413,
 - doi: 10.1111/j.1365-2966.2005.09131.x
- Bussmann, R. S., Dey, A., Borys, C., et al. 2009, ApJ, 705, 184, doi: 10.1088/0004-637X/705/1/184
- Bussmann, R. S., Dey, A., Lotz, J., et al. 2011, ApJ, 733, 21, doi: 10.1088/0004-637X/733/1/21
- Bussmann, R. S., Dey, A., Armus, L., et al. 2012, ApJ, 744, 150, doi: 10.1088/0004-637X/744/2/150
- Calzetti, D., Armus, L., Bohlin, R. C., et al. 2000, ApJ, 533, 682, doi: 10.1086/308692
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, ApJ, 345, 245, doi: 10.1086/167900
- Chabrier, G. 2003, PASP, 115, 763, doi: 10.1086/376392

- Ciesla, L., Charmandaris, V., Georgakakis, A., et al. 2015, A&A, 576, A10, doi: 10.1051/0004-6361/201425252
- Coatman, L., Hewett, P. C., Banerji, M., et al. 2017, MNRAS, 465, 2120, doi: 10.1093/mnras/stw2797
- Collin, S., Kawaguchi, T., Peterson, B. M., & Vestergaard, M. 2006, A&A, 456, 75,
 - doi: 10.1051/0004-6361:20064878
- Croom, S. M., Rhook, K., Corbett, E. A., et al. 2002, MNRAS, 337, 275,
 - doi: 10.1046/j.1365-8711.2002.05910.x
- Cutri, R. M. 2014, VizieR Online Data Catalog, 2328
- Dale, D. A., Helou, G., Magdis, G. E., et al. 2014, ApJ, 784, 83, doi: 10.1088/0004-637X/784/1/83
- Dawson, K. S., Schlegel, D. J., Ahn, C. P., et al. 2013, AJ, 145, 10, doi: 10.1088/0004-6256/145/1/10
- De Robertis, M. 1985, ApJ, 289, 67, doi: 10.1086/162865
- Denney, K. D., Peterson, B. M., Pogge, R. W., et al. 2009, ApJL, 704, L80, doi: 10.1088/0004-637X/704/2/L80
- Desai, V., Soifer, B. T., Dey, A., et al. 2009, ApJ, 700, 1190, doi: 10.1088/0004-637X/700/2/1190
- Dey, A., Soifer, B. T., Desai, V., et al. 2008, ApJ, 677, 943, doi: 10.1086/529516
- Dietrich, M., Hamann, F., Shields, J. C., et al. 2002, ApJ, 581, 912, doi: 10.1086/344410
- Ding, X., Silverman, J., Treu, T., et al. 2020, ApJ, 888, 37, doi: 10.3847/1538-4357/ab5b90
- Eales, S., Dunne, L., Clements, D., et al. 2010, PASP, 122, 499, doi: 10.1086/653086
- Eisenhardt, P. R. M., Wu, J., Tsai, C.-W., et al. 2012, ApJ, 755, 173, doi: 10.1088/0004-637X/755/2/173
- Eisenstein, D. J., Weinberg, D. H., Agol, E., et al. 2011, AJ, 142, 72, doi: 10.1088/0004-6256/142/3/72
- Feibelman, W. A. 1983, A&A, 122, 335
- Ferrarese, L., & Merritt, D. 2000, ApJL, 539, L9, doi: 10.1086/312838
- Fiore, F., Grazian, A., Santini, P., et al. 2008, ApJ, 672, 94, doi: 10.1086/523348
- Freudling, W., Romaniello, M., Bramich, D. M., et al. 2013, A&A, 559, A96, doi: 10.1051/0004-6361/201322494
- Gebhardt, K., Bender, R., Bower, G., et al. 2000, ApJL, 539, L13, doi: 10.1086/312840
- Goulding, A. D., Greene, J. E., Bezanson, R., et al. 2018, PASJ, 70, S37, doi: 10.1093/pasj/psx135

- Griffin, M. J., Abergel, A., Abreu, A., et al. 2010, A&A, 518, L3, doi: 10.1051/0004-6361/201014519
- Hamann, F., Zakamska, N. L., Ross, N., et al. 2017, MNRAS, 464, 3431, doi: 10.1093/mnras/stw2387
- Harrison, F. A., Craig, W. W., Christensen, F. E., et al. 2013, ApJ, 770, 103, doi: 10.1088/0004-637X/770/2/103
- Hopkins, P. F., Hernquist, L., Cox, T. J., & Kereš, D. 2008, ApJS, 175, 356, doi: 10.1086/524362
- Horne, K., Peterson, B. M., Collier, S. J., & Netzer, H. 2004, PASP, 116, 465, doi: 10.1086/420755
- Inoue, A. K. 2011, MNRAS, 415, 2920, doi: 10.1111/j.1365-2966.2011.18906.x
- Kashikawa, N., Aoki, K., Asai, R., et al. 2002, PASJ, 54, 819, doi: 10.1093/pasj/54.6.819
- Kawanomoto, S., Uraguchi, F., Komiyama, Y., et al. 2018, PASJ, 70, 66, doi: 10.1093/pasj/psy056
- Kinney, A. L., Rivolo, A. R., & Koratkar, A. P. 1990, ApJ, 357, 338, doi: 10.1086/168924
- Kollatschny, W., Ulbrich, K., Zetzl, M., Kaspi, S., & Haas, M. 2014, A&A, 566, A106, doi: 10.1051/0004-6361/201423901
- Komossa, S., Xu, D., Zhou, H., Storchi-Bergmann, T., & Binette, L. 2008, ApJ, 680, 926, doi: 10.1086/587932
- Kormendy, J., & Ho, L. C. 2013, ARA&A, 51, 511, doi: 10.1146/annurev-astro-082708-101811
- Leitherer, C., Li, I. H., Calzetti, D., & Heckman, T. M. 2002, ApJS, 140, 303, doi: 10.1086/342486
- Li, Y.-R., Wang, J.-M., Ho, L. C., Du, P., & Bai, J.-M. 2013, ApJ, 779, 110, doi: 10.1088/0004-637X/779/2/110
- Magorrian, J., Tremaine, S., Richstone, D., et al. 1998, AJ, 115, 2285, doi: 10.1086/300353
- Marconi, A., & Hunt, L. K. 2003, ApJL, 589, L21, doi: 10.1086/375804
- McCully, C., & Tewes, M. 2019, Astro-SCRAPPY: Speedy Cosmic Ray Annihilation Package in Python. http://ascl.net/1907.032
- Melbourne, J., Peng, C. Y., Soifer, B. T., et al. 2011, AJ, 141, 141, doi: 10.1088/0004-6256/141/4/141
- Melbourne, J., Soifer, B. T., Desai, V., et al. 2012, AJ, 143, 125, doi: 10.1088/0004-6256/143/5/125
- Miyazaki, S., Komiyama, Y., Kawanomoto, S., et al. 2018, PASJ, 70, S1, doi: 10.1093/pasj/psx063
- Monadi, R., & Bird, S. 2022, MNRAS, 511, 3501, doi: 10.1093/mnras/stac294
- Moran, E. C., Halpern, J. P., & Helfand, D. J. 1996, ApJS, 106, 341, doi: 10.1086/192341
- Netzer, H. 2015, ARA&A, 53, 365, doi: 10.1146/annurey-astro-082214-122302
- Niida, M., Nagao, T., Ikeda, H., et al. 2016, ApJ, 832, 208, doi: 10.3847/0004-637X/832/2/208

- Noboriguchi, A., Nagao, T., Toba, Y., et al. 2019, ApJ, 876, 132. doi: 10.3847/1538-4357/ab1754
- Noll, S., Burgarella, D., Giovannoli, E., et al. 2009, A&A, 507, 1793, doi: 10.1051/0004-6361/200912497
- Pancoast, A., Brewer, B. J., Treu, T., et al. 2014, MNRAS, 445, 3073, doi: 10.1093/mnras/stu1419
- Perrotta, S., Hamann, F., Zakamska, N. L., et al. 2019, MNRAS, 488, 4126, doi: 10.1093/mnras/stz1993
- Pilbratt, G. L., Riedinger, J. R., Passvogel, T., et al. 2010, A&A, 518, L1, doi: 10.1051/0004-6361/201014759
- Poglitsch, A., Waelkens, C., Geis, N., et al. 2010, A&A, 518, L2, doi: 10.1051/0004-6361/201014535
- Pope, A., Bussmann, R. S., Dey, A., et al. 2008, ApJ, 689, 127, doi: 10.1086/592739
- Ross, N. P., Hamann, F., Zakamska, N. L., et al. 2015, MNRAS, 453, 3932, doi: 10.1093/mnras/stv1710
- Sanders, D. B., Soifer, B. T., Elias, J. H., et al. 1988, ApJ, 325, 74, doi: 10.1086/165983
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525, doi: 10.1086/305772
- Shakura, N. I., & Sunyaev, R. A. 1973, A&A, 500, 33
- Shen, Y. 2013, Bulletin of the Astronomical Society of India, 41, 61. https://arxiv.org/abs/1302.2643
- Shen, Y., Richards, G. T., Strauss, M. A., et al. 2011, ApJS, 194, 45, doi: 10.1088/0067-0049/194/2/45
- Stalevski, M., Ricci, C., Ueda, Y., et al. 2016, MNRAS, 458, 2288, doi: 10.1093/mnras/stw444
- Toba, Y., Bae, H.-J., Nagao, T., et al. 2017a, ApJ, 850, 140, doi: 10.3847/1538-4357/aa918a
- Toba, Y., & Nagao, T. 2016, ApJ, 820, 46, doi: 10.3847/0004-637X/820/1/46
- Toba, Y., Nagao, T., Strauss, M. A., et al. 2015, PASJ, 67, 86, doi: 10.1093/pasj/psv057
- Toba, Y., Nagao, T., Kajisawa, M., et al. 2017b, ApJ, 835, 36, doi: 10.3847/1538-4357/835/1/36
- Toba, Y., Yamashita, T., Nagao, T., et al. 2019, ApJS, 243, 15, doi: 10.3847/1538-4365/ab238d
- Toba, Y., Yamada, S., Ueda, Y., et al. 2020, ApJ, 888, 8, doi: 10.3847/1538-4357/ab5718
- Toba, Y., Liu, T., Urrutia, T., et al. 2022, A&A, 661, A15, doi: 10.1051/0004-6361/202141547
- Treister, E., Schawinski, K., Urry, C. M., & Simmons,
 B. D. 2012, ApJL, 758, L39,
 doi: 10.1088/2041-8205/758/2/L39
- Tremaine, S., Gebhardt, K., Bender, R., et al. 2002, ApJ, 574, 740, doi: 10.1086/341002
- Valiante, E., Smith, M. W. L., Eales, S., et al. 2016, MNRAS, 462, 3146, doi: 10.1093/mnras/stw1806
- van Dokkum, P. G. 2001, PASP, 113, 1420, doi: 10.1086/323894

- Vanden Berk, D. E., Richards, G. T., Bauer, A., et al. 2001, AJ, 122, 549, doi: 10.1086/321167
- Véron-Cetty, M. P., Véron, P., & Gonçalves, A. C. 2001, A&A, 372, 730, doi: 10.1051/0004-6361:20010489
- Vestergaard, M., & Osmer, P. S. 2009, ApJ, 699, 800, doi: 10.1088/0004-637X/699/1/800
- Vestergaard, M., & Peterson, B. M. 2006, ApJ, 641, 689, doi: 10.1086/500572
- Vietri, G., Piconcelli, E., Bischetti, M., et al. 2018, A&A, 617, A81, doi: 10.1051/0004-6361/201732335
- Villar Martín, M., Perna, M., Humphrey, A., et al. 2020, A&A, 634, A116, doi: 10.1051/0004-6361/201937086
- Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, AJ, 140, 1868,
 - doi: 10.1088/0004-6256/140/6/1868

- Wu, J., Tsai, C.-W., Sayers, J., et al. 2012, ApJ, 756, 96, doi: 10.1088/0004-637X/756/1/96
- Wu, J., Jun, H. D., Assef, R. J., et al. 2018, ApJ, 852, 96, doi: 10.3847/1538-4357/aa9ff3
- Yang, G., Boquien, M., Buat, V., et al. 2020, MNRAS, 491, 740, doi: 10.1093/mnras/stz3001
- York, D. G., Adelman, J., Anderson, Jr., J. E., et al. 2000, AJ, 120, 1579, doi: 10.1086/301513
- $\begin{aligned} & \text{Zamfir, S., Sulentic, J. W., Marziani, P., \& Dultzin, D.} \\ & 2010, \text{ MNRAS, } 403, \text{ } 1759, \end{aligned}$
 - doi: 10.1111/j.1365-2966.2009.16236.x
- Zou , F., Brandt, W. N., Vito, F., et al. 2020, MNRAS, 499, 1823, doi: 10.1093/mnras/staa2930