Dark Comets? Unexpectedly Large Nongravitational Accelerations on a Sample of Small Asteroids

DARRYL Z. SELIGMAN ^(D),¹ DAVIDE FARNOCCHIA ^(D),² MARCO MICHELI ^(D),³ DAVID VOKROUHLICKÝ ^(D),⁴ ASTER G. TAYLOR ^(D),⁵ STEVEN R. CHESLEY ^(D),² JENNIFER B. BERGNER ^(D),⁶ PETER VEREŠ ^(D),⁷ OLIVIER R. HAINAUT ^(D),⁸ KAREN J. MEECH ^(D),⁹ MAXIME DEVOGELE ^(D),¹⁰ PETR PRAVEC,¹¹ ROB MATSON,¹² SAM DEEN,¹² DAVID J. THOLEN,¹³ ROBERT WERYK ^(D),¹⁴ EDGARD G. RIVERA-VALENTÍN ^(D),¹⁵ AND BENJAMIN N. L. SHARKEY ^(D),¹⁶

¹Department of Astronomy and Carl Sagan Institute, Cornell University, 122 Sciences Drive, Ithaca, NY, 14853, USA

² Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Dr., Pasadena, CA 91109, USA

³ESA NEO Coordination Centre, Largo Galileo Galilei 1, I-00044 Frascati (RM), Italy

⁴Institute of Astronomy, Charles University, V Holešovičkách 2, CZ-18000 Prague 8, Czech Republic

⁵Dept. of Astronomy and Astrophysics, University of Chicago, 5640 S Ellis Ave, Chicago, IL 60637, USA

⁶ University of Chicago, Department of the Geophysical Sciences, Chicago, IL 60637, USA

⁷Harvard-Smithsonian Center for Astrophysics, Minor Planet Center, 60 Garden Street, Cambridge, MA 02138, USA

⁸ European Southern Observatory, Karl-Schwarzschild-Strasse 2, Garching bei München, D-85748, Germany

⁹Institute for Astronomy, University of Hawaii, 2680 Woodlawn Dr., Honolulu, HI 96822, USA

¹⁰Arecibo Observatory, University of Central Florida, HC-3 Box 53995, Arecibo, PR 00612, USA

¹¹Astronomical Institute, Academy of Sciences of the Czech Republic, Fričova 1, CZ-25165 Ondřejov, Czech Republic

 $^{12} Independent \ Researcher$

¹³Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822, USA

¹⁴Physics and Astronomy, The University of Western Ontario, 1151 Richmond Street, London ON N6A 3K7, Canada

¹⁵ Johns Hopkins University Applied Physics Laboratory, 11100 Johns Hopkins Road, Laurel, MD 20723, USA.

¹⁶Lunar and Planetary Laboratory, University of Arizona, 1629 E University Blvd, Tucson, AZ 85721, USA

ABSTRACT

We report statistically significant detections of non-radial nongravitational accelerations based on astrometric data in the photometrically inactive objects 1998 KY₂₆, 2005 VL₁, 2016 NJ₃₃, 2010 VL₆₅, 2016 RH_{120} , and 2010 RF_{12} . The magnitudes of the nongravitational accelerations are greater than those typically induced by the Yarkovsky effect and there is no radiation-based, non-radial effect that can be so large. Therefore, we hypothesize that the accelerations are driven by outgassing, and calculate implied H_2O production rates for each object. We attempt to reconcile outgassing induced acceleration with the lack of visible comae or photometric activity via the absence of surface dust and low levels of gas production. Although these objects are small and some are rapidly rotating, surface cohesive forces are stronger than the rotational forces and rapid rotation alone cannot explain the lack of surface debris. It is possible that surface dust was removed previously, perhaps via outgassing activity that increased the rotation rates to their present day value. We calculate dust production rates of order $\sim 10^{-4}$ g s⁻¹ in each object assuming that the nuclei are bare, within the upper limits of dust production from a sample stacked image of 1998 KY₂₆ of $\dot{M}_{\text{Dust}} < 0.2 \text{ g s}^{-1}$. This production corresponds to brightness variations of order $\sim 0.0025\%$, which are undetectable in extant photometric data. We assess the future observability of each of these targets, and find that the orbit of 1998 KY_{26} — which is also the target for the extended Hayabusa2 mission — exhibits favorable viewing geometry before 2025.

Keywords: Asteroids (72) — Comets (280)

1. INTRODUCTION

A typical — albeit simplistic — classification of Solar System small bodies is to categorize the populations based on volatile-driven activity. In this classical pic-

Corresponding author: Darryl Z. Seligman dzs9@cornell.edu

ture, comets are defined as icy objects that produce dusty comae and presumably formed in distant regions of the Solar System, while asteroids lack volatiles due to prolonged exposure to solar irradiation. However, recent advances have demonstrated that this simplistic classification may not be an accurate depiction of the census of small bodies within the Solar System. A subset of objects on cometary orbits lack detectable activity, while some asteroids have displayed volatile-induced comae. These intriguing continuum objects could offer insights into little understood processes such as cometary fading (Brasser & Wang 2015) and volatile delivery to terrestrial planets (Chyba 1990; Owen & Bar-Nun 1995; Albarède 2009).

It is generally thought that asteroids can be grouped into classes based on their sizes. The typical timescale for a ~ 10 km scale asteroid to experience a catastrophic impact is approximately the age of the Solar System (Bottke et al. 2005, 2015). It is therefore believed that asteroids larger than ~ 10 km are intact primordial remnants, while smaller objects with diameter between 200 m to ~ 10 km are re-accumulated rubblepiles held together by self-gravity (Harris 1979; Harris & Burns 1979; Harris 1996; Walsh 2018). As evidence, the ~ 17 km equivalent diameter asteroid Eros visited by NEAR-Shoemaker displayed geologic properties consistent with being a damaged but intact primordial remnant (Cheng et al. 2002). Notable examples of extremely porous rubble-pile asteroids are (25143) Itokawa (Fujiwara et al. 2006), (162173) Ryugu (Watanabe et al. 2019), and (101955) Bennu (Barnouin et al. 2019).

A subset of asteroids are known to exhibit activity, also called active asteroids (Jewitt 2012; Hsieh 2017; Jewitt & Hsieh 2022). Almost all active asteroids known to date have diameters consistent with being rubble-piles, with the exception of the large objects (1) Ceres and (493) Griseldis. A subset of active asteroids are the Main Belt Comets (MBCs) — objects that reside in the main asteroid belt between Mars and Jupiter and display cometary activity driven by the sublimation of volatiles (Hsieh & Jewitt 2006). The first MBC discovered was Comet 133P/(7968)Elst-Pizarro (Elst et al. 1996; Boehnhardt et al. 1996; Toth 2000; Hsieh et al. 2004) and several others have 238P/Read, 259P/Garradd, since been identified: 288P/(300163) 2006 VW139, 313P/Gibbs, 324P/La Sagra, 358P/PANSTARRS, 107P/(4015)Wilson-Harrington, and 433P/(248370) 2005 QN173 (for images, see Figure 13 in Jewitt & Hsieh 2022).

Targeted, homogeneous searches for MBCs imply occurrence rates for these objects within the total population between < 1/500 and $\sim 1/300$ (Sonnett et al. 2011; Bertini 2011; Snodgrass et al. 2017; Ferellec et al. 2022). Indirect measurements of activity can come from anisotropic mass-loss-driven nongravitational accelerations (Whipple 1950, 1951). For example, Hui & Jewitt (2017) reported statistically significant nongravitational accelerations in the active asteroids 313P/Gibbs, 324P/La Sagra, and (3200) Phaethon.

Non-sublimation effects such as impacts (Snodgrass et al. 2010) and rotational effects (Jewitt et al. 2014) can also cause activity in asteroids. The ~ 6 km sized object (3200) Phaethon was first determined to be active when its association with the Geminid meteoroid stream was identified (Gustafson 1989; Williams & Wu 1993). Subsequent observations of (3200) Phaethon near its perihelion revealed a small tail with micron sized dust production rates of $\sim 3 \text{ kg s}^{-1}$ (Jewitt & Li 2010; Jewitt et al. 2013; Li & Jewitt 2013; Hui & Li 2017). Because these production rates are not large enough to explain the meteoroid stream, it has been suggested that processes such as repeated thermally induced stresses (Jewitt & Li 2010), sublimation of minerologically bound sodium (Masiero et al. 2021), rotational effects (Ansdell et al. 2014; Nakano & Hirabayashi 2020) and geometric effects (Hanuš et al. 2016; Taylor et al. 2019) also contribute to the activity of (3200) Phaethon. The Japan Aerospace Exploration Agency (JAXA) DESTINY+ mission will visit (3200) Phaethon and is expected to launch in 2024 (Arai et al. 2021). DESTINY+ will be equipped with a mass spectrometer which would detect elemental compositions of dust released by the object (Krüger et al. 2019).

(101955) Bennu is another intriguing case of an active asteroid. The OSIRIS-REx spacecraft observed repeated periods of particle ejection from the ~ 500 m rubble pile (Lauretta et al. 2019; Hergenrother et al. 2019). The average mass loss rate of dust was measured to be only $\dot{M}_{\rm Dust} \sim 10^{-4}$ g s⁻¹ (Hergenrother et al. 2020), but the source of the activity is unclear (Bottke et al. 2020; Molaro et al. 2020; Chesley et al. 2020).

A somewhat related but distinct family of minor bodies are the *inactive* comets. For example, "Manxcomets" are objects with Long Period Comet (LPC) trajectories that exhibit little or no cometary activity. The Manx-comet C/2014 S3 (PANSTARRS) displayed levels of cometary activity that were 5-6 orders of magnitude less than typical comets, and spectral features similar to S- type asteroids (Meech et al. 2016). The origin of the Manx-comets are unclear, because Jupiter scatters objects within the H₂O snowline primarily into the interstellar medium and not into the Oort cloud (Hahn & Malhotra 1999; Shannon et al. 2015). Other inactive comets are Damocloids (named after the first object (5335) Damocles; Asher et al. 1994; Jewitt 2005) and Asteroids on Cometary Orbits (ACOs), defined as inactive objects on Halley-type comet orbits or Jupiter Family Comet (JFC) orbits, respectively. Damocloids and ACOs are believed to be comets that have little or no activity due to cometary fading (Wang & Brasser 2014; Brasser & Wang 2015), or the depletion of volatiles or mantling (Podolak & Herman 1985; Prialnik & Bar-Nun 1988). Observed spectral features and surface colors of Damocloids and ACOs indicate that they are likely extinct or dormant comets (Jewitt 2005; Licandro et al. 2018).

The recently discovered population of interstellar objects also appear to exhibit a continuum of apparent activity. The first interstellar object 11/'Oumuamua was detected on UT 2017 October 19 (Williams et al. 2017) and exhibited a mixture of cometary and asteroidal properties. On the other hand, the 0.2-0.5 km radius scale (Jewitt et al. 2020) 2I/Borisov displayed a visible cometary tail (Jewitt & Luu 2019; Guzik et al. 2020; Kim et al. 2020; Cremonese et al. 2020; Hui et al. 2020; Bodewits et al. 2020; Yang et al. 2021). The ~ 120 m diameter 11/'Oumuamua displayed an extreme light curve amplitude (Knight et al. 2017; Bolin et al. 2018; Fraser et al. 2018; McNeill et al. 2018; Belton et al. 2018; Mashchenko 2019; Drahus et al. 2018), reddened color (Masiero 2017; Fitzsimmons et al. 2018; Bannister et al. 2017; Ye et al. 2017) and incoming trajectory consistent with the local standard of rest (Mamajek 2017; Gaidos et al. 2017; Feng & Jones 2018; Almeida-Fernandes & Rocha-Pinto 2018; Hallatt & Wiegert 2020; Hsieh et al. 2021a). Of particular interest and relevance to this paper, 'Oumuamua exhibited no evidence for an extended dust coma in deep stacked composite images (Meech et al. 2017; Jewitt et al. 2017). Moreover, it exhibited no obvious infrared fluorescence of carbon-based outgassing species (e.g., CO or CO_2) based on observations obtained with the Spitzer Space Telescope (Trilling et al. 2018). However, the trajectory of the object was affected by a significant nongravitational acceleration (Micheli et al. 2018). Micheli et al. (2018) argued that the most likely cause of this acceleration — which was primarily in the radial direction opposite the direction of the sun — was cometary outgassing. The physical properties of the first two interstellar interlopers are readily summarized; see Jewitt & Seligman (2022) and Moro-Martín (2022) for recent reviews.

Recently, Farnocchia et al. (submitted 2022) presented observations that revealed that the object 2003 RM exhibited a significant transverse nongravitational acceleration incompatible with the Yarkovsky effect. However, this object, like 'Oumuamua, lacked cometary



Figure 1. Nongravitational accelerations measured in asteroids versus their absolute magnitude H. Measured radial nongravitational acceleration A_1 are due to solar radiation pressure (pink) and transverse acceleration A_2 are due to the Yarkovsky effect (blue). The out-of-plane nongravitational acceleration A_3 measured in the objects presented in this paper are shown in orange diamonds. The blue and pink dashed lines show lines of best fit, assuming that the acceleration is proportional to the inverse of the nuclear size, and computed using the relationship in Equation 2 assuming an albedo p = 0.1. The log-scaled intercepts are -15.858 for the radiation pressure, and -17.440 for the Yarkovsky effect. Dark comet candidates with anomalous A_2 values, 2003 RM and 2006 RH₁₂₀, are shown in purple diamonds.

activity clearly visible in photometric images. They found that outgassing can cause the acceleration and still escape photometric detection. In this paper, we report identification of similar nongravitational accelerations in five other small bodies on asteroid orbits.

2. DETECTIONS OF NONGRAVITATIONAL ACCELERATION

2.1. Possible Nongravitational Perturbations on Small Bodies

The trajectories of small bodies are typically estimated from observational datasets, primarily optical astrometry and sometimes radar delay and Doppler measurements. As the observational arcs extend in time and the accuracy of the data improves, the resulting orbits become better constrained. Therefore, the requirements on the fidelity of the force model to compute the orbit become more stringent.

It has long been established that the motion of comets can be significantly perturbed by nongravitational accelerations due to outgassing in addition to typical gravitational forces. Marsden et al. (1973) introduced a parametric model where the nongravitational acceleration is described as:

$$\mathbf{a}_{\rm NG} = \left(A_1\hat{\mathbf{r}} + A_2\hat{\mathbf{t}} + A_3\hat{\mathbf{n}}\right)g(r)\,.\tag{1}$$

In Equation 1, $\hat{\mathbf{r}}$, $\hat{\mathbf{t}}$, $\hat{\mathbf{n}}$ are the orbital radial, transverse, and out-of-plane directions, g(r) is a function based on the H₂O sublimation profile capturing the dependence on the heliocentric distance r, and A_1 , A_2 , and A_3 are free parameters that give the acceleration components that the comet would experience at a heliocentric distance of r = 1 au.

For asteroids, the two main nongravitational perturbations are the Yarkovsky effect (Vokrouhlický et al. 2015) and solar radiation pressure (Vokrouhlický & Milani 2000). By setting $q(r) = (1 \text{ au}/r)^2$, these two perturbations can be modeled as purely transverse $A_2g(r)$ and purely radial $A_1q(r)$ accelerations using the Marsden et al. (1973) formalism (e.g., Farnocchia et al. 2015). Despite being smaller than those on comets, these two perturbations can cause detectable deviations from a gravity-only trajectory for sufficiently long observational data arcs. In particular, the Yarkovsky effect has been detected on more than 200 near-Earth asteroids (e.g., Greenberg et al. 2020) shown in blue points in Figure 1. Solar radiation pressure is a radial acceleration and is therefore less effective at producing significant orbital deviations. As such, it has been measured only on a handful of small asteroids (MPEC 2008-D12,¹ Micheli et al. 2012, 2013, 2014; Mommert et al. 2014a,b; Farnocchia et al. 2017; Fedorets et al. 2020) shown in pink points in Figure 1. As a testament of the fidelity of the modeling of the force model for asteroids, including these nongravitational forces, Farnocchia et al. (2021) were able to reconstruct the trajectory of asteroid (101955) Bennu by matching ground-based optical and radar astrometry from 1999 to 2018 as well as meterlevel ranging measurements from the OSIRIS-REx mission proximity operations from January 2019 to October 2020.

2.2. Asteroids with Excess Nongravitational Acceleration



In the accompanying paper Farnocchia et al. (submitted 2022), we report the detection of an anomalously large nongravitational acceleration in the object 2003 RM. In that paper, we conclude that outgassing could have caused the acceleration of 2003 RM while not producing a detectable signature in photometric data. Further, Chesley et al. (2016) reported that the object 2006 RH120, which was temporarely captured by the Earth for about a year starting in June 2006 (Granvik et al. 2012), exhibited a transverse nongravitational acceleration that was inconsistent with typical forces observed acting on asteroids. In this paper, we report the detection of out-of-plane nongravitational accelerations in a sample of five other objects that we measured from a combination of serendipitous (and uncoordinated) astrometric observations: 1998 KY_{26} , 2005 VL_1 , 2016 NJ_{33} , 2010 VL_{65} , and 2010 RF_{12} , all identified as point source asteroidal objects with no extended dust coma or evidence for outgassing. We also detect significant out-ofplane perturbations for 2006 RH_{120} . There have been no reports of clear detection of cometary activity for any of these small objects to the MPC to date, and they are all classified as asteroids.

In December 2020, we observed asteroid 1998 KY_{26} — a potential Yarkovsky detection candidate (Vokrouhlický et al. 2000) and the target of the Havabusa 2 extended mission (Hirabayashi et al. 2021) — with the VLT (MPEC 2020-X181)² with the purpose of improving the measurement of the orbit. At the same time, we also submitted two previously unreported observations from Mauna Kea in 2002 to the Minor Planet Center. While the Yarkovsky effect was included in the fit and detected, the VLT observations showed a -0.2'' bias in declination that was larger than the astrometric uncertainties. The bias was confirmed in additional observations from La Palma (MPEC 2021-A42)³ and Mauna Kea (MPEC 2021-G127)⁴. To ensure that the cause for this bias was not problematic data during the discovery apparition in 1998, we remeasured observations from the Modra Astronomical and Geophysical Observatory and Ondřejov, which, together with radar (Ostro et al. 1999), were the only observations from 1998 we included in the fit. Despite our dataset revision, the bias still persisted. Adding solar radiation pressure through the A_1 parameter was unfruitful, but adding an out-ofplane acceleration $A_3(1 \text{ au}/r)^2$ removed the bias in the 2020 data.

 $^{\rm 4} \rm \ https://www.minorplanetcenter.net/mpec/K21/K21GC7.html$

 $^{^{2}\ \}rm https://www.minorplanetcenter.net/mpec/K20/K20XI1.html$

 $^{^{3} \ \}rm https://www.minorplanetcenter.net/mpec/K21/K21A42.html$

Subsequently, we observed 2005 VL₁ in 2021 (MPEC 2021-X95)⁵ and 2016 NJ₃₃ in 2022 (MPECs 2022-N88, 2022-O09)⁶⁷ as Yarkovsky candidates. These observations were obtained with the 4.3 m Lowell Discovery Telescope in Arizona. Again, the collected data revealed a bias that could only be removed by adding an out-of-plane acceleration.

Another case arose when we noticed the similarity between the orbits of 2010 VL₆₅⁸ and 2021 UA12⁹. The objects can be readily linked by fitting both datasets together, but only if A_3 is estimated as part of the fit.

The last case is 2010 RF_{12} , which we observed in 2022 from Mauna Kea with the Canada France Hawaii Telescope and from Cerro Paranal with the VLT with the purpose of ruling out impact solutions detected by impact monitoring systems such as Sentry¹⁰ and ESA's own system¹¹. The observations we collected, in combination with an archival Mauna Kea set of images from 2011 taken with the Subaru Telescope (MPEC 2022-S77)¹² once again revealed a bias that could only be removed by estimating A_3 while fitting the data.

Given these results, we decided to review the fit for 2006 RH_{120} and found that in that case, there is also significant detection of an out-of-plane acceleration in addition to the radial and transverse ones.

In Table 1, we list each component, A_1 , A_2 , and A_3 of the best-fit nongravitational accelerations and their associated uncertainties. In order to avoid underestimating the uncertainty in the A_1 , A_2 , and A_3 parameters, we estimated all three parameters even when some of them (especially A_1) are not significantly detected. For each of these fits, we adopted $g(r) = (1 \text{ au}/r)^2$. As examples, we show the right ascension and declination residuals of the astrometric fits with and without nongravitational accelerations for 2010 VL₆₅ and 2005 VL₁ in Figure 2. The improvement in the fit is evident from the figures. For VL₆₅, the gravity-only fit results in residuals as large as 10 arcsec. For 2005 VL₁, it is clear how the bias in 2021 (but also systematic errors in 2005) is removed by adding nongravitational accelerations.

In Figure 1, we show the best-fit A_2 and A_3 parameter for the non-radial nongravitational acceleration for each object. We compare the estimated A_2 and A_3 to the measured nongravitational accelerations in asteroids. We show both A_1 , which is related to solar radiation pressure, and A_2 , which is related to the Yarkovsky effect. Nominal by-eye fits of the acceleration, being inversely proportional to the nuclear radius computed using Equation 2, are shown. There is a linear correlation between $\log |A_2|$ and H (blue dashed line), as is expected from the Yarkovsky relation (Vokrouhlický et al. 2015). The scatter around the mean is presumably due to different albedo, density, obliquity, or surface thermal inertia values. As a rule of thumb, the $\log |A_1|$ vs H (pink dashed line) again shows a linear trend with the same slope but an approximately order of magnitude larger normalization (Vokrouhlický & Milani 2000). The nonradial component of the detected accelerations appears to be more consistent with the extrapolation of the A_1 values. However, solar radiation pressure is mostly radial, and only a minor fraction would project in the outof-plane direction (Vokrouhlický & Milani 2000). Specifically, Vokrouhlický & Milani (2000) demonstrated that small off-radial components of nongravitational accelerations could emerge from radiation pressure because of variable albedo or non-spherical shapes. This suggests that these accelerations are inconsistent with radiation effects.

Based on this analysis, we conclude that there are significant out-of-plane accelerations for all of these objects. If the out-of-plane accelerations in these objects are in fact caused by outgassing, it is possible that these are manifestations of nearly-polar jets with spin axis orthogonal to the orbit plane.

The radial A_1 values are all consistent with zero acceleration, with the possible exception of 2016 NJ_{33} . Moreover, the A_2 values are much smaller in magnitude than the A_3 values and are generally consistent with the Yarkovsky effect. In the case of 2003 RM and 2006 RH120, the A_2 values are anomalously high and presumably due to outgassing (Chesley et al. 2016). Therefore, in the rest of the paper we focus on A_3 only. We note, however, that if high latitude jets are in fact causing the nongravitational accelerations, this would be somewhat distinct from typical comets that exhibit outgassing events isotropically over their surfaces. Moreover, incident stellar irradiation and subsequent surface temperature variations should not only affect the outof-plane acceleration. This analysis can be generalized to larger accelerations if A_1 were to be statistically detected with longer data arcs.

⁵ https://www.minorplanetcenter.net/mpec/K21/K21X95.html

 $^{^{6}\} https://www.minorplanetcenter.net/mpec/K22/K22N88.html$

⁷ https://www.minorplanetcenter.net/mpec/K22/K22O09.html

 $^{^{8}}$ https://ssd.jpl.nasa.gov/tools/sbdb_lookup.html#/?sstr= 2010vl65

 $^{^{9}}$ https://ssd.jpl.nasa.gov/tools/sbdb_lookup.html#/?sstr=2021ua12

 $^{^{10}\; \}rm https://cneos.jpl.nasa.gov/sentry/$

 $^{^{11}\;\}rm https://neo.ssa.esa.int/risk-list$

 $^{^{12}\;\}mathrm{https://www.minorplanetcenter.net/mpec/K22/K22S77.html}$



Figure 2. The (signed) residuals of the astrometric fits (in arcseconds) for the right ascension and declination of 2010 VL_{65} (top) and 2005 VL_1 (bottom) with (pink points) and without (blue points) nongravitational accelerations. The gravity-only solution does not allow for outlier rejection in the astrometric fits.



Figure 3. The location in semi-major axis and eccentricity space of the currently known active asteroids (pentagons, Table 3) and candidate dark comets (diamonds Table 1). Active asteroids are color coded based on their activity sources: sublimation driven activity (blue), non-sublimation driven activity (pink), and unknown activity source (unfilled). The semi-major axis of Mars and the Earth and orbits with perihelia equal to that of Mars and the Earth are indicated with dashed lines.

Table 1 summarizes the observed orbital properties, absolute magnitudes and rotational periods (where measured) for each of these objects. We use the following equation (Pravec & Harris 2007) to estimate the sizes for all objects except 1998 KY₂₆, 2003 RM and 2006 RH₁₂₀

$$2R_{\rm Nuc} = \left(\frac{1329}{\sqrt{p}}\right) 10^{-0.2\,H},\tag{2}$$

where H is the absolute magnitude, p is the geometric albedo and R_{Nuc} is the radius in kilometers. In Table 1, we assume p = 0.1 to estimate the size of objects where this value has not been previously measured. The trajectories of the objects are unremarkable. In Figure 3, we show the location of these candidate "dark comets" as well as the location of all of the currently known active asteroids, whose properties are listed in Table 3.

2.3. Lack of Cometary Activity Detection

None of the objects studied in this paper have reported cometary activity. Out of all of our candidate targets, 1998 KY_{26} was the only one for which we had meaningful data that could provide limits on the dust production. The 1998 KY_{26} images were obtained using the using the Unit Telescope 1 of the ESO Very Large Telescope (VLT) on Mount Paranal, Chile, with the FOcal Reducer and low dispersion Spectrograph 2 (Appenzeller et al. 1998, FORS2). The instrument was used without filter (white light), in order to reach the deepest possible limiting magnitude on asteroids and dust. FORS2 was equipped with a mosaic of two MIT/LL CCD detectors. of which only the main chip, A, or CCID20-14-5-3, was used. The CCD was read with a 2×2 binning, resulting in a 0.252'' on sky. The de-biased and flat-fielded images were aligned on the position of the target. In Figure 4, we show a stack of the images from December 2020 where no background object was too close from the object. This image contains 120 exposures of 30 s each, resulting in a 3600 s exposure.

The object was rather faint (26.1 for the first set, 25.5)for the second), requiring that the astrometric measurements be performed on stacks. The S/N in the individual images is barely above 1. This implies that the photometry would also require stacks. The object is the blob at the centre, outlined with a red circle. Although the object is very faint, the stacked images produced a solid detection with a combined $S/N \sim 12$. Furthermore, 1998 KY_{26} is clearly visible in various sub-stacks, confirming without doubt the reality of the image and its identification with the object (the image is $64" \times 64"$). There is no evidence for an extended tail in the deep image: the variations observed in the sky around the object can be identified with residual of the various background objects passing behind the object. In order to further quantify the lack of dust around the target, we show in Figure 5 its photometric profile.

The selection of the images composing the stack used to generate the profile ensured that no background object would contaminate the area of the profile. We obtained it by integrating the instrumental fluxes converted to surface brightnesses in a set of concentric circular apertures centered on the object. We use a conversion to magnitudes with a zero point of 27.8 (from Hainaut et al. 2021). To scale the brightness of the object for comparison, we used the profile of a field star. This analysis does not yield a stringent constraint because the noise and the background dominate very quickly. Because this object is so faint, the quantity of dust that could be undetected in the image is not well constrained. We estimate a coma mass upper limit of ~ 3 kg of dust by integrating the corresponding flux from 0.5'' to 2''(assuming 1 micron sized dust grains, with an albedo p = 0.2, density $\rho = 3000$ kg m³). This corresponds to a magnitude for the dust of 24.0, which would be a 5σ detection from 0.5 to 2" from the object. The un-



Figure 4. A stacked and cleaned image of 1998 KY₂₆ with VLT which spans 2 sets of 60 exposures, each of which is 30 s. The resulting temporal coverage is ~ 1 h in Dec 2020. The image is $64 \times 64^{\circ}$. This is a linear greyscale from 0 to 4 adu/s/pix. 1998 KY₂₆ is faint but visible in the final image and indicated with the red circle. There is no evidence for an extended cometary tail.



Figure 5. A standard profile analysis for dust activity in the deep stacked and cleaned image with VLT of 1998 KY₂₆ shown in Figure 4. The upper limit of dust present in the image is ~ 3 kg. The solid line indicates the profile of our chosen field star scaled to the brightness of the object. The solid points indicate the integrated flux in concentric rings centered on the target. This analysis assumes that the dust is composed of 1 micron sized dust grains, with an albedo p = 0.2 and density $\rho = 3000$ kg m³, with ~ 1 order of magnitude uncertainty. The x-axis increases to the right.

certainty is ~ 1 order of magnitude because of all the assumptions required to convert from the magnitudes to the mass of dust (grain size, albedo, density). The corresponding upper limit on the dust production rate is $\dot{M}_{\rm Dust} < 0.2$ g s⁻¹. However, as discussed in the next section, outgassing could still explain the acceleration even if the upper limit was orders of magnitude higher.

3. IMPLIED PRODUCTION RATES

Based on the results presented in the previous section, it is clear that no mass loss has been detected in any of these dark comet candidates. However, the magnitudes of the nongravitational accelerations are inconsistent with being caused by the Yarkovsky effect or radiation pressure. Therefore, for the remainder of this paper, we hypothesize that the non-radial nongravitational accelerations are caused by outgassing. We calculate the implied production rates of H_2O using only the dominant, nonradial accelerations.

The production rate Q(X) for a given species denoted by X may be calculated using

$$Q(X) = \left(\frac{M_{\rm Nuc}}{m_{\rm X}}\right) \left(\frac{|A_i|}{v_{\rm Gas}\zeta}\right).$$
(3)

In Equation 3, $M_{\rm Nuc}$ is the mass of the nucleus, and $m_{\rm X}$ and $v_{\rm Gas}$ are the mass and velocity of the outgassing species. A_i is the dominant component of the nongravitational acceleration used in the calculation. The variable ζ indicates the isotropy of the outflow, where ζ = 1 corresponds to a collimated outflow and ζ = 0.5 corresponds to an isotropic hemispherical outflow. The velocity of the gas can be related to the temperature of the outflow, $T_{\rm Gas}$, using

$$v_{\rm Gas} = \left(\frac{8k_B T_{\rm Gas}}{\pi m_{\rm X}}\right)^{1/2}.$$
 (4)

We calculate the implied production rates of H₂O outgassing using Equations 3 and 4 at perihelion (Table 1). We assume that $\zeta = 1$ and that the nuclei of the objects are spherical with radii listed in Table 1, which in turn are calculated assuming an albedo of 0.1 and a bulk density of $\rho_{\text{Bulk}} = 1 \,\text{g}\,\text{cm}^{-3}$. These production rates, $\sim 10^{21}$ molec s⁻¹, are several orders of magnitude lower than typical cometary production rates, which are on the order of $\sim 10^{26}$ molec s⁻¹ (see Tables 1-3 in Harrington-Pinto et al. 2022). The implied mass production of H₂O of 1998 KY₂₆ is ~ 0.03 g s⁻¹, compared to $\lesssim 1 \text{ kg s}^{-1}$ for MBCs and $\sim 10^2 - 10^3 \text{ kg}$ s^{-1} for comets (Jewitt 2012; Jewitt & Hsieh 2022). If 1998 KY_{26} has been outgassing continuously at this rate since its discovery, the total mass lost would be $\sim 10^7$ g, $\sim 0.1\%$ of its total mass. The analogous calculations for

the other dark comet candidates yield similarly small mass loss fractions, implying that outgassing is an allowable mechanism. The production of MBCs has not led to detectable gas production (Snodgrass et al. 2017), except for a tentative detection of one candidate by Ferellec et al. (2022).

It should be noted that the case of 2016 NJ₃₃ requires a $\sim 3\sigma$ significant A_1 radial acceleration to fit the astrometric data. We calculate the implied production rate using only the A_3 acceleration. However, if the A_1 is as large as the fit suggests, then the implied production of H₂O would be slightly higher.

The characteristic mass-loss timescale τ_M for which this level of activity is sustainable is approximately given by

$$\tau_M \simeq M_{\rm Nuc} \left/ \left(\frac{dM_{\rm Nuc}}{dt} \right) \simeq \left(\frac{v_{\rm Gas} \zeta}{|A_i|} \right)$$
(5)

A scaled relationship version of Equation 5 gives

$$\tau_M = 2.8 \times 10^4 \,\mathrm{yr} \,\left(\frac{v_{\mathrm{Gas}}}{350 \,\mathrm{m/s}}\right) \,\left(\frac{2 \times 10^{-11} \,\mathrm{au/day}^2}{|A_i|}\right). \tag{6}$$

The orbits in our sample are similar to those of typical evolved NEOs and may have had long residency times. From Equation 6, it is evident that these objects could not have been outgassing at these inferred rates for longer than $\sim 10^4$ yrs. Therefore, if these objects are outgassing, then they either (i) were not outgassing at this rate in the recent past or (ii) were only recently emplaced on these orbits. Objects with the trajectories similar to those of our sample dark comet candidates are thought to evolve from the main belt. Specifically, there is currently no known dynamical pathway to inject these objects from the JFC or LPC populations (see, e.g., Figures 4-6 in Granvik et al. 2018). Moreover, these objects all have the highest probability of being injected onto their current trajectories from the ν_6 region in the innermost part of the asteroid belt. If our interpretation of the nongravitational acceleration based on outgassing is correct, it might have interesting population-wise implications on very small objects in the inner main belt. Although outside of the scope of this paper, future investigation of the dynamical history of these objects would be informative.

4. POSSIBLE EXPLANATION FOR LACK OF VISIBLE COMA

4.1. Cohesive Versus Rotational Forces For Surface Dust Retention

The objects in our sample are between $\sim 3 - 15$ m in radius, smaller than typical cometary nuclei that have been measured (Jewitt & Seligman 2022). Because small objects tend to rotate faster, surface dust will experience stronger rotational forces. However, when considering typical cohesive forces on the surface regoliths of asteroids, it is evident that the rapid rotation of these objects is not sufficient to explain the lack of comae.

Consider a particle at the equator and on the surface of a spherical nucleus with a radius $R_{\rm Nuc}$ and a uniform density $\rho_{\rm Bulk}$. In the absence of cohesive surface forces, the particle will be removed if its velocity is greater than the escape velocity, $v_{\rm Esc}$, which is given by

$$v_{\rm Esc} = R_{\rm Nuc} \sqrt{\frac{8\pi}{3}} \, G\rho_{\rm Bulk} \, . \tag{7}$$

A scaled version of the same equation gives the following relationship:

$$v_{\rm Esc} = 7.48 \,\mathrm{cm/s} \left(\frac{R_{\rm Nuc}}{1 \,\mathrm{km}}\right) \left(\frac{\rho_{\rm Bulk}}{1.0 \,\mathrm{g \, cm^{-3}}}\right)^{1/2}.$$
 (8)

The rotational velocity $v_{\rm Rot}$ at the surface of a spherical nucleus rotating with rotational period $P_{\rm Rot}$ is $v_{\rm Rot} = 2\pi R_{\rm Nuc}/P_{\rm Rot}$. If the rotational velocity is greater than the escape velocity — if $v_{\rm Rot} > v_{\rm Esc}$ — then cohesionless particles will not remain on the surface. It is critical to note that this criteria only holds at the equator, and is diminished at higher latitudes. This criterion reduces to the following expression,

$$P_{\rm Rot} < \sqrt{\frac{3\pi}{2 \, G \rho_{\rm Bulk}}} \,. \tag{9}$$

The critical rotational period, $P_{\rm Crit}$ — analogous to the rotation limit for a cohesion-less rubble-pile object (Harris 1979; Harris & Burns 1979; Harris 1996; Warner et al. 2009) — is given by

$$P_{\rm Crit} = 2.33 \,\mathrm{h} \left(\frac{1 \,\mathrm{g \, cm^{-3}}}{\rho_{\rm Bulk}} \right)^{1/2} \,.$$
 (10)

If an object has a rotational period less than the critical period, $P_{\rm Rot} < P_{\rm Crit}$, particles without cohesive forces will not be retained on the surface. It has been established that the rotation periods of comets (when measured) are slower than those of asteroids (Binzel et al. 1992; Jewitt 2021; Jewitt & Hsieh 2022). In Figure 6 we show the rotational period vs perihelion for all comets for which the rotational period of the nucleus has been measured. The comets all have slow nuclear rotational periods and therefore weak rotational forces operating on their surfaces. Clearly, 1998 KY₂₆, 2006 RH₁₂₀ and 2016 NJ₃₃ are rotating faster than this critical period. The comet data shown in Figure 6 is listed in Table 2, with references for the cometary rotation period measurements.

However, it is worth noting that smaller dust may be retained on the surface by cohesion given by molecular forces, or some electrostatic interaction (Rozitis et al. 2014). In fact — and somewhat counterintuitively smaller objects tend to have more dominant cohesive forces (Scheeres et al. 2010; Sánchez & Scheeres 2020). The rotational force acting on the equatorial particle is $f_{\rm Rot} = m_{\rm D} R_{\rm Nuc} \omega_{\rm Nuc}$ (where $\omega_{\rm Nuc}$ is rotational frequency of the comet nucleus and $m_{\rm D}$ is the mass of the dust grain). The cohesive forces can be estimated as $f_{\rm Coh} = \sigma_{\rm C} \Delta A_{\rm D}$, where $\sigma_{\rm C}$ is the strength of the regolith and $\Delta A_{\rm D}$ is the surface area of the grain (as in Section 2 in Sánchez & Scheeres 2020). Typical values of $\sigma_{\rm C}$ are $\sim 0 - 100$ Pa (Hirabayashi et al. 2014). Ignoring the effects of self gravity, the cohesive forces will dominate over the rotational forces when the fraction $f_{\text{Coh}} > f_{\text{Rot}}$. Assuming that $m_{\rm D} \simeq \rho_{\rm D} R_{\rm D}^3$ and $\Delta A_{\rm D} \simeq R_{\rm D}^2$ for grain size $R_{\rm D}$ and density $\rho_{\rm D}$, the ratio of cohesive to rotational forces can be written as:

$$\frac{f_{\rm Coh}}{f_{\rm Rot}} = 3283 \left(\frac{\sigma_{\rm C}}{10\,{\rm Pa}}\right) \left(\frac{1\,{\rm g\,cm^{-3}}}{\rho_{\rm D}}\right) \\
\left(\frac{1\,{\rm mm}}{R_{\rm D}}\right) \left(\frac{1\,{\rm km}}{R_{\rm Nuc}}\right) \left(\frac{P_{\rm Rot}}{1\,{\rm hr}}\right)^2.$$
(11)

In Figure 7, we show a 2-dimensional histogram of the diameter and rotational period of all asteroids for which these properties have been measured. The diameter ranges for which asteroids are believed to be rubble piles and primordial remnants as outlined in Walsh (2018) are indicated. We also show the location of all of the currently known active asteroids. It is evident that 1998 KY_{26} and 2016 NJ_{23} — and presumably the remaining candidate dark comets — are distinct at the population level from active and inactive asteroids. The dashed line shows the critical rotation period given by Equation 10 and the dotted line shows where the cohesive forces equal the rotational forces computed with Equation 11, assuming mm-sized dust grains, $\sigma_{\rm C}$ = 10 Pa, and $\rho_{\rm D} = 1 \text{ g cm}^{-3}$. It is clear that although the candidate dark comets rotate rapidly, typical cohesive forces will retain dust regolith on the surface. Therefore, the rapid rotation periods alone cannot explain the lack of detectable dust activity for these objects.

4.2. Dependence of Spin-Up Timescale on Object Size

It is straightforward to show that, for a given sublimation torque, smaller objects spin up faster (the same holds for the radiation induced torques, aka the YORP effect, which are however less important in the case of



Figure 6. Nuclear rotational periods of comets for which this values has been measured versus their perihelia. The nuclear diameter is shown in color for objects where this quantity has been measured. The location of 1998 KY₂₆, 2006 RH₁₂₀ and 1I/'Oumuamua are indicated, as well as the upper estimated rotation period of 2016 NJ₃₃. The dashed line shows the critical rotation period where self gravity balances with rotational forces. Data for each comet is listed in Table 2.

our sample of objects). Again, we consider a spherical nucleus with radius $R_{\rm Nuc}$. If the outgassing induces average anisotropic rotational forces on the nucleus, $\vec{F}_{\rm Gas}$, the resulting torque is $\vec{\tau}_{\rm Gas}$,

$$\vec{\tau}_{\rm Gas} \sim \vec{R}_{\rm Nuc} \times \vec{F}_{\rm Gas} \,.$$
(12)

This torque produces a change of rotational frequency of the comet nucleus, $\vec{\omega}_{Nuc}$

$$\vec{\tau}_{\rm Gas} \sim M_{\rm Nuc} R_{\rm Nuc}^2 \frac{d\vec{\omega}_{\rm Nuc}}{dt} \,.$$
(13)

By combining the previous two equations, the rate of change of the spin of the nucleus is given by (neglecting the $\sin(\theta)$ term from the cross product)

$$\frac{d\omega_{\rm Nuc}}{dt} \propto \frac{|F_{\rm Gas}|}{M_{\rm Nuc}R_{\rm Nuc}} \,. \tag{14}$$

And, although this is a very general argument, surface forces on small bodies should be proportional to the surface area, i.e. $F_{\text{Gas}} \propto R_{\text{Nuc}}^2$. Thus, for a given bulk density, we obtain

$$\frac{d\omega_{\rm Nuc}}{dt} \propto \frac{1}{R_{\rm Nuc}^2} \,. \tag{15}$$

Therefore, smaller objects will be particularly susceptible to rapid spin up from outgassing torques. In Figure 7, the most rapidly rotating objects are preferentially small (≤ 1 km) in diameter. It is important to note,



Figure 7. A 2-dimensional histogram showing the distribution of rotational periods vs. diameters of asteroids. The critical period is shown in a dashed line, and the relationship where cohesive forces are equal to rotational forces (Equation 11 assuming $\sigma_{\rm C} = 10$ Pa, $\rho_{\rm D} = 1$ g cm⁻³, $R_{\rm D} = 1$ mm) is indicated with a dotted line. These data are from Warner et al. (2009). Vertical dashed lines delineate size scales of rubble piles and primordial intact remnants. 1998 KY₂₆, 2016 NJ₃₃ and 2006 RH₁₂₀ are indicated with diamonds, while the other candidate dark comets in our sample do not have rotation periods measured. Active asteroids with sizes and rotational periods measured (Table 3) are indicated with pentagons.

however, that the magnitude of outgassing torques scales with the production, which could depend on the size of the nucleus with a different power than two assumed here. Based on the ratio of cohesive to rotational forces, it is possible that the rapid rotation of these small objects contributed to the removal of dust *if* the objects were spun up via outgassing, and that same outgassing removed dust previously. However, the rotational forces alone, even for these rapidly rotating objecs, are not sufficiently strong to remove surface dust.

4.3. Subsurface Dust Production

Given that our candidate dark comets do not exhibit dust comae, if their accelerations are caused by outgassing then they must not have retained significant dust on their surfaces. It is not clear what mechanism removed the dust from of the surface of the nuclei, given that rotation is not sufficient to overcome typical cohesive forces. It is possible that previous outgassing activity removed surface dust, and that the preferential rapid rotation in these dark comets is a leftover signature of this process. Processes such as meteorite bombardment (as with (101955) Bennu) and fatigue due to thermal cycling will replenish surface dust, so for the nuclei to remain bare, the removal rate of dust via outgassing must be larger than the generation rate. Moreover, the episodic nature of these mechanisms militates against detection. It is also possible that significant surface material was cleared rapidly at birth, and the present-day dust creation rate is low enough to avoid apparent comae detection.

If these small objects are unable to retain surface material, the only possible source of dust activity is from subsurface dust, which is entrained during ice sublimation. Assuming the nongravitational acceleration is due to volatile outgassing, we calculate the maximum amount of dust that could be produced via such entrainment, assuming a dust-to-gas mass ratio $\mathcal{Z} = 0.01$ typically assumed in the ISM and protoplanetary disks (Birnstiel et al. 2010). The dust production rate $\dot{M}_{\rm Dust}$ for a given outgassing species X can be calculated from the gas production rate (Table 1) by multiplying the total gas mass production of the outflow by the dustto-gas ratio. Assuming H₂O driven gas production, the dust production can be calculated using

$$\dot{M}_{\text{Dust}}(X) = 3 \times 10^{-4} \,\text{g}\,\text{s}^{-1} \left(\frac{\mathcal{Z}}{0.01}\right)$$

$$\left(\frac{Q(X)}{10^{21} \text{molec}\,\text{s}^{-1}}\right) \left(\frac{m_{\text{H}_2\text{O}}}{m_X}\right).$$
(16)

Due to the small gas production rates required to power the accelerations of order $10^{19} - 10^{21}$ molec s⁻¹. the corresponding dust production rates are extremely small. For instance, in the case of H_2O outgassing from 1998 KY_{26} , the dust production rate would be $\sim 10^{-4}$ g s⁻¹. For context, (101955) Bennu exhibited $\dot{M}_{\rm Dust} \sim 10^{-4} {\rm g s}^{-1}$ which was only measurable in situ (Hergenrother et al. 2020). By comparison, typical observable cometary dust production rates are $\gtrsim 10^4$ g s^{-1} (Delsemme 1976). As a point source analogue, micron sized dust production limits from deep stacked optical images of 1I/'Oumuamua were $M_{\text{Dust}} \leq 0.2 - 2$ g s⁻¹ (Jewitt et al. 2017; Meech et al. 2017), $\gtrsim 10^3 \times$ greater than these dust production rates. Therefore, in the absence of surface dust, small, rapidly rotating, and weakly outgassing bodies should not produce detectable dust activity. It is not surprising that these objects have not been identified as active asteroids (Jewitt & Hsieh 2022).

In the absence of extended tails in images, activity may be detected via drastic brightness changes in light curves. For the case of (101955) Bennu, which displayed no tail, Hergenrother et al. (2020) estimated that the dust production $\dot{M}_{\rm Dust} \sim 10^{-4} {\rm g s}^{-1}$ would not be detectable in light curves. They estimated the detection threshold of ground based photometric surveys as 0.1 magnitudes, or 9.6% of its absolute magnitude. Given (101955) Bennu's average projected surface area of 1.9×10^9 cm², this sensitivity limit corresponds to the release of dust particles with total area 1.8×10^8 cm^2 . The largest ejection event on 6 January produced a surface area of $\sim 170-190$ cm². Therefore, even the largest production event recorded for (101955) Bennu would have been undetectable with ground based observations by $\sim 10^6$ orders of magnitude. And so, again, it is not surprising that activity has not been detected in these objects, since the implied dust production is comparable to that of (101955) Bennu. It is important to note that (101955) Bennu does not have a significant nongravitational acceleration due to particle ejection, and the activity is thought to stem from other sources such as meteoroid bombardment with a granular surface prepped by thermal fracturing (Chesley et al. 2020).

We perform an analogous calculation for 1998 KY₂₆. The projected surface area of 1998 KY₂₆ with $R_{\rm Nuc} =$ 15 m is ~ 7 × 10⁶ cm². We assume that the dust produced had the same properties as the particles ejected from (101955) Bennu, with a resulting projected surface area of 170-190 cm². Because 1998 KY₂₆ is smaller than (101955) Bennu, the change in brightness would be ~ 2.5 × 10⁻⁵ or ~ 0.0025% of its absolute magnitude. This is orders of magnitude below the uncertainty level (0.1 mags) of the photometric measurements obtained during the 1998 apparition (Figure 3 in Ostro et al. 1999). For the remaining objects, extant photometric data (some of which is reported without error bars) is not sufficient to capture such brightening events (not shown). Although there has been no activity reported for any of our candidate dark comets, the dust production levels estimated here are not detectable in extant data by orders of magnitude. Moreover, even if the photometric measurements did have the absolute accuracy required to detect this level of activity, the *in*trinsic variation due to (i) rotation, (ii) phase variation, and (iii) changing of geometry would mask this relatively small activity signature. To detect this level of activity photometrically we would first require a high-precision shape model with high fidelity information regarding the scattering properties of the surface.

5. FOLLOW-UP OBSERVATIONS

The peculiar behavior of these objects compared to known comets and asteroids makes them prime targets for follow-up observations. Given that volatile outgassing is a plausible mechanism to explain the out-ofplane nongravitational accelerations, spectroscopic observations are needed to detect or place limits on the presence of candidate volatiles (H_2O, CO_2, CO) around these objects. Given the low production rates, the sensitivity of the James Webb Space Telescope (JWST) will likely be required to obtain meaningful constraints. Additionally, deep imaging is needed to improve the limits on dust activity levels. Moreover, additional astrometric observations would help to constrain the radial acceleration A_1 , and may show that the nongravitational accelerations of these objects are higher than those reported in this paper.

In Figure 8, we show the solar elongation, visual magnitude and rate of motion on the sky for 1998 KY_{26} . Based on the observing constraints of JWST, 1998 KY_{26} is the best candidate for observations in the near future (through 2024). Improved constraints on the nature of this objects will help to resolve whether they are indeed part of a previously undescribed class of Solar System bodies.

It was announced in September 2020 that Hayabusa2 will visit 1998 KY_{26} after performing two swing-by maneuvers at Earth in the extended mission (Hirabayashi et al. 2021). The spacecraft is equipped (Watanabe et al. 2017) with a suite of instruments, the first being an Optical Navigation Camera (ONC) with one telescopic and two wide-angle cameras with seven filters at 0.39 microns (ul-band), 0.48 microns (b-band), 0.55 microns (v-band), 0.59 microns (Na), 0.70 microns (x-band), 0.86



Figure 8. The observability of the 1998 KY₂₆ with JWST from July 2023-2024. We show the Sun-Observer-Target apparent solar elongation angle as seen from the observers' location (top), the apparent visual magnitude (middle) and the rate of motion on the sky (bottom). The other candidate dark comets are not observable during this time period. The pink shaded region corresponds to nominal observability criteria for JWST — specifically $85^{\circ} < \text{Elongation} < 135^{\circ}$, V-mag < 27 mag and Rate of motion < 75 mas s⁻¹. The vertical dotted lines indicate the time window between February and March 2024 when all three observability criteria will be satisifed.

microns (w-band), and 0.95 microns (p-band) (Kameda et al. 2015, 2017; Suzuki et al. 2018; Tatsumi et al. 2019). The Thermal Infrared Imager (TIR) is capable of measuring surface roughness and thermal emmissivity and inertia via high resolution images covering 8-12 microns of thermal infrared emission which capture surface temperature ranges from 150-460 K (Arai et al. 2017; Okada et al. 2017; Takita et al. 2017). The Laser Altimeter (LI-DAR) (Mizuno et al. 2017; Senshu et al. 2017; Yamada et al. 2017) can detect dust grains in the vicinity of an asteroid, with quoted sensitivity to asteroidal dust grains similar to the Hayabusa sample surrounding the target if the number density is $> 10^5$ m⁻³ (Senshu et al. 2017). It is also equipped with a Near-Infrared Spectrometer (NIRS3) capable of obtaining spectra at near infrared wavelengths of 1.8 to 3.2 microns, designed to measure reflectance spectra of absorption bands of hydrated and hydroxide minerals (Iwata et al. 2017). Therefore, the nature of the nongravitational acceleration of 1998 KY₂₆ should be identified definitively by Hayabusa2.

6. CONCLUSIONS

In this paper, we identified five inactive objects lacking visible coma, which nonetheless showed significant nongravitational acceleration in the out of the plane direction. These objects are part of the NEO population, and are all characterized by non-peculiar orbits (e.g. typical semi-major axes, eccentricity, and inclination). As a rule, these objects are small ($R_{\rm Nuc} \sim 3 - 16$ m) and have rapid rotation periods (when measured).

While the nature of these objects remains uncertain, we showed that an outgassing mechanism can plausibly explain the out-of-plane non-gravitation acceleration without producing a visible dust coma. This is largely due to the small size of the bodies, which means that (i) relatively low gas production rates are needed to explain the accelerations, and (ii) the nuclei may not have surface dust possibly due to continual cleansing from outgassing. For objects that have lost their surface dust, the only contribution to dust activity is the entrainment of dust from the subsurface during ice sublimation. Based on the gas production rates inferred for the bodies, we show that the production of dust via entrainment is extremely small and well within the observational limits. It is possible that these objects are Solar System analogues to 11/'Oumuamua, which also exhibited a nongravitational acceleration and lack of cometary activity.

Follow-up observations are critical to understanding the nature of these unusual objects, potentially members of a new class of "dark comets" in the Solar System. In particular, observations with JWST would reveal whether their nongravitational accelerations are in fact due to volatile outgassing (e.g. H_2O , CO_2 , or CO), or whether a new mechanism is needed to explain their peculiar properties.

7. ACKNOWLEDGMENTS

We thank Dave Jewitt, John Noonan, Masatoshi Hirabayashi, Dong Lai, Fred Adams, Nikole Lewis, Konstantin Batygin, Samantha Trumbo, Gregory Laughlin, Juliette Becker, Mike Brown, Ngoc Truong, J.T. Laune, and Jonathan Lunine for useful conversations and suggestions. We thank Dave Jewitt and Henry Hsieh for compiling the data in Table 3 and for granting permission to reproduce it. We thank the scientific editor Faith Vilas for securing constructive and rapid referee reports. We thank the two anonymous reviewers for insightful comments and constructive suggestions that strengthened the scientific content of this manuscript.

DZS acknowledges financial support from the National Science Foundation Grant No. AST-17152, NASA Grant No. 80NSSC19K0444 and NASA Contract NNX17AL71A from the NASA Goddard Spaceflight Center. Part of this research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration (80NM0018D0004). DV was supported by the Czech Science Foundation, grant GA21-11058S. This work was partially supported by NASA grant number NNX17AH06G (PI N. Moskovitz) issued through the Near-Earth Object Observations program to the Mission Accessible Near-Earth Object Survey (MANOS). DJT acknowledges support from NASA Grant 80NSSC21K0807. The Arecibo planetary radar observation presented is this work was supported by the National Aeronautics and Space Administration's (NASA's) Near-Earth Object Observations program through grant No. NNX13AQ46G awarded to Universities Space Research Association (USRA). Reduction of the data was performed under grant Nos. 80NSSC18K1098 awarded to the University of Central Florida (UCF). The Arecibo Observatory is a facility of the National Science Foundation (NSF) operated under cooperative agreement by UCF, Yang Enterprises, Inc., and Universidad Ana G. Méndez. This paper is partially based on observations collected at the European Southern Observatory under ESO programme 105.202E.002.

REFERENCES

- Abell, P. A., Fernández, Y. R., Pravec, P., et al. 2005, Icarus, 179, 174, doi: 10.1016/j.icarus.2005.06.012
- Agarwal, J., Kim, Y., Jewitt, D., et al. 2020, A&A, 643, A152, doi: 10.1051/0004-6361/202038195
- A'Hearn, M. F., Campins, H., Schleicher, D. G., & Millis,
 R. L. 1989, ApJ, 347, 1155, doi: 10.1086/168204
- A'Hearn, M. F., Belton, M. J. S., Delamere, W. A., et al. 2005, Science, 310, 258, doi: 10.1126/science.1118923

Albarède, F. 2009, Nature, 461, 1227, doi: 10.1038/nature08477

- Almeida-Fernandes, F., & Rocha-Pinto, H. J. 2018, MNRAS, 480, 4903, doi: 10.1093/mnras/sty2202
- Ansdell, M., Meech, K. J., Hainaut, O., et al. 2014, ApJ, 793, 50, doi: 10.1088/0004-637X/793/1/50
- Appenzeller, I., Fricke, K., Fürtig, W., et al. 1998, The Messenger, 94, 1
- Arai, T., Nakamura, T., Tanaka, S., et al. 2017, SSRv, 208, 239, doi: 10.1007/s11214-017-0353-9

- Arai, T., Yoshida, F., Kobayashi, M., et al. 2021, in 52nd Lunar and Planetary Science Conference, Lunar and Planetary Science Conference, 1896
- Asher, D. J., Bailey, M. E., Hahn, G., & Steel, D. I. 1994, MNRAS, 267, 26, doi: 10.1093/mnras/267.1.26
- Bannister, M. T., Schwamb, M. E., Fraser, W. C., et al. 2017, ApJL, 851, L38, doi: 10.3847/2041-8213/aaa07c
- Barnouin, O. S., Daly, M. G., Palmer, E. E., et al. 2019, Nature Geoscience, 12, 247, doi: 10.1038/s41561-019-0330-x

Belton, M. J. S., Thomas, P., Li, J.-Y., et al. 2013, Icarus, 222, 595, doi: 10.1016/j.icarus.2012.06.037

- Belton, M. J. S., Hainaut, O. R., Meech, K. J., et al. 2018, ApJL, 856, L21, doi: 10.3847/2041-8213/aab370
- Benner, L. A., Ostro, S. J., Magri, C., et al. 2008, Icarus, 198, 294
- Bertini, I. 2011, Planet. Space Sci., 59, 365, doi: 10.1016/j.pss.2011.01.014
- Binzel, R. P., Xu, S., Bus, S. J., & Bowell, E. 1992, Science, 257, 779, doi: 10.1126/science.257.5071.779

- Birnstiel, T., Dullemond, C. P., & Brauer, F. 2010, A&A, 513, A79, doi: 10.1051/0004-6361/200913731
- Bodewits, D., Noonan, J. W., Feldman, P. D., et al. 2020, Nature Astronomy, 4, 867, doi: 10.1038/s41550-020-1095-2
- Boehnhardt, H., Schulz, R., Tozzi, G. P., Rauer, H., & Sekanina, Z. 1996, IAUC, 6495, 2
- Bolin, B. T., Weaver, H. A., Fernandez, Y. R., et al. 2018, ApJL, 852, L2, doi: 10.3847/2041-8213/aaa0c9
- Bottke, W. F., Brož, M., O'Brien, D. P., et al. 2015, in Asteroids IV, 701–724, doi: 10.2458/azu_uapress_9780816532131-ch036
- Bottke, W. F., Durda, D. D., Nesvorný, D., et al. 2005, Icarus, 175, 111, doi: 10.1016/j.icarus.2004.10.026
- Bottke, W. F., Moorhead, A. V., Connolly, H. C., et al. 2020, Journal of Geophysical Research (Planets), 125, e06282, doi: 10.1029/2019JE006282
- Brasser, R., & Wang, J. H. 2015, A&A, 573, A102, doi: 10.1051/0004-6361/201423687
- Cheng, A. F., Izenberg, N., Chapman, C. R., & Zuber,
 M. T. 2002, M&PS, 37, 1095,
 doi: 10.1111/j.1945-5100.2002.tb00880.x
- Chesley, S. R., Farnocchia, D., Pravec, P., & Vokrouhlický,
 D. 2016, in Asteroids: New Observations, New Models,
 ed. S. R. Chesley, A. Morbidelli, R. Jedicke, &
 D. Farnocchia, Vol. 318, 250–258,
 doi: 10.1017/S1743921315008790
- Chesley, S. R., French, A. S., Davis, A. B., et al. 2020, Journal of Geophysical Research (Planets), 125, e06363, doi: 10.1029/2019JE006363
- Chyba, C. F. 1990, Nature, 343, 129, doi: 10.1038/343129a0
- Cremonese, G., Fulle, M., Cambianica, P., et al. 2020, ApJL, 893, L12, doi: 10.3847/2041-8213/ab8455
- Delsemme, A. H. 1976, in Interplanetary Dust and Zodiacal Light, ed. H. Elsaesser & H. Fechtig, Vol. 48, 314, doi: 10.1007/3-540-07615-8_499
- Dermawan, B., Nakamura, T., & Yoshida, F. 2011, PASJ, 63, 555, doi: 10.1093/pasj/63.sp2.S555
- Devogèle, M., Ferrais, M., Jehin, E., et al. 2021, MNRAS, 505, 245, doi: 10.1093/mnras/stab1252
- Drahus, M., Guzik, P., Waniak, W., et al. 2018, Nature Astronomy, 2, 407, doi: 10.1038/s41550-018-0440-1
- Drahus, M., Waniak, W., Tendulkar, S., et al. 2015, ApJL, 802, L8, doi: 10.1088/2041-8205/802/1/L8
- Eisner, N., Knight, M. M., & Schleicher, D. G. 2017, AJ, 154, 196, doi: 10.3847/1538-3881/aa8b0b
- Elst, E. W., Pizarro, O., Pollas, C., et al. 1996, IAUC, 6456, 1

- Farnocchia, D., Chesley, S. R., Milani, A., Gronchi, G. F., & Chodas, P. W. 2015, in Asteroids IV, 815–834, doi: 10.2458/azu_uapress_9780816532131-ch041
- Farnocchia, D., Tholen, D. J., Micheli, M., et al. 2017, in AAS/Division for Planetary Sciences Meeting Abstracts #49, 100.09
- Farnocchia, D., Chesley, S. R., Takahashi, Y., et al. 2021, Icarus, 369, 114594, doi: 10.1016/j.icarus.2021.114594
- Fedorets, G., Micheli, M., Jedicke, R., et al. 2020, AJ, 160, 277, doi: 10.3847/1538-3881/abc3bc
- Feng, F., & Jones, H. R. A. 2018, ApJL, 852, L27, doi: 10.3847/2041-8213/aaa404
- Ferellec, L., Snodgrass, C., Fitzsimmons, A., et al. 2022, MNRAS, doi: 10.1093/mnras/stac3199
- Fernández, Y. R., Jewitt, D. C., & Sheppard, S. S. 2005a, AJ, 130, 308, doi: 10.1086/430802
- Fernández, Y. R., Lowry, S. C., Weissman, P. R., et al. 2005b, Icarus, 175, 194, doi: 10.1016/j.icarus.2004.10.019
- Fernández, Y. R., McFadden, L. A., Lisse, C. M., Helin, E. F., & Chamberlin, A. B. 1997, Icarus, 128, 114, doi: 10.1006/icar.1997.5728
- Fitzsimmons, A., Snodgrass, C., Rozitis, B., et al. 2018, Nature Astronomy, 2, 133, doi: 10.1038/s41550-017-0361-4
- Fraser, W. C., Pravec, P., Fitzsimmons, A., et al. 2018, Nature Astronomy, 2, 383, doi: 10.1038/s41550-018-0398-z
- Fujiwara, A., Kawaguchi, J., Yeomans, D. K., et al. 2006, Science, 312, 1330, doi: 10.1126/science.1125841
- Gaidos, E., Williams, J., & Kraus, A. 2017, Research Notes of the American Astronomical Society, 1, 13, doi: 10.3847/2515-5172/aa9851
- Granvik, M., Vaubaillon, J., & Jedicke, R. 2012, Icarus, 218, 262, doi: 10.1016/j.icarus.2011.12.003
- Granvik, M., Morbidelli, A., Jedicke, R., et al. 2018, Icarus, 312, 181, doi: 10.1016/j.icarus.2018.04.018
- Greenberg, A. H., Margot, J.-L., Verma, A. K., Taylor,
 P. A., & Hodge, S. E. 2020, AJ, 159, 92,
 doi: 10.3847/1538-3881/ab62a3

Gustafson, B. A. S. 1989, A&A, 225, 533

- Guzik, P., Drahus, M., Rusek, K., et al. 2020, Nature Astronomy, 4, 53, doi: 10.1038/s41550-019-0931-8
- Hahn, J. M., & Malhotra, R. 1999, AJ, 117, 3041, doi: 10.1086/300891
- Hainaut, O. R., Micheli, M., Cano, J. L., et al. 2021, A&A, 653, A124, doi: 10.1051/0004-6361/202141519
- Hainaut, O. R., Kleyna, J. T., Meech, K. J., et al. 2019, A&A, 628, A48, doi: 10.1051/0004-6361/201935868
- Hallatt, T., & Wiegert, P. 2020, AJ, 159, 147, doi: 10.3847/1538-3881/ab7336

Hanuš, J., Delbo', M., Vokrouhlický, D., et al. 2016, A&A, 592, A34, doi: 10.1051/0004-6361/201628666

Harmon, J. K., Nolan, M. C., Howell, E. S., Giorgini, J. D., & Taylor, P. A. 2011, ApJL, 734, L2, doi: 10.1088/2041-8205/734/1/L2

Harrington-Pinto, O., Womack, M., Fernandez, Y., & Bauer, J. 2022, PSJ, 3, 247, doi: $10.3847/\mathrm{PSJ/ac960d}$

Harris, A. W. 1979, Icarus, 40, 145, doi: 10.1016/0019-1035(79)90059-9

Harris, A. W. 1996, in Lunar and Planetary Science Conference, Vol. 27, Lunar and Planetary Science Conference, 493

Harris, A. W., & Burns, J. A. 1979, Icarus, 40, 115, doi: 10.1016/0019-1035(79)90058-7

Hergenrother, C. W., Maleszewski, C. K., Nolan, M. C., et al. 2019, Nature Communications, 10, 1291, doi: 10.1038/s41467-019-09213-x

Hergenrother, C. W., Maleszewski, C., Li, J. Y., et al. 2020, Journal of Geophysical Research (Planets), 125, e06381, doi: 10.1029/2020JE006381

Hicks, M., & Thackeray, B. 2016, The Astronomer's Telegram, 8905, 1

Hirabayashi, M., Scheeres, D. J., Sánchez, D. P., & Gabriel, T. 2014, ApJL, 789, L12,

doi: 10.1088/2041-8205/789/1/L12

Hirabayashi, M., Mimasu, Y., Sakatani, N., et al. 2021, Advances in Space Research, 68, 1533, doi: 10.1016/j.asr.2021.03.030

Hsieh, C.-H., Laughlin, G., & Arce, H. G. 2021a, ApJ, 917, 20, doi: 10.3847/1538-4357/ac0729

Hsieh, H. H. 2017, Philosophical Transactions of the Royal Society of London Series A, 375, 20160259, doi: 10.1098/rsta.2016.0259

Hsieh, H. H., Ishiguro, M., Knight, M. M., et al. 2018, AJ, 156, 39, doi: 10.3847/1538-3881/aac81c

Hsieh, H. H., & Jewitt, D. 2006, Science, 312, 561, doi: 10.1126/science.1125150

Hsieh, H. H., Jewitt, D., & Fernández, Y. R. 2009, ApJL, 694, L111, doi: 10.1088/0004-637X/694/2/L111

Hsieh, H. H., Jewitt, D. C., & Fernández, Y. R. 2004, AJ, 127, 2997, doi: 10.1086/383208

Hsieh, H. H., Meech, K. J., & Pittichová, J. 2011, ApJL, 736, L18, doi: 10.1088/2041-8205/736/1/L18

Hsieh, H. H., & Sheppard, S. S. 2015, MNRAS, 454, L81, doi: 10.1093/mnrasl/slv125

Hsieh, H. H., Hainaut, O., Novaković, B., et al. 2015, ApJL, 800, L16, doi: 10.1088/2041-8205/800/1/L16

Hsieh, H. H., Chandler, C. O., Denneau, L., et al. 2021b, ApJL, 922, L9, doi: 10.3847/2041-8213/ac2c62 Hui, M.-T., & Jewitt, D. 2017, AJ, 153, 80, doi: 10.3847/1538-3881/153/2/80

Hui, M.-T., & Li, J. 2017, AJ, 153, 23, doi: 10.3847/1538-3881/153/1/23

Hui, M.-T., Ye, Q.-Z., Föhring, D., Hung, D., & Tholen,
D. J. 2020, AJ, 160, 92, doi: 10.3847/1538-3881/ab9df8

Ishiguro, M., Hanayama, H., Hasegawa, S., et al. 2011, ApJL, 740, L11, doi: 10.1088/2041-8205/740/1/L11

Iwata, T., Kitazato, K., Abe, M., et al. 2017, SSRv, 208, 317, doi: 10.1007/s11214-017-0341-0

Jewitt, D. 2005, AJ, 129, 530, doi: 10.1086/426328

--. 2012, AJ, 143, 66, doi: 10.1088/0004-6256/143/3/66

—. 2021, AJ, 161, 261, doi: 10.3847/1538-3881/abf09c

Jewitt, D., Agarwal, J., Li, J., et al. 2014, ApJL, 784, L8, doi: 10.1088/2041-8205/784/1/L8

Jewitt, D., & Hsieh, H. H. 2022, arXiv e-prints, arXiv:2203.01397. https://arxiv.org/abs/2203.01397

Jewitt, D., Hui, M.-T., Kim, Y., et al. 2020, ApJL, 888, L23, doi: 10.3847/2041-8213/ab621b

Jewitt, D., & Li, J. 2010, AJ, 140, 1519, doi: 10.1088/0004-6256/140/5/1519

Jewitt, D., Li, J., & Agarwal, J. 2013, ApJL, 771, L36, doi: 10.1088/2041-8205/771/2/L36

Jewitt, D., & Luu, J. 1989, AJ, 97, 1766, doi: 10.1086/115118

---. 2019, ApJL, 886, L29, doi: 10.3847/2041-8213/ab530b

Jewitt, D., Luu, J., Rajagopal, J., et al. 2017, ApJL, 850, L36, doi: 10.3847/2041-8213/aa9b2f

Jewitt, D., & Seligman, D. Z. 2022, arXiv e-prints, arXiv:2209.08182. https://arxiv.org/abs/2209.08182

Jewitt, D., Weaver, H., Mutchler, M., et al. 2018, AJ, 155, 231, doi: 10.3847/1538-3881/aabdee

Jewitt, D., Kim, Y., Rajagopal, J., et al. 2019, AJ, 157, 54, doi: 10.3847/1538-3881/aaf563

Kameda, S., Suzuki, H., Cho, Y., et al. 2015, Advances in Space Research, 56, 1519, doi: 10.1016/j.asr.2015.06.037

Kameda, S., Suzuki, H., Takamatsu, T., et al. 2017, SSRv, 208, 17, doi: 10.1007/s11214-015-0227-y

Kim, Y., Jewitt, D., Mutchler, M., et al. 2020, ApJL, 895, L34, doi: 10.3847/2041-8213/ab9228

Knight, M. M., Protopapa, S., Kelley, M. S. P., et al. 2017, ApJL, 851, L31, doi: 10.3847/2041-8213/aa9d81

Knight, M. M., Schleicher, D. G., Farnham, T. L., Schwieterman, E. W., & Christensen, S. R. 2012, AJ, 144, 153, doi: 10.1088/0004-6256/144/5/153

Kokotanekova, R., Snodgrass, C., Lacerda, P., et al. 2017, MNRAS, 471, 2974, doi: 10.1093/mnras/stx1716

Krüger, H., Strub, P., Srama, R., et al. 2019, Planet. Space Sci., 172, 22, doi: 10.1016/j.pss.2019.04.005

- Küppers, M., O'Rourke, L., Bockelée-Morvan, D., et al. 2014, Nature, 505, 525, doi: 10.1038/nature12918
- Kwiatkowski, T., Kryszczyńska, A., Polińska, M., et al. 2009, A&A, 495, 967, doi: 10.1051/0004-6361:200810965
- Lauretta, D. S., Hergenrother, C. W., Chesley, S. R., et al. 2019, Science, 366, eaay3544,
 - doi: 10.1126/science.aay3544
- Li, J., & Jewitt, D. 2013, AJ, 145, 154, doi: 10.1088/0004-6256/145/6/154
- Licandro, J., Popescu, M., de León, J., et al. 2018, A&A, 618, A170, doi: 10.1051/0004-6361/201832853
- Licandro, J., Campins, H., Kelley, M., et al. 2009, A&A, 507, 1667, doi: 10.1051/0004-6361/200913116
- Lowry, S. C., & Weissman, P. R. 2007, Icarus, 188, 212, doi: 10.1016/j.icarus.2006.11.014
- MacLennan, E. M., & Hsieh, H. H. 2012, ApJL, 758, L3, doi: 10.1088/2041-8205/758/1/L3
- Mainzer, A., Read, M. T., Scottii, J. V., Ryan, W. H., & Ryan, E. V. 2010, IAUC, 9117, 1
- Mamajek, E. 2017, Research Notes of the American Astronomical Society, 1, 21, doi: 10.3847/2515-5172/aa9bdc
- Marsden, B. G., Sekanina, Z., & Yeomans, D. K. 1973, AJ, 78, 211, doi: 10.1086/111402
- Mashchenko, S. 2019, MNRAS, 489, 3003, doi: 10.1093/mnras/stz2380
- Masiero, J. 2017, arXiv e-prints. https://arxiv.org/abs/1710.09977
- Masiero, J. R., Davidsson, B. J. R., Liu, Y., Moore, K., & Tuite, M. 2021, PSJ, 2, 165, doi: 10.3847/PSJ/ac0d02
- McNeill, A., Trilling, D. E., & Mommert, M. 2018, ApJL, 857, L1, doi: 10.3847/2041-8213/aab9ab
- Meech, K. J., Yang, B., Kleyna, J., et al. 2016, Science Advances, 2, e1600038, doi: 10.1126/sciadv.1600038
- Meech, K. J., Weryk, R., Micheli, M., et al. 2017, Nature, 552, 378, doi: 10.1038/nature25020
- Micheli, M., Tholen, D. J., & Elliott, G. T. 2012, NewA, 17, 446, doi: 10.1016/j.newast.2011.11.008
- —. 2014, ApJL, 788, L1, doi: 10.1088/2041-8205/788/1/L1
 Micheli, M., Farnocchia, D., Meech, K. J., et al. 2018,
- Nature, 559, 223, doi: 10.1038/s41586-018-0254-4
- Mizuno, T., Kase, T., Shiina, T., et al. 2017, SSRv, 208, 33, doi: 10.1007/s11214-015-0231-2
- Molaro, J. L., Hergenrother, C. W., Chesley, S. R., et al. 2020, Journal of Geophysical Research (Planets), 125, e06325, doi: 10.1029/2019JE006325
- Mommert, M., Hora, J. L., Farnocchia, D., et al. 2014a, ApJ, 786, 148, doi: 10.1088/0004-637X/786/2/148

- Mommert, M., Farnocchia, D., Hora, J. L., et al. 2014b, ApJL, 789, L22, doi: 10.1088/2041-8205/789/1/L22
- Moreno, F., Licandro, J., Cabrera-Lavers, A., Morate, D., & Guirado, D. 2021, MNRAS, 506, 1733, doi: 10.1093/mnras/stab1841
- Moreno, F., Licandro, J., Cabrera-Lavers, A., & Pozuelos,
 F. J. 2016a, ApJ, 826, 137,
 doi: 10.3847/0004-637X/826/2/137
- —. 2016b, ApJL, 826, L22, doi: 10.3847/2041-8205/826/2/L22
- Moreno, F., Pozuelos, F. J., Novaković, B., et al. 2017, ApJL, 837, L3, doi: 10.3847/2041-8213/aa6036
- Moro-Martín, A. 2022, arXiv e-prints, arXiv:2205.04277. https://arxiv.org/abs/2205.04277
- Mottola, S., Lowry, S., Snodgrass, C., et al. 2014, A&A, 569, L2, doi: 10.1051/0004-6361/201424590
- Nakano, R., & Hirabayashi, M. 2020, ApJL, 892, L22, doi: 10.3847/2041-8213/ab7d36
- Novaković, B., Pavela, D., Hsieh, H. H., & Marčeta, D. 2022, MNRAS, 516, 757, doi: 10.1093/mnras/stac2250
- Okada, T., Fukuhara, T., Tanaka, S., et al. 2017, SSRv, 208, 255, doi: 10.1007/s11214-016-0286-8
- Ostro, S. J., Pravec, P., Benner, L. A. M., et al. 1999, Science, 285, 557, doi: 10.1126/science.285.5427.557
- Owen, T., & Bar-Nun, A. 1995, Icarus, 116, 215, doi: 10.1006/icar.1995.1122
- Park, R. S., Konopliv, A. S., Bills, B. G., et al. 2016, Nature, 537, 515, doi: 10.1038/nature18955
- Podolak, M., & Herman, G. 1985, Icarus, 61, 267, doi: 10.1016/0019-1035(85)90108-3
- Pravec, P., & Harris, A. W. 2007, Icarus, 190, 250, doi: 10.1016/j.icarus.2007.02.023
- Prialnik, D., & Bar-Nun, A. 1988, Icarus, 74, 272, doi: 10.1016/0019-1035(88)90042-5
- Rozitis, B., Maclennan, E., & Emery, J. P. 2014, Nature, 512, 174, doi: 10.1038/nature13632
- Russell, C. T., Aroian, R., Arghavani, M., & Nock, K. 1984, Science, 226, 43, doi: 10.1126/science.226.4670.43
- Sánchez, P., & Scheeres, D. J. 2020, Icarus, 338, 113443, doi: 10.1016/j.icarus.2019.113443
- Scheeres, D. J., Hartzell, C. M., Sánchez, P., & Swift, M. 2010, Icarus, 210, 968, doi: 10.1016/j.icarus.2010.07.009
- Senshu, H., Oshigami, S., Kobayashi, M., et al. 2017, SSRv, 208, 65, doi: 10.1007/s11214-016-0242-7
- Shannon, A., Jackson, A. P., Veras, D., & Wyatt, M. 2015, MNRAS, 446, 2059, doi: 10.1093/mnras/stu2267
- Sheppard, S. S., & Trujillo, C. 2015, AJ, 149, 44, doi: 10.1088/0004-6256/149/2/44
- Sierks, H., Barbieri, C., Lamy, P. L., et al. 2015, Science, 347, aaa1044, doi: 10.1126/science.aaa1044

- Snodgrass, C., Tubiana, C., Vincent, J.-B., et al. 2010, Nature, 467, 814, doi: 10.1038/nature09453
- Snodgrass, C., Agarwal, J., Combi, M., et al. 2017, A&A Rv, 25, 5, doi: 10.1007/s00159-017-0104-7
- Sonnett, S., Kleyna, J., Jedicke, R., & Masiero, J. 2011, Icarus, 215, 534, doi: 10.1016/j.icarus.2011.08.001
- Suzuki, H., Yamada, M., Kouyama, T., et al. 2018, Icarus, 300, 341, doi: 10.1016/j.icarus.2017.09.011
- Takita, J., Senshu, H., & Tanaka, S. 2017, SSRv, 208, 287, doi: 10.1007/s11214-017-0336-x
- Tatsumi, E., Kouyama, T., Suzuki, H., et al. 2019, Icarus, 325, 153, doi: 10.1016/j.icarus.2019.01.015
- Taylor, P. A., Rivera-Valentín, E. G., Benner, L. A. M., et al. 2019, Planet. Space Sci., 167, 1, doi: 10.1016/j.pss.2019.01.009
- Tedesco, E. F., Noah, P. V., Noah, M., & Price, S. D. 2004, NASA Planetary Data System, IRAS
- Tholen, D. J., Sheppard, S. S., & Trujillo, C. A. 2015, in AAS/Division for Planetary Sciences Meeting Abstracts, Vol. 47, AAS/Division for Planetary Sciences Meeting Abstracts #47, 414.03
- Toth, I. 2000, A&A, 360, 375
- Trilling, D. E., Mommert, M., Hora, J. L., et al. 2018, AJ, 156, 261, doi: 10.3847/1538-3881/aae88f
- Urakawa, S., Okumura, S.-i., Nishiyama, K., et al. 2011, Icarus, 215, 17, doi: 10.1016/j.icarus.2011.06.044
- Virkki, A. K., Marshall, S. E., Venditti, F. C. F., et al. 2022, PSJ, 3, 222, doi: 10.3847/PSJ/ac8b72
- Vokrouhlický, D., Bottke, W. F., Chesley, S. R., Scheeres, D. J., & Statler, T. S. 2015, in Asteroids IV, 509–531, doi: 10.2458/azu_uapress_9780816532131-ch027
- Vokrouhlický, D., & Milani, A. 2000, A&A, 362, 746
- Vokrouhlický, D., Milani, A., & Chesley, S. R. 2000, Icarus, 148, 118, doi: 10.1006/icar.2000.6469

- Walsh, K. J. 2018, ARA&A, 56, 593, doi: 10.1146/annurev-astro-081817-052013
- Wang, J. H., & Brasser, R. 2014, A&A, 563, A122, doi: 10.1051/0004-6361/201322508
- Warner, B. D. 2016, Minor Planet Bulletin, 43, 240
- Warner, B. D., Harris, A. W., & Pravec, P. 2009, Icarus, 202, 134, doi: 10.1016/j.icarus.2009.02.003
- Watanabe, S., Hirabayashi, M., Hirata, N., et al. 2019, Science, 364, 268, doi: 10.1126/science.aav8032
- Watanabe, S.-i., Tsuda, Y., Yoshikawa, M., et al. 2017, SSRv, 208, 3, doi: 10.1007/s11214-017-0377-1
- Weryk, R., Wainscoat, R., Ramanjooloo, Y., Woodworth, D., & Micheli, M. 2018, Central Bureau Electronic Telegrams, 4548, 1
- Weryk, R., Wainscoat, R., Woodworth, D., & Prunet, S. 2019, Central Bureau Electronic Telegrams, 4598, 1
- Weryk, R., Wainscoat, R. J., Wipper, C., & Micheli, M. 2017, Central Bureau Electronic Telegrams, 4448, 1
- Whipple, F. L. 1950, ApJ, 111, 375, doi: 10.1086/145272
- -. 1951, ApJ, 113, 464, doi: 10.1086/145416
- Williams, G. V., Sato, H., Sarneczky, K., et al. 2017, Central Bureau Electronic Telegrams, 4450, 1
- Williams, I. P., & Wu, Z. 1993, MNRAS, 262, 231, doi: 10.1093/mnras/262.1.231
- Wisniewski, W. Z. 1990, Icarus, 86, 52, doi: 10.1016/0019-1035(90)90197-H
- Yamada, R., Senshu, H., Namiki, N., et al. 2017, SSRv, 208, 49, doi: 10.1007/s11214-016-0240-9
- Yang, B., Li, A., Cordiner, M. A., et al. 2021, Nature Astronomy, doi: 10.1038/s41550-021-01336-w
- Ye, Q.-Z., Zhang, Q., Kelley, M. S. P., & Brown, P. G. 2017, ApJL, 851, L5, doi: 10.3847/2041-8213/aa9a34



Figure 9. A radar CW spectrum of 2016 NJ₃₃ taken with the Arecibo Observatory. The CW spectrum shows that the minimum rotation period should be between 0.41 and 1.99 h. Because we have no information regarding the pole orientation of the object, it is possible that the rotational period is much more rapid. The low SC/OC ratio exclude an E-type classification (Benner et al. 2008). Reproduced from https://www.naic.edu/~pradar/asteroids/2016NJ33/2016NJ33.2016Jul15.s0p25Hz.cw.gif

APPENDIX

A. ROTATION PERIOD OF 2016 NJ₃₃

For 2016 NJ₃₃, we estimate the rotational period using one continuous wave (CW) observation obtained with the Arecibo Observatory planetary radar system, shown in Figure 9. The CW spectrum shows a bandwidth of 0.50 ± 0.02 Hz. Using the relation linking the bandwidth of a CW spectrum and the rotation period (see Virkki et al. (2022) for more information on CW radar observation and their analysis) we estimate that the rotation period of 2016 NJ₃₃ is between 0.41 and 1.99 h with a mean of 1.43 h with 3 sigma confidence. We obtained this result by simulating random pole orientations and a normal distribution of the bandwidth (0.50 ± 0.02 Hz) and the diameter (32 ± 3 m). This estimation of the rotational period should be treated with caution because (i) we only have one observation of the object, (ii) an observation very close to pole-on geometry would result in a very fast rotation period, (iii) we assume a spherical nucleus and if the object is highly elongated, a very different cross-section would be allowable. However, there is a low probability of randomly observing the object pole-on.

B. DATA FOR ACTIVE ASTEROIDS AND COMETARY NUCLEI

Table 1 is reproduced from Jewitt & Hsieh (2022). In Table 2, we show relevant data and references for comets with measured nuclei rotation rates and sizes. In Table 3, we list active asteroid properties.

| component for each ol assuming a | t, onl bject an alt | y 201 in ou sedo (| 6 NJ If sat of $p =$ | [₃₃ , 2(mple = 0.1 | 003 R at pe and t | M and rihelic hat tł | d 2006 R. m. We c re molecu | H ₁₂₀ have robu alculate this u des are released | ist det sing tl 1 at a | ections. We calcu le most significan temperature of 10 | ilate i it com 0 K. | mplied producti uponent of the a | ion rate uccelera | es of H ₂ O ation, and |
|--|---------------------------|--------------------------|----------------------------|---------------------------------------|-------------------------|----------------------------|-----------------------------------|---|------------------------------|--|---------------------------|--|----------------------|--------------------------------------|
| Object | а | υ | i | d | Η | $R_{ m Nuc}$ | $P_{ m Rot}$ | A_1 | Signif. | A_2 | Signif | A_3 | Signif. | $Q({ m H_2O})$ |
| | [au] | | [0] | [au] | [mag] | [m] | [µ] | $[10^{-10} \text{ au } \mathrm{d}^{-2}]$ | $\left[\sigma \right]$ | $[10^{-10} \text{ au } \mathrm{d}^{-2}]$ | [σ] | $[10^{-10} \text{ au } \mathrm{d}^{-2}]$ | σ | $[10^{20} \text{ molec s}^{-1}]$ |
| 1998 KY_{26} | 1.23 | 0.20 | 1.48 | 0.98 | 25.60 | 15 | 0.178 | 1.73 ± 0.91 | 2 | -0.00126 ± 0.00061 | 7 | 0.320 ± 0.115 | e | 9.19 |
| 2005 VL_1 | 0.89 | 0.23 | 0.25 | 0.69 | 26.45 | 11 | | -6.66 ± 8.02 | $^{\prime}$ | -0.00711 ± 0.00592 | 1 | -0.240 ± 0.041 | 9 | 5.5 |
| 2016 NJ_{33} | 1.31 | 0.21 | 6.64 | 1.04 | 25.49 | 16 | 0.41 - 1.99 | 9.28 ± 2.96 | 3 | -0.00566 ± 0.00193 | 33 | 0.848 ± 0.163 | ъ | 24.8 |
| 2010 VL_{65} | 1.07 | 0.14 | 4.41 | 0.91 | 29.22 | 3 | | 6.57 ± 13.0 | $^{\prime}$ | -0.00146 ± 0.00534 | \sim | -0.913 ± 0.130 | 7 | 0.24 |
| $2010~{\rm RF}_{12}$ | 1.06 | 0.19 | 0.88 | 0.86 | 28.42 | 4 | | $0.488 {\pm} 0.597$ | $^{\prime}$ | -0.00136 ± 0.00286 | \sim | -0.168 ± 0.021 | x | 0.12 |
| 2006 RH_{120} | 0 1.00 | 0.04 | 0.31 | 0.96 | 29.50 | 2-7 | 0.046 | 1.38 ± 0.08 | 18 | -0.507 ± 0.0637 | x | -0.130 ± 0.032 | 4 | 1.5 |
| 2003 RM | 2.92 | 0.60 | 10.86 | 1.17 | 19.80 | 230 | | -1.045 ± 1.217 | \sim | 0.0215 ± 0.0004 | 56 | 0.0156 ± 0.0543 | $^{\prime}$ | 1600 |

eters for each object in our sample, with formal $(1-\sigma)$ uncertainties. The statistical significance is given for each acceleration and calculated as the ratio of the magnitude of the acceleration to its uncertainty. We also include the object 2003 RM, for

Table 1. Objects with Anomalous Nongravitational Accelerations. The best-fit nongravitational acceleration param-

Statistically robust results $(\sigma \ge 3)$ are found for the A₃ component of all newly reported objects in this paper. For the A_2

which we report anomalously large nongravitational acceleration in the accompanying paper Farnocchia et al. (submitted 2022).

| rotational period of 1998 KY ₂₆ was first reported by Ostro et al. (1999). The rotational period of | from radar CW spectral observations taken with the Arecibo Observatory planetary radar system (see | lear radius and period of 2006 RH ₁₂₀ were reported by Kwiatkowski et al. (2009). The remaining objects | neasurements of size or rotational period. We estimate the remaining sizes with Equation 2 assuming a | = 0.1. |
|--|--|--|---|---------------------------------|
| NOTES: The size and rotational period of 199 | 2016 NJ_{33} is estimated from radar CW spectral | Appendix A). The nuclear radius and period of | do not have reported measurements of size or r | geometric albedo of $p = 0.1$. |

| Object Name | q [au] | $i \ [^{\circ}]$ | Classification | Diameter [km] | $P_{\rm Rot}$ [h] | Reference |
|-----------------------------|--------|------------------|----------------|---------------|-------------------|------------------------------|
| P/2006 HR30 (Siding Spring) | 1.226 | 31.88 | HTC | | 70.7 | |
| C/2001 OG108 (LONEOS) | 0.994 | 80.25 | HTC | 13.6 | 57.12 | Abell et al. (2005) |
| | | | | | | Fernández et al. $(2005a)$ |
| 9P/Tempel 1 | 1.542 | 10.47 | JFC | 6 | 40.7 | A'Hearn et al. (2005) |
| P/2016 BA14 (PANSTARRS) | 1.009 | 18.92 | JFC | | 36.6 | Warner (2016) |
| 162P/Siding Spring | 1.233 | 27.82 | JFC | | 32.853 | Kokotanekova et al. (2017) |
| 333P/LINEAR | 1.115 | 131.88 | JFC | | 21.04 | Hicks & Thackeray (2016) |
| 94P/Russell 4 | 2.24 | 6.18 | ETC | | 20.7 | Kokotanekova et al. (2017) |
| 93P/Lovas 1 | 1.7 | 12.2 | JFC | | 18.2 | Kokotanekova et al. (2017) |
| 103P/Hartley 2 | 1.059 | 13.62 | JFC | 1.6 | 18.1 | Belton et al. (2013) |
| | | | | | | Harmon et al. (2011) |
| 47P/Ashbrook-Jackson | 2.802 | 13.05 | JFC | 5.6 | 15.6 | Kokotanekova et al. (2017) |
| 49P/Arend-Rigaux | 1.424 | 19.05 | JFC | 8.48 | 13.45 | Eisner et al. (2017) |
| 67P/Churyumov-Gerasimenko | 1.241 | 7.05 | JFC | 3.4 | 12.76129 | Mottola et al. (2014) |
| | | | | | | Sierks et al. (2015) |
| 149P/Mueller 4 | 2.647 | 29.75 | JFC | | 11.88 | Kokotanekova et al. (2017) |
| 2P/Encke | 0.336 | 11.78 | ETC | 4.8 | 11.083 | Fernández et al. $(2005b)$ |
| | | | | | | Lowry & Weissman (2007) |
| 110P/Hartley 3 | 2.465 | 11.7 | JFC | 4.3 | 10.153 | Kokotanekova et al. (2017) |
| 14P/Wolf | 2.729 | 27.94 | JFC | 4.66 | 9.02 | Kokotanekova et al. (2017) |
| 10P/Tempel 2 | 1.421 | 12.03 | JFC | 10.6 | 8.93 | A'Hearn et al. (1989) |
| | | | | | | Jewitt & Luu (1989) |
| | | | | | | Wisniewski (1990) |
| | | | | | | Knight et al. (2012) |
| 169P/NEAT | 0.607 | 11.3 | JFC | | 8.369 | |
| 137P/Shoemaker-Levy2 | 1.933 | 4.85 | JFC | 5.8 | 7.7 | Kokotanekova et al. (2017) |
| 123P/West-Hartley | 2.126 | 15.35 | JFC | | 3.7 | Kokotanekova et al. (2017) |
| C/2003 WT42 (LINEAR) | 5.191 | 31.41 | HYP | | 3.31 | Dermawan et al. (2011) |

Table 2. Comets with nuclear rotation periods measured. The data is drawn from JPL small body database.

| Table 3. Properties of the currently known active asteroids. | This table is reproduced | from Jewitt & Hsieh | (2022) with |
|--|--------------------------|---------------------|-------------|
| permission from the authors. | | | |

| Object Name | a [au] | e | $i~[^\circ]$ | $H \;[\mathrm{mag}]$ | $R_{\rm Nuc} [{\rm m}]$ | $P_{\rm Rot}$ [h] | Reference |
|--------------------------------------|--------|-------|--------------|----------------------|--------------------------|-------------------|----------------------------|
| (1) Ceres | 2.766 | 0.078 | 10.588 | 3.53 | 469.7 | 9.07 | Küppers et al. (2014) |
| | | | | | | | Park et al. (2016) |
| (493) Griseldis | 3.116 | 0.176 | 15.179 | 10.97 | 20.78 | 51.94 | Tholen et al. (2015) |
| (596) Scheila | 2.929 | 0.163 | 14.657 | 8.93 | 79.86 | 15.85 | Ishiguro et al. (2011) |
| (2201) Oljato | 2.174 | 0.713 | 2.522 | 15.25 | 0.90 | >26. | Russell et al. (1984) |
| | | | | | | | Tedesco et al. (2004) |
| (3200) Phaethon | 1.271 | 0.890 | 22.257 | 14.32 | 3.13 | 3.60 | Jewitt & Li (2010) |
| | | | | | | | Ansdell et al. (2014) |
| (6478) Gault | 2.305 | 0.193 | 22.812 | 14.81 | 2.8 | 2.49 | Devogèle et al. (2021) |
| (62412) 2000 SY ₁₇₈ | 3.159 | 0.079 | 4.738 | 13.74 | 5.19 | 3.33 | Sheppard & Trujillo (2015) |
| (101955) Bennu | 1.126 | 0.204 | 6.035 | 20.21 | 0.24 | 4.29 | Lauretta et al. (2019) |
| 107P/(4015) Wilson-Harrington | 2.625 | 0.632 | 2.799 | 16.02 | 3.46 | 7.15 | Fernández et al. (1997) |
| | | | | | | | Licandro et al. (2009) |
| | | | | | | | Urakawa et al. (2011) |
| 133P/(7968) Elst-Pizarro | 3.165 | 0.157 | 1.389 | 15.49 | 1.9 | 3.47 | Hsieh et al. (2009) |
| 176P/(118401) LINEAR | 3.194 | 0.193 | 0.235 | 15.10 | 2.0 | 22.23 | Hsieh et al. (2009) |
| $233P/La Sagra (P/2005 JR_{71})$ | 3.033 | 0.411 | 11.279 | 16.6 | 1.5 | — | Mainzer et al. (2010) |
| 238P/Read (P/2005 U1) | 3.162 | 0.253 | 1.266 | 19.05 | 0.4 | — | Hsieh et al. (2011) |
| 259P/Garradd (P/2008 R1) | 2.727 | 0.342 | 15.899 | 19.71 | 0.30 | | MacLennan & Hsieh (2012) |
| $288P/(300163) 2006 VW_{139}$ | 3.051 | 0.201 | 3.239 | 17.8, 18.2 | $0.9,\!0.6$ | — | Agarwal et al. (2020) |
| 311P/PANSTARRS (P/2013 P5) | 2.189 | 0.116 | 4.968 | 19.14 | 0.2 | > 5.4 | Jewitt et al. (2018) |
| 313P/Gibbs (P/2014 S4) | 3.154 | 0.242 | 10.967 | 17.1 | 1.0 | — | Hsieh et al. (2015) |
| $324\mathrm{P/La}$ Sagra (P/2010 R2) | 3.098 | 0.154 | 21.400 | 18.4 | 0.55 | — | Hsieh & Sheppard (2015) |
| 331P/Gibbs (P/2012 F5) | 3.005 | 0.042 | 9.739 | 17.33 | 1.77 | 3.24 | Drahus et al. (2015) |
| 354P/LINEAR (P/2010 A2) | 2.290 | 0.125 | 5.256 | | 0.06 | 11.36 | Snodgrass et al. (2010) |
| 358P/PANSTARRS (P/2012 T1) | 3.155 | 0.236 | 11.058 | 19.5 | 0.32 | — | Hsieh et al. (2018) |
| 426P/PANSTARRS (P/2019 A7) | 3.188 | 0.161 | 17.773 | 17.1 | 1.2 | — | |
| 427P/ATLAS (P/2017 S5) | 3.171 | 0.313 | 11.849 | 18.91 | 0.45 | 1.4 | Jewitt et al. (2019) |
| 432P/PANSTARRS (P/2021 N4) | 3.045 | 0.244 | 10.067 | >18.2 | < 0.7 | — | |
| $433P/(248370) 2005 QN_{173}$ | 3.067 | 0.226 | 0.067 | 16.32 | 1.6 | — | Hsieh et al. $(2021b)$ |
| | | | | | | | Novaković et al. (2022) |
| P/2013 R3 (Catalina-PANSTARRS) | 3.033 | 0.273 | 0.899 | — | ~ 0.2 | — | Jewitt et al. (2014) |
| P/2015 X6 (PANSTARRS) | 2.755 | 0.170 | 4.558 | >18.2 | $<\!0.7$ | — | Moreno et al. (2016a) |
| P/2016 G1 (PANSTARRS) | 2.583 | 0.210 | 10.968 | — | < 0.4 | — | Moreno et al. $(2016b)$ |
| | | | | | | | Hainaut et al. (2019) |
| P/2016 J1-A/B (PANSTARRS) | 3.172 | 0.228 | 14.330 | — | $<\!0.4,\!<\!0.9$ | — | Moreno et al. (2017) |
| P/2017 S9 (PANSTARRS) | 3.156 | 0.305 | 14.138 | > 17.8 | < 0.8 | — | Weryk et al. (2017) |
| P/2018 P3 (PANSTARRS) | 3.007 | 0.416 | 8.909 | > 18.6 | $<\!0.6$ | — | Weryk et al. (2018) |
| P/2019 A3 (PANSTARRS) | 3.147 | 0.265 | 15.367 | >19.3 | < 0.4 | — | Weryk et al. (2019) |
| P/2019 A4 (PANSTARRS) | 2.614 | 0.090 | 13.319 | — | 0.17 | — | Moreno et al. (2021) |
| P/2020 O1 (Lemmon-PANSTARRS) | 2.647 | 0.120 | 5.223 | > 17.7 | < 0.9 | | |
| P/2021 A5 (PANSTARRS) | 3.047 | 0.140 | 18.188 | _ | 0.15 | — | Moreno et al. (2021) |
| P/2021 L4 (PANSTARRS) | 3.165 | 0.119 | 16.963 | >15.8 | $<\!2.2$ | | |
| P/2021 R8 (Sheppard) | 3.019 | 0.294 | 2.203 | _ | | — | |