Forward Orthogonal Deviations GMM and the Absence of Large Sample Bias

Robert F. Phillips^{*}

Department of Economics George Washington University

July 2024

Abstract

It is well known that generalized method of moments (GMM) estimators of dynamic panel data regressions can have significant bias when the number of time periods (T) is not small compared to the number of cross-sectional units (n). The bias is attributed to the use of many instrumental variables. This paper shows that if the maximum number of instrumental variables used in a period increases with T at a rate slower than $T^{1/2}$, then GMM estimators that exploit the forward orthogonal deviations (FOD) transformation do not have asymptotic bias, regardless of how fast T increases relative to n. This conclusion is specific to using the FOD transformation. A similar conclusion does not necessarily apply when other transformations are used to remove fixed effects. Monte Carlo evidence illustrating the analytical results is provided.

Keywords: Asymptotic bias; dynamic panel data; first difference; forward orthogonal deviations; generalized method of moments

^{*2115} G Street, NW, Suite 340, Washington DC, 20052; phone: 202-994-8619; fax: 202-994-6147; email: rphil@gwu.edu

1 Introduction

In an influential paper, Alvarez and Arellano (2003) examined the asymptotic properties of a generalized method of moments (GMM) estimator that relied on forward orthogonal deviations (FOD) to remove fixed effects (FOD GMM). They showed that an FOD GMM estimator of the autoregression parameter in a first-order autoregressive (AR(1)) panel data model has a bias term in its asymptotic distribution if the number of time periods (T) increases too quickly relative to the number of crosssectional units (n)—that is, if T/n does not converge to zero. Though the model considered by Alvarez and Arellano (2003) was a special case, the paper's finding had an important implication. This is because if an estimator, say $\hat{\delta}$, of parameter δ has a bias term in the asymptotic distribution of $\sqrt{nT}(\hat{\delta} - \delta)$, then even though the estimator may be consistent, large sample confidence intervals and test statistics based on the estimator will be inaccurate.

Alvarez and Arellano's conclusion that the FOD GMM estimator they studied has asymptotic bias if T/n does not converge to zero depends on using all available instrumental variables. If T is not small, the number of instrumental variables can be large if all of them are used, and it is well-known that a GMM estimator that exploits many instrumental variables can be biased. Consequently, in practice researchers often resort to using fewer instrumental variables than all that are available. But left unanswered is the question: how many instrumental variables can be used while still avoiding bias when T is not small compared to n? This paper addresses that question.

Like this paper, Anderson et al. (2011), Bekker (1994), Bun and Kiviet (2006), Chao and Swanson (2005), Hansen et al. (2008), Hayakawa et al. (2019), and Koenker and Machado (1999), among others, study how the properties of estimators are affected by the number of moment restrictions that are exploited. However, the papers most closely related to this paper are Alvarez and Arellano (2003) and Hsiao and Zhou (2017). As already noted, Alvarez and Arellano (2003) considered estimation of the AR(1) panel data model:

$$y_{i,t} = \beta y_{i,t-1} + \eta_i + v_{i,t}, \qquad |\beta| < 1.$$
(1)

Assuming all available moment restrictions are exploited, Alvarez and Arelleno found that an FOD GMM estimator, say $\hat{\beta}$, of the autoregression parameter, β , has a bias term in the asymptotic distribution of $\sqrt{nT}(\hat{\beta} - \beta)$ if $T/n \to c > 0$, as $n, T \to \infty$, but it does not have a bias term if $T/n \to 0$. Hsiao and Zhou (2017), on the other hand, investigated the effect of using fewer than all available moment restrictions. In particular, Hsiao and Zhou (2017) analyzed FOD GMM estimation of the model in (1), but, instead of focusing on only using all available instrumental variables, Hsiao and Zhou also considered estimation based on a single instrumental variable per period. Upon doing so, they obtained results indicating that the FOD GMM estimators they considered—when based on a single instrumental variable per period—have no bias in their asymptotic distributions, as $n, T \to \infty$, regardless of what happens to T/n.¹ The papers by Alvarez and Arellano (2003) and Hsiao and Zhou (2017), therefore, indicate that whether or not there is a bias term in the asymptotic distribution of an FOD GMM estimator depends not just on what happens to T/n, as $n, T \to \infty$, but also on how many per-period instruments are used.

Using more general conditions than previously considered, this paper shows that whether or not an FOD GMM estimator is asymptotically biased can be summarized in terms of the largest number of instrumental variables used in a period. Specifically, I show that the FOD GMM estimator has no asymptotic bias regardless of what happens to T/n, if the maximum number of instruments used in a period increases with T at a rate slower than $T^{1/2}$ increases. This conclusion is specific to using the FOD transformation. It does not necessarily carry over to other transformations that may be used to remove fixed effects.

Moreover, the conclusion is robust in the sense that the large sample bias result holds regardless of how n and T increase. Specifically, the conclusion is based on taking joint limits, which are limits obtained by letting n and T increase simultaneously.

A second less robust result is also provided: an asymptotic distribution result is provided that is based on taking limits sequentially. With a sequential limit, one index—n or T—is taken to infinity, and then the other goes to infinity. Taking limits sequentially is often a more tractable tactic for obtaining results than joint limits, and sequential limit results may require weaker conditions than joint limit results. These advantages may explain why limits are sometimes taken sequentially in the literature. Hsiao and Zhou (2017), for example, used sequential limits to derive asymptotic distribution results for their FOD GMM estimators of the autoregression

¹On the other hand, Hsiao and Zhou (2017) also argued that GMM based on first differences (FD GMM) and a single instrumental variable per period is asymptotically biased if $T/n \rightarrow c > 0$.

parameter in the model in (1). A few other papers that exploit sequential limits are Kapetanios (2008) and Hsiao and Zhang (2015).

However, sequential limits are not guaranteed to yield the same results one gets from joint limit analysis (Phillips and Moon, 1999, 2000). Indeed, I show that this is the case when all available instrumental variables are used. On the other hand, when fewer than all available instruments are used, Monte Carlo evidence indicates that the normal approximation obtained by taking limits sequentially appears to work well for constructing confidence intervals when T—in addition to n—is not small, provided the FOD transformation is used to remove fixed effects.

2 FOD GMM

2.1 The model and estimator

The regression model studied in this paper is

$$y_{i,t} = \mathbf{x}'_{i,t} \boldsymbol{\beta} + \eta_i + v_{i,t}$$
 $(t = 1, \dots, T, i = 1, \dots, n).$

In this regression, $\mathbf{x}'_{i,t} := (x_{i,t,1}, \ldots, x_{i,t,K})$ and $\boldsymbol{\beta}' := (\beta_1, \ldots, \beta_K)$ are vectors of regressors and parameters. Some or all of the $x_{i,t,k}$ s may be lagged values of $y_{i,t}$. The term η_i is an unobserved individual-specific or fixed effect, and $v_{i,t}$ is an error term.

In order to estimate β , the first step is to remove the fixed effect η_i by transforming the dependent and explanatory variables. This paper studies transforming the variables using forward orthogonal deviations—the FOD transformation. The FOD transformation subtracts from each variable its within-group average over future periods. For example, the FOD transformed explanatory variables for the *i*th individual in the *t*th period are $\ddot{\boldsymbol{x}}_{i,t} := c_t (\boldsymbol{x}_{i,t} - \bar{\boldsymbol{x}}_{i,t})$, where $\bar{\boldsymbol{x}}_{i,t} := (1/(T-t)) \sum_{s=1}^{T-t} \boldsymbol{x}_{i,t+s}$ and $c_t^2 := (T-t)/(T-t+1)$. Similarly, the transformed value of the (i, t)th observation on the dependent variable is $\ddot{y}_{i,t} := c_t (y_{i,t} - \bar{y}_{i,t})$, where $\bar{y}_{i,t} := (1/(T-t)) \sum_{s=1}^{T-t} y_{i,t+s}$. The constant c_t ensures the transformed errors (the $\ddot{v}_{i,t}$ s) are conditionally homoskedastic and uncorrelated if the original errors (the $v_{i,t}$ s) are conditionally homoskedastic and uncorrelated (Arellano, 2003, p. 17).

Now let $\boldsymbol{z}_{i,t}$ denote a $q_t \times 1$ vector of instrumental variables for the *i*th individual in the *t*th period $(t = 1, \ldots, T-1)$. Also, set $\boldsymbol{Z}'_t := (\boldsymbol{z}_{1,t}, \ldots, \boldsymbol{z}_{n,t}), \ \boldsymbol{\ddot{X}}'_t := (\boldsymbol{\ddot{x}}_{1,t}, \ldots, \boldsymbol{\ddot{x}}_{n,t}), \ \boldsymbol{\ddot{y}}'_t := (\boldsymbol{\ddot{y}}_{1,t}, \ldots, \boldsymbol{\ddot{y}}_{n,t}),$ and finally $\boldsymbol{P}_t := \boldsymbol{Z}_t (\boldsymbol{Z}'_t \boldsymbol{Z}_t)^{-1} \boldsymbol{Z}'_t$. Then, the FOD GMM esti-

mator can be written as

$$\widehat{\boldsymbol{\beta}} := \left(\sum_{t=1}^{T-1} \ddot{\boldsymbol{X}}_t' \boldsymbol{P}_t \ddot{\boldsymbol{X}}_t\right)^{-1} \sum_{t=1}^{T-1} \ddot{\boldsymbol{X}}_t' \boldsymbol{P}_t \ddot{\boldsymbol{y}}_t$$
(2)

(see, e.g., Arellano, 2003, p. 154).

As an alternative to Eq. (2), the FOD GMM estimator can also be expressed as a two-stage least squares (TSLS) estimator after removing fixed effects with the FOD transformation. To see this, let $\mathbf{Z}_{d,i}$ be the block-diagonal matrix given by

$$\boldsymbol{Z}_{d,i} := \begin{pmatrix} \boldsymbol{z}_{i,1}' & \boldsymbol{0} & \cdots & \boldsymbol{0} \\ \boldsymbol{0} & \boldsymbol{z}_{i,2}' & \cdots & \boldsymbol{0} \\ \vdots & \vdots & \ddots & \vdots \\ \boldsymbol{0} & \boldsymbol{0} & \cdots & \boldsymbol{z}_{i,T-1}' \end{pmatrix}.$$
(3)

1

Also, define $\dot{\mathbf{X}}'_i := (\ddot{\mathbf{x}}_{i,1}, \dots, \ddot{\mathbf{x}}_{i,T-1})$ and $\dot{\mathbf{y}}'_i := (\ddot{y}_{i,1}, \dots, \ddot{y}_{i,T-1})$. Then, note that $\sum_{t=1}^{T-1} \ddot{\mathbf{X}}'_t \mathbf{P}_t \ddot{\mathbf{X}}_t = \sum_{i=1}^n \dot{\mathbf{X}}'_i \mathbf{Z}_{d,i} (\sum_{i=1}^n \mathbf{Z}'_{d,i} \mathbf{Z}_{d,i})^{-1} \sum_{i=1}^n \mathbf{Z}'_{d,i} \dot{\mathbf{X}}_i$. Similarly, $\sum_{t=1}^{T-1} \ddot{\mathbf{X}}'_t \mathbf{P}_t \ddot{\mathbf{y}}_t = \sum_{i=1}^n \dot{\mathbf{X}}'_i \mathbf{Z}_{d,i} (\sum_{i=1}^n \mathbf{Z}'_{d,i} \mathbf{Z}_{d,i})^{-1} \sum_{i=1}^n \mathbf{Z}'_{d,i} \dot{\mathbf{y}}_i$. From this and Eq. (2), we see that the FOD GMM estimator can be expressed as a TSLS estimator after applying the FOD transformation to the dependent and explanatory variables and upon using block-diagonal instrument matrices:

$$egin{aligned} \widehat{oldsymbol{eta}} &= \left[\sum_{i=1}^n \dot{oldsymbol{X}}_i' oldsymbol{Z}_{d,i} \left(\sum_{i=1}^n oldsymbol{Z}_{d,i}' oldsymbol{Z}_{d,i}
ight)^{-1} \sum_{i=1}^n oldsymbol{Z}_{d,i}' \dot{oldsymbol{X}}_i
ight]^{-1} \ & imes \sum_{i=1}^n \dot{oldsymbol{X}}_i' oldsymbol{Z}_{d,i} \left(\sum_{i=1}^n oldsymbol{Z}_{d,i}' oldsymbol{Z}_{d,i}
ight)^{-1} \sum_{i=1}^n oldsymbol{Z}_{d,i}' \dot{oldsymbol{y}}_i. \end{aligned}$$

It is well-known that TSLS is efficient GMM when the errors are conditionally homoskedastic and uncorrelated. Moreover, as already noted, if the $v_{i,t}$ s are conditionally homoskedastic and uncorrelated, then the transformed errors (the $\ddot{v}_{i,t}$ s) are conditionally homoskedastic and uncorrelated. Hence, if the $v_{i,t}$ s are conditionally homoskedastic and uncorrelated, the FOD GMM estimator is the asymptotically efficient GMM estimator given the moment restrictions $E(\mathbf{z}_{i,t}\ddot{v}_{i,t}) = \mathbf{0}$ $(t = 1, \ldots, T - 1)$. This is a total of $\sum_{t=1}^{T-1} q_t$ moment restrictions, which can be a large number when Tis large, especially if $q_T^* := \max_{1 \le t \le T-1} q_t$ —i.e., the maximum per-period number of instrumental variables—increases with T. Therefore, although the FOD GMM estimator is efficient when the errors are conditionally homoskedastic and uncorrelated, we might anticipate it to be biased when T is not small.

2.2 When there is no asymptotic bias

Alvarez and Arellano (2003) provide conditions under which the distribution of an FOD GMM estimator, $\hat{\beta}$, of the autoregression parameter, β , in the model in (1) has an asymptotic bias term if $T/n \to c > 0$, as $n, T \to \infty$. In particular, note that

$$\sqrt{nT}\left(\widehat{\beta} - \beta\right) - \theta_{n,T} = \frac{b_{n,T} - E(b_{n,T})}{a_{n,T}},\tag{4}$$

where $\theta_{n,T} := E(b_{n,T})/a_{n,T}$, $a_{n,T} := (1/(nT)) \sum_{t=1}^{T-1} \ddot{\boldsymbol{y}}_{t-1}' \boldsymbol{P}_t \ddot{\boldsymbol{y}}_{t-1}$, $b_{n,T} := (1/\sqrt{nT}) \sum_{t=1}^{T-1} \ddot{\boldsymbol{y}}_{t-1}' \boldsymbol{P}_t \ddot{\boldsymbol{v}}_t$, $\ddot{\boldsymbol{y}}_{t-1} := c_t(\boldsymbol{y}_{t-1} - \overline{\boldsymbol{y}}_{t-1})$, $\boldsymbol{y}'_{t-1} := (y_{1,t-1}, \dots, y_{n,t-1})$, and $\overline{\boldsymbol{y}}_{t-1} := (1/(T-t)) \sum_{s=1}^{T-t} \boldsymbol{y}_{t-1+s}$. If $a_{n,T}$ converges in probability to a fixed non-zero limit and $b_{n,T} - E(b_{n,T})$ has a limit distribution, then $\sqrt{nT}(\widehat{\beta} - \beta)$ has an asymptotic distribution that is centered at zero only if $\theta_{n,T} \xrightarrow{p} 0$, as $n, T \to \infty$. Alvarez and Arellano (2003) provide conditions that imply

$$a_{n,T} \xrightarrow{p} \frac{\sigma^2}{1-\beta^2}$$
 and $E(b_{n,T}) - \sqrt{\frac{T}{n}} \left(\frac{\sigma^2}{\beta-1}\right) \to 0$ $(n, T \to \infty),$

with $\sigma^2 := \operatorname{var}(v_{i,t})$ (see Alvarez and Arellano, 2003, p. 1128, Lemma 2). Using these results, they showed that if $\lim(T/n) = c$, with $0 \le c < \infty$, then

$$\sqrt{nT}\left(\widehat{\beta}-\beta\right) - \left[-\sqrt{T/n}\left(1+\beta\right)\right] \stackrel{d}{\to} N(0, 1-\beta^2) \qquad (n, T \to \infty).$$

(see Alvarez and Arellano, 2003, p.1129, Theorem 2). Therefore, for the AR(1) panel data model, $\sqrt{nT}(\hat{\beta} - \beta)$ has an asymptotic bias term of $\theta := -\sqrt{c}(1 + \beta)$, which is zero only when c = 0.

On the other hand, if $T/n \to c > 0$, then $\theta_{n,T} \xrightarrow{p} \theta \neq 0$, and

$$\sqrt{nT}\left(\widehat{\beta}-\beta\right) \xrightarrow{d} N(\theta, 1-\beta^2) \qquad (n, T \to \infty).$$
 (5)

The fact that $\sqrt{nT}(\hat{\beta}-\beta)$ has a bias term in its asymptotic distribution does not imply $\hat{\beta}$ is inconsistent. In fact, it is a consistent estimator of β (see Alvarez and Arellano,

2003, p.1129, Theorem 2).² However, because the asymptotic distribution of $\sqrt{nT}(\hat{\beta} - \beta)$ is not centered at zero, large sample confidence intervals and test statistics will be misleading. Specifically, the actual coverage of a confidence interval and the size of a test will differ from what they would be if the distribution of $\sqrt{nT}(\hat{\beta} - \beta)$ were centered at zero.

The case considered by Alvarez and Arellano is instructive for two reasons. First, it illustrates that $\theta_{n,T} \xrightarrow{p} 0$, as $n, T \to \infty$, is required for the asymptotic distribution of $\sqrt{nT}(\hat{\beta} - \beta)$ to be centered at zero. The second reason the example is instructive is less obvious and is the point of the first result (Theorem 1) provided here. The fact that $\theta_{n,T} \to \theta \neq 0$ when c > 0 is due to the number of instrumental variables that are used. Alvarez and Arellano (2003) assumed all available moment restrictions are exploited, in which case the maximum number of instrumental variables used in a period increases at the rate T increases, and the total number of moment restrictions increases at the rate T^2 increases. It has long been known that using many moment restrictions has deleterious effects on the bias of a GMM estimator. However, Theorem 1 sheds light on how many moment restrictions can be used without leading to large sample bias.

Theorem 1 provides conditions under which a generalization of $\theta_{n,T}$ converges in probability to a vector of zeros. Specifically, let $\mathbf{A}_{n,T} := (1/(nT)) \sum_{t=1}^{T-1} \ddot{\mathbf{X}}_t' \mathbf{P}_t \ddot{\mathbf{X}}_t$ and $\mathbf{b}_{n,T} := (1/\sqrt{nT}) \sum_{t=1}^{T-1} \ddot{\mathbf{X}}_t' \mathbf{P}_t \ddot{\mathbf{v}}_t$, with $\ddot{\mathbf{v}}_t := c_t (\mathbf{v}_t - \overline{\mathbf{v}}_t), \mathbf{v}_t' := (v_{1,t}, \dots, v_{n,t})$, and $\overline{\mathbf{v}}_t := (1/(T-t)) \sum_{s=1}^{T-t} \mathbf{v}_{t+s}$. Also, set $\boldsymbol{\theta}_{n,T} := \mathbf{A}_{n,T}^{-1} E(\mathbf{b}_{n,T})$. Then, analogous to (4), we have

$$\sqrt{nT}\left(\widehat{\boldsymbol{\beta}}-\boldsymbol{\beta}\right)-\boldsymbol{\theta}_{n,T}=\boldsymbol{A}_{n,T}^{-1}\left(\boldsymbol{b}_{n,T}-E(\boldsymbol{b}_{n,T})\right).$$

For every *n* and *T*, the distribution of $\mathbf{b}_{n,T} - E(\mathbf{b}_{n,T})$ is centered at a vector of zeros, **0**. It follows that if $\mathbf{A}_{n,T} \xrightarrow{p} \mathbf{A} > 0$,³ the distribution of $\sqrt{nT} \left(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta} \right) - \boldsymbol{\theta}_{n,T}$ becomes centered at **0** as the sample size grows. Therefore, whether or not the distribution of $\sqrt{nT}(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta})$ is centered at **0** depends on whether or not $\boldsymbol{\theta}_{n,T} \xrightarrow{p} \mathbf{0}$. If the distribution of $\sqrt{nT}(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta})$ is not centered at **0** in large samples, then the estimator $\hat{\boldsymbol{\beta}}$ will henceforth be described as having asymptotic bias. Theorem 1 shows that whether

²This conclusion follows from (5). This is because the limit distribution in (5) implies that, for large nT, $\hat{\beta}$ has an approximate normal distribution that is centered at $\beta + \theta/\sqrt{nT}$ with a variance of $(1 - \beta)/(nT)$. Hence, the limit distribution of $\hat{\beta}$, as $n, T \to \infty$, is degenerate at β , which implies $\hat{\beta}$ converges in probability to β .

³The notation $\boldsymbol{B} > 0$, when \boldsymbol{B} is a matrix, means \boldsymbol{B} is a positive definite matrix.

or not an FOD GMM estimator has asymptotic bias depends not just on the relative sizes of n and T but also on the number of instrumental variables used per period.

A few more definitions are needed in order to state Theorem 1. Let $\boldsymbol{w}_{i,t}$ be a column vector consisting of all of the distinct entries in $\{(\boldsymbol{x}'_{i,s}, \boldsymbol{z}'_{i,s}); s = 1, \ldots, t\}$. That is, $\boldsymbol{w}_{i,t}$ contains all of the distinct values of the explanatory variables and instrumental variables for the *i*th individual from the first period up to the *t*th period. Also, set $\boldsymbol{u}'_i := (\boldsymbol{w}'_{i,T}, v_{i,1}, \ldots, v_{i,T}, \eta_i)$, and let $\gamma_{k,t,s} := \operatorname{cov}(v_{1,t}, x_{1,t+s,k} | \boldsymbol{w}_{1,t})$.

In addition to these definitions, Theorem 1 relies on several conditions:

- A1: the u_i s are independent and identically distributed (i.i.d.) across i;
- A2: rank $(\mathbf{Z}_t) = q_t$ with probability 1 (wp1);
- A3: $E(v_{1,t}|\boldsymbol{w}_{1,t}) = 0;$
- A4: $\boldsymbol{A}_{n,T} \xrightarrow{p} \boldsymbol{A} > 0$; and
- A5: $\left|\sum_{s=1}^{S} \gamma_{k,t,s}\right| \leq M$ wp1, for some finite M and all $t \geq 1, S \geq 1$, and $k = 1, \dots, K$.

These conditions are weak enough to be satisfied by many dynamic regressions. For example, the estimation problems studied by Alvarez and Arellano (2003) and Hsiao and Zhou (2017), among others (e.g., Bun and Kiviet, 2006; Okui, 2009), are special cases of the estimation problem examined here.

The first four assumptions are relatively straightforward. Nevertheless, it is worthwhile to review them to clarify what is—and what is not—required. For example, Assumption A1 does not require that the errors be conditionally homoskedastic or time series homoskedastic. Moreover, the i.i.d. assumption across i can possibly be relaxed, but unfortunately relaxing the assumption would appear to require more technical detail.

The second assumption implies n must increase with T if the largest number of instrumental variables per period increases with T. Specifically, it implies $n \ge q_T^*$, and, therefore, if q_T^* increases with T, then n must likewise increase at least as fast as q_T^* increases. On the other hand, Assumption A2 does not rule out cases where nis fixed. For example, if at most q instruments are used each period and q does not increase with T, then n can be fixed while T increases provided $n \ge q$.

The third assumption implies the error in period t $(v_{i,t})$ is uncorrelated with the current and past explanatory variables. The assumption also implies the current error is uncorrelated with the instruments used in period t. However, it imposes no

additional restrictions on the instruments. The entries in $z_{i,t}$ can consist of, but need not be restricted to, current and past explanatory variables. Moreover, they can be in levels, differenced, or transformed in some other way. Regardless of what instrumental variables are used or how they are constructed, Assumption A3 allows for dynamic panel data regressions with additional predetermined explanatory variables, provided the $v_{i,t}$ s are uncorrelated across time.

As for the fourth assumption, it implicitly entails some familiar conditions which are typically taken for granted. To see this, first note that Assumption A2 ensures $[(1/n)\mathbf{Z}'_t\mathbf{Z}_t]^{-1}$ is defined (wp 1). This, in turn, implies $\Psi_{n,t} :=$ $(1/n)\ddot{\mathbf{X}}'_t\mathbf{Z}_t[(1/n)\mathbf{Z}'_t\mathbf{Z}_t]^{-1}(1/n)\mathbf{Z}'_t\ddot{\mathbf{X}}_t$ and $\mathbf{A}_{n,T} = (1/T)\sum_{t=1}^{T-1}\Psi_{n,t}$ are defined. But in order for $\Psi_{n,t}$ to be positive definite, the rank of $(1/n)\mathbf{Z}'_t\ddot{\mathbf{X}}_t$ must be K. That, in turn, requires the well-known necessary condition that $q_t \geq K$. In other words, it requires that we use at least as many instrumental variables each period as explanatory variables. Moreover, if the $\Psi_{n,t}$ s are positive definite, then the average of these positive definite matrices—that is, $\mathbf{A}_{n,T}$ —is positive definite, then $\hat{\boldsymbol{\beta}}$ is not defined. Therefore, it is not unreasonable to assume the average of the $\Psi_{n,t}$ s $(\mathbf{A}_{n,T})$ converges (in probability), and that the limit (\mathbf{A}) is—like $\mathbf{A}_{n,T}$ —positive definite. An example in which the condition is satisfied is provided by Alvarez and Arellano (2003).

The fifth assumption is a characterization of weak dependence. It is likely unfamiliar, but, like the other conditions, it does not appear to impose significant limitations on the types of panel data models to which the conclusion of Theorem 1 applies. Assumption A5 characterizes the linear association between the error in period tand the kth regressor s periods in the future relative to that period, conditional on the available information at time t. Because the conditional covariance between $v_{1,t}$ and $x_{1,t+s,k}$ is not restricted to be zero, Assumption A5 allows for predetermined regressors. Moreover, if the error term and future values of the explanatory variables are suitably weakly dependent, the bound on the sum of covariances in A5 will be satisfied.

This fact is illustrated by a stationary Kth order autoregressive (AR(K)) panel data model; see Lemma 1. Proofs are provided in the appendix.

Lemma 1. Let

 $\beta(L)y_{1,t} = \eta_1 + v_{1,t}$ $(t = 0, \pm 1, \pm 2, \ldots),$

⁴Not all of the $\Psi_{n,t}$ s need be positive definite in order for $A_{n,T}$ to be positive definite.

where $\beta(L) := 1 - \beta_1 L - \beta_2 L^2 - \cdots - \beta_K L^K$, and L is the lag operator. Also, let $w'_{1,t} = (y_{1,1-K}, \ldots, y_{1,t-1})$ $(t = 1, 2, \ldots, T)$. Assume the roots of the characteristic equation $1 - \beta_1 z - \beta_2 z^2 - \cdots - \beta_K z^K = 0$ all lie outside the unit circle. Also, assume the $v_{1,t}s$ are i.i.d. across t, with mean 0 and variance σ^2 , and η_1 and $v_{1,t}$ are independent for all t. Then Assumption A5 is satisfied.⁵

The AR(K) panel data model is just one example of the models that satisfy Assumption A5. In the Supplemental Appendix (Phillips, 2024), I show that A5 is satisfied by a variety of models. These models include—but are not limited to—panel vector autoregressions and other panel data models with predetermined regressors. The panel data models considered by Alvarez and Arellano (2003), Bun and Kiviet (2006), Hsiao and Zhou (2017), and Okui (2009), among others, are special cases of models that satisfy A5.

Theorem 1 can now be stated.

Theorem 1. Assume A1 through A5 are satisfied and that $T \ge 3$. If $q_T^* = O(T^{\alpha})$, then $\boldsymbol{\theta}_{n,T} \xrightarrow{p} \mathbf{0}$ as $T^{2\alpha}(\ln T)^2/(nT) \to 0$.

Theorem 1 shows that the absence of asymptotic bias depends on the relative rate of increase in n and T and on the largest number of instrumental variables used per period (q_T^*) . The maximum per-period number of instruments may depend on T. A bound on how fast q_T^* increases with T is parameterized in the theorem by α .

The case $\alpha = 1$ has been widely studied in the literature. If all available instrumental variables are used, and past lags of explanatory variables are viable instruments, then the maximum number of instruments used in a period increases with T at the rate T increases. This corresponds to $\alpha = 1$. Then $T^{2\alpha}(\ln T)^2/(nT) = T(\ln T)^2/n$, which clearly does not go to zero for all sequences of n and T. Instead, it only goes to zero for sequences of n and T for which either T does not increase or it increases slowly enough relative to n that $T(\ln T)^2/n \to 0$. The set of sequences for which $T(\ln T)^2/n \to 0$ is smaller than what Alvarez and Arellano (2003) found, for they found that there was no bias term if $T/n \to 0$. However, Alvarez and Arellano (2003) established their result for an AR(1) panel data model with i.i.d. homoskedastic errors. Theorem 1, on the other hand, provides sufficient conditions for how fast

⁵Because variables are assumed to be identically distributed across i, this lemma is stated in terms of a representative cross-sectional unit—the first (i = 1).

T can increase relative to n for more general panel data regressions and under weaker assumptions about the errors.

Moreover, although $\boldsymbol{\theta}_{n,T} \xrightarrow{p} \mathbf{0}$ is only guaranteed for some sequences of n and Twhen $\alpha = 1$, we get $T^{2\alpha}(\ln T)^2/(nT) \to 0$ for all sequences of n and T if $\alpha < 1/2$. Hence, Theorem 1 shows that $\sqrt{nT}(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta})$ has no asymptotic bias regardless of what happens to T/n, provided the maximum number of instrumental variables used in a period increases with T more slowly than $T^{1/2}$ increases.

An important special case that satisfies the restriction $\alpha < 1/2$ is when the number of instrumental variables used per period never exceeds some fixed number, say q. Then, $\alpha = 0$. Thus, if a researcher uses a fixed number of instruments each period, the FOD GMM estimator has no asymptotic bias regardless of how n and T increase. In fact, although it is unusual for n to be small while T is large, Theorem 1 nevertheless tells us there will be little bias in the distribution of $\sqrt{nT}(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta})$ even when n is relatively small provided T is large and $n \geq q$. This conclusion is similar to what is true for the fixed effects estimator when the explanatory variables are all strictly exogenous. That estimator, like the FOD GMM estimator, relies on eliminating fixed effects by subtracting from each variable a within cross-sectional average.

2.3 A sequential limit result

The absence of bias in the distribution of $\sqrt{nT}(\hat{\beta} - \beta)$ tells us only that. It does not tell us what distribution approximates the distribution of $\sqrt{nT}(\hat{\beta} - \beta)$ when n and Tare both large. To address that question, Hsiao and Zhou (2017) studied a choice of q_t for which $\alpha = 0$ in some detail. Specifically, they obtained asymptotic distribution results for FOD GMM estimators of the autoregression parameter in the AR(1) panel data model using a single instrumental variable per period. In doing so, they used sequential limits—specifically, they let $n \to \infty$, and then $T \to \infty$.⁶

Phillips and Moon remark that "Sequential limit theory is easy to derive and generally leads to quick results for a variety of model configurations" (Phillips and Moon, 1999, p. 1059). Moreover, sequential limit results can sometimes be obtained under weaker assumptions than those needed for joint limit results (Phillips and Moon, 1999, 2000). These advantages may explain why sequential limits have been used

⁶An alternative sequential-limit approach is to take limits in the order $T \to \infty$, then $n \to \infty$ (see, e.g., Phillips and Moon, 1999; Phillips and Moon, 2000; Kapetanios, 2008).

to analyze estimator properties when both n and T are large (see, e.g., Kapetanios, 2008; Hsiao and Zhang, 2015; Hsiao and Zhou, 2017).

However, taking limits sequentially has a drawback: sequential limit results are not always robust. Specifically, a sequential limit is not, in general, guaranteed to be equal to a joint limit (Phillips and Moon, 2000).⁷ Indeed, in the present application, letting $n \to \infty$ first, and then $T \to \infty$, always leads to the conclusion that there is no asymptotic bias in the distribution of $\sqrt{nT}(\hat{\beta} - \beta)$ regardless of q_T^* . This is illustrated by Theorem 2

Before stating Theorem 2, some additional assumptions and definitions are needed. Specifically, assume

A6: the entries in $\boldsymbol{x}_{1,t}$ have finite second-order moments, while $\ddot{v}_{1,t}$ and the entries in $\boldsymbol{z}_{1,t}$ have finite fourth-order moments for all t.

Assumption A6 implies the entries in the matrix $C_t := E(\boldsymbol{z}_{1,t} \boldsymbol{\ddot{x}}_{1,t})$ are finite. Similarly, the entries in the matrices $\boldsymbol{Q}_t := E(\boldsymbol{z}_{1,t} \boldsymbol{z}_{1,t}')$, and $E(\boldsymbol{\ddot{v}}_{1,s} \boldsymbol{\ddot{v}}_{1,t} \boldsymbol{z}_{1,s} \boldsymbol{z}_{1,t}')$ are finite. Moreover, given Assumption A2 implies \boldsymbol{Q}_t is positive definite, if Assumptions A6 and A2 are both satisfied, we can define $\boldsymbol{\Sigma}_{t,t+s} := \boldsymbol{\Pi}_t' E(\boldsymbol{\ddot{v}}_{1,t} \boldsymbol{\ddot{v}}_{1,t+s} \boldsymbol{z}_{1,t+s}) \boldsymbol{\Pi}_{t+s}$, where $\boldsymbol{\Pi}_t := \boldsymbol{Q}_t^{-1} \boldsymbol{C}_t$. Also, let $\boldsymbol{\Omega}_T = (1/T) \sum_{t=1}^{T-1} \boldsymbol{\Sigma}_{t,t} + (1/T) \sum_{t=1}^{T-2} \sum_{s=1}^{T-1-t} (\boldsymbol{\Sigma}_{t,t+s} + \boldsymbol{\Sigma}_{t+s,t})$. Then the last set of assumptions for Theorem 2 are

A7: $\mathbf{\Omega} := \lim_{T \to \infty} \mathbf{\Omega}_T$ exists; and

A8:
$$\boldsymbol{A} := \lim_{T \to \infty} (1/T) \sum_{t=1}^{T-1} \boldsymbol{C}'_t \boldsymbol{Q}_t^{-1} \boldsymbol{C}_t > 0$$
 exists

Theorem 2. Assume A1 through A3 and A6 through A8 are satisfied. Then

$$\sqrt{nT}\left(\widehat{\boldsymbol{\beta}}-\boldsymbol{\beta}\right) \stackrel{d}{\rightarrow} N\left(\boldsymbol{0}, \boldsymbol{A}^{-1}\boldsymbol{\Omega}\boldsymbol{A}^{-1}\right) \qquad (n, T \to \infty)_{seq}.$$

The notation " $(n, T \to \infty)_{seq}$ " means $n \to \infty$, then $T \to \infty$.

Theorem 2 illustrates the problem with taking limits sequentially in the present context. Specifically, Theorem 2 says the asymptotic distribution of $\sqrt{nT}(\hat{\beta} - \beta)$ is centered at **0**, and it does so without referring to how many instrumental variables

⁷Phillips and Moon (1999) provide conditions that, if satisfied, allows sequential limit results to be strengthened to joint limit results. However, their paper focuses on taking sequential limits with $T \to \infty$, and then $n \to \infty$. Moreover, even for limits are taken in this order, it can be difficult to verify the conditions that imply a sequential limit equals a joint limit.

are used. However, as has already been noted, Alvarez and Arellano (2003) showed that if all instrumental variables are used, the FOD GMM estimator they considered has a nonzero bias term in its asymptotic distribution if $T/n \rightarrow c > 0$, as $n, T \rightarrow \infty$. Alvarez and Arellano examined estimation of the AR(1) panel data model, which is a special case of the models covered by Theorem 2. Therefore, Theorem 2 contradicts the result provided in Alvarez and Arellano (2003).

The contradiction is not the fault of Alvarez and Arellano. Instead, it arises from how limits are executed. Taking a limit with $n \to \infty$ first, implicitly holds T fixed initially, and GMM estimators do not generally have asymptotic bias, as $n \to \infty$, in the fixed T case. Hence, the bias term is gone by the time the second step—which consists of letting $T \to \infty$ —is reached.

On the other hand, under the conditions of Theorem 1, there is also no bias term regardless of how n and T increase, provided the maximum number of per-period instruments, q_T^* , is selected so that $\alpha < 1/2$. In the next section, the accuracy of the normal approximation, for large n and T and $\alpha = 0$, is investigated with Monte Carlo experiments.

3 Simulations

Phillips (2020) provides Monte Carlo evidence that supports the claim that the distribution of $\sqrt{nT}(\hat{\beta} - \beta)$ is centered at **0** when, for example, $\alpha = 0$ and T is not quite small compared to n. That paper reports Monte Carlo evidence illustrating that when the per-period number of instrumental variables is fixed, FOD GMM generally has less bias and is more efficient than one-step first difference GMM (FD GMM). In those experiments, the FOD GMM estimator was also less biased and usually more efficient than a two-step FD GMM in the presence of heteroskedastic errors, provided T is not particularly small (e.g., T = 40, n = 200).⁸

This section, on the other hand, reports on Monte Carlo experiments that were used to investigate the reliability of confidence intervals based on the FOD GMM estimator. When q_T^* was chosen so that $\alpha = 0$, the coverage of the confidence intervals based on the FOD GMM estimator turned out to be consistent with the conclusion of Theorem 2. Specifically, for these experiments, the coverage of the FOD GMM confidence intervals, which relied on the normal approximation in Theorem 2, was

⁸See Arellano and Bond (1991) for a description of one-step and two-step FD GMM.

accurate for large T. In addition to confidence intervals, this section compares the estimator's precision, as measured by root mean squared error (RMSE), to the precision of the efficient GMM estimator.

3.1 Samples

Monte Carlo samples were generated using a sampling scheme similar to one of the sampling schemes described in Bun and Kiviet (2006).⁹ In particular, the dependent variable in the regression model was generated according to

$$y_{i,t} = \beta_1 y_{i,t-1} + \beta_2 x_{i,t} + \eta_i + v_{i,t} \qquad (t = -49, \dots, -1, 0, 1, \dots, T, \ i = 1, \dots, n),$$

with $\beta_1 \in \{0.25, 0.75\}$ and $\beta_2 = 1 - \beta_1$. The start-up value for $y_{i,-50}$ was zero. Moreover, the error components $v_{i,t}$ and η_i were generated independently of one another as i.i.d. standard normal variates. As for the explanatory variable $x_{i,t}$, it was generated as

$$x_{i,-50} = \kappa_1 \eta_i + \varepsilon_{i,-50},$$

$$x_{i,-49} = \kappa_1 \eta_i + \frac{1}{1 - \rho L} (\varepsilon_{i,-49} + \phi_1 v_{i,-50})$$

$$x_{i,t} = \kappa_1 \eta_i + \frac{1}{1 - \rho L} \varepsilon_{i,t} + \phi_1 v_{i,t-1} \qquad (t = -48, \dots, -1, 0, +1, \dots, T),$$

with $\rho \in \{0.50, 0.95\}$, $\kappa_1 \in \{-1, 0, +1\}$, and $\phi_1 \in \{-1, 0, +1\}$. Moreover, the $\varepsilon_{i,t}$ s were generated as $\varepsilon_{i,t} \stackrel{i.i.d.}{\sim} U(-\sqrt{12}/2, \sqrt{12}/2)$ independently of the $v_{i,t}$ s and η_i s.

I considered a total of 36 experimental designs, where a design is a combination of parameter values. Table 1 lists Designs 1 through 18. For these designs, I set $\beta_1 = 0.25$. Designs 19 through 36 were identical to Designs 1 through 18, except that β_1 was set to 0.75 for Designs 19 through 36. The latter designs will henceforth be described as the weak instruments designs, for it is well-known that lagged values of the dependent variable become weaker instruments as β_1 approaches one (see, e.g., Blundell and Bond, 1998).

The effect of how each series was initialized was eliminated by dropping the first 50 values of each generated time series. As a result, estimation was based on the

⁹Bun and Kiviet (2006) considered two sampling schemes, and sampling schemes similar to both were initially considered. However, the results were qualitatively similar across the two schemes. Therefore, for the sake of brevity, the results for only one of them are reported here.

				Do	zione				
	1	2	3	4 Dec	5 signs	6	7	8	9
	-	-	0	Ĩ	0	Ŭ	·	0	Ū
ρ	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50
ϕ_1	-1.00	-1.00	-1.00	0.00	0.00	0.00	1.00	1.00	1.00
κ_1	-1.00	0.00	1.00	-1.00	0.00	1.00	-1.00	0.00	1.00
				-					
				Des	signs				
	10	11	12	13	14	15	16	17	18
ρ	0.95	0.95	0.95	0.95	0.95	0.95	0.95	0.95	0.95
ϕ_1	-1.00	-1.00	-1.00	0.00	0.00	0.00	1.00	1.00	1.00
κ_1	-1.00	0.00	1.00	-1.00	0.00	1.00	-1.00	0.00	1.00

Table 1: Designs 1 through 18 ($\beta_1 = 0.25$).

values $(x_{i,0}, y_{i,0}), (x_{i,1}, y_{i,1}), \ldots, (x_{i,T}, y_{i,T})$ $(i = 1, \ldots, n)$. Moreover, I set n = 200. The number of time periods, T, was set equal to 20, then 40, and finally 100 in order to examine the effect of increasing T/n on confidence intervals and estimator precision. Finally, for each sample size and parameter design, 5,000 samples were generated. Estimates were calculated for each of these 5,000 samples.¹⁰

3.2 Estimators

After a sample was generated, three estimates of $\boldsymbol{\beta} = (\beta_1, \beta_2)'$ were calculated: an FOD GMM estimate, an FD GMM estimate, and an efficient GMM estimate. For the FD and FOD GMM estimates, I set $\boldsymbol{z}'_{i,1} = (y_{i,0}, x_{i,0}, x_{i,1})$ and $\boldsymbol{z}'_{i,t} = (y_{i,t-2}, y_{i,t-1}, x_{i,t-2}, x_{i,t-1}, x_{i,t})$ $(t = 2, \ldots, T - 1, i = 1, \ldots, n)$. Given these instrumental variables, FOD GMM estimates were calculated using the formula in (2).

¹⁰The data were generated using GAUSS. Moreover, all calculations were performed with GAUSS.

The FD GMM estimates, on the other hand, were calculated using the formula

$$\widetilde{\boldsymbol{\beta}} := \left[\sum_{i=1}^{n} \widetilde{\boldsymbol{X}}_{i}^{\prime} \boldsymbol{Z}_{d,i} \left(\sum_{i=1}^{n} \boldsymbol{Z}_{d,i}^{\prime} \boldsymbol{G} \boldsymbol{Z}_{d,i} \right)^{-1} \sum_{i=1}^{n} \boldsymbol{Z}_{d,i}^{\prime} \widetilde{\boldsymbol{X}}_{i} \right]^{-1} \\ \times \sum_{i=1}^{n} \widetilde{\boldsymbol{X}}_{i}^{\prime} \boldsymbol{Z}_{d,i} \left(\sum_{i=1}^{n} \boldsymbol{Z}_{d,i}^{\prime} \boldsymbol{G} \boldsymbol{Z}_{d,i} \right)^{-1} \sum_{i=1}^{n} \boldsymbol{Z}_{d,i}^{\prime} \widetilde{\boldsymbol{y}}_{i}.$$

$$(6)$$

In this formula, $\tilde{\boldsymbol{y}}'_i := (y_{i,2} - y_{i,1}, \dots, y_{i,T} - y_{i,T-1})$; $\tilde{\boldsymbol{X}}_i$ stacks $\tilde{\boldsymbol{x}}'_{i,t+1} := \boldsymbol{x}'_{i,t+1} - \boldsymbol{x}'_{i,t}$, with $\boldsymbol{x}'_{i,t} = (y_{i,t-1}, x_{i,t})$ $(t = 1, \dots, T-1)$; and $\boldsymbol{Z}_{d,i}$ is defined in (3). Also, \boldsymbol{G} is a $(T-1) \times (T-1)$ matrix with twos running down the main diagonal, minus ones just above and below the main diagonal, and zeros everywhere else (see, e.g., Arellano and Bond, 1991).

Finally, efficient GMM estimates were also calculated. The efficient GMM estimator exploited all available instrumental variables. For this estimator, I set $\mathbf{z}'_{i,1} = (y_{i,0}, x_{i,0}, x_{i,1})$ and $\mathbf{z}'_{i,t} = (\mathbf{z}'_{i,t-1}, y_{i,t-1}, x_{i,t})$ $(t = 2, \ldots, T - 1, i = 1, \ldots, n)$. To calculate efficient estimates, either the formula in (2) or the formula in (6) can be used, because when all available instrumental variables are used, the two formulas give numerically identical results (see, e.g., Phillips, 2019). Although (2) and (6) give the same estimate in this case, the formula in (2) was used to calculate estimates because it is more efficient computationally (Phillips, 2020). Moreover, efficient estimates were not calculated for T = 100, because the requirement that n be no smaller than q_t was not satisfied—and hence $(\mathbf{Z}'_t \mathbf{Z}_t)^{-1}$ was not defined—for all t when Tis this large relative to n and all available instruments are used. Finally, because FD GMM and FOD GMM are the same when all available instruments are used, the efficient GMM estimator will henceforth be referred to as the FD/FOD GMM estimator.

In order to construct confidence intervals, standard errors were also calculated. Their calculation was simplified by the fact that the experimental $v_{i,t}$ s were conditionally homoskedastic and uncorrelated. In this case, the variance-covariance matrix of $\sqrt{nT}(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta})$ simplifies to $\sigma^2 \boldsymbol{A}^{-1}$. Therefore, standard errors for the FOD GMM estimates were calculated by taking the square roots of the diagonal entries of $\hat{\sigma}^2 \left(\sum_{t=1}^{T-1} \ddot{\boldsymbol{X}}'_t \boldsymbol{P}_t \ddot{\boldsymbol{X}}_t \right)^{-1}$, where $\hat{\sigma}^2 := \sum_{t=1}^{T-1} (\ddot{\boldsymbol{y}}_t - \ddot{\boldsymbol{X}}_t \hat{\boldsymbol{\beta}})'(\ddot{\boldsymbol{y}}_t - \ddot{\boldsymbol{X}}_t \hat{\boldsymbol{\beta}})/[n(T-1)]$. Moreover, when all available instruments were used, standard errors for FD/FOD estimates were calculated similarly. On the other hand, the standard errors for the FD estimator, $\tilde{\boldsymbol{\beta}}$, were calculated using the square roots of the diagonal entries in $\tilde{\sigma}^2 \left[\sum_{i=1}^n \tilde{\boldsymbol{X}}'_i \boldsymbol{Z}_{d,i} \left(\sum_{i=1}^n \boldsymbol{Z}'_{d,i} \boldsymbol{G} \boldsymbol{Z}_{d,i} \right)^{-1} \sum_{i=1}^n \boldsymbol{Z}'_{d,i} \tilde{\boldsymbol{X}}_i \right]^{-1}$, where $\tilde{\sigma}^2 := \sum_{i=1}^n (\tilde{\boldsymbol{y}}_i - \tilde{\boldsymbol{X}}_i \tilde{\boldsymbol{\beta}})/[2n(T-1)]$.

3.3 Simulation results

3.3.1 Confidence intervals

Tables 2 through 5 report the coverage of 95 percent confidence intervals for β_1 and β_2 using the FOD, FD, and FD/FOD GMM estimators, their respective standard errors, and the normal approximation—specifically, each interval extends from 1.96 standard errors below a regression parameter estimate to 1.96 standard errors above it. Tables 2 and 3 provide the coverage of the confidence intervals for β_1 and β_2 , respectively, for Designs 1 through 18, and Tables 4 and 5 provide the coverage of the confidence intervals for β_1 and β_2 for Designs 19 through 36.

Coverage estimates for the FD/FOD confidence intervals are only provided for T = 20. This is the best case for the FD/FOD GMM estimator because the estimator was developed for situations where T is small compared to n. Nevertheless, even in this case the coverage of the FD/FOD confidence intervals is often unreliable. For example, consider the FD/FOD GMM intervals for β_1 given weak instruments (Table 4). For Designs 19 through 36, the coverage of the FD/FOD GMM intervals for β_1 ranged from a low of 51.8 percent (Design 27) to a high of 82.7 percent (Design 28). Unsurprisingly, increasing T to 40 made for even worse coverage. This result is consistent with Alvarez and Arellano's finding of non-zero asymptotic bias in the distribution of the FOD GMM estimator of the AR(1) panel data model when T is not small compared to n and all available moment restrictions are exploited.

Using fewer than all available instrumental variables improved the reliability of the confidence intervals, as expected, but by how much depended on how the data were transformed to remove fixed effects. Usually, the coverage of the FOD confidence intervals approximated 95 percent at least as well, if not better, than did the FD confidence intervals. Moreover, confidence intervals based on the FOD GMM estimator were no less reliable for T = 100 than for T = 20. Indeed, the reliability of the FOD confidence intervals appears to improve as T increases relative to n; see, for example, the FOD confidence intervals for β_1 and Designs 19 through 36 (Table 4). On the other hand, the FD confidence intervals often under estimated 95 percent, and their

Estimator	T										
		Designs:	1	2	3	4	5	6	7	8	9
FD/FOD	20		95.0	95.7	94.5	90.6	90.9	91.5	90.3	89.0	88.1
FD:	20		95.2	95.3	94.5	93.3	94.1	94.0	93.8	93.3	92.9
FD:	40		94.7	95.0	94.0	93.4	93.8	92.8	93.0	92.7	92.4
FD	100		95.2	94.9	94.7	91.1	90.2	91.5	89.8	89.0	88.8
FOD:	20		95.0	95.4	94.7	94.7	95.4	94.6	95.2	94.5	94.5
FOD:	40		94.6	95.0	94.8	94.9	95.6	94.8	94.8	95.1	94.5
FOD	100		95.1	94.9	94.9	94.5	94.8	95.2	94.9	94.9	95.0
		Designs:	10	11	12	13	14	15	16	17	18
FD/FOD	20		95.0	95.4	94.7	91.9	91.2	91.0	90.7	90.4	89.7
FD:	20		94.9	94.6	94.9	94.2	94.5	94.1	94.3	93.2	93.9
FD:	40		94.4	94.5	95.4	93.1	93.6	94.2	93.4	92.1	93.1
FD	100		94.3	95.2	95.0	92.9	92.5	92.7	90.9	91.1	91.0
FOD:	20		95.2	95.5	94.7	95.1	95.3	94.6	94.6	94.3	94.9
FOD:	40		94.9	95.3	95.4	95.0	94.8	95.4	95.1	94.7	94.4
FOD	100		94.7	95.2	95.4	94.9	94.7	94.7	94.9	94.4	94.6

Table 2: Coverage of 95-percent confidence intervals for β_1 , Designs 1–18 (n = 200).

Estimator	T										
		Designs:	1	2	3	4	5	6	7	8	9
FD/FOD	20		94.1	93.9	93.2	94.8	95.2	95.2	95.2	94.7	94.6
FD:	20		94.8	94.5	94.7	95.3	94.7	94.9	95.4	94.7	94.6
FD:	40		94.0	94.2	94.0	94.9	95.2	94.6	94.8	95.2	94.9
FD	100		93.1	93.3	93.4	94.5	94.9	94.5	95.0	95.3	95.6
FOD:	20		95.1	94.8	94.8	95.7	95.3	94.8	95.4	94.9	94.5
FOD:	40		94.9	94.7	95.1	95.3	95.0	95.1	94.7	95.4	95.2
FOD	100		95.0	94.9	95.6	94.9	94.2	94.9	95.1	95.7	95.5
		Designs:	10	11	12	13	14	15	16	17	18
FD/FOD	20		93.4	92.4	92.8	94.4	94.3	94.5	94.6	94.8	94.7
FD:	20		95.0	94.0	94.3	95.1	95.1	94.8	94.7	95.0	95.0
FD:	40		94.0	94.0	94.6	94.8	94.4	95.5	95.0	94.6	94.7
FD	100		93.8	93.7	93.6	94.5	94.5	94.6	94.6	94.9	94.4
FOD:	20		94.8	94.7	94.2	95.0	94.9	95.1	94.4	95.0	95.0
FOD:	40		95.0	95.5	94.4	94.7	95.0	95.5	95.3	94.7	94.9
FOD	100		95.3	95.1	95.0	95.1	94.8	94.8	95.0	94.9	94.1

Table 3: Coverage of 95-percent confidence intervals for β_2 , Designs 1–18 (n = 200).

Estimator	T										
		Designs:	19	20	21	22	23	24	25	26	27
FD/FOD	20		67.6	62.6	61.4	62.7	56.2	52.0	66.1	58.8	51.8
FD:	20		87.8	82.9	79.4	85.9	84.4	82.1	86.7	84.8	82.0
FD:	40		87.5	86.4	84.6	86.1	85.3	83.7	87.7	86.5	85.4
FD	100		83.1	83.1	82.9	80.7	79.2	79.7	81.8	81.2	81.3
FOD:	20		93.5	92.1	91.2	93.3	92.3	91.3	93.8	93.0	91.5
FOD:	40		94.3	94.3	94.0	94.0	93.8	93.7	94.6	93.9	93.7
FOD	100		95.2	94.5	94.3	94.9	94.9	94.4	94.7	94.7	95.1
		Designs:	28	29	30	31	32	33	34	35	36
FD/FOD	20		82.7	81.8	81.7	70.8	72.2	70.6	69.1	67.5	65.3
FD:	20		92.4	91.5	91.6	89.6	89.9	89.2	89.7	89.0	88.4
FD:	40		93.1	92.0	92.3	90.5	90.3	90.6	90.1	89.9	90.0
FD	100		91.7	91.7	92.3	88.4	88.1	88.1	87.7	87.0	87.4
FOD:	20		94.5	94.3	94.1	93.8	94.3	93.5	94.1	93.6	93.5
FOD:	40		94.6	94.9	94.6	94.0	94.0	94.8	94.6	94.2	94.5
FOD	100		94.7	95.0	94.7	94.9	94.7	94.4	94.9	94.9	94.7

Table 4: Coverage of 95-percent confidence intervals for β_1 , Designs 19–36 (n = 200).

Estimator	T										
		Designs:	19	20	21	22	23	24	25	26	27
FD/FOD	20		92.2	87.3	83.8	94.1	94.4	92.1	94.9	94.3	94.2
FD:	20		92.2	88.0	83.6	94.9	93.5	91.7	95.4	95.0	94.0
FD:	40		93.6	91.6	90.5	95.0	94.6	93.9	94.5	94.9	95.1
FD	100		92.4	92.3	91.6	95.2	95.0	95.0	94.9	94.5	94.4
FOD:	20		94.8	93.0	92.3	94.9	94.6	94.2	95.3	94.9	94.7
FOD:	40		94.7	94.5	94.3	94.5	94.7	95.1	94.5	95.0	94.8
FOD	100		95.0	94.7	94.5	95.5	95.3	94.6	95.7	94.9	94.9
		Designs:	28	29	30	31	32	33	34	35	36
FD/FOD	20		94.2	94.7	94.7	93.9	93.9	93.5	94.4	94.7	94.6
FD:	20		94.6	93.9	94.4	94.6	94.1	93.9	94.4	94.9	94.5
FD:	40		95.4	94.8	94.8	94.1	95.0	95.1	94.6	94.1	94.2
FD	100		94.7	94.7	95.0	94.2	94.8	94.8	93.8	94.2	94.8
FOD:	20		94.8	94.9	95.3	95.2	94.9	94.4	94.5	95.0	94.9
FOD: FOD:	$\begin{array}{c} 20\\ 40 \end{array}$		$94.8 \\ 94.7$	$94.9 \\ 95.1$	$95.3 \\ 94.7$	$95.2 \\ 94.7$	$\begin{array}{c} 94.9\\ 94.3\end{array}$	$94.4 \\ 95.2$	$94.5 \\ 94.7$	$95.0 \\ 94.7$	$94.9 \\ 94.4$
FOD: FOD: FOD	$20 \\ 40 \\ 100$		94.8 94.7 94.7	$94.9 \\ 95.1 \\ 94.9$	95.3 94.7 94.9	95.2 94.7 95.2	94.9 94.3 94.8	94.4 95.2 94.4	94.5 94.7 94.5	$95.0 \\ 94.7 \\ 94.9$	94.9 94.4 95.5

Table 5: Coverage of 95-percent confidence intervals for β_2 , Designs 19–36 (n = 200).

reliability did not improve as T was increased relative to n; see, for example, Table 4.¹¹

Moreover, using critical values from the standard normal distribution, 90 percent and 50 percent confidence intervals were also calculated. For the sake of brevity, the coverage of these intervals is omitted and is only summarized here: their reliability was similar to that of the 95 percent confidence intervals. For example, for large nand T, the coverage of the 90 and 50 percent FOD confidence intervals approximated 90 and 50 percent, respectively. Apparently for these experiments, the FOD GMM estimator has little bias and the normal approximation works well when n and T are both large and q = 0. On the other hand, the coverage of the 90 and 50 percent FD confidence intervals often under estimated 90 and 50 percent, especially for the weak instrument designs (Designs 19 through 36).

3.3.2 Relative precision

Although confidence intervals based on the FOD GMM estimator have superior coverage accuracy, the estimator sacrifices estimation efficiency to achieve this superior accuracy. In the experiments considered here, the FD and FOD GMM estimators use at most five instrumental variables each period. The FD/FOD GMM estimator, on the other hand, exploits all available moment restrictions. Therefore, the FD and FOD GMM estimators are inefficient relative to the FD/FOD GMM estimator. However, for $\alpha < 1/2$, the FOD GMM estimator does not have asymptotic bias, and, therefore, it can improve on the FD and FD/FOD GMM estimators in terms of precision, as measured by RMSE. The Monte Carlo experiments were also used to investigate this possibility.

Figures 1 through 4 are line plots of relative precision estimates for the 36 designs. A relative precision estimate is an RMSE ratio. Specifically, an FD relative precision estimate is an FD RMSE divided by the corresponding FD/FOD RMSE, and a FOD relative precision estimate is an FOD RMSE divided by the FD/FOD RMSE. Figures 1 through 4 plot the FD and FOD GMM relative precision estimates for the 36 designs. Red line plots graph the FD relative precisions, whereas blue line plots graph the FOD relative precisions. Figures 1 and 2 provide relative precisions for β_1 for T = 20 and

¹¹The finding that the coverage of the FD confidence intervals did not improve as T was increased is consistent with analytical results Hsiao and Zhou (2017) provide for FD GMM estimation of the AR(1) panel data model.



Figure 1: RMSE of Estimator of β_1 Over RMSE of FD/FOD Estimator (T = 20).

Figure 2: RMSE of Estimator of β_1 Over RMSE of FD/FOD Estimator (T = 40).





Figure 3: RMSE of Estimator of β_2 Over RMSE of FD/FOD Estimator (T = 20).

Figure 4: RMSE of Estimator of β_2 Over RMSE of FD/FOD Estimator (T = 40).



T = 40, whereas Figures 3 and 4 give the same for β_2 . For all estimates, n = 200.

The plots make conclusions about relative precision obvious visually. If an estimator's relative precision plot is above one, the estimator is less precise than the FD/FOD GMM estimator in terms of RMSE. If the estimator's plot is below one, the estimator is more precise than the FD/FOD estimator.

The plots illustrate the effect of the absence, or presence, of bias on precision. For example, although the FOD GMM estimator of β_1 has a larger variance than the corresponding FD/FOD GMM estimator, the bias of the FOD estimator is less—in fact, much less, especially as T increases relative to n. Hence, the FOD estimator of β_1 is usually more precise than the FD/FOD estimator. On the other hand, the FD/FOD GMM estimator of β_2 generally had smaller RMSE than the FOD GMM estimator of β_2 for T = 20. The FOD estimator of β_2 , however, became more competitive relative to the FD/FOD estimator in terms of RMSE for T = 40. Again, finite sample bias, or its absence, provides the explanation. The finite sample bias of the FD/FOD GMM estimator was less pronounced when estimating β_2 than it was when β_1 was estimated, especially for T = 20.

On the other hand, the FD GMM estimators of β_1 and β_2 have larger variances than the FD/FOD GMM estimators, but, unlike the FOD estimators, they do not benefit from having much less bias than the FD/FOD estimators. Consequently, the FD estimators of β_1 and β_2 are less precise than the FD/FOD estimators of β_1 and β_2 .

4 Conclusion

When using panel data to estimate a dynamic regression, researchers typically remove fixed effects by transforming the data. Many transformations are available, but—due to historical precedence—differencing the data became a widely adopted approach. However, when not all of the available instrumental variables are used, how the data are transformed matters, and differencing may not be the best transformation.

The results provided in this paper show that transforming the data using forward orthogonal deviations produces a GMM estimator, $\hat{\beta}$, such that whether or not the distribution of $\sqrt{nT}(\hat{\beta} - \beta)$ is centered at a vector of zeros, as $n, T \to \infty$, depends on how fast the largest number of instruments used in a period increases with T. For the absence of large sample bias, this number must increase more slowly than $T^{1/2}$ increases. This observation is important because the reliability of large sample confidence intervals and test statistics based on an FOD GMM estimator depends on the distribution of $\sqrt{nT}(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta})$ being centered at a vector of zeros.

Appendix: Proofs

The proof of Theorem 1 draws on Lemma A.1.

Lemma A.1. Assume $rank(\mathbf{Z}_t) = q_t$ wp1 and define $p_{n,t} := n \mathbf{z}'_{1,t} (\mathbf{Z}'_t \mathbf{Z}_t)^{-1} \mathbf{z}_{1,t}$. If $\mathbf{z}_{i,t}$ is identically distributed across *i*, then $E(p_{n,t}) = q_t$.

Proof: By a well-known result for the trace of a projection matrix, we have $\operatorname{tr}(\boldsymbol{P}_t) = \operatorname{rank}(\boldsymbol{Z}_t)$. And $\operatorname{rank}(\boldsymbol{Z}_t) = q_t$ wp1 by assumption. But $\operatorname{tr}(\boldsymbol{P}_t) = \sum_{i=1}^n \boldsymbol{z}'_{i,t} (\boldsymbol{Z}'_t \boldsymbol{Z}_t)^{-1} \boldsymbol{z}_{i,t}$. Hence, $\sum_{i=1}^n E[\boldsymbol{z}'_{i,t} (\boldsymbol{Z}'_t \boldsymbol{Z}_t)^{-1} \boldsymbol{z}_{i,t}] = q_t$. Therefore, if $\boldsymbol{z}_{i,t}$ is identically distributed across i, then $nE[\boldsymbol{z}'_{1,t} (\boldsymbol{Z}'_t \boldsymbol{Z}_t)^{-1} \boldsymbol{z}_{1,t}] = q_t$.

Theorem 1 proof

First note that

$$\sqrt{nT}\boldsymbol{b}_{n,T} = \sum_{t=1}^{T-1} c_t^2 \boldsymbol{X}_t' \boldsymbol{P}_t \left(\boldsymbol{v}_t - \overline{\boldsymbol{v}}_t \right) - \sum_{t=1}^{T-1} c_t^2 \overline{\boldsymbol{X}}_t' \boldsymbol{P}_t \left(\boldsymbol{v}_t - \overline{\boldsymbol{v}}_t \right).$$
(7)

Moreover, Assumption A1 implies $(\boldsymbol{w}'_{i,t}, v_{i,t+s})$ is independent of $\boldsymbol{w}_{j,t}$ for $i \neq j$. Thus, if we set $\boldsymbol{w}'_t := (\boldsymbol{w}'_{1,t}, \dots, \boldsymbol{w}'_{n,t})$, then $E(v_{i,t+s}|\boldsymbol{w}_t) = E(v_{i,t+s}|\boldsymbol{w}_{i,t})$. And, A1, A3, and the law of iterated expectations imply $E(v_{i,t+s}|\boldsymbol{w}_{i,t}) = E[E(v_{i,t+s}|\boldsymbol{w}_{i,t+s})|\boldsymbol{w}_{i,t}] =$ $E(0|\boldsymbol{w}_{i,t}) = 0$ ($s \geq 0$). Hence, $E(\boldsymbol{v}_t - \overline{\boldsymbol{v}}_t|\boldsymbol{w}_t) = \boldsymbol{0}$, and thus $E[\boldsymbol{X}'_t\boldsymbol{P}_t(\boldsymbol{v}_t - \overline{\boldsymbol{v}}_t)] =$ $E[\boldsymbol{X}'_t\boldsymbol{P}_tE(\boldsymbol{v}_t - \overline{\boldsymbol{v}}_t|\boldsymbol{w}_t)] = \boldsymbol{0}$. From this observation and Eq. (7), it follows that

$$\sqrt{nT}E(\boldsymbol{b}_{n,T}) = -\sum_{t=1}^{T-1} \frac{1}{T-t+1} \left(\boldsymbol{s}_{1,T-t} - \boldsymbol{s}_{2,T-t} \right),$$
(8)

where $\boldsymbol{s}_{1,T-t} := \sum_{s=1}^{T-t} E\left(\boldsymbol{X}_{t+s}' \boldsymbol{P}_t \boldsymbol{v}_t\right)$ and $\boldsymbol{s}_{2,T-t} := \sum_{s=1}^{T-t} E\left(\boldsymbol{X}_{t+s}' \boldsymbol{P}_t \overline{\boldsymbol{v}}_t\right)$.

In order to evaluate $s_{1,T-t}$, note that the kth entry of $E(X'_{t+s}P_tv_t)$ is

$$\sum_{i=1}^{n} E\left(x_{i,t+s,k}\boldsymbol{z}_{i,t}'(\boldsymbol{Z}_{t}'\boldsymbol{Z}_{t})^{-1}\boldsymbol{Z}_{t}'\boldsymbol{v}_{t}\right) = \sum_{i=1}^{n} E\left[\boldsymbol{z}_{i,t}'(\boldsymbol{Z}_{t}'\boldsymbol{Z}_{t})^{-1}\sum_{j=1}^{n}\boldsymbol{z}_{j,t}E\left(v_{j,t}x_{i,t+s,k}|\boldsymbol{w}_{t}\right)\right]$$
(9)

And, by the law of iterated expectations, $E(v_{j,t}x_{i,t+s,k}|\boldsymbol{w}_t) = E[x_{i,t+s,k}E(v_{j,t}|\boldsymbol{w}_t,\boldsymbol{x}_{i,t+s})|\boldsymbol{w}_t]$. Given $(\boldsymbol{w}'_{j,t},v_{j,t})$ is independent of $\boldsymbol{w}_{i,T}$ for $j \neq i$, it follows that $E(v_{j,t}|\boldsymbol{w}_t,\boldsymbol{x}_{i,t+s}) = E(v_{j,t}|\boldsymbol{w}_{j,t})$ for $j \neq i$. Moreover, $E(v_{j,t}|\boldsymbol{w}_{j,t}) = 0$ by A1 and A3. Hence, $E(v_{j,t}x_{i,t+s,k}|\boldsymbol{w}_t) = 0$ for $j \neq i$. On the other hand, $E(v_{j,t}x_{i,t+s,k}|\boldsymbol{w}_t) = E(v_{i,t}x_{i,t+s,k}|\boldsymbol{w}_t)$ for j = i. Hence, $\sum_{j=1}^{n} \boldsymbol{z}_{j,t} E(v_{j,t}x_{i,t+s,k}|\boldsymbol{w}_t) = \boldsymbol{z}_{i,t} E(v_{i,t}x_{i,t+s,k}|\boldsymbol{w}_t)$, which implies the right-hand side of (9) is

$$\sum_{i=1}^{n} E\left[\boldsymbol{z}_{i,t}'(\boldsymbol{Z}_{t}'\boldsymbol{Z}_{t})^{-1}\boldsymbol{z}_{i,t}E\left(\boldsymbol{v}_{i,t}\boldsymbol{x}_{i,t+s,k}|\boldsymbol{w}_{t}\right)\right] = E\left[p_{n,t}E\left(\boldsymbol{v}_{1,t}\boldsymbol{x}_{1,t+s,k}|\boldsymbol{w}_{t}\right)\right], \quad (10)$$

where the equality in Eq. (10) follows from the fact that the \boldsymbol{u}_i s are identically distributed and $p_{n,t} = n \boldsymbol{z}'_{1,t} (\boldsymbol{Z}'_t \boldsymbol{Z}_t)^{-1} \boldsymbol{z}_{1,t}$. Moreover, A1 implies $E(v_{1,t}x_{1,t+s,k}|\boldsymbol{w}_t) = E(v_{1,t}x_{1,t+s,k}|\boldsymbol{w}_{1,t})$. Furthermore, from A3 and the definition of conditional covariance, we have $\gamma_{k,t,s} = E(v_{1,t}x_{1,t+s,k}|\boldsymbol{w}_{1,t})$. The preceding observations imply the kth entry of $E(\boldsymbol{X}'_{t+s}\boldsymbol{P}_t\boldsymbol{v}_t)$ is $E(p_{n,t}\gamma_{k,t,s})$. This conclusion and the definition of $\boldsymbol{s}_{1,T-t}$ implies the kth entry in $\boldsymbol{s}_{1,T-t}$ is $s_{1,T-t,k} = E(p_{n,t}\sum_{s=1}^{T-t}\gamma_{k,t,s})$ ($1 \leq t \leq T-1$). Moreover,

$$|s_{1,T-t,k}| \le E\left(p_{n,t}\left|\sum_{s=1}^{T-t} \gamma_{k,t,s}\right|\right) \le E\left(p_{n,t}\right)M = q_tM,\tag{11}$$

where the second inequality in (11) follows from A5 and the equality on the far right-hand side follows from Lemma A.1.

Next, the entries in $\mathbf{s}_{2,T-t}$ are evaluated. To that end, note that by arguments similar to those used to establish Eq. (10), the *k*th entry in $E\left(\mathbf{X}'_{t+s}\mathbf{P}_t\overline{\mathbf{v}}_t\right)$ is $E\left(p_{n,t}\overline{\mathbf{v}}_{1,t}x_{1,t+s,k}\right)$. Moreover, $E\left(p_{n,t}v_{1,t+r}x_{1,t+s,k}\right) = E\left[p_{n,t}x_{1,t+s,k}E\left(v_{1,t+r}|\mathbf{w}_{t+s}\right)\right] =$ 0 if $r \geq s$. Hence, for s = 1, we get $(T-t) E\left(p_{n,t}\overline{v}_{1,t}x_{1,t+s,k}\right) =$ $E\left(p_{n,t}x_{1,t+1,k}\sum_{r=1}^{T-t}v_{1,t+r}\right) = 0$. And, when t = T - 1, it must be that s = 1. Thus, for t = T - 1 and s = 1, we get $(T - t) E\left(p_{n,t}\overline{v}_{1,t}x_{1,t+s,k}\right) = E\left(p_{n,T-1}v_{1,T}x_{1,T,k}\right) = 0$. On the other hand, for $1 \le t \le T - 2$ and $s \ge 2$, we have

$$(T-t)E(p_{n,t}\overline{v}_{1,t}x_{1,t+s,k}) = E\left(p_{n,t}x_{1,t+s,k}\sum_{r=1}^{T-t}v_{1,t+r}\right)$$

= $E\left(p_{n,t}x_{1,t+s,k}\sum_{r=1}^{s-1}v_{1,t+r}\right)$
= $E\left[p_{n,t}\sum_{r=1}^{s-1}E(v_{1,t+r}x_{1,t+r+(s-r),k}|\boldsymbol{w}_{t+r})\right]$
= $E\left[p_{n,t}\sum_{r=1}^{s-1}\gamma_{k,t+r,s-r}\right].$ (12)

Moreover, for $T \geq 3$,

$$\sum_{s=2}^{T-t} \sum_{r=1}^{s-1} \gamma_{k,t+r,s-r} = \gamma_{k,t+1,1} \qquad (s=2) + \gamma_{k,t+1,2} + \gamma_{k,t+2,1} \qquad (s=3) + \cdots \qquad (13) + \gamma_{k,t+1,T-t-1} + \gamma_{k,t+2,T-t-2} + \cdots + \gamma_{k,T-1,1} \qquad (s=T-t) = \sum_{r=1}^{T-t-1} \sum_{j=1}^{T-t-1} \gamma_{k,t+r,j}.$$

It follows from Eq.s (12) and (13), $E(p_{n,T-1}v_{1,T}x_{1,T,k}) = 0$, and the definition of $\mathbf{s}_{2,T-t}$ that the *k*th entry in $\mathbf{s}_{2,T-t}$ is $s_{2,T-t,k} = (T-t)^{-1}E(p_{n,t}\sum_{r=1}^{T-t-1}\sum_{j=1}^{T-t-r}\gamma_{k,t+r,j})$ for $1 \le t \le T-2$, and $s_{2,T-t,k} = 0$ for t = T-1. Hence, $|s_{2,T-t,k}| = 0$ for t = T-1, and

$$|s_{2,T-t,k}| \le \frac{1}{T-t} E\left(p_{n,t} \sum_{r=1}^{T-t-1} \left| \sum_{j=1}^{T-t-r} \gamma_{k,t+r,j} \right| \right) \le E\left(p_{n,t} \right) \frac{1}{T-t} \sum_{r=1}^{T-t-1} M \le q_t M$$
(14)

for $1 \le t \le T - 2$.

Let $b_{n,T,k}$ denote the kth entry in $b_{n,T}$ (k = 1, ..., K). Expression (8), an obvious

inequality, and the inequalities in (11) and (14) imply

$$|E(b_{n,T,k})| \le \frac{1}{\sqrt{nT}} \sum_{t=1}^{T-1} \frac{1}{T-t+1} \left(|s_{1,T-t,k}| + |s_{2,T-t,k}| \right) \le \frac{2}{\sqrt{nT}} M \sum_{t=1}^{T-1} \frac{q_t}{T-t+1}.$$

Moreover, given $q_t \leq q_T^*$ for all t, we have $\sum_{t=1}^{T-1} q_t / (T-t+1) \leq q_T^* \sum_{t=1}^{T-1} 1 / (T-t+1)$. And, $\sum_{t=1}^{T-1} 1 / (T-t+1) = \sum_{t=1}^{T} 1 / t - 1$. Furthermore, $\sum_{t=1}^{T} 1 / t - 1 \leq \ln T$ for $T \geq 1$ (see, e.g., Havil, 2003, p. 47). The foregoing implies that, for $T \geq 3$,

$$|E(b_{n,T,k})| \le 2M \frac{q_T^*}{\sqrt{nT}} \ln T \qquad (k = 1, \dots, K).$$

$$(15)$$

Eq. (15) and $q_T^* = O(T^{\alpha})$ imply $E(b_{n,T,k}) = O\left((T^{\alpha} \ln T)/\sqrt{nT}\right)$ for $k = 1, \ldots, K$. The conclusion of the theorem follows from this observation and Assumption A4.

Lemma 1 proof

Assumption A5 will be verified for $y_{1,t-1}$. Verification of A5 for $y_{1,t-2}, \ldots, y_{1,t-K}$ is similar.

In order to verify A5 for $y_{1,t-1}$, first note that if the roots of the characteristic equation all lie outside the unit circle, then $y_{1,t}$ can be expressed as

$$y_{1,t} = \mu_1 + \sum_{j=0}^{\infty} \psi_j v_{1,t-j},$$

where $\mu_1 = \eta_1 / (1 - \beta_1 - \beta_2 - \dots - \beta_K)$ and $\sum_{j=0}^{\infty} |\psi_j| < \infty$ (see, e.g., Hamilton, 1994, pp. 58–59). Hence,

$$E(v_{1,t}y_{1,t+s-1}|\boldsymbol{w}_{1,t}) = E(v_{1,t}\mu_1|\boldsymbol{w}_{1,t}) + \sum_{j=0}^{\infty} \psi_j E(v_{1,t}v_{1,t+s-1-j}|\boldsymbol{w}_{1,t})$$

$$= \sum_{j=0}^{\infty} \psi_j E(v_{1,t}v_{1,t+s-1-j}|\boldsymbol{w}_{1,t}).$$
(16)

where the second equality follows from the fact that $E(v_{1,t}\mu_1|\boldsymbol{w}_{1,t}) = E[\mu_1 E(v_{1,t}|\eta_1, \boldsymbol{w}_{1,t})|\boldsymbol{w}_{1,t}] = E[\mu_1 E(v_{1,t})|\boldsymbol{w}_{1,t}] = E(\mu_1 0) = 0.$

In order to evaluate $E(v_{1,t}v_{1,t+s-1-j}|\boldsymbol{w}_{1,t})$, recall that $\boldsymbol{w}'_{1,t} = (y_{1,1-K}, \dots, y_{1,t-1})$

is past information. Moreover, if $s \geq j + 1$, then $v_{1,t+s-1-j}$ is a current or future error, and because current and future errors are independent of past information, we have $E(v_{1,t}v_{1,t+s-1-j}|\mathbf{w}_{1,t}) = E(v_{1,t}v_{1,t+s-1-j})$. Also, $E(v_{1,t}v_{1,t+s-1-j}) = \sigma^2$ or 0 according as j = s - 1 or $j \neq s - 1$. On the other hand, suppose s < j + 1, in which case $v_{1,t+s-1-j}$ is a past error. To evaluate $E(v_{1,t}v_{1,t+s-1-j}|\mathbf{w}_{1,t})$ for such cases, let $\mathcal{F}_{1,t-1}$ be the σ -field generated by $\mathbf{w}_{1,t} = (y_{1,1-K}, \ldots, y_{1,t-1})'$. Then $E(v_{1,t}v_{1,t+s-1-j}|\mathbf{w}_{1,t}) = E(v_{1,t}v_{1,t+s-1-j}|\mathcal{F}_{1,t-1})$. Next, let \mathcal{F}_{t-1} denote the σ -field generated by η_1 and the past $y_{1,t}$ s, i.e., $\mathcal{F}_{t-1} := \sigma(\ldots, y_{1,t-2}, y_{1,t-1}, \eta_1)$. By the law of iterated expectations, $E(v_{1,t}v_{1,t+s-1-j}|\mathcal{F}_{1,t-1}) = E[E(v_{1,t}v_{1,t+s-1-j}|\mathcal{F}_{t-1})|\mathcal{F}_{1,t-1}]$. Moreover, given we are now considering the cases for which s - 1 - j = $-1, -2, \ldots$, we get $E(v_{1,t}v_{1,t+s-1-j}|\mathcal{F}_{t-1}) = (\beta(L)y_{1,t+s-1-j} - \eta_1)E(v_{1,t}|\mathcal{F}_{t-1}) =$ $(\beta(L)y_{1,t+s-1-j} - \eta_1)E(v_{1,t}) = 0$. Hence, $E(v_{1,t}v_{1,t+s-1-j}|\mathbf{w}_{1,t}) = 0$ for s < j + 1. The preceding shows $E(v_{1,t}v_{1,t+s-1-j}|\mathbf{w}_{1,t}) = \sigma^2$ or 0 according as j = s - 1 or $j \neq s - 1$.

The conclusion of the last paragraph and (16) imply $E(v_{1,t}y_{1,t+s-1}|\boldsymbol{w}_{1,t}) = \sigma^2\psi_{s-1}$. Moreover, because $E(v_{1,t}|\boldsymbol{w}_{1,t}) = 0$, we have $\operatorname{cov}(v_{1,t}, y_{1,t+s-1}|\boldsymbol{w}_{1,t}) = E(v_{1,t}y_{1,t+s-1}|\boldsymbol{w}_{1,t})$. Hence, $\operatorname{cov}(v_{1,t}, y_{1,t+s-1}|\boldsymbol{w}_{1,t}) = \sigma^2\psi_{s-1}$ for $s \geq 1$. And $\sum_{j=0}^{\infty} |\psi_j| < \infty$ by assumption. Therefore, Assumption A5 is satisfied for the first regressor $y_{1,t-1}$. Similar arguments verify A5 for $y_{1,t-2}, \ldots, y_{1,t-K}$.

Theorem 2 proof

Note that $\mathbf{\Pi}_{t} = \mathbf{Q}_{t}^{-1}\mathbf{C}_{t}$ are the linear projection parameters in the K reduced form equations $\ddot{\mathbf{x}}_{i,t} = \mathbf{\Pi}_{t}'\mathbf{z}_{i,t} + \mathbf{r}_{i,t}$, where $\mathbf{r}_{i,t}' = (r_{i,t,1}, \ldots, r_{i,t,K})$ are reduced form errors. Hence, $\ddot{\mathbf{X}}_{i,t} = \mathbf{Z}_{t}\mathbf{\Pi}_{t} + \mathbf{R}_{t}$, where $\mathbf{R}_{t}' = (\mathbf{r}_{1,t}, \ldots, \mathbf{r}_{n,t})$. Therefore, $\mathbf{b}_{n,T} = (1/\sqrt{nT})\sum_{t=1}^{T-1}\mathbf{\Pi}_{t}'\mathbf{Z}_{t}'\ddot{\mathbf{v}}_{t} + (1/\sqrt{nT})\sum_{t=1}^{T-1}\mathbf{R}_{t}'\mathbf{P}_{t}\ddot{\mathbf{v}}_{t}$. Moreover, $(1/\sqrt{nT})\sum_{t=1}^{T-1}\mathbf{R}_{t}'\mathbf{P}_{t}\ddot{\mathbf{v}}_{t} = (1/\sqrt{T})\sum_{t=1}^{T-1}(1/n)\mathbf{R}_{t}'\mathbf{Z}_{t}[(1/n)\mathbf{Z}_{t}'\mathbf{Z}_{t}]^{-1}(1/\sqrt{n})\mathbf{Z}_{t}'\ddot{\mathbf{v}}_{t}$. Because $E(\mathbf{r}_{i,t}\mathbf{z}_{i,t}') = E(\ddot{\mathbf{x}}_{i,t}\mathbf{z}_{i,t}') - \mathbf{\Pi}_{t}'E(\mathbf{z}_{i,t}\mathbf{z}_{i,t}') = \mathbf{C}_{t}' - \mathbf{C}_{t}'\mathbf{Q}_{t}^{-1}\mathbf{Q}_{t} = \mathbf{0}$, we have that $(1/n)\mathbf{R}_{t}'\mathbf{Z}_{t} = (1/n)\sum_{i=1}^{n}\mathbf{r}_{i,t}\mathbf{z}_{i,t}' \xrightarrow{p} \mathbf{0}$, as $n \to \infty$. Moreover, Assumption A3 implies $E(\mathbf{z}_{1,t}\ddot{\mathbf{v}}_{1,t}) = \mathbf{0}$, and Assumption A6 implies that $\operatorname{Var}(\mathbf{z}_{1,t}\ddot{\mathbf{v}}_{1,t})$ has finite entries. Therefore, by the central limit theorem for i.i.d. random vectors, we have $(1/\sqrt{n})\mathbf{Z}_{t}'\ddot{\mathbf{v}}_{t} \xrightarrow{d} N(\mathbf{0}, \operatorname{Var}(\mathbf{z}_{1,t}\ddot{\mathbf{v}}_{1,t}))$ as $n \to \infty$. Hence, $(1/\sqrt{n})\mathbf{Z}_{t}'\ddot{\mathbf{v}}_{t} = O_{p}(1)$ for each t. It follows that $\mathbf{b}_{n,T} - (1/\sqrt{nT})\sum_{t=1}^{T-1}\mathbf{\Pi}_{t}'\mathbf{Z}_{t}'\ddot{\mathbf{v}}_{t} \xrightarrow{p} \mathbf{0}$ as $n \to \infty$. Therefore, the asympton totic distribution of $\boldsymbol{b}_{n,T}$, as $n \to \infty$, is the same as the asymptotic distribution of $(1/\sqrt{nT}) \sum_{t=1}^{T-1} \Pi'_t \boldsymbol{Z}'_t \boldsymbol{\ddot{v}}_t$, as $n \to \infty$.

To evaluate the latter, set $\boldsymbol{\epsilon}_{i,T} := (1/\sqrt{T}) \sum_{t=1}^{T-1} \mathbf{\Pi}'_t \boldsymbol{z}_{i,t} \ddot{\boldsymbol{v}}_{i,t}$. Note that $E(\boldsymbol{\epsilon}_{1,T}) = \mathbf{0}$ and $\operatorname{Var}(\boldsymbol{\epsilon}_{1,T}) = \boldsymbol{\Omega}_T$. By the central limit theorem for i.i.d. random vectors, we have that $(1/\sqrt{n}) \sum_{i=1}^n \boldsymbol{\epsilon}_{i,T} \stackrel{d}{\to} N(\mathbf{0}, \boldsymbol{\Omega}_T)$ $(n \to \infty)$. But $(1/\sqrt{n}) \sum_{i=1}^n \boldsymbol{\epsilon}_{i,T} = (1/\sqrt{nT}) \sum_{t=1}^{T-1} \mathbf{\Pi}'_t \boldsymbol{Z}'_t \ddot{\boldsymbol{v}}_t$, and recall from the last paragraph that $\boldsymbol{b}_{n,T} - (1/\sqrt{nT}) \sum_{t=1}^{T-1} \mathbf{\Pi}'_t \boldsymbol{Z}'_t \ddot{\boldsymbol{v}}_t$ $(n \to \infty)$. Hence, $\boldsymbol{b}_{n,T} \stackrel{d}{\to} \boldsymbol{b}_T \sim N(\mathbf{0}, \boldsymbol{\Omega}_T)$ $(n \to \infty)$.

The characteristic function of \boldsymbol{b}_T is $\phi_T(\boldsymbol{\lambda}) := \exp(-(1/2)\boldsymbol{\lambda}'\boldsymbol{\Omega}_T\boldsymbol{\lambda})$. Moreover, Assumption A7 implies $\lim_{T\to\infty} \phi_T(\boldsymbol{\lambda}) = \exp(-(1/2)\boldsymbol{\lambda}'\boldsymbol{\Omega}\boldsymbol{\lambda})$. The latter limit is the characteristic function of a multivariate normal vector with mean **0** and variancecovariance matrix $\boldsymbol{\Omega}$. Hence, $\boldsymbol{b}_T \stackrel{d}{\to} \boldsymbol{b} \sim N(\mathbf{0}, \boldsymbol{\Omega}) \ (T \to \infty)$.

The preceding verifies $\boldsymbol{b}_{n,T} \stackrel{d}{\rightarrow} \boldsymbol{b} \sim N(\boldsymbol{0}, \boldsymbol{\Omega}) \ (n, T \rightarrow \infty)_{\text{seq}}$. We have that $\boldsymbol{A}_{n,T} \stackrel{p}{\rightarrow} \boldsymbol{A}_T := (1/T) \sum_{t=1}^{T-1} \boldsymbol{C}'_t \boldsymbol{Q}_t^{-1} \boldsymbol{C}_t \ (n \rightarrow \infty)$ by the law of large numbers, and $\boldsymbol{A}_T \rightarrow \boldsymbol{A} > 0 \ (T \rightarrow \infty)$ by Assumption A8. Hence, $\sqrt{nT}(\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}) = \boldsymbol{A}_{n,T}^{-1} \boldsymbol{b}_{n,T} \stackrel{d}{\rightarrow} N(\boldsymbol{0}, \boldsymbol{A}^{-1} \boldsymbol{\Omega} \boldsymbol{A}^{-1}) \ (n, T \rightarrow \infty)_{\text{seq}}$.

References

- Alvarez, J., & Arellano, M. (2003). The time series and cross-section asymptotics of dynamic panel data estimators. *Econometrica* 71, 1121–1159.
- Anderson, T. W., Kunitomo, N., & Matsushita, Y. (2011). On finite sample properties of alternative estimators of coefficients in a structural equation with many instruments. *Journal of Econometrics* 165, 58–69.
- Arellano, M. (2003). Panel Data Econometrics. Oxford University Press, Oxford.
- Arellano, M., & Bond, S. (1991). Some tests of specification for panel data: Monte Carlo evidence and an application to employment equations. The Review of Economic Studies 58, 277–297.
- Bekker, P. A. (1994). Alternative approximations to the distributions of instrumental variables estimators. *Econometrica* 62, 657-681.
- Blundell, R., & Bond, S. (1998). Initial conditions and moment restrictions in dynamic panel data models. *Journal of Econometrics* 87, 115–143.

- Bun, M. J. G., & Kiviet, J. F. (2006). The effects of dynamic feedbacks on LS and MM estimator accuracy in panel data models. *Journal of Econometrics 132*, 409–444.
- Chao, J. C., & Swanson, N. R. (2005). Consistent estimation with a large number of weak instruments. *Econometrica* 73, 1673–1692.
- Hamilton, J. D. (1994). Time Series Analysis. Princeton University Press, Princeton, New Jersey.
- Hansen, C., Hausman, J., & Newey, W. (2008). Estimation with many instrumental variables. *Journal of Business & Economic Statistics*, 26, 398–422.
- Havil, J. (2003). *Gamma: Exploring Euler's Constant*. Princeton University Press, Princeton, New Jersey.
- Hayakawa, K., Qi, M., & Breitung, J. (2019). Double filter instrumental variable estimation of panel data models with weakly exogenous variables. *Econometric Reviews*, 38, 1055–1088.
- Hsiao, C. & Zhou, Q. (2017). First difference or forward demeaning: Implications for the method of moments estimators. *Econometric Reviews* 36, 883–897.
- Hsiao, C. & Zhang, J. (2015). IV, GMM or likelihood approach to estimate dynamic panel models when either N or T or both are large. *Journal of Econometrics 187*, 312–322.
- Kapetanios, G. (2008). A bootstrap procedure for panel data sets with many crosssectional units. *Econometrics Journal* 11, 377–395.
- Koenker, R. & Machado, J. A. F. (1999). GMM inference when the number of moment conditions is large. *Journal of Econometrics* 93, 327–344.
- Okui, R. (2009). The optimal choice of instruments in dynamic panel data models. Journal of Econometrics 151, 1–16.
- Phillips, P. C. B. & Moon, H. R. (1999). Linear regression limit theory for nonstationary panel data. *Econometrica* 67, 1057–1111.

- Phillips, P. C. B. & Moon, H. R. (2000). Nonstationary panel data analysis: an overview of some recent developments. *Econometric Reviews 19*, 263–286.
- Phillips, R. F. (2019). A numerical equivalence result for generalized method of moments. *Economics Letters 179*, 13–15.
- Phillips, R. F. (2020). Quantifying the advantages of forward orthogonal deviations for long time series. *Computational Economics* 55, 653-672.

Phillips, R. F. (2024). Supplemental Appendix.