



Weighted and Branching Bisimilarities from Generalized Open Maps

Jérémy Dubut¹ and Thorsten Wisßmann²

¹ National Institute of Advanced Industrial Science and Technology, Tokyo, Japan
jeremy.dubut@aist.go.jp

² Radboud University, Nijmegen, the Netherlands thorsten.wissmann@ru.nl

Abstract. In the open map approach to bisimilarity, the paths and their runs in a given state-based system are the first-class citizens, and bisimilarity becomes a derived notion. While open maps were successfully used to model bisimilarity in non-deterministic systems, the approach fails to describe quantitative system equivalences such as probabilistic bisimilarity. In the present work, we see that this is indeed impossible and we thus generalize the notion of open maps to also accommodate weighted and probabilistic bisimilarity. Also, extending the notions of strong path and path bisimulations into this new framework, we show that branching bisimilarity can be captured by this extended theory and that it can be viewed as the history preserving restriction of weak bisimilarity.

Keywords: Open maps · Weighted Bisimilarity · Probabilistic Bisimilarity · Branching Bisimilarity · Weak Bisimilarity

1 Introduction

The theory of open maps is a categorical framework to reason about systems and their bisimilarities [16]. Given a category of systems and a description of the shape of the executions and how to extend them, open maps are morphisms with lifting properties with respect to those extensions. Intuitively, open maps are morphisms which preserve and reflect transitions of systems, that is, they are morphisms whose graphs are bisimulations. The theory covers various classical notions of bisimilarity. For example, two LTSs are strongly bisimilar if and only if there is a span of open maps between them. Varying the category of models and the execution shapes allows describing weak bisimilarity, timed bisimilarity, probabilistic Larsen and Skou bisimilarity, and history-preserving bisimilarity of event structures (see [16,3,12] for examples).

Another categorical framework for bisimilarity is coalgebra [22]. This time, given a category and an endofunctor describing respectively the type of state spaces and the type of transitions, a ‘system’ is understood as a coalgebra for this functor. Coalgebra homomorphisms are then very similar to open maps in spirit: they also are morphisms that preserve and reflect transitions. This intuition has been made formal by transformations between the categorical frameworks in both ways; from open maps to coalgebra [19], and conversely [25]. However, the

latter suggests that open maps are only adapted to modeling non-deterministic systems and would struggle with other types of branchings, such as probabilistic.

In coalgebra, there are no particular difficulties in modeling weighted systems, and by extension, discrete probabilistic systems [17]. There is also some work for continuous probabilities, although the theory is much more complicated [5,4]. As we will explain more precisely later, there have been some attempts to do so with open maps in [3,5], but the result is somewhat disappointing.

Conversely, coalgebra is not adapted to bisimilarities for systems where transitions are not history-preserving, that is, for which the behavioral equivalence does not just depend on the transitions at a given state, but on the whole history of the execution that led to this state. That is the case for example for branching bisimilarity [23]. Branching bisimilarity arose precisely to make weak bisimilarity history-preserving. In [3], weak bisimilarity has been described using open maps by carefully choosing the underlying category, with a general theory developed in [9] using presheaf models. Branching bisimilarity has also been studied using open maps in [1,2], but indirectly, through a translation into presheaves.

To resume, the goal of this paper is to capture weighted and branching bisimilarities using a generalization of open maps. Concretely, the contributions are:

1. a proof that it is impossible to appropriately model probabilistic system using standard open maps (Section 3.2),
2. a faithful extension of the theory of open maps and (strong) path bisimulations (Section 4),
3. a generalized open map situation capturing weighted and probabilistic bisimilarities (Section 5),
4. a generalized open map situation where strong path bisimulations correspond to stuttering branching bisimulations, open map bisimilarity to branching bisimilarity, and path bisimulations to weak bisimulations (Section 6).

In addition, Section 2 introduces some theoretical requirements.

2 From path categories to bisimilarity

Before discussing weighted bisimilarity, let us first recall the main ideas of modeling bisimilarity via open maps, as introduced by Joyal et al. [16]. The definition is parametric in a functor $J: \mathbb{P} \rightarrow \mathbb{M}$, from a category \mathbb{P} of paths to a category \mathbb{M} of models or systems of interest. In the prime example, \mathbb{M} is the category of labelled transition systems LTS as defined next:

Definition 2.1. *For a fixed set A of labels, the category LTS contains:*

1. *Objects: a labelled transition system (X, \rightarrow, x_0) is a set X of states, a transition relation $\rightarrow \subseteq X \times A \times X$ and a distinguished initial state $x_0 \in X$. We write $x \xrightarrow{a} x'$ to denote that $(x, a, x') \in \rightarrow$ and simply refer to the LTS as X if \rightarrow and x_0 are clear from the context. For disambiguation, we use \rightarrow for morphisms and \rightarrow for transitions.*

2. *Morphisms:* a functional simulation $f: (X, \rightarrow, x_0) \rightarrow (Y, \rightarrow, y_0)$ is a function $f: X \rightarrow Y$ with $f(x_0) = y_0$ and for all $x \xrightarrow{a} x'$ in X , we have $f(x) \xrightarrow{a} f(x')$.

A functional simulation $f: X \rightarrow Y$ intuitively means that the system Y has at least the transitions of X , but possibly more. A special case of a functional simulation is the *run* of a word in a system:

Definition 2.2. For the label set A , let (A^*, \leq) be the partially ordered set of words, ordered by the prefix ordering. The functor $J: (A^*, \leq) \rightarrow \text{LTS}$ sends a word $w \in A^*$ to the LTS $Jw = (\{v \mid v \leq w\}, \rightarrow, \varepsilon)$ of all prefixes of w with $v \xrightarrow{a} va$ for all $a \in A$, $va \leq w$.

This functor J (or more precisely, its image) is often called *path category* of LTS: the possible runs of a word $w \in A^*$ in (X, \rightarrow, x_0) correspond precisely to the functional simulations $Jw \rightarrow (X, \rightarrow, x_0)$ in LTS.

On the abstract level, for a general functor $J: \mathbb{P} \rightarrow \mathbb{M}$, we understand the set of morphisms $r: Jw \rightarrow X$ for $w \in \mathbb{P}$ and $X \in \mathbb{M}$ as the runs of the path w in the model X . We can already make the trivial observation that all morphisms $f: X \rightarrow Y$ in \mathbb{M} preserve runs: given a run $r: Jw \rightarrow X$ of some path $w \in \mathbb{P}$ in X , there is a run $f \cdot r: Jw \rightarrow Y$ of w in Y .

The converse does not hold for a general $f: X \rightarrow Y$ in \mathbb{M} : given a run of w in Y , there is not necessarily a run of w in X . If f reflects runs, it is called *open*:

Definition 2.3. For a functor $J: \mathbb{P} \rightarrow \mathbb{M}$, a morphism $f: X \rightarrow Y$ in \mathbb{M} is called *open* if f satisfies the following lifting property for all $e: v \rightarrow w$ in \mathbb{P} :

$$\text{for all } \begin{array}{ccc} Jv & \xrightarrow{r} & X \\ \downarrow J_e & \circlearrowleft & \downarrow f \\ Jw & \xrightarrow{s} & Y \end{array} \quad \text{there is } d: Jw \rightarrow X \text{ with } \begin{array}{ccc} Jv & \xrightarrow{r} & X \\ \downarrow J_e & \circlearrowleft & \downarrow f \\ Jw & \xrightarrow{s} & Y \end{array}$$

That is, for all commutative squares $(s \cdot J_e = f \cdot r)$, there is $d: Jw \rightarrow X$ in \mathbb{M} that makes both triangles on the right commute ($f \cdot d = s$ and $d \cdot J_e = r$).

By construction, we can only make statements about states that are reachable via some run. Thus, one often restricts \mathbb{M} beforehand to contain only models in which all states are reachable from the initial state.

For LTSs in which all states are reachable from the initial state, open maps are related to strong bisimulations [20]: open maps are precisely functions whose graph relation $\{(x, fx) \mid x \in X\}$ is a strong bisimulation. Reformulated in the context of allegories [10], open maps are precisely the maps in the allegory of relations that are strong bisimulations. It is then natural to recover bisimulations as tabulations of open maps, that is:

Definition 2.4. For a functor $J: \mathbb{P} \rightarrow \mathbb{M}$, we say that two models X and Y are *J-bisimilar*, if there exist another model Z and two *J-open* maps $f: Z \rightarrow X$ and $g: Z \rightarrow Y$, that is, if there is a span of *J-open* maps between them.

Of course, J -bisimilarity is a reflexive (identities are open maps) and symmetric (by permuting f and g in the definition) relation on models, but it is not transitive in general. It is when the category \mathbb{M} has pullbacks [16].

Given a functor $J: \mathbb{P} \rightarrow \mathbb{M}$, there are more classical ways of defining bisimilarities given in [16]. The first one is (*strong*) *path bisimulations*, which are relations on runs (similar to history-preserving bisimulations) satisfying the usual bisimilarity conditions. The second one is by using a modal logic similar to the Hennessy-Milner theorem. In the case of LTSs with strong bisimilarity, all those notions describe the same notion of bisimilarity, but that is not true for general $J: \mathbb{P} \rightarrow \mathbb{M}$: it can only be proved that J -bisimilarity implies the existence of a (strong) path bisimulation, which itself implies that the two models satisfy the same formulas of the modal logic. In [6], some mild sufficient conditions in terms of trees (i.e., colimits of paths in \mathbb{M}) are given for those three notions to coincide. In particular, all the examples of bisimilarities covered by open maps cited earlier satisfy these conditions.

We use coalgebra for uniform statements about state-based systems of different branching type (including non-deterministic and probabilistic branching):

Definition 2.5. *For an object 1 of a category \mathcal{C} and an endofunctor $F: \mathcal{C} \rightarrow \mathcal{C}$, a pointed coalgebra is a pair of morphisms of \mathcal{C} of the form $1 \xrightarrow{i} X \xrightarrow{\xi} FX$.*

For example, LTSs can be modeled as pointed coalgebras with $\mathcal{C} = \text{Set}$, 1 any singleton, and $F = \mathcal{P}(A \times _)$, where \mathcal{P} is the power set functor. The usual notion of morphisms of coalgebras can be spelt out as follows:

Definition 2.6. *A (proper) homomorphism of pointed coalgebras from (X, ξ, i) to (Y, ζ, j) is a morphism $f: X \rightarrow Y$ of \mathcal{C} such that the diagram on the right commutes.*

$$\begin{array}{ccccc} 1 & \xrightarrow{i} & X & \xrightarrow{\xi} & FX \\ & \searrow & \circlearrowleft & \downarrow f & \circlearrowleft & \downarrow Ff \\ & & j & \searrow & Y & \xrightarrow{\zeta} & FY \end{array}$$

Pointed coalgebras and proper homomorphisms always form a category, but in the case of LTSs as described above, this category is not equivalent to the category LTS. Indeed, proper homomorphisms are not just morphisms that preserve transitions, but similarly to open maps, they also reflect them. In [25], the authors proved that for a large class of endofunctors, whose coalgebras basically are non-deterministic, proper homomorphisms precisely correspond to J -open maps for a certain functor J . To model morphisms that are only required to preserve transitions, homomorphisms have to be made lax as follows (see [25]):

Definition 2.7. *Assume a relation \sqsubseteq on every Hom-set $\mathcal{C}(X, FY)$. A lax homomorphism of pointed coalgebras from (X, ξ, i) to (Y, ζ, j) is a morphism $f: X \rightarrow Y$ of \mathcal{C} such that the diagram on the right laxly commutes, that is, $f \cdot i = j$ and $Ff \cdot \xi \sqsubseteq \zeta \cdot f$ in $\mathcal{C}(X, FY)$.*

$$\begin{array}{ccccc} 1 & \xrightarrow{i} & X & \xrightarrow{\xi} & FX \\ & \searrow & \circlearrowleft & \downarrow f & \sqcap & \downarrow Ff \\ & & j & \searrow & Y & \xrightarrow{\zeta} & FY \end{array}$$

In the case of the functor $\mathcal{P}(A \times _)$, we can consider the pointwise inclusion on every Hom-set $\text{Set}(X, \mathcal{P}(A \times Y))$. With this, pointed coalgebras and lax

homomorphisms form a category which is isomorphic to the category LTS. However, it is not true in general that they form a category, as a compatibility of \sqsubseteq with the composition is needed as follows:

Definition 2.8. *A partial order on F is a collection of partial orders \sqsubseteq , one for each Hom-set of the form $\mathcal{C}(X, FY)$ such that*

$$\forall X \xrightarrow{f_1, f_2} FY, X' \xrightarrow{g} X, Y \xrightarrow{h} Y': f_1 \sqsubseteq f_2 \Rightarrow Fh \cdot f_1 \cdot g \sqsubseteq Fh \cdot f_2 \cdot g.$$

This is equivalent to the requirement that the Hom-functor $\mathcal{C}(_, F_)$ factors through partially ordered sets: $\mathcal{C}(_, F_) : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \text{Pos}$.

Remark 2.9. The present definition subsumes the definition of order on a Set-functor established by Hughes and Jacobs [11, Def 2.1] (details in the appendix).

Lemma 2.10 [25]. *When \sqsubseteq is a partial order on F , pointed coalgebras and lax homomorphisms form a category, which we denote by $\text{LCoalg}(1, F)$.*

Much as with open maps, many flavors of bisimilarity can be recovered using spans of proper homomorphisms:

Definition 2.11. *We say that two pointed coalgebras are coalgebraically bisimilar if there is a span of proper homomorphisms between them.*

There are many ways of defining bisimilarities in coalgebra (see [13] for an overview), but they coincide for the purpose of the present paper.

3 Weighted Bisimilarity and Open Maps

In this section, we describe known attempts to model weighted systems, and particularly probabilistic ones, using open maps. They all work with some variations of the (discrete) distribution functor on Set . We will denote this functor, which maps a set X to the set

$$\mathcal{D}X = \left\{ f : X \rightarrow [0, 1] \mid f^{-1}((0, 1]) \text{ is finite and } \sum_{x \in X} f(x) = 1 \right\},$$

by \mathcal{D} and the variation where the condition $= 1$ is replaced by ≤ 1 by $\mathcal{D}_{\leq 1}$ (i.e. $\mathcal{D}_{\leq 1}X := \mathcal{D}(X + 1)$). We will prove that, even though Larsen-Skou bisimulations for reactive systems can be modeled with open maps, that is impossible for bisimulations for generative systems.

3.1 Larsen-Skou bisimilarity for reactive systems using open maps

In [3], Cheng et al. describe an open map situation for Probabilistic Transition Systems (PTSs), which corresponds to coalgebras for the functor $(\mathcal{D}(_) + 1)^A$. In this setting, they consider Partial PTSs (PPTS) which are coalgebras for $(\mathcal{D}_{\leq 1}^\varepsilon(_) + 1)^A$ where the sub-probability distributions can have values in hyper-reals, allowing infinitesimals ε . The category of PTSs embeds in that of PPTSs,

and the path category is the full subcategory of PPTSs consisting of finite linear systems whose probabilities of transitions are infinitesimals. It is then proved that J -bisimilarity, restricted to PTSs, for this path category corresponds to Larsen-Skou's probabilistic bisimilarity [18].

This open map situation has been reformulated in [7] in terms of coreflections: the obvious functor from PPTSs to TSs is a coreflection whose left-adjoint maps a LTS T to the PPTS whose underlying LTS is T and where all transitions have infinitesimal probabilities. In general, given a coreflection $F : \mathcal{C} \rightarrow \mathcal{D}$ with left-adjoint G and a path category J on \mathcal{D} , one automatically has the path category $G \circ J$ on \mathcal{C} , and this construction preserves good properties of J . In particular, one has that two systems A and B are $(G \circ J)$ -bisimilar if and only if FA and FB are J -bisimilar. Cheng et al.'s path category is obtained in this manner with the coreflection above and the standard path category on LTSs. In particular, it means that two PPTSs are bisimilar if and only if their underlying TSs are strongly bisimilar.

3.2 Impossibility result for generative systems

In [5], Desharnais et al. describe several bisimilarities for generative probabilistic systems, that is, coalgebras for the functor $\mathcal{D}_{\leq 1}(A \times _)$, in a coalgebraic way. They pointed out that their efforts to model those bisimilarities using open maps failed [5, p. 188]. In the following, we see that it is in fact not possible. We will show that for generative probabilistic systems modeled by the category $M := \text{LCoalg}(1, \mathcal{D}_{\leq 1}(A \times _))$, there is no open map characterization of the coalgebraic bisimilarity. Actually, the argument here is valid for many other types of weights and is not limited to reals.

Here, for two functions $f, g : X \rightarrow \mathcal{D}_{\leq 1}(Y)$, $f \sqsubseteq g$ means that for all $x \in X$, for all $y \in Y$, $f(x)(y) \leq g(x)(y)$, where \leq is the usual ordering on $[0, 1]$.

In this situation:

Theorem 3.1. *For $M := \text{LCoalg}(1, \mathcal{D}_{\leq 1}(A \times _))$ there is no category \mathbb{P} and no functor $J : \mathbb{P} \rightarrow M$ such that for every $h : X \rightarrow Y$ with reachable X the following equivalence holds:*

$$h \text{ is } J\text{-open} \iff h \text{ is a proper homomorphism}$$

and there is no \mathbb{P} and no functor J such that for every X and Y :

$$X \text{ and } Y \text{ are } J\text{-bisimilar} \iff X \text{ and } Y \text{ are coalgebraically bisimilar.}$$

Proof (Sketch). By contradiction, assume that there is such a J . We prove that there is a proper homomorphism of the form:

$$X = \boxed{\begin{array}{c} \rightarrow r \begin{array}{l} \nearrow \frac{1}{n}, a \ x_1 \\ \rightarrow \vdots \\ \searrow \frac{1}{n}, a \ x_n \end{array} \end{array}} \xrightarrow{h} Y = \boxed{\rightarrow r \xrightarrow{1, a} y}$$

which cannot be J -open. Consider first the unique lax homomorphism $0_{\mathbb{M}} \rightarrow Y$ where $0_{\mathbb{M}}$ consists in one state and no transition. This is not a proper homomorphism, so it is not open by assumption. That is there is a square:

$$\begin{array}{ccc} JP & \xrightarrow{p} & 0_{\mathbb{M}} \\ \downarrow J\phi & & \downarrow !_Y \\ JQ & \xrightarrow{q} & Y \end{array}$$

with no lifting. It is mechanical to check that $JP \simeq 0_{\mathbb{M}}$ and JQ has at least one transition from its initial state to another state $r \xrightarrow{w, a} z$ with $w \neq 0$. With $n = 2 \cdot \lceil \frac{1}{w} \rceil$, the proper homomorphism h above is not open: there cannot be a morphism from JQ to X because $w > \frac{1}{n}$. \square

4 Generalized Open Maps

The main argument of the proof of impossibility is the fact that sometimes, a transition with some probability w in the codomain comes from probabilities w_1, \dots, w_n with $\sum_i w_i = w$ in the domain, which makes a lifting morphism impossible with the current framework of open maps.

In this section, we will extend the open map framework with the main intuition that the lifting morphism *splits* the probability w into smaller parts w_1, \dots, w_n . After defining these generalized open maps, we show some basic properties of the bisimilarity generated by them.

4.1 Generalized Open Maps Situation

Here, we describe our extension of the open maps framework. The data is similar: we start with a category of models \mathbb{M} , but we need more than just a functor $J: \mathbb{P} \rightarrow \mathbb{M}$. Assume:

- a set V together with a function $J: V \rightarrow \text{ob}(\mathbb{M})$,
- two small categories \mathbb{E} and \mathbb{S} whose sets of objects are V ,
- two functors $J_{\mathbb{E}}: \mathbb{E} \rightarrow \mathbb{M}$ and $J_{\mathbb{S}}: \mathbb{S} \rightarrow \mathbb{M}$ coinciding with J on objects.

The classical open maps situation $J: \mathbb{P} \rightarrow \mathbb{M}$ fits in this extension as follows. The category \mathbb{E} is given by \mathbb{P} with the intention that they model path shapes and their *extensions*. The functor $J_{\mathbb{E}}$ is given by J . The category \mathbb{S} is given by the discrete category $|\mathbb{P}|$, that is, the category whose objects are those of \mathbb{P} and whose morphisms are only identities. The functor $J_{\mathbb{S}}$ is the only possible one respecting the conditions of the definition above.

In the general context of this extension, the interpretation is a bit different. Now V is meant to be a set of trees labelled by alphabets and weights. \mathbb{E} still consists in extensions, extending trees into trees with longer branches. \mathbb{S} then consists in *merging morphisms*, similar to the description above: for the example of weighted systems, those morphisms are allowed to merge states into one,

as long as they sum up the weights of the in-going branches. Generally, those morphisms are allowed to perform some merges that are harmless for bisimilarity.

With this data, we can define generalized open maps:

Definition 4.1. *A morphism $f: X \rightarrow Y$ in \mathbb{M} is called (\mathbb{E}, \mathbb{S}) -open if it satisfies the following lifting property for all $e: v \rightarrow w$ in \mathbb{E} :*

$$\begin{array}{ccc}
 \text{for all} & \begin{array}{ccc} Jv & \xrightarrow{x} & X \\ J_{\mathbb{E}}e \downarrow & \circlearrowleft & \downarrow f \\ Jw & \xrightarrow{y} & Y \end{array} & \text{there is} & \begin{array}{ccc} Jv & \xrightarrow{x} & X \\ \downarrow & \swarrow J_{\mathbb{E}}e' \circlearrowleft & \searrow x' \dashrightarrow \\ J_{\mathbb{E}}e & \circlearrowleft & Ju \\ \downarrow & \swarrow J_{\mathbb{S}}s \circlearrowleft & \searrow \\ Jw & \xrightarrow{y} & Y \end{array}
 \end{array}$$

The interpretation starts the same as in usual open maps. Assume that we have a tree y in Y extending the image by f of the tree x in X . If f is open, there should be a tree x' extending x and whose image by f is y . However, x' may have a different shape than y , since it might be necessary to split transitions. That is what u and s are modeling: w is obtained from u by merging some states.

The connection with the classical open maps can be formulated as follows

Proposition 4.2. *Given a functor $J: \mathbb{P} \rightarrow \mathbb{M}$ and a morphism $f: X \rightarrow Y$,*

$$f \text{ is } J\text{-open if and only if } f \text{ is } (\mathbb{P}, |\mathbb{P}|)\text{-open.}$$

Again, bisimilarity can be defined as the existence of a span of open maps

Definition 4.3. *We say that X and Y are (\mathbb{E}, \mathbb{S}) -bisimilar if there is a span of (\mathbb{E}, \mathbb{S}) -open maps between them.*

4.2 Basic Properties

In this section, we will prove general properties of (\mathbb{E}, \mathbb{S}) -bisimilarity similar to the classical case. First, we show that if \mathbb{M} has pullbacks, then (\mathbb{E}, \mathbb{S}) -bisimilarity is an equivalence relation. Secondly, we describe two notions of path bisimulations, both implied by (\mathbb{E}, \mathbb{S}) -bisimilarity. Finally, we prove that it is enough to check openness on some generators of \mathbb{E} .

In order to see when (\mathbb{E}, \mathbb{S}) -bisimilarity is an equivalence relation, we need to check symmetry, reflexivity, and transitivity. *Symmetry* always holds because we can always swap the legs of the span. For *reflexivity*, it is enough to prove that identities are open which is valid because \mathbb{S} is a category and $J_{\mathbb{S}}$ is a functor, as shown in the diagram on the right. The proof of *transitivity* relies on composition and pullbacks:

$$\begin{array}{ccc}
 Jv & \xrightarrow{x} & X \\ \downarrow & \swarrow J_{\mathbb{E}}e \circlearrowleft & \searrow y \dashrightarrow \\ J_{\mathbb{E}}e & \circlearrowleft & Jw \\ \downarrow & \swarrow J_{\mathbb{S}} \text{id} = \text{id} \circlearrowleft & \searrow \\ Jw & \xrightarrow{y} & Y \end{array}$$


Lemma 4.4. *(\mathbb{E}, \mathbb{S}) -open maps are closed under composition and pullbacks.*

Theorem 4.5. *If \mathbb{M} has pullbacks, then (\mathbb{E}, \mathbb{S}) -bisimilarity is a transitive relation, and thus is an equivalence relation.*

Generalized Path Bisimulations. In the classical open map setup [16], another notion of bisimilarity can be defined by using path extensions directly: so-called strong path and path bisimulations, which can be generalized as follows. Like originally [16], we assume that there is an element $0 \in V$, such that $J0$ is an initial object of \mathbb{M} (note that 0 is not required to be initial in \mathbb{E} or \mathbb{S}). The intuition is that the unique morphism $!_X : J0 \rightarrow X$ points to the initial state of X . For example, $J0$ can be given by $(1, \text{id}_1, \perp)$ in a category of pointed coalgebras if 1 is the final object of \mathcal{C} and if $\mathcal{C}(1, F1)$ has the least element $\perp : 1 \rightarrow F1$ (those conditions hold in the cases of interest).


Definition 4.6. A path simulation from A to B in \mathbb{M} is a set R of spans of the form $A \xleftarrow{a} Jv \xrightarrow{b} B$ (for $v \in V$) satisfying the following two properties

- **initial condition:** the span $A \xleftarrow{!_A} J0 \xrightarrow{!_B} B$ belongs to R .
- **forward closure:** for all spans $A \xleftarrow{a} Jv \xrightarrow{b} B$ in R , all $e : v \rightarrow w \in \mathbb{E}$ and all $a' : Jw \rightarrow A \in \mathbb{M}$ such that $a = a' \cdot J_{\mathbb{E}}e$, there are $e' : v \rightarrow u \in \mathbb{E}$, $s : u \rightarrow w \in \mathbb{S}$, and $b' : Ju \rightarrow B \in \mathbb{M}$ such that $J_{\mathbb{E}}e = J_{\mathbb{S}}s \cdot J_{\mathbb{E}}e'$, $b = b' \cdot J_{\mathbb{E}}e'$, and the span $A \xleftarrow{a' \cdot J_{\mathbb{S}}s} Ju \xrightarrow{b'} B$ belongs to R .



We say that R is a strong path simulation if it additionally satisfies the following:

- **backward closure:** for all spans $A \xleftarrow{a} Jv \xrightarrow{b} B$ in R and all $e : w \rightarrow v \in \mathbb{E}$, we have that the span $A \xleftarrow{a \cdot J_{\mathbb{E}}e} Jw \xrightarrow{b \cdot J_{\mathbb{E}}e} B$ belongs to R .




We say that R is a (strong) path bisimulation from A to B if R and $R^\dagger = \{B \xleftarrow{b} Jv \xrightarrow{a} A \mid A \xleftarrow{a} Jv \xrightarrow{b} B \in R\}$ are (strong) path simulations.

Remark that this version of (strong) path bisimulations has the same type as the one by Joyal et al. [16], but satisfies more general conditions. In particular, when \mathbb{S} is a discrete category, the formulation above is exactly the one from [16]. Obviously, a strong path bisimulation is a path bisimulation.

The main result of this section is the following.

Theorem 4.7. Assume two models A and B in \mathbb{M} . If there is a span $A \xleftarrow{f} C \xrightarrow{g} B$ where g is a morphism of \mathbb{M} and f is an (\mathbb{E}, \mathbb{S}) -open map, then the following set is a strong path simulation:

$$R_{f,g} := \{A \xleftarrow{a} Jv \xrightarrow{b} B \mid \exists c : Jv \rightarrow C \text{ with } a = f \cdot c \text{ and } b = g \cdot c\}$$



Consequently, if A and B are (\mathbb{E}, \mathbb{S}) -bisimilar, then there is strong path bisimulation between them.

As in the classical case of [16], there is no reason for the converse to be true in general: there might be a strong path bisimulation between two models, but no span of generalized open maps. However, conditions from [6] could be accommodated to describe a general framework in which the converse holds. Since this is not the main focus of this paper, we will not do it here, but will show a particular case in Section 6.

Generators of the Category of Extensions. In the first example of open maps for LTSs introduced in Section 2, the path category was described as the poset of words with the prefix order. Consequently, to prove that a functional simulation is J -open, we have to prove the lifting property of Definition 4.1 with respect to all pairs $w \leq w'$. However, it is sufficient to check the lifting property for extensions by one letter: $w' = w.a$ for some $a \in A$. The general reason is that, as a category, (A^*, \leq) is generated by the morphisms $w \leq w.a$, and verifying the lifting property with respect to generators of the category \mathbb{P} is enough to obtain J -openness. This can be extended to generalized open maps, with additional care.

Proposition 4.8. *Assume a subgraph \mathbb{E}' of \mathbb{E} that generates \mathbb{E} , that is, every morphism of \mathbb{E} is a finite composition of morphisms of \mathbb{E}' . Assume additionally, that for every $e \in \mathbb{E}'$ and $s \in \mathbb{S}$ for which $J_{\mathbb{E}}e \cdot J_{\mathbb{S}}s$ is well-defined, there are $s' \in \mathbb{S}$ and $e' \in \mathbb{E}'$ such that $J_{\mathbb{E}}e \cdot J_{\mathbb{S}}s = J_{\mathbb{S}}s' \cdot J_{\mathbb{E}}e'$.*

In that case, if a morphism of \mathbb{M} satisfies the lifting property of Definition 4.1 for all morphisms in \mathbb{E}' , then it is (\mathbb{E}, \mathbb{S}) -open. Also, if a set of spans satisfies the conditions of Definition 4.6, where \mathbb{E} is replaced by \mathbb{E}' , then it is a (strong) path bisimulation.

The first condition is satisfied when \mathbb{E} is a free category and \mathbb{E}' is its class of generators. The second condition is satisfied for e.g. $\mathbb{E} = \mathbb{P}$ and $\mathbb{S} = |\mathbb{P}|$.

5 Open Maps for Weighted Systems

In this section, we will prove that weighted systems can be captured by this generalized open map theory for a large variety of weights, including those needed to capture probabilistic systems.

5.1 Category of Coalgebras for Weighted Systems

In this section, we will consider weighted functors as follows.

Definition 5.1. *Given a commutative monoid $(K, +, e)$, the K -weighted functor $(K, +, e)^{(-)}: \mathbf{Set} \rightarrow \mathbf{Set}$ is defined as follows on sets and maps:*

$$\begin{aligned} \text{sets:} \quad X &\mapsto (K, +, e)^{(X)} = \{\mu: X \rightarrow K \mid \mu^{-1}(K \setminus \{e\}) \text{ is finite}\} \\ \text{maps:} \quad f: X \rightarrow Y &\mapsto (K, +, e)^{(f)}(\mu) = (y \in Y \mapsto \sum \{\mu(x) \mid x \in X, f(x) = y\}) \end{aligned}$$

An element μ of $(K, +, e)^{(X)}$ is a finite distributions sending each $x \in X$ to a weight in K . Whenever a map $f: X \rightarrow Y$ identifies elements $f(x_1) = f(x_2) = \dots$, then the functor action turns μ into a distribution on Y by adding up the weights $\mu(x_1) + \mu(x_2) + \dots$ as elements of X are sent to the same element in Y . Since μ is finite and K is commutative, this addition is well-defined.

Given a commutative monoid $(K, +, e)$ and an alphabet A , we want to consider weighted systems as coalgebras for the functor $(K, +, e)^{(A \times -)}$. As described in Section 2, we want to be able to talk about lax homomorphisms, so we need an order on $(K, +, e)^{(A \times -)}$ as in Definition 2.8. For that, we need to assume an ordered commutative monoid $(K, +, e, \sqsubseteq)$, that is, a monoid $(K, +, e)$ with a partial order \sqsubseteq such that $+$ is monotone in both its arguments.

Lemma 5.2. *Given an ordered commutative monoid $(K, +, e, \sqsubseteq)$, then for all sets X and Y , the relation on the hom-set $\text{Set}(X, (K, +, e)^{(A \times Y)})$ defined by*

$$f_1 \sqsubseteq f_2 \iff \forall x \in X, \forall y \in Y, \forall a \in A, f_1(x)(a, y) \sqsubseteq f_2(x)(a, y)$$

is an order on $(K, +, e)^{(A \times -)}$.

So, we have a category $\text{LCoalg}\left(1, (K, +, e)^{(A \times -)}\right)$ of pointed coalgebras and lax homomorphisms. The goal of this section is to design a generalized open maps situation for which (\mathbb{E}, \mathbb{S}) -bisimilarity characterizes coalgebraic bisimilarity and more precisely for which (\mathbb{E}, \mathbb{S}) -openness characterizes proper homomorphisms.

In the course of the constructions and proofs, we will need additional assumptions that we list here.

Definition 5.3. *We call an ordered commutative monoid $(K, +, e, \sqsubseteq)$ a rearrangement monoid if it satisfies the additional requirement that if $n, m \geq 1$ and*

$$\sum_{i=1}^n x_i \sqsubseteq \sum_{j=1}^m y_j,$$

then there exists a family $(u_{i,j})_{1 \leq i \leq n, 1 \leq j \leq m}$ such that

$$\text{for all } j, \sum_{i=1}^n u_{i,j} \sqsubseteq y_j \quad \text{and} \quad \text{for all } i, \sum_{j=1}^m u_{i,j} = x_i.$$

In addition, we say that a rearrangement monoid is strict if the condition above holds also when replacing \sqsubseteq with $=$.

The intuition is as follows. We have some weights arranged as x_1, \dots, x_n . We want to be able to decompose those weights into smaller weights, the $u_{i,j}$ s, and by rearranging those small weights obtaining weights smaller than the y_j . This condition states that this is possible when there is enough weight in total. The special case of strictness is called the *row-column property* in [17].

Lemma 5.4. *For any subgroup G of the real numbers $(\mathbb{R}^n, +, -, 0)$ such that for all x, y in G $(\min(x_1, y_1), \dots, \min(x_n, y_n)) \in G$, the monoids $(G, +, 0, \leq)$ and $(G_{\geq 0}, +, 0, \leq)$, where \leq is the usual order on \mathbb{R}^n , are strict rearrangement monoids.*

For any lattice with bottom element $(L, \leq, \sqcup, \sqcap, \perp)$, (L, \sqcup, \perp, \leq) is a rearrangement monoid if and only if $(L, \leq, \sqcup, \sqcap)$ is distributive. Furthermore, in that case, it is always strict.

Another property is a form of positivity: we say that an ordered monoid is *positively ordered* if e is the bottom element for \sqsubseteq , that is, for all $k \in K$, $e \sqsubseteq k$.

Example 5.5. The positive real line $(\mathbb{R}_+, +, 0, \leq)$ is a positively ordered strict rearrangement monoid and it is necessary to define probabilistic systems. Another example is the monoid of natural numbers $(\mathbb{N}, +, 0, \leq)$, which defines the bag functor. Finally, any distributive lattice with bottom element (L, \sqcup, \perp, \leq) , typically powerset lattices $(\mathcal{P}(X), \cup, \emptyset, \subseteq)$, is too. On the contrary, $(\mathbb{R}, +, 0, \leq)$ and $(\mathbb{Z}, +, 0, \leq)$ are strict rearrangement monoids but are not positively ordered. Conversely $(\mathbb{N}_{\geq 1}, \times, 1, \leq)$ is positively ordered but not a rearrangement monoid. Indeed, it is impossible to rearrange the inequality $2 \times 5 \leq 3 \times 4$.

5.2 Generalized Open Maps Situation for Weighted Systems

Let $(K, +, e, \sqsubseteq)$ be a commutative ordered monoid. Elements of V_K are

- either words on $A \times (K \setminus \{e\})$, $w = (a_1, k_1), \dots, (a_n, k_n)$,
- or triples (w_1, b, w_2) of a word w_1 on $A \times (K \setminus \{e\})$, a letter $b \in A$, and a non-empty word w_2 on $(K \setminus \{e\})$.

The function J_K maps

- a word $w = (a_1, k_1), \dots, (a_n, k_n)$ to the system

$$Jw = \boxed{\rightarrow 0 \xrightarrow{(a_1, k_1)} 1 \xrightarrow{(a_2, k_2)} \dots \xrightarrow{(a_n, k_n)} n}$$

that is, to the coalgebra $Jw: \{0, \dots, n\} \rightarrow K^{(A \times \{0, \dots, n\})}$ such that if $b = a_{i+1}$ and $j = i + 1$ then $Jw(i)(b, j) = k_{i+1}$, else $= e$.

- a triple (w_1, b, w_2) with $w_1 = (a_1, k_1), \dots, (a_n, k_n)$ and $w_2 = l_1, \dots, l_m$ is mapped to the system

$$J(w_1, b, w_2) = \boxed{\rightarrow 0 \xrightarrow{(a_1, k_1)} 1 \xrightarrow{(a_2, k_2)} \dots \xrightarrow{(a_n, k_n)} n \begin{array}{l} \xrightarrow{(b, l_1)} (n+1, 1) \\ \vdots \\ \xrightarrow{(b, l_m)} (n+1, m) \end{array}}$$

that is, $J(w_1, b, w_2)(n)(n+1, i) = (b, l_i)$.

The category \mathbb{E}_K is defined as follows. For every w_1, b , and w_2 , there is a unique edge e from w_1 to (w_1, b, w_2) . The functor then maps this edge e to $J_{\mathbb{E}}e$, the obvious injection.

The category \mathbb{S}_K has two types of morphisms:




Fig. 1. Example of a lifting of a path extension in \mathbb{R}_+ -weighted systems and for a singleton label alphabet $|A| = 1$, thus omitting action labels.

- identities on words w_1 ,
- morphisms from (w_1, b, w'_2) to (w_1, b, w_2) , with $w'_2 = l'_1, \dots, l'_{m'}$ and $m \leq m'$, which are given by surjective monotone functions $s: \{1, \dots, m'\} \rightarrow \{1, \dots, m\}$ such that for all $i \leq m$, $l_i = \sum_{\{j|s(j)=i\}} l'_j$.

The functor $J_{\mathbb{S}}$ then maps s of the second type to the proper homomorphism $J_{\mathbb{S}}s$ which maps i to i and $(n+1, j)$ to $(n+1, s(j))$.

As a piece of notation, for a morphism $x: Jw_1 \rightarrow X$, with w_1 of length n we denote $x(n) \in X$ by $\text{end}(x)$. We then say that a state p of X is reachable if there is a morphism of type $x: Jw_1 \rightarrow X$ with $\text{end}(x) = p$. By extension, we say that X is reachable if all its states are reachable.

5.3 Equivalence between Open Maps and Proper Homomorphisms

An example of an (\mathbb{E}, \mathbb{S}) -open map h is provided in Figure 1, together with a path extension that is lifted. Like it is often the case in the non-deterministic systems, the lifting map d is not unique. Hence, only existence (and no uniqueness) is required in the lifting property. Since h is a proper homomorphism, it provides a lifting for all extensions, as we show in general:

Theorem 5.6. *Assume a lax homomorphism $f: X \rightarrow Y$. If f is $(\mathbb{E}_K, \mathbb{S}_K)$ -open, X is reachable, and K is positively ordered, then f is a proper homomorphism. Conversely, if f is a proper homomorphism and K is a rearrangement monoid, then f is $(\mathbb{E}_K, \mathbb{S}_K)$ -open. In particular, if K is a positively ordered rearrangement monoid, two weighted systems X and Y are $(\mathbb{E}_K, \mathbb{S}_K)$ -bisimilar if and only if they are coalgebraically bisimilar.*

For an endofunctor on \mathbf{Set} , to prove that coalgebraic bisimilarity is an equivalence relation it is enough to show that the functor preserves weak-pullbacks. In the case of the weighted functor, this is given by strictness (see also [17]):

Corollary 5.7. *If K is a positively ordered strict rearrangement monoid, then $(\mathbb{E}_K, \mathbb{S}_K)$ -bisimilarity is an equivalence relation.*

5.4 About Sub-distribution Functor

Until now, we have not dealt with probabilistic systems, that is, coalgebras for the sub-distribution functor $\mathcal{D}_{\leq 1}$. Those coalgebras are particular cases of coalgebras for the weighted functor $X \mapsto (\mathbb{R}_+, +)^{(X)}$. We want to show in this section that it is equivalent to consider coalgebras for $X \mapsto \mathcal{D}_{\leq 1}(A \times X)$ as coalgebras for $X \mapsto (\mathbb{R}_+, +)^{(A \times X)}$, in the sense that, two coalgebras for the former are bisimilar if and only if they are bisimilar when seen as coalgebras for the latter. The main ingredient is the following remark.

Lemma 5.8. *Assume a pointed coalgebra $1 \xrightarrow{i} X \xrightarrow{c} \mathcal{D}_{\leq 1}(A \times X)$ and assume given a lax (resp. proper) homomorphism f from $1 \xrightarrow{j} Y \xrightarrow{d} (\mathbb{R}_+, +)^{(A \times Y)}$ to $1 \xrightarrow{i} X \xrightarrow{c} \mathcal{D}_{\leq 1}(A \times X) \subseteq (\mathbb{R}_+, +)^{(A \times X)}$. Then $Y \xrightarrow{d} \mathcal{D}_{\leq 1}(A \times Y)$ and f is a lax (resp. proper) homomorphism from $1 \xrightarrow{j} Y \xrightarrow{d} \mathcal{D}_{\leq 1}(A \times Y)$ to $1 \xrightarrow{i} X \xrightarrow{c} \mathcal{D}_{\leq 1}(A \times X)$.*

Remark that this property is not true for the proper distribution functor \mathcal{D} . This suggests that we can define a generalized open maps situation $\mathbb{E}_{\mathcal{D}}, \mathbb{S}_{\mathcal{D}}$ for coalgebras for the functor $X \mapsto \mathcal{D}_{\leq 1}(A \times X)$ by considering $\mathbb{E}_{(\mathbb{R}_+, +)}, \mathbb{S}_{(\mathbb{R}_+, +)}$ as defined in Section 5.2, and restricting it to those v such that Jv is a coalgebra for $X \mapsto \mathcal{D}_{\leq 1}(A \times X)$.

Corollary 5.9. *A lax homomorphism from $1 \xrightarrow{j} Y \xrightarrow{d} \mathcal{D}_{\leq 1}(A \times Y)$ to $1 \xrightarrow{i} X \xrightarrow{c} \mathcal{D}_{\leq 1}(A \times X)$ is $(\mathbb{E}_{\mathcal{D}}, \mathbb{S}_{\mathcal{D}})$ -open if and only if it is $(\mathbb{E}_{(\mathbb{R}_+, +)}, \mathbb{S}_{(\mathbb{R}_+, +)})$ -open. Furthermore, two $\mathcal{D}_{\leq 1}(A \times _)$ -coalgebras are $(\mathbb{E}_{\mathcal{D}}, \mathbb{S}_{\mathcal{D}})$ -bisimilar if and only if they are $(\mathbb{E}_{(\mathbb{R}_+, +)}, \mathbb{S}_{(\mathbb{R}_+, +)})$ -bisimilar.*

Finally, the main result of this section:

Theorem 5.10. *Let $f: X \rightarrow Y$ be a lax homomorphism between $\mathcal{D}_{\leq 1}(A \times _)$ -coalgebras (X, c, i) and (Y, d, j) . If (X, c, i) is reachable and f is $(\mathbb{E}_{\mathcal{D}}, \mathbb{S}_{\mathcal{D}})$ -open, then f is a proper homomorphism. Conversely, if f is a proper homomorphism, then it is $(\mathbb{E}_{\mathcal{D}}, \mathbb{S}_{\mathcal{D}})$ -open. Moreover, two $\mathcal{D}_{\leq 1}(A \times _)$ -coalgebras (X, c, i) and (Y, d, j) are $(\mathbb{E}_{\mathcal{D}}, \mathbb{S}_{\mathcal{D}})$ -bisimilar if and only if they are coalgebraically bisimilar.*

6 Open Maps for Branching Bisimilarity

In this section, we present a new way of modeling branching and weak bisimulations using our generalized framework of open maps. Using this additional flexibility, we do not need to rely on weak morphisms anymore, but on a slight modification of the morphism described in Definition 2.1. Concretely, we build a

generalized open map situation such that stuttering branching bisimulations coincide with strong path bisimulations, and that in this case, they precisely characterize (\mathbb{E}, \mathbb{S}) -bisimilarity. In addition, in this framework, path bisimulations precisely correspond to weak bisimulations, witnessing branching bisimilarity as the history-preserving analogue to weak bisimilarity.

6.1 LTSs with Internal Moves, Category and Bisimilarities

Definition 6.1. For a fixed set A of labels with a particular element τ (called internal move), the category **WLTS** contains the same objects as **LTS**, and its morphisms $f: (X, \rightarrow, x_0) \rightarrow (Y, \rightarrow, y_0)$ are functions $f: X \rightarrow Y$ such that $f(x_0) = y_0$ and for all $x \xrightarrow{a} x'$ in X , we have $f(x) \xrightarrow{a} f(x')$, or $a = \tau$ and $f(x) = f(x')$.

LTS is a (non-full) subcategory of **WLTS**, and in fact the **LTS**-morphisms will be used later in the paper. For easier distinction, we use the terminology *strong morphisms* for **WLTS**-morphisms that are also in **LTS** (alluding to *strong bisimulations* which were the bisimulation notion in **LTS**). Another notion of morphisms are so-called *weak morphisms* [3]:

- if $x \xrightarrow{a} x'$ in X , then $f(x) \xrightarrow{\tau}^* \xrightarrow{a} \xrightarrow{\tau}^* f(x')$ in Y ,
- if $x \xrightarrow{\tau} x'$ in X , then $f(x) \xrightarrow{\tau}^* f(x')$ in Y .

Though we do not use weak morphisms in the following development of the paper, it is worth mentioning the **WLTS**-morphisms form a proper subclass of the weak morphisms.

Definition 6.2. A branching bisimulation from (X, \rightarrow_X, i_X) to (Y, \rightarrow_Y, i_Y) is a relation $R \subseteq X \times Y$ such that $(i_X, i_Y) \in R$, and for $(x, y) \in R$:

- if $x \xrightarrow{a} x'$ then
 - $a = \tau$ and $(x', y) \in R$, or
 - $y \xrightarrow{\tau} y_1 \xrightarrow{\tau} \dots \xrightarrow{\tau} y_n \xrightarrow{a} z_1 \xrightarrow{\tau} \dots \xrightarrow{\tau} z_m$ such that (x, y_n) , (x', z_1) , and $(x', z_m) \in R$.
- symmetrically when $y \xrightarrow{a} y'$.

If furthermore in the second condition (x, y_i) , $(x', z_i) \in R$ for all i (and symmetrically in the third condition), then R is said to be stuttering.

It is known from [23] that the largest branching bisimulation is stuttering, so that both notions generate the same bisimilarity. In the following, we will prove that strong path bisimulations are more naturally related to stuttering branching bisimulations thanks to their backward closure.

Definition 6.3. A weak bisimulation from (X, \rightarrow_X, i_X) to (Y, \rightarrow_Y, i_Y) is a relation $R \subseteq X \times Y$ such that $(i_X, i_Y) \in R$, and for $(x, y) \in R$:


- if $x \xrightarrow{\tau} x'$, then there is y' such that $(x', y') \in R$ and $y \xrightarrow{\tau}^* y'$,
- if $x \xrightarrow{a} x'$ with $a \neq \tau$, then there is y' such that $(x', y') \in R$ and $y \xrightarrow{\tau}^* \xrightarrow{a} \xrightarrow{\tau}^* y'$.
- symmetrically when $y \xrightarrow{\tau} y'$ or $y \xrightarrow{a} y'$.

It is clear that a (stuttering) branching bisimulation is a weak bisimulation.

6.2 Generalized Open Maps for Branching Bisimulations

In this section, we describe the generalized open maps situation that captures branching bisimulation. Like for plain LTSs (Def. 2.2), elements of V will be words on A , representing a finite linear LTS labelled by this word. However, to emphasize the particularity of the internal move τ , we will provide another presentation here.

Here, V is the set of sequences of the form: $v = n_1, a_1, n_2, \dots, n_k, a_k, n_{k+1}$ such that $a_i \in A \setminus \{\tau\}$ and $n_i \in \mathbb{N}$, e.g. $\tau a \tau b c \tau \hat{=} 2, a, 1, b, 0, c, 1$. The natural numbers $n_i \in \mathbb{N} \cong \{\tau\}^*$ represent the number of internal moves between two observable moves. Then, J maps this sequence to the usual linear LTS:



Elements of \mathbb{E} append *at most* one observable (i.e. non- τ) move:

- **Only internal moves:** for sequences $v = n_1, a_1, \dots, a_k, n_{k+1}$ and $w = n_1, a_1, \dots, a_k, n'_{k+1}$ with $n_{k+1} \leq n'_{k+1}$ there is a unique edge $e_\tau: v \rightarrow w$ in \mathbb{E} , e.g. $e_\tau: 2, a, 1, b, 0, c, 1 \rightarrow 2, a, 1, b, 0, c, 3$
- **One observable move:** for sequences $v = n_1, a_1, \dots, a_k, n_{k+1}$ and $w = n_1, a_1, \dots, a_k, n'_{k+1}, a, n_{k+2}$ with $n_{k+1} \leq n'_{k+1}$ there is a unique edge $e_a: v \rightarrow w$ in \mathbb{E} .

The graph morphism $J_{\mathbb{E}}: \mathbb{E} \rightarrow \mathbb{M}$ maps those edges to the obvious inclusion, mapping state (i, j) of Jv to the same state in Jw .

Strictly speaking, \mathbb{E} is not a category, but just a graph, because we have $a \xrightarrow{e_b} ab$ and $ab \xrightarrow{e_c} abc$, but there is no morphism from a to abc . To fit in the framework of Section 4, we take the free category $\text{Free}(\mathbb{E})$ generated by this graph and the unique functor extending the graph homomorphism $J_{\mathbb{E}}$. By Proposition 4.8, it is equivalent to consider $\text{Free}(\mathbb{E})$ and \mathbb{E} for openness and path bisimulations, so we will talk of (\mathbb{E}, \mathbb{S}) -openness, when we mean $(\text{Free}(\mathbb{E}), \mathbb{S})$ -openness, and all the statements and proofs will be done using \mathbb{E} only.

Elements of \mathbb{S} are trickier to describe. The intuition is that they are morphisms that merge states. In the context of LTSs with internal moves, merging happens when the source and the target of a τ -transition are mapped to the same state. This is crucial for the open maps we want to describe: to lift one τ -transition, it might be necessary to use several τ -transitions. With this knowledge, elements of \mathbb{S} are as follows.

- **Merging internal moves:** morphisms in \mathbb{S} from $v = n_1, a_1, \dots, a_k, n_{k+1}$ to $w = n'_1, a_1, \dots, a_k, n'_{k+1}$ with $n_i \geq n'_i$ are $(k+1)$ -tuples $s = (s_1, \dots, s_{k+1})$ of monotone surjective functions $s_i: \{0 < 1 < \dots < n_i\} \rightarrow \{0 < 1 < \dots < n'_i\}$.

For example, there are two morphisms from $a\tau\tau b \hat{=} 0, a, 2, b, 0$ to $a\tau b \hat{=} 0, a, 1, b, 0$, one for each τ that can be dropped. The functor $J_{\mathbb{S}}$ then maps s to the morphism from Jv to Jw defined by $J_{\mathbb{S}}(s)(i, j) = (s_j(i), j)$.

As a piece of notation, for a morphism $x: J(n_1, a_1, \dots, a_k, n_{k+1}) \rightarrow X$, we denote $x(n_{k+1}, k+1) \in X$ by $\text{end}(x)$.

6.3 Equivalence of Bisimilarities

In this section, we prove that (\mathbb{E}, \mathbb{S}) -bisimilarity indeed coincides with branching bisimilarity. To do so, we prove first that for the present instance of \mathbb{E} and \mathbb{S} (Sec. 6.2), (\mathbb{E}, \mathbb{S}) -bisimilarity coincides with strong path bisimilarity. In general, (\mathbb{E}, \mathbb{S}) -bisimilarity implies strong path bisimilarity (Theorem 4.7), so it remains to show the converse direction for the present instance. To this end, we start by internalizing strong path bisimulations into objects of LTS/WLTS, in order to relate it them to open maps:

Definition 6.4. *For a strong path bisimulation R from X to Y , define the LTS $\tilde{R} = (R, \rightarrow_R, (X \xleftarrow{!} J0 \xrightarrow{!} Y))$ to have transitions*

$$(X \xleftarrow{x} Jv \xrightarrow{y} Y) \xrightarrow{a}_R (X \xleftarrow{x'} Jw \xrightarrow{y'} Y)$$

- for $a \neq \tau$ with $v = (n_1, a_1, \dots, a_k, n_{k+1})$, $w = (n_0, a_1, \dots, a_k, n_{k+1}, a, 0)$, $x' = x \cdot J_{\mathbb{E}}e_a$, and $y' = y \cdot J_{\mathbb{E}}e_a$ (for the unique $e_a: v \rightarrow w$);
- for $a = \tau$ with $v = (n_1, a_1, \dots, a_k, n_{k+1})$, $w = (n_1, a_1, \dots, a_k, n_{k+1} + 1)$, $x' = x \cdot J_{\mathbb{E}}e_\tau$, and $y' = y \cdot J_{\mathbb{E}}e_\tau$ (for the unique $e_\tau: v \rightarrow w$).

As a first observation, we describe runs in \tilde{R} in terms of projection maps:

Lemma 6.5. *In WLTS, we have projection maps $X \xleftarrow{\pi_X} \tilde{R} \xrightarrow{\pi_Y} Y$ given by $\pi_X: (X \xleftarrow{x} Jv \xrightarrow{y} Y) \mapsto \text{end}(x)$ and $\pi_Y: (X \xleftarrow{x} Jv \xrightarrow{y} Y) \mapsto \text{end}(y)$. For every strong morphism $r: Jv \rightarrow \tilde{R}$ (i.e. $r \in \text{LTS}$),*

$$\text{end}(r) \text{ is of the form } (X \xleftarrow{\pi_X \cdot r} Jv \xrightarrow{\pi_Y \cdot r} Y).$$

Remark that in this statement, we require r to be strong and not just a morphism of WLTS. With a morphism of WLTS, the statement would become that there is $s: v' \rightarrow v \in \mathbb{S}$ such that $\pi_X \cdot r = x \cdot J_{\mathbb{S}}s$ instead. For the characterization of open maps in WLTS, it suffices for our needs to restrict to strong morphisms:

Lemma 6.6. *For $f: X \rightarrow Y$ in WLTS to be (\mathbb{E}, \mathbb{S}) -open, it is sufficient to verify the lifting in Definition 4.1 in the special case of x being a strong morphism.*

We use this simplification to prove that the projection maps π_X, π_Y are open:

Proposition 6.7. *For a strong path bisimulation R from X to Y , the projections $X \xleftarrow{\pi_X} \tilde{R} \xrightarrow{\pi_Y} Y$ are both (\mathbb{E}, \mathbb{S}) -open.*

The next step is to prove the equivalence between strong path and stuttering branching bisimulations.

Table 1. Equivalences of bisimilarity notions in LTSs with τ -actions $X, Y \in \text{WLTS}$

branching bisimilarity	\iff strong path bisimilarity (Theorem 6.8)
	\iff (\mathbb{E}, \mathbb{S}) -bisimilarity (Proposition 6.7 & Theorem 4.7)
weak bisimilarity	\iff path bisimilarity (Theorem 6.9)

Theorem 6.8. *If R is a stuttering branching bisimulation from X to Y , then*

$$\bar{R} = \{X \xleftarrow{x} Jv \xrightarrow{y} Y \mid v = (n_1, a_1, \dots, n_{k+1}) \wedge \forall i, j. (x(i, j), y(i, j)) \in R\}$$

is a strong path bisimulation. Conversely, if R is a strong path bisimulation, then

$$\check{R} = \{(end(x), end(y)) \mid (X \xleftarrow{x} Jv \xrightarrow{y} Y) \in R\}$$

is a stuttering branching bisimulation.

The same reasoning can be made for weak and path bisimulations:

Theorem 6.9. *If R is a weak bisimulation from X to Y , then*

$$\hat{R} = \{X \xleftarrow{x} Jv \xrightarrow{y} Y \mid (end(x), end(y)) \in R\}$$

is a path bisimulation. If R is a path bisimulation, then \check{R} is a weak bisimulation.

In total, we can describe branching and weak bisimilarity by categorical bisimilarity notions, as summarized in Table 1.

7 Conclusions and Future Work

In this paper, we investigate bisimilarities of weighted and probabilistic systems through the theory of open maps. After showing that the usual theory cannot capture weights, we provide a faithful extension of the theory by the notion of mergings. The new theory has similar properties (equivalence relation, characterization as sets of spans, restriction to generators) as classical open maps but also captures bisimilarity of weighted systems and even branching bisimilarity.

The new instances come at the cost of more parameters to the theory. It remains for future work whether the parameters \mathbb{E}, \mathbb{S} can be combined in a single path category with two morphism classes and morphism factorizations. It would also be illuminating to know whether this new theory satisfies the axioms of a *class of open maps* from [15], in particular for toposes of coalgebras [14].

For the framework as presented, we would like to formally relate it to coalgebra – as this has been done for non-deterministic systems [19,25]. Furthermore, we would like to investigate how system semantics of true concurrency, such as Higher Dimensional Automata [21] can be integrated. Designing open maps for them turned out to be complicated (see [8]), but a hope would be that the addition of mergings would allow modeling homotopy more naturally.

Finally, it would be interesting to see whether our theory capture quantitative extensions of systems classically modeled by open maps, such as probabilistic and quantum extensions of petri nets and event structures (see [24] for example).

References

1. Beohar, H., Cuijpers, P.J.L.: Open Maps in Concrete Categories and Branching Bisimulation for Prefix Orders. *Electronic Notes in Theoretical Computer Science* **319**, 51–66 (2015)
2. Beohar, H., Küpper, S.: Bisimulation Maps in Presheaf Categories. *Electronic Notes in Theoretical Computer Science* **347**, 5–24 (2019)
3. Cheng, A., Nielsen, M.: Open Maps (at) Work. B R I C S Report Series (RS-95-23) (1995)
4. Danos, V., Desharnais, J., Laviolette, F., Panangaden, P.: Bisimulation and cocongruence for probabilistic systems. *Information and Computation* **204**(4), 503–523 (2006)
5. Desharnais, J., Edalat, A., Panangaden, P.: Bisimulation for Labelled Markov Processes. *Information and Computation* **179**(2), 163–193 (2003)
6. Dubut, J., Goubault, E., Goubault-Larrecq, J.: Bisimulations and unfolding in P-accessible categorical models. In: Desharnais, J., Jagadeesan, R. (eds.) 27th International Conference on Concurrency Theory, CONCUR 2016. LIPIcs, vol. 59, pp. 25:1–25:14. Schloss Dagstuhl - Leibniz-Zentrum für Informatik (2016)
7. Dubut, J., Hasuo, I., Katsumata, S., Sprunger, D.: Quantitative bisimulations using coreflections and open morphisms (2018), arXiv:1809.09278
8. Fahrenberg, U., Legay, A.: History-Preserving Bisimilarity for Higher-Dimensional Automata via Open Maps. *Electronic Notes in Theoretical Computer Science* **298**, 165–178 (2013)
9. Fiore, M., Cattani, G.L., Winskel, G.: Weak bisimulation and open maps. In: Proceedings. 14th Symposium on Logic in Computer Science. pp. 67–76 (1999)
10. Freyd, P., Scedrov, A.: *Categories, Allegories*, Mathematical Library, vol. 39. North-Holland (1990)
11. Hughes, J., Jacobs, B.: Simulations in coalgebra. *Theor. Comput. Sci.* **327**(1–2), 71–108 (2004). <https://doi.org/10.1016/j.tcs.2004.07.022>, <https://doi.org/10.1016/j.tcs.2004.07.022>
12. Hune, T., Nielsen, M.: Timed bisimulations and open maps. In: Brim, L., Gruska, J., Zlatuška, J. (eds) *Mathematical Foundations of Computer Science 1998*. MFCS 1998. *Lecture Notes in Computer Science*, vol. 1450. Springer, Berlin, Heidelberg (1998)
13. Jacobs, B.: *Introduction to Coalgebra: Towards Mathematics of States and Observation*, Cambridge Tracts in Theoretical Computer Science, vol. 59. Cambridge University Press (2016)
14. Johnstone, P., Power, J., Tsujishita, T., Watanabe, H., Worrell, J.: On the structure of categories of coalgebras. *Theoretical Computer Science* **260**, 87–117 (2001)
15. Joyal, A., Moerdijk, I.: A completeness theorem for open maps. *Annals of Pure and Applied Logic* **70**, 51–86 (1994)
16. Joyal, A., Nielsen, M., Winskel, G.: Bisimulation from Open Maps. *Information and Computation* **127**, 164–185 (1996)
17. Klin, B.: *Semantics and Algebraic Specification*, *Lecture Notes in Computer Science*, vol. 5700, chap. *Structural Operational Semantics for Weighted Transition Systems*, pp. 121–139. Springer, Berlin, Heidelberg (2009)
18. Larsen, K.G., Skou, A.: Bisimulations through Probabilistic Testing. *Information and Computation* **94**, 1–28 (1991)
19. Lasota, S.: Coalgebra morphisms subsume open maps. *Theoretical Computer Science* **280**(1), 123 – 135 (2002). [https://doi.org/10.1016/S0304-3975\(01\)00023-8](https://doi.org/10.1016/S0304-3975(01)00023-8), coalgebraic Methods in Computer Science

20. Park, D.: Concurrency and automata on infinite sequences. In: Proceedings of the 5th GI-Conference on Theoretical Computer Science. Lecture Notes in Computer Science, vol. 104, pp. 167–183. Springer (1981)
21. Pratt, V.: Higher dimensional automata revisited. *Math. Struct. Comp. Sci.* (2005)
22. Rutten, J.: Universal coalgebra: a theory of systems. *Theoretical Computer Science* **249**(1), 3 – 80 (2000)
23. van Glabbeek, R.J., Weijland, W.P.: Branching Time and Abstraction in Bisimulation Semantics. *Journal of the ACM* **43**(3), 555–600 (1996)
24. Winskel, G.: Distributed probabilistic and quantum strategies. *Electronic Notes in Theoretical Computer Science* **298**, 403–425 (2013)
25. Wißmann, T., Dubut, J., Katsumata, S., Hasuo, I.: Path category for free. In: Bojańczyk, M., Simpson, A. (eds.) *Foundations of Software Science and Computation Structures (FoSSaCS 2019)*. pp. 523–540. Springer International Publishing, Cham (04 2019). https://doi.org/10.1007/978-3-030-17127-8_30

A Details for Section 2

Details for Remark 2.9

Proof. Hughes and Jacobs [11] work with preorders instead of partial orders, but adapting their definition [11, Def. 2.1] to partial orders yields: An order on a Set-functor $F: \text{Set} \rightarrow \text{Set}$ is a functor $\leq: \text{Set} \rightarrow \text{Pos}$ making the following diagram commute:

$$\begin{array}{ccc} & & \text{Pos} \\ & \nearrow \leq & \downarrow U \\ \text{Set} & \xrightarrow{F} & \text{Set} \end{array}$$

Every order on a functor \sqsubseteq in our sense (Definition 2.8) for $\mathcal{C} := \text{Set}$ induces such a functor $\leq: \text{Set} \rightarrow \text{Pos}$, because for every set Y , we can put $\leq_Y \in \text{Pos}$ to be the order \sqsubseteq on $\text{Set}(1, FY)$, identifying elements of FY with maps $1 \rightarrow FY$.

Conversely, given $\leq: \text{Set} \rightarrow \text{Pos}$ in the sense of Hughes and Jacobs, we can define the order on $f_1, f_2 \in \text{Set}(X, FY)$ by

$$f_1 \sqsubseteq f_2 : \iff \forall x \in X: f_1(x) \leq_Y f_2(x). \quad (1)$$

However, it might be possible that not all orders \sqsubseteq on functors in the sense of Definition 2.8 satisfy (1). \square

B Proof of Section 3

Recall that an object of the category $\text{LCoalg}(1, \mathcal{D}_{\leq 1}(A \times _))$ is a pointed coalgebra, that is, a pair of functions of the form

$$1 \xrightarrow{i} X \xrightarrow{\xi} \mathcal{D}_{\leq 1}(A \times X).$$

In the following, we will identify i with the element $x_0 \in X$, such that the image of i is $\{x_0\}$, and we will denote the pointed coalgebra above by (X, ξ, x_0) instead.

Proof of Theorem 3.1

Proof. Assume some J with the equivalence, and note that wlog we can assume that JP is reachable.³ For $Y = \boxed{\rightarrow r \xrightarrow{1, a} y}$ (implicitly defining $\zeta: Y \rightarrow \mathcal{D}_{\leq 1}(A \times Y)$ by $\zeta(r)(a, y) = 1$), consider the initial morphism $!_Y: 0_{\mathbb{M}} \rightarrow Y$, where $0_{\mathbb{M}} = \boxed{\rightarrow r}$. Here $0_{\mathbb{M}}$ has only one state and no transition. Since $!_Y$ is not a proper coalgebra homomorphism, it is not a J -open morphism. Hence there is some commutative square

$$\begin{array}{ccc} JP & \xrightarrow{p} & 0_{\mathbb{M}} \\ \downarrow J\phi & & \downarrow !_Y \\ JQ & \xrightarrow{q} & Y \end{array}$$

³ if there was such a J , then $R \cdot J$ is also such a candidate, where $R: \mathbb{M} \rightarrow \mathbb{M}_{\text{reach}}$ restrict a pointed coalgebra to its reachable states.

for which no diagonal lifting exists. Since no diagonal exists, $J\phi$ is no isomorphism. Since p is a lax coalgebra homomorphism, JP has no edges and hence $JP \cong 0_{\mathbb{M}}$, p is an iso. This implies that JQ has at least one edge, because $JQ \not\cong JP$, and it must be between distinct elements and labelled with a because q is a lax coalgebra homomorphism. Hence, there are $r, z \in JQ$, with $r \neq z$ and $r \xrightarrow{w, a} z$, $0 < w \leq 1$. Define $n = 2 \cdot \lceil \frac{1}{w} \rceil$ and

$$X = \boxed{r \begin{array}{l} \xrightarrow{1/n, a} x_1 \\ \xrightarrow{\quad \quad} \cdot \\ \xrightarrow{1/n, a} x_n \end{array}}, \text{ i.e. } X = \{r, x_1, \dots, x_n\}, \xi(r)((a, x_k)) = \frac{1}{n}, 1 \leq k \leq n.$$

We have a proper coalgebra homomorphism $h: X \rightarrow Y$ given by $h(r) = r$ and $h(x_k) = y$, $1 \leq k \leq n$. It is a proper coalgebra homomorphism, because $\zeta(r)(y) = 1 = \sum_{1 \leq k \leq n} \frac{1}{n} = \sum_{1 \leq k \leq n} \xi(r)(x_k)$. By initiality of $0_{\mathbb{M}}$ and so of JP , the following diagram commutes:

$$\begin{array}{ccc} JP & \xrightarrow{!_X} & X \\ \downarrow J\phi & & \downarrow h \\ JQ & \xrightarrow{q} & Y \end{array}$$

By the assumed equivalence, h is J -open, and thus, there is some diagonal lifting $d: JQ \rightarrow X$. Now, d being a lax coalgebra homomorphism leads to a contradiction, because d maps the edge $r \xrightarrow{w, a} z$ in JQ to some $d(r) \xrightarrow{w', a} d(z)$ with $w' \geq w$ in X . However, there is no such edge, because every edge in X has weight $\frac{1}{n}$ which is less than w' , which prove the first impossibility. Now, since h is a proper coalgebra homomorphism, X and Y are coalgebraically bisimilar. Let us prove they are not J -bisimilar. Assume that there is a span $Y \xleftarrow{f} Z \xrightarrow{g} X$ of J -open morphisms. By a similar argument as earlier, f being open means that there is a transition $r' \xrightarrow{w', a} z'$ in Z with $w' \geq w$. Since g is a lax homomorphism, g will map this transition on a transition of the form $r \xrightarrow{w'', a} x_i$ with $w'' \geq w' \geq w$, which is again impossible. \square

C Proofs of Section 4

Proof of Lemma 4.4

Proof. Let us prove that they are closed under composition first. So start with the following situation

$$\begin{array}{ccc} Jv & \xrightarrow{x} & X \\ \downarrow & & \downarrow f \\ J_e e & & Y \\ \downarrow & & \downarrow g \\ Jw & \xrightarrow{z} & Z \end{array}$$

reorganizing the arrows, we obtain the following square on which we apply the openness of g

$$\begin{array}{ccc}
 Jv & \xrightarrow{f \cdot x} & Y \\
 \downarrow & \dashv J_{\mathbb{E}}e' \circlearrowleft & \nearrow d' \\
 J_{\mathbb{E}}e & \circlearrowleft & Ju \\
 \downarrow & \dashv J_{\mathbb{S}}s \circlearrowleft & \downarrow g \\
 Jw & \xrightarrow{z} & Z
 \end{array}$$

We then transform the top triangle into a square on which we apply the openness of f

$$\begin{array}{ccc}
 Jv & \xrightarrow{x} & X \\
 \downarrow & \dashv J_{\mathbb{E}}e'' \circlearrowleft & \nearrow d' \\
 J_{\mathbb{E}}e' & \circlearrowleft & Ju' \\
 \downarrow & \dashv J_{\mathbb{S}}s' \circlearrowleft & \downarrow f \\
 Ju & \xrightarrow{d} & Y
 \end{array}$$

Putting things together, we obtain the desired lifting

$$\begin{array}{ccc}
 Jv & \xrightarrow{x} & X \\
 \downarrow & \dashv J_{\mathbb{E}}e'' \circlearrowleft & \nearrow d' \\
 J_{\mathbb{E}}e & \circlearrowleft & Ju' \\
 \downarrow & \dashv J_{\mathbb{S}}s' \circlearrowleft & \downarrow f \\
 Ju & \xrightarrow{d} & Y \\
 \downarrow & \dashv J_{\mathbb{S}}s \circlearrowleft & \downarrow g \\
 Jw & \xrightarrow{z} & Z
 \end{array}
 \quad \text{that is} \quad
 \begin{array}{ccc}
 Jv & \xrightarrow{x} & X \\
 \downarrow & \dashv J_{\mathbb{E}}e'' \circlearrowleft & \nearrow d' \\
 J_{\mathbb{E}}e & \circlearrowleft & Ju' \\
 \downarrow & \dashv J_{\mathbb{S}}(s \cdot s') \circlearrowleft & \downarrow g \cdot f \\
 Jw & \xrightarrow{z} & Z
 \end{array}$$

Now, let us prove they are closed under pullback, that is, if the following

$$\begin{array}{ccc}
 D & \xrightarrow{k} & C \\
 \downarrow h & \circlearrowleft & \downarrow g \\
 B & \xrightarrow{f} & A
 \end{array}$$

is a pullback square and f is (\mathbb{E}, \mathbb{S}) -open, then k is (\mathbb{E}, \mathbb{S}) -open. Start with the following situation:

$$\begin{array}{ccc}
 Jv & \xrightarrow{d} & D \\
 \downarrow J_{\mathbb{E}}e & \circlearrowleft & \downarrow k \\
 Jw & \xrightarrow{c} & C
 \end{array}$$

Composing with the pullback square, we obtain the following on which we apply the openness of f

$$\begin{array}{ccc}
 Jv & \xrightarrow{h \cdot d} & B \\
 \downarrow & \dashrightarrow^{J_{\mathbb{E}}e'} \circlearrowright & \downarrow \\
 J_{\mathbb{E}}e & \circlearrowright & Ju \\
 \downarrow & \dashrightarrow^{J_{\mathbb{S}}s} \circlearrowright & \downarrow \\
 Jw & \xrightarrow{g \cdot c} & A
 \end{array}$$

Reorganizing the arrows of the bottom right corner, we obtain the following square

$$\begin{array}{ccc}
 Ju & \xrightarrow{c \cdot J_{\mathbb{S}}s} & C \\
 \downarrow b & \circlearrowright & \downarrow g \\
 B & \xrightarrow{f} & A
 \end{array}$$

Applying the universality of the pullback in the assumption, we obtain the existence of a morphism $d' : Ju \rightarrow D$ such that

$$k \cdot d' = c \cdot J_{\mathbb{S}}s \quad \text{and} \quad h \cdot d' = b.$$

Let us prove that the following lifting works

$$\begin{array}{ccc}
 Jv & \xrightarrow{d} & D \\
 \downarrow & \dashrightarrow^{J_{\mathbb{E}}e'} \circlearrowright & \downarrow \\
 J_{\mathbb{E}}e & \circlearrowright & Ju \\
 \downarrow & \dashrightarrow^{J_{\mathbb{S}}s} \circlearrowright & \downarrow \\
 Jw & \xrightarrow{c} & C
 \end{array}$$

The bottom right corner ($k \cdot d' = c \cdot J_{\mathbb{S}}s$) and the left triangle ($J_{\mathbb{S}}s \cdot J_{\mathbb{E}}e' = J_{\mathbb{E}}e$) are already given. Let us prove $d' \cdot J_{\mathbb{E}}e' = d$. We already know that

$$h \cdot d' \cdot J_{\mathbb{E}}e' = b \cdot J_{\mathbb{E}}e' = h \cdot d.$$


and that

$$k \cdot d' \cdot J_{\mathbb{E}}e' = c \cdot J_{\mathbb{S}}s \cdot J_{\mathbb{E}}e' = c \cdot J_{\mathbb{E}}e = k \cdot d.$$

Then, by universality of the pullback square, $d' \cdot J_{\mathbb{E}}e' = d$. □

Proof of Theorem 4.5

Proof. To finish with transitivity, assume given two spans of open maps as follows (in plain)




Forming the pullback of g along h (dashed), which are both open, and using Lemma 4.4, l and m are also open. Using Lemma 4.4 again, $f \cdot l$ and $k \cdot m$ are open, and we have a span of open maps between A and C . \square


Proof of Theorem 4.7

Proof. Let us prove the three properties of the definition.


- **initial condition:** By the universal property of the initial object $J0$ the following diagram commutes:




- **forward closure:** Assume given the following situation:



Making the two triangles on the left into a square and applying the openness of f , we obtain:



We then obtain:



- **backward closure:** obvious.

\square

Proof of Proposition 4.8

Proof. It is enough to prove the following.

1. Assume that we have the following situation:

$$\begin{array}{ccc}
 Jv & \xrightarrow{b} & B \\
 \downarrow J_{\mathbb{E}}e & & \downarrow f \\
 Jw & \circlearrowleft & \\
 \downarrow J_{\mathbb{E}}e' & & \downarrow \\
 Jw' & \xrightarrow{a} & A
 \end{array}$$

where $e, e' \in \mathbb{E}'$. Applying the assumption on the following situation (in plain):

$$\begin{array}{ccc}
 Jv & \xrightarrow{b} & B \\
 \downarrow J_{\mathbb{E}}e & \dashrightarrow J_{\mathbb{E}}e'' \circlearrowleft & \downarrow f \\
 Jw & \circlearrowleft & Ju \\
 \downarrow J_{\mathbb{E}}e' & \dashrightarrow J_{\mathbb{S}}s \circlearrowleft & \downarrow \\
 Jw & \xrightarrow{a \cdot J_{\mathbb{E}}e'} & A
 \end{array}$$

we obtain the dashed data with $e'' \in \mathbb{E}'$. By assumption, there are $\tilde{e}' \in \mathbb{E}'$ and $\tilde{s} \in \mathbb{S}$ such that

$$J_{\mathbb{E}}e' \cdot J_{\mathbb{S}}s = J_{\mathbb{S}}\tilde{s} \cdot J_{\mathbb{E}}\tilde{e}'.$$

Rearranging the bottom-right triangle, we obtain the following situation (in plain)

$$\begin{array}{ccc}
 Ju & \xrightarrow{b'} & B \\
 \downarrow J_{\mathbb{E}}\tilde{e}' & \dashrightarrow J_{\mathbb{E}}e''' \circlearrowleft & \downarrow f \\
 Jw & \circlearrowleft & J\tilde{w} \\
 \downarrow J_{\mathbb{E}}e' & \dashrightarrow J_{\mathbb{S}}s' \circlearrowleft & \downarrow \\
 Jw & \xrightarrow{a \cdot J_{\mathbb{S}}\tilde{s}} & A
 \end{array}$$

and applying the assumption, we obtain the dashed data with $e''' \in \mathbb{E}'$. In total, we have obtained:

$$\begin{array}{ccc}
 Jv & \xrightarrow{b} & B \\
 \downarrow J_{\mathbb{E}}(e' \cdot e) & \dashrightarrow J_{\mathbb{E}}(e''' \cdot e'') \circlearrowleft & \downarrow f \\
 Jw & \circlearrowleft & J\tilde{w} \\
 \downarrow J_{\mathbb{E}}e' & \dashrightarrow J_{\mathbb{S}}(\tilde{s} \cdot s') \circlearrowleft & \downarrow \\
 Jw & \xrightarrow{a} & A
 \end{array}$$

with $e''' \cdot e'' \in \mathbb{E}$ and $\tilde{s} \cdot s' \in \mathbb{S}$.

D Proofs of Section 5

Proof of Lemma 5.2

Proof. Assume given $f_1, f_2: X \rightarrow (K, +, e)^{(A \times Y)}$, $g: X' \rightarrow X$, and $h: Y \rightarrow Y'$ with $f_1 \sqsubseteq f_2$. Let us prove that $(K, +, e)^{(A \times h)} \cdot f_1 \cdot g \sqsubseteq (K, +, e)^{(A \times h)} \cdot f_2 \cdot g$, that is, for all x', y', a

$$\sum_{\{y|h(y)=y'\}} f_1(g(x'))(a, y) \sqsubseteq \sum_{\{y|h(y)=y'\}} f_2(g(x'))(a, y).$$

If $\{y \mid h(y) = y'\}$ is empty, this means proving $e \sqsubseteq e$, which is true by reflexivity. Otherwise, for all y in this set,

$$f_1(g(x'))(a, y) \sqsubseteq f_2(g(x'))(a, y)$$

by assumption. Then, we conclude by monotonicity of $+$. \square

Proof of Lemma 5.4

Proof. For $x = (x_1, \dots, x_d)$ and $y = (y_1, \dots, y_d)$ in \mathbb{R}^d , we will denote

$$(\min(x_1, y_1), \dots, \min(x_d, y_d))$$

by $x \sqcap y$. Let us prove it for $(G_{\geq 0}, +, 0, \leq)$ first. Define

$$K_x = \{(i, k) \mid x_i = (x_{i,1}, \dots, x_{i,p}) \wedge x_{i,k} > 0\}$$

and k_x the cardinal of K_x . We define K_y and k_y similarly. Let us prove it by induction on $k_x + k_y$.

- if $k_x = 0$, then $u_{i,j} = 0$ works.
- if $k_y = 0$, then $\sum_{i=1}^n x_i = 0$, and since $x_i \geq 0$ for all i , then $x_i = 0$ for all i . This means that $k_x = 0$ and we are back to the previous case.

- if $k_x > 0$ and $k_y > 0$, then let i_0 and k such that $x_{i_0,k} > 0$. Since $\sum_{i=1}^n x_i \leq \sum_{j=1}^m y_j$, then there exists j_0 such that $y_{j_0,k} > 0$. By assumption, $z_0 = x_{i_0,k} \sqcap y_{j_0,k}$, $x'_{i_0} = x_{i_0} - z_0$, and $y'_{j_0} = y_{j_0} - z_0$ belong to $G_{\geq 0}$. For $i \neq i_0$, define $x'_i = x_i$, and for $j \neq j_0$, define $y'_j = y_j$. Then we have $\sum_{i=1}^n x'_i \leq \sum_{j=1}^m y'_j$. By construction, if $x_{i_0,l} = 0$ then $x'_{i_0,l} = 0$, if $y_{j_0,l} = 0$ then $y'_{j_0,l} = 0$, and ($x'_{i_0,k} = 0$ or $y'_{j_0,k} = 0$), so then $k'_x + k'_y < k_x + k_y$. By induction hypothesis, there is a family $(u'_{i,j})$ in $G_{\geq 0}$ such that

$$\text{for all } j, \sum_{i=1}^n u'_{i,j} \leq y'_j \text{ and for all } i, \sum_{j=1}^m u'_{i,j} = x'_i.$$

Define $u_{i,j} = u'_{i,j}$ if $i \neq i_0$ and $j \neq j_0$, $u_{i_0,j_0} + z_0$ otherwise.

For $(G, +, 0, \leq)$, we reduce the problem $\sum_{i=1}^n x_i \leq \sum_{j=1}^m y_j$ in G to a problem in $G_{\geq 0}$ as follows. First, for $x \in G$, define $x^- = -(0 \sqcap x)$ and $x^+ = x + x^-$. By assumption, x^- and x^+ belong to $G_{\geq 0}$ and $x = x^+ - x^-$. Then, defining $x'_i = x_i^+$ if $i \leq n$, y_{i-n}^- otherwise, and $y'_j = y_j^+$ if $j \leq m$, x_{j-m}^- otherwise, we have the following problem in $G_{\geq 0}$

$$\sum_{i=1}^{n+m} x'_i \leq \sum_{j=1}^{m+n} y'_j.$$

By the previous case, we have $(u'_{i,j})$ in $G_{\geq 0}$ such that

- for $i \leq n$, $\sum_{j=1}^{m+n} u_{i,j} = x_i^+$,
- for $i > n$, $\sum_{j=1}^{m+n} u_{i,j} = y_{i-n}^-$,
- for $j \leq m$, $\sum_{i=1}^{n+m} u_{i,j} \leq y_j^+$, and
- for $j > m$, $\sum_{i=1}^{n+m} u_{i,j} \leq x_{j-m}^-$.

Define

$$u_{i,j} = \begin{cases} u'_{i,j} - u'_{n+j,m+i} & \text{if } i > 1 \text{ and } j > 1 \\ u'_{1,j} - u'_{n+j,m+1} \\ \quad + \sum_{j'=1}^m (u'_{n+j',j} - u'_{n+j,j'}) & \text{if } i = 1 \text{ and } j > 1 \\ u'_{i,1} - u'_{n+1,m+i} + \sum_{i'=1}^n u'_{i,m+i'} \\ \quad - \left(x_i^- - \sum_{j'=1}^m u'_{n+j',m+i} \right) & \text{if } i > 1 \text{ and } j = 1 \\ u'_{1,1} - u'_{n+1,m+1} + \sum_{j'=1}^m (u'_{n+j',1} - u'_{n+1,j'}) \\ \quad + \sum_{i'=1}^n u'_{1,m+i'} - \left(x_1^- - \sum_{j'=1}^m u'_{n+j',m+1} \right) & \text{if } i = 1 \text{ and } j = 1 \end{cases}$$

Let us check that this is a solution of the original problem. First, all the $u_{i,j}$ belong to G by assumption. Next, for $i > 1$

$$\sum_{j=1}^m u_{i,j} = \sum_{j=1}^{m+n} u'_{i,j} - x_i^- = x_i^+ - x_i^- = x_i.$$

For $j > 1$

$$\sum_{i=1}^n u_{i,j} = \sum_{i=1}^{n+m} u'_{i,j} - \sum_{j'=1}^{m+n} u'_{n+j,j'} \leq y_j^+ - y_j^- = y_j.$$

For $i = 1$

$$\sum_{j=1}^m u_{1,j} = x_1 + \sum_{j=1}^m \sum_{j'=1}^m u'_{n+j,j'} - \sum_{j=1}^m \sum_{j'=1}^m u'_{n+j',j} = x_1.$$

For $j = 1$

$$\sum_{i=1}^n u_{i,1} \leq y_1 + \sum_{i=1}^n \sum_{i'=1}^n u'_{i,m+i'} - \sum_{i=1}^n \left(x_i^- - \sum_{j'=1}^m u'_{n+j',m+i} \right).$$

We know that for all i , $x_i^- - \sum_{j'=1}^m u'_{n+j',m+i} \geq \sum_{i'=1}^n u'_{i',m+i}$, which means

$$\sum_{i=1}^n u_{i,1} \leq y_1 + \sum_{i=1}^n \sum_{i'=1}^n u'_{i,m+i'} - \sum_{i=1}^n \sum_{i'=1}^n u'_{i',m+i} = y_1.$$

For strictness, let us prove something stronger, namely that if K is a rearrangement monoid and $+$ is strictly monotone (which is the case in \mathbb{R}^d), then it is strict. Indeed, assume $\sum_{i=1}^n x_i = \sum_{j=1}^m y_j$, then we have by rearrangement that

$$\sum_{i=1}^n u_{i,j} \leq y_j \text{ and } \sum_{j=1}^m u_{i,j} = x_i, \text{ so}$$

$$\sum_{j=1}^m \sum_{i=1}^n u_{i,j} = \sum_{i=1}^n x_i = \sum_{j=1}^m y_j.$$

Since the sum is strictly monotone, if $\sum_{i=1}^n u_{i,j} < y_j$ for some j then, $\sum_{j=1}^m \sum_{i=1}^n u_{i,j} < \sum_{j=1}^m y_j$ which is a contradiction. Consequently, $\sum_{i=1}^n u_{i,j} = y_j$ for all j .

Assume given a distributive lattice with bottom element. Define $u_{i,j} = x_i \sqcap y_j$. For all j , $u_{i,j} \leq y_j$, and thus $\bigsqcup_{i=1}^n u_{i,j} \leq y_j$, which is the first condition. By the assumption, for all i

$$x_i \leq \bigsqcup_{j=1}^m y_j.$$

But since L is distributive

$$x_i = x_i \sqcap \left(\bigsqcup_{j=1}^m y_j \right) = \bigsqcup_{j=1}^m (x_i \sqcap y_j) = \bigsqcup_{j=1}^m u_{i,j},$$

which is the second condition.

If furthermore $\sum_{i=1}^n x_i = \sum_{j=1}^m y_j$, then we can prove symmetrically, using distributivity, that $y_j = \bigsqcup_{i=1}^n u_{i,j}$ which proves strictness. Conversely, assume that L

is a rearrangement monoid. We then want to prove that for all x, y_1, y_2 ,

$$x \sqcap (y_1 \sqcup y_2) = (x \sqcap y_1) \sqcup (x \sqcap y_2).$$

Let us start proving it when $x \leq y_1 \sqcup y_2$. In that case, we can use the rearrangement condition and obtain u_1 and u_2 such that $x = u_1 \sqcup u_2$ and $u_j \sqcup y_j$. This implies that $u_j \leq x \sqcap y_j$, so then:

$$x \sqcap (y_1 \sqcup y_2) = x = u_1 \sqcup u_2 \leq (x \sqcap y_1) \sqcup (x \sqcap y_2) \leq x,$$

which implies what we want. Now, let us assume any x, y_1, y_2 . Apply the previous case with $x \sqcap (y_1 \sqcup y_2), y_1, y_2$. We then obtain

$$\begin{aligned} x \sqcap (y_1 \sqcup y_2) &= (x \sqcap (y_1 \sqcup y_2)) \sqcap (y_1 \sqcup y_2) \\ &= (x \sqcap (y_1 \sqcup y_2) \sqcap y_1) \sqcup (x \sqcap (y_1 \sqcup y_2) \sqcap y_2) \\ &= (x \sqcap y_1) \sqcup (x \sqcap y_2). \end{aligned}$$

□

Proof of Theorem 5.6

Proof. Assume that $f: (X, c, i) \rightarrow (Y, d, j)$ is $(\mathbb{E}_K, \mathbb{S}_K)$ -open, X is reachable, and K is positive. Proving that f is a proper homomorphism means proving

$$\begin{array}{ccc} 1 & \xrightarrow{i} & X & \xrightarrow{c} & K^{(A \times X)} \\ & \searrow \circlearrowleft & \downarrow f & \searrow \circlearrowleft & \downarrow K^{(A \times f)} \\ & & Y & \xrightarrow{d} & K^{(A \times Y)} \end{array}$$

The left triangle is given by the fact that f is a lax homomorphism. The right square means that for all $x \in X$, for all $y \in Y$, and for $a \in A$,

$$d(f(x))(a, y) = \sum_{\{x' \mid f(x')=y\}} c(x)(a, x').$$

Since f is a lax homomorphism, we already know that

$$d(f(x))(a, y) \sqsupseteq \sum_{\{x' \mid f(x')=y\}} c(x)(a, x').$$

and since \sqsupseteq is antisymmetric, it is enough to prove the other inequality. If $d(f(x))(a, y) = 0$, then by positivity, we are done. Let us assume given x, y , and a such that $d(f(x))(a, y) > 0$. Since x is reachable, there is a morphism $\alpha: Jw \rightarrow X$ where w is a word $(a_1, k_1), \dots, (a_n, k_n)$ on $A \times (K \setminus \{e\})$ and $\text{end}(\alpha) = x$. Consider $v = (w, a, d(f(x))(a, y))$ so that there is an edge $e: w \rightarrow v$ in \mathbb{E} . There is a lax

homomorphism $\beta: Jv \rightarrow Y$ defined as $\beta(i) = f \cdot \alpha(i)$ for $i \leq n$ and $\beta(n+1, 1) = y$. So we have the following situation (in plain)

$$\begin{array}{ccc}
 Jw & \xrightarrow{\alpha} & (X, c, i) \\
 \downarrow & \dashrightarrow^{J_{\mathbb{E}e}} & \circlearrowright \dashrightarrow^{\alpha'} \\
 J_{\mathbb{E}e} & \circlearrowright & Ju \\
 \downarrow & \dashleftarrow^{J_{\mathbb{S}s}} & \circlearrowright \\
 Jv & \xrightarrow{\beta} & (Y, d, j)
 \end{array}$$

Since f is $(\mathbb{E}_K, \mathbb{S}_K)$ -open, then we obtain that dashed data. In particular, $u = (w, a, w_2)$ with $w_2 = l_1, \dots, l_m$ with $m \geq 1$ and

$$d(f(x))(a, y) = \sum_{i=1}^m l_i.$$

Furthermore, for all i , $\beta \cdot J_{\mathbb{S}s}(n+1, i) = y$. Since α' is a lax morphism, for all i

$$c(x)(a, i) \cong \sum_{\{i' | \alpha'(i')=i\}} l_{i'}.$$

Consequently, since for all i , $f \cdot \alpha'(n+1, i) = \beta \cdot J_{\mathbb{S}s}(n+1, i) = y$

$$d(f(x))(a, y) = \sum_{i=1}^m l_i \cong \sum_{\{x' | \exists i. \alpha'(n+1, i)=x'\}} c(x)(a, x') \cong \sum_{\{x' | f(x')=y\}} c(x)(a, x'),$$

the last inequality being by positivity.

Conversely, assume that f is a proper homomorphism and that K is a rearrangement monoid. Let us prove that f is $(\mathbb{E}_K, \mathbb{S}_K)$ -open. So we start with the following situation (in plain)

$$\begin{array}{ccc}
 Jw & \xrightarrow{\alpha} & (X, c, i) \\
 \downarrow & \dashrightarrow^{J_{\mathbb{E}e}} & \circlearrowright \dashrightarrow^{\alpha'} \\
 J_{\mathbb{E}e} & \circlearrowright & Ju \\
 \downarrow & \dashleftarrow^{J_{\mathbb{S}s}} & \circlearrowright \\
 Jv & \xrightarrow{\beta} & (Y, d, j)
 \end{array}$$

with $w = (a_1, k_1), \dots, (a_n, k_n)$ being a word on $A \times K \setminus \{e\}$ and v being of the form (w, a, w_2) with $w_2 = l_1, \dots, l_m$ a word on $K \setminus \{e\}$. Let us call $x = \text{end}(\alpha)$ and $L = \{\beta(n+1, i) \mid 1 \leq i \leq m\} \subseteq Y$. For y in L , define $V_y = \{i \mid \beta(n+1, i) = y$. Since β is a lax homomorphism,

$$d(f(x))(a, y) \cong \sum_{i \in V_y} l_i.$$

Denote the set $\{x' \in X \mid f(x') = y \wedge c(x)(a, x') \neq e\}$ by X_y . Since f is a proper homomorphism,

$$d(f(x))(a, y) = \sum_{x' \in X_y} c(x)(a, x').$$

Since V_y and X_y are finite subsets of K , then since K is a rearrangement monoid, there is a family $(u_{i,x'})_{i \in V_y, x' \in X_y}$ such that

$$\sum_{x' \in X_y} u_{i,x'} = l_i \text{ and } \sum_{i \in V_y} u_{i,x'} \sqsubseteq c(x)(a, x').$$

Define $U_i = \{x' \in X_y \mid u_{i,x'} \neq e\} = \{x_1, \dots, x_{m_i}\}$ for $i \in V_y$, and m_i its cardinal. Since $l_i \neq e$, U_i is non-empty. Now, define $u = (w, a, w'_2)$ with $w'_2 = l_{1,1}, \dots, l_{1,m}, \dots, l_{m,1}, \dots, l_{m,m_m}$ with $l_{i,j} = u_{i,x_j} \neq e$. There is an edge $e: w \rightarrow u$ in \mathbb{E} . Define $s: \{1, \dots, \sum_{i=1}^m m_i\} \rightarrow \{1, \dots, m\}$ as $s(k) = j$ if $\sum_{i=1}^{j-1} m_i < k \leq \sum_{i=1}^j m_i$. This function is monotone (easy) and surjective (since U_i is non-empty). Now for any j

$$\sum_{\{k \mid s(k + \sum_{i=1}^{j-1} m_i) = j\}} l_{j,k} = \sum_{k=1}^{m_j} l_{j,k} = \sum_{k=1}^{m_j} u_{i,x_k} = l_j,$$

so that s is a morphism from v to u in \mathbb{S} . It is clear that $J_{\mathbb{S}}s \cdot J_{\mathbb{E}}e = J_{\mathbb{E}}e$.

Define $\alpha': Ju \rightarrow X$ as $\alpha'(i) = \alpha(i)$ for $i \leq n$, and $\alpha'(n+1, k) = x_s$ for $\sum_{i=1}^{j-1} m_i < k \leq \sum_{i=1}^j m_i$ and $s = k - \sum_{i=1}^{j-1} m_i$. By construction, α' is a lax morphism and $\alpha' \cdot J_{\mathbb{E}}e$. Then, for k with $s(k) \in V_y$, $f \cdot \alpha'(n+1, k) = f(x')$ with $x' \in X_y$ and

$$\beta \cdot J_{\mathbb{S}}s(n+1, k) = y = f \cdot \alpha'(n+1, k).$$

Furthermore, since $f \cdot \alpha = \beta \cdot J_{\mathbb{E}}e$, we obtain that

$$\beta \cdot J_{\mathbb{S}}s = f \cdot \alpha',$$

which concludes that f is $(\mathbb{E}_K, \mathbb{S}_K)$ -open.

For the last statement, by the previous two points, the only thing to prove is that when there is a span $(X, c, i) \xleftarrow{f} (Z, e, k) \xrightarrow{g} (Y, d, j)$ of $(\mathbb{E}_K, \mathbb{S}_K)$ -open maps, then there is one where (Z, e, k) is reachable.

Define $(\tilde{Z}, \tilde{e}, k)$ with \tilde{Z} being the set of states of Z that are reachable. In particular, k is reachable. Define $\tilde{e}(z_1)(a, z_2) = e(z_1)(a, z_2)$ for z_1, z_2 reachable. It is enough to prove that the injection $\iota: \tilde{Z} \rightarrow Z$ is a proper morphism. Indeed, in that case, by what precedes, ι is $(\mathbb{E}_K, \mathbb{S}_K)$ -open, and since open maps are closed under composition by Lemma 4.4, then

$$(X, c, i) \xleftarrow{f \cdot \iota} (\tilde{Z}, \tilde{e}, k) \xrightarrow{g \cdot \iota} (Y, d, j)$$

is a span of open maps whose tip is reachable. Now, ι is a proper morphism for the following reason. For z reachable and z' non-reachable, $e(z_1)(a, z_2) = 0$

for any a , otherwise z' would be reachable. Consequently, for z reachable, z' non-reachable

$$\begin{aligned} e(\iota(z))(a, z') = 0 &= \sum_{a, z'' \in \emptyset} \tilde{e}(z)(a, z'') \\ &= \sum_{a \in A, z'' \text{ reach.}, \iota(z'')=z} \tilde{e}(z)(a, z'') \\ &= K^{(A \times \iota)} \cdot \tilde{e}(z)(a, z') \end{aligned}$$

and for z' reachable

$$e(\iota(z))(a, z') = e(\iota(z))(a, \iota(z')) = \tilde{e}(z)(a, z') = K^{(A \times \iota)} \cdot \tilde{e}(z)(a, z').$$

□

Proof of Corollary 5.7

Proof. It is enough to prove that strictness of K implies that the weighted functor preserves weak pullbacks. That is, assume given a pullback in **Set**:

$$\begin{array}{ccc} W & \xrightarrow{k} & Y \\ \downarrow h & \circlearrowleft & \downarrow g \\ X & \xrightarrow{f} & Z \end{array}$$

and form this other pullback:

$$\begin{array}{ccc} W' & \xrightarrow{k'} & K^{(A \times Y)} \\ \downarrow h' & \circlearrowleft & \downarrow K^{(A \times g)} \\ K^{(A \times X)} & \xrightarrow{K^{(A \times f)}} & K^{(A \times Z)} \end{array}$$

Since **Set** has the external axiom of choice (that is, that every epi is split), it is enough to prove that the unique morphism $e: K^{(A \times W)} \rightarrow W'$ such that

$$h' \cdot e = h \quad \text{and} \quad k' \cdot e = k$$

is a surjective function. Let us describe more concretely W and W' :

$$W = \{(x, y) \in X \times Y \mid f(x) = g(y)\}$$

$$W' = \left\{ (\mu, \rho) \mid \forall (a, z) \in A \times Z. \sum_{\{x \in X \mid f(x)=z\}} \mu(a, x) = \sum_{\{y \in Y \mid g(y)=z\}} \rho(a, y) \right\}.$$

Now the function e is given by

$$\theta \mapsto \left((a, x) \mapsto \sum_{\{y \in Y \mid (x, y) \in W\}} \theta(a, (x, y)), (a, y) \mapsto \sum_{\{x \in X \mid (x, y) \in W\}} \theta(a, (x, y)) \right)$$

Fix $(\mu, \rho) \in W'$. We want to construct $\theta \in K^{(A \times W)}$ such that $e(\theta) = (\mu, \rho)$. By definition of the the weighted functor, the set

$$\mathcal{N} = \{(a, z) \mid \exists x. f(x) = z \wedge \mu(a, x) \neq e\}$$

is finite. Then, for fixed $(a, z) \in \mathcal{N}$, the sets

$$\mathcal{N}_{a,z,X} = \{x \mid f(x) = z \wedge \mu(a, x) \neq e\}$$

and

$$\mathcal{N}_{a,z,Y} = \{y \mid g(y) = z \wedge \rho(a, y) \neq e\}$$

are finite. Consequently, the condition that $(\mu, \rho) \in W'$ can be reformulated as

$$\sum_{x \in \mathcal{N}_{a,z,X}} \mu(a, x) = \sum_{y \in \mathcal{N}_{a,z,Y}} \rho(a, y).$$

Now, by strictness of K , this means we have $(u_{x,y}^a)_{x \in \mathcal{N}_{a,z,X}, y \in \mathcal{N}_{a,z,Y}}$ family of elements of K such that:

$$\sum_{x \in \mathcal{N}_{a,z,X}} u_{x,y}^a = \rho(a, y) \quad \text{and} \quad \sum_{y \in \mathcal{N}_{a,z,Y}} u_{x,y}^a = \mu(a, x)$$

Define θ as

$$(a, (x, y)) \in W \mapsto \begin{cases} u_{x,y}^a & \text{if } z = f(x), (a, z) \in \mathcal{N}, x \in \mathcal{N}_{a,z,X}, \\ & \text{and } y \in \mathcal{N}_{a,z,Y} \\ 0 & \text{otherwise} \end{cases}$$

By finiteness of \mathcal{N} , $\mathcal{N}_{a,z,X}$, and $\mathcal{N}_{a,z,Y}$, $\theta \in K^{(A \times W)}$, and by construction $e(\theta) = (\mu, \rho)$. \square

Proof of Lemma 5.8

Proof. Let $y \in Y$, so that $d(y) \in (\mathbb{R}_+, +)^{(A \times Y)}$. Let us prove that $d(y) \in \mathcal{D}_{\leq 1}(A \times Y)$. Let us consider the case where f is a lax homomorphism (the other is similar). Then for every $x' \in X$, $\sum_{\{y' \mid f(y')=x'\}} d(y)(y') \leq c(f(y), x')$. This means that:

$$\sum_{y' \in Y} d(y)(y') = \sum_{x' \in X} \sum_{\{y' \mid f(y')=x'\}} d(y)(y') \leq \sum_{x' \in X} c(f(y), x') \leq 1,$$

and $d(y) \in \mathcal{D}_{\leq 1}(A \times Y)$. \square

Proof of Corollary 5.9

Proof. Let us assume that it is $(\mathbb{E}_{\mathcal{D}}, \mathbb{S}_{\mathcal{D}})$ -open, and prove that it is $(\mathbb{E}_{(\mathbb{R}_+, +)}, \mathbb{S}_{(\mathbb{R}_+, +)})$ -open. So we have the following situation in $(\mathbb{R}_+, +)^{(_)}\text{-coalgebras}$

(in plain):

$$\begin{array}{ccc}
 Jv & \xrightarrow{y} & (Y, d, j) \\
 \downarrow & \dashrightarrow^{J_{\mathbb{E}}e'} & \circlearrowleft \\
 J_{\mathbb{E}}e & \circlearrowleft & Ju \\
 \downarrow & \dashrightarrow^{J_{\mathbb{S}}s} & \circlearrowleft \\
 Jw & \xrightarrow{x} & (X, c, i)
 \end{array}$$

By Lemma 5.8 on x and y , this situation is also in $\mathcal{D}_{\leq 1}(A \times _)$ -coalgebras. Then by $(\mathbb{E}_{\mathcal{D}}, \mathbb{S}_{\mathcal{D}})$ -openness, we open the dashed data in $\mathcal{D}_{\leq 1}(A \times _)$ -coalgebras, which are also $(\mathbb{R}_+, +)^{(-)}$ -coalgebras. Conversely, we start with the same plain situation as above, in $\mathcal{D}_{\leq 1}(A \times _)$ -coalgebras this time, so also in $(\mathbb{R}_+, +)^{(-)}$ -coalgebras. By $(\mathbb{E}_{(\mathbb{R}_+, +)}, \mathbb{S}_{(\mathbb{R}_+, +)})$ -openness, we obtain the dashed data in $(\mathbb{R}_+, +)^{(-)}$ -coalgebras. By Lemma 5.8 on y' , the dashed data is also in $\mathcal{D}_{\leq 1}(A \times _)$ -coalgebras.

Finally, if (X, c, i) and (Y, d, j) are $(\mathbb{E}_{\mathcal{D}}, \mathbb{S}_{\mathcal{D}})$ -bisimilar, then they are $(\mathbb{E}_{(\mathbb{R}_+, +)}, \mathbb{S}_{(\mathbb{R}_+, +)})$ -bisimilar by what precedes. For the converse, if we have a span $(X, c, i) \xleftarrow{f} (Z, e, k) \xrightarrow{g} (Y, d, j)$ of $(\mathbb{E}_{(\mathbb{R}_+, +)}, \mathbb{S}_{(\mathbb{R}_+, +)})$ -open morphisms, to conclude by what precedes, we have to make sure that (Z, e, k) is a $\mathcal{D}_{\leq 1}(A \times _)$ -coalgebra, which is again by Lemma 5.8. \square

E Proofs of Section 6

Proof of Lemma 6.5

Proof. Let us prove first that π_X is a morphism. So we start with a transition

$$(X \xleftarrow{x} Jv \xrightarrow{y} Y) \xrightarrow{a}_R (X \xleftarrow{x'} Jw \xrightarrow{y'} Y)$$

of \tilde{R} , and distinguish the two cases.

- Assume $a \neq \tau$, $v = (n_1, a_1, \dots, a_k, n_{k+1})$, $w = (n_0, a_1, \dots, a_k, n_{k+1}, a, 0)$, $x' = x \cdot J_{\mathbb{E}}e_a$, and we want to prove that there is a transition

$$\text{end}(x) = x(n_{k+1}, k+1) \xrightarrow{a} x'(0, k+2) = \text{end}(x').$$

Since $x' = x \cdot J_{\mathbb{E}}e_a$, then $x'(n_{k+1}, k+1) = x(n_{k+1}, k+1)$, and since x' is a morphism, there is a transition as wanted.

- The proof of the other case is the same.

For $r: Jv \rightarrow \tilde{R}$, $\text{end}(r) \in \tilde{R}$, i.e. by definition, $\text{end}(r)$ is a span of the form

$$(X \xleftarrow{x} Jv \xrightarrow{y} Y).$$

The statement then follows from a simple induction on v .

Proof of Lemma 6.6

Proof. Let us start with the following square

$$\begin{array}{ccc}
 Jv & \xrightarrow{-x} & X \\
 \downarrow J_{\mathbb{E}}e & \circlearrowleft & \downarrow f \\
 Jw & \xrightarrow{-y} & Y
 \end{array}$$

for which we construct a new square with a strong morphism $x': Jv' \rightarrow X$. The lifting in the new square will then give rise to a lifting in the above square.

Let us start by constructing $s: v \rightarrow v' \in \mathbb{S}$ and $x': Jv' \rightarrow X$ such that $x = x' \cdot J_{\mathbb{S}}s$. Assume that v is of the form n_1, a_1, \dots, n_{k+1} and v' will be of the form $n'_1, a_1, \dots, n'_{k+1}$ with $n'_i \leq n_i$. Define $f_i: \{0 < \dots < n_i\} \rightarrow \{0 < \dots < n'_i\}$ as

$$f_i(j) = \min \{k \leq j \mid \forall k' \leq k' \leq j. x(k', i) = x(j, i)\}.$$

Then n'_i is the number of elements in the image of f_i , that is, n'_i is the number of τ -transitions from $(0, i)$ to (n_i, i) that are mapped to τ -transitions by x . If we denote by g_i the unique monotone bijection from the image of f_i to $\{0 < \dots < n'_i\}$, then $s_i = g_i \cdot f_i$. It is clear that $s_i: \{0 < \dots < n_i\} \rightarrow \{0 < \dots < n'_i\}$ is monotone and surjective, so $s = (s_1, \dots, s_{k+1})$ is a morphism from v to v' in \mathbb{S} . Now, define $x': Jv' \rightarrow X$ by

$$x'(i, j) = x(g_j^{-1}(i), j).$$

Let us prove that x' is a strong morphism. In Jv' , we have two types of transitions:

- $(i, j) \xrightarrow{-\tau} (i+1, j)$. By construction of v' , it means that we have $(g_j^{-1}(i), j) \xrightarrow{-\tau} (g_j^{-1}(i)+1, j) \xrightarrow{-\tau} \dots \xrightarrow{-\tau} (g_j^{-1}(i)+k, j) \xrightarrow{-\tau} (g_j^{-1}(i+1), j)$ in Jv for some k , and that $x(g_j^{-1}(i)+k', j) = x(g_j^{-1}(i), j)$ for all $k' \leq k$, and $x(g_j^{-1}(i)+k, j) \xrightarrow{-\tau} x(g_j^{-1}(i+1), j)$ in X . This means that $x'(i, j) \xrightarrow{-\tau} x'(i+1, j)$ in X .
- $(n'_j, i) \xrightarrow{-a} (0, j+1)$. Similarly, we can prove that $x'(n'_j, i) \xrightarrow{-a} x'(0, j+1)$ in X .

Now

$$x' \cdot J_{\mathbb{S}}s(i, j) = x'(s_j(i), j) = x(f_j(i), j) = x(i, j).$$

The second step is to obtain a diagram of the following shape

$$\begin{array}{ccccc}
 & & \xrightarrow{x} & & \\
 & & \circlearrowleft & & \\
 Jv & \xrightarrow{-J_{\mathbb{S}}s} & Jv' & \xrightarrow{-x'} & X \\
 \downarrow J_{\mathbb{E}}e & \circlearrowleft & \downarrow J_{\mathbb{E}}e' & \circlearrowleft & \downarrow f \\
 Jw & \xrightarrow{-J_{\mathbb{S}}s'} & Jw' & \xrightarrow{-y'} & Y \\
 & & \circlearrowleft & & \\
 & & \xrightarrow{y} & &
 \end{array}$$

For now, let us assume is e_a (the case $e = e_\tau$ is similar). In that case, w is of the shape $n_1, a_1, \dots, n''_{k+1}, a, n_{k+2}$ with $n_{k+1} \leq n''_{k+1}$. We can then define w' as $n'_1, a_1, \dots, n'_{k+1} + n''_{k+1} - n_{k+1}, a, n_{k+2}$ and $s' = (s'_1, \dots, s'_{k+2})$ as

$$s'_j(i) = \begin{cases} s_j(i) & \text{if } j \leq k \text{ or } (j = k + 1 \text{ and } i \leq n_{k+1}) \\ n'_{k+1} + i - n_{k+1} & \text{if } j = k + 1 \text{ and } i > n_{k+1} \\ i & \text{if } j = k + 2 \end{cases}$$

Then s' is a morphism from w to w' in \mathbb{S} . Now, by construction of v' and w' , there is a unique $e' = e_a : v' \rightarrow w'$ in \mathbb{E} , and it is easy to see that $J_{\mathbb{E}}e' \cdot J_{\mathbb{S}}s = J_{\mathbb{S}}s' \cdot J_{\mathbb{E}}e$. Finally, we want to construct $y' : Jw' \rightarrow Y$ as

$$y'(i, j) = \begin{cases} y(g_j^{-1}(i), j) & \text{if } j \leq k \text{ or } (j = k + 1 \text{ and } i \leq n'_{k+1}) \\ y(n_{k+1} + i - n'_{k+1}, k + 1) & \text{if } j = k + 1 \text{ and } i > n'_{k+1} \\ y(i, k + 2) & \text{if } j = k + 2 \end{cases}$$

Now:

$$y' \cdot J_{\mathbb{E}}e'(i, j) = y(g_j^{-1}(i), j) = f \cdot x(g_j^{-1}(i), j) = f \cdot x'(i, j),$$

which proves the commutativity of the right square. Also, let us prove the bottom commutativity. The only interesting part is when $j \leq k$ or $(j = k + 1$ and $i \leq n_{k+1})$. In that case,

$$y' \cdot J_{\mathbb{S}}s' = y(f_j(i), j) = f \cdot x(f_j(i), j) = f \cdot x(i, j) = y(i, j).$$

Using that x' , f , and y are morphisms, it is easy to prove that y' is also a morphism.

Using the assumption on the right square, since x' is a strong morphism, we obtain the following data

$$\begin{array}{ccccc} Jv' & \xrightarrow{x'} & & \longrightarrow & X \\ & \searrow J_{\mathbb{E}}e'' & \circlearrowright & \nearrow x'' & \downarrow \\ J_{\mathbb{E}}e' & \circlearrowleft & Ju' & & f \\ & \searrow J_{\mathbb{S}}s'' & \circlearrowright & & \downarrow \\ Jw' & \xrightarrow{y'} & & \longrightarrow & Y \end{array}$$

Since $J_{\mathbb{S}}s'' \cdot J_{\mathbb{E}}e'' = J_{\mathbb{E}}e'$ and $e' = e_a$, we know the following:

- $e'' = e_a$,
- u' is of the form $n'_1, a_1, \dots, n'''_{k+1}, a, n'_{k+2}$ with $n'_{k+1} \leq n'_{k+1} + n'''_{k+1} - n_{k+1} \leq n'''_{k+1}$ and $n_{k+2} \leq n'_{k+2}$,
- for all $j \leq k$ or $(j = k + 1$ and $i \leq n'_{k+1})$, $s''_j(i) = i$.

To conclude, we want the following

$$\begin{array}{ccccc} Jv & \xrightarrow{x} & & \longrightarrow & X \\ & \searrow J_{\mathbb{E}}e''' & \circlearrowright & \nearrow x''' & \downarrow \\ J_{\mathbb{E}}e & \circlearrowleft & Ju & & f \\ & \searrow J_{\mathbb{S}}s''' & \circlearrowright & & \downarrow \\ Jw & \xrightarrow{y} & & \longrightarrow & Y \end{array}$$

We first define u as

$$n_1, a_1, \dots, n''_{k+1} + n_{k+1} - n'_{k+1}, a, n'_{k+2}$$

Since $n''_{k+1} + n_{k+1} - n'_{k+1} \geq n_{k+1}$, there is a unique $e''' = e_a : v \rightarrow u$ in \mathbb{E} . Next, s''' is defined as

$$s'''_j(i) = \begin{cases} i & \text{if } j \leq k \text{ or } (j = k + 1 \text{ and } i \leq n_{k+1}) \\ s''_{k+1}(n'_{k+1} + i - n_{k+1}) + & \\ \quad n'_{k+1} - n_{k+1} & \text{if } j = k + 1 \text{ and } i > n_{k+1} \\ s''_{k+2}(i) & \text{if } j = k + 2 \end{cases}$$

which is a morphism from u to w in \mathbb{S} with $J_{\mathbb{S}}s''' \cdot J_{\mathbb{E}}e''' = J_{\mathbb{E}}e$. We now need to construct $x''' : u \rightarrow X$ as

$$x'''(i, j) = \begin{cases} x(i, j) & \text{if } j \leq k \text{ or } (j = k + 1 \text{ and } i \leq n_{k+1}) \\ x''(n'_{k+1} + i - n_{k+1}, k + 1) & \text{if } j = k + 1 \text{ and } i > n_{k+1} \\ x''(i, k + 2) & \text{if } j = k + 2 \end{cases}$$

Since x'' and x are morphisms, x''' is also a morphism. Furthermore, the commutativity of the top triangle is by construction of x''' . Finally, let us prove the commutativity of the bottom right triangle. For $j \leq k$ or $(j = k + 1$ and $i \leq n_{k+1})$

$$f \cdot x'''(i, j) = f \cdot x(i, j) = y(i, j) = y \cdot J_{\mathbb{S}}s'''(i, j).$$

For $j = k + 1$ and $i > n_{k+1}$,

$$\begin{aligned} f \cdot x'''(i, k + 1) &= f \cdot x''(n'_{k+1} + i, k + 1) \\ &= y'' \cdot s''_{k+1}(n'_{k+1} + i - n_{k+1}) \\ &= y(n'_{k+1} + s''_{k+1}(n'_{k+1} + i - n_{k+1}) - n_{k+1}, k + 1) \\ &= y \cdot J_{\mathbb{S}}s'''(i, k + 1). \end{aligned}$$

Finally, for $j = k + 2$

$$f \cdot x'''(i, k + 2) = f \cdot x''(i, k + 2) = y' \cdot J_{\mathbb{S}}s''(i, k + 2) = y \cdot J_{\mathbb{S}}s'''(i, k + 2).$$

□

Proof of Proposition 6.7

Proof. Let us prove that π_X is (\mathbb{E}, \mathbb{S}) -open. We then start with the following square

$$\begin{array}{ccc} Jv & \xrightarrow{r} & \tilde{R} \\ \downarrow J_{\mathbb{E}}e & \circlearrowleft & \downarrow \pi_X \\ Jw & \xrightarrow{x} & X \end{array}$$

and we can assume that r is a strong morphism by Lemma 6.6. There are two cases, depending on whether $e = e_a$ or e_τ . Let us prove the case for e_a , the other

case being similar. So $v = n_1, a_1, \dots, a_k, n_{k+1}$ and $w = n_1, a_1, \dots, a_k, n'_{k+1}, a, n_{k+2}$ with $n_{k+1} \leq n'_{k+1}$. By Lemma 6.5, $\text{end}(r)$ is of the form $(X \xleftarrow{x'} Jv \xrightarrow{y'} Y)$ and $\pi_X \cdot r = x'$. In particular, $(X \xleftarrow{x'} Jv \xrightarrow{y'} Y)$ belongs to R .

In summary, we have the following situation (in plain)

$$\begin{array}{ccccc}
 Jw & \leftarrow \text{-----} & J_{\mathbb{S}S} & \text{-----} & Ju \\
 \downarrow & \swarrow J_{\mathbb{E}}e & \circlearrowleft & \searrow J_{\mathbb{E}}e' & \downarrow \\
 x & \circlearrowleft & Jv & \circlearrowleft & y \\
 \downarrow & \swarrow x' & \in R & \searrow y' & \downarrow \\
 X & & & & Y
 \end{array} \in R$$

and since R is a path bisimulation, we obtain the rest of the diagram (dashed). In particular, $(X \xleftarrow{x \cdot J_{\mathbb{S}S}} Jv \xrightarrow{y} Y)$ belongs to R . To conclude, it is enough to construct a morphism $r': Ju \rightarrow \tilde{R}$ such that

$$\begin{array}{ccc}
 Jv & \xrightarrow{\quad r \quad} & \tilde{R} \\
 \downarrow & \swarrow J_{\mathbb{E}}e' & \circlearrowleft \searrow r' \\
 J_{\mathbb{E}}e & \circlearrowleft & Ju \\
 \downarrow & \swarrow J_{\mathbb{S}S} & \circlearrowleft \searrow \\
 Jw & \xrightarrow{\quad x \quad} & X
 \end{array}$$

By construction, u is of the form $n_1, a_1, \dots, a_k, n''_{k+1}, a, n'_{k+2}$ with $n_{k+1} \leq n'_{k+1} \leq n''_{k+1}$ and $n_{k+2} \leq n'_{k+2}$. For every (i, j) state of Ju , the restriction of Ju up to (i, j) is obtained as $Ju_{i,j}$ where $u_{i,j} = n_1, a_1, \dots, a_{j-1}, i$. Let us denote $e_{i,j}$ the injection of $Ju_{i,j}$ into Ju , which is a strict morphism as a composition of morphisms of the form $J_{\mathbb{E}}e_a$ and $J_{\mathbb{E}}e_\tau$. Then define r' as follows:

$$r'(i, j) = (X \xleftarrow{x \cdot J_{\mathbb{S}S} \cdot e_{i,j}} Ju_{i,j} \xrightarrow{y \cdot e_{i,j}} Y).$$

First, r' is with values in \tilde{R} , because $(X \xleftarrow{x \cdot J_{\mathbb{S}S} \cdot e_{i,j}} Ju_{i,j} \xrightarrow{y \cdot e_{i,j}} Y)$ belongs to R by induction using the backward closure of R being a strong path bisimulation. It is also easy to see that r' is a strict morphism. For the equation $r' \cdot J_{\mathbb{E}}e' = r$, this is a consequence of $x' = x \cdot J_{\mathbb{E}}e = x \cdot J_{\mathbb{S}S} \cdot J_{\mathbb{E}}e'$. Finally, $\pi_X \cdot r' = x \cdot J_{\mathbb{S}S}$ is by construction and Lemma 6.5. \square

Proof of Theorem 6.8

Proof. Assume that R is a stuttering branching bisimulation. Let us prove that \bar{R} is a strong path bisimulation. The initial condition and the backward closure are obvious. Let us then prove the forward closure. We then have the following

situation (in plain)

$$\begin{array}{ccccc}
 Jw & \xleftarrow{\quad J_S s \quad} & & \xrightarrow{\quad} & Ju \\
 \downarrow & \swarrow J_E e & \circlearrowleft & \searrow J_E e' & \downarrow \\
 x' & & \circlearrowleft & & y' \\
 \downarrow & \swarrow x & \in \bar{R} & \searrow y & \downarrow \\
 X & & & & Y
 \end{array} \in \bar{R}$$

and want to construct the dashed data. Let us assume that $e = e_a$, the other case being simpler. So v is of the form n_1, a_1, \dots, n_{k+1} and w of the form $n_1, a_1, \dots, n'_{k+1}, a, n_{k+2}$ with $n_{k+1} \leq n'_{k+1}$. We also have:

- $(x(n_{k+1}, k+1), y(n_{k+1}, k+1)) \in R$,
- $x(n_{k+1}, k+1) = x'(n_{k+1}, k+1)$ (since $x = x' \cdot J_E e$),
- for every $i \geq n_{k+1}$, $(x'(i+1, k+1) = x'(i, k+1)$ or $x'(i, k+1) \xrightarrow{\tau} x'(i+1, k+1)$),
- $x'(n'_{k+1}, k+1) \xrightarrow{a} x'(0, k+2)$,
- for every i , $(x'(i+1, k+2) = x'(i, k+2)$ or $x'(i, k+2) \xrightarrow{\tau} x'(i+1, k+2)$),

the last three points holding since x' is a morphism. Since R is a branching bisimulation with the stuttering, we can build $(y_{i,j})_{n_{k+1} \leq j \leq n'_{k+1}, 1 \leq i \leq m_j}$ states of y such that:

- $y_{1, n_{k+1}} = y(n_{k+1}, k+1)$,
- for all i, j , $(x'(j, k+1), y_{i,j}) \in R$,
- $y_{i,j} = y_{i+1,j}$ or $y_{i,j} \xrightarrow{\tau} y_{i+1,j}$,
- $y_{m_j, j} = y_{1, j+1}$ or $y_{m_j, j} \xrightarrow{\tau} y_{1, j+1}$.

by induction on j .

- **base case** $j = n_{k+1}$: $m_j = 1$, $y_{1, n_{k+1}} = y(n_{k+1}, k+1)$.
- **inductive case**: $(y_{i,j})_{n_{k+1} \leq j \leq s, 1 \leq i \leq m_j}$ has been constructed as wanted. In particular, $(x'(s, k+1), y_{m_s, s}) \in R$. Two cases:
 - $x'(s+1, k+1) = x'(s, k+1)$. In that case, $m_{s+1} = 1$ and $y_{1, s+1} = y_{m_s, s}$.
 - or $x'(s, k+1) \xrightarrow{\tau} x'(s+1, k+1)$. Since R is a stuttering branching bisimulation, two cases again:
 - * $(x'(s+1, k+1), y_{m_s, s})$. In that case, $m_{s+1} = 1$ and $y_{1, s+1} = y_{m_s, s}$.
 - * or $y_{m_s, s} \xrightarrow{\tau} y_1 \xrightarrow{\tau} \dots \xrightarrow{\tau} y_d \xrightarrow{\tau} z_1 \xrightarrow{\tau} \dots \xrightarrow{\tau} z_{d'}$, with $(x'(s, k+1), y_{d'}) \in R$, $(x'(s+1, k+1), z_{d'}) \in R$, and $(x'(s+1, k+1), z_{d''}) \in R$ for all d'' . Then the following family $(y'_{i,j})_{n_{k+1} \leq j \leq s+1, 1 \leq i \leq m'_j}$ works:
 - for $j < s$, $m'_j = m_j$ and $y'_{i,j} = y_{i,j}$,
 - $m'_s = m_s + d$, $y'_{i,s} = y_{i,s}$ for $i \leq m_s$, and $y'_{i,s} = y_{i-m_s}$ otherwise,
 - $m'_{s+1} = d'$ and $y'_{i, s+1} = z_i$.

Repeating this process with the transition $x'(n'_{k+1}, k+1) \xrightarrow{a} x'(0, k+2)$ and the transitions $x'(i, k+2) \xrightarrow{\tau} x'(i+1, k+2)$, we can build two families $(y_{i,j})_{n_{k+1} \leq j \leq n'_{k+1}, 1 \leq i \leq m_j}$ and $(y'_{i,j})_{0 \leq j \leq n_{k+2}, 1 \leq i \leq m'_j}$ such that:

- $y_{1,n_{k+1}} = y(n_{k+1}, k + 1)$,
- for all i, j , $(x'(j, k + 1), y_{i,j}) \in R$,
- $y_{i,j} = y_{i+1,j}$ or $y_{i,j} \xrightarrow{\tau} y_{i+1,j}$,
- $y_{m_j,j} = y_{1,j+1}$ or $y_{m_j,j} \xrightarrow{\tau} y_{1,j+1}$,
- $y_{m_{n'_{k+1}},n'_{k+1}} \xrightarrow{a} y'_{1,0}$,
- for all i, j , $(x'(j, k + 2), y'_{i,j}) \in R$,
- $y'_{i,j} = y'_{i+1,j}$ or $y'_{i,j} \xrightarrow{\tau} y'_{i+1,j}$,
- $y'_{m_j,j} = y'_{1,j+1}$ or $y'_{m_j,j} \xrightarrow{\tau} y'_{1,j+1}$.

Now, define u as $n_1, a_1, \dots, n_{k+1} + \sum_{j=n_{k+1}}^{n'_{k+1}} m_j, a, \sum_{j=0}^{n_{k+2}} m'_j$. By construction, there is a unique $e' = e_a : v \rightarrow u$ in \mathbb{E} . Next $s = (s_1, \dots, s_{k+2})$ is defined as follows:

$$s_j(i) = \begin{cases} i & \text{if } j \leq k \text{ or } (j = k + 1 \text{ and } i \leq n_{k+1}) \\ s & \text{if } j = k + 1 \text{ and} \\ & n_{k+1} + \sum_{j=n_{k+1}}^{s-1} m_j < i \leq n_{k+1} + \sum_{j=n_{k+1}}^s m_j \\ s & \text{if } j = k + 2 \text{ and } \sum_{j=0}^{s-1} m'_j < i \leq \sum_{j=0}^s m'_j. \end{cases}$$

By construction, $s : u \rightarrow w$ is a morphism in \mathbb{S} and $J_{\mathbb{E}}e = J_{\mathbb{S}}s \cdot J_{\mathbb{E}}e'$. Finally, we can define y' as

$$y'(i, j) = \begin{cases} y(i, j) & \text{if } j \leq k \text{ or } (j = k + 1 \text{ and } i \leq n_{k+1}) \\ y_{i-(n_{k+1} + \sum_{j=n_{k+1}}^{s-1} m_j), s} & \text{if } j = k + 1 \text{ and} \\ & n_{k+1} + \sum_{j=n_{k+1}}^{s-1} m_j < i \leq n_{k+1} + \sum_{j=n_{k+1}}^s m_j \\ y'_{i-(\sum_{j=0}^{s-1} m'_j), s} & \text{if } j = k + 2 \text{ and } \sum_{j=0}^{s-1} m'_j < i \leq \sum_{j=0}^s m'_j. \end{cases}$$

By construction, $y' \cdot J_{\mathbb{E}}e' = y$. It only remains to prove that $(X \xleftarrow{x' \cdot J_{\mathbb{S}}s} Ju \xrightarrow{y'} Y) \in \overline{R}$, that is, for all i, j , $(x' \cdot J_{\mathbb{S}}s(i, j), y'(i, j)) \in R$:

- if $j \leq k$ or $(j = k + 1 \text{ and } i \leq n_{k+1})$, $x' \cdot J_{\mathbb{S}}s(i, j) = x(i, j)$, $y'(i, j) = y(i, j)$, and $(x(i, j), y(i, j)) \in R$.
- if $j = k + 1$ and $n_{k+1} + \sum_{j=n_{k+1}}^{s-1} m_j < i \leq n_{k+1} + \sum_{j=n_{k+1}}^s m_j$, $x' \cdot J_{\mathbb{S}}s(i, j) = x'(s, k + 1)$ and $y'(i, j) = y_{i-(n_{k+1} + \sum_{j=n_{k+1}}^{s-1} m_j), s}$, and by construction, $(x'(s, k + 1), y_{i-(n_{k+1} + \sum_{j=n_{k+1}}^{s-1} m_j), s}) \in R$.

- if $j = k + 2$ and $\sum_{j=0}^{s-1} m'_j < i \leq \sum_{j=0}^s m'_j$, $x' \cdot J_{\mathbb{S}}s(i, j) = x'(s, k + 2)$ and $y'(i, j) = y_{i - (\sum_{j=0}^{s-1} m'_j), s}$, and by construction, $(x'(s, k + 2), y_{i - (\sum_{j=0}^{s-1} m'_j), s}) \in R$.

Conversely, assume that R is a strong path bisimulation and let us prove that \check{R} is stuttering branching bisimulation. So we start with $(X \xleftarrow{x} Jv \xrightarrow{y} Y) \in R$ so that $(\text{end}(x), \text{end}(y)) \in \check{R}$. Let us assume that $\text{end}(x) \xrightarrow{a} p$. Let us do the case when $a = \tau$, the other case being similar. Let us assume that v is of the form n_1, a_1, \dots, n_{k+1} . Define $w = n_1, a_1, \dots, n_{k+1} + 1$ so that $e = e_\tau: v \rightarrow w \in \mathbb{E}$. We can define $x': Jw \rightarrow X$ as $x'(i, j) = p$ if $i = n_{k+1} + 1$ and $j = k + 1$; $x'(i, j) = x(i, j)$ otherwise. This is a morphism since x is a morphism and by the assumption on p . Now, we have the following situation (in plain)

$$\begin{array}{ccccc}
 Jw & \xleftarrow{\text{dashed}} & J_{\mathbb{S}}s & \xrightarrow{\text{dashed}} & Ju \\
 \downarrow & \swarrow J_{\mathbb{E}}e & \circlearrowleft & \searrow J_{\mathbb{E}}e' & \downarrow \text{dashed} \\
 x' & \circlearrowleft & Jv & \circlearrowleft & y' \\
 \downarrow & \swarrow x & \in R & \searrow y & \downarrow \\
 X & & & & Y
 \end{array} \in R$$

and since R is path bisimulation, we obtain the dashed data. In particular, $e' = e_\tau$ and u is of the form $n_1, a_1, \dots, n'_{k+1}$ with $n_{k+1} + 1 \leq n'_{k+1}$. Two cases:

- for every $i \geq n_{k+1}$, $y'(i, k + 1) = \text{end}(y)$, then $(\text{end}(x'), \text{end}(y)) \in \check{R}$,
 – or there is a sequence $i_0 = n_{k+1} < i_1 < \dots < i_m < n'_{k+1} = i_{m+1}$ such that

$$\text{end}(y) \xrightarrow{\tau} y'(i_1, k + 1) \xrightarrow{\tau} \dots \xrightarrow{\tau} y'(i_m, k + 1) \xrightarrow{\tau} \text{end}(y')$$

Now, there is necessarily $n_{k+1} \leq i' < n'_{k+1}$ such that

- for all $i \leq i'$, $J_{\mathbb{S}}s(i, k + 1) = n_{k+1}$,
- for all $i > i'$, $J_{\mathbb{S}}s(i, k + 1) = n_{k+1} + 1$.

Let l such that $i_l \leq i' < i_{l+1}$. If we denote n_1, a_1, \dots, i_h by u_h for all h , then $e_\tau: u_h \rightarrow u$ belongs to \mathbb{E} and since $(X \xleftarrow{x' \cdot J_{\mathbb{S}}s} Jv \xrightarrow{y'} Y) \in R$ and R is backward closed, for all h , $(X \xleftarrow{x'_h = x' \cdot J_{\mathbb{S}}s \cdot J_{\mathbb{E}}e_\tau} Ju_h \xrightarrow{y'_h = y' \cdot J_{\mathbb{E}}e_\tau} Y) \in R$. Now:

- for $h \leq l$, $(\text{end}(x), y'(i_h, k + 1)) = (\text{end}(x_h), \text{end}(y_h)) \in \check{R}$,
- for $h > l$, $(\text{end}(x'), y'(i_h, k + 1)) = (\text{end}(x_h), \text{end}(y_h)) \in \check{R}$.

which is what we had to prove. □

Proof of Theorem 6.9

Proof. The constructions and the reasonings are the same as for Theorem 6.8, except that we do not have to care about intermediate states to be bisimilar (stuttering and backward closure). □