

GAPS BETWEEN CONSECUTIVE EIGENVALUES FOR COMPACT METRIC GRAPHS

DAVID BORTHWICK, EVANS M. HARRELL II, AND HAOZHE YU

ABSTRACT. On a compact metric graph, we consider the spectrum of the Laplacian defined with a mix of standard and Dirichlet vertex conditions. A Cheeger-type lower bound on the gap $\lambda_2 - \lambda_1$ is established, with a constant that depends only on the total length of the graph and minimum edge length. We also prove some improvements of known upper bounds for eigenvalue gaps and ratios for metric trees and extensions to certain other types of graphs.

1. INTRODUCTION

Estimating the gaps between eigenvalues is a classical problem in spectral theory, dating back to the upper bounds established for planar domains by Payne, Pólya, and Weinberger [17]. For the Dirichlet problem on a bounded domain $\Omega \subset \mathbb{R}^d$, their “universal bound” on eigenvalue gaps generalizes to

$$\lambda_{n+1} - \lambda_n \leq \frac{4}{nd} \sum_{j=1}^n \lambda_j.$$

For Ω convex, a lower bound $\lambda_2 - \lambda_1 \geq 4\pi^2 / \text{diam}(\Omega)^2$ was proven using gradient methods in Singer-Wong-Yau-Yau [18]. For the Dirichlet problem on a compact manifold with boundary, $\lambda_2 - \lambda_1$ was bounded below in terms of a weighted Cheeger constant in Cheng-Oden [5] and Oden-Sung-Wang [16].

In this paper we study the corresponding problems for the Laplacian on a connected, compact metric graph Γ , with either standard (Kirchhoff-Neumann) or Dirichlet conditions imposed at each vertex. More specifically, the vertex set \mathcal{V} of Γ is subdivided as $\mathcal{V}_N \cup \mathcal{V}_0$, and $-\Delta$ is the self-adjoint operator associated to the quadratic form $\|u'\|^2$ with domain $\{u \in H^1(\Gamma) : u = 0 \text{ on } \mathcal{V}_0\}$. This definition assigns standard vertex conditions to the elements of \mathcal{V}_N . Because Dirichlet conditions have the effect of separating edges, all vertices in \mathcal{V}_0 are assumed to have degree 1. Vertices of degree 2 are effectively invisible under standard vertex conditions, so for \mathcal{V}_N we assume the degree is either 1 or ≥ 3 .

The spectrum of $-\Delta$ is given by

$$0 \leq \lambda_1 < \lambda_2 \leq \lambda_3 \leq \dots$$

Our main concern is the spectral gap $\lambda_2 - \lambda_1$, but first let us review some of the existing eigenvalue estimates in this context.

In the case of fully standard vertex conditions, where $\mathcal{V}_0 = \emptyset$, a variety of interesting eigenvalue estimates are known. Following the notation of [2, 3], we

Date: January 19, 2023.

2010 Mathematics Subject Classification. 34B45, 81Q35.

denote the eigenvalues under standard vertex conditions by λ_j^N . In terms of the total length $L := |\Gamma|$, the standard eigenvalues satisfy

$$(1.1) \quad \lambda_k^N \geq \frac{k^2 \pi^2}{4L^2}$$

for all $k \geq 2$. This lower bound was proven for $k = 2$ by Nicaise [15, Thm. 3.1], for $k \geq 2$ by Friedlander [7, Thm. 1], and independently by Kurasov-Naboko [14] for k even. The bound is sharp, and equality for some k implies that Γ is a segment if $k = 2$ and an equilateral k -star if $k > 2$.

Nicaise also established a Cheeger-type lower bound,

$$(1.2) \quad \lambda_2^N \geq \frac{1}{4} h(\Gamma)^2,$$

where the metric graph version of the Cheeger constant is defined by

$$(1.3) \quad h(\Gamma) := \inf \frac{\#S}{\min(|Y_1|, |Y_2|)},$$

with the infimum taken over finite sets S such that $\Gamma \setminus S$ is a disjoint union of non-empty open sets Y_1 and Y_2 . The Cheeger constant satisfies a trivial lower bound

$$(1.4) \quad h(\Gamma) \geq \frac{2}{L},$$

but this gives a much weaker result than (1.1). If $D := \text{diam}(\Gamma)$ satisfies $D \leq L/2$, then Kennedy et al. [11, Thm. 7.2] showed that

$$\lambda_2^N \geq \frac{1}{2DL}.$$

This improves on (1.1) for graphs of sufficiently small diameter.

Upper bounds for λ_2^N can be proven by a combination of min-max and surgery principles. For example, if the number of edges of Γ is $E \geq 2$, then [11, Thm. 4.2] gives the sharp upper bound

$$\lambda_2^N \leq \frac{\pi^2 E^2}{L^2}.$$

Estimates based on other geometric properties of Γ can be found, for example, in [1, 2, 3, 4, 11, 13].

Returning to the general case, if \mathcal{V}_0 is not empty then the arguments leading to (1.1) imply a lower bound

$$(1.5) \quad \lambda_k \geq \frac{k^2 \pi^2}{4L^2}.$$

This was proven for $k = 1$ in [15, Thm. 3.1] and is implicit in the proof of [7, Thm. 1] for $k \geq 1$. Berkolaiko-Kennedy-Kurasov-Mugnolo [1, Thm. 4.7] showed that, if Γ is not a cycle the bound can be improved for $k \geq E - V_0 + 1$ to

$$\lambda_k \geq \left(k - \frac{1}{2}(E - V_0 + 1)\right)^2 \frac{\pi^2}{L^2},$$

where $V_0 := \#\mathcal{V}_0$. The argument for the Cheeger estimate (1.2) from [15, Thm. 3.2] also yields

$$\lambda_1 \geq \frac{1}{4} h(\Gamma)^2$$

if $\mathcal{V}_0 \neq \emptyset$.

As for upper bounds, a simple test function argument yields

$$(1.6) \quad \lambda_1 \leq \frac{\pi^2}{\ell_{\max}^2},$$

where ℓ_{\max} is the maximum edge length. Similarly, Berkolaiko-Kennedy-Kurasov-Mugnolo [3, Thm. 1.3] proved an upper bound

$$(1.7) \quad \lambda_1 \leq \frac{\pi^2}{\text{girth}(\Gamma)^2},$$

where the *girth* is defined as the minimum cycle length of the graph formed from Γ by identifying all Dirichlet vertices. For higher eigenvalues, the same authors also proved upper bounds in terms of the Betti number $\beta = E - V + 1$, which counts the number of independent cycles in Γ . In our notation, this yields [1, Thm. 4.9]

$$(1.8) \quad \lambda_k \leq \left(k - \frac{1}{2} + \frac{3}{2}E - V_N - \frac{1}{2}V_0\right)^2 \frac{\pi^2}{L^2}.$$

With this context established, we turn now to the question of estimates on $\lambda_2 - \lambda_1$ when Γ has at least one Dirichlet vertex. As far as we are aware, there are no lower bounds in the literature for this case. The combination of (1.5) and (1.7) gives no bound on the gap, because the girth is obviously bounded by the total length.

By adapting the argument of Cheng and Oden [5, Prop. 1.3], we establish in §2 a lower bound for $\lambda_2 - \lambda_1$ in terms of a weighted Cheeger constant, with weight given by the first eigenfunction ϕ_1 . To make use of this bound, we prove a Harnack inequality for ϕ_1 in §3, with a constant depending only L and ℓ_0 , the minimum edge length of Γ .

The Harnack inequality (Proposition 3.1) implies an envelope estimate for ϕ_1 , which is perhaps of independent interest. Define the function

$$(1.9) \quad \Upsilon(q) := \begin{cases} 1, & \text{dist}(q, \mathcal{V}_0) \geq \ell_0/2, \\ \sin(\frac{\pi}{\ell_0} \text{dist}(q, \mathcal{V}_0)), & \text{dist}(q, \mathcal{V}_0) < \ell_0/2. \end{cases}$$

Theorem 1.1. *Let Γ be a compact metric graph with total length L and minimum edge length ℓ_0 . Suppose $\mathcal{V}_0 \neq \emptyset$ and the first eigenfunction is normalized so that $\phi_1 \geq 0$ and $\|\phi_1\| = 1$. There exists a constant $c_1(L, \ell_0) > 0$ such that*

$$c_1(L, \ell_0)\Upsilon \leq \phi_1 \leq \sqrt{\frac{2}{\ell_0}}\Upsilon.$$

The combination of the weighted Cheeger inequality and the estimates of ϕ_1 lead to the following result, whose proof is given in §4:

Theorem 1.2. *Let Γ be a metric graph with $\mathcal{V}_0 \neq \emptyset$. There exists a constant $C(L, \ell_0) > 0$, depending only on the total length L and minimum edge length ℓ_0 , such that*

$$(1.10) \quad \lambda_2 - \lambda_1 \geq C(L, \ell_0).$$

For fully standard boundary conditions, $\lambda_1 = 0$ and the corresponding gap estimate is (1.1), depending on L alone. The following cases show that dependence on both L and ℓ_0 is required for graphs with at least one Dirichlet vertex.

Example 1.3. Let Γ be a star graph with four edges, two of length 1 and two of length $a < 1$, as shown in Figure 1. Dirichlet conditions are imposed at the four




FIGURE 1. A 4-star graph as in Example 1.3.




FIGURE 2. A comparison of the first two eigenfunctions for the 4-star graph of Example 1.3.




FIGURE 3. A star graph as in Example 1.4.

external vertices. Note that the total length $L = 2(1 + a) < 4$, $\ell_0 = a$, and the diameter is a constant $D = 2$.

Define a coordinate x so that the longer edges are parametrized by $x \in [0, 1]$ and the shorter edges by $x \in [1, 1+a]$. In terms of this coordinate, the first eigenfunction can be written as

$$(1.11) \quad \phi_1(x) = \sin\left(\frac{\pi x}{1+a}\right),$$

yielding

$$\lambda_1 = \frac{\pi^2}{(1+a)^2}.$$

The second eigenfunction vanishes on the short edges and is proportional to $\sin(\pi x)$ on the long edges, so that $\lambda_2 = \pi^2$. The spectral gap is thus

$$\lambda_2 - \lambda_1 = \pi^2 \left(1 - \frac{1}{(1+a)^2}\right),$$

which is $\sim 2\pi^2 a$ as $a \rightarrow 0$.

Example 1.4. Suppose Γ is a star graph with one edge of length 2 and k edges of length 1, with Dirichlet vertex conditions on all external vertices, as shown in Figure 3. Here $L = 2 + k$, $\ell_0 = 1$, and $D = 3$.

For the two lowest eigenvalues, it suffices to consider eigenfunctions which do not vanish at the central vertex, and thus must take the same values on each smaller edge. We can thus use the linear coordinate $x \in [0, 3]$, with the interior vertex




FIGURE 4. Comparison of the first two eigenfunctions from Example 1.4.

located at $x = 2$. Suppose an eigenfunction is given by

$$(1.12) \quad \phi(x) = \begin{cases} \sin(\sigma x), & x \in [0, 2], \\ c \sin \sigma(3 - x), & x \in [2, 3]. \end{cases}$$

The continuity and vertex conditions give $\sin(2\sigma) = c \sin \sigma$ and $\cos(2\sigma) = kc \cos \sigma$. With double-angle formulas, this reduces to $(\tan \sigma)^2 = 2k + 1$, yielding

$$\sigma \in \arctan \sqrt{2k + 1} + \pi \mathbb{Z}.$$

On the other hand, eigenfunctions which vanish at the central vertex have eigenvalues in $(\pi \mathbb{N})^2$. The first two eigenvalues are thus given by

$$\lambda_1 = \left(\arctan \sqrt{2k + 1} \right)^2, \quad \lambda_2 = \left(\pi - \arctan \sqrt{2k + 1} \right)^2,$$

with values just below and above $\pi^2/4$. For large k , we have $\lambda_2 - \lambda_1 \sim 2\pi/\sqrt{k}$. In this example, the quantities $\ell_0 = 1$ and $D = 3$ are fixed, while $L = k + 2$.

Now let us turn to the issue of upper bounds on the gaps or ratios between consecutive eigenvalues. For a general graph Γ with $\mathcal{V}_0 \neq \emptyset$, we have at least a bound deduced from (1.5) and (1.8):

$$(1.13) \quad \frac{\lambda_{n+1}}{\lambda_n} \leq (2n + 1 + 3E - 2V_N - V_0)^2,$$

provided Γ is not a cycle graph. The ratio λ_2/λ_1 can indeed be unbounded in general, as is demonstrated, for instance, in this paraphrase of [6, Example 1.2]:

Example 1.5. Consider an equilateral “balloon” graph, consisting of a pumpkin graph with k edges of length 1, with a pendant edge attached at one vertex, as illustrated in Figure 5. Dirichlet conditions are imposed at the endpoint of the free edge. The lowest two eigenvalues are equal on all edges of the pumpkin, so we can use the linear parameter $x \in [0, 2]$, with the range $[0, 1]$ corresponding to the free edge. For an eigenfunction of the form

$$\phi(x) = \begin{cases} \sin(\sigma x), & x \in [0, 1], \\ c \cos \sigma(2 - x), & x \in [1, 2], \end{cases}$$

the continuity and vertex conditions give $(\tan \sigma)^2 = 1/k$. The lowest two solutions give

$$\lambda_1 = \arctan(1/\sqrt{k})^2, \quad \lambda_2 = (\pi - \arctan(1/\sqrt{k}))^2,$$

and hence

$$\frac{\lambda_2}{\lambda_1} = \left(\frac{\pi}{\arctan(1/\sqrt{k})} - 1 \right)^2.$$

This gives $\lambda_2/\lambda_1 = 25$ for $k = 3$, and $\lambda_2/\lambda_1 \sim \pi^2 k$ as $k \rightarrow \infty$.




FIGURE 5. An equilateral balloon graph with $k = 6$, for which $\lambda_2/\lambda_1 \approx 50$.




FIGURE 6. Comparison of the first two eigenfunctions from Example 1.5.

In addition to showing that λ_2/λ_1 can be arbitrarily large, Example 1.5 shows the need for the cycle restriction in (1.13), as the balloon is a cycle graph. The right side of (1.13) reduces to $(3k + 1)^2$ for this case, and so the inequality fails as the ratio approaches π^2 for large k .

For a metric tree Γ with Dirichlet vertices on the external vertices, Nicaise [15, Thm. 4.3] proved

$$(1.14) \quad \frac{\lambda_2}{\lambda_1} \leq 2 + \sqrt{5},$$

along with a bound analogous to Payne-Pólya-Weinberger [17]: for $n \geq 2$,

$$(1.15) \quad \lambda_{n+1} - \lambda_n \leq \frac{4}{n} \sum_{j=1}^n \lambda_j.$$

The bound (1.15) is derived using the approach that Hile-Protter [10] developed for Dirichlet eigenvalues of domains in \mathbb{R}^n . As in that source, the bound (1.15) is the corollary of a more complicated but sharper estimate. Again, for a metric tree Γ with Dirichlet vertices on the external vertices,

$$(1.16) \quad \lambda_{n+1} \leq \sigma,$$

where σ denotes the unique root in $[\lambda_n, \infty)$ of the equation

$$(1.17) \quad \sum_{j=1}^n \frac{\lambda_j}{\sigma - \lambda_j} = \frac{n}{4}.$$

The bound (1.15) is deduced from (1.16) by using $\lambda_{n+1} - \lambda_n \leq \sigma - \lambda_j$ in (1.17).

In §5, we will show that the estimation techniques of Harrell-Stubbe [9], based on commutator formulas, can be applied to metric trees by combining ideas from Nicaise [15] and Demirel-Harrell [6]. This gives a more general bound which includes (1.16) as a special case.

In the final section, §6, we consider extensions of the upper bounds on the ratio λ_2/λ_1 to metric graphs which are not trees. In particular, we will see that the bound $\lambda_2/\lambda_1 \leq 5$ continues to hold for graphs consisting of trees to which pendant




FIGURE 7. A Cheeger cut dividing the graph into two components.

graphs are attached, provided the pendants are sufficiently small and include at least one Dirichlet vertex.

2. WEIGHTED CHEEGER CONSTANT

As in the introduction, Γ denotes a compact, connected metric graph, and the Laplacian $-\Delta$ is defined with vertex conditions according to the decomposition $\mathcal{V} = \mathcal{V}_N \cup \mathcal{V}_0$. The eigenvalues $\{\lambda_j\}$ are written in ascending order, starting from $\lambda_1 \geq 0$. We may assume that the corresponding eigenfunctions ϕ_j are real and that $\phi_1 > 0$ away from \mathcal{V}_0 .

In this section we will establish a lower bound for $\lambda_2 - \lambda_1$ by adapting the weighted Cheeger constant from Cheng and Oden [5] to the metric graph setting. Given a continuous function $\phi : \Gamma \rightarrow \mathbb{R}$, we define

$$(2.1) \quad h_\phi(\Gamma) := \inf \frac{\sum_S \phi^2}{\min(\int_{Y_1} \phi^2, \int_{Y_2} \phi^2)},$$

where $S \subset \Gamma$ is a finite subset such that $\Gamma \setminus S$ is a disjoint union of non-empty open sets Y_1 and Y_2 . This decomposition, called a Cheeger cut, is illustrated in Figure 7. Note that (2.1) agrees with (1.3) when ϕ is constant.

Theorem 2.1. *In the setting described above,*

$$\lambda_2 - \lambda_1 \geq \frac{1}{4} h_{\phi_1}(\Gamma)^2.$$

Note that this result reduces to (1.2) in the case $\mathcal{V}_0 = \emptyset$. The key step in the proof is the following estimate.

Lemma 2.2. *Let f be a piecewise C^1 function on Γ , and for $\phi : \Gamma \rightarrow \mathbb{R}$ continuous, suppose that*

$$(2.2) \quad \int_{\Gamma} f \phi^2 = 0.$$

Then

$$\|f' \phi\| \geq \frac{1}{2} h_\phi(\Gamma) \|f \phi\|.$$

Proof. Set

$$k := \sup \left\{ t : \int_{\{f \leq t\}} \phi^2 \leq \frac{1}{2} \int_{\Gamma} \phi^2 \right\},$$

and define

$$g_+ = (f - k)^2 \chi_{\{f > k\}}.$$

The level set $\{g_+ = t\}$ is finite for almost every $t > 0$, and the co-area formula gives

$$(2.3) \quad \int_{\Gamma} |g'_+| \phi^2 = \int_0^\infty \left(\sum_{\{g_+ = t\}} \phi^2 \right) dt.$$

Note also that, by the choice of k ,

$$\int_{\{g_+ > t\}} \phi^2 \leq \int_{\{g_+ < t\}} \phi^2$$

for all $t > 0$. Thus, for t such that $\{g_+ = t\}$ is finite,

$$\sum_{\{g_+ = t\}} \phi^2 \geq h_\phi(\Gamma) \int_{\{g_+ > t\}} \phi^2.$$

Plugging this back into (2.3) gives

$$(2.4) \quad \begin{aligned} \int_\Gamma |g'_+| \phi^2 &\geq h_\phi(\Gamma) \int_0^\infty \left(\int_{\{g_+ > t\}} \phi^2 \right) dt \\ &= h_\phi(\Gamma) \int_\Gamma g_+ \phi^2. \end{aligned}$$

Similarly, for $g_- := (f - k)^2 \chi_{\{f < k\}}$ we obtain

$$(2.5) \quad \int_\Gamma |g'_-| \phi^2 \geq h_\phi(\Gamma) \int_\Gamma g_- \phi^2.$$

Since $(f - k)^2 = g_+ + g_-$, adding (2.4) and (2.5) gives

$$(2.6) \quad \int_\Gamma (|g'_+| + |g'_-|) \phi^2 \geq h_\phi(\Gamma) \int_\Gamma (f - k)^2 \phi^2.$$

After computing

$$|g'_+| + |g'_-| = 2|f - k| |f'|,$$

we can apply Cauchy-Schwarz to (2.6) to obtain

$$\begin{aligned} h_\phi(\Gamma) \|(f - k)\phi\|^2 &\leq 2 \int_\Gamma |f - k| |f'| \phi^2 \\ &\leq 2 \|(f - k)\phi\| \|f'\phi\|. \end{aligned}$$

Hence

$$\|f'\phi\| \geq \frac{1}{2} h_\phi(\Gamma) \|(f - k)\phi\|.$$

The final step is to note that the hypothesis (2.2) implies that

$$\|(f - k)\phi\|^2 = \|f\phi\|^2 + k^2 \|\phi\|^2 \geq \|f\phi\|^2.$$

□

To deduce the weighted Cheeger bound from Lemma 2.2, we call upon a variational principle from the folklore. In our notation,

$$(2.7) \quad \lambda_2 - \lambda_1 = \inf \left\{ \frac{\|f'\phi_1\|^2}{\|f\phi_1\|^2} : \int_\Gamma f\phi_1^2 = 0 \right\},$$

for f a piecewise C^1 function on Γ , with the minimum attained at $f = \phi_2/\phi_1$. This formula is quite general and has been proven independently in a variety of contexts. The earliest reference that we are aware of is Thompson-Kac [19, eq. (3.10)]. The short proof for metric graphs is included in the argument below.

Proof of Theorem 2.1. Let $f = \phi_2/\phi_1$, which is smooth away from the vertices and satisfies

$$\int_{\Gamma} f \phi_1^2 = \langle \phi_2, \phi_1 \rangle = 0.$$

Assuming the eigenfunctions are normalized,

$$\begin{aligned} \lambda_2 &= \|\phi_2'\|^2 \\ &= \|f'\phi_1 + f\phi_1'\|^2 \\ (2.8) \quad &= \|f'\phi_1\|^2 + \|f\phi_1'\|^2 + \frac{1}{2} \int_{\Gamma} (f^2)'(\phi_1^2)'. \end{aligned}$$

Since both f^2 and ϕ_1^2 satisfy the vertex conditions, we can integrate by parts to obtain

$$\begin{aligned} \frac{1}{2} \int_{\Gamma} (f^2)'(\phi_1^2)' &= -\frac{1}{2} \int_{\Gamma} f^2 \Delta(\phi_1^2) \\ &= \int_{\Gamma} (\lambda_1 \phi_1^2 - f^2(\phi_1')^2) \\ &= \lambda_1 - \|f\phi_1'\|^2. \end{aligned}$$

This simplifies (2.8) to

$$(2.9) \quad \lambda_2 - \lambda_1 = \|f'\phi_1\|^2,$$

which is the minimum case of (2.7). The result then follows from Lemma 2.2, since $f\phi_1 = \phi_2$. \square

3. ESTIMATES OF THE FIRST EIGENFUNCTION

To make use of Theorem 2.1 in the case $\mathcal{V}_0 \neq \emptyset$, we need some control over the range of the first eigenfunction ϕ_1 . In particular, we will establish a lower bound on a subset that excludes the Dirichlet vertices.

For each vertex $v_j \in \mathcal{V}_0$, parametrize the edge incident to v_j by $x_j \in [0, \ell_j]$, with $x_j = 0$ at v_j . Within this edge define the interval,

$$(3.1) \quad I_j := \sup\{x_j \in [0, \ell_j] : \phi_1'(x_j) > 0\},$$

which includes the full interior of the edge unless ϕ_1 has a local maximum. Then let

$$(3.2) \quad \Gamma_1 := \Gamma \setminus (\cup_{v_j \in \mathcal{V}_0} I_j).$$

If ϕ_1 has no local maxima in external edges, then Γ_1 is the subgraph obtained by trimming from Γ all edges incident on \mathcal{V}_0 , as shown in Figure 8. If ϕ_1 does have local maxima within an outer edge, then we would need to first add artificial vertices at these maxima before trimming.

Our goal in this section is to prove an explicit Harnack inequality for the restriction of ϕ_1 to Γ_1 . A general Harnack inequality for eigenfunctions on quantum graphs was proven in Harrell-Maltsev [8, Thm. 2.2]. The difference here is that we will produce a constant that depends only on L and ℓ_0 .

By construction, the maximum value of the first eigenfunction,

$$(3.3) \quad M_1 := \max_{\Gamma} \phi_1,$$




FIGURE 8. The subgraph Γ_1 created by trimming external edges.

occurs at a point in Γ_1 , and the minimum over Γ_1 ,

$$(3.4) \quad m_1 := \min_{\Gamma_1} \phi_1,$$

is strictly positive. Note that it is possible for Γ_1 to consist of a single point. In this case, Γ has $k = \#\mathcal{V}_0$ edges, each connecting v to a Dirichlet vertex. The continuity condition at v implies that all of these edges have equal length. Hence Γ is a half-Dirichlet interval if $k = 1$, a full Dirichlet interval for $k = 2$, and an equilateral star graph for $k \geq 3$.

If Γ_1 contains more than one point, then m_1 occurs at a vertex $v \in \mathcal{V}_N$, by the concavity of ϕ_1 on edges. The standard vertex condition implies that the outgoing derivative of ϕ_1 at v is ≤ 0 on at least one incident edge e in Γ . By concavity, ϕ_1 is strictly decreasing in the interior of e , and so e must lie outside Γ_1 . Therefore, the minimum m_1 is achieved at a vertex in \mathcal{V}_N which is adjacent to a vertex in \mathcal{V}_0 .

The fact that ϕ_1 is strictly decreasing on an edge incident on a Dirichlet vertex implies that $\sigma\ell \leq \pi/2$, where ℓ is the length of this edge and $\sigma := \sqrt{\lambda_1}$. This gives an upper bound


$$(3.5) \quad \lambda_1 \leq \frac{\pi^2}{4\ell_0^2},$$

provided Γ_1 contains more than one point. For the exceptional cases where Γ_1 contains a single point, we see explicitly that $\lambda_1 = \pi^2/\ell_0^2$ for the full Dirichlet interval and $\lambda_1 = \pi^2/4\ell_0^2$ for all other cases. Hence the Dirichlet interval is the only exception to the bound (3.5).

The estimate (3.5) is implied by the girth estimate (1.7) if \mathcal{V}_0 contains more than one vertex, or if Γ contains a cycle with at least two edges. But it does cover a few additional cases, such as a tree or tadpole graph with a single Dirichlet vertex.

Proposition 3.1. *Assuming that Γ_1 contains more than one point, there exists a constant $c(L, \ell_0) > 0$, depending only on $L = |\Gamma|$ and ℓ_0 the minimum edge length, such that*

$$(3.6) \quad \frac{m_1}{M_1} \geq c(L, \ell_0).$$

FIGURE 9. The parametrization (3.7) with phases α_j, β_j .

Proof. Let $\lambda_1 = \sigma^2$ for $\sigma > 0$. On each edge e_j of Γ_1 , ϕ_1 is given by a positive arc of the sine function. Thus, we can choose phases $\alpha_j, \beta_j \in (0, \pi)$ such that

$$(3.7) \quad \phi_1|_{e_j}(x) = A_j \sin(\sigma x),$$

for a parametrization of e_j by

$$x \in \left[\frac{\alpha_j}{\sigma}, \frac{(\pi - \beta_j)}{\sigma} \right].$$

These phases are illustrated in Figure 9. By switching the orientation if necessary, we can assume that $\alpha_j \leq \beta_j$.

The change in $\log \phi_1$ across e_j is given by

$$\log \phi_1 \Big|_{\alpha_j/\sigma}^{(\pi-\beta_j)/\sigma} = \log \frac{\sin \beta_j}{\sin \alpha_j}.$$

If we define

$$\delta_0 := \min_{e_j \subset \Gamma_1} \alpha_j,$$

then for each edge e_j in Γ_j ,

$$(3.8) \quad \left| \log \phi_1 \Big|_{\alpha_j/\sigma}^{(\pi-\beta_j)/\sigma} \right| \leq -\log \sin \delta_0.$$

Note that $\delta_0 < \pi/2$, since $\alpha_j + \beta_j < \pi$ on each edge.

Suppose that the vertices of Γ_1 where the minimum and maximum of ϕ_1 occur can be joined by a path with at most q edges. The estimate (3.8) then gives

$$(3.9) \quad \log \frac{M_1}{m_1} \leq -q \log \sin \delta_0,$$

and the problem is now reduced to finding a lower bound for δ_0 .

Suppose that the minimal phase δ_0 occurs at a vertex v_0 of Γ_1 which is the $x = 0$ endpoint of an edge e_j parametrized by $x \in [0, \ell_j]$. In this parametrization,

$$\phi_1(x) = A \sin(\sigma x + \delta_0).$$

Because $\delta_0 < \pi/2$, ϕ_1 does not have a local maximum at v_0 , which implies that $v_0 \in \mathcal{V}_N$.

The outward derivative into e_j from v_0 is given by

$$\phi_1'(0^+) = A \cos \delta_0.$$

Let e_k be the edge of Γ incident to v_0 for which the inward-pointing derivative at v_0 is maximal. We can parametrize e_k by $x \in [-\ell_k, 0]$, and then continue the eigenfunction as

$$(3.10) \quad \phi_1(x) = B \sin(\alpha_k - \sigma x), \quad \text{for } x \leq 0,$$




FIGURE 10. Bounds on the first eigenfunction for a star graph as in Example 1.4, with $k = 100$.

for some phase $\alpha_k \in (0, \pi)$. By continuity, the amplitudes satisfy

$$(3.11) \quad A \sin \delta_0 = B \sin \alpha_k.$$

Because the vertex condition at v_0 is standard, and by the choice of e_k , we can estimate

$$\phi_1'(0^+) \leq (d_0 - 1)\phi_1'(0^-),$$

where d_0 is the degree of v_0 . This gives

$$A \cos \delta_0 \leq (d_0 - 1)B \cos \alpha_k.$$

Combining this with (3.11) yields a lower bound

$$(3.12) \quad \tan \delta_0 \geq \frac{1}{d-1} \tan \alpha_k.$$

To obtain a lower bound on the phase α_k from (3.10), note that since ϕ_1 cannot vanish in the interior of e_k , $\alpha_k \geq \sigma \ell_k$. From (3.12), we thus obtain

$$(3.13) \quad \tan \delta_0 \geq \frac{1}{d_0 - 1} \tan(\sigma \ell_k).$$

If we denote the right side of (3.13) by b , then this gives

$$\sin \delta_0 \geq \frac{b}{\sqrt{1+b^2}},$$

and (3.9) implies that

$$(3.14) \quad m_1 \geq \left(\frac{b}{\sqrt{1+b^2}} \right)^q M_1.$$

To complete the proof, note that $\ell_k \geq \ell_0$, both q and d_0 are bounded by L/ℓ_0 and σ was bounded below by $\pi/2L$ in (1.5). \square

Example 1.4 gives an interesting test case for the sharpness of the eigenfunction estimate. The first eigenfunction is given by (1.12) with $\sigma = \arctan \sqrt{2k+1}$. In this normalization $\max \phi_1 = 1$, which occurs at $x = \pi/2\sigma$. The subgraph Γ_1 consists of the segment $[\pi/2\sigma, 2]$, and the minimum of ϕ_1 on this segment occurs at $x = 2$. Thus

$$\begin{aligned} \frac{m_1}{M_1} &= \sin(2\sigma) \\ &= \frac{\sqrt{2k+1}}{k}. \end{aligned}$$

For this graph, the right side of (3.13) is

$$\frac{1}{k} \tan \sigma = \frac{\sqrt{2k+1}}{k},$$

and $q = 1$. Hence, the lower bound (3.14) is asymptotically sharp as $k \rightarrow \infty$ for this case.

We can extend the bound (3.6) beyond Γ_1 . On each interval $I_j = [0, \ell_j)$ from (3.1), the eigenfunction takes the form $\phi_1(x_j) = A_j \sin(\sigma x_j)$. The points $x_j = \ell_j$ lie in Γ_1 , so

$$A_j \sin(\sigma \ell_j) \geq m_1.$$

Therefore,

$$(3.15) \quad \phi_1(x_j) \geq \frac{m_1}{\sin(\sigma \ell_j)} \sin(\sigma x_j)$$

on each interval I_j .

To prove Theorem 1.1, we first note that in the notation used above, the variable x_j used to parametrize I_j is equal to $\text{dist}(\cdot, \mathcal{V}_0)$ for $x_j \leq \ell_0/2$. The envelope function Υ defined in (1.9) is equal to 1 on Γ_1 and on the outer edges satisfies

$$(3.16) \quad \Upsilon|_{I_j}(x_j) = \begin{cases} 1, & x_j \geq \ell_0/2, \\ \sin(\frac{\pi}{\ell_0} x_j), & x_j < \ell_0/2. \end{cases}$$

Proof of Theorem 1.1. We can assume that Γ_1 consists of more than one point, since ϕ_1 is easily computed explicitly in the exceptional cases.

From (3.15), (3.16), and the fact that $\sigma > \pi/2L$, we can immediately deduce the inequality,

$$(3.17) \quad \phi_1 \geq m_1 \sin(\pi \ell_0/4L) \Upsilon,$$

on all of Γ . On the other hand, since ϕ_1 has the form $A_j \sin(\sigma x_j)$ on the outer edges and $\sigma \leq \pi/\ell_0$, it follows that

$$(3.18) \quad \phi_1 \leq M_1 \Upsilon.$$

The constant M_1 satisfies the trivial inequality

$$(3.19) \quad M_1 \geq \sqrt{\frac{2}{L}}.$$

To complete the argument, we need to estimate M_1 from above. Suppose that the maximum value of ϕ_1 is achieved at a point $q_1 \in \Gamma_1$. By the concavity of ϕ_1 and the vertex conditions, there exists a segment of Γ , parametrized by $y \in [0, \pi/2\sigma]$ with $y = 0$ at q_1 , on which

$$\phi_1(y) \geq M_1 \cos(\sigma y).$$

Integrating ϕ_1^2 over this segment gives the inequality

$$\pi M_1^2/4\sigma \leq 1.$$

Since $\sigma \leq \pi/2\ell_0$ by (3.5), this proves

$$(3.20) \quad M_1 \leq \sqrt{\frac{2}{\ell_0}},$$

Using the inequalities (3.17), (3.18), (3.19), and (3.20), Theorem 1.1 now follows from Proposition 3.1. \square

4. ESTIMATION OF THE WEIGHTED CHEEGER CONSTANT

To complete the proof of Theorem 1.2, we will use Proposition 3.1 to estimate $h_{\phi_1}(\Gamma)$. We will assume $\mathcal{V}_0 \neq \emptyset$ throughout this discussion, since the bound (1.1) already covers the case of standard vertex conditions.

As we noted at the beginning of §3, if Γ_1 consists of a single point v , then Γ has k equilateral edges connecting v to Dirichlet vertices. If a denotes the edge length, then in all cases

$$(4.1) \quad \lambda_1 = \frac{\pi^2}{4a^2}, \quad \lambda_2 = \frac{\pi^2}{a^2}.$$

Since $\ell_0 = 2a$ for $k = 2$ and $\ell_0 = a$ for $k \neq 2$, the conclusion of Theorem 1.2 holds trivially for these cases.

In the general case, our goal is to estimate $h_{\phi_1}(\Gamma)$ in terms of the unweighted Cheeger constant $h(\Gamma)$. Since the latter satisfies the trivial bound $h(\Gamma) \geq 2/L$, this will complete the proof.

If Γ is not an interval (and $\mathcal{V}_0 \neq \emptyset$), then its Cheeger constant also satisfies a trivial upper bound,

$$(4.2) \quad h(\Gamma) \leq \frac{1}{\ell_0}.$$

To see this, take a cut S given by a single point on an outer edge of Γ , such that Y_1 is a segment of length ℓ_0 .

We can make a similar estimate for $h_{\phi_1}(\Gamma)$, in terms of the function

$$(4.3) \quad f(x) := \frac{\sin^2(\sigma x)}{\int_0^x \sin^2(\sigma t) dt} = \frac{2 \sin^2(\sigma x)}{x - \frac{1}{2\sigma} \sin(2\sigma x)},$$

defined for $x \in (0, \pi/\sigma)$. It is easy to check that f is decreasing on this interval.

Lemma 4.1. *If Γ_1 contains more than one point, then*

$$(4.4) \quad h_{\phi_1}(\Gamma) \leq f(\ell_0),$$

Proof. Taking a Cheeger cut S consisting of a single point $x_j = \ell_0$ in one of the intervals I_j from (3.1) gives the ratio

$$(4.5) \quad \frac{\sum_S \phi_1^2}{\int_{Y_1} \phi_1^2} = f(\ell_0),$$

where $Y_1 = \{0 \leq x_j < \ell_0\}$ and Y_2 is the other component of $\Gamma \setminus S$. If \mathcal{V}_0 contains more than one point, then we can choose j so as to minimize $\int_{Y_1} \phi_1^2$ among all I_j , guaranteeing that

$$(4.6) \quad \int_{Y_1} \phi_1^2 \leq \int_{Y_2} \phi_1^2.$$

On the other hand, if $\#\mathcal{V}_0 = 1$, then, assuming that Γ_1 is not a single point, Γ contains at least one interior edge on which $\phi_1 \geq m_1$, implying that (4.6) holds also in this case. From (4.5) we thus obtain (4.4). \square

Proposition 4.2. *If Γ_1 contains more than one point, then*

$$h_{\phi_1}(\Gamma) \geq \left(\frac{m_1 \sin(\sigma \ell_0 / 2)}{M_1} \right)^2 h(\Gamma),$$

where $\sigma = \sqrt{\lambda_1}$ and M_1, m_1 are the upper and lower bounds on ϕ_1 from §3.

Proof. By Lemma 4.1, to estimate $h_{\phi_1}(\Gamma)$ we may limit our attention to Cheeger cuts S for which

$$(4.7) \quad \frac{\sum_S \phi_1^2}{\min(\int_{Y_1} \phi_1^2, \int_{Y_2} \phi_1^2)} \leq f(\ell_0).$$

By relabeling if necessary, we can also assume that the minimum in the denominator is the Y_1 integral, i.e.,

$$(4.8) \quad \int_{Y_1} \phi_1^2 \leq \int_{Y_2} \phi_1^2.$$

Let

$$W := \{q \in \Gamma : \text{dist}(q, \mathcal{V}_0) \geq \ell_0/2\}.$$

Case 1: Suppose that $S \cap W = \emptyset$. In the notation of (3.1),

$$\Gamma \setminus W = \cup_j \{0 \leq x_j \leq \ell_0/2\}.$$

Because ϕ_1 is increasing as a function of x_j , the convention (4.8) implies that $Y_1 \subset \Gamma \setminus W$ and $Y_2 \supset W$. Let \mathcal{J} be the set of indices j for which S intersects I_j , and set

$$s_j = \max S \cap I_j.$$

If the restriction of ϕ_1 to I_j is written as $A_j \sin(\sigma x_j)$, then

$$\begin{aligned} \sum_S \phi_1^2 &= \sum_{j \in \mathcal{J}} A_j^2 \sin^2(\sigma s_j) \\ &= \sum_{j \in \mathcal{J}} f(s_j) \int_0^{s_j} A_j^2 \sin^2(\sigma x) dx. \end{aligned}$$

Since f is decreasing and $s_j \leq \ell_0/2$, this implies

$$(4.9) \quad \sum_S \phi_1^2 \geq f(\ell_0/2) \int_{Y_1} \phi_1^2,$$

which contradicts the assumption (4.7).

Case 2: Suppose that $S \cap W \neq \emptyset$ and Y_1 contains an interval $J \subset \Gamma \setminus W$. If S' denotes the corresponding cut with the endpoints of J deleted, then Y_1 is reduced to a component $Y_1' = Y_1 \setminus J$, while $Y_2' = Y_2 \cup J$. The inequality (4.8) is still satisfied after the replacement, and

$$(4.10) \quad \sum_{\partial J} \phi_1^2 \geq f(\ell_0/2) \int_J \phi_1^2,$$

by the same argument used for (4.9). By the assumption (4.7), and the fact that f is strictly decreasing, we obtain

$$\frac{\sum_S \phi_1^2}{\int_{Y_1} \phi_1^2} < \frac{\sum_{\partial J} \phi_1^2}{\int_J \phi_1^2}$$

A comparison of cross-ratios then shows that

$$\frac{\sum_S \phi_1^2}{\int_{Y_1} \phi_1^2} > \frac{\sum_{S'} \phi_1^2}{\int_{Y_1'} \phi_1^2}.$$

That is, cutting the interval J from Y_1 will reduce the Cheeger ratio.

Case 3: Suppose that $S \cap W \neq \emptyset$ and Y_1 contains no interval in $\Gamma \setminus W$. We continue to assume that the components Y_j satisfy (4.7) and (4.8), which together imply that

$$(4.11) \quad \frac{\sum_S \phi_1^2}{\int_{Y_1} \phi_1^2} \leq f(\ell_0).$$

For a cut S with these properties, the points of $S \setminus W$ (if any) correspond, in the parametrization (3.1), to $x_j = b_j < \ell_0/2$, such that $[0, b_j] \subset Y_2$. If we set

$$Z := \bigcap_{b_j \in S \setminus W} [0, b_j],$$

then the set $S'' := S \cap W$ separates Γ into components $Y_1'' := Y_1 \cup Z$ and $Y_2'' = Y_2 \setminus Z$.

By (3.15), ϕ_1 satisfies a lower bound

$$\min_W \phi_1 \geq m_1 \sin(\sigma \ell_0/2).$$

We thus have

$$(4.12) \quad \begin{aligned} \frac{\sum_S \phi_1^2}{\int_{Y_1} \phi_1^2} &\geq \frac{\sum_{S''} \phi_1^2}{\int_{Y_1''} \phi_1^2} \\ &\geq \left(\frac{m_1 \sin(\sigma \ell_0/2)}{M_1} \right)^2 \frac{\#S''}{|Y_1''|}. \end{aligned}$$

For Y_2 we can write the ratio as

$$(4.13) \quad \frac{\sum_S \phi_1^2}{\int_{Y_2} \phi_1^2} = \frac{\sum_{S''} \phi_1^2 + \sum_{S \setminus W} \phi_1^2}{\int_{Y_2''} \phi_1^2 + \int_Z \phi_1^2}.$$

By (4.11), and the fact

$$\sum_{S \setminus W} \phi_1^2 \geq f(\ell_0/2) \int_Z \phi_1^2,$$

the decomposition (4.13) shows that

$$\frac{\sum_{S''} \phi_1^2}{\int_{Y_2''} \phi_1^2} \leq \frac{\sum_S \phi_1^2}{\int_{Y_1} \phi_1^2}.$$

The estimates of ϕ_1 then yield

$$(4.14) \quad \frac{\sum_S \phi_1^2}{\int_{Y_1} \phi_1^2} \geq \left(\frac{m_1 \sin(\sigma \ell_0/2)}{M_1} \right)^2 \frac{\#S''}{|Y_2''|}.$$

Combining (4.12) and (4.14) gives

$$(4.15) \quad \frac{\sum_S \phi_1^2}{\int_{Y_1} \phi_1^2} \geq \left(\frac{m_1 \sin(\sigma \ell_0/2)}{M_1} \right)^2 h(\Gamma).$$

To summarize, Case 1 is ruled out by (4.7) and Case 2 can be reduced to Case 3 with a reduction in the weighted Cheeger ratio. Hence the bound (4.15) applies to $h_{\phi_1}(\Gamma)$. \square

We can now complete the proof of Theorem 1.2. As noted above, the result follows from (4.1) if Γ_1 consists of a single point. Otherwise, Propositions 3.1 and 4.2 together show that

$$\begin{aligned} h_{\phi_1}(\Gamma) &\geq \left(c(L, \ell_0) \sin\left(\frac{\sigma\ell_0}{2}\right)\right)^2 h(\Gamma) \\ &\geq \left(c(L, \ell_0) \sin\left(\frac{\pi\ell_0}{4L}\right)\right)^2 \frac{2}{L}. \end{aligned}$$

By Theorem 2.1, this proves (1.10) with

$$C(L, \ell_0) = \frac{1}{L^2} \left(c(L, \ell_0) \sin\left(\frac{\pi\ell_0}{4L}\right)\right)^4.$$

5. UPPER BOUNDS FOR TREES

The upper bounds for Dirichlet domains in \mathbb{R}^n mentioned in the introduction are proven by creating a family of test functions from the products of eigenfunctions with an affine function on \mathbb{R}^n . The results of Nicaise [15, Thm. 4.1] and Demirel-Harrell [6, Thm. 2.5] use a similar construction, where an *affine function* on a metric graph is interpreted as a continuous function which is linear on each edge. Throughout this section, we assume that Γ is a metric tree, with \mathcal{V}_0 consisting of the external vertices, meaning of degree one.

Our goal in this section is to adapt the techniques of Harrell-Stubbe [9] to produce a general eigenvalue estimate which generalizes the Nicaise bound (1.16). Our first step is to produce a family of affine functions whose derivatives cover Γ uniformly in an average sense.

Let $\mathcal{A}(\Gamma)$ denote the space of affine functions on Γ which satisfy standard vertex conditions at the points of \mathcal{V}_N . No vertex condition are imposed at the points of \mathcal{V}_0 . If we interpret Γ as an electric circuit, with each edge assigned a resistance equal to its length, then functions $\mathcal{A}(\Gamma)$ corresponds precisely to a voltage function satisfying the Ohm's and Kirchhoff's circuit laws. As Kirchhoff [12] demonstrated in 1847, there exists a voltage function for any combination of external voltages applied at the points in \mathcal{V}_0 . Hence, $\mathcal{A}(\Gamma)$ contains non-constant functions provided \mathcal{V}_0 contains at least two points.

From Nicaise [15, Lemma 4.2] we quote the following result. The original did not include a proof, so we will give one here. A similar result was derived independently in Demirel-Harrell [6, Thm. 2.9], but without the restriction to three functions.

Lemma 5.1. *For a metric tree Γ , there exist functions $g_\alpha \in \mathcal{A}(\Gamma)$ for $\alpha \in \{1, 2, 3\}$ such that on all edges, $|g'_\alpha|$ equals 0 or 1 and*

$$(5.1) \quad \sum_{\alpha=1}^3 |g'_\alpha| = 2.$$

Proof. Let us refer to a subgraph of Γ consisting of a vertex with two adjoining external edges (leaves) as a *leaf-pair*. Trimming a leaf pair from a vertex of degree ≥ 3 reduces its degree by 2. If all possible leaf-pairs are trimmed from a given vertex, the result either an artificial (degree 2) or an external vertex (degree 1). By carrying out this trimming process as far as possible at each vertex, we eventually reduce Γ to a single segment. Hence Γ can be constructed by starting from a single segment and attaching leaf-pairs successively. Each leaf-pair is added by gluing its

vertex to any point on the graph, which could be an existing vertex or an edge point.

The family $\{g_\alpha\}$ is constructed by induction, using this decomposition of Γ . For the initial segment, we may choose an arbitrary parametrization x and set $g_1(x) = x$, $g_2(x) = -x$, and $g_3(x) = 0$.

Now suppose the family $\{g_\alpha\}$ has been defined with the desired properties for a tree Γ . Let $\tilde{\Gamma}$ be a graph obtained by adding a single leaf-pair to Γ . The extensions \tilde{g}_α may be defined as follows:

- (1) Suppose the leaf-pair is attached at an internal vertex of Γ (possibly artificial), so that each g_α already satisfies standard vertex conditions at this point. We can extend the family so that both \tilde{g}'_1 and \tilde{g}'_2 alternate ± 1 on edges of the leaf-pair, while \tilde{g}_3 is constant on these edges.
- (2) If the leaf-pair is attached an an external vertex of Γ , then by construction two of the g_α , say $\alpha = 1, 2$ will have derivatives ± 1 at this vertex and g_3 will be constant. We extend the family so that \tilde{g}'_1 is zero on one leaf of the pair, \tilde{g}'_2 is zero on the other, and the derivatives on the other leaves are chosen to satisfy the vertex condition. The third function, whose derivative vanishes into the vertex, is extended so that \tilde{g}'_3 alternates ± 1 on the new leaves.

At every stage of the induction, each function satisfies the vertex conditions and $|g'_\alpha| = 1$ for two values of α and $|g'_\alpha| = 0$ for the third. \square

Using the collection $\{g_\alpha\}$, we can prove the main result of this section, a version of the general eigenvalue inequality from Harrell-Stubbe [9, Thm. 5].

Theorem 5.2. *Let Γ be a metric tree with Dirichlet conditions on external vertices. For $z \in (\lambda_n, \lambda_{n+1}]$, suppose that f is a positive function on the set $\{\lambda_1, \dots, \lambda_n\}$ such that $f(\lambda_j)/(z - \lambda_j)^2$ is nondecreasing with j . Then*

$$\sum_{j=1}^n f(\lambda_j) \leq 4 \sum_{j=1}^n \frac{f(\lambda_j)}{z - \lambda_j} \lambda_j.$$

Proof. Let G_α be the multiplication operator on $L^2(\Gamma)$ associated to g_α . We define a corresponding set of first-order differential operators

$$D_\alpha := \frac{1}{2}[\Delta, G_\alpha].$$

On an edge parametrized by x , D_α acts as $(\partial_x g_\alpha)\partial_x$. Since g'_α takes values in $\{0, \pm 1\}$, the commutator $[D_\alpha, G_\alpha]$ is the projection onto the support of g'_α . By the construction in Lemma 5.1,

$$\sum_{\alpha=1}^3 [D_\alpha, G_\alpha] = 2.$$

Thus we can write

$$\begin{aligned} \sum_{j=1}^n f(\lambda_j) &= \text{tr}(P_n f(-\Delta)) \\ &= \frac{1}{2} \sum_{\alpha=1}^3 \text{tr}(P_n f(-\Delta)[D_\alpha, G_\alpha]), \end{aligned}$$

where P_n denotes the spectral projection onto the eigenspace for $\{\lambda_1, \dots, \lambda_n\}$. Expanding the trace, just as in the proof of [9, Thm. 1], then gives

$$\begin{aligned} \sum_{j=1}^n f(\lambda_j) &= - \sum_{\alpha=1}^3 \sum_{j=1}^n \sum_{\substack{m=1 \\ \lambda_m \neq \lambda_j}}^n \frac{f(\lambda_j) - f(\lambda_m)}{\lambda_j - \lambda_m} |\langle D_\alpha \phi_j, \phi_m \rangle|^2 \\ &\quad + 2 \sum_{\alpha=1}^3 \sum_{j=1}^n \sum_{q=n+1}^{\infty} \frac{f(\lambda_j)}{\lambda_q - \lambda_j} |\langle D_\alpha \phi_j, \phi_q \rangle|^2. \end{aligned}$$

By the hypotheses on f , the argument from proof of [9, Thm. 5] then applies directly to give

$$\sum_{j=1}^n f(\lambda_j) \leq 2 \sum_{\alpha=1}^3 \sum_{j=1}^n \frac{f(\lambda_j)}{z - \lambda_j} \|D_\alpha \phi_j\|^2.$$

By the construction of g_α ,

$$\sum_{\alpha=1}^3 \|D_\alpha \phi_j\|^2 = 2 \|\phi_j'\|^2 = 2\lambda_j,$$

which completes the proof. \square

Taking $f = 1$ in Theorem 5.2 yields the Hile-Protter bound (1.16) obtained by Nicaise. Using $f(\lambda) = (z - \lambda)^2$ gives an inequality

$$(5.2) \quad \sum_{j=1}^n (z - \lambda_j)(z - 5\lambda_j) \leq 0,$$

for $z \in [\lambda_n, \lambda_{n+1}]$, which was obtained previously by Demirel-Harrell [6, Eq. (3.15)]. The discriminant of the quadratic polynomial on the left side of (5.2) is positive and the roots must lie outside the interval $(\lambda_n, \lambda_{n+1})$. This yields the following result, analogous to [9, Prop. 6]:

Theorem 5.3. *For Γ a metric tree with Dirichlet vertices on the external vertices, the quantity*

$$D_n := \left(\frac{3}{n} \sum_{j=1}^n \lambda_j \right)^2 - \frac{5}{n} \sum_{j=1}^n \lambda_j^2$$

satisfies $D_n \geq 0$ for all $n \geq 2$. Furthermore, the eigenvalues satisfy the inequalities

$$\lambda_n \geq \frac{3}{n} \sum_{j=1}^n \lambda_j - \sqrt{D_n}, \quad \lambda_{n+1} \leq \frac{3}{n} \sum_{j=1}^n \lambda_j + \sqrt{D_n},$$

and hence

$$\lambda_{n+1} - \lambda_n \leq 2\sqrt{D_n}.$$

For $n = 1$ we have $D_1 = 4\lambda_1^2$, so the estimate reduces to $\lambda_2/\lambda_1 \leq 5$, equivalent to (1.15) and (1.16) but weaker than (1.14). To compare the estimates for λ_3 , let us define the moments for the first pair of eigenvalues,

$$a_1 := \frac{1}{2}(\lambda_1 + \lambda_2), \quad a_2 := \frac{1}{2}(\lambda_1^2 + \lambda_2^2).$$

Then the estimate from (5.3) reads

$$(5.3) \quad \lambda_3 \leq 3a_1 + \sqrt{9a_1^2 - 5a_2}.$$

On the other hand, the Hile-Protter type bound (1.16) reduces to

$$\lambda_3 \leq 3a_1 + \sqrt{9a_1^2 - 5\lambda_1\lambda_2}.$$

Since $\lambda_1\lambda_2 \leq a_2$, by the geometric mean inequality, the bound (5.3) is stronger.

6. EXTENSIONS OF THE UPPER BOUND

In this section we investigate the possibility for extending the upper bounds discussed in §5 to graphs which are modifications of trees. We will consider two possibilities: adding edges between existing vertices of the tree and attaching pendant graphs.

We cannot expect to produce a set of affine functions satisfying (5.1) on a general graph. However, it is worth noting that the existence of a single function $h \in \mathcal{A}(\Gamma)$ gives a bound in terms of ϕ_1 . Returning to the electric circuit analogy from §5, let us define a (scalar) *current* as a function $\eta : \Gamma \rightarrow [0, \infty)$ which is constant on each edge and which, under some choice of edge orientations, satisfies Kirchhoff's current law. This is equivalent to the condition

$$(6.1) \quad \eta = |h'| \text{ for some } h \in \mathcal{A}(\Gamma).$$

The following inequality is implicit in the proofs of the Payne-Pólya-Weinberger estimate $\lambda_2/\lambda_1 \leq 5$ in various cases. For convenience we include a direct proof.

Lemma 6.1. *If Γ is a metric graph that admits a non-zero current function η , then*

$$(6.2) \quad \lambda_2 - \lambda_1 \leq 4 \frac{\|\eta\phi_1'\|^2}{\|\eta\phi_1\|^2}.$$

Proof. Given η , choose h according to (6.1). By shifting h by a constant if necessary, we can assume that $u := h\phi_1$ is orthogonal to ϕ_1 . Then min-max gives the estimate

$$(6.3) \quad \lambda_2 \leq \frac{\|u'\|^2}{\|u\|^2}.$$

Because u satisfies the vertex conditions, by the assumptions on h , we can integrate by parts to compute

$$\begin{aligned} \|u'\|^2 &= \langle u, -\Delta u \rangle \\ &= \langle u, \lambda_1 u - 2h'\phi_1' \rangle \\ &= \lambda_1 \|u\|^2 - 2\langle u, h'\phi_1' \rangle. \end{aligned}$$

By (6.3) this gives

$$(6.4) \quad \lambda_2 - \lambda_1 \leq \frac{A}{\|u\|^2},$$

where

$$A := -2\langle u, h'\phi_1' \rangle.$$

The Cauchy-Schwarz estimate gives

$$(6.5) \quad A^2 \leq 4\|u\|^2 \|\eta\phi_1'\|^2.$$




FIGURE 11. A saguaro graph.

On the other hand, we can compute using integration by parts,

$$\begin{aligned} A &= -\frac{1}{2} \int_{\Gamma} (h^2)' (\phi_1^2)' \\ &= \frac{1}{2} \int_{\Gamma} (h^2)'' \phi_1^2 \\ &= \|\eta \phi_1\|^2. \end{aligned}$$

By (6.5) this yields

$$\frac{A}{\|u\|^2} \leq \frac{\|\eta \phi_1'\|^2}{\|\eta \phi_1\|^2},$$

and the result follows from (6.4). \square

As noted in §5, any graph Γ for which \mathcal{V}_0 has at least two vertices will admit non-zero current functions. However, to obtain universal bounds from (6.2) we need either some uniformity in the choice of η or some control over the behavior of ϕ_1 .

6.1. Saguaro graphs. The first observation is that the conclusion of Lemma 5.1 continues to hold for graphs constructed from trees by adding edges uniformly. That is, suppose Γ is constructed from a tree by replacing each internal edge by a pumpkin with $k > 1$ edges of the same length, and each external edge by a star with k edges of the same length. We will call the result a *saguaro* graph, after the tree-like cactus.

Given a saguaro graph Γ based on the tree Γ_T , we can apply Lemma 5.1 to produce a trio of affine functions $\{g_\alpha\}$ on Γ_T . These functions can be extended to elements of $\mathcal{A}(\Gamma)$ satisfying (5.1) by simply replicating the values on parallel edges. The existence of this tree implies that all of the universal tree bounds carry over to Γ , including

$$\frac{\lambda_2}{\lambda_1} \leq 2 + \sqrt{5}, \quad \lambda_{n+1} - \lambda_n \leq \frac{4}{n} \sum_{j=1}^n \lambda_j,$$

from Nicaise [15, §4], along with the general bounds from Theorem 5.2.

We can also consider the case of a modified saguaro graph, where the number of edges of each pumpkin or star varies between values k_{\min} and k_{\max} . We continue




FIGURE 12. An ornamented tree with two pendants.

to assume that the edges have equal lengths within each pumpkin or star. Suppose $\{g_\alpha\}$ denotes the trio of affine functions produced by Lemma 5.1 for the underlying tree Γ_T . On Γ we can produce a corresponding set of currents $\{\eta_\alpha\}$ by subdividing the current g'_α among the parallel edges in each segment. That is, on a segment (pumpkin or star) with k edges, we set $\eta_\alpha := |g'_\alpha|/k$. Because $\sum_\alpha |g'_\alpha|^2 = 1$ on Γ_T , this construction gives

$$\frac{1}{k_{\max}^2} \leq \sum_{\alpha=1}^3 \eta_\alpha^2 \leq \frac{1}{k_{\min}^2}.$$

By summing the inequality

$$(\lambda_2 - \lambda_1) \|\eta_\alpha \phi_1\|^2 \leq 4 \|\eta_\alpha \phi_1'\|^2$$

over α , we obtain

$$\frac{\lambda_2}{\lambda_1} \leq 1 + 4 \frac{k_{\max}^2}{k_{\min}^2}.$$

6.2. Ornamented trees. Suppose Γ is constructed from a tree graph Γ_T by attaching pendant graphs P_j , $j = 1, \dots, m$, to internal (and possibly artificial) vertices of Γ_T . Let $q_j \in \mathcal{V}_N$ denote the attachment vertex for P_j . See Figure 12 for an illustration of this *ornamented tree* graph.

Proposition 6.2. *Let Γ be an ornamented tree consisting of metric tree Γ_T , with Dirichlet conditions at exterior vertices, and pendants P_1, \dots, P_m each containing at least one Dirichlet vertex. Suppose that*

$$\lambda_1 \leq \lambda_1(P_j)$$

for each j , where $\{\lambda_n\}$ denotes the spectrum of Γ and $\lambda_1(P_j)$ is the first eigenvalue of P_j defined by assigning standard boundary conditions at the attachment point q_j . Then

$$\frac{\lambda_2}{\lambda_1} \leq 5.$$

In particular, this inequality holds if

$$(6.6) \quad |P_j| \leq \frac{1}{2} \ell_{\max}(\Gamma_T)$$

for each j .

Proof. Let $\{g_\alpha\} \in \mathcal{A}(\Gamma_T)$ be the trio of functions obtained by Lemma 5.1. We define corresponding current functions on Γ by setting $\eta_\alpha = |g'_\alpha|$ on edges of Γ_T and $\eta_\alpha = 0$ on each pendant P_j . This gives

$$\frac{1}{2} \sum_{\alpha=1}^3 \|\eta_\alpha \phi_1\|^2 = \int_{\Gamma_T} \phi_1^2,$$

and

$$\frac{1}{2} \sum_{\alpha=1}^3 \|\eta_\alpha \phi'_1\|^2 = \int_{\Gamma_T} (\phi'_1)^2.$$

Thus, by Lemma 6.1,

$$(6.7) \quad \lambda_2 - \lambda_1 \leq 4 \frac{\int_{\Gamma_T} (\phi'_1)^2}{\int_{\Gamma_T} \phi_1^2}.$$

An integration by parts gives

$$(6.8) \quad \int_{\Gamma_T} (\phi'_1)^2 = \lambda_1 \int_{\Gamma_T} \phi_1^2 + \sum_{j=1}^m \phi_1(q_j) \partial_{P_j} \phi_1(q_j),$$

where $\partial_{P_j} \phi_1(q_j)$ denotes the sum of derivatives of ϕ_1 at q_j into the incident edges of P_j .

Now for each j let u_j denote the first eigenfunction of P_j , with standard vertex conditions imposed at q_j , so that $u_j \geq 0$ on P_j and

$$-\Delta u_j = \lambda_1(P_j) u_j.$$

By Green's identity, and the fact that $\partial_{P_j} u_j(q_j) = 0$,

$$\int_{P_j} (-\phi_1 \Delta u_j + u_j \Delta \phi_1) = -u_j(q_j) \partial_{P_j} \phi_1(q_j).$$

On the other hand,

$$\int_{P_j} (-\phi_1 \Delta u_j + u_j \Delta \phi_1) = (\lambda_1(P_j) - \lambda_1) \int_{P_j} \phi_1 u_j.$$

Since the eigenfunctions are positive, we conclude that $\lambda_1(P_j) - \lambda_1 \geq 0$ implies that

$$\partial_{P_j} \phi_1(q_j) \leq 0.$$

Under this assumption, (6.8) gives

$$\int_{\Gamma_T} (\phi'_1)^2 \leq \lambda_1 \int_{\Gamma_T} \phi_1^2,$$

and it follows from (6.7) that $\lambda_2/\lambda_1 \leq 5$.

By the general bounds (1.1) and (1.6), the eigenvalue condition will hold provided

$$|P_j| \leq \frac{1}{2} \ell_{\max}(\Gamma).$$

Clearly it suffices to compute ℓ_{\max} over Γ_T rather than Γ . \square

REFERENCES

- [1] G. Berkolaiko, J. B. Kennedy, P. Kurasov, and D. Mugnolo, Edge connectivity and the spectral gap of combinatorial and quantum graphs, *J. Phys. A* **50** (2017), 365201, 29.
- [2] G. Berkolaiko, J. B. Kennedy, P. Kurasov, and D. Mugnolo, Surgery principles for the spectral analysis of quantum graphs, *Trans. Amer. Math. Soc.* **372** (2019), 5153–5197.
- [3] G. Berkolaiko, J. B. Kennedy, P. Kurasov, and D. Mugnolo, Impediments to diffusion in quantum graphs: geometry-based upper bounds on the spectral gap, preprint, [arXiv:2206.10046v1](https://arxiv.org/abs/2206.10046v1), 2022.
- [4] D. Borthwick, L. Corsi, and K. Jones, Sharp diameter bound on the spectral gap for quantum graphs, *Proc. Amer. Math. Soc.* **149** (2021), 2879–2890.
- [5] S.-Y. Cheng and K. Oden, Isoperimetric inequalities and the gap between the first and second eigenvalues of an Euclidean domain, *J. Geom. Anal.* **7** (1997), 217–239.
- [6] S. Demirel and E. M. Harrell, II, On semiclassical and universal inequalities for eigenvalues of quantum graphs, *Rev. Math. Phys.* **22** (2010), 305–329.
- [7] L. Friedlander, Extremal properties of eigenvalues for a metric graph, *Ann. Inst. Fourier (Grenoble)* **55** (2005), 199–211.
- [8] E. M. Harrell, II and A. V. Maltsev, Localization and landscape functions on quantum graphs, *Trans. Amer. Math. Soc.* **373** (2020), 1701–1729.
- [9] E. M. Harrell, II and J. Stubbe, On trace identities and universal eigenvalue estimates for some partial differential operators, *Trans. Amer. Math. Soc.* **349** (1997), 1797–1809.
- [10] G. N. Hile and M. H. Protter, Inequalities for eigenvalues of the Laplacian, *Indiana Univ. Math. J.* **29** (1980), 523–538.
- [11] J. B. Kennedy, P. Kurasov, G. Malenová, and D. Mugnolo, On the spectral gap of a quantum graph, *Ann. Henri Poincaré* **17** (2016), 2439–2473.
- [12] G. R. Kirchhoff, Ueber die Auflösung der Gleichungen, auf welche man bei der Untersuchung der linearen Vertheilung galvanischer Ströme geführt wird, *Pogg. Annal. Phys. u. Chemie* **72** (1847), Reprinted as pp. 22–33 in Gustav Robert Kirchhoff, *Gesammelte Abhandlungen*, Edition Classic Verlag Dr. Müller. Saarbrücken: VDM Verlag Dr. Müller, 2006.
- [13] P. Kurasov, G. Malenová, and S. Naboko, Spectral gap for quantum graphs and their edge connectivity, *J. Phys. A* **46** (2013), 275309, 16.
- [14] P. Kurasov and S. Naboko, Rayleigh estimates for differential operators on graphs, *J. Spectr. Theory* **4** (2014), 211–219.
- [15] S. Nicaise, Spectre des réseaux topologiques finis, *Bull. Sci. Math. (2)* **111** (1987), 401–413.
- [16] K. Oden, C.-J. Sung, and J. Wang, Spectral gap estimates on compact manifolds, *Trans. Amer. Math. Soc.* **351** (1999), 3533–3548.
- [17] L. E. Payne, G. Pólya, and H. F. Weinberger, On the ratio of consecutive eigenvalues, *J. Math. and Phys.* **35** (1956), 289–298.
- [18] I. M. Singer, B. Wong, S.-T. Yau, and S. S.-T. Yau, An estimate of the gap of the first two eigenvalues in the Schrödinger operator, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **12** (1985), 319–333.
- [19] C. J. Thompson and M. Kac, Phase transition and eigenvalue degeneracy of a one dimensional anharmonic oscillator, *Studies in Appl. Math.* **48** (1969), 257–264.

DEPARTMENT OF MATHEMATICS, EMORY UNIVERSITY, ATLANTA, GA 30322

Email address: dborthw@emory.edu

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332

Email address: harrell@math.gatech.edu

DEPARTMENT OF MATHEMATICS, EMORY UNIVERSITY, ATLANTA, GA 30322

Email address: haozhe.yu@emory.edu