Redshift evolution of cosmic birefringence in CMB anisotropies

Matteo Galaverni, $^{1,\,2,\,*}$ Fabio Finelli, $^{3,\,4,\,\dagger}$ and Daniela Paoletti $^{2,\,4,\,\ddagger}$

¹Specola Vaticana (Vatican Observatory), V-00120, Vatican City State

²INAF/OAS Bologna, Via Gobetti 101, I-40129 Bologna, Italy

³INAF/OAS Bologna, via Gobetti 101, I-40129 Bologna, Italy

⁴INFN, Sezione di Bologna, Via Irnerio 46, I-40127 Bologna, Italy

(Dated: May 2, 2023)

We study the imprints of a cosmological redshift-dependent pseudoscalar field ϕ on the rotation of cosmic microwave background (CMB) linear polarization generated by a coupling $\phi F^{\mu\nu} \tilde{F}_{\mu\nu}$. We show how either phenomenological or theoretically motivated redshift dependence of the pseudoscalar field, such as those in models of Early Dark Energy, Quintessence or axion-like dark matter, lead to CMB polarization and temperature-polarization power spectra which exhibit a multipole dependence which goes beyond the widely adopted approximation in which the redshift dependence of the linear polarization angle is neglected. Because of this multipole dependence, the isotropic birefringence effect due to a general coupling $\phi F^{\mu\nu} \tilde{F}_{\mu\nu}$ is not degenerate with a systematic calibration angle uncertainty. By taking this multipole dependence into account, we calculate the parameters of these phenomenological and theoretical redshift dependence of the pseudoscalar field which can be detected by future CMB polarization experiments on the basis of a χ^2 analysis for a Wishart likelihood. As a final example of our approach, we compute by Markov Chain MonteCarlo (MCMC) the minimal coupling g_{ϕ} in Early Dark Energy which could be detected by future experiments, with or without marginalizing on a systematic rotation angle uncertainty.

I. INTRODUCTION

When the electromagnetic tensor $F_{\mu\nu}$ is coupled to a pseudoscalar field $\phi(x)$ a new term appears in the Lagrangian density:

$$\mathcal{L} \supset -\frac{g_{\phi}}{4} \phi F_{\mu\nu} \,\widetilde{F}^{\mu\nu} \,, \tag{1}$$

where g_{ϕ} is a model dependent coupling constant and $\tilde{F}^{\mu\nu} \equiv \frac{1}{2} \epsilon^{\mu\nu\rho\sigma} F_{\rho\sigma}$ is the dual of the electromagnetic tensor. The plane of linear polarization of a single photon propagating in this evolving cosmological pseudoscalar field background undergoes a rotation given by [1, 2]:

$$\frac{g_{\phi}}{2} \left[\phi(x) - \phi(x_{\rm em}) \right] \,, \tag{2}$$

where $\phi(x_{\rm em})$ is the value of the pseudoscalar field when light is emitted. This effect is called *cosmological birefringence*.

First upper limits on the coupling constant g_{ϕ} were based on optical imaging polarimetry of radio galaxies [2– 7]. Soon after it was realized that also CMB polarization could be used to study this interaction which induces a rotation of the plane of linear polarization [8–10] to leading order in g_{ϕ} as well as circular polarization to the next-to-leading order [11, 12].

Either the redshift dependence [11, 13–18] and the inhomogeneities [18–23] of the cosmological pseudo-scalar field contribute to cosmological birefringence. In this paper we study the imprints of the isotropic redshift dependence of ϕ along the line-of-sight from last scattering surface to the observer into CMB parity even and odd power spectra. We consider either phenomenological or theoretically motivated redshift dependence of the pseudoscalar field, such as those in models of Early Dark Energy (EDE), Quintessence (DE) or axion-like dark matter (DM) [13, 14, 24–31].

As already pointed out [11, 13, 14, 16–18], this redshift dependence induces a multipole dependence of the cosmological birefringence effect in the CMB power spectra which goes beyond the widely used approximation for which the rotation angle is assumed constant in redshift [8]. Although this approximation is a key working assumption for deriving constraints from CMB polarization data [32–43] and forecast the capabilities of future experiments [44–47], we believe it is timely to fully exploit the theoretical predictions of isotropic cosmological birefringence for two main reasons.

Firstly, by assuming the isotropic birefringence angle as independent on the multipoles an exact degeneracy between the cosmological birefringence effect and the uncertainty in the calibration angle which would be otherwise absent opens up. We explicitly show how taking into account the redshift dependence of cosmological birefringence mitigate this degeneracy (see also [16, 17]).

As a second point, we stress that the advance in data analysis and in the increasingly precision of CMB polarization data shrank error bars approximately by a factor 3 from the *Planck* analysis on data release 2 [36]: hints of isotropic cosmic birefringence within the constant angle approximation were claimed with *Planck* data release 3 (DR3) [48], *Planck* data release 4 (DR4) [49], and more recently with *WMAP 9-year* and *Planck* data-processing

^{*} matteo.galaverni@gmail.com

[†] fabio.finelli@inaf.it

[‡] daniela.paoletti@inaf.it

pipeline called NPIPE [50] ($\alpha = 0.342^{+0.094}_{-0.091}$ deg [51]). For a recent review see [52]. We will indeed show that the differences between a physical model and the constant angle approximation are important and within the reach of future CMB polarization experiments.

The paper is organized as follows: in Sect. II we review the Boltzmann equation in presence of an isotropic redshift-dependent birefringence. We compare the power spectra for some phenomenological models with the widely used approximation where the time dependence of the linear polarization angle is neglected. The study of theoretically motivated redshift dependence of the pseudoscalar field is presented in Sect. III: Early Dark Energy, Quintessence and axion-like dark matter. In Sect. IV we present the forecasts for CMB experiments, focusing in particular on *LiteBIRD* on the basis of a χ^2 analysis for a Wishart likelihood and we perform few exploratory runs exploring the whole cosmological and birefringence parameter space using the Markov Chain MonteCarlo code cosmonc. We conclude in Sect. V.

In this work, we use natural units, $\hbar = c = 1$, and assume flat ΛCDM cosmological model with *Planck* 2018 estimates of cosmological parameters [53]: $\Omega_b h^2 = 0.02237$, $\Omega_c h^2 = 0.120$, $\tau = 0.0544$, $n_s = 0.9649$, $\ln (10^{10}A_s) = 3.044$, $H_0 = 100 h \,\mathrm{km \, s^{-1} \, Mpc^{-1}} = 67.36 \,\mathrm{km \, s^{-1} \, Mpc^{-1}}$.

II. EFFECTS OF REDSHIFT EVOLUTION OF THE BIREFRINGENCE FIELD

The linear polarization rotation for a of a CMB photon is described by:

$$Q = Q_{\rm rec} \cos(2\beta) + U_{\rm rec} \sin(2\beta) , \qquad (3)$$

$$U = U_{\rm rec} \cos(2\beta) - Q_{\rm rec} \sin(2\beta) , \qquad (4)$$

where $Q_{\rm rec}$ and $U_{\rm rec}$ are the Stokes parameters at recombination, when CMB photons are last scattered.¹

In the case of isotropic time-dependent birefringence angle induced by a cosmological pseudoscalar field the Boltzmann equation for linear polarization contains an additional term proportional to $g_{\phi}\phi'(\eta)$, where ϕ' is the derivative of ϕ with respect conformal time η [11, 13, 14, 57]:

$$\begin{aligned} \Delta'_{Q\pm iU}(k,\eta) + ik\mu\Delta_{Q\pm iU}(k,\eta) \\ &= -n_e\sigma_T a(\eta) \left[\Delta_{Q\pm iU}(k,\eta) \right. \\ &+ \sum_m \sqrt{\frac{6\pi}{5}}_{\pm 2} Y_2^m S_P^{(m)}(k,\eta) \right] \\ &\mp ig_\phi \phi'(\eta) \Delta_{Q\pm iU}(k,\eta) \,. \end{aligned}$$
(5)

The the cosine of the angle between the CMB photon direction and the Fourier wave vector is indicated by μ , n_e is the number density of free electrons, σ_T is the Thomson cross section, ${}_{s}Y_2^m$ are spherical harmonics with spinweight s, and $S_P^{(m)}(k,\eta)$ is the source term generating linear polarization.

In order to integrate along the line-of-sight we note that [14]:

$$\Delta'_{Q\pm iU}(k,\eta) + [ik\mu + \tau'(\eta) \pm ig_{\phi}\phi'(\eta)] \Delta_{Q\pm iU}(k,\eta)$$
$$= e^{-ik\mu\eta} e^{\tau(\eta)} e^{\mp i2\alpha(\eta)}$$
$$\frac{d}{d\eta} \left[e^{ik\mu\eta} e^{-\tau(\eta)} e^{\pm i2\alpha(\eta)} \Delta_{Q\pm iU}(k,\eta) \right], \qquad (6)$$

where we introduced the differential optical depth $\tau'(\eta) \equiv n_e \sigma_T a(\eta)$ and we formally integrated $\phi'(\eta)$ defining $\alpha(\eta) = \frac{g_{\phi}}{2} \phi(\eta)$, defined up to a constant. Therefore, the Boltzmann Eq. (5) can be re-written as:

$$e^{-ik\mu\eta}e^{\tau(\eta)}e^{\mp i2\alpha(\eta)}$$
$$\frac{d}{d\eta}\left[e^{ik\mu\eta}e^{-\tau(\eta)}e^{\pm i2\alpha(\eta)}\Delta_{Q\pm iU}(k,\eta)\right]$$
$$= -\tau'(\eta)\sum_{m}\sqrt{\frac{6\pi}{5}}\pm 2Y_2^m S_P^{(m)}(k,\eta).$$
(7)

Following the integration along the line-of-sight methodology [58], we obtain these expressions for the polarization C_{ℓ} auto- and cross-spectra:

$$C_{\ell}^{XY} = (4\pi)^2 \frac{9(\ell+2)!}{16(\ell-2)!} \int k^2 dk \, [\Delta_{X,\ell}(k,\eta_0)\Delta_{Y,\ell}(k,\eta_0)] , \qquad (8)$$
$$C_{\ell}^{TX} = (4\pi)^2 \sqrt{\frac{9(\ell+2)!}{16(\ell-2)!}} \int k^2 dk \, \Delta_{T,\ell}(k,\eta_0)\Delta_{X,\ell}(k,\eta_0) , \qquad (9)$$

where X, Y can be either E or B. The integrals defining the polarization scalar perturbations $\Delta_{T,\ell}$, $\Delta_{E,\ell}$ and $\Delta_{B,\ell}$ are:

$$\Delta_{T,\ell}(k,\eta_0) = \int_{\eta_{\rm rec}}^{\eta_0} d\eta \, g(\eta) S_T(k,\eta) j_\ell(k\eta_0 - k\eta) \eta_0$$

$$\Delta_{E,\ell}(k,\eta_0) = \int_{\eta_{\rm rec}}^{\eta_0} d\eta \, g(\eta) S_P^{(0)}(k,\eta) \frac{j_\ell(k\eta_0 - k\eta)}{(k\eta_0 - k\eta)^2} \cos 2 \left[\alpha(\eta) - \alpha(\eta_0)\right], \quad (11)$$

$$\Delta_{B,\ell}(k,\eta_0) = \int_{\eta_{\rm rec}}^{\eta_0} d\eta \, g(\eta) S_P^{(0)}(k,\eta) \frac{j_\ell(k\eta_0 - k\eta)}{(k\eta_0 - k\eta)^2} \sin 2 \left[\alpha(\eta) - \alpha(\eta_0)\right]. \quad (12)$$

here $S_T(k,\eta) [S_P^{(0)}(k,\eta)]$ is the source term for temperature [scalar polarization] anisotropies, and j_ℓ is the spherical Bessel function of order ℓ .

Note that Δ_E and Δ_B are sensitive to cosmic birefringence through a term proportional to $\alpha(\eta) - \alpha(\eta_0)$, where

¹ We follow CMB-HEALPix coordinate conventions: the linear polarization angle increases clockwise looking toward the source [54–56].

 $\alpha(\eta)$ describes linear polarization rotation from recombination (η_{rec}) to time η :

$$\alpha(\eta) = \int_{\eta_{\rm rec}}^{\eta} \alpha'(\eta_1) d\eta_1 \,. \tag{13}$$



FIG. 1. CAMB visibility function (g_{CAMB}) as a function of redshift z, blue band: 7 < z < 12, green band: $z = z_{rec} \pm 100 (z_{rec} \simeq 1059)$.

The visibility function $g(\eta) = \tau'(\eta)e^{-\tau(\eta)}$ [58] is not constant for photons propagating from last scattering to nowadays: it reaches its maximum at recombination and a second peak is present at reionization epoch, but it is several orders of magnitude smaller. The visibility function of the Boltzmann code CAMB [59] is plotted in Fig. 1 as a function of redshift.

Since the visibility function is highly peaked at recombination a widely used approximation consists in evaluating the new term $\alpha(\eta) - \alpha(\eta_0)$ appearing in Eqs. (11)-(12) at recombination [24]. In this approximation:

$$\alpha(\eta) - \alpha(\eta_0) \simeq \alpha(\eta_{\rm rec}) - \alpha(\eta_0) \equiv \bar{\alpha} \,, \tag{14}$$

the constant terms $\cos(2\bar{\alpha})$ and $\sin(2\bar{\alpha})$ exit integration over time in Eqs. (11) - (12) and the following expressions for the power spectra as a function of the power spectra at recombination (rec) are obtained (assuming at recombination both $C_{\ell}^{BB,\text{rec}} = 0$ and vanishing parity odd power spectra $C_{\ell}^{TB,\text{rec}} = C_{\ell}^{EB,\text{rec}} = 0$) [8, 9, 14, 60, 61]:

$$C_{\ell}^{TE,\text{const}} = C_{\ell}^{TE,\text{rec}}\cos(2\bar{\alpha}), \qquad (15)$$

$$C_{\ell}^{TB,\text{const}} = C_{\ell}^{TE,\text{rec}} \sin(2\bar{\alpha}), \qquad (16)$$

$$C_{\ell}^{EE,\text{const}} = C_{\ell}^{EE,\text{rec}} \cos^2(2\bar{\alpha}), \qquad (17)$$

$$C_{\ell}^{BB,\text{const}} = C_{\ell}^{EE,\text{rec}} \sin^2(2\bar{\alpha}), \qquad (18)$$

$$C_{\ell}^{EB,\text{const}} = \frac{1}{2} C_{\ell}^{EE,\text{rec}} \sin(4\bar{\alpha}) \,. \tag{19}$$

The main purpose of this paper is to compare the results of Eqs. (8)-(9) with the constant approximation of Eqs. (15) - (19).

In order to study in a more detailed way the effects of isotropic cosmic birefringence $\alpha(\eta)$ we modified

source terms in the Boltzmann code CAMB [59] following Eqs. (11) and (12). In Fig. 2 we compare the effects on the power spectra of a sudden/instantaneous rotation $\bar{\alpha} = 1$ deg occurring at different epochs. The initial value of the linear polarization angle is always $\alpha(\eta_{\rm rec}) = 0$ deg, then α drops to -1 deg, but at different epochs. We consider in particular $\alpha(\eta) = -1/2 \{ 1 + \tanh [10^4 (x - x_*)] \}$ deg, where $x \equiv$ $(\eta - \eta_{\text{rec}}) / (\eta_0 - \eta_{\text{rec}})$ and $x_* = \{0.005, 0.25, 0.5, 0.996\}.$ To give an idea of the numbers involved in ΛCDM we have $x(\eta_{\rm rec}) = 0$, $x(\eta_{z=100}) = 0.08$, $x(\eta_{z=10}) = 0.31$, $x(\eta_{z=5}) = 0.44, x(\eta_0) = 1$. If the change of the linear polarization angle happens nowadays (at $\eta \simeq \eta_0$, or $x_* \simeq 1$) then we clearly have $\alpha(\eta) - \alpha(\eta_0) = 1$ deg during all integration along the line-of-sight. In this case the power spectra obtained using the modified Boltzmann code exactly coincide with the analytic expressions of Eqs. (15)- (19) fixed $\bar{\alpha} = 1$ deg. Note that a miscalibration of the orientation of the detector is assimilable to a rotation at present time of the linear polarization vector and gives an analogue effect on the power spectra [61-63]. We clearly see that earlier in time the rotation happens, smaller are the effects on the power spectra (in particular the difference is larger at lower ℓ). For BB we also plot the power spectra induced by lensing (black dotted line) and tensor perturbations assuming a tensor-to-scalar ratio $r = 10^{-2}$ (dashed black line) and $r = 10^{-3}$ (dot-dashed black line). For a detailed discussion of the impact of lensing on cosmic birefringence we refer to [64].

In Fig. 3 we show the output of the modified Boltzmann code considering a rotation $\bar{\alpha}$ of linear polarization (i) localized only at*early times* (from last scattering to reionization) $\alpha(\eta) = \bar{\alpha} \left\{ \frac{\tanh[10^4(x-0.001)]+1}{2} - \frac{\tanh[10^4(x-0.25)]+1}{2} \right\}, \text{ or}$ (ii) only at *late times* (from reionization to nowadays) $\alpha(\eta) = \bar{\alpha} \left\{ \frac{\tanh[10^4(x-0.25)]+1}{2} - \frac{\tanh[10^4(x-0.996)]+1}{2} \right\}$ Since the linear polarization rotation is not constant over time, the effects on the power spectra are different. If α is rotated only at early times the effects on the power spectra are localized at $\ell \gtrsim 10$. Otherwise if the linear polarization angle rotates after reionization (late times), the effects are visible only at $\ell \lesssim 10$.

Interestingly we stress that birefringence effects on the power spectra can be present even if $\alpha(\eta_{\rm rec}) = \alpha(\eta_0)$, differently from what stated in Ref. [29]. In this case according to the analytic expressions of Eqs. (15)-(19) there should be no effects since $\bar{\alpha} = 0$. On the contrary there are evident effects on the power spectra using the modified CAMB code based on the Boltzmann equation for cosmic birefringence, see in particular Fig. 4. See also Appendix A for other interesting phenomenological cases with $\alpha(\eta_{\rm rec}) = \alpha(\eta_0)$.



FIG. 2. (a) Evolution of $\alpha(\eta) - \alpha(\eta_0)$ as a function of conformal time $(\eta - \eta_{\rm rec}) / (\eta_0 - \eta_{\rm rec})$; α always starts at $\alpha(\eta_{\rm rec}) = 0$ deg and ends at $\alpha(\eta_0) = -1$ deg, but the "sudden/instantaneous rotation" happens at different times: purple line corresponds to a rotation near present time $x_* \equiv (\eta_* - \eta_{\rm rec}) / (\eta_0 - \eta_{\rm rec}) = 0.996$, orange line line corresponds to a rotation occurring at $x_* = 0.5$, yellow line line corresponds to a rotation occurring at $x_* = 0.25$, and green line line corresponds to a rotation near last scattering surface $x_* = 0.005$ - the CAMB visibility function $g_{\rm CAMB}$ is plotted in red (on a different scale); angular power spectra obtained with the modified version of CAMB are compared in (b) $C_{\ell}^{EE} - C_{\ell}^{EE}(\bar{\alpha} = 0)$, (c) C_{ℓ}^{BB} , here we plot for comparison also the signal induced by gravitational lensing (black dotted line), primordial signal for $r = 10^{-2}$ (black dashed line), and primordial signal for $r = 10^{-3}$ (black dot-dashed line) (d) $C_{\ell}^{TE} - C_{\ell}^{TE}(\bar{\alpha} = 0)$, (e) C_{ℓ}^{BB} .



FIG. 3. (a) Evolution of the birefringence angle α as a function of conformal time $(\eta - \eta_{\rm rec}) / (\eta_0 - \eta_{\rm rec})$: rotation angle equal to 1 deg (0.1 deg) from last scattering to reionization green continuous (dash-dotted line) line, rotation angle equal to 1 deg (0.1 deg) from reionization to nowadays blue continuous (dash-dotted line) line - the CAMB visibility function $g_{\rm CAMB}$ is plotted in red (on a different scale); angular power spectra obtained with the modified version of CAMB are compared in (b) $C_{\ell}^{EE} - C_{\ell}^{EE}(\bar{\alpha} = 0)$, (c) C_{ℓ}^{BB} , (d) $C_{\ell}^{TE} - C_{\ell}^{TE}(\bar{\alpha} = 0)$, (e) C_{ℓ}^{TB} , (f) C_{ℓ}^{EB} .

III. THEORY MODELING

In an expanding universe a spatially homogeneous scalar field obeys:

$$\ddot{\phi} + 3H\dot{\phi} - \frac{dV}{d\phi} = 0, \qquad (20)$$



FIG. 4. (a) Oscillating birefringence angle with $\bar{\alpha} \equiv \alpha(\eta_{\rm rec}) - \alpha(\eta_0) = 0$ deg, the linear polarization angle oscillates between 1 deg and -1 deg: $\alpha(\eta) = \sin(20\pi x)$ (green line), we plot for comparison also a the case of a sudden rotation of +/-1 deg (+/-0.1 deg) occurring at present time, see continuous (dot-dashed) red/blue line; (b) $C_{\ell}^{EE} - C_{\ell}^{EE}(\bar{\alpha} = 0)$, (c) C_{ℓ}^{BB} , (d) $C_{\ell}^{TE} - C_{\ell}^{TE}(\bar{\alpha} = 0)$, (e) C_{ℓ}^{TB} , (f) C_{ℓ}^{EB} .

where $\dot{}$ denotes derivative respect to cosmic time t. In this Section we specify the potential $V(\phi)$ for: (a) axionlike Early Dark Energy (Sect. III A), (b) Quintessence (Sect. III B), and (c) axion-like dark matter (Sect. III C). From the evolution of $\phi(t)$ we estimate the effects on CMB power spectra using a modified version of CAMB [59].

A. Axion-like as Early Dark Energy

Early Dark Energy was proposed in order to solve the tension between the local and the cosmological measurements of the Hubble parameter [27, 30, 31, 65, 66]. In this case we consider a potential of the form:

$$V(\phi) = \Lambda^4 \left(1 - \cos\frac{\phi}{f}\right)^n, \qquad (21)$$

describing the spontaneous breaking of a continuous symmetry at scale f. The evolution of the pseudoscalar field ϕ is determined by the following system of equations:

$$\begin{cases} \ddot{\phi} + 3H\dot{\phi} + \frac{n\Lambda^4}{f} \left(1 - \cos\frac{\phi}{f}\right)^{n-1} \sin\frac{\phi}{f} = 0, \\ H^2 = \frac{1}{3M_{\rm pl}^2} \left(\rho_{\rm RAD} + \rho_{\rm MAT} + \rho_{\Lambda} + \rho_{\phi}\right), \end{cases}$$
(22)

where $M_{\rm pl} = 2.43 \times 10^{18}$ GeV is the reduced Planck mass. Initially the field is frozen and acts as a cosmological constant, and it begins to oscillate when the effective mass becomes of the order of H. In practice, we solve numerically this system in the new variable $x \equiv \ln t/t_i$, from a fixed point t_i in radiation dominated era to nowadays (t_0) :

$$\begin{cases} \frac{d\Theta}{dx^{2}} + \left(\frac{3}{a}\frac{da}{dx} - 1\right)\frac{d\Theta}{dx} \\ +t_{i}^{2}e^{2x}\frac{n\Lambda^{4}}{f^{2}}\left(1 - \cos\Theta\right)^{n-1}\sin\Theta = 0, \\ \frac{da}{dx} = t_{i}e^{x}H_{i}a\left[\Omega_{\text{RAD},i}\left(\frac{a_{i}}{a}\right)^{4} + \Omega_{\text{MAT},i}\left(\frac{a_{i}}{a}\right)^{3} \\ +\Omega_{\Lambda,i} + \frac{1}{6}\frac{f^{2}}{H_{i}^{2}M_{\text{pl}}^{2}t_{i}^{2}}e^{-2x}\left(\frac{d\Theta}{dx}\right)^{2} \\ +\frac{1}{3}\frac{\Lambda^{4}}{H_{i}^{2}M_{\text{pl}}^{2}}\left(1 - \cos\Theta\right)^{n}\right]^{1/2}, \end{cases}$$
(23)

where $\Theta(t) \equiv \phi(t)/f$. In the oscillating regime, for n = 2, we approximate the evolution of Θ as a function of cosmic time with an elliptic sine (sn), see [67–69]. In particular, fixed $\Lambda = 0.417$ eV, $f = 0.05 M_{\rm pl} = 1.22 \times 10^{17}$ GeV, $\Theta_i = 1$ and $\dot{\Theta}_i = 0$ the following numerical fit for Θ is obtained [70]:

$$\Theta(\eta) \simeq \left(-6.49 \times 10^{-3} + 2.15 \times 10^{-3} \frac{\eta_0}{\eta}\right) \\ * \operatorname{sn}\left(6.35 \times 10^{-1} + 5.18 \times 10^2 \frac{\eta}{\eta_0}, \frac{1}{\sqrt{2}}\right) (24)$$

In Fig. 5 we plot this function for Θ as a function of redshift z, from recombination to z = 0.

In Fig. 6 we plot the power spectra using the evolution of $\Theta(\eta)$ directly provided by CAMB code. Note that for this model $\Theta(\eta_0)$ is negligible compared to to the values at $\eta_{\rm rec}$, since the field is quickly oscillating at z = 0. We consider two values of the coupling constant: $g_{\phi} =$



FIG. 5. Early dark energy, evolution of Θ as a function of z from recombination to nowadays fixed n = 2, $\Lambda = 0.417$ eV, $f = 0.05 M_{\rm pl} = 1.22 \times 10^{17}$ GeV, $\Theta_i = 1$ and $\dot{\Theta}_i = 0$: numerical fit of Eq. (24) (green line) and the evolution provided by CAMB-1.3.2 (blue line); in this version of CAMB EDE is implemented following [71].

 $8.17\times 10^{-18}{\rm GeV}^{-1}$ - corresponding to $\bar{\alpha}=1.15~{\rm deg}$ - and $g_{\phi}=3.5\times 10^{-19}{\rm GeV}^{-1}$ - corresponding to $\bar{\alpha}=0.05~{\rm deg}.$ For EE and TE we decided to plot the difference with the standard un-rotated spectra, $C_{\ell}^{EE}-C_{\ell}^{EE}(\bar{\alpha}=0)$ and $C_{\ell}^{TE}-C_{\ell}^{TE}(\bar{\alpha}=0),$ in order to underline the differences.

B. Axion-like as dark energy

We consider dark energy driven by an axion-like pseudo-scalar, as suggested in [72], with a potential:

$$V(\phi) = M^4 \left(1 + \cos\frac{\phi}{f}\right).$$
⁽²⁵⁾

We solve numerically the system in the new variable x, as in the previous Subsection:

$$\begin{cases} \frac{d\Theta}{dx^2} + \left(\frac{3}{a}\frac{da}{dx} - 1\right)\frac{d\Theta}{dx} - t_i^2 e^{2x}\frac{M^4}{f^2}\sin\Theta = 0,\\ \frac{da}{dx} = t_i e^x H_i a \left[\Omega_{\text{RAD},i}\left(\frac{a_i}{a}\right)^4 + \Omega_{\text{MAT},i}\left(\frac{a_i}{a}\right)^3 + \frac{1}{6}\frac{f^2}{H_i^2 M_{\text{pl}}^2 t_i^2} e^{-2x}\left(\frac{d\Theta}{dx}\right)^2 + \frac{1}{3}\frac{M^4}{H_i^2 M_{\text{pl}}^2}\left(1 + \cos\Theta\right) \right]^{1/2}.\end{cases}$$

For $M \sim 10^{-3}$ eV and $f \lesssim M_{\rm pl}$ the pseudoscalar field mimics the cosmological constant contribution. There are indications from string theory that f cannot be larger than $M_{\rm pl}$ [73, 74]. In the future, when the expansion rate of the universe will become smaller, the field will start to oscillate and the universe will become cold dark matter dominated.

The pseudoscalar field becomes dynamical only recently. By fixing $M = 1.95 \times 10^{-3}$ eV, $f = 0.265 M_{\rm pl}$, $\Theta_i = 0.25$ and $\dot{\Theta}_i = 0$ we use the following numerical fit for $\Theta(\eta)$:

$$\Theta(\eta) \simeq 0.25 + 1.468 \times 10^{-4} \exp\left[8.857 \left(\frac{\eta - \eta_{\rm rec}}{\eta_0 - \eta_{\rm rec}}\right)\right]$$

Differently from the Early Dark Energy model, discussed in the previous Subsection, here the field is not oscillating at z = 0 and it is important to consider $\Theta(\eta_0)$. Using this numerical fit we evaluate the linear polarization angular power spectra for $g_{\phi} = 1.8 \times 10^{-20} \text{GeV}^{-1}$, corresponding to a total rotation angle today $\alpha(\eta_0) =$ 0.35 deg. In some models the coupling constant between the pseudoscalar field is assumed to be proportional to the inverse of the energy breaking scale f [75, 76]:

$$\mathcal{L} \supset -\frac{C \,\alpha_{\rm EM}}{2\pi f} \phi F_{\mu\nu} \,\widetilde{F}^{\mu\nu} \,, \tag{27}$$

where $C \simeq \mathcal{O}(1)$ is a model dependent constant. Therefore we discuss also the case: $g_{\phi} = \frac{2 \alpha_{\rm EM}}{f \pi} \simeq 7.2 \times 10^{-21} \text{GeV}^{-1}$ - corresponding to $\alpha(\eta_0) = 0.14$ deg.

See Fig. 7 for the power spectra $C_{\ell}^{EE} - C_{\ell}^{EE}(\bar{\alpha} = 0)$, C_{ℓ}^{BB} , $C_{\ell}^{TE} - C_{\ell}^{TE}(\bar{\alpha} = 0)$, C_{ℓ}^{TB} , and C_{ℓ}^{EB} evaluated using CAMB.

C. Axion-like as dark matter

For axion-like field acting as dark matter [76–78] we consider the potential:

$$V(\phi) = m^2 \frac{f^2}{N^2} \left(1 - \cos\frac{\phi N}{f}\right), \qquad (28)$$

in the regime where the pseudoscalar field oscillates near the minimum. The field evolves according to [11, 79]:

$$\phi(t) = \sqrt{6\Omega_{\text{MAT}}} \frac{H_0 M_{\text{pl}}}{ma^{3/2}(t)}$$
$$\sin\left[mt\sqrt{1 - (1 - \Omega_{\text{MAT}})\left(\frac{3H_0}{2m}\right)^2}\right], (29)$$

where the evolution of the scale factor is [80]:

$$a(t) = \left(\frac{\Omega_{\text{MAT}}}{1 - \Omega_{\text{MAT}}}\right)^{\frac{1}{3}} \left[\sinh\left(\frac{3}{2}\sqrt{1 - \Omega_{\text{MAT}}}H_0t\right)\right]^{\frac{2}{3}}.$$
 (30)

Since the Boltzmann CAMB code works in conformal time we fit numerically the relation between cosmic and conformal time from recombination to today; for the ACDM model cosmological model with *Planck* 2018 estimates of cosmological parameters we obtain [53]:

$$t \simeq \frac{\eta_0}{3.5041} \left(\frac{\eta}{\eta_0}\right)^{3.09358}$$
 (31)

As in the Early dark Energy case, the field quickly oscillates at z = 0 therefore we can assume $\phi(\eta_0) \simeq 0$.



FIG. 6. Early Dark Energy (a) evolution for $\Theta \equiv \phi/f$ as a function of redshift (green line) provided by the code CAMB fixed n = 2, $\Lambda = 0.417$ eV, $f = 0.05 M_{\rm pl} = 1.22 \times 10^{17}$ GeV, $\Theta_i = 1$ and $\dot{\Theta}_i = 0$; the CAMB visibility function $g_{\rm CAMB}$ is plotted in red (on a different scale). Assuming $g_{\phi} = 8.17 \times 10^{-18} \text{GeV}^{-1}$ or $g_{\phi} = 3.5 \times 10^{-19} \text{GeV}^{-1}$ we plot the angular power spectra for (b) $C_{\ell}^{EE} - C_{\ell}^{EE}(\bar{\alpha} = 0)$, (c) C_{ℓ}^{BB} , (d) $C_{\ell}^{TE} - C_{\ell}^{TE}(\bar{\alpha} = 0)$, (e) C_{ℓ}^{EB} .



FIG. 7. Axion-like dark energy (a) evolution of $\Theta(\eta) - \Theta(\eta_0)$ as a function of conformal time $(\eta - \eta_{\rm rec}) / (\eta_0 - \eta_{\rm rec})$ is plotted in green fixed $M = 1.95 \times 10^{-3}$ eV, $f = 0.265 M_{\rm pl}$, $\Theta_i = 0.25$ and $\dot{\Theta}_i = 0$; the CAMB visibility function $g_{\rm CAMB}$ is plotted in red (on a different scale). Assuming $g_{\phi} = 1.8 \times 10^{-20} \text{GeV}^{-1}$ and $g_{\phi} = \frac{2 \,\alpha_{\rm EM}}{f \,\pi} \simeq 7.2 \times 10^{-21} \text{GeV}^{-1}$, we plot the angular power spectra for (b) $C_{\ell}^{EE} - C_{\ell}^{EE}(\bar{\alpha} = 0)$, (c) C_{ℓ}^{BB} , (d) $C_{\ell}^{TE} - C_{\ell}^{TE}(\bar{\alpha} = 0)$, (e) C_{ℓ}^{TB} , (f) C_{ℓ}^{EB} .

We plot $g_{\phi}\phi(\eta)/2$ as a function of time chosen a particular value for $m = 10^{-22} \text{eV}$ and $g_{\phi} = 10^{-14} \text{GeV}^{-1}$, see Fig. 8a. Once the source terms for scalar perturbations

in the Boltzmann code are modified inserting the new terms proportional to $\alpha(\eta) - \alpha(\eta_0)$, see Eqs. (11)-(12), the rotated power spectra are obtained. In Fig. 8 we



FIG. 8. Axion-like dark matter: (a) evolution of $g_{\phi}\phi(\eta)/2$ is plotted in green as a function of conformal time $(\eta - \eta_{\rm rec})/(\eta_0 - \eta_{\rm rec})$ fixed $m = 10^{-22} \text{eV}$ and $g_{\phi} = 10^{-14} \text{GeV}^{-1}$ (green continuous line); the CAMB visibility function $g_{\rm CAMB}$ is plotted in red (on a different scale). Assuming $g_{\phi} = 10^{-14} \text{GeV}^{-1}$ (green continuous line), $g_{\phi} = 2 \times 10^{-15} \text{GeV}^{-1}$ (light-green dashed line), we plot the angular power spectra for: (b) $C_{\ell}^{EE} - C_{\ell}^{EE}(\bar{\alpha} = 0)$, (c) C_{ℓ}^{BB} , (d) $C_{\ell}^{TE} - C_{\ell}^{TE}(\bar{\alpha} = 0)$, (e) C_{ℓ}^{TB} , (f) C_{ℓ}^{EB} .

plot $C_{\ell}^{EE} - C_{\ell}^{EE}(\bar{\alpha} = 0)$, C_{ℓ}^{BB} , $C_{\ell}^{TE} - C_{\ell}^{TE}(\bar{\alpha} = 0)$, C_{ℓ}^{TB} , and C_{ℓ}^{EB} for $m = 10^{-22} \text{eV}$ and for two different values of the coupling constant: $g_{\phi} = 10^{-14} \text{GeV}^{-1}$ - corresponding to a total rotation angle $\alpha(\eta_0) = 0.52$ deg, and $g_{\phi} = 2 \times 10^{-15} \text{GeV}^{-1}$ - corresponding to a total rotation angle $\alpha(\eta_0) = 0.103$ deg.

IV. CURRENT MEASUREMENTS AND

FORECASTS FOR FUTURE EXPERIMENTS

In this Section we discuss the status of current measurements and forecast the science capabilities of future experiments in the context of cosmological birefringence.

We will analyze various cosmological models with differ-

ent approximations by providing effective $\Delta \chi^2$ and poste-

rior probabilities for parameters by Monte Carlo Markov

The parity violating nature of the interaction generates nonzero parity-odd correlators $(C_{\ell}^{TB} \text{ and } C_{\ell}^{EB})$. We therefore consider the full theoretical data covariance ma-

Chain (MCMC) exploration.

trix:

inverse-variance weighted sum of the noise sensitivity convolved with a Gaussian beam window function for each frequency channel ν [81]:

$$N_{\ell}^{XX} = \left[\sum_{\nu} \frac{1}{N_{l\nu}^{XX}}\right]^{-1}, \qquad (33)$$

with:

(32)

$$N_{l\nu}^{XX} = \Delta_{X\nu}^2 \exp\left[l\left(l+1\right)\frac{\theta_{\rm FWHM\,\nu}^2}{8\ln 2}\right],$$
 (34)

here $X = \{T, E, B\}$, $\Delta_{X\nu}$ is the detector noise level, and $\theta_{\text{FWHM}\nu}$ is the full width half maximum (FWHM) for a given frequency channel ν .

ν [GHz]	$\theta_{\rm FWHM \nu}[{\rm arcmin}]$	$\Delta_{T\nu} \ [\mu K \operatorname{arcm.}]$	$\Delta_{P\nu} [\mu K \operatorname{arcm.}]$
78	39	9.56	13.5
89	35	8.27	11.7
100	29	6.50	9.2
119	25	5.37	7.6
140	23	4.17	5.9
166	21	4.60	6.5
195	20	4.10	5.8

TABLE I. Experimental specification for LiteBIRD: the full width half maximum $(\theta_{\text{FWHM}\nu})$ and the detector noise levels for different frequency channels [82, 83].

The noise power spectra are obtained by considering an

$$\begin{split} \bar{\mathbf{C}}_l &= \begin{pmatrix} \bar{C}_{\ell}^{TT} & \bar{C}_{\ell}^{TE} & \bar{C}_{\ell}^{TB} \\ \bar{C}_{\ell}^{TE} & \bar{C}_{\ell}^{EE} & \bar{C}_{\ell}^{EB} \\ \bar{C}_{\ell}^{TB} & \bar{C}_{\ell}^{EB} & \bar{C}_{\ell}^{BB} \end{pmatrix} \\ &= \begin{pmatrix} C_{\ell}^{TT} + N_{\ell}^{TT} & C_{\ell}^{TE} & C_{\ell}^{TB} \\ C_{\ell}^{TE} & C_{\ell}^{EE} + N_{\ell}^{EE} & C_{\ell}^{EB} \\ C_{\ell}^{TB} & C_{\ell}^{EB} & C_{\ell}^{BB} + N_{\ell}^{BB} \end{pmatrix} \end{split}$$

Following [84–86], we consider a Wishart likelihood and



FIG. 9. Estimated noise power spectrum $N_{l\nu}^{BB}$ for LiteBIRD (purple continuous line), compared with the signal induced by gravitational lensing (black dotted line), primordial signal for $r = 10^{-2}$ (black dashed line), primordial signal for $r = 10^{-3}$ (black dot-dashed line), signal induced by cosmic birefringence for $\bar{\alpha} = \pm 1 \deg$ (red continuous line), signal induced by cosmic birefringence for $\bar{\alpha} = \pm 0.1 \deg$ (red dot-dashed line).

introduce the effective χ^2_{eff} :

$$\chi_{\rm eff}^2 = \sum_{\ell} \left(2\ell + 1 \right) f_{\rm sky} \left(\frac{A}{|\bar{C}|} + \ln \frac{|\bar{C}|}{|\hat{C}|} - 3 \right) \,, \quad (35)$$

where f_{sky} denotes the observed fraction of the sky, A is defined as:

$$A = \hat{C}_{\ell}^{TT} \left(\bar{C}_{\ell}^{EE} \bar{C}_{\ell}^{BB} - (\bar{C}_{\ell}^{EB})^2 \right) + \hat{C}_{\ell}^{TE} \left(\bar{C}_{\ell}^{TB} \bar{C}_{\ell}^{EB} - \bar{C}_{\ell}^{TE} \bar{C}_{\ell}^{BB} \right) + \hat{C}_{\ell}^{TB} \left(\bar{C}_{\ell}^{TE} \bar{C}_{\ell}^{EB} - \bar{C}_{\ell}^{TB} \bar{C}_{\ell}^{EE} \right) + \hat{C}_{\ell}^{TE} \left(\bar{C}_{\ell}^{TB} \bar{C}_{\ell}^{EB} - \bar{C}_{\ell}^{TE} \bar{C}_{\ell}^{BB} \right) + \hat{C}_{\ell}^{EE} \left(\bar{C}_{\ell}^{TT} \bar{C}_{\ell}^{BB} - (\bar{C}_{\ell}^{TB})^2 \right) + \hat{C}_{\ell}^{EB} \left(\bar{C}_{\ell}^{TE} \bar{C}_{\ell}^{TB} - \bar{C}_{\ell}^{TT} \bar{C}_{\ell}^{EB} \right) + \hat{C}_{\ell}^{TB} \left(\bar{C}_{\ell}^{TE} \bar{C}_{\ell}^{EB} - \bar{C}_{\ell}^{EE} \bar{C}_{\ell}^{TB} \right) + \hat{C}_{\ell}^{EB} \left(\bar{C}_{\ell}^{TE} \bar{C}_{\ell}^{TB} - \bar{C}_{\ell}^{TT} \bar{C}_{\ell}^{EB} \right) + \hat{C}_{\ell}^{BB} \left(\bar{C}_{\ell}^{TT} \bar{C}_{\ell}^{EE} - (\bar{C}_{\ell}^{TE})^2 \right) ,$$
(36)

 $|\bar{C}|$ is the determinant of the theoretical covariance matrix, see Eq. (32):

$$|\bar{C}| = \bar{C}_{\ell}^{TT} \bar{C}_{\ell}^{EE} \bar{C}_{\ell}^{BB} + 2\bar{C}_{\ell}^{TE} \bar{C}_{\ell}^{TB} \bar{C}_{\ell}^{EB} - \bar{C}_{\ell}^{TT} \left(\bar{C}_{\ell}^{EB}\right)^2 - \bar{C}_{\ell}^{EE} \left(\bar{C}_{\ell}^{TB}\right)^2 - \bar{C}_{\ell}^{BB} \left(\bar{C}_{\ell}^{TE}\right)^2 , \qquad (37)$$

and $|\hat{C}|$ is the determinant of the observed covariance matrix:

$$|\hat{C}| = \hat{C}_{\ell}^{TT} \hat{C}_{\ell}^{EE} \hat{C}_{\ell}^{BB} + 2\hat{C}_{\ell}^{TE} \hat{C}_{\ell}^{TB} \hat{C}_{\ell}^{EB} - \hat{C}_{\ell}^{TT} \left(\hat{C}_{\ell}^{EB}\right)^2 - \hat{C}_{\ell}^{EE} \left(\hat{C}_{\ell}^{TB}\right)^2 - \hat{C}_{\ell}^{BB} \left(\hat{C}_{\ell}^{TE}\right)^2 .$$
(38)

As a representative example for the next generation of CMB polarization experiments we consider Lite (Light) satellite for the study of B-mode polarization and Inflation from cosmic background Lite Background Radiation Detection (LiteBIRD) [82, 87], selected by the Japan Aerospace Exploration Agency (JAXA) as a strategic large class mission. In Tab. I we report the LiteBIRD-like experimental specifications that we use for our forecasts. We produce simulated data for T, E by considering the inverse noise weighting of the central frequency channels in Tab. I and by assuming that the lowest and highest frequencies are used to separate the foreground emission as done in [87] (see also [88]). For the *B*-mode polarization (in addition to the instrumental noise) we include the following two sources of confusion: the lensing signal and a contribution which mimics the foreground residuals, as also done in [89]. We compare these LiteBIRD-like noise power spectrum N_{ℓ}^{BB} with the signal induced by cosmic birefringence in Fig. 9. With these settings we

consider $f_{sky} = 0.7$ and $\ell_{max} = 1350$.

As first step we consider which constant birefringence angle $\bar{\alpha}$ could be detected with this LiteBIRD-like configuration. We consider a covariance matrix \bar{C} obtained with $\bar{\alpha} = \{1, 0.5, 0.35, 0.2, 0.1, 0.01\}$ deg and using the power spectra obtained from of Eqs. (15) - (19) we estimated for some values of χ^2_{eff} , see Tab. II . The observed power spectra \hat{C}_{ℓ} correspond to the case without cosmic birefringence ($\alpha = 0$). We add, both to the theoretical and to the observed power spectra, the noise power spectra N_{ℓ}^{TT} , N_{ℓ}^{EE} and N_{ℓ}^{BB} from Eq. (33); for C_{ℓ}^{BB} we consider also the contribution of lensing and foregrounds.

A. Limits for axion-like as Early Dark Energy

We find that an axion-like field acting as Early Dark Energy (EDE) could produce a signal similar to the detection of $\bar{\alpha} = 0.35 \text{ deg}$ [48] by taking into account the

\bar{C}_{ℓ} theoretical $(\bar{\alpha}) + N_{\ell}$	\hat{C}_{ℓ} observed $+ N_{\ell}$	$\chi^2_{ m eff}$
$C_{\ell}(\bar{\alpha} = 1 \deg)$	$C_{\ell}(\alpha = 0 \deg)$	3.03×10^4
$C_{\ell}(\bar{\alpha} = 0.5 \mathrm{deg})$	$C_{\ell}(\alpha = 0 \deg)$	7.57×10^3
$C_{\ell}(\bar{\alpha} = 0.35 \mathrm{deg})$	$C_{\ell}(\alpha = 0 \deg)$	3.71×10^{3}
$C_{\ell}(\bar{\alpha} = 0.2 \deg)$	$C_{\ell}(\alpha = 0 \deg)$	1.12×10^{3}
$C_{\ell}(\bar{\alpha} = 0.1 \mathrm{deg})$	$C_{\ell}(\alpha = 0 \deg)$	3.03×10^{2}
$C_{\ell}(\bar{\alpha} = 0.01 \mathrm{deg})$	$C_{\ell}(\alpha = 0 \text{ deg})$	3.03

TABLE II. $\chi^2_{\rm eff}$, see Eq. (35), for different values of $\bar{\alpha}$: the theoretical power spectra are obtained using the analytic approximation of Eqs. (15) - (19) and they are compared to the un-rotated case ($\alpha = 0$). We assume $f_{\rm sky} = 0.7$ and $\ell_{max} = 1350$ for LiteBIRD.

redshift dependence of the scalar field with:

$$g_{\phi} \sim 1.65 \times 10^{-18} \text{GeV}^{-1}$$
, (39)

assuming, as in Section III A, n = 2, $\Lambda = 0.417$ eV, $f = 0.05 M_{\rm pl}$, $\Theta_i = 1$ and $\dot{\Theta}_i = 0$. We determine this value of g_{ϕ} by finding the C_{ℓ} obtained when taking into account the redshift dependence of the axion-like field acting as EDE which best mimics the $C_{\ell}^{\rm const}(\bar{\alpha} = 0.35 \, {\rm deg})$ in Eq. (15)-(19) by considering a minimization of $\Delta \chi^2_{\rm eff}$ in presence of the lensing BB.

Let us now turn to future experiments such as Lite-BIRD. It is important to note that a LiteBIRD-like experiment can in principle distinguish between the EDE signal induced by $g_{\phi} \sim 1.65 \times 10^{-18} \text{GeV}^{-1}$ and $C_{\ell}^{\text{const}}(\bar{\alpha} = 0.35 \text{ deg})$ at very high statistical significance, with $\chi_{\text{eff}}^2 = 67.3$ according to Tab. III. This capability of future experiments opens up the possibility to understand the physical mechanism of cosmological birefringence.

From Tab. III we retrieve other two important information. We report a value of $g_{\phi} \sim 1.4 \times 10^{-19} \text{GeV}^{-1}$ as the smallest value of the coupling which can be distinguished by a $C_{\ell}(\bar{\alpha})$ with $\bar{\alpha} = \alpha(\eta_{\text{rec}}) - \alpha(\eta_0)$ in Eq. (14); and $g_{\phi} \sim 6.0 \times 10^{-20} \text{GeV}^{-1}$ as the 95 % upper bound which a LiteBIRD-like experiment as the one we adopt can achieve.

Our results improve those obtained in Tab. I of [27] for LiteBIRD: $g_{\phi} \simeq 1.45 \times 10^{-16} \text{GeV}^{-1}$ (considering the power spectrum of the rotation angle $C_{\ell}^{\alpha\alpha}$) and $g_{\phi} \simeq 7.8 \times 10^{-17} \text{GeV}^{-1}$ (considering the cross-correlation between the rotation angle and the temperature $C_{\ell}^{\alpha T}$). See also the constraints for axionlike particles acting as Early Dark Energy discussed in [31].

B. Limits for axion-like for dark energy

In the case of axion-like field acting as dark energy we find that we should consider a coupling constant of the order:

$$g_{\phi} \sim -1.8 \times 10^{-20} \text{GeV}^{-1}$$
, (40)

in order to best mimic a birefringence signal $C_{\ell}^{\text{const}}(\bar{\alpha} = 0.35 \text{ deg})$ [48], assuming $M = 1.95 \times 10^{-3} \text{ eV}$, $f = 0.265 M_{\rm pl}$, $\Theta_i = 0.25$ and $\dot{\Theta}_i = 0$, as in Section III B.

\bar{C}_{ℓ} theoretical (EDE) $+N_{\ell}$	\hat{C}_{ℓ} observed $+N_{\ell}$	$\chi^2_{ m eff}$
$C_{\ell}(g_{\phi} = 8.17 \times 10^{-18} \text{GeV}^{-1})$	$C_{\ell}(\alpha = 0 \deg)$	1.10×10^{5}
$C_{\ell}(g_{\phi} = 1.51 \times 10^{-18} \text{GeV}^{-1})$	$C_{\ell}(\alpha = 0 \deg)$	3.81×10^3
$C_{\ell}(g_{\phi} = 4.35 \times 10^{-19} \text{GeV}^{-1})$	$C_{\ell}(\alpha = 0 \deg)$	3.21×10^{2}
$C_{\ell}(g_{\phi} = 3.5 \times 10^{-19} \text{GeV}^{-1})$	$C_{\ell}(\alpha = 0 \deg)$	2.09×10^{2}
$C_{\ell}(g_{\phi} = 6.0 \times 10^{-20} \text{GeV}^{-1})$	$C_{\ell}(\alpha = 0 \deg)$	10.5
$C_{\ell}(g_{\phi} = 1.65 \times 10^{-18} \text{GeV}^{-1})$	$C_{\ell}(\bar{\alpha}=0.35\mathrm{deg})$	67.3
$C_{\ell}(g_{\phi} = 1.4 \times 10^{-19} \text{GeV}^{-1})$	$C_{\ell}(\bar{\alpha} = 0.02 \mathrm{deg})$	9.62

TABLE III. Early Dark energy: $\chi^2_{\rm eff}$, see Eq. (35), for different values of g_{ϕ} , fixed n = 2, $\Lambda = 0.417 \, {\rm eV}$, $f = 0.05 \, M_{\rm pl} = 1.22 \times 10^{17}$ GeV, $\Theta_i = 1$ and $\dot{\Theta}_i = 0$. The theoretical power spectra are obtained using the modified version of CAMB. The observed power spectra correspond to the case without cosmic birefringence ($\alpha = 0$), except the last two lines where we consider a rotation $\bar{\alpha}$, see Eqs. (15)-(19). We assume $f_{\rm sky} = 0.7$ and $\ell_{max} = 1350$ for LiteBIRD.

We always consider a minimization of $\Delta \chi^2_{\rm eff}$ in presence of the lensing BB.

In this dark energy case the pseudoscalar field ϕ becomes dynamic at late times and therefore the linear polarization angle rotates at low redshift. The power spectra are quite similar to those obtained using the analytic approximation of Eqs. (15) - (19). The smallest coupling g_{ϕ} that a LiteBIRD-like experiment can distinguish from $C_{\ell}^{\rm const}(\bar{\alpha})$ is ~ 8.0 × 10⁻¹⁹ GeV⁻¹, always assuming $\bar{\alpha} = \alpha(\eta_{\rm rec}) - \alpha(\eta_0)$. As 95% upper bound we find $g_{\phi} \sim 9.0 \times 10^{-22} \text{GeV}^{-1}$ (see Tab. IV).

\bar{C}_{ℓ} theoretical (DE) $+N_{\ell}$	\hat{C}_{ℓ} observed $+N_{\ell}$	$\chi^2_{ m eff}$
$C_{\ell}(g_{\phi} = 1.8 \times 10^{-20} \text{GeV}^{-1})$	$C_{\ell}(\alpha = 0 \deg)$	3.78×10^{3}
$C_{\ell}(g_{\phi} = 5.2 \times 10^{-21} \text{GeV}^{-1})$	$C_{\ell}(\alpha = 0 \deg)$	3.15×10^2
$C_{\ell}(g_{\phi} = 7.2 \times 10^{-21} \text{GeV}^{-1})$	$C_{\ell}(\alpha = 0 \deg)$	6.04×10^{2}
$C_{\ell}(g_{\phi} = 9.0 \times 10^{-22} \text{GeV}^{-1})$	$C_{\ell}(\alpha = 0 \deg)$	9.4
$C_{\ell}(g_{\phi} = 8.0 \times 10^{-19} \text{GeV}^{-1})$	$C_{\ell}(\bar{\alpha} = -15.7 \mathrm{deg})$	9.8
$C_{\ell}(g_{\phi} = 1.8 \times 10^{-20} \text{GeV}^{-1})$	$C_{\ell}(\bar{\alpha} = -0.35 \mathrm{deg})$	0.30

TABLE IV. Axion like dark energy: $\chi^2_{\rm eff}$, see Eq. (35), for different values of g_{ϕ} , fixed $M = 1.95 \times 10^{-3}$ eV, $f = 0.265 M_{\rm pl}$, $\Theta_i = 0.25$ and $\dot{\Theta}_i = 0$. The theoretical power spectra are obtained using the modified version of CAMB. The observed power spectra correspond to the case without cosmic birefringence ($\alpha = 0$), except the last two lines where we consider a rotation $\bar{\alpha}$, see Eqs. (15)-(19). We assume $f_{\rm sky} = 0.7$ and $\ell_{max} = 1350$ for LiteBIRD.

C. Limits for axion-like for dark matter

For a pseudoscalar field acting as dark matter we find that

$$g_{\phi} \sim 1.37 \times 10^{-14} \text{GeV}^{-1}$$
, (41)

is needed to reproduce a signal similar to $\bar{\alpha} = 0.35$ deg [48] for $m = 10^{-22} \text{eV}$, as in Section III C, by considering a minimization of $\Delta \chi^2_{\text{eff}}$ in presence of the lensing BB.

A LiteBIRD-like experiment can easily distinguish between a birefringence signal induced by dark matter with a coupling constant of this order of magnitude and $C_{\ell}^{\rm const}(\bar{\alpha} = 0.35 \, {\rm deg})$ at very high statistical significance $\chi_{\rm eff}^2 = 69.8$, see Tab. V.

From this Table we report also a value of $g_{\phi} \sim 7.5 \times 10^{-15} \text{GeV}^{-1}$ as the smallest value of the coupling which can be distinguished by a $C_{\ell}(\bar{\alpha})$ with $\bar{\alpha} = \alpha(\eta_{\text{rec}}) - \alpha(\eta_0)$ in Eq. (14); and $g_{\phi} \sim 8.1 \times 10^{-16} \text{GeV}^{-1}$ as the 95 % upper bound which a LiteBIRD-like experiment as the one we adopt can achieve.

The above results depend on the mass of the axion. If we consider a heavier mass for the pseudoscalar field, as $m = 10^{-20}$ eV, the coupling that mimics $C_{\ell}^{\text{const}}(\bar{\alpha} = 0.35 \text{ deg})$ is smaller than the one reported in Eq. (41) and is:

$$g_{\phi} \sim 8.0 \times 10^{-13} \text{GeV}^{-1}$$
, (42)

On the contrary, for smaller masses (e.g. $m = 10^{-20}$ eV), we have to consider a smaller coupling constant of the order:

$$g_{\phi} \sim 2.7 \times 10^{-16} \text{GeV}^{-1}$$
 (43)

In Fig. 10 we compare the limits on the axion-photon coupling obtained from isotropic cosmic birefringence with the other limits present in literature [90, 91]. CMB cosmic birefringence nicely complements other experimental/astrophysical tests [92].

\bar{C}_{ℓ} theoretical (DM) $+N_{\ell}$	\hat{C}_{ℓ} observed $+N_{\ell}$	$\chi^2_{\rm eff}$
$C_{\ell}(g_{\phi} = 1.5 \times 10^{-14} \text{GeV}^{-1})$	$C_{\ell}(\alpha = 0 \deg)$	3.75×10^{3}
$C_{\ell}(g_{\phi} = 1.0 \times 10^{-14} \text{GeV}^{-1})$	$C_{\ell}(\alpha = 0 \deg)$	1.70×10^{4}
$C_{\ell}(g_{\phi} = 4.3 \times 10^{-15} \text{GeV}^{-1})$	$C_{\ell}(\alpha = 0 \deg)$	3.05×10^{2}
$C_{\ell}(g_{\phi} = 2.0 \times 10^{-15} \text{GeV}^{-1})$	$C_{\ell}(\alpha = 0 \deg)$	66.0
$C_{\ell}(g_{\phi} = 8.1 \times 10^{-16} \text{GeV}^{-1})$	$C_{\ell}(\alpha = 0 \deg)$	10.4
$C_{\ell}(g_{\phi} = 1.37 \times 10^{-14} \text{GeV}^{-1})$	$C_{\ell}(\bar{\alpha} = 0.35 \mathrm{deg})$	69.8
$C_{\ell}(g_{\phi} = 7.5 \times 10^{-15} \text{GeV}^{-1})$	$C_{\ell}(\bar{\alpha} = 0.17 \mathrm{deg})$	10.4

TABLE V. Axion like Dark Matter: $\chi^2_{\rm eff}$, see Eq. (35), for different values of g_{ϕ} , fixed $m = 10^{-22} {\rm eV}$. The theoretical power spectra are obtained using the modified version of CAMB. The observed power spectra correspond to the case without cosmic birefringence ($\alpha = 0$), except the last two lines where we consider a rotation $\bar{\alpha}$, see Eqs. (15)-(19). We assume $f_{\rm sky} = 0.7$ and $\ell_{max} = 1350$ for LiteBIRD.

D. Markov Chain MonteCarlo results

We perform few exploratory runs exploring the whole cosmological and birefringence parameter space using the MCMC code cosmomc [93, 94]. We use the exact Wishart likelihood with mock data generated following Eq. (35), with an effective sky fraction of 70% for all channels T, E, B and including the same instrumental noise (see Fig. 9) and foreground residuals in BB. In order to include the full contribution of cosmic birefringence we extended the standard exact likelihood to include also the odd cross correlators TB and EB. We perform first an idealistic case where we vary only the coupling together with the six standard parameters of the Λ CDM:dark matter density $\Omega_c h^2$, baryon density $\Omega_{\rm B} h^2$, angular diameter distance to the last scattering surface θ , optical depth τ , the scalar spectral index n_s and the amplitude of primordial fluctuations A_s . The resulting posterior probability distribution is shown in blue Fig. 11 and the 68% error bar is $\sigma(g_{\phi}M_{\rm pl}/2) = 0.032$ with a fiducial $g_{\phi} = 0$ (corresponding to $|g_{\phi}| \leq 5.3 \times 10^{-20} {\rm GeV}^{-1}$ at 2σ). Note that the degradation of a factor of few in the constraints in g_{ϕ} with respect to those quoted in Section IV is due to the variation of all the cosmological parameters in the MCMC exploration.

The result above mentioned represents an ideal case because we have not included the uncertainty due to the miscalibration angle related to the uncertainty in the calibration of polarization angles. In order to account for such uncertainty we add an isotropic rotation of the spectra due to an isotropic angle that we call $\alpha_{miscalib}$. This mimics the confusion created by not knowing the calibration angle when the birefringence signal arrives at the detectors. We vary this additional parameter assuming a Gaussian prior. We consider two cases: optimistic with a width of the prior of 0.00175 rad (= 0.1 deg = 6 arcmin) and pessimistic with a width of the prior of 0.0035 rad (= 0.2 deg = 12 arcmin). The resulting one dimensional posteriors are presented in red and black in Fig. 11, we note how with respect to the blue ideal curve we have a slight degradation of the constraints on the coupling whose uncertainty increase by roughly one order of magnitude to $\sigma(g_{\phi}M_{\rm pl}/2) = 0.35$ (corresponding to $|g_{\phi}| \lesssim 5.7 \times 10^{-19} \text{GeV}^{-1}$ at 2σ) for the optimistic and $\sigma(g_{\phi}M_{\rm pl}/2) = 0.41$ (corresponding to $|g_{\phi}| \lesssim 6.7 \times 10^{-19} \text{GeV}^{-1}$ at 2σ) for the pessimistic case. Note that the constraints on g_{ϕ} obtained by taking into account the miscalibration angle degrade by roughly one order of magnitude.

In Fig. 12 we show the correlation between the miscalibration angle and the coupling. Note that the degeneracy the miscalibration angle and the coupling is not exact as would be with the birefringence angle when the redshift dependence of the rotation angle is neglected.

V. CONCLUSIONS

We studied isotropic cosmological birefringence induced by a cosmological redshift-dependent pseudoscalar field with a coupling $g_{\phi}\phi F^{\mu\nu}\tilde{F}_{\mu\nu}$. We showed how time evolution of the background pseudoscalar field imprints in general a non-trivial multipole dependence in the observed CMB angular power spectra which is not captured by the widely adopted approximation in which the redshift dependence of the rotation angle is neglected. This effect could be important in interpreting reported hints of birefringence in the Planck CMB spectra [48, 49, 51, 52]. Beyond considering phenomenological redshift evolution for the rotation angle induced by a pseudo-scalar field,



FIG. 10. Limits on the coupling constant with photons g_{ϕ} for pseudoscalar field acting as dark matter as a function of mass m (coloured regions are excluded). The light blue dotted line corresponds to CMB birefringence of the order of 0.35 deg [48] obtained by taking into account the redshift dependency of the birefringence angle, compared to CMB birefringence limits presented in [11] (dark blue region) and in [29] (blue dashed line). Plot created with the AxionLimits code [91], we refer to online documentation for references on the other constraints.



FIG. 11. One dimensional posterior distribution for the coupling and the miscalibration angle. In blue the ideal case where only the coupling is considered. In red the optimistic case with the miscalibration angle and in black the pessimistic case.

we also considered the theoretical prediction for Early Dark Energy, Quintessence and axion-like matter.

As consequences, not only the total rotation $\bar{\alpha} = \alpha(\eta_{\rm rec}) - \alpha(\eta_0)$ determines the final CMB spectra, but also when the rotation occurs compared to the main changes in the visibility function, i.e. recombination and reionization. Moreover, non-vanishing parity violating effects occur also in the particular case $\alpha(\eta_{\rm rec}) = \alpha(\eta_0)$.

Due to the non-trivial multipole dependence induced



FIG. 12. Two dimensional posterior distribution for the miscalibration angle vs the coupling. In green the pessimistic case and in red the optimistic one.

by the redshift evolution of the pseudo-scalar field, the resulting isotropic birefringence is not degenerate with a polarization rotation angle independent on the multipoles, which is connected to a systematic calibration angle uncertainty.

For the theoretical models of EDE, DE, and axion-like DM we estimated the size of the couplings which will be detected by a LiteBIRD-like experiment by a χ^2 calculation. Moreover, always for these models and by a χ^2 calculation, we also computed at which level our theoretical predictions can be distinguished by the widely adopted approximation in which the redshift dependence of the rotation angle is neglected for a LiteBIRD-like experiment. Finally, we have explicitly shown by MCMC the reduction of the degeneracy between the isotropic birefringence effect for Early Dark Energy and the miscalibration angle by allowing all the cosmological parameters to vary, always for a LiteBIRD-like experiment.

As a next step, we will add the effects due inhomogeneities in the pseudoscalar field, i.e. anisotropic birefringence, to complete the theoretical predictions of interesting models with a pseudo-scalar field, such as Early Dark Energy, Quintessence and axion-like matter.

ACKNOWLEDGMENTS

We would like to thank A. Gruppuso and E. Komatsu for useful discussions. FF and DP acknowledge financial support by ASI Grant 2016-24-H.0 and the agreement n. 2020-9-HH.0 ASI-UniRM2. We acknowledge the use of the INAF-OAS HPC cluster.

Appendix A: Additional phenomenological power spectra

In this Appendix we discuss other phenomenological examples of redshift dependence of the linear polarization angle.

First we consider "instantaneous rotation at present time" (see Fig. 13). In this case $\bar{\alpha} = \alpha(\eta_{\rm rec}) - \alpha(\eta_0)$ is exactly constant during integration along the line-ofsight therefore the power spectra obtained using the modified CAMB code (coloured lines) exactly coincide with the $C_{\ell}^{\rm obs}$ given by the analytic expressions of Eqs. (15)-(19) (coloured regions).

The comparison between Fig. 2 and Fig. 14 assures us that the effects on the power spectra are not dominated by the slope of the tanh function describing the transition between two different values of the linear polarization angle. Instead it is very important when this transition occurs: earlier in time the rotation happens, smaller are the effects on the power spectra.

In Fig. 4 we already showed that power spectra are influenced by cosmic birefringence also when $\bar{\alpha} = \alpha(\eta_{\rm rec}) - \alpha(\eta_{\rm rec})$ $\alpha(\eta_0) = 0$. In Fig. 15 we always focus on an oscillating behaviour with $\bar{\alpha} = 0$, but we compare different frequencies. Effects on the power spectra at high- ℓ seem to increase at higher oscillating frequencies $(\alpha(\eta) = \sin(10\pi x) \text{ and } \alpha(\eta) = \sin(100\pi x))$, but for a linear polarization angle oscillating extremely quickly (e.g. $\alpha(\eta) = \sin(1000\pi x)$) the effects cancel out. Since the visibility function reaches its maximum at recombination and a second peak at reionization - see Section II for more details - the overall effects highly depend on the value of $\alpha(\eta) - \alpha(\eta_0)$ at these two epochs. In the case $\alpha(\eta) = \sin(\pi x)$ the birefringence angle is too small at recombination to modify the source terms of Eqs. (11)-(12), but effects are visible at reionization (lower- ℓ). On the contrary for $\alpha(\eta) = \sin(10\pi x)$ and $\alpha(\eta) = \sin(100\pi x)$ the effects at recombination (high- ℓ) are quite important, while the effects at low- ℓ (reionization) are wiped out by the rapid oscillations of the birefringence angle; for even faster oscillations, $\alpha(\eta) = \sin(1000\pi x)$, the effects are deleted also at recombination.

Finally, see Fig. 16, we compare three different oscillating behaviours of the linear polarization angle. In one case $\alpha(\eta_{\rm rec}) = \alpha(\eta_0)$, while in the other two cases $\alpha(\eta_{\rm rec}) \neq \alpha(\eta_0)$. When there is a difference between the value of α at recombination and today this clearly dominates the effects on the power spectra.

- S. M. Carroll, G. B. Field and R. Jackiw, "Limits on a Lorentz and Parity Violating Modification of Electrodynamics," Phys. Rev. D 41, 1231 (1990) doi:10.1103/PhysRevD.41.1231.
- [2] D. Harari and P. Sikivie, "Effects of a Nambu-Goldstone boson on the polarization of radio galaxies and the cosmic microwave background," Phys. Lett. B 289 (1992) 67.
- [3] S. M. Carroll and G. B. Field, "The Einstein equivalence principle and the polarization of radio galaxies," Phys. Rev. D 43, 3789 (1991) doi:10.1103/PhysRevD.43.3789
- [4] A. Cimatti, S. di Serego Alighieri, G. B. Field and R. A. E. Fosbury, "Stellar and scattered light in a radio galaxy at z = 2.63," Astrophys. J. **422** (1994) 562 doi:10.1086/173749.
- [5] S. M. Carroll and G. B. Field, "Is there evidence for cosmic anisotropy in the polarization of dis-

tant radio sources?," Phys. Rev. Lett. **79**, 2394-2397 (1997) doi:10.1103/PhysRevLett.79.2394 [arXiv:astro-ph/9704263 [astro-ph]].

- S. M. Carroll, "Quintessence and the rest of the world," Phys. Rev. Lett. 81, 3067-3070 (1998) doi:10.1103/PhysRevLett.81.3067 [arXiv:astroph/9806099 [astro-ph]].
- [7] S. di Serego Alighieri, F. Finelli and M. Galaverni, "Limits on Cosmological Birefringence from the UV Polarization of Distant Radio Galaxies," Astrophys. J. 715 (2010), 33-38 doi:10.1088/0004-637X/715/1/33 [arXiv:1003.4823[astro-ph.CO]].
- [8] A. Lue, L. M. Wang and M. Kamionkowski, "Cosmological signature of new parity violating interactions," Phys. Rev. Lett. 83, 1506-1509 (1999) doi:10.1103/PhysRevLett.83.1506 [arXiv:astro-



FIG. 13. (a) Evolution of $\alpha(\eta) - \alpha(\eta_0)$ as a function of conformal time: during propagation $\bar{\alpha} = +/-1$ deg (continue red/blue line), +/-0.5 deg (dashed red/blue line), +/-0.1 deg (dotted red/blue line). In this case the output of the modified CAMB code (coloured lines) coincides with the analytic approximations of Eqs. (15)-(19) (coloured regions): (b) $C_{\ell}^{EE} - C_{\ell}^{EE}(\bar{\alpha} = 0)$, (c) C_{ℓ}^{BB} , (d) $C_{\ell}^{TE} - C_{\ell}^{TE}(\bar{\alpha} = 0)$, (e) C_{ℓ}^{TB} , (f) C_{ℓ}^{EB} .



FIG. 14. Comparison between different hyperbolic tangents: (a) evolution of $\alpha(\eta) - \alpha(\eta_0)$ as a function of conformal time $x \equiv (\eta - \eta_{\rm rec}) / (\eta_0 - \eta_{\rm rec})$: $\alpha(\eta) = -1/2 \{1 + \tanh \left[10^4(x - 0.005)\right]\}$ (green continuous line), $\alpha(\eta) = -1/2 \{1 + \tanh \left[10^3(x - 0.005)\right]\}$ (green dotted line), $\alpha(\eta) = -1/2 \{1 + \tanh \left[10^4(x - 0.25)\right]\}$ (green time), $\alpha(\eta) = -1/2 \{1 + \tanh \left[20(x - 0.25)\right]\}$ (green dotted line), $\alpha(\eta) = -1/2 \{1 + \tanh \left[10^4(x - 0.25)\right]\}$ (green continuous line), $\alpha(\eta) = -1/2 \{1 + \tanh \left[20(x - 0.25)\right]\}$ (green dotted line) - the CAMB visibility function $g_{\rm CAMB}$ is plotted in red (on a different scale); angular power spectra obtained with the modified version of CAMB are compared in (b) $C_{\ell}^{EE} - C_{\ell}^{EE}(\bar{\alpha} = 0)$, (c) C_{ℓ}^{BB} , (d) $C_{\ell}^{TE} - C_{\ell}^{TE}(\bar{\alpha} = 0)$, (e) C_{ℓ}^{TB} , (f) C_{ℓ}^{EB} .

ph/9812088 [astro-ph]].

[9] B. Feng, H. Li, M. z. Li and X. m. Zhang, "Gravitational leptogenesis and its signatures in CMB," Phys. Lett. B **620**, 27-32 (2005) doi:10.1016/j.physletb.2005.06.009 [arXiv:hep-ph/0406269 [hep-ph]].

[10] B. Feng, M. Li, J. Q. Xia, X. Chen and X. Zhang,



FIG. 15. (a) Evolution of the birefringence angle α as a function of conformal time $x \equiv (\eta - \eta_{\rm rec}) / (\eta_0 - \eta_{\rm rec})$: $\alpha(\eta) = \sin(\pi x)$ (green continuous line), $\alpha(\eta) = \sin(10\pi x)$ (green dotted line), $\alpha(\eta) = \sin(100\pi x)$ (green dotted line), $\alpha(\eta) = \sin(100\pi x)$ (green dotted line), $\alpha(\eta) = \sin(100\pi x)$ (green dotted line), note that in all tree cases $\alpha(\eta_{\rm rec}) = \alpha(\eta_0)$ ($\bar{\alpha} = 0$ deg) - the CAMB visibility function $g_{\rm CAMB}$ is plotted in red (on a different scale); (b) $C_{\ell}^{EE} - C_{\ell}^{EE}(\bar{\alpha} = 0)$, (c) C_{ℓ}^{BB} , (d) $C_{\ell}^{TE} - C_{\ell}^{TE}(\bar{\alpha} = 0)$, (e) C_{ℓ}^{TB} , (f) C_{ℓ}^{EB} .



FIG. 16. (a) Evolution of the birefringence angle α as a function of conformal time $x \equiv (\eta - \eta_{\rm rec}) / (\eta_0 - \eta_{\rm rec})$: $\alpha(\eta) = \sin(9.5\pi x)$ (green continuous line, $\bar{\alpha} = 1$ deg), $\alpha(\eta) = \sin(9.75\pi x)$ (green dotted line, $\bar{\alpha} = 0.71$ deg), $\alpha(\eta) = \sin(10\pi x)$ (green dashed line, $\bar{\alpha} = 0$ deg) - we plot for comparison also a the case of a sudden rotation of +/-1 deg (+/-0.1 deg) occurring at present time, see continuous (dot-dashed) red/blue line; (b) $C_{\ell}^{EE} - C_{\ell}^{EE}(\bar{\alpha} = 0)$, (c) C_{ℓ}^{BB} , (d) $C_{\ell}^{TE} - C_{\ell}^{TE}(\bar{\alpha} = 0)$, (e) C_{ℓ}^{TB} .

"Searching for CPT Violation with Cosmic Microwave Background Data from WMAP and BOOMERANG," Phys. Rev. Lett. **96**, 221302 (2006) doi:10.1103/PhysRevLett.96.221302 [arXiv:astroph/0601095 [astro-ph]].

[11] F. Finelli and M. Galaverni, "Rotation of Linear Polarization Plane and Circular Polarization from Cosmological Pseudo-Scalar Fields," Phys. Rev. D **79** (2009), 063002 doi:10.1103/PhysRevD.79.063002 [arXiv:0802.4210 [astro-ph]].

- [12] S. Alexander, J. Ochoa and A. Kosowsky, "Generation of Circular Polarization of the Cosmic Microwave Background," Phys. Rev. D 79, 063524 (2009) doi:10.1103/PhysRevD.79.063524
- [13] G. C. Liu, S. Lee and K. W. Ng, "Effect on cosmic microwave background polarization of coupling of quintessence to pseudoscalar formed from the electromagnetic field and its dual," Phys. Rev. Lett. 97, 161303 (2006) doi:10.1103/PhysRevLett.97.161303 [arXiv:astroph/0606248 [astro-ph]].
- [14] G. Gubitosi, M. Martinelli and L. Pagano, "Including birefringence into time evolution of CMB: current and future constraints," JCAP 12, 020 (2014) doi:10.1088/1475-7516/2014/12/020 [arXiv:1410.1799 [astro-ph.CO]].
- [15] S. Chigusa, T. Moroi and K. Nakayama, "Signals of Axion Like Dark Matter in Time Dependent Polarization of Light," Phys. Lett. B 803, 135288 (2020) doi:10.1016/j.physletb.2020.135288 [arXiv:1911.09850 [astro-ph.CO]].
- [16] B. D. Sherwin and T. Namikawa, "Cosmic birefringence tomography and calibration-independence with reionization signals in the CMB," [arXiv:2108.09287 [astroph.CO]].
- [17] H. Nakatsuka, T. Namikawa and E. Komatsu, "Is cosmic birefringence due to dark energy or dark matter? A tomographic approach," Phys. Rev. D 105, no.12, 123509 (2022) doi:10.1103/PhysRevD.105.123509 [arXiv:2203.08560 [astro-ph.CO]].
- [18] A. Greco, N. Bartolo and A. Gruppuso, "Probing Axions through Tomography of Anisotropic Cosmic Birefringence," [arXiv:2211.06380 [astro-ph.CO]].
- [19] M. Li and X. Zhang, "Cosmological CPT violating effect on CMB polarization," Phys. Rev. D 78, 103516 (2008) doi:10.1103/PhysRevD.78.103516 [arXiv:0810.0403 [astro-ph]].
- [20] M. Pospelov, A. Ritz and C. Skordis, "Pseudoscalar perturbations and polarization of the cosmic microwave background," Phys. Rev. Lett. **103**, 051302 (2009) doi:10.1103/PhysRevLett.103.051302 [arXiv:0808.0673 [astro-ph]].
- M. Kamionkowski, "How to De-Rotate the Cosmic Microwave Background Polarization," Phys. Rev. Lett. 102, 111302 (2009) doi:10.1103/PhysRevLett.102.111302
 [arXiv:0810.1286 [astro-ph]].
- [22] H. Cai and Y. Guan, "Computing microwave background polarization power spectra from cosmic birefringence," Phys. Rev. D 105, no.6, 063536 (2022) doi:10.1103/PhysRevD.105.063536 [arXiv:2111.14199 [astro-ph.CO]].
- [23] A. Greco, N. Bartolo and A. Gruppuso, "Cosmic birefrigence: cross-spectra and cross-bispectra with CMB anisotropies," JCAP 03, no.03, 050 (2022) doi:10.1088/1475-7516/2022/03/050 [arXiv:2202.04584 [astro-ph.CO]].
- [24] G. Gubitosi, L. Pagano, G. Amelino-Camelia, A. Melchiorri and A. Cooray, "A Constraint on Planck-scale Modifications to Electrodynamics with CMB polarization data," JCAP 08, 021 (2009) doi:10.1088/1475-7516/2009/08/021 [arXiv:0904.3201 [astro-ph.CO]].
- [25] G. C. Liu and K. W. Ng, "Axion Dark Matter Induced Cosmic Microwave Background B-modes," Phys. Dark

Univ. **16**, 22-25 (2017) doi:10.1016/j.dark.2017.02.004 [arXiv:1612.02104 [astro-ph.CO]].

- [26] G. Sigl and P. Trivedi, "Axion-like Dark Matter Constraints from CMB Birefringence," [arXiv:1811.07873 [astro-ph.CO]].
- [27] L. M. Capparelli, R. R. Caldwell and A. Melchiorri, "Cosmic birefringence test of the Hubble tension," Phys. Rev. D 101, no.12, 123529 (2020) doi:10.1103/PhysRevD.101.123529 [arXiv:1909.04621 [astro-ph.CO]].
- [28] T. Fujita, Y. Minami, K. Murai and H. Nakatsuka, "Probing axionlike particles via cosmic microwave background polarization," Phys. Rev. D 103, no.6, 063508 (2021) doi:10.1103/PhysRevD.103.063508 [arXiv:2008.02473 [astro-ph.CO]].
- [29] M. A. Fedderke, P. W. Graham and S. Rajendran, "Axion Dark Matter Detection with CMB Polarization," Phys. Rev. D 100, no.1, 015040 (2019) doi:10.1103/PhysRevD.100.015040 [arXiv:1903.02666 [astro-ph.CO]].
- [30] K. Murai, F. Naokawa, T. Namikawa and E. Komatsu, "Isotropic cosmic birefringence from early dark energy," [arXiv:2209.07804 [astro-ph.CO]].
- [31] T. Fujita, K. Murai, H. Nakatsuka and S. Tsujikawa, "Detection of isotropic cosmic birefringence and its implications for axionlike particles including dark energy," Phys. Rev. D 103, no.4, 043509 (2021) doi:10.1103/PhysRevD.103.043509 [arXiv:2011.11894 [astro-ph.CO]].
- [32] E. Komatsu *et al.* [WMAP Collaboration], "Seven-year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Interpretation," Astrophys. J. Suppl. **192**, 18 (2011) doi:10.1088/0067-0049/192/2/18 [arXiv:1001.4538 [astro-ph.CO]].
- [33] A. Gruppuso, P. Natoli, N. Mandolesi, A. De Rosa, F. Finelli and F. Paci, "WMAP 7 year constraints on CPT violation from large angle CMB anisotropies," JCAP 02, 023 (2012) doi:10.1088/1475-7516/2012/02/023 [arXiv:1107.5548 [astro-ph.CO]].
- [34] F. Finelli, A. De Rosa, A. Gruppuso and D. Paoletti, "Cosmological Parameters from a re-analysis of the WMAP-7 low resolution maps," Mon. Not. Roy. Astron. Soc. 431, 2961 (2013) doi:10.1093/mnras/stt142 [arXiv:1207.2558 [astro-ph.CO]].
- [35] A. Gruppuso, M. Gerbino, P. Natoli, L. Pagano, N. Mandolesi, A. Melchiorri and D. Molinari, "Constraints on cosmological birefringence from Planck and Bicep2/Keck data," JCAP 06, 001 (2016) doi:10.1088/1475-7516/2016/06/001 [arXiv:1509.04157 [astro-ph.CO]].
- [36] N. Aghanim *et al.* [Planck], "Planck intermediate results. XLIX. Parity-violation constraints from polarization data," Astron. Astrophys. **596**, A110 (2016) doi:10.1051/0004-6361/201629018 [arXiv:1605.08633 [astro-ph.CO]].
- [37] P. A. R. Ade *et al.* [POLARBEAR], "A Measurement of the Cosmic Microwave Background *B*-Mode Polarization Power Spectrum at Sub-Degree Scales from 2 years of POLARBEAR Data," Astrophys. J. **848**, no.2, 121 (2017) doi:10.3847/1538-4357/aa8e9f [arXiv:1705.02907 [astro-ph.CO]].
- [38] W. L. K. Wu, L. M. Mocanu, P. A. R. Ade, A. J. Anderson, J. E. Austermann, J. S. Avva, J. A. Beall, A. N. Bender, B. A. Benson and F. Bianchini, *et al.*

"A Measurement of the Cosmic Microwave Background Lensing Potential and Power Spectrum from 500 deg² of SPTpol Temperature and Polarization Data," Astrophys. J. **884**, 70 (2019) doi:10.3847/1538-4357/ab4186 [arXiv:1905.05777 [astro-ph.CO]].

- [39] A. Gruppuso, D. Molinari, P. Natoli and L. Pagano, "Planck 2018 constraints on anisotropic birefringence and its cross-correlation with CMB anisotropy," JCAP 11, 066 (2020) doi:10.1088/1475-7516/2020/11/066 [arXiv:2008.10334 [astro-ph.CO]].
- [40] P. A. R. Ade *et al.* [BICEP/Keck], "BICEP/Keck XII: Constraints on axionlike polarization oscillations in the cosmic microwave background," Phys. Rev. D 103, no.4, 042002 (2021) doi:10.1103/PhysRevD.103.042002 [arXiv:2011.03483 [astro-ph.CO]].
- [41] P. A. R. Ade *et al.* [BICEP/Keck], "BICEP/Keck XIV: Improved constraints on axionlike polarization oscillations in the cosmic microwave background," Phys. Rev. D **105**, no.2, 022006 (2022) doi:10.1103/PhysRevD.105.022006 [arXiv:2108.03316 [astro-ph.CO]].
- [42] T. Namikawa, Y. Guan, O. Darwish, B. D. Sherwin, S. Aiola, N. Battaglia, J. A. Beall, D. T. Becker, J. R. Bond and E. Calabrese, *et al.* "Atacama Cosmology Telescope: Constraints on cosmic birefringence," Phys. Rev. D **101**, no.8, 083527 (2020) doi:10.1103/PhysRevD.101.083527 [arXiv:2001.10465 [astro-ph.CO]].
- [43] M. Bortolami, M. Billi, A. Gruppuso, P. Natoli and L. Pagano, "Planck constraints on cross-correlations between anisotropic cosmic birefringence and CMB polarization," JCAP 09, 075 (2022) doi:10.1088/1475-7516/2022/09/075 [arXiv:2206.01635 [astro-ph.CO]].
- [44] K. N. Abazajian *et al.* [CMB-S4], "CMB-S4 Science Book, First Edition," [arXiv:1610.02743 [astro-ph.CO]].
- [45] D. Molinari, A. Gruppuso and P. Natoli, "Constraints on parity violation from ACTpol and forecasts for forthcoming CMB experiments," Phys. Dark Univ. 14, 65-72 (2016) doi:10.1016/j.dark.2016.09.006 [arXiv:1605.01667 [astro-ph.CO]].
- [46] S. Hanany *et al.* [NASA PICO], "PICO: Probe of Inflation and Cosmic Origins," [arXiv:1902.10541 [astroph.IM]].
- [47] L. Pogosian, M. Shimon, M. Mewes and B. Keating, "Future CMB constraints on cosmic birefringence and implications for fundamental physics," Phys. Rev. D 100, no.2, 023507 (2019) doi:10.1103/PhysRevD.100.023507 [arXiv:1904.07855 [astro-ph.CO]].
- [48] Y. Minami and E. Komatsu, "New Extraction of the Cosmic Birefringence from the Planck 2018 Polarization Data," Phys. Rev. Lett. **125**, no.22, 221301 (2020) doi:10.1103/PhysRevLett.125.221301 [arXiv:2011.11254 [astro-ph.CO]].
- [49] P. Diego-Palazuelos, J. R. Eskilt, Y. Minami, M. Tristram, R. M. Sullivan, A. J. Banday, R. B. Barreiro, H. K. Eriksen, K. M. Górski and R. Keskitalo, *et al.* "Cosmic Birefringence from the Planck Data Release 4," Phys. Rev. Lett. **128**, no.9, 091302 (2022) doi:10.1103/PhysRevLett.128.091302 [arXiv:2201.07682 [astro-ph.CO]].
- [50] Y. Akrami et al. [Planck], "Planck intermediate results. LVII. Joint Planck LFI and HFI data processing," Astron. Astrophys. 643, A42 (2020) doi:10.1051/0004-6361/202038073 [arXiv:2007.04997 [astro-ph.CO]]

- [51] J. R. Eskilt and E. Komatsu, "Improved constraints on cosmic birefringence from the WMAP and Planck cosmic microwave background polarization data," Phys. Rev. D 106, no.6, 063503 (2022) doi:10.1103/PhysRevD.106.063503 [arXiv:2205.13962 [astro-ph.CO]].
- [52] E. Komatsu, "New physics from polarised light of the cosmic microwave background," Nature Rev. Phys. 4, no.7, 452-469 (2022) doi:10.1038/s42254-022-00452-4 [arXiv:2202.13919 [astro-ph.CO]].
- [53] N. Aghanim *et al.* [Planck], "Planck 2018 results. VI. Cosmological parameters," Astron. Astrophys. **641**, A6 (2020) [erratum: Astron. Astrophys. **652**, C4 (2021)] doi:10.1051/0004-6361/201833910 [arXiv:1807.06209 [astro-ph.CO]].
- [54] M. Galaverni, G. Gubitosi, F. Paci and F. Finelli, "Cosmological birefringence constraints from CMB and astrophysical polarization data," JCAP 08 (2015), 031 doi:10.1088/1475-7516/2015/08/031 [arXiv:1411.6287 [astro-ph.CO]].
- [55] M. Galaverni, "Testing New Physics with Polarized Light: Cosmological Birefringence," Astrophys. Space Sci. Proc. 51, 165-174 (2018) doi:10.1007/978-3-319-67205-2.11.
- [56] S. di Serego Alighieri, "The conventions for the polarization angle," Exper. Astron. 43, no.1, 19-22 (2017) doi:10.1007/s10686-016-9517-y [arXiv:1612.03045 [astroph.IM]].
- [57] A. Kosowsky and A. Loeb, "Faraday rotation of microwave background polarization by a primordial magnetic field," Astrophys. J. 469 (1996), 1-6 doi:10.1086/177751 [arXiv:astro-ph/9601055 [astro-ph]].
- [58] U. Seljak and M. Zaldarriaga, "A Line of sight integration approach to cosmic microwave background anisotropies," Astrophys. J. 469 (1996), 437-444 doi:10.1086/177793 [arXiv:astro-ph/9603033 [astro-ph]].
- [59] A. Lewis, A. Challinor and A. Lasenby, "Efficient computation of CMB anisotropies in closed FRW models," Astrophys. J. 538, 473-476 (2000) doi:10.1086/309179 [arXiv:astro-ph/9911177 [astro-ph]].
- [60] A. Gruppuso, G. Maggio, D. Molinari and P. Natoli, "A note on the birefringence angle estimation in CMB data analysis," JCAP 05, 020 (2016) doi:10.1088/1475-7516/2016/05/020 [arXiv:1604.05202 [astro-ph.CO]].
- [61] Y. Minami, H. Ochi, K. Ichiki, N. Katayama, E. Komatsu and T. Matsumura, "Simultaneous determination of the cosmic birefringence and miscalibrated polarization angles from CMB experiments," PTEP 2019, no.8, 083E02 (2019) doi:10.1093/ptep/ptz079 [arXiv:1904.12440 [astro-ph.CO]].
- [62] B. G. Keating, M. Shimon, and A. P. S. Yadav, "Selfcalibration of Cosmic Microwave Background Polarization Experiments," Astrophys. J. Letters **762** (2013) L23 doi:10.1088/2041-8205/762/2/L23.
- [63] Y. Minami and E. Komatsu, "Simultaneous determination of the cosmic birefringence and miscalibrated polarization angles II: Including cross frequency spectra," PTEP 2020, no.10, 103E02 (2020) doi:10.1093/ptep/ptaa130 [arXiv:2006.15982 [astroph.CO]]
- [64] T. Namikawa, "CMB mode coupling with isotropic polarization rotation," Mon. Not. Roy. Astron. Soc. 506, no.1, 1250-1257 (2021) doi:10.1093/mnras/stab1796 [arXiv:2105.03367 [astro-ph.CO]].

- [65] V. Poulin, T. L. Smith, D. Grin, T. Karwal and M. Kamionkowski, "Cosmological implications of ultralight axionlike fields," Phys. Rev. D 98, no.8, 083525 (2018) doi:10.1103/PhysRevD.98.083525 [arXiv:1806.10608 [astro-ph.CO]].
- [66] V. Poulin, T. L. Smith, T. Karwal and M. Kamionkowski, "Early Dark Energy Can Resolve The Hubble Tension," Phys. Rev. Lett. **122**, no.22, 221301 (2019) doi:10.1103/PhysRevLett.122.221301 [arXiv:1811.04083 [astro-ph.CO]].
- [67] P. B. Greene, L. Kofman, A. D. Linde and A. A. Starobinsky, "Structure of resonance in preheating after inflation," Phys. Rev. D 56, 6175-6192 (1997) doi:10.1103/PhysRevD.56.6175 [arXiv:hepph/9705347 [hep-ph]].
- [68] F. Finelli and R. H. Brandenberger, "Parametric amplification of gravitational fluctuations during reheating," Phys. Rev. Lett. 82, 1362-1365 (1999) doi:10.1103/PhysRevLett.82.1362 [arXiv:hepph/9809490 [hep-ph]].
- [69] M. Abramowitz and I. A. Stegun, "Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables," 1964 New York, USA Dover, 1964 New York.
- [70] In order to insert the elliptic sine $\operatorname{sn}\left(x, \frac{1}{\sqrt{2}}\right)$ in the CAMB code we applied twice descending Landen transformation [69, Sect. 16.12]:

$$\operatorname{sn}(x,m) = \frac{\left(1 + \mu_1^{1/2}\right) \operatorname{sn}\left(\frac{x}{1 + \mu_1^{1/2}}, \mu_1\right)}{1 + \mu_1^{1/2} \operatorname{sn}^2\left(\frac{x}{1 + \mu_1^{1/2}}, \mu_1\right)}$$

where:

$$\mu_1 \equiv \left(\frac{1-\sqrt{1-m}}{1+\sqrt{1-m}}\right)^2$$

Starting from $m = 1/\sqrt{2} \simeq 0.707$ we obtained first $\mu_1 \simeq 0.089$, and applying the Landen transformation a second time $\mu_2 \simeq 0.00053$. Finally, since we reduced to a case where $m \ll 1$, we used the approximation of the elliptic sine in terms of circular functions [69, Sect. 16.12]:

$$\operatorname{sn}(x,m) \simeq \sin x - \frac{m}{4} (x - \sin x \cos x) \cos x$$

- [71] T. L. Smith, V. Poulin and M. A. Amin, "Oscillating scalar fields and the Hubble tension: a resolution with novel signatures," Phys. Rev. D 101, no.6, 063523 (2020) doi:10.1103/PhysRevD.101.063523 [arXiv:1908.06995 [astro-ph.CO]].
- [72] J. A. Frieman, C. T. Hill, A. Stebbins and I. Waga, "Cosmology with ultralight pseudo Nambu-Goldstone bosons," Phys. Rev. Lett. **75**, 2077-2080 (1995) doi:10.1103/PhysRevLett.75.2077 [arXiv:astroph/9505060 [astro-ph]].
- [73] M. Dine, "Dark matter and dark energy: A physicist's perspective," arXiv:hep-th/0107259.
- [74] T. Banks, M. Dine, P. J. Fox and E. Gorbatov, "On the possibility of large axion decay constants," JCAP 0306, 001 (2003) [arXiv:hep-th/0303252].
- [75] P. Sikivie, "Experimental Tests of the Invisible Axion," Phys. Rev. Lett. 51, 1415-1417 (1983) [erratum: Phys. Rev. Lett. 52, 695 (1984)] doi:10.1103/PhysRevLett.51.1415

- [76] G. G. Raffelt, "Stars as laboratories for fundamental physics: The astrophysics of neutrinos, axions, and other weakly interacting particles," Chicago Univ. Pr., 1996.
- [77] E. W. Kolb and M. S. Turner, "The Early Universe," Front. Phys. 69, 1-547 (1990) doi:10.1201/9780429492860.
- [78] P. Sikivie, "Axion Cosmology," Lect. Notes Phys.
 741, 19-50 (2008) doi:10.1007/978-3-540-73518-2_2
 [arXiv:astro-ph/0610440 [astro-ph]].
- [79] M. Galaverni and F. Finelli, "Rotation of linear polarization plane from cosmological pseudoscalar fields," Nucl. Phys. B Proc. Suppl. **194**, 51-56 (2009) doi:10.1016/j.nuclphysbps.2009.07.021.
- [80] A. Gruppuso and F. Finelli, "Analytic results for a flat universe dominated by dust and dark energy," Phys. Rev. D 73, 023512 (2006) doi:10.1103/PhysRevD.73.023512 [arXiv:astro-ph/0512641 [astro-ph]].
- [81] F. Finelli *et al.* [CORE], "Exploring cosmic origins with CORE: Inflation," JCAP **04**, 016 (2018) doi:10.1088/1475-7516/2018/04/016 [arXiv:1612.08270 [astro-ph.CO]].
- [82] M. Hazumi, P. A. R. Ade, Y. Akiba, D. Alonso, K. Arnold, J. Aumont, C. Baccigalupi, D. Barron, S. Basak and S. Beckman, *et al.* "LiteBIRD: A Satellite for the Studies of B-Mode Polarization and Inflation from Cosmic Background Radiation Detection," J. Low Temp. Phys. **194**, no.5-6, 443-452 (2019) doi:10.1007/s10909-019-02150-5.
- [83] D. Paoletti and F. Finelli, "Constraints on primordial magnetic fields from magnetically-induced perturbations: current status and future perspectives with Lite-BIRD and future ground based experiments," JCAP 11, 028 (2019) doi:10.1088/1475-7516/2019/11/028 [arXiv:1910.07456 [astro-ph.CO]].
- [84] R. Easther, W. H. Kinney and H. Peiris, "Observing trans-Planckian signatures in the cosmic microwave background," JCAP 05, 009 (2005) doi:10.1088/1475-7516/2005/05/009 [arXiv:astro-ph/0412613 [astro-ph]].
- [85] J. Q. Xia, H. Li, G. B. Zhao and X. Zhang, "Probing for the Cosmological Parameters with PLANCK Measurement," Int. J. Mod. Phys. D 17, 2025-2048 (2009) doi:10.1142/S0218271808013698 [arXiv:0708.1111 [astroph]].
- [86] J. Q. Xia, H. Li, X. l. Wang and X. m. Zhang, "Testing CPT Symmetry with CMB Measurements," Astron. Astrophys. 483, 715-718 (2008) doi:10.1051/0004-6361:200809410 [arXiv:0710.3325 [hep-ph]].
- [87] E. Allys *et al.* [LiteBIRD], "Probing Cosmic Inflation with the LiteBIRD Cosmic Microwave Background Polarization Survey," [arXiv:2202.02773 [astro-ph.IM]].
- [88] F. Finelli *et al.* [CORE], "Exploring cosmic origins with CORE: Inflation," JCAP **04**, 016 (2018) doi:10.1088/1475-7516/2018/04/016 [arXiv:1612.08270 [astro-ph.CO]].
- [89] D. Paoletti, F. Finelli, J. Valiviita and M. Hazumi, "Planck and BICEP/Keck Array 2018 constraints on primordial gravitational waves and perspectives for future B-mode polarization measurements," [arXiv:2208.10482 [astro-ph.CO]].
- [90] R. L. Workman *et al.* [Particle Data Group], "Review of Particle Physics," PTEP **2022**, 083C01 (2022) doi:10.1093/ptep/ptac097.
- [91] C. O'Hare, Zenodo (2020), https://cajohare.github. io/AxionLimits/ doi:10.5281/zenodo.3932430.

[arXiv:astro-

- [92] M. Galaverni, "Redshift dependent cosmic birefringence from axion-like dark matter," PoS ICHEP2022, 126 doi:10.22323/1.414.0126.
- [93] A. Lewis and S. Bridle, "Cosmological parameters from CMB and other data: A Monte Carlo approach," Phys. Rev. D 66 (2002), 103511

doi:10.1103/PhysRevD.66.103511 ph/0205436 [astro-ph]].

 [94] A. Lewis, "Efficient sampling of fast and slow cosmological parameters," Phys. Rev. D 87 (2013) no.10, 103529 doi:10.1103/PhysRevD.87.103529 [arXiv:1304.4473 [astro-ph.CO]].