# RATIONAL CURVES IN A QUADRIC THREEFOLD VIA AN $SL(2, \mathbb{C})$ -REPRESENTATION

## KIRYONG CHUNG, SUKMOON HUH, AND SANG-BUM YOO

ABSTRACT. In this paper, we regard the smooth quadric threefold  $Q_3$  as Lagrangian Grassmannian and search for fixed rational curves of low degree in  $Q_3$  with respect to a torus action, which is the maximal subgroup of the special linear group  $SL(2, \mathbb{C})$ . Most of them are confirmations of very well-known facts. If the degree of a rational curve is 3, it is confirmed using the Lagrangian's geometric properties that the moduli space of twisted cubic curves in  $Q_3$  has a specific projective bundle structure. From this, we can immediately obtain the cohomology ring of the moduli space.

## 1. Introduction

1.1. **Motivation and results.** By definition, a *prime Fano threefold* X is a smooth projective variety of dimension 3 with  $\operatorname{Pic}_{\mathbb{Z}}(X) = \langle K_X \rangle$  and the anti-canonical divisor  $-K_X$  is an ample divisor. Such threefolds X are classified to be one of the following:

$$\mathbb{P}^3$$
,  $Q_3$ ,  $V_5$ , and  $V_{22}$ .

It is well known that the first three  $\mathbb{P}^3$ ,  $Q_3$  and  $V_5$  are rigid but the last one  $V_{22}$  forms a 6-dimensional family; see the excellent paper [Don08, Section 5.3]. All of theses varieties have the same cohomology groups as  $\mathbb{P}^3$  does. Furthermore, the  $SL(2,\mathbb{C})$ -orbit descriptions of these prime Fano threefolds are well studied in the papers [MU83, AF93].

The reason for writing this paper is to fill in the missing list for the space of rational curves. The moduli space of rational curves of degree 3 or less in the prime Fano variety has been studied very diversely ([MU83, PS85, EPS87, Ili94, Sch01, VX02, KS04, CK11, CHK12, San14]). In the case of  $\mathbb{P}^3$ , starting from the analysis of the Hilbert scheme [PS85], it served as a key example of enumerative geometry in the 80s and 90s ([ES96]) and created many variant spaces related with birational geometry ([VX02, EPS87]). In the case of quintic del Pezzo threefold, the description of the space of rational curves was possible from the Grassmannian geometry of lines ([Ili94, San14, CHL18]). Lastly, in the case of Mukai variety, its construction is very fantastic ([MU83, Sch01]), and it plays a very decisive role in relation to the existence of the Kähler-Einstein metric on Fano manifold. For specific conclusions, refer to the papers [Don08] and [CS18]. In this paper, we examine

<sup>2020</sup> Mathematics Subject Classification. 14E05, 14E15, 14M15.

Key words and phrases. Rational curves, Torus invariant curve, Lagrangian Grassmannian, Projective bundle.

the moduli space of twisted cubic curves in a quadric threefold. The authors think that a description of its moduli space is just not written down in the literature because it is well-known and easy to experts. However, it seems necessary to record global geometry and cohomological ring in order to serve as concrete examples required by enumerative geometry ([Cao19, CMT18, CLW21]). Now let us introduce the main contents of this paper. Let X be a projective variety with a fixed embedding  $X \subset \mathbb{P}^r$ .

- Let  $S_d(X)$  be the moduli space of *stable* sheaves F in X with Hilbert polynomial  $\chi(F(m)) = dm + 1$ .
- Let  $\mathbf{H}_d(X)$  be the Hilbert scheme of curves C in X with  $\chi(\mathcal{O}_C(m)) = dm + 1$ .

In this paper, we focus on the case that  $X = Q_3$  is a quadric threefold.  $Q_3$  is homogeneous and thus one can apply the main results of [CK11, CHK12]. Specially,  $\mathbf{S}_d(Q_3)$  is smooth for  $d \leq 3$ . Also the quadric threefold  $Q_3$  does not contain any plane and so  $\mathbf{S}_3(Q_3)$  is isomorphic to  $\mathbf{H}_3(Q_3)$ . We find the torus invariant rational curves of the lower degrees and extend it into the cubic curve case. It turns out that the torus fixed loci are isolated until degree 2, but not in degree 3.

**Proposition 3.11** (Proposition 3.11 and Proposition 3.15). The torus invariant cubic curves in  $Q_3$  consists of isolated ones (32) and two connected components (isomorphic to  $\mathbb{P}^1$ ).

Through the computation of torus invariant cubic curves, one realize that the cubic curves in the hyperplane section of  $Q_3$  need to be ordered depending on its linear class. By using the universal family of the Fano scheme of lines in  $Q_3$ , we obtain a global description of  $S_3(Q_3)$  as follows.

**Theorem 1.2** (Proposition 4.4). The moduli space  $S_3(Q_3)$  of stable sheaves in  $Q_3$  is isomorphic to a projectivized rank 6 bundle over Gr(2,4).

The key point of the proof of Theorem 1.2 is to give the algebraic meaning of the ruling line of the hyperplane section of  $Q_3$ . From Theorem 1.2, one can easily obtain the cohomology ring of  $S_3(Q_3)$  (Corollary 4.6).

The method of the proof of Theorem 1.2 is a very natural approach to study the space of curves in prime Fano threefolds. For example, the moduli space of rational quartic curves in  $Q_3$  is birational to the space of cubic curves in  $\mathbb{P}^3$  by using Gauss map or tangent developable surfaces ([FGP19, the proof of Proposition 4.17]). The first named author is willing to study of birational relations among these moduli spaces in the following paper.

1.2. **Organization of the paper.** In Section 2, we collect the well-known facts for finding torus invariant curves. In Section 3, we apply the Białynicki–Birula theorem to the moduli space  $S_d(Q_3)$  for  $d \leq 3$ . Lastly, in Section 4, we describe the moduli space  $S_3(Q)$  as a projective bundle.

## Notations and conventions.

- Let us denote by Gr(k, n) the Grassmannian variety parameterizing k-dimensional subspaces in a fixed vector space V with  $\dim V = n$ .
- When no confusion can arise, we do not distinguish the moduli point  $[x] \in \mathcal{M}$  from the object x parameterized by [x].
- The set of fixed points of X is denoted by  $X^{\mathbb{C}^*}$  under the  $\mathbb{C}^*$ -action.

**Acknowledgements.** The authors gratefully acknowledge the many helpful suggestions and comments of Jeong-Seop Kim, Yeongrak Kim, Wanseok Lee, Kyeong-Dong Park, Joonyeong Won during the preparation of the paper. A part of the paper has been initiated in the workshop (held in Jinju, Korea, Feb. 2022) aimed for finding research topics through arXiv preprint survey.

#### 2. Preliminary

In this section, we introduce a general theory about a geometric structure of a smooth projective variety with a torus action. Also we collect the well-known facts about the multiple structure on Cohen-Macaulay curves.

2.1. **Białynicki—Birula (BB) Theorem.** Let X be a smooth projective variety with a  $\mathbb{C}^*$ -action. Then the  $\mathbb{C}^*$ -fixed locus of X decomposes into

$$X^{\mathbb{C}^*} = \bigsqcup_{i} Y_i$$

such that each component  $Y_i$  is connected. Note that  $Y_i$  is smooth ([Ive72]). For each  $Y_i$ , the  $\mathbb{C}^*$ -action on the tangent bundle  $TX|_{Y_i}$  provides a decomposition as

$$TX|_{Y_i} = T^+ \oplus T^0 \oplus T^-$$

where  $T^+$ ,  $T^0$  and  $T^-$  are the subbundles of  $TX|_{Y_i}$  such that the group  $\mathbb{C}^*$  acts with positive, zero and negative weights respectively. Under the local linearization,  $T^0 \cong TY_i$  and

$$T^+ \oplus T^- = N_{Y_i/X} = N^+(Y_i) \oplus N^-(Y_i)$$

is the decomposition of the normal bundle  $N_{Y_i/X}$  of  $Y_i$  in X.

A fundamental result in a theory of  $\mathbb{C}^*$ -action on X has been provided by A. Białynicki-Birula. Let

$$X^+(Y_i) = \left\{ x \in X \mid \lim_{t \to 0} t \cdot x \in Y_i \right\} \quad \text{and} \quad X^-(Y_i) = \left\{ x \in X \mid \lim_{t \to \infty} t \cdot x \in Y_i \right\}.$$

**Theorem 2.1** ([BB73]). *Under the above assumptions and notations,* 

(1) 
$$X = \bigsqcup_{i} X^{\pm}(Y_i);$$

2

(2) for each connected component  $Y_i$ , there are  $\mathbb{C}^*$ -equivariant isomorphism  $X^{\pm}(Y_i) \cong N^{\pm}(Y_i)$  over  $Y_i$  where  $N^{\pm}(Y_i) \to Y_i$  is a Zariski locally trivial fiberation with the affine space  $\mathbb{C}^{\mu^{\pm}(Y_i)}$  of dimension  $\mu^{\pm}(Y_i) := \operatorname{rank} N^{\pm}(Y_i)$ .

Throughout this article, we denote a smooth quadric threefold simply by  $Q_3 = Q$ .

**Proposition 2.2.** The moduli space  $S_d(Q) \cong H_d(Q)$  is smooth for  $d \leq 3$ .

*Proof.* The case d=1 is easily verified by the normal bundle sequence of  $L \subset Q \subset \mathbb{P}^4$ . The cases d=2 and d=3 have been proved in [CHK12, Theorem 3.7 and Proposition 4.13].

- 2.2. Non-reduced cubic curves in a quadric threefold. Note that Q is defined by a quadratic polynomial and does not contain any plane. Hence each curve C with Hilbert polynomial  $\chi(\mathcal{O}_C(m)) = 3m+1$  is Cohen-Macaulay (CM). Let us start by recalling the list of the non-reduced CM curve C in a quadric threefold Q; see [EH82] and [Chu22, Lemma 2.1]. Let  $p_a(C) := \dim H^1(C, \mathcal{O}_C)$  be the *arithmetic genus* of the curve C.
  - (i) (Triple thickness) The structure sheaf  $\mathcal{O}_C$  with [C]=3[L] fits into the non-split extension

$$0 \to \mathcal{O}_L(-1) \oplus \mathcal{O}_L(-1) \to \mathcal{O}_C \to \mathcal{O}_L \to 0.$$

Moreover, such C's are parameterized by the GIT-quotient:

- (1)  $\mathbb{P}(\operatorname{Ext}_{O}^{1}(\mathcal{O}_{L}, \mathcal{O}_{L}(-1) \oplus \mathcal{O}_{L}(-1)))^{s} / / \operatorname{SL}(2) \cong \operatorname{Gr}(2, \operatorname{dim}\operatorname{Ext}_{O}^{1}(\mathcal{O}_{L}, \mathcal{O}_{L}(-1))).$
- (ii) (Triple line lying on a quadric cone) The structure sheaf  $\mathcal{O}_C$  with [C]=3[L] fits into the non-split extension

$$0 \to \mathcal{O}_L(-1) \to \mathcal{O}_C \to \mathcal{O}_{L^2} \to 0$$

where  $p_a(L^2) = 0$ .

(iii) (Pair of lines) The structure sheaf  $\mathcal{O}_C$  with [C]=2[L]+[L'] fits into the non-split extension

$$0 \to \mathcal{O}_L(-1) \to \mathcal{O}_C \to \mathcal{O}_{L \cup L'} \to 0$$
,

where  $p_a(L^2) = 0$  or -1.

**Remark 2.3.** More generally, let G be a stable sheaf on a projective space  $\mathbb{P}^n$  with  $\chi(G(m))=(d-1)m+1$  and L be a line. Then every sheaf F fitting into the non-split short exact sequence

$$0 \to \mathcal{O}_L(-1) \to F \to G \to 0$$

is stable; see [CCM14, Lemma 4.7].

## 2.3. **Deformation theory.** We address an exact sequence which we will use later.

**Lemma 2.4.** Let  $Y \stackrel{i}{\hookrightarrow} X$  be a smooth, closed subvariety of the smooth variety X. If F and  $G \in Coh(Y)$ , then there is an exact sequence

(2) 
$$0 \to \operatorname{Ext}_{Y}^{1}(F,G) \to \operatorname{Ext}_{X}^{1}(i_{*}F, i_{*}G) \to \operatorname{Hom}_{Y}(F, G \otimes N_{Y/X}) \\ \to \operatorname{Ext}_{Y}^{2}(F,G) \to \operatorname{Ext}_{X}^{2}(i_{*}F, i_{*}G).$$

*Proof.* This is the base change spectral sequence in [McC00, Theorem 12.1].

#### 3. APPLICATION OF THE BB-THEOREM ON A QUADRIC THREEFOLD

In this section we find the fixed rational curves and its weights in Q under the  $\mathbb{C}^*$ -action which inherits from the  $\mathrm{SL}_2 := \mathrm{SL}(2,\mathbb{C})$ -representation. Let us regard the quadric threefold Q as a hyperplane section of  $\mathrm{Gr}(2,4)$ . The  $\mathbb{C}^*$ -actions on Q and curves in there come from an action on its related vector space.

Let  $V_d = \operatorname{Sym}^d(\mathbb{C}^2)$  with the  $\operatorname{SL}_2$ -action induced from the left multiplication of  $\operatorname{SL}_2$  on  $\mathbb{C}^2$ . Then the maximal torus  $\{\operatorname{diag}(t^{-1},t)\mid t\in\mathbb{C}^*\}\cong\mathbb{C}^*$  acts on the basis vectors

$$\{v_d = x^d, v_{d-2} = x^{d-1}y, \dots, v_{-1} = xy^{2-d}, v_{-d} = y^d\}$$

with weights  $(d, d-2, \ldots, 2-d, -d)$ . The infinitesimal  $\mathfrak{sl}_2$ -action on  $V_d$  is given by

$$e = x\partial_y, \quad f = y\partial_x, \quad h = x\partial_x - y\partial_y$$

for the standard basis  $\langle e, f, h \rangle$  for  $\mathfrak{sl}_2$ . Indeed, we have the following equations for  $0 \le i \le d$ :

$$\begin{cases} e \cdot v_{d-2i} = i v_{d-2(i-1)}, \\ f \cdot v_{d-2i} = (d-i) v_{d-2(i+1)}, \\ h \cdot v_{d-2i} = (d-2i) v_{d-2i}. \end{cases}$$

Now fix d=3 and set  $V=V_3$ . Setting  $v_{a,b}:=v_a^*\wedge v_b^*\in \wedge^2V^*$  and  $W:=\wedge^2V$ , we get the Plücker coordinates for  $i:\operatorname{Gr}(2,V)\hookrightarrow \mathbb{P}(W)$ 

$$v_{3,1}, v_{3,-1}, v_{3,-3}, v_{1,-1}, v_{1,-3}, v_{-1,-3}$$

with weights (4, 2, 0, 0, -2, -4) and the defining equation of Gr(2, V) is given by the Klein relation:

(3) 
$$v_{3,1}v_{-1,-3} - v_{3,-1}v_{1,-3} + v_{3,-3}v_{1,-1} = 0.$$

Note that for  $g \in \mathfrak{sl}_2$ ,  $g(v_{i,j}) = (g \cdot v_i^*) \wedge v_j^* + v_i^* \wedge (g \cdot v_j^*)$  for  $i \neq j$ . For example, g = e acts on the basis of W by

$$e \cdot v_{3,1} = 0,$$
  $e \cdot v_{3,-1} = 2v_{3,1},$   $e \cdot v_{3,-3} = 3v_{3,-1}$   $e \cdot v_{1,-1} = v_{3,-1},$   $e \cdot v_{1,-3} = v_{3,-3} + 3v_{1,-1},$   $e \cdot v_{-1,-3} = 2v_{1,-3}.$ 

By the action of  $h \in \mathfrak{sl}_2$ , each  $v_i$  is an eigenvector of V so that we get a decomposition

$$V = V(3) \oplus V(1) \oplus V(-1) \oplus V(-3)$$

such that  $V(i) = \langle v_i \rangle$  for each  $i \in \{3, 1, -1, -3\}$ . Similarly we get a decomposition for W:

$$W = W(4) \oplus W(2) \oplus W(0) \oplus W(-2) \oplus W(-4),$$

where  $W(4) = \langle v_{3,1}^* \rangle$ ,  $W(2) = \langle v_{3,-1}^* \rangle$ ,  $W(0) = \langle v_{3,-3}^*, v_{1,-1}^* \rangle$  and  $W(-2) = \langle v_{1,-3}^* \rangle$  and  $W(-4) = \langle v_{-1,-3}^* \rangle$ .

**Remark 3.1.** Let  $H = \mathbb{P}(W_1) \cong \mathbb{P}^4$  be the  $\mathrm{SL}_2$ -invariant hyperplane of  $\mathbb{P}(W)$  for a five-dimensional subspace  $W_1 \subset W$ . Then it must be defined by an equation  $av_{3,-3} + bv_{1,-1} = 0$  for some  $a, b \in \mathbb{C}$ . In particular, we have

$$e \cdot (av_{3,-3} + bv_{1,-1}) = (3a+b)v_{3,-1} = 0.$$

Thus *H* is defined by an linear equation  $v_{3,-3} - 3v_{1,-1} = 0$ .

Unless otherwise stated, let us define the smooth quadric threefold Q by  $Q = Gr(2, V) \cap H$  for the hyperplane H in Remark 3.1.

3.1. **Fixed points and lines in** Q**.** The fixed points  $Q^{\mathbb{C}^*}$  of Q are the following four points:

(4) 
$$p_{3,1} = [1:0:0:0:0:0], \quad p_{3,-1} = [0:1:0:0:0:0], \\ p_{1,-3} = [0:0:0:0:1:0], \quad p_{-1,-3} = [0:0:0:0:0:1].$$

Note that the fixed points  $H^{\mathbb{C}^*}$  consists of 5 points, including  $Q^{\mathbb{C}^*}$ . From the defining equation of H, the fifth fixed point must be

(5) 
$$q_0 = [0:0:3:1:0:0].$$

Therefore one can get a decomposition  $W=W_1\oplus W_1^\perp$  so that

$$\mathbb{P}(W_1) = \langle p_{3,1}, p_{3,-1}, p_{1,-3}, p_{-1,-3}, q_0 \rangle,$$

where  $\langle S \rangle$  denotes the linear span for a subset  $S \subset \mathbb{P}(W)$ .

**Lemma 3.2.** There exist exactly four lines on Q, fixed by the action of  $\mathbb{C}^*$ :

- (a)  $\overline{p_{3,1}p_{3,-1}} = [s:t:0:0:0:0]$ ,
- (b)  $\overline{p_{3,1}p_{1,-3}} = [s:0:0:0:t:0],$
- (c)  $\overline{p_{3,-1}p_{-1,-3}} = [0:s:0:0:0:t]$ , and
- (d)  $\overline{p_{1,-3}p_{-1,-3}} = [0:0:0:0:s:t].$

*Proof.* Let  $L(\cong \mathbb{P}^1)$  be a line in Q fixed by the  $\mathbb{C}^*$ -action. Then one can regard that the torus  $\mathbb{C}^*$  acts on L. But  $\chi(L)=2$  and thus two points on L are fixed. So we obtain the four lines as listed.

Type	Weights of	Weights of	Number of negative weights of	
	$T_{[L]}\mathrm{Gr}(2,W_1)$	$\mathrm{H}^0(\mathcal{O}_L(2))$	$T_{[L]}\mathbf{S}_1(Q)$	
(a)	-8, -6, -6, -4, -4, -2	-8, -6, -4	3	
(b)	-8, -4, -2, -2, 2, 4	-8, -2, 4	2	
(c)	-4, -2, 2, 2, 4, 8	-4, 2, 8	1	
(d)	2, 4, 4, 6, 6, 8	4, 6, 8	0	

TABLE 1. Weights of  $T_{[L]}\mathbf{S}_1(Q)$ 

From the inclusion  $L \subset Q \subset \mathbb{P}(W_1)$  and Lemma 2.4, we have an exact sequence:

(6) 
$$0 \to T_{[L]}\mathbf{S}_1(Q) \to T_{[L]}\mathrm{Gr}(2, W_1) \to \mathrm{H}^0(\mathcal{O}_L(2)) \to 0$$

for each line  $L \subset Q$ . The last term is zero since  $S_1(Q)$  is smooth with dim  $S_1(Q) = 3$ .

**Remark 3.3** (Table 1). One can compute the number  $\delta([L])$  of negative weights  $T_{[L]}\mathbf{S}_1(Q)$  for each case in Lemma 3.2.

(a) The weights of the fixed line  $L = \mathbb{P}(\mathbb{C}^2)$  are (4, 2). From the isomorphism

$$T_{[L]}\mathrm{Gr}(2,W_1) \cong \mathrm{Hom}(\mathbb{C}^2,W_1/\mathbb{C}^2) = (\mathbb{C}^2)^{\vee} \otimes (W_1/\mathbb{C}^2)$$

we get that the weight of  $T_{[L]}Gr(2, W_1)$  is given by (-8, -6, -6, -4, -4, -2). Since  $H^0(\mathcal{O}_L(2)) \cong \operatorname{Sym}^2(\mathbb{C}^2)^\vee$  has the weight (-8, -6, -4), the weight of  $T_{[L]}\mathbf{S}_1(Q)$  is (-6, -4, -2). In particular, we get  $\delta(L) = 3$ .

- (b) Similarly as in (a), we get that the weight of the fixed line  $L = \mathbb{P}(\mathbb{C}^2)$  is (4, -2) and so the weight of  $T_{[L]}\mathbf{S}_1(Q)$  is (-4, -2, 2) and so  $\delta(L) = 2$ .
- (c) and (d): Again as in the above, we get  $\delta(L) = 1$  (resp. 0) for the case (c) (resp. (d)).

Let  $P(X,t) = \sum_{i=0}^{2 \dim X} h^i(X,\mathbb{C}) t^i$  be the Poincaré polynomial of a smooth projective variety X.

**Corollary 3.4.** The Poincaré polynomial of  $S_1(Q)$  is

$$P(\mathbf{S}_1(Q), t) = 1 + t^2 + t^4 + t^6.$$

*Proof.* The proof is straightforward by Theorem 2.1 and the result of Table 1.

## 3.2. Fixed conics in Q and its weights.

**Lemma 3.5.** There exist exactly 10 conics on Q, fixed under the action of  $\mathbb{C}^*$ . Furthermore, the defining equation of fixed conics are given by

(1-a) 
$$\langle v_{1,-3}, v_{-1,-3}, v_{3,-3} - 3v_{1,-1}, v_{3,-3}v_{1,-1} \rangle$$
,

(1-b) 
$$\langle v_{3,-1}, v_{-1,-3}, v_{3,-3} - 3v_{1,-1}, v_{3,-3}v_{1,-1} \rangle$$
,

(1-c) 
$$\langle v_{3,1}, v_{1,-3}, v_{3,-3} - 3v_{1,-1}, v_{3,-3}v_{1,-1} \rangle$$
,




FIGURE 1. Fixed conics in *Q* 

$$(1-d) \langle v_{3,1}, v_{3,-1}, v_{3,-3} - 3v_{1,-1}, v_{3,-3}v_{1,-1} \rangle,$$

$$(2-a) \langle v_{-1,-3}, v_{3,-3}, v_{1,-1}, v_{3,-1}v_{1,-3} \rangle$$

$$(2-b) \langle v_{1,-3}, v_{3,-3}, v_{1,-1}, v_{3,1}v_{-1,-3} \rangle$$
,

$$(2-c) \langle v_{3,-1}, v_{3,-3}, v_{1,-1}, v_{3,1}v_{-1,-3} \rangle$$
,

$$(2-d) \langle v_{3,1}, v_{3,-3}, v_{1,-1}, v_{3,-1}v_{1,-3} \rangle$$
,

(3-a) 
$$\langle v_{3,1}, v_{-1,-3}, v_{3,-3} - 3v_{1,-1}, v_{3,-1}v_{1,-3} - v_{3,-3}v_{1,-1} \rangle$$
, and

(3-b) 
$$\langle v_{3,-1}, v_{1,-3}, v_{3,-3} - 3v_{1,-1}, v_{3,1}v_{-1,-3} + v_{3,-3}v_{1,-1} \rangle$$
.

*Proof.* Let  $[C] \in \mathbf{S}_2(Q)$  be a fixed conic. Then the linear span  $\langle C \rangle \cong \mathbb{P}^2$  is also invariant under  $\mathbb{C}^*$ -action. The  $\mathbb{C}^*$ -fixed planes are generated by the point  $q_0$  in equation (5) and two different points in  $Q^{\mathbb{C}^*}$ , or three different points in  $Q^{\mathbb{C}^*}$ . From these planes, one obtain the list.

(1-a), (1-b), (1-c) and (1-d) of the list of Lemma 3.5 are the *unique* double lines supported on (a), (b), (c) and (d) respectively in Lemma 3.2. Also (2-a), (2-b), (2-c) and (2-d) in Lemma 3.5 are pairs of two lines supported on the fixed lines in Lemma 3.2. Lastly, (3-a) (resp. (3-b)) in Lemma 3.5 are smooth conics passing through the points  $p_{3,-1}$  and  $p_{1,-3}$  (resp.  $p_{3,1}$  and  $p_{-1,-3}$ ). As combining these ones, we obtain its configuration (Figure 1).

Note that  $\mathbf{H}_2(W_1)$  is isomorphic to  $\mathbb{P}(\mathrm{Sym}^2(\mathcal{U}^{\vee}))$  where  $\mathcal{U}$  is the universal subbundle of  $\mathrm{Gr}(3,W_1)$ . Hence one can apply Lemma 2.4 to get an exact sequence

(7) 
$$0 \to T_{[C]}\mathbf{S}_2(Q) \to T_{[C]}\mathbb{P}(\operatorname{Sym}^2(\mathcal{U}^{\vee})) \to \operatorname{H}^0(\mathcal{O}_C(2)) \to 0$$

for any conic  $C \subset Q$ . In equation (7), the last term is zero since  $S_2(Q)$  is smooth and  $\dim S_2(Q) = 6$ .

Type	Weights of	Weights of	Number of negative weights of	
	$T_{[C]}\mathbb{P}(\operatorname{Sym}^2(\mathcal{U}^{\vee}))$	$\mathrm{H}^0(\mathcal{O}_C(2))$	$T_{[C]}\mathbf{S}_2(Q)$	
(1-a)	-8,-6,-6,-4,-4,-2,-8,-6,-4,-4,-2	-8,-6,-4,-4,-2	6	
(1-b)	-8,-4,-2,-2,2,4,-8,-4,-2,2,4	-8,-4,-2,2,4	4	
(1-c)	-4,-2,2,2,4,8,-4,-2,2,4,8	-4,-2,2,4,8	2	
(1-d)	2,4,4,6,6,8,2,4,4,6,8	2,4,4,6,8	0	
(2-a)	-8,-6,-4,-2,-2,2,-8,-6,-4,-2,4	-8,-6,-4,-2,4	5	
(2-b)	-6,-4,-4,-2,2,4,-8,-6,-4,2,8	-8,-6,-4,2,8	4	
(2-c)	-4,-2,2,4,4,6,-8,-2,4,6,8	-8,-2,4,6,8	2	
(2-d)	-2,2,2,4,6,8,-4,2,4,6,8	-4,2,4,6,8	1	
(3-a)	-6,-4,-2,2,4,6,-4,-2,0,2,4	-4,-2,0,2,4	3	
(3-b)	-6,-2,-2,2,2,6,-8,-4,0,4,8	-8,-4,0,4,8	3	

TABLE 2. Weights of  $T_{[C]}\mathbf{S}_2(Q)$ 

**Remark 3.6.** The case (3-a) of Lemma 3.5:  $\mathcal{U}_C = \langle v_{3,-1}, 3v_{3,-3} + v_{1,-1}, v_{1,-3} \rangle$  has weights (2,0,-2) and

(8) 
$$T_{[C]}\mathbb{P}(\operatorname{Sym}^{2}(\mathcal{U}^{\vee})) = \operatorname{Hom}(\mathcal{U}_{C}, W_{1}/\mathcal{U}_{C}) \oplus (\operatorname{Sym}^{2}(\mathcal{U}_{C}^{\vee})/\mathbb{C}).$$

The first (resp. second) term in equation (8) has weights (-6, -4, -2, 2, 4, 6) (resp. (-4, -2, 0, 2, 4)). On the other hand, from the short exact sequence  $0 \to \mathcal{O}_{\mathbb{P}(\mathcal{U}_C)} \to \mathcal{O}_{\mathbb{P}(\mathcal{U}_C)}(2) \to \mathcal{O}_C(2) \to 0$ , we have an exact sequence

$$0 \to \mathrm{H}^0(\mathcal{O}_{\mathbb{P}(\mathcal{U}_C)}) \to \mathrm{H}^0(\mathcal{O}_{\mathbb{P}(\mathcal{U}_C)}(2)) \to \mathrm{H}^0(\mathcal{O}_C(2)) \to 0.$$

Since  $H^0(\mathcal{O}_{\mathbb{P}(\mathcal{U}_C)}(2)) = \operatorname{Sym}^2(\mathcal{U}_C^*)$ ,  $H^0(\mathcal{O}_C(2))$  has weights (-4, -2, 0, 2, 4). Therefore  $T_{[C]}\mathbf{S}_2(Q)$  has weights (-6, -4, -2, 2, 4, 6). The other cases are presented in Table 2.

**Corollary 3.7.** The Poincaré polynomial of  $S_2(Q)$  is

$$P(\mathbf{S}_2(Q), t) = 1 + t^2 + 2t^4 + 2t^6 + 2t^8 + t^{10} + t^{12}.$$

*Proof.* The proof is straightforward by Theorem 2.1 and the result of Table 2.  $\Box$ 

3.3. **Fixed components of**  $S_3(Q)$ : **Degenerated case.** In this subsection, the fixed degenerate cubic curve is found using the fixed lines and conics (Lemma 3.2 and Lemma 3.5). Furthermore, we also find fixed smooth cubics component by analyzing cubic curves lying in the quadric cone (Proposition 3.15).

**Lemma 3.8.** Every Cohen-Macaulay curve  $C \in \mathbf{S}_3(Q)$  with [C] = 3[L] is a triple line lying on a quadric cone (Subsection 2.2).

*Proof.* Since dimExt $_Q^1(\mathcal{O}_L, \mathcal{O}_L(-1)) = \dim H^0(N_{L|Q}(-1)) = 1$  for any line L in Q by Lemma 2.4, the GIT-quotient in equation (1) is an empty set and so the assertion follows.

**Lemma 3.9.** Let L be a  $\mathbb{C}^*$ -invariant line and  $L^2$  be the unique planar double line in Q supporting L. Then we have

$$\dim \operatorname{Ext}_Q^1(\mathcal{O}_{L^2}, \mathcal{O}_L(-1)) = 2.$$

Furthermore, the induced action on the extension space is non-trivial.

*Proof.* The dimension of the extension space can be easily checked by using Macaulay2 ([GS]). By taking  $\operatorname{Ext}_Q^{\bullet}(-,\mathcal{O}_L(-1))$  into the short exact sequence  $0 \to \mathcal{O}_L(-1) \to \mathcal{O}_{L^2} \to \mathcal{O}_L \to 0$ , we have a long exact sequence

(9) 
$$0 \cong \operatorname{Hom}_{Q}(\mathcal{O}_{L^{2}}, \mathcal{O}_{L}(-1)) \to \operatorname{Hom}_{Q}(\mathcal{O}_{L}(-1), \mathcal{O}_{L}(-1)) \stackrel{\cong}{\to} \operatorname{Ext}_{Q}^{1}(\mathcal{O}_{L}, \mathcal{O}_{L}(-1)) \\ \to \operatorname{Ext}_{Q}^{1}(\mathcal{O}_{L^{2}}, \mathcal{O}_{L}(-1)) \to \operatorname{Ext}_{Q}^{1}(\mathcal{O}_{L}(-1), \mathcal{O}_{L}(-1)) \to \cdots.$$

The equality of the first term in (9) holds due to the stability. Also the second isomorphism in (9) holds by the dimension counting. But the last term in (9) is isomorphic to the tangent space  $T_{[L]}\mathbf{S}_1(Q)$ . By Table 1, there does not exist the reduplication in the space  $\mathrm{Ext}^1_O(\mathcal{O}_L(-1),\mathcal{O}_L(-1))$  and thus the induced action is non-trivial.

**Lemma 3.10.** Let  $C = L_1 \cup L_2$  be a fixed conic in Q. Then we have

$$\dim \operatorname{Ext}_Q^1(\mathcal{O}_C, \mathcal{O}_{L_2}(-1)) = 2.$$

Furthermore, the induced action on the extension space is non-trivial.

*Proof.* One can prove the claim by the same method in the proof of Lemma 3.9.

From this discussion, one can present all of the  $\mathbb{C}^*$ -invariant degenerated cubic curves in  $S_3(Q)$  as follows.

**Proposition 3.11.** The number of the  $\mathbb{C}^*$ -invariant, degenerated cubic curves in  $\mathbf{S}_3(Q)$  is 36.

*Proof.* We read the invariant cubics from Figure 1. The reduced cubic cases are a tree of three lines or a conic attached a line. From Figure 1, one can read such cases are  $4+2^3=12$ . When the support of cubic is a pair of lines, then our choice is  $4 \times 2 = 8$ . For each case, by Lemma 3.10, there are two cases. Hence the possible case is 16. Lastly, the support of triple line lying on a quadric cone has 4 cases and by Lemma 3.9, we have two cases for each of them. After of all, we have 12+16+8=36.

3.4. **Fixed components of**  $S_3(Q)$ **: Smooth case.** To find the smooth invariant cubic curves, we consider the *enveloping map* 

$$\xi: \mathbf{S}_3(Q) \to \mathrm{Gr}(4, W_1)$$

defined by  $[C] \mapsto \langle C \rangle \cong \mathbb{P}^3 \subset \mathbb{P}(W_1) = H$ . As did in the case of  $\mathbf{S}_2(Q)$ , we can use the five fixed points  $Q^{\mathbb{C}^*} \cup \{q_0\}$  to obtain the  $\mathbb{C}^*$ -fixed points in  $\mathrm{Gr}(4,W_1)$ . The result is the second row of the Table 3. Let us find the fixed twisted cubics in  $\xi^{-1}(H_0) \subset \mathbf{S}_3(Q)$  for a  $\mathbb{C}^*$ -invariant element  $[H_0] \in \mathrm{Gr}(4,W_1)$ . That is, we seek the  $\mathbb{C}^*$ -invariant cubics in the

(i)	(ii)	(iii)	(iv)	(v)
$v_{3,-3} = v_{1,-1} = 0$	$v_{-1,-3} = H = 0$	$v_{1,-3} = H = 0$	$v_{3,-1} = H = 0$	$v_{3,1} = H = 0$
Smooth quadric surface		Quadric cone		

TABLE 3. Fixed hyperplane section.

quadric surface  $S_0 := Q \cap H_0$ . Since the cases (ii) and (iii) is symmetric to (iv) and (v) in Table 3, we treat the first two cases only.

## • Case (i) in the Table 3

**Proposition 3.12.** Under the notations in the above, the  $\mathbb{C}^*$ -fixed cubics with  $S_0$  smooth, are always reducible.

*Proof.* We fall into the case (i) of Table 3, in which we can let

$$[v_{3,1}:v_{3,-1}:v_{1,-3}:v_{-1,-3}]$$

be the homogenous coordinates of  $\mathbb{P}^3$ . Then the smooth quadric surface  $S_0 \subset \mathbb{P}^3$  is defined by  $v_{3,1}v_{-1,-3}-v_{3,-1}v_{1,-3}=0$  induced from the Klein relation of  $\mathrm{Gr}(2,V)$ . Set

$$\mathbb{P}^1 \times \mathbb{P}^1 \to S_0 \subset \mathbb{P}^3$$
,  $[s:t] \times [v:w] \mapsto [sv:sw:tv:tw]$ 

be the  $\mathbb{C}^*$ -equivariant map where  $\mathbb{C}^*$  acts on  $\mathbb{P}^1 \times \mathbb{P}^1$  diagonally with weights (1,-1,3,-3). Then we get  $\mathbf{S}_3(S_0) \cong |\mathcal{O}_{S_0}(1,2)| \sqcup |\mathcal{O}_{S_0}(2,1)|$ . The first component of the latter space is isomorphic to the projective space  $\mathbb{P}\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(1) \boxtimes \mathcal{O}_{\mathbb{P}^1}(2))$ , and by the Künneth formula we get

$$\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(1)\boxtimes\mathcal{O}_{\mathbb{P}^1}(2))\cong\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(1))\otimes\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(2))=\mathbb{C}_{s,t}\otimes\mathrm{Sym}^2\mathbb{C}_{v,w}.$$

Hence the weights for  $|\mathcal{O}_{S_0}(1,2)|$  are  $(\pm 7, \pm 5, \pm 1)$  and thus the fixed points are coordinate points; we get a similar description for the second component.

On the other hand, let  $\{sv^2, svw, sw^2, tv^2, tvw, tw^2\}$  be a basis for  $H^0(\mathcal{O}_{S_0}(1,2))$  and its dual basis by  $\{h_0, h_1, \dots, h_5\}$ . Then each element in the system  $|\mathcal{O}_{S_0}(1,2)|$  is written as

$$F := h_0 s v^2 + h_1 s v w + h_2 s w^2 + h_3 t v^2 + h_4 t v w + h_5 t w^2.$$

Using the homogenous coordinates of  $\mathbb{P}^3$ 

$$v_{3,1} = sv, v_{3,-1} = sw, v_{1-3} = tv, v_{-1,-3} = tw,$$

we can write

$$sF = h_0(sv)^2 + h_1s^2vw + h_2(sw)^2 + h_3(sv)(tv) + h_4stvw + h_5(sw)(tw)$$

$$= h_0v_{3,1}^2 + h_1v_{3,1}v_{3,-1} + h_2v_{3,-1}^2 + h_3v_{3,1}v_{1-3} + h_4v_{3,-1}v_{1-3} + h_5v_{3,-1}v_{-1,-3};$$

$$tF = h_0(sv)(tv) + h_1tsvw + h_2(sw)(tw) + h_3(tv)^2 + h_4t^2vw + h_5(tw)^2$$

$$= h_0v_{3,1}v_{1-3} + h_1v_{3,-1}v_{1-3} + h_2v_{3,-1}v_{-1,-3} + h_3v_{1-3}^2 + h_4v_{1-3}v_{-1,-3} + h_5v_{-1,-3}^2.$$

Thus the defining equation of a cubic curve in  $S_0$  is given by  $\langle sF, tF \rangle$ . In particular, one can read that  $\mathbb{C}^*$ -fixed cubics are not irreducible.

#### • Case (ii) and (iii) in the Table 3

In these cases, the quadric surface  $S_0$  is singular. A very natural invariant curves can be obtained by taking the closure of the  $\mathbb{C}^*$ -orbit of the point  $[1:a:3:1:3a^{-1}:0]$ ,  $a \neq 0$ . That is, one parameter family of twisted cubics

(10) 
$$C_a := \{ [u^3 : au^2v : 3uv^2 : uv^2 : 3a^{-1}v^3 : 0] \times (a) \mid [u : v] \in \mathbb{P}^1, a \in \mathbb{C}^* \} \subset \mathbb{P}^5 \times \mathbb{C}_a^* \}$$

parameterized by  $a \in \mathbb{C}_a^*$  is invariant under the  $\mathbb{C}^*$ -action. In fact, such a family gives a connected component in  $\mathbf{S}_3(Q)^{\mathbb{C}^*}$ . Note that  $\mathbb{C}_a^*$  is compactified as  $\mathbb{P}^1 \subset \mathbf{S}_3(Q)$  by adding the two flat limits: A triple line lying on quadric cone and a conic attached a line.

**Example 3.13.** One can easily see that the family  $C_a$  in equation (10) lies in the case (ii) (i.e.,  $S_0 = Q \cap \{v_{-1,-3} = 0\}$ ) of Table 3. We compute the number of zero-weights of  $T_{[C]}\mathbf{S}_3(Q)$  at  $C := \mathcal{C} \mid_{a=1}$ . Let us denote the number of zero-weights of the global section  $H^0(F)$  of a locally free sheaf F by  $w_0(F)$ . Let  $f : \mathbb{P}^1 \to Q \subset \mathbb{P}(W_1) \subset \mathbb{P}(W)$  be a morphism defined by

$$[u:v] \mapsto [u^3:u^2v:3uv^2:uv^2:3v^3:0].$$

If we put the weights (1,-1) of the coordinate of the domain curve and shift the weights of the codomain by -1, then the map f is a  $\mathbb{C}^*$ -equivariant one. We compute  $w_0(f^*T_Q)$  of the space  $H^0(f^*T_Q)$  which is the first deformation space of the graph space  $Map(\mathbb{P}^1,Q)$ . Since the map f is an embedding, we have a short exact sequence

$$(11) 0 \to f^*T_Q \to f^*T_{\mathbb{P}^4} \to f^*N_{Q|\mathbb{P}^4} \to 0.$$

But we also have an exact sequence

$$0 \to f^* \mathcal{O}_{\mathbb{P}^4} \to f^* \mathcal{O}_{\mathbb{P}^4}(1) \otimes \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^4}(1))^* \to f^* T_{\mathbb{P}^4} \to 0$$

by pulling back the Euler sequence along the map f. Therefore the number of zero-weights of  $f^*T_{\mathbb{P}^4}$  is  $w_0(f^*T_{\mathbb{P}^4})=3$ . Also  $w_0(f^*N_{Q|\mathbb{P}^4})=w_0(\mathcal{O}_{\mathbb{P}^1}(6))=1$ . By plugging into equation (11), we have  $w_0(f^*T_Q)=2$ . Finally our space is the quotient space by  $\mathrm{SL}_2=\mathrm{Aut}(\mathbb{P}^1)$ . Since  $w_0(T_{\mathbb{P}^1})=1$ , the number of zero-weights of  $T_{[C]}\mathbf{S}_3(Q)$  is one.

We prove now that smooth cubic curves fixed by  $\mathbb{C}^*$ -action are only the given one in equation (10). Our strategy is to find all of the fixed curves in the resolved surface where the resolution map is  $\mathbb{C}^*$ -equivariant. Then one can find the fixed cubics in singular quadric  $S_0$ . The following contents before the end of this section are well studied in [Rei95, Section 2], [Hos90, Section 1] and [Cos06, Section 2]. The resolution of the quadric cone  $S_0$  at the cone point p is isomorphic to the Hirzebruch surface  $\mathbb{F}_2 := \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(2))$ . Let  $C_0$  (resp. F) be the canonical section class (resp. fiber class) in the divisor class  $\mathrm{Div}(\mathbb{F}_2) = \langle C_0, F \rangle$ . Then  $S_0$  is given by the image of the complete linear system  $|C_0 + 2F| (\cong \mathbb{P}^3)$ . Also

the surface  $\mathbb{F}_2$  embeds in the complete linear system  $|C_0 + 3F| (\cong \mathbb{P}^5)$ . That is, we have a commutative diagram:

(12) 
$$\mathbb{F}_{2} \xrightarrow{|C_{0}+3F|} \mathbb{P}^{5}$$

$$\downarrow^{C_{0}+2F|} \downarrow^{\pi}$$

$$S_{0} \subset \mathbb{P}^{3}$$

Furthermore, the right vertical map  $\pi$  in (12) is a linear projection.

Remark 3.14 (cf. [CPS19, Lemma 4.12]). Let  $[C] \in \mathbf{S}_3(S_0)$ . Note that C is a CM-curve. Let p be the cone point of  $S_0$ . Let  $\operatorname{mult}_p(C)=1$ . Then by the projection formula, the strict transform  $\tilde{C}$  of C along the map  $\pi_{\mathbb{F}_2}:\mathbb{F}_2\to S_0$  lies in the linear system  $[\tilde{C}]\in |C_0+3F|$ . In fact,  $\tilde{C}\cdot(C_0+2F)=C\cdot H_{\mathbb{P}^3}=3$  by the projection formula. Also,  $\tilde{C}\cdot C_0=1$ . Hence  $\tilde{C}=C_0+3F$ . Let  $\operatorname{mult}_p(C)>1$  and  $\tilde{C}=\pi_{\mathbb{F}_2}^{-1}(C)$  be the scheme theoretic inverse image of C along the map  $\pi_{\mathbb{F}_2}$ . Then  $[\tilde{C}]=[C_0]+[\tilde{C}_s]$  where  $\tilde{C}_s$  is the strict transform of C such that  $[\tilde{C}_s]=3[F]$ . Therefore  $\tilde{C}$  lies in the linear system  $|C_0+3F|$ .

We interpret  $\mathbb{F}_2$  and  $S_0$  as *scrolls* to make all relevant maps of the diagram (12) a  $\mathbb{C}^*$ -equivariant one. By definition, the *rational normal scroll* S(p,q) is the *join variety* of the rational normal curves of degree p and q; here, we allow p or q to be zero. In our case,  $\mathbb{F}_2 \cong S(1,3)$  and  $S_0 \cong S(0,2)$ . Then it is well-known that the map  $\pi$  in (12) is a successive projection from a point on the ruling line and cubic curve. Let us present  $\pi$  in (12) as homogeneous coordinates. To do this, let us define S(1,3) and S(0,2) as the closure of the image of 4-dimensional complex torus  $(\mathbb{C}^*)^4$  as below.

(13) 
$$(\mathbb{C}^*)^4 \xrightarrow{(t_0, t_1, u_0, u_1) \mapsto [t_0^3 u_0 : t_0^2 t_1 u_0 : t_0 t_1^2 u_0 : t_1^3 u_0 : t_0 u_1 : t_1 u_1]} \to \mathbb{P}^5$$

$$\downarrow \qquad \qquad \downarrow \qquad$$

Then the scroll S(1,3) (resp. S(0,2)) is defined by the maximal minors of the *catalecticant* matrix

(14) 
$$\begin{bmatrix} z_0 & z_1 & z_2 & z_4 \\ z_1 & z_2 & z_3 & z_5 \end{bmatrix} \quad (\text{resp.} \begin{bmatrix} z_0 & z_1 \\ z_1 & z_2 \end{bmatrix}).$$

Under this setting, the map  $\pi$  in (13) is

$$\pi([z_0:z_1:z_2:z_3:z_4:z_5]) = [z_0:z_1:z_2:z_4].$$

Now we are ready to state our main proposition.

**Proposition 3.15.** *In the case (ii) and (iii) (and thus (iv) and (v)) of Table 3, the fixed cubic curves are classified by the followings.* 

- (1) If  $S_0 = Q \cap \{v_{-1,-3} = 0\}$ , then the  $\mathbb{C}^*$ -fixed, irreducible cubic curve in a quadric cone  $S_0$  is defined by the family  $\mathcal{C}_a$  in equation (10).
- (2) If  $S_0 = Q \cap \{v_{1,-3} = 0\}$ , then every  $\mathbb{C}^*$ -fixed cubic curve in a quadric cone  $S_0$  is isolated and degenerated one.

*Proof.* Case (1): Since  $S_0$  is defined by  $v_{3,-1}v_{1,-3} - 3v_{1,-1}^2 = 0$ , we may assume that the weights of  $z_0$ ,  $z_1$ ,  $z_2$  and  $z_4$  are 2, 0, -2 and 4 (after rescaling the coordinate  $z_1$ ). If we let the weights  $z_3$  and  $z_5$  by -4 and 2 to be invariant of equations (14), then the map  $\pi$  is a  $\mathbb{C}^*$ -equivariant one. This induces a  $\mathbb{C}^*$ -action on the complete linear system  $|C_0 + 3F|$  which regards as the space of quartic curves in  $\mathbb{F}_2$ . Hence the repeat of the weight 2 gives us one parameter family of quartic curves in  $\mathbb{F}_2$  and thus rational cubic curves in  $S_0$  (Remark 3.14). Note that the other case is isolated and the corresponding curve is a degenerated one by substituting in equation (14). Obviously, the family  $C_a$  in (10) is invariant under  $\mathbb{C}^*$ -action and thus we proved the claim.

Case (2): The similar computation as in the case (1) shows that if we let the weights of  $z_0, \dots, z_4$  and  $z_5$  by 4, 0, -4, -8, 2 and -2, then  $\pi$  is  $\mathbb{C}^*$ -equivariant. Thus  $\mathbb{C}^*$ -fixed curves in  $\mathbb{F}_2$  are isolated and degenerated one. Thus the same thing holds on  $S_0$ .

It seems possible to compute the weights by using the result in [ES96] and Lemma 2.4. Since we explicitly describe the moduli space  $S_3(Q)$  in terms of a projective bundle in the following section, we omit it.

#### 4. CUBIC CURVES SPACE AS A PROJECTIVE BUNDLE

In this section, we describe the moduli space  $S_3(Q)$  as a projective bundle over a Grassmannian (Proposition 4.4). As a corollary, we give the cohomology ring structure of  $S_3(Q)$  (Corollary 4.6).

4.1. Sheaf theoretic description of  $S_1(Q)$ . Let us recall the global description of  $S_1(Q)$ . Let L be a line in Q. It is known that the locally free resolution of the ideal sheaf  $\mathcal{I}_{L,Q}$  is given by

(15) 
$$0 \to \mathcal{O}_Q(-1) \to \mathcal{U}_Q \to \mathcal{I}_{L,Q} \to 0,$$

where  $\mathcal{U}_Q$  is the restriction of the universal subbundle of Grassmannian Gr(2,4); see [Fae14, Section 2.1.1]. Also, every line in Q arises in this fashion, so called the Hartshorne-Serre correspondence ([Arr07]). Thus we get the following.

**Proposition 4.1.** Let  $S_1(Q)$  be the Hilbert scheme of lines in Q. Then,

$$\mathbf{H}_1(Q) \cong \mathbb{P}\mathrm{Hom}(\mathcal{O}_Q(-1), \mathcal{U}_Q) \cong \mathbb{P}\mathrm{H}^0(\mathcal{U}_Q(1)) \cong \mathbb{P}^3.$$

4.2. **Proof of Theorem 1.2.** The moduli space of representations of a Kronecker quiver parametrizes the isomorphism classes of stable sheaf homomorphisms

(16) 
$$\mathcal{O}_Q^{\oplus 2} \longrightarrow \mathcal{U}_Q(1)$$

up to the natural action of the automorphism group  $\mathbb{C}^* \times \operatorname{GL}_2/\mathbb{C}^* \cong \operatorname{GL}_2$ . For two vector spaces E and F of dimension 2 and 1 respectively and  $V^* := \operatorname{H}^0(Q, \mathcal{U}_Q(1))$ , the moduli space is constructed as  $\mathbf{G} := \operatorname{Hom}(E, V^* \otimes F) /\!/ \operatorname{GL}_2 \cong V^* \otimes F \otimes E^* /\!/ \operatorname{GL}_2$  with an appropriate linearization; see [Kin94]. Note that since the  $\operatorname{GL}_2$  acts as a row operation on the space of  $(2 \times 4)$ -matrices, and thus  $\mathbf{G} \cong \operatorname{Gr}(2,4)$ .

Let  $p_1 : \mathbf{G} \times Q \to \mathbf{G}$  and  $p_2 : \mathbf{G} \times Q \to Q$  be the two projections, and write  $\mathcal{A} \boxtimes \mathcal{B} := p_1^* \mathcal{A} \otimes p_2^* \mathcal{B}$  for  $\mathcal{A} \in \mathsf{Coh}(\mathbf{G})$  and  $\mathcal{B} \in \mathsf{Coh}(Q)$ . If  $\mathcal{U}_{\mathbf{G}}$  is the universal subbundle over  $\mathbf{G}$ , then there is a *universal morphism* 

(17) 
$$\phi: \mathcal{U}_{\mathbf{G}} \boxtimes \mathcal{O}_{Q} \longrightarrow \mathcal{O}_{\mathbf{G}} \boxtimes \mathcal{U}_{Q}(1);$$

see [Kin94]. Set  $\mathcal{J} := \operatorname{coker}(\phi)$  and denote  $\mathcal{J}_s := \mathcal{J}|_{\{s\} \times Q}$  for each point  $s \in \mathbf{G}$ .

**Proposition 4.2.** For the cokernel sheaf  $\mathcal{J}$  of the map in (17), we have the following.

- (1) For each point  $s \in \mathbf{G}$ , the restriction  $\mathcal{J}_s$  is isomorphic to a twisted ideal sheaf  $\mathcal{J}_s \cong \mathcal{I}_{L,S}(2)$  for some hyperplane section  $S := Q \cap H$  and a line  $L \subset S$ .
- (2)  $\mathcal{J}$  is flat over G and thus the universal morphism  $\phi$  in (17) is injective.

*Proof.* If  $\mathcal{K} := \ker(\phi_s)$  is non-zero, then it is a reflexive sheaf of rank one on  $\{s\} \times Q \cong Q$ . By the semistability of  $\mathcal{O}_Q^{\oplus 2}$  and  $\mathcal{U}_Q(1)$ , the sheaf  $\mathcal{K}$  is isomorphic to a line bundle  $\mathcal{K} \cong \mathcal{O}_Q(k)$  for some  $k \in \mathbb{Z}$  and the slope condition of  $\operatorname{Im}(\phi_s)$  gives  $0 \le -k \le \frac{1}{2}$ . Thus we get k = 0 and so  $\phi_s$  is not stable, i.e.,  $\operatorname{rank}(H^0(\phi_s)) = 1$ . Therefore  $\phi_s$  is injective.

Now, for the inclusion  $i_1: \mathcal{O}_Q \to \mathcal{O}_Q^{\oplus 2}$  into the first component, we have a commutative diagram:

Here,  $\xi:=\phi_s\circ i_1:\mathcal{O}_Q\to\mathcal{U}_Q(1)$  is the composition map. Then we get  $\operatorname{coker}(\xi)\cong\mathcal{I}_{L,Q}(1)$  for some line  $L\subset Q$  by Proposition 4.1. The composite map  $i\circ\overline{\phi}_s:\mathcal{O}_Q\stackrel{\overline{\phi}}{\to}\mathcal{I}_{L,Q}(1)\stackrel{i}{\hookrightarrow}\mathcal{O}_Q(1)$  is not a zero-map and thus it is injective. Hence we get  $\operatorname{coker}(i\circ\overline{\phi}_s)\cong\mathcal{O}_{Q\cap H}(1)$  for some hyperplane  $H\subset\mathbb{P}^4$ . Let  $S:=Q\cap H$ . Then the top row in the diagram (18) becomes

$$0 \to \mathcal{O}_Q \cong \mathcal{I}_{S,Q}(1) \xrightarrow{\overline{\phi}_s} \mathcal{I}_{L,Q}(1) \to \mathcal{I}_{L,S}(1) \to 0$$

and so we get  $\mathcal{J}_s = \mathcal{I}_{L,S}(1)$ , proving the assertion (1). Now  $\mathcal{J}_s$  has a constant Hilbert polynomial by the result of (1) and thus  $\mathcal{J}$  is flat over G, which confirms (2).

**Remark 4.3.** By the proof of (1) of Proposition 4.2, each hyperplane section  $Q \cap H$  is parameterized by G. Hence one can consider the universal family of hyperplane sections in  $G \times Q$ . Furthermore,  $G \cong Gr(2,4)$  is the space of lines in  $\mathbb{P}H^0(\mathcal{U}_Q(1)) = \mathbb{P}^3$ . On the other hand, the latter space  $\mathbb{P}^3$  is the Fano scheme of lines in Q; see Proposition 4.1. Let  $\mathcal{C} \subset \mathbb{P}^3 \times Q$  be the universal lines over  $\mathbb{P}^3$  with  $p: \mathcal{C} \to \mathbb{P}^3$  and  $q: \mathcal{C} \to Q$  the projection maps. Then it can be easily checked that the transform  $p(q^{-1}(S \cap H))$  of the hyperplane section  $Q \cap H$  becomes a line  $L \subset \mathbb{P}^3$ , i.e.,  $[L] \in G$ .

From the proof of Proposition 4.2, we obtain the exact sequence

(19) 
$$0 \to \mathcal{U}_{\mathbf{G}} \boxtimes \mathcal{O}_{\mathcal{O}} \to \mathcal{O}_{\mathbf{G}} \boxtimes \mathcal{U}_{\mathcal{O}}(1) \to \mathcal{J} \to 0.$$

By applying the functor  $R^{\bullet}p_{1,*}((-)\boxtimes \mathcal{O}_Q(1))$  to the exact sequence (19), we have

(20) 
$$0 \to \mathcal{U}_{\mathbf{G}} \otimes \mathrm{H}^{0}(\mathcal{O}_{Q}(1)) \to \mathcal{O}_{\mathbf{G}} \otimes \mathrm{H}^{0}(\mathcal{U}_{Q}(2)) \to p_{1,*}(\mathcal{J} \boxtimes \mathcal{O}_{Q}(1)) \to 0,$$

since  $\mathrm{H}^1(\mathcal{O}_Q(1))=0$ . Since  $\mathrm{h}^0(\mathcal{O}_Q(1))=5$  and  $\mathrm{h}^0(\mathcal{U}_Q(2))=16$ , we see that the direct image sheaf  $p_{1,*}(\mathcal{J}\boxtimes\mathcal{O}_Q(1))$  in (20) is a vector bundle of rank 6 on G.

**Proposition 4.4.** The moduli space  $\mathbf{S}_3(Q)$  is isomorphic to the projective bundle  $\mathbb{P}(p_{1,*}(\mathcal{J} \boxtimes \mathcal{O}_Q(1)))$ .

*Proof.* Let  $\mathcal{G} := p_{1,*}(\mathcal{J} \boxtimes \mathcal{O}_Q(1))$  and  $\pi : \mathbb{P}(\mathcal{G}) \to \mathbf{G}$  be the bundle morphism. Then, there exists a commutative diagram:

$$\mathbf{G} \times Q \stackrel{\pi \times i}{\longleftarrow} \mathbb{P}(\mathcal{G}) \times Q$$

$$\downarrow^{p_1} \qquad \qquad \downarrow_{\overline{p}_1}$$

$$\mathbf{G} \stackrel{}{\longleftarrow} \mathbb{P}(\mathcal{G}),$$

where  $\overline{p}_1:\mathbb{P}(\mathcal{G}) imes Q o \mathbb{P}(\mathcal{G})$  is the projection map. Let

$$c: \overline{p}_1^* \mathcal{O}_{\mathbb{P}(G)}(-1) \to (\pi \times i)^* (\mathcal{J} \boxtimes \mathcal{O}_{\mathcal{O}}(1))$$

be the composition of the pullback of the tautological map

(21) 
$$\mathcal{O}_{\mathbb{P}(\mathcal{G})}(-1) \to \pi^* \mathcal{G} \cong \overline{p}_{1,*}(\pi \times i)^* (\mathcal{J} \boxtimes \mathcal{O}_Q(1))$$

and the natural map

$$\overline{p}_{1}^{*}\overline{p}_{1,*}(\pi \times i)^{*}(\mathcal{J} \boxtimes \mathcal{O}_{Q}(1)) \to (\pi \times i)^{*}(\mathcal{J} \boxtimes \mathcal{O}_{Q}(1)).$$

Let  $\overline{\mathcal{Z}}:=(\pi\times i)^*\mathcal{Z}$  be the pull-back of the universal hyperplane section  $\mathcal{Z}\subset\mathbf{G}\times Q$  parameterized by  $\mathbf{G}$ ; see Remark 4.3. We claim that the local extension space  $\mathcal{E}xt^1_{\overline{\mathcal{Z}}}(\operatorname{coker}(c),\mathcal{O}_{\overline{\mathcal{Z}}})$  is a flat family of stable sheaves over  $\mathbb{P}(\mathcal{G})$  with Hilbert polynomial 3m+1. Note that it is enough to check the claim fiberwise. Over a point  $x\in\mathbb{P}(\mathcal{G})$ , the exact sequence

$$0 \to \operatorname{Im}(c) \to (\pi \times i)^*(\mathcal{J} \boxtimes \mathcal{O}_Q(1)) \to \operatorname{coker}(c) \to 0$$

becomes a short exact sequence

$$0 \to \mathcal{O}_S \to \mathcal{I}_{L,S}(2) \to \mathcal{T} \to 0$$

where  $\mathcal{T} := \operatorname{coker}(c)_x$ . By applying the dual functor  $\mathcal{H}om_S(-, \mathcal{O}_S)$  to it, we have

$$0 \cong \mathcal{H}om_{S}(\mathcal{T}, \mathcal{O}_{S}) \to \mathcal{H}om_{S}(\mathcal{I}_{L,S}(2), \mathcal{O}_{S}) \to \mathcal{H}om_{S}(\mathcal{O}_{S}, \mathcal{O}_{S}) \cong \mathcal{O}_{S}$$
$$\to \mathcal{E}xt_{S}^{1}(\mathcal{T}, \mathcal{O}_{S}) \to \mathcal{E}xt_{S}^{1}(\mathcal{I}_{L,S}(2), \mathcal{O}_{S}) \cong 0.$$

The first term vanishes because  $\mathcal{T}$  is one-dimensional. Also, the last vanishing is a special case of a more general vanishing  $\mathcal{E}xt_S^{i\geq 1}(\mathcal{I}_{L,S}(2),\mathcal{O}_S)\cong 0$ , which is obvious when S is smooth. If S is not smooth, one can check this by using Macaulay2 ([GS]). By computing the Hilbert polynomial of  $\mathcal{H}om_S(\mathcal{I}_{L,S}(2),\mathcal{O}_S)$  and  $\mathcal{O}_S$ , we conclude that  $\mathcal{E}xt_S^1(\mathcal{T},\mathcal{O}_S)=\mathcal{O}_C$  for some twisted cubic curve  $C\subset Q$ . Hence by the universal property of moduli space  $\mathbf{S}_3(Q)$ , there exists a morphism

(22) 
$$\Phi: \mathbb{P}(\mathcal{G}) \longrightarrow \mathbf{S}_3(Q)$$

induced by the flat family  $\mathcal{E}xt^{1}_{\overline{Z}}(\operatorname{coker}(c), \mathcal{O}_{\overline{Z}})$  over  $\mathbb{P}(\mathcal{G})$ .

Lastly, we prove that the induced map  $\Phi$  in (22) is an isomorphism. By Lemma 4.5 below and Zariski main theorem, it is enough to check that the map  $\Phi$  is generically one-to-one. Let us choose a smooth twisted cubic  $C \subset Q$  such that, for the linear span  $H_0 := \langle C \rangle \cong \mathbb{P}^3$ , the hyperplane section  $Q \cap H_0 =: S_0$  is a smooth quadric surface. In  $C \subset S_0$ , the curve class [C] in  $S_0$  is automatically determined. Hence the inverse image  $\Phi^{-1}([\mathcal{O}_C])$  is a unique point in  $\mathbb{P}(\mathcal{G})$ .

## **Lemma 4.5.** The rank of the Picard group $Pic(S_3(Q))$ is 2.

*Proof.* Following the blowing-up/down diagram in [CHK12], we know that the rank of Picard group of  $S_3(Q)$  is the same as that of  $M_3(Q)$ . Here  $M_3(Q)$  is the moduli space of stable maps of degree 3 in Q. But the Picard group of  $M_3(Q)$  is generated by the boundary divisor of reducible curves and the locus of stable maps whose images meet a fixed line in Q (compare with the result in [Opr05]).

**Corollary 4.6.** The cohomology ring of  $S_3(Q)$  is given by

$$\mathrm{H}^*(\mathbf{S}_3(Q),\mathbb{Q})\cong\mathbb{Q}[c_1,c_2,h]/I,$$

$$I = \langle c_1^3 - 2c_1c_2, c_1^4 - 3c_1^2c_2 + c_2^2, h^6 - 5c_1h^5 + (15c_1^2 - 5c_2)h^4 - 40c_1c_2h^3 + 50c_2^2h^2 \rangle$$

where  $c_1 = c_1(\mathcal{U}_G)$ ,  $c_2 = c_2(\mathcal{U}_G)$  and  $h = c_1(\mathcal{O}_{\mathbb{P}}(1))$  is the hyperplane class of the projective bundle in Proposition 4.4.

*Proof.* By the exact sequence (20) and the presentation of the cohomology ring of Gr(2, 4) in [EH16, Theorem 5.2], one can obtain the result.

## REFERENCES

- [AF93] Paolo Aluffi and Carol Faber. Linear orbits of *d*-tuples of points in  $\mathbb{P}^1$ . 1993(445):205–220, 1993. 1
- [Arr07] Enrique Arrondo. A home-made Hartshorne-Serre correspondence. *Revista Matemática Complutense*, 20(2):423–443, 2007. 14
- [BB73] A. Białynicki-Birula. Some theorems on actions of algebraic groups. *Annals of Mathematics*, 98(3):480–497, 1973. 3
- [Cao19] Yalong Cao. Counting conics on sextic 4-folds. *Mathematical Research Letters*, 26(5):1343–1357, 2019.
- [CCM14] Jinwon Choi, Kiryong Chung, and Mario Maican. Moduli of sheaves supported on quartic space curves. arXiv:1409.1449, 2014. 4
- [CHK12] Kiryong Chung, Jaehyun Hong, and Young-Hoon Kiem. Compactified moduli spaces of rational curves in projective homogeneous varieties. *J. Math. Soc. Japan*, 64(4):1211–1248, 2012. 1, 2, 4, 17
- [CHL18] Kiryong Chung, Jaehyun Hong, and SangHyeon Lee. Geometry of moduli spaces of rational curves in linear sections of Grassmannian Gr(2,5). *Journal of Pure and Applied Algebra*, 222(4):868 888, 2018. 1
- [Chu22] Kiryong Chung. Desingularization of Kontsevich's compactification of twisted cubics in  $V_5$ . Manuscripta Mathematica, 2022. 4
- [CK11] Kiryong Chung and Young-Hoon Kiem. Hilbert scheme of rational cubic curves via stable maps. *Amer. J. Math.*, 133(3):797–834, 2011. 1, 2
- [CLW21] Kiryong Chung, Sanghyeon Lee, and Joonyeong Won. Correspondence of Donaldson-Thomas and Gopakumar-Vafa invariants on local Calabi-Yau 4-folds over  $V_5$  and  $V_{22}$ . arXiv:2109.02087, 2021.
- [CMT18] Yalong Cao, Davesh Maulik, and Yukinobu Toda. Genus zero Gopakumar-Vafa type invariants for Calabi-Yau 4-folds. *Advances in Mathematics*, 338:41–92, 2018. 2
- [Cos06] Izzet Coskun. Degenerations of surface scrolls and the Gromov-Witten invariants of Grassmannians. *J. Algebraic Geom.*, 15(2):223–284, 2006. 12
- [CPS19] Ivan Cheltsov, Victor Przyjalkowski, and Constantin Shramov. Fano threefolds with infinite automorphism groups. *Izvestiya: Mathematics*, 83, 08 2019. 13
- [CS18] Ivan Cheltsov and Constanti Shramov. Kähler-Einstein Fano threefolds of degree 22. arXiv:1803.02774, 2018. 1
- [Don08] S.K. Donaldson. Kähler geometry on toric manifolds, and some other manifolds with large symmetry. arXiv:0803.0985, 2008. 1
- [EH82] D. Eisenbud and J. Harris. Curves in Projective Space. Presses de L'Université de Montréal, 1982. 4
- [EH16] D. Eisenbud and J. Harris. 3264 and All That: A Second Course in Algebraic Geometry. Cambridge: Cambridge University Press, 2016. 18

- [EPS87] Geir Ellingsrud, Ragni Piene, and Stein Arild Strømme. On the variety of nets of quadrics defining twisted cubics. In *Space curves* (*Rocca di Papa, 1985*), volume 1266 of *Lecture Notes in Math.*, pages 84–96. Springer, Berlin, 1987. 1
- [ES96] Geir Ellingsrud and Stein Arild Strømme. Bott's formula and enumerative geometry. *J. Amer. Math. Soc.*, 9(1):175–193, 1996. 1, 14
- [Fae14] Daniele Faenzi. Even and odd instanton bundles on Fano threefolds of Picard number one. *manuscripta math*, 144:199–239, 2014. 14
- [FGP19] Andrea Fanelli, Laurent Gruson, and Nicolas Perrin. Rational curves on  $V_5$  and rational simple connectedness. https://arxiv.org/abs/1901.06930, 2019. 2
- [GS] Daniel R. Grayson and Michael E. Stillman. Macaulay2, a software system for research in algebraic geometry. Available at http://www.math.uiuc.edu/Macaulay2/. 10, 17
- [Hos90] Toshio Hosoh. Stable sheaves of rank 2 on a 3-dimensional rational scroll. *Tokyo Journal of Mathematics*, 13(2):301 326, 1990. 12
- [Ili94] Atanas Iliev. The Fano surface of the Gushel threefold. Compositio Mathematica, 94(1):81–107, 1994. 1
- [Ive72] Birger Iversen. A fixed point formula for action of tori on algebraic varieties. *Inventiones mathematicae*, 16(3):229–236, 1972. 3
- [Kin94] A. D. King. Moduli of representations of finite-dimensional algebras. *Quart. J. Math. Oxford Ser.* (2), 45(180):515–530, 1994. 15
- [KS04] J. Koll´ar and F.-O. Schreyer. Real Fano 3-folds of type  $V_{22}$ . Dipartimento di Matematica dell'Universit´a di Torino, v 62, 2004. 1
- [McC00] John McCleary. A User's Guide to Spectral Sequences. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2 edition, 2000. 5
- [MU83] Shigeru Mukai and Hiroshi Umemura. Minimal rational threefolds. In Michel Raynaud and Tetsuji Shioda, editors, *Algebraic Geometry*, pages 490–518, Berlin, Heidelberg, 1983. Springer Berlin Heidelberg. 1
- [Opr05] Dragos Oprea. Divisors on the moduli spaces of stable maps to flag varieties and reconstruction. *J. Reine Angew. Math.*, 2005(586):169–205, 2005. 17
- [PS85] Ragni Piene and Michael Schlessinger. On the Hilbert scheme compactification of the space of twisted cubics. *Amer. J. Math.*, 107(4):761–774, 1985. 1
- [Rei95] M. Reid. *Chapters on Algebraic Surfaces*. Warwick preprints. University of Warwick. Mathematics Institute, 1995. 12
- [San14] Giangiacomo Sanna. Rational curves and instantons on the Fano threefold  $V_5$ . arXiv:1411.7994, 2014. 1
- [Sch01] Frank-Olaf Schreyer. Geometry and algebra of prime Fano 3-folds of genus 12. *Compositio Mathematica*, 127(3):297–319, 2001. 1
- [VX02] I. Vainsencher and F. Xavier. A compactification of the space of twisted cubics. *Math. Scand.*, 91(2):221–243, 2002. 1

Department of Mathematics Education, Kyungpook National University, 80 Daehakro, Bukgu, Daegu, 41566, Republic of Korea

Email address: krchung@knu.ac.kr

Department of Mathematics, Sungkyunkwan University, 2066 Seobu-ro, Suwon, 16419, Republic of Korea

Email address: sukmoonh@skku.edu

DEPARTMENT OF MATHEMATICS EDUCATION, GONGJU NATIONAL UNIVERSITY OF EDUCATION, 27 UNGJIN-RO, GONGJU-SI, CHUNGCHEONGNAM-DO, 32553, REPUBLIC OF KOREA

Email address: sbyoo@gjue.ac.kr