

Adding a Tail in Classes of Perfect Graphs^{*}

Anna Mpanti^{1,2}, Stavros D. Nikolopoulos^{1,3}, and Leonidas Palios^{1,4}

¹ Dept. of Computer Science and Engineering, University of Ioannina, Greece

² ampanti@cs.uoi.gr

³ 0000-0001-6684-8459; stavros@cs.uoi.gr

⁴ 0000-0001-8630-3835; palios@cs.uoi.gr

Abstract. Consider a graph G which belongs to a graph class \mathcal{C} . We are interested in connecting a node $w \notin V(G)$ to G by a single edge uw where $u \in V(G)$; we call such an edge a *tail*. As the graph resulting from G after the addition of the tail, denoted $G + uw$, need not belong to the class \mathcal{C} , we want to compute a minimum \mathcal{C} -completion of $G + w$, i.e., the minimum number of non-edges (excluding the tail uw) to be added to $G + uw$ so that the resulting graph belongs to \mathcal{C} .

In this paper, we study this problem for the classes of split, quasi-threshold, threshold, and P_4 -sparse graphs and we present linear-time algorithms by exploiting the structure of split graphs and the tree representation of quasi-threshold, threshold, and P_4 -sparse graphs.

Keywords: edge addition · completion · split graph · quasi-threshold graph · threshold graph · P_4 -sparse graph

1 Introduction

Given a graph G , an edge connecting a vertex $w \notin V(G)$ to a vertex u of G is a *tail* added to G ; let us denote the resulting graph as $G + uw$. If G belongs to a class \mathcal{C} of graphs, this may not hold for the graph $G + uw$. Hence, we are interested in computing a minimum \mathcal{C} -completion of $G + uw$, i.e., the minimum number of non-edges (excluding the tail uw) to be added to $G + uw$ so that the resulting graph belongs to \mathcal{C} ; such non-edges are called *fill edges*. The above problem is an instance of the more general $(\mathcal{C}, +k)$ -MinEdgeAddition problem [22] in which we add k given non-edges in a graph belonging to a class \mathcal{C} and we want to compute a minimum \mathcal{C} -completion of the resulting graph.

Computing a minimum completion of an arbitrary graph into a specific graph class is an important and well studied problem with applications in areas involving graph modeling with missing edges due to lacking data, e.g., molecular biology and numerical algebra [9,19,25]. Unfortunately, minimum completions

^{*} Research at the University of Ioannina supported by the Hellenic Foundation for Research and Innovation (H.F.R.I.) under the “First Call for H.F.R.I. Research Projects to support Faculty members and Researchers and the procurement of high-cost research equipment grant”, Project FANTA (eFficient Algorithms for NeTwork Analysis), number HFRI-FM17-431.

into many interesting graph classes, such as split graphs, chordal graphs and cographs, are NP-hard to compute [2,8,16,30]. This led the researchers towards the computation of minimal completions, the solution of problems with restricted input, and approximation or parameterized algorithms.

A related field is that of the dynamic recognition (or on-line maintenance) problem on graphs: a series of requests for the addition or the deletion of an edge or a vertex (potentially incident on a number of edges) are submitted and each is executed only if the resulting graph remains in the same class of graphs. Several authors have studied this problem for different classes of graphs and have given algorithms supporting some or all the above operations; we mention the edges-only fully dynamic algorithm of Ibarra [13] for chordal and split graphs, and the fully dynamic algorithms of Hell et al. [11] for proper interval graphs, of Shamir and Sharan [26] for cographs, of Heggenes and Mancini for split graphs [10], and of Nikolopoulos et al. for P_4 -sparse graphs [23].

In this paper, we exploit the structure of split graphs and the tree representation of quasi-threshold, threshold, and P_4 -sparse graphs in order to present algorithms for computing a minimum completion of a given graph G in each of these classes to which we have added a tail. Given the (K, S) -partition of a given split graph or the tree representation of a given quasi-threshold, threshold, or P_4 -sparse graph, our algorithms run in optimal $O(n)$ time where n is the number of vertices of G . These algorithms are a first step towards the solution of the $(\mathcal{C}, +1)$ -MinEdgeAddition problem [22] for each of these four classes \mathcal{C} of graphs.

2 Theoretical Framework

We consider finite undirected graphs with no loops or multiple edges. For a graph G , we denote by $V(G)$ and $E(G)$ the vertex set and edge set of G , respectively. Let S be a subset of the vertex set $V(G)$ of a graph G . Then, the subgraph of G induced by S is denoted by $G[S]$. The *neighborhood* $N_G(x)$ of a vertex x of the graph G is the set of all the vertices of G which are adjacent to x . The *closed neighborhood* of x is defined as $N_G[x] := N_G(x) \cup \{x\}$. The *degree* of a vertex x in G , denoted $deg(x)$, is the number of vertices adjacent to x in G ; thus, $deg(x) = |N_G(x)|$. A vertex of a graph is *universal* if it is adjacent to all other vertices of the graph. We extend this notion to a subset of the vertices of a graph G and we say that a vertex is *universal in a set* $S \subseteq V(G)$, if it is universal in the induced subgraph $G[S]$. Finally, C_k (P_k resp.) denotes the chordless cycle (chordless path resp.) on k vertices; in each P_4 , the unique edge incident on its first or last vertex is called a *wing*.

3 Split Graphs

The split graphs are of wide theoretical interest and have been the focus of many research papers. An undirected graph G is *split* if its vertex set $V(G)$ admits a partition into a clique K and an independent set S [7]; the partition into K, S




Fig. 1: The structure of the tree representation of a threshold graph [20]

can be computed in time proportional to the size of the graph. It also holds that a graph is split if and only if it contains no induced C_4 , C_5 , or $2K_2$.

Lemma 1. *Let G be a split graph with vertex partition into a clique K and an independent set S , u a vertex of G , uw a tail, and $K_s = \{x \in K \mid N_G(x) \cap S \neq \emptyset\}$. Then, there exists a split-completion for the graph $G + uw$ in which the number of fill edges needed is 0 if $u \in K$ and $|K_s| - \deg_G(u)$ if $u \in S$.*

Proof. If $u \in K$, no fill edge (in addition to uw) is needed, which is optimal, since $G + uw$ is a split graph with clique K and independent set $S \cup \{w\}$.

Now consider that $u \in S$. A split completion of $G + uw$ can be obtained by connecting u to all its non-neighbors in K_s ; the resulting graph is split with clique $K_s \cup \{u\}$ and independent set $S \cup (K - K_s) \cup \{w\}$. To prove its optimality, suppose for contradiction that there existed a split completion of $G + uw$ that uses fewer than $|K_s| - \deg_G(u)$ fill edges. Then, there would exist a vertex $a \in K_s \setminus N_G(u)$ which is not incident on any fill edge. If there existed one more vertex $b \in K_s \setminus (N_G(u) \cup \{a\})$ not incident on any fill edge as well, then the edges ab and uw would form a $2K_2$, a contradiction. Then, all the fill edges would be incident on the vertices in $K_s \setminus (N_G(u) \cup \{a\})$. But then, if z is a neighbor of a in S , the edges az and uw would form a $2K_2$, a contradiction. \square

Since the vertices in the clique K are all pairwise adjacent, we note that $|K \setminus K_s| \leq 1$. The lemma directly implies that given the set K_s , the minimum number of fill edges can be computed in $O(|V(G)|)$ time otherwise the time complexity is $O(|V(G)| + |E(G)|)$.

4 Threshold and Quasi-threshold Graphs

Threshold Graphs. A well-known subclass of perfect graphs called threshold graphs are those whose independent vertex set subsets can be distinguished by using a single linear inequality. A graph G is *threshold* if there exists a threshold




Fig. 2: The tree representation of a quasi-threshold graph.

assignment $[\alpha, t]$ consisting of a labeling α of the vertices by non-negative integers and an integer threshold t such that: a set $S \subseteq V(G)$ is independent if and only if $\alpha(v_1) + \alpha(v_2) + \dots + \alpha(v_p) \leq t$ where $v_i \in S, 1 \leq i \leq p$. Chvátal and Hammer [3] first proposed threshold graphs in 1973 and have proved that the threshold graphs are precisely the graphs that contain no induced C_4 , P_4 , or $2K_2$.

Nikolopoulos [20] proved that every threshold graph admits a unique rooted tree representation as shown in Figure 1: each tree node is associated with a vertex set $V_{i,j}$ (these sets partition the vertex set of the graph) with each $V_{i,1}$ inducing a clique and each of the remaining sets containing a single vertex (note that the tree nodes associated with these singleton sets have no descendants) and the vertices in the union of the sets associated with the nodes on a path of tree edges from a tree node to any one of its descendants induce a clique. Thus, the vertices in $V_{k,1}$ are adjacent to all the vertices in $\bigcup_{i>k} V_{i,j}$, the vertices in $\bigcup_i \bigcup_{j \geq 2} V_{i,j}$ form an independent set, and the vertices in $\bigcup_i V_{i,1}$ induce a clique.

Let G be a threshold graph and consider adding a tail uw to G where $u \in V(G)$. Then we can show the following lemma.

Lemma 2. *Let G be a threshold graph and let its tree representation T_G consist of nodes associated with sets $V_{i,j}$ where $0 \leq i \leq h$ and $1 \leq j \leq k_i$ (Figure 1). Consider the addition of a tail uw where $u \in V(G)$. Then, there exists a minimum threshold completion of the graph $G + uw$ using f fill edges (excluding uw) where:*

- (i) *If $u \in V_{i,1}$, then $f = \min_{0 \leq \ell \leq i} \left\{ \left(\sum_{s=\ell+1}^i \sum_{t=2}^{k_s} |V_{s,t}| \right) + \sum_{s=0}^{\ell} |V_{s,1}| \right\}$;*
- (ii) *If $u \in V_{i,j}$ where $2 \leq j \leq k_i$, then $f = \min\{f_1, f_2\}$ where*

$$f_1 = \sum_{r=i}^h |V_{r,1}| + \min_{i \leq \ell \leq h} \left\{ \left(\sum_{s=\ell+1}^h \sum_{t=2}^{k_s} |V_{s,t}| \right) + \sum_{s=0}^{\ell} |V_{s,1}| \right\} \text{ and}$$

$$f_2 = \sum_{r=i}^h |V_{r,1}| + \min_{0 \leq \ell \leq i-1} \left\{ \left(\sum_{s=\ell+1}^h \sum_{t=2}^{k_s} |V_{s,t}| \right) + \sum_{s=0}^{\ell} |V_{s,1}| \right\} - 1.$$

Quasi-threshold Graphs. A graph G is called *quasi-threshold*, or QT-graph for short, if G contains no induced C_4 or P_4 [6,18,27,28]. The class of quasi-threshold-




Fig. 3: The forbidden subgraphs of the class of P_4 -sparse graphs [15].

graphs is a subclass of the class of cographs and properly contains the class of threshold graphs [4,5,7,29]. Nikolopoulos and Papadopoulos [24] have shown, among other properties, a unique rooted tree representation of QT-graphs which is a generalization of the tree representation of threshold graphs (Figure 2): the tree nodes are associated with disjoint vertex subsets each inducing a clique and the vertex sets associated with the tree nodes in a path from a tree node to any of its descendants induce a clique. It has been proven that a graph is QT-graph if and only if it admits such a tree representation [17,21]. Then, by generalizing the approach for threshold graphs, we can show the following lemma.

Lemma 3. *Let $G = (V, E)$ be a QT-graph, and let T_G be its tree representation. Consider the addition of a tail uw incident on a node u of G and suppose without loss of generality that $u \in V_{i,1}$ and that the vertex sets associated with the tree nodes on the path from the root of T_G to the node associated with $V_{i,1}$ are in order $V_{0,1}, V_{1,1}, \dots, V_{i,1}$. Then, there exists a minimum QT completion of the graph $G + uw$, and the minimum number of fill edges needed (excluding the tail uw) is $\min_{0 \leq \ell \leq i} \left\{ \left(\sum_{s=\ell+1}^i \sum_{t=2}^{k_s} |V_{s,t}| \right) + \sum_{s=0}^{\ell} |V_{s,1}| \right\}$*

5 P_4 -sparse Graphs

The P_4 -sparse graphs are defined as the graphs for which every set of 5 vertices induces at most one P_4 [12] (Figure 3 depicts the 7 forbidden subgraphs for the class of P_4 -sparse graphs). The P_4 -sparse graphs are perfect and also perfectly orderable [12], and properly contain many graph classes, such as, the cographs, the P_4 -reducible graphs, etc. (see [1,14,15]). They have received considerable attention in recent years and find applications in applied mathematics and computer science (e.g., communications, transportation, clustering, scheduling, computational semantics) in problems that deal with graphs featuring “local density” properties.

For a P_4 -sparse graph either the graph or its complement is disconnected with the connected components inducing P_4 -sparse graphs, or induces a spider. A graph H is called a *spider* if its vertex set $V(H)$ admits a partition into sets S, K, R such that:

- the set S is an independent set, the set K is a clique, and $|S| = |K| \geq 2$;
- every vertex in R is adjacent to every vertex in K and to no vertex in S ;




Fig. 4: (left) A thin spider; (right) a thick spider.




Fig. 5

- there exists a bijection $f : S \rightarrow K$ such that for each vertex $s \in S$ either $N_G(s) \cap K = \{f(s)\}$ or $N_G(s) \cap K = K - \{f(s)\}$; in the former case, the spider is *thin*, in the latter it is *thick* (see Figure 4).

Note that for $|S| = |K| = 2$, the spider is simultaneously thin and thick. To avoid ambiguity, in the following, for thick spiders we assume that $|K| \geq 3$.

In [15], Jamison and Olariu showed that each P_4 -sparse graph G admits a unique tree representation, up to isomorphism, called the P_4 -sparse tree $T(G)$ of G , which is a rooted tree such that:

- (i) each internal node of $T(G)$ has at least 2 children provided that $|V(G)| \geq 2$;
- (ii) the internal nodes are labelled by one of 0, 1, or 2 (0-, 1-, 2-nodes, resp.) and the parent-node of each 0- or 1-node t has a different label than t ;
- (iii) the leaves of the P_4 -sparse tree are in a 1-to-1 correspondence with the vertices of G ; if the least common ancestor of the leaves corresponding to two vertices v_i, v_j of G is a 0-node (1-node, resp.) then the vertices v_i, v_j are non-adjacent (adjacent, resp.) in G , whereas the vertices corresponding to the leaves of a subtree rooted at a 2-node induce a spider.

The structure of the P_4 -sparse tree implies the following lemma.

Lemma 4. *Let G be a P_4 -sparse graph and let $H = (S, K, R)$ be a thin spider of G . Moreover, let $s \in S$ and $k \in K$ be vertices that are adjacent in the spider.*

P1. *Every vertex of the spider is adjacent to all vertices in $N_G(s) \setminus \{k\}$.*

P2. *Every vertex $z \in K \setminus \{k\}$ is adjacent to all vertices in $N_G(k) \setminus \{s, z\}$.*

Let G be a given graph to which we want to add the tail uw with $u \in V(G)$. Let $t_0 t_1 \dots t_h u$ be the path from the root t_0 of the P_4 -sparse tree T_G of G to the leaf associated with u . Moreover, let V_i ($0 \leq i < h$) be the set of vertices associated with the leaves of the subtrees rooted at the children of t_i except for t_{i+1} and V_h be the set of vertices associated with the leaves of the subtrees rooted at the children of t_h except for the leaf associated with u (see Figure 5). The sets V_0, V_1, \dots, V_h form a partition of $V(G) \setminus \{u\}$.

We show that there always exists a minimum P_4 -sparse completion of the graph $G + uw$ exhibiting one of a small number of different formations for u, w .




Fig. 6: (left) Formation 1; (right) Formation 2 where t is a 1- or a 2-node. Formation 1 is a special case of Formation 2 when $Z = \emptyset$.

Lemma 5. *Let G be a P_4 -sparse graph and T_G be its P_4 -sparse tree. Consider the addition of a tail uw incident on a node u of G . Then, there exists a minimum P_4 -sparse completion G' of the graph $G + uw$ such that for the P_4 -sparse tree $T_{G'}$ of G' , one of the following three cases holds:*

1. *The nodes u, w in $T_{G'}$ have the same parent-node which is a 2-node corresponding to a thin spider (S, K, R) with $u \in K$ and $w \in S$.*
2. *The P_4 -sparse tree $T_{G'}$ results from T_G by replacing the leaf for u by the 3-treenode Formation 1 shown in Figure 6(left).*
3. *The P_4 -sparse tree $T_{G'}$ results from T_G by removing the leaf for u and replacing an 1- or a 2-node t in the path from the root of T_G to the leaf for u by the 5-treenode Formation 2 in Figure 6(right).*

Proof. Let G_{OPT} be a minimum P_4 -sparse completion of the graph $G + uw$ and let T_{OPT} be its P_4 -sparse tree. We consider the following cases:

A. *The leaves associated with u, w in T_{OPT} do not have the same parent-node:* Let T_R be the P_4 -sparse tree obtained from T_{OPT} by using Formation 2 just above the least common ancestor t of w and u in T_{OPT} (Figure 7); let G_R be the P_4 -sparse graph corresponding to the tree T_R . Then, G_R uses no more fill edges than T_{OPT} . To see this, let t' be the child of t that is an ancestor of the leaf for u (note that t' may coincide with the leaf for u). Since u, w are adjacent in G_{OPT} , t is a 1- or a 2-node. In either case, w is adjacent to all vertices corresponding to the leaves of the subtree of T_{OPT} rooted at t' and all these edges, except for the tail uw , are fill edges. If t is a 1-node, then u is adjacent to all vertices in X (Figure 7) and thus G_R uses no more fill edges. If t is a 2-node then u is adjacent to all the vertices in the clique K_X of the corresponding spider (which includes w). Moreover, because $w \in K_X$, w is adjacent to all the vertices in $K_X \setminus \{w\}$ and to at least 1 vertex in the independent set for a total of $|K_X|$ fill edges; these fill edges can be used to connect u to the vertices in the independent set of the spider and thus G_R uses no more fill edges in this case too.

Now, in the P_4 -sparse tree T_R in Figure 7(right), let $A = V(G) \setminus (Z \cup \{u\})$. Recall that in the P_4 -sparse tree T_G of G , the path from the root t_0 to u is




Fig. 7: (left) The P_4 -sparse tree T_{OPT} in which the leaves for u, w do not have the same parent-node and have node t as their least common ancestor; (right) The P_4 -sparse tree T_R obtained by using Formation 2 just above node t which results in no more fill edges than those in G_{OPT} .

$t_0 t_1 \cdots t_h u$ and V_i ($0 \leq i \leq h$) is the set of vertices associated with the leaves of the subtrees rooted at the children of t_i except for t_{i+1} (where t_{h+1} is the leaf associated with u); see Figure 5.

We first observe that the induced subgraph $G_R[Z]$ induced by the set of vertices Z corresponding to the leaves of the subtree of T_R rooted at node t coincides with the induced subgraph $G[Z]$ (otherwise G_{OPT} would include fill edges that could be removed in contradiction to its optimality); then, let $t = t_k$. It also holds that node t in T_R is a 1- or a 2-node, since node t was a 1- or a 2-node in T_{OPT} , as well. Let $A = V(G) \setminus (Z \cup \{u\})$. Note that there is no set V_j such that $x, y \in V_j$, x is a neighbor of u in G , $x \in V_j \cap A$ and $y \in V_j \cap Z$, otherwise we can move x to Z along with y ; because y is in Z , all adjacencies from y to all the vertices in $V(G) \setminus (V_j \cup \{u\})$ in G are maintained and this will also hold for x , and the fill edge xw will be removed, a contradiction. Similarly, there is no set V_j such that $x, y \in V_j$, y is a non-neighbor of u in G and $x \in V_j \cap A$ and $y \in V_j \cap Z$ otherwise we can move y to A along with x . This implies that for each $i = 0, 1, \dots, h$, either $V_i \subseteq A$ or $V_i \subseteq Z$, and since $t = t_k$, $V_k \subseteq Z$.

Finally, there exists no $j > k$ such that $V_j \subseteq A$. Suppose that there existed such a V_j and let j be the largest such index. Then, because $t = t_k$ is a 1- or a 2-node and $k < j$, there would exist vertex $z \in V_k$ which would be adjacent to all vertices in V_j . This implies that in T_R , the least common ancestor of z and the vertices in V_k would be a 1-node. But then, if we moved V_j to Z then we would have fewer fill edges, a contradiction. Therefore, the tree T_R is as described in Case 3 of the statement of the lemma.

B. *The leaves associated with u, w in T_{OPT} have the same parent-node p : Then, since u, w are adjacent, the parent-node p is either an 1-node or a 2-node.*




Fig. 8: A transformation that reduces the number of fill edges.

- (i) *The parent-node p of u, w in T_{OPT} is an 1-node:* Then, the leaves associated with u and w are the only children of p (Formation 1), otherwise we can use Formation 2 as shown in Figure 8 which requires fewer fill edges. Then, w will be adjacent to all neighbors of u in T_{OPT} ; this and the optimality of G_{OPT} imply that T_{OPT} results from T_G by replacing the leaf for u by Formation 1.
- (ii) *The parent-node p of u, w in T_{OPT} is a 2-node:* Let $H = (S, K, R)$ be the corresponding spider. If H is thick (thus $|K| \geq 3$), then no matter whether the tail uw is an S - K , K - K , or R - K edge, the sum of degrees of u, w in H (excluding uw) is at least $|V(H)| - 3 + |K| - 2$ (consider an S - K edge). However, we would have added no more fill edges if we have made u universal in $G[V(H) \setminus \{w\}]$ and then applied Formation 2 at the parent of the leaf for u (then $Z = V(H) \setminus \{u, w\}$) using $V(H) - 2 \leq V(H) + |K| - 5$ fill edges. In the same way, we show that we would have added no more fill edges if H is a thin spider and the tail uw is a K - K or K - R edge. If uw is an S - K edge with $u \in S$ and $w \in K$, then we exchange u and w for the same total number of fill edges and get a thin spider with $u \in K$ and $w \in S$.

□

5.1 Adding a Tail to a Spider

In this section, we consider adding a tail uw to a spider $H = (S_H, K_H, R_H)$ where $u \in V(H)$. In the following two lemmas, we address the cases of a thin or a thick spider H respectively.

Lemma 6. *Consider the addition of a tail uw to a thin spider $H = (S_H, K_H, R_H)$ where u is a vertex of H . Then, for the number f of fill edges (excluding the tail uw) in a minimum P_4 -sparse completion of the graph $H + uw$, it holds:*

1. if $u \in S_H$, $f = |K_H| - 1$ if $R_H = \emptyset$ and $f = |K_H|$ otherwise;
2. if $u \in K_H$, $f = |K_H| - 1$;
3. if $u \in R_H$, then $f = \min\{|R_H \setminus N_H[u]|, |K_H| + f'\}$ where f' is the number of fill edges (excluding uw) in a minimum P_4 -sparse completion of the graph $H[R_H] + uw$.

Proof. 1. Let $v \in K_H$ be the neighbor of u in H . Then, we can get a P_4 -sparse graph as follows: if $R_H = \emptyset$, we connect u to all vertices in $K_H \setminus \{v\}$ (we get a thin spider with $S = (S_H \setminus \{u\}) \cup \{w\}$, $K = (K_H \setminus \{v\}) \cup \{u\}$, and $R = \{v\}$, that is, the tail uw is a wing of a P_4 of a thin spider), otherwise we connect v to all vertices in $\{w\} \cup (S_H \setminus \{u\})$, which makes v universal in $V(H) \cup \{w\}$ and u, w form a separate connected component in $G[V(G) \setminus \{v\}]$; the total number of fill edges (excluding the tail uw) is precisely $|K_H| - 1$ if $R_H = \emptyset$ and K_H otherwise.

Moreover, this is the minimum number of fill edges (excluding uw) needed. First, we note that for each pair k_i, s_i where $k_i \in K_H \setminus \{v\}$ and $s_i \in S_H \setminus \{u\}$, the vertices v, u, w, k_i, s_i define an F_5 or an F_3 depending on whether the vertices v, w are adjacent or not, which implies that at least $|K_H| - 1$ fill edges (excluding uw) are needed. Then, if there is a way of getting a P_4 -sparse graph by adding fewer than the number of fill edges mentioned in Case 1 of the statement of the lemma, it has to be the case that (i) $R_H \neq \emptyset$, (ii) each pair k_i, s_i where $k_i \in K_H \setminus \{v\}$ and $s_i \in S_H \setminus \{u\}$ is incident on exactly 1 fill edge, and (iii) no more fill edges exist. Let $r \in R_H$ and $k \in K_H \setminus \{v\}$. Then, the vertices v, u, w, k, r induce a forbidden subgraph (an F_5 if k is non-adjacent to both u, w , or an F_6 (F_1 , resp.) if k becomes adjacent to u (w , resp.) by means of a fill edge); thus, at least $|K_H|$ fill edges are needed in this case.

2. Let $v \in S_H$ be the neighbor in H of $u \in K_H$. Then, by connecting u to all vertices in $S_H \setminus \{v\}$ (which makes u universal in H) or by connecting w to all vertices in $K_H \setminus \{u\}$ yields a P_4 -sparse graph. Moreover, this is the minimum number of fill edges (excluding the tail uw) that need to be added. Suppose, for contradiction, that we get a P_4 -sparse graph after having added fewer than $|K_H| - 1$ fill edges (excluding uw) to the thin spider H . Then, there exists a pair of adjacent vertices s, k with $s \in S_H \setminus \{v\}$ and $k \in K_H \setminus \{u\}$ such that neither s nor k is incident on a fill edge. Then the vertices u, v, w, s, k induce a forbidden subgraph F_5 or F_3 if w and v are adjacent or not, respectively, a contradiction.

3. The term $R_H \setminus N_H[u]$ corresponds to making u universal in $H[R_H]$ in which case the resulting graph is P_4 -sparse (it is a thin spider with $S = S_H \cup \{w\}$, $K = K_H \cup \{u\}$, and $R = R_H \setminus \{u\}$). The term $|K_H| + f'$ corresponds to adding $|K_H|$ fill edges connecting w to the vertices in K_H and then computing a minimum P_4 -sparse completion of the graph $H[R_H] + uw$. Note that no minimum P_4 -sparse completion of $H + uw$ exists with u not being universal in R_H and with using fewer than $|K_H|$ fill edges incident on the vertices in $S_H \cup K_H$: if there were such a minimum P_4 -sparse completion H' of $H + uw$, then in H' , there would exist a non-neighbor $r \in R_H$ and a pair of adjacent vertices s, k where $s \in S_H$ and $k \in K_H$ such that neither s nor k would be incident on a fill edge; but then, in H' , the vertices u, w, r, s, k induce an F_4 or an F_3 if w, r have been connected by a fill edge or not, respectively, which leads to a contradiction. In turn, if H' has at least $|K_H|$ fill edges incident on vertices in $S_H \cup K_H$ then $H'[R_H \cup \{w\}]$ would be P_4 -sparse using fewer than f' fill edges in contradiction to the minimality of f' . \square

Lemma 7. *Consider the addition of a tail uw to a thick spider $H = (S_H, K_H, R_H)$ where u is a vertex of H . Then, for the number f of fill edges (excluding the tail uw) in a minimum P_4 -sparse completion of the graph $H + uw$, it holds:*

1. if $u \in S_H$,

$$f = \begin{cases} |K_H| - 1 = 2 & \text{if } |K_H| = 3 \text{ and } R_H = \emptyset \\ |K_H| = 3 & \text{if } |K_H| = 3 \text{ and } |R_H| = 1 \\ |K_H| + 1 = 4 & \text{if } |K_H| = 3 \text{ and } |R_H| \geq 2 \\ |K_H| & \text{if } |K_H| \geq 4 \text{ and } R_H = \emptyset \\ |K_H| + 1 & \text{if } |K_H| \geq 4 \text{ and } |R_H| \geq 1; \end{cases}$$

2. if $u \in K_H$, $f = 1$;
3. if $u \in R_H$, then $f = |K_H| + f'$ where f' is the number of fill edges (excluding uw) in a minimum P_4 -sparse completion of the graph $H[R_H] + uw$.

Proof. 1. Let $v \in K_H$ be the non-neighbor of u in H . Let us first consider the case $|K_H| = 3$. If $|R_H| \leq 2$, we can get a P_4 -sparse graph after having added the fill edges vu and vw (this implies that v becomes universal in $(V(H) \setminus \{v\}) \cup \{w\}$) and those connecting u to the vertices in R_H if R_H is non-empty; then the vertices in $(V(H) \setminus \{v\}) \cup \{w\}$ induce a thin spider with $K = (K_H \setminus \{v\}) \cup \{u\}$, $S = (S_H \setminus \{u\}) \cup \{w\}$, and $R = R_H$, for a total of $|K_H| - 1 + |R_H|$ fill edges (excluding the tail uw). If $|R_H| \geq 2$, a P_4 -sparse graph is obtained after in addition to the tail uw , we add the fill edges vu , vw (again v is universal in $(V(H) \setminus \{v\}) \cup \{w\}$), and the fill edges connecting w to the vertices in $K_H \setminus \{v\}$ (then the vertices in $(V(H) \setminus \{v\}) \cup \{w\}$ induce a thin spider with $K = K_H \setminus \{v\}$, $S = S_H \setminus \{u\}$, and $R = R_H \cup \{u, w\}$), for a total of $|K_H| + 1$ fill edges (excluding uw).

Now, consider the case that $|K_H| \geq 4$. If $|R_H| \leq 1$, we get a P_4 -sparse graph after having made u universal by connecting it to the remaining vertices in S_H by using $|K_H| - 1$ fill edges, and adding the fill edge uv , and those connecting u to the vertices in R_H if R_H is non-empty, for a total of $|K_H| + |R_H|$ fill edges (excluding uw). If $|R_H| \geq 1$, a P_4 -sparse graph is obtained after having made v universal (by adding the fill edges vu and vw) and after having connected w to all vertices in $K_H \setminus \{v\}$ (then the vertices in $(V(H) \setminus \{v\}) \cup \{w\}$ induce a thick spider with $K = K_H \setminus \{v\}$, $S = S_H \setminus \{u\}$, and $R = R_H \cup \{u, w\}$) for a total of $|K_H| + 1$ fill edges (excluding uw).

Below we show the minimality of this solution. Recall that $v \in K_H$ is the non-neighbor of u in H . We consider each of the five cases.

- (i) $|K_H| = 3$ and $R_H = \emptyset$: Suppose, for contradiction, that there is a P_4 -sparse completion of $H + uw$ with at most $|K_H| - 2 = 1$ fill edge (excluding uw). If v is incident on the unique fill edge (which connects v to u or w), then the vertices in $S \cup \{v, w\}$ induce an F_3 . Now suppose that the fill edge is not incident on v . Moreover, there exists at least one vertex $s \in S_H \setminus \{u\}$ that is not incident on the fill edge either. Then, the vertices u, v, w, s, k (where $k \in K_H$ is the non-neighbor of s in H) induce an F_5 if k, w are connected by the fill edge, or an F_2 otherwise.

- (ii) $|K_H| = 3$ and $|R_H| = 1$: Let $R_H = \{r\}$. Suppose, for contradiction, that there is a P_4 -sparse completion of $H + uw$ with at most $|K_H| - 1 = 2$ fill edges (excluding uw). We distinguish three cases depending on whether v is incident on 0, 1, or 2 fill edges:
- *v is not incident on a fill edge*: If there exists a pair s, k of non-neighbors with $s \in S_H \setminus \{u\}$ and $k \in K_H \setminus \{v\}$ such that none of s, k is incident on a fill edge to u or w , the vertices u, v, w, s, k induce an F_2 . Otherwise, since the number of such pairs is 2, for each such pair s, k , exactly one of s, k is incident on a fill edge to u or w , and no other fill edges exist. If there exists a vertex $k \in K_H \setminus \{v\}$ not incident on a fill edge to w , the vertices u, v, w, k, r induce an F_5 , otherwise each of the fill edges connects each of the vertices in $K_H \setminus \{v\}$ to w , and then u, v, w, s, k (for any pair s, k of non-neighbors with $s \in S_H \setminus \{u\}$ and $k \in K_H \setminus \{v\}$) induce an F_5 .
 - *v is incident on 1 fill edge (to u or w)*: Then, there is 1 more fill edge; hence, there exist 2 vertices in the set $(S_H \setminus \{u\}) \cup \{r\}$ that are not incident on a fill edge connecting them to u or w , and let these vertices be p_1, p_2 . Then, the vertices u, v, w, p_1, p_2 induce an F_5 if p_1, p_2 are connected by a fill edge or an F_3 otherwise.
 - *v is incident on 2 fill edges connecting it to u and w* : Then, there is no other fill edge. Then, the vertices u, w, k, k', r (where $\{k, k'\} = K_H \setminus \{v\}$) induce an F_6 .
- (iii) $|K_H| = 3$ and $|R_H| \geq 2$: Let r_1, r_2 be two vertices in R_H . Suppose, for contradiction, that there is a P_4 -sparse completion of $H + uw$ with at most $|K_H| = 3$ fill edges (excluding uw). Again, we distinguish three cases depending on whether v is incident on 0, 1, or 2 fill edges:
- *v is not incident on a fill edge*: Consider the case that there exists a vertex $k \in K_H \setminus \{v\}$ that is not incident on a fill edge to w . Let $s \in S_H$ be the non-neighbor of k in H and $A = (S_H \setminus \{u, s\}) \cup \{r_1, r_2\}$; the set A contains 3 vertices which are common neighbors of v, k . If at least one of these 3 vertices (say, p) is not incident on a fill edge to u, w , then the vertices u, v, w, k, p induce an F_5 , otherwise all 3 of these vertices are incident on a fill edge to u, w (then these are all the fill edges) and the vertices u, v, w, s, k induce an F_2 . On the other hand, if no such vertex k exists, then both vertices in $K_H \setminus \{v\}$ are incident on a fill edge to w , accounting for 2 of the 3 fill edges; then there exists a vertex $s' \in S_H \setminus \{u\}$ which is not incident on a fill edge to w , and the vertices u, v, w, s', k' (where $k' \in K_H$ is the non-neighbor of s') induce an F_5 .
 - *v is incident on 1 fill edge (to u or w)*: There are 2 more fill edges; hence, there exist 2 vertices in the set $(S_H \setminus \{u\}) \cup \{r_1, r_2\}$ that are not incident on a fill edge connecting them to u or w , and let these vertices be p_1, p_2 . Then, the vertices u, v, w, p_1, p_2 induce an F_5 if p_1, p_2 are connected by a fill edge or an F_3 otherwise.
 - *v is incident on 2 fill edges connecting it to u and w* : Then, there is 1 more fill edge; hence, there exists a vertex $k \in K_H \setminus \{v\}$ that is not incident on the fill edge. Moreover, there exist 2 vertices in the set $(S_H \setminus \{u, s\}) \cup \{r_1, r_2\}$ that are not incident on a fill edge connecting them to

- u or w (where $s \in S_H$ is the non-neighbor of k); let these vertices be p_1, p_2 . Then, the vertices u, w, k, p_1, p_2 induce an F_5 if p_1, p_2 are adjacent or an F_3 otherwise.
- (iv) $|K_H| \geq 4$ and $R_H = \emptyset$: Suppose, for contradiction, that there is a P_4 -sparse completion of $H + uw$ with at most $|K_H| - 1$ fill edges (excluding the tail uw). Again, we distinguish three cases depending on whether v is incident on 0, 1, or 2 fill edges:
- v is not incident on a fill edge: If there exists a vertex $s \in S_H \setminus \{u\}$ not incident on a fill edge to u, w or to its non-neighbor $k \in K_H$ in H , the vertices u, v, w, s, k induce an F_5 if k, w are connected by a fill edge, or an F_2 otherwise; if all vertices in $S_H \setminus \{u\}$ are incident on a fill edge to u, w , or their non-neighbor in K_H , then there are no more fill edges and the vertices u, v, w, k, k' (for any $k, k' \in K_H \setminus \{v\}$) induce an F_6 .
 - v is incident on 1 fill edge (to u or w): Then, the remaining fill edges are at most $|K_H| - 2$ in total. If there exist two vertices $s_1, s_2 \in S_H \setminus \{u\}$ not incident on a fill edge to u or w , the vertices u, v, w, s_1, s_2 induce an F_5 or an F_3 depending on whether s_1, s_2 are connected by a fill edge or not. Thus, there cannot be two such vertices s_1, s_2 ; this implies that the remaining fill edges are precisely $|K_H| - 2$, and they connect all but one vertex in $S_H \setminus \{u\}$ to u or w ; let that vertex be s . Then, the vertices u, v, w, s, k' (where $k' \in K_H \setminus \{v\}$ is a neighbor of s in H) induce an F_6 or an F_1 if the fill edge incident on v connects it to u or w respectively.
 - v is incident on 2 fill edges connecting it to u and w : Then, the remaining fill edges are at most $|K_H| - 3$ in total; hence, there exist two pairs of non-adjacent vertices s_1, k_1 and s_2, k_2 with $s_1, s_2 \in S_H \setminus \{u\}$ and $k_1, k_2 \in K_H \setminus \{v\}$ such that none of s_1, s_2, k_1, k_2 is incident on a fill edge to u or w . Let $A = S_H \setminus \{u, s_1, s_2\}$; the set A is the set of $|K_H| - 3$ common neighbors of k_1, k_2 in S_H other than u . If there exists a vertex $s \in A$ not incident on a fill edge to u or w , then the vertices u, w, k_1, k_2, s induce an F_6 , otherwise, the remaining fill edges are precisely $|K_H| - 3$ and they connect each of the vertices in A to u or w , that is, none of the vertices in $K_H \setminus \{v\}$ is incident on a fill edge. Then, the vertices u, w, s_1, s_2, k (where k is any vertex in $K_H \setminus \{v, k_1, k_2\}$) induce an F_3 .
- (v) $|K_H| \geq 4$ and $|R_H| \geq 1$: Let $r \in R_H$. Suppose, for contradiction, that there is a P_4 -sparse completion of $H + uw$ with at most $|K_H|$ fill edge (excluding the tail uw). Again, we distinguish three cases depending on whether v is incident on 0, 1, or 2 fill edges:
- v is not incident on a fill edge: If there exists a vertex $s \in S_H \setminus \{u\}$ not incident on a fill edge to u, w , or to its non-neighbor $k \in K_H$ in H , the vertices u, v, w, s, k induce an F_5 if k, w are connected by a fill edge, or an F_2 otherwise; if all vertices in $S_H \setminus \{u\}$ are incident on a fill edge to u, w , or their non-neighbor in K_H , which account for the $|K_H| - 1$ of the $|K_H|$ fill edges, there exist vertices $k, k' \in K_H \setminus \{v\}$ which are not incident on a fill edge and then the vertices u, v, w, k, k' induce an F_6 .
 - v is incident on 1 fill edge (to u or w): Then, the remaining fill edges are at most $|K_H| - 1$ in total. If all vertices in $K_H \setminus \{v\}$ are incident on a fill

edge to w , then no more fill edges exist and the vertices u, v, w, s_1, s_2 (for any $s_1, s_2 \in S_H \setminus \{u\}$) induce an F_3 . Thus, there exists $k \in K_H \setminus \{v\}$ which is not incident on a fill edge to w . The number of common neighbors of v, k in $S_H \cup r$ is $|K_H| - 1$. If each of these vertices is incident on a fill edge to u or w , then no more fill edges exist and the vertices u, v, w, s, k' induce an F_6 or an F_1 depending on whether the fill edge incident on v connects it to u or w , respectively, where $s \in S_H$ is the non-neighbor of k and k' is any vertex in $K_H \setminus \{v, k\}$; hence, there exists a common neighbor p not incident on a fill edge to u or w and the vertices u, v, w, k, p induce an F_6 or an F_1 depending on whether the fill edge incident on v connects it to u or w , respectively.

- *v is incident on 2 fill edges connecting it to u and w :* Then, the remaining fill edges are at most $|K_H| - 2$ in total; hence, there exists a pair of non-adjacent vertices s, k (where $s \in S_H \setminus \{u\}$ and $k \in K_H \setminus \{v\}$) which are not incident on a fill edge to u or w . Let $A = (S_H \setminus \{u, s\}) \cup \{r\}$; the set A is a set of $|K_H| - 1$ neighbors of k other than u . Then, there exists a vertex p_1 in A which is not incident on a fill edge to u or w . If there exists a second vertex p_2 in A not incident on a fill edge to u or w , then the vertices u, w, k, p_1, p_2 induce an F_5 if p_1, p_2 are connected by a fill edge or an F_3 otherwise. If each vertex in $A \setminus \{p_1\}$ is incident on a fill edge to u or w , then the fill edges incident on these vertices account for the remaining $|K_H| - 2$ fill edges and the vertices u, w, s, k_1, k_2 (for any vertices $k_1, k_2 \in K_H \setminus \{v, k\}$) induce an F_6 .

Therefore, if we use fewer than the stated number of fill edges, in each case, the resulting graph contains an induced forbidden subgraph, a contradiction.

2. Let $v \in S_H$ be the non-neighbor of u in H . Then, we get a P_4 -sparse graph by connecting u to v ; thus, u becomes universal in $V(H) \cup \{w\}$. This is the minimum number of fill edges (excluding the tail uw) that need to be added since for any pair of non-neighbors s, k with $s \in S_H \setminus \{v\}$ and $k \in K_H \setminus \{u\}$, the vertices u, v, w, s, k induce a forbidden subgraph F_3 , a contradiction.

3. By connecting w to all vertices in K_H and then computing a minimum P_4 -sparse completion of $H[R_H \cup \{w\}]$, we get a P_4 -sparse graph and the number of fill edges needed is $|K_H| + f'$.

To prove the minimality of this number of fill edges, suppose, for contradiction, that we can get a P_4 -sparse graph from $H + uw$ after having added at most $|K_H| - 1$ fill edges incident on vertices in $S_H \cup K_H$ (excluding the tail uw). Then, there exists a pair s_1, k_1 of non-neighbors in H with $s_1 \in S_H$ and $k_1 \in K_H$ none of which is incident on a fill edge to u or w . We distinguish the following two cases that cover all possibilities.

- *Each of the vertices in $K_H \setminus \{k_1\}$ is incident on a fill edge to w .* These are precisely all the $|K_H| - 1$ fill edges; hence none of the vertices in $S_H \setminus \{s_1\}$ is incident on a fill edge. Then, the vertices u, w, k_1, s_2, s_3 (for any $s_2, s_3 \in S_H \setminus \{s_1\}$) induce an F_3 .

- *There exists at least one vertex in $K_H \setminus \{k_1\}$ that is not incident on a fill edge to w .* Let that vertex be k_2 . Then, if there exists another vertex $k_3 \in K_H \setminus \{k_1, k_2\}$ that is not incident on a fill edge to w as well, the vertices u, w, k_2, k_3, s_1 induce an F_6 . On the other hand, if each of the vertices in $K_H \setminus \{k_1, k_2\}$ is incident on a fill edge to w (which implies that k_3 is adjacent to w), then these fill edges are $|K_H| - 2$ in total, with only 1 remaining. If the non-neighbor s_3 of k_3 in S_H is not incident on a fill edge to u or w , then the vertices u, w, k_1, k_2, s_3 induce an F_6 whereas if it is adjacent to u or w , then there are no more fill edges. In particular, if s_3 is adjacent to u , the vertices u, k_1, k_3, s_1, s_3 induce an F_6 and if it is adjacent to w , the vertices u, w, k_2, s_1, s_3 induce an F_4 .

In each case, we get a contradiction. Thus every minimum P_4 -sparse completion of $H + uw$ requires at least $|K_H|$ fill edges incident on vertices of $S_H \cup K_H$. Now, if there exists a minimum P_4 -sparse completion H' of $H + uw$ having fewer than $|K_H| + f'$ fill edges, then the fact that at least $|K_H|$ of them are incident on vertices in $S_H \cup K_H$ implies that $H'[R_H \cup \{w\}]$ is P_4 -sparse using fewer than f' fill edges in contradiction to the minimality of f' . \square

If the (thin or thick) spider H belongs to a more general P_4 -sparse graph, then Lemmas 6 and 7 imply the following result.

Corollary 1. *Let u be a vertex of a P_4 -sparse graph to which we add the tail uw . Let $t_0 \cdots t_h u$ be the path in the P_4 -sparse tree of G from the the root t_0 to the leaf for u and let V_0, \dots, V_h be the corresponding vertex sets as mentioned before. Then, if the parent t_h of u is a 2-node corresponding to a spider H , the number of fill edges needed for a minimum P_4 -sparse completion of the graph $G + uw$ (excluding the tail uw) does not exceed the minimum between*

- (i) *the minimum number given by Lemmas 6 and 7 (if H is thin or thick, respectively) augmented by $|N_G(u) \cap (V_0 \cup \dots \cup V_{h-1})|$ and*
- (ii) $\min_{t_i=1\text{- or }2\text{-node}} \{ |N_G(u) \cap (V_0 \cup \dots \cup V_{i-1})| + |(V_i \cup \dots \cup V_h) \setminus N_G(u)| \}$.

Case (i) of Corollary 1 corresponds to doing a minimum P_4 -completion of the graph $H + uw$ and not changing the rest of the P_4 -sparse tree T_G of G whereas Case (ii) corresponds to making u universal in H and then trying Formation 2 above each 1-node or 2-node in the path $t_0 \cdots t_h$ of T_G .

5.2 The Algorithm

Recall that $t_0 t_1 \cdots t_h u$ is the path in the P_4 -sparse tree T_G of G from the root t_0 to the leaf for u , and V_i ($0 \leq i < h$) is the set of vertices associated with the leaves of the subtrees rooted at the children of t_i except for t_{i+1} and V_h is the set of vertices associated with the leaves of the subtrees rooted at the children of t_h except for the leaf corresponding to u . See Figure 5.

Next we prove the conditions under which a minimum P_4 -sparse completion of the graph $G + uw$ uses fewer fill edges than when using Formation 1 or 2.

Lemma 8. *There exists a minimum P_4 -sparse completion G_{OPT} of the graph $G + uw$ which uses fewer fill edges than when using Formation 1 or 2 if and only if uw is a wing of a P_4 in G_{OPT} which implies that (i) either u is a vertex of a spider in G (Lemmas 6 and 7 apply) (ii) or there exists j ($0 \leq j < h$) such that t_j is a 1-node, t_{j+1} is a 0-node, and there exist vertices a, b such that $a \in V_j$ is universal in $G[V_j]$ and $b \in V_{j+1}$ is isolated in $G[V_{j+1}]$.*

Then, in G_{OPT} , the vertices u, w, a, b induce a P_4 in a spider (S, K, R) with $S = \{w, b\}$, $K = \{u, a\}$ and $R = (V_{j+1} \setminus \{b\}) \cup V_{j+2} \cup \dots \cup V_h$.

Proof. If Formation 1 or Formation 2 is not to be used then Lemma 5 implies that in G_{OPT} , uw is the wing of a P_4 . If u is a vertex of a spider, then Lemmas 6 and 7 apply. So, in the following, assume that u is not a vertex of a spider.

For the tail uw to be the wing of a P_4 in G_{OPT} , we can show that there exist vertices x, y such that uxy is a P_3 in the graph G : if u, x, y do not all belong to the same connected component of G , then we could add the tail uw to the connected component of G to which u belongs, thus using fewer fill edges than in G_{OPT} , a contradiction; if u, x, y belong to the same connected component of G but do not form a P_3 , then because u, y are not adjacent in G_{OPT} and thus neither in G , u, y are at distance 2 in G and there exists a P_3 uay in G (note that u, y cannot be at distance ≥ 4 in G since then G would contain an induced $P_5 = F_2$, and they cannot be at distance 3 either since then there exists a P_4 $uaby$ in G and u would be a vertex of a spider in G).

Therefore, in the following, consider that the minimum P_4 -sparse completion G_{OPT} of $G + uw$ contains an induced P_4 $wuab$ such that the graph G contains the induced P_3 uab . So, since u, b are not adjacent in G_{OPT} , then they are not adjacent in G either, and thus their least common ancestor t_k in the P_4 -sparse tree T_G of G is a 0-node; it cannot be a 2-node since then u would be a vertex of a spider. Moreover, a is a common neighbor of both u, b and thus the least common ancestor t_j of a, u in T_G is a 1- or a 2-node (in the latter case, a is a vertex of the clique of the spider) and $j < k$.

Let us now try forming the P_4 $wuab$, which clearly will belong to a spider, say $W = (S_W, K_W, R_W)$. We show that $|S_W| = |K_W| = 2$. First, note that the edge ab cannot belong to a spider in G , since then u would belong to that spider as well (note that the vertices of G not belonging to a spider are either adjacent to all vertices of the spider or to none of them), So, suppose for contradiction that the spider W has $|S_W| = |K_W| \geq 3$ and let $w, b, s \in S_W$ and $u, a, k \in K_W$ with the corresponding S - K pairs being w and u , b and a , and s and k . The spider W can be thin or thick.

- *The spider W is thin.* Then, $ba \in E(G)$ otherwise the removal of ba would produce a P_4 -sparse graph with fewer fill edges (b is isolated in $G[V(W)]$), a contradiction; similarly, $sk \in E(G)$. Moreover, $ak \in E(G)$: as above, if a, k do not belong to the same connected component of the induced sub-graph $G[V(W)]$, then by adding the tail uw to the connected component of $G[V(W)]$ to which u belongs would result into fewer fill edges; if a, k belong to the same connected component of $G[V(W)]$ then there exists a chordless

path ρ connecting them in the subgraph $G[K_W \cup R_W]$ and the vertices in $V(\rho) \cup \{b, s\}$ induce a P_ℓ with $\ell \geq 5$, in contradiction to the P_4 -sparseness of G . But then, G contains the P_4 *baks* and ab belongs to a spider.

- *The spider W is thick.* Then, $w \in S_W$ is incident on the tail uw and $|K_W| - 2 \geq 1$ fill edges. Since we can make u universal in $G[V(W) \setminus \{w\}]$ by using a single fill edge and then use Formation 2, it is clear that building the spider W does not result into fewer fill edges.

Thus, G_{OPT} with a spider W with $|K_W| \geq 3$ has no fewer fill edges than if we use Formation 2. Therefore, the P_4 *wuab* belongs to a spider with clique size equal to 2, which thus is thin. Then, Property P1 in Lemma 4 implies that w , u , and a are adjacent to all the neighbors of b except for a in G_{OPT} and thus at least to the neighbors of b in G ; thus, in G_{OPT} ,

- fill edges connect vertex w to the vertices in $((V_0 \cup \dots \cup V_{k-1}) \setminus \{a\}) \cap N_G(b) = [(V_0 \cup \dots \cup V_{k-1}) \setminus \{a\}] \cap N_G(u)$;
- vertex u and w are adjacent to all neighbors of b in V_k , that is, to the vertices in $(V_k \cap N_G(b)) \setminus N_G(u)$;
- vertex a is adjacent to all the vertices in $(V_j \cap N_G(b))$ and thus fill edges connect a to all vertices in $(V_j \cap N_G(b)) \setminus N_G[a] = (V_j \cap N_G(u)) \setminus N_G[a]$.

Additionally, Property P2 in Lemma 4 implies that because a is adjacent to all the vertices in $V_{j+1} \cup \dots \cup V_h$ and to the vertices in $V_j \cap N_G(a)$ in G_{OPT} (because it is adjacent to them in G), then so must be vertex u in G_{OPT} ; thus, in G_{OPT} , fill edges connect u to all the vertices in $(V_{j+1} \cup \dots \cup V_h) \setminus N_G(u)$ and the vertices in $(V_j \cap N_G(a)) \setminus N_G(u)$ (the set $(V_j \cap N_G(a)) \setminus N_G(u)$ is non-empty if and only if t_j is a 2-node).

Now, let us consider using Formation 2 right below node t_j in the P_4 -sparse tree T_G of G ; then, the number of fill edges is $|(V_{j+1} \cup \dots \cup V_h) \setminus N_G(u)| + |(V_0 \cup \dots \cup V_j) \cap N_G(u)|$; the former term corresponds to fill edges incident on u , the latter to fill edges incident on w . Then, because $j < k$ and $|((V_0 \cup \dots \cup V_{k-1}) \setminus \{a\}) \cap N_G(u)| = |(V_0 \cup \dots \cup V_{k-1}) \cap N_G(u)| - 1$, the only possibility for G_{OPT} to use fewer fill edges than using Formation 2 after node t_j requires that

1. $k = j + 1$;
2. $(V_j \cap N_G(u)) \setminus N_G[a] = \emptyset$;
3. $(V_k \cap N_G(b)) \setminus N_G(u) = \emptyset$ which implies that b is isolated in $G[V_k]$;
4. $(V_j \cap N_G(a)) \setminus N_G(u) = \emptyset$ which implies that t_j is a 1-node.

Requirement 4 implies that $V_j \cap N_G(u) = V_j$ which together with Requirement 2 imply that $N_G[a] = V_j$, that is, a is universal in $G[V_j]$, and we have the second case in the statement of the lemma. \square

Now we are ready to describe our algorithm for counting the number of fill edges in a minimum P_4 -sparse completion of the graph $G + uw$.

Algorithm P_4 -sparse-Tail-Addition

Input: a P_4 -sparse graph G , a vertex $u \in V(G)$, and a tail uw to be added to G .

Output: the minimum number of fill edges (excluding the tail uw) needed in a P_4 -sparse completion of the graph $G + uw$.

if $|V(G)| = 1$ **then** $\{V(G) = \{u\} \implies \text{the graph } G + uw \text{ is } P_4\text{-sparse}\}$
return(0);

compute the path $t_0 t_1 \dots t_h$ ($h \geq 1$) from the root t_0 of the P_4 -sparse tree of G to the parent-node t_h of the leaf corresponding to u ;

compute the sets of vertices V_i , $0 \leq i \leq h$ (see Figure 5);

$min \leftarrow |N_G(u)|$; $\{\text{corresponds to Formation 1}\}$

$\{\text{check for Formation 2 (Lemma 5(iii) and Case (ii) of Corollary 1)}\}$

for each t_i ($i = 0, 1, \dots, h$) that is a 1- or a 2-node **do**

$\ell \leftarrow |N_G(u) \cap (V_0 \cup \dots \cup V_{i-1})| + |(V_i \cup \dots \cup V_h) \setminus N_G(u)|$;

update min if $\ell < min$;

$\{\text{check for new } P_4 \text{ formation (Lemma 8)}\}$

for each $i = 0, 1, \dots, h - 1$ such that t_i is a 1-node and t_{i+1} is a 0-node **do**

if there exist vertex $a \in V_i$ such that a is universal in V_i **and**

vertex $b \in V_{i+1}$ such that b has no neighbors in V_{i+1} **then**

$\ell \leftarrow |N_G(u) \cap (V_0 \cup \dots \cup V_{i-1})| + |V_i \setminus \{a\}| + |V_{i+1} \setminus \{b\}| +$
 $| (V_{i+2} \cup \dots \cup V_h) \setminus N_G(u) |$;

update min if $\ell < min$;

$\{\text{check the cases if } t_h \text{ is a 2-node and apply case (i) of Corollary 1}\}$

if t_h is a 2-node **then**

$\ell \leftarrow$ number of fill edges according to the cases of Lemmas 6 or 7;

$\ell \leftarrow \ell + |N_G(u) \cap (V_0 \cup \dots \cup V_{h-1})|$; $\{\text{Case (i) of Corollary 1}\}$

update min if $\ell < min$;

return(min);

Algorithm P_4 -sparse-Tail-Addition can be easily augmented to return a minimum cardinality set of fill edges. The correctness of the algorithm follows from Lemmas 5, 6, 7, 8, and Corollary 1. Let G be the given graph and let n be the number of its vertices. If the P_4 -sparse tree T_G of G is given, an $O(n)$ -time traversal of the tree enables us to compute the path $t_0 t_1 \dots t_h u$ and the number of neighbors and non-neighbors of u in each of the sets V_0, \dots, V_h ; additionally, the height of T_G is $O(n)$ and thus $h = O(n)$. Since the conditions of Lemmas 6 and 7 can be checked in $O(1)$ -time, the entire algorithm runs in $O(n)$ time.

Theorem 1. *Let G be a P_4 -sparse graph on n vertices and let uw be tail attached at node u of G . If the P_4 -sparse tree of G is given, Algorithm P_4 -sparse-Tail-Addition computes the minimum number of edges to be added to $G + uw$ so that the resulting graph is P_4 -sparse in $O(n)$ time.*

If the P_4 -sparse tree T_G of G is not given, then it can be computed in $O(n+m)$ time where m is the number of edges of G [14], and the entire algorithm takes $O(n+m)$ time.

References

1. Brandstädt, A., Le, V.B., Spinrad, J.: Graph Classes – a Survey. SIAM Monographs in Discrete Mathematics and Applications, SIAM, Philadelphia (1999)
2. Burzyn, P., Bonomo, F., Durán, G.: NP-completeness results for edge modification problems. *Discrete Appl. Math.* **154**, 1824–1844 (2006)
3. Chvátal, V., Hammer, P.L.: Set-packing and threshold graphs, Research Report CORR 73-21, University of Waterloo, 1973.
4. Corneil, D.G., Lerches, H., Burlingham, L.: Complement reducible graphs, *Discrete Appl. Math.* **3** 163–174 (1981).
5. Corneil, D.G., Perl, Y., Stewart, L.K.: A linear recognition algorithm for cographs, *SIAM J. Comput.* **14** 926–934 (1985).
6. Golubic, M.C.: Trivially perfect graphs, *Discrete Math.* **24** 105–107 (1978).
7. Golubic, M.C.: Algorithmic Graph Theory and Perfect Graphs, Academic Press, New York, 1980.
8. El-Mallah, E., Colbourn, C.: The complexity of some edge deletion problems. *IEEE Trans. Circuits Systems* **35**, 354–362
9. Goldberg, P.W., Golubic, M.C., Kaplan, H., Shamir, R.: Four strikes against physical mapping of DNA. *J. Comput. Bio.* **2**(1), 139–152 (1995)
10. Heggernes, P., Mancini, F.: Dynamically maintaining split graphs, *Discrete Appl. Math.* **157**, 2047–2069 (2009)
11. Hell, P., Shamir, R., Sharan, R.: A fully dynamic algorithm for recognizing and representing proper interval graphs. *SIAM J. Comput.* **31**, 289–305 (2002)
12. Hoáng, C.: Perfect graphs. Ph.D. Thesis, McGill University, Montreal, Canada (1985)
13. Ibarra, L.: Fully dynamic algorithms for chordal graphs and split graphs. *ACM Trans. Algorithms* **4**, Article 40 (2008)
14. Jamison, B., Olariu, S.: Recognizing P_4 -sparse graphs in linear time. *SIAM J. Comput.* **21**, 381–406 (1992)
15. Jamison, B., Olariu, S.: A tree representation for P_4 -sparse graphs. *Discrete Appl. Math.* **35**, 115–129 (1992)
16. Kashiwabara, T., Fujisawa, T.: An NP-complete problem on interval graphs. In: *IEEE Symp. of Circuits and Systems*, pp. 82–83 (1979)
17. Kano, M., Nikolopoulos, S.D.: On the structure of A-free graphs. Part II, TR-25-99, Department of Computer Science, University of Ioannina, 1999.
18. Ma, S., Wallis, W.D., Wu, J.: Optimization problems on quasi-threshold graphs, *J. Comb. Inform. System Sci.* **14** 105–110 (1989).
19. Natanzon, A., Shamir, R., Sharan, R.: Complexity classification of some edge modification problems. *Discrete Appl. Math.* **113**, 109–128 (2001)
20. Nikolopoulos, S.D.: Recognizing cographs and threshold graphs through a classification of their edges. *Information Processing Letters* **74.3-4** (2000): 129-139.
21. Nikolopoulos, S.D.: Parallel algorithms for Hamiltonian problems on quasi-threshold graphs. *Parallel and Distributed Computing* **64**, 48–67 (2004).
22. Nikolopoulos, S.D., Palios, L.: Adding an Edge in a Cograph. In: Kratsch D. (ed.) *WG 2005, LNCS*, vol. 3787, pp. 214–226, Springer (2005).
https://doi.org/10.1007/11604686_19
23. Nikolopoulos, S.D., Palios, L., Papadopoulos, C.: A fully-dynamic algorithm for the recognition of P_4 -sparse graphs. *Theor. Comp. Science* **439**, 41-57 (2012).
24. Nikolopoulos, S.D., Papadopoulos, C.: The number of spanning trees in K_n -complements of quasi-threshold graphs. *Graphs and Combinatorics* **20.3**: 383-397 (2004).

25. Rose, D.J.: A graph-theoretic study of the numerical solution of sparse positive definite systems of linear equation. In: Read, R.C.(ed.) Graph Theory and Computing, pp. 183–217. Academic Press, New York (1972)
26. Shamir, R., Sharan, R.: A fully dynamic algorithm for modular decomposition and recognition of cographs. *Discrete Appl. Math.* **136**, 329–340 (2004)
27. Wolk, E.S.: The comparability graph of a tree. *Proc. Amer. Math. Soc.* 3 789–795 (1962) .
28. Wolk, E.S.: A note of the comparability graph of a tree. *Proc.Amer. Math. Soc.* 16 17–20 (1965).
29. Veldman, H.J.: A result on Hamiltonian line graphs involving restrictions on induced subgraphs. *J. Graph Theory* 12 413–420 (1988).
30. Yannakakis, M.: Computing the minimum fill-in is NP-complete. *SIAM J. Alg. Disc. Meth.* **2**, 77–79 (1981)