# Rescattering effects in nucleon-to-meson form factors and application to tau-lepton-induced proton decay

Andreas Crivellin<sup>a,b</sup>, Martin Hoferichter<sup>c</sup>

<sup>a</sup>Paul Scherrer Institut, 5232 Villigen PSI, Switzerland  $^bP$ hysik-Institut, Universität Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland <sup>c</sup>Albert Einstein Center for Fundamental Physics, Institute for Theoretical Physics, University of Bern, Sidlerstrasse 5, 3012 Bern, Switzerland

#### Abstract

Nucleon decays put extremely stringent bounds on baryon-number-violating interactions. However, in case the corresponding operators involve only  $\tau$  leptons, the direct two-body decays, e.g.,  $p \to \pi^0 \tau^+$ , are kinematically not allowed and nucleon decay operators involve only  $\tau$  leptons, the direct two-body decays, e.g.,  $p \to \pi^0 \tau^+$ , are kinematically not allowed and nucleon decay can only proceed via an off-shell  $\tau$ , leading to  $p \to \pi^0 \ell^+ \nu_\ell \bar{\nu}_\tau$ . To calculate such processes, the momentum dependence of the form factors for nucleon-to-meson transitions, which describe the hadronization of the underlying process at the quark level, is needed. In this work, we point out new isospin and Fierz relations among such proton–kaon matrix elements and calculate the momentum dependence of the nucleon-to-meson form factors from the universal final-state interactions in terms of pion–nucleon or kaon–nucleon scattering phase shifts. We use these results to derive novel limits on the Wilson coefficients of the baryon-numberviolating dimension-6 operators involving a  $\tau$  lepton, which were previously unconstrained.

The potential number is one of the Standard Monton decay in graceleon decay with some channels of the Hyper-Kamioka by another orde  $p \to \pi^0 e^+$  and while competitionalso expected for tantly, these intrive two-particl  $p \to e^+ X$  majo One can intro of effective of the SM effective of The potential simultaneous violation of baryon and lepton number is one of the frontiers of searches for physics beyond the Standard Model (BSM). Motivated by the prediction of proton decay in grand unified theories [1-3], limits on the nucleon decay width for various modes have been set that for some channels exceed 10<sup>34</sup> years, see Table 1. In the future, Hyper-Kamiokande [14] will be able to improve these limits by another order of magnitude, projecting  $7.8 \times 10^{34}$  years for  $p \rightarrow \pi^0 e^+$  and  $3.2 \times 10^{34}$  years for  $p \rightarrow K^+ \bar{\nu}$  (at 90% C.L.), while competitive new limits, especially for kaon modes, are also expected from DUNE [15] and JUNO [16]. Most importantly, these improvements not only pertain to the most sensitive two-particle channels, but also for generic searches such as  $p \rightarrow e^+ X$  major advances are projected.

One can interpret such limits model independently in terms of effective operators [17, 18], which are a subset of the full SM effective field theory (SMEFT) Lagrangian [19, 20]. The resulting constraints from direct two-body decays such as  $p \rightarrow$  $\pi^0 \ell$  are in general extremely stringent and push the BSM scale to at least 1014 GeV in case of tree-level effects with order-one couplings. However, in case only  $\tau$  leptons are involved, the decay of a nucleon is kinematically not allowed, so that the related SMEFT coefficients have not been constrained so far.<sup>1</sup>

Instead, since nucleon decay can be mediated by an off-shell  $\tau$ , processes such as  $p \to P\ell^+\nu_\ell\bar{\nu}_\tau$  with  $\ell = e, \mu$  and  $P = \pi, K, \eta$ do probe these couplings. They can be constrained by the inclusive limits given in the last panel of Table 1, but require knowledge of the momentum dependence of the hadronic part of the matrix elements. In fact, the value of such inclusive limits as generic tests of baryon number violation has already been stressed in Ref. [22]. In either case, to constrain the underlying Wilson coefficients at the partonic level, the hadronization of the effective operators must be calculated. Some relations among these nucleon-to-meson from factors can be derived using chiral perturbation theory (ChPT) [23-27], but especially estimates of their normalization have remained model dependent for a long time [28–30]. In the last years, calculations in lattice QCD [27, 31–35] have matured to the extent that these normalizations have now been reliably determined for many operators [35], thus rendering limits derived from two-body decays such as  $p \to \pi^0 e^+$  much more robust.

In this work, we focus on the momentum dependence of the form factors as required for the interpretation of the four-body decays sensitive to  $\tau$ -mediated proton decay. As a first step, we obtain new relations among the proton-to-kaon matrix elements from isospin symmetry and Fierz identities. Next, we analyze the unitarity relations that are produced by meson-nucleon rescattering effects, and show that the corresponding constraints allow one to calculate the momentum dependence from known meson-nucleon phase shifts. This strategy generalizes a similar approach for baryon-number-conserving nucleon form factors, in which case rescattering corrections in the two-meson system are resummed [36–42]. Finally, we apply these results to derive novel limits on the Wilson coefficients of the operators involving  $\tau$  leptons from four-body decays.

<sup>&</sup>lt;sup>1</sup>Note that because lepton flavor is exactly conserved in the SM with massless neutrinos, and only marginally violated by the light active neutrino masses, the renormalization group (RG) evolution from the BSM scale to the nucleon scale does not generate operators involving muons or electrons if these coupling are absent at the matching scale. Therefore, contrary to the case with heavy quarks [21], these operators do not contribute to proton decay even at the loop level.

Channel	Limit [10 <sup>30</sup> y]	Reference
$p \to \pi^0 e^+$	$2.4 \times 10^{4}$	[4]
$p\to\pi^0\mu^+$	$1.6 \times 10^{4}$	[4]
$p\to \pi^+\bar{\nu}$	$3.9 \times 10^{2}$	[5]
$p \to K^0 e^+$	$1.0 \times 10^{3}$	[6]
$p\to K^0\mu^+$	$3.6 \times 10^{3}$	[7]
$p \to K^+ \bar{\nu}$	$5.9 \times 10^{3}$	[8]
$p \rightarrow \eta e^+$	$1.0 \times 10^{4}$	[9]
$p \to \eta \mu^+$	$4.7 \times 10^{3}$	[9]
$n \to \pi^- e^+$	$5.3 \times 10^{3}$	[9]
$n\to \pi^-\mu^+$	$3.5 \times 10^{3}$	[9]
$n\to\pi^0\bar{\nu}$	$1.1 \times 10^{3}$	[5]
$n \to K^0 \bar{\nu}$	$1.3 \times 10^{2}$	[6]
$n \to \eta \bar{\nu}$	$1.6 \times 10^{2}$	[10]
$p, n \rightarrow e^+ X$	0.6	[11]
$p, n \to \mu^+ X$	12	[12]
·	•	

Table 1: Limits on nucleon lifetimes for various decay channels (all at 90% C.L.) [13].

### 2. Operators and matrix elements

As baryon number violation is experimentally extremely well constrained, it is reasonable to assume that it originates from a very high BSM scale. At this scale electroweak symmetry is unbroken and one can use SMEFT [19, 20] to reduce the number of possible *B*-violating operators to four, see Appendix A. After electroweak symmetry breaking, the decomposition of the lepton doublets results in the operators

$$\begin{aligned} Q_{udu}^{AB} &= \left[\bar{u}^c P_A d\right] P_B u, & Q_{udd}^{AB} &= \left[\bar{u}^c P_A d\right] P_B d, \\ Q_{usu}^{AB} &= \left[\bar{u}^c P_A s\right] P_B u, & Q_{usd}^{AB} &= \left[\bar{u}^c P_A s\right] P_B d, \\ Q_{dsu}^{AB} &= \left[\bar{d}^c P_A s\right] P_B u, & Q_{uds}^{AB} &= \left[\bar{u}^c P_A d\right] P_B s, \end{aligned} \tag{1}$$

whose matrix elements are calculated within lattice QCD [35]. In all cases, the contraction of the color indices with a Levi-Civita tensor is implied, the charge-conjugated fields are denoted by the superscript c, and projectors A, B = L, R are defined by  $P_{L/R} = (\mathbbm{1} \mp \gamma_5)/2$ . The corresponding Wilson coefficients are defined via  $\mathcal{L}_{\ell} = C_{\ell,\alpha}^{AB} \bar{\ell}_{\ell} Q_{\alpha}^{AB}$ ,  $\mathcal{L}_{\nu} = C_{\ell,\alpha}^{AB} \bar{\nu}_{\ell} Q_{\alpha}^{AB}$ , depending on whether the charge of  $\alpha = \{udu, \ldots\}$  requires a charged lepton or a neutrino.

Following the notation from Ref. [35], we label the matrix elements relevant for proton decay as

$$\langle \pi^{0} | \left[ \bar{u}^{c} P_{A} d \right] u_{B} | p \rangle = \frac{1}{\sqrt{2}} \langle \pi^{+} | \left[ \bar{u}^{c} P_{A} d \right] d_{B} | p \rangle \equiv \frac{1}{\sqrt{2}} U_{1}^{AB},$$

$$\langle K^{0} | \left[ \bar{u}^{c} P_{A} s \right] u_{B} | p \rangle \equiv S_{1}^{AB}, \qquad \langle K^{+} | \left[ \bar{u}^{c} P_{A} s \right] d_{B} | p \rangle \equiv S_{2}^{AB},$$

$$\langle K^{+} | \left[ \bar{u}^{c} P_{A} d \right] s_{B} | p \rangle \equiv S_{3}^{AB}, \qquad \langle K^{+} | \left[ \bar{d}^{c} P_{A} s \right] u_{B} | p \rangle \equiv S_{4}^{AB},$$

$$\langle \eta | \left[ \bar{u}^{c} P_{A} d \right] u_{B} | p \rangle \equiv S_{5}^{AB}, \qquad (2)$$

$X_i$	$W_0^{X_{iL}}(0)$	$W_1^{X_{iL}}(0)$	$W_0^{X_{iR}}(0)$	$W_1^{X_{iR}}(0)$
$U_1$	0.151(31)	-0.134(18)	-0.159(35)	0.169(37)
$S_1$	0.043(4)	0.028(7)	0.085(12)	-0.026(4)
$S_2$	0.028(4)	-0.049(7)	-0.040(6)	0.053(7)
$S_3$	0.101(11)	-0.075(13)	-0.109(19)	0.080(17)
$S_4$	-0.072(8)	0.024(6)	-0.044(5)	-0.026(6)
S <sub>1+2+4</sub>	0.000(0)	0.000(0)	0.000(0)	0.000(0)
$S_{2-3-4}$	0.000(0)	0.000(0)	0.112(15)	0.000(12)

Table 2: Nucleon-to-meson form-factor normalizations from Ref. [35], in units of GeV<sup>2</sup> at  $\overline{\rm MS}$  scale  $\mu=2$  GeV. All uncertainties have been combined in quadrature. Reference [35] also quotes results at  $q^2=m_\mu^2$ , but the observed variation compared to  $q^2=0$  is negligible within uncertainties. Results for the linear combinations from Eq. (3) are given in the last two lines [43].  $S_{5A}$  was not considered in Ref. [35] due to disconnected diagrams. Note that due to parity invariance  $X_{iL} \equiv X_i^{LL} = X_i^{RR}$ ,  $X_{iR} \equiv X_i^{RL} = X_i^{LR}$ .

where the first relation is known to follow from isospin symmetry [33], i.e., applies in the limit of equal up and down quark masses (isospin symmetry also determines the corresponding neutron matrix elements, see Appendix B). Since two of the four chirality combinations are redundant due to parity invariance of the strong interactions,  $X_i^{LL} = X_i^{RR}$  and  $X_i^{RL} = X_i^{LR}$ , we follow Ref. [35] and fix the second index to B = L, defining  $X_{iA} \equiv X_i^{AL}$ , X = U, S. In addition to the known isospin relation, we find that also the kaon matrix elements are not all independent:

$$S_{1A} + S_{2A} + S_{4A} = 0,$$
  
 $S_{2L} - S_{3L} - S_{4L} = 0,$  (3)

where the first equation follows from isospin symmetry and the second from Fierz identities. We have checked that these novel relations are indeed fulfilled both by the tree-level ChPT amplitudes from Ref. [27] and the most recent set of lattice-QCD results [35, 43].

# 3. Rescattering corrections

By virtue of Lorentz symmetry and parity, the hadronic matrix elements in Eq. (2), of the *B*-violating operators  $Q_{\alpha}^{AB}$  defined in Eq. (1), between a pseudoscalar meson *P* with momentum p' and a nucleon *N* with momentum p, can be decomposed as

$$X_i^{AB} = P_B \Big[ W_0^{X_i^{AB}}(s) + \frac{q}{m_N} W_1^{X_i^{AB}}(s) \Big] u_N(p), \tag{4}$$

where the two form factors  $W_{0,1}^{X_0^{AB}}$  depend on the momentum transfer  $s=q^2=(p-p')^2$ . Since the strong final-state interactions are universal, depending only on the isospin in a given channel, we will drop the chirality and operator indices for the remainder of this section. While the normalization of the form factors requires a lattice-QCD calculation, see Table 2, their momentum dependence is strongly constrained by unitarity, in




Figure 1: Unitarity diagram for the proton-decay matrix elements. The solid and dashed lines denote nucleons (N, N') and pseudoscalar mesons (P, P'), respectively, while the cross refers to an insertion of the B-violating operator  $O^B$  and the gray blob to the meson-nucleon scattering amplitude  $T_{PN}$ . The short-dashed line indicates the cut that produces the imaginary parts in  $W_{0,1}(s)$ .

terms of the imaginary part that is generated by the mesonnucleon rescattering, see Fig. 1.

We write the corresponding amplitude for  $N(p)P(k) \rightarrow N(p')P(k')$  as

$$T_{PN} = \bar{u}_N(p') \left[ A(s,t) + \frac{k + k'}{2} B(s,t) \right] u_N(p), \tag{5}$$

with Mandelstam variables

$$s = (p + k)^2, t = (p - p')^2,$$
 (6)

where the latter is related to the scattering angle z via

$$t = -2|\mathbf{k}|^2(1-z), \qquad |\mathbf{k}|^2 = \frac{\lambda(s, m_N^2, M_P^2)}{4s},$$
 (7)

and  $\lambda(a, b, c) = a^2 + b^2 + c^2 - 2(ab + ac + bc)$ . The partial waves are obtained from the projection [44]

$$f_{l\pm}(s) = \frac{1}{16\pi\sqrt{s}} \Big\{ (E + m_N) [A_l(s) + (\sqrt{s} - m_N)B_l(s)] + (E - m_N) [-A_{l\pm 1}(s) + (\sqrt{s} + m_N)B_{l\pm 1}(s)] \Big\},$$
(8)

where  $E = (s + m_N^2 - M_P^2)/(2\sqrt{s})$ , l is the orbital angular momentum,  $j = |l \pm 1/2|$ , and

$$Y_l(s) = \int_{-1}^1 dz \, P_l(z) Y(s, t) \Big|_{t = -2|\mathbf{k}|^2 (1-z)},\tag{9}$$

with  $Y \in \{A, B\}$  and Legendre polynomials  $P_l(z)$ .

The unitarity constraint from the rescattering diagram in Fig. 1 becomes shortest once expressed in terms of the linear combinations

$$W_{\pm}(s) = W_0(s) \pm \frac{\sqrt{s}}{m_N} W_1(s),$$
 (10)

both of which fulfill unitarity relations<sup>2</sup>

Im 
$$W_+(s) = |\mathbf{k}|W_+(s)[f_{0+}(s)]^*$$
,

$$\operatorname{Im} W_{-}(s) = |\mathbf{k}|W_{-}(s)[f_{1-}(s)]^{*}, \tag{11}$$

that can be solved explicitly in terms of an Omnès representation [45]. To this end, one writes the partial waves in terms of phase shifts  $\delta_{l+}^I$ ,

$$f_{l\pm}^{I}(s) = \frac{e^{i\delta_{l\pm}^{I}}}{|\mathbf{k}|} \sin \delta_{l\pm}^{I},\tag{12}$$

where I refers to the isospin of the PN system. Diagonalizing the unitarity relation in isospin space, we obtain that the solutions for  $U_{1A}$  and  $S_{5A}$  have I=1/2,  $S_{1A}$  and  $S_{2A}+S_{4A}$  have I=1, and  $S_{3A}$  and  $S_{2A}-S_{4A}$  have I=0, see Appendix B for details. These isospin numbers match the baryon poles  $(B=N,\Sigma,\Lambda,$  respectively) that can be present in the matrix elements. The ChPT analysis from Ref. [27] shows that such poles only occur in  $W_-$ , with residues

$$\operatorname{Res} W_{-}(s)\Big|_{s=m_{B}^{2}} = \frac{2m_{B}^{3}}{m_{N}}W_{1}(0). \tag{13}$$

The rescattering is described by Omnès factors

$$\Omega_{l\pm}^{I}(s) = \exp\left\{\frac{s}{\pi} \int_{s_{th}}^{\infty} ds' \frac{\delta_{l\pm}^{I}(s')}{s'(s'-s)}\right\},\tag{14}$$

where  $s_{th} = (m_N + M_P)^2$ , and if it were not for the kinematic singularity due to  $\sqrt{s}$  in Eq. (10), the solution for  $W_{\pm}(s)$  would simply become a polynomial times  $\Omega^I_{l\pm}(s)$ . Such kinematic singularities often arise in the transition from a set of scalar functions for which dispersion relations can be written,  $W_{0,1}(s)$ , to the one for which unitarity constraints are simplest,  $W_{\pm}(s)$ , see Refs. [46-51] for extensive discussions of this point in the context of hadronic light-by-light scattering. In this case, a solution can be found by diagonalizing the kernel functions in the dispersion integrals [50], but here this strategy fails because  $W_{+}$ are unitarized with different phase shifts, in such a way that the linear combinations of Eq. (11) that reproduce  $W_{0,1}$  no longer take the form of a simple Omnès problem. Instead, we construct a solution as follows: first, we observe that in the linear combination  $W_0(s) = (W_+(s) + W_-(s))/2$  the square-root singularity drops out. Since  $\Omega_{\pm}^{I}$  takes care of the right-hand cut, we can therefore make the ansatz<sup>3</sup>

$$W_0(s) = W_0(0) \Big[ (1 - \alpha) \Omega_{0+}^I(s) + \alpha \frac{m_B^2}{m_B^2 - s} \Omega_{1-}^I(s) \Big], \tag{15}$$

where we added the single-baryon pole that originates from  $W_{-}(s)$ . Next, we observe that the product

$$W_{+}(s)W_{-}(s) = \left[W_{0}(s)\right]^{2} - \frac{s}{m_{N}^{2}}\left[W_{1}(s)\right]^{2}$$
 (16)

<sup>&</sup>lt;sup>2</sup>These relations are obtained by evaluating the unitarity diagram in Fig. 1. Since  $W_{\pm}$  are unitarized with different  $\pi N$  phase shifts, the result depends critically on the sign convention for  $W_1$ ; we follow the choice of Refs. [27, 35].

<sup>&</sup>lt;sup>3</sup>Unitarity and analyticity determine the Omnès solutions up to a polynomial, whose degree can often be constrained by the asymptotic behavior of the amplitude or form factor, see, e.g., Ref. [52] for the case of the electromagnetic form factor of the pion. Given the limited kinematic range accessible here, we instead construct the solutions with the minimal amount of polynomial freedom required to satisfy all available constraints.

is again free of kinematic singularities, and fulfills an Omnès representation using the sum of the two phase shifts. In this case, we can thus make the ansatz

$$W_{+}(s)W_{-}(s) = [W_{0}(0)]^{2}\Omega_{0+}^{I}(s)\Omega_{1-}^{I}(s)\frac{m_{B}^{2}}{m_{P}^{2}-s}(1+\beta s), \quad (17)$$

where we kept a non-trivial order in the polynomial to be able to impose the normalization of  $W_1(s)$ . This yields

$$\beta = (1 - 2\alpha) \left[ \dot{\Omega}_{0+}^{I} - \dot{\Omega}_{1-}^{I} - \frac{1}{m_B^2} \right] - \frac{\left[ W_1(0) \right]^2}{m_N^2 \left[ W_0(0) \right]^2}, \tag{18}$$

where  $\dot{\Omega}_{l\pm}^I = d\Omega_{l\pm}^I/ds|_{s=0}$ . Finally, we determine  $\alpha$  from the residue (13). Combining  $W_0(s)$  from Eq. (15) and  $W_+(s)W_-(s)$  from Eq. (17) gives  $W_1(s)$  via Eq. (16), where the sign needs to be chosen in accordance with the overall signs determined in lattice QCD, see Table 2. In particular,  $W_\pm(s)$  can be identified by the fact that the single-baryon pole only contributes to  $W_-$ . Matching its residue to Eq. (13) gives

$$\alpha = -\frac{m_B}{m_N} \frac{W_1(0)}{W_0(0)},\tag{19}$$

which concludes the derivation of our dispersive representation for  $W_{0,1}(s)$ . The resulting expression implements the normalizations from lattice QCD, fulfills the unitarity constraint (11), and in the limit  $\Omega_{l\pm}^I(s)=1$  reduces to the tree-level ChPT result [27]

$$W_0^{\text{ChPT}}(s) = W_0(0) \left[ 1 - \frac{m_B}{m_N} \frac{W_1(0)}{W_0(0)} \frac{s}{m_B^2 - s} \right],$$

$$W_1^{\text{ChPT}}(s) = W_1(0) \frac{m_B^2}{m_P^2 - s},$$
(20)

in such a way that the dispersive result can be considered a unitarized version of Eq. (20).

Our numerical results are shown in Fig. 2 for a representative set of form factors. The low-energy phase shifts for the  $\pi N$  case are known to high precision from Roy-Steiner equations [53–56], in combination with experimental input from pionic atoms [57-62] or cross sections [63], while modern phaseshift analyses extend to 2.5 GeV [64, 65]. Since the main uncertainties arise from the high-energy continuation of the phase shift in the evaluation of  $\Omega^{I}_{l\pm}$ —compared to which other uncertainties, e.g., from higher intermediate states or higher orders in the matching to ChPT will be subdominant—we use the latter solution, and estimate the uncertainties from the variation observed when keeping the phase constant above a matching point chosen within [1.8, 2.5] GeV in the evaluation of the integral in the Omnès factors. Proceeding similarly for the KN case [66, 67] leads to the bands in Fig. 2, in comparison to the dashed lines representing the tree-level ChPT result (20). As expected, one sees that the unitarity corrections are most relevant for the decay into a pion, because the phase space in the kaon modes is much more limited.




Figure 2: Form factors  $\bar{W}_i^{X_{iL}}(s) = W_i^{X_{iL}}(s)/W_i^{X_{iL}}(0)$  for representative cases involving  $\pi N$ , I=1/2 (top), KN, I=1 (middle), and KN, I=0 (bottom) rescattering. The dashed line indicates the result without these unitarity corrections, which corresponds to the tree-level ChPT result (20). The results are shown for the entire range  $s \leq (m_N - M_\pi)^2$  relevant for  $p \to \pi^0 \ell^+ \bar{\nu}_\ell \nu_\tau$ , while the dotted line indicates the kinematic limit  $s = (m_N - M_K)^2$  for the kaon modes.

#### 4. Tau-induced proton decay

The rate for the two-body decay  $N \to P\bar{\ell}$  can be written as [35]

$$\Gamma[N \to P\bar{\ell}] = \frac{|\mathbf{q}|E_{\ell}}{8\pi m_N} \left[ \sum_{A=L,R} |W_A^{\ell}|^2 - \frac{2m_{\ell}}{E_{\ell}} \operatorname{Re}(W_L^{\ell} W_R^{\ell*}) \right], \quad (21)$$

where the form factors

$$W_{L}^{\ell}(s) = \sum_{\alpha,A} \left[ C_{\alpha}^{AL} W_{0}^{X^{AL}}(s) - \frac{s}{m_{N} m_{\ell}} C_{\alpha}^{AR} W_{1}^{X^{AR}}(s) \right],$$

$$W_{R}^{\ell}(s) = \sum_{\alpha,A} \left[ C_{\alpha}^{AR} W_{0}^{X^{AR}}(s) - \frac{s}{m_{N} m_{\ell}} C_{\alpha}^{AL} W_{1}^{X^{AL}}(s) \right], \tag{22}$$

are evaluated at  $s = m_{\ell}^2$  and

$$E_{\ell} = \frac{m_N^2 + m_{\ell}^2 - M_P^2}{2m_N}, \quad |\mathbf{q}| = \frac{\lambda^{1/2}(m_N^2, m_{\ell}^2, M_P^2)}{2m_N}.$$
 (23)

The sums in Eq. (22) run over all contributing operators  $\alpha$  and the two possible parity assignments A=L,R, see Appendix C for the explicit expressions in a given decay channel. Using the lifetime limits from Table 1, stringent constraints  $|C_{\alpha}| \lesssim (10^{-15}/\,\text{GeV})^2$  on most Wilson coefficients (given at the nucleon scale) can be obtained. An exception concerns those operators that only contribute to the processes with a charged




Figure 3: Decay  $N \to P\ell^+\nu_\ell\bar{\nu}_\tau$  via an off-shell  $\tau$ .  $O^B$  and  $G_F$  indicate the insertion of the *B*-violating operator and the electroweak vertex, respectively.

lepton, since the decay with a  $\tau$  final state is kinematically forbidden. Therefore, no limits on the corresponding  $\tau$ -flavored Wilson coefficients can be extracted from two-body decays.

However, the  $\tau$  lepton can be off-shell, thus inducing four-body decays, see Fig. 3, whose width can be expressed as

$$\Gamma[N \to P\ell^+ \nu_{\ell} \bar{\nu}_{\tau}] = \int_0^{s_{\text{max}}} ds \frac{G_F^2 s^2 \lambda^{1/2} (m_N^2, s, M_P^2)}{6144 \pi^5 m_N^3 (m_{\tau}^2 - s)^2} \times \left[ (m_N^2 + s - M_P^2) \left( m_{\tau}^2 |W_R^{\tau}|^2 + s |W_L^{\tau}|^2 \right) - 4 s m_N m_{\tau} \text{Re} \left( W_L^{\tau} W_R^{\tau*} \right) \right], \tag{24}$$

where  $s_{\text{max}} = (m_N - M_P)^2$ . For simplicity, Eq. (24) is given in the limit  $m_\ell = 0$ , see Appendix D for the full expression. Numerically, we find for the special case  $C_{\tau\alpha}^{AB} \equiv C_{\tau\alpha}^{LL} = C_{\tau\alpha}^{LR} = C_{\tau\alpha}^{RR}$ 

$$\Gamma[p \to \pi^0 \ell^+ \nu_\ell \bar{\nu}_\tau] = 14.3(1.8) \left| \frac{C_{\tau u d u}^{AB}}{(10^{10} \,\text{GeV})^{-2}} \right|^2 \frac{1}{10^{30} \,\text{yr}}, 
\Gamma[n \to \pi^- \ell^+ \nu_\ell \bar{\nu}_\tau] = 26.2(3.3) \left| \frac{C_{\tau u d u}^{AB}}{(10^{10} \,\text{GeV})^{-2}} \right|^2 \frac{1}{10^{30} \,\text{yr}}, 
\Gamma[p \to K^0 \ell^+ \nu_\ell \bar{\nu}_\tau] = 0.81(1) \left| \frac{C_{\tau u s u}^{AB}}{(10^{10} \,\text{GeV})^{-2}} \right|^2 \frac{1}{10^{30} \,\text{yr}},$$
(25)

where the uncertainty corresponds to the bands in Fig. 2 (the coefficients without rescattering corrections are 7.2, 13.3, and 0.84, respectively). As expected, the resulting numerical limits are significantly weaker than in the two-body decay,  $|C_{\alpha}| \lesssim (10^{-10}/\,\text{GeV})^2$  at the nucleon scale—due to a combination of the weaker inclusive limits, see Table 1, and the suppression from  $G_F$  as well as the four-particle phase space—but we emphasize that the corresponding Wilson coefficients were previously completely unconstrained. A comprehensive analysis in SMEFT, including the pertinent RG corrections [68], will be given in Ref. [69].

## 5. Conclusions

Processes such as  $p \to \pi^0 \ell^+ \nu_\ell \bar{\nu}_\tau$  are phenomenologically interesting, as they can probe Wilson coefficients that are not accessible in simpler two-body decays. I.e., while the decay to

 $\tau$  leptons is kinematically forbidden, *B*-violating operators involving a  $\tau$  lepton do become constrained by such four-body processes. However, for calculating the corresponding decay widths, the momentum dependence of the related hadronic matrix elements is necessary.

In this work, we examined these form factors for nucleon-to-meson transitions in dispersion theory and derived a representation that allows one to calculate the momentum dependence of the corresponding form factors including unitarity corrections from the meson–nucleon rescattering. With normalizations determined from lattice QCD, our results allowed us to calculate semileptonic nucleon decays involving multiple leptons in the final state [70–72] and to derive novel bounds on the Wilson coefficients of previously unconstrained dimension-6 operators involving  $\tau$  leptons.

#### Acknowledgments

We thank Bastian Kubis and Peter Stoffer for helpful discussions, and Taku Izubuchi and Sergey Syritsyn for communication regarding Ref. [35]. Financial support from the SNSF (Project Nos. PP00P21\_76884 and PCEFP2\_181117) is gratefully acknowledged.

#### Appendix A. Operators in SMEFT

The four  $SU(2)_L$ -invariant dimension-6 operators giving rise to nucleon decay are

$$Q_{duq} = \varepsilon^{\alpha\beta\gamma} \varepsilon_{jk} \left[ \left( d_p^{\alpha} \right)^T C u_r^{\beta} \right] \left[ \left( q_s^{\gamma j} \right)^T C L_t^{k} \right],$$

$$Q_{qqu} = \varepsilon^{\alpha\beta\gamma} \varepsilon_{jk} \left[ \left( q_p^{\alpha j} \right)^T C q_r^{\beta k} \right] \left[ \left( u_s^{\gamma} \right)^T C e_t \right],$$

$$Q_{qqq} = \varepsilon^{\alpha\beta\gamma} \varepsilon_{jn} \varepsilon_{km} \left[ \left( q_p^{\alpha j} \right)^T C q_r^{\beta k} \right] \left[ \left( q_s^{\gamma m} \right)^T C L_t^{n} \right],$$

$$Q_{duu} = \varepsilon^{\alpha\beta\gamma} \left[ \left( d_p^{\alpha} \right)^T C u_r^{\beta} \right] \left[ \left( u_s^{\gamma} \right)^T C e_t \right], \tag{A.1}$$

where q/L denote the quark/lepton  $SU(2)_L$  doublets, u, d, and e the quark and lepton singlets,  $p, \ldots / j, \ldots / \alpha, \ldots$  are generation/ $SU(2)_L$ /color indices, and C denotes the charge conjugation operator. After electroweak symmetry breaking, we can write the resulting effective Lagrangian below the weak scale as

$$\begin{split} \mathcal{L} &= \bar{\ell}_{t}^{c} \Big[ - C_{qqq}^{111t} Q_{udu}^{LL} + 2 C_{qqu}^{111t} Q_{udu}^{LR} - C_{duq}^{111t} Q_{udu}^{RL} \\ &- C_{duu}^{111t} Q_{udu}^{RR} - C_{qqq}^{121t} Q_{usu}^{LL} + (C_{qqu}^{121t} + C_{qqu}^{211t}) Q_{usu}^{LR} \\ &- C_{duq}^{211t} Q_{usu}^{RL} - C_{duu}^{211t} Q_{usu}^{RR} \Big] \\ &+ \bar{\nu}_{t}^{c} \Big[ C_{qqq}^{111t} Q_{udd}^{LL} + C_{duq}^{111t} Q_{udd}^{LL} + C_{qqq}^{211t} Q_{usd}^{LL} \\ &+ C_{duq}^{211t} Q_{usd}^{RL} + (C_{qqq}^{121t} - C_{qqq}^{211t}) Q_{dsu}^{LL} \\ &+ C_{qqq}^{112t} Q_{uds}^{LL} + C_{duq}^{112t} Q_{uds}^{RL} \Big], \end{split} \tag{A.2}$$

where the Wilson coefficients refer to the respective flavor combinations in Eq. (A.1), the quark part of the effective operators is defined in Eq. (1), and of course RG corrections need to be

added to relate low and high scale. Note that assuming that B violation is realized far above the electroweak scale reduces the number of operators, otherwise, also different flavor combinations with  $\Delta B \neq \Delta L$  appear [73, 74].

### Appendix B. Isospin symmetry

In addition to the isospin relations given in the main text, isospin symmetry also determines the neutron matrix elements [33]

$$\langle \pi^{-} | \left[ \bar{u}^{c} P_{A} d \right] u_{L} | n \rangle = -\sqrt{2} \langle \pi^{0} | \left[ \bar{u}^{c} P_{A} d \right] d_{L} | n \rangle = U_{1A},$$

$$\langle K^{+} | \left[ \bar{d}^{c} P_{A} S \right] d_{L} | n \rangle = -S_{1A},$$

$$\langle K^{0} | \left[ \bar{u}^{c} P_{A} S \right] d_{L} | n \rangle = -S_{4A},$$

$$\langle K^{0} | \left[ \bar{u}^{c} P_{A} d \right] s_{L} | n \rangle = S_{3A},$$

$$\langle K^{0} | \left[ \bar{d}^{c} P_{A} S \right] u_{L} | n \rangle = -S_{2A},$$

$$\langle \eta | \left[ \bar{u}^{c} P_{A} d \right] d_{L} | n \rangle = S_{5A}.$$
(B.1)

To work out the isospin quantum numbers of the rescattering amplitudes, we sum up all possible intermediate states in each charge channel and use the isospin relations (B.1) to obtain a closed system of unitarity relations

$$\operatorname{Im} U_{1A} = \left[ \langle \pi^0 p | T | \pi^0 p \rangle - \sqrt{2} \langle \pi^0 p | T | \pi^+ n \rangle \right]^* U_{1A},$$

$$\operatorname{Im} U_{1A} = \left[ \langle \pi^- p | T | \pi^- p \rangle - \frac{1}{\sqrt{2}} \langle \pi^- p | T | \pi^0 n \rangle \right]^* U_{1A},$$

$$\operatorname{Im} S_{1A} = \langle \bar{K}^0 p | T | \bar{K}^0 p \rangle^* S_{1A},$$

$$\operatorname{Im} S_{3A} = \left[ \langle K^- p | T | K^- p \rangle + \langle K^- p | T | \bar{K}^0 n \rangle \right]^* S_{3A},$$

$$\operatorname{Im} \begin{bmatrix} S_{2A} \\ S_{4A} \end{bmatrix} = \begin{bmatrix} \langle K^- p | T | K^- p \rangle & -\langle K^- p | T | \bar{K}^0 n \rangle \\ -\langle K^- p | T | \bar{K}^0 n \rangle & \langle K^- p | T | K^- p \rangle \end{bmatrix}^* \begin{bmatrix} S_{2A} \\ S_{4A} \end{bmatrix},$$

$$\operatorname{Im} S_{5A} = \langle \eta p | T | \eta p \rangle^* S_{5A}. \tag{B.2}$$

Expressing each amplitude in terms of isospin quantum numbers and diagonalizing the relation for  $S_{2A}$ ,  $S_{4A}$ , we obtain

$$\operatorname{Im} U_{1A} = [T^{1/2}]^* U_{1A},$$

$$\operatorname{Im} S_{5A} = [T^{1/2}]^* S_{5A},$$

$$\operatorname{Im} S_{1A} = [T^1]^* S_{1A},$$

$$\operatorname{Im} (S_{2A} + S_{4A}) = [T^1]^* (S_{2A} + S_{4A}),$$

$$\operatorname{Im} S_{3A} = [T^0]^* S_{3A},$$

$$\operatorname{Im} (S_{2A} - S_{4A}) = [T^0]^* (S_{2A} - S_{4A}),$$
(B.3)

which determines the isospin quantum numbers of the rescattering corrections as stated in the main text. Moreover, this analysis in terms of isospin symmetry leads one in a straightforward way to the relations (3), since the corresponding matrix elements carry the same isospin. Further relations could be derived from SU(3) flavor symmetry, but would only become relevant in the study of hyperon decays.

#### Appendix C. Two-body decay

The explicit decomposition of  $W_A$  in Eq. (22) for the different decay channels in terms of SMEFT Wilson coefficients reads

$$\begin{split} W_{L}[p \to \pi^{0}\ell^{+}] &\simeq -\frac{1}{\sqrt{2}} (C_{qqq}^{111t}W_{0}^{U_{1L}} + C_{duq}^{111t}W_{0}^{U_{1R}}), \\ W_{R}[p \to \pi^{0}\ell^{+}] &\simeq \frac{1}{\sqrt{2}} (2C_{qqq}^{111t}W_{0}^{U_{1R}} - C_{duq}^{111t}W_{0}^{U_{1L}}), \\ W_{L}[p \to \pi^{+}\bar{\nu}] &\simeq C_{qqq}^{111t}W_{0}^{U_{1L}} + C_{duq}^{111t}W_{0}^{U_{1R}}, \\ W_{L}[p \to K^{0}\ell^{+}] &\simeq -C_{qqq}^{121t}W_{0}^{S_{1L}} - C_{duq}^{211t}W_{0}^{S_{1R}}, \\ W_{R}[p \to K^{0}\ell^{+}] &\simeq (C_{qqq}^{121t} + C_{qqq}^{211t})W_{0}^{S_{1R}} - C_{duu}^{211t}W_{0}^{S_{1L}}, \\ W_{L}[p \to K^{+}\bar{\nu}] &\simeq (C_{qqq}^{12t} + C_{qqq}^{21t})W_{0}^{S_{3R}} - C_{duu}^{211t}W_{0}^{S_{3L}}, \\ W_{L}[p \to \eta\ell^{+}] &\simeq -C_{qqq}^{111t}W_{0}^{S_{2R}} - C_{duq}^{111t}W_{0}^{S_{2R}}, \\ W_{R}[p \to \eta\ell^{+}] &\simeq -C_{qqq}^{111t}W_{0}^{S_{3R}} - C_{duq}^{111t}W_{0}^{S_{3R}}, \\ W_{L}[n \to \pi^{-}\ell^{+}] &\simeq -C_{qqq}^{111t}W_{0}^{U_{1L}} - C_{duq}^{111t}W_{0}^{U_{1R}}, \\ W_{R}[n \to \pi^{-}\ell^{+}] &\simeq 2C_{qqu}^{111t}W_{0}^{U_{1R}} - C_{duq}^{111t}W_{0}^{U_{1L}}, \\ W_{L}[n \to \pi^{0}\bar{\nu}] &\simeq -\frac{1}{\sqrt{2}} (C_{qqq}^{111t}W_{0}^{U_{1L}} + C_{duq}^{111t}W_{0}^{U_{1R}}), \\ W_{L}[n \to K^{0}\bar{\nu}] &\simeq (C_{qqq}^{112t} + C_{qqq}^{211t})W_{0}^{S_{3L}} - C_{qqq}^{121t}W_{0}^{S_{2L}} - C_{duq}^{211t}W_{0}^{S_{3R}}, \\ W_{L}[n \to \eta\bar{\nu}] &\simeq C_{qqq}^{111t}W_{0}^{S_{4R}} + C_{duq}^{112t}W_{0}^{S_{3R}}, \\ W_{L}[n \to \eta\bar{\nu}] &\simeq C_{qqq}^{111t}W_{0}^{S_{4R}} + C_{duq}^{111t}W_{0}^{S_{5R}}, \\ (C.1) \end{aligned}$$

where the dependence of  $W_0$  on s has been suppressed and we only give the  $W_0$  term since the correction from  $W_1$  follows directly from Eq. (22). Equation (3) has been used to simplify the expressions for  $W_L[p \to K^+\bar{\nu}]$  and  $W_L[n \to K^0\bar{\nu}]$ . The notation using  $\simeq$  is meant to indicate that the expression in terms of SMEFT coefficients does not apply at the low scale without accounting for RG corrections.

# Appendix D. Four-body decay

Denoting momenta as  $p(p) \to \tau^+(q)P(p')$ ,  $\tau^+(q) \to \bar{\nu}_{\tau}(q_1)\ell^+(q_2)\nu_{\ell}(q_3)$ , we find as intermediate result for the spin-averaged squared matrix element:

$$|\bar{M}|^{2} = \frac{64G_{F}^{2}q_{1} \cdot q_{2}}{(m_{\tau}^{2} - q^{2})^{2}} \Big[ \Big( m_{\tau}^{2} |W_{R}^{\tau}|^{2} - q^{2} |W_{L}^{\tau}|^{2} \Big) p \cdot q_{3} + 2 \Big( |W_{L}^{\tau}|^{2} p \cdot q - m_{N} m_{\tau} \operatorname{Re} (W_{L}^{\tau} W_{R}^{\tau*}) \Big) q \cdot q_{3} \Big].$$
 (D.1)

We perform the phase-space integration following Refs. [75, 76] by subsequent reduction to two-body decays. This gives

$$\Gamma[N \to P\ell^+ \nu_{\ell} \bar{\nu}_{\tau}] = \frac{G_F^2}{512\pi^5 m_N^3} \int_{m_{\ell}^2}^{s_{\text{max}}} ds \frac{\lambda^{1/2} (m_N^2, s, M_P^2)}{s^2 (m_{\tau}^2 - s)^2} \times \left[ (m_N^2 + s - M_P^2) (m_{\tau}^2 |W_R^{\tau}|^2 + s|W_L^{\tau}|^2) - 4s m_N m_{\tau} \text{Re} \left( W_L^{\tau} W_R^{\tau*} \right) \right] I_{\ell}(s), \tag{D.2}$$

where

$$I_{\ell}(s) = \int_{m_{\ell}^{2}}^{s} dk^{2} \frac{(k^{2} - m_{\ell}^{2})^{2}(s - k^{2})^{2}}{k^{2}}$$

$$= \frac{1}{12} \left(s^{4} - 8s^{3}m_{\ell}^{2} + 8sm_{\ell}^{6} - m_{\ell}^{8}\right) + s^{2}m_{\ell}^{4} \log \frac{s}{m_{\ell}^{2}}$$
(D.3)

coincides with the standard phase-space factor for the leptonic decay of the muon, see, e.g., Refs. [77, 78].

#### References

- [1] J. C. Pati and A. Salam, Phys. Rev. D 8, 1240 (1973).
- [2] H. Georgi and S. L. Glashow, Phys. Rev. Lett. 32, 438 (1974).
- [3] H. Fritzsch and P. Minkowski, Annals Phys. 93, 193 (1975).
- [4] A. Takenaka et al. (Super-Kamiokande), Phys. Rev. D 102, 112011 (2020), arXiv:2010.16098 [hep-ex].
- K. Abe et al. (Super-Kamiokande), Phys. Rev. Lett. 113, 121802 (2014), arXiv:1305.4391 [hep-ex].
- [6] K. Kobayashi et al. (Super-Kamiokande), Phys. Rev. D 72, 052007 (2005), arXiv:hep-ex/0502026.
- [7] R. Matsumoto et al. (Super-Kamiokande), Phys. Rev. D 106, 072003 (2022), arXiv:2208.13188 [hep-ex].
- K. Abe et al. (Super-Kamiokande), Phys. Rev. D 90, 072005 (2014), arXiv:1408.1195 [hep-ex].
- [9] K. Abe et al. (Super-Kamiokande), Phys. Rev. D 96, 012003 (2017), arXiv:1705.07221 [hep-ex].
- [10] C. McGrew et al., Phys. Rev. D 59, 052004 (1999).
- [11] J. Learned, F. Reines, and A. Soni, Phys. Rev. Lett. 43, 907 (1979), [Erratum: Phys. Rev. Lett. 43, 1626 (1979)].
- [12] M. L. Cherry, M. Deakyne, K. Lande, C. K. Lee, R. I. Steinberg, and B. T. Cleveland, Phys. Rev. Lett. 47, 1507 (1981).
- [13] R. L. Workman et al. (Particle Data Group), PTEP 2022, 083C01 (2022).
- [14] K. Abe et al. (Hyper-Kamiokande), (2018), arXiv:1805.04163 [physics.ins-det].
- [15] B. Abi et al. (DUNE), (2020), arXiv:2002.03005 [hep-ex].
- [16] A. Abusleme et al. (JUNO), Prog. Part. Nucl. Phys. 123, 103927 (2022), arXiv:2104.02565 [hep-ex].
- F. Wilczek and A. Zee, Phys. Rev. Lett. 43, 1571 (1979).
- [18] S. Weinberg, Phys. Rev. Lett. 43, 1566 (1979).
- [19] W. Buchmüller and D. Wyler, Nucl. Phys. B 268, 621 (1986).
- [20] B. Grzadkowski, M. Iskrzyński, M. Misiak, and J. Rosiek, JHEP 10, 085 (2010), arXiv:1008.4884 [hep-ph].
- [21] J. C. Helo, M. Hirsch, and T. Ota, Phys. Rev. D 99, 095021 (2019), arXiv:1904.00036 [hep-ph].
- J. Heeck and V. Takhistov, Phys. Rev. D 101, 015005 (2020), arXiv:1910.07647 [hep-ph].
- [23] M. Claudson, M. B. Wise, and L. J. Hall, Nucl. Phys. B 195, 297 (1982).
- [24] S. Chadha and M. Daniel, Nucl. Phys. B 229, 105 (1983).
- [25] S. Chadha and M. Daniel, Phys. Lett. B 137, 374 (1984).
- [26] O. Kaymakcalan, C.-H. Lo, and K. C. Wali, Phys. Rev. D 29, 1962 (1984).
- [27] S. Aoki et al. (JLQCD), Phys. Rev. D 62, 014506 (2000), arXiv:heplat/9911026.
- [28] M. B. Gavela, A. Le Yaouanc, L. Oliver, O. Pene, and J. C. Raynal, Phys. Rev. D 23, 1580 (1981).
- [29] T. Okazaki and K. Fujii, Phys. Rev. D 27, 188 (1983).

arXiv:hep-lat/0402026.

- [30] A. Martin and G. C. Stavenga, Phys. Rev. D 85, 095010 (2012), arXiv:1110.2188 [hep-ph].
- Y. Hara, S. Itoh, Y. Iwasaki, and T. Yoshie, Phys. Rev. D 34, 3399 (1986).
- [32] N. Tsutsui et al. (CP-PACS, JLQCD), Phys. Rev. D 70, 111501 (2004),
- [33] Y. Aoki, E. Shintani, and A. Soni, Phys. Rev. D 89, 014505 (2014), arXiv:1304.7424 [hep-lat].
- [34] Y. Aoki, T. Izubuchi, E. Shintani, and A. Soni, Phys. Rev. D 96, 014506 (2017), arXiv:1705.01338 [hep-lat].
- [35] J.-S. Yoo, Y. Aoki, P. Boyle, T. Izubuchi, A. Soni, and S. Syritsyn, Phys. Rev. D 105, 074501 (2022), arXiv:2111.01608 [hep-lat].
- [36] G. Höhler, E. Pietarinen, I. Sabba Stefanescu, F. Borkowski, G. G. Simon, V. H. Walther, and R. D. Wendling, Nucl. Phys. B 114, 505 (1976).

- [37] J. Gasser, H. Leutwyler, and M. E. Sainio, Phys. Lett. B 253, 260 (1991).
- [38] M. Hoferichter, C. Ditsche, B. Kubis, and U.-G. Meißner, JHEP 06, 063 (2012), arXiv:1204.6251 [hep-ph].
- M. Hoferichter, J. Ruiz de Elvira, B. Kubis, and U.-G. Meißner, Phys. Rev. Lett. 115, 092301 (2015), arXiv:1506.04142 [hep-ph].
- M. Hoferichter, B. Kubis, J. Ruiz de Elvira, H.-W. Hammer, and U.-G. Meißner, Eur. Phys. J. A 52, 331 (2016), arXiv:1609.06722 [hep-ph].
- V. Cirigliano, A. Crivellin, and M. Hoferichter, Phys. Rev. Lett. 120, 141803 (2018), arXiv:1712.06595 [hep-ph].
- [42] M. Hoferichter, B. Kubis, J. Ruiz de Elvira, and P. Stoffer, Phys. Rev. Lett. 122, 122001 (2019), [Erratum: Phys. Rev. Lett. 124, 199901 (2020)], arXiv:1811.11181 [hep-ph].
- S. Syritsyn, (2023), private communication.
- [44] W. R. Frazer and J. R. Fulco, Phys. Rev. 119, 1420 (1960).
- [45] R. Omnès, Nuovo Cim. 8, 316 (1958).
- [46] G. Colangelo, M. Hoferichter, M. Procura, and P. Stoffer, JHEP 09, 091 (2014), arXiv:1402.7081 [hep-ph].
- [47] G. Colangelo, M. Hoferichter, M. Procura, and P. Stoffer, JHEP 09, 074 (2015), arXiv:1506.01386 [hep-ph].
- G. Colangelo, M. Hoferichter, M. Procura, and P. Stoffer, Phys. Rev. Lett. 118, 232001 (2017), arXiv:1701.06554 [hep-ph].
- [49] G. Colangelo, M. Hoferichter, M. Procura, and P. Stoffer, JHEP 04, 161 (2017), arXiv:1702.07347 [hep-ph].
- [50] M. Hoferichter and P. Stoffer, JHEP 07, 073 (2019), arXiv:1905.13198 [hep-ph].
- M. Hoferichter and P. Stoffer, JHEP 05, 159 (2020), arXiv:2004.06127 [hep-ph].
- [52] G. Colangelo, M. Hoferichter, and P. Stoffer, JHEP 02, 006 (2019), arXiv:1810.00007 [hep-ph].
- [53] C. Ditsche, M. Hoferichter, B. Kubis, and U.-G. Meißner, JHEP 06, 043 (2012), arXiv:1203.4758 [hep-ph].
- M. Hoferichter, J. Ruiz de Elvira, B. Kubis, and U.-G. Meißner, Phys. Rept. 625, 1 (2016), arXiv:1510.06039 [hep-ph].
- [55] M. Hoferichter, J. Ruiz de Elvira, B. Kubis, and U.-G. Meißner, Phys. Rev. Lett. 115, 192301 (2015), arXiv:1507.07552 [nucl-th].
- [56] D. Siemens, J. Ruiz de Elvira, E. Epelbaum, M. Hoferichter, H. Krebs, B. Kubis, and U.-G. Meißner, Phys. Lett. B 770, 27 (2017), arXiv:1610.08978 [nucl-th].
- T. Strauch et al., Eur. Phys. J. A 47, 88 (2011), arXiv:1011.2415 [nucl-ex].
- [58] M. Hennebach et al., Eur. Phys. J. A 50, 190 (2014), [Erratum: Eur. Phys. J. A 55, 24 (2019)], arXiv:1406.6525 [nucl-ex].
- [59] A. Hirtl et al., Eur. Phys. J. A 57, 70 (2021).
- [60] V. Baru, C. Hanhart, M. Hoferichter, B. Kubis, A. Nogga, and D. R. Phillips, Phys. Lett. B 694, 473 (2011), arXiv:1003.4444 [nucl-th].
- V. Baru, C. Hanhart, M. Hoferichter, B. Kubis, A. Nogga, and D. R. Phillips, Nucl. Phys. A 872, 69 (2011), arXiv:1107.5509 [nucl-th].
- [62] M. Hoferichter, J. Ruiz de Elvira, B. Kubis, and U.-G. Meißner, Phys. Lett. B 843, 138001 (2023), arXiv:2305.07045 [hep-ph].
- [63] J. Ruiz de Elvira, M. Hoferichter, B. Kubis, and U.-G. Meißner, J. Phys. G 45, 024001 (2018), arXiv:1706.01465 [hep-ph].
- R. A. Arndt, W. J. Briscoe, I. I. Strakovsky, and R. L. Workman, Phys. Rev. C 74, 045205 (2006), arXiv:nucl-th/0605082.
- [65] R. L. Workman, R. A. Arndt, W. J. Briscoe, M. W. Paris, and I. I. Strakovsky, Phys. Rev. C 86, 035202 (2012), arXiv:1204.2277 [hep-ph].
- [66] J. S. Hyslop, R. A. Arndt, L. D. Roper, and R. L. Workman, Phys. Rev. D 46, 961 (1992).
- R. A. Arndt, I. I. Strakovsky, and R. L. Workman, Phys. Rev. C 68, 042201 (2003), [Erratum: Phys. Rev. C 69, 019901 (2004)], arXiv:nuclth/0308012.
- [68] R. Alonso, H.-M. Chang, E. E. Jenkins, A. V. Manohar, and B. Shotwell, Phys. Lett. B 734, 302 (2014), arXiv:1405.0486 [hep-ph].
- [69] A. Crivellin, M. Hoferichter, and P. Stoffer, In preparation.
- [70] C. Faroughy, S. Prabhu, and B. Zheng, JHEP 06, 073 (2015), arXiv:1409.5438 [hep-ph].
- T. Hambye and J. Heeck, Phys. Rev. Lett. 120, 171801 (2018), arXiv:1712.04871 [hep-ph].
- [72] I. Doršner, S. Fajfer, and O. Sumensari, JHEP 05, 183 (2022), arXiv:2202.08287 [hep-ph].
- [73] A. Kobach, Phys. Lett. B 758, 455 (2016), arXiv:1604.05726 [hep-ph].
- E. E. Jenkins, A. V. Manohar, and P. Stoffer, JHEP 03, 016 (2018), arXiv:1709.04486 [hep-ph].

- [75] F.-K. Guo, B. Kubis, and A. Wirzba, Phys. Rev. D 85, 014014 (2012),
- arXiv:1111.5949 [hep-ph]. [76] M. Zanke, M. Hoferichter, and B. Kubis, JHEP **07**, 106 (2021), arXiv:2103.09829 [hep-ph].
- [77] A. Sirlin and A. Ferroglia, Rev. Mod. Phys. **85**, 263 (2013),
- arXiv:1210.5296 [hep-ph].

  [78] A. Ferroglia, C. Greub, A. Sirlin, and Z. Zhang, Phys. Rev. D 88, 033012 (2013), arXiv:1307.6900 [hep-ph].