

Precise Empirical Determination of Metallicity Dependence of Near-infrared Period-Luminosity Relations for RR Lyrae Variables

ANUPAM BHARDWAJ,^{1,*} MARCELLA MARCONI,¹ MARINA REJKUBA,² RICHARD DE GRIJS,^{3,4} HARINDER P. SINGH,⁵
VITTORIO F. BRAGA,⁶ SHASHI KANBUR,⁷ CHOW-CHOONG NGEOW,⁸ VINCENZO RIPEPI,¹ GIUSEPPE BONO,^{6,9}
GIULIA DE SOMMA,^{1,10} AND MASSIMO DALL'ORA¹

¹*INAF-Osservatorio Astronomico di Capodimonte, Via Moiariello 16, 80131 Napoli, Italy*

²*European Southern Observatory, Karl-Schwarzschild-StraÙe 2, 85748, Garching, Germany*

³*School of Mathematical and Physical Sciences, Macquarie University, Balaclava Road, Sydney NSW 2109, Australia*

⁴*Research Centre for Astronomy, Astrophysics and Astrophotonics, Macquarie University, Balaclava Road, Sydney NSW 2109, Australia*

⁵*Department of Physics and Astrophysics, University of Delhi, Delhi-110007, India*

⁶*INAF-Osservatorio Astronomico di Roma, via Frascati 33, 00040 Monte Porzio Catone, Italy*

⁷*Department of Physics, State University of New York, Oswego, NY 13126, USA*

⁸*Graduate Institute of Astronomy, National Central University, 300 Zhongda Road, 32001 Zhongli, Taiwan*

⁹*Dipartimento di Fisica, Università di Roma Tor Vergata, via della Ricerca Scientifica 1, 00133 Roma, Italy*

¹⁰*INFN-Sez. di Napoli, Compl. Univ. di Monte S. Angelo, Edificio G, Via Cinthia, 80126 Napoli, Italy*

(Accepted Feb 9, 2023)

Submitted to ApJL

ABSTRACT

RR Lyrae variables are excellent population II distance indicators thanks to their well-defined period-luminosity relations (PLRs) at infrared wavelengths. We present results of near-infrared (NIR) monitoring of Galactic globular clusters to empirically quantify the metallicity dependence of NIR PLRs for RR Lyrae variables. Our sample includes homogeneous, accurate, and precise photometric data for 964 RR Lyrae variables in 11 globular clusters covering a large metallicity range ($\Delta[\text{Fe}/\text{H}] \sim 2$ dex). We derive JHK_s band period-luminosity-metallicity (PLZ) and period-Wesenheit-metallicity (PWZ) relations anchored using 346 Milky Way field RR Lyrae stars with *Gaia* parallaxes, and simultaneously solved for independent distances to globular clusters. We find a significant metallicity dependence of ~ 0.2 mag/dex in JHK_s band PLZ and PWZ relations for RR Lyrae stars independent of the adopted metallicity scale. The metallicity coefficients and the zero-points of the empirical PLZ and PWZ relations are in excellent agreement with the predictions from the horizontal branch evolution and pulsation models. Furthermore, RR Lyrae based distances to our sample of globular clusters are also statistically consistent with other independent measurements in the literature. Our recommended empirical JHK_s band PLZ relations for RR Lyrae stars with periods of fundamental mode pulsation (P_f) are:

$$M_J = -0.44 (\pm 0.03) - 1.83 (\pm 0.02) \log(P_f) + 0.20 (\pm 0.02) [\text{Fe}/\text{H}] \quad (\sigma = 0.05 \text{ mag})$$

$$M_H = -0.74 (\pm 0.02) - 2.29 (\pm 0.02) \log(P_f) + 0.19 (\pm 0.01) [\text{Fe}/\text{H}] \quad (\sigma = 0.05 \text{ mag})$$

$$M_{K_s} = -0.80 (\pm 0.02) - 2.37 (\pm 0.02) \log(P_f) + 0.18 (\pm 0.01) [\text{Fe}/\text{H}] \quad (\sigma = 0.05 \text{ mag})$$

1. INTRODUCTION

RR Lyrae variables are old (> 10 Gyr), low-mass ($\sim 0.5-0.8M_\odot$) stars that are located at the intersection of the horizontal branch and the classical instability strip in the Hertzsprung-Russell diagram. These pulsating stars have long been used as distance indicators thanks to the correlation of their visual magnitudes with metallicity (see the reviews by Sandage & Tammann 2006;

Corresponding author: Anupam Bhardwaj
anupam.bhardwaj@gmail.com; anupam.bhardwaj@inaf.it

* Marie-Curie Fellow

Catelan & Smith 2015; Beaton et al. 2018; Bhardwaj 2020, and references therein). Since the pioneering work by Longmore et al. (1986) it has been known that RR Lyrae stars follow a well-defined period-luminosity relation (PLR) in K_s -band. Near-infrared (NIR) PLRs offer advantages over visual magnitude-metallicity relation for distance determinations due to smaller dependence on interstellar extinction, smaller variability amplitude, and less sensitivity to evolutionary effects (Bono et al. 2001; Catelan et al. 2004; Marconi et al. 2015).

Stellar evolution and pulsation models of RR Lyrae stars have predicted a strong effect of metallicity on NIR PLRs (~ 0.18 mag/dex in JHK_s , Catelan et al. 2004; Marconi et al. 2015). The empirical studies however have reported a wide range for the metallicity coefficient of period-luminosity-metallicity (PLZ) relations ($0.03 - 0.23$ mag/dex in K_s , Sollima et al. 2006, 2008; Bono et al. 2011; Muraveva et al. 2015, 2018; Braga et al. 2018; Layden et al. 2019; Neeley et al. 2019; Muhie et al. 2021; Cusano et al. 2021, and references therein). Most of these empirical investigations were carried out only in K_s -band, and very few of them included all three JHK_s filters. Neeley et al. (2019) used Milky Way (MW) field RR Lyrae stars with parallaxes from *Gaia* data release 2 (Gaia Collaboration et al. 2016, 2018) to derive PLZ relations in NIR bands. These authors found a metallicity coefficient which is consistent with theoretical predictions but with a four times larger scatter. Bhardwaj et al. (2021b) showed that the lack of homogeneous NIR photometry and spectroscopic metallicities limits the precision of PLZ relation for field RR Lyrae stars in the MW despite the improvements in their parallax measurements with the early *Gaia* data release 3 (Gaia Collaboration et al. 2021). The latest *Gaia* parallaxes of MW field RR Lyrae stars have also been used to calibrate PLZ relations at mid-infrared wavelengths based on Wide-field Infrared Survey Explorer data (e.g., Gilligan et al. 2021; Mullen et al. 2023).

RR Lyrae variables in globular clusters (GCs) of different mean metallicities provide a unique opportunity to quantify the metallicity coefficient of PLZ relations. A previous such study was carried out by Sollima et al. (2006) in K_s -band using literature photometry of 538 RR Lyrae stars in 16 GCs with $\Delta[\text{Fe}/\text{H}] = 1.3$ dex, which resulted in a weak constraint on the metallicity coefficient (0.08 ± 0.11 mag/dex). This work aims to use homogeneous JHK_s time-series photometry of RR Lyrae variables in GCs from our ongoing observational program to empirically calibrate their PLZ relations and extinction-free period-Wesenheit-metallicity (PWZ) relations. The empirical calibrations of RR Lyrae PLZ




Figure 1. Histograms of period distributions (top) of 964 GC RR Lyrae variables and the distribution of mean metallicities (bottom) for their parent GCs. The peak and the width of the metallicity distribution represent the mean metallicity and its scatter from Carretta et al. (2009), respectively.

and PWZ relations at NIR wavelengths will be crucial for stellar astrophysics and distance scale studies based on population II standard candles. The well-established infrared PLZ relations for RR Lyrae stars will be particularly important for the James Webb Space Telescope, upcoming Nancy Grace Roman Space Telescope, and ground-based thirty meter class telescopes, which will operate predominantly at infrared wavelengths and enable observations of a large number of RR Lyrae stars in galaxies beyond the Local Group.

2. THE DATA

We use NIR photometry of RR Lyrae variables in GCs from our ongoing observational programs using the WIRCam instrument at the Canada-France-Hawaii Telescope (CFHT) and the FLAMINGOS-2 instrument at the Gemini South telescope¹. All the photometric data for GCs have been calibrated to Two Micron All Sky Survey (2MASS, Skrutskie et al. 2006) photometric system. Time-series photometry has been published for NGC 5272, NGC 5024, NGC 7078, and NGC 6441 (Bhardwaj et al. 2020, 2021b,a, 2022), and will be presented for NGC 5904, NGC 6402, NGC 6934, and NGC 7089 in a subsequent study (Bhardwaj et al. in prep.).

¹ Interested readers are referred to Bhardwaj et al. (2020) and Bhardwaj et al. (2022) for details regarding the observing strategy, the data reduction, and analysis of the data from CFHT and Gemini telescopes, respectively.

Table 1. Globular clusters with JHK_s time-series photometric data.

GC	N		Ref. ^a	[Fe/H] ^b	E(B-V) ^c	μ_{GC}^d	μ_{TW}^e	
	RRab	RRc						
NGC 6441	-	21	9	B22	-0.44 ± 0.07	0.47	15.52 ± 0.03	15.48 ± 0.02
NGC 6121	M4	30	12	S14	-1.18 ± 0.02	0.35	11.34 ± 0.02	11.29 ± 0.02
NGC 5904	M5	67	32	B23	-1.33 ± 0.02	0.03	14.37 ± 0.02	14.30 ± 0.02
NGC 6402	M14	47	47	B23	-1.39 ± 0.09	0.60	14.80 ± 0.06	14.75 ± 0.02
NGC 5272	M3	171	57	B20	-1.50 ± 0.05	0.01	15.04 ± 0.02	15.02 ± 0.02
NGC 6934	-	63	12	B23	-1.56 ± 0.09	0.10	15.98 ± 0.02	15.95 ± 0.02
NGC 5139	ω Cen	83	97	B18	-1.64 ± 0.09	0.12	13.67 ± 0.02	13.67 ± 0.02
NGC 7089	M2	22	15	B23	-1.66 ± 0.07	0.06	15.34 ± 0.02	15.31 ± 0.03
NGC 5024	M53	28	31	B21a	-2.06 ± 0.09	0.02	16.34 ± 0.02	16.35 ± 0.03
NGC 7078	M15	48	61	B21b	-2.33 ± 0.02	0.10	15.15 ± 0.02	15.14 ± 0.03
NGC 6341	M92	8	3	D05	-2.35 ± 0.05	0.02	14.65 ± 0.02	14.65 ± 0.04

NOTE—^aReferences: B22 - Bhardwaj et al. (2022), S14 - Stetson et al. (2014), B20 - Bhardwaj et al. (2020), B18 - Braga et al. (2018), B21a - Bhardwaj et al. (2021a), B21b - Bhardwaj et al. (2021b), D05 - Del Principe et al. (2005), B23 - Bhardwaj et al. (in prep.).

^bThe mean metallicities were taken from Carretta et al. (2009).

^cThe reddening values were adopted from Harris (2010) with an average uncertainty of 0.02 mag, which represents the scatter in $E(B - V)$ values from Harris (2010) and Green et al. (2019).

^dThe homogeneous distance moduli (μ_{GC}) are from the catalog of Baumgardt & Vasiliev (2021).

^eThe average distance moduli to GCs derived in this work (μ_{TW}) using NIR PLZ and PWZ relations in the form of equation (1) for the whole sample of 1310 RR Lyrae stars.

We also included homogeneous JHK_s mean magnitudes for RR Lyrae variables in NGC 6341 (Del Principe et al. 2005), NGC 6121 (Stetson et al. 2014), and NGC 5139 (Braga et al. 2018). Our sample consists of 964 stars including 588 fundamental mode (RRab), 345 first-overtone (RRc), and 31 mixed-mode RR Lyrae (RRd) variables. The top panel of Fig. 1 shows period distributions of 964 GC RR Lyrae stars and the bottom panel shows the distribution of mean metallicities for their parent GCs. Table 1 lists the GCs which host RR Lyrae with JHK_s time series photometric data together with their homogeneous distances, mean metallicities, and reddening values.

To calibrate the absolute zero-point of NIR PLZ relations, we also included MW field RR Lyrae stars with *Gaia* data and JHK_s photometry from 2MASS. The initial sample of ~ 400 stars with positions, metallicities, and reddening values from the literature was taken from Muraveva et al. (2018). This sample was cross-matched, within a $1''$ search radius, with the 2MASS catalog and the *Gaia* distances and RR Lyrae variability catalogs (Bailer-Jones et al. 2021; Clementini et al. 2022). The combined sample of 382 stars was further restricted to have metallicity range ($-0.3 < [\text{Fe}/\text{H}] < -2.5$ dex) similar to that of cluster variables and have 2MASS quality flag ‘A’ in JHK_s filters (346 stars). We did not apply random-phase corrections to single-epoch 2MASS magnitudes because no proper quantification of amplitude




Figure 2. Near-infrared PLRs for MW field RR Lyrae variables in J (top), H (middle) and K_s (bottom) bands. The solid black and dashed red lines represent theoretical PLZ relations for $[\text{Fe}/\text{H}] = -0.4$ dex and -2.4 dex, respectively. The representative error bars on the left display $\pm 3\sigma$ scatter in the theoretical relations (Marconi et al. 2015).

ratios or phase lags is available between G and JHK_s bands, and these observations are not contemporaneous. A comparison of intensity-averaged magnitudes of 342 RR Lyrae stars in GCs with single-epoch 2MASS

magnitudes (with quality flag ‘A’ in JHK_s) results in a median offset of 0.026/0.015/0.051 mag and a scatter of 0.13/0.11/0.12 mag in $J/H/K_s$. These systematic offsets were propagated to the photometric uncertainties on 2MASS magnitudes of MW field stars.

Fig. 2 displays JHK_s PLRs for MW field RR Lyrae stars using the final sample of 346 field variables (321 RRab and 25 RRc stars). The 2MASS JHK_s magnitudes were corrected for reddening assuming the reddening law of Cardelli et al. (1989) with an $R_V = 3.23$, and the total-to-selective absorption ratios in NIR filters as: $A_{J/H/K_s} = 0.94/0.58/0.39E(B - V)$. The periods and classification of RR Lyrae variables from the *Gaia* catalog (Clementini et al. 2022), literature reddening estimates (Muraveva et al. 2018) and the *Gaia* based distances (Bailer-Jones et al. 2021) were used to obtain absolute magnitudes for individual stars. A large scatter in the PLZ relations (0.20/0.17/0.16 mag in $J/H/K_s$) for MW field stars is seen in Fig. 2 as compared to the theoretical relations (0.06/0.04/0.04 mag in $J/H/K_s$, Marconi et al. 2015), which is not only due to random-phase magnitudes but also due to parallax uncertainties, heterogeneous metallicity and reddening estimates. Nevertheless, the slopes of NIR PLRs for RR Lyrae variables in GCs are well constrained (Bhardwaj 2022), which will allow an accurate determination of absolute zero-points of PLZ and PWZ relations at JHK_s wavelengths using MW field stars. This analysis will use the combined sample of 1310 RR Lyrae stars from GCs (964 sources) and MW field (346 sources).

3. LINEAR REGRESSION FORMALISM

Our goal is to simultaneously solve for a common slope and a metallicity coefficient, and a separate zero-point for each GC. This approach does not require a prior assumption of an independent distance to a given GC, which can be obtained simultaneously given an absolute zero-point based on field RR Lyrae stars. In the mathematical form, the extinction-corrected apparent magnitude or the Wesenheit magnitude of the j^{th} RR Lyrae star in i^{th} GC at wavelength λ is defined as:

$$m_{\lambda,i,j} = \mu_i + a_\lambda + b_\lambda \log(P_{i,j}) + c_\lambda [\text{Fe}/\text{H}]_{i,j}, \quad (1)$$

where, $P_{i,j}$ is the period in days and $[\text{Fe}/\text{H}]_{i,j}$ is the metallicity of RR Lyrae stars. For a given cluster, the mean metallicity from Carretta et al. (2009), listed in Table 1, was adopted for all RR Lyrae members. The term $(\mu_i + a_\lambda)$ also gives the apparent zero-point of the PLZ/PWZ relation in a given cluster. In the case of field variables, the absolute magnitude of a given RR Lyrae in the above equation is:

$$M_{\lambda_j} = a_\lambda + b_\lambda \log(P_j) + c_\lambda [\text{Fe}/\text{H}]_j. \quad (2)$$

To solve equations (1) and (2) simultaneously, we setup a matrix formalism where vector Y includes all the magnitude measurements. The set of equations are given by a matrix L of n_Y rows and n_Q columns. Here, n_Y is the total number of magnitude measurements in Y and n_Q is the number of free parameters (11 GC distances, zero-point, slope, and metallicity coefficient) in the vector Q . The χ^2 statistics is given as $\chi^2 = (Y - LQ)^T C^{-1} (Y - LQ)$, where C is the error matrix. The maximum-likelihood parameters are obtained using $Q = (L^T C^{-1} L)^{-1} L^T C^{-1} Y$, while the standard errors on the free parameters come from the covariance matrix $(L^T C^{-1} L)^{-1}$. This setup of matrix equations is similar to those described in detail in Riess et al. (2016, in the appendix). This matrix formalism is employed to obtain free parameters Q , which are used to compute residuals of the best-fitting linear-regression. This process is iterated by removing the single largest outlier in each iteration until all residuals are within 3σ , where σ is the standard deviation of the global fit. We created 10^3 random realizations by varying the magnitudes, adopted mean-metallicity and reddening values within their uncertainties to obtain final coefficients and their associated uncertainties.

4. PERIOD-LUMINOSITY-METALLICITY RELATIONS

We used JHK_s mean magnitudes for 964 RR Lyrae variables in GCs and 2MASS single epoch magnitudes for 346 MW field stars jointly to derive the PLZ and PWZ relations. Three different samples of RR Lyrae stars were considered separately: RRab, RRc+RRd, and the combined sample of all 1310 RR Lyrae variables. The mixed-mode RRd variables have the dominant first-overtone periods as seen in Fig. 1. Therefore, the first-overtone periods of RRc and RRd stars were fundamentalized using the equation: $\log(P_f) = \log(P_o) + 0.127$ (e.g., Coppola et al. 2015; Braga et al. 2022) for the combined sample of all 1310 RR Lyrae stars.

The intensity-averaged JHK_s magnitudes for variables in GCs were corrected for reddening assuming the previously mentioned reddening law of Cardelli et al. (1989) and absorption ratios, and using $E(B - V)$ values provided by Harris (2010) listed in Table 1. The extinction-free Wesenheit magnitudes (Madore 1982) in NIR filters were defined as in Bhardwaj et al. (2016): $W_{J,H} = H - 1.83(J - H)$, $W_{J,K_s} = K - 0.69(J - K_s)$, $W_{H,K_s} = K - 1.92(H - K_s)$.

We simultaneously solved for individual GC distances and derived PLZ relations for the global sample of 1310




Figure 3. Near-infrared PLZ and PWZ relations for RR Lyrae variables in GCs in JHK_s bands. In each case, the color-coded solid lines in the top panels correspond to $[\text{Fe}/\text{H}] = [-0.5, -1.5, -2.5]$ dex. The solid line in the bottom panels shows the best-fitting linear regression while the dashed line displays the predicted relation from Marconi et al. (2015). The average residuals (ΔJ , ΔH , ΔK_s) in a period bin of $\log(P) = 0.02$ days are shown for visualization purposes. The data points are color-coded according to the color bar.

RR Lyrae stars. In the case of RRc variables, we also added 52 RRc models with $0.25 < P < 0.45$ days (Marconi et al. 2015) to the calibrator sample because there are only 25 MW field RRc stars. These RRc models allowed us to break the degeneracy between the metallicity coefficient and the zero-points. Table 2 lists the results of the best-fitting linear regression formalism discussed in Section 3. Fig. 3 displays the PLZ/PWZ relations for the combined sample of RR Lyrae variables in GCs. The metallicity effect on RR Lyrae magnitudes is distinctly evident in the top panels showing that the metal poor stars are brighter. Tighter PLZ/PWZ relations are seen in the bottom panels where the metallicity term is taken into account, and the residuals do not exhibit any significant trend as a function of period or metallicity. These empirical metallicity coefficients are the most precisely determined in JHK_s filters, and are in excellent agreement with the horizontal branch evolution and pulsation model predictions. There is a hint of increasing metallicity dependence at shorter wavelengths, but the differences are not statistically significant. We did not find any statistically significant changes in the PLZ relations for different metallicity scales and reddening values as discussed in the appendix A.

Figure 3 also shows predicted PLZ/PWZ relations from Marconi et al. (2015, corresponding to $[\text{Fe}/\text{H}] = -0.5, -1.5$ and -2.5 dex) which suggest a good agreement with our empirical relations. The last three columns of Table 2 list the differences in the slopes, zero-points, and the metallicity coefficients between theoretical (Marconi et al. 2015) and empirical PLZ/PWZ relations derived in this work. While the metallicity coefficients are very similar, there are some differences in the slopes and zero-points for some specific sample and/or filter combinations. In particular, the slope and zero-point of RRc stars in J -band are significantly shallower and fainter than the model predictions, respectively. We note that the error in the predicted slope for RRc stars in J -band is 0.08 mag (Marconi et al. 2015), and therefore, the difference of -0.32 mag in the slopes between predicted and empirical relations is just beyond 3σ of their combined uncertainties. Similarly, the largest difference in zero-point (-0.18 ± 0.06 mag) also occurs for RRc stars in the J -band, but it is within ± 0.05 mag for most sample and/or filter combinations.

4.1. Distance moduli to globular clusters

Table 1 also lists the average distance moduli to the GCs derived after solving equations (1) and (2), for PLZ and PWZ relations for the sample of all 1310 RR Lyrae stars. Fig. 4 in the appendix B shows no obvious

trend in the difference between the distance moduli derived in this work and those from Baumgardt & Vasiliev (2021) as a function of metallicity. While the residual offset in average distance moduli is statistically small for all RR Lyrae sample (-0.026 ± 0.025 mag), the RR Lyrae distances to GCs are systematically smaller using empirical calibrations for all/RRab sample and larger using semi-empirical calibrations for RRc+RRd sample (see $\Delta\mu$ in Table 2). Bhardwaj et al. (2021b) also found a similar result with absolute zero-points of $-0.76/ -0.84$ mag in K_s using empirical/theoretical calibrations. The empirical calibrations use *Gaia* distances taking into account the parallax zero-point offset from Lindegren et al. (2021). A fainter absolute zero-point of empirical PLZ/PWZ relation, which results in smaller distance moduli, suggests an over-correction in parallaxes because a smaller parallax zero-point offset than the Lindegren et al. (2021) correction will result in a brighter zero-point for RR Lyrae stars (see Bhardwaj et al. 2021b). A large scatter in the PLZ relations for MW field stars (Fig. 2) precludes us from quantifying this small offset in parallaxes in the latest *Gaia* data. Comparison of distance moduli listed in Table 1 also suggests that the difference is larger for metal-rich GCs while the metal-poor GCs show excellent agreement. This difference can be attributed to a small number of both RR Lyrae stars and GCs for $[\text{Fe}/\text{H}] < -1.2$ dex (see Fig. 1), which could affect the accuracy of the apparent zero-points of PLZ relations, and in turn distance moduli, for the relatively metal-rich GCs.

4.2. Validation of empirical period-luminosity-metallicity relations

To validate the accuracy of our empirically derived PLZ and PWZ relations, we determined a distance to the LMC using available NIR photometry in the literature. We use ~ 18500 RR Lyrae stars (13611 RRab, 4848 RRc) from Cusano et al. (2021) with JK_s photometry from the VMC survey together with their photometric metallicity and reddening estimates. Our empirical calibrations result in an average distance modulus to the LMC of 18.54 ± 0.13 , 18.57 ± 0.12 , and 18.60 ± 0.15 mag in K_s band using the whole sample, RRab, and RRc+RRd stars, respectively. The quoted values and their uncertainties represent the peak and the width of the Gaussian distribution taking into account the uncertainties on the metallicity scale and on individual metallicities. When using theoretical relations of Marconi et al. (2015), we obtained a distance of 18.60 ± 0.13 , and 18.59 ± 0.15 mag for all/RRab and RRc+RRd samples, respectively, which reflects the dif-

Table 2. NIR period-luminosity-metallicity and period-Wesenheit-metallicity relations.

Band	Type	a	b	c	σ	N_f	χ_{dof}^2	$\Delta\mu$	$\Delta\mathbf{a}$	$\Delta\mathbf{b}$	$\Delta\mathbf{c}$
<i>J</i>	RRL	-0.44 ± 0.03	-1.83 ± 0.02	0.20 ± 0.02	0.05	1049	1.03	0.02 ± 0.05	-0.06	-0.07	-0.02
<i>H</i>	RRL	-0.74 ± 0.02	-2.29 ± 0.02	0.19 ± 0.01	0.05	1037	0.98	-0.01 ± 0.04	-0.02	0.06	-0.01
<i>K_s</i>	RRL	-0.80 ± 0.02	-2.37 ± 0.02	0.18 ± 0.01	0.05	1077	1.02	-0.01 ± 0.03	-0.02	0.12	-0.00
<i>W_{J,H}</i>	RRL	-1.20 ± 0.02	-2.96 ± 0.02	0.18 ± 0.01	0.07	989	1.07	-0.05 ± 0.04	0.01	0.20	0.02
<i>W_{J,K_s}</i>	RRL	-1.06 ± 0.03	-2.73 ± 0.02	0.16 ± 0.02	0.06	1096	0.99	-0.04 ± 0.03	0.01	0.23	0.02
<i>W_{H,K_s}</i>	RRL	-0.93 ± 0.02	-2.45 ± 0.02	0.16 ± 0.01	0.07	1033	1.17	-0.01 ± 0.04	-0.01	0.13	0.02
<i>J</i>	RRab	-0.46 ± 0.02	-1.97 ± 0.04	0.20 ± 0.01	0.05	651	0.83	0.03 ± 0.05	-0.05	-0.01	0.01
<i>H</i>	RRab	-0.74 ± 0.02	-2.28 ± 0.03	0.19 ± 0.01	0.05	667	0.88	-0.01 ± 0.04	-0.02	0.04	-0.00
<i>K_s</i>	RRab	-0.81 ± 0.03	-2.39 ± 0.03	0.18 ± 0.02	0.05	709	0.99	-0.02 ± 0.04	-0.01	0.12	0.00
<i>W_{J,H}</i>	RRab	-1.19 ± 0.03	-2.91 ± 0.04	0.18 ± 0.02	0.07	632	0.88	-0.06 ± 0.04	0.02	0.24	0.02
<i>W_{J,K_s}</i>	RRab	-1.07 ± 0.03	-2.69 ± 0.04	0.16 ± 0.02	0.06	714	0.84	-0.05 ± 0.05	0.04	0.20	0.03
<i>W_{H,K_s}</i>	RRab	-0.93 ± 0.03	-2.41 ± 0.04	0.16 ± 0.01	0.07	680	0.99	0.00 ± 0.06	0.01	0.07	0.01
<i>J</i>	RRc+RRd	-0.89 ± 0.04	-2.14 ± 0.05	0.17 ± 0.03	0.05	437	1.02	0.03 ± 0.04	-0.18	-0.32	-0.02
<i>H</i>	RRc+RRd	-1.23 ± 0.04	-2.51 ± 0.05	0.17 ± 0.03	0.05	428	0.93	0.04 ± 0.02	-0.08	-0.19	-0.01
<i>K_s</i>	RRc+RRd	-1.30 ± 0.04	-2.60 ± 0.05	0.17 ± 0.03	0.05	435	1.03	0.04 ± 0.03	-0.08	-0.12	-0.02
<i>W_{J,H}</i>	RRc+RRd	-1.72 ± 0.04	-2.92 ± 0.06	0.18 ± 0.03	0.06	392	1.08	0.09 ± 0.05	0.00	-0.16	-0.01
<i>W_{J,K_s}</i>	RRc+RRd	-1.55 ± 0.04	-2.88 ± 0.05	0.18 ± 0.03	0.05	422	1.07	0.06 ± 0.03	-0.03	-0.02	-0.02
<i>W_{H,K_s}</i>	RRc+RRd	-1.37 ± 0.04	-2.71 ± 0.05	0.18 ± 0.03	0.06	410	1.22	0.07 ± 0.04	-0.11	-0.04	-0.03

NOTE—The zero-point (*a*), slope (*b*), metallicity coefficient (*c*), dispersion (σ), number of stars (N_f), and the chi-square per degree of freedom (χ_{dof}^2) of the final PLR fits are tabulated. $\Delta\mu$ represents the mean and the standard deviation of residuals between distance moduli from this work and those from Baumgardt & Vasiliev (2021). The last three columns show the difference in the zero-points, the slopes, and the metallicity coefficients between predicted (Marconi et al. 2015) and empirical relations derived in this study. In the case of RRc+RRd stars only, the calibrator sample of MW field stars also includes 52 models from Marconi et al. (2015) as mentioned in the text.

ference in the slopes and zero-points between empirical and theoretical calibrations. The difference in the LMC distance modulus with the percent-level precise value of 18.477 ± 0.026 mag (Pietrzyński et al. 2019) is likely due to the differences in the calibrator and target PLZ relations, and due to uncertainties in photometric metallicities and reddening, and geometrical effects, which contribute to three times larger scatter in the PLZ relations for RR Lyrae stars in the LMC.

5. CONCLUSIONS

We presented the results of a NIR monitoring program of GCs to quantify the metallicity dependence on RR Lyrae PLZ and PWZ relations. The largest sample of 964 cluster RR Lyrae stars with homogeneous JHK_s mean magnitudes based on time-series data in 11 GCs of different mean metallicities was used in this analysis. When anchored using additional 346 MW field RR Lyrae stars with *Gaia* parallaxes, we obtained purely empirical PLZ and PWZ relations that agree well with the theoretical predictions based on horizontal branch evolution and pulsation models. The metallicity coefficient of NIR PLZ and PWZ relations is ~ 0.2 mag/dex, in excellent agreement with the predicted relations. We simultaneously obtained distance moduli to individual

GCs and found those to be consistent with most recent and independent measurements in the literature based on *Gaia* data. While the metallicity coefficients and the slopes are firmly constrained, the absolute zero-points of PLZ/PWZ relations are sensitive, at the percent-level precision, to the uncertainties in parallaxes, parallax zero-point offset, lack of homogeneous time-series photometry and spectroscopic metallicities for MW field stars. Furthermore, any possible variations in the slopes of the underlying PLZ/PWZ relations in different stellar environments may also affect the zero-point, and this basic assumption of universality of the PLRs will be investigated in detail in a future study. Nevertheless, our accurate and precise empirical calibrations of PLZ/PWZ relations will be crucial for RR Lyrae based distance measurements and the calibration of the distance scale based on population II standard candles in the era of space telescopes and next-generation ground-based facilities operating mainly at infrared wavelengths.

ACKNOWLEDGMENTS

We thank the anonymous referee for the constructive referee report that helped improve the manuscript. This project has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 886298. This research was supported in part by the Australian Research Council Centre of Excellence for All Sky Astrophysics in 3 Dimensions (ASTRO 3D), through project number CE170100013. HPS acknowledges grant 03(1428)/18/EMR-II from Council of Scientific and Industrial Research (CSIR), India. CCN thanks the funding from the National Science and Technology Council (Taiwan) under the contract 109-2112-M-008-014-MY3. Access to the CFHT was made possible by the Institute of Astronomy and Astrophysics, Academia Sinica.

Facility: CFHT-WIRCam, Gemini FLAMINGOS-2

Software: IDL (Landsman 1993), Astropy (Astropy Collaboration et al. 2013)

APPENDIX

A. IMPACT OF DIFFERENT METALLICITY SCALES AND REDDENING VALUES

We tested for possible variations in the PLZ relations due to different metallicity scales and reddening values in the literature using the full sample of 1310 RR Lyrae stars. Our calibrator sample of 346 RR Lyrae stars was adopted from Muraveva et al. (2018) which was based on the homogenized compilation of spectroscopic and photometric metallicities by Dambis et al. (2013) in Zinn & West (1984) metallicity scale. Crestani et al. (2021) published high resolution spectroscopic metallicities for 208 MW field RR Lyrae stars. There are 166 RR Lyrae stars in the Crestani et al. (2021) sample after restricting that to the same metallicity range and 2MASS quality flag criteria as in Section 2. The coefficients of the PLZ relations derived using these 166 stars together with 964 GCs variables are statistically consistent with those listed in Table 2 within 1σ of their quoted uncertainties. This was expected considering a small median offset of $\Delta[\text{Fe}/\text{H}] = 0.09 \pm 0.02$ dex for 133 common stars in the two calibrator samples of MW field stars.

In the case of mean metallicities of GCs, the two sets of independent estimates were obtained from the catalogs of Harris (2010) and Dias et al. (2016). The mean difference in the $[\text{Fe}/\text{H}]$ values of 11 GCs is 0.03 dex between the estimates from Harris (2010) and Carretta et al. (2009). This difference in the $[\text{Fe}/\text{H}]$ values is 0.08 dex between the measurements of Dias et al. (2016) and Carretta et al. (2009). Given these small mean differences with our adopted metallicities, the coefficients of PLZ relations were found to be in agreement with the values listed in Table 2 within 0.5σ of their uncertainties, when using the aforementioned two independent sets of mean $[\text{Fe}/\text{H}]$ values.

To investigate the impact of the adopted extinction law and reddening values, we obtained $E(B - V)$ values from the Baystar19 3D dust map (Green et al. 2019). The Baystar19 map does not cover NGC 6441 or NGC 5139, for which $E(B - V)$ values were adopted from Schlegel et al. (1998). The $E(B - V)$ values from these maps are in the same units and the corresponding absorption ratios in JHK_s were taken from Green et al. (2019, Table 1). After applying these independent reddening corrections, we derive JHK_s band PLZ relations which exhibit the same slope as listed in Table 2. The metallicity coefficients increase by 0.01 dex in all three NIR filters and the zero-points are fainter by 0.04/0.02/0.02 mag in $J/H/K_s$ bands. These differences in the J and HK_s band zero-points are within 1.4σ and 1σ of their associated uncertainties, respectively.

B. DIFFERENCE IN THE DISTANCE MODULI TO GLOBULAR CLUSTERS

Fig. 4 displays the difference in the distance moduli listed in the last two columns of Table 1.




Figure 4. The difference between the distance moduli (μ_{TW}) resulting from equation (1) and those from Baumgardt & Vasiliev (2021, μ_{GC}), as listed in Table 1. The mean and the standard deviation of residuals are quoted at the bottom left. The dashed and dotted lines represent the mean and three times the standard deviation of the residuals, respectively.

REFERENCES

- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33, doi: [10.1051/0004-6361/201322068](https://doi.org/10.1051/0004-6361/201322068)
- Bailer-Jones, C. A. L., Rybizki, J., Fouesneau, M., Demleitner, M., & Andrae, R. 2021, *AJ*, 161, 147, doi: [10.3847/1538-3881/abd806](https://doi.org/10.3847/1538-3881/abd806)
- Baumgardt, H., & Vasiliev, E. 2021, *MNRAS*, 505, 5957, doi: [10.1093/mnras/stab1474](https://doi.org/10.1093/mnras/stab1474)
- Beaton, R. L., Bono, G., Braga, V. F., et al. 2018, *Space Science Reviews*, 214, 113, doi: [10.1007/s11214-018-0542-1](https://doi.org/10.1007/s11214-018-0542-1)
- Bhardwaj, A. 2020, *Journal of Astrophysics and Astronomy*, 41, 23, doi: [10.1007/s12036-020-09640-z](https://doi.org/10.1007/s12036-020-09640-z)
- . 2022, *Universe*, 8, 122, doi: [10.3390/universe8020122](https://doi.org/10.3390/universe8020122)
- Bhardwaj, A., Kanbur, S. M., Macri, L. M., et al. 2016, *AJ*, 151, 88, doi: [10.3847/0004-6256/151/4/88](https://doi.org/10.3847/0004-6256/151/4/88)
- Bhardwaj, A., Kanbur, S. M., Rejkuba, M., et al. 2022, *A&A*, 668, A59, doi: [10.1051/0004-6361/202244728](https://doi.org/10.1051/0004-6361/202244728)
- Bhardwaj, A., Rejkuba, M., de Grijs, R., et al. 2020, *AJ*, 160, 220, doi: [10.3847/1538-3881/abb3f9](https://doi.org/10.3847/1538-3881/abb3f9)
- Bhardwaj, A., Rejkuba, M., Sloan, G. C., Marconi, M., & Yang, S.-C. 2021a, *ApJ*, 922, 20, doi: [10.3847/1538-4357/ac214d](https://doi.org/10.3847/1538-4357/ac214d)
- Bhardwaj, A., Rejkuba, M., de Grijs, R., et al. 2021b, *ApJ*, 909, 200, doi: [10.3847/1538-4357/abdf48](https://doi.org/10.3847/1538-4357/abdf48)
- Bono, G., Dall’Ora, M., Caputo, F., et al. 2011, in *RR Lyrae Stars, Metal-Poor Stars, and the Galaxy*, ed. A. McWilliam, Vol. 5, 1. <https://arxiv.org/abs/1108.5372>
- Bono, G., Gieren, W. P., Marconi, M., Fouqué, P., & Caputo, F. 2001, *ApJ*, 563, 319, doi: [10.1086/323834](https://doi.org/10.1086/323834)
- Braga, V. F., Stetson, P. B., Bono, G., et al. 2018, *AJ*, 155, 137, doi: [10.3847/1538-3881/aaadab](https://doi.org/10.3847/1538-3881/aaadab)
- Braga, V. F., Fiorentino, G., Bono, G., et al. 2022, *MNRAS*, 517, 5368, doi: [10.1093/mnras/stac2813](https://doi.org/10.1093/mnras/stac2813)
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245, doi: [10.1086/167900](https://doi.org/10.1086/167900)
- Carretta, E., Bragaglia, A., Gratton, R., D’Orazi, V., & Lucatello, S. 2009, *A&A*, 508, 695, doi: [10.1051/0004-6361/200913003](https://doi.org/10.1051/0004-6361/200913003)
- Catelan, M., Pritzl, B. J., & Smith, H. A. 2004, *ApJS*, 154, 633, doi: [10.1086/422916](https://doi.org/10.1086/422916)
- Catelan, M., & Smith, H. A. 2015, *Pulsating Stars* (Wiley-VCH)
- Clementini, G., Ripepi, V., Garofalo, A., et al. 2022, *A&A*, arXiv:2206.06278, doi: [10.1051/0004-6361/202243964](https://doi.org/10.1051/0004-6361/202243964)
- Coppola, G., Marconi, M., Stetson, P. B., et al. 2015, *ApJ*, 814, 71, doi: [10.1088/0004-637X/814/1/71](https://doi.org/10.1088/0004-637X/814/1/71)
- Crestani, J., Braga, V. F., Fabrizio, M., et al. 2021, *ApJ*, 914, 10, doi: [10.3847/1538-4357/abfa2310.48550/arXiv.2104.08113](https://doi.org/10.3847/1538-4357/abfa2310.48550/arXiv.2104.08113)
- Cusano, F., Moretti, M. I., Clementini, G., et al. 2021, *MNRAS*, 504, 1, doi: [10.1093/mnras/stab901](https://doi.org/10.1093/mnras/stab901)
- Dambis, A. K., Berdnikov, L. N., Kniazev, A. Y., et al. 2013, *MNRAS*, 435, 3206, doi: [10.1093/mnras/stt1514](https://doi.org/10.1093/mnras/stt1514)
- Del Principe, M., Piersimoni, A. M., Bono, G., et al. 2005, *AJ*, 129, 2714, doi: [10.1086/430148](https://doi.org/10.1086/430148)
- Dias, B., Barbuy, B., Saviane, I., et al. 2016, *A&A*, 590, A9, doi: [10.1051/0004-6361/201526765](https://doi.org/10.1051/0004-6361/201526765)
- Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016, *A&A*, 595, A1, doi: [10.1051/0004-6361/201629272](https://doi.org/10.1051/0004-6361/201629272)
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, *A&A*, 616, A1, doi: [10.1051/0004-6361/201833051](https://doi.org/10.1051/0004-6361/201833051)
- . 2021, *A&A*, 649, A1, doi: [10.1051/0004-6361/202039657](https://doi.org/10.1051/0004-6361/202039657)
- Gilligan, C. K., Chaboyer, B., Marengo, M., et al. 2021, *MNRAS*, 503, 4719, doi: [10.1093/mnras/stab857](https://doi.org/10.1093/mnras/stab857)
- Green, G. M., Schlafly, E., Zucker, C., Speagle, J. S., & Finkbeiner, D. 2019, *ApJ*, 887, 93, doi: [10.3847/1538-4357/ab5362](https://doi.org/10.3847/1538-4357/ab5362)

- Harris, W. E. 2010, arXiv:1012.3224, arXiv:1012.3224.
<https://arxiv.org/abs/1012.3224>
- Landsman, W. B. 1993, in ASP Conference Series, Vol. 52, Astronomical Data Analysis Software and Systems II, ed. R. J. Hanisch, R. J. V. Brissenden, & J. Barnes, 246
- Layden, A. C., Tiede, G. P., Chaboyer, B., Bunner, C., & Smitka, M. T. 2019, AJ, 158, 105, doi: [10.3847/1538-3881/ab2e10](https://doi.org/10.3847/1538-3881/ab2e10)
- Lindgren, L., Klioner, S. A., Hernández, J., et al. 2021, A&A, 649, A2, doi: [10.1051/0004-6361/202039709](https://doi.org/10.1051/0004-6361/202039709)
- Longmore, A. J., Fernley, J. A., & Jameson, R. F. 1986, MNRAS, 220, 279, doi: [10.1093/mnras/220.2.279](https://doi.org/10.1093/mnras/220.2.279)
- Madore, B. F. 1982, ApJ, 253, 575, doi: [10.1086/159659](https://doi.org/10.1086/159659)
- Marconi, M., Coppola, G., Bono, G., et al. 2015, ApJ, 808, 50, doi: [10.1088/0004-637X/808/1/50](https://doi.org/10.1088/0004-637X/808/1/50)
- Muhie, T. D., Dambis, A. K., Berdnikov, L. N., Kniazev, A. Y., & Grebel, E. K. 2021, MNRAS, 502, 4074, doi: [10.1093/mnras/stab219](https://doi.org/10.1093/mnras/stab219)
- Mullen, J. P., Marengo, M., Martínez-Vázquez, C. E., et al. 2023, arXiv e-prints, arXiv:2301.03777.
<https://arxiv.org/abs/2301.03777>
- Muraveva, T., Delgado, H. E., Clementini, G., Sarro, L. M., & Garofalo, A. 2018, MNRAS, 481, 1195, doi: [10.1093/mnras/sty2241](https://doi.org/10.1093/mnras/sty2241)
- Muraveva, T., Palmer, M., Clementini, G., et al. 2015, ApJ, 807, 127, doi: [10.1088/0004-637X/807/2/127](https://doi.org/10.1088/0004-637X/807/2/127)
- Neeley, J. R., Marengo, M., Freedman, W. L., et al. 2019, MNRAS, 490, 4254, doi: [10.1093/mnras/stz2814](https://doi.org/10.1093/mnras/stz2814)
- Pietrzyński, G., Graczyk, D., Gallenne, A., et al. 2019, Nature, 567, 200, doi: [10.1038/s41586-019-0999-4](https://doi.org/10.1038/s41586-019-0999-4)
- Riess, A. G., Macri, L. M., Hoffmann, S. L., et al. 2016, ApJ, 826, 56, doi: [10.3847/0004-637X/826/1/56](https://doi.org/10.3847/0004-637X/826/1/56)
- Sandage, A., & Tammann, G. A. 2006, ARA&A, 44, 93, doi: [10.1146/annurev.astro.43.072103.150612](https://doi.org/10.1146/annurev.astro.43.072103.150612)
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525, doi: [10.1086/305772](https://doi.org/10.1086/305772)
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, AJ, 131, 1163, doi: [10.1086/498708](https://doi.org/10.1086/498708)
- Sollima, A., Cacciari, C., Arkharov, A. A. H., et al. 2008, MNRAS, 384, 1583, doi: [10.1111/j.1365-2966.2007.12804.x](https://doi.org/10.1111/j.1365-2966.2007.12804.x)
- Sollima, A., Cacciari, C., & Valenti, E. 2006, MNRAS, 372, 1675, doi: [10.1111/j.1365-2966.2006.10962.x](https://doi.org/10.1111/j.1365-2966.2006.10962.x)
- Stetson, P. B., Braga, V. F., Dall’Ora, M., et al. 2014, PASP, 126, 521, doi: [10.1086/677195](https://doi.org/10.1086/677195)
- Zinn, R., & West, M. J. 1984, ApJs, 55, 45, doi: [10.1086/190947](https://doi.org/10.1086/190947)